

Construction of the automatic Carcinologic Speech Severity Index (C2SI) score



Mathieu Balaguer^(1,2), Jérôme Farinas⁽¹⁾, Julien Pinquier⁽¹⁾, Virginie Woisard^(2,3)


(1) IRIT, CNRS, Université Paul Sabatier Toulouse III, Toulouse, France

(2) CHU Larrey, Toulouse, France

(3) Laboratoire Octogone-Lordat, Toulouse, France



Institut de Recherche
en Informatique de Toulouse
CNRS - INP - UT3 - UT1 - UT2J




Context

Upper aerodigestive tract cancer:

14,700 new cases every year in France

massive impact on communication and on the quality of life



Fex, S. (1992). Perceptual evaluation. *Journal of Voice*, 6(2), 155–158.

Ghio, A., Giusti, L., Blanc, Pinto, S., Lalain, M., Robert, D., ... Orl, S. (2016). Quels tests d'intelligibilité pour évaluer les troubles de production de la parole ?, 4(1).

Middag, C. (2013). Automatische analyse van pathologische spraak Automatic Analysis of Pathological Speech.

Borggreven, P. A., Verdonck-De Leeuw, I. M., Muller, M. J., Heiligers, M. L. C. H., De Bree, R., Aaronson, N. K., & Leemans, C. R. (2007). Quality of life and functional status in patients with cancer of the oral cavity and oropharynx: Pretreatment values of a prospective study. *European Archives of Oto-Rhino-Laryngology*, 264(6), 651–657. <https://doi.org/10.1007/s00405-007-0249-5>


Objectives

Evaluate the validity
of different automatic speech disorder severity measures
in patients treated for upper aerodigestive tract cancer,
in order to build an automated global score

Materials and Methods

Database from the C2SI project
(Carcinologic Speech Severity Index)

Grant 2014-135 from the Institut National pour le Cancer in 2014, "Sciences Humaines et Sociales, épidémiologie et santé publique" call, led by Pr Virginie Woisard at the University of Toulouse



Materials and Methods

Study sample

Patients:



Controls:



Inclusion criteria	<ul style="list-style-type: none">▪ Oral and/or oropharyngeal cancer (TNM: T2 to T4)▪ Treatment: surgery and/or radiotherapy and/or chemotherapy▪ At least 6 months since the end of treatment	<ul style="list-style-type: none">▪ Paired by age and gender with patient group
Non-inclusion criteria	<ul style="list-style-type: none">▪ Additional potential causes of speech disorders (e.g. stuttering)▪ Cognitive or visual impairment that might affect the tasks	

Materials and Methods

Tasks

	Perceptual assessment		Automatic measures
	Visual analog scales (VAS)	Recognition rate	
Intelligibility	Reading (RED) Picture description (DES)	Acoustic-Phonetic Decoding (APD) Prosodic tasks: <i>Modality</i> (MOD) <i>Focus</i> (FOC) <i>Syntax</i> (SYN)	Sustained [a] AAA APD RED
Comprehensibility / Severity	Reading (RED) Picture description (DES)	<i>Comprehensibility</i> (SVT)	

Miller, N. (2013). Measuring up to speech intelligibility. Int J Lang Commun Disord, 48(6), 601-612. <https://doi.org/10.1111/1460-6984.12061>

Materials and Methods

Validity and reliability of the automatic measures

Clinical validity

- Can the measures discriminate between subjects and controls?
- *Mann-Whitney U test*

Criterion validity

- Do the measures correlate with the reference score?
- *Spearman's correlation coefficients*

Reliability

- Are the automatic measures reproducible?
- *Ensured by the use of automatic measurements*



Modeling

Results

Study sample characteristics



87 patients



Median age: 65.8 years old
With surgery: 73%



42 controls

Median age: 62 years old

Population




Astésano, C., Balaguer, M., Farinas, J., Fredouille, C., Gaillard, P., Ghio, A., ... Woisard, V. (2018). Carcinologic Speech Severity Index Project: A Database of Speech Disorder Productions to Assess Quality of Life Related to Speech After Cancer. Language Resources and Evaluation Conference (LREC)

Results

Perceptual assessment:

Which reference ?



Patients	Mean	Median	Min	Max	Interquartile range	ICC	95% confidence interval
Severity (RED) / 10	6.51	7.02	1.03	10	5.03 – 8.1	0.69	0.61 – 0.77
Severity (DES) / 10	6.06	6.33	0.58	9.67	4.2 – 8	0.69	0.62 – 0.77
Intelligibility (DES) / 10	7.61	8.67	1.2	10	6.83 – 9.5	0.77	0.70 – 0.83
Intelligibility (RED) / 10	8.69	9	3.5	10	8.4 – 9.5	0.43	0.32 – 0.55

Results

Automatic measures: Clinical and Criterion Validity


	Task	Clinical validity (U-test)	Criterion validity (Spearman – Severity DES)
Voice dimension			
Fundamental frequency - interquartile range	AAA	p = 0.028	-0.29
Harmonic poorness		p = 0.009	0.05
Quality of glottal attack		p = 0.005	0.01
Jitter - median		p = 0.001	-0.14
Jitter - interquartile range		p = 0.001	-0.11
Pitch instability		p = 0.001	-0.25
Amplitude instability		p = 0.072	0.01
log-HNR - median		p = 0.432	0.23
Harmonics-to-noise ratio		p = 0.005	-0.06
Speech dimension			
Mean number of automatically computed deviation features per phoneme	APD	p < 0.001	-0.62
Distance stimulus - most probable identified logatome (Cumulative rank)		p < 0.001	-0.81
Anomaly rate		p < 0.001	-0.64
Automatic likelihood score		p < 0.001	0.68
Automatic likelihood score	RED	p < 0.001	0.85

Results

C2SI score modeling:

Task	Perceptual severity	Bivariate analysis		C2SI		
		Coef.	p-value	Coef.	95% CI	p-value
	Constant			11,26482	9,57 ; 12,96	<0,001*
AAA	Fundamental frequency – interquartile range	-0,29	0,01*	-0,0049184	-0,02 ; 0,01	0,45
	Pitch instability	-0,25	0,04*	-0,0946604	-0,29 ; 0,10	0,34
RED	Automatic likelihood score	0,85	< 0,001*	1,391981	0,84 ; 1,94	<0,001*
APD	Automatic likelihood score	0,68	< 0,001*	-0,147016	-0,77 ; 0,48	0,64
	Cumulative rank	-0,81	< 0,001*	-2,09e-06	-3,86e-06 ; -3,07e-07	0,02*
	Anomaly rate	-0,64	< 0,001*	-0,0111486	-0,05 ; 0,02	0,52
				$R^2 = 0,7565$ Adjusted R² = 0,7285		

Discussion



Discussion

C2SI score modeling:


Tasks	Perceptual severity	Bivariate analysis		C2SI FULL VERSION			C2SI REDUCED VERSION		
				Multivariate analysis			Multivariate analysis		
		Coef.	p-value	Coef.	95% CI	p-value	Coef.	95% CI	p-value
	Constant			11.26482	9.57 ; 12.96	<0.001*	11.48726	10.50 ; 12.47	<0.001*
AAA	Fundamental frequency – interquartile range	-0.29	0.01*	-0.0049184	-0.02 ; 0.01	0.45			
	Pitch instability	-0.25	0.04*	-0.0946604	-0.29 ; 0.10	0.34			
RED	Automatic likelihood score	0.85	< 0.001*	1.391981	0.84 ; 1.94	<0.001*	1.52926	1.16 ; 1.90	<0.001*
APD	Automatic likelihood score	0.68	< 0.001*	-0.147016	-0.77 ; 0.48	0.64			
	Cumulative rank	-0.81	< 0.001*	-2.09e-06	-3.86e-06 ; -3.07e-07	0.02*	-1.94e-06	-3.36e-06 ; -5.12e-07	0.008*
	Anomaly rate	-0.64	< 0.001*	-0.0111486	-0.05 ; 0.02	0.52			
				R² = 0.7565 Adjusted R² = 0.7285 Correlation with reference: 0.87			R² = 0.7618 Adjusted R² = 0.7550 Correlation with reference: 0.87		

Discussion

A measure to assess the functional impact of the speech disorder?

Speech production disorder

- Need to include automatic analysis of comprehensibility tasks
- Other signal parameters?



Functional impact on communication

- Self-assessment of speech impairment
- Communicational needs
- Associated factors: *physical and functional deficits, cognitive impairment, anxiety/depression,...*

Care adaptation
based on
the automatic score:

Speech deficit
improvement

+

Functional
improvement

Thank you for your attention

