



HAL
open science

Existence of a traveling wave solution in a free interface problem with fractional order kinetics

Claude-Michel Brauner, Robert Roussarie, Peipei Shang, Linwan Zhang

► **To cite this version:**

Claude-Michel Brauner, Robert Roussarie, Peipei Shang, Linwan Zhang. Existence of a traveling wave solution in a free interface problem with fractional order kinetics. *Journal of Differential Equations*, 2021, 281, pp.105-147. 10.1016/j.jde.2021.01.034 . hal-02979187

HAL Id: hal-02979187

<https://hal.science/hal-02979187>

Submitted on 27 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXISTENCE OF A TRAVELING WAVE SOLUTION IN A FREE INTERFACE PROBLEM WITH FRACTIONAL ORDER KINETICS

CLAUDE-MICHEL BRAUNER, ROBERT ROUSSARIE, PEIPEI SHANG, LINWAN ZHANG

ABSTRACT. In this paper we consider a system of two reaction-diffusion equations that models diffusional-thermal combustion with stepwise ignition-temperature kinetics and fractional reaction order $0 < \alpha < 1$. We turn the free interface problem into a scalar free boundary problem coupled with an integral equation. The main intermediary step is to reduce the scalar problem to the study of a non- C^1 vector field in dimension 2. The latter is treated by qualitative topological methods based on the Poincaré-Bendixson Theorem. The phase portrait is determined and the existence of a stable manifold at the origin is proved. A significant result is that the settling time to reach the origin is finite, meaning that the trailing interface is finite in contrast to the case $\alpha = 1$, but in accordance with $\alpha = 0$. Finally, the integro-differential system is solved via a fixed-point method.

1. INTRODUCTION AND MAIN RESULTS

A flame spreading through a motionless gas may be described by a system of two parabolic nonlinear equations for the normalized temperature, T , and the concentration of deficient reactant, Y . This system, in non-dimensional form, reads:

$$\begin{cases} T_t = \Delta T + W(T, Y), \\ Y_t = \Lambda \Delta Y - W(T, Y), \end{cases} \quad (1.1)$$

where Λ (the ratio of mass diffusivity and thermal diffusivity) is the inverse of the Lewis number and $W(T, Y)$ is the reaction rate.

In this paper, the reaction rate is assumed to be of stepwise ignition type, that is, $W(T, Y) \equiv 0$ when the normalized temperature T is below the ignition temperature $\theta \in (0, 1)$, and $W(T, Y)$ depends only on the concentration Y when the temperature is above θ . The recent renewed interest in stepwise ignition temperature kinetics is due to its applicability to the studies of overall effective chemical kinetics of certain reactive mixtures (see Brailovsky, Gordon, Kagan and Sivashinsky [8] and references therein).

In [8, Section 2], two classes of stepwise kinetics are considered.

(i) Zero-order stepwise temperature kinetics with reaction rate given by

$$W_0(T, Y) = \begin{cases} 1, & \text{if } T \geq \theta \text{ and } Y > 0, \\ 0, & \text{if } T < \theta \text{ and/or } Y = 0. \end{cases} \quad (1.2)$$

(ii) First-order stepwise temperature kinetics with reaction rate given by

$$W_1(T, Y) = \begin{cases} Y, & \text{if } T \geq \theta, \\ 0, & \text{if } T < \theta, \end{cases} \quad (1.3)$$

* Corresponding author: Peipei SHANG (shang@tongji.edu.cn).

or, equivalently, $W_1(T, Y) = YH(T - \theta)$ where H is the Heaviside function.

It was shown in [8] that problem (1.1) is equivalent to a free interface problem in both cases. However, the resulting free interface problems for these two cases are very different. To clarify this difference, it is convenient to consider problem (1.1) in a two-dimensional strip $\mathbb{R} \times (-\ell/2, \ell/2)$; the spatial coordinates are denoted by (x, y) , $t > 0$ is the time.

First, in the case of the zero-order kinetics (1.2), there are two free interfaces: the *ignition interface* $x = f(t, y)$, also called the flame front, defined by

$$T(t, f(t, y), y) = \theta, \quad (1.4)$$

and the *trailing interface* $x = g(t, y)$, $g(t, y) > f(t, y)$, defined by

$$Y(t, g(t, y), y) = 0. \quad (1.5)$$

Then, the system for (T, Y) reads as follows

$$\begin{cases} T_t = \Delta T, Y_t = \Lambda \Delta Y, & x < f(t, y), \\ T_t = \Delta T + 1, Y_t = \Lambda \Delta Y - 1, & f(t, y) < x < g(t, y), \\ T_t = \Delta T, Y(t, x, y) = 0, & x > g(t, y). \end{cases} \quad (1.6)$$

The functions T and Y and their normal derivatives are continuous across the ignition and trailing free interfaces. As $x \rightarrow \pm\infty$, the following holds:

$$T(t, -\infty, y) = 0, \quad Y(t, -\infty, y) = T(t, +\infty, y) = 1. \quad (1.7)$$

Second, in the case of first-order stepwise kinetics, the system reads:

$$\begin{cases} T_t = \Delta T, Y_t = \Lambda \Delta Y, & x < f(t, y), \\ T_t = \Delta T + Y, Y_t = \Lambda \Delta Y - Y, & x > f(t, y). \end{cases} \quad (1.8)$$

Therefore, only the ignition interface is involved, at which T and Y and their normal derivatives are continuous. As $x \rightarrow \pm\infty$,

$$T(t, -\infty, y) = Y(t, +\infty, y) = 0, \quad Y(t, -\infty, y) = T(t, +\infty, y) = 1. \quad (1.9)$$

Such drastic difference in qualitative properties of traveling front solution for zero- and first-order reaction models raises a natural question: why two free interfaces are generated in the case of zero-order stepwise kinetics (1.2) and just one in first-order kinetics (1.3). We aim to understand the role of the reaction rate in the process; to this end, we consider the gamut of intermediate combustion systems when the order of the reaction (which we will denote by α) is between 0 and 1. It is important to note that this question is interesting not only from mathematical viewpoint but also from perspective of applications in combustion. Indeed, as evident from experimental observation, the overall reaction order can change quite substantially with the equivalence ratio (i.e. the ratio of fuel and oxidizer in the mixture) as well as pressure. For example, in the not too extreme case of hydrogen-air flame with equivalence ratio 3, the overall reaction rate of $\alpha = 1$ at pressure 20 atm drops to $\alpha = 0.3$ at 50 atm (see [20, p. 280]).

The main goal of this paper is to understand the role of the reaction order $0 < \alpha < 1$ on qualitative properties of planar traveling fronts for free interface problems as well as their connection with the limiting systems associated with the cases $\alpha = 0$ and $\alpha = 1$. Mathematically,

we fill the gap between the zero-order and first-order kinetics. Mimicking (1.3), we define an “ α -order reaction rate” for $0 < \alpha < 1$:

$$W_\alpha(T, Y) = Y^\alpha H(T - \theta). \quad (1.10)$$

At least formally, we state that $W_\alpha(T, Y) \rightarrow W_0(T, Y)$ when α tends to 0^+ , and $W_1(T, Y)$ obviously is retrieved as $\alpha \rightarrow 1^-$.

To facilitate this study, we limit ourselves to a special class of solutions, namely, one-dimensional (planar) traveling wave solutions (u, v) of the free interface problem which travel at a constant negative velocity $-c$; $c > 0$ is to be determined. In the moving frame coordinate, $\xi = x + ct$, the ignition interface is fixed at $\xi = 0$, taking advantage of the translation invariance. The trailing front is at an unknown position, $\xi = R$, $0 < R < +\infty$.

In the case of zero-order kinetics, we recall that the one-dimensional free interface problem for $(c, R, u(\xi), v(\xi))$ in the moving frame coordinate reads:

$$\begin{cases} u_{\xi\xi} - cu_\xi = 0, & \Lambda v_{\xi\xi} - cv_\xi = 0, & \xi < 0, \\ u_{\xi\xi} - cu_\xi = -1, & \Lambda v_{\xi\xi} - cv_\xi = 1, & 0 < \xi < R, \\ u_{\xi\xi} - cu_\xi = 0, & v = 0, & \xi > R. \end{cases} \quad (1.11)$$

The functions u and v are continuously differentiable on the real line such that, according to (1.7), $u(-\infty) = 0$, $v(-\infty) = 1$ and $u(+\infty) = 1$. At the interfaces, $u(0) = \theta$ and $v(R) = 0$, respectively. It is readily seen that $u(\xi) = 1$ for $\xi \geq R$; hence, $u_\xi(R) = v_\xi(R) = 0$. In this respect, the trailing interface is a degenerate free boundary in the sense of [10] (in contrast to the ignition interface), which creates further difficulty in the stability analysis (see [9], [1]). Naturally, solving (1.11) is an elementary exercise; it happens that $c = R$ is the unique strictly positive solution of the transcendental equation

$$e^{c^2} = \frac{1}{1 - c^2\theta}, \quad 0 < \theta < 1. \quad (1.12)$$

(Note that c is independent of Λ ; see Subsection 8.2 below).

For the first-order kinetics, the system for the triplet $(c, u(\xi), v(\xi))$ reads:

$$\begin{cases} u_{\xi\xi} - cu_\xi = 0, & \Lambda v_{\xi\xi} - cv_\xi = 0, & \xi < 0, \\ u_{\xi\xi} - cu_\xi = -v, & \Lambda v_{\xi\xi} - cv_\xi = v, & \xi > 0. \end{cases} \quad (1.13)$$

The functions u and v are continuously differentiable, such that $u(-\infty) = 0$, $v(-\infty) = 1$ and $u(+\infty) = 1$, $v(+\infty) = 0$. At the ignition interface, $u(0) = \theta$. This time, the speed c is given explicitly by the formula:

$$c = \left(\frac{\theta}{1 - \theta} + \Lambda \left(\frac{\theta}{1 - \theta} \right)^2 \right)^{-\frac{1}{2}}, \quad 0 < \theta < 1, \Lambda > 0. \quad (1.14)$$

Here, there is no trailing interface R . More specifically, the trailing interface is such that $R = +\infty$ (see [6]).

In the case of a “ α -order kinetic” (1.10), the main difficulty arises from the non-Lipschitz nonlinearity W_α . Similar issues have recently been addressed in the literature, e.g. a non-Lipschitz modification of the classical Michaelis-Menten law in enzyme kinetics in the analysis of healthy immune system dynamics from the perspective of Finite-Time Stability (FTS) (see [23]). In this regard, the theory of FTS of continuous, but non-Lipschitz systems has been part of

numerous papers particularly associated with optimal control (see, e.g., [3], [22]). It is to be seen that the free interface R may also be viewed as a finite time.

In this paper, we prove the following result:

Theorem 1.1. *Let $0 < \alpha < 1$, $0 < \theta < 1$, $\Lambda > 0$ be fixed. There exist $c > 0$ and $0 < R < +\infty$ such that the free interface problem*

$$\begin{cases} T_t = \Delta T, Y_t = \Lambda \Delta Y, & x < f(t, y), \\ T_t = \Delta T + Y^\alpha, Y_t = \Lambda \Delta Y - Y^\alpha, & f(t, y) < x < g(t, y), \\ T_t = \Delta T, Y(t, x, y) = 0, & x > g(t, y), \end{cases} \quad (1.15)$$

with (1.4), (1.5) and (1.7), admits a one-dimensional traveling wave solution $(c, R, u(\xi), v(\xi))$, $\xi = x + ct$, which verifies the free interface problem

$$\begin{cases} u_{\xi\xi} - cu_\xi = 0, & \xi < 0, \\ \Lambda v_{\xi\xi} - cv_\xi = 0, & \xi < 0, \\ u_{\xi\xi} - cu_\xi = -v^\alpha, & 0 < \xi < R, \\ \Lambda v_{\xi\xi} - cv_\xi = v^\alpha, & 0 < \xi < R, \\ u_{\xi\xi} - cu_\xi = 0, & \xi > R, \\ v = 0, & \xi > R. \end{cases} \quad (1.16)$$

The functions u and v are continuously differentiable on the real line such that

$$u(-\infty) = 0, \quad v(-\infty) = 1, \quad u(+\infty) = 1. \quad (1.17)$$

At the ignition and trailing interfaces, respectively, placed at $\xi = 0$ and $\xi = R$, it holds

$$u(0) = \theta, \quad v(R) = v_\xi(R) = 0. \quad (1.18)$$

Moreover, denoting by R_α the position of the trailing interface at fixed α , we have:

- (i) $0 < R_\alpha < +\infty$, $0 < \alpha < 1$;
- (ii) as $\alpha \rightarrow 1$, $R_\alpha \rightarrow +\infty$;
- (iii) as $\alpha \rightarrow 0$, $R_\alpha \rightarrow R = c$, solution of (1.12).

As is seen, the most significant result is that R is finite for $0 \leq \alpha < 1$. As a consequence, the ‘‘cut-off’’ exponent for the existence of a finite trailing interface is $\alpha = 1$ (see [6] for a comprehensive study of the case $\alpha = 1$ and also the case of a n -order reaction rate where n is an integer greater than 1).

The paper is organized as follows:

Firstly, we point out in Section 2 that the free interface problem (1.16)-(1.18) is equivalent to the following one-phase free boundary problem:

$$\begin{cases} u_{\xi\xi} - cu_\xi = -v^\alpha, & 0 < \xi < R, \\ \Lambda v_{\xi\xi} - cv_\xi = v^\alpha, & 0 < \xi < R, \end{cases} \quad (1.19)$$

with boundary conditions at $\xi = 0$

$$u(0) = \theta, \quad u'(0) = c\theta, \quad v'(0) = -\frac{c}{\Lambda}(1 - v(0)), \quad (1.20)$$

and free boundary conditions at $\xi = R$:

$$v(R) = v'(R) = 0, \quad u(R) = 1, \quad u'(R) = 0. \quad (1.21)$$

The unknowns are: the free boundary R ; the velocity c which is a kind of eigenvalue of the problem; the functions $u(\xi)$ and $v(\xi)$ which are smooth on the interval $[0, R)$, but whose second derivatives are Hölder continuous at $\xi = R$. It is immediately apparent that the main quantity in (1.19)-(1.21) is the value $v(0)$ that we henceforth take as a parameter and denote by $v_0 \in (0, 1)$. In Section 3, we formulate a scalar free boundary problem as follows:

$$\begin{cases} \Lambda v''(\xi) - cv'(\xi) = v^\alpha(\xi), & 0 < \xi < R, \\ v(0) = v_0, v'(0) = -\frac{c}{\Lambda}(1 - v_0), \\ v(R) = v'(R) = 0, \end{cases} \quad (1.22)$$

whose study is the main feature of the paper. There are two main results for problem (1.22): Theorem 3.1 is about the existence and uniqueness of a solution $(c(v_0), R(v_0), v(v_0; \xi))$; Corollary 3.2 is about the continuous dependence upon v_0 . Their proofs, based on topological methods, occupies a large part of this paper, namely Sections 4 to 6.

The scheme is as follows: we reformulate problem (1.22) as a shooting problem

$$\begin{cases} \Lambda v''(\xi) - cv'(\xi) = v^\alpha(\xi), & \xi > 0, \\ v(0) = v_0, v'(0) = -\frac{c}{\Lambda}(1 - v_0), \end{cases} \quad (1.23)$$

which is equivalent to finding a trajectory tending towards the origin (a stable manifold at the origin) of a vector field X_c , defined in the quadrant $Q = \{x \geq 0, y \leq 0\}$, which means looking at a solution $(x(t), y(t))$ of its differential equation:

$$X_c : \begin{cases} x'(t) = y(t), \\ \Lambda y'(t) = cy(t) + x^\alpha(t), \end{cases} \quad (1.24)$$

with initial conditions

$$x(0) = v_0, \quad y(0) = -\frac{c}{\Lambda}(1 - v_0). \quad (1.25)$$

Section 4 is devoted to the topological study of (1.24) for a fixed value of c considered as a parameter, in the first place the phase portrait (see Subsection 4.1). As the vector field X_c is not C^1 and the origin is not an hyperbolic singularity when $\alpha < 1$, it is impossible to apply Hartman-Grobman Theorem; however, the result stated in Proposition 4.3 can be compared with Hartman-Grobman. The proof of Proposition 4.3 is deferred to Appendix B. In the following Subsections 4.2 to 4.4, we prove the existence and uniqueness of the stable manifold $y_c(x)$ of (1.24). We use primarily the Poincaré-Bendixson Theorem (see a brief introduction in Appendix A), which places the origin at the boundary of the domain (refer to [25] for a comprehensive study).

Section 5 is devoted to a series of technical lemmata, which enlightens the dependence of the stable manifold y_c upon parameter c . In particular, we give an asymptotic expression of y_c in Subsection 5.2, which eventually provides the optimal Hölder regularity of $v(\xi)$ (see Lemma 6.5).

In Section 6, we proceed to the proof of Theorem 3.1, per se. At fixed v_0 , we prove in Subsection 6.1 the existence of a unique $c(v_0)$, such that

$$y_{c(v_0)}(v_0) = -\frac{c(v_0)}{\Lambda}(1 - v_0).$$

In Subsection 6.3, we prove that the settling time $R(v_0)$, i.e. the time to proceed from the initial condition (1.25) to the origin on the stable manifold $y_{c(v_0)}$ is finite and, based on the asymptotic

expansion of the stable manifold, we estimate $R(v_0)$:

$$\frac{(2(1+\alpha)\Lambda)^{1/2}}{1-\alpha} v_0^{\frac{1-\alpha}{2}} < R(v_0) < \frac{2\Lambda^{1/2}A(\alpha)}{1-\alpha} \frac{v_0^{\frac{1-\alpha}{2}}}{1-v_0}. \quad (1.26)$$

Not surprisingly, this estimate explodes when $\alpha \rightarrow 1$ which is coincident with the result given in [6]. Eventually, we return to the free boundary problem (1.22) and prove Corollary 3.2.

In Section 7, we revisit the free boundary problem (1.19)-(1.21). For v_0 in some interval $I \subset (0, 1)$ to be determined, we reformulate (1.19)-(1.21) as a fixed point problem for the system (1.22), i.e.,

$$\begin{cases} \Lambda v'' - cv' = v^\alpha, & 0 < \xi < R, \\ v(0) = v_0, v'(0) = -\frac{c}{\Lambda}(1-v_0), \\ v(R) = v'(R) = 0, \end{cases} \quad (1.27)$$

coupled with an integral equation which reads:

$$\theta + \Lambda v_0 = 1 - c(1-\Lambda) \int_0^R e^{-cs} v(s) ds \quad \text{or} \quad \theta + v_0 = 1 - (1-\Lambda) \int_0^R e^{-cs} v'(s) ds. \quad (1.28)$$

The interval I depends only on Λ and θ , especially whether Λ is smaller or larger than 1 (in other words, the Lewis number is larger or smaller than unity). We prove the existence of a fixed point in Theorems 7.4 and 7.5, taking advantage of the continuity of $(c(v_0), R(v_0), v(v_0; \xi))$ w.r.t. v_0 .

The last section of the paper is devoted to a gamut of special cases. In particular, we consider the situation of a solid combustion where the Lewis number is $+\infty$, i.e. $\Lambda = 0$. The latter case is mathematically relevant, because it allows explicit computations that yield the uniqueness of the solution to the free boundary problem. Finally, we consider the limit cases $\alpha \rightarrow 1$ and $\alpha \rightarrow 0$, which complete the proof of Theorem 1.1.

As already mentioned, the paper has two appendices: Appendix A regarding the Poincaré-Bendixson Theorem and Appendix B devoted to the proof of Proposition 4.3.

2. EQUIVALENCE WITH A ONE-PHASE FREE BOUNDARY PROBLEM

For given $0 < \alpha < 1$ and $\Lambda > 0$, let us examine the free interface problem (1.16)-(1.18): it is easy to integrate (1.16) for $\xi < 0$, therefore

$$u(\xi) = \theta e^{c\xi}, \quad v(\xi) = 1 - (1-v(0))e^{\frac{c}{\Lambda}\xi}, \quad \xi < 0.$$

Because we look for u and v continuously differentiable, it comes

$$u(0) = \theta, \quad u'(0) = c\theta, \quad v'(0) = -\frac{c}{\Lambda}(1-v(0)),$$

and for the same reason, it follows that $u(\xi) = 1$ for all $\xi \geq R$. This suggests the following simplification.

Proposition 2.1. *Let $0 < \alpha < 1$ and $\Lambda > 0$ be fixed. The free interface problem (1.16)-(1.18) is equivalent to the one-phase free boundary problem: find $c > 0$, a free boundary $0 < R < +\infty$, u and v in $C^\infty([0, R)) \cap C^1([0, R])$, such that $(c, R, u(\xi), v(\xi))$ verifies*

$$\begin{cases} u_{\xi\xi} - cu_\xi = -v^\alpha, & 0 < \xi < R, \\ \Lambda v_{\xi\xi} - cv_\xi = v^\alpha, & 0 < \xi < R, \end{cases} \quad (2.1)$$

with the following boundary conditions at $\xi = 0$

$$u(0) = \theta, \quad u'(0) = c\theta, \quad 0 < v(0) < 1, \quad v'(0) = -\frac{c}{\Lambda}(1 - v(0)), \quad (2.2)$$

and the free boundary conditions at $\xi = R$

$$v(R) = v'(R) = 0, \quad u(R) = 1, \quad u'(R) = 0. \quad (2.3)$$

Moreover, the following integral relation holds

$$c = \int_0^R v^\alpha(\xi) d\xi. \quad (2.4)$$

Proof. (i) Assume there exists $c > 0$, $R > 0$, $u \in C^1(\mathbb{R})$, $v \in C^1(\mathbb{R})$ such that $(c, R, u(\xi), v(\xi))$ verifies the free interface problem (1.16)-(1.18). It is clear that u is smooth on $(-\infty, 0) \cup (0, R) \cup (R, +\infty)$, v is smooth on $(-\infty, 0) \cup (0, R)$ (recall that v is identically zero on $[R, +\infty)$). Solving (1.16) for $\xi < 0$ and taking (1.17) and (1.18) into account, we obtain

$$\begin{cases} u(\xi) = \theta e^{c\xi}, & \xi < 0, \\ v(\xi) = 1 - (1 - v(0))e^{\frac{c}{\Lambda}\xi}, & \xi < 0. \end{cases} \quad (2.5)$$

Thus, (2.2) follows immediately. Next, solving $u_{\xi\xi} - cu_\xi = 0$ for $\xi > R$ with $u(+\infty) = 1$ yields $u(\xi) = 1$ for $\xi \geq R$. Because u and v are continuously differentiable, the free boundary conditions (2.3) are verified. Finally, $(c, R, u|_{[0,R]}, v|_{[0,R]})$ satisfies the free boundary problem (2.1)-(2.3). Note that in (2.3) the conditions $u(R) = 1$ and $u'(R) = 0$ are equivalent.

(ii) Now, assume there exist $c > 0$, $R > 0$, u and v in $C^\infty([0, R]) \cap C^1([0, R])$ such that $(c, R, u(\xi), v(\xi))$ is a solution to (2.1)-(2.3). Then, we extend u and v , respectively, by \tilde{u} and \tilde{v} to the whole line as follows: we define \tilde{u} and \tilde{v} for $\xi < 0$ via (2.5); for $\xi > R$ we set $\tilde{u} \equiv 1$, $\tilde{v} \equiv 0$. By construction, \tilde{u} and \tilde{v} are in $C^1(\mathbb{R})$. Finally, it is easy to check that $(c, R, \tilde{u}(\xi), \tilde{v}(\xi))$ is a solution of (1.16)-(1.18).

(iii) We easily obtain formula (2.4) by integrating the equation $\Lambda v_{\xi\xi} - cv_\xi = v^\alpha$ between 0 and R , and taking (2.2) and (2.3) into account. Note that $c = R$ when $\alpha = 0$ (see (1.11)). \square

3. THE SCALAR FREE BOUNDARY PROBLEM

It transpires from Proposition 2.1 that the main parameter of the free boundary problem (2.1)-(2.3) is $v(0)$. We first focus on the scalar problem for v with free boundary R , assuming $v(0) = v_0$ given in the interval $(0, 1)$. Throughout the next sections, we intend to prove the following theorem:

Theorem 3.1. *Let $0 < \alpha < 1$, $\Lambda > 0$ be fixed. Then, for each $v_0 \in (0, 1)$, there exists a unique solution $(c(v_0), R(v_0), v(v_0; \xi))$ to the one-phase free boundary problem*

$$\begin{cases} \Lambda v''(\xi) - cv'(\xi) = v^\alpha(\xi), & 0 < \xi < R, \\ v(0) = v_0, \quad v'(0) = -\frac{c}{\Lambda}(1 - v_0), \\ v(R) = v'(R) = 0, \end{cases} \quad (3.1)$$

such that $c(v_0) > 0$, $0 < R(v_0) < +\infty$, $v \in C^\infty([0, R]) \cap C^{2+[\beta], \beta-[\beta]}([0, R])$, $\beta = \frac{2\alpha}{1-\alpha}$. Moreover,

$$v(\xi) > 0, \quad v'(\xi) < 0, \quad v''(\xi) > 0, \quad \forall \xi \in [0, R]. \quad (3.2)$$

Corollary 3.2. *Let $0 < \alpha < 1$, $\Lambda > 0$ be fixed. Set $I = [a, b]$, $0 < a < b < 1$. For $v_0 \in I$, let $(c(v_0), R(v_0), v(v_0; \xi))$ denote the solution of system (3.1).*

(i) There exists a finite R_{max} depending only upon α , Λ , a and b such that $0 < R(v_0) \leq R_{max}$ for all $v_0 \in I$.

(ii) The mapping $v_0 \mapsto (c(v_0), R(v_0), \tilde{v}(v_0))$ from I to $(\mathbb{R}^+)^2 \times C^1([0, R_{max}])$ is continuous, where $\tilde{v}(v_0; \xi)$ is the extension by 0 of the function $v(v_0; \xi)$ to the interval $[0, R_{max}]$.

3.1. Formal a priori estimates for $c(v_0)$. At the outset, it is easy to establish a priori estimates for $c > 0$, assuming that $(c(v_0), R(v_0), v(v_0; \xi))$ is a solution to (3.1). Multiplying formally the equation by v' and integrating from 0 to R , using the boundary and free boundary conditions in (3.1), we arrive at

$$\frac{1}{2} \frac{c^2}{\Lambda} (v_0 - 1)^2 + c \int_0^R (v')^2 d\xi = \frac{1}{1 + \alpha} v_0^{1+\alpha}. \quad (3.3)$$

Thus, we immediately achieve the upper bound $c < c_+$ with

$$c_+^2 = \frac{2\Lambda}{1 + \alpha} \frac{v_0^{1+\alpha}}{(v_0 - 1)^2}. \quad (3.4)$$

Likewise, we multiply the equation by v . Integrating from 0 to R and using again the boundary and free boundary conditions in (3.1), we obtain

$$-cv_0(v_0 - 1) - \Lambda \int_0^R (v')^2 d\xi + \frac{c}{2} v_0^2 = \int_0^R v^{1+\alpha} d\xi, \quad (3.5)$$

which together with (3.3) gives the lower bound $c > c_-$ with

$$c_-^2 = \frac{2\Lambda}{1 + \alpha} v_0^{1+\alpha}. \quad (3.6)$$

Summarizing, we have the following a priori estimates:

$$0 < c_-(v_0) < c(v_0) < c_+(v_0). \quad (3.7)$$

These bounds are proved via a different method (see Subsection 6.2 below).

3.2. Proof of uniqueness. Here, we give a direct proof of uniqueness in Theorem 3.1. We find an alternate proof in Section 4 (see Lemma 4.8).

Lemma 3.3. *Let $v_0 \in (0, 1)$ be fixed. There exists a unique solution $(c(v_0), R(v_0), v(v_0; \xi))$ to system (3.1).*

Proof. Assume that $(c_i(v_0), R_i(v_0), v_i(v_0; \xi))$, $i = 1, 2$, are two different solutions to (3.1), namely

$$\begin{cases} \Lambda v_i''(\xi) - c_i v_i'(\xi) = v_i^\alpha(\xi), & 0 < \xi < R_i, \\ v_i(0) = v_0, v_i'(0) = -\frac{c_i}{\Lambda}(1 - v_0), \\ v_i(R_i) = v_i'(R_i) = 0. \end{cases} \quad (3.8)$$

Without loss of generality, we may assume that $c_2 > c_1$.

(i) First, we prove that

$$R_2 < R_1 \quad \text{and} \quad v_2(\xi) < v_1(\xi), \quad \forall \xi \in (0, R_2). \quad (3.9)$$

Set $\varphi = v_1 - v_2$, $\varphi(0) = 0$. As $|v_2'(0)| > |v_1'(0)|$, one has $\varphi'(0) > 0$, thus $\varphi > 0$. Assume by contradiction that the two solutions intersect: let $\xi_0 \in (0, \bar{R})$, $\bar{R} = \min\{R_1, R_2\}$, be the first

point such that $\varphi(\xi_0) = 0$, i.e. $\varphi(\xi) > 0, \forall \xi \in (0, \xi_0)$. Noticing that $\varphi \in C^2([0, \bar{R}])$, by Rolle's theorem there exists a $\bar{\xi} \in (0, \xi_0)$ such that $\varphi'(\bar{\xi}) = 0$. From (3.8), one has

$$\Lambda\varphi'' - c_1\varphi' + v_2'(c_2 - c_1) = v_1^\alpha - v_2^\alpha. \quad (3.10)$$

Integrating (3.10) from 0 to $\bar{\xi}$ and using $\varphi'(\bar{\xi}) = \varphi(0) = 0$, we obtain

$$-\Lambda\varphi'(0) - c_1\varphi(\bar{\xi}) + (c_2 - c_1)(v_2(\bar{\xi}) - v_2(0)) = \int_0^{\bar{\xi}} (v_1^\alpha - v_2^\alpha) d\xi. \quad (3.11)$$

Thanks to $\varphi'(0) > 0$, $\varphi(x) > 0$ and $v_2'(\xi) < 0, \forall \xi \in (0, \bar{\xi})$, it is found that the left-hand side of (3.11) is negative while the right-hand side is positive, thus, a contradiction.

(ii) Next, according to formula (2.4), it holds

$$c_1 = \int_0^{R_1} v_1^\alpha d\xi \quad \text{and} \quad c_2 = \int_0^{R_2} v_2^\alpha d\xi, \quad (3.12)$$

hence from (3.9)

$$c_2 = \int_0^{R_2} v_2^\alpha d\xi < \int_0^{R_2} v_1^\alpha d\xi < \int_0^{R_1} v_1^\alpha d\xi = c_1,$$

which contradicts with the assumption $c_2 > c_1$; the uniqueness is thus proved. \square

3.3. Scheme of the proof of Theorem 3.1. Let $0 < \alpha < 1$, $\Lambda > 0$, $0 < v_0 < 1$ be fixed. For $c > 0$, we consider the system

$$\begin{cases} \Lambda v''(\xi) - cv'(\xi) = v^\alpha(\xi), & \xi > 0, \\ v(0) = v_0, v'(0) = -\frac{c}{\Lambda}(1 - v_0), \end{cases} \quad (3.13)$$

and we look for solution v such that $v(\xi) > 0$, $v'(\xi) < 0$. Now, we set:

$$t = \xi, \quad x(t) = v(\xi), \quad y(t) = v'(\xi). \quad (3.14)$$

Therefore, problem (3.13) is equivalent to the following differential system in the quadrant $Q = \{x \geq 0, y \leq 0\}$

$$X_c : \begin{cases} x'(t) = y(t), \\ \Lambda y'(t) = cy(t) + x^\alpha(t), \end{cases} \quad (3.15)$$

with initial conditions

$$x(0) = v_0, \quad y(0) = -\frac{c}{\Lambda}(1 - v_0). \quad (3.16)$$

The origin $O = (0, 0)$ is the unique critical point of the vector field X_c given by (3.15). Clearly, Theorem 3.1 demonstrates the existence of $c > 0$ and a trajectory $(x(t), y(t))$ in Q , defined on $[0, R)$ with initial conditions $(x(0), y(0))$ given by (3.16), which tends towards O for $t \rightarrow R_-$. (It also can be said that O is the ω -limit set of the trajectory; $(x(t), y(t))$ extends continuously on $[0, R]$ by $(x(R), y(R)) = (0, 0)$).

We prove Theorem 3.1 by using primarily the Poincaré-Bendixson theorem (see Appendix A), and, from time to time, by using some other simple qualitative arguments. As mentioned above, the main difficulty is that the vector field is not of class C^1 along the axis $\{x = 0\}$ when $0 < \alpha < 1$. This is overcome by putting the origin at the boundary of the domain.

Section 4 is devoted to the topological study of the differential equation (3.15), for a fixed value of c , especially the phase portrait and the existence of a stable manifold. The dependence

on c w.r.t. v_0 is studied in Section 5. In Section 6, we approach the proof of Theorem 3.1, per se. The critical result (see Lemma 6.2) is that the settling time $R(v_0)$ to proceed from the initial condition to the origin is finite. Eventually, we return to the notation of the free boundary problem (3.1) and prove Corollary 3.2.

4. A TOPOLOGICAL APPROACH (c FIXED)

In this section, the parameter $c \geq 0$ is fixed; α is such that $0 \leq \alpha \leq 1$, and we assume $v_0 > 0$ (note that these hypotheses go beyond the physical framework). We intend to study the flow of the vector field X_c with differential equation (3.15), that reads:

$$\begin{cases} x' = y, \\ y' = \frac{1}{\Lambda}(cy + x^\alpha), \end{cases} \quad (4.1)$$

in a topological way. We recall that the vector field X_c is a first order differential operator $y \frac{\partial}{\partial x} + \frac{1}{\Lambda}(cy + x^\alpha) \frac{\partial}{\partial y}$. We write $X_c \cdot f$ its action on a function f :

$$X_c \cdot f(x, y) = y \frac{\partial f}{\partial x}(x, y) + \frac{1}{\Lambda}(cy + x^\alpha) \frac{\partial f}{\partial y}(x, y).$$

This vector field is considered in the quadrant $Q = \{x \geq 0, y \leq 0\}$. We denote its flow in Q by $\varphi_c(t, m)$. (For convenience, some information about vector fields is given in Appendix A).

In the first subsection, we state a general topological result: the phase portrait of the vector field X_c , extended to the whole plane, is independent of c , α and λ . The proof of this result (given in Appendix B), is based in a crucial way on the existence and uniqueness of the stable manifold. The latter result is established below in Subsections 4.2, 4.3 and 4.4.

4.1. Phase portrait. We extend the vector field X_c to the whole plane by defining:

$$X_c^E(x, y) = y \frac{\partial}{\partial x} + \frac{1}{\Lambda}(cy + \text{sign}(x)|x|^\alpha) \frac{\partial}{\partial y}. \quad (4.2)$$

Remark 4.1. *To have a non-ambiguous definition, even for $\alpha = 0$, we consider that $\text{sign}(0) = 0$ (and ± 1 otherwise). When $\alpha = 0$, X_c^E is well defined, but discontinuous along the Oy -axis, outside the origin, which is a singular point.*

As $X_c^E|_Q = X_c$, this new field is an extension of X_c . It is observed that

$$X_c^E(-x, -y) = -X_c^E(x, y).$$

It follows that the orbits for $\{x \leq 0\}$ are obtained from the orbits for $\{x \geq 0\}$ by using the symmetry $(x, y) \mapsto (-x, -y)$ and next changing the time orientation (see Figure 1). The trajectories of X_c^E cross in a regular way the Oy -axis at points different from the origin, and the origin is an equilibrium point.

In this subsection we present a topological description for the vector field X_c^E , based on the following definition (see [2], [17], [19], [15]):

Definition 4.2. *Two vector fields X and Y defined on the same manifold M , are said topologically equivalent if there exists a homeomorphism H of M , which sends the orbits of X onto the orbits of Y (preserving time-orientation). To be topologically equivalent is clearly an equivalence relation (in the space of vector fields on M). An equivalence class for the topological equivalence is called a phase portrait.*

That is to say, two vector fields are topologically equivalent if they have the same phase portrait. We have the following global result whose proof is given in Appendix B:

Proposition 4.3. *The phase portrait of X_c^E is independent of c, α and even of $\Lambda > 0$. In other words, the general vector field X_c^E is topologically equivalent on \mathbb{R}^2 to the linear saddle type vector field $X_0 = y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y}$ corresponding to $\Lambda = 1, \alpha = 1, c = 0$.*




FIGURE 1. Phase portrait of X_c^E for $\alpha < 1$. It fulfills some differentiable features of the flow such as the contact of orbits at the origin.

The result stated in Proposition 4.3 can be compared with the Hartman-Grobman Theorem (see, e.g. [26]). Of course, as the vector field X_c^E is not C^1 , and the origin is not an hyperbolic singularity when $\alpha < 1$, it is impossible to apply the Hartman-Grobman Theorem.¹ In Appendix B we construct directly “by hands” the homeomorphism H of equivalence, beginning with the construction on quadrant Q on which we consider X_c . Next, H is extended in the whole plane in a rather direct way. The fact that X_c has a unique stable manifold in Q is crucial for this construction.

4.2. Existence of a local stable manifold. For $c = 0$, the vector field X_0 is Hamiltonian with Hamiltonian function $H(x, y) = \frac{1}{2}y^2 - \frac{1}{\Lambda(1+\alpha)}x^{1+\alpha}$. This vector field has stable manifold in Q at the origin O :

$$L_0 := \left\{ y = y_0(x) = - \left(\frac{2}{\Lambda(1+\alpha)} \right)^{1/2} x^{\frac{1+\alpha}{2}} \right\}. \quad (4.3)$$

For any $c \geq 0$, we have that:

$$X_c \cdot H(x, y) = -\frac{1}{\Lambda}yx^\alpha + \frac{1}{\Lambda}(cy + x^\alpha)y = \frac{c}{\Lambda}y^2.$$

¹Moreover, the time to proceed from one point on the stable manifold to the origin is infinite for X_0 but finite for X_c^E when $\alpha < 1$, see Subsection 6.3.

This implies that, for $c > 0$, the vector field X_c is transverse to L_0 and directed downwards all along L_0 , outside O .

Now, for any $v_0 > 0$, we consider in \mathbb{R}^2 the region $\mathcal{T}_{v_0} = \{0 \leq x \leq v_0, y_0(x) \leq y \leq 0\}$. This compact region is a curved triangle that we call a *trapping triangle* for a reason that will become apparent. This triangle has the following three corners: O , $A_{v_0} = (v_0, 0)$ and $B_{v_0} = (v_0, y_0(v_0))$; and three sides denoted as follows: $[OA_{v_0}]$, $[A_{v_0}B_{v_0}]$ and $[OB_{v_0}]$. The vector field X_c is transverse and has an upward direction along $(OA_{v_0}) = [OA_{v_0}] \setminus \{O\}$. It is transverse and has a left direction along the open interval $(A_{v_0}B_{v_0}) = [A_{v_0}B_{v_0}] \setminus \{A_{v_0}\} \cup \{B_{v_0}\}$. As already mentioned, X_c is transverse and has a downward direction along $(OB_{v_0}) = [OB_{v_0}] \setminus \{O\}$ (see Figure 2).




FIGURE 2. Local stable manifold

The following lemma proves the existence of a local stable manifold contained inside \mathcal{T}_{v_0} :

Lemma 4.4. *There exist points $m \in (A_{v_0}B_{v_0})$ whose trajectory $\varphi_c(t, m)$ remains in \mathcal{T}_{v_0} for all times and tends toward O when $t \rightarrow \tau^+(m)$. Let S be the set of points in $(A_{v_0}B_{v_0})$ whose trajectory tends toward O . Trajectories starting at points $(A_{v_0}B_{v_0}) \setminus S$ cut the side (OA_{v_0}) or the side (OB_{v_0}) after a finite time.*

Proof. We consider any point $m \in (A_{v_0}B_{v_0})$ and look at its trajectory in \mathcal{T}_{v_0} . More precisely, to remain in the interior Ω_{v_0} of \mathcal{T}_{v_0} , we start at a point $\varphi_c(T_m, m)$ for $T_m > 0$ small enough, and we consider the positive trajectory of this point inside the open set Ω_{v_0} . The trajectory is defined on a time interval $[T_m, \tau^+(m))$.

Using Corollary A.4, we know that there exists a sequence $(t_n) \rightarrow \tau^+(m)$ such that the sequence of points $\varphi_c(t_n, m)$ tends toward a point $p \in \partial\mathcal{T}_{v_0}$. Clearly, p cannot belong to $(A_{v_0}B_{v_0})$, as the vector field is repulsive along this side. If $p \in (OA_{v_0})$, then it is easy to see, using a flow box centered at p , that the trajectory arrives at p at the finite time $\tau^+(m)$ and crosses the axis Ox at the regular point $p = \varphi(\tau^+(m), m)$. The same remark is valid if $p \in (OB_{v_0})$. In this case, the trajectory reaches a regular point on $L_0 \setminus \{O\}$ in a finite time.

The set \mathcal{O}_{up} of points m for which the trajectory arrives on (OA_{v_0}) is open in $(A_{v_0}B_{v_0})$, as the transversality of this trajectory at each end to $[A_{v_0}B_{v_0}]$ and Ox respectively is an open condition. Moreover, \mathcal{O}_{up} is non-empty, as this set contains points near A_{v_0} . In a similar way, it is seen that the set \mathcal{O}_{down} of points m for which the trajectory arrives on (OB_{v_0}) is a non-empty open subset of $(A_{v_0}B_{v_0})$. However, as $(A_{v_0}B_{v_0})$ is connected, it cannot be the union of the two non-empty and disjoint open sets \mathcal{O}_{up} and \mathcal{O}_{down} . This means that there are points $m \in (A_{v_0}B_{v_0})$ for which the limit point p is the origin O .

Then, consider such a point $m \in (A_{v_0}B_{v_0})$ with the property that there exists a time sequence $(t_n) \rightarrow \tau^+(m)$ such that $\varphi_c(t_n, m) \rightarrow O$. This means that for each $\varepsilon > 0$ small enough, there exists a $n(\varepsilon)$ such that $\varphi_c(t_{n(\varepsilon)}, m) \in \mathcal{T}_\varepsilon$, where \mathcal{T}_ε is the small triangle obtained by replacing v_0 by ε . However, as the vertical side of \mathcal{T}_ε contained in $\{x = \varepsilon\}$ is repulsive (the vector field is directed toward the left) and as the trajectory of m remains in \mathcal{T}_{v_0} by definition (we consider the trajectories only into Ω_{v_0}), the trajectory of m is trapped in \mathcal{T}_ε ; hence, the point $\varphi_c(t, m)$ belongs to \mathcal{T}_ε for all $t \geq t_{n(\varepsilon)}$. As the diameter of \mathcal{T}_ε tends to zero with ε , it is implied that $\varphi_c(t, m) \rightarrow O$ for $t \rightarrow \tau^+(m)$. Therefore, the trajectory of m is a local stable manifold contained in \mathcal{T}_{v_0} . \square

In the next subsections we use more general trapping triangles:

Definition 4.5. *A trapping triangle \mathcal{T}_{v_0} for the vector field X_c is a curved triangle in Q with a corner at O and two other corners A_{v_0}, B_{v_0} on a vertical line $\{x = v_0\}$, for some $v_0 > 0$. The side $[OA_{v_0}]$ is contained in a graph $L^u := \{y = y^u(x)\}$, the side $[OB_{v_0}]$ in a graph $L^d := \{y = y^d(x)\}$ and the side $[A_{v_0}B_{v_0}]$ is a vertical interval in $\{x = v_0\}$. We assume that L^u is located above L^d , i.e. $y^d(x) < y^u(x) \leq 0$ for all $x > 0$ (and by hypothesis: $y^d(0) = y^u(0) = 0$). We also assume that X_c is transverse in the upward direction all along $(OA_{v_0}) = [OA_{v_0}] \setminus \{O\}$ and is transverse in the downward direction all along (OB_{v_0}) . It is clear that X_c is transverse to (A_{v_0}, B_{v_0}) in the left direction (see Figure 3).*




FIGURE 3. Trapping triangle

The triangle used in the proof of Lemma 4.4 is obviously a trapping triangle. Any trapping triangle verifies the statement of Lemma 4.4. As the proof is exactly the same, we state the result without giving a new proof:

Lemma 4.6. *Let \mathcal{T}_{v_0} be a trapping triangle as in Definition 4.5. There exist points $m \in (A_{v_0}B_{v_0})$ whose trajectory $\varphi_c(t, m)$ remains into \mathcal{T}_{v_0} for all times and tends toward O for $t \rightarrow \tau^+(m)$. Trajectories starting at other points of $(A_{v_0}B_{v_0})$ cut the side (OA_{v_0}) or the side (OB_{v_0}) after a finite time.*

4.3. Existence of global stable manifold. In this subsection we see how to extend a local stable manifold found in Subsection 4.2 into a global one. Such a global stable manifold is a trajectory contained in the quadrant Q and converging toward the origin. We use again Corollary A.4, but now in an easier way, as the vector field X_c is analytic in a neighborhood of the domain under consideration.

Lemma 4.7. *There exists a global stable manifold at the origin, given by a trajectory of X_c in Q converging toward the origin in the positive direction of time. The orbit defined by this trajectory (its geometrical image) is the graph of a function $y = y(x)$ with $x \in (0, +\infty)$. This function is analytic for $x > 0$ and extends continuously at 0 with the value $y(0) = 0$. Its graph is located above L_0 , more precisely, $y_0(x) < y(x) < 0$ for all $x > 0$ ($y = y_0(x)$ is the equation of the curve L_0 defined by (4.3)).*

Proof. We start with a local stable manifold found in Lemma 4.4, which is defined by a graph $\{y = y(x)\}$ for $0 \leq x \leq v_0$ where $v_0 > 0$. As v_0 can take any positive value, the idea might be to make v_0 go to $+\infty$. But the problem is that local stable manifolds may depend on v_0 . Therefore, we proceed differently to obtain the desired global manifold, using again Corollary A.4.

A value $v_0 > 0$ being chosen, we consider the sequence $v_n = nv_0$. For each $n \in \mathbb{N}$, we consider the curved square S_n defined by $S_n = \{(x, y) \mid v_n \leq x \leq v_{n+1}, y_0(x) \leq y \leq 0\}$. S_n has four corners: $A_n = (v_n, 0)$, $B_n = (v_n, y_0(v_n))$, $B_{n+1} = (v_{n+1}, y_0(v_{n+1}))$ and $A_{n+1} = (v_{n+1}, 0)$; and four sides: the horizontal linear segment $[A_n A_{n+1}]$, the two vertical linear segments $[A_n B_n]$, $[A_{n+1} B_{n+1}]$ and the curved segment $[B_n B_{n+1}]$ located on the curve L_0 . The vector field X_c is analytic and without singularities in a neighborhood of S_n . Moreover, the vector field is transverse along each side, with an entering direction along the side $[A_{n+1} B_{n+1}]$ and a leaving direction along the three over sides (see Figure 4).

Now, applying Corollary A.4 in a neighborhood of S_n to the vector field $-X_c$, we have that if m_n is a point in the interior of the segment $[A_n B_n]$ (i.e., $m_n = (v_n, y_n)$ with $y_0(v_n) < y_n < 0$), then there exists a $\hat{t}_n < 0$ such that $\varphi(\hat{t}_n, m_n)$ belongs to the interior of the segment $[A_{n+1} B_{n+1}]$.

Let $m_0 = (v_0, y_0)$ be the endpoint of a local stable manifold founded by Lemma 4.4 in the trapping triangle \mathcal{T}_{v_0} . We have that m_0 belongs to the interior of $[A_0 B_0]$. Now, using the above result as an induction step in the construction of a global stable manifold, we find a decreasing sequence of times $t_0 = 0 > t_1 > t_2 > \dots > t_n \dots$, in the time interval of the trajectory $\varphi_c(t, m_0)$ of X_c , such that the segment of trajectory $\varphi_c([t_n, t_{n+1}], m_1)$ connects inside S_n a point $m_n \in (A_n B_n)$ to a point $m_{n+1} \in (A_{n+1} B_{n+1})$. As the vector field X_c has a norm greater than a positive value if $x \geq v_0$, the time sequence (t_i) tends to $-\infty$ (this means that $\tau_-(m_0) = -\infty$). Now, as the vector field X_c has a negative horizontal component in the interior $\{x > 0, y < 0\}$ of the quadrant Q , the orbit of m_0 has a regular projection onto the positive axis $Ox^+ = \{x > 0\}$ (the orbit is a covering of Ox^+). As Ox^+ is simply connected, this covering must be a diffeomorphism. The orbit is a graph of a function $y = y(x)$ defined for $x > 0$. As X_c is analytic, the trajectory is analytic and also the function $y(x)$ for $x > 0$. Taking into account the properties of the local stable manifold, we already know that $y(x)$ has a continuous extension at 0 by $y(0) = 0$. Finally, we have by construction that $y_0(x) < y(x) < 0$ for all $x > 0$. \square

4.4. Uniqueness of the stable manifold. Consider any stable manifold in Q , i.e. any trajectory in Q converging toward the origin. As seen above, such a stable manifold exists, and we want to prove that only one exists.

Lemma 4.8. *There exists a unique trajectory in Q which converges toward the origin in positive times.*

Proof. Let Γ_1 and Γ_2 be two such orbits. With the same arguments as those used in Lemma 4.7, it is easily seen that there exists a $v_0 > 0$ such that these orbits are the graphs of two




FIGURE 4. Construction of a global invariant manifold

functions $y_1(x)$ and $y_2(x)$ respectively, defined on the same interval $(0, v_0]$, verifying $y_1(x) < 0$ and $y_2(x) < 0$ for $0 < x \leq v_0$ and with a continuous extension $y_1(0) = y_2(0) = 0$. We want to prove that these two functions coincide on $[0, v_0]$, which implies that Γ_1 and Γ_2 also coincide globally.

Assume on the contrary, for instance, that $y_1(v_0) < y_2(v_0)$. As the two graphs are different half-orbits, this inequality persists for any $x \in (0, v_0]$, which means that $y_1(x) < y_2(x) < 0$ for any $x \in (0, v_0]$. Moreover, each function is a solution of the differential equation $\frac{dy}{dx} = \frac{c}{\Lambda} + \frac{1}{\Lambda} \frac{x^\alpha}{y}$. Then, we have the following differential equation:

$$\frac{d}{dx}(y_2 - y_1) = \frac{1}{\Lambda} \left(\frac{1}{y_2} - \frac{1}{y_1} \right) x^\alpha. \quad (4.4)$$

As $y_1(x) < y_2(x) < 0$, we have that $\frac{1}{y_2(x)} - \frac{1}{y_1(x)} < 0$ for all $0 < x \leq v_0$. It follows from (4.4) that the function $x \mapsto (y_2 - y_1)(x)$ is decreasing and, in particular:

$$(y_2 - y_1)(x) > (y_2 - y_1)(v_0) > 0$$

for all $x \in (0, v_0]$. This contradicts the fact that $(y_2 - y_1)(x) \rightarrow 0$ when $x \rightarrow 0$. \square

The unique stable manifold is the one obtained in Lemma 4.7. This manifold is the graph of a unique well-defined function we denote by $y_c(x)$. More specifically, we call it *the stable manifold of X_c at the origin and write it in the form $\{y = y_c(x)\}$* . We recall that $y_c(x)$ is analytic for $x > 0$, continuous at 0 with value $y_c(0) = 0$; it is located in Q above the curve L_0 , i.e., it verifies $y_0(x) < y_c(x) < 0$ for all $x > 0$. The new definition of $y_0(x)$ as the function $y_c(x)$ for $c = 0$, coincides with the former one.

5. SOME TECHNICAL LEMMATA

In Section 4, we defined a c -family of analytic functions $y_c(x)$. Now, considering $c \geq 0$ as a parameter, we study how this family depends on c and prove some useful technical lemmata.

5.1. Dependence of the stable manifold on parameter c . The vector field X_c rotates in function of the parameter c , in the sense of Duff (see [14]). We take advantage of this property in the following lemma:

Lemma 5.1. *For each $x > 0$, the map $c \mapsto y_c(x)$ is increasing.*

Proof. We consider two values of the parameter: $c_0 < c_1$. The stable manifolds are $L_{c_0} := \{y = y_{c_0}(x)\}$ and $L_{c_1} := \{y = y_{c_1}(x)\}$, respectively. Take any value $x > 0$; then, the vector

$$X_{c_1}(x, y_{c_1}(x)) = \left(y_{c_1}(x), \frac{c_1}{\Lambda} y_{c_1}(x) + \frac{1}{\Lambda} x^\alpha \right)$$

is tangent to L_{c_1} at the point $(x, y_{c_1}(x))$ for all $x > 0$. At the same point $(x, y_{c_1}(x))$, the value of the vector field X_{c_0} is :

$$X_{c_0}(x, y_{c_1}(x)) = \left(y_{c_1}(x), \frac{c_0}{\Lambda} y_{c_1}(x) + \frac{1}{\Lambda} x^\alpha \right).$$

As $\frac{c_0}{\Lambda} y_{c_1}(x) > \frac{c_1}{\Lambda} y_{c_1}(x)$, the vector field X_{c_0} is, for any $x > 0$, transverse to the curve L_{c_1} and is directed upward. Then, we construct trapping triangles for any v_0 (see Definition 4.5), by using the curves $L^d = L_0$ and $L^u = L_{c_1}$. As a consequence of Lemma 4.6, the unique stable manifold for the vector field X_{c_0} , i.e. L_{c_0} , is located strictly between L_0 and L_{c_1} meaning that, for any $x > 0$, $y_0(x) < y_{c_0}(x) < y_{c_1}(x)$. \square

Now look at the continuity of the stable manifold with respect to c .

Lemma 5.2. *For each $x > 0$, the map $c \mapsto y_c(x)$ is continuous.*

Proof. As the map $c \mapsto y_c(x)$ increases, left- and right-hand limits exist at any \bar{c} , namely $y_{\bar{c}}^-(x) = \lim_{c \rightarrow \bar{c}^-} y_c(x)$ and $y_{\bar{c}}^+(x) = \lim_{c \rightarrow \bar{c}^+} y_c(x)$, with the property that $y_{\bar{c}}^-(x) \leq y_{\bar{c}}(x) \leq y_{\bar{c}}^+(x)$. To prove that the function y_c is continuous at \bar{c} , we must check that $y_{\bar{c}}^-(x) = y_{\bar{c}}^+(x)$.

We claim that the positive half-orbit of $X_{\bar{c}}$ by $y_{\bar{c}}^-(x)$ is a local stable manifold. If this is not the case, it follows from Lemma 4.4 that the trajectory must cut (transversally) $L_0^+ = L_0 \setminus \{O\}$ or $Ox^+ = Ox \setminus \{O\}$ after a finite time t_0 . Choosing some $t_1 > t_0$, we apply Theorem A.2 at the segment of trajectory $\varphi([0, t_1], (x, y_{\bar{c}}^-(x)), \bar{c})$. Theorem A.2 implies that the map $c \mapsto \{t \in [0, t_1] \mapsto \varphi(t, (x, y_c(x)), c)\}$ from $(\bar{c} - \delta, \bar{c}]$ to $C^1([0, t_1], \mathbb{R}^2)$ exists and is continuous at the endpoint \bar{c} if δ small enough. Then, the orbit of X_c by the point $(x, y_c(x))$ for $c \in (\bar{c} - \delta, \bar{c})$ and δ small enough, cuts also L_0^+ or Ox^+ . However, this is impossible because $\{y = y_c(x)\}$ is the stable manifold of X_c . Then, the only possibility for the trajectory through the point $(x, y_{\bar{c}}^-(x))$ is the invariant manifold of $X_{\bar{c}}$. A similar argument proves that the trajectory through the point $(x, y_{\bar{c}}^+(x))$ has also to be the invariant manifold of $X_{\bar{c}}$. As the invariant manifold of $X_{\bar{c}}$ is unique, we obtain $y_{\bar{c}}^-(x) = y_{\bar{c}}^+(x)$. \square

Finally, the family of maps $y_c(x)$ enjoys the following global properties:

Lemma 5.3. *The mapping $(x, c) \in [0, +\infty) \times [0, +\infty) \mapsto y_c(x) \in \mathbb{R}$ is continuous as a function of two variables and for all c , one has $y_c(0) = 0$. For all c , the function $x \mapsto y_c(x)$ is analytic for $x > 0$. Recall that, for all $x > 0$, the map $c \mapsto y_c(x)$ increases continuously.*

Proof. Consider a fixed value $x_0 > 0$ and write $y_0 = y_{c_0}(x_0)$. The trajectory X_{c_0} by the point (x_0, y_0) cuts transversally the vertical line d_x passing by any $(x, 0)$ with $x > 0$, at the point $(x, y_{c_0}(x))$. Then, if (x_1, y_1, c) is near (x_0, y_0, c_0) , the trajectory of X_c by the point (x_1, y_1) cuts also d_x transversally. This defines a local function $y = F(x_1, y_1, x, c)$ in a neighborhood of the point $(x_0, y_0, x, c) \in \mathbb{R}^4$. Using Theorem A.2 and the Implicit Function Theorem we have F is analytic. Now, we observe that $y_c(x) = F(x_0, y_c(x_0), x, c)$, where x_0 is fixed. By Lemma 5.2, the function $c \mapsto y_c(x_0)$ is continuous. Then, the composed function $(x, c) \mapsto y_c(x)$ is also continuous. This proves the continuity at (x, c) for any $x > 0$. As $y_0(x) \leq y_c(x) < 0$, the function $y_c(x)$ converges uniformly in c toward 0 when $x \rightarrow 0_+$.

The function $x \mapsto y_c(x) = F(x_0, y_c(x_0), x, c)$ is analytic as a partial function of F . \square

5.2. An asymptotic expression for $y_c(x)$.

Lemma 5.4. *For all $c > 0$ and $x > 0$, the following holds:*

$$-\left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} x^{\frac{1+\alpha}{2}} < y_c(x) < -\left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} x^{\frac{1+\alpha}{2}} + \frac{c}{\Lambda}x. \quad (5.1)$$

Then, y_c has the following asymptotic expansion at the origin:

$$y_c(x) = -\left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} x^{\frac{1+\alpha}{2}} + O(x), \quad (5.2)$$

where the term $O(x)$ is uniform in c when c belongs to some compact subset of $[0, +\infty)$.

Proof. We claim that along the curve

$$l_c := \{y = \tilde{y}_c(x) = -\left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} x^{\frac{1+\alpha}{2}} + \frac{c}{\Lambda}x\}, \quad (5.3)$$

the vector field X_c is transverse, with an upward direction. To prove this claim, we consider the field $N_1(x)$ obtained by rotating the tangent field to the graph of l_c at the point $(x, \tilde{y}_c(x))$ by the angle $\pi/2$:

$$N_1(x) = \left(\left(\frac{1+\alpha}{2\Lambda}\right)x^{\frac{\alpha-1}{2}} - \frac{c}{\Lambda}, 1\right),$$

and we compute its scalar product with the vector $X_c(x) = X_c(x, \tilde{y}_c(x))$ which is equal to:

$$X_1(x) = \left(-\left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} x^{\frac{1+\alpha}{2}} + \frac{c}{\Lambda}x, -\frac{c}{\Lambda}\left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} x^{\frac{1+\alpha}{2}} + \frac{c^2}{\Lambda^2}x + \frac{1}{\Lambda}x^\alpha\right).$$

We obtain

$$\langle X_1(x), N_1(x) \rangle = \frac{c}{\Lambda} \left(\frac{1+\alpha}{2\Lambda}\right)^{1/2} x^{\frac{1+\alpha}{2}},$$

which implies the upward transversality all along $l_c \setminus \{0\}$.

Notice that the curve l_c cuts the Ox -axis at the value $x_1(c) = \left(\frac{2\Lambda}{(1+\alpha)c^2}\right)^{\frac{1}{1-\alpha}} > 0$ and remains in the quadrant Q only for $x \in [0, x_1(c)]$. Nevertheless, we can construct trapping triangles, using the curves L_0 and l_c , with a vertical side in $\{x = v_0\}$ when $0 < v_0 \leq x_1(c)$. It follows from Lemma 4.6 that $y_c(x)$ verifies (5.1) for $0 < x \leq x_1(x)$. As the graph of l_c is above the Ox axis, (5.1) is trivially verified if $x \geq x_1(c)$.

The asymptotic formula (5.2) follows directly from (5.1). \square

5.3. Estimate from below. The following estimate is significant in view of Lemma 6.6.

Lemma 5.5. *The stable manifold is located above the curve $\{y = -\frac{1}{c}x^\alpha\}$. This means that for all $x > 0$ it holds $y_c(x) > -\frac{1}{c}x^\alpha$.*

Proof. Along the curve $\{y = -\frac{1}{c}x^\alpha\}$, the vector field X_c is equal to $y \frac{\partial}{\partial x}$: it is horizontal and directed toward the left for all $x > 0$. As the tangent vector field to the curve $\{y = -\frac{1}{c}x^\alpha\}$ has a non-zero vertical component equal to $-\alpha x^{\alpha-1}$ for all $x > 0$, it follows that X_c is transverse to the curve $\{y = -\frac{1}{c}x^\alpha\}$ and directed downward all along it. Then, for any $v_0 > 0$, we can construct a trapping triangle \mathcal{T}_{v_0} , by using the curves $L^d = \{y = -\frac{1}{c}x^\alpha\}$ and the Ox -axis as L^u .

By Lemma 4.6, it follows that the stable trajectory is trapped inside \mathcal{T}_{v_0} for any $v_0 > 0$. This concludes the proof. \square

6. PROOF OF THEOREM 3.1

We are now in the position of proving Theorem 3.1. Let $0 < \alpha < 1$, $\Lambda > 0$ be fixed, $0 < v_0 < 1$ according to the physical framework.

6.1. Existence of $c(v_0)$. Recall that the initial condition (3.16) reads

$$x(0) = v_0, \quad y(0) = -\frac{c}{\Lambda}(1 - v_0). \quad (6.1)$$

With the notation of Section 4, fulfilling the initial condition (6.1) is equivalent to the existence of a $c = c(v_0)$ such that

$$y_{c(v_0)}(v_0) = -\frac{c(v_0)}{\Lambda}(1 - v_0). \quad (6.2)$$

The difficulty seems to be that $c(v_0)$ appears in both sides of the equation (6.2), i.e., in the expression of the stable manifold and in the initial condition itself. However, taking into account the above properties of $y_c(x)$, this is not really an issue, as we will see below.

Lemma 6.1. *For any v_0 , $0 < v_0 < 1$, there exists a unique value $c(v_0)$ such that*

$$y_{c(v_0)}(v_0) = -\frac{c(v_0)}{\Lambda}(1 - v_0).$$

Moreover, the functions $v_0 \mapsto c(v_0)$ and $v_0 \mapsto y_{c(v_0)}(x)$ are continuous on $(0, 1)$.

Proof. For simplicity, we denote $y(c) = y_c(v_0)$ and $\bar{y}(c) = -\frac{c}{\Lambda}(1 - v_0)$ for a fixed v_0 . At $c = 0$ it holds $y(0) = y_0(v_0) = -\left(\frac{2}{(1 + \alpha)\Lambda}\right)^{1/2} v_0^{\frac{1 + \alpha}{2}} < \bar{y}(0) = 0$. When c increases, $y(c)$ increases and $\bar{y}(c)$ decreases. When $\bar{y}(c)$ reaches the value $y_0(v_0)$, $y(c)$ is larger than $y_0(v_0)$.

Consider the function

$$\psi(v_0, c) = y_c(v_0) + \frac{c}{\Lambda}(1 - v_0).$$

To prove the existence and uniqueness of $c(v_0)$, we just need to apply the Intermediate Value Theorem to the increasing map $c \mapsto \psi(v_0, c)$, which rises from a negative value for $c = 0$ to a positive one when c is large enough.

We also use the function ψ to prove the continuity of the map $v_0 \mapsto c(v_0)$. To this end, we take any $v_0^0 \in (0, 1)$ and any $\varepsilon > 0$ such that $[c(v_0^0) - \frac{\varepsilon}{2}, c(v_0^0) + \frac{\varepsilon}{2}] \subset \mathbb{R}^+$. We have that

$$\psi\left(v_0^0, c(v_0^0) - \frac{\varepsilon}{2}\right) < \psi\left(v_0^0, c(v_0^0)\right) = 0 < \psi\left(v_0^0, c(v_0^0) + \frac{\varepsilon}{2}\right).$$

As the map $v_0 \mapsto \psi(v_0, c)$ is continuous by Lemma 5.3, a $\delta(\varepsilon) > 0$ exists, such that, if $|v_0 - v_0^0| < \delta(\varepsilon)$, we have that $\psi\left(v_0, c(v_0^0) - \frac{\varepsilon}{2}\right) < 0$ and $\psi\left(v_0, c(v_0^0) + \frac{\varepsilon}{2}\right) > 0$. Then, as a consequence of the Intermediate Value Theorem, the unique solution $c(v_0)$ of the equation $\psi(v_0, c) = 0$ is contained in the open interval $(c(v_0^0) - \frac{\varepsilon}{2}, c(v_0^0) + \frac{\varepsilon}{2})$. As ε can be taken arbitrarily small, this completes the proof of the continuity of $c(v_0)$ at the point v_0^0 . The continuity of the map $v_0 \mapsto y_{c(v_0)}(v_0)$ follows from the continuity of the map $c(v_0)$ and the continuity of the map $(c, v_0) \mapsto y_c(v_0)$ proved in Lemma 5.3. \square

6.2. Estimate of $c(v_0)$. To estimate $c(v_0)$, it is interesting to introduce the control line l_c defined by (5.3) which we have already used in the previous subsection. We now consider the motions of the three points $y(c) = y_c(v_0)$, $\tilde{y}(c) = \tilde{y}_c(v_0)$ and $\bar{y}(c)$. We have that $y(c) < \tilde{y}(c)$. Recall that $\bar{y}(c)$ is decreasing and $y(c)$, $\tilde{y}(c)$ are increasing, with $y(c) < \tilde{y}(c)$ for $c > 0$.

At $c = 0$ we have the inequality $y(0) = \tilde{y}(0) < \bar{y}(0) = 0$. Subsequently, $\bar{y}(c)$ crosses $\tilde{y}(c)$ at a value $c = c_-(v_0)$ at which it holds $y(c_-(v_0)) < \tilde{y}(c_-(v_0)) = \bar{y}(c_-(v_0))$. When c eventually reaches the value c_+ where $\bar{y}(c_+) = y_0(v_0)$, we have the inequality $\bar{y}(c_+) < y(c_+) < \tilde{y}(c_+)$. This yields the following bounds for $c(v_0)$:

$$c_-(v_0) < c(v_0) < c_+(v_0), \quad (6.3)$$

where $c_+ = c_+(v_0)$ is given by

$$-\frac{c_+}{\Lambda}(1 - v_0) = -\left(\frac{2}{(1 + \alpha)\Lambda}\right)^{1/2} v_0^{\frac{1+\alpha}{2}}, \quad (6.4)$$

and $c_- = c_-(v_0)$ by

$$-\frac{c_-}{\Lambda}(1 - v_0) = -\left(\frac{2}{(1 + \alpha)\Lambda}\right)^{1/2} v_0^{\frac{1+\alpha}{2}} + \frac{c_-}{\Lambda} v_0. \quad (6.5)$$

We find that $c_+(v_0) = \left(\frac{2\Lambda}{1+\alpha}\right)^{1/2} \frac{v_0^{\frac{1+\alpha}{2}}}{1-v_0}$ and $c_-(v_0) = \left(\frac{2\Lambda}{1+\alpha}\right)^{1/2} v_0^{\frac{1+\alpha}{2}}$. These values coincide with the a priori estimates (3.4) and (3.6) obtained by a different method in Subsection 3.1.

Finally, we observe that Lemma 5.5 provides us with the following new estimate:

$$\frac{c}{\Lambda}(v_0 - 1) > -\frac{1}{c} v_0^\alpha \quad \text{or equivalently} \quad c(v_0) < \left(\frac{\Lambda}{1 - v_0}\right)^{1/2} v_0^{\frac{\alpha}{2}}. \quad (6.6)$$

It is readily seen that $\left(\frac{\Lambda}{1 - v_0}\right)^{1/2} v_0^{\frac{\alpha}{2}} > c_-(v_0) = \left(\frac{2\Lambda}{1 + \alpha}\right)^{1/2} v_0^{\frac{1+\alpha}{2}}$, because it holds $v_0(1 - v_0) < (1 + \alpha)/2$ as $0 < v_0 < 1$ and $0 < \alpha < 1$. Therefore, combining (6.6) with the bounds (6.3) reveals that

$$c_-(v_0) < c(v_0) < \min\left(c_+(v_0), \left(\frac{\Lambda}{1 - v_0}\right)^{1/2} v_0^{\frac{\alpha}{2}}\right). \quad (6.7)$$

6.3. Estimations for the time $R(v_0)$. Taking into account the existence and uniqueness of the stable manifold $L_c := \{y = y_c\}$ obtained in Subsection 5.1 (see Lemma 5.3), it remains to prove that the settling time $R(v_0)$, along the stable manifold, to proceed from the initial condition $(v_0, y_{c(v_0)}(v_0))$ to the origin is finite.

We first consider the settling time $T(x, c)$ to go from $(x, y_c(x))$ to the origin, along the stable manifold L_c of X_c .

Lemma 6.2. *The settling time $T(x, c)$ is finite for any $(x, c) \in \mathbb{R}_+^2 = [0, +\infty) \times [0, +\infty)$. Moreover, $T(x, c)$ is continuous on \mathbb{R}_+^2 and such that $T(0, c) = 0$. In the interval $0 < x < x_1(c) = \left(\frac{2\Lambda}{(1 + \alpha)c^2}\right)^{\frac{1}{1-\alpha}}$, we have the following bounds for $T(x, c)$*

$$\frac{(2(1 + \alpha)\Lambda)^{1/2}}{1 - \alpha} x^{\frac{1-\alpha}{2}} < T(x, c) < \frac{(2(1 + \alpha)\Lambda)^{1/2}}{1 - \alpha} x^{\frac{1-\alpha}{2}} \left(1 - c \left(\frac{1 + \alpha}{2\Lambda}\right)^{1/2} x^{\frac{1-\alpha}{2}}\right)^{-1}, \quad (6.8)$$

which gives the following asymptotic expansion at $(0, c)$:

$$T(x, c) = \frac{(2(1+\alpha)\Lambda)^{1/2}}{1-\alpha} x^{\frac{1-\alpha}{2}} + O(cx^{1-\alpha}), \quad (6.9)$$

where the remainder is uniform in c when c is restricted to a compact subset of $[0, +\infty)$.

Proof. The settling time $T(x, c)$ is obtained by integrating the equation $dt = \frac{dx}{y_c(x)}$ from x to 0. Changing the sense of integration, we arrive that $T(x, c) = \int_0^x \frac{ds}{|y_c(s)|}$. This integral is improper at $x = 0$, but as by (5.2) the function $|y_c(x)|$ is equivalent to $\left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} x^{\frac{1+\alpha}{2}}$, with $\frac{1+\alpha}{2} < 1$, the integral converges and is continuous in x . To have continuity in (x, c) , we require the above equivalence to be uniform in c when c belongs to a compact subset of $[0, +\infty)$.

We have that

$$\left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} s^{\frac{1+\alpha}{2}} - \frac{c}{\Lambda} s < |y_c(s)| < \left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} s^{\frac{1+\alpha}{2}}. \quad (6.10)$$

Moreover, if $s \leq x < x_1(c) = \left(\frac{2\Lambda}{(1+\alpha)c^2}\right)^{\frac{1}{1-\alpha}}$, we have:

$$\left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} s^{\frac{1+\alpha}{2}} - \frac{c}{\Lambda} s > \left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} s^{\frac{1+\alpha}{2}} \left(1 - c\left(\frac{1+\alpha}{2\Lambda}\right)^{1/2}\right) x^{\frac{1-\alpha}{2}}.$$

As $\int_0^x s^{-\frac{1+\alpha}{2}} ds = \frac{2}{1-\alpha} x^{\frac{1-\alpha}{2}}$, the last inequality and (6.10) implies the estimates (6.8). The asymptotic expansion follows directly from (6.8). \square

We now consider the time $R(v_0) = T(v_0, c(v_0))$ corresponding to the initial conditions (6.1). A direct consequence of Lemma 6.2 and the continuity of the mapping $v_0 \mapsto c(v_0)$ is the following:

Lemma 6.3. *The time $R(v_0)$ to proceed from $\left(v_0, -\frac{c}{\Lambda}(1-v_0)\right)$ to the origin, along the stable manifold of $X_{c(v_0)}$, is finite for any $v_0 \in [0, 1)$. Moreover, $R(v_0)$ is continuous on $[0, 1)$ and $R(0) = 0$.*

Proof. As $c(v_0) < c_+(v_0) = \left(\frac{2\Lambda}{1+\alpha}\right)^{1/2} \frac{v_0^{\frac{1+\alpha}{2}}}{1-v_0}$, we have that $c(v_0) = O\left(v_0^{\frac{1+\alpha}{2}}\right)$. Then, $c(v_0)$ is continuous on $[0, 1)$ with $c(0) = 0$. It follows that $R(v_0)$ is continuous on $[0, 1)$ with the value $R(0) = 0$. \square

As the curve L_c is above the curve L_0 , the inequality $\frac{(2(1+\alpha)\Lambda)^{1/2}}{1-\alpha} x^{\frac{1-\alpha}{2}} < T(x, c)$ in (6.8) is in fact true for all $x > 0$. Then we have the following lower bound for $R(v_0)$:

$$R(v_0) > \frac{(2(1+\alpha)\Lambda)^{1/2}}{1-\alpha} v_0^{\frac{1-\alpha}{2}} \quad (6.11)$$

for $0 < v_0 < 1$ and $0 \leq \alpha < 1$.

To obtain an upper bound for $R(v_0)$ which would be valid for any $0 < v_0 < 1$, it would be better to replace the curve $l_{c(v_0)}$ by one which remains below $\{y = 0\}$. The simplest solution is a

curve $d_k := \{y = \bar{y}_k = -kx^{\frac{1+\alpha}{2}}\}$, with $k = k(v_0)$ chosen so that the vector field $X_{c(v_0)}$ is upward transverse to d_k , at least for $x \in (0, v_0]$. A first condition on k is clearly d_k above L_0 , i.e.

$$k^2 < \frac{2}{(1+\alpha)\Lambda}. \quad (6.12)$$

To express the transversality condition, we write the vector field X_c along d_k :

$$X(x) = \left(-kx^{\frac{1+\alpha}{2}}, -k\frac{c}{\Lambda}x^{\frac{1+\alpha}{2}} + \frac{1}{\Lambda}x^\alpha\right),$$

and the orthogonal vector field to d_k in the upward direction, given by :

$$N(x) = \left(\frac{1+\alpha}{2}kx^{\frac{\alpha-1}{2}}, 1\right).$$

Then, we compute their scalar product :

$$\langle X(x), N(x) \rangle = \left(\frac{1}{\Lambda} - \frac{1+\alpha}{2}k^2\right)x^\alpha - k\frac{c}{\Lambda}x^{\frac{1+\alpha}{2}}.$$

Upon factoring, it comes:

$$\langle X(x), N(x) \rangle = \left(\frac{1}{\Lambda} - \frac{1+\alpha}{2}k^2\right)x^\alpha \left[1 - k\frac{c}{\Lambda}\left(\frac{1}{\Lambda} - \frac{1+\alpha}{2}k^2\right)^{-1}x^{\frac{1-\alpha}{2}}\right].$$

The condition (6.12) implies that the factor $\left(\frac{1}{\Lambda} - \frac{1+\alpha}{2}k^2\right)$ is positive. Then, for the scalar product to be positive for $c = c(v_0) < c_+(v_0)$ and $x \leq v_0$, it is sufficient that the following condition be fulfilled:

$$k\frac{c_+(v_0)}{\Lambda}\left(\frac{1}{\Lambda} - \frac{1+\alpha}{2}k^2\right)^{-1}v_0^{\frac{1-\alpha}{2}} < 1.$$

Taking into account the expression of $c_+(v_0)$ this condition reads as follows:

$$k\left(\frac{2\Lambda}{1+\alpha}\right)^{1/2} < \frac{1-v_0}{v_0}\left(1 - \frac{(1+\alpha)\Lambda}{2}k^2\right). \quad (6.13)$$

From now on, we also assume that

$$k \leq \left(\frac{1}{(1+\alpha)\Lambda}\right)^{1/2}, \quad (6.14)$$

which implies the condition (6.12). As $\frac{1}{v_0} > 1$, to have (6.13), assuming that condition (6.14) is fulfilled, it suffices to have the following condition:

$$k \leq \frac{1}{2}\left(\frac{1+\alpha}{2\Lambda}\right)^{1/2}(1-v_0). \quad (6.15)$$

As

$$A(\alpha) = \frac{2^{3/2}}{(1+\alpha)^{1/2}} > (1+\alpha)^{1/2}, \quad (6.16)$$

it is easy to see that conditions (6.14) and (6.15) are verified if

$$k = \frac{1-v_0}{A(\alpha)\Lambda^{1/2}}. \quad (6.17)$$

For this value of k , the vector field $X_{c(v_0)}$ is transverse in the upward direction along the curve d_k for all $x \in (0, v_0]$. Accordingly, we construct a trapping triangle, which implies that, for all

$x \in (0, v_0]$, it holds $y_{c(v_0)} < \bar{y}_k(x) = -kx^{\frac{1+\alpha}{2}}$. This implies that $R(v_0)$ has an upper bound equal to $\frac{1}{k} \frac{1}{1-\alpha} v_0^{\frac{1-\alpha}{2}}$ for all $v_0 \in [0, 1)$. Thanks to (6.11) and (6.17), we have the following bounds:

Lemma 6.4. *For $0 < v_0 < 1$ and $0 \leq \alpha < 1$, we have that*

$$\frac{(2(1+\alpha)\Lambda)^{1/2}}{1-\alpha} v_0^{\frac{1-\alpha}{2}} < R(v_0) < \frac{2\Lambda^{1/2}A(\alpha)}{1-\alpha} \frac{v_0^{\frac{1-\alpha}{2}}}{1-v_0}, \quad (6.18)$$

where $A(\alpha)$ is defined by (6.16).

6.4. End of the proof of Theorem 3.1 and Corollary 3.2. To complete the proof of Theorem 3.1, the two following points remain to be established. We return to the original notation of the free boundary problem (3.1), see (3.14).

Lemma 6.5. *For $0 < \alpha < 1$, it holds $v \in C^\infty([0, R]) \cap C^{2+[\beta], \beta-[\beta]}([0, R])$, $\beta = \frac{2\alpha}{1-\alpha}$.*

Proof. It is clear that v is smooth as long as it does not vanish. We need to establish the Hölder regularity of the function $v(\xi)$ near $\xi = R$. It follows from Subsection 5.2 that near the origin, it holds

$$y_c(x) \sim -\left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} x^{\frac{1+\alpha}{2}}, \quad (6.19)$$

that is, near $\xi = R_-$,

$$v'(\xi) \sim -\left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} v^{\frac{1+\alpha}{2}}(\xi). \quad (6.20)$$

An elementary integration yields

$$v(\xi) \sim \frac{1-\alpha}{2} \left(\frac{2}{(1+\alpha)\Lambda}\right)^{1/2} (R-\xi)^{\frac{2}{1-\alpha}}, \quad \frac{2}{1-\alpha} = 2 + \beta. \quad (6.21)$$

□

Lemma 6.6. *The function $v(\xi)$ is decreasing, convex on the interval $[0, R)$.*

Proof. On one hand, it follows from Lemma 5.5 that the stable manifold y_c never cuts the curve $y = -\frac{1}{c}x^\alpha$, that is, y'_c does not vanish. In other words, $v(\xi)$ has no inflection points. On the other hand, Lemma 6.5 yields that $v(\xi)$ is convex near $\xi = R$, $\xi < R$, hence the convexity of $v(\xi)$ on $(0, R)$. □

Finally, we prove Corollary 3.2. Let $0 < a < b < 1$, $I = [a, b]$. According to Lemma 6.1 and Lemma 6.3, respectively, the mappings $v_0 \mapsto c(v_0)$ and $v_0 \mapsto R(v_0)$ are continuous from $I \rightarrow \mathbb{R}$. Thanks to Lemma 6.4, we have the uniform estimate:

$$\frac{(2(1+\alpha)\Lambda)^{1/2}}{1-\alpha} a^{\frac{1-\alpha}{2}} < R(v_0) \leq R_{max} = \frac{2\Lambda^{1/2}A(\alpha)}{1-\alpha} \frac{b^{\frac{1-\alpha}{2}}}{1-a}. \quad (6.22)$$

Let $\tilde{v}(v_0)$ be the extension by 0 of the function $\xi \mapsto v(v_0; \xi)$ to the interval $[0, R_{max}]$, clearly $\tilde{v}(v_0) \in C^1([0, R_{max}])$.

Lemma 6.7. *The mapping $v_0 \mapsto \tilde{v}(v_0)$ is continuous from I to $C^1([0, R_{max}])$.*

Proof. The mapping $\tilde{v}(v_0)$ can be seen as the function of two variables $\tilde{v}(v_0; \xi)$. The mapping $v_0 \mapsto \tilde{v}(v_0)$ is continuous from $I \rightarrow C^1([0, R_{max}])$ if and only if $\tilde{v}(v_0; \xi)$ is continuous on the rectangle $T = [0, R_{max}] \times I$, which is cut by the curve $l_R := v_0 \rightarrow R(v_0)$ into two closed subsets $T_1 = \{v_0 \in I, 0 \leq \xi \leq R(v_0)\}$ and $T_2 = \{v_0 \in I, R(v_0) \leq \xi \leq R_{max}\}$. The function $\tilde{v}(v_0; \xi)$ is continuous on T_2 where it is identically 0 and on T_1 as it is equal to the x -component $x(\xi, (v_0, y_{c(v_0)}(v_0)))$ of the trajectory of $X_{c(v_0)}$ with initial conditions $(v_0, y_{c(v_0)}(v_0))$. These two definitions have the same value 0 along $l_R = T_1 \cup T_2$. Then, they define a continuous function on T . \square

7. THE FREE BOUNDARY PROBLEM AS AN INTEGRO-DIFFERENTIAL SYSTEM

Here, we revisit the one-phase free boundary problem (2.1)-(2.3) that we rewrite in the form of two coupled subsystems:

$$\begin{cases} \Lambda v'' - cv' = v^\alpha, & 0 < \xi < R, \\ 0 < v(0) < 1, v'(0) = -\frac{c}{\Lambda}(1 - v(0)), \\ v(R) = v'(R) = 0, \end{cases} \quad (7.1)$$

and

$$\begin{cases} u'' - cu' = -v^\alpha, & 0 < \xi < R, \\ u(0) = \theta, u'(0) = c\theta, \\ u(R) = 1, u'(R) = 0. \end{cases} \quad (7.2)$$

Remark 7.1. *A first sight, subsystems (7.1) and (7.2) seem uncoupled because v appears in the right hand side of (7.2). However, the linkage is through c (see the explicit computation in Subsection 8.2 when $\Lambda = 0$).*

7.1. Equivalence with an integro-differential system. We have the following proposition.

Proposition 7.2. *Assume there is a solution $(c, R, u(\xi), v(\xi))$ to system (7.1)-(7.2) such that $c > 0$, $0 < R < +\infty$, u and v in $C^\infty([0, R]) \cap C^1([0, R])$. Then, $(c, R, v(\xi))$ verifies the system consisting of (7.1) and one of the following integral equations:*

$$\theta + \Lambda v(0) = 1 - c(1 - \Lambda) \int_0^R e^{-cs} v(s) ds, \quad (7.3)$$

$$\theta + v(0) = 1 - (1 - \Lambda) \int_0^R e^{-cs} v'(s) ds. \quad (7.4)$$

Furthermore, the three systems (7.1)-(7.2), (7.1)&mathcal{E}(7.3) and (7.1)&mathcal{E}(7.4) are equivalent.

Proof. (i) Let $(c, R, u(\xi), v(\xi))$ be a solution to (7.1)-(7.2) as above. Set $w = u + v$, one infers from system (7.1)-(7.2) that w satisfies

$$\begin{cases} w'' - cw' = (1 - \Lambda)v'', & 0 < \xi < R, \\ w(R) = 1, w'(R) = 0. \end{cases} \quad (7.5)$$

For any $\xi \in (0, R)$, we integrate (7.5) from ξ to R . Using the free boundary conditions in (7.1) and in (7.5), we arrive at

$$w' - cw = (1 - \Lambda)v' - c, \quad w(R) = 1. \quad (7.6)$$

It is easy to solve (7.6) and obtain:

$$w(\xi) = e^{-c(R-\xi)} - \int_{\xi}^R e^{-c(s-\xi)} ((1-\Lambda)v'(s) - c) ds. \quad (7.7)$$

Taking $\xi = 0$ in (7.7), it is not difficult to recover (7.4) using the boundary conditions in (7.1) and (7.2). Likewise, setting $\tilde{w} = u + \Lambda v$, one proceeds in the same manner to obtain (7.3) (or equivalently we may integrate by part the integral in (7.4)).

(ii) Assume that $(c, R, v(\xi))$ verifies (7.1), $v \in C^\infty([0, R]) \cap C^1([0, R])$, and in addition (7.4) holds. We construct a unique solution to (7.5) as in (7.7). Next, setting $u = w - v$, u is in $C^\infty([0, R]) \cap C^1([0, R])$ and satisfies the equation in (7.2). From the free boundary conditions in (7.1) and (7.5), we obtain $u(R) = 1, u'(R) = 0$. Letting $\xi = 0$ in (7.7), one has

$$\begin{aligned} w(0) &= e^{-cR} - (1-\Lambda) \int_0^R e^{-cs} v'(s) ds + \int_0^R ce^{-cs} ds \\ &= 1 - (1-\Lambda) \int_0^R e^{-cs} v'(s) ds, \end{aligned} \quad (7.8)$$

which together with (7.4) gives $u(0) = \theta$. Finally, we easily retrieve $u'(0) = c\theta$. \square

Remark 7.3. $w = u + v$ and $\tilde{w} = u + \Lambda v$ are the only linear relations between u and v (see [6, Remark 8.5]). The physical meaning of w is the normalized enthalpy (see, e.g., [12]). In the equidiffusion case, i.e. $\Lambda = 1$, $w \equiv 1$ is a Shvab-Zeldovich variable which eliminates $u = 1 - v$ (see Subsection 8.1).

7.2. Existence of a solution. Based on Proposition 7.2, we need to focus only on system (7.1)&(7.3) or (7.1)&(7.4) to study system (7.1)-(7.2). More specifically, we will prove the existence of a solution via a fixed point method.

Let $a < b$ and $I = [a, b] \subset \mathbb{R}$. We denote by P the projection from \mathbb{R} to I defined by $P(x) = x$ if $x \in \text{int}I$, $P(x) = a$ if $x \leq a$ and $P(x) = b$ if $x \geq b$. Obviously, P is a continuous mapping.

7.2.1. Case $0 < \Lambda < 1$. We first consider the case where $0 < \Lambda < 1$ and prove by a fixed point argument that there exists a solution to system (7.1)&(7.4).

For $0 < \theta < 1, 0 < \Lambda < 1$, we set $I = [1 - \theta, 1 - \Lambda\theta]$. For $v_0 \in I$, let $(c(v_0), R(v_0), v(v_0; \xi))$ be the unique solution of system (3.1) given by Theorem 3.1. We define the mapping $\Phi : I \rightarrow I$ by

$$\Phi(v_0) = P \left(1 - \theta + (1 - \Lambda) \int_0^{R(v_0)} e^{-c(v_0)s} (-v'(v_0; s)) ds \right), \quad (7.9)$$

where P is the projection from \mathbb{R} to I as above.

Theorem 7.4. *Let $0 < \theta < 1, 0 < \Lambda < 1$ and $I = [1 - \theta, 1 - \Lambda\theta]$. Then, the mapping $\Phi : I \rightarrow I$ defined by (7.9) has a fixed point v_0^* which solves (7.1)&(7.4) with $v(0) = v_0^*$.*

Proof. It follows from Corollary 3.2 that the mapping $\Phi : I \rightarrow I$ is continuous, hence it has a fixed point v_0^* . We denote by $c^* = c(v_0^*), R^* = R(v_0^*)$ and $v^*(\xi) = v(v_0^*, \xi)$ the solution to (3.1) associated with v_0^* . We must prove that (7.4) holds, namely

$$v_0^* = 1 - \theta + (1 - \Lambda) \int_0^{R^*} e^{-c^*s} (-v^{*'}(s)) ds. \quad (7.10)$$

From the definition of the projection P on I , there are 3 cases:

(i) If

$$1 - \theta + (1 - \Lambda) \int_0^{R^*} e^{-c^*s}(-v^{*'}(s)) ds \in (1 - \theta, 1 - \Lambda\theta),$$

then it holds that $v_0^* \in (1 - \theta, 1 - \Lambda\theta)$ and P is the identity. Therefore, (7.10) is satisfied.

(ii) If

$$1 - \theta + (1 - \Lambda) \int_0^{R^*} e^{-c^*s}(-v^{*'}(s)) ds \leq 1 - \theta,$$

it comes obviously

$$(1 - \Lambda) \int_0^{R^*} e^{-c^*s}(-v^{*'}(s)) ds \leq 0,$$

which is in contradiction to Theorem 3.1 which states that $v^{*'}(\xi) < 0$, for all $\xi \in [0, R^*)$. Then, case (ii) is ruled out.

(iii) Finally, let us consider the case where

$$1 - \theta + (1 - \Lambda) \int_0^{R^*} e^{-c^*s}(-v^{*'}(s)) ds \geq 1 - \Lambda\theta, \quad (7.11)$$

hence $v_0^* = 1 - \Lambda\theta$. It follows from Theorem 3.1 that v^* is decreasing and convex, hence $|v^{*'}(\xi)| \leq |v^{*'}(0)|$, $\forall \xi \in [0, R^*)$, therefore:

$$\theta \leq \int_0^{R^*} e^{-cs}(-v^{*'}(s)) ds \leq \int_0^{R^*} \frac{c^*}{\Lambda}(1 - v_0^*)e^{-c^*s} ds < \frac{1 - v_0^*}{\Lambda}. \quad (7.12)$$

It comes $\Lambda\theta < 1 - v_0^*$ which is in contradiction to $v_0^* = 1 - \Lambda\theta$. Then, case (iii) is also ruled out. \square

7.2.2. Case $\Lambda > 1$. Likewise, we prove via a fixed point argument that there exists a solution to system (7.1)&(7.3) in the case $\Lambda > 1$. Here, we take $I = [(1 - \theta)/\Lambda, 1 - \theta/\Lambda]$. This time, we define the mapping $\Psi : I \rightarrow I$ by

$$\Psi(v_0) = P \left(\frac{1}{\Lambda} \left[1 - \theta - c(v_0)(1 - \Lambda) \int_0^{R(v_0)} e^{-c(v_0)s} v(v_0; s) ds \right] \right). \quad (7.13)$$

Theorem 7.5. *Let $0 < \theta < 1$, $\Lambda > 1$ and $I = [(1 - \theta)/\Lambda, 1 - \theta/\Lambda]$. Then, the mapping $\Psi : I \rightarrow I$ defined by (7.13) has a fixed point v_0^* which solves (7.1)&(7.4) with $v(0) = v_0^*$.*

Proof. The proof follows the same pattern as the proof of Theorem 7.4. As before, we denote by (c^*, R^*, v^*) the solution to (3.1) associated with v_0^* . This time, we must prove that (7.3) holds, which is equivalent to

$$v_0^* = \frac{1}{\Lambda} \left[1 - \theta - c^*(1 - \Lambda) \int_0^{R^*} e^{-c^*s} v^*(s) ds \right]. \quad (7.14)$$

(i) If

$$\frac{1}{\Lambda} \left[1 - \theta - c^*(1 - \Lambda) \int_0^{R^*} e^{-c^*s} v^*(s) ds \right] \in \left(\frac{1 - \theta}{\Lambda}, 1 - \frac{\theta}{\Lambda} \right),$$

then it holds that $v_0^* \in (\frac{1 - \theta}{\Lambda}, 1 - \frac{\theta}{\Lambda})$ and P is the identity. Therefore, (7.14) is satisfied.

(ii) If

$$\frac{1}{\Lambda} \left[1 - \theta - c^*(1 - \Lambda) \int_0^{R^*} e^{-c^*s} v^*(s) ds \right] \leq \frac{1 - \theta}{\Lambda},$$

it comes

$$c^*(\Lambda - 1) \int_0^{R^*} e^{-c^*s} v^*(s) ds \leq 0,$$

in contradiction to Theorem 3.1 which states that $v^*(\xi) > 0$, for all $\xi \in [0, R^*)$. Case (ii) is then ruled out.

(iii) If

$$\frac{1}{\Lambda} \left[1 - \theta - c(1 - \Lambda) \int_0^{R^*} e^{-c^*s} v^*(s) ds \right] \geq 1 - \frac{\theta}{\Lambda}, \quad (7.15)$$

then it holds $v_0^* = 1 - \frac{\theta}{\Lambda}$. From (7.15), we infer that

$$c \int_0^{R^*} e^{-c^*s} v^*(s) ds \geq 1. \quad (7.16)$$

On the other hand, noticing from Theorem 3.1 that $v^{*'}(\xi) < 0$, for all $\xi \in [0, R^*)$, it comes

$$c^* \int_0^{R^*} e^{-c^*s} v^*(s) ds < c^* \int_0^{R^*} e^{-c^*s} v^*(0) ds = c^* \int_0^{R^*} e^{-c^*s} \left(1 - \frac{\theta}{\Lambda}\right) ds = \left(1 - \frac{\theta}{\Lambda}\right) (1 - e^{-c^*R^*}) < 1,$$

in contradiction to (7.15). □

7.3. Final results. We are now in position to summarize the results of Subsections 7.1 and 7.2 for $0 < \Lambda < 1$ and $\Lambda > 1$ (see Subsection 8.1 for $\Lambda = 1$).

Theorem 7.6. *Let $0 < \alpha < 1$, $0 < \theta < 1$ and $\Lambda > 0$ be fixed. We define the interval $I \subset (0, 1)$ as: $I = [1 - \theta, 1 - \Lambda\theta]$ if $0 < \Lambda < 1$, $I = [(1 - \theta)/\Lambda, 1 - \theta/\Lambda]$ if $\Lambda > 1$. Then, the one-phase free boundary problem (2.1)-(2.3) has a solution $(c, R, u(\xi), v(\xi))$ such that $c > 0$, $0 < R < +\infty$, $v(0) \in I$, u and v in $C^\infty([0, R]) \cap C^{2+[\beta], \beta-[\beta]}([0, R])$, $\beta = 2\alpha/(1 - \alpha)$.*

Proof. According to the value of Λ against 1, Theorem 7.4 and Theorem 7.5 provide the existence of a solution to system (7.1)&(7.4), or to system (7.1)&(7.3), such that $v(0) \in I$. We proved in Proposition 7.2 that these systems are equivalent to the free boundary problem (2.1)-(2.3).

Next, we apply Theorem 3.1 which gives the (optimal) Hölder regularity of $v(\xi)$ at the free boundary R . More precisely, this result follows from Lemma 6.5 which is related to the asymptotic expansion of the stable manifold near the origin. Thus, the same regularity result holds for $u(\xi)$. □

Finally, we are able to prove the existence of a solution to the free interface problem (1.16)-(1.18) thanks to Proposition 2.1. The last part of the proof of Theorem 1.1 concerning the limit cases where $\alpha \rightarrow 1$ and $\alpha \rightarrow 0$ is deferred to Subsections 8.3 and 8.4.

8. LIMIT CASES

8.1. Equidiffusion. In the case $\Lambda = 1$ (Lewis number equal to unity), equation (7.5) yields that the normalized enthalpy $w = u + v$ is equal to 1; hence $u = 1 - v$. The system for v reads as follows:

$$\begin{cases} v'' - cv' = v^\alpha, & 0 < \xi < R, \\ v(0) = 1 - \theta, & v'(0) = -c\theta, \\ v(R) = v'(R) = 0. \end{cases} \quad (8.1)$$

It follows from Theorem 3.1, taking $v_0 = 1 - \theta$, that system (8.1) has a unique solution $(c, R, v(\xi))$.

8.2. Infinite Lewis number. Here, we consider the limit case $\Lambda = 0$, namely an infinite Lewis number. In this case $v(0) = 1$. Integrating $cv' + v^\alpha = 0$ yields explicitly $R = c/(1 - \alpha)$ and

$$v(\xi) = \left(1 - \xi \frac{1 - \alpha}{c}\right)^{\frac{1}{1 - \alpha}}.$$

It comes the formula

$$\theta = 1 - c \int_0^{\frac{c}{1 - \alpha}} e^{-cs} \left(1 - s \frac{1 - \alpha}{c}\right)^{\frac{1}{1 - \alpha}} ds, \quad (8.2)$$

which coincides with (7.3) when $\Lambda = 0$.

Example 8.1. (i) In the case $\alpha = 0$, the formula (8.2) yields (see (1.12))

$$\theta = \frac{1 - e^{-c^2}}{c^2}. \quad (8.3)$$

(ii) In the case $\alpha = 1/2$, the formula becomes

$$\theta = \frac{1}{2}c^{-4}e^{-2c^2} - c^{-2} + \frac{1}{2}c^{-4}. \quad (8.4)$$

However, for general $\alpha \in (0, 1)$, the integral in (8.2) can not be computed explicitly. For $c > 0$, $0 < \alpha < 1$ and $0 < \theta < 1$, let us define:

$$f(c, \alpha, \theta) := \theta - 1 + c \int_0^{\frac{c}{1 - \alpha}} e^{-cs} \left(1 - s \frac{1 - \alpha}{c}\right)^{\frac{1}{1 - \alpha}} ds. \quad (8.5)$$

We have

Lemma 8.2. For fixed $0 < \alpha < 1$ and $0 < \theta < 1$, the equation $f(c, \alpha, \theta) = 0$ has a unique positive root.

Proof. (i) We first look for the existence of a root $c > 0$. It is easy to check that

$$\lim_{c \rightarrow 0^+} f(c, \alpha, \theta) = \theta - 1. \quad (8.6)$$

Next, let us prove that

$$\lim_{c \rightarrow +\infty} f(c, \alpha, \theta) = \theta. \quad (8.7)$$

Indeed, with the change of variable $y = 1 - s(1 - \alpha)/c$, (8.5) becomes

$$f(c, \alpha, \theta) = \theta - 1 + \frac{c^2}{1 - \alpha} \int_0^1 e^{-\frac{c^2(1-y)}{1-\alpha}} y^{\frac{1}{1-\alpha}} dy. \quad (8.8)$$

Next, let $z = c^2 y / (1 - \alpha)$. It comes:

$$f(c, \alpha, \theta) = \theta - 1 + \left(\frac{1 - \alpha}{c^2} \right)^{\frac{1}{1-\alpha}} e^{-\frac{c^2}{1-\alpha}} \int_0^{\frac{c^2}{1-\alpha}} e^z z^{\frac{1}{1-\alpha}} dz.$$

By using L'Hospital's rule, one obtains

$$\lim_{c \rightarrow +\infty} \left(\frac{1 - \alpha}{c^2} \right)^{\frac{1}{1-\alpha}} e^{-\frac{c^2}{1-\alpha}} \int_0^{\frac{c^2}{1-\alpha}} e^z z^{\frac{1}{1-\alpha}} dz = 1.$$

Thus, (8.7) holds.

(ii) Next, to prove the uniqueness of a positive root, we compute the sign of the derivative $f_c(c, \alpha, \theta)$. To this end, we make the convenient change of variable $z = c^2(1 - y)/(1 - \alpha)$ in (8.8):

$$f(c, \alpha, \theta) = \theta - 1 + \int_0^{\frac{c^2}{1-\alpha}} e^{-z} \left[1 - \frac{(1 - \alpha)z}{c^2} \right]^{\frac{1}{1-\alpha}} dz.$$

Then,

$$f_c(c, \alpha, \theta) = \int_0^{\frac{c^2}{1-\alpha}} e^{-z} \left[1 - \frac{(1 - \alpha)z}{c^2} \right]^{\frac{\alpha}{1-\alpha}} \frac{2z}{c^3} dz > 0. \quad (8.9)$$

□

8.3. Limit $\alpha \rightarrow 1$. Here, $\Lambda > 0$ and $\theta \in (0, 1)$ are fixed. With each α , $0 < \alpha < 1$, we associate a solution $(c, R, u(\xi), v(\xi))$ of the free boundary problem (2.1)-(2.3). However, we recall that problem (2.1)-(2.3) is equivalent to the system

$$\begin{cases} \Lambda v'' - cv' = v^\alpha, & \xi > 0, \\ v(0) \in I, v'(0) = -\frac{c}{\Lambda}(1 - v(0)), \\ v(R) = v'(R) = 0, \end{cases} \quad (8.10)$$

together with

$$\theta + \Lambda v(0) = 1 - c(1 - \Lambda) \int_0^R e^{-cs} v(s) ds, \quad (8.11)$$

where the interval $I = [a, b]$ is given in Theorem 7.4 or 7.5, according to the value of Λ , and depends only on Λ and θ .

Lemma 8.3. *Let $\Lambda > 0$ and $0 < \theta < 1$ be fixed. Let $(\alpha_n)_{n \in \mathbb{N}}$ be an increasing sequence such that $0 < \alpha_n < 1$, $\alpha_n \rightarrow 1$ as $n \rightarrow \infty$; the corresponding solution of (8.10)-(8.11) is denoted by $(c_n, R_n, v_n(\xi))$. Then, $R_n \rightarrow +\infty$ as $n \rightarrow \infty$.*

Proof. Assume by contradiction that the sequence $(R_n)_{n \in \mathbb{N}}$ is bounded, namely there exists $A > 0$ such that $R_n \leq A$ for all $n \in \mathbb{N}$. In view of $0 \leq v_n(\xi) < 1$ for all $0 \leq \xi \leq R_n$ and formula (2.4), it holds

$$c_n = \int_0^{R_n} v_n^{\alpha_n}(\xi) d\xi,$$

hence $c_n \leq R_n \leq A$. In addition, according to (3.7),

$$c_n^2 \geq \frac{2\Lambda}{1 + \alpha_n} v_n^{1+\alpha_n}(0),$$

$v_n(0) \geq a = \inf I > 0$. Hence, the sequence $(c_n)_{n \in \mathbb{N}}$ is bounded from below by $a\sqrt{\Lambda}$.

By abuse of notation, we also denote by $v_n(\xi)$ its extension by 0 to the interval $[0, A]$. The system for $(c_n, v_n(\xi))$ reads

$$\begin{cases} \Lambda v_n'' - c_n v_n' = v_n^{\alpha_n}, & 0 < \xi < A, \\ v_n(0) \in I, & v_n'(0) = -\frac{c_n}{\Lambda}(1 - v_n(0)), \\ v_n(A) = v_n'(A) = 0. \end{cases} \quad (8.12)$$

Then, it is easy to see that the sequence $(v_n)_{n \in \mathbb{N}}$ is bounded in the space $H^2([0, A])$. We extract a convergent subsequence $(c_{n'}, v_{n'})_{n' \in \mathbb{N}}$ such that $c_{n'} \rightarrow c_\infty > 0$, $v_{n'} \rightarrow v_\infty$ in the space $C^1([0, A])$. We pass to the limit as $n' \rightarrow \infty$, $\alpha_{n'} \rightarrow 1$: there exists a non-negative function χ such that $(v_{n'})^{\alpha_{n'}} \rightarrow \chi$ weakly in $L^2(0, A)$. At the limit it holds in the distribution sense in the interval $(0, A)$:

$$\Lambda v_\infty'' - c_\infty v_\infty' = \chi.$$

Moreover, at the limit,

$$v_\infty(0) \in I, \quad v_\infty'(0) = -(c_\infty/\Lambda)(1 - v_\infty(0)), \quad v_\infty(A) = v_\infty'(A) = 0.$$

We observe that v_∞ belongs to $C^1([0, A])$ and is convex, non-increasing: there exists $0 < R_\infty \leq A$ such that $v_\infty(\xi) > 0$ for $0 < \xi < R_\infty$, $v_\infty(\xi) = 0$ for $R_\infty \leq \xi \leq A$. At fixed $\xi \in (0, R_\infty)$, $(v_{n'}(\xi))^{\alpha_{n'}} \rightarrow v_\infty(\xi)$ as $n' \rightarrow \infty$. Therefore, it comes $\chi(\xi) = v_\infty(\xi)$ whenever $0 < \xi < R_\infty$. The system eventually reads:

$$\begin{cases} \Lambda v_\infty'' - c_\infty v_\infty' = v_\infty, & 0 < \xi < R_\infty, \\ v_\infty(0) \in I, & v_\infty'(0) = -\frac{c_\infty}{\Lambda}(1 - v_\infty(0)), \\ v_\infty(R_\infty) = v_\infty'(R_\infty) = 0, \end{cases} \quad (8.13)$$

which has no solution for finite R_∞ (see also [6, Lemma 3.6 case (i)]).

As a consequence, the sequence $(R_n)_{n \in \mathbb{N}}$ is unbounded when $\alpha_n \rightarrow 1$. We may extract a subsequence $n'' \rightarrow \infty$ such that $R_{n''} \rightarrow +\infty$. Finally, repeating the process, $(R_n)_{n \in \mathbb{N}}$ goes to $+\infty$. \square

8.4. Limit $\alpha \rightarrow 0$. As in the previous subsection, $\Lambda > 0$ and $\theta \in (0, 1)$ are fixed. With each α , $0 < \alpha < 1$, we associate a solution $(c, R, v(\xi))$ of the problem (8.10)-(8.11).

Lemma 8.4. *Let $\Lambda > 0$ and $0 < \theta < 1$ be fixed. Let $(\alpha_n)_{n \in \mathbb{N}}$ be a decreasing sequence such that $0 < \alpha_n < 1$ and $\alpha_n \rightarrow 0$ as $n \rightarrow \infty$; the corresponding solution of (8.10)-(8.11) is denoted by $(c_n, R_n, v_n(\xi))$. Then, $(c_n, R_n, v_n(\xi))_{n \in \mathbb{N}}$ converges to the unique solution $(c, R, v(\xi))$ of the limit system*

$$\begin{cases} \Lambda v'' - cv' = 1, & 0 < \xi < R, \\ v(0) \in I, & v'(0) = -\frac{c}{\Lambda}(1 - v(0)), \\ v(R) = v'(R) = 0, \end{cases} \quad (8.14)$$

together with

$$\theta + \Lambda v(0) = 1 - c(1 - \Lambda) \int_0^R e^{-cs} v(s) ds, \quad (8.15)$$

where $c = R$ is uniquely given by (1.12), see Subsection 8.2 (i).

Proof. The proof follows most of the lines of the proof of Lemma 8.3. This time, we must show that the sequence $(R_n)_{n \in \mathbb{N}}$ is bounded. We take advantage of formula (6.18) which reads:

$$\frac{(2(1 + \alpha_n)\Lambda)^{1/2}}{1 - \alpha_n} a^{\frac{1-\alpha_n}{2}} < R_n < \frac{2\Lambda^{1/2}A(\alpha_n)b^{\frac{1-\alpha_n}{2}}}{1 - \alpha_n} \frac{1-b}{1-a}. \quad (8.16)$$

Here, $A(\alpha_n) = 2^{3/2}/(1 + \alpha_n)^{1/2} < 2^{3/2}$. For simplicity we assume $0 < \alpha_n \leq 1/2$ for all $n \in \mathbb{N}$. It comes the new definition of A :

$$(2a\Lambda)^{1/2} < R_n < A = 2^{7/2}\Lambda^{1/2} \frac{b^{1/4}}{1-a}. \quad (8.17)$$

Hence, the sequence $(c_n)_{n \in \mathbb{N}}$ is also bounded from above and from below, as in Lemma 8.3.

Next, we mimic the proof of Lemma 8.3. By abuse of notation, we also denote by $v_n(\xi)$ its extension by 0 to the interval $[0, A]$. The system for $(c_n, v_n(\xi))$ reads

$$\begin{cases} \Lambda v_n'' - c_n v_n' = v_n^{\alpha_n}, & 0 < \xi < A, \\ v_n(0) \in I, & v_n'(0) = -\frac{c_n}{\Lambda}(1 - v_n(0)), \\ v_n(A) = v_n'(A) = 0. \end{cases} \quad (8.18)$$

The sequence $(v_n)_{n \in \mathbb{N}}$ is bounded in the space $H^2([0, A])$. We extract a convergent subsequence $(c_{n'}, v_{n'})_{n' \in \mathbb{N}}$ such that $c_{n'} \rightarrow c > 0$, $v_{n'} \rightarrow v$ in the space $C^1([0, A])$. We pass to the limit as $n' \rightarrow \infty$, $\alpha_{n'} \rightarrow 0$: there exists a non-negative function χ such that $(v_{n'})^{\alpha_{n'}} \rightharpoonup \chi$ weakly in $L^2(0, A)$. At the limit it holds in the distribution sense in the interval $(0, A)$:

$$\Lambda v'' - cv' = \chi.$$

As in the proof in Lemma 8.3, there exists $0 < R \leq A$ such that $v(\xi) > 0$ for $0 < \xi < R$, $v(\xi) = 0$ for $R \leq \xi \leq A$. The only difference is the following: at fixed $\xi \in (0, R)$, $(v_{n'}(\xi))^{\alpha_{n'}} \rightarrow 1$ as $n' \rightarrow \infty$. Therefore, $\chi(\xi) = 1$ whenever $0 < \xi < R$. The system eventually reads:

$$\begin{cases} \Lambda v'' - cv' = 1, & 0 < \xi < R, \\ v(0) \in I, & v'(0) = -\frac{c}{\Lambda}(1 - v(0)), \\ v(R) = v'(R) = 0. \end{cases} \quad (8.19)$$

The next step is to take the limit in the formula

$$\theta + \Lambda v_{n'}(0) = 1 - c_{n'}(1 - \Lambda) \int_0^A e^{-c_{n'}s} v_{n'}(s) ds.$$

By Lebesgue's Theorem, it comes:

$$\theta + \Lambda v(0) = 1 - c(1 - \Lambda) \int_0^R e^{-cs} v(s) ds. \quad (8.20)$$

It follows from the uniform convergence of $v_{n'}(\xi)$ to $v(\xi)$ on the interval $[0, A]$ that $R_{n'} \rightarrow R$. Finally, because the system (8.19)-(8.20) has a unique solution, the entire sequence converges to $(c, R, v(\xi))$.

The proof is thus complete. \square

9. ACKNOWLEDGMENT

The authors want to thank Peter Gordon for bringing Chung K. Law's book to our attention and for fruitful discussions. P.S. and L.Z. would like to thank the National Natural Science Foundation of China (No. 11771336).

APPENDIX A. ABOUT THE POINCARÉ-BENDIXSON THEOREM

There are many textbooks on the qualitative theory of vector fields (see, e.g., Hartman [17], Lefschetz [21], Arnold [2]). First versions of the Poincaré-Bendixson Theorem were given in the historical works of Poincaré (see [24]) and Bendixson (see [5]). Here, we state a simplified modern version of the Poincaré-Bendixson Theorem (see Theorem A.3 below). A more complete version may be found in [21].

To begin with, for the convenience of the reader, we recall some basic definitions.

Let us consider a topological flow $\varphi(t, m)$ on a manifold M . For each $m \in M$ the curve $t \mapsto \varphi(t, m)$, parametrized by the time t , is the trajectory by m defined on a maximum open interval $(-\tau_-(m), +\tau_+(m))$ neighborhood of 0 and such that $\varphi(0, m) = m$ (one may have $\tau_+(m) = +\infty$ and/or $-\tau_-(m) = -\infty$). The *orbit* γ_m is the oriented geometrical image of the trajectory by m (the orbit is positively oriented by the time direction but the time parametrization is omitted). We can also consider the positive half-orbit $\gamma_m^+ = \varphi([0, \tau_+(m)), m)$ and the negative half-orbit $\gamma_m^- = \varphi([0, -\tau_-(m)), m)$.

The following limit sets have been intensively studied and used in the qualitative theory of dynamical systems (see, e.g., [21], [2], [4], [13]):

Definition A.1. *Let $\gamma = \gamma_m$ be an orbit of a continuous flow $\varphi(t, m)$ on a manifold M . The ω -limit set of γ is the set $\omega(\gamma)$ of points $p \in M$ such that, for $m \in \gamma$, there exists a sequence of times $(t_i) \rightarrow +\tau_+(m)$ with the property that $\varphi(t_i, m) \rightarrow p$, for $i \rightarrow +\infty$. In a similar way, one defines the α -limit set $\alpha(\gamma)$ by considering the sequences $(t_i) \rightarrow -\tau_-(m)$. One can also write $\omega(\gamma_m^+)$ and $\alpha(\gamma_m^-)$ for $\omega(\gamma)$ and $\alpha(\gamma)$ respectively. These limit sets does not depend on the choice of a particular point $m \in \gamma$.*

We recall that a vector field X on a n -manifold M is defined, on each open chart Ω with coordinates $x = (x_1, \dots, x_n)$, as a first order differential operator:

$$X(x) = A_1(x) \frac{\partial}{\partial x_1} + \dots + A_n(x) \frac{\partial}{\partial x_n}.$$

The vector field is said of class C^k if the components $A_i(x)$ are of this class on each chart Ω . A vector field may depend on a parameter λ , when the components in each chart depend on this parameter. A family of vector fields X_λ is of class C^k , if the components $A_i(x, \lambda)$ are of this class as functions of (x, λ) .

We also recall a basic result which is a corollary of the (real) Cauchy Theorem:

Theorem A.2. *Let X_λ be a family of vector fields defined on an open set Ω in \mathbb{R}^n , with $\lambda \in W$, an open set of \mathbb{R}^p . This family is assumed of class of differentiability \mathcal{C}^k , $k \geq 1$, or of analytic class (we mean that it is represented by a map $\Omega \times W \rightarrow \mathbb{R}^n$ of this given class). We call $\varphi(t, m, \lambda)$ its flow and consider $(t_0, m_0, \lambda_0) \in \mathbb{R} \times \Omega \times W$ such that the segment of $\varphi([0, t_0], m_0, \lambda_0)$ is contained in Ω . Then the flow $\varphi(t, m, \lambda)$ is defined in a neighborhood U of (t_0, m_0, λ_0) in $\mathbb{R} \times \Omega \times W$ and has on U the same class of differentiability as the family X_λ .*

Theorem A.3. (Poincaré-Bendixson Theorem) *One considers a C^1 vector field X defined on an open set Ω of \mathbb{R}^2 . One assumes that a positive half-orbit γ_m^+ of the flow has a compact closure in Ω . Then, $\omega(\gamma_m^+)$ contains a singular point or is a closed (periodic) orbit. As the α -limit sets of X are the ω -limit sets of $-X$, one has a similar result for the negative half-orbits γ_m^- of X .*

We have the following corollary of the Poincaré-Bendixson Theorem which will be the version used in this paper:

Corollary A.4. *One considers a C^1 vector field X defined on an open set Ω of \mathbb{R}^2 , with flow $\varphi(t, m)$. One assumes that Ω has a compact closure $\bar{\Omega}$ in \mathbb{R}^2 and we write $\partial\Omega = \bar{\Omega} \setminus \Omega$ for its boundary. One also assumes that Ω is simply connected and that X has no singular point (in Ω). Let m be any point in Ω . Then there exists a sequence $t_i \rightarrow \tau_+(m)$ and a point $p \in \partial\Omega$ such that $\varphi(t_i, m) \rightarrow p$ when $i \rightarrow +\infty$. A similar result is true for the negative times.*

We refer to [25] for a detailed proof of Corollary A.4.

APPENDIX B. PROOF OF PROPOSITION 4.3

The vector field X_c^E has a unique singular point at the origin O , two trajectories whose ω -limit and two trajectories whose α -limit set is the origin O . Each other trajectory goes from infinity to infinity. One can expect that these properties imply that X_c^E is topologically equivalent to an hyperbolic linear saddle vector field. Unfortunately, this is not true in general as that was already detected by Bendixson in [5]. Then, in the proof given below, we will have to use also more specific properties of the vector field X_c^E .

B.1. Construction of the equivalence on Q . In order to simplify the notation, and as c, α, Λ are assumed to be fixed, we will write X for the vector X_c . We will denote by L the unique stable manifold of X and $y(x)$ the function whose graph is L ; the latter function reads $y_c(x)$ in Section 4. We also denote by L^0 the stable manifold of the linear field $X^0 = y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y}$, restricted to the quadrant Q , which is the graph $\{y = y^0(x) = -x\}$.

The stable manifold $L := \{y = y(x)\}$ splits Q into two invariant regions: $Q \setminus L = Q^+ \cup Q^-$ where $Q^+ = \{x > 0, y(x) < y \leq 0\}$ and $Q^- = \{x \geq 0, y < y(x)\}$. To begin with, we want to establish some properties for the orbits of X , which are trivially verified by the orbits of the linear vector field X^0 .

Lemma B.1. *We consider the vector field X defined on Q .*

- (1) *Any orbit γ contained into Q^+ is a graph $\{y = y_\gamma(x)\}$ over an interval $[x_\gamma, +\infty)$ for some x_γ such that $0 < x_\gamma < x$.*
- (2) *Any orbit contained into Q^- is a graph $\{y = y_\gamma(x)\}$ over $[0, +\infty)$. Moreover if $(x_0, y_0) \in \gamma$, we have that: $y_0 - c/\Lambda x_0 \leq y_\gamma(0) < 0$.*

Proof. The point (1) follows from the results of Section 4: a trajectory starting at a point $(x, y) \in Q^+$ cannot tend toward O because L is the unique stable manifold in Q by Lemma 4.7; then, as a consequence of Lemma 4.4, this trajectory must attain a point $(x_\gamma, 0)$ with $0 < x_\gamma < x$. Finally, as in the proof used in Lemma 4.7 for the stable manifold, it is easy to prove that the corresponding orbit γ is a graph $\{y = y_\gamma(x)\}$ above $[x_\gamma, +\infty)$.

We now consider the orbits in Q^- . We first claim that, in the interior of the whole quadrant Q , the vector field X is transverse to the direction of the vector $Z = (1, c/\Lambda)$ and directed toward

the left. To see this, we consider the orthogonal vector field $Z^\perp = (-c/\Lambda, 1)$ and compute the scalar product

$$\langle Z^\perp, X \rangle = -\frac{c}{\Lambda}y + \left(\frac{c}{\Lambda}y + \frac{1}{\Lambda}x^\alpha\right) = \frac{1}{\Lambda}x^\alpha,$$

which is strictly positive if $x > 0$. The claim follows.

Consider now an orbit γ in Q^- . By an argument similar to the one used in Lemma 4.7 it is easy to see that γ is a graph above some maximal interval I in the Ox -axis. We want firstly to prove that if $(x_0, y_0) \in \gamma$, then $[0, x_0] \subset I$ (this means that I is closed on the left with 0 as end point). To show this point we consider the *curved rectangle* $\mathcal{R} = [O, A, B, C]$, where $A = (x_0, y(x_0))$, $B = (x_0, y_0)$ and C belongs to the Oy -axis (observe that necessarily we have that $y_0 < y(x_0)$). The side $[O, A]$ is an arc in L , the three other sides are linear segments and $[B, C]$ as a direction parallel to the vector Z above, see Figure 5. The vector field X is tangent to $[O, A]$, transverse toward the left along $[A, B]$, $[B, C]$ and $[C, O]$: this means that X is entering \mathcal{R} along $[A, B] \cup [B, C)$ and is going out \mathcal{R} along $[C, O)$. Now, as in the point (1), we can use the results of Section 4 to obtain that the trajectory of the point (x_0, y_0) must reach a point of $[C, O)$ in a finite time.

This implies that γ is a graph $\{y = y_\gamma(x)\}$ over a maximal interval $[0, x_+)$. Moreover, as the ordinate of C is equal to $y_0 - c/\Lambda x_0$ that is less than $y_\gamma(0)$, we have that $y_\gamma(0) \geq y_0 - \frac{c}{\Lambda}x_0$, as stated in the Lemma.

We prove now that $x_+ = +\infty$ by using the same idea as in the proof of Lemma 4.8. We have that $d(y(x) - y_\gamma(x))/dx < 0$ on $(0, x_+)$, the common domain of existence of $y(x)$ and $y_\gamma(x)$. This inequality implies that, for all $x < x_+$, we have that the function $y(x) - y_\gamma(x)$ is decreasing and then that:

$$y_\gamma(x) > y(x) - y(x_0) + y_\gamma(x_0).$$

Passing to the limit x_+ this implies that the α -limit set of γ is contained in the closed interval $J = \{x_+\} \times [y(x_+) - y(x_0) + y_\gamma(x_0), 0]$ and then is non-empty and must contain a singular point, as consequence of the Poincaré-Bendixson Theorem. As X has no singular point in a neighborhood of J , this is impossible. \square




FIGURE 5. *Rectangle* \mathcal{R}

Lemma B.1 allows to define two transition functions: for $(x, y) \in Q^+$, the function $T_x(y)$ which is the value x_γ defined in Lemma B.1 for the orbit γ through the point (x, y) ; for $(x, y) \in Q^-$, the function $S_x(y)$ where $(0, S_x(y))$ is the point reached on the axis Oy by the orbit γ through (x, y) (see Figure 6). Clearly, these functions are analytic on Q^+ and Q^- respectively. The following lemma shows that they can be continuously extended on L with the common value 0:

Lemma B.2. *Let $\bar{Q}^+ = Q^+ \cup L = \{x \geq 0, y(x) \leq y \leq 0\}$ and $\bar{Q}^- = Q^- \cup L = \{x \geq 0, y \leq y(x)\}$ be the closure of Q^+ and Q^- . Then $T_x(y)$ and $S_x(y)$ can be continuously extended by 0 on L .*

Proof. To prove the continuity along L , we will proceed in three steps.

Step 1. We fix a value $x_0 > 0$ and prove that $T_{x_0}(y)$ and $S_{x_0}(y)$ tends toward 0 when y tends toward $y(x_0)$. We begin with $T_{x_0}(y)$. Let us observe that $y \mapsto T_{x_0}(y)$ is decreasing from $T_{x_0}(0) = x_0$ when y decreases from $y = 0$. Let $x_T \geq 0$ be the limit when $y \rightarrow y(x_0)$. We claim that $x_T = 0$. If not, the orbit of $(x_T, 0)$ for the vector field $-X$ reaches the vertical axis $\{x = x_0\}$ at some point (x_0, y_T) with $y(x_0) < y_T < 0$. This is clearly in contradiction with the definition of x_T . As a conclusion, the function $y \mapsto T_{x_0}(y)$ extends continuously by 0 at $y = y(x_0)$. We have exactly the same proof for $S_{x_0}(y)$.

Step 2. The extensions are continuous on $L \setminus \{O\}$ as functions of the two variables (x, y) . Let us prove the continuity of the extension of $T_x(y)$ at a point $(x_0, y(x_0)) \in L$ with $x_0 > 0$. The idea is that we can compute T_x in terms of T_{x_0} and the transitions along the flow of X . If we denote by Σ_x the vertical section by x (parameterized by y), there is an analytic transition along the flow of X : $y \rightarrow T_x^{x_0}(y)$ from Σ_x to Σ_{x_0} such that $T_x^{x_0}(y(x)) = T_x(y(x_0))$. Moreover it is clear that

$$T_x(y) = T_{x_0} \circ T_x^{x_0}(y).$$

As the extension of T_{x_0} is continuous in y at $y(x_0)$ and $T_x^{x_0}(y)$ is analytic in (x, y) at $(x_0, y(x_0))$, the above formula shows that the extension of $T_x(y)$ is continuous in (x, y) at $(x_0, y(x_0))$. Exactly the same proof can be made for $S_x(y)$.

Step 3. *Continuity of the extension at the origin O .* The fact that $T_x(y) \rightarrow 0$ for $(x, y) \rightarrow (0, 0)$ in \bar{Q}^+ follows from the inequality $0 \leq T_x(y) \leq x$. For the function $S_x(y)$, we apply the inequality $|S_x(y)| \leq |y| + \frac{c}{\lambda}|x|$ which is an equivalent form of the inequality states in the point (2) of Lemma B.1. \square

From now on and for simplicity, we will also write $T_x(y)$ and $S_x(y)$ for the continuous extensions of these functions on \bar{Q}^+ and \bar{Q}^- , i.e. along the curve L . For each $x > 0$ and as it is monotonic, the map $y \mapsto T_x(y)$ is an homeomorphism from the interval $[0, y(x)]$ to the interval $[x, 0]$. Also, for each $x \geq 0$, the map $y \mapsto S_x(y)$ is an homeomorphism from the interval $(-\infty, y(x))$ to the interval $(-\infty, 0]$. Moreover we have proved in Lemma B.2 that these x -families of homeomorphisms are continuous with the limits $S_0(y) \equiv y$ and $T_0(0) = 0$ (the interval $[0, y(0)]$ is reduced to the point $\{0\}$). This allows to define the inverse homeomorphisms $T_x^{-1}(z)$ and $S_x^{-1}(z)$ defined respectively for $0 \leq z \leq x$ and $z \in (-\infty, 0]$. These x -families are clearly analytic for $z \neq 0$. we prove now that they are continuous in (x, z) at $z = 0$:

Lemma B.3. *The inverse maps $T_x^{-1}(z)$ and $S_x^{-1}(z)$ have a continuous extension for $z = 0$ by $T_0^{-1}(0) = 0$ and $S_0^{-1}(z) \equiv z$.*

Proof. The proof is indeed much easier than the one given in Lemma B.2 for the direct maps. Now, we can take advantage of the contraction property of the vertical direction, along the flow followed in the x -direction, already proved in Lemma 4.8 and used above in Lemma B.1. This

property implies that $|T_x^{-1}(z) - y(x)| \leq |y(z)|$ and that $|S_x^{-1}(z) - y(x)| \leq |z|$. The result follows directly for $z \rightarrow 0$. \square

As X^0 is just a particular case, we have transition functions $T_x^0(y)$ and $S_x^0(y)$ for it, which verify the statements of Lemmas B.1, B.2 and B.3 (in fact, as $T_x^0(y) = \sqrt{x^2 - y^2}$ and $S_x^0(y) = -\sqrt{y^2 - x^2}$, we can verify directly their properties). Using the transitions functions of X^0 and X we can now construct on Q a topological equivalence between these vector fields:

Proposition B.4. *We define a function $h_x(y)$ on Q by $h_x(y) = T_x^{-1} \circ T_x^0(y)$ if $(x, y) \in \bar{Q}^+$ and by $h_x(y) = S_x^{-1} \circ S_x^0(y)$ if $(x, y) \in \bar{Q}^-$ (see Figure 6). Then, the map $H(x, y) = (x, h_x(y))$ is an homeomorphism of Q , which is the identity on the boundary ∂Q ($H(x, 0) = (x, 0)$ for $x \geq 0$ and $H(0, y) = (0, y)$ for $y \leq 0$). Moreover, H is a topological equivalence between X^0 and X , i.e. H sends the orbits of X^0 onto the orbits of X .*

Proof. As T_x^0 and T_x^{-1} are homeomorphisms from $[0, y_0(x)]$ to $[x, 0]$ and from $[x, 0]$ to $[0, y(x)]$ respectively, the map $h_x(y)$ is an homeomorphism from $[0, y_0(x)]$ to $[0, y(x)]$. For similar reasons we have that $h_x(y)$ is an homeomorphism from $(-\infty, y_0(x)]$ to $(-\infty, y(x)]$. As these two definitions of $h_x(y)$ coincide at $y_0(x)$, they define a global homeomorphism also denoted by $h_x(y)$ from $(-\infty, 0]$ to itself. The continuity on Q of the map $(x, y) \mapsto h_x(y)$ follows from the continuity of the transition functions and their inverse in terms of their two variables, proved in Lemmas B.2 and B.3. As $h_x^{-1}(y) = (T_x^0)^{-1} \circ T_x(y)$ on \bar{Q}^+ and $h_x^{-1}(y) = (S_x^0)^{-1} \circ S_x(y)$ on \bar{Q}^- , the same argument shows that $(x, y) \mapsto h_x^{-1}(y)$ is a continuous family of homeomorphisms from $(-\infty, 0]$ to itself. Clearly, the map $(x, y) \mapsto (x, h_x^{-1}(y))$ is the inverse of the map $H(x, y) = (x, h_x(y))$. As these two maps are continuous, we have that H is an homeomorphism of Q to itself. Moreover it is clear that H is equal to the identity on ∂Q .

By definition of the transition maps, the homeomorphism H sends each orbit of X^0 onto the orbits of X (see Figure 6; in particular H sends L^0 onto L). We have proved that H is a topological equivalence on Q . \square




FIGURE 6. *The topological equivalence H*

B.2. Extension of the equivalence to \mathbb{R}^2 . We can obtain very similar properties for X_c on the quadrant $Q' = \{x \geq 0, y \geq 0\}$ as the ones proved for X_c on Q , in Section 4 and in the above subsection B.1. A noticeable difference is that the slope of X_c is greater than c/Λ everywhere in Q' . As a consequence, we can find a unique unstable manifold at the origin which is a graph $x = x_c(y)$ and more generally each orbit γ in Q' is a graph $x_\gamma(y)$ above an interval $[y_\gamma, +\infty)$ with $y_\gamma \geq 0$.

We can repeat the proof given in Subsection B.1 in order to construct an homeomorphism H' of Q of the form $(h'_y(x), y)$ where $h'_y(x)$ and $(h'_y)^{-1}(x)$ are continuous families of homeomorphisms and such that H' is equal to the identity on $\partial Q'$. This homeomorphism can be glued up with H to give a topological equivalence \hat{H} of X^0 and X_c on the half-plane $\{x \geq 0\}$, which is equal to the identity along the Oy -axis.

To extend \hat{H} to the whole plane, we use the symmetry property of X_c^E , already mentioned in Subsection 4.1: $X_c^E(-x, -y) = -X_c(x, y)$. For each $x \leq 0$, we define $\hat{H}(x, y)$ by $\hat{H}(x, y) = -\hat{H}(-x, -y)$. This extends \hat{H} in the half-space $\{x \leq 0\}$ and the two definitions coincide along $\{x = 0\}$ with the identity. Then, this defines a global topological equivalence \hat{H} on \mathbb{R}^2 between X^0 and X_c^E (this construction takes into account the fact that the orbits of X^0 and X_c^E are crossing the Oy -axis at the points $(0, y)$ such that $y \neq 0$).

REFERENCES

- [1] D. Addona, C.-M. Brauner, L. Lorenzi and W. Zhang, Instabilities in a combustion model with two free interfaces, J. Differ. Equ. **268** (2020), 396–4016.
- [2] V.I. Arnold, Ordinary Differential Equations, MIT Press, Cambridge, MA, 1991.
- [3] S.P. Bhat and D.S. Bernstein, Finite-time stability of continuous autonomous systems, SIAM J. Control Optim. **38** (2000), 751-766.
- [4] N.P. Bathia and G.P. Szegő, Stability Theory of Dynamical Systems, Classics in Mathematics, Springer, 1st ed., 2002.
- [5] I. Bendixson, Sur les courbes définies par des équations différentielles, Acta Math., **24** (1901), 1-88.
- [6] H. Berestycki, B. Nicolaenko and B. Scheurer, Traveling wave solutions to combustion models and their singular limits, SIAM Journal on Mathematical Analysis, **16** (1985), 1207-1242.
- [7] H. Berestycki, P.-L. Lions and L.A. Peletier, An ODE approach to the existence of positive solutions for semilinear problems in \mathbb{R}^n , Indiana Univ. Math. J., **30** (1981), 141-157.
- [8] I. Brailovsky, P.V. Gordon, L. Kagan and G.I. Sivashinsky, Shape-thermal instabilities in premixed flames: stepwise ignition-temperature kinetics, Combustion and Flame **162** (2015), 2077-2086
- [9] C.-M. Brauner, P.V. Gordon and W. Zhang, An ignition-temperature model with two free interfaces in premixed flames, Combust. Theory Model., **20** (2016), 976-994.
- [10] C.-M. Brauner, J. Hulshof, A. Lunardi, A general approach to stability in free boundary problems, J. Differential Equations **164** (2000), 16-48.
- [11] C.-M. Brauner, L. Lorenzi and M. Zhang, Stability analysis and Hopf bifurcation for large Lewis number in a combustion model with free interface. Ann. Inst. H. Poincaré Anal. Non Linéaire (2020). <https://doi.org/10.1016/j.anihpc.2020.01.002>.
- [12] J.D. Buckmaster and G.S.S. Ludford, Theory of laminar flames, Cambridge University Press, Cambridge, 1982.
- [13] Y.N. Dowker and F.G. Friedlander, On limit sets in dynamical systems, Proc. London Math. Soc (3) **4**, (1954), 168–176.
- [14] G.F. Duff, Limit cycles and rotated vector fields, Annals of Math. (2) **57**, (1953), 15-31.
- [15] F. Dumortier, J. Llibre and J.C. Artés, Qualitative Theory of Planar Differentiable Systems, Universitext, Springer-Verlag (2006), 15-31.
- [16] D. Grobman, Homeomorphisms of systems of differential equations (english translation), Doklady Akademii Nauk SSSR, vol. **128** (1959), 880-881.

- [17] P. Hartman, Ordinary Differential Equations, Wiley, New York (1954).
- [18] P. Hartman, A lemma in the theory of structural stability of differential equations, Proc. Amer. Math. Soc., vol. **11**, $n^{\circ}4$ (1960) 610-620.
- [19] M. Hirsch and S. Smale, Differential equations, dynamical systems and linear algebra, Pure and Applied Mathematics, vol. **60**, Academic Press, New York London (1974).
- [20] C.K. Law, Combustion Physics, Cambridge University Press, Cambridge, 2010.
- [21] S. Lefschetz, Differential equations: Geometric theory, 2nd edition, Dover Publications Inc., New York, 1977.
- [22] S.G. Nersesov, W. M. Haddad and Q. Hui, Finite-time stabilization of nonlinear dynamical systems via control vector Lyapunov functions, J. Franklin Institute **345** (2008), 819–837.
- [23] H.B. Oza, S.K. Spurgeon and N.V. Valeyev, Non-Lipschitz growth functions as a natural way of modelling finite time behaviour in auto-immune dynamics, IFAC Proceedings Volumes **47** 2014, 11611-11616.
- [24] H. Poincaré, Sur les courbes définies par des équations différentielles, Journ. de Math. Pures et Appl., **s.3 t.VII** (1881)=[Oeuvres I, 3–44]; (Second part) Id., s. 3, t. VIII (1882)=[Oeuvres I, 44–84]; (Third part) Id., s. 4, t. I (1885)=[Oeuvres I, 90–158]; (Fourth part) Id., s. 4, t. II (1886)=[Oeuvres I, 167–222], Oeuvres de Henri Poincaré, **I-XI**, Gauthier-Villars (new impression 1950–1965).
- [25] R. Roussarie, Some applications of the Poincaré-Bendixson Theorem, in preparation.
- [26] M. Shub, Global stability of dynamical systems, Springer-Verlag, New York, 1987.

SCHOOL OF MATHEMATICAL SCIENCES, TONGJI UNIVERSITY, 1239 SIPING RD., SHANGHAI 200092, CHINA, AND INSTITUT DE MATHÉMATIQUES DE BORDEAUX UMR CNRS 5251, UNIVERSITÉ DE BORDEAUX, 33405 TALENCE CEDEX, FRANCE.

INSTITUT DE MATHÉMATIQUE DE BOURGOGNE UMR CNRS 5584, UNIVERSITÉ DE BOURGOGNE-FRANCHE COMTÉ, B.P. 47870 21078 DIJON CEDEX, FRANCE

SCHOOL OF MATHEMATICAL SCIENCES, TONGJI UNIVERSITY, 1239 SIPING RD., SHANGHAI 200092, CHINA.

SCHOOL OF MATHEMATICAL SCIENCES, TONGJI UNIVERSITY, 1239 SIPING RD., SHANGHAI 200092, CHINA.

E-mail address: `claude-michel.brauner@u-bordeaux.fr`

E-mail address: `Robert.Roussarie@u-bourgogne.fr`

E-mail address: `shang@tongji.edu.cn`

E-mail address: `1810879@tongji.edu.cn`