



**HAL**  
open science

## Disseminated invasive aspergillosis in patients with severe influenza infection

Grégoire Monseau, Mathilde Wairy, Marie Flores, Kevin Brunet, Faustine Reynaud, Damien Marie, Florence Boissier, Delphine Chatellier, Anne Veinstein, Jean-Pierre Frat, et al.

### ► To cite this version:

Grégoire Monseau, Mathilde Wairy, Marie Flores, Kevin Brunet, Faustine Reynaud, et al.. Disseminated invasive aspergillosis in patients with severe influenza infection. *IDCASSES*, 2020, 20, pp.e00777. 10.1016/j.idcr.2020.e00777 . hal-02980678

**HAL Id: hal-02980678**

**<https://hal.science/hal-02980678>**

Submitted on 22 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

## **Disseminated invasive aspergillosis in patients with severe influenza infection**

Grégoire Monseau MD<sup>1</sup>, Mathilde Wairy<sup>1</sup>, Marie Flores MD<sup>2</sup>, Kevin Brunet MD<sup>3</sup>, Faustine Reynaud MD<sup>1</sup>, Damien Marie MD<sup>1</sup>, Florence Boissier MD, PhD<sup>1</sup>, Delphine Chatellier MD<sup>1</sup>, Anne Veinstein MD<sup>1</sup>, Jean-Pierre Frat MD<sup>1-4</sup>, René Robert MD, PhD<sup>1-4</sup>, Arnaud W. Thille MD, PhD<sup>1-4</sup>

### ***Affiliations***

<sup>1</sup>CHU de Poitiers, Service de Médecine Intensive et Réanimation, Poitiers, France.

<sup>2</sup> CHU de Poitiers, Service d'Anatomo-pathologie, Poitiers, France.

<sup>3</sup> CHU de Poitiers, Laboratoire de Parasitologie et Mycologie Médicale, Poitiers, France.

<sup>4</sup> INSERM CIC 1402 ALIVE, Université de Poitiers, Poitiers, France.

### ***Corresponding author:***

**Arnaud W. Thille, Médecine Intensive et Réanimation, CHU de Poitiers**

**2 rue de la Milétrie**

**86000 Poitiers, FRANCE**

**[arnaud.thille@chu-poitiers.fr](mailto:arnaud.thille@chu-poitiers.fr)**

***Author contribution:*** GM, MW and AWT wrote the majority of the manuscript; MF, KB, FR, DM, FB, DC, AV, JPF and RR participated in writing the manuscript and approved its final version; GM, KB and MF realized the pictures

***Fundings:*** French Health Ministry and Research Ministry

***Conflict of Interest:*** None

***Manuscript word count:*** 413 words, 2 figures.

***Glossary:*** ARDS = Acute Respiratory Distress Syndrome, BAL = Bronchoalveolar Lavage, ECMO = Extracorporeal membrane oxygenation, HHV6 = Human Herpes Virus 6, ICU = Intensive Care Units, PCR = Polymerase Chain Reaction.

A female 66 year-old patient, not immunocompromised, was admitted in ICU for severe influenza complicated by severe ARDS leading to extra-corporeal membrane oxygenation (ECMO). Given respiratory disease severity, bronchoalveolar lavage (BAL) was not performed during this period, but iterative searches for aspergillosis using serum galactomannan test and cultures on tracheal aspirates were performed. Because delayed recovery of awakening, cerebral computed tomographic scan and subsequent magnetic resonance imaging showed a non-specific right fronto-temporal collection (Figure 1). CSF revealed meningitis with predominance of lymphoid-cells, and PCR for HHV6 was positive in the cerebrospinal fluid, in serum and in the bone marrow aspirate examination, probably reflecting an immunocompromised state induced by severe influenzae, and leading to initiation of Ganciclovir. Subsequently, she had transient cardiac arrest with ST-segment elevation myocardial infarction, and coronarography revealed acute occlusion of the anterior interventricular artery. Despite a successful revascularization procedure, the patient died of refractory shock. Given the uncertainty of diagnosis, a clinical autopsy was performed. Histological examination found disseminated mycelial filaments in the form of abscesses in the lungs and the brain, but also disseminated in the kidney, colon, spleen and heart, strongly suggesting aspergillosis. Coronary artery occlusion was directly due to angio-invasive mycelial filaments (Figure 2). *Aspergillosis fumigatus* PCR was performed twice but paraffin inclusion seems to be responsible for the failure. Our last culture for aspergillosis on tracheal aspirate was performed more than one week before death and we cannot know whether negative culture was due to rapid progression of aspergillosis over the last week or due to low sensitivity of culture.

Recent literature has reported cases of invasive aspergillosis in the non-immunocompromised host with severe influenza infection(1-2). Indeed invasive pulmonary aspergillosis was diagnosed early in the course of severe influenza aspergillosis (3 days after ICU admission in median)(2). Therefore, BAL would probably help to diagnose these patients and should be discussed, even in patients with severe ARDS. However, according to usual algorithms for aspergillosis diagnosis(3-4), more than one-third of patients were not classifiable because they had a positive galactomannan test but negative culture on BAL. A recent study found high sensitivity to detect aspergillosis by combining galactomannan test and PCR aspergillosis on BAL(5). The authors reported that a combination of galactomannan test and PCR aspergillosis on BAL with 1-3-beta-D-glucan on serum resulted in 100% sensitivity.

In conclusion, aspergillosis may occur in severe influenza and BAL should be performed whenever possible for diagnosis, using culture, galactomannan and PCR on BAL, even in the most severe acute respiratory failure.

## ***References***

1. Vanderbeke L, Spriet I, Breynaert C, Rijnders BJA, Verweij PE, Wauters J. Invasive pulmonary aspergillosis complicating severe influenza: epidemiology, diagnosis and treatment. *Curr Opin Infect Dis* 2018; 31(6): 471-80.
2. Schauwvlieghe A, Rijnders BJA, Philips N, et al. Invasive aspergillosis in patients admitted to the intensive care unit with severe influenza: a retrospective cohort study. *Lancet Respir Med* 2018; 6(10): 782-92.
3. De Pauw B, Walsh TJ, Donnelly JP, et al. Revised definitions of invasive fungal disease from the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. *Clin Infect Dis* 2008; 46(12): 1813-21.
4. Blot SI, Taccone FS, Van den Abeele AM, et al. A clinical algorithm to diagnose invasive pulmonary aspergillosis in critically ill patients. *Am J Respir Crit Care Med* 2012; 186(1): 56-64.
5. Boch T, Reinwald M, Spiess B, et al. Detection of invasive pulmonary aspergillosis in critically ill patients by combined use of conventional culture, galactomannan, 1-3-beta-D-glucan and *Aspergillus* specific nested polymerase chain reaction in a prospective pilot study. *J Crit Care* 2018; 47: 198-203.

***Legend***

*Figure 1*

Cerebral MRI Diffusion sequence: right frontal-temporal collection (Arrow).

*Figure 2*

Coronary vessel x 10, Grocott coloration: coronary vessel with endoluminal occlusion by mycelian filament (Arrow).



