



The Hausdorff dimension of the harmonic measure for relatively hyperbolic groups

Matthieu Dussaule, Wenyuan Yang

► To cite this version:

Matthieu Dussaule, Wenyuan Yang. The Hausdorff dimension of the harmonic measure for relatively hyperbolic groups. 2020. hal-02985049

HAL Id: hal-02985049

<https://hal.science/hal-02985049>

Preprint submitted on 1 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE HAUSDORFF DIMENSION OF THE HARMONIC MEASURE FOR RELATIVELY HYPERBOLIC GROUPS

MATTHIEU DUSSAULE AND WENYUAN YANG

ABSTRACT. The paper studies the Hausdorff dimension of harmonic measures on various boundaries of a relatively hyperbolic group which are associated with random walks driven by a probability measure with finite first moment. With respect to the Floyd metric and the shortcut metric, we prove that the Hausdorff dimension of the harmonic measure equals the ratio of the entropy and the drift of the random walk.

If the group is infinitely-ended, the same dimension formula is obtained for the end boundary endowed with a visual metric. In addition, the Hausdorff dimension of the visual metric is identified with the growth rate of the word metric. These results are complemented by a characterization of doubling visual metrics for accessible infinitely-ended groups : the visual metrics on the end boundary is doubling if and only if the group is virtually free. Consequently, there are at least two different bi-Hölder classes (and thus quasi-symmetric classes) of visual metrics on the end boundary.

1. INTRODUCTION

Relatively hyperbolic groups can admit several interesting compactifications such as the Floyd [18], Bowditch [5] and Freudenthal [19] (also called end) compactification. The resulting boundaries are compact metrizable spaces on which the groups act and have rich dynamics in terms of convergence actions. This point of view has found many applications [3] [23]. In addition, these boundaries can be endowed with two well-known classes of (quasi-)conformal [8] and harmonic measures [35] so they are examples of metric measured spaces (X, d, ν) . The Hausdorff dimension $Hdim_d(\nu)$ of the triple (X, d, ν) is the infimum of the Hausdorff dimensions of ν -full subsets, so it is a measurement of the largeness of the measure class of ν . Comparison of the conformal and the harmonic measures has been an active research problem with origins in dynamic systems, see [34] [26] [1] [25] [16] [20] [48] to just name a few.

This paper is devoted to computing the Hausdorff dimension formula for harmonic measures on various boundaries associated with a random walk with finite first moment on a non-elementary relatively hyperbolic group. Precisely, we compute the Hausdorff dimension of the harmonic measure ν on the Floyd and Bowditch boundaries endowed respectively with the Floyd and the Floyd shortcut distances. We also compute this Hausdorff dimension on the end boundary endowed with a visual distance whenever the group is infinitely ended. Up to a parameter depending only on the chosen distance, we show that

$$(1) \quad Hdim(\nu) = \frac{h}{l}$$

where h is the asymptotic entropy and l is the rate of escape in the word metric of the random walk. See Theorem 1.1 and Theorem 1.3 for accurate statements.

This formula (1) was first obtained by Kaimanovich [32] and Ledrappier [38] on free groups and then by Le Prince [47] on general hyperbolic groups. A substantial generalization of Le Prince's results was given by Tanaka [53] to any proper action with exponential growth on proper hyperbolic spaces with bounded geometry, and any acylindrical action on possibly improper hyperbolic spaces. In particular, his results do apply to both the geometrical finite action of a relatively hyperbolic group on a proper hyperbolic space, and the acylindrical action on its relative Cayley graph [42].

However, the former proper action under bounded geometry assumption forces peripheral subgroups to be virtually nilpotent [11]. This makes a serious limitation of Tanaka's result to be applied in this rather general class of groups. Moreover, visual distances on the Bowditch boundary are very non-canonical, for they can depend heavily on the choice of a hyperbolic space X on which the group acts, see [28]. On the other hand, the acylindrical action on the relative Cayley graph does yield the formula (1) for the harmonic measure on the Gromov boundary of the relative Cayley graph and where the drift is computed in the relative metric.

In a certain point of view, this is unsatisfactory since the relative metric on the group is non-proper and the (non-compact) Gromov boundary here is only a part of the compact Bowditch boundary.

One of our main contributions here is to consider the drift for the word metric and to compute the Hausdorff dimension associated to the Floyd shortcut distance on the Bowditch boundary, which only depends on the group and the choice of a word distance.

1.1. Hausdorff dimension of harmonic measures. Let μ be a probability measure on a finitely generated group G such that the support $\text{supp}(\mu)$ generates G as a semi-group. We call such a measure *admissible*. The measure defines a μ -random walk with step transitions given by $p(x, y) = \mu(x^{-1}y)$ for $x, y \in G$. Let

$$\Omega := \{\mathbf{x} = (\omega_n)_{n \geq 0} : \omega_n \in G\}$$

be the trajectory space of the μ -random walk with the probability measure \mathbb{P} , which is the pushforward of the product measure $(G^{\mathbb{N}}, \mu^{\mathbb{N}})$ under the product map $G^{\mathbb{N}} \rightarrow \Omega$ given by

$$(s_1, s_2, \dots, s_n, \dots) \mapsto (1, s_1, s_1 s_2, \dots, \omega_n, \dots),$$

where $\omega_n = s_1 \cdots s_n$.

Denote by

$$L(\mu) = \sum_{g \in G} d(1, g) \mu(g)$$

the expectation of $d(1, \omega_1)$ and more generally by

$$L(\mu^{*n}) = \sum_{g \in G} d(1, g) \mu^{*n}(g)$$

the expectation of $d(1, \omega_n)$ where μ^{*n} is the n -th convolution power of μ . Equivalently, μ^{*n} is the law of the random variable $\mathbf{x} \mapsto \omega_n$. The sequence $L(\mu^{*n})$ is sub-additive and so $\frac{1}{n} L(\mu^{*n})$ has a well defined limit l called the *rate of escape* (or *drift*). Whenever μ has *finite first moment* (i.e. $L(\mu) < \infty$), Kingman's sub-additive ergodic theorem shows that almost surely,

$$(2) \quad l := \lim_{n \rightarrow \infty} \frac{d(1, \omega_n)}{n} < +\infty.$$

Also define $H(\mu) = \sum_{g \in G} \mu(g) \log \mu(g)$ and $H(\mu^{*n}) = \sum_{g \in G} \mu^{*n}(g) \log \mu^{*n}(g)$. The sequence $H(\mu^{*n})$ also is sub-additive and so $\frac{1}{n} H(\mu^{*n})$ has a well defined limit h called the asymptotic *entropy*. Again, whenever $H(\mu)$ is finite, an application of the ergodic Theorem (see [13] or [34]) shows that

$$(3) \quad h := \lim_{n \rightarrow \infty} \frac{-\log(\mu^{*n}(\omega_n))}{n} < +\infty.$$

The groups G under consideration are assumed to be relatively hyperbolic throughout. There are many equivalent ways to formulate this notion. To motivate our results, we use the dynamical definition. An action of G by homeomorphism on a compact metrizable space M is called *convergence* if the induced action on the space of triple points is properly discontinuously. Let (Γ, d) denote the Cayley graph of G with respect to a finite generating set. Then G is called *relatively hyperbolic* if there exists a Hausdorff compact space M compactifying every Cayley graph Γ so that the left multiplication of G extends to a minimal geometrically finite action on the boundary M . See precise definitions in Section 2.1. The compact space M denoted by $\partial_{\mathcal{B}} G$ later on is called *Bowditch boundary* of G . A relatively hyperbolic group G is called *non-elementary* if its Bowditch boundary contains more than two points. Equivalently, G fixes no finite subset of $\partial_{\mathcal{B}} G$. In such a case, G is non-amenable. In particular, the rate of escape and the asymptotic entropy are positive.

There are two natural classes of metrics on the Bowditch boundary. First, by Yaman [58], $\partial_{\mathcal{B}} G$ can be realized as the Gromov boundary of a proper hyperbolic space X on which G acts via a geometrically finite action. Thus, we can endow $\partial_{\mathcal{B}} G$ with the visual metric constructed using the hyperbolic geometry of X . Second, the Floyd (shortcut) metric is obtained from the Cayley graph (Γ, d) as follows. We fix a parameter $\lambda \in (0, 1)$ and a basepoint $o \in G$. Rescaling the length of every edge e of Γ to $\lambda^{d(o, e)}$ induces a new length metric called the *Floyd metric* δ_λ on Γ . The Cauchy metric completion of (Γ, δ_λ) is the so-called Floyd compactification of Γ on which G acts as a convergence action [36]. By the work of Gerasimov and Potyagailo [22], [23], there exists λ_0 such that whenever $\lambda \in [\lambda_0, 1)$, the Bowditch boundary is an equivariant quotient of the Floyd boundary where the non-trivial fibers are possible only on bounded parabolic points.

The Floyd metric can thus be pushforwarded to obtain the so-called shortcut metric $\bar{\delta}_\lambda$ on $\partial_B G$ [23]. See more details in Section 2.2.

In [35], Karlsson proved that almost every trajectory converges to a limit point in the Floyd boundary, and the *hitting measure* or *harmonic measure* $\nu_{\mathcal{F}}$ on the Floyd boundary gives a model of the Poisson boundary of the μ -random walk [31]. The same discussion applies to the Bowditch boundary which gives another model of the Poisson boundary when endowed with a harmonic measure denoted by ν_B . We are now ready to state the first main result.

Theorem 1.1. *[Theorem 3.1] Suppose G is a non-elementary relatively hyperbolic group and fix a finite generating set for G . Also suppose that μ is an admissible probability measure with finite first moment on G . Then there exists $\lambda_0 \in (0, 1)$ such that for every $\lambda \in [\lambda_0, 1)$,*

$$Hdim_{\delta_\lambda}(\nu_{\mathcal{F}}) = Hdim_{\bar{\delta}_\lambda}(\nu_B) = \frac{-1}{\log \lambda} \frac{h}{l}.$$

Moreover, the measure ν_B on $(\partial_B G, \bar{\delta}_\lambda)$ and the measure $\nu_{\mathcal{F}}$ on $(\partial_{\mathcal{F}} G, \delta_\lambda)$ are exact dimensional.

Denote by $S_n = \{g \in G : d(1, g) = n\}$ the sphere of radius n in the Cayley graph (Γ, d) . We define the *growth rate* of (the Cayley graph of) G as

$$v := \lim_{n \rightarrow \infty} \frac{\log \#S_n}{n}$$

The two quantities l and v both depending on the word metric of Γ are related to the entropy h which only depends on the measure μ by the following fundamental inequality

$$\frac{h}{l} \leq v$$

also called the Guivarc'h inequality [12]. This inequality holds for any μ -random walk.

In [45], it is proved that $\frac{-v}{\log \lambda}$ is the Hausdorff dimension of the Floyd and Bowditch boundaries. The following corollary thus follows from the strictness of the fundamental inequality established in [16, Theorem 1.3, Theorem 1.6] for certain relatively hyperbolic groups. Recall that μ has *finite super-exponential moment* if $\sum_{g \in G} \exp(cd(1, g))\mu(g)$ is finite for every $c > 0$.

Corollary 1.2. *Suppose G is a non-elementary relatively hyperbolic group and μ is an admissible probability measure on G with finite super-exponential moment. If one of the parabolic subgroups is virtually abelian of rank at least 2, or if the Bowditch boundary is homeomorphic to a sphere of dimension at least 2, then*

$$Hdim_{\bar{\delta}_\lambda}(\nu_B) < Hdim_{\bar{\delta}_\lambda}(\partial_B G).$$

1.2. Hausdorff dimension of the end boundary and of harmonic measures with respect to visual metrics. According to a celebrated result of Stallings [51], [52], any infinitely ended group G splits nontrivially as an amalgamated product $A *_C B$ or an HNN extension $A *_C$, where C is a finite group. The action on the corresponding Bass-Serre tree satisfies the conditions of [5, Definition 2] and so G is relatively hyperbolic. Moreover, in the former case, the maximal parabolic subgroups are exactly the conjugates of A and the conjugates of B . In the later case, they are exactly the conjugates of A . In addition to the Floyd and Bowditch boundary, G can be compactified using the Freudenthal (or end) boundary introduced by Freudenthal [19]. The interplay between the end boundary and asymptotic properties of random walks is a well-studied subject, see for instance [43], [44], [57].

Let $\partial_{\mathcal{E}} G$ be the end boundary of an infinitely ended group G . The topology of the end boundary is independent of the choice of Cayley graph. For every $\lambda \in (0, 1)$, we define a *visual metric*, extending the definition of Candellero, Gilch and Müller [6] for free products. This metric was independently studied by Cornuier in [10]. Fixing a generating set, let Γ be the Cayley graph of G and let ρ_λ be the visual metric defined on the end compactification of $\partial_{\mathcal{E}} G \cup \Gamma$. Precisely, define the distance $\rho_\lambda(\xi, \eta) = \lambda^n$ between two ends ξ, η if n is the minimal radius of the ball at a basepoint separating ξ and η . A quasi-isometry f induces a homeomorphism \tilde{f} between end boundaries. Moreover, this induced homeomorphism is *bi-Hölder*: for some $\alpha, \beta > 0, c > 1$ and for all pair (ξ, η) of points,

$$c^{-1} \rho_\lambda(\xi, \eta)^\alpha \leq \rho_\lambda(\tilde{f}(\xi), \tilde{f}(\eta)) \leq c \rho_\lambda(\xi, \eta)^\beta$$

which holds more generally for sublinearly bi-Lipschitz equivalence by [9, Corollary 1.4].

Our second main result is analogous to Theorem 1.1 and computes the Hausdorff dimension of the end boundary and of harmonic measures with respect to visual metrics. Let us denote by $\nu_{\mathcal{E}}$ the harmonic measure on the end boundary and h, l the entropy and drift of the μ -random walk. Also, let us denote by v the growth rate of G .

Theorem 1.3. *[Theorem 4.7, Theorem 5.1] Suppose G is a finitely generated group with infinitely many ends. Also suppose that μ is an admissible probability measure with finite first moment on G . Then, for every Cayley graph Γ and every $\lambda \in (0, 1)$,*

$$Hdim_{\rho_{\lambda}}(\nu_{\mathcal{E}}) = \frac{-1}{\log \lambda} \frac{h}{l}$$

and

$$Hdim_{\rho_{\lambda}}(\partial_{\mathcal{E}}G) = \frac{-v}{\log \lambda}.$$

Remark 1.4. The second equality was proved for free products with a standard generating set in [6]. Our proof is very different and applies to any infinitely ended group with any finite generating set.

The archetypal groups with infinitely many ends are free products of the form $A * B$, where $A \neq \mathbb{Z}/2\mathbb{Z}$ or $B \neq \mathbb{Z}/2\mathbb{Z}$. Using again the strictness of the fundamental inequality established in [16, Theorem 1.5] for certain free products, we can state the following corollary.

Corollary 1.5. *Suppose G is a free product $A * B$ and assume that A is a non-virtually cyclic nilpotent group. Let μ be a probability measure on G with finite super-exponential moment. Then,*

$$Hdim_{\rho_{\lambda}}(\nu_{\mathcal{E}}) < Hdim_{\rho_{\lambda}}(\partial_{\mathcal{E}}G).$$

1.3. Doubling property of the end boundary. The study of the harmonic measure associated with a random walk is closely related to the doubling property of the involved boundary, see [53, Section 4]. We also refer to [27, Proposition 4.12], where the doubling property is combined with the so-called shadow lemma to give a short proof of the formula $Hdim(\nu) = h/l$ in the setting of hyperbolic groups. It is not known whether the Floyd boundary equipped with the Floyd distance and the Bowditch boundary equipped with the shortcut Floyd distance are doubling, see [45, Question 1.7]. On the other hand, to give a more complete picture, we clarify the situation for the end boundary equipped with a visual distance.

A metric space (X, d) is called *doubling* if there exists a constant $N > 0$ such that every ball of radius $\delta > 0$ can be covered by at most N balls of radius $\delta/2$. Equivalently, for any $\theta \in (0, 1)$, there exists $N(\theta) > 0$ such that every ball of radius δ can be covered by at most $N(\theta)$ balls of radius $\theta\delta$. The doubling property is known to be a bi-Hölder invariant. A measure μ on (X, d) is called *doubling* if there exists a constant C such that $\mu(2B) \leq C\mu(B)$ for every ball B , where $2B$ denotes the ball with the same center and a radius twice as large as B . The existence of a doubling measure on a metric space (X, d) implies that the metric space (X, d) is doubling. The converse is true for complete metric spaces.

It is well-known that the Patterson-Sullivan measure on the Gromov boundary of a hyperbolic group is Ahlfors regular, hence doubling, for the Gromov's visual metric, so the Hausdorff dimension of the Patterson-Sullivan measure equals that of the whole boundary, see [8]. One can construct, as in [59], a class of Patterson-Sullivan measures on the end boundary through the action on the Cayley graph. Those measures yield quasi-conformal densities without atoms. This motivates the question whether the Patterson-Sullivan measure is Ahlfors regular or at least doubling on the end boundary endowed with a visual metric.

Our third main result gives a characterization of the doubling property of the end boundary for *accessible* infinitely-ended groups. Such groups admit a splitting as a graph of groups over finite edge groups so that the vertex groups are either finite or one-ended. It is a famous result of Dunwoody [14] that finitely presented group are accessible.

Theorem 1.6. *Let G be a finitely generated, accessible, infinitely-ended group. Let $\lambda \in (0, 1)$ and endow the end boundary $\partial_{\mathcal{E}}G$ of G with the visual metric ρ_{λ} . Then, $(\partial_{\mathcal{E}}G, \rho_{\lambda})$ is doubling if and only if G is virtually free.*

From the above discussion, the following two corollaries are immediate.

Corollary 1.7. *The Patterson-Sullivan measure on the end boundary of an accessible infinitely-ended group is doubling for the visual metric if and only if the group is virtually free.*

The following one addresses an analogous question [53, Question 4.2] in our setup.

Corollary 1.8. *If an accessible infinitely-ended group is not virtually free, then the harmonic measure ν_ε on the end boundary endowed with a visual metric is not doubling.*

A result of David-Semmes [29, Theorem 15.5] says that a metric space is quasi-symmetric to the standard Cantor ternary set if and only if it is compact, doubling, uniformly perfect and uniformly disconnected. Precise definitions of the latter two can be found in §11.1 and §14.24 in [29]. It is known that a doubling space is uniformly perfect and a ultrametric space is uniformly disconnected. The end boundary of a virtually free group with visual metric is doubling by [8] and thus quasi-symmetric to the standard Cantor ternary set.

By [29, Corollary 11.3], quasi-symmetric maps between uniformly perfect compact spaces are bi-Hölder. However, the converse is not true: bi-Hölder homeomorphisms are not necessarily quasi-symmetric. Interesting examples are given by the boundaries of the real hyperbolic space \mathbb{H}^4 and the complex hyperbolic space \mathbb{CH}^2 equipped with Gromov's visual metrics, which are bi-Hölder but not quasi-symmetric, since they can be distinguished by their conformal dimension. Thus, a bi-Hölder classification of visual metrics on the end boundary is a reasonable and interesting problem. We refer the reader to [9, Introduction 1D] for further discussion. Recalling that the doubling property is a bi-Hölder invariant, we obtain the following corollary, which answers positively a question of Cornuier [10, Question 1.27].

Corollary 1.9. *The end boundary of an accessible infinitely-ended group is bi-Hölder equivalent to the standard Cantor ternary set if and only if it is virtually free.*

In particular, the end boundaries of a free group and the free product of two 1-ended groups are not bi-Hölder equivalent.

Inaccessible groups do exist by work of Dunwoody [15]. The proof of Theorem 1.6 fails generally for inaccessible groups, but we can still give a result for such groups under some additional assumptions, see precisely Proposition 6.1.

Finally, let us compare the visual metric on the end boundary with the Gromov's visual metric coming from an action of G on a hyperbolic space X . Fix a splitting of an infinitely-ended group G over finite groups as a finite graph of groups. As explained above, by [5, Definition 2], this splitting makes G hyperbolic relative to the set of vertex groups. Hence, by Yaman [58], G acts via a geometrically finite action on a proper hyperbolic space X so that the Bowditch boundary is homeomorphic to the Gromov boundary of X . If G is accessible and the splitting is terminal, then the end boundary is equivariantly homeomorphic to the Bowditch boundary, see Section 4.3 for more details. We can thus endow the end boundary with the Gromov's visual metric coming from the hyperbolic space X on which G acts. According to [11] and [2], X can be chosen so that this metric is doubling if and only if the parabolic subgroups are virtually nilpotent. In such a situation, by Theorem 1.6 the two possible metrics on the end boundary cannot be in the same bi-Hölder class unless the group is virtually free, for one is doubling and the other is not.

Corollary 1.10. *Assume that G is not virtually free and splits over finite groups as a finite graph of virtually nilpotent groups. Then, there exists a hyperbolic space X on which G acts via a geometrically finite action such that the visual metric on the end boundary is not bi-Hölder equivalent to the Gromov's visual metric coming from this action.*

Remark 1.11. Similarly as in Corollary 1.9, we can derive the following result from [11]. Let $G_1 = H_1 * \mathbb{Z}$ and $G_2 = H_2 * \mathbb{Z}$, where H_1 is virtually nilpotent and H_2 is one-ended but not virtually nilpotent. Then G_1 and G_2 admit geometrically finite actions on proper hyperbolic spaces X_1 and X_2 with bounded geometry so that their Gromov boundaries endowed with Gromov's visual metric are homeomorphic to the Cantor sets but are not bi-Hölder equivalent, for one boundary is doubling and not the other.

Overview and organisation of the paper. In Section 2, we review all the preliminary results we will need in the following. We recall the definition of relatively hyperbolic groups, the Bowditch boundary, the Floyd distance and the Floyd boundary. We also give more details about the Hausdorff dimension of a finite measure on a metric space.

Section 3 is devoted to the proof of Theorem 1.1, which treats separately the upper bound (Proposition 3.6) and lower bound (Proposition 3.14) for the Hausdorff dimension. The lower bound follows a strategy similar to the one developed by Tanaka [53]. One of the main tools in [53] is that the random walk sub-linearly

tracks geodesics $[1, \omega_\infty]$ on the hyperbolic space X on which the group G acts, where ω_∞ is the limit of the random walk in the Gromov boundary of X .

In our situation, the Cayley graph of a relatively hyperbolic group G is generally not hyperbolic anymore, so it cannot be expected that the random walk stay close to any point on a geodesic $[1, \omega_\infty]$ in the Cayley graph. However, using Maher-Tiozzo [39] and Tiozzo [55], we can prove that the random walk sub-linearly tracks word geodesics along transition points, which are the points that are not deep in parabolic subgroups, see Definition 2.2. Along the way, we also prove that the random walk spends at most sub-linear time in parabolic subgroups. Precisely, we prove the following result. For a given word geodesic α , let $\text{Tr } \alpha$ be the set of transition points on α .

Theorem 1.12. *[Proposition 3.2, Corollary 3.4] Suppose G is a non-elementary relatively hyperbolic group and fix a finite generating set for G . Also suppose that μ is an admissible probability measure with finite first moment on G . Then,*

$$(4) \quad \mathbb{P}(\sup d(\omega_n, \text{Tr } \alpha) = o(n)) = 1,$$

where the supremum is taken over all geodesics α from 1 to the limit ω_∞ of the random walk in the Bowditch boundary of G . Moreover,

$$(5) \quad \mathbb{P}\left(\sup_{U \in \mathcal{P}} d_U(1, \omega_n) = o(n)\right) = 1,$$

where \mathcal{P} is the set of all left cosets of a chosen full family of conjugacy classes of parabolic subgroups and where $d_U(x, y)$ is the distance between the projections of x and y on U .

This last result is a weak version (under a finite moment condition) of the results in [50] for finitely supported random walks. Indeed, it is proved in [50, Theorem 2.3] that such a random walk spends at most logarithmic time in parabolic subgroups.

With the sublinear tracking of transition points at hand, we use the estimates of Floyd disks by shadows in [45] to obtain the lower bound of Hausdorff dimension.

In Sections 4 and 5, we prove Theorem 1.3. We first extend the definition of visual metrics introduced in [6] to any group with infinitely many ends and we show that

$$\text{Hdim}_{\rho_\lambda}(\nu_\mathcal{E}) = \frac{-1}{\log \lambda} \frac{h}{l}$$

in Section 4. Our proof is again similar to the proof in [53]. We introduce the notion of bottleneck : a set V is a bottleneck between two points x and y if any path from x to y has to pass through a fixed neighborhood of V . We then replace the sub-linear tracking of transition points by the sub-linear tracking of bottlenecks, see precisely Proposition 4.9.

We then prove that

$$\text{Hdim}_{\rho_\lambda}(\partial_\mathcal{E} G) = \frac{-v}{\log \lambda}$$

in Section 5. The proof follows the outline of an analogous result for the Bowditch and Floyd boundaries endowed with the Floyd (shortcut) distance [45]. It is well-known that there exists a continuous and surjective map from $\partial_\mathcal{F} G$ to $\partial_\mathcal{E} G$ (see [36] and [20]). We further observe that the Floyd metric dominates the visual metric through the map. This gives the desired upper bound of $\partial_\mathcal{E} G$ by the same bound in [45] on the Hausdorff dimension of $\partial_\mathcal{F} G$. Moreover, inspired by transition points, we use an enhanced version of the notion of bottlenecks introduced above to construct a sequence of free semi-subgroups of G whose set of ends has Hausdorff dimension arbitrarily close to $\frac{-v}{\log \lambda}$. This proves the lower bound of $\partial_\mathcal{E} G$. A technical result in its proof is Proposition 5.13 saying that endowed with visual metric of corrected parameter depending on λ , the end boundary $\partial_\mathcal{E} T$ of free semi-groups is bi-Lipschitz embedded into $\partial_\mathcal{E} G$ with visual metric ρ_λ . This greatly simplifies the arguments in [45, Section 3] using Patterson-Sullivan measure to estimate the Hausdorff dimension of $\partial_\mathcal{E} T$.

Finally, Section 6 deals with Theorem 1.6. We consider an accessible infinitely ended group G . Then, G admits a splitting over finite edge groups as a finite graph of groups \mathcal{G} , so that the vertex groups either are finite or one-ended. If every vertex group is finite, then G is virtually free so that we can assume that one of the vertex group is one-ended. Denote by H such a vertex group. The unique end ξ of H embeds into the end boundary of G . We then show that for some fixed θ , the ball of radius λ^n centered at ξ cannot be

covered by $N(n)$ balls of radius $\theta\lambda^n$, where $N(n)$ goes to infinity, as n tends to infinity, which concludes the proof.

Acknowledgments. W.Y. is grateful to Giulio Tiozzo for explaining the results of [55] to him and also thanks Yves Cornuier for many helpful comments and corrections about the Hölder structure of the end boundary. W. Y. is supported by the National Natural Science Foundation of China (No. 11771022).

2. PRELIMINARIES

2.1. Relatively hyperbolic groups. We now properly define relatively hyperbolic groups and recall several tools and results that will be used in the paper. Let G be a finitely generated group. The action of G on a compact Hausdorff space T is called a *convergence action* if the induced action on triples of distinct points of T is properly discontinuous. Since G is countable, T must be metrizable by [21, Main Theorem]. Equivalently, the action $G \curvearrowright T$ is convergence if and only if every sequence of distinct elements g_n in G contains a subsequence g_{n_k} such that $g_{n_k} \cdot x \rightarrow a$ and for all $x \in X$ with at most perhaps one exceptional point.

The set of accumulation points ΛG of any orbit $G \cdot x$ ($x \in T$) is called the *limit set* of the action. As long as ΛG has more than two points, it is uncountable and it is then the unique minimal closed G -invariant subset of T . The action is then said to be *non-elementary*. In this case, the orbit of every point in ΛG is infinite. The action is *minimal* if $\Lambda G = T$.

A point $\zeta \in \Lambda G$ is called *conical* if there is a sequence g_n of G and distinct points $\alpha, \beta \in \Lambda G$ such that $g_n \cdot \zeta \rightarrow \alpha$ and $g_n \cdot \eta \rightarrow \beta$ for all $\eta \in T \setminus \{\zeta\}$. The point $\zeta \in \Lambda G$ is called *bounded parabolic* if it is the unique fixed point of its stabilizer in G , which is infinite and acts cocompactly on $\Lambda G \setminus \{\zeta\}$. The stabilizers of bounded parabolic points are called *maximal parabolic subgroups*. The convergence action $G \curvearrowright T$ is called *geometrically finite* if every point of $\Lambda G \subset T$ is either conical or bounded parabolic.

Definition 2.1. Let \mathcal{P} be a collection of subgroups of G . We say that G is *hyperbolic relative to \mathcal{P}* if there exists some compact Hausdorff space T on which G acts minimally and geometrically finitely and such that the maximal parabolic subgroups are exactly the elements of \mathcal{P} .

In this situation, Yaman [58] proved that there exists a proper geodesic hyperbolic space X on which G acts such that the Gromov boundary of X equivariantly coincides with T . Further, Bowditch [5] proved that the Gromov boundary of such a space X is unique up to homeomorphism, hence so is T . We call T the *Bowditch boundary* of G and we will denote it by $\partial_{\mathcal{B}}G$ in the following. The union $G \cup \partial_{\mathcal{B}}G$ is called the Bowditch compactification.

Following Osin [41], we define the relative Cayley graph as follows. We start with the Cayley graph Γ associated with a finite generating set S . We choose a system \mathcal{P}_0 of representatives of conjugacy classes of maximal parabolic subgroups. Such a system is finite by [5, Proposition 6.10]. The relative Cayley graph \hat{G} is obtained by adding one edge of length 1 between every two elements in the same left coset of a parabolic subgroup in \mathcal{P}_0 . In other words, setting $\mathcal{P}_0 = \{P_1, \dots, P_N\}$, the relative Cayley graph \hat{G} is the Cayley graph associated with the generating set $S \cup P_1 \cup \dots \cup P_N$. A (quasi-)geodesic in the relative Cayley graph \hat{G} is called a relative (quasi-)geodesic. This graph \hat{G} is quasi-isometric to the coned-off graph introduced by Farb [17] and is hyperbolic in the sense of Gromov.

A sequence g_n in G converges to a point ξ in the Gromov boundary $\partial \hat{G}$ of \hat{G} if and only if it converges to a conical limit point in the Bowditch compactification [3, Section 8]. We can thus identify $\partial \hat{G}$ with the set of conical limit points. We refer to [56] for more details on the comparison of these two boundaries.

A very useful tool when studying the geometry of a relatively hyperbolic group is the notion of transition points on a geodesic.

Definition 2.2. Let γ be a (finite or infinite) geodesic in the Cayley graph of G . A point v on γ is said to be (ϵ, R) -deep if there exists $g \in \Gamma$ and $P \in \mathcal{P}_0$ such that the R -neighborhood of v in γ is contained in the ϵ -neighborhood of gP . A point v on γ is called an (ϵ, R) -transition point if it is not (ϵ, R) -deep.

The following result of Hruska relates transition points and points on relative geodesics.

Lemma 2.3. [30, Proposition 8.13] For every large enough ϵ and R , there exists C such that the following holds. Let α be a (finite or infinite) geodesic in G and let $\hat{\alpha}$ be a relative geodesic with the same endpoints.

Then, any point on $\hat{\alpha}$ is within a distance at most C of an (ϵ, R) -transition point on α . Conversely, any (ϵ, R) -transition point on α is within a distance at most C of a point on $\hat{\alpha}$.

2.2. The Floyd distance and the Floyd boundary. We first recall the definition of the Floyd distance and the Floyd boundary and their relation with the Bowditch boundary. This boundary was introduced by Floyd in [18] and we also refer to [35] and [36] for more details.

Let G be a finitely generated group and let Γ denote its Cayley graph associated with a finite generating set. Let $f : \mathbb{N} \rightarrow \mathbb{R}$ be a function satisfying that the sum $\sum_{n \geq 0} f_n$ is finite and that there exists $\lambda \in (0, 1)$ such that $1 \geq f_{n+1}/f_n \geq \lambda$ for all $n \in \mathbb{N}$. The function f is then called the *rescaling function* or the *Floyd function*. In the following, we will always choose an exponential Floyd function, that is the function f will be of the form $f(n) = \lambda^n$ for some $\lambda \in (0, 1)$. Fix a basepoint $o \in \Gamma$ and rescale Γ by declaring the length of an edge σ to be $f(d(o, \sigma))$. The induced shortpath metric on Γ is called the *Floyd distance* with respect to the basepoint o and Floyd function f and is denoted by $\delta_{f,o}(\cdot, \cdot)$. Whenever f is of the form $f(n) = \lambda^n$, we will write $\delta_{\lambda,o} = \delta_{f,o}$ and if $o = 1$, $\delta_\lambda = \delta_{f,o}$.

The Floyd compactification $\bar{\Gamma}^{\mathcal{F}}$ of Γ is the Cauchy completion of Γ endowed with the Floyd metric. The Floyd boundary is then defined as $\partial_{\mathcal{F}}\Gamma = \bar{\Gamma}^{\mathcal{F}} \setminus \Gamma$. Different choices of basepoints yield bi-Lipschitz homeomorphisms of the Floyd compactifications. However, the topology may depend on the choice of the generating set and the rescaling function. Keeping in mind $f(n) = \lambda^n$ and a choice of Cayley graph, we will also call $G \cup \partial_{\mathcal{F}}\Gamma$ the Floyd compactification of G by abuse of language and we will write $\partial_{\mathcal{F}}G = \partial_{\mathcal{F}}\Gamma$.

The cardinality of the Floyd boundary is 0, 1, 2 or uncountable. Moreover, it is 2 if and only if the group G is virtually infinite cyclic, see [36, Proposition 7]. Following Karlsson, we say that the Floyd boundary is trivial if it is finite. We will only have to deal with groups with non-trivial Floyd boundary.

Finally, as mentioned in the introduction, whenever the Floyd boundary is non-trivial, G acts on it as a convergence action, see [36, Theorem 2]. Also, whenever the Floyd boundary is non-trivial, for any probability measure μ with finite first moment on G and whose support generates G as a semi-group, the random walk driven by μ almost surely converges to a point in the Floyd boundary. Letting $\nu_{\mathcal{F}}$ be the law of the limit point, the pair $(\partial_{\mathcal{F}}G, \nu_{\mathcal{F}})$ is a model for the Poisson boundary, see [35, Section 6, Corollary].

We now assume that G is non-elementary relatively hyperbolic. We denote by $\partial_{\mathcal{B}}G$ its Bowditch boundary. The following is due to Gerasimov.

Theorem 2.4. [22, Map Theorem] *There exists $\lambda_0 \in (0, 1)$ such that for every $\lambda \in [\lambda_0, 1)$, the identity of G extends to a continuous and equivariant surjection ϕ from the Floyd compactification to the Bowditch compactification of G .*

Actually, Gerasimov only stated the existence of the map ϕ for one Floyd function $f_0 = \lambda_0^n$, but then Gerasimov and Potyagailo proved that the same result holds for any Floyd function $f \geq f_0$, see [23, Corollary 2.8]. They also proved that the preimage of a conical limit point is reduced to a single point and described the preimage of a parabolic limit point in terms of the action of G on $\partial_{\mathcal{F}}G$, see precisely [23, Theorem A]. From now on, the parameter λ will always be assumed to be contained in $[\lambda_0, 1)$.

The Floyd distance can be transferred to a distance on the Bowditch boundary using the map ϕ . The resulting distance is called the *shortcut metric* and we denote it by $\bar{\delta}_\lambda$. It is the largest distance on the Bowditch boundary satisfying that for every $\xi, \zeta \in \partial_{\mathcal{F}}G$,

$$(6) \quad \bar{\delta}_\lambda(\phi(\xi), \phi(\zeta)) \leq \delta_\lambda(\xi, \zeta).$$

We refer to [24, Section 4] for more details on its construction. The next couple of lemmas will be used later on.

Lemma 2.5 (Visibility lemma). [36] *For every fixed λ, c , there exists a function $\varphi : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ such that for any $v \in G$ and any (λ, c) -quasi-geodesic γ in Γ , the following holds. If $\delta_{\lambda,v}(\gamma) \geq \kappa$, then $d(v, \gamma) \leq \varphi(\kappa)$.*

Note that [36] only deals with geodesics, but the proof applies to quasi-geodesic with fixed parameters, see [23] where this and more general cases are discussed.

The *big shadow* $\Pi(g, R)$ at g is the set of boundary points ξ in the Bowditch boundary such that there exists a geodesic ray $[1, \xi]$ intersecting the ball $B(g, R)$.

Lemma 2.6. [45, Lemma 3.14, Lemma 3.15] *For every large enough ϵ and R , the following holds. Let ξ be a conical limit point in the Bowditch boundary. Consider a geodesic γ between 1 and ξ and consider any point g on this geodesic. There exists C_1, C_2 such that*

$$\Pi(g, R) \subset B_{\delta_\lambda}(\xi, C_1 r)$$

and if, in addition, g is a (ϵ, R) -transition point on γ , then

$$B_{\delta_\lambda}(\xi, C_2 r) \subset \Pi(g, R),$$

where $r = \lambda^{d(1, g)}$.

Actually, [45, Lemma 3.16] also states that the above statement is true for the Floyd distance, but we will only need to use it for the shortcut distance in estimating the lower bound of Hausdorff dimensions (see Lemma 3.9 and Proposition 3.14).

Lemma 2.7. *For every $g \in G$, there exists a constant $c_g > 0$ such that the following inclusions hold. For every point ξ in the Floyd boundary and for every $r \geq 0$,*

$$B_{\delta_\lambda}(g\xi, c_g^{-1}r) \subset gB_{\delta_\lambda}(\xi, r) \subset B_{\delta_\lambda}(g\xi, c_g r).$$

For every point ξ in the Bowditch boundary,

$$B_{\delta_\lambda}(g\xi, c_g^{-1}r) \subset gB_{\delta_\lambda}(\xi, r) \subset B_{\delta_\lambda}(g\xi, c_g r).$$

Proof. First, a change a base point induces a bi-Lipschitz inequality for the Floyd distance : for every x, y in the Floyd compactification, for any basepoints $o, o' \in G$,

$$\lambda^{d(o, o')} \leq \frac{\delta_{\lambda, o}(x, y)}{\delta_{\lambda, o'}(x, y)} \leq \lambda^{-d(o, o')},$$

see [45, (2)]. Now, by definition, the same is true of the shortcut distance, see precisely [45, (3)].

We only give the proof of the lemma for the Floyd distance, the proof for the shortcut distance is exactly the same. Let $\zeta \in B_{\delta_\lambda}(\xi, r)$. We need to prove that $g\zeta \in B_{\delta_\lambda}(g\xi, c_g r)$ for some c_g . Note that $\delta_{\lambda, g}(g\zeta, g\xi) = \delta_{\lambda, 1}(\zeta, \xi)$, so that by the above discussion,

$$\lambda^{d(1, g)} \leq \frac{\delta_{\lambda, 1}(g\xi, g\zeta)}{\delta_{\lambda, 1}(\xi, \zeta)} \leq \lambda^{-d(1, g)}.$$

This proves that $\delta_{\lambda, 1}(g\xi, g\zeta) \leq \lambda^{-d(1, g)}r$ and so the right inclusion in the lemma holds for $c_g = \lambda^{-d(1, g)}$. We immediately deduce the left inclusion, using g^{-1} . \square

2.3. Hausdorff dimension of measures. Let (X, d) be a metric space and κ be a Borel measure on X .

Definition 2.8. *The Hausdorff dimension of κ is the smallest possible Hausdorff dimension of a set of full κ -measure :*

$$Hdim(\kappa) = \inf\{Hdim(E), \kappa(E^c) = 0\}.$$

When we want to insist on the choice of the distance, we will write $Hdim_d(\kappa)$.

Evaluating the Hausdorff dimension of a set can be a difficult task, so the following characterization of the $Hdim(\kappa)$ as the essential supremum of the local dimensions of κ is very useful. Recall that the essential supremum $\kappa - \sup f$ of a function f is defined as the infimum of the constants C such that $f \leq C$ κ -almost everywhere.

Proposition 2.9. [40, Corollary 8.2] *Let κ be a Borel measure on a metric space X . Then,*

$$Hdim(\kappa) = \kappa - \sup \liminf_{r \rightarrow 0} \frac{\log \kappa(B(x, r))}{\log r}.$$

Definition 2.10. *A measure κ on a metric space X is exact dimensional if for κ -almost every x , the above \liminf is a limit, that is, for κ -almost every x ,*

$$\lim_{r \rightarrow 0} \frac{\log \kappa(B(x, r))}{\log r} = Hdim(\kappa).$$

We will also use the following notation :

$$\overline{Hdim}(\kappa) = \kappa - \sup_{r \rightarrow 0} \limsup \frac{\log \kappa(B(x, r))}{\log r}.$$

By definition, $Hdim(\kappa) \leq \overline{Hdim}(\kappa)$. Let us say a few words about our strategy for evaluating the Hausdorff dimensions of harmonic measures ν . We will first prove that for ν -almost every x , we have $\limsup_{r \rightarrow 0} \frac{\log \nu(B(x, r))}{\log r} \leq \frac{-1}{\log \lambda} \frac{h}{l}$, so that $\overline{Hdim}(\nu) \leq \frac{-1}{\log \lambda} \frac{h}{l}$. We will then prove that for ν -almost every x , we have $\frac{-1}{\log \lambda} \frac{h}{l} \leq \liminf_{r \rightarrow 0} \frac{\log \nu(B(x, r))}{\log r}$, so that $\frac{-1}{\log \lambda} \frac{h}{l} \leq Hdim(\nu)$. This will both prove that $Hdim(\nu) = \frac{-1}{\log \lambda} \frac{h}{l}$ and that ν is exact dimensional.

3. HARMONIC MEASURES ON THE FLOYD AND THE BOWDITCH BOUNDARIES

Let G be a finitely generated non-elementary relatively hyperbolic group and let μ be a probability measure with finite first moment on G . Throughout this section, we consider the harmonic measure denoted by $\nu_{\mathcal{B}}$ on the Bowditch boundary $\partial_{\mathcal{B}}G$ equipped with the shortcut distance $\bar{\delta}_{\lambda}$ and the harmonic measure $\nu_{\mathcal{F}}$ on the Floyd boundary $\partial_{\mathcal{F}}G$ with the Floyd distance δ_{λ} .

Our goal in this section is to prove the following theorem.

Theorem 3.1. *For any $\lambda \in [\lambda_0, 1)$ with $\lambda_0 \in (0, 1)$ given by Theorem 2.4, we have*

$$Hdim_{\bar{\delta}_{\lambda}}(\nu_{\mathcal{B}}) = \frac{-1}{\log \lambda} \frac{h}{l}$$

and, for any $\lambda \in (0, 1)$,

$$Hdim_{\delta_{\lambda}}(\nu_{\mathcal{F}}) = \frac{-1}{\log \lambda} \frac{h}{l}.$$

Moreover, those two measures are exact-dimensional.

3.1. Sublinear deviation from transition points. We denote by \hat{G} the relative Cayley graph of G . We fix large enough $\epsilon > 0$ and $R > 0$ satisfying the conclusions of Lemma 2.3 and Lemma 2.6. Whenever α is a geodesic in the Cayley graph of G , we denote by $\text{Tr}_{\epsilon, R} \alpha$, or simply by $\text{Tr} \alpha$ the set of (ϵ, R) -transition points on α . We denote by ω_n the random walk driven by μ at time n and by ω_{∞} its almost sure limit in the Bowditch boundary.

Proposition 3.2. *With these notations, we have*

$$\mathbb{P}(\sup d(\omega_n, \text{Tr} \alpha) = o(n)) = 1,$$

where the supremum is taken over all geodesics α from 1 to ω_{∞} .

Proof. We follow the strategy of [55]. We introduce the function f defined by

$$f(\omega) = \sup d(1, \text{Tr} \hat{\alpha}),$$

where the supremum is taken over all geodesics $\hat{\alpha}$ between $\omega_{-\infty}$ and ω_{∞} , where $\omega_{-\infty}$ is the limit of the reflected random walk, see [39, Section 4.1].

Claim 3.3. *The function f is measurable and is almost surely finite.*

Proof of the claim. First, according to [39, Theorem 1.1], the exit points ω_{∞} and $\omega_{-\infty}$ are conical limit points and their law ν and $\tilde{\nu}$ are non-atomic. Since bounded parabolic points are countable, it follows that ω_{∞} and $\omega_{-\infty}$ almost-surely are distinct and so there exists a bi-infinite relative geodesic joining them. Hence, the distance (in the Cayley graph) between 1 and such a relative geodesic is finite. According to Lemma 2.3, any point on a relative geodesic is within a finite (and actually uniformly bounded) distance of a transition point on a geodesic in the Cayley graph, so f is almost surely finite.

We now prove that f is measurable. Recall that $\partial \hat{G}$ is the Gromov boundary of the relative Cayley graph that we identify with the set of conical limit points in the Bowditch boundary $\partial_{\mathcal{B}}G$. We just need to prove that the function

$$\tilde{f} : (\xi, \zeta) \in \partial \hat{G} \times \partial \hat{G} \mapsto \sup d(1, \text{Tr} \alpha_{\xi, \zeta})$$

is measurable, where the supremum is taken over all word geodesics $\alpha_{\xi, \zeta}$ from ξ to ζ . We follow the proof of [55, Lemma 12]. Note that \tilde{f} takes the value $+\infty$ when $\xi = \zeta$. Since there are no atoms at conical points, it

can be extended to a function that we still denote by \tilde{f} on the double Bowditch boundary $\partial_{\mathcal{B}}G \times \partial_{\mathcal{B}}G$. To prove that \tilde{f} is measurable, we just need to prove that for every $R \geq 0$, the set

$$\{(\xi, \zeta) \in \partial_{\mathcal{B}}G \times \partial_{\mathcal{B}}G, \tilde{f}(\xi, \zeta) > R\}$$

is measurable.

Recall the Bowditch compactification is a metrizable compact space containing the group G as an open and dense set. Choosing an arbitrary metric and taking a finite cover of $\partial_{\mathcal{B}}G$ made of balls of radius $1/k$, $k \in \mathbb{N}$, we construct a countable collection of open sets U_n such that the sets $U_n \cap \partial_{\mathcal{B}}G$ form a countable base for the topology of $\partial_{\mathcal{B}}G$. Moreover, for each $R \geq 0$, only finitely many sets U_n intersect the ball $B(1, R)$ and for each sequence n_k going to infinity, the intersection $\cap_k U_{n_k}$ contains at most one point. For fixed $R \geq 0$, say that a pair of open sets (U, V) avoids the ball $B(1, R)$ if there exist $u \in U \cap G$ and $v \in V \cap G$ and there exists a geodesic γ from u to v such that the ball $B(1, R)$ does not intersect $\text{Tr } \gamma$. Let us define $\mathcal{S}_R = \{(U_n, U_m) \text{ such that } (U_n, U_m) \text{ avoids the ball } B(1, R)\}$. This is a countable collection of pairs of open sets. By definition,

$$\{(\xi, \zeta) \in \partial_{\mathcal{B}}G \times \partial_{\mathcal{B}}G, \tilde{f}(\xi, \zeta) > R\} \subset \bigcap_{N \geq 1} \bigcup_{\substack{\min(n, m) \geq N \\ (U_n, U_m) \in \mathcal{S}_R}} U_n \times U_m.$$

Conversely, consider (ξ, ζ) in this intersection and assume that $\tilde{f}(\xi, \zeta) < +\infty$. Then, there are sequences of points g_{n_k} , respectively h_{m_k} , converging to ξ , respectively ζ and there is a geodesic γ_k from g_{n_k} to h_{m_k} such that $B(1, R)$ does not intersect $\text{Tr } \gamma_k$. Now, since $\tilde{f}(\xi, \zeta) < +\infty$, there exists a geodesic from ξ to ζ intersecting some big ball $B(1, R')$ for some $R' = R'_{\xi, \zeta}$. Moreover, up to taking R' large enough, all geodesics γ_k also enter $B(1, R')$. Thus, Arzelà-Ascoli Theorem allows us to choose a sub-sequence of geodesics γ_{k_l} converging to a geodesic γ from ξ to ζ , as l tends to infinity. We can also assume that the sub-geodesic of γ_{k_l} contained in $B(1, R')$ is constant. In particular, the limit geodesic γ also satisfies that $B(1, R)$ does not intersect $\text{Tr } \gamma$. Hence, $\tilde{f}(\xi, \zeta) > R$. This proves that

$$\{(\xi, \zeta) \in \partial_{\mathcal{B}}G \times \partial_{\mathcal{B}}G, \tilde{f}(\xi, \zeta) > R\} = \bigcap_{N \geq 1} \bigcup_{\substack{\min(n, m) \geq N \\ (U_n, U_m) \in \mathcal{S}_R}} U_n \times U_m$$

and so $\{(\xi, \zeta) \in \partial_{\mathcal{B}}G \times \partial_{\mathcal{B}}G, \tilde{f}(\xi, \zeta) > R\}$ is measurable. \square

Note that $f(T^n \omega) = d(\omega_n, \text{Tr}[\omega_{-\infty}, \omega_{\infty}])$, so that $|f(T\omega) - f(\omega)| \leq d(1, \omega_1)$. We deduce that the function $\omega \mapsto f(T\omega) - f(\omega)$ is integrable, since the random walk has finite first moment. Thus, [55, Lemma 7] shows that $\frac{1}{n}f(T^n \omega)$ almost surely converges to 0. Finally, consider a geodesic α_0 from 1 to ω_{∞} and a geodesic $\hat{\alpha}_0$ from $\omega_{-\infty}$ to ω_{∞} . Then, with probability one, there exists a transition point x_n on $\hat{\alpha}_0$ such that $\frac{1}{n}d(\omega_n, x_n)$ converges to 0. We now use that geodesic triangles are thin along transition points. Precisely, according to [16, Lemma 2.4], x_n is within a uniformly bounded distance of a transition point either on α_0 , or on a geodesic from 1 to $\omega_{-\infty}$. Note that ω_n converges to ω_{∞} and that $\frac{1}{n}d(1, \omega_n)$ almost surely converges to l . Hence, x_n also converges to ω_{∞} and so for large enough n , it cannot be within a bounded distance of a geodesic from 1 to $\omega_{-\infty}$. This proves that $\frac{1}{n}d(\omega_n, \text{Tr } \alpha_0)$ also almost surely converges to 0. Using again [16, Lemma 2.4], we see that $\frac{1}{n} \sup_{\alpha} d(\omega_n, \text{Tr } \alpha) \leq \frac{1}{n}d(\omega_n, \text{Tr } \alpha_0) + C$ for some uniform C . This concludes the proof. \square

We now deduce that the projection on parabolic subgroup is almost surely sublinear. We choose a full subset \mathcal{P}_0 of representatives of conjugacy classes of maximal parabolic subgroups. According to [5, Proposition 6.10], such a set \mathcal{P}_0 is finite. In the following, we will denote by \mathcal{P} the set of all left cosets of elements of \mathcal{P}_0 . Let $U \in \mathcal{P}$. Whenever $x, y \in G$, we set $d_U(x, y) = d(\pi_U(x), \pi_U(y))$, where π_U is the projection on U and where d is the distance in the Cayley graph of G . Our goal is to prove the following corollary.

Corollary 3.4. *With the above notations, we have*

$$\mathbb{P} \left(\sup_{U \in \mathcal{P}} d_U(1, \omega_n) = o(n) \right) = 1.$$

Before proving this corollary, note the following.

Lemma 3.5. *The sequence $f_n = \sup_{U \in \mathcal{P}} d_U(1, \omega_n)$ is sub-additive. That is, for every $n, m \geq 1$, we have*

$$f_{n+m} \leq f_n + f_m \circ T^n.$$

Proof. Consider some $U \in \mathcal{P}$. Then, $d_U(1, \omega_{n+m}) \leq d_U(1, \omega_n) + d_U(\omega_n, \omega_{n+m})$, which we can rewrite $d_U(1, \omega_{n+m}) \leq d_U(1, \omega_n) + d_{\omega_n^{-1}U}(1, \omega_n^{-1}\omega_{n+m})$. In particular,

$$d_U(1, \omega_{n+m}) \leq \sup_{U \in \mathcal{P}} d_U(1, \omega_n) + \sup_{U \in \mathcal{P}} d(1, \omega_n^{-1}\omega_{n+m}) = f_n + f_m \circ T^n.$$

This is true for all $U \in \mathcal{P}$, which concludes the proof. \square

According to Kingman's Theorem, $\frac{1}{n}f_n$ almost surely converges to some constant $l_{\mathcal{P}}$ that we call the parabolic linear drift. We just need to prove that $l_{\mathcal{P}} = 0$ to prove Corollary 3.4.

Proof of Corollary 3.4. By definition,

$$\sup_{U \in \mathcal{P}} d_U(\omega_n, \omega_{2n}) = \sup_{U \in \mathcal{P}} d_{\omega_n^{-1}U}(1, \omega_n^{-1}\omega_{2n}) = \sup_{U \in \mathcal{P}} d_U(1, \omega_n^{-1}\omega_{2n})$$

and since $\omega_n^{-1}\omega_{2n}$ and ω_n follow the same law, $\frac{1}{n} \sup_{U \in \mathcal{P}} d_U(\omega_n, \omega_{2n})$ also almost surely converges to $l_{\mathcal{P}}$.

Combining this with Proposition 3.2, we see that for every positive ϵ and η , there exists $N_{\epsilon, \eta}$ such that with probability at least $1 - \epsilon$, for all $n \geq N_{\epsilon, \eta}$, we have simultaneously

- (a) $\sup_{\alpha} d(\omega_n, \text{Tr } \alpha) \leq \eta n$, where the supremum is taken over all geodesics α from 1 to ω_{∞} ,
- (b) $|\sup_{U \in \mathcal{P}} d_U(\omega_n, \omega_{2n}) - l_{\mathcal{P}} n| \leq \eta n$,
- (c) $|\sup_{U \in \mathcal{P}} d_U(1, \omega_n) - l_{\mathcal{P}} n| \leq \eta n$.

Fix ϵ and η and set $N = N_{\epsilon, \eta}$. For every $n \geq N$, there exists a transition point (with fixed parameters) x_n on a geodesic α_n from 1 to ω_{∞} , such that $d(\omega_n, x_n) \leq \eta n$. Then, since for every $U \in \mathcal{P}$ and for every x, y , we have $d_U(x, y) \leq d(x, y) + c$ for some fixed c ,

$$\sup_{U \in \mathcal{P}} d_U(1, x_n) \leq \sup_{U \in \mathcal{P}} d_U(1, \omega_n) + d(\omega_n, x_n) + c \leq l_{\mathcal{P}} n + 2\eta n + c$$

and similarly,

$$\sup_{U \in \mathcal{P}} d_U(x_n, x_{2n}) \leq l_{\mathcal{P}} n + 3\eta n + 2c.$$

Also,

$$\sup_{U \in \mathcal{P}} d_U(1, \omega_{2n}) \leq \sup_{U \in \mathcal{P}} d_U(1, x_{2n}) + \eta n + c.$$

By definition, x_{2n} is a transition point on a geodesic α_{2n} from 1 to ω_{∞} . Let $\delta > 0$ and consider $U \in \mathcal{P}$ such that $d_U(1, x_{2n}) \geq \sup_{U \in \mathcal{P}} d_U(1, x_{2n}) - \delta$. Then, according to [49, Lemma 1.13, Lemma 1.15], the geodesic α_{2n} enters a fixed neighborhood of U and the first point, respectively last point in this neighborhood is within a bounded distance of $\pi_U(1)$, respectively $\pi_U(x_{2n})$. Now, x_n also is a transition point on α_n with the same endpoints as α_{2n} , so it is within a uniformly bounded distance of a transition point \tilde{x}_n on α_{2n} . There are two possibilities : the geodesic α_{2n} enters the neighborhood of U either before or after \tilde{x}_n . In the former case, we see that $d_U(1, x_{2n}) \leq d_U(1, x_n) + c'$ for some fixed constant c' . In the later case, we see that $d_U(1, x_{2n}) \leq d_U(x_n, x_{2n}) + c'$. In any case, we have that

$$d_U(1, x_{2n}) \leq \max\{d_U(1, x_n), d_U(x_n, x_{2n})\} + c' \leq l_{\mathcal{P}} n + 3\eta n + 2c + c'.$$

Since this is true for every $\delta > 0$, we have $\sup_{U \in \mathcal{P}} d_U(1, x_{2n}) \leq l_{\mathcal{P}} n + 3\eta n + 2c + c'$, hence finally, with probability at least $1 - \epsilon$,

$$\sup_{U \in \mathcal{P}} d_U(1, \omega_{2n}) \leq l_{\mathcal{P}} n + 4\eta n + 3c + c'.$$

Since ϵ and η are arbitrary, this proves that almost surely,

$$\limsup_{n \rightarrow \infty} \frac{1}{2n} \sup_{U \in \mathcal{P}} d_U(1, \omega_{2n}) \leq \frac{1}{2} l_{\mathcal{P}}.$$

Therefore, $l_{\mathcal{P}} \leq \frac{1}{2} l_{\mathcal{P}}$ and so $l_{\mathcal{P}} = 0$. \square

3.2. Upper-bound for the Hausdorff dimension. In this subsection, we first prove the following.

Proposition 3.6. *Under the assumptions of Theorem 3.1, we have*

$$\overline{Hdim}_{\bar{\delta}_\lambda}(\nu_{\mathcal{B}}) \leq \frac{-1}{\log \lambda} \frac{h}{l}$$

and

$$\overline{Hdim}_{\delta_\lambda}(\nu_{\mathcal{F}}) \leq \frac{-1}{\log \lambda} \frac{h}{l}.$$

The proof is inspired from the proof of Le Prince for hyperbolic groups [46]. We first show that we only need to deal with the harmonic measure on the Floyd boundary. Let ϕ be the map from the Floyd boundary to the Bowditch boundary given by Theorem 2.4. This map is surjective, equivariant and continuous. Moreover, the preimage of a conical limit point consists of a singleton.

Lemma 3.7. *With the same notations, we have $\phi_*\nu_{\mathcal{F}} = \nu_{\mathcal{B}}$.*

Proof. Recall that a measure κ on a set X endowed with an action of G is called μ -stationary if it satisfies that for every measurable set $A \subset X$,

$$\kappa(A) = \sum_{g \in G} \kappa(g^{-1}A)\mu(g).$$

Combining [35, Section 6, Theorem] with [33, Theorem 2.4], we get that the harmonic measure $\nu_{\mathcal{F}}$ on the Floyd boundary is μ -stationary. Since the map ϕ is equivariant, $\phi_*\nu_{\mathcal{F}}$ also is μ -stationary. Now, $\nu_{\mathcal{B}}$ is the only μ -stationary measure on the Bowditch boundary, so that $\phi_*\nu_{\mathcal{F}} = \nu_{\mathcal{B}}$. This can be seen by verifying that the Bowditch compactification satisfies the conditions of [33, Theorem 2.4], which is done for some examples of relatively hyperbolic groups in [33, Section 9]. This can also be done directly by using results of Maher and Tiozzo [39]. First, we prove by contradiction that any μ -stationary measure on the Bowditch boundary has no atom. The following fact is well-known and we include a proof for completeness.

Claim 3.8. *Let G be a group acting on a measured space (X, κ) such that κ is μ -stationary and G does not fix any finite set on X . Then, κ has not atom.*

Proof of the claim. If this were the case, then we would choose an atom $x \in X$ of maximal measure. Since the support $\text{supp}(\mu)$ generates the group G as a semi-group, we would necessarily have $\kappa(g^{-1}x) = \kappa(x)$ for every g , for κ is μ -stationary, so

$$\kappa(x) = \sum_{g \in G} \kappa(g^{-1}x)\mu(g) \leq \sum_{g \in G} \kappa(x)\mu(g).$$

Hence, the orbit of x would be finite, which is impossible since the group G does not fix any finite set on X . \square

We can apply this claim to our situation because G is non-elementary and so it does not fix any finite set on the Bowditch boundary. Thus, any μ -stationary measure is uniquely defined by its restriction to the set of conical limit points and now [39, Theorem 1.1] shows that there is a unique μ -stationary probability measure on this set, which concludes the proof. \square

We deduce the following lemma.

Lemma 3.9. *With the above notations, we have*

$$\overline{Hdim}_{\bar{\delta}_\lambda}(\nu_{\mathcal{B}}) \leq \overline{Hdim}_{\delta_\lambda}(\nu_{\mathcal{F}})$$

and

$$Hdim_{\bar{\delta}_\lambda}(\nu_{\mathcal{B}}) \leq Hdim_{\delta_\lambda}(\nu_{\mathcal{F}}).$$

Proof. By (6), $\bar{\delta}_\lambda \leq \delta_\lambda$ on the set of conical limit points. Hence, for every point ξ such that $\phi(\xi)$ is a conical limit point,

$$\nu_{\mathcal{B}}(\phi(B(\xi, r))) \leq \nu_{\mathcal{B}}(B(\phi(\xi), r)).$$

Since $\nu_{\mathcal{B}} = \phi_*\nu_{\mathcal{F}}$, $\nu_{\mathcal{B}}(\phi(B(\xi, r))) = \nu_{\mathcal{F}}(\phi^{-1} \circ \phi(B(\xi, r)))$. We thus get

$$\nu_{\mathcal{F}}(B(\xi, r)) \leq \nu_{\mathcal{F}}(\phi^{-1} \circ \phi(B(\xi, r))) \leq \nu_{\mathcal{B}}(B(\phi(\xi), r)).$$

This is true for every point ξ such that $\phi(\xi)$ is a conical limit point. Since the map ϕ is surjective and $\nu_{\mathcal{B}}$ gives full measure to the set of conical limit points, this concludes the proof. \square

We thus only need to show that

$$\overline{Hdim}_{\delta_\lambda}(\nu_{\mathcal{F}}) \leq \frac{-1}{\log \lambda} \frac{h}{l}$$

to prove Proposition 3.6.

We denote by ω_n the random walk driven by μ at time n . Let $\Omega = G^{\mathbb{N}_{\geq 0}}$ be the trajectory space of the random walk. We also denote by $\mathbf{x} = (e, \omega_1, \dots, \omega_n, \dots) \in \Omega$ a sample path, i.e. a trajectory of the random walk starting from the identity 1. Recall that the random walk ω_n almost surely converges to a point ω_∞ in the Floyd boundary.

Lemma 3.10. *For \mathbb{P} -almost every $\mathbf{x} = (\omega_n)$, we have that*

$$\frac{1}{n} d(\omega_n, \omega_{n+1}) \xrightarrow{n \rightarrow +\infty} 0.$$

Proof. Note that $d(\omega_n, \omega_{n+1}) = d(1, g_{n+1})$ where g_n are the increments of the random walk. In particular, the random variables $d(\omega_n, \omega_{n+1})$ are independent, identically distributed and are integrable, since μ has finite first moment. It follows from the law of large numbers that

$$\frac{1}{n} \sum_{k=0}^n d(\omega_k, \omega_{k+1}) \xrightarrow{n \rightarrow \infty} L = \mathbb{E}(d(1, \omega_1)) < +\infty.$$

In particular,

$$\frac{1}{n} d(\omega_n, \omega_{n+1}) = \frac{1}{n} \sum_{k=0}^n d(\omega_k, \omega_{k+1}) - \frac{1}{n} \sum_{k=0}^{n-1} d(\omega_k, \omega_{k+1}) \xrightarrow{n \rightarrow \infty} L - L = 0.$$

This concludes the proof. \square

As a corollary, we have the following.

Lemma 3.11. *For \mathbb{P} -almost every $\mathbf{x} = (\omega_n)$, we have that for every geodesic γ_n from ω_n to ω_{n+1} ,*

$$\frac{1}{n} d(1, \gamma_n) \xrightarrow{n \rightarrow \infty} l.$$

Proof. Let γ_n be such a geodesic. Then, we have $d(1, \gamma_n) \leq d(1, \omega_n) + d(\omega_n, \omega_{n+1})$ and similarly, we have $d(1, \omega_n) \leq d(1, \gamma_n) + d(\omega_n, \omega_{n+1})$. Since $\frac{1}{n} d(1, \omega_n)$ almost surely converges to l by (2) and $\frac{1}{n} d(\omega_n, \omega_{n+1})$ almost surely converges to 0 by Lemma 3.10, we have that $\frac{1}{n} d(1, \gamma_n)$ also almost surely converges to l . \square

For every $\epsilon, N > 0$, we let Ω_ϵ^N be the set of trajectories \mathbf{x} such that for every $n \geq N$, we have

- (a) $d(1, \gamma_n) \geq (l - \epsilon)n$
- (b) $-\log \mu^{*n}(\omega_n) \leq (h + \epsilon)n$, and
- (c) $d(\omega_n, \omega_{n+1}) \leq n$

Observe that for every $\delta > 0$, there exists $N_{\epsilon, \delta}$ such that

$$\mathbb{P}(\Omega_\epsilon^{N_{\epsilon, \delta}}) \geq 1 - \delta.$$

Indeed, this follows immediately from the almost sure convergence of the following limits

$$\frac{1}{n} d(1, \gamma_n) \xrightarrow{n \rightarrow \infty} l, \quad \frac{1}{n} d(\omega_n, \omega_{n+1}) \xrightarrow{n \rightarrow \infty} 0, \quad \frac{-1}{n} \log \mu^{*n}(\omega_n) \xrightarrow{n \rightarrow \infty} h$$

where the last one is called the Shannon-McMillan-Breiman Theorem and is given by (3).

To simplify, we set $\Omega_{\epsilon, \delta} = \Omega_\epsilon^{N_{\epsilon, \delta}}$. Also, for fixed \mathbf{x} , we let $C_{\mathbf{x}}^n$ be the set of trajectories \mathbf{x}' such that $\omega'_n = \omega_n$. Finally, for ξ in the Floyd boundary and $r > 0$, we set

$$D(\xi, r) = \{\mathbf{x} : \delta_\lambda(\omega_\infty, \xi) \leq r\}.$$

Lemma 3.12. *There exists a set $\Lambda_{\epsilon,\delta} \subset \Omega_{\epsilon,\delta}$ of measure at least $1 - 2\delta$ on which the quantity*

$$\frac{\mathbb{P}(C_{\mathbf{x}}^n \cap \Omega_{\epsilon,\delta})}{\mu^{*n}(\omega_n)}$$

admits a positive limit. In particular, for every $\mathbf{x} = (\omega_n) \in \Lambda_{\epsilon,\delta}$, we have

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \mathbb{P}(C_{\mathbf{x}}^n \cap \Omega_{\epsilon,\delta}) = \limsup_{n \rightarrow \infty} \frac{1}{n} \log \mu^{*n}(\omega_n).$$

The proof can be found within the proof of [32, Theorem 1.4.1]. We rewrite it for convenience.

Proof. First note that $\mathbb{P}(C_{\mathbf{x}}^n) = \mu^{*n}(\omega_n)$. So $\frac{\mathbb{P}(C_{\mathbf{x}}^n \cap \Omega_{\epsilon,\delta})}{\mu^{*n}(\omega_n)}$ is the conditional probability of $\Omega_{\epsilon,\delta}$ with respect to $C_{\mathbf{x}}^n$, that we denote by $\mathbb{P}(\Omega_{\epsilon,\delta} | C_{\mathbf{x}}^n)$. We introduce the σ -algebra $\mathcal{A}^{\geq n}$ of sample paths determined by the coordinates ω_k , $k \geq n$. The tail σ -algebra \mathcal{A}^{∞} is then the intersection of the non-increasing sequence of σ -algebras $\mathcal{A}^{\geq n}$. There is a projection from the path space to the tail boundary that we denote by **tail**. We refer to [31] for more details. We let $\mathbb{P}(A | \mathbf{tail}(\mathbf{x}))$ be the conditional probability $\mathbb{P}(A | \mathcal{A}^{\infty})$ evaluated at the sample path \mathbf{x} .

The Markov property and the conditional probabilities convergence theorem show that $\mathbb{P}(\Omega_{\epsilon,\delta} | C_{\mathbf{x}}^n)$ converges to $\mathbb{P}(\Omega_{\epsilon,\delta} | \mathbf{tail}(\mathbf{x}))$, see [32, (1.4.4)]. Now, let

$$\Lambda_{\epsilon,\delta} = \{\mathbf{x} \in \Omega_{\epsilon,\delta}, \mathbb{P}(\Omega_{\epsilon,\delta} | \mathbf{tail}(\mathbf{x})) > 0\}.$$

We have

$$\mathbb{E}(\mathbb{P}(\Omega_{\epsilon,\delta} | \mathbf{tail}(\mathbf{x}))) = \mathbb{P}(\Omega_{\epsilon,\delta}).$$

Moreover,

$$\mathbb{E}(\mathbb{P}(\Omega_{\epsilon,\delta} | \mathbf{tail}(\mathbf{x}))) = \mathbb{E}(\mathbb{P}(\Omega_{\epsilon,\delta} | \mathbf{tail}(\mathbf{x})) \cdot 1_{\Lambda_{\epsilon,\delta}}) + \mathbb{E}(\mathbb{P}(\Omega_{\epsilon,\delta} | \mathbf{tail}(\mathbf{x})) \cdot 1_{\Omega_{\epsilon,\delta}^c}).$$

Since $\mathbb{P}(\Omega_{\epsilon,\delta} | \mathbf{tail}(\mathbf{x})) \leq 1$ for \mathbb{P} -almost every \mathbf{x} , we get

$$\mathbb{P}(\Omega_{\epsilon,\delta}) = \mathbb{E}(\mathbb{P}(\Omega_{\epsilon,\delta} | \mathbf{tail}(\mathbf{x}))) \leq \mathbb{P}(\Lambda_{\epsilon,\delta}) + 1 - \mathbb{P}(\Omega_{\epsilon,\delta})$$

that we rewrite $\mathbb{P}(\Lambda_{\epsilon,\delta}) \geq 2\mathbb{P}(\Omega_{\epsilon,\delta}) - 1$. Therefore, $\mathbb{P}(\Lambda_{\epsilon,\delta}) \geq 1 - 2\delta$, which concludes the proof. \square

Lemma 3.13. *There exists $M \geq 0$ such that the following holds. For fixed ϵ and δ , for every $\mathbf{x} \in \Lambda_{\epsilon,\delta}$ and every $n \geq N_{\epsilon,\delta}$, we have*

$$C_{\mathbf{x}}^n \cap \Omega_{\epsilon,\delta} \subset D(\omega_{\infty}, Mn\lambda^{n(l-\epsilon)}).$$

Proof. Fix $\mathbf{x} \in \Lambda_{\epsilon,\delta}$ and let $\mathbf{x}' \in C_{\mathbf{x}}^n \cap \Omega_{\epsilon,\delta}$. Then, $\omega_n = \omega'_n$. Moreover, for every $m \geq N_{\epsilon,\delta}$, letting γ'_m be a geodesic from ω'_m to ω'_{m+1} , we have $d(1, \gamma'_m) \geq (l - \epsilon)m$. Thus there exists a path of length $d(\omega'_m, \omega'_{m+1})$ which stays at distance at least $(l - \epsilon)m$ from 1. Since we also have $d(\omega_m, \omega_{m+1}) \leq m$, this proves that

$$(7) \quad \delta_{\lambda}(\omega'_m, \omega'_{m+1}) \leq d(\omega'_m, \omega'_{m+1})\lambda^{(l-\epsilon)m} \leq m\lambda^{(l-\epsilon)m}.$$

Consequently, for any $m > n$,

$$\delta_{\lambda}(\omega'_n, \omega'_m) \leq \sum_{k=n}^{m-1} k\lambda^{(l-\epsilon)k} \leq \sum_{k \geq 0} (n+k)\lambda^{(l-\epsilon)(n+k)}.$$

Note that

$$\sum_{k \geq 0} (n+k)\lambda^{(l-\epsilon)(n+k)} = n\lambda^{(l-\epsilon)n} \sum_{k \geq 0} \lambda^{(l-\epsilon)k} + \lambda^{(l-\epsilon)n} \sum_{k \geq 0} k\lambda^{(l-\epsilon)k}.$$

Since $\lambda < 1$, both sums $\sum_{k \geq 0} \lambda^{(l-\epsilon)k}$ and $\sum_{k \geq 0} k\lambda^{(l-\epsilon)k}$ are finite. Hence, there exists M_0 such that

$$\sum_{k \geq 0} (n+k)\lambda^{(l-\epsilon)(n+k)} \leq M_0 n \lambda^{(l-\epsilon)n},$$

so that $\delta_{\lambda}(\omega_n, \omega_m) \leq M_0 n \lambda^{(l-\epsilon)n}$. This holds for every $m > n$, so we have

$$\delta_{\lambda}(\omega'_n, \omega'_{\infty}) \leq M_0 n \lambda^{(l-\epsilon)n}.$$

Clearly, we also have $\mathbf{x}' \in C_{\mathbf{x}}^n \cap \Omega_{\epsilon,\delta}$ and so $\delta_{\lambda}(\omega_n, \omega_{\infty}) \leq M_0 n \lambda^{(l-\epsilon)n}$. Finally, since by definition of $C_{\mathbf{x}}^n$, $\omega_n = \omega'_n$, we get

$$\delta_{\lambda}(\omega_{\infty}, \omega'_{\infty}) \leq 2M_0 n \lambda^{(l-\epsilon)n}.$$

This proves the lemma, setting $M = 2M_0$. \square

We can now finish the proof of Proposition 3.6.

Proof. Our goal is to prove that for $\nu_{\mathcal{F}}$ -almost every ξ ,

$$(8) \quad \limsup_{r \rightarrow 0} \frac{\log \nu_{\mathcal{F}}(B(\xi, r))}{\log r} \leq \frac{-1}{\log \lambda} \frac{h}{l}.$$

It is thus enough to prove that for \mathbb{P} -almost every trajectory $\mathbf{x} = (\omega_n)$, we have

$$\limsup_{r \rightarrow 0} \frac{\log \nu_{\mathcal{F}}(B(\omega_{\infty}, r))}{\log r} \leq \frac{-1}{\log \lambda} \frac{h}{l},$$

since $\nu_{\mathcal{F}}$ is the law of the random variable ω_{∞} . This shall follow from the statement that for every ϵ and δ , for every $\mathbf{x} \in \Lambda_{\epsilon, \delta}$, we have

$$\limsup_{r \rightarrow 0} \frac{\log \nu_{\mathcal{F}}(B(\omega_{\infty}, r))}{\log r} \leq \frac{-1}{\log \lambda} \frac{h + \epsilon}{l - \epsilon}.$$

Indeed, $\mathbb{P}(\Lambda_{\epsilon, \delta}) \geq 1 - 2\delta$ and ϵ and δ can be chosen arbitrarily small.

Let $\mathbf{x} \in \Lambda_{\epsilon, \delta}$. We need to estimate $\limsup_{r \rightarrow 0} \frac{\log \nu_{\mathcal{F}}(B(\omega_{\infty}, r))}{\log r}$. Since the function $x \in \mathbb{R} \mapsto Mx\lambda^{(l-\epsilon)x}$ is eventually decreasing, we can replace r with $Mn\lambda^{(l-\epsilon)n}$ and make n go to infinity. Thus, it suffices to prove that

$$\limsup_{n \rightarrow \infty} \frac{\log \mathbb{P}(D(\omega_{\infty}, Mn\lambda^{(l-\epsilon)n}))}{\log \lambda(l - \epsilon)n} \leq \frac{-1}{\log \lambda} \frac{h + \epsilon}{l - \epsilon}.$$

In other words, we will prove that for every $\mathbf{x} \in \Lambda_{\epsilon, \delta}$,

$$\limsup_{n \rightarrow \infty} \frac{\log \mathbb{P}(D(\omega_{\infty}, Mn\lambda^{(l-\epsilon)n}))}{n} \geq -(h + \epsilon).$$

Using Lemma 3.13, we see that

$$\limsup_{n \rightarrow \infty} \frac{\log \mathbb{P}(D(\omega_{\infty}, Mn\lambda^{(l-\epsilon)n}))}{n} \geq \limsup_{n \rightarrow \infty} \frac{\log \mathbb{P}(C_{\mathbf{x}}^n \cap \Omega_{\epsilon, \delta})}{n}$$

and using Lemma 3.12, we see that

$$\limsup_{n \rightarrow \infty} \frac{\log \mathbb{P}(C_{\mathbf{x}}^n \cap \Omega_{\epsilon, \delta})}{n} = \limsup_{n \rightarrow \infty} \frac{\log \mu^{*n}(\omega_n)}{n}.$$

Finally, since $\mathbf{x} \in \Lambda_{\epsilon, \delta} \subset \Omega_{\epsilon, \delta}$, for large enough n , $\frac{\log \mu^{*n}(x_n)}{n} \geq -(h + \epsilon)$, so finally

$$\limsup_{n \rightarrow \infty} \frac{\log \mathbb{P}(D(\omega_{\infty}, Mn\lambda^{(l-\epsilon)n}))}{n} \geq -(h + \epsilon),$$

which concludes the proof. \square

3.3. Lower-bound for the Hausdorff dimension. The goal of this subsection is the following.

Proposition 3.14. *Under the assumptions of Theorem 3.1, we have*

$$Hdim_{\delta_{\lambda}}(\nu_{\mathcal{B}}) \geq \frac{-1}{\log \lambda} \frac{h}{l}$$

and

$$Hdim_{\delta_{\lambda}}(\nu_{\mathcal{F}}) \geq \frac{-1}{\log \lambda} \frac{h}{l}.$$

Proof. We only prove the result for the measure $\nu_{\mathcal{B}}$ on the Bowditch boundary. According to Lemma 3.9, the result for the harmonic measure on the Floyd boundary will then follow.

Proposition 3.2 shows there almost surely exists a transition point on a geodesic between 1 and ω_{∞} such that $\frac{1}{n}d(\omega_n, x_n)$ converges to 0. Also, $\frac{-1}{n} \log \mu^{*n}(\omega_n)$ almost surely converges to h by (3) and $\frac{1}{n}d(1, \omega_n)$ almost surely converges to l by (2). For every $\epsilon > 0$ and N , we let Ω_{ϵ}^N be the set of trajectories \mathbf{x} such that for every $n \geq N$,

- (a) $d(\omega_n, x_n) \leq \epsilon n$
- (b) $(l - \epsilon)n \leq d(1, x_n) \leq (l + \epsilon)n$
- (c) $-\log \mu^{*n}(\omega_n) \geq (h - \epsilon)n$

Then, for every ϵ , $\mathbb{P}(\cup_N \Omega_\epsilon^N) = 1$. Hence, there exists N_ϵ such that $\mathbb{P}(\Omega_\epsilon^{N_\epsilon}) \geq 1 - \epsilon$. We set $\Omega_\epsilon = \Omega_\epsilon^{N_\epsilon}$.

We will both need to deal with the set Ω_ϵ and the sets Ω_ϵ^N in the following. We fix N and we fix a trajectory $\mathbf{x} \in \Omega_\epsilon^N$ and so we also fix the corresponding sequence of transition points x_n on the geodesic between 1 and ω_∞ . Let $\Pi(g, R)$ be the big shadow at g , that is, the set of boundary points ξ in the Bowditch boundary such that there exists a geodesic ray $[1, \xi]$ intersecting $B(g, R)$.

First of all, observe that for all n large enough (i.e. bigger than N and N_ϵ),

$$\mathbb{P}(\mathbf{x}' \in \Omega_\epsilon \cap \{\omega'_\infty \in \Pi(x_n, R)\}) \leq \mathbb{P}(\mathbf{x}' \in \Omega_\epsilon : \omega'_n \in B(x_n, 2R + 3n\epsilon)).$$

Indeed, if the limiting point $\xi' = \omega'_\infty$ lies in the shadow $\Pi(x_n, R)$, then there is a point g_n on a geodesic from 1 to ξ' entering the ball $B(x_n, R)$, so $d(1, g_n)$ is between $(l - \epsilon)n - R$ and $(l + \epsilon)n + R$. For every $\mathbf{x}' \in \Omega_\epsilon^N$ with transition points x'_n on $[1, \omega'_\infty]$, we also have that $d(1, x'_n)$ is between $(l - \epsilon)n$ and $(l + \epsilon)n$, so we deduce that $d(x'_n, g_n) \leq R + 2\epsilon n$. By (a), $d(\omega'_n, x'_n) \leq \epsilon n$, so we must have $d(\omega'_n, x_n) \leq d(\omega'_n, g_n) + d(g_n, x_n) \leq 2R + 3\epsilon n$ as desired.

By the defining property (c) of Ω_ϵ , we have $\mu^{*n}(\omega'_n) \leq \exp(-n(h - \epsilon))$ for any $n > N_\epsilon$, so

$$\begin{aligned} & \mathbb{P}(\mathbf{x}' \in \Omega_\epsilon, \omega'_n \in B(x_n, 2R + 3n\epsilon)) \\ & \leq \mathbb{P}(\mu^{*n}(\omega'_n) \leq \exp(-n(h - \epsilon)), \omega'_n \in B(x_n, 2R + 3n\epsilon)) \end{aligned}$$

and since μ^{*n} is the law of ω'_n , the right-hand side of this inequality can be written as the following sum

$$\sum_{\substack{u \in B(x_n, 2R + 3n\epsilon), \\ \mu^{*n}(u) \leq \exp(-n(h - \epsilon))}} \mu^{*n}(u).$$

We thus get

$$\mathbb{P}(\mathbf{x}' \in \Omega_\epsilon, \omega'_n \in B(x_n, 2R + 3n\epsilon)) \leq \#B(1, 2R + 3n\epsilon) \exp(-n(h - \epsilon))$$

and since balls grow at most exponentially, there exists v such that

$$\mathbb{P}(\mathbf{x}' \in \Omega_\epsilon, \omega'_n \in B(x_n, 2R + 3n\epsilon)) \leq C \exp(3n\epsilon v) \exp(-n(h - \epsilon)).$$

Hence, for any N and any fixed \mathbf{x} in Ω_ϵ^N , we have

$$(9) \quad \liminf_{n \rightarrow \infty} \frac{\log \mathbb{P}(\Omega_\epsilon \cap \Pi(x_n, R))}{-n} \geq h - \epsilon - 3v\epsilon$$

Following [53], we use conditional expectations with respect to the σ -algebra \mathcal{S} , which is the smallest σ -algebra such that the map $\mathbf{bnd} : \mathbf{x} \mapsto \omega_\infty$ is measurable. Beware that the σ -algebra \mathcal{S} and the σ -algebra \mathcal{A}^∞ that we used in the proof of Proposition 3.6 can be different. We refer to the discussion in the proof of [32, Theorem 1.4.1] for more details. We let $\mathbb{P}(A|\mathbf{bnd}(\mathbf{x}))$ be the conditional probability $\mathbb{P}(A|\mathcal{S})$ evaluated at the sample path \mathbf{x} . This is denoted by $\mathbb{P}_{\mathbf{bnd}(\mathbf{x})}(A)$ in [53]. Since the harmonic measure $\nu_{\mathcal{B}}$ is the pushforward of the measure \mathbb{P} by the map \mathbf{bnd} , one can define a family of conditional probabilities $\xi \in \partial_{\mathcal{B}}G \mapsto \mathbb{P}(A|\xi)$ such that for every measurable set $A \in \partial_{\mathcal{B}}G$ and every measurable set B in the path-space of the random walk,

$$\mathbb{P}(B|\pi^{-1}(A)) = \frac{1}{\nu(A)} \int_A \mathbb{P}(B|\xi) d\nu_{\mathcal{B}}(\xi).$$

We then define the set

$$F_\epsilon = \{\xi \in \partial_{\mathcal{B}}G, \mathbb{P}(\Omega_\epsilon|\xi) \geq \epsilon\}.$$

Desintegrating along $\nu_{\mathcal{B}}$, we have

$$\begin{aligned} 1 - \epsilon & \leq \mathbb{P}(\Omega_\epsilon) = \int_{\partial_{\mathcal{B}}G} \mathbb{P}(\Omega_\epsilon|\xi) d\nu_{\mathcal{B}}(\xi) \\ & = \int_{F_\epsilon} \mathbb{P}(\Omega_\epsilon|\xi) d\nu_{\mathcal{B}}(\xi) + \int_{F_\epsilon^c} \mathbb{P}(\Omega_\epsilon|\xi) d\nu_{\mathcal{B}}(\xi) \\ & \leq \nu_{\mathcal{B}}(F_\epsilon) + \epsilon. \end{aligned}$$

Therefore, $\nu_{\mathcal{B}}(F_\epsilon) \geq 1 - 2\epsilon$. We now evaluate $\nu_{\mathcal{B}}(F_\epsilon \cap \Pi(x_n, R))$:

$$\begin{aligned} \nu_{\mathcal{B}}(F_\epsilon \cap \Pi(x_n, R)) & = \mathbb{P}(\omega_\infty \in F_\epsilon \cap \Pi(x_n, R)) \\ & \leq \mathbb{P}(\Omega_\epsilon \cap \Pi(x_n, R)) + \mathbb{P}(\Omega_\epsilon^c \cap \{\omega_\infty \in F_\epsilon \cap \Pi(x_n, R)\}) \end{aligned}$$

By definition, for $\xi \in F_\epsilon$, we have $\mathbb{P}(\Omega_\epsilon^c | \xi) \leq 1 - \epsilon$, so

$$\begin{aligned} \mathbb{P}(\Omega_\epsilon^c \cap \{\omega_\infty \in F_\epsilon \cap \Pi(x_n, R)\}) &= \int_{F_\epsilon \cap \Pi(x_n, R)} \mathbb{P}(\Omega_\epsilon^c | \xi) d\nu_{\mathcal{B}}(\xi) \\ &\leq (1 - \epsilon) \nu_{\mathcal{B}}(F_\epsilon \cap \Pi(x_n, R)). \end{aligned}$$

Consequently,

$$\epsilon \cdot \nu_{\mathcal{B}}(F_\epsilon \cap \Pi(x_n, R)) \leq \mathbb{P}(\Omega_\epsilon \cap \Pi(x_n, R)).$$

Thus, (9) yields for every N and every $\mathbf{x} \in \Omega_\epsilon^N$

$$\liminf_{n \rightarrow \infty} \frac{\log \nu_{\mathcal{B}}(F_\epsilon \cap \Pi(x_n, R))}{-n} \geq h - \epsilon - 3v\epsilon.$$

According to Lemma 2.6, there exists some constant $C > 0$ such that

$$B(\omega_\infty, Cr_n) \subset \Pi(x_n, R)$$

where $r_n = \lambda^{d(1, x_n)}$ satisfies $\lambda^{(l+\epsilon)n} \leq r_n \leq \lambda^{n(l-\epsilon)}$. Therefore,

$$\liminf_{n \rightarrow \infty} \frac{\log \nu_{\mathcal{B}}(F_\epsilon \cap B(\omega_\infty, Cr_n))}{\log Cr_n} \geq \frac{h - \epsilon - 3v\epsilon}{-\log \lambda(l + \epsilon)}.$$

Thus, for any N and any fixed \mathbf{x} in Ω_ϵ^N , we have

$$\liminf_{r \rightarrow 0} \frac{\log \nu_{\mathcal{B}}(F_\epsilon \cap B(\omega_\infty, r))}{\log r} \geq \frac{h - \epsilon - 3v\epsilon}{-\log \lambda(l + \epsilon)}.$$

This is true for all N and $\mathbb{P}(\cup_N \Omega_\epsilon^N) = 1$. Hence, for $\nu_{\mathcal{B}}$ -almost every ξ ,

$$(10) \quad \liminf_{r \rightarrow 0} \frac{\log \nu_{\mathcal{B}}(F_\epsilon \cap B(\xi, r))}{\log r} \geq \frac{h - \epsilon - 3v\epsilon}{-\log \lambda(l + \epsilon)}.$$

We can now conclude the proof, exactly like Theorem 1.1 is deduced from Theorem 3.3 in [53]. Let us give the details for completeness.

Consider the restriction $\nu_{\mathcal{B}, \epsilon}$ of $\nu_{\mathcal{B}}$ to F_ϵ . Then, (10) yields

$$Hdim(\nu_{\mathcal{B}, \epsilon}) \geq \frac{h - \epsilon - 3v\epsilon}{-\log \lambda(l + \epsilon)}$$

and so

$$Hdim(\nu_{\mathcal{B}}) \geq \frac{h - \epsilon - 3v\epsilon}{-\log \lambda(l + \epsilon)}.$$

In particular, the set

$$G_\epsilon = \left\{ \xi \in \partial_{\mathcal{B}} G, \liminf_{r \rightarrow 0} \frac{\log \nu_{\mathcal{B}}(B(\xi, r))}{\log r} \geq \frac{h - \epsilon - 3v\epsilon}{-\log \lambda(l + \epsilon)} - \epsilon \right\}$$

has positive $\nu_{\mathcal{B}}$ -measure. We show that G_ϵ is G -invariant. First, recall that the measure $\nu_{\mathcal{B}}$ is μ -stationary. Since μ generates G as a semi-group, for any $g \in G$, there exists n such that $\mu^{*n}(g^{-1}) > 0$ and $\mu^{*n}(g) > 0$. Note that $\nu_{\mathcal{B}}$ also is μ^{*n} stationary, so that for any measurable set A ,

$$\sum_{g \in A} \nu_{\mathcal{B}}(gA) \mu^{*n}(g) = \nu_{\mathcal{B}}(A).$$

In particular, there exists $c_{g, \mu} > 0$ such that

$$c_{g, \mu}^{-1} \nu_{\mathcal{B}} \leq g^{-1} \nu_{\mathcal{B}} \leq c_{g, \mu} \nu_{\mathcal{B}}.$$

Together with Lemma 2.7, this implies that

$$\nu_{\mathcal{B}}(B(\xi, r)) \leq c_{g, \mu} \nu_{\mathcal{B}}(B(g\xi, c_g r))$$

for any $\xi \in G_\epsilon$. By taking the limit inf, the constant $c_{g, \mu}$ disappears and thus $g\xi \in G_\epsilon$. Hence, G_ϵ is indeed G -invariant.

The measure $\nu_{\mathcal{B}}$ is ergodic and $\nu_{\mathcal{B}}(G_\epsilon) > 0$. Consequently, the set G_ϵ has full measure. We thus get that for $\nu_{\mathcal{B}}$ -almost every ξ ,

$$\liminf_{r \rightarrow 0} \frac{\log \nu_{\mathcal{B}}(B(\xi, r))}{\log r} \geq \frac{h - \epsilon - 3v\epsilon}{-\log \lambda(l + \epsilon)} - \epsilon.$$

Since ϵ is arbitrary, this shows that for $\nu_{\mathcal{B}}$ -almost every ξ ,

$$(11) \quad \liminf_{r \rightarrow 0} \frac{\log \nu_{\mathcal{B}}(B(\xi, r))}{\log r} \geq \frac{-1}{\log \lambda} \frac{h}{l}$$

and thus concludes the proof. \square

We now conclude the proof of Theorem 3.1.

Proof. By Propositions 3.6 and 3.14, it remains to show the exact dimensionality of $\nu_{\mathcal{B}}$ and $\nu_{\mathcal{F}}$. According to (8) and Lemma 3.7, for $\nu_{\mathcal{F}}$ -almost every ξ ,

$$\limsup_{r \rightarrow 0} \frac{\log \nu_{\mathcal{B}}(B(\phi(\xi), r))}{\log r} \leq \limsup_{r \rightarrow 0} \frac{\log \nu_{\mathcal{F}}(B(\xi, r))}{\log r} \leq \frac{-1}{\log \lambda} \frac{h}{l}.$$

Similarly, (11) shows that for $\nu_{\mathcal{B}}$ -almost every ξ ,

$$\liminf_{r \rightarrow 0} \frac{\log \nu_{\mathcal{F}}(B(\xi, r))}{\log r} \geq \liminf_{r \rightarrow 0} \frac{\log \nu_{\mathcal{B}}(B(\phi(\xi), r))}{\log r} \geq \frac{-1}{\log \lambda} \frac{h}{l}.$$

This shows that both $\nu_{\mathcal{B}}$ and $\nu_{\mathcal{F}}$ are exact dimensional. \square

4. GROUPS WITH INFINITELY MANY ENDS

In this section, we compute the Hausdorff dimension of the harmonic measure on the end boundary equipped with a visual metric

4.1. The end compactification. Let Γ be an infinite, connected, locally finite graph. For every edge e in Γ , we denote by e^0 its endpoints. More generally, for every set of edges E , we denote by E^0 the set of vertices that are endpoints of an edge in E . We will also write Γ^0 for the set of all vertices of Γ .

Let E be a finite set of edges. Denote by $\Gamma \setminus E$ the spanning graph of the vertex set of $\Gamma^0 \setminus E^0$: removing all edges sharing one vertex with one edge in E . Let $\mathcal{C}(E)$ be the set of infinite components of $\Gamma \setminus E$. By definition, there exists at least one edge e for every component $C \in \mathcal{C}(E)$ such that $e^0 \cap C \neq \emptyset$ and $e^0 \cap E^0 \neq \emptyset$. Note that two components C_1, C_2 in $\mathcal{C}(E)$ are distinct if and only if every path between any two points $x \in C_1, y \in C_2$ intersects E^0 . Finally, two points x, y are called *separated* by E if x and y lie in distinct components of $\Gamma \setminus E$.

We can define the *end compactification* of Γ as follows. Consider the directed system $\mathcal{F}(\Gamma)$ of all finite set of edges in Γ with $E < F$ if $E \subset F$. There is a natural map from $\mathcal{C}(F)$ to $\mathcal{C}(E)$ induced by inclusions of infinite components. The *end boundary* $\partial_{\mathcal{E}}\Gamma$ is the inverse limit of the directed system $\mathcal{C}(E)$ over all finite set of edges E in Γ . By definition, a point $\xi \in \partial_{\mathcal{E}}\Gamma$ is a collection of infinite components $C_E(\xi)$ of $\Gamma \setminus E$ for every $E \in \mathcal{F}(\Gamma)$, such that $C_E(\xi) \cap C_{E'}(\xi)$ is infinite for any two E, E' . We call ξ an *end* of Γ . For every E , the component $C_E(\xi)$ of $\mathcal{C}(E)$ is uniquely determined by ξ and by abuse of language, we say that $C_E(\xi)$ contains ξ , that we denote by $\xi \in C_E(\xi)$. We can then extend the definition of separated pair of points to ends. Two ends $\xi \neq \zeta$ are *separated* by E if $C_E(\xi) \neq C_E(\zeta)$, or equivalently if $\xi \notin C_E(\zeta)$. Note that any two distinct ends are necessarily separated by some $E \in \mathcal{F}(\Gamma)$.

The end boundary $\partial_{\mathcal{E}}\Gamma$ defines a compactification of Γ in the following way. One can extend the discrete topology on the set of vertices Γ^0 to a metrizable topology on $\bar{\Gamma}^{\mathcal{E}} = \Gamma \cup \partial_{\mathcal{E}}\Gamma$ which makes it a compact space and such that Γ is dense in $\bar{\Gamma}^{\mathcal{E}}$. Moreover, a sequence of points $x_n \in \Gamma$ converges to an end $\xi \in \partial_{\mathcal{E}}\Gamma$ if and only if for every $E \in \mathcal{F}(\Gamma)$, we have that x_n lies in $C_E(\xi)$ for all but finitely many n .

The topological closure $\bar{C}_E(\xi)$ of a component $C_E(\xi)$ in the compactification $\Gamma \cup \partial_{\mathcal{E}}\Gamma$ is the union of $C_E(\xi)$ with all $\zeta \in \partial_{\mathcal{E}}\Gamma$ for which $C_E(\xi) = C_E(\zeta)$. Hence, a component C contains ξ if $\xi \in \bar{C}$. Let $C_{E_i}(\xi)$ be a sequence of strictly shrinking components, that is, $C_{E_{i+1}}(\xi) \subset C_{E_i}(\xi)$. Then their closures $\bar{C}_{E_i}(\xi)$ yield a neighborhood basis of ξ .

Let ξ be an end. By definition, there exists a sequence of finite subsets E_n such that $C_{E_{n+1}}(\xi) \subset C_{E_n}(\xi)$. Following Woess [57], we say that ξ is a *thin end* if the sets E_n can be chosen so that $\sup_n \text{diam}(E_n)$ is finite. We say that it is *M-thin* if $\sup_n \text{diam}(E_n) \leq M$.

Then end compactification of a finitely generated group G is the end compactification of its Cayley graph with respect to a finite generating set. The quasi-isometry extends to a homeomorphism between end boundary, so the topology of the end boundary does not depend on the choice of this generating set and so

it is well defined. We denote by $\partial_{\mathcal{E}}G$ the end boundary of G . This compactification was first introduced by Freudenthal and is also called the Freudenthal compactification. We refer to [19] for more details.

An infinitely-ended group is called *accessible* if it admits a splitting over finite edge groups as a finite graph of one ended or finite vertex groups. Finitely presented group are accessible [14]. The accessibility is a quasi-isometric invariant by the following graphical characterization [54]. An infinitely-ended group is accessible if and only if there exists $k > 0$ such that any two distinct ends are separated by k edges in a Cayley graph.

Taking limits of geodesics and using Arzelà-Ascoli Theorem, we see that the end boundary is a visual boundary: any end is connected to any point x in the group by an infinite geodesic and any two distinct ends are connected by a bi-infinite geodesic. Finally, by results of Stallings [52], G acts on $\partial_{\mathcal{E}}G$ as a convergence group, see also [4, Lemma 5.1].

4.2. Visual metrics. Let G be a finitely generated group with infinitely many ends. In [6], Candelero, Gilch and Müller defined a *visual metric* on the set of ends of a free product. We extend their definition to our situation. Fix a basepoint $o \in G$ and fix $0 < \lambda < 1$. We define a distance $\rho_{o,\lambda}$ on $\partial_{\mathcal{E}}G$ as follows. Consider the sequence of finite edge sets B_n , where B_n is the edge set of the subgraph spanned by the vertices in the closed ball of radius n around o . The inverse limit of the directed system $\{\mathcal{C}(B_n)\}$ is homeomorphic to $\partial_{\mathcal{E}}\Gamma$.

Let ξ, ζ be two distinct ends. Let n be the minimal integer such that ξ and ζ belongs to different components in $\mathcal{C}(B_n)$, then define $\rho_{\lambda,o}(\xi, \zeta) = \lambda^n$. When the basepoint is the neutral element 1 of the group, we will simply write $\rho_{\lambda}(\xi, \zeta)$.

By definition, the visual metric is *ultrametric*: for any triple of points $x, y, z \in \partial_{\mathcal{E}}\Gamma$,

$$\rho_{\lambda}(x, y) \leq \max\{\rho_{\lambda}(x, z), \rho_{\lambda}(z, y)\}$$

The next lemma follows directly from the definition.

Lemma 4.1. *For every $x, y \in \partial_{\mathcal{E}}\Gamma$ in the end boundary and for any choice of basepoints $o, o' \in G$,*

$$\lambda^{d(o,o')} \leq \frac{\rho_{\lambda,o}(x, y)}{\rho_{\lambda,o'}(x, y)} \leq \lambda^{-d(o,o')}.$$

As a consequence, we get the following, which is proved exactly like Lemma 2.7.

Lemma 4.2. *For every $g \in G$, there exists a constant c_g such that for every end ξ and for every $r \geq 0$,*

$$B_{\rho_{\lambda}}(g\xi, c_g^{-1}r) \subset gB_{\rho_{\lambda}}(\xi, r) \subset B_{\rho_{\lambda}}(g\xi, c_g r).$$

It is well known that the Floyd boundary covers the end boundary, see for example [20, Proposition 11.1] and [36]. The following lemma allows us to compare the Floyd distance with the visual distance.

Lemma 4.3. *The identity of G extends to a 1-Lipschitz surjective equivariant map ϕ from the Floyd boundary to the end boundary with the same parameter $\lambda \in (0, 1)$:*

$$\rho_{\lambda}(\phi(\xi), \phi(\zeta)) \leq \delta_{\lambda}(\xi, \eta)$$

for any $\xi, \zeta \in \partial_{\mathcal{F}}G$.

Proof. Consider $x, y \in G$ such that $\rho_{\lambda}(x, y) = \lambda^n$, where n is the minimal integer such that x, y are contained in distinct components of $\mathcal{C}(B_n)$. Then any path from x to y has to intersect $B(1, n)$, so $\delta_{\lambda}(x, y) \geq \lambda^n$. Hence, for any $x, y \in G$,

$$(12) \quad \delta_{\lambda}(x, y) \geq \rho_{\lambda}(x, y).$$

Consider now $\xi \in \partial_{\mathcal{F}}G$ and let x_n be a sequence in G converging to ξ . Then, x_n is Cauchy for the Floyd distance and so (12) shows it is also Cauchy for the visual distance. This proves that x_n converges to a point $\phi(\xi) \in \partial_{\mathcal{E}}G$, which is uniquely determined by ξ . By construction, (12) extends to points in the boundary and ϕ is equivariant. \square

4.3. The end boundary of accessible groups. As explained in Introduction, any infinitely-ended group is relatively hyperbolic. If it is accessible, the action on its end boundary is geometrically finite. Precisely, the peripheral structure is given by the “terminal” splitting of the accessible group as a finite graph of groups over finite edge groups so that every vertex groups are either one-ended or finite. So for accessible groups, the end boundary is homeomorphic to the Bowditch boundary, whose construction is briefly recalled below.

Let T be the corresponding Bass-Serre tree of the above terminal splitting. Following Bowditch [3], we can put a compact metrizable topology on $T^0 \cup \partial_{\mathcal{E}} T$, for T is a fine hyperbolic graph. A similar construction is also given for any (non-)locally finite graph by Cartwright-Soardi-Woess [7].

As a perfect compact space, the Bowditch boundary or the end boundary is homeomorphic to the subspace of $T^0 \cup \partial_{\mathcal{E}} T$ minus the isolated points coming from the vertices with finite stabilizer. According to the definition of a geometrical finite action, every point in the Bowditch boundary is either conical or bounded parabolic. The set of conical points are exactly $\partial_{\mathcal{E}} T$ and bounded parabolic points are the vertices in T with infinite stabilizer, which are the subset of ends in $\partial_{\mathcal{E}} G$ corresponding to the left cosets of stabilizer in the Cayley graph of G . Finally, let us remark that these two types of limit points are precisely thin end and thick end introduced and studied in [54]. The proof of this fact can be found in [20, Proposition 7.8]

4.4. Hausdorff dimension of the harmonic measure. In this section, we consider a finitely generated group G with infinitely many ends and we consider a probability measure μ with finite first moment on G . We denote by ω_n the random walk driven by μ . It is a classical fact that ω_n almost surely converges to an end ω_{∞} , and that denoting by $\nu_{\mathcal{E}}$ the law of ω_{∞} , the end boundary $(\partial_{\mathcal{E}} G, \nu_{\mathcal{E}})$ is a model for the Poisson boundary. See for example [33, Theorem 8.4]. We call $\nu_{\mathcal{E}}$ the harmonic measure on $\partial_{\mathcal{E}} G$.

Proposition 4.4. *There exists $M > 0$ such that $\nu_{\mathcal{E}}$ gives full measure to the set of M -thin ends.*

Proof. It is proved in [57, Theorem 4.1] that the set of ends of a locally finite graph with infinitely many ends can be decomposed into the union $\Omega_0 \cup \Omega'$, where Ω_0 is a dense set in the set of ends consisting of M -thin ends. When the graph is the Cayley graph of a finitely generated group, the set Ω_0 can be constructed as follows. As explained in the introduction, the group G splits as an amalgamated product $A *_C B$ or an HNN extension $A *_C$. In the former case, G is hyperbolic relative to the conjugates of A and B and in the later case, it is hyperbolic relative to the conjugates of A . In both cases, every element of G can be written with elements of A , B and C with a normal form, see [57, (9.2), (9.4)] and Ω_0 can be described as the set of infinite words with respect to this normal form, see [57, (9.3), (9.5)]. Moreover, the set Ω' is constructed as the union of translates of the set of ends of A and B , see the remarks after [57, (9.3), (9.5)]. We can thus construct a continuous, surjective and equivariant map ψ from the set of ends $\partial_{\mathcal{E}} G$ to the Bowditch boundary of G with respect to the relatively hyperbolic structure described above. The map ψ is obtained by collapsing the translate $g\partial_{\mathcal{E}} A$ of the set of ends of A to the point $g\alpha$ in the Bowditch boundary, where α is the parabolic limit point fixed by A and similarly with B . It follows from this construction that Ω_0 is mapped to conical limit points. Note that the measure $\nu_{\mathcal{E}}$ is μ -stationary and since ψ is equivariant, the pushforward $\psi_* \nu_{\mathcal{E}}$ also is μ -stationary on the Bowditch boundary. As explained in the proof of Lemma 3.7, the harmonic measure $\nu_{\mathcal{B}}$ on the Bowditch boundary is the unique μ -stationary measure, hence $\psi_* \nu_{\mathcal{E}} = \nu_{\mathcal{B}}$. Moreover, $\nu_{\mathcal{B}}$ gives full measure to conical limit points. Consequently,

$$\nu_{\mathcal{E}}(\Omega') \leq \nu_{\mathcal{E}}(\psi^{-1}(\psi(\Omega'))) = \nu_{\mathcal{B}}(\psi(\Omega')) = 0.$$

This concludes the proof. \square

The following definition is inspired by the work of Derriennic [12] in free groups.

Definition 4.5. *Let $x, y \in G \cup \partial_{\mathcal{E}} G$ and let $M \geq 0$. We say that a set $U \subset G$ is a M -bottleneck between x and y if $\text{diam}(U) \leq M$ and any path from x to y has to pass through U .*

Similar sets are called transitional sets by Derriennic in [12]. However, to avoid confusion with the terminology “transition points” in relatively hyperbolic groups, we used the name bottleneck, which will also be more suited to our use later. The next lemma follows from our definitions.

Lemma 4.6. *Let ξ be a M -thin end for $M > 0$. Then there exists an infinite sequence of distinct sets E_n in G with $\sup_{n \geq 1} \{\text{diam}(E_n)\} \leq M$ such that any path from 1 to ξ has to pass successively through each E_n .*

Moreover, for any $x \neq \xi \in G \cup \partial_{\mathcal{E}} G$, there exists $n_0 > 0$ such that any path from x to ξ has to pass successively through each E_n , for $n \geq n_0$. In particular, for all but finitely many $n > 0$, the sets E_n are M -bottlenecks between x and ξ .

Our goal in the remainder of this section is to compute the Hausdorff dimension of the harmonic measure with respect to the visual distance. Precisely, we prove the following theorem.

Theorem 4.7. *Let $(\partial_{\mathcal{E}}G, \nu_{\mathcal{E}})$ be as above and let h, l be the entropy and the rate of escape of the μ -random walk on G . Then, for any $\lambda \in (0, 1)$,*

$$Hdim_{\rho_{\lambda}}(\nu_{\mathcal{E}}) = \frac{-1}{\log \lambda} \frac{h}{l}.$$

Moreover, $\nu_{\mathcal{E}}$ is exact dimensional.

We will follow the strategy that we used for the harmonic measures on the Bowditch and the Floyd boundaries. We first give an upper bound.

Proposition 4.8. *For $\nu_{\mathcal{E}}$ -almost every ξ in $\partial_{\mathcal{E}}G$,*

$$\limsup_{r \rightarrow 0} \frac{\log \nu_{\mathcal{E}}(B_{\rho_{\lambda}}(\xi, r))}{\log r} \leq \frac{-1}{\log \lambda} \frac{h}{l}.$$

Proof. The proof of Proposition 3.6 for the Floyd distance again applies here. The only place where the Floyd distance is used there is in the estimate (7), which states that whenever x, x' are joined by a geodesic which stays at distance at least m_1 from 1 and which satisfies that $d(x, x') \leq m_2$, we have

$$\delta_{\lambda}(x, x') \leq m_2 \lambda^{m_1}.$$

This is again true replacing the Floyd distance δ_{λ} by the visual distance ρ_{λ} , since we have the better estimate

$$\rho_{\lambda}(x, x') \leq \lambda^{m_1}.$$

Also note that for $\lambda \geq \lambda_0$, we can give a direct proof. We use the map ϕ given by Lemma 4.3. By [33, Theorem 8.3], $\nu_{\mathcal{E}}$ is the only μ -stationary measure on $\partial_{\mathcal{E}}G$. Hence, $\phi_*\nu_{\mathcal{F}} = \nu_{\mathcal{E}}$. The result thus follows from the same result for the measure $\nu_{\mathcal{F}}$, which is given by (8). \square

To obtain a lower bound, we will use the following result, which says that the random walk almost surely sublinearly tracks bottlenecks.

Proposition 4.9. *There exists $M \geq 0$ such that for \mathbb{P} -almost every $\mathbf{x} = (\omega_n) \in \Omega$, there exists a sequence of M -bottlenecks U_n between 1 and ω_{∞} satisfying that*

$$\frac{1}{n} d(\omega_n, U_n) \xrightarrow{n \rightarrow \infty} 0.$$

Proof. We fix $M \geq 0$ as in Proposition 4.4. For every $x, y \in G \cup \partial_{\mathcal{E}}G$, we denote by $\text{Bn}(x, y)$ the set of M -bottlenecks between x and y . We introduce the function f defined by $f(\omega) = d(1, \text{Bn}(\omega_{-\infty}, \omega_{\infty}))$, where $\omega_{-\infty}$ is the limit of the reflected random walk in $\partial_{\mathcal{E}}G$. Since the measure $\nu_{\mathcal{E}}$ is stationary, it is non-atomic. Indeed, we can apply the Claim 3.8, for G cannot fix any finite set on $\partial_{\mathcal{E}}G$ since it is non-amenable, see [57, Theorem 2.3]. Hence, $\omega_{-\infty}$ and ω_{∞} are almost surely distinct. According to Proposition 4.4, $\omega_{-\infty}$ and ω_{∞} are almost surely M -thin ends, so Lemma 4.6 shows that $\text{Bn}(\omega_{-\infty}, \omega_{\infty})$ is non-empty. The proof of the Claim 3.3 thus shows that f is measurable and is almost surely finite.

Note that $f(T^n \omega) = d(\omega_n, \text{Bn}(\omega_{-\infty}, \omega_{\infty}))$ and so $|f(T\omega) - f(\omega)| \leq d(1, \omega_1)$. Hence, [55, Lemma 7] shows that $\frac{1}{n} f(T^n \omega)$ almost surely converges to 0. Thus, there almost surely exists a sequence of M -bottlenecks U_n between $\omega_{-\infty}$ and ω_{∞} such that $\frac{1}{n} d(\omega_n, U_n)$ converges to 0.

To conclude, we just need to show that for large enough n , U_n is also a M -bottleneck between 1 and ω_{∞} . Since $d(1, \omega_n)$ almost surely converges to l , we can assume that for large enough n , $d(1, \omega_n) \geq (l - \epsilon)n$ and so $d(1, U_n)$ goes to infinity. Fix a path α from $\omega_{-\infty}$ to 1. Then for large enough n , say $n \geq n_0$, α does not intersect U_n . Consider now any path β from 1 to ω_{∞} . Concatenating α and β yields a path from $\omega_{-\infty}$ to ω_{∞} which thus crosses U_n . By construction, we necessarily have that β intersects U_n for $n \geq n_0$. \square

We can now prove the following result.

Proposition 4.10. *For $\nu_{\mathcal{E}}$ -almost every ξ in $\partial_{\mathcal{E}}G$,*

$$\liminf_{r \rightarrow 0} \frac{\log \nu_{\mathcal{E}}(B_{\rho_{\lambda}}(\xi, r))}{\log r} \geq \frac{-1}{\log \lambda} \frac{h}{l}.$$

Proof. We choose a sequence of points $x_n \in U_n$, where U_n is a sequence of M -bottlenecks between 1 and ω_∞ given by Proposition 4.9. For every $\epsilon > 0$ and N , we let Ω_ϵ^N be the set of trajectories $\mathbf{x} = (\omega_n) \in \Omega$ such that for every $n \geq N$,

- (a) $d(\omega_n, x_n) \leq \epsilon n$,
- (b) $(l - \epsilon)n \leq d(1, x_n) \leq (l + \epsilon)n$, and
- (c) $-\log \mu^{*n}(\omega_n) \geq (h - \epsilon)n$.

Almost surely, $\frac{1}{n}d(\omega_n, x_n)$ converges to 0, $\frac{1}{n}d(1, \omega_n)$ converges to l by (2), and $\frac{1}{n}\log \mu^{*n}(\omega_n)$ converges to h by (3). Hence, there exists N_ϵ such that $\mathbb{P}(\Omega_\epsilon^{N_\epsilon}) \geq 1 - \epsilon$. We set $\Omega_\epsilon = \Omega_\epsilon^{N_\epsilon}$.

We fix N and we fix a trajectory $\mathbf{x} \in \Omega_\epsilon^N$ and so we also fix the corresponding sequence of points x_n . For $g \in G$, we define the *partial shadow* $\mathcal{U}(g, M)$ to be the set of $\xi \in \partial_\mathcal{E}G$ such that g lies in a M -bottleneck between 1 and ξ . For $n \geq N$ and $n \geq N_\epsilon$, we have

$$\mathbb{P}(\mathbf{x}' \in \Omega_\epsilon \cap \{\omega'_\infty \in \mathcal{U}(x_n, M)\}) \leq \mathbb{P}(\mathbf{x}' \in \Omega_\epsilon, \omega'_n \in B(x_n, 4M + 3n\epsilon)).$$

Indeed, assume that x_n lies in a M -bottleneck V between 1 and ω'_∞ . Fix a geodesic from 1 to ω'_∞ . This geodesic enters V at a point g_n which satisfies $d(g_n, x_n) \leq M$. In particular, $d(1, g_n)$ is between $(l - \epsilon)n - M$ and $(l + \epsilon)n + M$. There is also a point on this geodesic that enters the bottleneck U'_n at a point g'_n , satisfying $d(g'_n, x'_n) \leq M$ and so we also have that $d(1, g'_n)$ is between $(l - \epsilon)n - M$ and $(l + \epsilon)n + M$. Thus, $d(g_n, g'_n) \leq 2M + 2\epsilon n$ and since $d(\omega'_n, x'_n) \leq \epsilon n$, we get $d(\omega'_n, x_n) \leq 4M + 3\epsilon n$ as required. On Ω_ϵ , we have $\mu^{*n}(\omega'_n) \leq \exp(-n(h - \epsilon))$ for any $n > N_\epsilon$ and μ^{*n} is the law of ω'_n , so

$$\mathbb{P}(\mathbf{x}' \in \Omega_\epsilon, \omega'_n \in B(x_n, 4M + 3n\epsilon)) \leq \#B(1, 4M + 3n\epsilon) \exp(-n(h - \epsilon))$$

and since balls grow at most exponentially, there exists v such that

$$\mathbb{P}(\mathbf{x}' \in \Omega_\epsilon, \omega'_n \in B(x_n, 4M + 3n\epsilon)) \leq \exp(4vM) \exp(3n\epsilon v) \exp(-n(h - \epsilon)).$$

Hence, for any N and any fixed \mathbf{x} in Ω_ϵ^N , we have

$$(13) \quad \liminf_{n \rightarrow \infty} \frac{\log \mathbb{P}(\Omega_\epsilon \cap \mathcal{U}(x_n, M))}{-n} \geq h - \epsilon - 3v\epsilon$$

We can conclude exactly like in the proof of Proposition 3.14, replacing Lemma 2.7 by Lemma 4.2 and replacing Lemma 2.6 by the following result, which asserts that partial shadows are sandwiched by balls. \square

As before, we define the *big shadow* $\Pi(g, M)$ as the set of $\xi \in \partial_\mathcal{E}G$ such that there is a geodesic between 1 and ξ which intersects the ball of radius M centered at g . By definition, $\mathcal{U}(g, M) \subset \Pi(g, M)$ for $M > 0$.

Lemma 4.11. *Let $\xi \in \partial_\mathcal{E}G$ and $r \geq 0$. Let g be any point on a geodesic between 1 and ξ . Then for any $M \geq 0$ there exist $C_1, C_2 > 0$ such that*

$$\Pi(g, M) \subset B_{\rho_\lambda}(\xi, C_1 r)$$

and if, in addition, g lies in a M -bottleneck between 1 and ξ , then

$$B_{\rho_\lambda}(\xi, C_2 r) \subset \mathcal{U}(g, M),$$

where $r = \lambda^{d(1, g)}$.

Proof. Let $\zeta \in \Pi(g, M)$. Denote by h a point on a geodesic α from 1 to ζ such that $d(g, h) \leq M$. Also denote by β the geodesic from 1 to ξ in the statement of the lemma. Consider the path γ connecting ζ to ξ obtained by following α from ζ to h , then connecting h to g by a geodesic and following β from g to ξ . By construction, $d(1, \alpha) \geq d(1, g) - M$. Hence, ξ and ζ must lie in the same component of $\mathcal{C}(B_n)$ where $n = d(1, g) - M$. Therefore, $\rho_\lambda(\xi, \zeta) \leq \lambda^{n+1} \leq C_1 r$ where $C_1 = \lambda^{-M-1}$ and so $\zeta \in B_{\rho_\lambda}(\xi, C_1 r)$.

To prove the second inclusion, set $C_2 = \lambda^{M+1}$ and consider $\zeta \in B_{\rho_\lambda}(\xi, C_2 r)$. Write $\rho_\lambda(\xi, \zeta) = \lambda^n$, where n is the minimal integer such that ξ and ζ lie in different components of $\mathcal{C}(B_n)$, so that $n \geq d(1, g) + M + 1$. Then, there exists a path α from ξ to ζ that enters $B(1, n)$ but not $B(1, n - 1)$. In particular, we have $d(g, \alpha) \geq d(1, \alpha) - d(1, g) > M$. Let β be any path from 1 to ζ and consider the path γ obtained by concatenating α and β , which needs to cross $B(g, M)$, since g is in a bottleneck between 1 and ξ . By construction of α , β necessarily intersects $B(g, M)$, so $\zeta \in \mathcal{U}(g, M)$. \square

Finally, note that Proposition 4.10 yields

$$Hdim_{\rho_\lambda}(\nu_\mathcal{E}) \geq \frac{-1}{\log \lambda} \frac{h}{l}$$

and Proposition 4.8 yields

$$Hdim_{\rho_\lambda}(\nu_\mathcal{E}) \leq \overline{Hdim}_{\rho_\lambda}(\nu_\mathcal{E}) \leq \frac{-1}{\log \lambda} \frac{h}{l}.$$

Hence, $Hdim_{\rho_\lambda}(\nu_\mathcal{E}) = \frac{-1}{\log \lambda} \frac{h}{l}$. Moreover, combining these two propositions implies that for $\nu_\mathcal{E}$ -almost every ξ ,

$$\frac{\log \nu_\mathcal{E}(B(\xi, r))}{\log r} \xrightarrow{r \rightarrow 0} Hdim_{\rho_\lambda}(\nu_\mathcal{E})$$

and so $\nu_\mathcal{E}$ is exact dimensional. This concludes the proof of Theorem 4.7. \square

5. DIMENSION OF THE END BOUNDARY

Recall that G is a finitely generated group with infinitely many ends, and (Γ, d) is the Cayley graph with respect to a finite generating set. In this section, we compute the Hausdorff dimension of the end boundary $\partial_\mathcal{E}G$ endowed with a visual metric. Define the volume growth of a subgroup $H < G$ as

$$v_H = \limsup_{n \rightarrow \infty} \frac{\log \#\{g \in H : d(o, g) \leq n\}}{n}.$$

We prove the following.

Theorem 5.1. *Let G be a finitely generated group with infinitely many ends. Then, for every $\lambda \in (0, 1)$, we have*

$$Hdim_{\rho_\lambda}(\partial_\mathcal{E}G) = -\frac{v_G}{\log \lambda}.$$

By Lemma 4.3, we have $Hdim(\partial_\mathcal{E}G) \leq Hdim(\partial_\mathcal{F}G)$ and [45, Lemma 4.1] shows that for any $\lambda \in (0, 1)$, we have $Hdim(\partial_\mathcal{F}G) \leq -\frac{v_G}{\log \lambda}$. The remainder of the section is devoted to proving the following proposition, which will thus conclude the proof of Theorem 5.1.

Proposition 5.2. *With the same notations, we have*

$$Hdim_{\rho_\lambda}(\partial_\mathcal{E}G) \geq -\frac{v_G}{\log \lambda}.$$

5.1. Preparatory lemmas. Recall that by [4, Lemma 5.1], the action of G on the end boundary is a convergence action. Hence, we can consider conical points in $\partial_\mathcal{E}G$ and hyperbolic elements in the sense of [4] corresponding to this action. Recall that an element is elliptic if it has finite order and that an infinite order element is parabolic if it fixes exactly one point on $\partial_\mathcal{E}G$ and hyperbolic if it fixes exactly two points on $\partial_\mathcal{E}G$. Moreover, if g is parabolic and fixes ξ , then for any ζ , $g^n \cdot \zeta$ converges to ξ as $n \rightarrow \pm\infty$, see [3, Section 2] for more details. It follows from Lemma 4.3 that if g is parabolic for the action on the Floyd boundary, then it is parabolic for the action on the end boundary. Thus, an element $g \in G$ which is hyperbolic for the action on the end boundary is also hyperbolic for the action on the Floyd boundary. and by [60, Lemma 7.2], such an element is contracting.

Lemma 5.3. *Let f be a hyperbolic element with two fixed points $\xi_- \neq \xi_+ \in \partial_\mathcal{E}G$. Then there exists a finite set $E_f \in \mathcal{F}(\Gamma)$ of edges such that for every large enough $n_0 > 0$ and for every $n \geq 2n_0$, the two elements 1 and f^n are separated by $f^{n_0}E_f$. Moreover, $d(1, f^{n_0}E_f) \geq d(1, f^{n_0}) - c_0$ for some uniform constant c_0 .*

Proof. The two distinct ends $\xi_- \neq \xi_+$ are separated by a finite set $E_f \in \mathcal{F}(\Gamma)$, and since they are fixed by any power of f , $\xi_- \neq \xi_+$ are separated by $E_n := f^n E_f$ for any n . Note that $n \in \mathbb{Z} \mapsto f^n$ is a quasi-geodesic so that the two half-rays converge to the corresponding ends ξ_- and ξ_+ . Thus, whenever n_0 is large enough, f^{-n_0} and f^{n_0} are separated by E_f for any $n \geq 2n_0$. Up to translation, 1 and f^n are thus separated by E_{n_0} . Moreover, $d(1, f^{n_0}) \leq d(1, f^{n_0}E_f) + \sup_{x \in E_f} d(f^{n_0}x, f^{n_0})$ and since E_f is fixed, the second term in the right-hand side is uniformly bounded, which concludes the proof. \square

Let F be a set of three pairwise independent hyperbolic elements in G . We write $F^n = \{f^n : f \in F\}$ for any $n \geq 1$.

Lemma 5.4. *There exists an integer $n_1 > 0$ with the following property for every $n \geq n_1$. For any $g, h \in G$, there exist $f \in F$ and a finite set E_f of edges separating 1 and f^n and such that any path between g^{-1} and $f^n h$ has to cross E_f .*

Proof. Since hyperbolic elements are contracting, by [61, Lemma 2.14], there exist $n_1, \epsilon > 0$ with the following property: for any $n > n_1$ and $f^n \in F^n$ the points 1 and f^n stay within the ϵ -neighborhood of any geodesic $[g^{-1}, f^n h]$, so that the path $\gamma = [g^{-1}, 1][1, f^n][f^n, f^n h]$ is a $(1, 4\epsilon)$ -quasi-geodesic. Let E be provided by Lemma 5.3 for the element f . Hence, for large enough n_0 , assuming that $n \geq 2n_0$, the two elements 1 and f^n are separated by $E_f := f^{n_0} E$.

For given λ, c , we choose n_0 to satisfy $d(1, F^{n_0}) > c + \lambda \text{diam}(E_f) + c_0$.

Claim 5.5. *If γ is a (λ, c) -quasi-geodesic path containing $[1, f^n]$ for some $\lambda, c > 0$, then the endpoints of γ are separated by E_f .*

Proof of the Claim. Denote by γ_1 and γ_2 the corresponding subpaths before and after $[1, f^n]$. Observe that γ_1 and γ_2 are disjoint with E_f . Indeed, assume by contradiction that $x \in \gamma_1 \cap E \neq \emptyset$, the case for γ_2 being symmetric. By Lemma 5.3, the elements 1, f^n are separated by E_f and $d(1, E_f) \geq d(1, f^{n_0}) - c_0$. We then choose $y \in [1, f] \cap E_f$ so that $d(1, y) \geq d(1, f^{n_0}) - c_0$. This yields a subpath α of γ whose endpoints x and y are at most $\text{diam}(E_f)$ apart. Since $x, 1, y$ are aligned in this order on α , the quasi-geodesicity implies that $d(1, y) \leq \lambda d(x, y) + c$. This contradicts the choice of n_0 above.

As a consequence, any path between the two endpoints of γ pass through E_f . Indeed, if there was such a path disjoint with E_f , we would obtain a path from 1 to f^n disjoint with E_f , contradicting Lemma 5.3. \square

The proof is concluded by the Claim applied to the $(1, 4\epsilon)$ -quasi-geodesic $\gamma = [g^{-1}, 1][1, f^n][f^n, f^n h]$. \square

Let $r = \max\{\text{diam}(E_f) : f \in F\}$, where E_f is given by Lemma 5.4. The following definition refines the notion of bottleneck given in the last section.

Definition 5.6. *Let $x, y \in G \cup \partial_{\mathcal{E}} G$ and let n_1 be given by Lemma 5.4. A (r, F) -bottleneck point between x, y is a point $b \in G$ such that any path between x and y has to cross the r -neighborhood of $b[1, f]$ for some $f \in F^{n_1}$.*

Note that if b is a (r, F) -bottleneck point between x and y , then b is in $b[1, f]$ so it lies in a M -bottleneck for $M = (d(1, F^{n_1}) + 2r)$ in the sense of Definition 4.5. An immediate result follows by the same argument in Lemma 4.3.

Lemma 5.7. *If b is a (r, F) -bottleneck point between g, h , then the Floyd distance $\delta_{\lambda, b}(g, h)$ based at b is bounded below by a constant depending λ, r, F only.*

5.2. Construction of rooted geodesic trees with bottlenecking property. The strategy in proving Proposition 5.2 is similar to [45]. We shall construct a sequence of rooted quasi-geodesic trees so that the Hausdorff dimension of their ends tends to the Hausdorff dimension of $\partial_{\mathcal{E}} G$.

Before getting into the construction, we introduce the following definition of ends with the uniform bottlenecking property, which are uniform conical points in the sense of [45]. We fix r, F and n_1 as in Section 5.1.

Definition 5.8. *A path γ in Γ has the L -bottlenecking property for some $L > 0$ if there exists a sequence of (r, F^{n_1}) -bottleneck points $b_i \in \gamma$ between the endpoints of γ such that $\sup\{d(b_i, b_{i+1}) : i \geq 1\} \leq L$ and $d(x, \{b_{i+1} : i \geq 1\}) \leq L/2$ for any $x \in \gamma$.*

An end ξ has the L -bottlenecking property for some $L > 0$ if there exists a geodesic ray $[1, \xi]$ with the L -bottlenecking property.

Remark 5.9. By definition, an end with L -bottlenecking property must be thin, but the converse is false.

If H is a free semigroup with a free basis B , then the standard Cayley graph T of H with respect to B is a rooted tree at 1, where every edge with unit length is labeled by a letter in B .

The construction of rooted trees in the proof of Proposition 5.2 will be given by the standard Cayley graphs of a sequence of free semigroups described in the following lemma. Given $n, \Delta > 0$, define the annulus set

$$A(n, \Delta) = \{g \in G : |d(1, g) - n| \leq \Delta\}.$$

Lemma 5.10. *For any $0 < v < v_G$, there exists a free semigroup H with a free basis B and a constant $L = L(v) > 0$ with $\lim_{v \rightarrow v_G} L(v) = \infty$ so that the following hold.*

- (1) *There exists a quasi-isometric embedding map Φ from the standard Cayley graph T of H into the Cayley graph Γ of G , which is induced by the inclusion $H \subset G$ and such that each geodesic issuing from the identity in T is sent to a path with the L -bottlenecking property.*
- (2) *The map Φ extends to a topological embedding $\partial\Phi$ of the end boundary $\partial_\varepsilon H$ of H into $\partial_\varepsilon G$.*
- (3) *We have $v_H > v$ and $C_1 \exp(Lv_H) \leq \sharp H \cap A(1, n, \Delta) \leq C_2 \exp(Lv_H)$ for $n \geq 1$, where C_1 and C_2 depend on Δ .*

Proof. (1) We set $L_\pm = n \pm (\Delta + d(1, F^{n_1}))$. Let $\mathbb{W}(A)$ be the set of all finite words over an alphabet set A . Given a set $A \subset \Gamma$, we can define an *extension map* $\Phi : \mathbb{W}(A) \rightarrow G$ as follows: for any word $W = a_1 a_2 \cdots a_n \in \mathbb{W}(A)$, there exists a sequence $f_i \in F^{n_1}$ such that the path γ labeled by

$$(14) \quad \Phi(W) = a_1 \cdot f_1 \cdot a_2 \cdot f_2 \cdots a_{n-1} \cdot f_{n-1} \cdot a_n \cdot f_n \in G,$$

is a (λ, c) -quasi-geodesic for fixed constants $\lambda, c > 0$ depending only on F . Moreover, the path γ labeled by $\Phi(W)$ has the L_+ -bottlenecking property. The choice of the points f_i is made by iterating the construction given by Lemma 5.4. Precisely, by [61, Lemma 2.16], one can choose the f_i so that γ is a (λ, c) -quasi-geodesic with fixed λ and c . Then, by the Claim 5.5, the terminal endpoints of the segments labeled by a_i ($1 \leq i \leq n$) give n distinct (r, F) -bottleneck points between 1 and $\Phi(W)$ with pairwise distance between L_- and L_+ . Those bottlenecks shall be called *canonical*.

To generate a free semigroup, we need the following property, proved in [61, Lemma 2.19]. There exists a fixed constant $C_0 > 0$ such that for every large enough n and every $C \geq C_0$, there exists a C -separated set $A \subset A(n, \Delta)$ with the following properties:

- (a) $\sharp A \succ_C \sharp A(n, \Delta)$,
- (b) there is a common $f \in F^{n_1}$ for each pair $(a, a') \in A \times A$ in the path (14) with the L -bottlenecking property.

If C is taken sufficiently large, then the map $\Phi : (Af)^n \rightarrow G$ as above is injective. Indeed, consider two paths $\beta_1 = \Phi(W_1)$ and $\beta'_1 = \Phi(W'_1)$, where $W_1 = a_1 f \cdots a_m f$ and $W'_1 = a'_1 f \cdots a'_n f$. We shall prove the following stronger fact, used later on in Lemma 5.12. Denote by $(\beta_1)_+$, respectively $(\beta'_1)_+$ the terminal endpoint of β_1 , respectively β'_1 .

Claim 5.11. *If $d((\beta_1)_+, (\beta'_1)_+) \leq D$, then $d(a_1, a'_1) < C$ for some $C = C(D)$.*

By choosing $C \geq C(0)$ the claim proves the injectivity of Φ . Hence, the set $B := Af$ generates a free semi-group denoted by H .

Proof of the Claim. Write $g = \Phi(W_1) = (\beta_1)_+$ and $g' = \Phi(W'_1) = (\beta'_1)_+$. Since the point a_1 in β_1 is a (r, F) -bottleneck point between 1 and g , by Lemma 5.7 and Lemma 2.5 applied to the Floyd distance δ_{λ, a_1} based at a_1 , there exists a constant C depending only on r, F such that $d(a_1, [1, g]) < C$. Since $d(g, g') \leq D$, concatenating the geodesic $[1, g]$ with a geodesic from g to g' yields a quasi-geodesic α whose parameters only depend on D . Hence, Lemma 2.5 shows that $d(a'_1, \alpha) < C'$ where C' only depends on r, F and D . Using again that $d(g, g') \leq D$, we finally obtain that $d(a'_1, [1, g]) \leq C''$. By the choice of $a_1, a'_1 \in A \subset A(n, \Delta)$, we have $|d(1, a_1) - d(1, a'_1)| \leq 2\Delta$. We deduce that $d(a_1, a'_1) < C + C'' + 2\Delta$, which concludes the proof. \square

(2) Note that the image of each geodesic ray γ in T is a path with the bottlenecking property which converges to an end in $\partial_\varepsilon G$. We deduce that the map $\gamma \mapsto \phi(\gamma)$ induces the desired topological embedding.

(3) For a given v , one can choose n big enough such that $v_H > v$ for a free semi-group $H = \langle Af \rangle$ constructed as above. This is possible since $\sharp A \succ_C \sharp A(n, \Delta)$, and

$$v = \lim_{n \rightarrow \infty} \frac{\log \sharp A(n, \Delta)}{n}.$$

See [61, Section 3] for the details. The purely exponential growth

$$\sharp H \cap N(o, r) \asymp \exp(v_H r)$$

follows by standard arguments using the fact that the subgroup H is contracting, as stated there. See an argument in the proof of [45, Lemma 3.9]. \square

5.3. Completion of proof of Proposition 5.2. We shall first prove that the embedding $\partial\Phi : \partial_\varepsilon H \rightarrow \partial_\varepsilon G$ is bi-Lipschitz with respect to visual metrics for appropriate choice of parameters. The Hausdorff dimension of $\partial_\varepsilon G$ will then be bounded from below by that of the ends of the rooted tree T which will be easier to compute.

Recall T is the standard Cayley graph of H with respect to the free base $B = Af$ whose edges with unit length are labeled by a letter in B .

Lemma 5.12. *Under the assumption of Lemma 5.10, assume that $\Phi(p) = \xi$ and $\Phi(q) = \eta$ where $p, q \in \partial_\varepsilon T$. Let m be the length of the intersection $[1, p] \cap [1, q]$ where $[1, p], [1, q]$ are geodesic rays in T . Then*

$$\rho_\lambda(\xi, \eta) \asymp_L \lambda^{mL}.$$

Proof. In the proof, let us write

$$[1, p] = (a_1 f)(a_2 f) \cdots (a_m f)(a_{m+1} f) \cdots$$

and

$$[1, q] = (a'_1 f)(a'_2 f) \cdots (a'_m f)(a'_{m+1} f) \cdots$$

and denote $\beta_1 = \Phi([1, p])$ and $\beta_2 = \Phi([1, q])$. Those are two rays with the L -bottlenecking property ending at ξ and η . Let $[1, p] \cap [1, q] = s_1 \cdots s_m$ be a geodesic of length m where $s_i \in Af$. In other words, $a_i = a'_i$ for $1 \leq i \leq m$.

By definition, $\rho_\lambda(\xi, \eta) = \lambda^n$ where n is the minimal integer such that ξ and η belongs to different components of the complement of the ball B_n . Note that $n \geq mL/2$. Indeed, the path $\tau = \Phi(s_1 \cdots s_m)$ issuing from 1 has the L -bottlenecking property, ends at $u = s_1 \cdots s_m$ and is contained in both β_1 and β_2 . Thus, there are m bottleneck points between 1 and u with pairwise distance at least $L/2$, so that the geodesic $[1, u]$ in the Cayley graph Γ of G has to pass through them in order. We then derive that $d(1, u) \geq mL/2$. Since the path obtained from $\beta_1 \cup \beta_2 \setminus \tau$ connects ξ to η , the definition of n implies that $n > mL/2$.

We shall prove the upper-bound $n \leq R := (m+1)L + r$, which will conclude the proof. By definition, it suffices to show that ξ and η belongs to the distinct components of $G \setminus B_R$.

Let b be the second (r, F) -bottleneck point on β_1 after the initial subpath τ of β_1 : it is the terminal endpoint of a_{m+2} in β_1 . Thus, any path from 1 to ξ has to cross the r -neighborhood of $b[1, f]$. Let $C = C(r + d(1, F^{n_1}))$ be the constant given by the Claim 5.11. If A is chosen to be C -separated, then $d(b, \beta_2) > r + d(1, F^{n_1})$ and thus, $\beta_2 \cap N_r(b[1, f]) = \emptyset$.

It remains to prove that any path from ξ to η has to intersect the ball B_R . If not, let Q be a path from ξ to η so that $Q \cap B_R = \emptyset$. Recall that b is the $(m+1)$ -th (r, F) -bottleneck point and thus $d(1, bf) \leq (m+1)L$. The value of R implies that B_R contains $N_r(b[1, f])$. Also recall that $\beta_2 \cap N_r(b[1, f]) = \emptyset$, so concatenating Q and β_2 yields a path $Q \cdot \beta_2$ from ξ to 1 avoiding the set $N_r(b[1, f])$. This is a contradiction, since b is a (r, F) -bottleneck point between 1 and ξ . As desired, ξ and η lie in different components in the complement of B_R . The proof of the upper bound $n \leq R$ follows. \square

Proposition 5.13. *Let $\alpha = \lambda^L$ for given $\lambda \in (0, 1)$. Then the embedding $\partial\Phi$ from the ends boundary $(\partial_\varepsilon T, \rho_\alpha)$ into the ends boundary $(\partial_\varepsilon G, \rho_\lambda)$ is bi-Lipschitz.*

Proof. By Lemma 5.12, we see $\rho_\alpha(\xi, \eta) = \alpha^m = \lambda^{mL} \asymp_L \rho_\lambda(\Phi(p), \Phi(q))$. Thus, the map from $\partial_\varepsilon T$ is bi-Lipschitz onto its image in $\partial_\varepsilon G$. \square

Let T be any infinite rooted tree, with vertex set V partitioned by depth (distance from the root vertex): $V = \cup_{n=0}^\infty V_n$. Let N_n and M_n be increasing sequences of positive integers. Say that T is $\{M_n\}$ -regular relative to $\{N_n\}$ if for every $n \geq 1$ every vertex x at depth N_n has exactly M_{n+1} descendant vertices at depth N_{n+1} .

Lemma 5.14. [37, Lemma 2] *Let T be an infinite rooted tree with space of ends $\partial_\varepsilon T$ equipped with the visual metric of parameter $\alpha \in (0, 1)$. If there is an increasing sequence of integers N_n satisfying $\lim_{n \rightarrow \infty} \frac{N_{n+1}}{N_n} = 1$ such that, for some sequence $\{M_n\}$ of positive integers, T is $\{M_n\}$ -regular with respect to $\{N_n\}$, then the Hausdorff dimension of $\partial_\varepsilon T$ (relative to the visual metric ρ_α) is given by*

$$Hdim_{\rho_\alpha}(\partial_\varepsilon T) = \liminf_{n \rightarrow \infty} \frac{\log |V_{N_n}|}{-N_n \log \alpha} = \liminf_{n \rightarrow \infty} \frac{\log \prod_{j=1}^n M_j}{-N_n \log \alpha}.$$

We apply Lemma 5.14 to the standard Cayley graph T of H , where $M_n = \sharp Af$ and $N_n = n$. We obtain

$$Hdim_{\rho_\alpha}(\partial_\varepsilon T) = \liminf_{n \rightarrow \infty} \frac{\log \prod_{j=1}^n \sharp Af}{-n \log \alpha} = \frac{\log \sharp Af}{-L \log \lambda}.$$

By Lemma 5.10, $C_1 \exp(Lv_H) \leq \sharp Af \leq C_2 \exp(Lv_H)$, so

$$Hdim_{\rho_\alpha}(\partial T) \geq -\frac{\log C_2 + Lv_H}{L \log \lambda},$$

where $v < v_H \leq v_G$.

Since a bi-Lipschitz map preserves the Hausdorff dimension, by Proposition 5.13, we have

$$Hdim_{\rho_\lambda}(\partial_\varepsilon \Gamma) \geq -\frac{\log C_2 + Lv}{L \log \lambda}.$$

As v tends to v_G , L goes to infinity, so we finally obtain

$$Hdim_{\rho_\lambda}(\partial_\varepsilon \Gamma) \geq -\frac{v_G}{\log \lambda}.$$

The proof of Proposition 5.2 and thus Theorem 5.1 is complete. \square

6. CHARACTERIZING THE DOUBLING PROPERTY: PROOF OF THEOREM 1.6

It is well-known that a virtually free group is hyperbolic and that its end boundary endowed with a visual metric is bi-Hölder to its Gromov boundary endowed with a Gromov's visual metric. The latter has the doubling property, since the Patterson-Sullivan measure is doubling (even Ahlfors regular) by the work of Coorntart [8]. We prove here the following result.

Proposition 6.1. *If a finitely generated group admits a splitting over finite edge groups as a finite graph of groups with at least one one-ended vertex group, then the visual metric is not doubling.*

Theorem 1.6 is then a consequence of this proposition. Indeed, accessible groups admit a splitting over finite edge groups as a finite graph of groups \mathcal{G} , so that the vertex groups either are finite or one-ended. In the former case, the group is virtually free, in the later case, it satisfies the assumptions of Proposition 6.1.

Proof. Consider a splitting over finite edge groups as a finite graph of groups \mathcal{G} and let H be a one-ended vertex group. Notice that a metric space (X, d) is doubling if and only if for any (or some) $\theta \in (0, 1)$, there exists $N = N(\theta) > 0$ such that every ball of radius $s > 0$ can be covered by at most N balls of radius θs . To prove that the end boundary is not doubling, our strategy will be as follows. We will consider for every n the ball of radius $s = \lambda^n$ centered at the end ξ of G corresponding to the unique end of H . We will prove that for some fixed θ , this ball cannot be covered by $N(n)$ balls of radius $\theta s(n)$, where $N(n)$ goes to infinity. This will conclude the proof. We present the details below.

By the Bass-Serre theory, G is isomorphic to either an amalgamated product $H *_F K$ or a HNN extension $H *_F$ over a finite group F , where K might not be one-ended. In what follows, we only consider the amalgamated product case, the HNN extension case being similar. Let $G = H *_F K$ so that H and K are generated by two finite sets S and T respectively and assume for simplicity that both sets S and T contain F . The normal form given by [57, (9.2), (9.3)] shows that the Cayley graph Γ of G with respect to $S \cup T$ is obtained as the disjoint union

$$\bigsqcup_{g \in G} \{g \cdot \text{Cay}(H, S), g \cdot \text{Cay}(K, T)\}.$$

glued along cosets gF between $g \cdot \text{Cay}(H, S)$, $g \cdot \text{Cay}(K, T)$. In particular, for any end $\eta \neq \xi$, any path from η to ξ has to pass through a finite set F_η corresponding to some F -coset in H and any path from 1 to η also has to pass through F_η . Since we made the assumption that $F \subset S \cap T$, the diameter of F_η is at most 1, that is any two distinct vertices in F_η are connected by an edge.

Let $S_n := \{g \in G : d(1, g) = n\}$ be the n -sphere in the Cayley graph Γ of G . By definition of the visual metric, if $\rho_\lambda(\xi, \eta) = \lambda^n$ for $n \geq 1$, then every path from ξ and η stays within distance n of the identity and exits H at some vertex in F_η . Moreover there exists such a path from ξ and η which is disjoint from B_{n-1} . Recalling that the diameter of F_η is at most 1, we see that $d(1, F_\eta) \geq n - 1$. Assume by contradiction that $d(1, F_\eta) > n$. Then, the sub-path from η to F_η stays in the complement of the ball B_n of radius n . Indeed, if this were not the case, this sub-path would pass through a point $g \in B(1, n)$ and so concatenating

a path from 1 to g and the path from g to η would yield a path from 1 to η not passing through F_η . Now, since $\text{Cay}(H, S)$ is one-ended, one can join F_η with ξ by a path which also stays in the complement of B_n , contradicting the fact that any path from η to ξ needs to pass through this ball. Thus, $d(1, F_\eta) \leq n$. Similarly, if $d(1, F_\eta) \geq n$, then $\rho_\lambda(\xi, \eta) \leq \lambda^n$. Set $s = \lambda^n$. We can reformulate the above discussion by the following inclusions :

$$(15) \quad \{\eta \neq \xi \in \partial_\varepsilon \Gamma : d(1, F_\eta) \geq n\} \subseteq B^{\rho_\lambda}(\xi, s) \setminus \xi \subseteq \{\eta \neq \xi \in \partial_\varepsilon \Gamma : d(1, F_\eta) \geq n-1\}.$$

Since H is one-ended, it is infinite and it is not virtually cyclic. By Gromov's polynomial growth theorem, the growth function of H is at least quadratic, hence super-linear. Therefore, fixing $k \geq 2$, the number of elements in

$$S_{n+k}(H) := S_{n+k} \cap H$$

grows at least linearly in n and in particular goes to infinity as n goes to infinity.

For any point $h \in S_{n+k}(H)$, choose an end $\eta = \eta(h) \neq \xi \in \partial_\varepsilon \Gamma$ so that h lies in F_η . By definition of h , $d(1, F_\eta) \leq n+k$ and since the diameter of F_η is at most 1, we also have $d(1, F_\eta) \geq n+k-1$. According to (15), $\rho_\lambda(\xi, \eta) \leq \lambda^{n+k-1}$. Thus, $\eta \in B^{\rho_\lambda}(\xi, s)$.

Observe that for any two elements $h_1 \neq h_2 \in S_{n+k}(H)$ with $d(h_1, h_2) \geq 2$, we claim that

$$\rho_\lambda(\eta_1, \eta_2) \geq \lambda^{n+k+1}$$

where $\eta_1 = \eta(h_1)$, $\eta_2 = \eta(h_2)$. Indeed, $d(h_1, h_2) \geq 2$ implies that $F_{\eta_1} \cap F_{\eta_2} = \emptyset$. Notice that any path from η_1 to η_2 has to pass through both F_{η_1} and F_{η_2} , otherwise one would produce a path from 1 to η_i not passing through F_{η_i} for some $i \in \{1, 2\}$. Hence, any path from η_1 to η_2 has to intersect the ball B_{n+k+1} and the claim follows.

We are ready to finish the proof. Set $\theta = \lambda^{k+1}$ and let $N(n)$ be the maximal size of a set $\Sigma \subset S_{n+k}(H)$ such that any two elements h, h' in Σ satisfy that $d(h, h') \geq 2$. Then, $N(n)$ goes to infinity as n goes to infinity. We can thus produce $N(n)$ points which are θs -separated in $B_{\rho_\lambda}(\xi, s)$. In other words, the ball $B_{\rho_\lambda}(\xi, s)$ cannot be covered by $N(n)$ balls of radius θs . Since k is fixed and $N(n)$ goes to infinity as n goes to infinity, this provides a contradiction with the definition of the doubling property. This concludes the proof. \square

REFERENCES

- [1] Sébastien Blachère, Peter Haïssinsky, and Pierre Mathieu. Harmonic measures versus quasiconformal measures for hyperbolic groups. *Annales Scientifiques de l'École Normale Supérieure*, 44:683–721, 2011.
- [2] Mario Bonk and Oded Schramm. Embeddings of Gromov hyperbolic spaces. *Geometric and Functional Analysis*, 10:266–306, 2000.
- [3] Brian Bowditch. Cut points and canonical splittings of hyperbolic groups. *Acta Mathematica*, 180:145–186, 1998.
- [4] Brian Bowditch. Groups acting on Cantor sets and the end structure of graphs. *Pacific journal of mathematics*, 207:31–60, 2002.
- [5] Brian Bowditch. Relatively hyperbolic group. *International Journal of Algebra and Computation*, 22:66 pp, 2012.
- [6] Elisabetta Candellero, Lorenz Gilch, and Sebastian Müller. Branching random walks on free products of groups. *Proceedings of the London Mathematical Society*, 104:1085–1120, 2012.
- [7] Donald Cartwright, Paolo Soardi, and Wolfgang Woess. Martin and end compactifications for non-locally finite graphs. *Transactions of the American Mathematical Society*, 338:679–693, 1993.
- [8] Michel Coornaert. Mesures de Patterson-Sullivan sur le bord d'un espace hyperbolique au sens de Gromov. *Pacific Journal of Mathematics*, 159:241–270, 1993.
- [9] Yves Cornuier. On sublinear bilipschitz equivalence of groups. *Annales Scientifiques de l'École Normale Supérieure*, 52:1201–1242, 2019.
- [10] Yves Cornuier. On the space of ends of infinitely generated groups. *Topology and its Applications*, 263:279–298, 2019.
- [11] François Dahmani and Asli Yaman. Bounded geometry in relatively hyperbolic groups. *New York Journal of Mathematics*, 2:89–95, 2005.
- [12] Yves Derriennic. Marche aléatoire sur le groupe libre et frontière de Martin. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 32:261–276, 1975.
- [13] Yves Derriennic. Quelques applications du théorème ergodique sous-additif. In *Journées sur les marches aléatoires*, volume 74, pages 183–201. Astérisque, Société mathématique de France, 1980.
- [14] Martin Dunwoody. The accessibility of finitely presented groups. *Inventiones Mathematicae*, 81:449–457, 1985.
- [15] Martin Dunwoody. An inaccessible group. In *Geometric group theory, Vol. 1 (Sussex, 1991)*, volume 181, pages 75–78. Cambridge University Press, 1993.
- [16] Matthieu Dussaule and Ilya Gekhtman. Entropy and drift for word metric on relatively hyperbolic groups. arXiv:1811.10849, To appear in *Groups, Geometry, and Dynamics*, 2018.
- [17] Benson Farb. Relatively hyperbolic groups. *Geometric and Functional Analysis*, 8:810–840, 1998.

- [18] William Floyd. Group completions and limit sets of Kleinian groups. *Inventiones Mathematicae*, 57:205–218, 1980.
- [19] Hans Freudenthal. Über die enden diskreter räume und gruppen. *Commentarii Mathematici Helvetici*, 17:1–38, 1945.
- [20] Ilya Gekhtman, Victor Gerasimov, Leonid Potyagailo, and Wenyuan Yang. Martin boundary covers Floyd boundary. arXiv:1708.02133, 2017.
- [21] Victor Gerasimov. Expansive convergence groups are relatively hyperbolic. *Geometric and Functional Analysis*, 19:137–169, 2009.
- [22] Victor Gerasimov. Floyd maps for relatively hyperbolic groups. *Geometric and Functional Analysis*, 22:1361–1399, 2012.
- [23] Victor Gerasimov and Leonid Potyagailo. Quasi-isometric maps and Floyd boundaries of relatively hyperbolic groups. *Journal of the European Mathematical Society*, 15:2115–2137, 2013.
- [24] Victor Gerasimov and Leonid Potyagailo. Non-finitely generated relatively hyperbolic groups and Floyd quasiconvexity. *Groups, Geometry, and Dynamics*, 9:369–434, 2015.
- [25] Sébastien Gouëzel, Frédéric Mathéus, and François Maucourant. Entropy and drift in word hyperbolic groups. *Inventiones Mathematicae*, 211:1201–1255, 2018.
- [26] Yves Guivarc’h and Yves Le Jan. Asymptotic winding of the geodesic flow on modular surfaces and continued fractions. *Annales Scientifiques de l’ENS*, 26:23–50, 1993.
- [27] Peter Haïssinsky. Marches aléatoires sur les groupes hyperboliques. Course notes from the conference "Regards croisés sur les marches aléatoires et la géométrie des groupes, en l’honneur d’Emile Le Page", available on <http://web.univ-ubs.fr/lmam/matheus/emilfest.html>, 2011. French.
- [28] Brendan Burns Healy. Rigidity properties for hyperbolic generalizations. *Canadian Mathematical Bulletin*, 63:66–76, 2020.
- [29] Juha Heinonen. *Lectures on analysis on metric spaces*. Springer, 2001.
- [30] Geoffrey Hruska. Relative hyperbolicity and relative quasiconvexity for countable groups. *Algebra, Geometry and Topology*, 10:1807–1856, 2010.
- [31] Vadim Kaimanovich. Boundaries of invariant Markov operators: the identification problem. In *Ergodic theory of \mathbb{Z}^d actions*, pages 127–176. London Mathematical Society Lecture Note series, vol. 228, Cambridge University Press, 1996.
- [32] Vadim Kaimanovich. Hausdorff dimension of the harmonic measure on trees. *Ergodic Theory and Dynamical Systems*, 18:631–660, 1998.
- [33] Vadim Kaimanovich. The Poisson formula for groups with hyperbolic properties. *Annals of Mathematics*, 152:659–692, 2000.
- [34] Vadim Kaimanovich and Anatoly Vershik. Random walks on discrete groups: boundary and entropy. *Annals of Probability*, 11:457–490, 1983.
- [35] Anders Karlsson. Boundaries and random walks on finitely generated infinite groups. *Arkiv för Matematik*, 41:295–306, 2003.
- [36] Anders Karlsson. Free subgroups of groups with nontrivial Floyd boundary. *Communications in Algebra*, 31:5361–5376, 2003.
- [37] Steven Lalley and Thomas Sellke. An extension of Hawkes’ theorem on the Hausdorff dimension of a Galton-Watson tree. *Probability Theory and Related Fields*, 116:41–56, 2000.
- [38] François Ledrappier. Some asymptotic properties of random walks in free groups. In *Topics in Probability and Lie Groups: Boundary Theory*, pages 117–152. CRM Proceedings and Lecture Notes, 2001.
- [39] Joseph Maher and Giulio Tiozzo. Random walks on weakly hyperbolic groups. *Journal für die reine und angewandte Mathematik*, 742:187–239, 2018.
- [40] Richard Mauldin, Tomasz Szarek, and Mariusz Urbański. Graph directed Markov systems on Hilbert spaces. *Mathematical Proceedings of the Cambridge Philosophical Society*, 157:455–488, 2009.
- [41] Denis Osin. Relatively hyperbolic groups: Intrinsic geometry, algebraic properties, and algorithmic problems. *Memoirs of the American Mathematical Society*, 20, 2006.
- [42] Denis Osin. Acylindrically hyperbolic groups. *Transactions of the American Mathematical Society*, 368:851–888, 2016.
- [43] Massimo Picardello and Wolfgang Woess. Martin boundaries of random walks: ends of trees and groups. *Transactions of the American Mathematical Society*, 302:185–205, 1987.
- [44] Massimo Picardello and Wolfgang Woess. Harmonic functions and ends of graphs. *Proceedings of the Edinburgh Mathematical Society*, 31:457–461, 1988.
- [45] Leonid Potyagailo and Wenyuan Yang. Hausdorff dimension of boundaries of relatively hyperbolic groups. *Geometry and Topology*, 23:1779–1840, 2019.
- [46] Vincent Le Prince. Dimensional properties of the harmonic measure for a random walk on a hyperbolic group. *Transactions of the American Mathematical Society*, 359:2881–2898, 2007.
- [47] Vincent Le Prince. A relation between dimension of the harmonic measure, entropy and drift for a random walk on a hyperbolic space. *Electronic communications in Probability*, 13:45–53, 2008.
- [48] Anja Randecker and Giulio Tiozzo. Cusp excursion in hyperbolic manifolds and singularity of harmonic measure. arXiv:1904.11581, 2019.
- [49] Alessandro Sisto. Projections and relative hyperbolicity. *L’Enseignement Mathématique*, 59:165–181, 2013.
- [50] Alessandro Sisto and Sam Taylor. Largest projections for random walks and shortest curves for random mapping tori. arXiv:1909.02096, To appear in Mathematical Research Letters, 2019.
- [51] John Stallings. On torsion-free groups with infinitely many ends. *Annals of Mathematics*, 88:312–334, 1968.
- [52] John Stallings. *Group theory and three-dimensional manifolds*. Yale University Press, 1971.
- [53] Ryokichi Tanaka. Dimension of harmonic measures in hyperbolic spaces. *Ergodic Theory and Dynamical Systems*, 39:474–499, 2019.

- [54] Carsten Thomassen and Wolfgang Woess. Vertex-transitive graphs and accessibility. *Journal of Combinatorial Theory*, 58:248–268, 1993.
- [55] Giulio Tiozzo. Sublinear deviation between geodesics and samples paths. *Duke mathematical Journal*, 164:511–539, 2015.
- [56] Hung Cong Tran. Relations between various boundaries of relatively hyperbolic groups. *International Journal of Algebra and Computation*, 1-23:1551–1572, 2013.
- [57] Wolfgang Woess. Boundaries of random walks on graphs and groups with infinitely many ends. *Israel Journal of Mathematics*, 68:271–301, 1989.
- [58] Asli Yaman. A topological characterisation of relatively hyperbolic groups. *Journal for Pure and Applied Mathematics*, 566:41–89, 2004.
- [59] Wenyuan Yang. Patterson-Sullivan measures and growth of relatively hyperbolic groups. arXiv:1308.6326, 2013.
- [60] Wenyuan Yang. Growth tightness for groups with contracting elements. *Mathematical Proceedings of the Cambridge Philosophical Society*, 157:297–319, 2014.
- [61] Wenyuan Yang. Statistically convex-cocompact actions of groups with contracting elements. *International Mathematics Research Notices*, 23:7259–7323, 2019.

YOUR ADDRESS

Email address: `matthieu.dussaule@univ-nantes.fr`

BEIJING INTERNATIONAL CENTER FOR MATHEMATICAL RESEARCH (BICMR), BEIJING UNIVERSITY, NO. 5 YIHEYUAN ROAD, HAIDIAN DISTRICT, BEIJING, CHINA

Email address: `yabziz@gmail.com`