



**HAL**  
open science

## **Diagnostic Accuracy of the French Version of the Adult Attention Deficit / Hyperactivity Disorder Self-Report Screening Scale for DSM-5 (ASRS-5)**

Stéphanie Baggio, Sophie Bayard, Clémence Cabelguen, Martin Desseilles, Marie Gachet, Charlotte Kraemer, Hélène Richard-Lepouriel, Rosetta Nicastro, Stéphanie Bioulac, Anne Sauvaget, et al.

### ► To cite this version:

Stéphanie Baggio, Sophie Bayard, Clémence Cabelguen, Martin Desseilles, Marie Gachet, et al.. Diagnostic Accuracy of the French Version of the Adult Attention Deficit / Hyperactivity Disorder Self-Report Screening Scale for DSM-5 (ASRS-5). *Journal of Psychopathology and Behavioral Assessment*, 2020, 10.1007/s10862-020-09822-7 . hal-02985334

**HAL Id: hal-02985334**

**<https://hal.science/hal-02985334>**

Submitted on 31 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Diagnostic Accuracy of the French Version of the Adult Attention Deficit / Hyperactivity Disorder Self-Report Screening Scale for DSM-5 (ASRS-5)

BAGGIO, Stéphanie, *et al.*

### Abstract

Attention Deficit/Hyperactivity Disorder (ADHD) often persists into adulthood. However, few screening tools have been adapted to assess adult ADHD using the DSM-5 criteria. This study assessed the diagnostic accuracy of a French version of the ADHD Self-Report Screening for DSM-5 (ASRS-5). This multicentric cross-sectional study included 557 participants: 309 adult ADHD outpatients without bipolar disorder (BD)/borderline personality disorder (BPD) (n=236) or with BD/BPD (n=36) and 285 adults without ADHD who were either healthy volunteers (n=248) or outpatients with BD or BPD (n=37). Measures included ADHD diagnosis and the ASRS-5. The ASRS-5 was a good predictor of ADHD diagnosis (cut-off score  $\geq 13/24$ : sensitivity=84.3%, specificity=91.9%) in the sample of adult outpatients without comorbid disorders/healthy controls. Performances were lower with this cut-off score in some subgroups, notably low-severity ADHD symptomatology (sensitivity=63.5%) and participants with BD or BPD (sensitivity=91.7%, specificity=54.1%). The French ASRS-5 had acceptable screening [...]

### Reference

BAGGIO, Stéphanie, *et al.* Diagnostic Accuracy of the French Version of the Adult Attention Deficit / Hyperactivity Disorder Self-Report Screening Scale for DSM-5 (ASRS-5). *Journal of Psychopathology and Behavioral Assessment*, 2021, vol. 43, no. 2, p. 367-375

DOI : 10.1007/s10862-020-09822-7

Available at:

<http://archive-ouverte.unige.ch/unige:154392>

Disclaimer: layout of this document may differ from the published version.



UNIVERSITÉ  
DE GENÈVE

# **Diagnostic accuracy of the French version of the Adult Attention Deficit / Hyperactivity Disorder Self-Report Screening Scale for DSM-5 (ASRS-5)**

*Stéphanie Baggio<sup>1,2\*</sup>, Sophie Bayard<sup>3,4,5</sup>, Clémence Cabelguen<sup>6</sup>, Martin Desseilles<sup>7</sup>, Marie Gachet<sup>8</sup>, Charlotte Kraemer<sup>9</sup>, Hélène Richard-Lepouriel<sup>10</sup>, Rosetta Nicastrò<sup>10</sup>, Stéphanie Bouliac<sup>11</sup>, Anne Sauvaget<sup>12</sup>, Sébastien Weibel<sup>9,13</sup>, Nader Perroud<sup>10,14,15#</sup>, Régis Lopez<sup>16,17#</sup>*

<sup>1</sup> Division of Prison Health, Geneva University Hospitals and University of Geneva, Geneva, Switzerland

<sup>2</sup> Office of Corrections, Canton Zurich, Zurich, Switzerland

<sup>3</sup> University Paul Valéry Montpellier 3, Montpellier, France

<sup>4</sup> University of Montpellier, France

<sup>5</sup> EPSYLON EA 4556, Montpellier, France

<sup>6</sup> Department of Psychiatry, University Hospital of Nantes, Nantes, France.

<sup>7</sup> Department of Psychology, Faculty of Medicine, University of Namur, Belgium

<sup>8</sup> Department of Emergency Psychiatry & Acute Care, CHU Montpellier, Montpellier, France

<sup>9</sup> Department of Psychiatry, University Hospital of Strasbourg, Strasbourg, France

<sup>10</sup> Division of Psychiatric Specialties, Department of Psychiatry, University Hospitals of Geneva, Geneva, Switzerland

<sup>11</sup> Clinique du sommeil, CHU Pellegrin Boreaux, USR CNRS 3413, Université Bordeaux, France

<sup>12</sup> Nantes Université, CHU Nantes, Movement - Interactions - Performance, MIP, EA 4334, Nantes, France

<sup>13</sup> Inserm U1114, Strasbourg, France

<sup>14</sup> Department of Psychiatry, Faculty of Medicine, University of Geneva, Geneva, Switzerland

<sup>15</sup> Department of Psychiatry, Dalhousie University, Halifax, Nova Scotia, Canada

<sup>16</sup> Department of Neurology, CHU Montpellier, Montpellier, France

<sup>17</sup> Inserm U1061, Montpellier, France

\* **Corresponding author:** S. Baggio, Chemin du Petit Bel Air 2, 1226 Thônex, Switzerland, stephanie.baggio@hcuge.ch, +41 22 305 52 18

# These two authors contributed equally to the work.

**Conflict of Interest:** All authors declare that they have no conflict of interest.

## **Abstract**

Attention Deficit/Hyperactivity Disorder (ADHD) often persists into adulthood. However, few screening tools have been adapted to assess adult ADHD using the DSM-5 criteria. This study assessed the diagnostic accuracy of a French version of the ADHD Self-Report Screening for DSM-5 (ASRS-5). This multicentric cross-sectional study included 557 participants: 309 adult ADHD outpatients without bipolar disorder (BD)/borderline personality disorder (BPD) ( $n=236$ ) or with BD/BPD ( $n=36$ ) and 285 adults without ADHD who were either healthy volunteers ( $n=248$ ) or outpatients with BD or BPD ( $n=37$ ). Measures included ADHD diagnosis and the ASRS-5. The ASRS-5 was a good predictor of ADHD diagnosis (cut-off score  $\geq 13/24$ : sensitivity=84.3%, specificity=91.9%) in the sample of adult outpatients without comorbid disorders/healthy controls. Performances were lower with this cut-off score in some subgroups, notably low-severity ADHD symptomatology (sensitivity=63.5%) and participants with BD or BPD (sensitivity=91.7%, specificity= 54.1%). The French ASRS-5 had acceptable screening properties, even if its performance varied according to clinical variables. Further evidence is needed for patients with comorbid disorders having overlapping symptoms.

**KEYWORDS:** assessment; screening; validation.

# **Diagnostic accuracy of the French version of the Adult Attention Deficit / Hyperactivity Disorder Self-Report Screening Scale for DSM-5 (ASRS-5)**

## **Introduction**

There is increasing evidence that attention deficit/hyperactivity disorder (ADHD), a child-onset neurodevelopmental disorder, persists in adulthood (Caye et al. 2016; Kooij et al. 2010). It leads to functional impairment in several areas of life: psychosocial functioning, work, health care access and use, criminality and recidivism (Ginsberg et al. 2014; Mohr-Jensen and Steinhausen 2016; Philipp-Wiegmann et al. 2017; Young et al. 2015). Adult ADHD is also associated with comorbid psychiatric disorders, substance use, comorbid somatic diseases, and excess mortality because of the increase risk of suicide and accident (Asherson et al. 2016; Dalsgaard et al. 2015; Instanes et al. 2018; Shaw et al. 2012). Because of this long-term impairment, adult ADHD has become a major health concern (Asherson et al. 2016; Franke et al. 2018). Nevertheless, ADHD is often under-diagnosed and under-treated in adult populations (Weibel et al. 2020). Identifying adult ADHD with reliable screening scales has become crucial in primary care and for public health purposes.

To better take into account adult ADHD, the threshold of ADHD diagnosis has been revised in the fifth version of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5, APA 2013) compared to the fourth edition (DSM-IV, APA 2000). In the revised version, ADHD diagnosis requires a reduced number of symptoms during adult adulthood (five instead of six). In addition, changes in symptom wording for adolescents and adults have been added and are developmentally more appropriate. These changes better capture ADHD features in adulthood (Matte et al. 2015). Finally, although it does not specifically impact adult ADHD, the age of onset of ADHD symptoms has been changed (twelve instead of six).

To date, however, few screening scales have been adapted to assess adult ADHD using DSM-5 criteria. The widely used Adult ADHD Self-Report Screening Scale (ASRS v1.1), which is the World Health Organization's self-reported screening tool for ADHD in adults (Kessler et al. 2005; Kessler et al. 2007), has recently been updated (ASRS-5) to better match DSM-5 ADHD diagnosis (Ustun et al. 2017). The ASRS-5 yields good psychometric properties in a US sample from the general population (Ustun et al. 2017). The ASRS-5 includes four items from the ASRS v1.1 and two new non-DSM criteria (Kessler et al. 2010). To date, this screener has not been tested nor validated in any other language than English. In addition, even if some studies investigated the influence of other variables on the ASRS screening tool and compared psychometric performance across groups (Sonby et al. 2015; van de Glind et al. 2013), few studies investigated psychometric properties according to clinical characteristics. The impact of comorbidity on psychometric performance is also often neglected, even if ADHD is comorbid with some psychiatric disorders or share similar symptoms with others. For example, bipolar disorder (BD) and borderline personality disorder (BPD) are two highly comorbid disorders which have overlapping symptomatology with ADHD (Perroud et al. 2014; Weibel et al. 2018). The ASRS v1.1 showed reduced psychometric performances among patients with comorbid ADHD and BPD (sensitivity=72.8%, specificity=43.9%; Weibel et al. 2018). Similar results were obtained among BD patients (false positives=40%; Perroud et al. 2014) Therefore, including populations with comorbid disorders is crucial to achieve a better understanding of the psychometric performance of the ASRS-5.

Thus, the aim of our study was to examine the diagnostic accuracy of the French version of the ASRS-5. We also explored whether it was a reliable screening tool in detecting ADHD among different subgroups of the population, according to relevant clinical variables (severity, clinical presentation, and the presence of psychiatric comorbid disorders: BD and BPD).

## **Methods**

### ***Participants and procedure***

From 2018 to 2019 (10 months), 557 participants were included in the study: 309 adult ADHD outpatients without BD/BPD ( $n=236$ ) or with BD/BPD ( $n=36$ ) and 285 adults without ADHD who were either healthy volunteers ( $n=248$ , no formal ADHD diagnosis) or outpatients with BD or BPD ( $n=37$ ).

Adult outpatients with ADHD were recruited in six specialized French-speaking centers for the diagnosis and treatment of ADHD, located in France (Bordeaux, Montpellier [two centers], Nantes, and Strasbourg) and Switzerland (Geneva). All patients met the DSM-5 criteria for ADHD that require five symptoms or more in adulthood and six or more before the age of twelve, with significant impairment in at least two areas of daily life during the last six months prior to the interview and in childhood. All patients were assessed by clinicians having a long-lasting (more than five years) expertise in the assessment of adult ADHD and using structured clinical instruments (the ADHD evaluation adult version, ACE+, Young 2016) or the Diagnostic Interview for ADHD (DIVA 2.0, Kooij and Francken 2010) according to DSM-5 criteria. The presence of ADHD symptoms during childhood was systematically assessed using collateral reports and/or school records. Other mental health disorders, except BD (types 1 and 2) and BPD, were not considered in the study. The three ADHD clinical presentations were defined at the time of the interview: predominantly inattentive, predominantly hyperactive–impulsive, and combined presentation.

Healthy participants were community-dwelling adults who were recruited in Montpellier by means of advertisements, personal contacts, and through snowballing techniques. Three



participants were not included due to a past or actual diagnosis of ADHD. They self-completed the ASRS-5 at the Laboratory Epsilon (University Paul Valery Montpellier 3). The interviewers (trained final-year students) ensured that the questionnaire was adequately completed. After that, they had a face-to-face interview including medical information and socio-demographics.

BP/BPD patients without ADHD (BD:  $n=26$ , BPD:  $n=11$ ) were recruited in a specialized center for the treatment of these disorders in Geneva. Detailed information and descriptive statistics on the different centers are reported in **Table 1**.

All participants provided written informed consent to participate in the study. The study was approved by the local Ethics Committee of the Geneva University Hospitals (no. 2019-00991).

### ***Measures***

*ADHD diagnosis.* The diagnosis of ADHD was made or excluded by experienced clinicians, using validated and structured clinical interviews: the ADHD evaluation adult version (ACE+,(Young 2016) or the Diagnostic Interview for ADHD (DIVA 2.0, (Kooij and Francken 2010). The ACE+ is a semi-structured interview translated in French designed to assess ADHD in adults according to the DSM-5 criteria. It was used in one center (Geneva,  $n=128$ ). The remaining centers used the DIVA 2.0 ( $n=181$ ), a semi-structured clinical interview, originally designed to assess ADHD symptoms and related impairment according to the DSM-IV criteria. The DIVA 2.0 has been slightly modified to fulfil the DSM-5 criteria, by changing the age at onset cut-off from seven to twelve years of age and lowering the required number of symptoms in adulthood from six to five.

We also created groups according to ADHD severity using the diagnostic interview (DIVA 2.0 and ACE+). We summed up symptoms during adulthood and created groups corresponding to the four quartiles: low (5-9), moderate (10-11), high (12-14), and very high (15-18) severity.

*ASRS-5.* We used the World Health Organization Adult ADHD Self-Report Screening Scale for DSM-5 (ASRS-5, Ustun et al. 2017). Each of the six items is scored on a four-point Likert-type scale: 0 (*never*), 1 (*rarely*), 2 (*sometimes*), 3 (*often*), and 4 (*very often*), resulting in a scale with scores ranging from 0 to 24. In outpatient centers, the ASRS-5 was administered within one month after the clinical evaluation.

The translation process of the scale in French was as follows: (1) the scale was translated in French by the first author (SB); (2) NP translated the French version back into English; and (3) all discrepancies between the original English scale and the French translation were discussed between the two authors until an agreement was reached. The final version of the French ASRS-5 is presented in **Supplementary Table S1**.

*Demographic and clinical characteristics.* We also collected information on gender, age, level of education (*secondary education* vs. *higher education*), and having any prescription of psychotropic medication including psychostimulants (*yes/no*).

### ***Analytical strategy***

*Sample size calculation.* We used a non-inferiority test to define how many participants were needed to obtain a similar AUC as in the first validation study (Ustun et al. 2017). We used  $\alpha=5\%$ ,  $power=80\%$ , margin of equivalence= $10\%$  (Salazar et al. 2016),  $AUC=0.94$  (Ustun et al. 2017), and  $allocation\ ratio=1$ . A total of  $n=98$  (49 participants in each group) was needed. For the comorbid sample, we used  $AUC=0.83$  (Ustun et al. 2017). A total of  $n=48$  (24 participants in each group) was needed. The sample size was calculated for our main objective (diagnostic accuracy of the French version of the ASRS-5). The other objective was exploratory.

*Preliminary analyses.* We first computed descriptive statistics (means/standard deviations or percentages) for all variables and we compared socio-demographics between groups (ADHD

patients without BD/BPD vs. healthy control group and BPD/BD patients with ADHD vs. BPD/BD patients without ADHD) using Chi-square tests. We also investigated whether the ASRS-5 was normally distributed using indices of symmetry and skewness along with significance tests.

Then, we performed five sets of analyses:

*ASRS-5 in the whole sample.* We first used receiver operating characteristic (ROC) curves to test whether the ASRS-5 was a good indicator of ADHD, using ADHD patients without BD/BPD ( $n=236$ ) and the healthy control group ( $n=248$ ). The ROC curve is a probability curve of the diagnostic ability of a binary classifier.

*Comparisons between subgroups.* Then, we tested whether the ASRS-5 performed equally for different subgroups of the sample. We computed separate ROC curves for (1) ADHD severity and (2) ADHD presentations. The number of participants in each subgroup is reported in **Table 1**.

*Sensitivity analysis.* As psychotropic treatments (including stimulants) may affect ADHD symptoms, we ran a sensitivity analysis excluding participants who were on psychotropic medication in the non-BD/BPD ADHD group ( $n=60$ ). We also computed separated models for participants who had the DIVA 2.0 and the ACE+.

*ASRS-5 in BPD/BD patients with or without ADHD.* Also using ROC curves, we also tested whether the ASRS-5 was a good indicator of ADHD among participants with comorbid BPD and/or BD. We compared BPD/BD patients with ADHD ( $n=36$ ) with the group of BPD/BD without ADHD ( $n=37$ ).

For these five sets of analyses, we derived the area under the curve (AUC, which tells how much the model discriminate between those with and without the disease), sensitivity (true positive rate) and specificity (true negative rate). For AUC, the following criteria were used: 0.5: no discrimination, 0.7-0.8: acceptable, 0.8-0.9: excellent, and  $>0.9$ : outstanding

(Mandrekar 2010). For sensitivity and specificity, values close to 100% are ideal (van Stralen et al. 2009). The Youden's  $J$  statistic allows selecting the optimum cut-off point. It gives an equal weight to sensitivity and specificity, was used to select the best cut-off score:  $\max(\text{sensitivity} + \text{specificity} - 1)$ . It

Analyses were performed with Stata 15. The sample size calculation was performed with easyROC version 1.3.1 (<http://www.biosoft.hacettepe.edu.tr/easyROC>).

## **Results**

### ***Description of the population***

The description of the sample is reported in **Table 1**. Overall, there was 55.7% of females and 44.3% of males. The mean age of the participants was  $37.9 \pm 14.7$ . A total of 70.7% of participants had a secondary level of education. Participants in the BPD/BD samples (with and without ADHD) were less likely to have a secondary level of education (32.0%). Among ADHD participants, 47.3% had the inattentive presentation and 52.7% the combined/hyperactive presentation (seven participants had the hyperactive presentation). A total of 104 patients (20.9%) were on one psychotropic medication or more.

In the sample without comorbid disorders, participants with ADHD were significantly younger ( $p < .001$ ) and had a lower level of education ( $p < .001$ ) than healthy volunteers. There was no significant difference for gender ( $p = .215$ ). In the sample with comorbid disorders, participants with and without ADHD were not significantly different (gender:  $p = .740$ , age:  $p = .336$ , and level of education:  $p = .162$ ).

### ***Description of the French ASRS-5***

The distributions of the ASRS-5 score are reported in the **Supplementary Figure 1** separately for the ADHD groups (with and without BPD/BD) and the control groups (healthy participants

and BPD/BD without ADHD). The ASRS-5 was approximately normally distributed in each group, without floor or ceiling effects. Scores are more often below median for the control groups and above median for ADHD groups. The mean score was  $16.1 \pm 3.4$  in the ADHD groups and  $8.2 \pm 4.0$  in the control groups. Detailed information for all groups (two ADHD groups and two control groups) are reported in **Table 2**.

#### ***ASRS-5 performances in patients with ADHD but without BPD/BD and healthy controls***

The performances of the ASRS-5 in predicting ADHD diagnosis are reported in **Table 3**. The AUC was outstanding (.945), with the best cut-off score of 13/24 leading to a sensitivity of 84.3% and a specificity of 91.9%.

#### ***ASRS-5 performances according to clinical characteristics***

All results obtained in ADHD patients without BPD/BD and healthy controls are summarized in **Table 4**.

*ADHD severity.* The performances of the ASRS-5 were separately assessed in each severity group. The AUCs obtained in the four groups were different, with the best performances obtained in the very high-severity group ( $AUC=.977$ , best cut-off score=13/24 with  $sensitivity=95.2\%$ ). In the low-severity group, the AUC was .844, with the best cut-off score of 10/24 leading to a sensitivity of 88.5% and a specificity of 86.0%. For the cut-off score of 13, the sensitivity was rather low (63.5%, false negative rate=36.5%). The moderate- and high-severity groups displayed good psychometric properties ( $AUC=.958$  and .951,  $sensitivity=89.8\%$  and 87.3% with a cut-off score of 13).

*ADHD presentations.* Using the cut-off score of 13/24, the screener performed very well for patients with the combined/hyperactive presentation ( $AUC=.970$ ,  $sensitivity=92.2\%$ ). The

performance was a little lower for patients with the inattentive presentation ( $AUC=.918$ ,  $sensitivity=75.7\%$ ). A cut-off score of 12 was better ( $sensitivity=82.6\%$ ,  $specificity=85.5\%$ ).

*Sensitivity analysis.* In the final analysis excluding participants who were on psychotropic medication, the AUC was outstanding (.943) and the cut-off score of 13 yielded acceptable psychometric performances ( $sensitivity=83.9\%$ ,  $specificity=91.9\%$ ).

Performances were similar for participants who completed the DIVA 2.0 with the best cut-off score being 13 ( $n=163$ ,  $AUC=.951$ ,  $sensitivity=86.0\%$ ) and the ACE+ ( $n=73$ ,  $AUC=.931$ ,  $sensitivity=80.8\%$ ).

### ***ASRS-5 performances in BPD/BD patients with or without ADHD***

In patients with BD/BPD, the AUC obtained was fair (.801). The best cut-off score was also 13/24, with a sensitivity of 91.7%, but a specificity of 54.1% leading to a high proportion of false positives (45.9%) (see **Table 5**).

## **Discussion**

The purpose of this study was to confirm the diagnostic accuracy of the Adult ADHD Self-Report Screening Scale for DSM-5 (ASRS-5) in a sample of French-speaking adults with ADHD and non-ADHD controls.

### ***Performance in patients with ADHD but without BPD/BD and healthy controls***

The French version of the ASRS-5 did not perform as well as expected in our sample of French-speaking participants. In the first English validation study, the ASRS-5 offered acceptable performances in identifying ADHD patients in two separate samples, with a sensitivity ranging from 91.4% to 91.9% and a specificity ranging from 74.0 to 96.0% with a cut-off score of 14/24 (Ustun et al. 2017). Applying the same threshold, the sensitivity was 78.0% with a

specificity of 94.0% in our study. However, this threshold was not the best one, and we identified a better cut-off score ( $\geq 13$ ) with 8.1% of false positives and 15.7% of false negatives. The high rate of misclassification for false negatives was in line with studies reporting that adults are likely to be under-diagnosed (Ginsberg et al. 2014), contrariwise to children (Layton et al. 2018). Consequently, adults with ADHD are likely to be under-treated.

### ***Performance according to clinical characteristics***

The ASRS-5 did not perform equally among different subgroups of the population. The threshold to define ADHD should be adapted to keep acceptable sensitivity and specificity. This was the case for ADHD patients with a less severe symptomatology ( $\geq 10$ ) and ADHD patients with the inattentive presentation ( $\geq 12$ ). Thus, researchers and clinicians should be aware that the ASRS-5 probably misses individuals with less severe ADHD symptoms. Indeed, the cut-off score of 13 yielded 36.5% false negatives. This result contradicts previous findings suggesting that the psychometric properties of the ASRS are similar in different groups (van de Glind et al. 2013). This is an important issue, as even less severe symptomatology can lead to significant impairment in life. A screening tool should be useful for the whole range of the disorder, and not the most severe cases. Future research should focus on accurate identification of mild ADHD.

The ASRS-5 performed better to identify participants having the combined/hyperactive presentations. This might be because three out six items of the ASRS-5 rely to hyperactive/impulsive symptoms (items #2, #3 and #4). Furthermore, patients with combined presentation had a more severe ADHD symptomatology, as they could have 18 symptoms (maximum=9 symptoms for the inattentive presentation).

### ***Performance among patients with BPD/BD***

Finally, the ASRS-5 performed worse when participants had disorders that share common symptoms with ADHD. Indeed, the screener failed to identify ADHD among patients with BPD or BD, two frequent comorbid psychiatric disorders in ADHD which share common features and overlapping symptoms (i.e., impulsivity, emotional lability/dysregulation, interpersonal deficits, risk-taking behaviors, and inner restlessness, Perroud et al. 2014; Weibel et al. 2018). In participants with BPD or BD, the percentage of false positives was very high (45.9%), meaning that these patients were often misclassified as having ADHD. Therefore, clinicians should keep in mind that the ASRS-5 is not a reliable screener to detect ADHD for patients with these comorbid disorders, in line with previous results for the DSM-IV screener (Perroud et al. 2014; van de Glind et al. 2013; Weibel et al. 2018).

### ***Limitations***

This study has some limitations. A first shortcoming was that the translation/back-translation process involved two persons. More translators should have been involved, as well as the original author for the English back-translation. Second, patients with ADHD were recruited in specialized ADHD outpatient clinics, a setting that might not be representative of the whole ADHD population, thus limiting the generalization of our findings. For example, a lower frequency of inattentive presentation is usually observed in epidemiological samples (Simon et al. 2009), with more patients being treated by stimulants and psychotropic medications in clinical samples. To take this limitation into account, we ran a sensitivity analysis excluding participants who were on psychotropic medication (including psychostimulants). Findings were similar as those reported in this study. In addition, participants were younger and had a lower level of education in the ADHD group without BD/BPD in comparison with healthy volunteers. Third, the sample size was rather small in some cells (subgroups of participants with comorbidities). As investigating the diagnostic accuracy among subgroups was an



exploratory aim, it was not included in the sample size calculation. Because of these differences and issues, our results should be interpreted cautiously. Fourth, we did not assess the performance of the ASRS-5 according to other frequent comorbid conditions in adults with ADHD (e.g., depression, anxiety, substance use or sleep disorders). The comorbid disorders might have inflated sensitivity because other symptoms might increase the severity of ADHD or mimic ADHD symptoms. Another shortcoming of the study was that, in clinical participants, the ADHD assessment was made using two different structured clinical interviews (ACE+ and DIVA 2.0). Even if they rely on the same symptoms, the wording might be slightly different (e.g., examples for each symptom). Furthermore, these two widely used diagnostic tools have not been cross validated. However, they yielded similar psychometric performances in our sensitivity analyses. Nonetheless, future studies should rely on a single DSM-5 diagnostic interview. Other limitations are related to the subgroup analyses. When we used the sum score of symptoms to provide an overview of the severity, we automatically considered that symptoms were equivalent (same weight and same severity), which might not be the case. In addition, when we compared the different presentations (inattentive vs. combined/hyperactive), we should keep in mind that the inattentive presentation might be less severe, as the maximum number of symptoms was 9 (18 for the combined presentation). Finally, even if none of the healthy participants had previously received a diagnosis of ADHD, it was not formally ruled out by the standard diagnostic procedure. Given the low rate of diagnosis of ADHD in adult in the general population (Ginsberg et al. 2014), we therefore cannot exclude that some of them met criteria for ADHD. It might have inflated the rate of false negatives, meaning that the estimates given in the study are conservative.

## ***Conclusion***

The ASRS-5 appears as an acceptable tool to screen for ADHD in clinical settings. Nevertheless, researchers and clinicians should be aware of its limitations, namely a lower ability to detect ADHD among patients with a mild symptomatology or having the inattentive presentation. In addition, further evidence is needed for patients with comorbid disorders, and especially disorders with overlapping symptoms. To date, this was the first ADHD screener adapted for the DSM-5 criteria in French and this study will provide an extension to the available screening tools for French clinicians and thus a better detection of ADHD.

### **Compliance with Ethical Standards:**

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent: Informed consent was obtained from all individual participants included in the study.

### **References**

- APA (2000). *Diagnostic and Statistical Manual of Mental Disorders*, fourth ed. Washington, DC: American Psychiatric Association.
- APA (2013). *Diagnostic and Statistical Manual of Mental Disorders*, fifth ed. Arlington, VA: American Psychiatric Publishing.
- Asherson, P., Buitelaar, J., Faraone, S. V., & Rohde, L. A. (2016). Adult attention-deficit hyperactivity disorder: key conceptual issues. *The Lancet Psychiatry*, 3(6), 568-578, doi:10.1016/S2215-0366(16)30032-3.

- Caye, A., Swanson, J., Thapar, A., Sibley, M., Arseneault, L., Hechtman, L., et al. (2016). Life span studies of ADHD—Conceptual challenges and predictors of persistence and outcome. *Current Psychiatry Reports*, 18(12), 111, doi:10.1007/s11920-016-0750-x.
- Dalsgaard, S., Østergaard, S. D., Leckman, J. F., Mortensen, P. B., & Pedersen, M. G. (2015). Mortality in children, adolescents, and adults with attention deficit hyperactivity disorder: a nationwide cohort study. *Lancet (London, England)*, 385(9983), 2190-2196, doi:10.1016/S0140-6736(14)61684-6.
- Franke, B., Michelini, G., Asherson, P., Banaschewski, T., Bilbow, A., Buitelaar, J. K., et al. (2018). Live fast, die young? A review on the developmental trajectories of ADHD across the lifespan. *European Neuropsychopharmacology*, 28(10), 1059-1088, doi:10.1016/j.euroneuro.2018.08.001.
- Ginsberg, Y., Quintero, J., Anand, E., Casillas, M., & Upadhyaya, H. P. (2014). Underdiagnosis of attention-deficit/hyperactivity disorder in adult patients: A review of the literature. *The primary care companion for CNS disorders*, 16(3), doi:10.4088/PCC.13r01600.
- Instanes, J. T., Klungsøyr, K., Halmøy, A., Fasmer, O. B., & Haavik, J. (2018). Adult ADHD and comorbid somatic disease: A systematic literature review. *Journal of Attention Disorders*, 22(3), 203-228, doi:10.1177/1087054716669589.
- Kessler, R. C., Adler, L., Ames, M., Demler, O., Faraone, S., Hiripi, E., et al. (2005). The World Health Organization Adult ADHD Self-Report Scale (ASRS): a short screening scale for use in the general population. *Psychological Medicine*, 35(2), 245-256.
- Kessler, R. C., Adler, L., Gruber, M. J., Sarawate, C. A., Spencer, T., & Van Brunt, D. L. (2007). Validity of the World Health Organization Adult ADHD Self-Report Scale (ASRS) Screener in a representative sample of health plan members. *International Journal of Methods in Psychiatric Research*, 16(2), 52-65, doi:10.1002/mpr.208.

- Kessler, R. C., Green, J. G., Adler, L. A., Barkley, R. A., Chatterji, S., Faraone, S. V., et al. (2010). Structure and diagnosis of adult attention-deficit/hyperactivity disorder: analysis of expanded symptom criteria from the Adult ADHD Clinical Diagnostic Scale. *Archives of General Psychiatry*, 67(11), 1168-1178, doi:10.1001/archgenpsychiatry.2010.146.
- Kooij, J. J. S., & Francken, M. H. (2010). DIVA 2.0 Diagnostic Interview for ADHD in adults. The Hague, The Netherlands: DIVA Foundation.
- Kooij, S. J. J., Bejerot, S., Blackwell, A., Caci, H., Casas-Brugué, M., Carpentier, P. J., et al. (2010). European consensus statement on diagnosis and treatment of adult ADHD: The European Network Adult ADHD. *BMC Psychiatry*, 10(1), 67, doi:10.1186/1471-244X-10-67.
- Layton, T. J., Barnett, M. L., Hicks, T. R., & Jena, A. B. (2018). Attention deficit-hyperactivity disorder and month of school enrollment. *New England Journal of Medicine*, 379(22), 2122-2130, doi:10.1056/NEJMoa1806828.
- Mandrekar, J. N. (2010). Receiver operating characteristic curve in diagnostic test assessment. *Journal of Thoracic Oncology*, 5(9), 1315-1316, doi:10.1097/JTO.0b013e3181ec173d.
- Matte, B., Rohde, L. A., Turner, J. B., Fisher, P. W., Shen, S., Bau, C. H. D., et al. (2015). Reliability and validity of proposed DSM-5 ADHD symptoms in a clinical sample of adults. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 27(3), 228-236, doi:10.1176/appi.neuropsych.13060137.
- Mohr-Jensen, C., & Steinhausen, H.-C. (2016). A meta-analysis and systematic review of the risks associated with childhood attention-deficit hyperactivity disorder on long-term outcome of arrests, convictions, and incarcerations. *Clinical Psychology Review*, 48(Supplement C), 32-42, doi:10.1016/j.cpr.2016.05.002.

- Perroud, N., Cordera, P., Zimmermann, J., Michalopoulos, G., Bancila, V., Prada, P., et al. (2014). Comorbidity between attention deficit hyperactivity disorder (ADHD) and bipolar disorder in a specialized mood disorders outpatient clinic. *Journal of Affective Disorders, 168*, 161-166, doi:10.1016/j.jad.2014.06.053.
- Philipp-Wiegmann, F., Rösler, M., Clasen, O., Zinnow, T., Retz-Junginger, P., & Retz, W. (2017). ADHD modulates the course of delinquency: a 15-year follow-up study of young incarcerated man. *European Archives of Psychiatry and Clinical Neuroscience, 1-9*, doi:10.1007/s00406-017-0816-8.
- Salazar, A. J., Romero, J. A., Bernal, O. A., Moreno, A. P., Velasco, S. C., & Díaz, X. A. (2016). Noninferiority and equivalence evaluation of clinical performance among computed radiography, film, and digitized film for telemammography services. [Research article]. *International Journal of Telemedicine and Applications*.
- Shaw, M., Hodgkins, P., Caci, H., Young, S., Kahle, J., Woods, A. G., et al. (2012). A systematic review and analysis of long-term outcomes in attention deficit hyperactivity disorder: effects of treatment and non-treatment. *BMC medicine, 10*, 99, doi:10.1186/1741-7015-10-99.
- Simon, V., Czobor, P., Bálint, S., Mészáros, A., & Bitter, I. (2009). Prevalence and correlates of adult attention-deficit hyperactivity disorder: meta-analysis. *The British Journal of Psychiatry: The Journal of Mental Science, 194*(3), 204-211, doi:10.1192/bjp.bp.107.048827.
- Sonnby, K., Skordas, K., Olofsdotter, S., Vadlin, S., Nilsson, K. W., & Ramklint, M. (2015). Validation of the World Health Organization Adult ADHD Self-Report Scale for adolescents. *Nordic Journal of Psychiatry, 69*(3), 216-223, doi:10.3109/08039488.2014.968203.

- Ustun, B., Adler, L. A., Rudin, C., Faraone, S. V., Spencer, T. J., Berglund, P., et al. (2017). The World Health Organization Adult Attention-Deficit/Hyperactivity Disorder Self-Report Screening Scale for DSM-5. *JAMA Psychiatry*, 74(5), 520-526, doi:10.1001/jamapsychiatry.2017.0298.
- van de Glind, G., van den Brink, W., Koeter, M. W. J., Carpentier, P.-J., van Emmerik-van Oortmerssen, K., Kaye, S., et al. (2013). Validity of the Adult ADHD Self-Report Scale (ASRS) as a screener for adult ADHD in treatment seeking substance use disorder patients. *Drug and Alcohol Dependence*, 132(3), 587-596, doi:10.1016/j.drugalcdep.2013.04.010.
- van Stralen, K. J., Stel, V. S., Reitsma, J. B., Dekker, F. W., Zoccali, C., & Jager, K. J. (2009). Diagnostic methods I: sensitivity, specificity, and other measures of accuracy. *Kidney International*, 75(12), 1257-1263, doi:10.1038/ki.2009.92.
- Weibel, S., Menard, O., Ionita, A., Boumendjel, M., Cabelguen, C., Kraemer, C., et al. (2020). Practical considerations for the evaluation and management of Attention Deficit Hyperactivity Disorder (ADHD) in adults. *L'Encéphale*, 46(1), 30-40, doi:10.1016/j.encep.2019.06.005.
- Weibel, S., Nicastro, R., Prada, P., Cole, P., Rüfenacht, E., Pham, E., et al. (2018). Screening for attention-deficit/hyperactivity disorder in borderline personality disorder. *Journal of Affective Disorders*, 226, 85-91, doi:10.1016/j.jad.2017.09.027.
- Young, S. (2016). ACE+: A diagnostic interview of ADHD in adults. Psychology Services Limited.
- Young, S., Moss, D., Sedgwick, O., Fridman, M., & Hodgkins, P. (2015). A meta-analysis of the prevalence of attention deficit hyperactivity disorder in incarcerated populations. *Psychological Medicine*, 45(2), 247-258, doi:10.1017/S0033291714000762.



**Table 1. Descriptive statistics of the sample**

	ADHD without BPD/BD	Healthy control group	BPD/BD with ADHD	BPD/BD without ADHD
<b>n</b>	236	248	36	37
<b>Socio-demographics</b>				
<b>Gender</b>				
Male	50.4 (115)	44.8 (111)	25.0 (9)	21.6 (8)
Female	49.6 (113)	55.2 (137)	75.0 (27)	78.4 (29)
Missing values	n=8	-	-	-
<b>Age</b>				
Emerging adults (18-29 years)	48.0 (109)	23.0 (57)	27.8 (10)	24.3 (9)
Adults (30-49 years)	44.1 (100)	39.5 (98)	61.1 (22)	51.4 (19)
Ageing adults (50+ years)	7.9 (18)	37.5 (93)	11.1 (4)	24.3 (9)
Missing values	n=9	-	-	-
<b>Level of education</b>				
Secondary or lower	36.8 (75)	15.3 (38)	75.9 (22)	57.1 (12)
Tertiary	63.2 (129)	84.7 (210)	24.1 (7)	42.9 (9)
Missing values	n=32	-	n=7	n=16
<b>Clinical variables</b>				
<b>Any psychotropic medication</b>				
No	70.4 (143)	100 (248)	50.0 (17)	27.0 (10)
Yes	29.6 (60)	0 (0)	50.0 (17)	73.0 (27)
Missing values	n=33	-	-	-
<b>ADHD severity</b>				
Low (5-9 symptoms)	23.0 (52)	-	8.6 (3)	-
Moderate (10-11 symptoms)	21.7 (49)	-	11.4 (4)	-
High (12-14 symptoms)	27.9 (63)	-	34.3 (12)	-
Very high (15-18 symptoms)	27.4 (62)	-	45.7 (16)	-
Missing values	n=10	-	n=1	-
<b>ADHD presentation</b>				
Inattentive	50.4 (115)	-	27.8 (10)	-
Combined/hyperactive	49.6 (113)	-	72.2 (26)	-
Missing values	n=8	-	-	-
<b>Comorbid disorder</b>				
Bipolar disorder	-	-	44.4 (16)	70.3 (26)
Borderline personality disorder	-	-	55.6 (20)	29.7 (11)

Percentages and n (shown in brackets) are given.

ADHD: Attention Deficit hyperactivity disorder, BPD: borderline personality disorder, BD: bipolar disorder,

ASRS: Adult ADHD Self-Report Screening Scale.

<sup>1</sup> Education: 4 missing values.

<sup>2</sup> Education: 26 missing values.



**Table 2. Description of the ASRS-5**

	ADHD without BPD/BD	Healthy control group	BPD/BD with ADHD	BPD/BD without ADHD
<b>n</b>	236	248	36	37
<b>Items</b>				
1. Difficulties concentrating on what people say to you	2.88 (0.91)	1,37 (0.94)	2.78 (0.76)	1.84 (1.01)
2. Leave your seat in situations in which you are expected to remain seated	1.59 (1.29)	0.63 (0.97)	1.75 (1.11)	1.03 (0.87)
3. Difficulties unwinding and relaxing	2.79 (1.18)	1.47 (1.14)	3.31 (0.98)	2.35 (1.06)
4. Finish others' sentences	2.65 (1.19)	1.46 (1.08)	2.81 (1.09)	2.35 (1.21)
5. Put things off until the last minute	3.45 (0.83)	1.79 (1.12)	3.53 (0.69)	2.73 (1.31)
6. Depend on others to keep your life in order	2.58 (1.22)	0.86 (0.92)	2.69 (1.17)	2.73 (1.48)
<b>Total score</b>				
Sum-score	15.94 (3.47)	7.57 (3.58)	16.86 (2.84)	12.08 (4.62)
Skewness	-0.36 (p=.025)	0.46 (p=.004)	-0.29 (ns)	-0.05 (ns)
Kurtosis	2.98 (ns)	2.97 (ns)	2.75 (ns)	2.34 (ns)

ADHD: Attention Deficit hyperactivity disorder, BPD: borderline personality disorder, BD: bipolar disorder,

ASRS: Adult ADHD Self-Report Screening Scale.

**Table 3. Performance of the ASRS-5 compared to diagnosis of ADHD, general sample**

**(n=484)**

<b>Cut-off score</b>	<b>Sensitivity</b>	<b>Specificity</b>
Everybody	100	0.0
≥ 1	100	0.4
≥ 2	100	2.8
≥ 3	100	5.7
≥ 4	100	10.5
≥ 5	100	20.6
≥ 6	100	33.5
≥ 7	98.7	43.6
≥ 8	98.7	52.0
≥ 9	97.5	60.5
≥ 10	95.8	71.8
≥ 11	92.8	79.0
≥ 12	89.4	85.5
<b>≥ 13</b>	<b>84.3</b>	<b>91.9</b>
≥ 14	78.0	94.0
≥ 15	69.1	95.6
≥ 16	55.5	98.0
≥ 17	46.2	98.8
≥ 18	33.9	99.6
≥ 19	23.3	99.6
≥ 20	17.0	99.6
≥ 21	8.5	100
≥ 22	3.4	100
≥ 23	1.7	100
≥ 24	0.4	100
Nobody	0.0	100

ADHD: Attention deficit hyperactivity disorder, ASRS-5: Adult ADHD Self-Report Screening Scale according to the Diagnostic and Statistical Manual of Mental Disorders, fifth edition.

The best model is highlighted in bold.

**Table 4. Performances of the French version of the ASRS-5 according to demographic and clinical characteristics**

	Cut-off score	Sensitivity	Specificity	AUC
<b>Severity</b>				
Low severity (5-9 symptoms)	≥10	88.5%	71.8%	.884
Moderate severity (10-11 symptoms)	≥13	89.8%	91.9%	.958
High severity (12-14 symptoms)	≥13	87.3%	91.9%	.951
Very high severity (15-18 symptoms)	≥13	95.2%	91.9%	.977
<b>Clinical presentations</b>				
ADHD inattentive	≥12	82.6%	85.5%	.918
ADHD hyperactive/combined	≥13	92.9%	91.9%	.970

ADHD: Attention deficit hyperactivity disorder, ASRS-5: Adult ADHD Self-Report Screening Scale according to the Diagnostic and Statistical Manual of Mental Disorders, fifth edition, AUC: area under the curve.

Percentages and n under brackets are reported.

<sup>1</sup> Six participants had the hyperactive presentation.

**Table 5. Performance of the ASRS-5 compared to diagnosis of ADHD, clinical sample**


**(n=73)**

<b>Cut-off score</b>	<b>Sensitivity</b>	<b>Specificity</b>
Everybody	100	0.0
≥ 3	100	2.7
≥ 7	100	5.4
≥ 8	100	13.5
≥ 9	100	21.6
≥ 10	100	40.5
≥ 11	97.2	40.5
≥ 12	97.2	48.7
<b>≥ 13</b>	<b>91.7</b>	<b>54.1</b>
≥ 14	86.1	59.6
≥ 15	80.6	64.9
≥ 16	75.0	70.3
≥ 17	55.6	81.1
≥ 18	41.7	89.2
≥ 19	27.8	91.9
≥ 20	19.4	94.6
≥ 21	8.3	97.3
≥ 22	5.6	100
Nobody	0.0	100

ADHD: Attention deficit hyperactivity disorder, ASRS-5: Adult ADHD Self-Report Screening Scale according to the Diagnostic and Statistical Manual of Mental Disorders, fifth edition.

The best model is highlighted in bold.

**Supplementary Figure 1. Distributions of the French version of the ASRS-5 in the control and ADHD groups**



ADHD: Attention deficit hyperactivity disorder, ASRS-5: Adult ADHD Self-Report Screening Scale according to the Diagnostic and Statistical Manual of Mental Disorders, fifth edition.

**Supplementary Table S1. Adult ADHD Self-Report Screening Scale DSM-5, French version**

<p><i>Pour répondre, faites un 'X' dans la case qui correspond le plus à comment vous vous êtes senti ou conduit au cours des six derniers mois. Veuillez par la suite remettre cette liste, complétée, à votre professionnel de la santé lors de votre prochain rendez-vous pour discuter avec lui des résultats.</i></p>	Jamais	Rarement	Parfois	Souvent	Très souvent
1. À quelle fréquence avez-vous des difficultés à vous concentrer sur ce que les gens vous disent, même lorsqu'ils vous parlent directement ?					
2. À quelle fréquence vous levez-vous pendant des réunions ou d'autres situations dans lesquelles vous êtes censé rester assis ?					
3. À quelle fréquence avez-vous des difficultés à vous détendre et à vous relaxer pendant votre temps libre ?					
4. À quelle fréquence vous surprenez-vous terminant les phrases des autres dans une discussion avant qu'ils aient pu le faire eux-mêmes ?					
5. À quelle fréquence mettez-vous les choses de côté jusqu'à la dernière minute ?					
6. À quelle fréquence dépendez-vous des autres pour garder votre vie en ordre et s'occuper des détails ?					