



HAL
open science

Constructing discrete harmonic functions in wedges

Viet Hung Hoang, Kilian Raschel, Pierre Tarrago

► **To cite this version:**

Viet Hung Hoang, Kilian Raschel, Pierre Tarrago. Constructing discrete harmonic functions in wedges. 2020. hal-02987609v1

HAL Id: hal-02987609

<https://hal.science/hal-02987609v1>

Preprint submitted on 4 Nov 2020 (v1), last revised 18 Jan 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONSTRUCTING DISCRETE HARMONIC FUNCTIONS IN WEDGES

VIET HUNG HOANG, KILIAN RASCHEL, AND PIERRE TARRAGO

ABSTRACT. We propose a systematic construction of signed harmonic functions for discrete Laplacian operators with Dirichlet conditions in the quarter plane. In particular, we prove that the set of harmonic functions is an algebra generated by a single element, which conjecturally corresponds to the unique positive harmonic function.

CONTENTS

1. Introduction and main results	1
2. Study of the kernel	9
3. Boundary value problems for the generating functions	15
4. Proof of our main results (Theorems 1 and 2)	18
5. Various examples	25
References	35
Appendix A. Weierstrass' preparation and division theorems	38

1. INTRODUCTION AND MAIN RESULTS

Harmonic functions for the standard Laplacian. Let us first briefly recall some facts around the continuous Laplacian in \mathbb{R}^2

$$(1.1) \quad \Delta = \sigma_1 \frac{\partial^2}{\partial x^2} + 2\sigma_{1,2} \frac{\partial^2}{\partial x \partial y} + \sigma_2 \frac{\partial^2}{\partial y^2}.$$

Given a domain $D \subset \mathbb{R}^2$ with boundary ∂D , solving the Dirichlet problem on D for the operator Δ amounts to find the set $H(D)$ of functions $h : D \rightarrow \mathbb{R}$ which are harmonic on D , continuous on the closure \bar{D} of D , and zero on ∂D . This problem admits the unique solution $h = 0$ when D is bounded, but the situation is much richer when D is unbounded.

Assume first that Δ is the standard Laplacian (i.e., $\sigma_1 = \sigma_2 = 1$ and $\sigma_{1,2} = 0$ in (1.1)) and D is the upper half-space $\mathcal{H} = \{x + iy : y > 0\}$, and look at the associated Dirichlet problem. Defining $h(z) = -h(\bar{z})$ for z in the lower half-space, the classical Schwarz reflection principle implies that h can be extended to a harmonic function on \mathbb{C} . Hence h is solution to this Dirichlet problem if and only if there exists an analytic function

Date: November 4, 2020.

2010 Mathematics Subject Classification. Primary 31C35, 60G50; Secondary 60J45, 60J50, 31C20.

Key words and phrases. Discrete harmonic functions, conformal mappings.

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under the Grant Agreement No. 759702.

f on \mathbb{C} such that $h = \mathfrak{S}f$, and the Schwarz reflection implies that f is self-conjugate. The set $H(\mathcal{H})$ therefore admits the explicit description

$$\begin{aligned} H(\mathcal{H}) &= \{\mathfrak{S}f|_{\mathcal{H}} : f \text{ is analytic on } \mathbb{C} \text{ and } f(\bar{z}) = \overline{f(z)}\}, \\ &= \left\{ \sum_{n \geq 1} a_n \mathfrak{S}(x^n) : a_n \in \mathbb{R} \text{ and } |a_n|^{1/n} \rightarrow 0 \right\}. \end{aligned}$$

A similar description holds for other cones (not necessarily the half-space \mathcal{H}) and Laplacian operators (1.1) with general covariance matrix

$$(1.2) \quad \sigma = \begin{pmatrix} \sigma_1 & \sigma_{1,2} \\ \sigma_{1,2} & \sigma_2 \end{pmatrix}.$$

In particular, for a future use, when D is the positive quadrant \mathcal{Q} , we have

$$(1.3) \quad H(\mathcal{Q}) = \left\{ \sum_{n \geq 1} a_n h_n^\sigma : a_n \in \mathbb{R} \text{ and } |a_n|^{1/n} \rightarrow 0 \right\},$$

where the functions h_n^σ will be introduced later (see (4.1)) and σ denotes the covariance matrix (1.2).

A glimpse of our results. Our main objective in the present paper is to prove that in the discrete setting, a surprisingly simple equality of set holds, analogue to (1.3); see Theorem 2 for a precise statement.

A function $h : \mathbb{Z}^2 \rightarrow \mathbb{R}$ is discrete harmonic (or preharmonic) in a domain $D \subset \mathbb{Z}^2$ with respect to the discrete Laplacian operator

$$(1.4) \quad \Delta(h)(i, j) = \sum_{k, \ell} p_{k, \ell} h(i + k, j + \ell) - h(i, j),$$

the set of weights $\{p_{k, \ell}\}$ being fixed, if $\Delta(h)(i, j) = 0$ for all $(i, j) \in D$. In this article, we start from the analytic approach of [9] and propose a systematic construction of discrete harmonic functions in the quarter plane (i.e., $D = \mathbb{N}^2 = \{1, 2, 3, \dots\}^2$) which vanish on the boundary axes. We go beyond the existing literature, in the sense that our construction works:

- (i) for walks with arbitrary big (negative) jumps (see Figure 1),
- (ii) not only for positive, but also for signed discrete harmonic functions.

The two above features illustrate the robustness of our theory. The constructive aspect will follow from that we will obtain exact expressions for the generating functions of harmonic functions in terms of certain conformal mappings.

Construction of preharmonic functions. First of all, we would like to review some results in the literature dedicated to constructions of discrete harmonic functions. Let us first mention elementary constructions of preharmonic functions on \mathbb{Z}^d . Discrete polynomials and discrete exponential functions are constructed (mostly iteratively) in [21, 23, 27, 13]; see also [35] for a construction of preharmonic polynomials in terms of well-signed multinomials.

Further examples arise when preharmonic functions are defined on sets having certain rigid structures. Picardello and Woess [39] prove that discrete harmonic functions defined on Cartesian products of Markov chains have a product form. In the case of Weyl chambers of type A, Eichelsbacher and König [15] prove that preharmonic functions can be expressed

in terms of Vandermonde determinants. König and Schmid [29] demonstrate similar results for Weyl chambers of other types. Also in presence of a Weyl chamber structure, Biane, Bougerol and O’Connell [3] compute the (continuous) harmonic function (namely, the survival probability) with the help of the reflection principle.

Using the theory of analytic combinatorics in several variables [38], Courtiel et al. [10] study drifted simple random walks in various (two-dimensional) wedges and show that the harmonic functions obey to a rigid construction: namely, they are all obtained from a single function (again related to the reflection principle, see [10, Eq. (20)]), after some elementary operations (differentiation, division, evaluation). However, this technique seems to work only for random walks satisfying to (a certain algebraic version of) the reflection principle.

Roughly speaking, the Martin boundary theory aims at describing the set of all positive harmonic functions, given a Laplacian operator. See [30, 24, 26, 41, 42, 33] for examples when the Laplacian is related to random walks killed on the boundary of a half-plane or quadrant. This theory only rarely yields a construction of harmonic functions. In the previously cited articles, the computation of Martin boundary relies on an asymptotic study of quotients of Green functions. Let us underline, however, that Ignatiouk-Robert and Loree [24, 26] find an explicit formula for the exponential growth of all positive preharmonic functions in the Martin boundary.

In another direction, it is natural to ask whether discrete and classical harmonic functions are related, see for instance the papers [44, 45] by Varopoulos. In [11], Denisov and Wachtel prove that a certain natural preharmonic function (appearing as a prefactor in the persistence probability asymptotics) can be constructed by compensating the classical harmonic function, see [11, Eq. (5)].

Representation theory provides us with many examples of Markov processes for which explicit expressions of harmonic functions exist. For instance, preharmonic functions are expressed in terms of dimensions of irreducible representations in [2, 1]. Such results are quite powerful, but intrinsically limited to a few step sets.

We mention here a very last framework, which will be at the heart of the present work, and which relies on complex analysis techniques. As we shall see, the generating functions of preharmonic functions satisfy certain functional equations and, after further analysis, boundary value problems (BVPs). Solving these BVPs eventually yields exact expressions for the generating functions in terms of conformal mappings. The link between harmonic functions and conformal mappings is already illustrated in [42, 33], but we shall go much further here, by considering random walks with arbitrary big negative jumps (see item (i) above), and by constructing not only positive preharmonic functions (item (ii)); recall that in [42, 33], the attention is restricted to positive harmonic functions for small step walks.

Obviously, the list of constructive approaches to preharmonic functions could be continued, as the latter are all-present in probability theory and statistical physics, see the book [16] for recent illustrations.

Various analytic approaches. As we have seen above, some results on harmonic functions (structure of the Martin boundary, exact expressions, see [42, 33]) have already been obtained using the analytic approach of [19]. Let us now say a few words about this method. It was initially developed by Malyshev in [34], Fayolle and Iasnogorodski [18], to study the stationary distribution of random walks in the quarter plane reflected at the boundary. The stationary distribution generating functions are shown to satisfy BVPs.




FIGURE 1. Various step sets. From left to right: the simple random walk; the king walk (with jumps to all eight nearest neighbors); an example with big negative jumps; an example with big positive jumps; an arbitrary model.

From this, explicit expressions (typically, contour integral involving special functions) are deduced. See [19, Chap. 5] for full details.

Since then, the analytic approach of [19] has been applied to various contexts: queueing networks [31], potential theory [41, 42, 33], enumerative combinatorics (counting walks in the quarter plane) [6, 32, 12]. This approach is thus particularly fruitful and leads to precise results (both exact and asymptotic results).

However, the construction in [19] is restricted to the case of random walks with jumps to the eight nearest neighbors (see Figure 1). Its generalization to bigger jumps would require the precise understanding of the location of the branch points on a certain Riemann surface, see [20]. This is certainly possible on a few given examples, but the variety of possible behaviors makes us pessimistic to pursue in this direction to develop a general theory.

We will prefer here the alternative analytic approach of [9] by Cohen and Boxma (see also [8] by Cohen). The starting point is essentially the same (writing functional equations and BVPs for the generating functions) but remarkably, the construction can be done for arbitrary big (negative) jumps without increasing the level of complexity. The analytic approaches of [19] and [9] will be compared in Section 5.

Signed harmonic functions. In most of the literature cited above, the focus is put on positive harmonic functions, for clear probabilistic reasons, for example in relation with the concept of Doob transform, or because many probabilistic estimates use positive harmonic functions. In this paper, we go further and look at signed harmonic functions. Our motivation is fourfold: first, this will allow us to study the structure of the set of harmonic functions (for instance, we shall see that the vector space of harmonic functions having a bounded polynomial growth is finite-dimensional and will give a basis).

Our second motivation is that signed harmonic functions appear in various complete asymptotic expansions of relevant probabilistic or combinatorial quantities. To give a concrete example (related to this paper), let K be a given cone of \mathbb{R}^d and $\{Z(n)\}_{n \geq 0}$ be a zero-mean random walk (with some moment assumptions). Then it is shown in [11] that, as $n \rightarrow \infty$, the survival probability is asymptotically equivalent to

$$(1.5) \quad \mathbb{P}_x(\tau_K > n) \sim h(x)n^{-\alpha},$$

where $x \in K$ is the starting point, τ_K is the first exit time of $\{Z(n)\}_{n \geq 0}$ from the cone K , $h(x)$ is a harmonic function and α is a critical exponent. Assuming that (1.5) may be

refined as

$$(1.6) \quad \mathbb{P}_x(\tau_K > n) \sim \sum_i h_i(x) n^{-\alpha_i},$$

it is shown in [7] that the $h_i(x)$ may be constructed from certain polyharmonic functions as well as from signed harmonic functions.

Our third motivation comes from potential theory, where non-positive discrete harmonic functions turn out to be more difficult to study, compared to positive harmonic functions. Indeed, such classical tools as Harnack inequality (heavily used in [36, 43], for instance) do not hold anymore. Any construction of such functions becomes more relevant.

Finally, there might be some interesting features regarding the nodal domains of these non-positive harmonic functions. Indeed, within the connected components of the nodal lines, harmonic functions take a constant sign (by definition). Take an example: the function $h(i, j) = ij(i - j)(i + j)$ is discrete harmonic for the usual Laplacian in dimension 2 (probability $\frac{1}{4}$ to the four nearest neighbors):

$$(1.7) \quad \Delta(h)(i, j) = \frac{1}{4}(h(i + 1, j) + h(i, j - 1) + h(i - 1, j) + h(i, j + 1)) - h(i, j).$$

Its nodal lines are given by the two axes, the diagonal $\{i - j = 0\}$ and the anti-diagonal $\{i + j = 0\}$, see Figure 2. The same function h is also a positive harmonic function with Dirichlet boundary condition in the octant $\{(i, j) \in \mathbb{Z}^2 : 0 \leq i \leq j\}$. What would be the general structure of these nodal lines?




FIGURE 2. Nodal domains associated to the functions $ij(i - j)(i + j)$ and $ij(3i^4 - 10i^2j^2 + 3j^4 - 5i^2 - 5j^2 + 14)$, which are discrete harmonic for the operator (1.7). On the right picture, the blue cone is tangent to the nodal lines at infinity.

Preharmonic functions and their generating functions. Let $\{p_{k,\ell}\}_{(k,\ell) \in \mathbb{Z}^2}$ be non-negative weights (or transition probabilities) summing to 1, such that:

- (H1) The weights are symmetric, i.e., for all k and ℓ , $p_{k,\ell} = p_{\ell,k}$;
- (H2) Weights in the positive directions should be small, i.e., $p_{k,\ell} = 0$ if $k \geq 2$ or $\ell \geq 2$, while weights in the negative directions may be arbitrary large, see Figure 1;
- (H3) If $|\sum p_{k,\ell} x^k y^\ell| = 1$ and $|x| = |y| = 1$, then $x = y = 1$ (equivalently, the random walk on \mathbb{Z}^2 with increment distribution given by the $p_{k,\ell}$ is irreducible);
- (H4) The drift is zero, meaning that $\sum k p_{k,\ell} = \sum \ell p_{k,\ell} = 0$.

Consider now the associated discrete Laplacian operator (1.4), acting on complex-valued functions $h = \{h(i, j)\}_{(i,j) \in \mathbb{Z}^2}$. Our aim is to describe the functions h which satisfy to:

(H5) For all $(i, j) \in \mathbb{Z}^2$ with $i \leq 0$ and/or $j \leq 0$, $h(i, j) = 0$;

(H6) For all $i, j \geq 1$, $\Delta(h)(i, j) = 0$.

Such functions are harmonic for the random walk on \mathbb{Z}^2 killed when exiting the quadrant.

The generating function of the harmonic function h is

$$(1.8) \quad H(x, y) = \sum_{i, j \geq 1} h(i, j) x^{i-1} y^{j-1},$$

and its sections are

$$(1.9) \quad H(x, 0) = \sum_{i \geq 1} h(i, 1) x^{i-1} \quad \text{and} \quad H(0, y) = \sum_{j \geq 1} h(1, j) y^{j-1}.$$

Finally, the kernel is

$$(1.10) \quad K(x, y) = xy \left(1 - \sum p_{k, \ell} x^{-k} y^{-\ell} \right) = xy - \sum p_{k, \ell} x^{-k+1} y^{-\ell+1}.$$

The kernel is a polynomial due to hypothesis (H2) and is obviously fully characterized by the jumps $\{p_{k, \ell}\}$. The function $H(x, y)$ satisfies the functional equation (which simply reflects the harmonicity relations)

$$(1.11) \quad K(x, y)H(x, y) = K(x, 0)H(x, 0) + K(0, y)H(0, y) - K(0, 0)H(0, 0).$$

Main results. Equation (1.11) implies that any generating series $H(x, y)$ of a harmonic function has the form

$$(1.12) \quad H(x, y) = \frac{F(x) + G(y)}{K(x, y)}$$

for some power series $F, G \in \mathbb{C}[[t]]$. On the other hand, not any power series F and G are such that the right-hand side of (1.12) is a bivariate power series (indeed, in the case $p_{1,1} = 0$, notice that $K(0, 0) = 0$). From that point of view, we will answer the following questions:

- For which power series F and G is the function H in (1.12) a power series? (Remark that in the case $p_{1,1} \neq 0$, this is always the case, as $K(0, 0) \neq 0$.)
- In case the function H is a bivariate power series, is it analytic in a neighbourhood of $(0, 0)$? What is the associated radius of convergence?
- Which choice of $F, G \in \mathbb{C}(t)$ guarantees that the generating function H has positive coefficients?

In order to state our main result, we need to introduce a certain curve as well as two related conformal mappings. This step is the basis of our entire analysis and is inspired by the books [9, 8]. To do so, we first introduce the domain

$$(1.13) \quad \mathcal{K} = \{(x, y) \in \mathbb{C}^2 : K(x, y) = 0 \text{ and } |x| = |y| \leq 1\}.$$

As it turns out (more details are to come in Section 2), its projection along the first variable defines a curve \mathcal{S}_1 which is closed, non-intersecting, symmetric with respect to the real axis and contains 1; see Figure 3 for a few examples. The bounded (resp. unbounded) domain whose boundary is \mathcal{S}_1 is denoted by \mathcal{S}_1^+ (resp. \mathcal{S}_1^-). Let also \mathcal{H}^+ (resp. \mathcal{H}^-) denote the interior of the right (resp. left) half-plane. Let ψ_1 be a conformal mapping from




FIGURE 3. The curve \mathcal{S}_1 for the simple random walk (left). It is included in the parametric curve (2.6) with self-intersection (middle). On the right, the parametric curve for the (non-symmetric) walk with jumps $(-1, 1), (1, 0), (0, -1)$. The curve \mathcal{S}_1 is the intersection with the unit disk.

\mathcal{S}_1^+ to \mathcal{H}^+ with the conditions that $\psi_1(0) = p_{1,1}$ and $\psi_1'(0) > 0$. Finally, introduce for $n \geq 0$ the polynomial

$$(1.14) \quad P_n = (X^2 - p_{1,1}^2)^{\lfloor n/2 \rfloor} X^{n[2]},$$

where $n[2]$ stands for n modulo 2. The family $\{P_n\}_{n \geq 0}$ is a basis of $\mathbb{R}[X]$. The first few polynomials P_n are then:

$$P_0 = 1, \quad P_1 = X, \quad P_2 = X^2 - p_{1,1}^2, \quad P_3 = X^3 - p_{1,1}^2 X, \quad \text{etc.}$$

Theorem 1. For any $n \geq 1$, the function $\frac{P_n(\psi_1(x)) - P_n(-\psi_1(y))}{K(x,y)}$ defines a bivariate power series

$$(1.15) \quad H_n(x, y) = \sum_{i, j \geq 1} h_n(i, j) x^{i-1} y^{j-1} = \frac{P_n(\psi_1(x)) - P_n(-\psi_1(y))}{K(x, y)},$$

which satisfies the following properties:

- (1) Its Taylor coefficients $\{h_n(i, j)\}_{i, j \geq 1}$ form a discrete harmonic function for the Laplacian operator (1.4).
- (2) H_n is analytic for $x, y \in \mathcal{S}_1^+$.
- (3) Denote by h_n^σ the continuous harmonic function as in (1.3) or (4.1), with σ being the covariance matrix (1.2) associated to the transition probabilities $p_{k,\ell}$, and set

$$(1.16) \quad \theta = \arccos \frac{-\sigma_{12}}{\sqrt{\sigma_1 \sigma_2}} = \arccos \frac{-\sum k\ell p_{k,\ell}}{\sqrt{\sum k^2 p_{k,\ell}} \sqrt{\sum \ell^2 p_{k,\ell}}}.$$

One has $\frac{1}{m^{n\pi/\theta+1}} h_n(\lfloor mx \rfloor, \lfloor my \rfloor) \rightarrow h_n^\sigma(x, y)$ as m goes to infinity, in the sense of their Laplace transforms.

Our second main result shows that the harmonic functions h_n introduced in Theorem 1 actually describe the whole space of discrete harmonic functions.

Theorem 2. *The space of discrete harmonic functions is isomorphic to the vector space of formal power series $\mathbb{R}_0[[t]]$ with vanishing constant term, through the isomorphism*

$$\Phi : \begin{cases} \mathbb{R}_0[[t]] & \longrightarrow H(\mathbb{N}^2) \\ \sum_{n \geq 1} a_n t^n & \longmapsto \sum_{n \geq 1} a_n h_n. \end{cases}$$

Various examples illustrating Theorems 1 and 2 will be given in Section 5. Let us now describe the main features of these theorems:

- In the case $p_{1,1} = 0$, Theorems 1 and 2 imply that a function $\{h(i, j)\}_{i, j \geq 1}$ with generating function $H(x, y)$ is harmonic if and only if there exists a formal power series $F = \Phi^{-1}(h)$ such that

$$(1.17) \quad H(x, y) = \frac{F(\psi_1(x)) - F(-\psi_1(y))}{K(x, y)}.$$

In the case $p_{1,1} \neq 0$, the expression of H in terms of $\Phi^{-1}(h)$ is not so direct, and (1.17) should be replaced by (1.12) for some functions F and G depending on $\Phi^{-1}(h)$ in a non-trivial way.

In the framework of this article, we say that a function F characterizes a harmonic function h if its generating function can be written as (1.17).

- Part of the results of Theorem 2 is that for any formal power series F (even with a zero radius of convergence), the formula (1.17) defines formal power series (see our Lemmas 12 and 13) and eventually discrete harmonic functions.
- This correspondence between harmonic functions and formal power series may be refined as follows: writing $h(i, j) = \frac{h(i, j) + h(j, i)}{2} + \frac{h(i, j) - h(j, i)}{2}$, one may decompose a harmonic function as a sum of a symmetric harmonic function and another anti-symmetric harmonic function. Then in Theorem 2, symmetric (resp. anti-symmetric) harmonic functions correspond to even (resp. odd) power series $F(t) = \sum_{n \geq 1} a_n t^n$, see Section 4.4.
- In Equation (1.17), the only dependence in the model $\{p_{k, \ell}\}$ relies in the conformal mapping ψ_1 . The fact that we may take any power series F is independent of the model and therefore appears as a universal feature.
- It is known from [4, 25, 14] that there exists a unique function which is both positive and harmonic for the Laplacian operator (1.4). We conjecture (and bring some evidence) that this unique positive harmonic function corresponds to $F(t) = t$ in (1.17). In this sense, positivity of the harmonic function corresponds to minimality in terms of polynomial degree.

Conjecture 1. *The unique (up to multiplicative factors) positive harmonic function associated to the Laplacian operator (H1)–(H4) is h_1 in Theorem 1.*

- The polynomial growth of the unique positive harmonic function can be read off on the corner point of the curve \mathcal{S}_1 at 1 (see Figure 3), or equivalently on the singularity of the conformal mapping ψ_1 at 1.
- The vector space of harmonic functions with growth bounded by some constant g is finite-dimensional, and amounts to taking $F \in \mathbb{C}_n[t]$ for some n related to g .
- A much weaker statement of Theorem 1 (small step random walks and only $F(t) = t$) is given in [42]; see Section 5.2 for a more detailed comparison between our approach and the one in [42].

- The conformal mapping ψ_1 is uniquely characterized by the Riemann mapping theorem. We show how to obtain its explicit expression in the small step case, as well as in a few models with larger steps. However, in general, despite the particular structure of the set (1.13), there is little hope to derive concrete expressions for these conformal mappings.
- We will also explain how, given a (finite or infinite) number of boundary values, we may construct (and give a formula for) a harmonic function h having these values.
- Interestingly, by Theorem 2 one can transfer the algebra structure of $\mathbb{R}_0[[t]]$ into an algebra structure on the space of harmonic functions $H(\mathbb{N}^2)$. Formally, the multiplication is defined on $H(\mathbb{N}^2)$ by the formula

$$h \cdot h' = \Phi(\Phi^{-1}(h)\Phi^{-1}(h')), \quad \forall h, h' \in H(\mathbb{N}^2).$$

This implies in particular that $h_n \cdot h_m = h_{n+m}$, and that $H(\mathbb{N}^2)$ is generated as an algebra by the (conjecturally) unique positive harmonic function h_1 .

Acknowledgments. We would like to thank Éric Fusy for enlightening discussions concerning Section 5.1. We further thank Gerold Alsmeyer, Onno Boxma and François Chapon for interesting discussions.

2. STUDY OF THE KERNEL

The main objective of this section is the description of the set \mathcal{K} introduced in (1.13). To that purpose, in Section 2.1, we first recall from [9] some key properties of the two-variable polynomial K defined in (1.10). We also give a few elementary properties of the curve (1.13). In Section 2.2, we prove a new result on the existence of a corner point of the curve (1.13) and compute the associated angle. This angle turns out to be very important, as we will prove later that it is strongly related to the growth of harmonic functions, as stated in Theorem 1 (2).

Throughout the manuscript, \mathcal{C} will denote the unit circle and \mathcal{C}^+ (resp. \mathcal{C}^- , $\overline{\mathcal{C}^+}$) will stand for the interior (resp. exterior, resp. closed interior) domain, i.e., the open unit disk (resp. the complex plane deprived from the closed unit disk, resp. the closed unit disk). We shall also denote the bidisk by $\mathcal{U} = \mathcal{C}^+ \times \mathcal{C}^+ = \{(x, y) \in \mathbb{C}^2 : |x| < 1, |y| < 1\}$.

2.1. Preliminary results on the solutions of the kernel equation. To describe the domain \mathcal{K} in (1.13), we write

$$(2.1) \quad x = \eta s \quad \text{and} \quad y = \eta s^{-1},$$

with $\eta \in \mathbb{C}$ and $|s| = 1$. By (1.10), we have

$$(2.2) \quad K(\eta s, \eta s^{-1}) = \eta^2 - \sum_{k, \ell} p_{k, \ell} \eta^{-k-\ell+2} s^{-k+\ell}.$$

The lemma hereafter presents some properties of the roots of the kernel (2.2). Remark that in Lemma 1 below, and in many other statements as well, two cases will be considered separately, according to whether $p_{1,1} = 0$ or not.

Lemma 1. *Assume (H1)–(H4). For $|s| = 1$, the equation $K(\eta s, \eta s^{-1}) = 0$ admits exactly two solutions in $\overline{\mathcal{C}^+}$. Moreover, the following assertions hold:*

- (i) If $p_{1,1} = 0$, one solution is identically zero and the other one, denoted by $\eta(s)$, is real and varies in $[-1, 1]$. Furthermore, $\eta(1) = 1$, $\eta(-1) = -1$ and $|\eta(s)| < 1$ for all $|s| = 1$ with $s \neq \pm 1$.
- (ii) If $p_{1,1} \neq 0$, the two solutions, denoted by $\eta_1(s)$ and $\eta_2(s)$, are real and satisfy, for all $|s| = 1$, $\eta_2(s) = -\eta_1(-s)$. Further, $\eta_1(1) = 1$ and $\eta_1(s) \in (0, 1)$ for all $|s| = 1$ with $s \neq 1$.

Although the proof of Lemma 1 may be found in the book [9], we recall here the details, for the sake of completeness.

Proof. We first present the proof of (i). When $p_{1,1} = 0$, a term η may be factorised out of the kernel (2.2), so one of the roots is identically equal to zero. More precisely, we have $K(\eta s, \eta s^{-1}) = \eta \tilde{K}(\eta, s)$, where

$$(2.3) \quad \tilde{K}(\eta, s) = \eta - \sum p_{k,\ell} \eta^{-k-\ell+1} s^{-k+\ell}.$$

By Assumption (H3), one knows that if $(\eta s, \eta s^{-1})$ is a root of K with $|\eta| = |s| = 1$, then $\eta = s = 1$ or $\eta = s = -1$. Therefore, for all $|s| = 1$ with $s \neq \pm 1$, one has the following strict inequality:

$$(2.4) \quad \left| \sum p_{k,\ell} \eta^{-k-\ell+1} s^{-k+\ell} \right| < 1 = |\eta|.$$

It readily follows from Rouché's theorem that, as a function of η , $\tilde{K}(\eta, s)$ has a unique solution in \mathcal{C}^+ , denoted by $\eta(s)$. Since $\tilde{K}(1, s) \geq 0$ and $\tilde{K}(-1, s) \leq 0$, then $\eta(s) \in [-1, 1]$, hence the solution is real.

The previous argument does not work for $s = \pm 1$, because (2.4) becomes an equality at this point. We now consider $s = 1$. Fix $r \in (0, 1)$ and consider the polynomial

$$\tilde{K}_r(\eta) = \eta - r \sum p_{k,\ell} \eta^{-k-\ell+1}.$$

Rouché's theorem yields that \tilde{K}_r has a unique root $\eta_r(1) \in \mathcal{C}^+$. Further, Assumption (H4) ensures that \tilde{K}_r is increasing on $[0, 1]$, with $\tilde{K}_r(0) = -(p_{1,0} + p_{0,1}) < 0$ and $\tilde{K}_r(1) = 1 - r > 0$. Moreover, $\eta_r(1) \in (0, 1)$ and converges to 1 as $r \rightarrow 1$. By continuity arguments, one has $\eta(1) = 1$. At the point $s = -1$, notice that for all η and s , one has $\tilde{K}(-\eta, -s) = -\tilde{K}(\eta, s)$. This implies that $\eta(-1) = -1$.

We move to the proof of (ii). Fix $|s| = 1$ with $s \neq 1$. Since for all $|\eta| = 1$, one has

$$\left| \sum p_{k,\ell} \eta^{-k-\ell+2} s^{-k+\ell} \right| < 1 = |\eta^2|,$$

then by Rouché's theorem, $K(\eta s, \eta s^{-1})$ admits two solutions $\eta_1(s)$ and $\eta_2(s)$ in \mathcal{C}^+ . These quantities are real, as $K(\eta s, \eta s^{-1})$ is non-negative at 1 and -1 , but negative at 0. So one of them (say $\eta_1(s)$) is positive, and the other one ($\eta_2(s)$) is negative.

Looking now at the point $s = 1$, we set for $r \in (0, 1)$

$$K_r(\eta) = \eta^2 - r \sum p_{k,\ell} \eta^{-k-\ell+2}.$$

Rouché's theorem implies that K_r has two solutions in \mathcal{C}^+ , say $\eta_{1,r}(1)$ and $\eta_{2,r}(1)$. Notice that K_r is positive at ± 1 , negative at 0, so $\eta_{1,r}(1) \in (0, 1)$ and $\eta_{2,r}(1) \in (-1, 0)$.

Since K_r' is concave on $[0, 1]$, with $K_r'(0) = -r(p_{1,0} + p_{0,1}) \leq 0$ and $K_r'(1) = 2(1-r) > 0$, there exists $\eta_0 \in [0, 1)$ such that $K_r'(\eta_0) = 0$. Hence, K_r is decreasing on $[0, \eta_0]$, increasing

on $[\eta_0, 1]$, and $\eta_{1,r}(1) \in (\eta_0, 1)$. As $r \rightarrow 1$, then $\eta_{1,r}(1) \rightarrow 1$, while $\eta_{2,r}(1)$ goes to a point in $(-1, 0)$. Hence, K_1 has two solutions in $\overline{\mathcal{C}^+}$, which are $\eta_1(1) = 1$ and $\eta_2(1) \in (-1, 0)$.

To conclude, we prove that for all $|s| = 1$, $\eta_2(s) = -\eta_1(-s)$. Equation $K(\eta s, \eta s^{-1}) = 0$ may be rewritten as

$$(2.5) \quad \left(1 - \sum_{k+\ell \leq 0} p_{k,\ell} \eta^{-k-\ell} s^{-k+\ell} \right) \eta^2 - (p_{1,0} s^{-1} + p_{0,1} s) \eta - p_{1,1} = 0.$$

Notice that if $\eta(s)$ is a solution of (2.5), then $-\eta(-s)$ is the other solution. Hence, our claim follows. \square

It is worth mentioning that in the case $p_{1,1} \neq 0$, the domain (1.13) does not depend on the choice of $\eta_1(s)$ and $\eta_2(s)$. Hereafter, we shall write $\eta(s)$ for $\eta_1(s)$ in the case $p_{1,1} \neq 0$. Based on Lemma 1 and of the notation just introduced, we may now properly define the domain (1.13):

$$\mathcal{K} = \{(\eta(s)s, \eta(s)s^{-1}) : |s| = 1\} = \{(\eta(s)s, \overline{\eta(s)s}) : |s| = 1\},$$

with the bar denoting the complex conjugation. Introduce the curves

$$\mathcal{S}_1 = \{\eta(s)s : |s| = 1\} \quad \text{and} \quad \mathcal{S}_2 = \{\eta(s)s^{-1} : |s| = 1\},$$

which are obtained by taking the projection of \mathcal{K} along the first and second coordinates, respectively. See Figure 3 for a few examples. Obviously, an equivalent description of \mathcal{S}_1 (as a parametrized curve) is

$$\mathcal{S}_1 = \{(\eta(e^{it}) \cos t, \eta(e^{it}) \sin t) : t \in [0, 2\pi)\},$$

where we recall that $\eta(e^{it})$ is real, and that the interval $[0, 2\pi)$ may be replaced by $[0, \pi)$ in the case $p_{1,1} = 0$. For example, in the case of the simple random walk, one will have

$$(2.6) \quad \mathcal{S}_1 = \{(1 - \sin t, (1 - \sin t) \tan t) : t \in [0, \pi)\},$$

see Figures 3 and 4. We also emphasize that if the support of the $p_{k,\ell}$ is bounded, then the function $\eta(s)$ is an algebraic function of s .

Lemma 2. *Assume (H1)–(H4). The following assertions hold:*

- (i) *If $p_{1,1} = 0$, then $0 \in \mathcal{S}_1$ and for all $|s| = 1$, $\eta(-\bar{s}) = -\eta(s)$. The maps $s \mapsto \eta(s)s$ and $s \mapsto \eta(s)s^{-1}$ are two-to-one from \mathcal{C} to \mathcal{S}_1 and \mathcal{S}_2 , respectively.*
- (ii) *If $p_{1,1} \neq 0$, then $0 \in \mathcal{S}_1^+$ and for all $|s| = 1$, $\eta(s) > 0$ and $\eta(\bar{s}) = \eta(s)$. The maps $s \mapsto \eta(s)s$ and $s \mapsto \eta(s)s^{-1}$ are one-to-one from \mathcal{C} to \mathcal{S}_1 and \mathcal{S}_2 , respectively.*
- (iii) *The curves \mathcal{S}_1 and \mathcal{S}_2 are equal, non-self-intersecting and symmetric with respect to the real axis. If s traverses \mathcal{C} counterclockwise, then $\eta(s)s$ traverses \mathcal{S}_1 counterclockwise and $\eta(s)s^{-1}$ traverses \mathcal{S}_2 clockwise.*

Proof. Item (iii) is a direct consequence of (i) and (ii), and we start with the proof of (i). By Assumption (H1) and our notation (2.3), we have for $s = e^{i\lambda}$:

$$\tilde{K}(\eta, e^{i\lambda}) = \eta - \sum p_{k,\ell} \eta^{-k-\ell+1} \cos((-k+\ell)\lambda).$$

It is seen that if $\eta = \eta(e^{i\lambda})$ is a root of the above polynomial, then so is $-\eta(-e^{-i\lambda})$. By uniqueness of the solution, we must have $\eta(s) = -\eta(-\bar{s})$.




FIGURE 4. The curve \mathcal{S}_1 for the simple walk (left) and for the king walk (right), and definition of the angle θ . For the two models above, $\theta = \frac{\pi}{4}$.

Moving to the proof of (ii), we first have

$$(2.7) \quad K(\eta e^{i\lambda}, \eta e^{-i\lambda}) = \eta^2 - \sum p_{k,\ell} \eta^{-k-\ell+2} \cos((-k+\ell)\lambda),$$

see (2.2). It is seen that if $\eta(e^{i\lambda})$ is a solution of (2.7), then so is $\eta(e^{-i\lambda})$. Hence, $\eta(s) = \eta(\bar{s})$, being both positive (see Lemma 1 and its proof). \square

2.2. Corner point of the curve \mathcal{S}_1 at 1. Since it will be crucial in the next sections, we need to study the precise shape of the curve at the point 1. It should be noted that this single point 1 contains all the information about the growth of harmonic functions.

Throughout the paper, we write $V[K]$ for the set of zeros of K in \mathbb{C}^2 . We also recall here that a point of $V[K]$ is singular if and only if $\frac{\partial K}{\partial x}$ and $\frac{\partial K}{\partial y}$ simultaneously vanish at that point.

Lemma 3. *Assume (H1)–(H4). The curve \mathcal{S}_1 admits a corner point at 1 with interior angle θ given by (1.16), and is smooth elsewhere. Moreover, $\mathcal{K} \setminus \{(1, 1)\}$ consists of non-singular points of $V[K]$.*

In other words, the value of the angle at the corner point is given by the arccosine of the opposite of the correlation coefficient. See Figure 4 for an illustration of Lemma 3.

Proof. We start with the case $p_{1,1} = 0$ and first prove that $\mathcal{K} \setminus \{(1, 1)\}$ consists of smooth points of $V[K]$. Suppose that $\tilde{K}(\eta_0, s_0) = 0$, with $|\eta_0| < 1$, $|s_0| = 1$ and $s_0 \neq \pm 1$. Then, by Weierstrass preparation theorem for analytic functions in several variables (whose statement is recalled in Appendix A, see also [22]), there exist $r \geq 1$, a neighbourhood V around (η_0, s_0) (whose projection along the second coordinate is denoted by $P_s(V)$), r analytic functions g_0, \dots, g_{r-1} on $P_s(V)$ vanishing only at s_0 and a non-vanishing analytic function h on V such that

$$\tilde{K}(\eta, s) = h(\eta, s) \left((\eta - \eta_0)^r + g_{r-1}(s)(\eta - \eta_0)^{r-1} + \dots + g_0(s) \right).$$

Hence, for all s close to but different from s_0 , there are r distinct solutions to the equation $\tilde{K}(\eta, s) = 0$. Since for $s \in \mathcal{C}$ in a neighbourhood of s_0 , there is a unique solution to the

equation $\tilde{K}(\eta, s) = 0$, we must have $r = 1$. In particular,

$$\frac{\partial \tilde{K}}{\partial \eta}(\eta_0, s_0) = h(\eta_0, s_0) \neq 0,$$

and for $\eta_0 \neq 0$,

$$\frac{\partial K}{\partial \eta}(\eta_0 s_0, \eta_0 s_0^{-1}) = \eta_0 \frac{\partial \tilde{K}}{\partial \eta}(\eta_0, s_0) \neq 0.$$

Since

$$\frac{\partial K}{\partial \eta} = \frac{\partial x}{\partial \eta} \frac{\partial K}{\partial x} + \frac{\partial y}{\partial \eta} \frac{\partial K}{\partial y} = s \frac{\partial K}{\partial x} + s^{-1} \frac{\partial K}{\partial y},$$

we must have either $\frac{\partial K}{\partial x}$ or $\frac{\partial K}{\partial y}$ non-zero on $\mathcal{K} \setminus \{(1, 1)\}$. For $\eta_0 = 0$, it is easily seen that $\frac{\partial K}{\partial x}(0, 0) \neq 0$. The claim then follows.

We move to the proof of the smoothness of the curve $\mathcal{S}_1 \setminus \{1\}$. Differentiating the identity $\tilde{K}(\eta(s), s) = 0$, see (2.3), we obtain

$$(2.8) \quad \eta'(s) \frac{\partial \tilde{K}}{\partial \eta}(\eta(s), s) - \frac{\partial \tilde{K}}{\partial s}(\eta(s), s) = 0.$$

It is seen that for $|s| = 1$ and $s \neq \pm 1$, $\eta'(s)$ exists and is finite. Now we set $s = e^{i\lambda}$, with $\lambda \in (0, \pi)$. Since both λ and $\eta(e^{i\lambda})$ are real, then so is $(\eta(e^{i\lambda}))'$. Moreover, $\eta(e^{i\lambda})$ and $(\eta(e^{i\lambda}))'$ are not simultaneously zero, which leads to

$$(\eta(e^{i\lambda})e^{i\lambda})' = e^{i\lambda}((\eta(e^{i\lambda}))' + i\eta(e^{i\lambda})) \neq 0.$$

In other words, $\mathcal{S}_1 \setminus \{1\}$ is smooth.

We now deal with the case (1, 1). We differentiate again (2.8) and evaluate the new equation at $s = 1$, which leads to

$$(2.9) \quad \left(\sum p_{k,\ell}(k+\ell)^2 \right) \eta'(1)^2 + \sum p_{k,\ell}(k-\ell)^2 = 0.$$

Since the equation (2.9) has two distinct roots, then $\eta(s)$ is semi-differentiable at $s = 1$. Further, the roots of (2.9) represent the left and right derivatives of η at 1. Let $\partial_+ \eta(1)$ denote the right derivative of η at 1, i.e.,

$$\partial_+ \eta(1) = \lim_{\lambda \rightarrow 0^+} \frac{\eta(e^{i\lambda}) - \eta(1)}{e^{i\lambda} - 1}.$$

From Lemma 2, we know that $\eta(e^{i\lambda}) \rightarrow 1^-$ as $\lambda \rightarrow 0^+$. Hence, by (2.9), one has

$$\partial_+ \eta(1) = i \sqrt{\frac{\sum p_{k,\ell}(k-\ell)^2}{\sum p_{k,\ell}(k+\ell)^2}}.$$

We then have

$$(\partial_+ \eta(s)s)|_{s=1} = \partial_+ \eta(1) + \eta(1) = 1 + i \sqrt{\frac{\sum p_{k,\ell}(k-\ell)^2}{\sum p_{k,\ell}(k+\ell)^2}}.$$

This allows us to derive the interior angle θ of \mathcal{S}_1 at 1:

$$\cos \theta = 2 \sin^2((\arg \partial_+ \eta(s)s)|_{s=1}) - 1 = \frac{-\sum k\ell p_{k,\ell}}{\sqrt{\sum k^2 p_{k,\ell}} \sqrt{\sum \ell^2 p_{k,\ell}}}.$$

This completes the proof in the case $p_{1,1} = 0$.

In the case $p_{1,1} \neq 0$, one may apply the same arguments directly on K (instead of \tilde{K}) and derive the smoothness of the curves. Differentiating the identity $K(\eta(s)s, \eta(s)s^{-1}) = 0$, we obtain

$$\begin{aligned} \eta'(s) \left(2\eta(s) - \sum p_{k,\ell}(-k - \ell + 2)\eta(s)^{-k-\ell+1} s^{-k+\ell} \right) \\ - \left(\sum p_{k,\ell}(-k + \ell)\eta(s)^{-k-\ell+2} s^{-k+\ell-1} \right) = 0. \end{aligned}$$

Equation (2.9) and the rest of the proof follow in a similar way. \square

2.3. Non-existence of solutions to the kernel equation in $\mathcal{S}_1^+ \times \mathcal{S}_1^+$. In this part, we show that $K(x, y)$ does not have any zero in the domain $\mathcal{S}_1^+ \times \mathcal{S}_1^+$. This will allow us to deduce some regularity properties of $H(x, y)$ in the same domain, which will be used importantly in the next sections.

Lemma 4. *Assume (H1)–(H4). If $(x, y) \in V[K]$ are such that $|y| < |x| < 1$ (resp. $|x| < |y| < 1$), then $x \notin \mathcal{S}_1^+$ (resp. $y \notin \mathcal{S}_1^+$). As a consequence, $K(x, y)$ does not have any solution in $\mathcal{S}_1^+ \times \mathcal{S}_1^+$.*

Proof. We first consider the case $p_{1,1} \neq 0$ and set

$$\mathcal{V}_x = \{(x, y) \in V[K] : |y| < |x| < 1\}.$$

Reasoning by contradiction, let us suppose that there exists $(x, y) \in \mathcal{V}_x$ with $x \in \mathcal{S}_1^+$. Let γ be a path from x to 0 which avoids \mathcal{S}_1 and any point of $V[K]$ such that $\frac{\partial K}{\partial x}$ vanishes (there are only finitely many such points). Then, there exists a path $\gamma' : [0, 1] \rightarrow V[K]$ such that $\gamma'(0) = (x, y)$ and $P_x \circ \gamma' = \gamma$. Hence, since $p_{1,1} \neq 0$, $\gamma'(1) = (0, y')$ for some $y' \in \mathbb{C} \setminus \{0\}$. Since $P_x \circ \gamma' = \gamma$ never meets S_1 , γ' never meets \mathcal{K} and thus $\gamma'([0, 1]) \subset \mathcal{V}_x$. We should thus have $|y'| < 0$, which is a contradiction.

We move to the case $p_{1,1} = 0$, for which $(0, 0) \in \mathcal{K}$. By Weierstrass preparation theorem (recalled here in Appendix A) and coefficient identifications, we have

$$K(x, y) = h(x, y)(y + f(x)),$$

for (x, y) in a neighbourhood V of $(0, 0)$, with

$$f(x) = x + \frac{1 + 2p_{2,0} - p_{0,1}}{p_{1,0}} x^2 + o(x^2).$$

Since $\frac{1+2p_{2,0}-p_{0,1}}{p_{1,0}} > 0$, for x small enough with $x > 0$, there is a unique $y \in \mathbb{C}$ such that $(x, y) \in V[K] \cap V$, and moreover for x small enough

$$|y| = |f(x)| > |x|.$$

Hence, suppose that $(x, y) \in \mathcal{V}_x$ with $x \in \mathcal{S}_1^+$, and let γ be a path from x to 0 in \mathcal{S}_1^+ avoiding any singular point of $V[K]$ and such that $\gamma(t)$ is real for t close to 1. Then, there exists a path $\gamma'(x)$ such that $P_x \circ \gamma' = \gamma$. By the previous reasoning, for t close to 1 we have $|P_y \circ \gamma'(t)| > |P_x \circ \gamma'(t)|$. Since γ' is continuous and for t close to 0 we have $|P_y \circ \gamma'(t)| < |P_x \circ \gamma'(t)|$, there exists $t \in [0, 1)$ such that $\gamma'(t) \in \mathcal{K}$. This contradicts the fact that $P_x \circ \gamma'$ avoids \mathcal{S}_1 . \square

3. BOUNDARY VALUE PROBLEMS FOR THE GENERATING FUNCTIONS

The main objective of this section is to prove Proposition 8, which states a BVP for the generating functions $H(x, 0)$ and $H(0, y)$ defined in (1.9). The polynomial solutions to this BVP will be analyzed in Corollary 10. We first need to introduce conformal mappings for the bounded domain delimited by \mathcal{S}_1 .

We begin with the following notation. Let \mathcal{A} denote a non-intersecting curve separating the complex plane into two domains, \mathcal{A}^+ and \mathcal{A}^- . Let also f be meromorphic in $\mathbb{C} \setminus \mathcal{A}$. Then for t on the curve \mathcal{A} , $f^+(t)$ will denote $\lim_{x \rightarrow t, x \in \mathcal{A}^+} f(x)$, provided it exists. The notation $f^-(t)$ is defined similarly.

3.1. Conformal mappings.

Lemma 5. *For any $a \in \mathcal{S}_1^+ \cap \mathbb{R}$, there exists a unique conformal mapping π_1 from \mathcal{S}_1^+ onto \mathcal{C}^+ such that $\pi_1(a) = 0$ and $\pi_1'(a) > 0$. Moreover, $\pi_1^+(1) = 1$ and*

- (i) *In the case $p_{1,1} = 0$, $\pi_1^+(0) = -1$;*
- (ii) *In the case $p_{1,1} \neq 0$, $\pi_1(0) \in (-1, 1)$.*

Proof. The existence and uniqueness of π_1 is an immediate consequence of the classical Riemann mapping theorem.

We now prove that $\pi_1(x) = \overline{\pi_1(\bar{x})}$. Let $\chi(x)$ denote $\overline{\pi_1(\bar{x})}$ and a_0 denote the other intersection (not 1) of \mathcal{S}_1 and \mathbb{R} . Since $\pi_1(x)$ maps one-to-one from \mathcal{S}_1^+ onto \mathcal{C}^+ , which are both symmetric w.r.t. the real axis, then so is $\chi(x)$. Moreover, since $\chi(a) = \pi_1(a)$ and $\chi'(a) = \overline{\pi_1'(\bar{a})} > 0$ (since a is real and $\pi_1'(a) > 0$), then $\chi(x)$ maps conformally \mathcal{S}_1^+ onto \mathcal{C}^+ and $\chi = \pi_1$. This implies that $\pi(x) \in (-1, 1)$ for all $x \in (a_0, 1)$. Since π_1 is univalent and $\pi_1'(a) > 0$, then π_1 maps one-to-one from $(a_0, 1)$ onto $(-1, 1)$ in the same direction. Hence, $\pi_1^+(1) = 1$ and $\pi_1^+(a_0) = -1$. Since $0 = a_0$ in the case $p_{1,1} = 0$ and $0 \in (a_0, 1)$ in the case $p_{1,1} \neq 0$, then (i) and (ii) follow. \square

With π_1 as in Lemma 5, we introduce

$$(3.1) \quad \pi_2(z) = \frac{1}{\pi_1(z)},$$

which maps \mathcal{S}_1^+ conformally onto \mathcal{C}^- , the exterior of the unit disk.

Denote by \mathcal{H}^+ (resp. \mathcal{H}^-) the interior of the right (resp. left) half-plane and define the following mapping

$$(3.2) \quad \phi(z) = -\frac{z+1}{z-1}.$$

It is well known that ϕ maps conformally \mathcal{C}^+ (resp. \mathcal{C}^-) onto \mathcal{H}^+ (resp. \mathcal{H}^-). We finally introduce a few further conformal mappings:

- $\psi_1 = \phi \circ \pi_1 : \mathcal{S}_1^+ \rightarrow \mathcal{H}^+$;
- $\psi_2 = \phi \circ \pi_2 : \mathcal{S}_1^+ \rightarrow \mathcal{H}^-$;
- $\pi_{10}, \pi_{20}, \phi_0, \psi_{10}, \psi_{20}$ denote respectively the inverses of $\pi_1, \pi_2, \phi, \psi_1, \psi_2$.

The following lemma presents some crucial properties of ψ_{10} and ψ_{20} .

Lemma 6. *We have:*

- (i) $\psi_{10}(\infty) = 1$ and $\psi_{20}(\infty) = 1$;
- (ii) $\psi_{20}(t) = \psi_{10}(-t)$ for all $t \in \mathcal{H}^-$;

- (iii) $\psi_{10}(\bar{t}) = \overline{\psi_{10}(t)}$ for all $t \in \mathcal{H}^+$;
- (iv) $\psi_{20}(\bar{t}) = \overline{\psi_{20}(t)}$ for all $t \in \mathcal{H}^-$;
- (v) $\psi_{20}^-(t) = \psi_{10}^+(t)$ for all $t \in i\mathbb{R}$, i.e., $\psi_{10}^+ \times \psi_{20}^-(i\mathbb{R}) = \mathcal{H}$.

Proof. Item (i) follows by construction: $\psi_{10}(\infty) = \pi_{10} \circ \phi_0(\infty) = \pi_{10}(1) = 1$, see (3.2) and Lemma 5. Similar computations hold for $\psi_{20}(\infty)$.

In order to prove (ii), we show equivalently that for all $x \in \mathcal{S}_1^+$, $\psi_2(x) = -\psi_1(x)$. This comes from the construction of π_2 and ϕ . Indeed, for all $x \in \mathcal{S}_1^+$,

$$(3.3) \quad \psi_2(x) = \phi(\pi_2(x)) = \phi\left(\frac{1}{\pi_1(x)}\right) = -\phi(\pi_1(x)) = -\psi_1(x),$$

see (3.1) and (3.2).

We conclude by proving (v). We show equivalently that for all $x \in \mathcal{S}_1$, $\psi_2^+(x) = \psi_1^+(\bar{x})$. Notice that for $x \in \mathcal{S}_1$, since $\pi_2^+(x) = \frac{1}{\pi_1^+(x)} = \pi_1^+(\bar{x})$, then $\psi_2^+(x) = \psi_1^+(\bar{x})$. The proof is complete. \square

Lemma 7. *The following assertions hold true:*

- (i) *The asymptotic behavior of ψ_1 around 1 is*

$$\psi_1(x) \sim \frac{c}{(1-x)^{\pi/\theta}},$$

for some non-zero constant c and θ as in (1.16).

- (ii) *In the case $p_{1,1} = 0$, ψ_1 can be extended analytically around 0, such that $\psi_1(0) = 0$ and $\psi_1'(0) \neq 0$.*
- (iii) *In the case $p_{1,1} \neq 0$, $\psi_1(0) > 0$. Without loss of generality, we will assume that $\psi_1(0) = p_{1,1}$.*

Proof. We first prove (i) about the asymptotic behavior of ψ_1 around 1. Recall that π_{10} maps conformally \mathcal{C}^+ onto \mathcal{S}_1^+ and the interior angle of \mathcal{S}_1^+ at 1 is θ . Then by [40, Thm 3.11], there exists a non-zero constant c_1 such that as $z \rightarrow 1$,

$$\pi_{10}(z) = \pi_{10}(1) + (1-z)^{\theta/\pi}(c_1 + o(1)).$$

Hence as $x \rightarrow 1$, there exists $c \neq 0$ such that

$$\pi_1(x) = 1 + (1-x)^{\pi/\theta}(1/c_1 + o(1)) \quad \text{and} \quad \psi_1(x) = \phi \circ \pi_1(x) \sim \frac{c}{(1-x)^{\pi/\theta}}.$$

We now move to the proof of (ii). Carathéodory's theorem implies that π_1^+ maps one-to-one from \mathcal{S}_1 onto \mathcal{C} . Moreover, \mathcal{S}_1 is analytic around 0 (because the parametrization of \mathcal{S}_1 possesses derivatives of any order at 0). Hence, π_1 can be extended analytically around 0, see [37, p. 186], and by [40, Thm 3.9], $\pi_{10}'(0) \neq 0$.

We now prove (iii). Since $\pi_1(0) \in (-1, 1)$, then $\psi_1(0) > 0$. It is seen that if in Lemma 5 we choose two different points a_1 and a_2 , and consider the associated mappings ψ_1 and $\tilde{\psi}_{10}$, then there exists a constant $c > 0$ such that $\psi_1 = c\tilde{\psi}_1$. Indeed, since ϕ is a conformal mapping from \mathcal{C}^+ onto \mathcal{H}^+ , so is $c\phi$ for any $c > 0$. Hence, $c\tilde{\psi}_1$ is a conformal mapping from \mathcal{S}_1^+ onto \mathcal{H}^+ . One may choose $c = \frac{\psi_1(a_2)}{\tilde{\psi}_1(a_2)} = \psi_1(a_2) > 0$ and the proof is complete. \square

3.2. Boundary value problems. We now have collected enough material to construct a BVP, whose solutions relate to the generating functions $H(x, 0)$ and $H(0, y)$. For the sake of brevity, we shall denote $KH(x, 0) = K(x, 0) \times H(x, 0)$ and similarly $KH(0, y) = K(0, y) \times H(0, y)$. Define a function F on $\mathbb{C} \setminus i\mathbb{R}$ by

$$(3.4) \quad F(t) = \begin{cases} KH(\psi_{10}(t), 0) - \frac{KH(0,0)}{2} & \text{if } t \in \mathcal{H}^+, \\ -KH(0, \psi_{20}(t)) + \frac{KH(0,0)}{2} & \text{if } t \in \mathcal{H}^-, \end{cases}$$

and recall that the notation $F^\pm(t)$ has been introduced at the beginning of Section 3.

Proposition 8. *Assume (H1)–(H6), and assume in addition that the radius of convergence of $H(x, 0)$ and $H(0, y)$ are greater than or equal to one. Then F in (3.4) is sectionally analytic on $\mathbb{C} \setminus i\mathbb{R}$ and more specifically, F satisfies the following BVP:*

- (i) F is analytic on \mathcal{H}^+ and continuous on $\mathcal{H}^+ \cup i\mathbb{R}$.
- (ii) F is analytic on \mathcal{H}^- and continuous on $\mathcal{H}^- \cup i\mathbb{R}$.
- (iii) For all $t \in i\mathbb{R}$,

$$(3.5) \quad F^+(t) - F^-(t) = 0.$$

- (iv) If the associated harmonic function h is non-zero, then $F(\infty) = \infty$.

Proof. We start with the proof of (i). Since $H(x, 0)$ is assumed to be analytic in the unit disk, which contains \mathcal{S}_1^+ , then F is analytic on \mathcal{H}^+ . Moreover, with the exception of the point 1, the curve \mathcal{S}_1 is contained in the open unit disk, so the continuity on $\mathcal{H}^+ \cup i\mathbb{R}$ follows. Item (ii) would be proved along the same lines.

We now prove (iii). By Lemma 6 (v), we have $K(\psi_{10}^+(t), \psi_{20}^-(t)) = 0$ for all $t \in i\mathbb{R}$. So the identity (3.5) is just a consequence of the functional equation (1.11).

It remains to prove (iv). If F is bounded at infinity, then by Lemma 9 below, the function F should be constant, and actually even identically zero by the fact that if $x = y = 0$, then $KH(x, 0) = KH(0, y) = KH(0, 0)$. \square

Lemma 9. *Let F be a sectionally analytic on $\mathbb{C} \setminus i\mathbb{R}$, which satisfies (i), (ii) and (iii) of Proposition 8. If in addition F is bounded at infinity, then F is a constant function.*

Proof. Since F is sectionally analytic on $\mathbb{C} \setminus i\mathbb{R}$ and continuous on \mathbb{C} ($F^+(t) = F^-(t)$ on $i\mathbb{R}$ by (3.5)), then F is an entire function. Moreover, since F is bounded, then by Liouville's theorem, F is a constant function. \square

3.3. Polynomial solutions to the boundary value problems. As shown in Proposition 8 (iv), the solutions F to the BVP cannot be bounded at infinity for non-trivial harmonic functions. It is thus natural to look at functions F of (polynomial) order n at infinity, i.e., such that $F(t) = O(t^n)$, for some $n \geq 1$. Such functions will be called polynomial solutions and are studied in this section.

Corollary 10. *Given that F in (3.4) has a pole of order $n > 0$ at infinity, then F is a polynomial of degree n satisfying to the following conditions:*

$$(3.6) \quad F(p_{1,1}) = -F(-p_{1,1}) = \frac{KH(0,0)}{2} = -\frac{1}{2}p_{1,1}h(1, 1).$$

We then have, for $x, y \in \mathcal{S}_1^+$,

$$(3.7) \quad KH(x, 0) = F(\psi_1(x)) + \frac{KH(0, 0)}{2},$$

$$(3.8) \quad KH(0, y) = -F(-\psi_1(y)) + \frac{KH(0, 0)}{2},$$

$$(3.9) \quad H(x, y) = \frac{F(\psi_1(x)) - F(-\psi_1(y))}{K(x, y)}.$$

Proof. With a proof similar to that of Lemma 9, we derive that F is an entire function. Since F has a pole of order n at infinity, then by the extended version of Liouville's theorem, F is a polynomial.

The identity (3.6) is derived very naturally. First, in the case $p_{1,1} = 0$, since $\psi_{10}^+(0) = 0$ (see Lemma 7) and $KH(0, 0) = 0$, one must have $F(0) = KH(\psi_{10}^+(0), 0) = 0$. In the case $p_{1,1} \neq 0$, since $\psi_{10}(p_{1,1}) = \psi_{20}(-p_{1,1}) = 0$ (see again Lemma 7), one has

$$F(p_{1,1}) = -F(-p_{1,1}) = KH(\psi_{10}(p_{1,1}), 0) - \frac{KH(0, 0)}{2} = \frac{KH(0, 0)}{2}.$$

Applying into the solutions the inverses of ψ_{10} and ψ_{20} , which are ψ_1 and ψ_2 , we derive (3.7). Equation (3.9) is deduced from (3.7) together with the main functional equation (1.11). \square

4. PROOF OF OUR MAIN RESULTS (THEOREMS 1 AND 2)

This part is structured as follows: we successively prove Theorem 1 (1), Theorem 1 (2), Theorem 1 (3) and Theorem 2. Finally, in the independent Section 4.4, we study some features of symmetric and anti-symmetric harmonic functions.

4.1. Proof of Theorem 1 (1) and (2).

Proof. Let P_n be the family of polynomials introduced in (1.14) and H_n the associated bivariate function (1.15). Let us first prove that H_n defines a bivariate power series. This is obvious in the case $p_{1,1} \neq 0$, since $K(0, 0) \neq 0$. We therefore assume that $p_{1,1} = 0$. In this case $P_n(t) = t^n$, and one can rewrite (1.15) as

$$H_n(x, y) = \frac{\psi_1(x)^n - (-\psi_1(y))^n}{K(x, y)} = \frac{\psi_1(x) + \psi_1(y)}{K(x, y)} \sum_{k=0}^{n-1} \psi_1(x)^k (-\psi_1(y))^{n-1-k}.$$

By Lemma 7, ψ_1 is defined on a neighborhood of 0. Since $\psi_1(x) = -\psi_1(y)$ on a neighbourhood of $(0, 0)$ in the one-dimensional real variety \mathcal{K} in (1.13) (see Lemma 6 (v) and its proof), by analyticity of ψ_1 we have $\psi_1(x) = -\psi_1(y)$ on a neighbourhood of $(0, 0)$ in the one-dimensional complex variety where $K(x, y) = 0$.

Now recall that on a neighborhood of $(0, 0)$,

$$\frac{\partial}{\partial y}(\psi_1(x) + \psi_1(y)) \neq 0 \quad \text{and} \quad \frac{\partial}{\partial y}K(x, y) \neq 0.$$

Hence, by the Weierstrass preparation theorem for analytic functions in several variables (see Appendix A), there exist two functions $u(x, y)$ and $v(x, y)$ analytic in a neighborhood

of $(0, 0)$ and not vanishing at $(0, 0)$, as well as a function $g(x)$ analytic in a neighborhood of 0 and vanishing at 0, such that

$$\frac{\psi_1(x) + \psi_1(y)}{K(x, y)} = \frac{u(x, y)(y - g(x))}{v(x, y)(y - g(x))} = \frac{u(x, y)}{v(x, y)}.$$

This implies that $H_n(x, y)$ is analytic in a neighborhood of $(0, 0)$, i.e., $H_n(x, y)$ is the generating function of a function h_n on the quadrant. Since H_n satisfies the main functional equation (1.11), h_n is a harmonic function.

Since ψ_1 is defined on \mathcal{S}_1^+ and $K(x, y)$ does not have any solution in $\mathcal{S}_1^+ \times \mathcal{S}_1^+$ by Lemma 4, then $H_n(x, y)$ is analytic in $\mathcal{S}_1^+ \times \mathcal{S}_1^+$. The proof is complete. \square

4.2. Proof of Theorem 1 (3). Our main result in this part will be stated under Proposition 11 and appears as a refinement of Theorem 1 (3).

We first introduce some necessary notation. The Laplace transform of a discrete function f on \mathbb{N}^2 is defined as

$$\mathcal{L}f(x, y) = \sum_{u, v=0}^{\infty} f(u, v)e^{-(ux+vy)}.$$

For a measurable function f on the quadrant \mathcal{Q} , its Laplace transform is defined by

$$\mathcal{L}f(x, y) = \int_0^{\infty} \int_0^{\infty} f(u, v)e^{-(ux+vy)} dudv.$$

Remark that the above Laplace transforms are well defined (and analytic) on $\mathcal{H}^+ \times \mathcal{H}^+$ as soon as the growth of f at infinity is at most polynomial.

Finally, we introduce

$$(4.1) \quad h_n^{\sigma}(x, y) = \Im((x/\sin\theta + y \cot\theta + iy)^{n\pi/\theta}) = g_n(x/\sin\theta + y \cot\theta, y),$$

with $g_n(x, y) = \Im((x + iy)^{n\pi/\theta})$. It is easily checked that setting

$$(4.2) \quad \Delta = \frac{\partial^2}{\partial x^2} - 2 \cos\theta \frac{\partial^2}{\partial x \partial y} + \frac{\partial^2}{\partial y^2},$$

one has $\Delta(h_n^{\sigma}) = 0$. Notice that (4.2) is exactly (1.1) with $\sigma_1 = \sigma_2$ and $\theta = \arccos \frac{-\sigma_{12}}{\sqrt{\sigma_1 \sigma_2}}$, see (1.16).

Proposition 11. *Let h_n be the harmonic function with generating function H_n defined by (1.15) and h_n^{σ} be the continuous harmonic function defined in (4.1). Then there exists a positive constant c such that for all $x, y > 0$,*

$$\lim_{m \rightarrow \infty} \frac{c}{m^{n\pi/\theta+1}} \mathcal{L}h_n(mx, my) = \mathcal{L}h_n^{\sigma}(x, y).$$

Before embarking into the proof of Proposition 11, we provide a few remarks on the construction of the function h_n^{σ} as introduced in (4.1). We first consider a Dirichlet problem on the cone $D = \{(r \cos t, r \sin t) : r > 0, t \in (0, \theta)\}$. Recall from our introduction that the associated set of harmonic functions may be described as

$$H(D) = \left\{ \sum_{n \geq 1} a_n g_n(x, y) : a_n \in \mathbb{R} \text{ and } |a_n|^{1/n} \rightarrow 0 \right\},$$

with g_n as above. By the linear transformation $(x, y) \mapsto (x/\sin\theta + y \cot\theta, y)$ from the positive quadrant \mathcal{Q} onto the cone D , one can transform the above problem into a problem

on \mathcal{Q} with corresponding Laplacian (4.2). Naturally, the set of underlying continuous harmonic functions associated is described by (1.3).

Proof of Proposition 11. By definition of Laplace transforms and generating functions, one has

$$(4.3) \quad \begin{aligned} \mathcal{L}h_n(mx, my) &= \frac{e^{-(x+y)/m}}{m} H_n(e^{-x/m}, e^{-y/m}) \\ &= \frac{e^{-(x+y)/m}}{m} \frac{P(\psi_1(e^{-x/m})) - P(-\psi_1(e^{-y/m}))}{K(e^{-x/m}, e^{-y/m})}. \end{aligned}$$

Using now Lemma 7 (i), one deduces that as $m \rightarrow \infty$,

$$\begin{aligned} \psi_1(e^{-x/m}) &\sim \frac{c_0}{(1 - e^{-x/m})^{\pi/\theta}} \sim \frac{c_0}{(x/m)^{\pi/\theta}}, \\ P\left(\psi_1\left(e^{-\frac{x}{m}}\right)\right) &\sim c_1 \left(\left(\frac{m}{x}\right)^{\frac{2\pi}{\theta}} - p_{1,1} \right)^{\lfloor \frac{n}{2} \rfloor} \left(\frac{m}{x}\right)^{n[2]} \sim c_1 \left(\frac{m}{x}\right)^{\frac{n\pi}{\theta}}, \end{aligned}$$

where c_0 and c_1 are non-zero constants. On the other hand,

$$\begin{aligned} K(e^{-x/m}, e^{-y/m}) &= e^{-(x+y)/m} - \sum p_{k,\ell} e^{-(x(-k+1)+y(-\ell+1))/m} \\ &= -\frac{(x+y) \sum k p_{k,\ell}}{m} - \frac{x^2 \sum k^2 p_{k,\ell} + 2xy \sum k\ell p_{k,\ell} + y^2 \sum \ell^2 p_{k,\ell}}{2m^2} + o\left(\frac{1}{m^2}\right) \\ &= -\frac{\sum k^2 p_{k,\ell}}{2m^2} (x^2 - 2xy \cos \theta + y^2) + o\left(\frac{1}{m^2}\right). \end{aligned}$$

Going back to (4.3), we have for some non-zero constant c_2 that as $m \rightarrow \infty$,

$$\mathcal{L}h_n(mx, my) \sim c_2 m^{n\pi/\theta+1} \frac{(x^{-\pi/\theta})^n - (-y^{-\pi/\theta})^n}{x^2 - 2xy \cos \theta + y^2}.$$

We now move the Laplace transform of the continuous harmonic function h_n^σ :

$$\begin{aligned} \mathcal{L}h_n^\sigma(x, y) &= \int_0^\infty \int_0^\infty h_n^\sigma(u, v) e^{-(ux+vy)} du dv \\ &= \int_0^\infty \int_0^\infty g_n(u/\sin \theta + v \cot \theta, v) e^{-(ux+vy)} du dv \\ &= \int_0^\infty \int_{v' \cot \theta}^\infty g_n(u', v') e^{-(u' \sin \theta - v' \cos \theta)x - v'y} \sin \theta du' dv' \\ &= \sin \theta \int_0^\infty \int_0^\theta g_n(r \cos t, r \sin t) e^{-r(x \sin(\theta-t) + y \sin t)} r dt dr \\ &= \sin \theta \int_0^\theta \sin\left(n \frac{\pi}{\theta} t\right) \int_0^\infty r^{n \frac{\pi}{\theta} + 1} e^{-r(x \sin(\theta-t) + y \sin t)} dr dt \\ &= \Gamma(n\pi/\theta + 2) \sin \theta \int_0^\theta \frac{\sin(n \frac{\pi}{\theta} t)}{(x \sin(\theta-t) + y \sin t)^{n \frac{\pi}{\theta} + 2}} dt \\ &= \frac{\Gamma(n\pi/\theta + 2) \sin \theta}{n\pi/\theta + 1} \frac{-x \sin(\theta - (n \frac{\pi}{\theta} + 1)t) - y \sin((n \frac{\pi}{\theta} + 1)t)}{(x \sin(\theta-t) + y \sin t)^{n \frac{\pi}{\theta} + 1}} \Big|_0^\theta \end{aligned}$$




FIGURE 5. Illustrations of Lemmas 12 and 13. The harmonic function h_n is always zero on the axes. Left: in the case $p_{1,1} = 0$, the coefficients of h_n are zero at the blue points of the triangle, and non-zero just above (the red points). Right: in the case $p_{1,1} \neq 0$, the coefficients of h_n are zero at all blue points of a square, non-zero at the extremal red points.

$$= \frac{\Gamma(n\pi/\theta + 2)}{(n\pi/\theta + 1)(\sin \theta)^{n\pi/\theta - 1}} \frac{(x^{-\pi/\theta})^n - (-y^{-\pi/\theta})^n}{x^2 - 2xy \cos \theta + y^2}.$$

One can compare the above results and the proof is then complete. \square

Remark that one cannot deduce from Proposition 11 the asymptotics of $h_n(i, j)$ as n is fixed and $i + j \rightarrow \infty$. However, classically, Proposition 11 entails that h_n is converging locally in the L^1 -norm towards h_n^σ .

4.3. Proof of Theorem 2. As a first step, we need to prove that as n increases, the harmonic function h_n in (1.15) has more and more zero coefficients, in a sense which will be made precise in Lemma 12 (case $p_{1,1} = 0$) and Lemma 13 (case $p_{1,1} \neq 0$), see also Figure 5. Let us remark here that Lemma 13 is an a posteriori justification of our choice of the polynomials P_n in the definition (1.14).

Lemma 12. *Assuming $p_{1,1} = 0$, then h_n satisfies the following assertions:*

- For all $i, j \geq 1$ such that $i + j \leq n$, we have $h_n(i, j) = 0$;
- For all $i, j \geq 1$ such that $i + j = n + 1$, we have $h_n(i, j) \neq 0$.

Proof. As in (1.15), rewrite H_n as

$$\begin{aligned} H_n(x, y) &= \frac{\psi_1(x)^n - (-\psi_1(y))^n}{K(x, y)} = \frac{\psi_1(x) + \psi_1(y)}{K(x, y)} \sum_{k=0}^{n-1} \psi_1(x)^k (-\psi_1(y))^{n-1-k} \\ &= H_1(x, y) \sum_{k=0}^{n-1} \psi_1(x)^k (-\psi_1(y))^{n-1-k}. \end{aligned}$$

Recall from Lemma 7 that ψ_1 is analytic at 0, with $\psi_1(0) = 0$ and $\psi_1'(0) \neq 0$. Hence, we may write

$$\psi_1(x) = \sum_{i \geq 1} c_i x^i,$$

with $c_1 \neq 0$. We then obtain, using the bivariate expansion of H_1 , that

$$(4.4) \quad H_n(x, y) = \sum_{i, j \geq 1} h_1(i, j) x^{i-1} y^{j-1} \sum_{k=0}^{n-1} \left(\sum_{i \geq 1} c_i x^i \right)^k \left(- \sum_{j \geq 1} c_j y^j \right)^{n-1-k}.$$

By L'Hospital's rule, $\frac{\psi_1(x)}{K(x,0)} \rightarrow \frac{c_1}{p_{1,0}} \neq 0$. Hence, $h_1(1,1) \neq 0$ and we deduce from (4.4) the proof of Lemma 12. \square

Lemma 13. *If $p_{1,1} \neq 0$, then $h_1(1,1) \neq 0$ and for $k \geq 1$ and $\varepsilon \in \{0,1\}$, the harmonic function $h_{2k+\varepsilon}$ satisfies $h_{2k+\varepsilon}(i,j) = 0$ for all $1 \leq i, j \leq k$. Moreover, the following matrix is invertible:*

$$T_k = \begin{pmatrix} h_{2k}(k+1,1) & h_{2k+1}(k+1,1) \\ h_{2k}(1,k+1) & h_{2k+1}(1,k+1) \end{pmatrix}.$$

Proof. First, the proof that $h_1(1,1) \neq 0$ is similar to the end of the Lemma 12. Let $k \geq 1$ and $\varepsilon \in \{0,1\}$. Recall from Theorem 1 (1) that $h_{2k+\varepsilon}$ admits the generating function

$$H_{2k+\varepsilon}(x,y) = \frac{P_{2k+\varepsilon}(\psi_1(x)) - P_{2k+\varepsilon}(-\psi_1(y))}{K(x,y)},$$

with $P_{2k+\varepsilon}(t) = t^\varepsilon(t^2 - p_{1,1}^2)^k = t^\varepsilon(t^2 - \psi_1(0)^2)^k$, see (1.14). Hence we have

$$H_{2k+\varepsilon}(x,y) = \frac{1}{K(x,y)} \left(\psi_1(x)^\varepsilon (\psi_1(x) - \psi_1(0))^k (\psi_1(x) + \psi_1(0))^k - (-\psi_1(y))^\varepsilon (-\psi_1(y) - \psi_1(0))^k (-\psi_1(y) + \psi_1(0))^k \right).$$

Since $K(0,0) \neq 0$, $\frac{1}{K(x,y)}$ is analytic at $(0,0)$. Moreover, since $\frac{1}{K(x,y)}$ satisfies the functional equation (1.11), then it is also a generating function of a harmonic function $h_0(i,j)$, i.e.,

$$\frac{1}{K(x,y)} = \sum_{i,j \geq 1} h_0(i,j) x^{i-1} y^{j-1}.$$

Since ψ_1 is analytic at 0, which is in the interior of \mathcal{S}_1^+ , then it has an expansion

$$\psi_1(x) = \psi_1(0) + \sum_{i \geq 1} c_i x^i,$$

where $c_1 \neq 0$ since ψ_1 is a conformal map. Therefore,

$$H_{2k+\varepsilon}(x,y) = \left(\sum_{i,j \geq 1} h_0(i,j) x^{i-1} y^{j-1} \right) \times \left[\left(\sum_{i \geq 1} c_i x^i \right)^k \left(\psi_1(0) + \sum_{i \geq 1} c_i x^i \right)^\varepsilon \left(2\psi_1(0) + \sum_{i \geq 1} c_i x^i \right)^k - \left(-\sum_{j \geq 1} c_j y^j \right)^k \left(-\psi_1(0) - \sum_{j \geq 1} c_j y^j \right)^\varepsilon \left(-2\psi_1(0) - \sum_{j \geq 1} c_j y^j \right)^k \right].$$

Since $h_0(1,1) \neq 0$ and $c_1 \neq 0$, the bivariate power series expansion of the function above has zero coefficients for all monomials $x^i y^j$ with $i, j \leq k-1$, while the coefficient of the monomial x^k is

$$h_{2k+\varepsilon}(k,1) = (2c_1)^k h_0(1,1) \psi_1(0)^{k+\varepsilon} = (2c_1)^k h_0(1,1) p_{1,1}^{k+\varepsilon}$$

and that of y^k is

$$h_{2k+\varepsilon}(1, k) = (-1)^{\varepsilon-1} (2c_1)^k h_0(1, 1) p_{1,1}^{k+\varepsilon}.$$

Hence,

$$\begin{pmatrix} h_{2k}(k+1, 1) & h_{2k+1}(k+1, 1) \\ h_{2k}(1, k+1) & h_{2k+1}(1, k+1) \end{pmatrix} = (2c_1)^k h_0(1, 1) p_{1,1}^k \begin{pmatrix} 1 & p_{1,1} \\ -1 & p_{1,1} \end{pmatrix}.$$

The determinant of the latter matrix is $2^{k+1} c_1^k h_0(1, 1) p_{1,1}^{k+1} \neq 0$, which implies the second statement of Lemma 13. \square

Remark 1. As the proof of Lemma 13 shows, choosing instead of P_n in (1.14) the family of polynomials

$$P_{m,n} = (X - p_{1,1})^m (X + p_{1,1})^n$$

for $m, n \geq 1$, would yield harmonic functions vanishing on the rectangle $\{(i, j) : 1 \leq i \leq m, 1 \leq j \leq n\}$. In particular, there is no uniqueness in the choice of P_n .

Lemma 14. Given $p_{1,1} = 0$ and a sequence $\{c_n\}_{n \geq 1}$, there exists a unique sequence $\{a_n\}_{n \geq 1}$ such that the (harmonic) function $\sum_{n \geq 1} a_n h_n$ satisfies

$$(4.5) \quad \sum_{n \geq 1} a_n h_n(i, 1) = c_i, \quad \forall i \geq 1.$$

Moreover, $\{a_n\}_{n \geq 1}$ can be deduced from the (infinite) linear system of equations

$$M \cdot a = c,$$

where $a = (a_1, a_2, a_3, \dots)^\top$, $c = (c_1, c_2, c_3, \dots)^\top$ and M is a infinite non-singular lower triangular matrix.

By construction, the generating function of the harmonic function given in Lemma 14 is the function

$$H(x, y) = \frac{\sum_{n \geq 1} a_n \psi_1(x)^n - \sum_{n \geq 1} a_n (-\psi_1(y))^n}{K(x, y)} = \frac{F(\psi_1(x)) - F(-\psi_1(y))}{K(x, y)},$$

with $F(t) = \sum_{n \geq 1} a_n t^n$, in accordance with (1.17).

Proof. By Lemma 12, h_n vanishes on $(i, 1)$ for $1 \leq i \leq n-1$ and $h_n(n, 1) \neq 0$. Hence, the infinite matrix

$$L = (h_j(i, 1))_{1 \leq i, j \leq \infty}$$

is lower triangular, with non-zero diagonal coefficients. Hence, L is invertible, and for any vector $c = (c_1, c_2, c_3, \dots)^\top$, there exists a unique vector $a = (a_1, a_2, a_3, \dots)^\top$ such that

$$\begin{pmatrix} h_1(1, 1) & 0 & 0 & \dots & 0 & \dots \\ h_1(2, 1) & h_2(2, 1) & 0 & \dots & 0 & \dots \\ \vdots & \vdots & \ddots & \ddots & \ddots & \ddots \\ h_1(n, 1) & h_2(n, 1) & \dots & \dots & h_n(n, 1) & \dots \\ \vdots & \vdots & \ddots & \ddots & \ddots & \ddots \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \\ \vdots \end{pmatrix} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \\ \vdots \end{pmatrix}.$$

Hence, given $\{c_n\}_{n \geq 1}$, there exists a unique sequence $\{a_n\}_{n \geq 1}$ such that (4.5) holds. \square

Lemma 15. *Given $p_{1,1} \neq 0$ and two infinite sequences $\{c_n\}_{n \geq 1}$ and $\{d_n\}_{n \geq 2}$, there exists a unique sequence $\{a_n\}_{n \geq 1}$ such that (4.5) holds, as well as*

$$(4.6) \quad \sum_{n \geq 1} a_n h_n(1, i) = d_i, \quad \forall i \geq 2.$$

Moreover, $\{a_n\}_{n \geq 1}$ can be deduced from the linear system

$$M \cdot a = b,$$

where $a = (a_1, a_2, a_3, \dots)^\top$, $b = (c_1, c_2, d_2, c_3, d_3, \dots)^\top$ and M is an infinite block lower triangular matrix with invertible blocks of size 1 or 2 on the diagonal.

Proof. Let τ be the involution of $(\mathbb{N} \times \{1\}) \cup (\{1\} \times \mathbb{N})$ switching coordinates. The only fixed point of τ is $(1, 1)$. By Lemma 13, for fixed values of $k \geq 1$ and $\varepsilon \in \{0, 1\}$, $h_{2k+\varepsilon}$ vanishes at $(i, 1)$ and $(1, i)$ for all $i \leq k - 1$, and the matrix

$$T_k = (h_{2k+j-1}(\tau^{i-1}(k+1, 1)))_{1 \leq i, j \leq 2}$$

is invertible. Likewise, $h_1(1, 1) \neq 0$. Hence,

$$M = (h_j(\tau^{i[2]}([i/2] + 1, 1)))_{1 \leq i, j \leq \infty} = \begin{pmatrix} h_1(1, 1) & 0 & 0 & 0 & \dots \\ h_1(2, 1) & & & & \\ h_1(1, 2) & T_1 & 0 & \dots \\ \vdots & \vdots & T_2 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

with each T_i invertible. Thus M is invertible and for any vector $b = (c_1, c_2, d_2, c_3, d_3, \dots)^\top$, there exists a unique vector $a = (a_1, a_2, a_3, \dots)^\top$ such that $M \cdot a = b$. For such a vector a , we thus have conditions (4.5) and (4.6) satisfied. \square

Putting all the latter lemmas together yields the proof of Theorem 2, as follows:

Proof of Theorem 2. Let Φ be the function as in the statement of Theorem 2. First notice that by Lemma 12 when $p_{1,1} = 0$ and Lemma 13 for $p_{1,1} \neq 0$, the map Φ is well defined, since for any sequence $\{a_n\}_{n \geq 1}$ and all $i, j \geq 1$, $\sum_{n \geq 1} a_n h_n(i, j)$ is a finite sum and the harmonicity of $\sum a_n h_n$ is directly inherited by the harmonicity of each h_n .

Suppose first that $p_{1,1} > 0$. First, by Lemma 15, Φ is injective, and for any pair of formal power series F, G with $F(0) = G(0)$, there exists a sequence $\{a_n\}_{n \geq 1}$ such that

$$F(x) = \sum_{i \geq 1} \sum_{n \geq 1} a_n h_n(i, 1) x^{i-1} \quad \text{and} \quad G(y) = \sum_{j \geq 1} \sum_{n \geq 1} a_n h_n(1, j) y^{j-1}.$$

Since any harmonic function h is uniquely determined by its sectional generating functions

$$(4.7) \quad F(x) = \sum_{i \geq 1} h(i, 1) x^{i-1} \quad \text{and} \quad G(y) = \sum_{j \geq 1} h(1, j) y^{j-1}$$

through (1.11), this shows the surjectivity of Φ .

The proof is more involved in the case $p_{1,1} = 0$. By Lemma 14, Φ is injective. It remains to show that Φ is surjective. Let h be a harmonic function and set F as in (4.7). Then,

by Lemma 14, there exists $\{a_n\}_{n \geq 1}$ such that $\sum a_n h_n(i, 1) = h(i, 1)$ for all $i \geq 1$. Hence, the formal power series $F(x)$ and

$$\tilde{F}(x) = K(x, 0) \sum_{i \geq 1} \left(\sum_{n \geq 1} a_n h_n(i, 1) \right) x^{i-1}$$

coincide. Let H be the generating series of h and \tilde{H} the generating series of $\sum a_n h_n$. Then we have

$$H(x, y) = \frac{F(x) + G(y)}{K(x, y)} \quad \text{and} \quad \tilde{H}(x, y) = \frac{\tilde{F}(x) + \tilde{G}(y)}{K(x, y)} = \frac{F(x) + \tilde{G}(y)}{K(x, y)},$$

with $G, \tilde{G} \in \mathbb{R}[[t]]$. Since $K(0, 0) = 0$ and $\frac{\partial}{\partial x} K(0, 0) \neq 0$, by the Weierstrass preparation theorem (see Appendix A) applied to $K(x, y)$ with respect to x in the ring of formal power series $\mathbb{R}[[x, y]]$, one can write

$$K(x, y) = u(x, y)(x - v(y)),$$

with u invertible in $\mathbb{R}[[x, y]]$ and $v \in \mathbb{R}[[y]]$ satisfying $v(0) = 0$. In particular, $F(x) + G(y)$ and $F(x) + \tilde{G}(y)$ are both divisible by $(x - v(y))$ in $\mathbb{R}[[x, y]]$.

By the Weierstrass division theorem (see Appendix A) applied to the division of $F(x)$ by the Weierstrass polynomial of degree one $x - v(y)$ in $\mathbb{R}[[x, y]]$, there is a unique Weierstrass polynomial G of degree zero in $\mathbb{R}[[x, y]]$ (namely, $G \in \mathbb{R}[[y]]$ with $G(0) = 0$) such that $G(y) + F(x)$ is divisible by $x - v(y)$. Hence, $G(y) = \tilde{G}(y)$ and $H = \tilde{H}$. This shows the surjectivity of Φ . \square

4.4. Symmetric and anti-symmetric harmonic functions.

Proposition 16. *The harmonic function $h(i, j)$ is symmetric (resp. anti-symmetric) if and only if its characterizing series $F(t) = \Phi^{-1}(h)$ is odd (resp. even).*

Proof. It is seen that $h(i, j)$ is symmetric (resp. anti-symmetric) if and only if $H(x, 0) = H(0, x)$ (resp. $H(x, 0) = -H(0, x)$).

First, in the case where H is of type (3.7), the identity $H(x, 0) = H(0, x)$ is equivalent to $F(\psi_1(x)) = -F(-\psi_1(x))$. This implies that $h(i, j)$ is symmetric if and only if $F(t)$ is an odd function. The second statement in the anti-symmetric case would be proved similarly.

By (1.14), P_n is odd for n odd and even for n even, thus by the previous reasoning h_n is symmetric for n odd and anti-symmetric for n even. We deduce then from Theorem 2 that $\Phi(F)$ is symmetric if F is odd and anti-symmetric if F is even. Since the vector subspace of odd power series and the one of even power series are complementary in $\mathbb{R}_0[[t]]$, the result is deduced. \square

5. VARIOUS EXAMPLES

In this part, we provide a list of models for which one may compute the conformal map ψ_1 explicitly. We start with the example of simple random walks (Section 5.1). For this model, all generating functions happen to be rational functions, and most of the computations are rather easy to lead. We then move to arbitrary small step random walks (Section 5.2) and obtain an expression for the conformal mapping in terms of generalized Chebychev polynomials. Finally, we construct of family of walks with arbitrary large

jumps, in relation with plane bipolar orientations, for which one has a simple (algebraic) formula for the conformal mapping (Section 5.3).

5.1. The simple random walk. We present here a first example, which is the simple random walk with transition probabilities $p_{1,0} = p_{0,1} = p_{-1,0} = p_{0,-1} = 1/4$, see Figure 1 and (1.7) for the associated Laplacian operator. The kernel takes the form

$$K(x, y) = xy \left(1 - \frac{x + y + x^{-1} + y^{-1}}{4} \right) = -\frac{x(y-1)^2}{4} - \frac{y(x-1)^2}{4}.$$

Curve and conformal mappings. Setting $(x, y) = (\eta s, \eta s^{-1})$ and solving (in η) the equation $K(x, y) = 0$, one easily obtains

$$\mathcal{S}_1 = \left\{ is \frac{s-i}{s+i} : s = e^{it}, t \in [0, \pi) \right\},$$

which we may rewrite as

$$\mathcal{S}_1 = \{(1 - \sin t, (1 - \sin t) \tan t) : t \in [0, \pi)\},$$

as announced in (2.6). See Figures 3 and 4.

One can choose a conformal mapping π_1 as

$$\pi_1(x) = \frac{x - (x-1)^2}{x + (x-1)^2}.$$

Indeed, π_1 is analytic in \mathcal{S}_1^+ and for all $x = \eta(e^{it})e^{it} \in \mathcal{S}_1$,

$$\pi_1(x) = \frac{-(x + \frac{1}{x}) + 3}{(x + \frac{1}{x}) - 1} = \frac{1 + 2i \frac{\sin^2 t}{\cos t}}{1 - 2i \frac{\sin^2 t}{\cos t}} \in \mathcal{C}.$$

With the conformal mapping ϕ defined in (3.2), we finally obtain

$$(5.1) \quad \psi_1(x) = \phi \circ \pi_1(x) = \frac{x}{(1-x)^2}.$$

Polynomial harmonic functions. Using Equation (1.17) and (5.1), one has

$$H(x, y) = \frac{F\left(\frac{x}{(1-x)^2}\right) - F\left(-\frac{y}{(1-y)^2}\right)}{K(x, y)}.$$

Writing $X = \frac{x}{(1-x)^2}$, $Y = \frac{y}{(1-y)^2}$ and $G(t) = -4F(t)$, one may rewrite the above equation more symmetrically, as

$$(5.2) \quad H(x, y) = \frac{XY G(X) - G(-Y)}{xy(X+Y)}.$$

In particular, applying (5.2) with $G(t) = t$, we have

$$H(x, y) = \frac{1}{(1-x)^2(1-y)^2} = \sum_{i, j \geq 1} ij x^{i-1} y^{j-1}.$$

Recall that $h(i, j) = ij$ is the unique positive harmonic function for this model (unique up to multiplicative factors).

Characterization of the positive harmonic function. As explained in Section 4, the general Martin boundary theory implies that in the framework of this paper, there is a unique positive harmonic function, which in our construction corresponds to taking F as a one-degree polynomial in Theorem 2. However, we don't have any direct proof of this general fact, except precisely for the simple random walk, for which explicit, and in our opinion instructive computations may be done.

More specifically, the question we would like to address here is the following: prove that for all polynomials $G(t) = \sum_{m=1}^k a_m t^m$ with degree $k \geq 2$, there exists at least one coefficient of $H(x, y)$ in (5.2) above which is negative (and in fact infinitely many coefficients are then negative).

We first look at the case when G is a monomial. If $G(t) = t^{2k}$, then one has $H(x, y) = -H(y, x)$ (see (5.2)) and thus H must admit negative coefficients in its Taylor expansion, as the function itself takes negative values. The more interesting case is $G(t) = t^{2k+1}$. However, for a future use, we look at general (meaning non necessarily odd) exponents, i.e., $G(t) = t^k$, for some $k \geq 1$. Then with (5.2) one has

$$H(x, y) = \frac{1}{xy} \sum_{\substack{i+j=k+1 \\ i, j \geq 1}} (-1)^{j-1} X^i Y^j.$$

Moreover, as $n \rightarrow \infty$, one has

$$[x^n] X^i \sim \frac{n^{2i-1}}{(2i-1)!}.$$

So for large values of p and q ,

$$\begin{aligned} [x^{p+1} y^{q+1}] H(x, y) &\sim \sum_{\substack{i+j=k+1 \\ i, j \geq 1}} (-1)^{j-1} \frac{p^{2i-1}}{(2i-1)!} \frac{q^{2j-1}}{(2j-1)!} \\ &= pq \sum_{\substack{i+j=k-1 \\ i, j \geq 0}} (-1)^j \frac{p^{2i}}{(2i+1)!} \frac{q^{2j}}{(2j+1)!}. \end{aligned}$$

Consider now the general case where $G(t)$ is a polynomial $\sum_{m=1}^k a_m t^m$, with $a_k \neq 0$, and take $(p, q) = (r_1, r_2)n$, where r_1, r_2 are positive integers and n tends to infinity. Using the above estimate, we have

$$[x^{p+1} y^{q+1}] H(x, y) \sim n^{2k} r_1^{2k-1} r_2 \sum_{\substack{i+j=k-1 \\ i, j \geq 0}} (-1)^j \frac{(r_2/r_1)^{2j}}{(2i+1)!(2j+1)!}.$$

Replacing r_2/r_1 by x , our question is therefore equivalent to prove that for any fixed integer $k \geq 2$, there exists $x \in (0, \infty)$ such that

$$\sum_{\substack{i+j=k-1 \\ i, j \geq 0}} (-1)^j \frac{x^{2j}}{(2i+1)!(2j+1)!} < 0.$$

To that purpose, we first observe that

$$\begin{aligned} \sum_{\substack{i+j=k-1 \\ i,j \geq 0}} (-1)^j \frac{x^{2j}}{(2i+1)!(2j+1)!} &= \frac{(1+ix)^{2k} - (1-ix)^{2k}}{2(2k)!ix} \\ &= \frac{1}{(2k)!x} \Im((ix+1)^{2k}) \\ &= \frac{(1+x^2)^k}{(2k)!x} \sin(2k \arctan x). \end{aligned}$$

Clearly, given $k \geq 2$ one may fix $x \in (0, \infty)$ such that the above is negative. More precisely, this function admits $k-1$ sign changes.

A related example: the king walk. We continue with the king walk, which (see the second example on Figure 1) by definition admits the kernel

$$K(x, y) = xy \left(1 - \frac{xy + x + xy^{-1} + y^{-1} + x^{-1}y^{-1} + x^{-1} + x^{-1}y + y}{8} \right).$$

As the simple walk, its unique positive harmonic function is given by $h(i, j) = ij$. However, this example is a bit different as we now have $p_{1,1} \neq 0$. A few computations starting from the kernel yield

$$\mathcal{A}_1 = \left\{ \frac{u(t) - \sqrt{u(t)^2 - 4}}{2} e^{it} : t \in [0, 2\pi) \right\},$$

where $u(t) = -\cos t + \sqrt{12 - 3\cos^2 t}$. This curve is drawn on Figure 4. Computations similar to that of the simple walk lead to the following rather simple expressions for the conformal mappings:

$$\pi_1(x) = \frac{3x}{x^2 + x + 1} \quad \text{and} \quad \psi_1(x) = \frac{x^2 + 4x + 1}{(1-x)^2}.$$

Using Equation (1.17) with $F(t) = t/16$, then one can recover the generating function of the positive harmonic function

$$H(x, y) = \frac{F(\psi_1(x)) - F(-\psi_1(y))}{K(x, y)} = \frac{1}{(1-x)^2(1-y)^2} = \sum_{i,j \geq 1} ijx^{i-1}y^{j-1}.$$

Similar computations would lead to other harmonic functions.

5.2. Symmetric small step random walks and comparison between the analytic approaches of [9] and [19, 42]. In this part, we look at the class of symmetric, small step random walks, meaning that $p_{i,j} = 0$ as soon as $|i| \geq 2$ or $|j| \geq 2$. As illustrated by our bibliography, this class of models has been (and is still) widely studied in the literature, the main reason being that the zero set of the kernel is, in this case, a Riemann surface of genus 0 or 1, opening the way to explicit parametrizations in terms of rational or elliptic functions.

Here our main objective is twofold: first, we will derive an expression of the conformal mapping ψ_1 for small step random walks, see Proposition 17. Doing so, we will introduce some tools from complex analysis, which are close to the analytic approach developed in




FIGURE 6. The conformal mapping for the simple random walk: $\sigma(x)$ maps conformally $\mathcal{S}_1^+ \setminus [x_1, 1]$ (the light blue domain) onto the cone, and $\omega(s)$ maps conformally the cone onto the right half plane \mathcal{H}^+ .

[19, 42]. As a second step, we will compare the analytic approach used in this paper (inspired by [9]) with the one of [19, 42].

Explicit expression for the conformal mapping ψ_1 . We first introduce a function $T_a(x)$ that generalizes the classical Chebyshev polynomials of the first kind, obtained when $a \in \mathbb{N}$. This function is defined, for $x \in \mathbb{C} \setminus (-\infty, -1]$, by

$$(5.3) \quad T_a(x) = {}_2F_1\left(-a, a; \frac{1}{2}; \frac{1-x}{2}\right),$$

where ${}_2F_1$ is the Gauss hypergeometric function. Then T_a admits the following expansion, valid for $|x-1| < 2$:

$$T_a(x) = \sum_{n \geq 0} \frac{a}{a+n} \binom{a+n}{2n} 2^n (x-1)^n.$$

When $a \in \mathbb{N}$, then the above sum ranges from 0 to a , and $T_a(\cos t) = \cos(at)$, by definition of the classical Chebyshev polynomial. Another useful formula, valid for x in $\mathbb{C} \setminus (-\infty, -1]$, is

$$T_a(x) = \frac{1}{2} \left((x + \sqrt{x^2 - 1})^a + (x - \sqrt{x^2 - 1})^a \right).$$

Proposition 17. *Assume (H1)–(H4) and the small step hypothesis. Let θ as in (1.16) and*

$$\mu(x) = \frac{\mu_0 x - \mu_1}{2(x-1)},$$

with μ_0 and μ_1 defined by (5.4). Then the conformal map ψ_1 may be chosen such that

$$\psi_1(x) = 2T_{\pi/\theta}(\mu(x)),$$

where $T_{\pi/\theta}$ is the generalized Chebyshev polynomial (5.3) with $a = \pi/\theta$.

We now prove Proposition 17. The main idea of the proof is borrowed from [19, Sec. 6.5], see also [42]. For driftless, small step random walks, the zero set of the kernel

$$\{(x, y) \in (\mathbb{C} \cup \{\infty\})^2 : K(x, y) = 0\}$$

is a Riemann surface of genus 0, which can thus be parametrized with rational functions. As it turns out, the curves \mathcal{S}_1 and \mathcal{S}_2 become particularly simple in the uniformizing variable, from what we will deduce an expression for the conformal map.

Before stating the rational uniformisation of the above Riemann surface, we introduce a few notations. First, the kernel (1.10) may be rewritten as $K(x, y) = a(x)y^2 + b(x)y + c(x)$, where

$$\begin{cases} a(x) &= -(p_{-1,-1}x^2 + p_{0,-1}x + p_{1,-1}), \\ b(x) &= -(p_{-1,0}x^2 - x + p_{1,0}), \\ c(x) &= -(p_{-1,1}x^2 + p_{0,1}x + p_{1,1}). \end{cases}$$

Let also $d = b^2 - 4ac$ denote the discriminant of $K(x, y)$ in y . It is seen that d has degree 3 or 4, and that 1 is a double root. In the case that d has degree 3 (resp. 4), then the remaining root is denoted by x_1 (resp. the remaining roots are denoted by x_1 and x_4). It has been proved in [19] that $x_1 \in [-1, 1)$ and $x_4 \in (1, \infty) \cup (-\infty, -1]$. In the case of d having degree 3, we will denote $x_4 = \infty$. Now put

$$\begin{cases} s_0 &= \frac{2-(x_1+x_4)+2\sqrt{(1-x_1)(1-x_4)}}{x_4-x_1}, \\ s_1 &= \frac{x_1+x_4-2x_1x_4+2\sqrt{x_1x_4(1-x_1)(1-x_4)}}{x_4-x_1}, \end{cases}$$

as well as (with θ defined in (1.16))

$$(5.4) \quad \mu_0 = s_0 + \frac{1}{s_0}, \quad \mu_1 = s_1 + \frac{1}{s_1}, \quad \rho = e^{-i\theta}.$$

We may now state the rational uniformization; for a proof, we refer to [17, Sec. 2.3].

Lemma 18. *One has*

$$\{(x, y) \in (\mathbb{C} \cup \{\infty\})^2 : K(x, y) = 0\} = \{(x(s), y(s)) : s \in \mathbb{C} \cup \{\infty\}\},$$

where

$$x(s) = \frac{(s - s_1)(s - \frac{1}{s_1})}{(s - s_0)(s - \frac{1}{s_0})} \quad \text{and} \quad y(s) = \frac{(\rho s - s_1)(\rho s - \frac{1}{s_1})}{(\rho s - s_0)(\rho s - \frac{1}{s_0})}.$$

Moreover, the above rational functions admit the involutions $x(s) = x(1/s)$ and $y(s) = y(1/(\rho^2 s))$.

Then the set of complex points where $|x| = |y|$ (of which \mathcal{H} in (1.13) is a subset) is very simple, as in the s variable, it becomes simply the line $e^{i\frac{\theta}{2}}\mathbb{R}$. More precisely, defining the cone

$$\mathcal{E}^+ = \{re^{i\chi} : r > 0 \text{ and } \chi \in (0, \frac{\theta}{2})\},$$

one has the following result, which is illustrated on Figure 6:

Lemma 19. *The function $x(s)$ is one-to-one from $e^{i\frac{\theta}{2}}\mathbb{R}_+$ onto \mathcal{S}_1 . Moreover, $x(s)$ is a conformal mapping from \mathcal{E}^+ onto $\mathcal{S}_1^+ \setminus [x_1, 1]$.*

Proof. The crucial fact is that $x(s)$ maps one-to-one from \mathcal{H}^+ onto the whole plane \mathbb{C} cut some segments. Depending on the value of x_4 , $x(s)$ maps one-to-one from \mathcal{H}^+ onto

- $\mathbb{C} \setminus [x_1, x_4]$ if $x_4 > 1$ or $x_4 = \infty$;
- $\mathbb{C} \setminus ([x_1, \infty) \cup (-\infty, x_4])$ if $x_4 < -1$.

We first prove that $x(s)$ maps one-to-one from $e^{i\theta/2}\mathbb{R}_+$ onto \mathcal{S}_1 . Indeed, for a point $s = re^{i\frac{\theta}{2}}$ with $r > 0$, one has

$$\Im(\phi(x(s))) = \frac{2(r + \frac{1}{r})\cos\frac{\theta}{2} - (\mu_0 + \mu_1)}{\mu_1 - \mu_0} \geq \frac{4\cos\frac{\theta}{2} - (\mu_0 + \mu_1)}{\mu_1 - \mu_0} > 0,$$

where the last inequality follows from a direct computation. This means that $|x(s)| < 1$. Combining with the facts that $x(s) = \overline{y(s)}$ and $(x(s), y(s))$ is a solution of $K(x, y)$, we deduce that $x(s) \in \mathcal{S}_1$ for all $s \in e^{i\theta/2}\mathbb{R}_+$. The one-to-one property between two curves then follows. To conclude, it is easy to check that $x(s)$ also maps one-to-one $[0, 1]$ (resp. $[1, \infty)$) onto $[x_1, 1]$. Hence, $x(s)$ maps one-to-one from \mathcal{E}^+ onto $\mathcal{S}_1 \setminus [x_1, 1]$. \square

Let $\sigma(s)$ denote the inverse mapping of $x(s)$:

$$(5.5) \quad \sigma(x) = \frac{\mu_0 x - \mu_1}{2(x-1)} + \sqrt{\left(\frac{\mu_0 x - \mu_1}{2(x-1)}\right)^2 - 1}.$$

The branch cut of the square root in (5.5) is chosen such that $\sigma(s)$ maps conformally $\mathcal{S}_1^+ \setminus [x_1, 1]$ onto \mathcal{E}^+ .

Now consider the following mapping

$$\omega(s) = s^{\pi/\theta} + s^{-\pi/\theta},$$

which maps conformally \mathcal{E}^+ onto $\mathcal{H}^+ \setminus [2, \infty)$. It is seen that $\omega \circ \sigma(s)$ maps conformally $\mathcal{S}_1^+ \setminus [x_1, 1]$ onto $\mathcal{H}^+ \setminus [2, \infty)$ (see Figure 6). In fact, in the following lemma, a stronger statement can be deduced.

Lemma 20. *The function $\omega \circ \sigma$ maps conformally \mathcal{S}_1^+ onto \mathcal{H}^+ .*

Proof. By the specific form of $\omega \circ \sigma$, we know that $\omega \circ \sigma$ is analytic in \mathcal{S}_1^+ . Moreover, since $\omega \circ \sigma$ is univalent on $\mathcal{S}_1^+ \setminus [x_1, 1]$, then it remains to prove that $\omega \circ \sigma$ is injective on $[x_1, 1)$. This is true since σ (with suitable branch) maps one-to-one $[x_1, 1)$ onto $[1, \infty)$ and ω maps one-to-one $[1, \infty)$ onto $[2, \infty)$. Therefore, $\omega \circ \sigma$ is univalent on \mathcal{S}_1^+ . The proof is complete. \square

End of the proof of Proposition 17. We use Lemma 20 together with the expressions for $\omega(s)$ and $s(x)$. \square

Comparison between the analytic approaches of [9] and [19, 42]. Both approaches start with the same functional equation (1.11), and, as a second step, introduce subsets of \mathbb{C}^2 where this equation may be evaluated. The approaches differ in the choice of these subsets:

- In [9] (which we choose to follow in the present work), this subset is chosen to be \mathcal{H} in (1.13) (where we recall that $|x| = |y| \leq 1$);
- On the other hand, the set in [19] is $x \in [x_1, 1]$ (then $y = Y(x)$ is a solution to the kernel equation, and x_1 is the branch point introduced in the previous section).

In both cases, the method continues by stating (and solving) a BVP for the generating functions on curves obtained from the subsets above.

This short recap shows that the unique, but major difference in the two approaches lies in the choice of the domain of evaluation. Both choices are equally natural for small step random walks. However, the main advantage of the choice of [9] is that the domain \mathcal{H} may be defined without any difficulty for models admitting arbitrary big negative jumps, as in our paper. On the contrary, we did not find any canonical way to extend the definition of the segment $[x_1, 1]$ for large step models¹. Let us also underline the $x \leftrightarrow y$ symmetry of

¹For small step models, there is only one branch point interior to the unit disk (see again the previous section), so x_1 appears as the only possible choice. However, the number of branch points being increasing




FIGURE 7. On the left: the model studied in Section 5.3. It is inspired by a similar (non-symmetric) model analyzed in [28, 5], displayed on the right picture.

the domain \mathcal{X} , while this symmetry is broken when taking $x \in [x_1, 1]$ (indeed x is then real and y becomes non-real).

Finally, as shown in the last section, in the case of small steps, the two approaches are very similar: our curve \mathcal{S}_1 corresponds to the line $e^{i\theta/2}\mathbb{R}_+$, while the curve $[x_1, 1]$ would correspond to \mathbb{R}_+ ; one passes from one curve to the other simply by multiplying by a complex number.

5.3. A family of random walks with larger steps. Consider the model whose jumps and weights are given by

$$(5.6) \quad p_{k,\ell} = \begin{cases} z & \text{if } (k, \ell) = (1, 1), \\ z_r & \text{if } k + \ell + r = 0, \end{cases}$$

where the z_r satisfy $z + \sum_r (r+1)z_r = 1$ (so that the $p_{k,\ell}$ are transition probabilities) and $z = \sum_r z_r \frac{r(r+1)}{2}$ (so as to have a zero drift, see our hypothesis (H4) in Section 1). See Figure 7. For example, choosing $z = z_1 = \frac{1}{3}$ and all other $z_r = 0$ leads to Kreweras' step set $\{(1, 1), (-1, 0), (0, -1)\}$ with uniform weights.

Let us remark that the model (5.6) is not always irreducible. More precisely, it is reducible if and only if all steps (k, ℓ) have even size (by which we mean that all coordinate sums $k+\ell$ are even). Although our irreducibility assumption (H3) is not satisfied in general, we will show how to apply our main results in this slightly modified framework.

Our motivation to look at this particular family of model comes from bipolar orientations on planar maps, which, as shown in the papers [28] and [5], are in close correspondence with the model of walks confined to the first quadrant as on the right of Figure 7. Then our model (on the left on the same picture) is just obtained through a horizontal symmetry, so as in particular to have a model symmetric in the first diagonal. Because of the connection with bipolar orientations, both models represented on Figure 7 admit a very strong structure, which the results in this section will also illustrate.

with the amplitude of the big jumps, it is not clear at all in general what segment, or what union of segments might replace $[x_1, 1]$ in general.

Introduce

$$(5.7) \quad \rho(x) = z \sqrt{\frac{I_0(x) - I_0(t)}{I_0(x) - I_0(1)}},$$

where the function I_0 , introduced in [5], takes the value

$$I_0(x) = x + \frac{z}{x} - \sum z_r x^{r+1},$$

where t is the unique real point of \mathcal{S}_1 apart from 1, and where the branch cut of the square root function in (5.7) is chosen on \mathbb{R}_- . Equivalently, t is characterized by

$$(5.8) \quad I'_0(t) = 0 \quad \text{and} \quad t \in (-1, 1),$$

as it easily follows from (5.9). Our main result is the following:

Proposition 21. *The map ρ in (5.7) is a conformal map from \mathcal{S}_1^+ to \mathcal{H}^+ , sending 0 to z .*

The remainder of this section has the following structure: we will first provide the proof of Proposition 21, and then give a few consequences of it.

Proof of Proposition 21. We start by noticing that for $x \neq y$,

$$\begin{aligned} K(x, y) &= xy - z - xy \sum z_r \frac{y^{r+1} - x^{r+1}}{y - x} \\ &= \frac{xy}{y - x} \left(y - x - \frac{z}{x} + \frac{z}{y} - \sum z_r (y^{r+1} - x^{r+1}) \right) \\ &= \frac{xy}{y - x} (I_0(y) - I_0(x)), \end{aligned}$$

and for $x = y$

$$(5.9) \quad K(x, x) = x^2 - z - \sum (r + 1) z_r x^{r+2} = x^2 I'_0(x).$$

Denote $\mathbb{P}^1 = \mathbb{C} \cup \{\infty\}$.

Lemma 22. *If $(x, y) \in \mathcal{K}$, then $I_0(x) = I_0(y) \in \mathbb{R}$, and the map $(x, y) \mapsto I_0(x)$ is two-to-one from \mathcal{K} to an interval $[a, b]$ of \mathbb{R} , with $a = I_0(t) < 0$ and $b = I_0(1) > 0$. Moreover, I_0 maps the curve \mathcal{S}_1^+ onto $\mathbb{P}^1 \setminus [a, b]$.*

Proof. Note first that if $x = y$, then the condition $x = \bar{y}$ implied by the symmetry of K yields $x = \pm 1$, which implies $I_0(x) \in \mathbb{R}$. The point $(1, 1)$ always belongs to \mathcal{K} , and so does $(-1, -1)$ if and only if the walk is reducible (which, we recall, happens when all steps have even size).

If $x \neq y$, the above expression of K yields $I_0(x) = I_0(y)$. Since I_0 has real coefficients and $y = \bar{x}$, we also have $I_0(x) = I_0(\bar{x}) = \overline{I_0(x)}$, and $I_0(x) \in \mathbb{R}$.

By the equality $K(x, x) = x^2 I'_0(x)$, see (5.9), we get that $I'_0(x) = 0$ if and only if $K(x, x) = 0$. By Rouché's theorem, for $h \in (0, 1)$, the map

$$K_h(x) = x^2 - h \left(z + \sum (r + 1) z_r x^{r+2} \right)$$

has exactly two zeros in the unit disk, which are real. Hence, letting h go to one yields that $x \mapsto K(x, x)$ has at most two real zeros in the unit disk, plus eventual additional zeros

on the unit circle. Moreover, $K(x, x)$ cannot vanish on the unit circle except at $-1, 1$; this follows from (H3) in the irreducible case, and would follow from a similar argument in the reducible case. By the study of $x \mapsto K(x, x)$ on $[-1, 1]$ done in Section 2, there are only two zeros at the two real points of \mathcal{S}_1 , one being 1 and the other one being strictly negative (since $p_{1,1} > 0$). Hence, $I_0'(x)$ vanishes only at the two real points of \mathcal{S}_1 . Since $I_0(x) = I_0(\bar{x})$, we deduce that I_0 is two-to-one from \mathcal{S}_1 to an interval $[a, b]$ of \mathbb{R} . Since $z > \sum z_r$, we deduce that $I_0(1) > 0$ and $I_0(t) < 0$, where t is the unique real point of \mathcal{S}_1 apart from 1 (recall that t is negative by Lemma 2).

Since I_0 is analytic on \mathcal{S}_1^+ and continuous on the closure $\overline{\mathcal{S}_1^+}$,

$$\partial(I_0(\mathcal{S}_1^+)) \subset I_0(\partial\mathcal{S}_1^+) = I_0(\mathcal{S}_1) = [I_0(t), I_0(1)] = [a, b].$$

Hence, $\mathbb{P}^1 \setminus [a, b] \subset I_0(\mathcal{S}_1^+)$. Let $x \in \mathcal{S}_1^+ \setminus \mathbb{R}$. Then, $I_0(x) \notin [a, b]$, for otherwise $K(x, \bar{x}) = 0$ which would contradict the fact that $x \notin \mathcal{S}_1$. On $\mathbb{R} \cap \mathcal{S}_1^+$, $x^2 I_0'(x) = K(x, x) < 0$, thus I_0 is increasing. Since $I_0(0) = \infty$, we deduce that $I_0(\mathbb{R} \cap \mathcal{S}_1^+) = (\mathbb{R} \setminus [a, b]) \cup \{\infty\}$, and thus $I_0(\mathcal{S}_1^+) = \mathbb{P}^1 \setminus [a, b]$. \square

End of the proof of Proposition 21. First, we directly check that $\rho(0) = z\sqrt{1} = z$. Further, since

$$z \mapsto \sqrt{\frac{z-a}{z-b}}$$

maps conformally $\mathbb{P}^1 \setminus [a, b]$ to the right half-plane \mathcal{H}^+ , we deduce from Lemma 22 that ρ in (5.7) is analytic from \mathcal{S}_1^+ to \mathcal{H}^+ . As a second step, we will prove that ρ is continuous and one-to-one from \mathcal{S}_1 to $[a, b]$. As a direct consequence, ρ will be conformal from \mathcal{S}_1^+ to \mathcal{H}^+ , thereby concluding the proof of Proposition 21.

We thus show that ρ extends by continuity to an injective map from \mathcal{S}_1 to $[a, b] \subset \mathbb{R}$. Suppose that $\{x_n\}_{n \geq 1}$ converges to $\xi \in \mathcal{S}_1$ with $\Im \xi > 0$. Then, by continuity of I_0 , $I_0(x_n) \rightarrow I_0(\xi)$. Since $I_0'(x) < 0$ for $x \in [t, 1]$ and $I_0([t, 1]) \subset \mathbb{R}$, we deduce that $\Im I_0(x) < 0$ for x in a neighbourhood of $[1, t]$ in $i\mathcal{H}^+$. Taking into account that $I_0(\mathcal{S}_1^+ \cap i\mathcal{H}^+) \subset \mathbb{C} \setminus \mathbb{R}$ (for otherwise it would meet \mathcal{S}_1), we have $I_0(\mathcal{S}_1^+ \cap i\mathcal{H}^+) \subset -i\mathcal{H}^+$. Hence, $I_0(x_n) \rightarrow I_0(\xi) \in [a, b]$ while staying in $-i\mathcal{H}^+$. This implies that

$$\frac{I_0(x_n) - a}{I_0(x_n) - b} = 1 + \frac{b-a}{I_0(x_n) - b} \rightarrow \frac{I_0(\xi) - a}{I_0(\xi) - b},$$

while being after some rank in $i\mathcal{H}^+$. Thus, $\rho(x_n)$ goes towards $iz\sqrt{\frac{I_0(\xi)-a}{b-I_0(\xi)}}$, where $\sqrt{\frac{I_0(\xi)-a}{b-I_0(\xi)}}$ is the unique positive root of $X^2 = \frac{I_0(\xi)-a}{b-I_0(\xi)}$.

Similarly, if $\{x_n\}_{n \geq 1}$ converges to $\xi \in \mathcal{S}_1$ with $\Im \xi < 0$, then $\rho(x_n)$ goes to $-iz\sqrt{\frac{I_0(\xi)-a}{b-I_0(\xi)}}$. Hence, we can extend ρ by continuity to \mathcal{S}_1 with the value

$$\rho(\xi) = \pm iz \sqrt{\frac{I_0(\xi) - a}{b - I_0(\xi)}} \quad \text{if } \xi \in \pm i\mathcal{H}^+.$$

Notice that the above formula still holds when $\xi \in \mathbb{R}$, with $\rho(1) = \infty$ and $\rho(t) = 0$. Since I_0 is two-to-one from \mathcal{S}_1 to $[a, b]$ except at t and 1, we deduce that ρ is injective on \mathcal{S}_1 . \square

Applications of Proposition 21. Theorem 1 directly yields the following explicit expression for the harmonic functions h_n as introduced in (1.15).

Proposition 23. *Let t as in (5.8). Then, for each $n \geq 1$, the power series expansion at $(0, 0)$ of the bivariate series*

$$H_n(x, y) = \frac{(y-x)z^n \left(\left(\frac{I_0(x)-I_0(t)}{I_0(x)-I_0(1)} \right)^{n/2} + (-1)^{n-1} \left(\frac{I_0(y)-I_0(t)}{I_0(y)-I_0(1)} \right)^{n/2} \right)}{xy(I_0(y) - I_0(x))}$$

defines a harmonic function h_n for the Laplacian operator associated to the model (5.6).

Remark that for any even values of $n \geq 1$, H_n is a rational function.

For Kreweras' model, $I_0(x) = x + \frac{1}{3x} - \frac{x^2}{3}$. Hence, $I_0'(x) = 1 - \frac{1}{3x^2} - \frac{2x}{3}$, which admits the unique root $t = -\frac{1}{2}$ in $(-1, 1)$. Hence, after some computations, ρ in (5.7) simplifies into

$$\rho(x) = \frac{1+2x}{3} \sqrt{\frac{1-x/4}{(1-x)^3}}$$

and we have for example

$$\begin{aligned} H_1(x, y) &= \frac{(1+2x)\sqrt{\frac{1-x/4}{(1-x)^3}} + (1+2y)\sqrt{\frac{1-y/4}{(1-y)^3}}}{3xy(1 - \frac{1}{3}(\frac{1}{xy} + x + y))} \\ &= -\frac{1}{18} \left(1 + \frac{27}{16}x + \frac{27}{16}y + \frac{567}{256}x^2 + 3xy + \frac{567}{256}y^2 + \dots \right). \end{aligned}$$

We also have

$$H_2(x, y) = -\frac{9}{4} \frac{x-y}{(1-x)^3(1-y)^3} = -\frac{9}{4} (x-y + 3x^2 - 3y^2 + \dots).$$

As a second example, consider $z = \frac{1}{2}$, $z_2 = \frac{1}{6}$ and all other $z_r = 0$. Then the curve admits the parametrization

$$(5.10) \quad \mathcal{S}_1 = \left\{ \frac{1}{\sqrt{1 + \frac{2}{\sqrt{3}} \sin t}} e^{it} : t \in [0, 2\pi) \right\},$$

see Figure 8. In this case, the conformal mapping takes the form

$$\rho(x) = \frac{1}{2} \sqrt{\frac{(3-x)(1+x)^3}{(3+x)(1-x)^3}}$$

REFERENCES

- [1] Ph. Biane. Équation de Choquet-Deny sur le dual d'un groupe compact. *Probab. Theory Related Fields*, 94(1):39–51, 1992.
- [2] Philippe Biane. Quantum random walk on the dual of $SU(n)$. *Probab. Theory Related Fields*, 89(1):117–129, 1991.
- [3] Philippe Biane, Philippe Bougerol, and Neil O'Connell. Littelmann paths and Brownian paths. *Duke Math. J.*, 130(1):127–167, 2005.
- [4] Aymen Bouaziz, Sami Mustapha, and Mohamed Sifi. Discrete harmonic functions on an orthant in \mathbb{Z}^d . *Electron. Commun. Probab.*, 20:no. 52, 13, 2015.




FIGURE 8. The curve \mathcal{S}_1 in (5.10), for the model (5.6) with jumps $z = \frac{1}{2}$, $z_2 = \frac{1}{6}$ and all other $z_r = 0$.

- [5] Mireille Bousquet-Mélou, Éric Fusy, and Kilian Raschel. Plane bipolar orientations and quadrant walks. *Sém. Lothar. Combin.*, 81:Art. B81, 2020.
- [6] Mireille Bousquet-Mélou and Marni Mishna. Walks with small steps in the quarter plane. In *Algorithmic probability and combinatorics*, volume 520 of *Contemp. Math.*, pages 1–39. Amer. Math. Soc., Providence, RI, 2010.
- [7] François Chapon, Éric Fusy, and Kilian Raschel. Polyharmonic functions and random processes in cones. In Michael Drmota and Clemens Heuberger, editors, *31st International Conference on Probabilistic, Combinatorial and Asymptotic Methods for the Analysis of Algorithms (AofA 2020)*, volume 159 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 9:1–9:19, Dagstuhl, Germany, 2020. Schloss Dagstuhl–Leibniz-Zentrum für Informatik.
- [8] J. W. Cohen. *Analysis of random walks*, volume 2 of *Studies in Probability, Optimization and Statistics*. IOS Press, Amsterdam, 1992.
- [9] Jacob Willem Cohen and O. J. Boxma. *Boundary value problems in queueing system analysis*, volume 79 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1983.
- [10] J. Courtiel, S. Melzer, M. Mishna, and K. Raschel. Weighted lattice walks and universality classes. *J. Combin. Theory Ser. A*, 152:255–302, 2017.
- [11] Denis Denisov and Vitali Wachtel. Random walks in cones. *Ann. Probab.*, 43(3):992–1044, 2015.
- [12] Thomas Dreyfus, Charlotte Hardouin, Julien Roques, and Michael F. Singer. On the nature of the generating series of walks in the quarter plane. *Invent. Math.*, 213(1):139–203, 2018.
- [13] R. J. Duffin. Basic properties of discrete analytic functions. *Duke Math. J.*, 23:335–363, 1956.
- [14] Jetlir Duraj, Kilian Raschel, Pierre Tarrago, and Vitali Wachtel. Martin boundary of random walks in convex cones. *Preprint arXiv:2003.03647*, 2020.
- [15] Peter Eichelsbacher and Wolfgang König. Ordered random walks. *Electron. J. Probab.*, 13:no. 46, 1307–1336, 2008.
- [16] David Ellwood, Charles Newman, Vidas Sidoravicius, and Wendelin Werner, editors. *Probability and statistical physics in two and more dimensions*, volume 15 of *Clay Mathematics Proceedings*. American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2012.
- [17] G. Fayolle and K. Raschel. Random walks in the quarter-plane with zero drift: an explicit criterion for the finiteness of the associated group. *Markov Process. Related Fields*, 17(4):619–636, 2011.
- [18] Guy Fayolle and Roudolf Iasnogorodski. Two coupled processors: the reduction to a Riemann-Hilbert problem. *Z. Wahrsch. Verw. Gebiete*, 47(3):325–351, 1979.
- [19] Guy Fayolle, Roudolf Iasnogorodski, and Vadim Malyshev. *Random walks in the quarter plane*, volume 40 of *Probability Theory and Stochastic Modelling*. Springer, Cham, second edition, 2017.

- Algebraic methods, boundary value problems, applications to queueing systems and analytic combinatorics.
- [20] Guy Fayolle and Kilian Raschel. About a possible analytic approach for walks in the quarter plane with arbitrary big jumps. *C. R. Math. Acad. Sci. Paris*, 353(2):89–94, 2015.
 - [21] Jacqueline Ferrand. Fonctions préharmoniques et fonctions préholomorphes. *Bull. Sci. Math. (2)*, 68:152–180, 1944.
 - [22] R.C. Gunning and H. Rossi. *Analytic Functions of Several Complex Variables*. Ams Chelsea Publishing. Prentice-Hall, 1965.
 - [23] H. A. Heilbronn. On discrete harmonic functions. *Proc. Cambridge Philos. Soc.*, 45:194–206, 1949.
 - [24] Irina Ignatiouk-Robert. Martin boundary of a killed random walk on a half-space. *J. Theoret. Probab.*, 21(1):35–68, 2008.
 - [25] Irina Ignatiouk-Robert. Harmonic functions of random walks in a semigroup via ladder heights. *J. Theoret. Probab.*, 21, to appear.
 - [26] Irina Ignatiouk-Robert and Christophe Loree. Martin boundary of a killed random walk on a quadrant. *Ann. Probab.*, 38(3):1106–1142, 2010.
 - [27] Rufus Isaacs. Monodiffic functions. Construction and applications of conformal maps. In *Proceedings of a symposium*, National Bureau of Standards, Appl. Math. Ser., No. 18, pages 257–266. U. S. Government Printing Office, Washington, D. C., 1952.
 - [28] Richard Kenyon, Jason Miller, Scott Sheffield, and David B. Wilson. Bipolar orientations on planar maps and SLE_{12} . *Ann. Probab.*, 47(3):1240–1269, 2019.
 - [29] Wolfgang König and Patrick Schmid. Random walks conditioned to stay in Weyl chambers of type C and D. *Electron. Commun. Probab.*, 15:286–296, 2010.
 - [30] I. A. Kurkova and V. A. Malyshev. Martin boundary and elliptic curves. *Markov Process. Related Fields*, 4(2):203–272, 1998.
 - [31] I. A. Kurkova and Y. M. Suhov. Malyshev’s theory and JS-queues. Asymptotics of stationary probabilities. *Ann. Appl. Probab.*, 13(4):1313–1354, 2003.
 - [32] Irina Kurkova and Kilian Raschel. On the functions counting walks with small steps in the quarter plane. *Publ. Math. Inst. Hautes Études Sci.*, 116:69–114, 2012.
 - [33] Cédric Lecouvey and Kilian Raschel. t -Martin boundary of killed random walks in the quadrant. In *Séminaire de Probabilités XLVIII*, volume 2168 of *Lecture Notes in Math.*, pages 305–323. Springer, Cham, 2016.
 - [34] V. A. Malyšev. An analytic method in the theory of two-dimensional positive random walks. *Sibirsk. Mat. Ž.*, 13:1314–1329, 1421, 1972.
 - [35] B. H. Murdoch. A theorem on harmonic functions. *J. London Math. Soc.*, 39:581–588, 1964.
 - [36] Sami Mustapha. Gaussian estimates for spatially inhomogeneous random walks on \mathbf{Z}^d . *Ann. Probab.*, 34(1):264–283, 2006.
 - [37] Zeev Nehari. *Conformal mapping*. McGraw-Hill Book Co., Inc., New York, Toronto, London, 1952.
 - [38] Robin Pemantle and Mark C. Wilson. *Analytic combinatorics in several variables*, volume 140 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2013.
 - [39] Massimo A. Picardello and Wolfgang Woess. Martin boundaries of Cartesian products of Markov chains. *Nagoya Math. J.*, 128:153–169, 1992.
 - [40] Ch. Pommerenke. *Boundary behaviour of conformal maps*, volume 299 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1992.
 - [41] Kilian Raschel. Green functions for killed random walks in the Weyl chamber of $Sp(4)$. *Ann. Inst. Henri Poincaré Probab. Stat.*, 47(4):1001–1019, 2011.
 - [42] Kilian Raschel. Random walks in the quarter plane, discrete harmonic functions and conformal mappings. *Stochastic Process. Appl.*, 124(10):3147–3178, 2014. With an appendix by Sandro Franceschi.
 - [43] N. Th. Varopoulos. Potential theory in conical domains. *Math. Proc. Cambridge Philos. Soc.*, 125(2):335–384, 1999.
 - [44] Nicolas Th. Varopoulos. The discrete and classical Dirichlet problem. *Milan J. Math.*, 77:397–436, 2009.

[45] Nicolas Th. Varopoulos. The discrete and classical Dirichlet problem: Part II. *Milan J. Math.*, 83(1):1–20, 2015.

APPENDIX A. WEIERSTRASS' PREPARATION AND DIVISION THEOREMS

We recall here two important results in the study of analytic functions or formal power series in several variables. We will only state these results in the case where two variables are involved, even if they hold for an arbitrary (finite) number of variables. Let $\mathbb{K}[[x, y]]$ denote the formal ring of power series in two variables, with coefficients in the field \mathbb{K} . Recall that $f \in \mathbb{K}[[x, y]]$ is invertible in $\mathbb{K}[[x, y]]$ if and only if the constant term of f is non-zero.

Definition 1. *A Weierstrass polynomial with respect to y of degree n is a polynomial $P \in \mathbb{K}[[x]][y]$ of degree n with leading coefficient equal to 1 and with non-invertible lower coefficients. In other words, P can be written as*

$$P(x, y) = y^n + F_{n-1}(x)y^{n-1} + \cdots + F_0(x),$$

with $F_0, \dots, F_{n-1} \in \mathbb{K}[[x]]$ such that $F_i(0) = 0$ for all $0 \leq i \leq n-1$.

Then, we have the following main results, respectively called Weierstrass preparation theorem and Weierstrass division theorem.

Theorem 3 (Weierstrass preparation theorem). *Suppose that $f \in \mathbb{K}[[x, y]]$ is non-invertible and that for some $k \geq 1$, the coefficient in y^k of f is non-zero. Then there exists a unique Weierstrass polynomial P with respect to y and an invertible element of $\mathbb{K}[[x, y]]$ such that*

$$f = hP.$$

If f is the germ of an analytic function at $(0, 0)$, then h and P are also germs of analytic functions at $(0, 0)$.

Theorem 4 (Weierstrass division theorem). *Suppose that $f \in \mathbb{K}[[x, y]]$ and P is a Weierstrass polynomial of degree n with respect to y . Then there exists a unique pair (q, R) with $q \in \mathbb{K}[[x, y]]$ and R a Weierstrass polynomial with respect to y of degree less than n such that*

$$f = qP + R.$$

If f and P are germs of analytic functions at $(0, 0)$, then q and R are also germs of analytic functions at $(0, 0)$.

CNRS AND INSTITUT DENIS POISSON, UMR CNRS 7013, UNIVERSITÉ DE TOURS ET UNIVERSITÉ D'ORLÉANS, PARC DE GRANDMONT, 37200 TOURS, FRANCE

Email address: viet.hung-hoang@lmpt.univ-tours.fr

CNRS AND INSTITUT DENIS POISSON, UMR CNRS 7013, UNIVERSITÉ DE TOURS ET UNIVERSITÉ D'ORLÉANS, PARC DE GRANDMONT, 37200 TOURS, FRANCE

Email address: raschel@math.cnrs.fr

LABORATOIRE DE PROBABILITÉS, STATISTIQUE ET MODÉLISATION, UMR CNRS 8001, SORBONNE UNIVERSITÉ, 4 PLACE JUSSIEU, 75005 PARIS, FRANCE

Email address: pierre.tarrago@upmc.fr