



HAL
open science

Mediator complex subunit Med19 binds directly GATA transcription factors and is required with Med1 for GATA-driven gene regulation in vivo

Clément Immarigeon, Sandra Bernat-Fabre, Emmanuelle Guillou, Alexis Verger, Elodie Prince, Mohamed Benmedjahed, Adeline Payet, Marie Couralet, Didier Monté, Vincent Villeret, et al.

► To cite this version:

Clément Immarigeon, Sandra Bernat-Fabre, Emmanuelle Guillou, Alexis Verger, Elodie Prince, et al.. Mediator complex subunit Med19 binds directly GATA transcription factors and is required with Med1 for GATA-driven gene regulation in vivo. *Journal of Biological Chemistry*, 2020, 295 (39), pp.13617-13629. 10.1074/jbc.RA120.013728 . hal-02987861v1

HAL Id: hal-02987861

<https://hal.science/hal-02987861v1>

Submitted on 19 Nov 2020 (v1), last revised 1 Jul 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Mediator complex subunit Med19 binds directly GATA transcription factors and is required with Med1 for GATA-driven gene regulation *in vivo*

Clément Immarigeon¹, Sandra Bernat-Fabre¹, Emmanuelle Guillou¹, Alexis Verger² Elodie Prince^{1,3};
Mohamed A. Benmedjahed¹, Adeline Payet¹ Marie Couralet^{1,4} Didier Monte², Vincent Villeret²,
Henri-Marc Bourbon¹ and Muriel Boube^{1*}

¹Centre de Biologie Integrative CBD, UMR5547 CNRS/UPS, Université de Toulouse, 31062
Toulouse Cedex, France.

²Univ. Lille, Inserm, CHU Lille, Institut Pasteur de Lille, U 1167 – RID-AGE, CNRS ERL 9002
Integrative Structural Biology, F-59000 Lille, France

³present address : Institut de Biologie Valrose Université Nice Sophia Antipolis- Batiment de
Biochimie - Parc Valrose 06108 Nice Cedex 2

⁴present address : Institut de Pharmacologie Moléculaire et Cellulaire UMR7275 660 Rte de Lucioles
06560 Sophia Antipolis

*corresponding author: Muriel Boube
Email: muriel.boube-trey@univ-tlse3.fr

Running title: GATA transcription factors and Med19 Mediator complex subunit

Keywords : GATA transcription factors, Serpent, (Srp) Pannier (Pnr), Mediator complex, Med19, Med1, cofactor, *Drosophila* development, thoracic closure, DNA-binding domain, Activation domain

Abstract

The evolutionarily conserved multiprotein Mediator complex (MED) serves as an interface between DNA-bound transcription factors (TFs) and the RNA Pol II machinery. It has been proposed that each TF interacts with a dedicated MED subunit to induce specific transcriptional responses. But are these binary partnerships sufficient to mediate TF functions? We have previously established that the Med1 Mediator subunit serves as a cofactor of GATA TFs in *Drosophila*, as shown in mammals. Here, we observe mutant phenotype similarities between another subunit, Med19, and the *Drosophila* GATA TF Pannier (Pnr), suggesting functional interaction. We further show that Med19 physically interacts with the *Drosophila* GATA TFs, Pnr and Serpent (Srp), *in vivo* and *in vitro* through their conserved C-zinc finger domains. Moreover, Med19 loss of function experiments *in vivo* or *in cellulo* indicate that it is required for Pnr- and Srp-dependent gene expression, suggesting general GATA cofactor functions. Interestingly, Med19 but not Med1 is critical for the regulation of all tested GATA target genes, implying shared or differential use of MED subunits by GATAs depending on the target gene. Lastly, we show a direct interaction between Med19 and Med1 by GST-pull-down experiments indicating privileged contacts between these two subunits of the MED middle module. Together, these findings identify Med19/Med1 as a composite GATA TF interface and suggest that binary MED subunit - TF partnerships are probably oversimplified models. We propose several mechanisms to account for the transcriptional regulation of GATAs-targeted genes.

Introduction

Transcription, the first stage of gene expression, is a fundamental cellular process governed by the binding of sequence-specific transcription factors (TFs) at gene enhancers, inducing the recruitment/activation of the general RNA Polymerase II (Pol II) machinery at gene promoters. In eukaryotes, TFs do not bind directly the Pol II enzyme but instead

contact a multisubunit complex called Mediator (MED), serving as a physical and functional interface between DNA-bound TFs and PolII (for review see (1–3)). Whereas TF DNA-binding specificity has been largely decoded, how TFs interact with the Mediator complex has been less extensively studied, and it is not clear whether each TF binds a specific MED subunit or whether TF-MED interactions obey more complex rules.

Mediator is an evolutionarily-conserved complex composed of 25 to 30 distinct proteins distributed in four modules: head, middle, and tail forming the core MED, and a separable regulatory Cdk8 kinase module (CKM) (1). Despite a general role of the Mediator complex in regulating transcription, some MED subunits display striking functional specificities, as exemplified by their differential requirements for cell viability (4, 5), their involvement in specific human diseases (6, 7), or their roles in given developmental processes (8–10). It has been proposed that MED subunit specificity comes from their ability to contact specific transcription factors and mediate their regulatory activity (11, 12). For example, specific interactions have been demonstrated between Med15 and SMAD transcription factors in *Xenopus* (13), Med23 and RUNX2 in mice (14) Med12 and Gli3 in mammalian cells (15), Med19 and REST in mammals or Med19 and HOX developmental regulators in *Drosophila* (16, 17), or also between Med1 and hormone nuclear receptors or GATA TF families in mammalian cells (18, 19).

GATA transcription factors represent a good model to analyse interaction between TFs and Mediator subunits. The mammalian GATA TF family comprises 6 members (GATA1-6), shown to specifically interact with the Med1 Mediator subunit (20). They have conserved homologs among both vertebrates and invertebrates (21) and contain two highly-conserved zinc finger (ZF) domains. The C-terminal one (C-ZF) is both necessary and sufficient for sequence-specific DNA binding at WGATAR genomic sites, while the N-terminal ZF (N-ZF) appears only to modulate DNA-binding affinity (22) and has been involved in direct interactions with GATA cofactors (23–26). Mammalian GATAs are key

regulators of developmental processes: GATA1, -2 and -3 are crucial hematopoietic TFs while GATA4, -5 and -6 control cardiac development, among other functions (21). Interestingly, among the 5 GATA TFs encoded by the *Drosophila* genome, only Serpent (Srp), is a *bona fide* hematopoietic GATA factor, while Pannier (Pnr), is involved in cardiac development (27). Pnr activity is also crucial during central thorax patterning and dorso-central (DC) mechanosensory bristle formation, and it has been studied in depth in this context (28–30)(29). Within the wing imaginal disc, the Pnr TF directly activates proneural genes of the *achaete-scute (ac-sc)* complex in the dorso-central cluster, which gives rise to the DC bristles (28). In addition, Pnr activates the *wingless* gene in a strip of cells of the presumptive thorax (31).

In a genome-wide RNA interference screen in *Drosophila* cultured cells we identified a set of MED subunits as modulators of GATA/Serpent-induced transactivation, among which, Med12, Med13, Med1 and Med19 (32). This work further showed that Med12 and Med13 subunits are required *in vivo* for Srp-driven developmental processes, but we were unable to detect direct physical interaction with Srp *in vitro*, suggesting that GATA/Srp may recruit the Mediator complex by contacting other MED subunits. Indeed, we recently showed that Med1 mediates GATA TFs function in *Drosophila* (33). Med1 does interact physically with both Pnr and Srp GATA TFs, through their conserved zinc finger region. Furthermore, *in vivo* experiments showed that Med1 is involved in Srp-driven hematopoiesis and Pnr-driven thorax differentiation and is required for Srp and Pnr target gene expression in the corresponding tissues. These data established that the Med1 GATA cofactor activity is evolutionarily conserved and involves the GATA N- and C-zinc finger domains in both mammals and *Drosophila*. Nevertheless, we also showed that *Drosophila* Med1 is not critical for *wingless*-induced transactivation by Pnr, raising the possibility that other MED subunits could mediate some GATA TFs functions.

Here, we reveal that another MED subunit, Med19, also acts as a GATA coactivator. *Med19* mutants phenocopy *pnr* loss-of-

function, and extinguish the expression of both Pnr target genes *achaete* and *wg*, whereas Med1 mutants were previously shown to affect only *achaete* expression. Using Immunoprecipitation, pull-down and Bimolecular Fluorescence Complementation (BiFC) techniques, we establish that Med19 physically interacts with Pnr *in cellulo*, *in vivo* and *in vitro* through its C-ZF domain. Med19 also interacts physically with GATA/Srp, suggesting that Med19 acts as a generic GATA cofactor. Moreover, we show that both Med1 and Med19 jointly regulate a series of Srp target genes in *Drosophila* cultured cells. Finally, *in vitro* experiments revealed that Med1 and Med19 physically interact through the Med1 domain which is conserved throughout eukaryotes. Taken together, our results show that GATA-driven regulatory functions in *Drosophila* require two MED complex subunits, Med19 in all tested cases and Med1 in a majority. The evolutionary conservation of Med19 and GATA interacting domains suggests that Med19 may play a conserved GATA cofactor function in mammals.

Results:

***Drosophila* Med19 is required for notum morphogenesis, bristle development and GATA/Pannier target gene activation**

Our whole-genome dsRNA screen in *Drosophila* cultured cells identified Med19 as one MED subunit capable of modulating Srp TF-induced transactivation *ex vivo*. (32). This led us to ask whether and how Med19 could interact with GATA TFs *in vivo*. To this end, we generated *Med19* mutant clones in the larval wing imaginal disc, which gives rise to adult thoracic structures whose proper development depends on GATA/Pnr activity. Flies bearing *Med19* clones displayed specific phenotypes in the thorax, including thoracic cleft and loss of dorso-central (DC) mechanosensory macrochaetes (Fig.1A, D), typical of *pnr* loss-of-function (29, 30). We observed similar phenotypes upon expression of interfering RNAs (RNAi) against *Med19* in the *apterous (ap)* domain encompassing all the presumptive notum (Fig S1A-B). To investigate the functional relationship between Med19 and Pnr, we first examined *pnr* gene

expression in *Med19*-deficient wing discs by fluorescent *in situ* hybridization (FISH), and observed that *pnr* is expressed in *Med19*-depleted wing discs (Fig S1C-F). Thus, *Med19* mutant phenotypes cannot be explained by a loss of *pnr* expression. To further analyse the functional relationship between *Med19* and *Pnr*, we then examined GATA/*Pnr* TF activity in *Med19* loss-of-function clones by analyzing the expression of known *Pnr* target genes. Compared to wild-type cells shown in Fig. 1B-C", we observed that both *wingless* (*wg*) and *achaete* (*ac*) expression was cell autonomously lost in *Med19*^{-/-} cells (Fig.1E-F") indicating that *Med19* is required for the expression of both *Pnr* target genes. Note that *ac* expression has been visualized by a DC-*ac*-lacZ reporter gene which is directly activated upon *Pnr* binding to the DC *ac* enhancer (28).

These data show that *Med19* is cell autonomously required for *Pnr* activity but not for *Pnr* expression, suggesting that it could act as a GATA/*Pnr* cofactor.

***Drosophila Med19* interacts physically with the Pannier GATA TF**

We investigated whether GATA/*Pnr* transcription factor and *Med19* physically interact by using three independent experimental approaches: co-immunoprecipitation (co-IP) from cultured cells, *in vitro* pull-down and *in vivo* Bimolecular Fluorescence Complementation (BiFC) interaction tests. We first tested whether *Pnr*-MED complexes actually form within *Drosophila* cells by performing co-Immunoprecipitations (co-IP) experiments on total protein extracts from cultured cells expressing a functional Myc-tagged *Pnr* form. We observed that *Pnr* co-precipitated with endogenous *Drosophila Med19* (Fig.2A). In the reverse experiment, endogenous *Med19* protein co-precipitated with Myc-tagged *Pnr* protein (Fig.2B). These data provide complementary evidence for the formation of *Med19*-GATA complexes in *Drosophila* cells.

To investigate whether *Med19*-*Pnr* interaction is direct, we tested the ability of *Med19* and *Pnr* proteins to bind each other physically *in vitro* through pulldown assays with

Glutathione S-transferase (GST) fusion proteins. *In vitro*-produced *Med19* readily bound full-length recombinant GST-*Pnr* (Fig.2C), and *vice versa* (Fig.2D). These results show that *Med19* and *Pnr* can interact physically in the absence of any other *Drosophila* MED subunits.

We then used Bimolecular Fluorescence complementation (BiFC) (16, 34) to analyse *Med19*-GATA interaction *in vivo*. Based on fusing N- and C-terminal portions (VN and VC) of the GFP-variant Venus protein, with two proteins of interest respectively, this technique allows the reconstitution of a fluorescent Venus protein if the two candidate proteins are close enough within the cell. We used the *dpp*GAL4 driver (Gal4/UAS system (35)) to co-express VN-*Pnr* with either *Med19*-VC or another MED subunit fusion, *CycC*-VC, along the antero/posterior (A/P) frontier of the wing imaginal disc (Fig2.E, F). The co-expression of VN-*Pnr* and *Med19*-VC resulted in a clear BiFC signal, whereas the control VN-*Pnr* /*CycC*-VC combination gave a very low signal (Fig.2E'-F'), even though *CycC*-VC and *Med19*-VC proteins were expressed at similar levels (Fig. 2E-F). These data indicate that the BiFC technique discriminates specific interactions between different subunits within the MED complex and that *Med19* and *Pnr* are in close proximity in the nucleus of living cells. Of note, the BiFC signal was observed in the entire *dpp*GAL4 expression domain including the wing pouch where endogenous GATA/*Pnr* is not expressed, showing that the *Pnr*-*Med19* interaction can occur at ectopic locations independently of tissue-specific *Pnr* partners, thus providing further support for a direct molecular interaction *in vivo*.

Collectively, *in cellulo*, *in vitro* and *in vivo* data support a direct physical interaction between the GATA/*Pnr* transcription factor and the *Med19* Mediator subunit. Together with our previous results (33), these data suggest that the *Pnr* TF can interact with the entire MED complex via a direct molecular contact with the *Med19* subunit in addition to or in place of the *Med1* subunit.

Med19* Core and HIM domains bind the C-zinc finger domain of GATA/*Pnr

We previously showed that Med1 directly interacts with the dual Zinc Finger domains of Pnr (33). We therefore decided to characterize interacting domains within Pnr and Med19 in order to determine whether Med19 and Med1 interact with the same Pnr domain.

We first looked for the Med19-interacting domain(s) within the GATA/Pnr protein using full length GST-Med19 as a bait (Fig.3A). Pnr was split into three parts: the poorly evolutionarily-conserved N-terminal region (amino acids (aa) 1-137), the strongly-conserved central region spanning the 2 zinc fingers, N- and C-ZF, and the divergent C-terminal region containing two amphipathic alpha helices, H1 and H2. Only the ZF-containing region (aa 130-278) displayed significant binding. When cutting full-length Pnr into two halves separating the two zinc finger domains, binding was observed only with the C-ZF-containing part (Fig. 3A), suggesting that C-ZF mediates binding of Pnr to Med19. Consistently, the ability of the N-ZF-containing half of Pnr to bind Med19 was recovered when we added back the C-ZF proper (aa 220-253) containing the four zinc-chelating cysteines forming the finger structure. Interestingly, binding was increased when the C-ZF proper was extended by its neighboring C-terminal 25 amino acids (basic tail motif, aa 253-278; Fig. 3B). Sequence alignment of *Drosophila* and mammalian GATAs indicates that the C-ZF basic tail has been strongly conserved during evolution, especially at positions shown to participate in DNA binding (open circles in Fig. 3B, (36, 37)). Together, these experiments indicate that the entire Pnr C-ZF domain, zinc-finger proper and adjacent basic tail, is necessary for optimal Med19 binding.

In the reciprocal experiment, we identified the GATA/Pnr interacting domain within Med19. Our prior analysis of *Drosophila* Med19 function and evolutionary conservation within the eukaryotic kingdom (16, 38), allowed to define four structural domains: a conserved MED-anchoring “CORE” region, an animal-specific basic HOX homeodomain-interacting motif (HIM) and two less well conserved N- and C-terminal regions. To investigate which protein domain(s) is (are) required for Pnr binding, we tested the ability of *in vitro* translated Pnr¹⁻²⁹¹ - to bind a series of GST-

Med19 truncated forms (Fig.3C-G). A Med19 protein deleted for its evolutionarily-conserved CORE domain (Δ CORE) still bound Pnr¹⁻²⁹¹ (Fig.3D). Binding was also retained after truncating both C-ter and HIM domains, but was abolished if the deletion included the C-terminal end of the CORE domain (aa 126 to 165) (Fig.3E). Deletions starting from the Med19 N-terminus indicated that a truncated protein containing HIM and Cter domains also interacts with Pnr 1-291 (Fig.3F). Further deletions revealed that one fragment of HIM from aa 206 to 220 was critical for Pnr binding in the absence of the CORE domain. Taken together, our data suggest the presence of two Pnr binding sites within Med19, aa 126 to 165 of the CORE (BS1) and aa 206 to 220 of the HIM domain (BS2) (Fig 3C). To further assess their implication, we deleted both BS1 and BS2 in an otherwise full-length Med19 protein. As shown in Fig.3G, the Δ BS1-BS2 mutant no longer bound GATA/Pnr, indicating that the presence of either BS1 or BS2 is essential for interacting with Pnr.

In conclusion, our *in vitro* binding assays indicate that the GATA/Pnr C-zinc finger domain, including its basic tail, binds two separate domains within the evolutionarily-conserved Med19 CORE and HIM regions. Med19 appears only to bind the C-ZF, whereas Med1 has been shown to bind both GATA/Pnr C-ZF or N-ZF domains *in vitro* (33).

Med19 also physically interacts with GATA/Srp

To investigate whether Med19 is a general GATA cofactor, we tested if it is able to interact with Serpent (Srp), another *Drosophila* GATA TF family member. First, we used similar GST-pull-down assays (Fig.4A). They showed that recombinant GST-Med19 protein bound *in vitro*-translated full-length GATA/Srp protein. As previously shown for GATA/Pnr, when assaying Srp truncated forms, binding was only retained with the ZF-containing middle part. Splitting the Srp protein in two halves and separating both zinc finger domains indicated that only the C-ZF is involved in binding Med19 (Fig 4A), as it is also the case for GATA/Pnr (Fig 3E).

To test whether Med19-Srp interaction also occurs *in vivo*, we used again the BiFC experimental approach. Upon expression of Med19-VC with VN-tagged Srp in the *dpp* expression domain, we observed a strong BiFC signal, similar to what we obtained with Pnr-VN (Fig. 4B), indicating that Med19 and GATA Srp indeed interact *in vivo*.

Altogether, these results show that, Med19 interacts *in vivo* and, *in vitro* with both Pnr and Srp, suggesting that Med19 is a general GATA cofactor. This interactions occurs via the GATA family-defining C-zinc finger domain. .

Med19 shares GATA-cofactor functions with the Med1 Mediator subunit

We previously showed that Med1, another subunit of the MED middle module, is required for Pnr and Srp TF activities *in vivo* and interacts directly with Srp and Pnr, in this case through both their N- and C-zinc finger domains (33). Our new data showing that Med19 can also act as GATA cofactor thus raises the question of the respective roles of Med19 and Med1 in GATA-driven transcriptional regulation. Concerning Pnr-dependent transcriptional activity, we have shown that Med19 is cell autonomously required for DC-*ac-lacZ* reporter expression like Med1, whereas *wg* expression requires Med19, but not Med1. This prompted us to consider each GATA target gene as a particular case that could involve interaction with both MED subunits or with Med19 or possibly Med1 alone. Kuuluvainen and collaborators (39) identified a set of Srp target genes in *Drosophila* S2 cells, which can be used to test the impact of *Med1*- or *Med19* mRNA depletion. Here, we quantified the expression of 6 Srp target genes: *SrCl*, *CG14629*, *CG8157*, *arg*, and *CG34417* which are activated (positive targets), and *CG13252* which is repressed by Srp, using Real Time quantitative PCR (qRT-PCR) in control, *Med1*- or *Med19*-mRNA depleted S2 cells. Quantification of mRNAs coding for the Myosin light chain (*Mlc2*) served as a control for housekeeping transcription. As shown in Fig. 5A, in cells depleted for *Med19* mRNA, expression of the 5 activated Srp target genes *SrCl*, *CG14629*, *CG8157*, *arg* and *CG34417* was significantly down-regulated and the Srp-

repressed target gene *CG13252* was instead up regulated, indicating that Med19 is required *in cellulo* for GATA/Srp transcriptional activity. In cells depleted for the *Med1* transcript (Fig. 5B), expression of Srp target genes followed the same trend than after *Med19* mRNA depletion, although less efficiently.

Several conclusions can be drawn from these experiments: (i) contrary to other MED components, Med1 and Med19 are not required for general PolII-dependent transcription given that some genes are unchanged or even upregulated. (ii) Med19 and Med1 are both required for Srp-mediated gene regulation in cultured cells, seemingly on the same target genes, and (iii) both for activation and repression.

Med19 and Med1 can interact directly

Given the functional implication of both Mediator subunits in mediating GATA activity, we lastly asked whether Med1 and Med19 proteins are able to interact physically using GST pulldown assays. As shown in Fig.5C, a GST fusion of Med1 largest isoform A (Med1^A) bound *in vitro*-translated full-length Med19. In the reverse experiment, purified GST-Med19 also bound *in vitro* translated Med1^A (Fig.5D), showing that both proteins indeed bind to each other *in vitro*, in absence of other MED subunits.

We next sought to identify Med19-interacting domain(s) within the Med1 isoforms by analysing truncated proteins. As shown in Fig.5D, the Med19 interacting domain lies within the evolutionarily-conserved Med1 N-terminal part, which has been proposed to be required for its incorporation within the MED complex (40) and is shared by the three Med1 isoforms. Conversely, Med1 isoform-specific parts are not required for Med19 interaction.

Taken together, GST-pulldown data reveal a direct interaction between Med1 and Med19 which could not be anticipated from structural data (see Fig.6) given that Med1 and Med19 have been proposed to lie in opposite parts of the Mediator complex middle module (41).

Discussion

In this work, we establish using molecular, cellular and genetic analyses that *Drosophila*

GATA factors' transcriptional activity depends on the Mediator complex subunit Med19, in addition to the previously characterized Med1 cofactor (33). Four main conclusions that are discussed below can be drawn from our results (i) Med19 interacts with the GATA C-ZF domain which also serves as the GATA DNA-binding domain (ii) Med19 and GATA interacting domains are evolutionary conserved suggesting conserved Med19 cofactor functions in higher metazoans. (iii) Comparative analysis of Med19 and Med1 depletion indicates that Med19 but not Med1 is systematically required for GATA target gene expression suggesting a differential use of MED subunits by GATAs depending on the target gene. (iv) Med1 and Med19 interact *in vitro*. Taken together our data allows us to propose new models of Mediator complex mechanism of action.

Overlapping DNA-Binding and Activation domains of GATA TFs

TFs minimally contain two domains: the DNA binding domains (DBD), which have been extensively studied and allowed to define different TF families, and transcriptional activation domains (TAD), which link TFs to the RNA polymerase II machinery, and whose structure and characteristics are less well defined. GATA TFs are characterised by the presence of two ZFs which were, so far, thought to play distinct roles. While the C-ZF appeared to be dedicated to DNA binding, the N-ZF was shown to bind co-activators such as dLMO (42) and FOG (43). Our present data show that Med19 interacts specifically with the Pnr C-ZF (Fig. 6A). Full interaction requires both the zinc finger and its adjacent basic tail which also contributes to DNA binding (36, 37). It is the first evidence that the *Drosophila* GATA C-ZF may play a dual role, in DNA binding and as an interface with MED subunit(s). Interestingly, the analysis of GATA ZF evolutionary conservation indicates that N- and C-ZF domains comes from a duplication event of the C-ZF with its basic tail (44). Thus, this transactivation function of GATAs' DBD might represent an ancestral GATA function allowing minimal primitive GATAs, essentially composed of the DBD, to connect the MED complex and thus recruit the transcriptional machinery to regulate its target

genes. They provide rationale why slightly-extended GATA ZF domains are in some cases sufficient for transcriptional activities *in vivo* (45).

This dual activity of DBD is not restricted to GATA factors. We have previously shown that HOX TFs also contact Med19 through their DNA-binding homeodomain (16). Our data also corroborate results from a recent high-throughput approach, looking for trans-activation domains of *Drosophila* transcription factors. This work shows that trans-activation domains of several zinc-finger- (ZF-) and basic Helix-Loop-Helix- (bHLH-) TFs overlap structured DNA-binding domains (46). Altogether, these results identify a novel class of TF characterized by overlapping TAD and DBD and suggest an emerging Med19 property as a dedicated cofactor directly connecting these TFs DNA-binding domains to the general PolII transcriptional machinery.

How is this dual function of DBDs achieved? Do DNA binding and transactivation functions use distinct or shared molecular determinants? Recent improvements of electron microscopy analyses could allow characterizing GATA molecular residues involved in MED- versus DNA-binding to try to separate the GATA DNA binding- from GATA transactivation functions.

Evolutionarily-conserved GATA-coactivator functions of Med19?

While Med1 is a known GATA cofactor both in mammals and in *Drosophila* (19, 33), the role of Med19 in mediating GATA transcription regulatory properties had never been investigated until now. Here we show that *Drosophila* Med19 binds GATA factors, via motifs lying within the evolutionary-conserved Med19 CORE and HIM domains (16). Both of these domains bind to the C-ZF domain of GATAs, which is a hallmark of GATA TF family suggesting that interaction with Med19 is likely to be conserved in mammals. Yet, Med1 depletion experiments in mammalian cultured cells induces defects in only a subset of GATA1-activated genes and does not prevent GATA1-dependent repression (47, 48). Furthermore, in studies of the different blood cell types produced by conditional *Med1*

knock-out mice, Med1 appears to be critical for erythroid lineages which depend upon GATA1-function but is dispensable for hematopoietic stem cell production and T-cell development which require GATA2 and GATA3, respectively (49). Thus, despite being capable of binding all GATA factors *in vitro*, Med1 is not critical for all GATA functions, which suggests that (an) other MED subunit(s) also bind(s) GATAs to relay their regulatory signals to the PolII machinery. Considering the evolutionary-conservation of interaction motifs within both GATAs and Med19, we argue that Med19 is a strong candidate as a GATA cofactor in mammals.

New models for GATA – MED interactions

Our data show that most *Drosophila* GATA target genes require both Med19 and Med1. How does this work? We showed that Med19 only interacts with the C-ZF domain, but Med1 can bind both GATA zinc-finger domains (33), suggesting that Med1 and Med19 can simultaneously bind GATA factors (Fig. 6A). We thus propose that in the majority of cases where GATA-driven gene expression requires both Med19 and Med1, (Fig. 6B), enhancer-bound GATAs must directly contact both Med1 and Med19 subunits to recruit the Mediator complex and thus the PolII machinery at GATA target genes.

Some genes (e.g. wingless) require Med19 but not Med1. How does this kind of gene specificity occur mechanistically? We hypothesize that for these Med1-independent genes, other transcription factors might be involved in recruiting the MED (through other subunits) and hence overcome the necessity for Med1-GATA interaction (see Figure 6C).

A future challenge will be to test these models by site-directed mutagenesis *in vivo* to assess the functional contribution of each GATA-MED contact. Nevertheless, this task is complicated because of overlapping DNA- and MED- interacting domains within GATAs (see above). On the other hand, Med19 CORE domain has also a dual function of MED anchorage and interaction with GATA. It thus requires prior structural analysis of molecular contacts to specifically target GATA-MED interaction without affecting essential DNA-

binding activity of GATAs or Med19 ability to incorporate the MED-complex.

Another interpretation of our results could be that other subunits necessary for GATA target gene expression fall off from the complex when Med19 or Med1 are deleted or knockdown. However, structural analyses of MED complexes from yeast and mammalian cells lacking Med19 or Med1 indicate that global MED organization is unchanged (4, 50) and we therefore considered it very unlikely. Other lines of evidence indicate that complexes missing only Med19 can be isolated from Med19 depleted mammalian or yeast cells (51, 52). Altogether, these results suggest that *Drosophila* MED subunit loss is unlikely in Med19 or Med1 depletion conditions. Since we found a direct physical interaction between GATAs and Med1/Med19, we consider that the simplest explanation for our results is that the loss of either of these subunits is enough to abolish GATA regulatory signals.

Previous models of core MED structure-function analysis suggested that the middle and head modules contact the PolII enzyme and associated general transcription factors (GTFs) while the tail module interacts with sequence-specific TFs (2, 3). Our data show that two MED subunits of the middle module, Med1 and Med19, are able to bind GATA factors and are required for their function. They emphasize that MED should be viewed as a much more complex interface using multiple MED subunits to contact different TF combinations thus mediating specific transcriptional responses.

Unexpected direct interaction between Med1 and Med19 middle module subunits

Modelization of MED spatial organization indicate that Med1 and Med19 are most likely located at two opposite ends of the middle module, Med1 near the tail module and Med19 within the so called “hook” domain proposed to anchor the separable CDK8 module (CKM) (4, 53)(Fig 6.A). Nevertheless, our data indicate that Med1 and Med19 interact *in vitro*. Furthermore, this interaction occurs via the highly-conserved, N-terminal, MED-addressing domain of Med1 suggesting an evolutionary conservation. How, then,

reconcile the proposed MED architecture with our results showing a direct interaction between Med1 and Med19 subunits *in vitro*?

We propose two non-exclusive hypotheses: First, MED complexes could adopt different conformations, which would differ from the “canonical” architecture of the MED complex in isolation. This is supported by observations that the MED complex changes its overall shape when engaged in interactions with either TF, CKM or PolII (50). Perhaps when MED is recruited by GATA, Med1 - Med19 contacts within the MED complex could stabilize one of these “alternative” conformations.

A second possibility is that Med1-Med19 interactions do not occur within but between MED complexes and could thus stabilize “multi-MED” structures. It has been shown that master TFs control gene expression programs by establishing clusters of enhancers called super-enhancers, at genes with prominent roles in cell identity (54). Recent studies have revealed that, at super-enhancers, master TFs and the Mediator coactivator form phase-separated condensates, which compartmentalize and concentrate the PolII machinery to specific nuclear foci, to ensure high level of transcription (55–57). Interestingly, mammalian Med1 can form such phase-separated droplets that concentrate the transcription machinery at super-enhancers (56). Bringing together several MED complexes associated with TFs via Med1-Med19 trans-interaction might thus help phase-separated droplet formation at clustered gene enhancers and ensure high transcriptional level (Fig 6.D).

In conclusion, our work shows that 2 MED subunits physically bind GATAs and are required to relay the regulatory signals from common TFs. This argues against the generally admitted view of binary interaction between one MED subunit and one TF, which appears as an oversimplified model for MED action. The Mediator should be viewed as a complex interface allowing fine-tuned gene regulation by TFs through specific contacts with different MED subunit combinations. This study highlights the unexpected role of *Drosophila* Med19 as a GATA cofactor and Med1 interactor. This work sheds new light on the GATA-MED paradigm and suggests novel

means by which several MED subunits might collaborate to regulate gene transcription.

Experimental procedures:

Drosophila stocks, genetic mosaics and phenotypic analyses

No vertebrate animals were used in this study. No applicable regulations are available for *Drosophila* but this animal study followed guidelines from the Animal Ethic Committee of CNRS. Each experiment with *Drosophila* was performed with at least two independent replicates. There were no pre-determined exclusion criteria for animal work.

Stocks and crosses were raised at 25°C or 22°C and 12:12-h light–dark cycle on standard yeast-agar-cornmeal medium (Bloomington *Drosophila* Stock Center). Mitotic clones were generated using the Flp-FRT system with the *Med19*²null allele FRT80B chromosome (16). The Flp recombinase was expressed in the dorsal part of the wing using the *ap*-GAL4 driver recombined with UAS-Flp. The following stocks were used: *ap*GAL4 UASFlp/Cyo; *Med19*² FRT2A/TM6B, UbGFP M FRT2A/TM6B; *ap*GAL4^{#MD544} UASGFP/CyO, UAS dsRNA *Med19*^{#27559} and *dpp*GAL4 from BDSC, UAS VN-AbdA and UAS VN-Srp kindly provided by S. Merabet, DC-ac-lacZ kindly provided by P. Heitzler UAS-Med19VC (16) and UAS-Med1^ΔVC (33). Phenotypes of approximately 20 adults/genotype were analysed by scanning electron microscopy (Hitachi TM-1000 Tabletop model) of frozen adults.

Immunofluorescence and fluorescent *in situ* hybridization (FISH) of wing discs:

Approximately 30 third-instar larval imaginal discs were prepared and stained using standard procedures. Antibodies used were Rabbit anti-βgal (Cappel, 1:2500) for both DC-ac-LacZ and UAS-lacZ detection; Mouse anti Wg (1:200) (4D4 antibody from Developmental Studies Hybridoma Bank/Iowa University). For BiFC experiments, we used mouse anti-GFP from Roche (1:200) which only recognize the C terminal half of the GFP variant Venus (ΔVC).

For FISH, DIG-U labeled antisense RNA probes against *pannier*, spanning the whole Pnr ORF were used. Fast-RED revelation was optimal when carried out at 4°C overnight.

Image acquisition was performed on a Leica SP5 confocal microscope. Z-stacks were generated using ImageJ-related FiJi software.

BiFC assay:

This technique is based on expressing *in vivo* two candidate partner proteins with the N- and C-terminal portions (VN and VC) of the Venus protein in order to test the reconstitution of a functional fluorescent protein. UAS-Pnr-VN, UAS-VN-Pnr, and UAS CycC-VC lines were generated by inserting Pnr^A or CycC ORF in phase with VN173 (aa 1-172) or VC155 (aa 155-238) ORF in a recipient pUAST-attB plasmid, allowing site specific insertion. attP-carrying embryos expressing PhiC31 integrase in the presumptive germline were injected with these plasmids: Pnr constructions were inserted on the X chromosome (attP ZH-2A), and CycC constructions on the 2nd chromosome (attP 51D) to ensure identical expression of the different VN lines, and easy combination of VN and VC lines. Note that the BiFC constructions were functionally validated for their ability to rescue mutant lethality for Med19-VC, Med1-VC and CycC-VC or to produce typical gain-of-function phenotypes for VN-Pnr.

Crosses were carried out at 22°C for interaction tests to express candidate proteins at homogenous and relatively low levels in order to avoid non-specific signal. We used the *dpp*GAL4 driver (Gal4/UAS system) to direct co-expression of fusion proteins at the antero-posterior frontier of the wing imaginal disc both in the thorax where *pnr* is normally expressed and in the wing pouch region that does not express *pnr*.

Image acquisition was performed on a Leica SP5 using the same settings and number of z slices for the different genetic contexts. BiFC fluorescence was quantified using ImageJ software.

Co-immunoprecipitation experiments:

Cultured *Drosophila* S2 cells were grown in 10% serum containing Schneider's medium at 25°C, and transfected using FuGENE HD transfection reagent (Roche) following manufacturer recommendations. Transfections of 18 x 10⁶ cells per plate were carried out with *pActin-Myc-Pnr* (3µg) and *pActin-GAL4* (2µg) plasmids. The cell harvest, protein extraction and IP were performed as described in (33)

with the following modifications: the buffer used for protein extraction and subsequent IP contained 0,1% NP40 instead of 0,5% to increase purification of large complexes such as MED; 1mg of total protein extract was used for each IP instead of 1,5mg. Anti-Med19 and non-relevant (NR) IPs were performed with 5µl of decomplexed serum from a Med19-immunized guinea pig (16) or 5µl of the same animal's pre-immune serum, respectively. We used 10µl G-protein coupled Sepharose beads per IP (SIGMA, P3296). Anti-Myc IPs were performed with 10µl Anti-Myc-Agarose bead (SIGMA, A7470).

Med19 was revealed using a home-made polyclonal serum from guinea-pig (diluted 1:500) immunized with a full length Med19-GST fusion protein (16). Antibody specificity has been assessed by immunodetection from Med19-depleted (see Fig S1) or Med19 overexpressing tissues ; Myc-Pnr was detected with rabbit anti-Pnr (kind gift from G. Morata, diluted 1:1000). We used Lumi-Light^{PLUS} western blotting substrate (ROCHE, 12015196001) and high performance chemiluminescence film (Amersham HyperfilmTM ECL, GE healthcare, 28906837) for revelation.

GST-pulldown experiments:

Preparation of GST fusion proteins, ³⁵S-Methionine-labeled proteins and pulldown were performed essentially as described in (Mojica et al 2017). Med1, Pannier and Serpent proteins or sub-fragments have been produced from cDNA corresponding to Pnr^A (Romain et al 1993), Srp^B (Waltzer et al 2002) and Med1^A (Immarigeon et al 2019) by *in vitro* transcription/translation coupled reactions using rabbit reticulocyte extracts (TnT-Promega) isoforms labeled. cDNA encoding full-length dMed19 and deletion derivatives were amplified by PCR using appropriated oligonucleotides and inserted into the *Bam*HI/*Not*I site of the pGEX-6P1 vector (GE Healthcare). Pnr aa 1-291 fragment-encoding was amplified by PCR and cloned into pcDNA3 vector, with an HA-tag at the C-terminus and a Flag-tag at the N-terminus. All clones were verified by sequencing. Primers sequences and complete clone sequences are available upon request. Bacterial expression vectors pGEX-6P1 were transformed in competent *E. coli* strain BL21 (DE3). The transformed cells were plated in LB agar

containing 50 µg/ml of ampicillin. A single colony was grown overnight in 25 ml of LB medium containing ampicillin on a rotary shaker (180 rpm) at 37°C. Overnight starter culture was diluted 1:30 and bacteria were grown in 150 mL of LB medium containing ampicillin at 37°C to an optical density of 0.8-0.9 at 600 nm and expression was induced with 0.5 mM IPTG for 2 h at 37°C. Bacteria were pelleted by centrifugation and pellets were stored overnight at - 20°C. Pellet was resuspended in 15 ml lysis buffer (50 mM Tris-HCl, pH 8.0, 100 mM NaCl, 10 % w/v glycerol, 0.1 % Nonidet-P40) including one Complete™ EDTA-free protease inhibitor tablet and sonicated on ice. After centrifugation at 10 000 x g 45 minutes at 4°C, the supernatant was mixed 2 h at 4°C on a rotating platform with 2 ml Glutathione Sepharose 4B resin. Beads were washed four times with lysis buffer and stored at 4°C.

6µl of tagged Pnr 1-291-HA *in vitro* translation product was mixed with 50 µl of glutathione-agarose bead-GST Med19 derivatives in 200 µl of pull-down buffer (50 mM Tris-HCl, pH 8.0, 100 mM NaCl, 10 % w/v glycerol, 0.1 % Nonidet-P40, 10 µM ZnSO₄). The mixture was incubated for 2 h at 4°C, washed four times with 500 µl pull-down buffer. Protein complexes were eluted from the beads with 2X Laemmli sample buffer, boiled 5 min and separated by SDS-PAGE on Mini-PROTEAN® TGX™ precast gels (Bio-Rad). Bound Pnr 1-291 was identified by Western-blot (1:5000 rabbit anti-HA polyclonal) using an ECL kit (Amersham GE Healthcare Life Sciences) based on the manufacturer's recommendations. HRP-conjugated secondary antibodies were used at 1:5000 and were purchased from Amersham GE Healthcare Life Sciences.

Data Availability

All data are contained within this manuscript

Acknowledgements

We gratefully acknowledge P. Heitzler, S. Merabet and BDSC for plasmids and fly stocks, D. Jullien, A. Vincent and C. Monod for critical reading of the manuscript, V. Gobert and B. Augé for technical help, J. Favier, A. Destenabes and V. Nicolas for fly facilities and finally the Toulouse Rio Imaging platform.

Fundings and additional information

This work was supported by grants from the French “Ministère de l'Enseignement et de la Recherche” (C.I.), the Ligue Nationale contre le Cancer (C.I. and A.P.), the Association pour la Recherche sur le Cancer (ARC PJA 20141201932), the Agence Nationale de Recherche (ANR-16 CE12-0021-01), and we benefitted from the support of the Centre National de Recherche Scientifique (CNRS) and Toulouse III University.

Sciences. All the membranes were scanned on an ImageQuant LAS 500 (GE Healthcare Life Sciences).

Sequence conservation analysis

C-ZF domains from the *Drosophila* and human GATA family members were extracted from the NCBI web site (<https://www.ncbi.nlm.nih.gov/protein/>), aligned with the MAFFT software (<https://mafft.cbrc.jp/alignment/server/>), using default parameters, and amino acid conservation were visualized with Jalview 2.10.5 version (<http://www.jalview.org/>) using Clustal coloring.

RT-qPCR:

Two different dsRNA were used for *Med1*- or *Med19* mRNA depletion, only one for control GFP mRNA. The indicated dsRNAs (see Table I) is added at 2µg/ml to exponentially growing S2 cells, in an orbital shaker, at 2 10⁶ cell/ml in serum-free medium. After 40min, serum is added. 24h later a second addition of dsRNA is done at 1µg/ml. Cells are collected 5 days after the first dsRNA treatment.

For mRNA quantification, mRNAs were purified by RNeasy kit (Qiagen). Reverse transcription was done using SuperScript™ II Reverse Transcriptase (Thermo Fisher Scientific) and cDNA were quantified by real-time qPCR (CFX Bio-Rad) with specific oligonucleotides (Table I). Absolute quantification of each mRNA was normalized to *GAPDH* mRNA quantity in the same sample. mRNA measured in cells treated with a control dsRNA GFP was set at 100% to compare with cells treated with a dsRNA against Med1 or Med19.

Conflict of Interest: The authors declare no conflicts of interest in regards to this manuscript.

References

1. Soutourina, J. (2018) Transcription regulation by the Mediator complex. *Nat. Rev. Mol. Cell Biol.* **19**, 262–274
2. Jeronimo, C., and Robert, F. (2017) The Mediator Complex: At the Nexus of RNA Polymerase II Transcription. *Trends Cell Biol.* **27**, 765–783
3. Verger, A., Monté, D., and Villeret, V. (2019) Twenty years of Mediator complex structural studies. *Biochem. Soc. Trans.* **47**, 399–410
4. El Khattabi, L., Zhao, H., Kalchschmidt, J., Young, N., Jung, S., Van Blerkom, P., Kieffer-Kwon, P., Kieffer-Kwon, K. R., Park, S., Wang, X., Krebs, J., Tripathi, S., Sakabe, N., Sobreira, D. R., Huang, S. C., Rao, S. S. P., Pruett, N., Chauss, D., Sadler, E., Lopez, A., Nóbrega, M. A., Aiden, E. L., Asturias, F. J., and Casellas, R. (2019) A Pliable Mediator Acts as a Functional Rather Than an Architectural Bridge between Promoters and Enhancers. *Cell.* **178**, 1145–1158.e20
5. Boube, M., Faucher, C., Joulia, L., Cribbs, D. L., and Bourbon, H. M. (2000) Drosophila homologs of transcriptional mediator complex subunits are required for adult cell and segment identity specification. *Genes Dev.* **14**, 2906–2917
6. Schiano, C., Casamassimi, A., Rienzo, M., de Nigris, F., Sommese, L., and Napoli, C. (2014) Involvement of Mediator complex in malignancy. *Biochim. Biophys. Acta - Rev. Cancer.* **1845**, 66–83
7. Berk, A. J. (2012) Yin and yang of mediator function revealed by human mutants. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 19519–19520
8. Grants, J. M., Goh, G. Y. S., and Taubert, S. (2015) The Mediator complex of *Caenorhabditis elegans*: Insights into the developmental and physiological roles of a conserved transcriptional coregulator. *Nucleic Acids Res.* **43**, 2442–2453
9. Loncle, N., Boube, M., Joulia, L., Boschiero, C., Werner, M., Cribbs, D. L., and Bourbon, H. M. (2007) Distinct roles for Mediator Cdk8 module subunits in *Drosophila* development. *EMBO J.* **26**, 1045–1054
10. Malik, N., Agarwal, P., and Tyagi, A. (2017) Emerging functions of multi-protein complex Mediator with special emphasis on plants. *Crit. Rev. Biochem. Mol. Biol.* **52**, 475–502
11. Borggreffe, T., and Yue, X. (2011) Interactions between subunits of the Mediator complex with gene-specific transcription factors. *Semin. Cell Dev. Biol.* **22**, 759–768
12. Yin, J. W., and Wang, G. (2014) The Mediator complex: A master coordinator of transcription and cell lineage development. *Dev.* **141**, 977–987
13. Kato, Y., Habas, R., Katsuyama, Y., Näär, A. M., and He, X. (2002) A component of the ARC/mediator complex required for TGF β /nodal signalling. *Nature.* **418**, 641–646
14. Liu, Z., Yao, X., Yan, G., Xu, Y., Yan, J., Zou, W., and Wang, G. (2016) Mediator MED23 cooperates with RUNX2 to drive osteoblast differentiation and bone development. *Nat. Commun.* **7**, 1–11
15. Zhou, H., Kim, S., Ishii, S., and Boyer, T. G. (2006) Mediator Modulates Gli3-Dependent Sonic Hedgehog Signaling. *Mol. Cell. Biol.* **26**, 8667–8682
16. Boube, M., Hudry, B., Immarigeon, C., Carrier, Y., Bernat-Fabre, S., Merabet, S., Graba, Y., Bourbon, H. M., and Cribbs, D. L. (2014) *Drosophila melanogaster* Hox Transcription Factors Access the RNA Polymerase II Machinery through Direct Homeodomain Binding to a Conserved Motif of Mediator Subunit Med19. *PLoS Genet.* 10.1371/journal.pgen.1004303
17. Crona, F., Holmqvist, P. H., Tang, M., Singla, B., Vakifahmetoglu-Norberg, H., Fantur, K., and Mannervik, M. (2015) The Brakeless co-regulator can directly activate and repress transcription in early *Drosophila* embryos. *Dev. Biol.* **407**, 173–181
18. Fondell, J. D. (2013) The Mediator complex in thyroid hormone receptor action. *Biochim. Biophys. Acta - Gen. Subj.* **1830**, 3867–3875
19. Stumpf, M., Waskow, C., Krötschel, M., Van Essen, D., Rodriguez, P., Zhang, X., Guyot, B., Roeder, R. G., and Borggreffe, T. (2006) The mediator complex functions as a coactivator for GATA-1 in erythropoiesis via subunit Med1/TRAP220. *Proc. Natl. Acad. Sci. U. S. A.* **103**,

- 18504–18509
20. Crawford, S. E., Qi, C., Misra, P., Stellmach, V., Sambasiva Rao, M., Enge, J. D., Zhu, Y., and Reddy, J. K. (2002) Defects of the heart, eye, and megakaryocytes in peroxisome proliferator activator receptor-binding protein (PBP) null embryos implicate GATA family of transcription factors. *J. Biol. Chem.* **277**, 3585–3592
 21. Chlon, T. M., and Crispino, J. D. (2012) Combinatorial regulation of tissue specification by GATA and FOG factors. *Dev.* **139**, 3905–3916
 22. Whyatt, D. J., deBoer, E., and Grosveld, F. (1993) The two zinc finger-like domains of GATA-1 have different DNA binding specificities. *EMBO J.* **12**, 4993–5005
 23. Tsang, A. P., Visvader, J. E., Turner, C. A., Fujiwara, Y., Yu, C., Weiss, M. J., Crossley, M., and Orkin, S. H. (1997) FOG, a multitype zinc finger protein, acts as a cofactor for transcription factor GATA-1 in erythroid and megakaryocytic differentiation. *Cell.* **90**, 109–119
 24. Rodriguez, P., Bonte, E., Krijgsveld, J., Kolodziej, K. E., Guyot, B., Heck, A. J. R., Vyas, P., De Boer, E., Grosveld, F., and Strouboulis, J. (2005) GATA-1 forms distinct activating and repressive complexes in erythroid cells. *EMBO J.* **24**, 2354–2366
 25. Tripic, T., Deng, W., Cheng, Y., Zhang, Y., Vakoc, C. R., Gregory, G. D., Hardison, R. C., and Blobel, G. A. (2009) SCL and associated proteins distinguish active from repressive GATA transcription factor complexes. *Blood.* **113**, 2191–2201
 26. Wilkinson-White, L., Gamsjaeger, R., Dastmalchi, S., Wienert, B., Stokes, P. H., Crossley, M., Mackay, J. P., and Matthews, J. M. (2011) Structural basis of simultaneous recruitment of the transcriptional regulators LMO2 and FOG1/ZFPM1 by the transcription factor GATA1. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 14443–14448
 27. Sorrentino, R. P., Gajewski, K. M., and Schulz, R. A. (2005) GATA factors in Drosophila heart and blood cell development. *Semin. Cell Dev. Biol.* **16**, 107–116
 28. García-García, M. J., Ramain, P., Simpson, P., and Modolell, J. (1999) Different contributions of pannier and wingless to the patterning of the dorsal mesothorax of Drosophila. *Development.* **126**, 3523–3532
 29. Ramain, P., Heitzler, P., Haenlin, M., and Simpson, P. (1993) Pannier, a negative regulator of achaete and scute in Drosophila, encodes a zinc finger protein with homology to the vertebrate transcription factor GATA-1. *Development.* **119**, 1277–1291
 30. Heitzler, P., Haenlin, M., Ramain, P., Calleja, M., and Simpson, P. (1996) A genetic analysis of pannier, a gene necessary for viability of dorsal tissues and bristle positioning in Drosophila. *Genetics.* **143**, 1271–1286
 31. Sato, M., and Saigo, K. (2000) Involvement of pannier and u-shaped in regulation of Decapentaplegic-dependent wingless expression in developing Drosophila notum. *Mech. Dev.* **93**, 127–138
 32. Gobert, V., Osman, D., Bras, S., Augé, B., Boube, M., Bourbon, H.-M., Horn, T., Boutros, M., Haenlin, M., and Waltzer, L. (2010) A Genome-Wide RNA Interference Screen Identifies a Differential Role of the Mediator CDK8 Module Subunits for GATA/ RUNX-Activated Transcription in Drosophila. *Mol. Cell. Biol.* **30**, 2837–2848
 33. Immarigeon, C., Bernat-Fabre, S., Augé, B., Faucher, C., Gobert, V., Haenlin, M., Waltzer, L., Payet, A., Cribbs, D. L., Bourbon, H.-M. G., and Boube, M. (2019) Drosophila Mediator Subunit Med1 Is Required for GATA-Dependent Developmental Processes: Divergent Binding Interfaces for Conserved Coactivator Functions. *Mol. Cell. Biol.* **39**, 1–18
 34. Hu, C.-D., Grinberg, A. V., and Kerppola, T. K. (2005) Visualization of Protein Interactions in Living Cells Using Bimolecular Fluorescence Complementation (BiFC) Analysis. *Curr. Protoc. Cell Biol.* **29**, 21.3.1-21.3.21
 35. Brand, A. H., and Perrimon, N. (1993) Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development.* **118**, 401–415
 36. Ghirlando, R., and Trainor, C. D. (2003) Determinants of GATA-1 binding to DNA: The role of non-finger residues. *J. Biol. Chem.* **278**, 45620–45628
 37. Newton, A., Mackay, J., and Crossley, M. (2001) The N-terminal Zinc Finger of the Erythroid Transcription Factor GATA-1 Binds GATC Motifs in DNA. *J. Biol. Chem.* **276**, 35794–35801
 38. Bourbon, H. M. (2008) Comparative genomics supports a deep evolutionary origin for the

- large, four-module transcriptional mediator complex. *Nucleic Acids Res.* **36**, 3993–4008
39. Kuuluvainen, E., Hakala, H., Havula, E., Estimé, M. S., Rämetsä, M., Hietakangas, V., and Mäkelä, T. P. (2014) Cyclin-dependent kinase 8 module expression profiling reveals requirement of Mediator subunits 12 and 13 for transcription of Serpent-dependent innate immunity genes in *Drosophila*. *J. Biol. Chem.* **289**, 16252–16261
 40. Ge, K., Cho, Y.-W., Guo, H., Hong, T. B., Guermah, M., Ito, M., Yu, H., Kalkum, M., and Roeder, R. G. (2008) Alternative Mechanisms by Which Mediator Subunit MED1/TRAP220 Regulates Peroxisome Proliferator-Activated Receptor α -Stimulated Adipogenesis and Target Gene Expression. *Mol. Cell. Biol.* **28**, 1081–1091
 41. Soutourina, J. (2019) Mammalian Mediator as a Functional Link between Enhancers and Promoters. *Cell.* **178**, 1036–1038
 42. Asmar, J., Biryukova, I., and Heitzler, P. (2008) *Drosophila* dLMO-PA isoform acts as an early activator of achaete/scute proneural expression. *Dev. Biol.* **316**, 487–497
 43. Haenlin, M., Cubadda, Y., Blondeau, F., Heitzler, P., Lutz, Y., Simpson, P., and Romain, P. (1997) Transcriptional activity of Pannier is regulated negatively by heterodimerization of the GATA DNA-binding domain with a cofactor encoded by the u-shaped gene of *Drosophila*. *Genes Dev.* **11**, 3096–3108
 44. Lowry, J. A., and Atchley, W. R. (2000) Molecular evolution of the GATA family of transcription factors: Conservation within the DNA-binding domain. *J. Mol. Evol.* **50**, 103–115
 45. Visvader, J. E., Crossley, M., Hill, J., Orkin, S. H., and Adams, J. M. (1995) The C-terminal zinc finger of GATA-1 or GATA-2 is sufficient to induce megakaryocytic differentiation of an early myeloid cell line. *Mol. Cell. Biol.* **15**, 634–641
 46. Arnold, C. D., Nemčko, F., Woodfin, A. R., Wienerroither, S., Vlasova, A., Schleiffer, A., Pagani, M., Rath, M., and Stark, A. (2018) A high-throughput method to identify trans-activation domains within transcription factor sequences. *EMBO J.* 10.15252/embj.201798896
 47. Pope, N. J., and Bresnick, E. H. (2013) Establishment of a Cell-Type-Specific Genetic Network by the Mediator Complex Component Med1. *Mol. Cell. Biol.* **33**, 1938–1955
 48. Pope, N. J., and Bresnick, E. H. (2010) Differential coregulator requirements for function of the hematopoietic transcription factor GATA-1 at endogenous loci. *Nucleic Acids Res.* **38**, 2190–2200
 49. Stumpf, M., Yue, X., Schmitz, S., Luche, H., Reddy, J. K., and Borggrefe, T. (2010) Specific erythroid-lineage defect in mice conditionally deficient for Mediator subunit Med1. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 21541–21546
 50. Tsai, K. L., Tomomori-Sato, C., Sato, S., Conaway, R. C., Conaway, J. W., and Asturias, F. J. (2014) Subunit architecture and functional modular rearrangements of the transcriptional mediator complex. *Cell.* **157**, 1430–1444
 51. Baidooonso, S. M., Guidi, B. W., and Myers, L. C. (2007) Med19(Rox3) regulates intermodule interactions in the *Saccharomyces cerevisiae* Mediator complex. *J. Biol. Chem.* **282**, 5551–5559
 52. Ding, N., Tomomori-Sato, C., Sato, S., Conaway, R. C., Conaway, J. W., and Boyer, T. G. (2009) MED19 and MED26 are synergistic functional targets of the RE1 silencing transcription factor in epigenetic silencing of neuronal gene expression. *J. Biol. Chem.* **284**, 2648–2656
 53. Robinson, P. J., Trnka, M. J., Pellarin, R., Greenberg, C. H., Bushnell, D. A., Davis, R., Burlingame, A. L., Sali, A., and Kornberg, R. D. (2015) Molecular architecture of the yeast Mediator complex. *Elife.* 10.7554/eLife.08719
 54. Hnisz, D., Abraham, B., Lee, T., Lau, A., Saint-Andre, V., Sigova, A., Hoke, H., and Young, R. (2014) Transcriptional super-enhancers connected to cell identity and disease. *Cell.* **155**, 1–24
 55. Boija, A., Klein, I. A., Sabari, B. R., Dall’Agnese, A., Coffey, E. L., Zamudio, A. V., Li, C. H., Shrinivas, K., Manteiga, J. C., Hannett, N. M., Abraham, B. J., Afeyan, L. K., Guo, Y. E., Rimel, J. K., Fant, C. B., Schuijers, J., Lee, T. I., Taatjes, D. J., and Young, R. A. (2018) Transcription Factors Activate Genes through the Phase-Separation Capacity of Their Activation Domains. *Cell.* **175**, 1842–1855.e16
 56. Sabari, B. R., Dall’Agnese, A., Boija, A., Klein, I. A., Coffey, E. L., Shrinivas, K., Abraham,

- B. J., Hannett, N. M., Zamudio, A. V., Manteiga, J. C., Li, C. H., Guo, Y. E., Day, D. S., Schuijers, J., Vasile, E., Malik, S., Hnisz, D., Lee, T. I., Cisse, I. I., Roeder, R. G., Sharp, P. A., Chakraborty, A. K., and Young, R. A. (2018) Coactivator condensation at super-enhancers links phase separation and gene control. *Science* (80-.). **361**, eaar3958
57. Cho, W. K., Spille, J. H., Hecht, M., Lee, C., Li, C., Grube, V., and Cisse, I. I. (2018) Mediator and RNA polymerase II clusters associate in transcription-dependent condensates. *Science* (80-.). **361**, 412–415




Fig. 1: *Med19* is required for GATA/Pannier target gene expression.

Drosophila thoraces: (A) wild type or (D) displaying *Med19*^{-/-} clones. Arrowheads point to DC bristles in A and asterisks to their expected position in D. The vertical arrow points to the thoracic cleft. Scale bars, 200µm.

(B-F) Expression of Pannier target genes in the dorsal compartment of control wing discs (B-C) or in *Med19*^{-/-} mitotic clones (GFP-) (E-F) Scale bar, 200µm. . The expression of *DC-ac-LacZ* reporter and Wg protein are revealed with anti-βgal (red, C', F'), and anti-Wg (blue, C'', F'') antibodies. Magnifications of the DC region are shown (C and F). White scale bars (B,E), 200 µm, yellow scale bars (C-C'' and F-F''), 20 µm.




Fig. 2: Med19 physically interacts with GATA/Pnr in cellulo, in vitro and in vivo.

(A-B) Co-immunoprecipitation (IP) experiments from S2 cells transfected with *pAct-Myc-Pnr* using anti Med19 antibody or preimmune serum (control IP) (A). The reverse experiment (B) was performed using anti-Myc beads from control cells or cells transfected with *pAct-Myc-Pnr*. Western blot assays using α Med19, α Pnr or α Myc antibodies are shown. (C-D) Autoradiographs from GST pull-down assays between GST-Pnr and 35 S-labeled (*) *in vitro*-translated Med19 (C), and between GST-Med19 and *in vitro*-translated 35 S-Pnr (D). (E-F') Bimolecular Fluorescence Complementation (BiFC) assays using VN-Pnr and either Med19-VC (E) or CycC-VC (F). Expression of fusion proteins along the A/P boundary of the wing disc is under the control of *dppGAL4* driver. Immunostaining shows expression of VC constructs (magenta, E and F), and BiFC signals are shown in green (E' and F'). Scale bars, 200 μ m.




Fig. 3: The GATA/Pnr C-zinc finger domain interacts with two conserved Med19 domains.

(A) GST pulldown assays delimitating Med19 interacting domains within Pnr. Left, schematic representation of GATA/Pnr and Pnr fragments generated to probe for binding to full-length GST-Med19. N and C show proper zinc fingers (orange boxes) and their basic tails (hatched green and orange boxes). H1 and H2 show amphipathic alpha helices. Right, corresponding autoradiographs from GST pulldown experiments. The critical domain for strong binding is narrowed down to Pnr amino-acids 220-278 (Med19 BS dotted rectangle) comprising C-ZF proper zinc finger and its basic tail. (B) Sequence alignment shows that this domain is highly conserved in *Drosophila* GATA factors (Srp, Grn, GATAd and GATAe) as well as human GATA factors (GATA1 to 6). Open circles denote residues participating in DNA binding (1, 2). Level of each amino acid conservation is represented underneath. (C) Schematic representation of the full length Med19 protein and the Med19 subdomains generated as GST fusions. + and – summarize Pnr GST-pulldown results, based on HA-Pnr¹⁻²⁹¹ detection on Western blots shown in (D-G). The 2 distinct domains of Med19 which mediate binding of Pnr are squared in blue and denoted BS1 and BS2 in C.




Fig. 4: Med19 physically interacts with GATA/Srp.

(A) Left, schematic representation of GATA/Srp and the multiple Srp fragments generated to probe for binding to full-length GST-Med19. Autoradiographs from GST pulldown experiments are shown on the right. Again, the critical binding domain is restricted to the GATA C-ZF domain containing the zinc-finger proper (orange square) and its basic tail (hatched green and orange boxes). (B) Bimolecular Fluorescence Complementation (BiFC) assays using expression of Med19-VC with either VN-Pnr or VN-Srp under the control of the *dppGAL4* driver. Immunostaining shows similar expression of VC constructs (magenta), and strong BiFC signals (green) with Pnr and also with Srp. Scale bars, 200 μ m.




Fig. 5: Med19 and Med1 are both required for GATA/Srp target genes expression and interact physically.

(A-B) Real Time quantitative PCR analysis of mRNA expression in Med19-depleted cells (dsMed19, A), or Med1-depleted cells (dsMed1, B). The control cultured cells treated with dsGFP were used as reference (100%), and *gapdh1* as an internal normalization gene. Each dot represents the result of one of 3 independent biological experiment (yellow, green and blue dots), bar graph indicates the mean and the standard deviation. The experiment was also reproduced with a second dsRNA against Med1 and Med19. (C) GST pulldown with GST-Med1^A and full length Med19 reveal direct interaction between the two MED subunits. (D) Left, schematic representation of the three Med1 protein isoforms (Med1-A, -B and -C) and Med1 fragments generated to probe for binding to full-length GST-Med19. The N-ter region (darker grey rectangle) comprises short evolutionarily-conserved motifs (black boxes) and correspond essentially to the yeast Med1 orthologue (38). The middle (white) and C-ter (light grey) regions emphasizes the divergent long metazoan-specific extensions except for a conserved C-terminal α helix (hatched box). Right, GST-Pull down essays narrow the Med19-binding domain to the highly conserved N-terminal portion of Med1 proteins.




Fig. 6: Models of MED-GATA interaction involving both Med19 and Med1.

(A) In addition to Med1, another Mediator subunit belonging to the middle module, Med19, directly interacts with GATA factors. Whereas Med1 interacts with both the GATA N and C-ZF domains, Med19 only interacts with the GATA C-ZF, that serves both as transactivation domain and DNA-binding domain. (B) Given that both Med19 and Med1 are required for the expression of most tested GATA Pnr and GATA Srp target genes, we propose that DNA-bound GATAs recruit the Mediator complex through direct contacts with both Med19 and Med1, thus enabling PolII machinery recruitment at GATA target genes. (C) In some cases such as Pnr-dependent *wg* expression in *Drosophila* wing discs, Med1 is dispensable. We propose a model in which additional contacts between MED subunits and other TFs or GATA cofactors would render GATA-Med19-interaction sufficient to recruit the Mediator complex at GATA target genes. (D) The binding affinity observed *in vitro* between Med1 and Med19 MED subunits, located at distant positions according to MED structural analyses, could reflect inter-MED instead of intra MED interactions. Such interactions could promote the formation of MED complex condensates at phase-separated nuclear foci, previously shown to boost transcription at TF-bound super-enhancers.

Mediator complex subunit Med19 binds directly GATA transcription factors and is required with Med1 for GATA-driven gene regulation in vivo

Clément Immarigeon, Sandra Bernat-Fabre, Emmanuelle Guillou, Alexis Verger, Elodie Prince, Mohamed A. Benmedjahed, Adeline Payet, Marie Couralet, Didier Monte, Vincent Villeret, Henri-Marc Bourbon and Muriel Boube

J. Biol. Chem. published online July 31, 2020

Access the most updated version of this article at doi: [10.1074/jbc.RA120.013728](https://doi.org/10.1074/jbc.RA120.013728)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts