



HAL
open science

Rhizospheric Plant–Microbe Interactions

Harriet Middleton, Étienne Yergeau, Cécile Monard, Jean-Philippe Combier,
Abdelhak El Amrani

► **To cite this version:**

Harriet Middleton, Étienne Yergeau, Cécile Monard, Jean-Philippe Combier, Abdelhak El Amrani. Rhizospheric Plant–Microbe Interactions. Trends in Plant Science, 2021, 26 (2), pp.132-141. 10.1016/j.tplants.2020.09.005 . hal-02989014

HAL Id: hal-02989014

<https://hal.science/hal-02989014v1>

Submitted on 18 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Rhizospheric plant-microbe interactions: microRNAs, a key mediator**

2 Harriet Middleton^{1*}, Étienne Yergeau², Cécile Monard¹, Jean-Philippe Combier³, Abdelhak El
3 Amrani¹

4 ¹University of Rennes 1, CNRS/UMR 6553/OSUR, Ecosystems - Biodiversity - Evolution, 35042 Rennes
5 Cedex, France

6 ²Institut national de la recherche scientifique, Centre Armand-Frappier Santé Biotechnologie, Laval, H7V
7 1B7, Canada

8 ³Laboratoire de Recherche en Sciences Vegetales, UMR 5546, UPS, CNRS, Auzeville-Tolosane 31320,
9 France

10

11 *Correspondence: harriet.middleton@univ-rennes1.fr (H. Middleton)

12 Abstract:

13 The importance of microorganisms in plant development, nutrition and stress resistance
14 is consensus and has led to a more holistic approach of plant-microbe interactions,
15 under the holobiont concept. The structure of plant microbiota is often described as host-
16 driven, especially in the rhizosphere, where microbial communities are shaped by
17 diverse rhizodeposits. Gradually, this anthropogenic vision is fading and being replaced
18 by the idea that plants and microorganisms co-shape the plant microbiota. Through co-
19 evolution, plants and microbes have developed cross-kingdom communication channels.
20 We propose that microRNAs are crucial mediators of plant-microbe interactions and
21 microbiota shaping in the rhizosphere. Moreover, we suggest, as an alternative to
22 generally unsuccessful strategies based on microbial inoculants, microRNAs as a
23 promising tool for novel holobiont engineering.

24 Keywords: miRNA; extracellular vesicles (EVs); rhizospheric microbiota; inter-kingdom
25 communication; holobiont engineering

26

27

28

29 Highlights

- 30 ● Plants interact with a diversity of microbes, especially in the rhizosphere. The
31 plant and its associated microbes form a co-evolved unit called “holobiont”,
32 implying the emergence of bi-directional interkingdom communication channels.
- 33 ● MicroRNAs may be key mediators of plant-microbe interactions, as they are well-
34 conserved throughout the living, can be conveyed by extracellular vesicles on
35 long-distances and are part of interkingdom interactions.
- 36 ● Plant microRNAs are uptaken by gut microbiota and phytopathogens, resulting in
37 a modulation of microbial gene expression, hence of microbial community activity
38 and composition. In return, microbial microRNA-like RNAs target plant genes and
39 act on their fitness.

40 Cross-kingdom communication in the rhizosphere *via* microRNAs

41 Plant-associated microorganisms are essential for the well-being of plants: they play a
42 crucial role in development, nutrition and resistance to various stresses [1]. Notably, the
43 **rhizosphere** (see [Glossary](#)) provides a rich environment in which diverse microbial
44 communities, including plant beneficial microorganisms, live abundantly, in close relation
45 with the plant. Together, a plant and its associated microbes form a coordinated unit,
46 referred to as a **holobiont** [2]. In order for this association to remain coordinated, it is
47 necessary to have some form of regulation within the microbial communities and with
48 the plant. Currently, there is still a gap in our knowledge with regards to the assembly
49 and composition dynamics of the rhizospheric **microbiota**. Generally, it is thought that
50 the plant performs a selection of its microbial partners through the influence of its
51 **rhizodeposits**, and especially its **root exudates** [3]. However, this uni-directional

52 selection is starting to be questioned [4]. Indeed, the **co-evolution** of the plant-microbe
53 holobiont suggests a bi-directional communication, notably regarding the shaping of the
54 rhizospheric microbiota. The composition and function of these communities are
55 modulated by a variety of molecules and signals [5]. These past years, a growing
56 interest in small RNAs (sRNAs, [Box 1](#)) and more specifically microRNAs as mediators of
57 cross-kingdom communication has emerged [6]. Conveyed by **extracellular vesicles**
58 (EVs), mammal host microRNAs and dietary plant microRNAs have been shown to be
59 implicated in regulating the intestinal microbiota, by modulating both bacterial gene
60 expression and species abundance [7–9]. Regarding the commonalities between the gut
61 microbiota in animals and the rhizospheric microbiota in plants [10,11], combined with
62 their key role in holobiont nutrition and health, and the universality of the RNA
63 “language”, the implication of microRNAs in plant-rhizospheric microbiota **cross-**
64 **kingdom** interactions is very plausible, but completely understudied.

65
66 In this Opinion, we suggest that microRNAs, conveyed by EVs, constitute a cross-
67 kingdom bi-directional communication channel between the plant and its associated
68 rhizospheric microorganisms ([Figure 1, Key Figure](#)). Supported by experimental
69 evidence of the connection between plant microRNAs and the intestinal microbiota [9]
70 and between plant microRNAs and pathogens [12–15], we propose that microRNAs are
71 implicated in shaping the abundance, composition, functions and activities of the
72 rhizospheric microbiota.

73

74

75 **Who shapes the rhizospheric microbiota?**

76 From the very start of their lives, as a seed and then through all stages of development,
77 from root to shoot, plants are surrounded and populated inside and outside by
78 microorganisms. The significant role of these microorganisms in the survival and growth
79 of the plant is fully accepted and has been thoroughly reviewed [1]. Moreover, it has
80 been stated that plant-associated microbes allow the plant to better adapt to their
81 environment [16]. The plant microbiota can be acquired from its environment, through
82 horizontal transmission, but also directly in the seed from the parent, through vertical
83 transmission [17]. The environmental acquisition of microbes and the laws governing
84 their assembly in the plant holobiont are poorly understood. It is generally thought that
85 during root penetration in the bulk soil, the soil microbiota gradually differentiates into the
86 rhizospheric microbiota through contact with rhizodeposits, which greatly impact the
87 composition of the microbiota by having antimicrobial properties or by serving as a
88 **substrate**, for example [18]. After this first community shift, the microbiota composition
89 will be fine-tuned in a compartment-specific manner (i.e. endosphere, rhizoplane or
90 rhizosphere) [16], but also depending on host genotype. Indeed, characteristics
91 determined by plant genetic factors, such as root morphology and root exudate quality
92 and quantity were shown to shape the rhizospheric microbiota [3]. In addition, other
93 factors such as the plant immune system, plant developmental stage and season, were
94 shown to be greatly involved in shaping the rhizospheric microbiota [19]. The existence
95 of a core microbiota in certain plants, regardless of soil origin and fertilization regimes,

96 further suggests that the assembly and selection of these communities is at least partly
97 plant-driven [19].

98 Even though the assembly of rhizospheric microbial communities seems to be mainly
99 the result of **selection**, it is important to consider the other processes of community
100 assembly: **dispersal**, **drift** and **speciation** [20]. It is suggested that these processes
101 modulate, at varying degrees through plant development, the establishment of microbial
102 communities in the rhizosphere [20]. Changes in the microbiota and its functional
103 diversity are not the sole result of the acquisition of novel microbial strains, selected by
104 the host plant. Such variations in the holobiont are also the consequence of specific
105 **microbial amplification/reduction** and **horizontal gene transfer** (HGT) [21]. The
106 combination of these three processes allow rapid adaptation of the plant holobiont to
107 challenging and changing environments. These shifts from originally present microbial
108 communities, to communities specifically tailored to the situation, have been observed in
109 many studies. An example of microbial abundance modulation by the plant was
110 described in rice: under drought stress, the shift in composition of root microbial
111 communities was characterized by a specific enrichment of certain bacterial taxa, known
112 as plant growth promoting rhizobacteria (PGPR) and/or resisting drought stress [22].
113 The authors of this study suggested that the change in root exudation, due to drought
114 stress, could be the origin of the selective enrichment of certain microbes. In addition to
115 a change in microbial composition, stressful conditions can also impact microbial activity
116 in the rhizosphere. In hydrocarbon-contaminated soils, both microbial composition and
117 activity are modulated: for example, *via* a specialization of the microbiota, through a
118 selective enrichment of hydrocarbon degraders and PGPRs [23] or *via* a stimulation of

119 specific microbial activities like hydrocarbon degradation, in the microbes already
120 present [24].

121 The regulation of rhizospheric microbiota is not solely driven by plants, but also by the
122 microbes themselves. Indeed, they can modulate the plant's environment and even
123 reprogram the plant to their advantage *via* exudation of phytohormones, volatile organic
124 compounds, quorum-sensing molecules and antimicrobials [5]. Shaping of the
125 rhizospheric microbiota is also based on microbial metabolite substrate preference [25]
126 and interactions between microbial members. Microbial interactions can be based on
127 cooperative mechanisms, such as quorum-sensing and metabolite exchange under
128 nutrient-poor conditions, or competitive mechanisms, such as secretion of antimicrobial
129 compounds [19], but it remains unclear which type of interaction dominates
130 communities. Overall, microbial interactions can induce an alteration in gene expression
131 within the communicating microbes [3]: hence, microbial interactions modulate both
132 shape and function of the microbiota. Based on their role in regulating plant-associated
133 microbial communities, some argue that symbionts are ecological engineers of the
134 holobiont [4]. Thus, on top of plant regulation of the rhizospheric microbiota, microbial
135 interactions in the rhizosphere also play an essential role in structuring the plant
136 microbiota.

137 **The role of microRNAs in bi-directional cross-kingdom communication**

138 Between the host plant and the associated microorganisms, there is a constant two-way
139 dialogue, adapted to cross-kingdom communication (Box 2). A well-known example is
140 during the establishment of the rhizobium-legume symbiosis, in which the plant

141 produces flavonoids that induce bacterial *nod* genes, resulting in the secretion of Nod
142 factors, which in turn initiates nodule formation, that house the rhizobacteria that will
143 there fix atmospheric nitrogen and transfer it to the plant [26,27].

144 As discussed previously, a plethora of signalling molecules are exchanged in the
145 rhizosphere. Interestingly, a majority of microbial signalling molecules used to
146 communicate with the host plant are also part of inter-microbial communication [5].
147 Knowing that these compounds affect plants, in their development or immune system,
148 these signals may allow the microbiota to control the plant, or plants have simply
149 evolved to perceive them. With regards to the holobiont concept, these two possibilities
150 could co-exist, as there is constant bi-directional communication between host and
151 microbes.

152 In the past years, interkingdom communication studies have bloomed, especially
153 regarding host-microbiota interactions and their implications for human health. Amongst
154 the different possible mediators of such conversations, sRNAs, and more specifically
155 microRNAs, have sparked a growing interest [28] (Box 1).

156 MicroRNAs are a class of small non-coding RNAs (~21 nucleotides), implicated in
157 regulating gene expression: either by repressing the translation or by directly cleaving
158 the target mRNA. MicroRNAs were first discovered in nematodes [29,30] which then led
159 to their identification in fishes to flies and mammals, highlighting their conservation
160 across the animal kingdom [31]. Shortly after, plant microRNAs were described, for the
161 first time, in *Arabidopsis thaliana* [32] and then, microRNA-like RNAs (miRNAs) were
162 characterized in fungi [33] (Box 1).

163 It was previously thought that microRNAs had appeared with pluricellular organisms
164 and that their function was limited to intercellular communication within a single
165 organism. Since then, microRNAs or microRNA-like RNAs have also been found in
166 unicellular organisms: first, in *Chlamydomonas reinhardtii* [34,35], then in oomycetes
167 [36] and in bacteria [37], the latter being referred to as microRNA-size small RNA
168 (msRNA) [38]. The discovery of microRNA-like RNAs in unicellular organisms suggests
169 that microRNA-based gene regulation is an ancient mechanism and could have
170 appeared prior to pluricellular life [34]. MicroRNA, or at least very similar molecules, are
171 indeed evolutionarily well conserved and produced throughout all branches of life.

172
173 Moreover, microRNAs are not restricted to intercellular communication within a tissue,
174 an organ or an organism: they can be transported on long distances, between
175 individuals and from one species to another. With regards to other signalling
176 compounds, the ancient and widespread RNA “language” is particularly suited for cross-
177 kingdom communication, as it is employed by all organisms which use RNA-based
178 regulation: RNA provides a universally conserved, stable, communication channel [39].
179 MicroRNAs, as mediators of such conversations, are conveyed by nanovesicles,
180 generally named EVs, although Outer Membrane Vesicles (OMVs) is sometimes used
181 for Gram-negative bacteria [40]. In eukaryotic cells, EVs originate from the fusion of
182 multivesicular bodies with the plasma membrane, whereas bacterial EVs or OMVs are
183 produced through external budding of the bacterial outer membrane [40]. These vesicles
184 function as a protective transportation structure for RNA, but also proteins and lipids.
185 Before encapsulation, these molecules are selectively sorted into EVs or OMVs [40],

186 suggesting a functional targeting. Cross-kingdom microRNA exchanges have been
187 described for a multitude of species and interactions (Table 1). Many studies have
188 focused on host-pathogen interactions, often emphasizing on sRNA-mediated **RNA**
189 **interference** (RNAi) mechanisms which are conserved virulence mechanisms shared by
190 many pathogens to suppress host immunity [13], whereas the host may target virulence-
191 related genes in the pathogen [41]. Evidence of such host-pathogen interactions were
192 demonstrated between mammals and bacteria [42,43]; nematodes and mammals [44];
193 plants and fungi [12,14,15,41,45]; plants and insects [46] and between parasitic plants
194 and host plants [47]. Actual bi-directional cross-kingdom RNAi, through sRNA trafficking,
195 was first described between *Arabidopsis thaliana* and a fungal pathogen, *Botrytis*
196 *cinerea* [13].

197 However, more and more studies are now highlighting the role of microRNA exchanges
198 in non-pathogenic relationships, with many focusing on host-microbiota interactions in
199 the gut. Indeed, host microRNAs from the intestinal epithelium cells have been shown to
200 impact the composition of the gut microbiota by specifically entering certain bacteria and
201 regulating the transcription of a wide range of housekeeping and sugar degradation
202 genes, thereby impacting bacterial growth [7,8]. Interestingly, ingested plants release
203 EVs in the gut, where they are preferentially uptaken, along with their content in plant
204 microRNA, by certain target bacteria, inducing a modification of their gene expression
205 [9,48]. In turn, their production and secretion of metabolites may be altered, leading to
206 differential development of bacterial species interacting with these target bacteria [9].
207 Hence, ingested plants are capable of shaping the gut microbiota of mammals through

208 microRNA-based communication, partly explaining the effect of diet on the gut
209 microbiota.

210 Given these recent advances on the impact of host microRNAs and plant microRNAs on
211 the gut microbiota, we propose that plants and their rhizospheric microbiota also
212 communicate *via* microRNAs, regulating microbial composition and activity in the
213 rhizosphere. Besides, it has recently been shown that bacterial vesicles can carry their
214 DNA cargo into other bacteria [49], a mechanism called “vesiduction” [50], but also into
215 host eukaryotic cells [51]. Hence, it is easily conceivable that msRNA are also conveyed
216 in such a manner and may mediate cross-kingdom communication. Plant microRNAs
217 are tissue specific, and many hundreds are found in the root tissues [52], and it is thus
218 likely that a subset of these, selected through co-evolution with surrounding organisms,
219 would be excreted via EVs in the rhizosphere.

220 **Holobiont engineering and current RNAi applications**

221 Under the holobiont era, the role of microorganisms in shaping their own communities,
222 concurrently to plant influence, is becoming a subject of interest. In many fields, such as
223 ecological engineering, scientists have aimed to successfully engineer the plant’s
224 microbiota, in order to optimize the ecosystemic services provided by the plant
225 holobiont. Many biotech companies, worldwide, have made large efforts to create
226 specific **inocula** containing carefully selected microorganisms, in order to improve crop
227 production. However, many of these methods do not translate efficiently from lab to field:
228 inoculated microbes can be in competition with native communities or simply washed
229 away before being able to establish themselves stably. Inoculation of non-native species

230 also raises questions about potential invasive species. Hence, microbiome engineering
231 is slowly evolving into “holobiont engineering” [53], which is already applied in
232 agricultural practices such as grafting, organic or low-farming and plant breeding
233 strategies which include microbiota [54]. Holistic biotechnologies are also being
234 developed such as approaches based on soil memory [55] or synthetic microbiomes
235 [56]. We suggest that these holobiont engineering techniques could also be based on
236 cross-kingdom communication mediators such as microRNAs.

237 Currently, cross-kingdom RNAi technology for crop protection already exists and is
238 largely investigated, notably Host-Induced Gene Silencing (HIGS) which uses
239 genetically modified plants able to express specific sRNA or double-stranded RNAs (ds-
240 RNA) which target specific genes in pathogens [57]. However, HIGS requires the use of
241 genetically modified plants, which is controversial, and even banned in certain countries.
242 To counter these issues, environmental RNAi quickly emerged: sRNA or ds-RNA are
243 directly applied on the host plant, resulting in the silencing of specific genes in
244 pathogens and disease control [58]. These environmental RNAi methods are named
245 Spray-Induced Gene Silencing (SIGS). However, SIGS technologies are still in their
246 infancy and encounter many problems, such as penetration, application effects and
247 production costs: currently, there are no available SIGS-based products on the market
248 [59,60].

249 Alternatively, the use of microRNA-encoded peptides (miPEPs) [61,62], as a tool to
250 amplify the expression of specific plant microRNAs, could be an interesting perspective.
251 miPEPs are small regulatory peptides, encoded in plant pri-miRNAs (Box 1), capable of
252 enhancing the transcription of their corresponding microRNA, resulting in a stronger

253 repression of the target mRNA [61]. When synthetic miPEP171b was applied on
254 *Medicago truncatula* the expression of the microRNA171b was increased and led to
255 enhanced mycorrhization by *Rhizophagus irregularis* [63]. If our hypothesis, that
256 microRNAs are exchanged between the plant and its rhizospheric microbiota, is
257 confirmed, then the use of miPEPs to stimulate the synthesis and release of certain
258 beneficial plant microRNAs could be a promising tool for many fields in ecological
259 engineering. Indeed, miPEPs could indirectly modulate the rhizospheric microbiota,
260 without adding exogenous inocula. Moreover, the application of miPEPs is
261 straightforward and does not require genetic transformation [62], which makes miPEP
262 technology an exciting pathway for holobiont engineering.

263 **Concluding remarks and future perspectives**

264 Regarding our current knowledge about the role of host and plant microRNA on the
265 composition and activity of the gut microbiota, the universality of the RNA “language”,
266 combined with the potentiality of RNAi technology, our opinion is that microRNAs at
267 least partly shape the rhizospheric microbiota. Such a point of view implies that
268 microRNA-based microbial community manipulation has an enormous potential for
269 agriculture, forestry and environmental remediation. Evidently, more fundamental
270 research is needed to decrypt the implication of microRNAs in plant-microbe
271 communication and especially the impact of plant microRNAs on rhizospheric microbial
272 communities. If our hypotheses reveal to be correct, then a new field of perspectives
273 regarding our knowledge of plant-microbe interactions and holobiont engineering
274 possibilities will arise, with a plethora of **Outstanding Questions**.

275

276

277 [Glossary](#)

278 **Co-evolution:** process during which two or more species, sharing a close ecological
279 relationship, evolve simultaneously

280 **Cross-kingdom:** involves species from distinct taxonomic high ranks, i.e. animals,
281 plants, fungi, protists bacteria or archaea

282 **Dispersal:** refers to the movement of species between communities, from one site to
283 another

284 **Drift:** random change in population size due to births and deaths, which can happen in
285 small, low-diversity communities

286 **Extracellular Vesicles (EVs):** spherical structures with a lipid-membrane, comprising
287 microvesicles to exosomes, ranging from 30nm to 1000nm, secreted by a wide diversity
288 of organisms. Plant EVs range mostly from 20nm to 500nm. EVs are greatly involved in
289 intercellular communication as they contain and transport proteins, lipids, nucleic acids
290 and other molecules.

291 **Holobiont:** originally termed by Lynn Margulis as an organism, animal or plant, and its
292 symbiotic associations, “holobiont” now refers to a host macrobe and, more generally, its
293 numerous associated microbes

294 **Horizontal Gene Transfer (HGT):** a mechanism of gene exchange between bacterial
295 cells

296 **Inoculum** (pl. inocula): microbial sample destined to be added to a plant or soil, for
297 example

298 **Microbial amplification/reduction**: respectively, an increase/decrease in numbers of a
299 certain group of associated microbial species, due to environmental conditions, also
300 known as “enrichment/depletion”

301 **Microbiota**: community of microorganisms living in a specific region (e.g. human
302 intestine, skin or the rhizosphere)

303 **Rhizodeposits**: variety of carbon- and nitrogen-rich compounds released by roots into
304 their environment (root-cells, mucilage, root exudates, soluble lysates and volatile
305 compounds), responsible for shaping the rhizosphere microbiota [18]

306 **Rhizosphere**: portion of soil surrounding the root system, greatly influenced by the plant
307 notably *via* its rhizodeposits. Microorganisms living in this habitat vary from those living
308 on the rhizoplane (root surface) or in the endosphere (inside plant tissues).

309 **RNA interference (RNAi)**: Eukaryotic process of gene silencing, at the post-
310 transcriptional level, by small non-coding RNAs (siRNA, miRNA and piRNA).
311 Prokaryotes have similar mechanisms such as microRNA-like regulatory systems and
312 CRISPR-Associated System [64]

313 **Root exudates**: a class of rhizodeposits secreted by plants in the meristematic region of
314 root tips, such as sugars and amino acids, but also organic acids, nucleotides, peptides,
315 fatty acids and secondary metabolites. The composition of root exudates depends on
316 plant genotype, its developmental stage and environmental conditions [3].

317 **Selection:** result of biotic and abiotic influence on the reproductive success (i.e. fitness)
318 of an individual or species. This is the most studied process regarding microbial
319 communities composition differences, due to plant exudates or environment, whereas
320 the study of drift, dispersal and speciation in the microbial field is more complex to
321 approach, if even possible [20].

322 **Speciation:** evolutionary process through which novel species form

323 **Substrate:** nutritious substance used by microorganisms to grow and feed on to meet
324 their nutritional requirements

325

326 **Box 1. An overview of plant and microbial small RNAs**

327 Plants produce a wide variety of small RNAs (sRNAs), each capable of regulating gene
328 expression through different silencing pathways. These 21 to 24-nucleotide RNAs
329 originate from hairpin precursors or double-stranded RNA (dsRNA) which are processed
330 by an RNase III Dicer-like protein (DCL) forming sRNA duplexes, with 2'-O-methylation
331 at 3' ends, which in turn are loaded into Argonaute proteins (AGO) in order to function
332 and repress expression of target RNAs of complementary sequence [65]. Plant sRNAs
333 encompass small-interfering RNAs (siRNAs) and microRNAs (miRNAs).

334 siRNAs transcripts are converted by an RNA-Dependent RNA Polymerase (RDRP) into
335 long dsRNAs, which are then processed by DCL proteins into a siRNA duplex. Plant
336 siRNAs are categorized into secondary siRNAs and heterochromatic siRNAs
337 (hetsiRNAs). They are involved in post-transcriptional gene silencing (PTGS), but mostly
338 in RNA-directed DNA methylation (RdDM) and transcriptional gene silencing (TGS) [65].

339 MicroRNAs encoded by MIR genes are transcribed by RNA polymerase II (Pol II) into a
340 hairpin structure, the primary-miRNA (pri-miRNA). A DCL protein then processes pri-
341 miRNA into a smaller stem-loop structure, precursor-miRNA (pre-miRNA), which is
342 processed again by DCL protein forming a mature miRNA duplex (Figure 1). Once
343 loaded into AGO proteins, forming the multi-protein RNA-Induced Silencing Complex
344 (RISC), miRNAs can then repress target gene expression by mRNA cleavage or
345 translation inhibition [65].

346 Together, plant small RNAs, especially microRNAs, are involved in many regulatory
347 processes and play essential roles in plant development and resilience to stresses. The
348 conservation of microRNA or microRNA-like RNAs and their mechanisms across the
349 branches of life suggests their importance for many organisms and their potential role as
350 mediator of cross-kingdom communication.

351 For instance, in the rhizosphere, the plant interacts with diverse microorganisms,
352 capable of producing microRNA-like RNAs. Similarly to eukaryotic microRNA, bacterial
353 ~22nt RNA fragments, namely msRNA, obtained from hairpin structures have been
354 discovered [37,38,66] (Figure 1), but their function remains unknown, even though their
355 abundance suggests an important functional role. Aside msRNAs, bacteria possess a
356 variety of small RNAs, mainly characterized as trans-acting sRNA and cis-acting sRNA,
357 but many new bacterial small regulatory RNAs are to be defined [67].

358 Fungi are eukaryotic organisms and possess diverse small RNAs, however it is still
359 under debate if they produce microRNAs. Most studies have demonstrated the presence
360 of microRNA-like RNA (miRNA) in fungi, which are not produced by the same

361 biogenesis pathway as real microRNAs (Figure I) [33,68]. Interestingly, the evolutionarily
362 ancient lineage of oomycetes also possesses microRNAs [36,69].

363

364 **Box 2. Plant-microbe interactions under the hologenome theory**

365 The hologenome theory states that the combined genomes of the holobiont (e.g. the
366 plant genome and the associated microbial genomes), is one of the biological units
367 subjected to natural selection [21]. This concept is mainly based on: (i) all animals and
368 plants are associated with a diversity of microbes, (ii) a proportion of these microbes are
369 transmitted between holobiont generations, (iii) host-microbiota interactions impact the
370 holobiont fitness [70]. This theory could explain the speed at which genetic variation
371 happens in holobionts: the microbiota evolves at a fast rate due to short generation
372 times, horizontal gene transfer and arrival of new microbes in the community [21]. The
373 strong bonds established between a host and its microbiota, combined with the power of
374 microbial evolution, must have great consequences on co-evolution and macroevolution
375 mechanisms in the holobiont. In the case of the plant holobiont, co-evolution between
376 plant and microbes suggests the existence of cross-kingdom bi-directional
377 communication. Regarding the diversity and dynamics of the rhizospheric microbiota, it
378 is probable that the host plant co-evolves with only a portion of its microbiota, namely
379 the core microbiota, which is often characterized as stable and transmittable to offspring.
380 However, these two-way interactions allow regulation of the hologenome, by comprising
381 the essential roles of both plant and microbes in holobiont fitness.

382

383

Host	Interacting organism	Small RNA mobility	Small RNA exchanged	Small RNA vehicle	Targeted genes	Effect	Reference
Human & Mice	Gut microbiota (<i>F. nucleatum</i> , <i>E.coli</i> ...)	From mammal host to bacteria	microRNA	EVs	dnaK operon, 16S rRNA and rpoB (in <i>Fn</i>) & RNase P,	Regulation of bacterial growth	[7]

384

385 **Table 1. Examples of cross-kingdom communication through small RNAs**

					rutA, yegH and fucO (in <i>E. coli</i>)		
Human & Mice	Gut microbiota	From mammal host to bacteria	microRNA	EVs	β -galactosidase AMUC_RS06985	Upregulation of target gene, enrichment of <i>Akkermansia muciniphila</i> , increasing regulatory T cells & ameliorating multiple sclerosis	[8]
Ginger (<i>Zingiber officinale</i>)	Gut microbiota (<i>Lactobacillaceae</i>)	From plant to bacteria	microRNA	Exosome-Like Nanoparticles	Monoxygenase ycnE	Regulation of bacterial growth & increase in indole-3-carboxaldehyde (I3A)	[9]
Ginger, grapefruit (<i>Citrus paradisi</i>), carrot (<i>Daucus carota</i>), grape (<i>Vitis vinifera</i>)	Mice gut cells	From plant to mice	microRNA	Exosome-Like Nanoparticles	?	Induction of anti-inflammation genes expression	[48]
Soybean (<i>Glycine max</i>)	Rhizobia bacteria	From bacteria to plant	tRNA-derived small RNA fragments (tRFs)	?	Orthologs of <i>Arabidopsis</i> genes involved in root hair/plant development	Positive regulation of rhizobial infection and nodulation formation in soybean	[71]
<i>Arabidopsis thaliana</i>	<i>Botrytis cinerea</i>	From plant to fungal pathogen	sRNA	EVs	Vesicles-trafficking pathway, vacuolar protein sorting	Decrease in fungal virulence	[72]
Cotton plant (<i>Gossypium sp.</i>)	<i>Verticillium dahliae</i>	From plant to fungal pathogen	microRNA	?	Virulence genes: Ca ²⁺ -dependent cysteine protease & an isotrichodermin C-15 hydroxylase	Silencing fungal virulence genes	[12]
Wheat (<i>Triticum aestivum</i>)	<i>Puccinia striiformis f. sp. tritici</i> (Pst)	From fungal pathogen to plant	microRNA-like RNA (milRNA)	?	Pathogenesis-related 2 (PR2)	Suppression of plant defenses	[14]
<i>Arabidopsis thaliana</i> & <i>Nicotiana benthamiana</i>	Parasitic plant <i>Cuscuta campestris</i>	From parasitic plant to host plant	microRNA	?	TIR1, AFB2 and AFB3 (auxin receptors), BIK1 (kinase receptor), SEOR1 (phloem protein) and HSFB4 (transcriptional repressor)	mRNA cleavage & secondary siRNA production	[47]

386

387

389 [References](#)

- 390 1 Vandenkoornhuysen, P. *et al.* (2015) The importance of the microbiome of the plant holobiont. *New*
391 *Phytol.* 206, 1196–1206
- 392 2 Bordenstein, S.R. and Theis, K.R. (2015) Host Biology in Light of the Microbiome: Ten Principles of
393 Holobionts and Hologenomes. *PLOS Biol.* 13, e1002226
- 394 3 Sasse, J. *et al.* (2018) Feed Your Friends: Do Plant Exudates Shape the Root Microbiome? *Trends Plant*
395 *Sci.* 23, 25–41
- 396 4 Uroz, S. *et al.* (2019) Plant Symbionts Are Engineers of the Plant-Associated Microbiome. *Trends Plant*
397 *Sci.* 24, 905–916
- 398 5 Venturi, V. and Keel, C. (2016) Signaling in the Rhizosphere. *Trends Plant Sci.* 21, 187–198
- 399 6 Leitão, A.L. *et al.* (2020) Interspecies Communication in Holobionts by Non-Coding RNA Exchange. *Int.*
400 *J. Mol. Sci.* 21, 2333
- 401 7 Liu, S. *et al.* (2016) The Host Shapes the Gut Microbiota via Fecal MicroRNA. *Cell Host Microbe* 19, 32–
402 43
- 403 8 Liu, S. *et al.* (2019) Oral Administration of miR-30d from Feces of MS Patients Suppresses MS-like
404 Symptoms in Mice by Expanding *Akkermansia muciniphila*. *Cell Host Microbe* 26, 779-794.e8
- 405 9 Teng, Y. *et al.* (2018) Plant-Derived Exosomal MicroRNAs Shape the Gut Microbiota. *Cell Host Microbe*
406 24, 637-652.e8
- 407 10 Hacquard, S. *et al.* (2015) Microbiota and Host Nutrition across Plant and Animal Kingdoms. *Cell*
408 *Host Microbe* 17, 603–616
- 409 11 Ramírez-Puebla, S.T. *et al.* (2013) Gut and Root Microbiota Commonalities. *Appl. Environ.*
410 *Microbiol.* 79, 2–9
- 411 12 Zhang, T. *et al.* (2016) Cotton plants export microRNAs to inhibit virulence gene expression in a
412 fungal pathogen. *Nat. Plants* 2, 16153
- 413 13 Wang, M. *et al.* (2016) Bidirectional cross-kingdom RNAi and fungal uptake of external RNAs
414 confer plant protection. *Nat. Plants* 2, 16151
- 415 14 Wang, B. *et al.* (2017) *Puccinia striiformis* f. sp. *tritici* microRNA-like RNA 1 (*Pst*-miR1), an
416 important pathogenicity factor of *Pst*, impairs wheat resistance to *Pst* by suppressing the wheat
417 pathogenesis-related 2 gene. *New Phytol.* 215, 338–350
- 418 15 Wang, M. *et al.* (2017) *Botrytis* small RNA *Bc*-siR37 suppresses plant defense genes by cross-
419 kingdom RNAi. *RNA Biol.* 14, 421–428
- 420 16 Bulgarelli, D. *et al.* (2013) Structure and Functions of the Bacterial Microbiota of Plants. *Annu.*
421 *Rev. Plant Biol.* 64, 807–838
- 422 17 Shade, A. *et al.* (2017) Ecological patterns of seed microbiome diversity, transmission, and
423 assembly. *Curr. Opin. Microbiol.* 37, 15–22
- 424 18 Tian, T. *et al.* (2020) The role of rhizodeposits in shaping rhizomicrobiome. *Environ. Microbiol.*
425 *Rep.* 12, 160–172
- 426 19 Hassani, M.A. *et al.* (2018) Microbial interactions within the plant holobiont. *Microbiome* 6, 58
- 427 20 Dini-Andreote, F. and Raaijmakers, J.M. (2018) Embracing Community Ecology in Plant
428 Microbiome Research. *Trends Plant Sci.* 23, 467–469
- 429 21 Zilber-Rosenberg, I. and Rosenberg, E. (2008) Role of microorganisms in the evolution of animals
430 and plants: the hologenome theory of evolution. *FEMS Microbiol. Rev.* 32, 723–735

431 22 Santos-Medellín, C. *et al.* (2017) Drought Stress Results in a Compartment-Specific Restructuring
432 of the Rice Root-Associated Microbiomes. *mBio* 8, mBio.00764-17, e00764-17

433 23 Tardif, S. *et al.* (2016) The Willow Microbiome Is Influenced by Soil Petroleum-Hydrocarbon
434 Concentration with Plant Compartment-Specific Effects. *Front. Microbiol.* 7,

435 24 Yergeau, E. *et al.* (2014) Microbial expression profiles in the rhizosphere of willows depend on
436 soil contamination. *ISME J.* 8, 344–358

437 25 Zhalnina, K. *et al.* (2018) Dynamic root exudate chemistry and microbial substrate preferences
438 drive patterns in rhizosphere microbial community assembly. *Nat. Microbiol.* 3, 470–480

439 26 Redmond, J.W. *et al.* (1986) Flavones induce expression of nodulation genes in Rhizobium.
440 *Nature* 323, 632–635

441 27 Rossen, L. *et al.* (1985) The *nodD* gene of *Rhizobium leguminosarum* is autoregulatory and in the
442 presence of plant exudate induces the *nodA,B,C* genes. *EMBO J.* 4, 3369–3373

443 28 Weiberg, A. *et al.* (2015) Conversations between kingdoms: small RNAs. *Curr. Opin. Biotechnol.*
444 32, 207–215

445 29 Lee, R.C. *et al.* (1993) The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense
446 complementarity to *lin-14*. *Cell* 75, 843–854

447 30 Wightman, B. *et al.* (1993) Posttranscriptional regulation of the heterochronic gene *lin-14* by *lin-*
448 4 mediates temporal pattern formation in *C. elegans*. *Cell* 75, 855–862

449 31 Wahid, F. *et al.* (2010) MicroRNAs: Synthesis, mechanism, function, and recent clinical trials.
450 *Biochim. Biophys. Acta BBA - Mol. Cell Res.* 1803, 1231–1243

451 32 Reinhart, B.J. (2002) MicroRNAs in plants. *Genes Dev.* 16, 1616–1626

452 33 Lee, H.-C. *et al.* (2010) Diverse Pathways Generate MicroRNA-like RNAs and Dicer-Independent
453 Small Interfering RNAs in Fungi. *Mol. Cell* 38, 803–814

454 34 Zhao, T. *et al.* (2007) A complex system of small RNAs in the unicellular green alga
455 *Chlamydomonas reinhardtii*. *Genes Dev.* 21, 1190–1203

456 35 Molnár, A. *et al.* (2007) miRNAs control gene expression in the single-cell alga *Chlamydomonas*
457 *reinhardtii*. *Nature* 447, 1126–1129

458 36 Cui, J. *et al.* (2014) Prediction and validation of potential pathogenic microRNAs involved in
459 *Phytophthora infestans* infection. *Mol. Biol. Rep.* 41, 1879–1889

460 37 Kang, S.-M. *et al.* (2013) Identification of microRNA-Size, Small RNAs in *Escherichia coli*. *Curr.*
461 *Microbiol.* 67, 609–613

462 38 Lee, H.-J. and Hong, S.-H. (2012) Analysis of microRNA-size, small RNAs in *Streptococcus mutans*
463 by deep sequencing. *FEMS Microbiol. Lett.* 326, 131–136

464 39 Nelson, J.W. and Breaker, R.R. (2017) The lost language of the RNA World. *Sci. Signal.* 10,
465 eaam8812

466 40 Choi, J.-W. *et al.* (2017) Tiny RNAs and their voyage via extracellular vesicles: Secretion of
467 bacterial small RNA and eukaryotic microRNA. *Exp. Biol. Med.* 242, 1475–1481

468 41 Gabriel, A.F. *et al.* (2019) Si vis pacem para bellum: A prospective in silico analysis of miRNA-
469 based plant defenses against fungal infections. *Plant Sci.* 288, 110241

470 42 Blenkiron, C. *et al.* (2016) Uropathogenic *Escherichia coli* Releases Extracellular Vesicles That Are
471 Associated with RNA. *PLOS ONE* 11, e0160440

472 43 Koeppen, K. *et al.* (2016) A Novel Mechanism of Host-Pathogen Interaction through sRNA in
473 Bacterial Outer Membrane Vesicles. *PLOS Pathog.* 12, e1005672

474 44 Buck, A.H. *et al.* (2014) Exosomes secreted by nematode parasites transfer small RNAs to
475 mammalian cells and modulate innate immunity. *Nat. Commun.* 5, 5488

476 45 Weiberg, A. *et al.* (2013) Fungal Small RNAs Suppress Plant Immunity by Hijacking Host RNA
477 Interference Pathways. *Science* 342, 118–123

478 46 Zhu, K. *et al.* (2017) Plant microRNAs in larval food regulate honeybee caste development. *PLOS*
479 *Genet.* 13, e1006946

480 47 Shahid, S. *et al.* (2018) MicroRNAs from the parasitic plant *Cuscuta campestris* target host
481 messenger RNAs. *Nature* 553, 82–85

482 48 Mu, J. *et al.* (2014) Interspecies communication between plant and mouse gut host cells through
483 edible plant derived exosome-like nanoparticles. *Mol. Nutr. Food Res.* 58, 1561–1573

484 49 Grüll, M.P. *et al.* (2018) Small extracellular particles with big potential for horizontal gene
485 transfer: membrane vesicles and gene transfer agents. *FEMS Microbiol. Lett.* 365,
486 50 Soler, N. and Forterre, P. (2020) Vesiduction: the fourth way of HGT. *Environ. Microbiol.* 22,
487 2457–2460

488 51 Bitto, N.J. *et al.* (2017) Bacterial membrane vesicles transport their DNA cargo into host cells. *Sci.*
489 *Rep.* 7, 7072

490 52 Breakfield, N.W. *et al.* (2012) High-resolution experimental and computational profiling of tissue-
491 specific known and novel miRNAs in Arabidopsis. *Genome Res.* 22, 163–176

492 53 Dessaux, Y. *et al.* (2016) Engineering the Rhizosphere. *Trends Plant Sci.* 21, 266–278

493 54 Bettenfeld, P. *et al.* (2020) Woody Plant Declines. What’s Wrong with the Microbiome? *Trends*
494 *Plant Sci.* 25, 381–394

495 55 Lapsansky, E.R. *et al.* (2016) Soil memory as a potential mechanism for encouraging sustainable
496 plant health and productivity. *Curr. Opin. Biotechnol.* 38, 137–142

497 56 Vorholt, J.A. *et al.* (2017) Establishing Causality: Opportunities of Synthetic Communities for
498 Plant Microbiome Research. *Cell Host Microbe* 22, 142–155

499 57 Huang, C.-Y. *et al.* (2019) Small RNAs – Big Players in Plant-Microbe Interactions. *Cell Host*
500 *Microbe* 26, 173–182

501 58 Cai, Q. *et al.* (2018) Cross-kingdom RNA trafficking and environmental RNAi — nature’s blueprint
502 for modern crop protection strategies. *Curr. Opin. Microbiol.* 46, 58–64

503 59 Zhang, H. *et al.* (2013) Feasibility, limitation and possible solutions of RNAi-based technology for
504 insect pest control: *RNAi-based technology in insect pest control.* *Insect Sci.* 20, 15–30

505 60 Dalakouras, A. *et al.* (2016) Induction of Silencing in Plants by High-Pressure Spraying of In vitro-
506 Synthesized Small RNAs. *Front. Plant Sci.* 07,

507 61 Laressergues, D. *et al.* (2015) Primary transcripts of microRNAs encode regulatory peptides.
508 *Nature* 520, 90–93

509 62 Couzigou, J.-M. *et al.* (2015) miRNA-encoded peptides (miPEPs): A new tool to analyze the roles
510 of miRNAs in plant biology. *RNA Biol.* 12, 1178–1180

511 63 Couzigou, J.-M. *et al.* (2017) Positive Gene Regulation by a Natural Protective miRNA Enables
512 Arbuscular Mycorrhizal Symbiosis. *Cell Host Microbe* 21, 106–112

513 64 Shabalina, S. and Koonin, E. (2008) Origins and evolution of eukaryotic RNA interference. *Trends*
514 *Ecol. Evol.* 23, 578–587

515 65 Borges, F. and Martienssen, R.A. (2015) The expanding world of small RNAs in plants. *Nat. Rev.*
516 *Mol. Cell Biol.* 16, 727–741

517 66 Zhao, C. *et al.* (2017) Salmonella small RNA fragment Sal-1 facilitates bacterial survival in infected
518 cells via suppressing iNOS induction in a microRNA manner. *Sci. Rep.* 7, 16979


519 67 Carrier, M.-C. *et al.* (2018) Broadening the Definition of Bacterial Small RNAs: Characteristics and
520 Mechanisms of Action. *Annu. Rev. Microbiol.* 72, 141–161

521 68 Zhou, J. *et al.* (2012) Identification of microRNA-like RNAs in a plant pathogenic fungus
522 *Sclerotinia sclerotiorum* by high-throughput sequencing. *Mol. Genet. Genomics* 287, 275–282

523 69 Fahlgren, N. *et al.* (2013) Phytophthora Have Distinct Endogenous Small RNA Populations That
524 Include Short Interfering and microRNAs. *PLoS ONE* 8, e77181

525 70 Rosenberg, E. and Zilber-Rosenberg, I. (2016) Microbes Drive Evolution of Animals and Plants:
 526 the Hologenome Concept. *mBio* 7, e01395-15, /mbio/7/2/e01395-15.atom
 527 71 Ren, B. *et al.* (2019) Rhizobial tRNA-derived small RNAs are signal molecules regulating plant
 528 nodulation. *Science* 365, 919–922
 529 72 Cai, Q. *et al.* (2018) Plants send small RNAs in extracellular vesicles to fungal pathogen to silence
 530 virulence genes. *Science* 360, 1126–1129
 531

532
 533



534

535 **Key figure**

536 **Model of cross-kingdom interactions in the rhizosphere through exchanges of**
 537 **extracellular vesicles, containing small RNA**

538 **Figure 1.** A suggested representation of plant-microbe interactions in the rhizosphere:
539 the selective power of the host plant is demonstrated through the rhizodeposits (in blue)
540 which play a great part in shaping the microbiota. Rhizodeposits are diverse in form and
541 function, they allow signalling with other organisms and their environment. Microbe
542 development in the rhizosphere depends on these compounds and is somewhat
543 regulated by them, as microbes feed preferentially on certain rhizodeposits as
544 substrates. In turn, microbes produce a variety of signalling molecules (in orange) such
545 as volatile organic compounds, phytohormones, quorum-sensing molecules, etc....
546 which are perceived by the plant. We suggest that plants and microbes also
547 communicate *via* extracellular vesicles (EVs) which contain small RNAs, and notably
548 microRNAs or microRNA-like RNAs. Host plant EVs containing microRNA (in green) can
549 modulate microbial gene expression to its advantage by regulating community
550 composition and activity (bold green arrow, representing the effect of microRNA-
551 mediated regulation). In return, microbes can also regulate host plant gene expression
552 through their EVs containing microRNA (in purple), thus controlling their close
553 environment and plant fitness (bold purple arrow). Additionally, the rhizospheric
554 microbiota members also interact between themselves and with other microbial
555 communities (black dashed arrows), potentially through EVs containing small RNAs. Not
556 to scale.


557 **Outstanding questions**

- 558 • Are rhizospheric bacteria and fungi able to incorporate plant microRNAs,
559 conveyed or not by extracellular vesicles? If so, can microbial genes be
560 modulated by plant microRNAs? What genes are targeted? To what effect?

- 561
- Are members of the rhizospheric microbiota capable of producing extracellular vesicles? If so, do they contain small RNAs? microRNAs? microRNA-like RNAs? To who are destined these vesicles: the host plant? other members of the community?
- 562
- 563
- 564
- As microRNA-size small RNAs exist in bacteria, what are the bacterial mechanisms of RNA interference, aside from the CRISPR-mediated system?
- 565
- 566
- To which extent can plant/microbial microRNAs play a role in shaping the rhizospheric microbiota and its associated functions? What role do plant microRNA play in microbial transcriptomics? Can they affect community activity and/or composition?
- 567
- 568
- 569
- 570
- Do the secreted microRNAs vary depending on environmental conditions, such as abiotic stresses, in order to modulate accordingly the surrounding organisms? What are the selective mechanisms involved in microRNAs sorting in extracellular vesicles?
- 571
- 572
- 573
- 574
- How can we use this potential novel interkingdom communication channel to develop holobiont engineering tools, in the future, such as miPEPs? How can these modifications be durable and what are the implications for long-term use?
- 575
- 576
- 577

578

579



580

581 **Figure from Box 1**

582 **Figure I.** A comparison of microRNA and microRNA-like RNA biogenesis in plants and
 583 **microbes.** In this figure, microRNA production and interspecies exchanges (dashed
 584 arrows), in the rhizosphere, are compared. The biosynthetic pathway of plant
 585 microRNAs has been extensively studied and is mainly composed of DCL and AGO
 586 proteins. Fungi produce miRNA through at least four different mechanisms [33]: after
 587 transcription of miRNA gene, the processing of pri-miRNA into mature miRNA involves
 588 a combination of enzymes (represented as *) such as Dicer, QDE-2 (an Argonaute
 589 protein), MRPL3 (an RNase III) and the exonuclease QIP. Interestingly, there is a Dicer-
 590 independent microRNA biogenesis pathway, which opens up an opportunity to consider

591 microRNA production in bacteria and archaea which lack Dicer enzymes, but do have
592 Argonaute-like proteins [33]. Details on bacterial msRNAs biogenesis are still to be
593 discovered, however multiple studies have demonstrated the presence of msRNAs in
594 different bacterial strains and have predicted similar hairpin RNA precursors, validated
595 by both msRNA and msRNA*. It is suggested that an RNA-restriction enzyme, such as
596 MazF found in *E. coli*, may be involved in msRNA processing [38]. Once the mature
597 microRNA is synthesized, many questions remain unanswered as to their movement
598 and function between species: are they transported in EVs? in OMVs? Do they travel
599 bound to an AGO protein or freely? What is the role of plant microRNAs on the
600 microbiota? What microbial genes are targeted and for what purpose? How does this
601 regulation shape the microbiota? In return, fungal small RNAs have been shown to use
602 plant AGO proteins in order to silence host genes, however it is unknown if bacteria use
603 this mechanism.

604

605