



The global Gan-Gross-Prasad conjecture for unitary groups: the endoscopic case

Raphaël Beuzart-Plessis, Pierre-Henri Chaudouard, Michal Zydor

► To cite this version:

Raphaël Beuzart-Plessis, Pierre-Henri Chaudouard, Michal Zydor. The global Gan-Gross-Prasad conjecture for unitary groups: the endoscopic case. 2024. <hal-02989495>

HAL Id: hal-02989495

<https://hal.science/hal-02989495v1>

Preprint submitted on 9 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

The global Gan-Gross-Prasad conjecture for unitary groups: the endoscopic case

Raphaël Beuzart-Plessis, Pierre-Henri Chaudouard and Michał Zydor

Abstract

In this paper, we prove the Gan-Gross-Prasad conjecture and the Ichino-Ikeda conjecture for unitary groups $U_n \times U_{n+1}$ in all the endoscopic cases. Our main technical innovation is the computation of the contributions of certain cuspidal data, called $*$ -generic, to the Jacquet-Rallis trace formula for linear groups. We offer two different computations of these contributions: one, based on truncation, is expressed in terms of regularized Rankin-Selberg periods of Eisenstein series and Flicker-Rallis intertwining periods. The other, built upon Zeta integrals, is expressed in terms of functionals on the Whittaker model. A direct proof of the equality between the two expressions is also given. Finally several useful auxiliary results about the spectral expansion of the Jacquet-Rallis trace formula are provided.

Contents

1	Introduction	2
1.1	The endoscopic cases of the Gan-Gross-Prasad conjecture	2
1.2	The spectral expansion of the Jacquet-Rallis trace formula for the linear groups . .	5
1.3	On the $*$ -generic contribution for the Jacquet-Rallis trace formula for the linear groups	7
1.4	Outline of the paper	10
1.5	Acknowledgement	11
2	Preliminaries	12
2.1	General notation	12
2.2	Algebraic groups and adelic points	12
2.3	Haar measures	14
2.4	Norms and Harish-Chandra Ξ function	16
2.5	Spaces of functions	17
2.6	Estimates on Fourier coefficients	21
2.7	Automorphic forms and representations	23
2.8	Relative characters	25
2.9	Decomposition according to cuspidal data and automorphic kernels	27
3	The spectral expansion of the Jacquet-Rallis trace formula for general linear groups	34
3.1	Notations	34
3.2	The coarse spectral expansion for Schwartz functions	35
3.3	Auxiliary expressions for I_χ	36
3.4	Convergence of a truncated kernel	38
3.5	Asymptotic formulas	48
4	Flicker-Rallis period of some spectral kernels	50
4.1	Flicker-Rallis intertwining periods and related distributions	50
4.2	A spectral expansion of a truncated integral	53
4.3	The case of $*$ -generic cuspidal data	54

5	The *-generic contribution in the Jacquet-Rallis trace formula	57
5.1	Relative characters	57
5.2	The *-generic contribution	59
6	Spectral decomposition of the Flicker-Rallis period for certain cuspidal data	62
6.1	Notation	62
6.2	Statements of the main results	63
6.3	Proof of Theorem 6.2.5.1.2	64
6.4	Proof of Theorem 6.2.6.1	67
6.5	Convergence of Zeta integrals	69
7	Canonical extension of the Rankin-Selberg period for certain cuspidal data	71
7.1	Statements of the main results	71
7.2	Proof of Theorem 7.1.3.1.2	72
7.3	Proof of Theorem 7.1.4.1	73
7.4	Convergence of Zeta integrals	75
8	Contributions of certain cuspidal data to the Jacquet-Rallis trace formula: second proof	76
8.1	Main result	76
8.2	Proof of Theorem 8.1.4.1	79
9	Flicker-Rallis functional computation	81
9.1	Local comparison	81
9.2	Global comparison	86
10	Proofs of the Gan-Gross-Prasad and Ichino-Ikeda conjectures	87
10.1	Identities among some global relative characters	87
10.2	Proof of Theorem 1.1.5.1	90
10.3	Proof of Theorem 1.1.6.1	90
A	Topological vector spaces	93

1 Introduction

1.1 The endoscopic cases of the Gan-Gross-Prasad conjecture

1.1.1. One of the main motivation of the paper is the obtention of the remaining cases, the so-called “endoscopic cases”, of the Gan-Gross-Prasad and the Ichino-Ikeda conjectures for unitary groups. To begin with, we shall give the main statements we prove.

1.1.2. Let E/F be a quadratic extension of number fields and c be the non-trivial element of the Galois group $\text{Gal}(E/F)$. Let \mathbb{A} be the ring of adèles of F . Let $n \geq 1$ be an integer. Let \mathcal{H}_n be the set of isomorphism classes of non-degenerate c -Hermitian spaces h over E of rank n . For any $h_n \in \mathcal{H}_n$, we identify h_n with a representative and we shall denote by $U(h_n)$ its automorphisms group. Let $h_0 \in \mathcal{H}_1$ be the element of rank 1 given by the norm $N_{E/F}$.

We attach to any $h \in \mathcal{H}_n$ the following algebraic groups over F :

- the unitary group U'_h of automorphisms of h ;
- the product of unitary groups $U_h = U(h) \times U(h \oplus h_0)$ where $h \oplus h_0$ denoted the orthogonal sum.

We have an obvious diagonal embedding $U'_h \hookrightarrow U_h$.

1.1.3. Arthur parameter. — Let G_n be the group of automorphisms of the E -vector space E^n . We view G_n as an F -group by Weil restriction. By a *Hermitian Arthur parameter*¹ of G_n , we mean an irreducible automorphic representation Π for which there exists a partition $n_1 + \dots + n_r = n$ of n and for any $1 \leq i \leq r$ a cuspidal automorphic representation Π_i of $G_{n_i}(\mathbb{A})$ such that

1. each Π_i is conjugate self-dual and the Asai L -function $L(s, \Pi_i, As^{(-1)^{n_i+1}})$ has a pole at $s = 1$;
2. the representations Π_i are mutually non-isomorphic for $1 \leq i \leq r$;
3. the representation Π is isomorphic to the full induced representation $\text{Ind}_P^{G_n}(\Pi_1 \boxtimes \dots \boxtimes \Pi_r)$ where P is a parabolic subgroup of G_n of Levi factor $G_{n_1} \times \dots \times G_{n_r}$.

Remark 1.1.3.1. — It is well-known (see [Fli88]) that condition 1 above is equivalent to the fact that Π_i is $(GL_{n_i, F}, \eta^{n_i+1})$ -distinguished in the sense of §4.1.2 below.

The integer r and the representations $(\Pi_i)_{1 \leq i \leq r}$ are unique (up to a permutation). We set $S_\Pi = (\mathbb{Z}/2\mathbb{Z})^r$.

Let $G = G_n \times G_{n+1}$. By a *Hermitian Arthur parameter* of G , we mean an automorphic representation of the form $\Pi = \Pi_n \boxtimes \Pi_{n+1}$ where Π_i is a Hermitian Arthur parameter of G_i for $i = n, n+1$. For such a Hermitian Arthur parameter, we set $S_\Pi = S_{\Pi_n} \times S_{\Pi_{n+1}}$.

1.1.4. Let $h \in \mathcal{H}_n$ and σ be a cuspidal automorphic representation of $U_h(\mathbb{A})$. We say that a Hermitian Arthur parameter Π of G_n is a *weak base-change* of σ if for almost all places of F that split in E , the local component Π_v is the split local base change of σ_v . If this is the case, we write $\Pi = BC(\sigma)$.

Remark 1.1.4.1. — By the work of Mok [Mok15] and Kaletha-Minguez-Shin-White [KMSW], we know that if σ admits a weak base-change then it admits a *strong base-change* that is a Hermitian Arthur parameter Π of G_n such that Π_v is the base-change of σ_v for every place v of F (where the local base-change in the ramified case is also constructed in *loc. cit.* and characterized by certain local character relations). Moreover, this is the case if and only if σ has a *generic Arthur parameter* in the sense of *loc. cit.* Besides, a result of Ramakrishnan [Ram18] implies that a weak base-change is automatically a strong base-change. Therefore, we could have used the notion of strong base-change instead. However, we prefer to stick with the terminology of weak base-change in order to keep the statement of the next theorem independent of [Mok15] and [KMSW].

1.1.5. Gan-Gross-Prasad conjecture. — Our first main result is the global Gan-Gross-Prasad conjecture [GGP12, Conjecture 24.1] in the case of $U(n) \times U(n+1)$ and can be stated as follows.

Theorem 1.1.5.1. — *Let Π be a Hermitian Arthur parameter of G . The following two statements are equivalent:*

1. *The complete Rankin-Selberg L -function of Π (including Archimedean places) satisfies*

$$L\left(\frac{1}{2}, \Pi\right) \neq 0;$$

2. *There exists $h \in \mathcal{H}_n$ and an irreducible cuspidal automorphic subrepresentation σ of U_h such that Π is a weak base change of σ and the period integral \mathcal{P}_h defined by*

$$\mathcal{P}_h(\varphi) = \int_{[U'_h]} \varphi(h) dh$$

induces a non-zero linear form on the space of σ .

¹Strictly speaking, it is a *discrete* Arthur parameter. By simplicity, we shall omit the adjective *discrete*.

Remark 1.1.5.2. — If the Arthur parameter is moreover simple (that is if Π is cuspidal), the theorem is proved by Beuzart-Plessis-Liu-Zhang-Zhu (cf. [BLZZ19, Theorem 1.7]). Previous works had to assume extra local hypothesis on Π , which implied that Π was also simple (see [Zha14b], [Xue19], [Beu16] and [BP18b]) or only proved the direction $2. \Rightarrow 1.$ of the theorem ([GJR09], [IY19], [JZ]).

As observed in [Zha14b, Theorem 1.2] and [BLZZ19, Theorem 1.8] we can deduce from Theorem 1.1.5.1 the following statement (whose proof is word for word that of [Zha14b]):

Theorem 1.1.5.3. — *Let Π_{n+1} be a Hermitian Arthur parameter of G_{n+1} . Then there exists a simple Hermitian Arthur parameter Π_n of G_n such that the Rankin-Selberg L -function satisfies:*

$$L\left(\frac{1}{2}, \Pi_n \times \Pi_{n+1}\right) \neq 0.$$

1.1.6. Ichino-Ikeda conjecture. — Let $\sigma = \bigotimes'_v \sigma_v$ be an irreducible cuspidal automorphic representation of U_h that is tempered everywhere in the following sense: for every place v , the local representation σ_v is tempered. By [Mok15] and [KMSW], σ admits a weak (hence a strong) base-change Π to G . Set

$$\mathcal{L}(s, \sigma) = \prod_{i=1}^{n+1} L(s + i - 1/2, \eta^i) \frac{L(s, \Pi)}{L(s + 1/2, \sigma, \text{Ad})}$$

where η denotes the quadratic idele class character associated to the extension E/F , $L(s, \eta^i)$ is the completed Hecke L -function associated to η^i and $L(s, \sigma, \text{Ad})$ is the completed adjoint L -function of σ (defined using the local Langlands correspondence for G from [Mok15], [KMSW]). We denote by $\mathcal{L}(s, \sigma_v)$ the corresponding quotient of local L -factors. For each place v of F , we define a *local normalized period* $\mathcal{P}_{h, \sigma_v}^{\natural} : \sigma_v \times \sigma_v \rightarrow \mathbb{C}$ as follows. It depends on the choice of a Haar measure on $U'_h(F_v)$ as well as an invariant inner product $(\cdot, \cdot)_v$ on σ_v and is given by

$$\mathcal{P}_{h, \sigma_v}^{\natural}(\varphi_v, \varphi'_v) = \mathcal{L}\left(\frac{1}{2}, \sigma_v\right)^{-1} \int_{U'_h(F_v)} (\sigma_v(h_v) \varphi_v, \varphi'_v)_v dh_v, \quad \varphi_v, \varphi'_v \in \sigma_v,$$

where, thanks to the temperedness assumption, the integral is absolutely convergent [Har14, Proposition 2.1] and the local factor $\mathcal{L}(s, \sigma_v)$ has no zero (nor pole) at $s = \frac{1}{2}$. Moreover, by [Har14, Theorem 2.12], if $\varphi = \bigotimes'_v \varphi_v \in \sigma$, then for almost all places v we have

$$(1.1.6.1) \quad \mathcal{P}_{h, \sigma_v}^{\natural}(\varphi_v, \varphi_v) = \text{vol}(U'_h(\mathcal{O}_v))(\varphi_v, \varphi_v)_v.$$

We also recall that the global representation σ has a natural invariant inner product given by

$$(\varphi, \varphi)_{\text{Pet}} = \int_{[U_h]} |\varphi(g)|^2 dg, \quad \varphi \in \sigma.$$

Our second main result is the global Ichino-Ikeda conjecture for unitary groups formulated in [Har14, Conjecture 1.3] and can be stated as follows (this result can be seen as a refinement of Theorem 1.1.5.1, the precise relation requiring the local Gan-Gross-Prasad conjecture and Arthur's multiplicity formula for unitary groups will not be discussed here).

Theorem 1.1.6.1. — *Assume that σ is a cuspidal automorphic representation of U_h that is tempered everywhere and let $\Pi = \Pi_n \boxtimes \Pi_{n+1}$ be the weak (hence the strong) base-change of σ to G . Suppose that we normalize the period integral \mathcal{P}_h and the Peterssen inner product $(\cdot, \cdot)_{\text{Pet}}$ by choosing the invariant Tamagawa measures² $d_{\text{Tam}} h$ and $d_{\text{Tam}} g$ on $U'_h(\mathbb{A})$ and $U_h(\mathbb{A})$ respectively. Assume also that the local Haar measures dh_v on $U'_h(F_v)$ factorize the Tamagawa measure:*

²We warn the reader that our convention is to include the global normalizing L -values in the definition of Tamagawa measures, cf. Section 2.3 for precise definitions.

$d_{\text{Tam}}h = \prod_v dh_v$. Then, for every nonzero factorizable vector $\varphi = \otimes'_v \varphi_v \in \sigma$, we have

$$\frac{|\mathcal{P}_h(\varphi)|^2}{(\varphi, \varphi)_{\text{Pet}}} = |S_{\Pi}|^{-1} \mathcal{L}\left(\frac{1}{2}, \sigma\right) \prod_v \frac{\mathcal{P}_{h, \sigma_v}^{\natural}(\varphi_v, \varphi_v)}{(\varphi_v, \varphi_v)_v}$$

where we recall that S_{Π} denotes the finite group $S_{\Pi_n} \times S_{\Pi_{n+1}}$.

Note that the product over all places in the theorem is well-defined by (1.1.6.1). Moreover, once again, this theorem is proved in [BLZZ19] under the extra assumption that Π is cuspidal (in which case $|S_{\Pi}| = 4$). Previous results in that direction includes [Zha14a], [Beu16], [BP18b] where some varying local assumptions on σ entailing the cuspidality of Π were imposed. In a slightly different direction, the paper [GL06] establishes the above identity up to an unspecified algebraic number under some arithmetic assumptions on σ .

1.2 The spectral expansion of the Jacquet-Rallis trace formula for the linear groups

1.2.1. Motivations. — As in [Zha14b], [Zha14a], [Xue19], [Beu16], [BP18b] and [BLZZ19], our proofs of Theorems 1.1.5.1 and 1.1.6.1 follow the strategy of Jacquet and Rallis [JR11] and are thus based on a comparison of *relative trace formulas* on unitary groups U_h for $h \in \mathcal{H}_n$ and the group G . Let's recall that these trace formulas have two different expansions: one, called the geometric side, in terms of distributions indexed by geometric classes and the other, called the spectral side, in terms of distributions indexed by cuspidal data. As usual, the point is to get enough test functions to first compare the geometric sides which gives a comparison of spectral sides.

For specific test functions, the trace formula boils down to a simple and quite easy equality between a sum of relative regular orbital integrals and a sum of relative characters attached to cuspidal representations. This is the simple trace formula used by Zhang in [Zha14b] and [Zha14a] to prove special cases of Theorems 1.1.5.1 and 1.1.6.1. In return one has to impose restrictive local conditions on the representations one considers.

In [Zyd16], [Zyd18], [Zyd20], Zydor established general Jacquet-Rallis trace formulas. Besides, in [CZ], Chaudouard-Zydor proved the comparison of all the geometric terms for matching test functions, that is functions with matching local orbital integrals. Using these results, Beuzart-Plessis-Liu-Zhang-Zhu in [BLZZ19] proved 1.1.5.1 and 1.1.6.1 when Π is cuspidal. Their main innovation is a construction of Schwartz test functions only detecting certain cuspidal data. In this way, they were able to construct matching test functions for which the spectral expansions reduce to some relative characters attached to cuspidal representations.

1.2.2. In this paper, we also want to use the construction of Beuzart-Plessis-Liu-Zhang-Zhu. But for this, we need two extra ingredients. First we need the slight extension of Zydor's work to the space of Schwartz test functions. For the geometric sides, this was done in [CZ]. For the test functions we need, the spectral side of the trace formulas for unitary groups still reduces to relative characters attached to cuspidal representations and we need nothing more. But, for the group G , we shall extend the spectral side of the trace formula to the space of Schwartz functions. Second there is an even more serious question: since the representation Π is no longer assumed to be cuspidal, the spectral contribution associated to Π is much more involved. In this section, we shall explain alternative and somewhat more tractable expressions for the spectral contributions in the trace formula for G . For the specific cuspidal datum attached to Π , we get a precise result as we shall see in section 1.3 below.

1.2.3. The spectral expansion for the Schwartz space. — Let $\mathfrak{X}(G)$ be the set of cuspidal data of G (see § 2.9.1). To χ is associated a direct invariant factor $L_{\chi}^2([G])$ of $L^2([G])$ (see [MW89, Chap. II] or section 2.9 for a review). Let f be a function in the Schwartz space $\mathcal{S}(G(\mathbb{A}))$ (cf. §2.5.2 for a definition). Let K_f , resp. $K_{f, \chi}$, be the kernel associated to the action by right convolution of f on $L^2([G])$, resp. $L_{\chi}^2([G])$.

Following [Zyd20] (see §3.2.3), we introduce the modified kernel $K_{f,\chi}^T$ depending on a parameter T in a certain real vector space. Set $H = G_n$ and $G' = \mathrm{GL}_{n,F} \times \mathrm{GL}_{n+1,F}$ both seen as subgroups of G (the embedding $H \hookrightarrow G$ being the “diagonal” one where the inclusion $G_n \hookrightarrow G_{n+1}$ is induced by the identification of E^n with the hyperplane of E^{n+1} of vanishing last coordinate). The following theorem is an extension to Schwartz functions of [Zyd20, théorème 0.1].

Theorem 1.2.3.1. — (see theorem 3.2.4.1)

1. For any T in a certain positive Weyl chamber, we have

$$\sum_{\chi \in \mathfrak{X}(G)} \int_{[H]} \int_{[G']} |K_{f,\chi}^T(h, g') \eta_{G'}(g')| dg' dh < \infty$$

2. Let $\eta_{G'}$ be the quadratic character of $G'(\mathbb{A})$ defined in §3.1.8. For each $\chi \in \mathfrak{X}(G)$, the integral

$$(1.2.3.1) \quad \int_{[H]} \int_{[G']} K_{f,\chi}^T(h, g') \eta_{G'}(g') dg' dh$$

coincides with a polynomial-exponential function in T whose purely polynomial part is constant and denoted by $I_\chi(f)$.

3. The distributions I_χ are continuous, left $H(\mathbb{A})$ -equivariant and right $(G'(\mathbb{A}), \eta_{G'})$ -equivariant. Moreover the sum

$$(1.2.3.2) \quad I(f) = \sum_{\chi} I_\chi(f)$$

is absolutely convergent and defines a continuous distribution.

The (coarse) spectral expansion of the trace formula for G is precisely the expression (1.2.3.2).

1.2.4. The definition of I_χ given in theorem 1.2.3.1 is convenient to relate the spectral expansion to the geometric expansion. However, to get more explicit forms of the distributions I_χ , we shall use the following three expressions:

$$(1.2.4.3) \quad \int_{[H]} \int_{[G']} (\Lambda_r^T K_{f,\chi})(h, g') \eta_{G'}(g') dg' dh$$

$$(1.2.4.4) \quad \int_{[H]} F^{G_{n+1}}(h, T) \int_{[G']} K_{f,\chi}(h, g') \eta_{G'}(g') dg' dh$$

$$(1.2.4.5) \quad \int_{[G']} F^{G_{n+1}}(g'_n, T) \int_{[H]} K_{f,\chi}(h, g') dh \eta_{G'}(g') dg'$$

Essentially they are given by integration of the kernel $K_{f,\chi}$ along $[H] \times [G']$. However, to have a convergent expression for a general χ , one needs to use some truncation depending on the same parameter T as above. We introduce the Ichino-Yamana truncation operator, denoted by Λ_r^T , whose definition is recalled in §3.3.2. In (1.2.4.3), we apply it to the left-variable of $K_{f,\chi}$. But one can also use the Arthur characteristic function $F^{G_{n+1}}(\cdot, T)$ whose definition is recalled in 3.3.4. In (1.2.4.4), this function is evaluated at $h \in H(\mathbb{A})$ through the embedding $H = G_n \hookrightarrow G_{n+1}$. In (1.2.4.5), it is evaluated at the component g'_n of the variable $g' = (g'_n, g'_{n+1}) \in G'(\mathbb{A}) = \mathrm{GL}_n(\mathbb{A}) \times \mathrm{GL}_{n+1}(\mathbb{A})$.

The link with the distribution I_χ is provided by the following theorem (which is a combination of Propositions 3.3.3.1 and 3.3.5.1 and Theorem 3.3.7.1). Note that we shall not need the full strength of the theorem in this paper. However it will be used in a greater generality in a subsequent paper.

Theorem 1.2.4.1. — Let $f \in \mathcal{S}(G(\mathbb{A}))$ and $\chi \in \mathfrak{X}(G)$.

1. For any T in some positive Weyl chamber, the expressions (1.2.4.3), (1.2.4.4) and (1.2.4.5) are absolutely convergent.

2. Each of the three expressions is asymptotically equal (in the technical sense of §3.3.6) to a polynomial-exponential function of T whose purely polynomial term is constant and equal to $I_\chi(f)$.

1.3 On the *-generic contribution for the Jacquet-Rallis trace formula for the linear groups

1.3.1. From now on we assume that the cuspidal datum χ is relevant *-generic that is χ is the class of a pair (M, π) with the property that the normalized induction $\Pi := \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})}(\pi)$, where we have fixed a parabolic subgroup P with Levi component M , is a Hermitian Arthur parameter of G . To Π we associate, following [Zha14a, §3.4], a *relative character* I_Π . The precise definition of this object is recalled in §8.1.3. Let us just say here that it is associated to two functionals λ and β_η on the Whittaker model $\mathcal{W}(\Pi, \psi_N)$ of Π , where ψ_N is a certain generic automorphic character of the standard maximal unipotent subgroup N of G , that naturally show up in integrals of Rankin-Selberg type. More precisely, λ is the value at $s = \frac{1}{2}$ of a family of Zeta integrals, studied by Jacquet-Piatetski-Shapiro-Shalika [JPSS83], representing the Rankin-Selberg L -function $L(s, \Pi)$ whereas β_η is essentially the pole at $s = 1$ of another family of Zeta integrals, first introduced by Flicker [Fli88], representing the (product of) Asai L -functions $L(s, \Pi, \text{As}_G) := L(s, \Pi_n, \text{As}^{(-1)^{n+1}})L(s, \Pi_{n+1}, \text{As}^{(-1)^n})$. The relative character I_Π is then given in terms of these functionals by

$$(1.3.1.1) \quad I_\Pi(f) = \sum_{\varphi \in \Pi} \lambda(\Pi(f)W_\varphi) \overline{\beta_\eta(W_\varphi)}, \quad f \in \mathcal{S}(G(\mathbb{A})),$$

where the sum runs over an orthonormal basis of Π (for the Petersson inner product) and W_φ denotes the Whittaker function associated to the Eisenstein series $E(\varphi)$ (obtained, as usual, by integrating $E(\varphi)$ against ψ_N^{-1} over $[N]$).

The following is our main technical result whose proof occupies most part of the paper.

Theorem 1.3.1.1. — *Let χ be a cuspidal datum associated to a Hermitian Arthur parameter Π as above. Then, for every function $f \in \mathcal{S}(G(\mathbb{A}))$ we have*

$$I_\chi(f) = 2^{-\dim(A_M)} I_\Pi(f)$$

where A_M denotes the maximal central split torus of M .

Remark 1.3.1.2. — It is perhaps worth emphasizing that the contribution of χ is *purely discrete* in the Jacquet-Rallis trace formula. Such a phenomenon happens in Jacquet relative trace formula, see [Lap06]. By contrast, the contribution of the same kind of cuspidal datum χ to the Arthur-Selberg trace formula is purely continuous (unless, of course, if Π is cuspidal).

We shall provide two different proofs of theorem 1.3.1.1, one based on truncations, the other using integral representations of Asai and Rankin-Selberg L -functions. Let's explain separately the main steps of each approach.

1.3.2. A journey through truncations. — We first begin with the approach based on truncations. The first step is to get a spectral decomposition of the function

$$(1.3.2.2) \quad \int_{[G']} K_{f, \chi}(g, g') \eta_{G'}(g') dg'$$

of the variable $g \in [G]$.

The kernel itself $K_{f, \chi}$ has a well-known spectral decomposition based on the Langlands decomposition. Then the problem is basically to invert an adelic integral and a complex integral. It is solved by Lapid in [Lap06] (up to some non-explicit constants) but we will use a slightly different

method avoiding delicate Lapid's contour moving. Instead we replace the integral (1.3.2.2) by its truncated version

$$(1.3.2.3) \quad \int_{[G']} (K_{f,\chi} \Lambda_m^T)(g, g') \eta_{G'}(g') dg'.$$

where the *mixed truncation operator* Λ_m^T defined by Jacquet-Lapid-Rogawski [JLR99] is applied to the right variable of the kernel. We can recover (1.3.2.2) by taking the limit when $T \rightarrow +\infty$. It is easy to get the spectral decomposition of (1.3.2.3) (see Proposition 4.2.3.3). Using an analog of the famous Maaß-Selberg relations due to Jacquet-Lapid-Rogawski (see [JLR99] and Lemma 4.3.6.2 below), we get in proposition 4.3.6.1) that (1.3.2.3) is equal to a finite sum of contributions (up to an explicit constant) of the following type

$$(1.3.2.4) \quad \int_{i\mathfrak{a}_P^{G,*}} \sum_{Q \in \mathcal{P}(M)} J_{Q,\chi}(g, \lambda, f) \frac{\exp(-\langle \lambda, T_Q \rangle)}{\theta_Q(-\lambda)} d\lambda.$$

Here it suffices to say that $i\mathfrak{a}_P^{G,*}$ is some space of unramified unitary characters and that $J_{Q,\chi}(g, \lambda, f)$ is a certain relative character built upon Flicker-Rallis intertwining periods (introduced by Jacquet-Lapid-Rogawski). The integrand is a familiar expression of Arthur's theory of (G, M) -families with quite standard notations. It turns out that the family $(J_{Q,\chi}(g, \lambda, f))_{Q \in \mathcal{P}(M)}$ is indeed an Arthur (G, M) -family of Schwartz functions in the parameter λ . Let's emphasize that this Schwartz property relies in fact on deep estimates introduced by Lapid in [Lap06] and [Lap13]. By a standard argument, it is then easy to get the limit of (1.3.2.4) when $T \rightarrow +\infty$ which gives the spectral decomposition of (1.3.2.2) (see Theorem 4.3.3.1). Note that the spectral decomposition we get is already discrete at this stage.

From this result, one gets the equality

$$(1.3.2.5) \quad \int_{[H]} \int_{[G']} (\Lambda_r^T K_{f,\chi})(h, g') \eta_{G'}(g') dh dg' = 2^{-\dim(A_M)} I_{P,\pi}(f).$$

The left-hand side has been defined in §1.2.4 and the relative character $I_{P,\pi}$ is defined as follows:

$$\sum_{\varphi \in \Pi} I_{RS}(\Pi(f)\varphi) \cdot \overline{J_\eta(\varphi)}$$

where the sum is over an orthonormal basis, $I_{RS}(\varphi)$ is the regularized Rankin-Selberg period of the Eisenstein series $E(\varphi)$ defined by Ichino-Yamana and $J_\eta(\varphi)$ is a Flicker-Rallis intertwining period (for more detail we refer to §5.1.5).

In particular, the left-hand side of (1.3.2.5) does not depend on T . So Theorem 1.2.4.1 implies

Theorem 1.3.2.1. — (see theorem 5.2.1.1 for a slightly more precise statement)

$$I_\chi(f) = 2^{-\dim(A_M)} I_{P,\pi}(f).$$

Remark 1.3.2.2. — As the reading of Section 10.2 should make it clear, this statement suffices to prove the Gan-Gross-Prasad conjecture namely Theorem 1.1.5.1. However, to get the Ichino-Ikeda conjecture, namely Theorem 1.1.6.1, we will want to use statements about comparison of local relative characters written in terms of Whittaker functions. For this purpose, Theorem 1.3.1.1 will be more convenient.

The link between regularized Rankin-Selberg period of Eisenstein series and Whittaker functionals has been investigated by Ichino-Yamana (see [IY15]). The following theorem relates the Flicker-Rallis intertwining periods to the functional $\beta_\eta(W_\varphi)$ in (1.3.1.1). It uses a local unfolding method inspired from [FLO12, Appendix A] (see Chapter 9).

Theorem 1.3.2.3. — *For all $\varphi \in \Pi$, we have*

$$J_\eta(\varphi) = \beta_\eta(W_\varphi)$$

In this way, one proves the following theorem which implies Theorem 1.3.1.1.

Theorem 1.3.2.4. —

$$I_{P,\pi} = I_\Pi.$$

1.3.3. Second proof: the use of Zeta integrals. — The spectral decomposition of (1.3.2.2) essentially boils down to a spectral expansion of the period integral

$$P_{G',\eta}(\varphi) := \int_{[G']} \varphi(g') \eta_{G'}(g') dg'$$

for test functions $\varphi \in \mathcal{S}_\chi([G])$, where $\mathcal{S}_\chi([G])$ denotes the *Schwartz space* of $[G]$ consisting of smooth functions rapidly decaying with all their derivatives that are “supported on χ ” (see Section 2.5 for a precise definition). Choose a parabolic subgroup $P = MN_P$ with Levi component M . By Langlands L^2 spectral decomposition and of the special form of χ , any $\varphi \in \mathcal{S}_\chi([G])$ admits a spectral decomposition

$$(1.3.3.6) \quad \varphi = \int_{i\mathfrak{a}_M^*} E(\varphi_\lambda) d\lambda$$

where $i\mathfrak{a}_M^*$ denotes the real vector space of unramified unitary characters of $M(\mathbb{A})$ and φ_λ belongs to the normalized induction space $\text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})}(\pi \otimes \lambda)$ and $E(\varphi_\lambda)$ is the associated Eisenstein series.

Theorem 1.3.3.1. — *For every $\varphi \in \mathcal{S}_\chi([G])$, we have*

$$P_{G',\eta}(\varphi) = 2^{-\dim(A_M)} \beta_\eta(W_{\varphi_0}).$$

where W_{φ_0} stands for the Whittaker function of the Eisenstein series $E(\varphi_0)$.

The proof of Theorem 1.3.3.1 is close to the computation by Flicker [Fli88] of the Flicker-Rallis period of cusp forms in terms of an Asai L -function and local Zeta integrals. More precisely, we first realize $P_{G',\eta}(\varphi)$ as the residue at $s = 1$ of the inner product of the restriction $\varphi|_{[G']}$ with some Eisenstein series $E(s, \phi)$ (where ϕ is an auxiliary Schwartz function on $\mathbb{A}^n \oplus \mathbb{A}^{n+1}$). Mimicking the unfolding of *loc. cit.* we connect this inner product with an Eulerian Zeta integral $Z^{\text{FR}}(s, W_\varphi, \phi)$ involving the Whittaker function W_φ of φ (obtained as before by integration against ψ_N^{-1}). We should emphasize here that, since φ is not a cusp form, the unfolding gives us more terms but using the special nature of the cuspidal datum χ we are able to show that these extra terms do not contribute to the residue at $s = 1$. The formation of $Z^{\text{FR}}(s, W_\varphi, \phi)$ commutes with the spectral expansion (1.3.3.6) when $\Re(s) \gg 1$ and, as follows from the local theory, the Zeta integrals $Z^{\text{FR}}(s, W_{\varphi_\lambda}, \phi)$ for $\lambda \in i\mathfrak{a}_M^*$ are essentially Asai L -functions whose meromorphic continuations, poles and growths in vertical strips are known. Combining this with an application of the Phragmen-Lindelöf principle, we are then able to deduce Theorem 1.3.3.1.

Let us mention here that, as in the proof of (1.3.2.5), a key point is the fact (due to Lapid ([Lap13] or [Lap06])) that the spectral transform $\lambda \mapsto \varphi_\lambda$ is, in a suitable technical sense, “Schwartz” that is rapidly decreasing together with all its derivatives.

The second step is to integrate (1.3.2.2) over $g \in [H]$. To do so, we define a regularization of the integral over $[H]$ that doesn’t require truncation. More precisely, denoting by $\mathcal{T}([G])$ the space of functions of *uniform moderate growth* on $[G]$, we can define the “ χ -part” $\mathcal{T}_\chi([G])$ of $\mathcal{T}([G])$ (see Section 2.5) of which $\mathcal{S}_\chi([G])$ is a dense subspace. Moreover, starting with $\varphi \in \mathcal{T}([G])$ we can

also form its Whittaker function W_φ and consider the usual Rankin-Selberg integral $Z^{\text{RS}}(s, W_\varphi)$ that converges for $\Re(s) \gg 1$ and represents, when φ is an automorphic form, the Rankin-Selberg L -function for $G_n \times G_{n+1}$.

Theorem 1.3.3.2. — (see theorems 7.1.3.1 and 7.1.4.1) *The functional*

$$\varphi \in \mathcal{S}_\chi([G]) \mapsto \int_{[H]} \varphi(h) dh$$

extends by continuity to a functional on $\mathcal{T}_\chi([G])$ denoted by $\varphi \in \mathcal{T}_\chi([G]) \mapsto \int_{[H]}^ \varphi(h) dh$. Moreover, for every $\varphi \in \mathcal{T}_\chi([G])$, the Zeta function $s \mapsto Z^{\text{RS}}(s, W_\varphi)$ extends to an entire function on \mathbb{C} and we have*

$$\int_{[H]}^* \varphi(h) dh = Z^{\text{RS}}\left(\frac{1}{2}, W_\varphi\right).$$

The proof of this theorem is similar to that of Theorem 1.3.3.1: we first show that, for $\varphi \in \mathcal{S}_\chi([G])$ and $\Re(s) \gg 1$, we have

$$\int_{[H]} \varphi(h) |\det h|^s dh = Z^{\text{RS}}\left(s + \frac{1}{2}, W_\varphi\right)$$

by mimicking the usual unfolding for the Rankin-Selberg integral. Once again, as φ is not necessarily a cusp form, we get extra terms in the course of the unfolding but, thanks to the special nature of the cuspidal datum χ , we are able to show that they all vanish. At this point, we use the spectral decomposition (1.3.3.6) to express $Z^{\text{RS}}(s, W_\varphi)$ as the integral of $Z^{\text{RS}}(s, W_{\varphi_\lambda})$ when $\Re(s) \gg 1$. By Rankin-Selberg theory, $Z^{\text{RS}}(s, W_{\varphi_\lambda})$ is essentially a Rankin-Selberg L -function whose meromorphic continuation, location of the poles, control in vertical strips and functional equation are known. Combining this with another application of the Phragmen-Lindelöf principle, we are able to bound $Z^{\text{RS}}(\frac{1}{2}, W_\varphi) = \int_{[H]} \varphi(h) dh$ in terms of $Z^{\text{RS}}(s, W_\varphi)$ for $\Re(s) \gg 1$ and this readily gives the theorem.

One direct consequence of Theorem 1.3.3.2 is that the regularized period $\int_{[H]}^* E(h, \Pi(f)\varphi) dh$ coincides with $Z^{\text{RS}}(\frac{1}{2}, \Pi(f)W_\varphi) = \lambda(\Pi(f)W_\varphi)$. Thus by a combination of Theorems 1.3.3.1 and 1.3.3.2, we get

$$(1.3.3.7) \quad \int_{[H]}^* \int_{[G']} K_{f,\chi}(h, g') \eta_{G'}(g') dg' dh = 2^{-\dim(A_M)} I_\Pi(f).$$

Finally we have to show that the left-hand side is equal to $I_\chi(f)$. In fact, we show (see Theorem 8.1.4.1 and §8.2.3) that we have

$$\int_{[H]}^* \int_{[G']} K_{f,\chi}(h, g') \eta_{G'}(g') dg' dh = \int_{[G']} \int_{[H]} K_{f,\chi}(h, g') dh \eta_{G'}(g') dg'$$

where the right-hand side is (conditionally) convergent. We can conclude that it is equal to $I_\chi(f)$ by applying Theorem 1.2.4.1 to the expression (1.2.4.5).

1.4 Outline of the paper

We now give a quick outline of the content of the paper. Chapter 2 contains preliminary material. Notably, we fix most notation to be used in the paper, we explain our convention on normalization of measures, we introduce the various spaces of functions we need and we discuss several properties of Langlands decomposition along cuspidal data as well as kernel functions that are important

for us. Chapter 3 contains the statements and proofs concerning the spectral expansion of the Jacquet-Rallis trace formula for G that were discussed in Section 1.2 above.

In Chapter 4, we introduce the Flicker-Rallis intertwining periods and prove the spectral expansion of the Flicker-Rallis period of the kernel associated to $*$ -generic cuspidal datum. In Chapter 5 we deduce from it Theorem 1.3.2.1 namely the spectral expansion for $I_\chi(f)$. Chapters 6 and 7 are devoted to the proofs of Theorems 1.3.3.1 and 1.3.3.2 respectively. These two theorems are combined in Chapter 8 to give another proof of the spectral expansion of $I_\chi(f)$ (Theorem 1.3.1.1). In Chapter 9, we relate the Flicker-Rallis intertwining periods to the functional β_η . From this, we deduce Theorem 1.3.2.3 and Theorem 1.3.2.4. The final Chapter 10 explain the deduction of Theorems 1.1.5.1 and 1.1.6.1 from Theorem 1.3.1.1. Finally, we have gathered in Appendix A some useful facts on topological vector spaces and holomorphic functions valued in them. It contains in particular some variations on the theme of the Phragmen-Lindelöf principle for such functions that will be crucial for the proofs of Theorems 1.3.3.1 and 1.3.3.2.

1.5 Acknowledgement

The project leading to this publication of R. B.-P. has received funding from Excellence Initiative of Aix-Marseille University-A*MIDEX, a French “Investissements d’Avenir” programme.

2 Preliminaries

2.1 General notation

2.1.1. For f and g two positive functions on a set X , we say that f is *essentially bounded* by g and we write

$$f(x) \ll g(x), \quad x \in X,$$

if there exists a constant $C > 0$ such that $f(x) \leq Cg(x)$ for every $x \in X$. If we want to emphasize that the constant C depends on auxiliary parameters y_1, \dots, y_k , we will write $f(x) \ll_{y_1, \dots, y_k} g(x)$. We say that the functions f and g are *equivalent* and we write

$$f(x) \sim g(x), \quad x \in X,$$

if $f(x) \ll g(x)$ and $g(x) \ll f(x)$. If moreover f and g take values in the set of real numbers greater than 1, we say that f and g are *weakly equivalent*, and we write

$$f(x) \approx g(x), \quad x \in X,$$

if there exists $N > 0$ such that

$$g(x)^{1/N} \ll f(x) \ll g(x)^N, \quad x \in X.$$

2.1.2. For every $C, D \in \mathbb{R} \cup \{-\infty\}$ with $D > C$, we set $\mathcal{H}_{>C} = \{z \in \mathbb{C} \mid \Re(z) > C\}$ and $\mathcal{H}_{]C, D[} = \{z \in \mathbb{C} \mid C < \Re(z) < D\}$. A *vertical strip* is a subset of \mathbb{C} which is the closure of $\mathcal{H}_{]C, D[}$ for some $C, D \in \mathbb{R}$ with $D > C$.

When f is a meromorphic function on some open subset U of \mathbb{C} and $s_0 \in U$, we denote by $f^*(s_0)$ the leading term in the Laurent expansion of f at s_0 .

2.1.3. When G is a group and we have a space of functions on it invariant by right translation, we denote by R the corresponding representation of G . If G is a Lie group and the representation is differentiable, we will also denote by the same letter the induced action of the Lie algebra or of its associated enveloping algebra. If G is a topological group equipped with a bi-invariant Haar measure, we denote by $*$ the convolution product (whenever it is well-defined).

2.2 Algebraic groups and adelic points

2.2.1. Let F be a number field and \mathbb{A} its adèle ring. We write \mathbb{A}_f for the ring of finite adèles and $F_\infty = F \otimes_{\mathbb{Q}} \mathbb{R}$ for the product of Archimedean completions of F so that $\mathbb{A} = F_\infty \times \mathbb{A}_f$. Let V_F be the set of places of F and $V_{F, \infty} \subset V_F$ be the subset of Archimedean places. For every $v \in V_F$, we let F_v be the local field obtained by completion of F at v . We denote by $|\cdot|$ the morphism $\mathbb{A}^\times \rightarrow \mathbb{R}_+^\times$ given by the product of normalized absolute values $|\cdot|_v$ on each F_v . For any finite subset $S \subset V_F \setminus V_{F, \infty}$, we denote by \mathcal{O}_F^S the ring of S -integers in F .

2.2.2. Let G be an algebraic group defined over F . We denote by N_G the unipotent radical of G . Let $X^*(G)$ be the group of characters of G defined over F . Let $\mathfrak{a}_G^* = X^*(G) \otimes_{\mathbb{Z}} \mathbb{R}$ and $\mathfrak{a}_G = \text{Hom}_{\mathbb{Z}}(X^*(G), \mathbb{R})$. We have a canonical pairing

$$(2.2.2.1) \quad \langle \cdot, \cdot \rangle : \mathfrak{a}_G^* \times \mathfrak{a}_G \rightarrow \mathbb{R}.$$

We have also a canonical homomorphism

$$(2.2.2.2) \quad H_G : G(\mathbb{A}) \rightarrow \mathfrak{a}_G$$

such that $\langle \chi, H_G(g) \rangle = \log |\chi(g)|$ for any $g \in G(\mathbb{A})$. The kernel of H_G is denoted by $G(\mathbb{A})^1$. We define $[G] = G(F) \backslash G(\mathbb{A})$ and $[G]^1 = G(F) \backslash G(\mathbb{A})^1$.

We let \mathfrak{g}_∞ be the Lie algebra of $G(F_\infty)$, $\mathcal{U}(\mathfrak{g}_\infty)$ be the enveloping algebra of its complexification and $\mathcal{Z}(\mathfrak{g}_\infty) \subset \mathcal{U}(\mathfrak{g}_\infty)$ be its center.

2.2.3. From now on we assume that G is also reductive. We will mainly use the notations of Arthur's works. For the convenience of the reader, we recall some of them. Let P_0 be a parabolic subgroup of G defined over F and minimal for these properties. Let M_0 be a Levi factor of P_0 defined over F .

We call a parabolic (resp. and semi-standard, resp. and standard) subgroup of G a parabolic subgroup of G defined over F (resp. which contains M_0 , resp. which contains P_0). For any semi-standard parabolic subgroup P , we have a Levi decomposition $P = M_P N_P$ where M_P contains M_0 and we define $[G]_P = M_P(F) N_P(\mathbb{A}) \backslash G(\mathbb{A})$. We call a Levi subgroup of G (resp. semi-standard, resp. standard) a Levi factor defined over F of a parabolic subgroup of G (resp. semi-standard, resp. standard).

2.2.4. Let $K = \prod_{v \in V_F} K_v \subset G(\mathbb{A})$ be a "good" maximal compact subgroup in good position relative to M_0 . We write

$$K = K_\infty K^\infty$$

where $K_\infty = \prod_{v \in V_{F,\infty}} K_v$ and $K^\infty = \prod_{v \in V_F \setminus V_{F,\infty}} K_v$. We let \mathfrak{k}_∞ be the Lie algebra of K_∞ , $\mathcal{U}(\mathfrak{k}_\infty)$ be the enveloping algebra of its complexification and $\mathcal{Z}(\mathfrak{k}_\infty) \subset \mathcal{U}(\mathfrak{k}_\infty)$ be its center.

2.2.5. Let P be a semi-standard parabolic subgroup. We extend the homomorphism $H_P : P(\mathbb{A}) \rightarrow \mathfrak{a}_P$ (see (2.2.2.2)) into the Harish-Chandra map

$$H_P : G(\mathbb{A}) \rightarrow \mathfrak{a}_P$$

in such a way that for every $g \in G(\mathbb{A})$ we have $H_P(g) = H_P(p)$ where $p \in P(\mathbb{A})$ is given by the Iwasawa decomposition namely $g \in pK$.

2.2.6. Let A be a split torus over F . Then, A admits an unique split model over \mathbb{Q} (which is also the maximal split subtorus of $\text{Res}_{F/\mathbb{Q}}(A)$) and by abuse of notation we denote by $A(\mathbb{R})$ the group of \mathbb{R} -points of this model. In particular, this gives an embedding $\mathbb{R}^\times \subset F_\infty^\times \subset \mathbb{A}^\times$. We also write A^∞ for the neutral component of $A(\mathbb{R})$. Let A_G be the maximal central F -split torus of G . We define $[G]_0 = A_G^\infty G(F) \backslash G(\mathbb{A})$.

Let P be a semi-standard parabolic subgroup of G . We define $A_P = A_{M_P}$, $A_P^\infty = A_{M_P}^\infty$ and $[G]_{P,0} = A_P^\infty M_P(F) N_P(\mathbb{A}) \backslash G(\mathbb{A})$. The restrictions maps $X^*(P) \rightarrow X^*(M_P) \rightarrow X^*(A_P)$ induce isomorphisms $\mathfrak{a}_P^* \simeq \mathfrak{a}_{M_P}^* \simeq \mathfrak{a}_{A_P}^*$. Let $\mathfrak{a}_0^* = \mathfrak{a}_{P_0}^*$, $\mathfrak{a}_0 = \mathfrak{a}_{P_0}$, $A_0 = A_{P_0}$ and $A_0^\infty = A_{P_0}^\infty$.

2.2.7. For any semi-standard parabolic subgroups $P \subset Q$ of G , the restriction map $X^*(Q) \rightarrow X^*(P)$ induces maps $\mathfrak{a}_Q^* \rightarrow \mathfrak{a}_P^*$ and $\mathfrak{a}_P \rightarrow \mathfrak{a}_Q$. The first one is injective whereas the kernel of the second one is denoted by \mathfrak{a}_P^Q . The restriction map $X^*(A_P) \rightarrow X^*(A_Q)$ gives a surjective map $\mathfrak{a}_P^* \rightarrow \mathfrak{a}_Q^*$ whose kernel is denoted by $\mathfrak{a}_P^{Q,*}$. We get also an injective map $\mathfrak{a}_Q \rightarrow \mathfrak{a}_P$. In this way, we get dual decompositions $\mathfrak{a}_P = \mathfrak{a}_P^Q \oplus \mathfrak{a}_Q$ and $\mathfrak{a}_P^* = \mathfrak{a}_P^{Q,*} \oplus \mathfrak{a}_Q^*$. Thus we have projections $\mathfrak{a}_0 \rightarrow \mathfrak{a}_P^Q$ and $\mathfrak{a}_0^* \rightarrow \mathfrak{a}_P^{Q,*}$ which we will denote by $X \mapsto X_P^Q$.

We denote by $\mathfrak{a}_{P,\mathbb{C}}^{Q,*}$ and $\mathfrak{a}_{P,\mathbb{C}}^Q$ the \mathbb{C} -vector spaces obtained by extension of scalars from $\mathfrak{a}_P^{Q,*}$ and \mathfrak{a}_P^Q . We still denote by $\langle \cdot, \cdot \rangle$ the pairing (2.2.2.1) we get by extension of the scalars to \mathbb{C} . We have a decomposition

$$\mathfrak{a}_{P,\mathbb{C}}^{Q,*} = \mathfrak{a}_P^{Q,*} \oplus i \mathfrak{a}_P^{Q,*}$$

where $i^2 = -1$. We shall denote by \Re and \Im the associated projections and call them real and imaginary parts. The same holds for the dual spaces $\mathfrak{a}_{P,\mathbb{C}}^Q$. In the obvious way, we define the complex conjugate denoted by $\bar{\lambda}$ of $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^{Q,*}$.

2.2.8. Let Ad_P^Q be the adjoint action of M_P on the Lie algebra of $M_Q \cap N_P$. Let ρ_P^Q be the unique element of $\mathfrak{a}_P^{Q,*}$ such that for every $m \in M_P(\mathbb{A})$

$$|\det(\text{Ad}_P^Q(m))| = \exp(\langle 2\rho_P^Q, H_P(m) \rangle).$$

For $Q = G$, the exponent G is omitted. For every $g \in G(\mathbb{A})$, we set

$$\delta_P(g) = \exp(\langle 2\rho_P, H_P(g) \rangle)$$

so that, in particular, the restriction of δ_P to $P(\mathbb{A})$ coincides with the modular character of the latter.

2.2.9. Let $P'_0 = M_0 N_{P'_0}$ be a minimal semi-standard parabolic subgroup such that $P'_0 \subset P$. Let $\Delta_{P'_0}^P$ be the set of simple roots of A_0 in $M_P \cap P'_0$. We denote this set by Δ_0^P if $P'_0 = P_0$. Let Δ_P be the image of $\Delta_{P'_0}^P \setminus \Delta_{P'_0}^P$ (viewed as a subset of \mathfrak{a}_0^*) by the projection $\mathfrak{a}_0^* \rightarrow \mathfrak{a}_P^*$. It does not depend on the choice of P'_0 . More generally one defines Δ_P^Q . We have also the set of coroots $\Delta_P^{Q,\vee} \subset \mathfrak{a}_P^Q$. By duality, we get a set of simple weights $\hat{\Delta}_P^Q$. The sets Δ_P^Q and $\hat{\Delta}_P^Q$ determine open cones in \mathfrak{a}_0 whose characteristic functions are denoted respectively by τ_P^Q and $\hat{\tau}_P^Q$. If $Q = G$, the exponent G is omitted. We set

$$A_P^{\infty,+} = \{a \in A_P^\infty \mid \langle \alpha, H_P(a) \rangle \geq 0, \forall \alpha \in \Delta_P\}.$$

In the same way, we define \mathfrak{a}_P^+ and $\mathfrak{a}_P^{*,+}$ (this time using coroots).

2.2.10. Weyl group. — Let W be the Weyl group of (G, A_0) that is the quotient by M_0 of the normalizer of A_0 in $G(F)$. For $P = M_P N_P$ and $Q = M_Q N_Q$ two standard parabolic subgroups of G , we denote by $W(P, Q)$ the set of $w \in W$ such that $w\Delta_0^P = \Delta_0^Q$. For $w \in W(P, Q)$, we have $wM_P w^{-1} = M_Q$. When $P = Q$, the group $W(P, P)$ is simply denoted by $W(P)$. Sometimes, we shall also denote $W(P, Q)$ by $W(M_P, M_Q)$ if we want to emphasize the Levi components (and $W(M_P) = W(P)$).

2.2.11. Let M be a standard Levi subgroup of G . We denote by $\mathcal{P}(M)$ the set of semi-standard parabolic subgroups P of G such that $M_P = M$. There is an unique element $P \in \mathcal{P}(M)$ which is standard and the map

$$(2.2.11.3) \quad (Q, w) \mapsto w^{-1}Qw$$

induces a bijection from the disjoint union $\bigcup_Q W(P, Q)$ where Q runs over the set of standard parabolic subgroups of G onto $\mathcal{P}(M)$.

2.2.12. Truncation parameter. — We shall denote by T a point of \mathfrak{a}_0 such that $\langle \alpha, T \rangle$ is large enough for every $\alpha \in \Delta_0$. We do not want to be precise here. We just need that Arthur's formulas about truncation functions hold for the T 's we consider (see [Art78] §§5,6). The point T plays the role of a truncation parameter.

For any semi-standard parabolic subgroup P , we define a point $T_P \in \mathfrak{a}_P$ such that for any $w \in W$ such that $wP_0 w^{-1} \subset P$, the point T_P is the projection of $w \cdot T$ on \mathfrak{a}_P (this does not depend on the choice of w). The reader should be warned that it is not consistent with the notation of §2.2.7 since there T_P denotes instead the projection of T onto \mathfrak{a}_P (of course, the two conventions coincide when P is standard).

2.2.13. Let P be a standard parabolic subgroup. By a *Siegel domain* for $[G]_P$ we mean a subset of $G(\mathbb{A})$ of the form

$$\mathfrak{s}_P = \omega_0 \{a \in A_0^\infty \mid \langle \alpha, H_0(a) + T \rangle \geq 0, \forall \alpha \in \Delta_0^P\} K$$

where $T \in \mathfrak{a}_0$ and $\omega_0 \subset P_0(\mathbb{A})^1$ is a compact such that $G(\mathbb{A}) = M_P(F)N_P(\mathbb{A})\mathfrak{s}_P$.

2.3 Haar measures

2.3.1. We equip \mathfrak{a}_P with the Haar measure that gives a covolume 1 to the lattice $\text{Hom}(X^*(P), \mathbb{Z})$. The space $i\mathfrak{a}_P^*$ is then equipped with the dual Haar measure so that we have

$$\int_{i\mathfrak{a}_P^*} \int_{\mathfrak{a}_P} \phi(H) \exp(-\langle \lambda, H \rangle) dH d\lambda = \phi(0)$$

for all $\phi \in C_c^\infty(\mathfrak{a}_P)$. Note that this implies that the covolume of $iX^*(P)$ in $i\mathfrak{a}_P^*$ is given by

$$(2.3.1.1) \quad \text{vol}(i\mathfrak{a}_P^*/iX^*(P)) = (2\pi)^{-\dim(\mathfrak{a}_P)}.$$

The group A_P^∞ is equipped with the Haar measure compatible with the isomorphism $A_P^\infty \simeq \mathfrak{a}_P$ induced by the map H_P . The groups $\mathfrak{a}_P^G \simeq \mathfrak{a}_P/\mathfrak{a}_G$ and $i\mathfrak{a}_P^{G,*} \simeq i\mathfrak{a}_P^*/i\mathfrak{a}_G^*$ are provided with the quotient Haar measures. For any basis B of \mathfrak{a}_P^G we denote by $\mathbb{Z}(B)$ the lattice generated by B and by $\text{vol}(\mathfrak{a}_P^G/\mathbb{Z}(B))$ the covolume of this lattice. We have on \mathfrak{a}_0^* the polynomial function:

$$\theta_P(\lambda) = \text{vol}(\mathfrak{a}_P^G/\mathbb{Z}(\Delta_P^\vee))^{-1} \prod_{\alpha \in \Delta_P} \langle \lambda, \alpha^\vee \rangle.$$

2.3.2. Let H be a linear algebraic group over F . In this paper, we will always equip $H(\mathbb{A})$ with its right-invariant Tamagawa measure simply denoted dh . Let us recall how it is defined in order to fix some notation. We choose a right-invariant rational volume form ω_H on H as well as a non-trivial continuous additive character $\psi' : \mathbb{A}/F \rightarrow \mathbb{C}^\times$. For each place $v \in V_F$, the local component ψ'_v of ψ' induces an additive measure on F_v which is the unique Haar measure autodual with respect to ψ'_v . Then using local F -analytic charts, we associate to ω_H a right Haar measure $dh_v = |\omega_H|_{\psi'_v}$ on $H(F_v)$ as in [Wei82, §2.2]. By [Gro97], there exists an Artin-Tate L -function $L_H(s)$ such that, denoting by $L_{H,v}(s)$ the corresponding local L -factor and setting $\Delta_{H,v} = L_{H,v}(0)$, for any model of H over \mathcal{O}_F^S for some finite set $S \subseteq V_F \setminus V_{F,\infty}$, we have

$$(2.3.2.2) \quad \text{vol}(H(\mathcal{O}_v)) = \Delta_{H,v}^{-1}$$

for almost all $v \in V_F$. Setting $\Delta_H^* = L_H^*(0)$, the Tamagawa measure on $H(\mathbb{A})$ is defined as the product

$$(2.3.2.3) \quad dh = (\Delta_H^*)^{-1} \prod_v \Delta_{H,v} dh_v.$$

Although the local measures dh_v depend on choices, the global measure dh doesn't (by the product formula).

2.3.3. For $S \subseteq V_F$ a finite subset, we put $\Delta_H^{S,*} = L_H^{S,*}(0)$ where $L_H^S(s)$ stands for the corresponding partial L -function and we equip $H(F_S)$, $H(\mathbb{A}^S)$ with the right Haar measures $dh_S = \prod_{v \in S} dh_v$ and $dh^S = (\Delta_H^{S,*})^{-1} \prod_{v \notin S} \Delta_{H,v} dh_v$ respectively. Note that we have the decomposition

$$(2.3.3.4) \quad dh = dh_S \times dh^S.$$

In particular, this means that $H(F_v)$ is equipped with the right Haar measure dh_v for every $v \in V_F$.

2.3.4. We have $L_H(s) = L_{H_{\text{red}}}(s)$ where $H_{\text{red}} = H/H_u$ denotes the quotient of H by its unipotent radical H_u . When $H = N$ is unipotent we have $\text{vol}([N]) = 1$. For $H = \text{GL}_n$, the L -function $L_H(s)$ is given by

$$L_H(s) = \zeta_F(s+1) \dots \zeta_F(s+n)$$

where ζ_F stands for the (completed) zeta function of the number field F . In this case, we will take

$$\omega_H = (\det h)^{-1} \bigwedge_{1 \leq i, j \leq n} dh_{i,j}$$

so that (2.3.2.2) is satisfied for every non-Archimedean place v where ψ_v is unramified.

2.3.5. The homogeneous space $[G]$ (resp. $[G]^1 \simeq [G]_0$) is equipped with the quotient of the Tamagawa measure on $G(\mathbb{A})$ by the counting measure on $G(F)$ (resp. by the product of the counting measure on $G(F)$ with the Haar measure we fixed on A_G^∞). For P a standard parabolic

subgroup, we equip similarly $[G]_P$ with the quotient of the Tamagawa measure on $G(\mathbb{A})$ by the product of the counting measure on $M_P(F)$ with the Tamagawa measure on $N_P(\mathbb{A})$. Since the action by left translation of $a \in A_P^\infty$ on $[G]_P$ multiplies the measure by $\delta_P(a)^{-1}$, taking the quotient by the Haar measure on A_P^∞ induces a “semi-invariant” measure on $[G]_{P,0} = A_P^\infty \backslash [G]_P$ that is a positive linear form on the space of continuous functions $\varphi : [G]_P \rightarrow \mathbb{C}$ satisfying $\varphi(ag) = \delta_P(a)\varphi(g)$ for $a \in A_P^\infty$ and compactly supported modulo A_P^∞ .

2.4 Norms and Harish-Chandra Ξ function

2.4.1. Let X be an algebraic variety over F . We define following [Beu16, §A.1] a weak equivalence class of norms $\|\cdot\|_{X(\mathbb{A})} : X(\mathbb{A}) \rightarrow \mathbb{R}_{\geq 1}$. For all the algebraic varieties that we encounter in this paper, we implicitly fix a norm in this equivalence class and we set $\sigma_{X(\mathbb{A})}(x) = 1 + \log\|x\|_{X(\mathbb{A})}$, $x \in X(\mathbb{A})$. If the context is clear, we simply write $\|\cdot\|$ and σ for $\|\cdot\|_{G(\mathbb{A})}$ and $\sigma_{G(\mathbb{A})}$ respectively.

Let $P \subset G$ be a semi-standard parabolic subgroup. We set

$$\|g\|_{[G]_P} = \inf_{\gamma \in M_P(F)N_P(\mathbb{A})} \|\gamma g\| \quad \text{and} \quad \sigma_{[G]_P}(g) = \inf_{\gamma \in M_P(F)N_P(\mathbb{A})} \sigma(\gamma g)$$

for every $g \in [G]_P$. We have

$$(2.4.1.1) \quad \|mk\|_{[G]_P} \approx \|m\|_{[M_P]}, \quad m \in [M_P], k \in K.$$

Let \mathfrak{s}_P be a Siegel domain as in Section 2.2.13. By Proposition [Beu16, A.1.1 (viii)] and (2.4.1.1), we have

$$(2.4.1.2) \quad \|g\|_{[G]_P} \approx \|g\|_{G(\mathbb{A})}, \quad g \in \mathfrak{s}_P.$$

2.4.2. Let $\omega_G \subseteq G(\mathbb{A})$ be a compact subset with nonempty interior and set

$$\Xi^{[G]_P}(g) = \text{vol}_{[G]_P}(g\omega_G)^{-1/2}, \quad g \in [G]_P.$$

For another choice of compact subset $\omega'_G \subseteq G(\mathbb{A})$ with nonempty interior, the resulting functions are equivalent which is why we dropped the subset ω_G from the notation. We have

$$(2.4.2.3) \quad \Xi^{[G]_P}(g) \sim \exp(\langle \rho_0, H_0(g) \rangle), \quad g \in \mathfrak{s}_P.$$

Indeed, we readily check that $\Xi^{[G]_P}(mk) \sim \exp(\langle \rho_P, H_P(m) \rangle) \Xi^{[M_P]}(m)$ for every $m \in [M_P]$ and $k \in K$ so that we are reduced to the case $P = G$. By invariance we have $\text{vol}_{[G]}(g\omega_G) = \text{vol}_{[G]}(g\omega_G g^{-1})$ and it is easy to see that, if ω_G is chosen sufficiently small, there exists a compact subset $\omega'_G \subseteq G(\mathbb{A})$ such that $g\omega_G g^{-1} \subseteq N_0(F)\omega'_G$ for every $g \in \mathfrak{s}_G$. As $G(F)$ is discrete inside $G(\mathbb{A})$, the set

$$\{\gamma \in P_0(F) \backslash G(F) \mid \gamma \omega'_G \cap P_0(F)\omega'_G \neq \emptyset\}$$

is finite. Therefore, we have

$$\text{vol}_{[G]}(g\omega_G g^{-1}) \sim \text{vol}_{[G]_{P_0}}(g\omega_G g^{-1}) \sim \exp(-\langle 2\rho_0, H_0(p) \rangle)$$

for $g \in \mathfrak{s}_G$ and (2.4.2.3) follows.

By [Lap13, §2, (9)], we also have³

$$(2.4.2.4) \quad \text{There exists } d > 0 \text{ such that } \int_{[G]_P} \Xi^{[G]_P}(g)^2 \sigma_{[G]_P}(g)^{-d} dg < \infty;$$

From (2.4.1.2) and (2.4.2.3), we deduce the existence of $N \geq 1$ such that

$$(2.4.2.5) \quad \Xi^{[G]_P}(g) \ll \|g\|_{[G]_P}^N, \quad g \in [G]_P.$$

³Note that the definition of the Ξ function in *loc. cit.* coincides, up to equivalence, with ours by (2.4.2.3).

2.5 Spaces of functions

2.5.1. We say of a function $f : G(\mathbb{A}) \rightarrow \mathbb{C}$ that it is *smooth* if it is right invariant by a compact-open subgroup J of $G(\mathbb{A}_f)$ and for every $g_f \in G(\mathbb{A}_f)$ the function $g_\infty \in G(F_\infty) \mapsto f(g_f g_\infty)$ is C^∞ .

2.5.2. Let C be a compact subset of $G(\mathbb{A}_f)$ and let $J \subset K^\infty$ be a compact open subgroup. The left and right actions of the envelopping algebra $\mathcal{U}(\mathfrak{g}_\infty)$ are denoted by L and R respectively. Let $\mathcal{S}(G(\mathbb{A}), C, J)$ be the space of smooth functions $f : G(\mathbb{A}) \rightarrow \mathbb{C}$ which are biinvariant by J , supported in the subset $G(F_\infty) \times C$ and such that the semi-norms

$$\|f\|_{r,X,Y} = \sup_{g \in G(\mathbb{A})} \|g\|_{G(\mathbb{A})}^r |(R(X)L(Y)f)(g)|$$

are finite for every integer $r \geq 1$ and $X, Y \in \mathcal{U}(\mathfrak{g}_\mathbb{C})$. This family of semi-norms define a topology on $\mathcal{S}(G(\mathbb{A}), C, K_0)$ making it into a Fréchet space. The global *Schwartz space* $\mathcal{S}(G(\mathbb{A}))$ is the topological direct limit over all pairs (C, J) of the spaces $\mathcal{S}(G(\mathbb{A}), C, J)$. The Schwartz space is an algebra for the convolution product denoted by $*$. It contains the dense subspace $C_c^\infty(G(\mathbb{A}))$ of smooth and compactly supported functions. For an integer $r \geq 0$, we will also consider the space $C_c^r(G(\mathbb{A}))$ generated by products $f_\infty f^\infty$ where f_∞ is a compactly supported function on $G(F \otimes_\mathbb{Q} \mathbb{R})$ which admits derivatives up to the order r and f^∞ is a smooth compactly supported function on $G(\mathbb{A}_f)$.

For every integer $n \geq 1$, we define similarly the global Schwartz space $\mathcal{S}(\mathbb{A}^n)$ (the definition is the same as above up to replacing G by the additive group \mathbf{G}_a^n).

2.5.3. Let \mathcal{H} be a Hilbert space carrying a continuous representation of $G(\mathbb{A})$ (not necessarily unitary). A vector $v \in \mathcal{H}$ is *smooth* if it is invariant by a compact-open subgroup of $G(\mathbb{A}_f)$ and the function $g_\infty \in G(F_\infty) \mapsto g_\infty v \in \mathcal{H}$ is C^∞ . We denote by \mathcal{H}^∞ the subspace of smooth vectors. It is a $G(\mathbb{A})$ -invariant subspace carrying its own locally convex topology making it into a strict LF space: for every compact-open subgroup $J \subseteq G(\mathbb{A}_f)$, the subspace $(\mathcal{H}^\infty)^J$ of vectors fixed by J is equipped with the Fréchet topology associated to the semi-norms $v \mapsto |Xv|_\mathcal{H}$ where $|\cdot|_\mathcal{H}$ is the norm on \mathcal{H} , $X \in \mathcal{U}(\mathfrak{g}_\infty)$ and $Xv := R(X)(g_\infty \mapsto g_\infty v)|_{g_\infty=1}$.

2.5.4. Let P be a semi-standard parabolic subgroup of G . We denote by $L^2([G]_P)$ the space of L^2 -measurable functions on $[G]_P$. It is a Hilbert space when equipped with the scalar product

$$\langle \varphi_1, \varphi_2 \rangle_{[G]_P} = \int_{[G]_P} \varphi_1(g) \overline{\varphi_2(g)} dg$$

associated to the Tamagawa invariant measure on $[G]_P$. We denote similarly by $L^2([G]_{P,0})$ the Hilbert space of measurable functions φ on $[G]_P$ satisfying $\varphi(ag) = \delta_P(a)^{1/2} \varphi(g)$ for almost all $a \in A_P^\infty$ and such that $\int_{[G]_{P,0}} |\varphi(g)|^2 dg$ is convergent.

We can define more generally weighted L^2 spaces as follows. Let w be a weight on $[G]_P$ [Ber88, §3.1] that is a positive measurable function on $[G]_P$ such that for every compact subset $\omega_G \subset G(\mathbb{A})$ we have $w(xg) \ll w(x)$ for $x \in [G]_P$ and $g \in \omega_G$. The Hilbert space $L_w^2([G]_P) = L^2([G]_P, w(g)dg)$ is then equipped with a continuous (non-unitary) representation of $G(\mathbb{A})$ by right-translation. In particular, following the previous paragraph, we denote by $L_w^2([G]_P)^\infty$ its subspace of smooth vectors: it consists of smooth functions $\varphi : [G]_P \rightarrow \mathbb{C}$ such that $R(X)\varphi \in L_w^2([G]_P)$ for every $X \in \mathcal{U}(\mathfrak{g}_\infty)$. By the Sobolev inequality, see [Ber88, §3.4, Key Lemma], for every $\varphi \in L_w^2([G]_P)^\infty$ we have

$$(2.5.4.1) \quad |\varphi(g)| \ll \Xi^{[G]_P}(g) w(g)^{-1/2}, \quad g \in [G]_P.$$

In this paper, we will only use three kind of weights. First, for $N \in \mathbb{R}$, $\|\cdot\|_{[G]_P}^N$ is a weight and we set $L_N^2([G]_P) = L_{\|\cdot\|_{[G]_P}^N}^2([G]_P)$. Secondly, for $d > 0$, $\sigma_{[G]_P}^d$ is a weight and we set $L_{\sigma,d}^2([G]_P) = L_{\sigma_{[G]_P}^d}^2([G]_P)$. Thirdly, if λ is a weight on \mathfrak{a}_P , i.e. a real measurable function such that for every

compact $\omega \subset \mathfrak{a}_P$ we have $|\lambda(X + Y) - \lambda(X)| \ll 1$ for $X \in \mathfrak{a}_P$ and $Y \in \omega$ (e.g. $\lambda \in \mathfrak{a}_P^*$), then $\exp(\lambda \circ H_P)$ is a weight on $[G]_P$ and we set $L_\lambda^2([G]_P) = L_{\exp(\lambda \circ H_P)}^2([G]_P)$.

2.5.5. The *Schwartz space* $\mathcal{S}([G]_P)$ of $[G]_P$ is defined as the space of smooth functions $\varphi : [G]_P \rightarrow \mathbb{C}$ such that for every $N > 0$ and $X \in \mathcal{U}(\mathfrak{g}_\infty)$ we have

$$|(R(X)\varphi)(g)| \ll \|g\|_{[G]_P}^{-N}, \quad g \in [G]_P.$$

Then, $\mathcal{S}([G]_P)$ is naturally equipped with a locally convex topology making it into a strict LF space (it is the inductive limit of the Fréchet spaces $\mathcal{S}([G]_P)^J$ for J a compact-open subgroup of $G(\mathbb{A}_f)$). From (2.5.4.1), we have the alternative description

$$(2.5.5.2) \quad \mathcal{S}([G]_P) = \bigcap_{N>0} L_N^2([G]_P)^\infty.$$

2.5.6. The *Harish-Chandra Schwartz space* $\mathcal{C}([G]_P)$ of $[G]_P$ is defined as the space of smooth functions $\varphi : [G]_P \rightarrow \mathbb{C}$ such that for every $d > 0$ and $X \in \mathcal{U}(\mathfrak{g}_\infty)$ we have

$$|(R(X)\varphi)(g)| \ll \Xi^{[G]_P}(g) \sigma_{[G]_P}(g)^{-d}, \quad g \in [G]_P.$$

Once again, $\mathcal{C}([G]_P)$ is naturally equipped with a locally convex topology making it into a strict LF space. Alternatively, we have

$$(2.5.6.3) \quad \mathcal{C}([G]_P) = \bigcap_{d>0} L_{\sigma,d}^2([G]_P)^\infty.$$

2.5.7. The *space of functions of uniform moderate growth* $\mathcal{T}([G]_P)$ of $[G]_P$ is defined as the space of smooth functions $\varphi : [G]_P \rightarrow \mathbb{C}$ for which there exists $N > 0$ such that for every $X \in \mathcal{U}(\mathfrak{g}_\infty)$ we have

$$|(R(X)\varphi)(g)| \ll \|g\|_{[G]_P}^N, \quad g \in [G]_P.$$

For $N > 0$, we denote by $\mathcal{T}_N([G]_P)$ the subspace of functions $\varphi \in \mathcal{T}([G]_P)$ satisfying the above inequality for every $X \in \mathcal{U}(\mathfrak{g}_\infty)$. Then, $\mathcal{T}_N([G]_P)$ is naturally equipped with a locally convex topology making it into a strict LF space and $\mathcal{T}([G]_P) = \bigcup_{N>0} \mathcal{T}_N([G]_P)$ is a (non-strict) LF space. We also have the alternative description

$$(2.5.7.4) \quad \mathcal{T}([G]_P) = \bigcup_{N>0} L_{-N}^2([G]_P)^\infty.$$

2.5.8. The spaces $\mathcal{S}([G]_P)$, $\mathcal{C}([G]_P)$ and $\mathcal{T}([G]_P)$ are all topological representations of $G(\mathbb{A})$ for the action by right translation R . For $J \subseteq G(\mathbb{A}_f)$ a compact-open subgroup and $N > 0$, $\mathcal{S}([G]_P)^J$, $\mathcal{C}([G]_P)^J$ and $\mathcal{T}_N([G]_P)^J$ are even SF representations of $G(F_\infty)$ in the sense of [BK14]. It follows that the action of $G(\mathbb{A})$ on $\mathcal{S}([G]_P)$, $\mathcal{C}([G]_P)$ and $\mathcal{T}([G]_P)$ integrates to an action of the algebra $(\mathcal{S}(G(\mathbb{A})), *)$ (by right convolution). Moreover, by [Ber88, end of Section 3.5] (see also [Cas89, Corollary 2.6]) we have

(2.5.8.5) The Fréchet spaces $\mathcal{S}([G]_P)^J$ and $\mathcal{C}([G]_P)^J$ are nuclear.

Let $\mathcal{S}([G]_P)'$ be the topological dual of $\mathcal{S}([G]_P)$. By duality, it is also equipped with an action by convolution of $\mathcal{S}(G(\mathbb{A}))$. By the alternative descriptions (2.5.5.2) and (2.5.7.4), we have

(2.5.8.6) For every distribution $D \in \mathcal{S}([G]_P)'$ and $f \in \mathcal{S}(G(\mathbb{A}))$, the distribution $R(f)D$ is representable by a function in $\mathcal{T}([G]_P)$.

2.5.9. Assume that $G = G_1 \times G_2$ where G_1 and G_2 are two connected reductive groups over F . Let $J_1 \subset G_1(\mathbb{A}_f)$, $J_2 \subseteq G_2(\mathbb{A}_f)$ be two compact open subgroups and set $J = J_1 \times J_2$. By (2.5.8.5), (A.0.6.6) and a reasoning similar to (the proof of) [BP15, Proposition 4.4.1 (v)] we obtain:

(2.5.9.7) There are topological isomorphisms

$$\mathcal{C}([G_1])^{J_1} \widehat{\otimes} \mathcal{C}([G_2])^{J_2} \simeq \mathcal{C}([G])^J, \quad \mathcal{S}([G_1])^{J_1} \widehat{\otimes} \mathcal{S}([G_2])^{J_2} \simeq \mathcal{S}([G])^J$$

sending a pure tensor $\varphi_1 \otimes \varphi_2$ to the function $(g_1, g_2) \mapsto \varphi_1(g_1)\varphi_2(g_2)$.

By the above, given two continuous linear forms L_1, L_2 on $\mathcal{C}([G_1]), \mathcal{C}([G_2])$ respectively, the linear form $L_1 \otimes L_2$ on $\mathcal{C}([G_1]) \otimes \mathcal{C}([G_2])$ extends by continuity to a linear form on $\mathcal{C}([G])$ that we shall denote by $L_1 \widehat{\otimes} L_2$.

2.5.10. Constant terms and pseudo-Eisenstein series. — Let $Q \subset P$ be another standard parabolic subgroup. We have two continuous $G(\mathbb{A})$ -equivariant linear maps

$$\mathcal{T}([G]_P) \rightarrow \mathcal{T}([G]_Q), \quad \varphi \mapsto \varphi_Q \quad \text{and} \quad \mathcal{S}([G]_Q) \rightarrow \mathcal{S}([G]_P), \quad \varphi \mapsto E_Q^P(\varphi)$$

defined by

$$\varphi_Q(g) = \int_{[N_Q]} \varphi(ug) du \quad \text{and} \quad E_Q^P(\varphi, g) = \sum_{\gamma \in Q(F) \backslash P(F)} \varphi(\gamma g)$$

respectively.

Lemma 2.5.10.1. — *There is a constant $c > 0$ such that for every $N \geq 0$,*

$$f \mapsto \sup_{g \in [G]_P} \|g\|_{[G]_P}^N \delta_P(g)^{cN} |f_P(g)|$$

is a continuous semi-norm on $\mathcal{S}([G])$.

Proof. — As $[N_P]$ is compact, it suffices to show the existence of $N \geq 1$ and $c > 0$ such that

$$\delta_P(g)^c \|g\|_{[G]_P} \ll \|g\|_{[G]}^N$$

for every $g \in G(\mathbb{A})$. By (2.4.1.2), we are easily reduced to show the existence of $N \geq 1$ and $c > 0$ such that

$$\delta_P(a)^c \|a\| \ll \|ua\|_{[G]}^N$$

for every $(u, a) \in N_P(\mathbb{A}) \times A_0^\infty$. It is then equivalent to establish the following:

(2.5.10.8) For every algebraic character χ of A_0 , there exist $N \geq 1$ and $c > 0$ such that

$$\delta_P(a)^c |\chi(a)| \ll \|ua\|_{[G]}^N, \quad \text{for } (a, u) \in A_0^\infty \times [N_P].$$

Let χ be such a character and let δ_P^{alg} be the algebraic character of M_P given by the determinant of the adjoint action on the Lie algebra of N_P . We have $|\delta_P^{alg}(m)| = \delta_P(m)$ for every $m \in M_P(\mathbb{A})$ and we can find an integer $c \geq 1$, a rational representation (V, ρ) of G and a nonzero vector $v_0 \in V$ such that $\rho(ua)v_0 = \chi(a)\delta_P^{alg}(a)^c v_0$ for every $a \in A_0$ and $u \in N_P$. Fixing a basis v_1, \dots, v_d of V , for every $v = \lambda_1 v_1 + \dots + \lambda_d v_d \in V_{\mathbb{A}} := V \otimes_F \mathbb{A}$, we set

$$|v|_V = \prod_v \max(|\lambda_1|_v, \dots, |\lambda_d|_v).$$

Then, there exists $N \geq 1$ such that $|\rho(g)v_0|_V \ll \|g\|_{G(\mathbb{A})}^N$ for all $g \in G(\mathbb{A})$ and moreover we have $|v|_V \geq 1$ for every $v \in V \setminus \{0\}$. Therefore, we obtain

$$\|\gamma ua\|_{G(\mathbb{A})}^N \gg |\rho(\gamma ua)v_0|_V = |\chi(a)|\delta_P(a)^c |\rho(\gamma)v_0|_V \geq |\chi(a)|\delta_P(a)^c$$

for $\gamma \in G(F)$, $a \in A_0^\infty$ and $u \in N_P(\mathbb{A})$. The estimates (2.5.10.8) follows by replacing the left hand side in the inequality above by its the infimum over $\gamma \in G(F)$. \square

2.5.11. For $Q \subset P$ two standard parabolic subgroup, we let $(\mathfrak{a}_Q^*)^{P++}$ be the set of $\lambda \in \mathfrak{a}_Q^*$ such that $\langle \alpha^\vee, \lambda \rangle > 0$ for every $\alpha \in \Delta_0^P \setminus \Delta_0^Q$.

Proposition 2.5.11.1. — Let $\varphi \in \mathcal{T}([G]_P)$. Then, $\varphi \in \mathcal{S}([G]_P)$ if and only if for every standard parabolic subgroup $Q \subset P$ and every $\lambda \in (\mathfrak{a}_Q^*)^{P++}$ we have $\varphi_Q \in L_{2\rho_Q+\lambda}^2([G]_Q)^\infty$. Moreover, for each compact-open subgroup $J \subset G(\mathbb{A}_f)$, the semi-norms

$$\varphi \mapsto \|R(X)\varphi_Q\|_{L_{2\rho_Q+\lambda}^2}$$

for $Q \subset P$ a standard parabolic subgroup, $\lambda \in (\mathfrak{a}_Q^*)^{P++}$ and $X \in \mathcal{U}(\mathfrak{g}_\infty)$ generate the topology on $\mathcal{S}([G]_P)^J$.

Proof. — Let $\varphi \in \mathcal{S}([G]_P)$, $Q \subset P$ be a standard parabolic subgroup, $\lambda \in (\mathfrak{a}_Q^*)^{P++}$ and $X \in \mathcal{U}(\mathfrak{g}_\infty)$. By Cauchy-Schwarz we have

$$\int_{[G]_Q} |R(X)\varphi_Q(x)|^2 e^{\langle 2\rho_Q+\lambda, H_Q(x) \rangle} dx \leq \int_{[G]_P} |R(X)\varphi(x)|^2 \sum_{\gamma \in Q(F) \backslash P(F)} e^{\langle 2\rho_Q+\lambda, H_Q(\gamma x) \rangle} dx.$$

and by the convergence of Eisenstein series the inner sum above converges to a function which is essentially bounded by $\|x\|^N$ for some N . Therefore, $\varphi_Q \in L_{2\rho_Q+\lambda}^2([G]_Q)^\infty$ and this shows the direct implication.

We now prove the converse: let $\varphi \in \mathcal{T}([G]_P)$ be such that for every standard parabolic subgroup $Q \subset P$ and every $\lambda \in (\mathfrak{a}_Q^*)^{P++}$ we have $\varphi_Q \in L_{2\rho_Q+\lambda}^2([G]_Q)^\infty$. Applying this assumption to $Q = P$ and λ varying over fixed basis of \mathfrak{a}_P^* and its opposite, from (2.5.4.1) and (2.4.2.3) we get for every $N > 0$ the estimates

$$(2.5.11.9) \quad |\varphi(g)| \ll \exp(\langle \rho_0, H_0(g) \rangle - N\|H_P(g)\|), \quad g \in \mathfrak{s}_P.$$

Similarly, applying the hypothesis to the maximal parabolic subgroup $Q_\alpha \subset P$ associated to a simple root $\alpha \in \Delta_0^P$ and λ a multiple of $\rho_{Q_\alpha}^P$, from (2.5.4.1) and (2.4.2.3) for every $N > 0$ we obtain

$$(2.5.11.10) \quad |\varphi_{Q_\alpha}(g)| \ll \exp(\langle \rho_0, H_0(g) \rangle - N\langle \alpha, H_0(g) \rangle), \quad g \in \mathfrak{s}_Q.$$

Moreover, by the approximation property of the constant term [MW94, Lemma I.2.10], there exists $M > 0$ such that for every $N > 0$ we have

$$(2.5.11.11) \quad |\varphi(g) - \varphi_{Q_\alpha}(g)| \ll \exp(-N\langle \alpha, H_0(g) \rangle + M\|H_0(g)\|), \quad g \in \mathfrak{s}_P.$$

Finally, there exist $\epsilon > 0$ and $C > 0$ such that $\max\{\langle \alpha, H_0(g) \rangle \mid \alpha \in \Delta_0^P\} \cup \{\|H_P(g)\|\} \geq \epsilon\|H_0(g)\| - C$ for all $g \in \mathfrak{s}_P$. Combining this with (2.5.11.9), (2.5.11.10) and (2.5.11.11), gives the estimates

$$|\varphi(g)| \ll \|g\|^{-N}, \quad g \in \mathfrak{s}_P,$$

for every $N > 0$. Applying the same reasoning to derivatives of φ , this shows that $\varphi \in \mathcal{S}([G]_P)$.

For the last part of the statement, it suffices to notice that the linear map

$$\begin{aligned} \mathcal{S}(G(\mathbb{A})) &\rightarrow \prod_{Q \subset P, \lambda \in (\mathfrak{a}_Q^*)^{P++}} L_{2\rho_Q+\lambda}^2([G]_Q)^\infty \\ \varphi &\mapsto (\varphi_Q)_Q \end{aligned}$$

is injective with a closed image (by the previous characterization) hence is a topological embedding by the open mapping theorem. \square

2.6 Estimates on Fourier coefficients

2.6.1. Let P be a standard parabolic subgroup of G , $\psi : \mathbb{A}/F \rightarrow \mathbb{C}^\times$ be a non-trivial additive character and $\ell : N_P \rightarrow \mathbb{G}_a$ be an algebraic character. We set $\psi_\ell := \psi \circ \ell_{\mathbb{A}} : [N_P] \rightarrow \mathbb{C}^\times$. For $f \in C^\infty([G])$, we set

$$f_{N_P, \psi_\ell}(g) = \int_{[N_P]} f(ug) \psi_\ell(u)^{-1} du, \quad g \in G(\mathbb{A}).$$

Let $N_{P, \text{der}}$ denote the derived subgroup of N_P and set $N_{P, \text{ab}} = N_P / N_{P, \text{der}}$ (a vector space over F). Then, ℓ can be seen as an element in the dual space $N_{P, \text{ab}}^*$ that we also consider as an algebraic variety over F . The adjoint action of M_P on N_P induces one on $N_{P, \text{ab}}^*$ that we denote by Ad^* .

Lemma 2.6.1.1. —

1. There exists $c > 0$ such that for every $N_1, N_2 \geq 0$,

$$f \mapsto \sup_{m \in M_P(\mathbb{A})} \|\text{Ad}^*(m^{-1})\ell\|_{N_{P, \text{ab}}^*(\mathbb{A})}^{N_1} \|m\|_{[M_P]}^{N_2} \delta_P(m)^{cN_2} |f_{N_P, \psi_\ell}(m)|$$

is a continuous semi-norm on $\mathcal{S}([G])$.

2. For every $N_1, N_2 \geq 0$,

$$f \mapsto \sup_{m \in M_P(\mathbb{A})} \|\text{Ad}^*(m^{-1})\ell\|_{N_{P, \text{ab}}^*(\mathbb{A})}^{N_1} \|m\|_{[M_P]}^{-N_2} |f_{N_P, \psi_\ell}(m)|$$

is a continuous semi-norm on $\mathcal{T}_N([G])$.

Proof. — Bounding brutally under the integral sign, we have

$$|f_{N_P, \psi_\ell}(g)| \leq |f|_P(g)$$

for $f \in C^\infty([G])$ and $g \in G(\mathbb{A})$. Let $N \geq 0$ and $J \subseteq G(\mathbb{A}_f)$ be a compact-open subgroup. By Lemma 2.5.10.1 and (2.4.1.1), it suffices to show the existence of elements $X_1, \dots, X_M \in \mathcal{U}(\mathfrak{g}_\infty)$ such that

$$(2.6.1.1) \quad |f_{N_P, \psi_\ell}(m)| \leq \|\text{Ad}^*(m^{-1})\ell\|_{N_{P, \text{ab}}^*(\mathbb{A})}^{-N} \sum_{i=1}^M |(R(X_i)f)_{N_P, \psi_\ell}(m)|$$

for every $f \in C^\infty([G])^J$ and $m \in M_P(\mathbb{A})$. Let $u \in N_P(\mathbb{A})$. By definition of $\|\cdot\|_{N_{P, \text{ab}}^*(\mathbb{A})}$, we are readily reduced to show the existence of $X_1, \dots, X_M \in \mathcal{U}(\mathfrak{g}_\infty)$ such that

$$(2.6.1.2) \quad |f_{N_P, \psi_\ell}(m)| \leq \|\ell(\text{Ad}(m)u)\|_{\mathbb{A}}^{-1} \sum_{i=1}^M |(R(X_i)f)_{N_P, \psi_\ell}(m)|$$

for every $f \in C^\infty([G])^J$ and $m \in M_P(\mathbb{A})$. This last claim is a consequence of the two following facts whose proofs are elementary and left to the reader.

(2.6.1.3) For every non-Archimedean place v , there exists a constant $C_v \geq 1$ with $C_v = 1$ for almost all v such that $|\ell(\text{Ad}(m_v)u_v)|_v > C_v$ implies $f_{N_P, \psi_\ell}(m) = 0$ for every $f \in C^\infty([G])^J$ and $m \in M_P(\mathbb{A})$.

(2.6.1.4) Let v be an Archimedean place and let $X \in \mathfrak{g}_v$ be such that $u_v = e^X$. Then, we have $(R(X)f)_{N_P, \psi_\ell}(m) = d\psi_v(\ell(\text{Ad}(m)u)_v) f_{N_P, \psi_\ell}(m)$ for all $f \in C^\infty([G])$ and $m \in M_P(\mathbb{A})$ where $d\psi_v : F_v \rightarrow i\mathbb{R}$ is the differential of ψ_v at the origin.

□

2.6.2. Let $n \geq 1$ be a positive integer. We let GL_n acts on F^n by right multiplication and we denote by $e_n = (0, \dots, 0, 1)$ the last element of the standard basis of F^n . We also denote by \mathcal{P}_n the *mirabolic* subgroup of GL_n , that is the stabilizer of e_n in GL_n . We identify A_{GL_n} with \mathbf{G}_m , and thus $A_{\mathrm{GL}_n}^\infty$ with $\mathbb{R}_{>0}$, in the usual way. The next lemma will be used in conjunction with Lemma 2.6.1.1 to show the convergence of various Zeta integrals.

Lemma 2.6.2.1. — *Let $C > 1$. Then, for $N_1 \gg_C 1$ and $N_2 \gg_C 1$ the integral*

$$\int_{\mathcal{P}_n(F) \backslash \mathrm{GL}_n(\mathbb{A}) \times \mathbb{R}_{>0}} \|ag\|_{[\mathrm{GL}_n]}^{-N_1} \|e_n g\|_{\mathbb{A}^n}^{-N_2} |\det g|^s \, dadg$$

converges for $s \in \mathcal{H}_{1,C}$ uniformly on every (closed) vertical strip.

Proof. — The integral of the lemma can be rewritten as

$$(2.6.2.5) \quad \int_{[\mathrm{GL}_n]} \|g\|_{[\mathrm{GL}_n]}^{-N_1} \int_{\mathbb{R}_{>0}} \sum_{\xi \in F^n \setminus \{0\}} \|\xi ag\|_{\mathbb{A}^n}^{-N_2} |\det ag|^s \, dadg.$$

There exists $N_3 > 0$ such that $\|v\|_{\mathbb{A}^n} \ll \|vg\|_{\mathbb{A}^n}^{N_3} \|g\|_{\mathrm{GL}_n(\mathbb{A})}^{N_3}$ for $(v, g) \in \mathbb{A}^n \times \mathrm{GL}_n(\mathbb{A})$. Therefore, the inner integral above is essentially bounded by

$$|\det g|^{\Re(s)} \|g\|_{\mathrm{GL}_n(\mathbb{A})}^{N_2} \int_{\mathbb{R}_{>0}} \sum_{\xi \in F^n \setminus \{0\}} \|a\xi\|_{\mathbb{A}^n}^{-N_2/N_3} |a|^{ns} \, da$$

hence, for $1 < \Re(s) < C$, by

$$\|g\|_{\mathrm{GL}_n(\mathbb{A})}^{N_2+N_4} \int_{\mathbb{R}_{>0}} \sum_{\xi \in F^n \setminus \{0\}} \|a\xi\|_{\mathbb{A}^n}^{-N_2/N_3} |a|^{ns} \, da$$

for some $N_4 > 0$. However, since the inner integral in (2.6.2.5) is left invariant by $\mathrm{GL}_n(F)$, as a function of g , we may replace $\|g\|_{\mathrm{GL}_n(\mathbb{A})}$ in the estimate above by $\|g\|_{[\mathrm{GL}_n]}$. As for $N \gg 1$ we have

$\int_{[\mathrm{GL}_n]} \|g\|_{[\mathrm{GL}_n]}^{-N} dg < \infty$ [Beu16, Proposition A.1.1 (vi)], it only remains to show that for $N \gg 1$ the integral

$$\int_{\mathbb{R}_{>0}} \sum_{\xi \in F^n \setminus \{0\}} \|a\xi\|_{\mathbb{A}^n}^{-N} |a|^{ns} \, da$$

converges for $1 < \Re(s) < C$ uniformly in vertical strips. This is an easy consequence of the following claim:

(2.6.2.6) For every $k \geq n$, if N is sufficiently large we have

$$\sum_{\xi \in F^n \setminus \{0\}} \|a\xi\|_{\mathbb{A}^n}^{-N} \ll |a|^{-k}, \quad a \in \mathbb{R}_{>0}.$$

There exists $M_0 \geq 1$ such that

$$|a| = \max_{1 \leq i \leq n} (|a\xi_i|) \ll \|a\xi\|_{\mathbb{A}^n}^{M_0}$$

for $(a, \xi) \in \mathbb{R}_{>0} \times (F^n \setminus \{0\})$. Therefore, we just need to prove (2.6.2.6) when $k = n$. Let $C \subset \mathbb{A}^n$ be a compact subset which surjects onto \mathbb{A}^n/F^n . There exists $M_1 \geq 0$ such that

$$\|a\xi + av\|_{\mathbb{A}^n} \ll \|a\xi\|_{\mathbb{A}^n}^{M_1} \max(1, |a|)^{M_1} \ll \|a\xi\|_{\mathbb{A}^n}^{M_1+M_0M_1}$$

for $(a, \xi, v) \in \mathbb{R}_{>0} \times (F^n \setminus \{0\}) \times C$. Hence, for every $N' > 0$ if N is sufficiently large, we have

$$\begin{aligned} \sum_{\xi \in F^n \setminus \{0\}} \|a\xi\|_{\mathbb{A}^n}^{-N} &\ll \int_C \sum_{\xi \in F^n \setminus \{0\}} \|a\xi + av\|_{\mathbb{A}^n}^{-N'} dv \\ &\ll \int_{\mathbb{A}^n/F^n} \sum_{\xi \in F^n} \|a\xi + av\|_{\mathbb{A}^n}^{-N'} dv = |a|^{-n} \int_{\mathbb{A}^n} \|v\|_{\mathbb{A}^n}^{-N'} dv \end{aligned}$$

for $a \in \mathbb{R}_{>0}$. The last integral above is absolutely convergent when $N' \gg 1$ [Beu16, Proposition A.1.1 (vi)] and the claim (2.6.2.6) follows. \square

2.7 Automorphic forms and representations

2.7.1. Let P be a standard parabolic subgroup of G . The space $\mathcal{A}_P(G)$ of automorphic forms on $[G]_P$ is defined as the subspace of $\mathcal{Z}(\mathfrak{g}_\infty)$ -finite functions in $\mathcal{T}([G]_P)$. The subspace $\mathcal{A}_{P,\text{cusp}}(G)$ of cuspidal automorphic forms consists of the $\varphi \in \mathcal{A}_P(G)$ such that for every proper standard parabolic subgroup $Q \subsetneq P$ we have $\varphi_Q = 0$.

For $\mathcal{J} \subset \mathcal{Z}(\mathfrak{g}_\infty)$ an ideal of finite codimension, we denote by $\mathcal{A}_{P,\mathcal{J}}(G)$ the subspace of automorphic forms $\varphi \in \mathcal{A}_P(G)$ such that $R(z)\varphi = 0$ for every $z \in \mathcal{J}$ and we set $\mathcal{A}_{P,\text{cusp},\mathcal{J}}(G) = \mathcal{A}_{P,\mathcal{J}}(G) \cap \mathcal{A}_{P,\text{cusp}}(G)$. Then, there exists $N \geq 1$ such that $\mathcal{A}_{P,\mathcal{J}}(G)$ is a closed subspace of $\mathcal{T}_N([G]_P)$ and we equip $\mathcal{A}_{P,\mathcal{J}}(G)$ with the induced topology from $\mathcal{T}_N([G]_P)$ (this topology does not depend on the choice of N by the open mapping theorem). Similarly, $\mathcal{A}_{P,\text{cusp},\mathcal{J}}(G)$ is a closed subspace of $\mathcal{S}([G]_P)$ and we equip $\mathcal{A}_{P,\text{cusp},\mathcal{J}}(G)$ with the induced topology (which also coincides with the topology induced from $\mathcal{A}_{P,\mathcal{J}}(G)$). We have $\mathcal{A}_P(G) = \bigcup_{\mathcal{J}} \mathcal{A}_{P,\mathcal{J}}(G)$ and $\mathcal{A}_{P,\text{cusp}}(G) = \bigcup_{\mathcal{J}} \mathcal{A}_{P,\text{cusp},\mathcal{J}}(G)$ where \mathcal{J} runs over all ideals of finite codimensions in $\mathcal{Z}(\mathfrak{g}_\infty)$ and we equip these spaces with the convex inductive limit topology (these are strict LF spaces). We shall also consider the closed subspace $\mathcal{A}_{P,\text{disc}}(G)$ of $\mathcal{A}_P(G)$ of automorphic forms on $[G]_{P,0}$ that are square-integrable. Its topology is also induced from the strict LF space $L^2([G]_P)^\infty$.

For $P = G$, we simply set $\mathcal{A}(G) = \mathcal{A}_G(G)$, $\mathcal{A}_{\text{disc}}(G) = \mathcal{A}_{G,\text{disc}}(G)$ and $\mathcal{A}_{\text{cusp}}(G) = \mathcal{A}_{G,\text{cusp}}(G)$.

2.7.2. By a *cuspidal automorphic representation* (resp. discrete) σ of $M_P(\mathbb{A})$ we mean a topologically irreducible subrepresentation of $\mathcal{A}_{\text{cusp}}(M_P)$ (resp. $\mathcal{A}_{\text{disc}}(M_P)$). Let σ be a cuspidal or discrete automorphic representation of $M_P(\mathbb{A})$. For every $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^*$, the twist $\sigma_\lambda = \sigma \otimes \lambda$ is defined as the space of automorphic forms

$$m \in [M_P] \mapsto \exp(\langle \lambda, H_P(m) \rangle) \varphi(m)$$

for $\varphi \in \sigma$. If σ is cuspidal, σ_λ is again a cuspidal automorphic representation. We denote by $\mathcal{A}_\sigma(M_P)$ the σ -isotypic component of $\mathcal{A}_{\text{cusp}}(M_P)$ (resp. $\mathcal{A}_{\text{disc}}(M_P)$) i.e. the sum of all cuspidal (resp. discrete) automorphic representations of $M_P(\mathbb{A})$ that are isomorphic to σ . We let $\Pi = I_{P(\mathbb{A})}^{G(\mathbb{A})}(\sigma)$ (resp. $\mathcal{A}_{P,\sigma}(G) = I_{P(\mathbb{A})}^{G(\mathbb{A})}(\mathcal{A}_\sigma(M_P))$) be the *normalized smooth induction* of σ (resp. $\mathcal{A}_\sigma(M_P)$) that we identify with the space of forms $\varphi \in \mathcal{A}_P(G)$ such that

$$m \in [M_P] \mapsto \exp(-\langle \rho_P, H_P(m) \rangle) \varphi(mg)$$

belongs to σ (resp. $\mathcal{A}_\sigma(M_P)$) for every $g \in G(\mathbb{A})$. We have $\mathcal{A}_{P,\sigma}(G) \subset \mathcal{A}_{P,\text{cusp}}(G)$ if σ is cuspidal and $\mathcal{A}_{P,\sigma}(G) \subset \mathcal{A}_{P,\text{disc}}(G)$ if σ is discrete. The algebra $\mathcal{S}(G(\mathbb{A}))$ acts on $\mathcal{A}_{P,\sigma}(G)$ by right convolution. For every $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^*$, we denote by $I(\lambda)$ the action on $\mathcal{A}_{P,\sigma}(G)$ we get by transport from the action of $\mathcal{S}(G(\mathbb{A}))$ on $\mathcal{A}_{P,\sigma_\lambda}$ and the identification $\mathcal{A}_{P,\sigma} \rightarrow \mathcal{A}_{P,\sigma_\lambda}$.

The spaces of σ and Π naturally carry topologies making them into strict LF spaces. More precisely, for every compact-open subgroups $J_M \subset M_P(\mathbb{A}_f)$ and $J \subset G(\mathbb{A}_f)$, σ^{J_M} and Π^J are SF representations of $M_P(F_\infty)$ and $G(F_\infty)$ in the sense of [BK14] respectively and when $J'_M \subseteq J_M$, $J' \subseteq J$ are smaller compact-open subgroups the inclusions $\sigma^{J'_M} \subseteq \sigma^{J_M}$, $\Pi^{J'} \subseteq \Pi^J$ are closed embeddings.

If the central character of σ is unitary, we equip Π with the *Petersson inner product*

$$\|\varphi\|_{\text{Pet}}^2 = \langle \varphi, \varphi \rangle_{\text{Pet}} = \int_{[G]_{P,0}} |\varphi(g)|^2 dg, \quad \varphi \in \Pi.$$

2.7.3. Eisenstein series. — Let P be a standard parabolic subgroup of G . For any $\varphi \in \mathcal{A}_P(G)$, $g \in G(\mathbb{A})$ and $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^*$, we denote by

$$E(g, \varphi, \lambda) = \sum_{\delta \in P(F) \backslash G(F)} \exp(\langle \lambda, H_P(\delta g) \rangle) \varphi(\delta g)$$

the Eisenstein series where the left-hand side is obtained from the analytic continuation of the right-hand side which is only defined for $\Re(\lambda)$ in a suitable cone.

2.7.4. Let P and Q be standard parabolic subgroups of G . For any $w \in W(P, Q)$ and $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^*$, we have the intertwining operator

$$M(w, \lambda) : \mathcal{A}_P(G) \rightarrow \mathcal{A}_Q(G)$$

defined by analytic continuation from the integral

$$(M(w, \lambda)\varphi)(g) = \exp(-\langle w\lambda, H_P(g) \rangle) \int_{(N_Q \cap wN_P w^{-1})(\mathbb{A}) \backslash N_Q(\mathbb{A})} \exp(\langle \lambda, H_P(w^{-1}ng) \rangle) \varphi(w^{-1}ng) dn.$$

2.7.5. Assume that $G = G_1 \times G_2$ where G_1 and G_2 are connected reductive groups over F . This induces decompositions $P = P_1 \times P_2$, $M_P = M_{P_1} \times M_{P_2}$ and there exist two, uniquely determined, cuspidal automorphic representations σ_1, σ_2 of $M_{P_1}(\mathbb{A})$ and $M_{P_2}(\mathbb{A})$ respectively such that, setting $\Pi_1 = I_{P_1(\mathbb{A})}^{G_1(\mathbb{A})}(\sigma_1)$ and $\Pi_2 = I_{P_2(\mathbb{A})}^{G_2(\mathbb{A})}(\sigma_2)$, for every compact-open subgroups $J_1 \subseteq G_1(\mathbb{A}_f)$, $J_2 \subseteq G_2(\mathbb{A}_f)$ (resp. $J_1 \subseteq M_{P_1}(\mathbb{A}_f)$, $J_2 \subseteq M_{P_2}(\mathbb{A}_f)$), setting $J = J_1 \times J_2$, there is a topological isomorphism

$$(2.7.5.1) \quad \Pi_1^{J_1} \widehat{\otimes} \Pi_2^{J_2} \simeq \Pi^J \quad (\text{resp. } \sigma_1^{J_1} \widehat{\otimes} \sigma_2^{J_2} \simeq \sigma^J)$$

sending $\varphi_1 \otimes \varphi_2 \in \Pi_1^{J_1} \otimes \Pi_2^{J_2}$ (resp. $\varphi_1 \otimes \varphi_2 \in \sigma_1^{J_1} \otimes \sigma_2^{J_2}$) to the function $(g_1, g_2) \mapsto \varphi_1(g_1)\varphi_2(g_2)$. We then write $\Pi = \Pi_1 \boxtimes \Pi_2$ and $\sigma = \sigma_1 \boxtimes \sigma_2$ respectively.

2.7.6. Assume now that G is quasi-split. Let $\psi_N : N_0(\mathbb{A}) \rightarrow \mathbb{C}^\times$ be a continuous non-degenerate character which is trivial on $N_0(F)$. If the representation Π is ψ_N -generic, i.e. if it admits a continuous nonzero linear form $\ell : \Pi \rightarrow \mathbb{C}$ such that $\ell \circ \Pi(u) = \psi_N(u)\ell$ for every $u \in N(\mathbb{A})$, it is (abstractly) isomorphic to its *Whittaker model*

$$\mathcal{W}(\Pi, \psi_N) = \{g \in G(\mathbb{A}) \mapsto \ell(\Pi(g)\varphi) \mid \varphi \in \Pi\}.$$

We equip this last space with the topology coming from Π (thus it is a strict LF space).

If we are moreover in the situation of §2.7.5, there are decompositions $N_0 = N_{0,1} \times N_{0,2}$, $\psi_N = \psi_1 \boxtimes \psi_2$ and the isomorphism (2.7.5.1) induces one between Whittaker models

$$\mathcal{W}(\Pi_1, \psi_1)^{J_1} \widehat{\otimes} \mathcal{W}(\Pi_2, \psi_2)^{J_2} \simeq \mathcal{W}(\Pi, \psi_N)^J.$$

2.8 Relative characters

2.8.1. Let B a $G(F_\infty)$ -invariant nondegenerate symmetric bilinear form on \mathfrak{g}_∞ . We assume that the restriction of B to \mathfrak{k}_∞ is negative and the restriction of B to the orthogonal complement of \mathfrak{k}_∞ is positive. Let $(X_i)_{i \in I}$ be an orthonormal basis of \mathfrak{k}_∞ relative to $-B$. Let $C_K = -\sum_{i \in I} X_i^2$: this is a “Casimir element” of $\mathcal{U}(\mathfrak{k}_\infty)$.

2.8.2. Let \hat{K}_∞ and \hat{K} be respectively the sets of isomorphism classes of irreducible unitary representations of K_∞ and of K .

2.8.3. Let π be a discrete automorphic representation of M_P . For any $\tau \in \hat{K}$, let $\mathcal{A}_{P,\pi}(G, \tau)$ be the (finite dimensional) subspace of functions in $\mathcal{A}_{P,\pi}(G)$ which transform under K according to τ . A K -basis $\mathcal{B}_{P,\pi}$ of $\mathcal{A}_{P,\pi}(G)$ is by definition the union over of $\tau \in \hat{K}$ of orthonormal bases $\mathcal{B}_{P,\pi,\tau}$ of $\mathcal{A}_{P,\pi}(G, \tau)$ for the Petersson inner product.

2.8.4. Let

$$B : \mathcal{A}_{P,\pi}(G) \times \mathcal{A}_{P,\pi}(G) \rightarrow \mathbb{C}$$

be a continuous sesquilinear form.

Proposition 2.8.4.1. — *Let ω be a compact subset of $\mathfrak{a}_P^{G,*}$.*

1. *Let $f \in \mathcal{S}(G(\mathbb{A}))$ and $\mathcal{B}_{P,\pi}$ be a K -basis of $\mathcal{A}_{P,\pi}(G)$. The sum*

$$(2.8.4.1) \quad \sum_{\varphi \in \mathcal{B}_{P,\pi}} I_P(\lambda, f) \varphi \otimes \overline{\varphi}$$

converges absolutely in the completed projective tensor product $\mathcal{A}_{P,\pi}(G) \widehat{\otimes} \overline{\mathcal{A}_{P,\pi}(G)}$ uniformly for $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^{G,}$ such that $\Re(\lambda) \in \omega$. In particular, the sum*

$$(2.8.4.2) \quad J_B(\lambda, f) = \sum_{\varphi \in \mathcal{B}_{P,\pi}} B(I_P(\lambda, f) \varphi, \varphi)$$

is absolutely convergent uniformly for $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^{G,}$ such that $\Re(\lambda) \in \omega$. Moreover these sums do not depend on the choice of $\mathcal{B}_{P,\pi}$.*

2. *The map*

$$f \mapsto J_B(\lambda, f)$$

is a continuous linear form on $\mathcal{S}(G(\mathbb{A}))$. More precisely for $C \subset G(\mathbb{A}_f)$ a compact subset and K_0 as above, there exist $c > 0$ and a continuous semi-norm $\|\cdot\|$ on $\mathcal{S}(G(\mathbb{A}), C, K_0)$ such that for all $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^{G,}$ such that $\Re(\lambda) \in \omega$ and $f \in \mathcal{S}(G(\mathbb{A}), C, K_0)$ we have*

$$|J_B(\lambda, f)| \leq c \|f\|.$$

Remark 2.8.4.2. — An examination of the proof below show that the assertion 2 also holds *mutatis mutandis* if $f \in C_c^r(G(\mathbb{A}))$ with r large enough. The semi-norm is then taken among the norms $\|\cdot\|_{r,X,Y}$ for which the sum of the degrees of X and Y is less than r .

Proof. — By definition of the projective tensor product topology, it suffices to show the following: for every continuous semi-norm p on $\mathcal{A}_{P,\pi}(G)$, the series

$$\sum_{\varphi \in \mathcal{B}_{P,\pi}} p(I_P(\lambda, f) \varphi) p(\varphi)$$

is absolutely convergent uniformly for $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^{G,*}$ such that $\Re(\lambda) \in \omega$. Let $K_0 \subset K^\infty$ be a normal open compact subgroup by which f is biinvariant. The series above can be rewritten as

$$(2.8.4.3) \quad \sum_{\tau \in \hat{K}} \sum_{\varphi \in \mathcal{B}_{P,\pi,\tau}} p(I_P(\lambda, f)\varphi)p(\varphi).$$

and the only representations that contribute to (2.8.4.3) are the representations τ that admits K_0 -invariant vectors. Then the elements $\varphi \in \mathcal{B}_{P,\pi,\tau}$ are automatically K_0 -fixed. For any $\tau \in \hat{K}_\infty$, let $\mathcal{A}_{P,\pi}(G, K_0, \tau) \subset \mathcal{A}_{P,\pi}(G)$ be the subspace of functions that are right K_0 -invariant and transform under K_∞ according to τ . By [Wal92] §10.1, there exist $c > 0$ and an integer r such that for all $\varphi \in \mathcal{A}_{P,\pi}(G)^{K_0}$ we have

$$p(\varphi) \leq c \|R(1 + C_K)^r \varphi\|_{\text{Pet}}.$$

For any $\tau \in \hat{K}_\infty$ or \hat{K} , let $\lambda_\tau \geq 0$ be the eigenvalue of C_K acting on τ . We have

$$\begin{aligned} \|R(1 + C_K)^r I_P(\lambda, f)\varphi\|_{\text{Pet}} &= \|I_P(\lambda, L((1 + C_K)^r)f)\varphi\|_{\text{Pet}} \\ &= (1 + \lambda_\tau)^{-N} \|I_P(\lambda, f_{r,N})\varphi\|_{\text{Pet}} \end{aligned}$$

where $f_{r,N} = R((1 + C_K)^N)L((1 + C_K)^r)f$. Let $C \subset G(\mathbb{A}_f)$ be a compact subset. There exists $c_1 > 0$ and a semi-norm $\|\cdot\|$ on $\mathcal{S}(G(\mathbb{A}), C, K_0)$ (among those of §2.5.2) such that for any $f \in \mathcal{S}(G(\mathbb{A}), C, K_0)$ and $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^{G,*}$ such that $\Re(\lambda) \in \omega$ we have

$$\|I_P(\lambda, f_{r,N})\varphi\|_{P,\pi} \leq c_1 \|f\| \|\varphi\|_{\text{Pet}}.$$

Thereby we are reduced to prove for large enough N the convergence of

$$(2.8.4.4) \quad \sum_{\tau \in \hat{K}_\infty} (1 + \lambda_\tau)^{r-N} \dim(\mathcal{A}_{P,\pi}^\infty(G, K_0, \tau)).$$

However there exist $c_2 > 0$ and $m \geq 1$ such that $\dim(\mathcal{A}_{P,\pi}^\infty(G, K_0, \tau)) \leq c_2(1 + \lambda_\tau)^m$ (see the proof of [Mö0] lemma 6.1). So the convergence of (2.8.4.4) is reduced to that of $\sum_{\tau \in \hat{K}_\infty} (1 + \lambda_\tau)^{-N}$ which is well-known.

Finally it is easy to show that $J_B(f)$ does not depend on the choice of the basis $\mathcal{B}_{P,\pi,\tau}$. \square

Proposition 2.8.4.3. — *Let $K_0 \subset K^\infty$ be a normal open compact subgroup. For any integer $m \geq 1$ there exist $Z \in \mathcal{U}(\mathfrak{g}_\mathbb{C})$, $g_1 \in C_c^\infty(G(\mathbb{A}))$ and $g_2 \in C_c^m(G(\mathbb{A}))$ such that*

- Z , g_1 and g_2 are invariant under K_∞ -conjugation ;
- g_1 and g_2 are K_0 -biinvariant ;
- for any $f \in \mathcal{S}(G(\mathbb{A}))$ that is K_0 -biinvariant we have:

$$f = f * g_1 + (f * Z) * g_2.$$

For large enough m , we have

$$J_B(\lambda, f) = \sum_{\varphi \in \mathcal{B}_{P,\pi}} B(I_P(\lambda, f)\varphi, I_P(\bar{\lambda}, g_1^\vee)\varphi) + \sum_{\varphi \in \mathcal{B}_{P,\pi}} B(I_P(\lambda, f * Z)\varphi, I_P(\bar{\lambda}, g_2^\vee)\varphi)$$

where the sums are absolutely convergent and $g_i^\vee(x) = \overline{g_i(x^{-1})}$.

Proof. — The first part of the proposition is lemma 4.1 and corollary 4.2 of [Art78]. Once we have noticed that the operators $I_P(\lambda, g_i)$ preserve the spaces $\mathcal{A}_{P,\pi}(G, \tau)$, the second part results from an easy computation in a finite dimensional space. \square

2.9 Decomposition according to cuspidal data and automorphic kernels

2.9.1. Cuspidal data. — Let $\underline{\mathfrak{X}}(G)$ be the set of pairs (M_P, σ) where P is a standard parabolic subgroup of G and σ is an isomorphism class of cuspidal automorphic representations of $M_P(\mathbb{A})$ with central character trivial on A_P^∞ . We let $\mathfrak{X}(G)$ be the quotient of $\underline{\mathfrak{X}}(G)$ by the equivalence relation defined as follows: $(M_P, \sigma) \sim (M_Q, \tau)$ if there exists $w \in W(P, Q)$ such that $w\sigma w^{-1} \simeq \tau$. We call $\mathfrak{X}(G)$ the set of *cuspidal data* for G . For every standard parabolic subgroup P of G , the natural inclusion $\underline{\mathfrak{X}}(M_P) \subset \underline{\mathfrak{X}}(G)$ descends to a finite map $\mathfrak{X}(M_P) \rightarrow \mathfrak{X}(G)$. For $\chi \in \mathfrak{X}(G)$ represented by a pair (M_P, σ) , we denote by χ^\vee the cuspidal datum associated to (M_P, σ^\vee) where σ^\vee stands for the complex conjugate of σ .

2.9.2. Langlands decomposition For $(M_P, \sigma) \in \underline{\mathfrak{X}}(G)$, we let $\mathcal{S}_\sigma([G]_P)$ be the subspace of $\varphi \in \mathcal{S}([G]_P)$ such that the function

$$\varphi_\lambda(g) := \int_{A_P^\infty} \exp(-\langle \rho_P + \lambda, H_P(a) \rangle) \varphi(ag) da, \quad g \in [G]_P,$$

belongs to $\mathcal{A}_{P, \sigma_\lambda}(G)$ for every $\lambda \in \mathfrak{a}_{P, \mathbb{C}}^*$.

Let $P \subset G$ a standard parabolic subgroup, $\chi \in \mathfrak{X}(G)$ be a cuspidal datum and $\{(M_{Q_i}, \sigma_i) \mid i \in I\}$ be the (possibly empty but finite) inverse image of χ in $\underline{\mathfrak{X}}(M_P)$. We define $L_\chi^2([G]_P)$ as the closure of the subspace $\sum_{i \in I} E_{Q_i}^P(\mathcal{S}_{\sigma_i}([G]_{Q_i}))$ in $L^2([G]_P)$. More generally, for w be a weight on

\mathfrak{a}_P (see §2.5.4), we let $L_{w, \chi}^2([G]_P)$ be the closure of $\sum_{i \in I} E_{Q_i}^P(\mathcal{S}_{\sigma_i}([G]_{Q_i}))$ in $L_w^2([G]_P)$. We define

similarly a subspace $L_\chi^2([G]_{P,0}) \subset L^2([G]_{P,0})$. By Langlands (see e.g. [MW94, Proposition II.2.4]), we have Hilbert decompositions

$$(2.9.2.1) \quad L^2([G]_P) = \widehat{\bigoplus_{\chi \in \mathfrak{X}(G)} L_\chi^2([G]_P)} \text{ and } L^2([G]_{P,0}) = \widehat{\bigoplus_{\chi \in \mathfrak{X}(G)} L_\chi^2([G]_{P,0})}.$$

More generally, for every weight w on \mathfrak{a}_P we have a Hilbert decomposition

$$(2.9.2.2) \quad L_w^2([G]_P) = \widehat{\bigoplus_{\chi \in \mathfrak{X}(G)} L_{w, \chi}^2([G]_P)}.$$

Let $\mathfrak{X} \subseteq \mathfrak{X}(G)$ be a subset and w be a weight on \mathfrak{a}_P . We set

$$L_{w, \mathfrak{X}}^2([G]_P) := \widehat{\bigoplus_{\chi \in \mathfrak{X}} L_\chi^2([G]_P)}, \quad L_w^{2, \mathfrak{X}}([G]_P) := \widehat{\bigoplus_{\chi \in \mathfrak{X}(G) \setminus \mathfrak{X}} L_{w, \chi}^2([G]_P)}$$

and when $w = 0$, we just drop the index w . We have

$$(2.9.2.3) \quad \text{Let } w \text{ and } w' \text{ be two weights on } \mathfrak{a}_P. \text{ The orthogonal projections } L_w^2([G]_P) \rightarrow L_{w, \mathfrak{X}}^2([G]_P) \\ \text{and } L_{w'}^2([G]_P) \rightarrow L_{w', \mathfrak{X}}^2([G]_P) \text{ coincide on the intersection } L_w^2([G]_P) \cap L_{w'}^2([G]_P).$$

Indeed, we have $L_w^2([G]_P) \cap L_{w'}^2([G]_P) = L_{w''}^2([G]_P)$ where $w'' = \max(w, w')$ is a weight on \mathfrak{a}_P and this allows to restrict to the case where $w \leq w'$. The claim then follows from the fact, easy to see from the definition, that the natural inclusion $L_{w'}^2([G]_P) \subset L_w^2([G]_P)$ sends $L_{w', \mathfrak{X}}^2([G]_P)$ (resp. $L_{w'}^{2, \mathfrak{X}}([G]_P)$) into $L_{w, \mathfrak{X}}^2([G]_P)$ (resp. $L_w^{2, \mathfrak{X}}([G]_P)$).

We will denote by $\varphi \mapsto \varphi_{\mathfrak{X}}$ the orthogonal projection $L_w^2([G]_P) \rightarrow L_{w, \mathfrak{X}}^2([G]_P)$ (by (2.9.2.3) such a notation shouldn't lead to any confusion). These projections are $G(\mathbb{A})$ -equivariant and so preserve the subspaces of smooth vectors.

2.9.3. Let again $\mathfrak{X} \subseteq \mathfrak{X}(G)$ be a subset. We set

$$\mathcal{S}_{\mathfrak{X}}([G]_P) := \mathcal{S}([G]_P) \cap L_{\mathfrak{X}}^2([G]_P), \quad \mathcal{S}^{\mathfrak{X}}([G]_P) := \mathcal{S}([G]_P) \cap L^{2, \mathfrak{X}}([G]_P), \quad \mathcal{C}_{\mathfrak{X}}([G]_P) := \mathcal{C}([G]_P) \cap L_{\mathfrak{X}}^2([G]_P).$$

We also define $\mathcal{T}_{\mathfrak{X}}([G]_P)$ (resp. $\mathcal{T}_{N,\mathfrak{X}}([G]_P)$ for $N > 0$) as the orthogonal of $\mathcal{S}^{\mathfrak{X}}([G]_P)$ in $\mathcal{T}([G]_P)$ (resp. in $\mathcal{T}_N([G]_P)$). Then, $\mathcal{T}_{N,\mathfrak{X}}([G]_P)$ is a closed subspace of $\mathcal{T}_N([G]_P)$ hence is a strict LF space. We equip $\mathcal{T}_{\mathfrak{X}}([G]_P) = \bigcup_{N>0} \mathcal{T}_{N,\mathfrak{X}}([G]_P)$ with the inductive limit locally convex topology (it is a LF space).

Proposition 2.9.3.1. — *Let $\mathfrak{X} \subseteq \mathfrak{X}(G)$ be a subset and $P \subseteq G$ be a standard parabolic subgroup.*

1. *For every standard parabolic subgroup $Q \subset P$, we have $E_Q^P(\mathcal{S}_{\mathfrak{X}}([G]_Q)) \subset \mathcal{S}_{\mathfrak{X}}([G]_P)$ and $E_Q^P(\mathcal{S}^{\mathfrak{X}}([G]_P)) \subset \mathcal{S}^{\mathfrak{X}}([G]_P)$.*
2. *For every $\varphi \in \mathcal{S}([G]_P)$, we have $\varphi_{\mathfrak{X}} \in \mathcal{S}_{\mathfrak{X}}([G]_P)$. Moreover, the linear map $\varphi \in \mathcal{S}([G]_P) \mapsto \varphi_{\mathfrak{X}} \in \mathcal{S}_{\mathfrak{X}}([G]_P)$ is continuous and the series*

$$(2.9.3.4) \quad \sum_{\chi \in \mathfrak{X}(G)} \varphi_{\chi}$$

converges absolutely (see §A.0.2) to φ in $\mathcal{S}([G]_P)$.

3. *For every $\varphi \in \mathcal{T}([G]_P)$, there exists a unique function $\varphi_{\mathfrak{X}} \in \mathcal{T}_{\mathfrak{X}}([G]_P)$ such that $\langle \varphi, \psi_{\mathfrak{X}} \rangle_{[G]_P} = \langle \varphi_{\mathfrak{X}}, \psi \rangle_{[G]_P}$ for every $\psi \in \mathcal{S}([G]_P)$. Moreover, the linear map $\varphi \in \mathcal{T}([G]_P) \mapsto \varphi_{\mathfrak{X}} \in \mathcal{T}_{\mathfrak{X}}([G]_P)$ is $A_P^{\infty} \times G(\mathbb{A})$ -equivariant and continuous and for every parabolic subgroup $Q \subseteq P$ and $\varphi \in \mathcal{T}([G]_P)$ we have*

$$(2.9.3.5) \quad (\varphi_{\mathfrak{X}})_Q = (\varphi_Q)_{\mathfrak{X}}.$$

4. *For every compact-open subgroup $J \subset G(\mathbb{A}_f)$ and $N > 0$, there exists $M > N$ such that for $\varphi \in \mathcal{T}_N([G]_P)^J$ the series (2.9.3.4) converges absolutely to φ in $\mathcal{T}_M([G]_P)$.*

Proof. —

1. Up to replacing \mathfrak{X} by its complement in $\mathfrak{X}(G)$, the two inclusions are equivalent and so we just need to prove the first. By definition of $\mathcal{S}_{\mathfrak{X}}([G]_P)$, we need to establish that $E_Q^P(\mathcal{S}_{\mathfrak{X}}([G]_Q))$ is orthogonal to $\mathcal{S}^{\mathfrak{X}}([G]_P)$ or, by adjunction, that for every $\varphi \in \mathcal{S}^{\mathfrak{X}}([G]_P)$, φ_Q is orthogonal to $\mathcal{S}_{\mathfrak{X}}([G]_Q)$. Denote by $\{(M_{Q_i}, \sigma_i) \mid i \in I\}$ the (possibly infinite) inverse image of \mathfrak{X} in $\underline{\mathfrak{X}}(M_Q)$ and let $\varphi \in \mathcal{S}^{\mathfrak{X}}([G]_P)$. By adjunction again, and the definition of $\mathcal{S}^{\mathfrak{X}}([G]_P)$, φ_Q is orthogonal to $\sum_{i \in I} E_{Q_i}^Q(\mathcal{S}_{\sigma_i}([G]_{Q_i}))$. Let $\kappa \in C_c^{\infty}(\mathfrak{a}_Q)$. Then, $(\kappa \circ H_Q)E_{Q_i}^Q(\psi) = E_{Q_i}^Q((\kappa \circ H_Q)\psi)$ and $(\kappa \circ H_Q)\psi \in \mathcal{S}_{\sigma_i}([G]_{Q_i})$ for every $i \in I$ and $\psi \in \mathcal{S}_{\sigma_i}([G]_{Q_i})$. It follows that $(\kappa \circ H_Q)\varphi_Q$ is also orthogonal to $\sum_{i \in I} E_{Q_i}^Q(\mathcal{S}_{\sigma_i}([G]_{Q_i}))$. Besides, by Lemma 2.5.10.1, the function $(\kappa \circ H_Q)\varphi_Q$ belongs to $\mathcal{S}([G]_Q)$. Therefore, by definition of $L_{\mathfrak{X}}^2([G]_Q)$, $(\kappa \circ H_Q)\varphi_Q$ is orthogonal to $L_{\mathfrak{X}}^2([G]_Q)$ and in particular to $\mathcal{S}_{\mathfrak{X}}([G]_Q)$. Finally, there certainly exists a sequence $\kappa_n \in C_c^{\infty}(\mathfrak{a}_Q)$ such that $(\kappa_n \circ H_Q)\varphi_Q$ converges to φ_Q in $\mathcal{T}([G]_Q)$ and we conclude that φ_Q is indeed orthogonal to $\mathcal{S}_{\mathfrak{X}}([G]_Q)$.
2. We prove the first part by induction on $\dim(\mathfrak{a}_0) - \dim(\mathfrak{a}_P)$. Let $\varphi \in \mathcal{S}([G]_P)$, $Q \subset P$ be a standard parabolic subgroup, $\lambda \in (\mathfrak{a}_Q^*)^{P++}$. By Proposition 2.5.11.1, in order to show that $\varphi_{\mathfrak{X}} \in \mathcal{S}([G]_P)$ it suffices to check that $(\varphi_{\mathfrak{X}})_Q \in L_{2\rho_Q+\lambda}^2([G]_Q)^{\infty}$. Again by Proposition 2.5.11.1 and (2.9.2.3), this will follow from the equality

$$(2.9.3.6) \quad (\varphi_{\mathfrak{X}})_Q = (\varphi_Q)_{\mathfrak{X}}.$$

If $Q = P$ the above identity is tautological and this already settles the case $P = P_0$. If $Q \neq P$, by the induction hypothesis we have $\mathcal{S}([G]_Q) = \mathcal{S}_{\mathfrak{X}}([G]_Q) \oplus \mathcal{S}^{\mathfrak{X}}([G]_Q)$ and (2.9.3.6) is equivalent to

$$\langle (\varphi_{\mathfrak{X}})_Q, \psi \rangle_{[G]_Q} = \langle (\varphi_Q)_{\mathfrak{X}}, \psi \rangle_{[G]_Q}$$

for every $\psi \in \mathcal{S}_{\mathfrak{X}}([G]_Q) \cup \mathcal{S}^{\mathfrak{X}}([G]_Q)$. This last equality follows directly, by adjunction, from point 1.

The continuity of $\varphi \in \mathcal{S}([G]_P) \mapsto \varphi_{\mathfrak{X}} \in \mathcal{S}_{\mathfrak{X}}([G]_P)$ is easy to obtain from the closed graph theorem. We now show the absolute convergence of $\sum_{\chi \in \mathfrak{X}(G)} \varphi_{\chi}$. Fix a compact-open subgroup $J \subset G(\mathbb{A}_f)$. By Proposition 2.5.11.1 and (2.9.3.6), there exists an increasing family $(\|\cdot\|_n)_n$ of Hilbertian norms defining the topology of $\mathcal{S}([G]_P)^J$ such that for each n , denoting by \mathcal{H}_n the Hilbert completion of $\mathcal{S}([G]_P)^J$ with respect to $\|\cdot\|_n$, the linear maps $\varphi \mapsto \varphi_{\chi}$, $\chi \in \mathfrak{X}(G)$, extend to orthogonal projections onto two by two orthogonal subspaces of \mathcal{H}_n . In particular, for each n and $\varphi \in \mathcal{H}_n$ we have

$$\sum_{\chi \in \mathfrak{X}(G)} \|\varphi_{\chi}\|_n^2 < \infty.$$

On the other hand, the space $\mathcal{S}([G]_P)^J$ being nuclear (2.5.8.5), for each n there exists $m \geq n$ such that the induced linear map $\mathcal{H}_m \rightarrow \mathcal{H}_n$ is Hilbert-Schmidt. By the Cauchy-Schwarz inequality, this implies

$$\sum_{\chi \in \mathfrak{X}(G)} \|\varphi_{\chi}\|_n \leq \left(\sum_{\chi \in \mathfrak{X}(G)} \|\varphi_{\chi}\|_m^2 \right)^{1/2} \left(\sum_{\chi \in \mathfrak{X}(G)} \left\| \frac{\varphi_{\chi}}{\|\varphi_{\chi}\|_m} \right\|_n^2 \right)^{1/2} < \infty$$

for $\varphi \in \mathcal{H}_m$. Since this holds for every n , the absolute convergence of $\sum_{\chi \in \mathfrak{X}(G)} \varphi_{\chi}$ follows. That the sum necessarily converges to φ is obvious (e.g. because it converges to φ in $L^2([G]_P)$).

3. The first part follows from 2., (2.5.8.6) and the Dixmier-Malliavin theorem. The continuity of $\varphi \mapsto \varphi_{\mathfrak{X}}$ follows from the closed graph theorem and the equivariance with respect to the $A_P^{\infty} \times G(\mathbb{A})$ -action follows from uniqueness and equivariance of the projection $L^2([G]_P) \rightarrow L_{\mathfrak{X}}^2([G]_P)$. Also, (2.9.3.5) can be proven the same way as (2.9.3.6), using 1. and adjunction.
4. Let $(\|\cdot\|_n)_n$ be an increasing family of Hilbertian norms defining the topology of $\mathcal{S}([G]_P)^J$ satisfying the same property as before: denoting by \mathcal{H}_n the completion of $\mathcal{S}([G]_P)^J$ with respect to $\|\cdot\|_n$, for each n the maps $\varphi \mapsto \varphi_{\chi}$, $\chi \in \mathfrak{X}(G)$, extend to orthogonal projections onto two by two orthogonal subspaces of \mathcal{H}_n . We may also assume that the norms $\|\cdot\|_n$ are $G(F_{\infty})$ -continuous (that is, the $G(F_{\infty})$ -action extends to \mathcal{H}_n). Let \mathcal{H}_{-n} be the dual Hilbert space equipped with the dual Hilbertian norm $\|\cdot\|_{-n}$. The adjoint of the natural linear map $\mathcal{H}_m \rightarrow \mathcal{H}_n$, $m \geq n$, is a continuous $G(F_{\infty})$ -equivariant inclusion $\mathcal{H}_{-n} \hookrightarrow \mathcal{H}_{-m}$. Moreover, by (2.5.8.6) and the Dixmier-Malliavin theorem we have a natural inclusion $\mathcal{H}_{-n}^{\infty} \hookrightarrow \mathcal{T}([G]_P)^J$ and an equality $\mathcal{T}([G]_P)^J = \bigcup_n \mathcal{H}_{-n}^{\infty}$ of LF spaces (that this identification is a topological isomorphism follows from the open mapping theorem for LF spaces). Finally, there exists n such that $\mathcal{T}_N([G]_P)^J \subset \mathcal{H}_{-n}^{\infty}$ and for each m there exists M such that $\mathcal{H}_{-m}^{\infty} \subset \mathcal{T}_M([G]_P)^J$. Therefore, it suffices to show that for every n there exists $m \geq n$ such that the series (2.9.3.4) converges absolutely in \mathcal{H}_{-m} for every $\varphi \in \mathcal{H}_{-n}$. This follows from the same argument as before since for m large enough the inclusion $\mathcal{H}_{-n} \hookrightarrow \mathcal{H}_{-m}$ is Hilbert-Schmidt.

□

2.9.4. Let $\mathfrak{X} \subseteq \mathfrak{X}(G)$ be a subset. By the previous proposition, the projection $\varphi \mapsto \varphi_{\mathfrak{X}}$ maps $\mathcal{T}([G]_P)$ and $\mathcal{S}([G]_P)$ continuously onto $\mathcal{T}_{\mathfrak{X}}([G]_P)$ and $\mathcal{S}_{\mathfrak{X}}([G]_P)$ respectively. As $\mathcal{S}([G]_P)$ is dense in $\mathcal{T}([G]_P)$ this entails that

$$(2.9.4.7) \quad \mathcal{S}_{\mathfrak{X}}([G]_P) \text{ is dense in } \mathcal{T}_{\mathfrak{X}}([G]_P).$$

More precisely, let $N > 0$ and $J \subset G(\mathbb{A}_f)$ be a compact-open subgroup. There exists $M > L > N$ such that the closure of $\mathcal{S}([G]_P)^J$ in $\mathcal{T}_L([G]_P)^J$ contains $\mathcal{T}_N([G]_P)^J$ and the projection

$\varphi \mapsto \varphi_{\mathfrak{X}}$ restricts to a continuous linear mapping $\mathcal{T}_L([G]_P)^J \rightarrow \mathcal{T}_{M,\mathfrak{X}}([G]_P)^J$. Therefore:
(2.9.4.8)

There exists $M > 0$ such that the closure of $\mathcal{S}_{\mathfrak{X}}([G])^J$ in $\mathcal{T}_{M,\mathfrak{X}}([G])^J$ contains $\mathcal{T}_{N,\mathfrak{X}}([G])^J$.

2.9.5. Assume that $G = G_1 \times G_2$ where G_1 and G_2 are connected reductive groups over F . We have then a natural identification $\mathfrak{X}(G) = \mathfrak{X}(G_1) \times \mathfrak{X}(G_2)$. For subsets $\mathfrak{X}_i \subseteq \mathfrak{X}(G_i)$, $i = 1, 2$, setting $\mathfrak{X} = \mathfrak{X}_1 \times \mathfrak{X}_2$, the space $L_{\mathfrak{X}_1}^2([G_1]) \otimes L_{\mathfrak{X}_2}^2([G_2])$ is dense in $L_{\mathfrak{X}}^2([G])$ (this follows from the fact that for $(M_{P_i}, \sigma_i) \in \mathfrak{X}_i$, $i = 1, 2$, $\mathcal{S}_{\sigma_1}([G_1]_{P_1}) \otimes \mathcal{S}_{\sigma_2}([G_2]_{P_2})$ is dense in $\mathcal{S}_{\sigma_1 \boxtimes \sigma_2}([G]_{P_1 \times P_2})$). Using again (2.5.8.5) and (A.0.6.6), we see that for any compact-open subgroups $J_i \subset G_i(\mathbb{A}_f)$, $i = 1, 2$, setting $J = J_1 \times J_2$, the isomorphisms (2.5.9.7) restrict to isomorphisms

$$(2.9.5.9) \quad \mathcal{C}_{\mathfrak{X}_1}([G_1])^{J_1} \widehat{\otimes} \mathcal{C}_{\mathfrak{X}_2}([G_2])^{J_2} \simeq \mathcal{C}_{\mathfrak{X}}([G])^J, \quad \mathcal{S}_{\mathfrak{X}_1}([G_1])^{J_1} \widehat{\otimes} \mathcal{S}_{\mathfrak{X}_2}([G_2])^{J_2} \simeq \mathcal{S}_{\mathfrak{X}}([G])^J.$$

2.9.6. Generic cuspidal data. — Let $n \geq 1$. We say that a cuspidal datum $\chi \in \mathfrak{X}(\mathrm{GL}_n)$ is *generic* if it is represented by a pair (M_P, σ) with

$$M_P = \mathrm{GL}_{n_1} \times \dots \times \mathrm{GL}_{n_k}$$

a standard Levi subgroup of GL_n and

$$\sigma = \sigma_1 \boxtimes \dots \boxtimes \sigma_k$$

a cuspidal automorphic representation of $M_P(\mathbb{A})$ whose central character is trivial on A_P^∞ and such that $\sigma_i \neq \sigma_j$ for $1 \leq i < j \leq k$.

More generally, assume that G is a product of the form $\mathrm{Res}_{K_1/F} \mathrm{GL}_{n_1} \times \dots \times \mathrm{Res}_{K_r/F} \mathrm{GL}_{n_r}$, where K_1, \dots, K_r are finite extensions of F . Then a cuspidal datum $\chi \in \mathfrak{X}(G)$ can be seen as a r -uple (χ_1, \dots, χ_r) where $\chi_i \in \mathfrak{X}(\mathrm{GL}_{n_i, K_i})$, $i = 1, \dots, r$, and we say that χ is *generic* if each of the χ_i is so.

The following proposition is a consequence of Langlands's spectral decomposition [MW94, VI.2.2], the description of the discrete spectrum of GL_n by Mœglin-Waldspurger [MW89] (which implies that for χ a generic cuspidal datum and Q a parabolic subgroup, the discrete spectrum of $L_\chi^2([G]_Q)$ is entirely cuspidal), the computation of the constant terms of cuspidal Eisenstein series [MW94, Proposition II.1.7] and the fact that intertwining operators are unitary for purely imaginary arguments.

Proposition 2.9.6.1. — *Let $\chi \in \mathfrak{X}(G)$ be a generic cuspidal datum and let P be a standard parabolic subgroup of G . Then, for every $\varphi \in L_\chi^2([G])$ we have $\varphi_P \in L_\chi^2([G]_P)$.*

Corollary 2.9.6.2. — *Let $\chi \in \mathfrak{X}(G)$ be a generic cuspidal datum, P be a standard parabolic subgroup of G and χ_M be the inverse image of χ in $\mathfrak{X}(M_P)$. Then, for every $\varphi \in \mathcal{S}_\chi([G])$ and $s \in \mathcal{H}_{>0}$, the function*

$$\varphi_{P,s} : m \in [M_P] \mapsto \delta_P(m)^{s-1/2} \varphi_P(m)$$

belongs to $\mathcal{C}_{\chi_M}([M_P])$. Moreover, the family of linear maps

$$\mathcal{S}_\chi([G]) \rightarrow \mathcal{C}_{\chi_M}([M_P]), \varphi \mapsto \varphi_{P,s}$$

for $s \in \mathcal{H}_{>0}$ is holomorphic.

Proof. — Let $\varphi \in \mathcal{S}_\chi([G])$. Note that by Proposition 2.9.3.1.1, $\varphi_{P,s}$ is orthogonal to $\mathcal{S}^{\chi_M}([M_P])$ for every $s \in \mathbb{C}$. Hence, we just need to show that for every $s \in \mathcal{H}_{>0}$ the function $\varphi_{P,s}$ belongs to $\mathcal{C}([M_P])$ and the map $s \in \mathcal{H}_{>0} \mapsto \varphi_{P,s} \in \mathcal{C}([M_P])$ is holomorphic. As for every $X \in \mathfrak{m}_\infty$ we have $R(X)\varphi_{P,s} = (2s-1)\langle \rho_P, X \rangle \varphi_{P,s} + (R(X)\varphi)_{P,s}$ (where we consider ρ_P as an element of the dual space \mathfrak{m}_∞^*), by the equality (2.5.6.3), it suffices to show that for every $d > 0$ and $s \in \mathcal{H}_{>0}$ we have

$\varphi_{P,s} \in L^2_{\sigma,d}([M_P])$ and that the map $s \in \mathcal{H}_{>0} \mapsto \varphi_{P,s} \in L^2_{\sigma,d}([M_P])$ is holomorphic. By Lemma 2.5.10.1 and (2.4.2.5), there exists $c > 0$ such that $\varphi_{P,s} \in L^2_{\sigma,d}([M_P])$ for every $s \in \mathcal{H}_{>c}$ and $d > 0$. On the other hand, by Proposition 2.9.6.1, we have $\varphi_{P,0} \in L^2([M_P])$. By Hölder inequality, for every $s \in \mathcal{H}_{>0}$, $t > \Re(s)$ and $d > 0$, we have

$$\|\varphi_{P,s}\|_{L^2_{\sigma,d}} \leq \|\varphi_{P,0}\|_{L^2}^{1-\Re(s)/t} \|\varphi_{P,t}\|_{L^2_{\sigma,t d/\Re(s)}}^{\Re(s)/t}$$

and it follows that $\varphi_{P,s} \in L^2_{\sigma,d}([M_P])$ for every $s \in \mathcal{H}_{>0}$ and $d > 0$. The holomorphy of the map $s \in \mathcal{H}_{>0} \mapsto \varphi_{P,s} \in L^2_{\sigma,d}([M_P])$ is equivalent to the holomorphy of $s \in \mathcal{H}_{>0} \mapsto \langle \varphi_{P,s}, \psi \rangle_{[M_P]}$ for every $\psi \in L^2_{\sigma,-d}([M_P])$ but this follows from the inequality

$$|\varphi_{P,s}| \leq |\varphi_{P,t_1}| + |\varphi_{P,t_2}|$$

for every $s \in \mathbb{C}$ and $t_2 > \Re(s) > t_1$. \square

2.9.7. Let $\chi \in \mathfrak{X}(G)$ be a generic cuspidal datum represented by a pair $(M_P, \pi) \in \mathfrak{X}(G)$. Set $\Pi = I_{P(\mathbb{A})}^{G(\mathbb{A})}(\pi) = \mathcal{A}_{P,\pi}(G)$ for the normalized smooth induction of π . Let $\mathcal{B}_{P,\pi}$ be a K -basis of Π as in §2.8.3. For $\varphi \in \mathcal{S}([G])$ and $\lambda \in i\mathfrak{a}_P^*$ the series

$$(2.9.7.10) \quad \varphi_{\Pi_\lambda} = \sum_{\psi \in \mathcal{B}_{P,\pi}} \langle \varphi, E(\psi, \lambda) \rangle_{[G]} E(\psi, \lambda)$$

converges absolutely in $\mathcal{T}_N([G])$ for some N (that may a priori depend on λ). Indeed by [BL19, Corollary 6.5] and (A.0.4.3), there exists N such that the linear map $\psi \in \mathcal{A}_{P,\sigma}(G) \mapsto E(\psi, \lambda) \in \mathcal{T}_N([G])$ is continuous so that the claim follows from Proposition 2.8.4.1 and the Dixmier-Malliavin theorem. The next theorem is a slight restatement of (part of) the main result of [Lap13]⁴. We refer the reader to §A.0.8 for the notion of Schwartz function valued in a TVS.

Theorem 2.9.7.1. — *[(Lapid)] There exists $N > 0$ such that for $\varphi \in \mathcal{C}([G])$, the series (2.9.7.10) still makes sense (that is the scalar products $\langle \varphi, E(\psi, \lambda) \rangle_{[G]}$ are convergent) and converges in $\mathcal{T}_N([G])$ for every $\lambda \in i\mathfrak{a}_P^*$. Moreover, the function $\lambda \in i\mathfrak{a}_P^* \mapsto \varphi_{\Pi_\lambda} \in \mathcal{T}_N([G])$ is Schwartz and if $\varphi \in \mathcal{C}_\chi([G])$ we have the equality*

$$\varphi = \int_{i\mathfrak{a}_P^*} \varphi_{\Pi_\lambda} d\lambda$$

(the right hand-side being absolutely convergent in $\mathcal{T}_N([G])$).

Proof. — Note that G satisfies condition (HP) of [Lap13]: it is proven in *loc. cit.* that general linear groups satisfy (HP) and it is straightforward to check that products of groups satisfying (HP) again satisfy (HP). The first part of the theorem is then a consequence of [Lap13, Proposition 5.1]. Indeed, by Dixmier-Malliavin we may assume that $\varphi = R(f)\varphi'$ where $\varphi' \in \mathcal{C}([G])$ and $f \in C_c^\infty(G(\mathbb{A}))$. By *loc. cit.* the scalar product $\langle \varphi, E(\psi, \lambda) \rangle_{[G]}$ converges for every $\psi \in \mathcal{B}_{P,\pi}$ and there exists $N > 0$ such that $\psi \mapsto E(\psi, \lambda)$ factorizes through a continuous linear mapping $\Pi \rightarrow \mathcal{T}_N([G])$ for every $\lambda \in i\mathfrak{a}_P^*$. As

$$\langle \varphi, E(\psi, \lambda) \rangle_{[G]} = \langle \varphi', E(R(f^*)\psi, \lambda) \rangle_{[G]}, \quad \varphi \in \mathcal{B}_{P,\pi},$$

we deduce by Proposition 2.8.4.1 that the series (2.9.7.10) converges absolutely in $\mathcal{T}_N([G])$ for every $\lambda \in i\mathfrak{a}_P^*$. That the function $\lambda \in i\mathfrak{a}_P^* \mapsto \varphi_{\Pi_\lambda} \in \mathcal{T}_N([G])$ is Schwartz follows similarly from [Lap13, Corollary 5.7]. The last part of the theorem is a consequence of [Lap13, Theorem 4.5] since χ is generic and therefore $L_\chi^2([G])$ is included in the “induced from cuspidal part” $L_c^2([G])$ of $L^2([G])$, with the notation of *loc. cit.*, and moreover the stabilizer of the pair (M_P, π) in W is trivial. \square

⁴Note that in *loc. cit.* the Harish-Chandra Schwartz space $\mathcal{C}([G])$ is denoted by $\mathcal{S}(G(F)\backslash G(\mathbb{A}))$

2.9.8. Automorphic kernels. — The right convolution by $f \in \mathcal{S}(G(\mathbb{A}))$ on each space of the decompositions (2.9.2.1) gives integral operators whose kernels are respectively denoted by $K_f(x, y)$, $K_{f,\chi}(x, y)$, $K_f^0(x, y)$ and $K_{f,\chi}^0(x, y)$ where $x, y \in G(\mathbb{A})$. If the context is clear, we shall omit the subscript f in the notation. The kernels are related by the following equality for all $x, y \in G(\mathbb{A})$

$$K_\chi^0(x, y) = \int_{A_G^\infty} K_\chi(x, ay) da.$$

Lemma 2.9.8.1. — *For every $N > 0$, there exists $N' > 0$ such that*

$$(2.9.8.11) \quad \sum_{\chi \in \mathfrak{X}(G)} |K_\chi(x, y)| \leq \|x\|_{[G]}^{N'} \|y\|_{[G]}^{-N}, \quad x, y \in [G],$$

$$(2.9.8.12) \quad \sum_{\chi \in \mathfrak{X}(G)} |K_\chi^0(x, y)| \leq \|x\|_{[G]}^{N'} \|y\|_{[G]}^{-N}, \quad x, y \in [G]^1.$$

More generally, for every $N > 0$, there exists $N' > 0$ such that for each continuous semi-norm $\|\cdot\|_{N'}$ on $\mathcal{T}_{N'}([G])$

$$(2.9.8.13) \quad \sum_{\chi \in \mathfrak{X}(G)} \|K_\chi(\cdot, y)\|_{N'} \ll \|y\|_{[G]}^{-N}, \quad y \in [G].$$

Proof. — Obviously, (2.9.8.13) implies (2.9.8.11) and (2.9.8.11) implies (2.9.8.12). Let $N > 0$ and choose $L > N$. By Proposition 2.9.3.1.4 and the uniform boundedness principle, there exists $N' > L$ such that for every continuous semi-norm $\|\cdot\|_{N'}$ on $\mathcal{T}_{N'}([G])$ there exists a continuous semi-norm $\|\cdot\|_L$ on $\mathcal{T}_L([G])$ satisfying

$$\sum_{\chi \in \mathfrak{X}(G)} \|K_\chi(\cdot, y)\|_{N'} \leq \|K_f(\cdot, y)\|_L$$

for every $y \in [G]$. Therefore, it suffices to show that if L is large enough then

$$\|K_f(\cdot, y)\|_L \ll \|y\|_{[G]}^{-N}, \quad y \in [G].$$

As $R(X)K_f(\cdot, y) = K_{L(X)f}(\cdot, y)$ and by definition of the topology on $\mathcal{T}_L([G])$, it even suffices to prove that for L large enough

$$(2.9.8.14) \quad \sum_{\gamma \in G(F)} \|x^{-1}\gamma y\|^{-L} \ll \|x\|_{[G]}^L \|y\|_{[G]}^{-N}, \quad x, y \in [G].$$

There exists $N_0 > 0$ such that $\sum_{\gamma \in G(F)} \|\gamma\|^{-N_0} < \infty$. Fix such a N_0 . Then, as $\|y\|_{[G]} \leq \|\gamma y\|$ for every $\gamma \in G(F)$ and $y \in G(\mathbb{A})$, we have

$$\sum_{\gamma \in G(F)} \|x^{-1}\gamma y\|^{-2N_0-N} \ll \|x\|^{2N_0+N} \|y\|^{N_0} \|y\|_{[G]}^{-N-N_0} \sum_{\gamma \in G(F)} \|\gamma\|^{-N_0} \ll \|x\|^{2N_0+N} \|y\|^{N_0} \|y\|_{[G]}^{-N-N_0}$$

for $x, y \in G(\mathbb{A})$. Since the left hand side of the above inequality is invariant by left translations of both x and y by $G(F)$, we may replace $\|x\|^{2N_0+N}$ and $\|y\|^{N_0}$ in the right hand side by $\|x\|_{[G]}^{2N_0+N}$ and $\|y\|_{[G]}^{N_0}$ respectively. This gives (2.9.8.14) for $L = 2N_0 + N$ and this ends the proof of the lemma. \square

2.9.9. Let P be standard parabolic subgroup of G and let $M = M_P$. Let $\chi \in \mathfrak{X}(G)$ and

$$\mathcal{A}_{P,\chi}(G) = \oplus_{\pi} \mathcal{A}_{P,\pi}(G)$$

where the sum is over cuspidal representations π of M such that the image of (M, π) by the map $\mathfrak{X}(M) \rightarrow \mathfrak{X}(G)$ is χ . Let $\mathcal{B}_{P,\chi}$ be a K -basis of $\mathcal{A}_{P,\chi}(G)$ that is the union $\cup_{\pi} \mathcal{B}_{P,\pi}$ over π as above of K -bases of $\mathcal{A}_{P,\pi}(G)$ (see § 2.8.3). In the same way we define $\mathcal{B}_{P,\chi,\tau} = \cup_{\pi} \mathcal{B}_{P,\pi,\tau}$ for any $\tau \in \hat{K}$.

By a slight variant of [Art78] §4 and [Art80] section 3, we have the following lemma.

Lemma 2.9.9.1. — *(Arthur) Let $C \subset G(\mathbb{A}_f)$ be a compact subset and let $K_0 \subset K^\infty$ be a normal open compact subgroup. There exists a continuous semi-norm $\|\cdot\|$ on $\mathcal{S}(G(\mathbb{A}), C, K_0)$ and an integer N such that for all $X, Y \in \mathcal{U}(\mathfrak{g}_{\mathbb{C}})$, all $x, y \in G(\mathbb{A})^1$ and all $f \in \mathcal{S}(G(\mathbb{A}), C, K_0)$ we have*

$$\begin{aligned} \sum_{\chi \in \mathfrak{X}(G)} \sum_{P_0 \subset P} |\mathcal{P}(M_P)|^{-1} \int_{\mathfrak{ia}_P^{G,*}} \sum_{\tau \in \hat{K}} \left| \sum_{\varphi \in \mathcal{B}_{P,\chi,\tau}} (R(X)E)(x, I_P(\lambda, f)\varphi, \lambda) \overline{R(Y)E(y, \varphi, \lambda)} \right| d\lambda \\ \leq \|f\| \|x\|_{G(\mathbb{A})}^N \|y\|_{G(\mathbb{A})}^N. \end{aligned}$$

Moreover for all $x, y \in G(\mathbb{A})$ and all $\chi \in \mathfrak{X}(G)$ we have

$$K_{\chi}^0(x, y) = \sum_{P_0 \subset P} |\mathcal{P}(M_P)|^{-1} \int_{\mathfrak{ia}_P^{G,*}} \sum_{\varphi \in \mathcal{B}_{P,\chi}} E(x, I_P(\lambda, f)\varphi, \lambda) \overline{E(y, \varphi, \lambda)} d\lambda.$$

Proof. — One point is to remove the K_∞ -finiteness assumption in lemma 4.4 of [Art78]. This can be done by approximation by K_∞ -finite functions and this also enables us to put the sum over τ outside the absolute value. The other point is to remove the hypothesis about the compactness of the support of f . However the key point is in fact lemma 4.3 of [Art78] which can be replaced by Lemma 2.9.8.1. \square

3 The spectral expansion of the Jacquet-Rallis trace formula for general linear groups

This chapter has two goals. The first, accomplished in Theorem 3.4.3.2, is to extend the coarse spectral expansion $I = \sum_{\chi \in \mathfrak{X}(G)} I_\chi$ of the Jacquet-Rallis trace formula for linear groups G (as proved in [Zyd20]) to the Schwartz space. The second, given in Theorem 3.3.7.1, is to provide spectral expressions more suitable for explicit calculations.

3.1 Notations

3.1.1. Let E/F be a quadratic extension of number fields. For convenience, we will fix $\tau \in F^\times$ such that $E = F[\sqrt{\tau}]$. Let η be the quadratic character of \mathbb{A}_F^\times attached to E/F . Let $n \geq 1$ be an integer. Let $G'_n = GL_{n,F}$ be the algebraic group of F -linear automorphisms of F^n . Let $G_n = \text{Res}_{E/F}(G'_n \times_F E)$ be the F -group obtained by restriction of scalars from the algebraic group $GL_{n,E}$ of E -linear automorphisms of E^n . We denote by c the Galois involution. We have a natural inclusion $G'_n \subset G_n$ which induces an inclusion $A_{G'_n} \subset A_{G_n}$ which is in fact an equality. The restriction map $X^*(G_n) \rightarrow X^*(G'_n)$ gives an isomorphism $\mathfrak{a}_{G_n}^* \simeq \mathfrak{a}_{G'_n}^*$.

3.1.2. Let (B'_n, T'_n) be a pair where B'_n is the Borel subgroup G'_n of upper triangular matrices and T'_n is the maximal torus of G'_n of diagonal matrices. Let (B_n, T_n) be the pair deduced from (B'_n, T'_n) by extension of scalars to E and restriction to F : it is a pair of a minimal parabolic subgroup of G_n and its Levi factor.

Let $K_n \subset G_n(\mathbb{A})$ and $K'_n = K_n \cap G'_n(\mathbb{A}) \subset G'_n(\mathbb{A})$ be the “standard” maximal compact subgroups. Notice that we have $K'_n \subset K_n$.

3.1.3. The map $P' \mapsto P = \text{Res}_{E/F}(P' \times_F E)$ induces a bijection between the sets of standard parabolic subgroups of G'_n and G_n whose inverse bijection is given by

$$P \mapsto P' = P \cap G'_n.$$

Let P be a standard parabolic subgroup of G_n . The restriction map $X^*(P) \rightarrow X^*(P')$ identifies $X^*(P)$ with a subgroup of $X(P')$ of index $2^{\dim(\mathfrak{a}_P)}$. It also induces an isomorphism $\mathfrak{a}_{P'} \rightarrow \mathfrak{a}_P$ which fits into the commutative diagram:

$$\begin{array}{ccc} G'_n(\mathbb{A}) & \xrightarrow{H_{P'}} & \mathfrak{a}_{P'} \\ \downarrow & & \downarrow \\ G'_n(\mathbb{A}) & \xrightarrow{H_P} & \mathfrak{a}_P \end{array}$$

For any standard parabolic subgroups $P \subset Q$, the restriction of the function τ_P^Q to $\mathfrak{a}_{P'}$ coincides with the function $\tau_{P'}^{Q'}$. However we have for all $x \in G'(\mathbb{A})$

$$\langle \rho_P^Q, H_P(x) \rangle = 2 \langle \rho_{P'}^{Q'}, H_{P'}(x) \rangle.$$

Remark 3.1.3.1. — The map $\mathfrak{a}_{P'} \rightarrow \mathfrak{a}_P$ does not preserve Haar measures. In fact, the pull-back on $\mathfrak{a}_{P'}$ of the Haar measure on \mathfrak{a}_P is $2^{\dim(\mathfrak{a}_P)}$ times the Haar measure on $\mathfrak{a}_{P'}$. In particular, although the groups A_P^∞ and $A_{P'}^\infty$ can be canonically identified, the Haar measure on A_P^∞ is $2^{\dim(\mathfrak{a}_P)}$ times the Haar measure on $A_{P'}^\infty$.

3.1.4. We shall use the natural embeddings $G'_n \subset G'_{n+1}$ and $G_n \subset G_{n+1}$ where the smaller group is identified with the subgroup of the bigger one that fixes e_{n+1} and preserves the space generated by (e_1, \dots, e_n) where (e_1, \dots, e_{n+1}) denotes the canonical basis of F^{n+1} .

3.1.5. Let $G = G_n \times G_{n+1}$ and $G' = G'_n \times G'_{n+1}$. Thus G' is an F -subgroup of G . Let

$$\iota : G_n \hookrightarrow G_n \times G_{n+1}$$

be the diagonal embedding. Let H be the image of ι (so H is isomorphic to G_n).

3.1.6. Let $K = K_n \times K_{n+1}$: it is a maximal compact subgroup of $G(\mathbb{A})$. We define pairs $(P_0, M_0) = (B_n \times B_{n+1}, T_n \times T_{n+1})$ and $(P'_0, M'_0) = (B'_n \times B'_{n+1}, T'_n \times T'_{n+1})$ of minimal parabolic F -subgroups of G and G' with their Levi components. As in §3.1.3, we have a bijection denoted $P \mapsto P'$ between the sets of standard parabolic subgroups of G and G' .

3.1.7. In general, for a subgroup P (usually a parabolic subgroup) of G_n , G_{n+1} or G , we write P' for the intersection of P with G'_n , G'_{n+1} or G' respectively.

3.1.8. Let \det_n (resp. \det_{n+1}) be the morphism $G' \mapsto \mathbb{G}_{m,F}$ given by the determinant on the first (resp. second) component. Let $\eta_{G'}$ be the character $G'(\mathbb{A}) \rightarrow \{\pm 1\}$ given by

$$\eta_{G'}(h) = \eta(\det_n(h))^{n+1} \eta(\det_{n+1}(h))^n.$$

3.1.9. We set $\mathfrak{a}_{n+1} = \mathfrak{a}_{B_{n+1}}$ and $\mathfrak{a}_{n+1}^+ = \mathfrak{a}_{B_{n+1}}^+$ (see §2.2.9).

3.1.10. We let \mathcal{F}_{RS} to be the set of F -parabolic subgroups of G of the form $P = P_n \times P_{n+1}$ where P_n is a standard parabolic subgroup of G_n and P_{n+1} is a semi-standard parabolic subgroup of G_{n+1} such that $P_{n+1} \cap G_n = P_n$ (using the embedding $G_n \hookrightarrow G_{n+1}$).

3.1.11. For $P, Q \in \mathcal{F}_{RS}$ such that $P \subset Q$ we let $\epsilon_P^Q = (-1)^{\dim(\mathfrak{a}_{P_{n+1}}^{Q_{n+1}})}$.

3.2 The coarse spectral expansion for Schwartz functions

3.2.1. Let $f \in \mathcal{S}(G(\mathbb{A}))$ be a Schwartz test function (see §2.5.2).

3.2.2. Let P be a parabolic subgroup of G . The right convolution by f on $L^2([G]_P)$ gives an integral operator whose kernel is denoted by $K_{P,f}$. Let $\chi \in \mathfrak{X}(G)$. Replacing $L^2([G]_P)$ by its closed subspace $L_\chi^2([G]_P)$ (see (2.9.2.1)), we get a kernel denoted by $K_{P,\chi,f}$. If $P = G$, we omit the subscript P . Most of the time, we will also omit the subscript f . We have $K_P = \sum_{\chi \in \mathfrak{X}(G)} K_{P,\chi}$.

3.2.3. For $x \in H(\mathbb{A})$, $y = (y_n, y_{n+1}) \in G'_n(\mathbb{A}) \times G'_{n+1}(\mathbb{A})$, $\chi \in \mathfrak{X}(G)$ and $T \in \mathfrak{a}_{n+1}$ we set

$$(3.2.3.1) \quad K_\chi^T(x, y) = \sum_{P \in \mathcal{F}_{RS}} \epsilon_P^G \sum_{\gamma \in (P \cap H)(F) \backslash H(F)} \sum_{\delta \in P'(F) \backslash G'(F)} \hat{\tau}_{P_{n+1}}(H_{P_{n+1}}(\delta_n y_n) - T_{P_{n+1}}) K_{P,\chi}(\gamma x, \delta y).$$

where

- we write $\delta = (\delta_n, \delta_{n+1})$ and $y = (y_n, y_{n+1})$ according to the decomposition $G' = G'_n \times G'_{n+1}$;
- in the notation $H_{P_{n+1}}(\delta_n y_n)$, we consider $\delta_n y_n$ as an element of $G'_{n+1}(\mathbb{A})$ (via the embedding $G'_n \hookrightarrow G'_{n+1}$);
- $T_{P_{n+1}}$ is defined as in §2.2.12;

and the rest of notation is explained in §3.1.

Remark 3.2.3.1. — This is the kernel used in [Zyd20] for compactly supported functions. Since we are considering a Schwartz function f , the sums over γ and δ are not finite. However, the component δ_n may be taken in a *finite* set depending on y_n (see [Art78] Lemma 5.1). We can then easily show that the sums are absolutely convergent using majorization of Lemma 2.9.8.1.

3.2.4.

Theorem 3.2.4.1. — Let $T \in \mathfrak{a}_{n+1}^+$.

1. We have

$$\sum_{\chi \in \mathfrak{X}(G)} \int_{[H]} \int_{[G']} |K_{f,\chi}^T(h, g')| dg' dh < \infty$$

2. As a function of T , the integral

$$(3.2.4.2) \quad I_\chi^T(f) = \int_{[H]} \int_{[G']} K_{f,\chi}^T(h, g') \eta_{G'}(g') dg' dh$$

coincides with a polynomial-exponential function in T whose purely polynomial part is constant and denoted by $I_\chi(f)$.

3. The distributions I_χ are continuous, left $H(\mathbb{A})$ -invariant and right $(G'(\mathbb{A}), \eta_{G'})$ -equivariant.

4. The sum

$$(3.2.4.3) \quad I(f) = \sum_{\chi} I_\chi(f)$$

is absolutely convergent and defines a continuous distribution I .

Remark 3.2.4.2. — The last statement is the coarse spectral expansion of the Jacquet-Rallis trace formula for G .

Proof. — All the statements but the continuity and the extension to Schwartz functions are proved in [Zyd20, Theorems 3.1 and 3.9] for compactly supported functions. Assuming extension to Schwartz case, continuity is the result of the explicit formula of [Zyd20], Theorem 3.7 (which also holds for Schwartz functions). As for absolute convergence in the Schwartz case, we state and prove a twin theorem below (see Theorem 3.4.3.2) whose proof can easily be adapted to the current theorem, so we will not repeat the arguments here. \square

3.3 Auxiliary expressions for I_χ

3.3.1. The goal of this section is to provide new expressions for the distribution I_χ defined in Theorem 3.2.4.1. In this paper, we will use these expressions to explicitly compute I_χ . The main results are subsumed in Theorem 3.5.1.1. Before giving its statement, we have to explain the main objects. For this, we fix $f \in \mathcal{S}(G(\mathbb{A}))$ and we simply denote by K_χ the kernel $K_{f,\chi}$.

3.3.2. The Ichino-Yamana truncation operator. — Let $T \in \mathfrak{a}_{n+1}$. In [IY15], Ichino-Yamana defined a truncation operator which transforms functions of moderate growth on $[G_{n+1}]$ into rapidly decreasing functions on $[G_n]$. By applying it to the *right* component of $[G] = [G_n] \times [G_{n+1}]$, we get a truncation operator which we denote by Λ_r^T (the subscript r is for right). It associates to any function φ on $[G]$ the function on $[H]$ defined by the following formula: for any $h \in [H]$:

$$(3.3.2.1) \quad (\Lambda_r^T \varphi)(h) = \sum_{P \in \mathcal{F}_{RS}} \epsilon_P^G \sum_{\delta \in (P \cap H)(F) \backslash H(F)} \hat{\tau}_{P_{n+1}}(H_{P_{n+1}}(\delta h) - T_{P_{n+1}}) \varphi_{G_n \times P_{n+1}}(\delta h)$$

where we follow notations of §3.2.3. Note that in the expression $H_{P_{n+1}}(\delta h)$, we view δh as an element of $G_{n+1}(\mathbb{A})$ by the composition $H \hookrightarrow G \rightarrow G_{n+1}$ where the second map is the second projection. We denote by $\varphi_{G_n \times P_{n+1}}$ the constant term of φ along $G_n \times P_{n+1}$.

For properties of Λ_r^T we shall refer to [IY15]. However for our purposes it is convenient to state the following proposition.

Proposition 3.3.2.1. — *For any integers N and N' , any open compact subset $K_0 \subset G(\mathbb{A}_f)$, there is an integer $r \geq 0$ and a finite family $(X_i)_{i \in I}$ of elements of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ of degree $\leq r$ such that for any $\varphi \in C^r(G(F) \backslash G(\mathbb{A})/K_0)$ we have for all $h \in [H]$*

$$(\Lambda_r^T \varphi)(h) \leq \|h\|_{[H]}^{-N} \sum_{i \in I} \left(\sup_{x \in G(\mathbb{A})} \|x\|_{[G]}^{-N'} |(R(X_i)\varphi)(x)| \right)$$

Proof. — The result, a variant of Arthur's Lemma 1.4 of [Art80], is proven in [IY15], Lemma 2.4. \square

3.3.3. Convergence of a first integral. — It is given by the following proposition.

Proposition 3.3.3.1. — *Let $\chi \in \mathfrak{X}(G)$. The integral*

$$(3.3.3.2) \quad \int_{[H]} \int_{[G']} \Lambda_r^T K_\chi(x, y) \eta_{G'}(y) dx dy$$

is absolutely convergent.

Proof. — We can easily deduce from Lemma 2.9.8.1 that for all $r_1 \geq 0$, there exists a continuous semi-norm $\|\cdot\|$ on $\mathcal{S}(G(\mathbb{A}))$ and an integer $N \in \mathbb{N}$ such that for all $x \in [G]$, $y \in G(\mathbb{A})^1$, $a \in A_G^\infty$, $f \in \mathcal{S}(G(\mathbb{A}), C, K_0)$ we have

$$(3.3.3.3) \quad |K_{f,\chi}(x, ay)| \leq \|f\| \|a\|_{G(\mathbb{A})}^{-r_1} \|x\|_{[G]}^N.$$

In particular, if we restrict ourselves to $x \in [H]$, we have with the same hypothesis the existence of N and $\|\cdot\|$ such that

$$(3.3.3.4) \quad \|x\|_{[H]}^{-N} |K_{f,\chi}(x, ay)| \leq \|f\| \|a\|_{G(\mathbb{A})}^{-r_1}.$$

for all $x \in [H]$, $y \in G(\mathbb{A})^1$, $a \in A_G^\infty$ and $f \in \mathcal{S}(G(\mathbb{A}), C, K_0)$.

The right derivatives in the first variable of the kernel $K_\chi(x, y)$ can be expressed in terms of the kernel K_χ associated to left derivatives of f . Thus, taking into account Proposition 3.3.2.1, we see that for any $r_2 \geq 0$ there exists a continuous semi-norm $\|\cdot\|$ on $\mathcal{S}(G(\mathbb{A}), C, K_0)$ such that for $f \in \mathcal{S}(G(\mathbb{A}), C, K_0)$ we have

$$|\Lambda_r^T K_\chi(x, ay)| \leq \|f\| \|a\|_{G(\mathbb{A})}^{-r_1} \|x\|_{[H]}^{-r_2}$$

for all $a \in A_G^\infty$, $x \in [H]$ and $y \in G(\mathbb{A})^1$. The convergence is then obvious. \square

3.3.4. Arthur function $F^{G_{n+1}}(\cdot, T)$. — For $T \in \mathfrak{a}_{n+1}$ we shall use Arthur function $F^{G_{n+1}}(\cdot, T)$ (see [Art78] §6). Recall that this is the characteristic function of the set of $x \in G_{n+1}(\mathbb{A})$ for which there exists a $\delta \in G_{n+1}(F)$ such that $\delta x \in \mathfrak{s}_{G_{n+1}}$ (see § 2.2.13) and $\langle \varpi, H_0(\delta x) - T \rangle \leq 0$ for all $\varpi \in \widehat{\Delta}_{B_{n+1}}$. Recall also that $F^{G_{n+1}}(\cdot, T)$ descends to characteristic function of a compact subset of $Z_{n+1}(\mathbb{A})G_{n+1}(F) \backslash G_{n+1}(\mathbb{A})$.

3.3.5. Two other convergent integrals. —

Proposition 3.3.5.1. — *The following integrals are absolutely convergent:*

$$(3.3.5.5) \quad \int_{[H]} \int_{[G']} F^{G_{n+1}}(x, T) K_\chi(x, y) \eta_{G'}(y) dy dx$$

$$(3.3.5.6) \quad \int_{[H]} \int_{[G']} K_\chi(x, y) F^{G_{n+1}}(y_n, T) \eta_{G'}(y) dy dx$$

where $y = (y_n, y_{n+1}) \in G'_n(\mathbb{A}) \times G'_{n+1}(\mathbb{A})$.

Proof. — The convergence of the integral (3.3.5.5) is proved in the same way as in the proof of Proposition 3.3.3.1. The only point to observe is that the restriction of $F^{G_{n+1}}(\cdot, T)$ to $[H]$ is compactly supported.

The convergence of the integral (3.3.5.6) is a consequence of two facts: first the restriction of $F^{G_{n+1}}(\cdot, T)$ to $[G_n]$ is compactly supported; second for every N', N'' there exist $N > 0$ and a continuous semi-norm on $\mathcal{S}(G(\mathbb{A}))$ such that

$$|K_\chi(x, y)| \leq \|f\| \|y_n\|_{[G_n]}^N \|y_{n+1}\|_{[G_{n+1}]}^{-N'} \|x\|_{[H]}^{-N''}.$$

This majorization can be proved as in the proof of Lemma 2.9.8.1. It suffices to prove the same result for the whole kernel (without subscript χ). We can assume that $f = f_n \otimes f_{n+1}$ is a product. The kernel itself is then a product. We can use Lemma 2.9.8.1 assertion 1 to bound $K_{f_{n+1}}(x, y_{n+1})$ and get the negative power of $\|y_{n+1}\|_{[G_{n+1}]}$. Then we can bound $K_{f_n}(x, y_n)$ to get the negative power of $\|x\|_{[H]}$. \square

3.3.6. We say a functions p and q on \mathfrak{a}_{n+1} are asymptotically equal if for all $\varepsilon > 0$ and $m \geq 0$ there exists $c \geq 0$ such that for all $T \in \mathfrak{a}_{n+1}$ such that $\langle \alpha, T \rangle > \varepsilon \|T\|$ for all $\alpha \in \Delta_{B_{n+1}}$ we have

$$|p(T) - q(T)| \leq ce^{-m\|T\|}.$$

3.3.7. We can now state the main theorem of the section.

Theorem 3.3.7.1. — *Let $\chi \in \mathfrak{X}(G)$. Each of the three expressions (3.3.3.2) (see Proposition 3.3.3.1), (3.3.5.5) and (3.3.5.6) (see Proposition 3.3.5.1) is asymptotically equal (in the sense of §3.3.6) to a polynomial-exponential function of T whose purely polynomial term is constant and equal to $I_\chi(f)$.*

Proof. — The theorem is a simple combination of Theorem 3.2.4.1 above and Theorems 3.4.3.2 and 3.5.1.1 below. \square

3.4 Convergence of a truncated kernel

3.4.1. In this section, we give a first step in the proof of Theorem 3.3.7.1: we define a new truncated kernel and we prove that the integral over $[H] \times [G']$ of this kernel is absolutely convergent.

3.4.2. We fix $f \in \mathcal{S}(G(\mathbb{A}))$, $\chi \in \mathfrak{X}(G)$ and $T \in \mathfrak{a}_{n+1}^+$.

3.4.3. A new truncated kernel. — For $x \in [H]$ and $y \in [G']$ we set

$$\kappa_\chi^T(x, y) = \sum_{P \in \mathcal{F}_{RS}} \epsilon_P^G \sum_{\gamma \in (P \cap H)(F) \backslash H(F)} \sum_{\delta \in P'(F) \backslash G'(F)} \hat{\tau}_{P_{n+1}}(H_{P_{n+1}}(\gamma x) - T_{P_{n+1}}) K_{P, \chi}(\gamma x, \delta y).$$

The notations are those of §3.1. The expression $H_{P_{n+1}}(\gamma x)$ is interpreted as in the comments following (3.3.2.1).

Remark 3.4.3.1. — This is a version of the truncated kernel K_χ^T defined in §3.2.3. We will consider the connection between the two in Theorem 3.4.3.2. The sum over γ is finite whereas the sum over δ is convergent but not finite (see remark 3.2.3.1).

The following theorem is the main result of the section.

Theorem 3.4.3.2. —

1. We have

$$\sum_{\chi \in \mathfrak{X}(G)} \int_{[H]} \int_{[G']} |\kappa_\chi^T(x, y)| dy dx < \infty.$$

2. The integral

$$(3.4.3.1) \quad i_\chi^T(f) := \int_{[H]} \int_{[G']} \kappa_\chi^T(x, y) \eta_{G'}(y) dy dx$$

coincides with a polynomial-exponential function in T whose purely polynomial part is constant and denoted by $i_\chi(f)$.

3. The distribution i_χ is continuous. Moreover we have $i_\chi = I_\chi$ where the right-hand side is defined in Theorem 3.2.4.1.

The proof of assertion 1 of Theorem 3.4.3.2 will be given in §3.4.11 whereas the proof of assertions 2 and 3 will be given in §3.4.12. Before that, we must introduce additional notations and lemmas.

3.4.4. Until §3.4.11, we assume that G is a general reductive group with the notations of section 2.2. In particular, a maximal F -split torus A_0 is fixed as well as a minimal parabolic subgroup P_0 containing it.

3.4.5. Recall that we have defined $\mathfrak{a}_P^{*,+}$ in §2.2.9.

3.4.6. For semi-standard parabolic subgroups P, Q of G , such that $P \subset Q$ we have the function σ_P^Q as defined in §6 of [Art78]. It is a characteristic function of a region in \mathfrak{a}_P . We note the following formula satisfied by it (see proof of Theorem 7.1 in [Art78])

$$(3.4.6.2) \quad \tau_P^Q \hat{\tau}_Q = \sum_{R \supset Q} \sigma_P^R.$$

More generally, if $P \subset Q \subset R$ are semi-standard parabolic subgroups of G , we note $\sigma_P^{Q,R} := \sigma_{P \cap M_R}^{Q \cap M_R}$ the σ function with respect to the group M_R and its parabolic subgroups $P \cap M_R$ and $Q \cap M_R$.

Lemma 3.4.6.1. — *Let P, Q, R be standard parabolic subgroups of G with $P \subset Q \subset R$. Let $\alpha \in \Delta_P \setminus \Delta_P^R$. Suppose $H \in \mathfrak{a}_P$ satisfies $\sigma_P^{Q,R}(H) = 1$. Then, if we denote H_R the projection of H onto \mathfrak{a}_R , we have $\langle \alpha, H_R \rangle \geq \langle \alpha, H \rangle$.*

Proof. — Using (3.4.6.2) we have $\hat{\tau}_P^R(H) = \hat{\tau}_P^R(H - H_R) = 1$. This means that $H - H_R$ has positive coefficients in the coroot basis $\Delta_P^{R,\vee}$ of \mathfrak{a}_P^R . The result follows from the known fact that distinct coroots in Δ_0^\vee form obtuse angles and so do distinct elements of Δ_P^\vee (c.f. [LW13] Lemme 1.2.4). \square

3.4.7. For a standard parabolic subgroup P of G we fix representatives for the cosets $W^P \backslash W$ and W/W^P as follows:

$$W^P \backslash W := \{w \in W \mid w^{-1}\alpha > 0 \ \forall \alpha \in \Delta_0^P\}, \quad W/W^P := \{w \in W \mid w\alpha > 0 \ \forall \alpha \in \Delta_0^P\},$$

where $\beta > 0$, for a root β of (G, A_0) , means that it's a sum of elements of Δ_0 .

3.4.8. Norms

We use the definition of 2.4 to define the norm on $[G]_P$. As we will only use norms on automorphic quotients, we denote this norm simply by $\|\cdot\|$. We use the same symbol for a fixed W^G -invariant norm on \mathfrak{a}_0 .

3.4.9. Bound on Eisenstein Series. —

Lemma 3.4.9.1. — *Let $w \in W/W^P$ and $\lambda \in \mathfrak{a}_P^{*,+}$. For all $\varepsilon > 0$, there is an $N \geq 0$, independent of λ , such that we have*

$$\sum_{\delta \in (P \cap w P_0 w^{-1})(F) \backslash P_0(F)} e^{\langle \lambda + (2+\varepsilon)\rho_P, H_P(w^{-1}\delta x) \rangle} \ll \|x\|^N e^{\langle w\lambda, H_0(x) \rangle}.$$

for any x in the Siegel domain \mathfrak{s}_P (see §2.2.13).

Proof. — We have for $\delta \in P_0(F)$

$$\langle \lambda, H_P(w^{-1}\delta x) \rangle = \langle \lambda, H_0(w^{-1}\delta x) \rangle = \langle w\lambda, H_0(x) \rangle + \langle \lambda, H_0(w^{-1}n) \rangle$$

for some $n \in N_0(\mathbb{A})$ depending on x and δ . Since $\lambda \in \mathfrak{a}_P^{+,*}$ we have, by [LW13] Lemma 3.3.1, $\langle \lambda, H_0(w^{-1}n) \rangle \leq c$ for some constant c depending on λ but independent of n . We have then

$$\sum_{\delta \in (P \cap w P_0 w^{-1})(F) \setminus P_0(F)} e^{\langle \lambda + (2+\varepsilon)\rho_P, H_P(w^{-1}\delta x) \rangle} \ll \sum_{\delta \in (P \cap w P_0 w^{-1})(F) \setminus P_0(F)} e^{\langle (2+\varepsilon)\rho_P, H_P(w^{-1}\delta x) \rangle} e^{\langle w\lambda, H_0(x) \rangle}$$

and the result follows by moderate growth of Eisenstein series. \square

3.4.10. Auxiliary characteristic functions. — We introduce the function $\bar{\tau}_P$ as the characteristic function of $H \in \mathfrak{a}_P$ such that $\langle \alpha, H \rangle \leq 0$ for all $\alpha \in \Delta_P$. By Langlands Combinatorial Lemma ([LW13], Proposition 1.7.2) we have

$$(3.4.10.3) \quad \sum_{Q \supset P} \hat{\tau}_P^Q \bar{\tau}_Q = 1.$$

3.4.11. Proof of assertion 1 of Theorem 3.4.3.2. — Unless otherwise stated, all sums of the type Σ_P or $\Sigma_{P \subset Q}$ are over elements of \mathcal{F}_{RS} .

In §2.4 of [Zyd20] the operator $\Lambda_d^{T,P}$ is defined for all $P \in \mathcal{F}_{RS}$. For a function $\phi : P(F) \backslash G(\mathbb{A}) \rightarrow \mathbb{C}$, we have

$$(3.4.11.4) \quad \Lambda_d^{T,P} \phi(x) = \sum_{P \supset Q} \epsilon_Q^P \sum_{\delta \in (Q \cap H)(F) \setminus (P \cap H)(F)} \hat{\tau}_{Q_{n+1}}^{P_{n+1}}(H_{Q_{n+1}}(\delta x) - T_{Q_{n+1}}) \phi_Q(\delta x), \quad x \in (P \cap H)(F) \setminus H(\mathbb{A}).$$

Note that if $P = G$, the operator $\Lambda_d^{T,G}$ is close but not exactly equal to the operator Λ_r^T defined in (3.3.2.1). Indeed in the former we take the constant term along P whereas in the latter we take it along $G_n \times P_{n+1}$.

Using the inversion formula of Lemme 2.7 [Zyd20] together with the formula (3.4.6.2) we obtain that the integral is bounded by the sum over $P, Q \in \mathcal{F}_{RS}$ of

$$(3.4.11.5) \quad \sum_{\chi} \int_{(H \cap P)(F) \setminus H(\mathbb{A})} \int_{[G']} \sigma_{P_{n+1}}^{Q_{n+1}}(H_{P_{n+1}}(x) - T_{P_{n+1}}) |\Lambda_d^{T,P} \left(\sum_{P \subset \tilde{P} \subset Q} \epsilon_P^{P_1} \sum_{\delta' \in \tilde{P}'(F) \setminus G'(F)} K_{\tilde{P},\chi}(x, \delta' y) \right)| dy dx$$

where $\tilde{P} \in \mathcal{F}_{RS}$ in the alternating sum and $\Lambda_d^{T,P}$ is applied with respect to x . We consider P and Q fixed from now on. Without loss of generality we can assume they are standard in G .

For any standard parabolic subgroup S of G we have the mixed truncation operator $\Lambda_m^{T,S}$ of [JLR99]

$$\Lambda_m^{T,S} \phi(x) = \sum_{R \subset S} (-1)^{\dim(\mathfrak{a}_R^S)} \sum_{\delta \in R'(F) \setminus S'(F)} \hat{\tau}_R^S(H_R(\delta x) - T) \phi_R(\delta x), \quad x \in R'(F) \setminus S'(\mathbb{A}).$$

Using (3.4.10.3), the inversion formula (§6 (19) in [JLR99]) and (3.4.6.2), we bound the expression (3.4.11.5) by a sum over $R \subset S \subset \tilde{S}$, all standard parabolic subgroups of G , of

$$(3.4.11.6) \quad \sum_{\chi} \int_{(H \cap P)(F) \setminus H(\mathbb{A})} \int_{(G' \cap R)(F) \setminus G'(\mathbb{A})} \sigma_{P_{n+1}}^{Q_{n+1}}(H_{P_{n+1}}(x) - T) \bar{\tau}_{\tilde{S}}(H_{\tilde{S}}(y) - T') \sigma_R^{S,\tilde{S}}(H_R(y) - T') \\ |\Lambda_d^{T,P} \Lambda_m^{T',R} \left(\sum_{P^+ \subset \tilde{P} \subset Q^+} \epsilon_P^{\tilde{P}} K_{\tilde{P},\chi}(x, y) \right)| dy dx$$

where $\Lambda_m^{T',R}$ is applied with respect to y , P^+ is the smallest element of \mathcal{F}_{RS} containing R and P , Q^+ is the largest element of \mathcal{F}_{RS} contained in S and Q , the sum runs over $\tilde{P} \in \mathcal{F}_{RS}$, $T' \in \mathfrak{a}_{n+1}^+$

is any parameter, and we write T instead of $T_{P_{n+1}}$ because P is assumed standard so there is no difference.

Let $P_H = H \cap P$, it is a standard parabolic subgroup of H . Let $Z_P^\infty = A_P^\infty \cap A_{P_H}^\infty$. We also note $\mathfrak{z}_P = \mathfrak{a}_P \cap \mathfrak{a}_{P_H}$. Let $M(\mathbb{A})^{P,1}$ be the kernel of the composition of $H_{P_H} : M_{P_H}(\mathbb{A}) \rightarrow \mathfrak{a}_{P_H}$ with the orthogonal projection $\mathfrak{a}_{P_H} \rightarrow \mathfrak{z}_P$.

We assume that f is bi- K -invariant - it makes the subsequent computations and notation clearer. It is not a serious restriction, one should deal with the general case as in [Art80] or [Zyd20]. Using the Iwasawa decomposition and Propositions 2.3 and 2.8 of [Zyd20] we bound (3.4.11.7) by

$$\begin{aligned} & \sum_{\chi} \int_{Z_P^\infty} \int_{A_R^\infty} \int_{M_{P_H}(F) \backslash M(\mathbb{A})^{P,1}} \int_{[M_{R'}]^1} e^{\langle -2\rho_{P_H}, H_{P_H}(zm_1) \rangle} e^{\langle -2\rho_{R'}, H_{R'}(am_2) \rangle} \sigma_{P_{n+1}}^{Q_{n+1}}(H_{P_{n+1}}(zm_1) - T) \\ & \quad \bar{\tau}_{\tilde{S}}(H_{\tilde{S}}(a) - T') \sigma_R^{S, \tilde{S}}(H_R(a) - T') \|m_1\|^{-r_1} \|m_2\|^{-r_2} \sup_{m'_1 \in M_{P_H}(F) \backslash M(\mathbb{A})^{P,1}} \sup_{m'_2 \in M_{R'}(F) \backslash M_{R'}(\mathbb{A})^1} \\ & \quad \left| \int_{[N_P]} \int_{[N_R]} \left(\sum_{P^+ \subset \tilde{P} \subset Q^+} \epsilon_{\tilde{P}}^{\tilde{P}} K_{\tilde{P}, \chi}(n_1 zm'_1, n_2 am'_2) \right) dn_2 dn_1 \|m'_1\|^{-r'_1} \|m'_2\|^{-r'_2} dm_2 dm_1 dadz \right| \end{aligned}$$

for any r_1, r'_1, r_2, r'_2 . Here, we should replace the kernel K by a finite sum of kernels K_i , but to make the notation simpler we will ignore this detail. Making a few changes of variables we bound the above expression by

$$\begin{aligned} & (3.4.11.7) \\ & \sum_{\chi} \int_{Z_P^\infty} \int_{A_R^\infty} \int_{M_{P_H}(F) \backslash M(\mathbb{A})^{P,1}} \int_{[M_{R'}]^1} \sigma_{P_{n+1}}^{Q_{n+1}}(H_{P_{n+1}}(z) - T) \bar{\tau}_{\tilde{S}}(H_{\tilde{S}}(a) - T') \sigma_R^{S, \tilde{S}}(H_R(a) - T') \\ & \quad \|z\|^{r_0} \|a\|^{r_0} \|m_1\|^{-r_1} \|m_2\|^{-r_2} \sup_{m'_1 \in M_{P_H}(F) \backslash M(\mathbb{A})^{P,1}} \sup_{m'_2 \in M_{R'}(F) \backslash M_{R'}(\mathbb{A})^1} \\ & \quad \|m'_1\|^{-r'_1} \|m'_2\|^{-r'_2} \left| \int_{[N_P]} \int_{[N_R]} \left(\sum_{P^+ \subset \tilde{P} \subset Q^+} \epsilon_{\tilde{P}}^{\tilde{P}} K_{\tilde{P}, \chi}(n_1 zm'_1, n_2 am'_2) \right) dn_2 dn_1 \right| dm_2 dm_1 dadz \end{aligned}$$

where r_0 is a fixed number and r_1, r_2, r'_1, r'_2 are arbitrarily large.

We set

$$\Psi(z, a, m_1, m_2) = \sum_{\chi} \left| \int_{[N_P]} \int_{[N_R]} \left(\sum_{P^+ \subset \tilde{P} \subset Q^+} \epsilon_{\tilde{P}}^{\tilde{P}} K_{\tilde{P}, \chi}(n_1 zm_1, n_2 am_2) \right) dn_2 dn_1 \right|.$$

Given that the proof of Theorem 3.4.3.2 assertion 1 has been reduced to proving convergence of the integral (3.4.11.7), the following lemma will conclude the proof of the Theorem.

Lemma 3.4.11.1. — *For all $N \geq 0$, there exists an M such that*

$$\sigma_{P_{n+1}}^{Q_{n+1}}(H_{P_{n+1}}(z) - T) \bar{\tau}_{\tilde{S}}(H_{\tilde{S}}(a) - T') \sigma_R^{S, \tilde{S}}(H_R(a) - T') \Psi(z, a, m_1, m_2) \leq \|z\|^{-N} \|a\|^{-N} \|m_1\|^M \|m_2\|^M.$$

To prove 3.4.11.1 we first need a bound on $\Psi(z, a, m_1, m_2)$. It will be more natural to bound $\Psi(z, a, m_1, m_2)^2$.

Before we proceed let us make some remarks and establish some notation. Recall that $P = P_n \times P_{n+1}$ and $Q = Q_n \times Q_{n+1}$ with $P_{n+1} \cap G_n = P_n$ and $Q_{n+1} \cap G_n = Q_n$. We write $R = R_n \times R_{n+1}$, $S = S_n \times S_{n+1}$, $\tilde{S} = \tilde{S}_n \times \tilde{S}_{n+1}$. All parabolic subgroups are standard in their respective ambient groups.

Recall that B_n and B_{n+1} are fixed Borel subgroups of G_n and G_{n+1} with $B_{n+1} \cap G_n = B_n$. The inclusion $G_n \hookrightarrow G_{n+1}$ induces the inclusion $\Delta_{B_n}^\vee \hookrightarrow \Delta_{B_{n+1}}^\vee$. The latter inclusion induces therefore a natural inclusion $\iota : \widehat{\Delta}_{B_n} \hookrightarrow \widehat{\Delta}_{B_{n+1}}$. We let ϖ^{n+1} be the unique element of $\widehat{\Delta}_{B_{n+1}} \setminus \iota(\widehat{\Delta}_{B_n})$. Its restriction to \mathfrak{a}_{B_n} equals the determinant divided by $n+1$. Note that, since $P_{n+1} \cap G_n = P_n$, the set $\widehat{\Delta}_{P_{n+1}} \setminus \iota(\widehat{\Delta}_{P_n})$ is either empty or consists solely of ϖ^{n+1} .

Having introduced these, we can write $P^+ = P_n^+ \times P_{n+1}^+$ and $Q^+ = Q_n^+ \times Q_{n+1}^+$, both standard parabolic subgroups and elements of \mathcal{F}_{RS} . We have then

$$\widehat{\Delta}_{P_{n+1}^+} = \left(\iota(\widehat{\Delta}_{R_n}) \cup \{\varpi^{n+1}\} \right) \cap \widehat{\Delta}_{R_{n+1}} \cap \widehat{\Delta}_{P_{n+1}}, \quad \widehat{\Delta}_{Q_{n+1}^+} = \iota(\widehat{\Delta}_{S_n}) \cup \widehat{\Delta}_{S_{n+1}} \cup \widehat{\Delta}_{Q_{n+1}}.$$

For $w \in W$ we write $w = (w_n, w_{n+1}) \in W^{G_n} \times W^{G_{n+1}}$. Since W^{G_n} embeds naturally into $W^{G_{n+1}}$ we view w_n as element of $W^{G_{n+1}}$. For any $s \in W^{G_{n+1}}$ let $\widehat{\Delta}_s$ be the set of $\varpi \in \widehat{\Delta}_{B_{n+1}}$ stabilized by s .

Using an easy extension of Lemma 2.3 of [Art80] to Schwartz functions, for $\widetilde{P} \supset P$, we have

$$(3.4.11.8) \quad \int_{[N_P]} K_{\widetilde{P}, \chi}(nx, y) dn = \sum_{w \in W^{\widetilde{P}}/W^P} \sum_{\delta \in (P \cap w P_0 w^{-1})(F) \setminus P_0(F)} K_{P, \chi}(x, w^{-1} \delta y).$$

Let Ω' be the subset of $(w, w') \in W^{Q^+}/W^P \times W^{Q^+}/W^R$ satisfying

$$(3.4.11.9) \quad \widehat{\Delta}_{P_{n+1}^+} \cap \widehat{\Delta}_{w_n} \cap \widehat{\Delta}_{w_{n+1}} = \widehat{\Delta}_{Q_{n+1}^+}, \quad \widehat{\Delta}_{P_{n+1}^+} \cap \widehat{\Delta}_{w'_n} \cap \widehat{\Delta}_{w'_{n+1}} = \widehat{\Delta}_{Q_{n+1}^+}.$$

Using the above notation, applying (3.4.11.8), and its analogue for the group R , taking into consideration cancellations in alternating sums, we see that $\Psi(z, a, m_1, m_2)^2$ is bounded by a sum over $(w, w') \in \Omega'$ of

$$(3.4.11.10) \quad \left(\sum_{\chi} \sum_{\delta \in (P \cap w P_0 w^{-1})(F) \setminus P_0(F)} |K_{P, \chi}(z m_1, w^{-1} \delta a m_2)| \right) \left(\sum_{\chi} \sum_{\delta \in (R \cap w P_0 w^{-1})(F) \setminus P_0(F)} |K_{R, \chi}((w')^{-1} \delta z m_1, a m_2)| \right)$$

here, we drop the remaining compact unipotent integration as it won't affect the bounds.

Fix $(w, w') \in \Omega'$, it's enough to focus on one pair. We have the following natural variant of Lemma 2.9.8.1, equation (2.9.8.11): for all $N > 0$ there exists $N' > 0$ such that for all $m \geq 0$, $x, y \in [M_P]^1$, $z \in A_P^\infty$ and $\lambda \in \mathfrak{a}_P^*$ we have

$$(3.4.11.11) \quad \sum_{\chi} |K_{P, \chi}(x, zy)| \ll \|x\|^{-N} \|y\|^{N'} e^{-m \|H_G(z)\|} e^{\langle \lambda, H_P(z) \rangle}.$$

Using the bound (3.4.11.11) above and Lemma 3.4.9.1 we bound (3.4.11.10) by

$$(3.4.11.12) \quad \|m_1\|^{N'_2 - N_1} \|m_2\|^{N'_1 - N_2} \|a\|^{r_3} \|z\|^{r_3} e^{-r_4 \|H_G(a)\|} e^{\langle w \lambda_1, H_0(a) \rangle - \langle \lambda_1, H_0(z) \rangle} \cdot e^{\langle w' \lambda_2, H_0(z) \rangle - \langle \lambda_2, H_0(a) \rangle}$$

times a power of , where $\lambda_1 \in \mathfrak{a}_P^{*,+}$ and $\lambda_2 \in \mathfrak{a}_R^{*,+}$ are for us to be chosen appropriately and r_3 and r_4 are independent of λ_1 and λ_2 . Moreover, choosing r'_1 and r'_2 in (3.4.11.7) sufficiently large we don't have to worry about the constants N_1, N'_1, N_2, N'_2 (they play a role in the proof of Theorem 3.5.1.1). Note that for $a \in A_R^\infty$ the term $H_G(a)$ is not affected by truncation but thanks to the factor $e^{-r_4 \|H_G(a)\|}$ the integral over A_G^∞ is convergent, we can assume then that $a \in A_R^{G, \infty}$.

Recall the space $\mathfrak{z}_P \subset \mathfrak{a}_0$ which equals the diagonally embedded $\mathfrak{a}_{P_n} \cap \mathfrak{a}_{P_{n+1}}$. Our goal is then, for all $N \geq 0$, to choose $\lambda_1 \in \mathfrak{a}_P^{*,+}$ and $\lambda_2 \in \mathfrak{a}_R^{*,+}$ so that for $Z \in \mathfrak{z}_P$ and $H = (H_1, H_2) \in \mathfrak{a}_{R_n}^{G_n} \times \mathfrak{a}_{R_{n+1}}^{G_{n+1}}$ such that

$$(3.4.11.13) \quad \sigma_{P_{n+1}}^{Q_{n+1}}(Z - T) \sigma_{R_n}^{S_n, \widetilde{S}_n}(H_n - T'_n) \sigma_{R_{n+1}}^{S_{n+1}, \widetilde{S}_{n+1}}(H_{n+1} - T'_{n+1}) \bar{\tau}_{\widetilde{S}_n}(H_n - T'_n) \bar{\tau}_{\widetilde{S}_{n+1}}(H_{n+1} - T'_{n+1}) = 1$$

we have that

$$\langle \lambda_1 - w' \lambda_2, Z \rangle + \langle \lambda_2 - w \lambda_1, H \rangle \gg \|Z\|^N + \|H\|^N.$$

More specifically, since the natural projection of \mathfrak{z}_P onto $\mathfrak{a}_{P_{n+1}}^{G_{n+1}}$ is an isomorphism, using Corollary 6.2 of [Art78], it is enough to show the following lemma.

Lemma 3.4.11.2. — *There exist $\lambda_1 \in \mathfrak{a}_P^{*,+}$ and $\lambda_2 \in \mathfrak{a}_R^{*,+}$ such that under (3.4.11.13) we have*

$$\begin{aligned} & \langle \lambda_1 - w' \lambda_2, Z \rangle + \langle \lambda_2 - w \lambda_1, H \rangle \\ \gg_{T,T'} & \sum_{\alpha \in \Delta_{P_{n+1}}^{Q_{n+1}}} \langle \alpha, Z \rangle + \sum_{\alpha \in \Delta_{R_n}^{S_n}} \langle \alpha, H_n \rangle + \sum_{\alpha \in \Delta_{R_{n+1}}^{S_{n+1}}} \langle \alpha, H_{n+1} \rangle - \sum_{\alpha \in \Delta_{S_n}} \langle \alpha, H_n \rangle - \sum_{\alpha \in \Delta_{S_{n+1}}} \langle \alpha, H_{n+1} \rangle \end{aligned}$$

where by $\gg_{T,T'}$ we mean an inequality up to an additive constant that depends on T and T' .

Indeed, given λ_1, λ_2 as in Lemma 3.4.11.2 above, we can then take their arbitrary multiples ensuring, through a reasoning explained at the end of Theorem 2.2 of [Zyd16], the desired property of Lemma 3.4.11.1.

We focus on proving Lemma 3.4.11.2 from now on. We introduce the following notation that will save some space in what follows. Let

$$\underline{a} = \{a_\varpi\}_{\varpi \in \widehat{\Delta}_{P_n} \sqcup \widehat{\Delta}_{P_{n+1}} \sqcup \widehat{\Delta}_{R_n} \sqcup \widehat{\Delta}_{R_{n+1}}}$$

be a set of numbers, where the sets of weights are treated as disjoint for indexing purposes. Define then for $Z \in \mathfrak{z}_P$ and $H = (H_1, H_2) \in \mathfrak{a}_{R_n}^{G_n} \times \mathfrak{a}_{R_{n+1}}^{G_{n+1}}$

$$\begin{aligned} \lambda(\underline{a}, Z, H_n, H_{n+1}) := & \sum_{\varpi \in \widehat{\Delta}_{P_n}} a_\varpi(\langle \varpi, Z \rangle - \langle w_n \varpi, H_n \rangle) + \sum_{\varpi \in \widehat{\Delta}_{P_{n+1}}} a_\varpi(\langle \varpi, Z \rangle - \langle w_{n+1} \varpi, H_{n+1} \rangle) + \\ & \sum_{\varpi \in \widehat{\Delta}_{R_n}} a_\varpi(\langle \varpi, H_n \rangle - \langle w'_n \varpi, Z \rangle) + \sum_{\varpi \in \widehat{\Delta}_{R_{n+1}}} a_\varpi(\langle \varpi, H_{n+1} \rangle - \langle w'_{n+1} \varpi, Z \rangle). \end{aligned}$$

Lemma 3.4.11.3. — *Let $\alpha_0 \in \Delta_{P_{n+1}}^{Q_{n+1}}$ and let $\varpi_0 \in \widehat{\Delta}_{P_{n+1}}$ be the corresponding weight. There exists a set of positive constants*

$$\underline{a}_{\alpha_0} = \{a_{\alpha_0, \varpi}\}_{\varpi \in \widehat{\Delta}_{P_n} \sqcup \widehat{\Delta}_{P_{n+1}} \sqcup \widehat{\Delta}_{R_n} \sqcup \widehat{\Delta}_{R_{n+1}}}$$

such that, assuming (3.4.11.13), we have

$$\lambda(\underline{a}_{\alpha_0}, Z, H_n, H_{n+1}) \gg_{T,T'} \begin{cases} \langle \alpha_0, Z \rangle - \langle \alpha_0, H_n, \tilde{S}_n \rangle & \text{if } \varpi_0 \in (\widehat{\Delta}_{P_{n+1}} \setminus \widehat{\Delta}_{Q_{n+1}}) \cap \iota(\widehat{\Delta}_{\tilde{S}_n}) \\ \langle \alpha_0, Z \rangle - \langle \alpha_0, H_{n+1}, \tilde{S}_{n+1} \rangle & \text{if } \varpi_0 \in (\widehat{\Delta}_{P_{n+1}} \setminus \widehat{\Delta}_{Q_{n+1}}) \cap \widehat{\Delta}_{\tilde{S}_{n+1}} \\ \langle \alpha_0, Z \rangle & \text{else} \end{cases}$$

where H_n, \tilde{S}_n and H_{n+1}, \tilde{S}_{n+1} are projections of H_n and H_{n+1} onto $\mathfrak{a}_{\tilde{S}_n}$ and $\mathfrak{a}_{\tilde{S}_{n+1}}$ respectively.

Proof. — Let $\widehat{\Delta} = \widehat{\Delta}_{P_{n+1}} \setminus \widehat{\Delta}_{Q_{n+1}}$. Define

$$\begin{aligned} \widehat{\Delta}_1 &= \widehat{\Delta} \cap \iota(\widehat{\Delta}_{\tilde{S}_n}), \quad \widehat{\Delta}_2 = \widehat{\Delta} \cap \widehat{\Delta}_{\tilde{S}_{n+1}}, \quad \widehat{\Delta}_3 = \widehat{\Delta} \cap \iota((\widehat{\Delta}_{S_n} \setminus \widehat{\Delta}_{\tilde{S}_n})), \quad \widehat{\Delta}_4 = \widehat{\Delta} \cap (\widehat{\Delta}_{S_{n+1}} \setminus \widehat{\Delta}_{\tilde{S}_{n+1}}), \\ \widehat{\Delta}_5 &= \widehat{\Delta} \cap \iota(\widehat{\Delta}_{B_n} \setminus \widehat{\Delta}_{R_n}), \quad \widehat{\Delta}_6 = \widehat{\Delta} \cap (\widehat{\Delta}_{B_{n+1}} \setminus \widehat{\Delta}_{R_{n+1}}), \quad \widehat{\Delta}_7 = \widehat{\Delta} \cap \iota(\widehat{\Delta}_{R_n} \cup \{\varpi^{n+1}\} \setminus \widehat{\Delta}_{S_n}) \cap (\widehat{\Delta}_{R_{n+1}} \setminus \widehat{\Delta}_{S_{n+1}}). \end{aligned}$$

The union of $\widehat{\Delta}_i$ is $\widehat{\Delta}$. Suppose $\varpi_0 \in \widehat{\Delta}_i$ for $i = 2, 4, 6$. Using Lemma 6.1 and Corollary 6.2 of [Art78], We obtain constants $a_{\alpha_0, \varpi} \geq 0$ for $\varpi \in \widehat{\Delta}_{P_{n+1}} \sqcup \widehat{\Delta}_{R_{n+1}}$ directly from the proof of Lemma A.2 of [Zyd20] (equation (A.4) specifically), so that

$$\sum_{\varpi \in \widehat{\Delta}_{P_{n+1}}} a_{\alpha_0, \varpi}(\langle \varpi, Z \rangle - \langle w_{n+1} \varpi, H_{n+1} \rangle) + \sum_{\varpi \in \widehat{\Delta}_{R_{n+1}}} a_{\alpha_0, \varpi}(\langle \varpi, H_{n+1} \rangle - \langle w'_{n+1} \varpi, Z \rangle) \gg_{T,T'} \langle \alpha_0, Z \rangle - \langle \alpha_0, H_{n+1} \rangle.$$

We set the rest of the constants $a_{\alpha_0, \varpi}$ to zero. If $\varpi_0 \in \widehat{\Delta}_4 \cup \widehat{\Delta}_6$ we conclude directly from Lemma 6.1 of [Art78]. Otherwise, if $\varpi_0 \in \widehat{\Delta}_2$, the condition (3.4.11.13) gives $\sigma_{R_{n+1}}^{S_{n+1}, \tilde{S}_{n+1}}(H_{n+1} - T'_{n+1}) = 1$ and Lemma 3.4.6.1 implies $\langle \alpha_0, H_{n+1}, \tilde{S}_{n+1} \rangle \gg_{T'} \langle \alpha_0, H_{n+1} \rangle$.

If $\varpi_0 \in \widehat{\Delta}_i$ for $i = 1, 3, 5$ we obtain constants $a_{\alpha_0, \varpi}$ in the same fashion. One just has to observe that elements of $\Delta_{P_n}^{Q_n}$ act identically on \mathfrak{z}_P as their counterparts in $\Delta_{P_{n+1}}^{Q_{n+1}}$.

Suppose now that $\varpi_0 \in \widehat{\Delta}_7$. By the condition (3.4.11.9), we have $\varpi_0 \notin \widehat{\Delta}_{w'_n}$ or $\varpi_0 \notin \widehat{\Delta}_{w'_{n+1}}$. Suppose the latter holds. Using the results recalled in the beginning of Appendix A of [Zyd20], as well as Lemma 6.1 of [Art78], we have $\langle \varpi_0 - w'_{n+1}\varpi_0, Z \rangle \gg_T c_0 \langle \alpha_0, Z \rangle$ for some $c_0 > 0$ and $\langle \varpi_0 - w_{n+1}\varpi_0, H_{n+1} \rangle \gg_{T'} 0$. This identifies the the desired constants in this case. We deal with the other case in the same fashion, which completes the proof of Lemma 3.4.11.3. \square

Remark 3.4.11.4. — In the above proof, we do not use the fact that $T \in \mathfrak{a}_{0, n+1}^+$. Assuming this, we simply have $\langle \varpi_0 - w'_{n+1}\varpi_0, Z \rangle \gg c_0 \langle \alpha_0, Z \rangle$ when $\varpi_0 \in \widehat{\Delta}_7$, and similarly, Lemma A.2 in [Zyd20] invoked several times yields inequalities independent of T , as is clear by inspection of its proof in loc. cit. Therefore, if $T \in \mathfrak{a}_{0, n+1}^+$, the implied additive constant in Lemma 3.4.11.3 is independent of T .

The proof of the following lemma is similar and easier to that of the preceding one.

Lemma 3.4.11.5. — *Let $\alpha_n \in \Delta_{R_n}^{S_n}$, $\alpha_{n+1} \in \Delta_{R_{n+1}}^{S_{n+1}}$ and let $\varpi_n \in \widehat{\Delta}_{R_n}$ and $\varpi_{n+1} \in \widehat{\Delta}_{R_{n+1}}$ be their corresponding weights. There exists positive constants*

$$\underline{a}_{\alpha_n} = \{a_{\alpha_n, \varpi}\}_{\varpi \in \widehat{\Delta}_{P_n} \sqcup \widehat{\Delta}_{P_{n+1}} \sqcup \widehat{\Delta}_{R_n} \sqcup \widehat{\Delta}_{R_{n+1}}}$$

and

$$\underline{a}_{\alpha_{n+1}} = \{a_{\alpha_{n+1}, \varpi}\}_{\varpi \in \widehat{\Delta}_{P_n} \sqcup \widehat{\Delta}_{P_{n+1}} \sqcup \widehat{\Delta}_{R_n} \sqcup \widehat{\Delta}_{R_{n+1}}}$$

such that, assuming (3.4.11.13), we have

$$\lambda(\underline{a}_{\alpha_n}, Z, H_n, H_{n+1}) \gg_{T, T'} \begin{cases} \langle \alpha_n, H_n \rangle - \langle \alpha_n, Z \rangle & \text{if } \varpi_n \in (\widehat{\Delta}_{R_n} \setminus \widehat{\Delta}_{S_n}) \cap (\widehat{\Delta}_{P_n} \setminus \widehat{\Delta}_{Q_n}) \\ \langle \alpha_n, H_n \rangle & \text{else} \end{cases}$$

and

$$\lambda(\underline{a}_{\alpha_{n+1}}, Z, H_n, H_{n+1}) \gg_{T, T'} \begin{cases} \langle \alpha_{n+1}, H_{n+1} \rangle - \langle \alpha_{n+1}, Z \rangle & \text{if } \varpi_{n+1} \in (\widehat{\Delta}_{R_{n+1}} \setminus \widehat{\Delta}_{S_{n+1}}) \cap (\widehat{\Delta}_{P_{n+1}} \setminus \widehat{\Delta}_{Q_{n+1}}) \\ \langle \alpha_{n+1}, H_{n+1} \rangle & \text{else.} \end{cases}$$

Finally, we have the following result

Lemma 3.4.11.6. — *Let $\alpha_n \in \Delta_{\tilde{S}_n}$, $\alpha_{n+1} \in \Delta_{\tilde{S}_{n+1}}$ and let $\varpi_n \in \widehat{\Delta}_{\tilde{S}_n}$ and $\varpi_{n+1} \in \widehat{\Delta}_{\tilde{S}_{n+1}}$ be their corresponding weights. Then, there exists positive constants*

$$\underline{a}_{\alpha_n} = \{a_{\alpha_n, \varpi}\}_{\varpi \in \widehat{\Delta}_{P_n} \sqcup \widehat{\Delta}_{P_{n+1}} \sqcup \widehat{\Delta}_{R_n} \sqcup \widehat{\Delta}_{R_{n+1}}}$$

and

$$\underline{a}_{\alpha_{n+1}} = \{a_{\alpha_{n+1}, \varpi}\}_{\varpi \in \widehat{\Delta}_{P_n} \sqcup \widehat{\Delta}_{P_{n+1}} \sqcup \widehat{\Delta}_{R_n} \sqcup \widehat{\Delta}_{R_{n+1}}}$$

such that, assuming (3.4.11.13), $\varpi_n \notin (\widehat{\Delta}_{P_n} \setminus \widehat{\Delta}_{Q_n})$ and $\varpi_{n+1} \notin (\widehat{\Delta}_{P_{n+1}} \setminus \widehat{\Delta}_{Q_{n+1}})$ we have

$$\lambda(\underline{a}_{\alpha_n}, Z, H_n, H_{n+1}) \gg_{T, T'} -\langle \alpha_n, H_n \rangle$$

and

$$\lambda(\underline{a}_{\alpha_{n+1}}, Z, H_n, H_{n+1}) \gg_{T, T'} -\langle \alpha_{n+1}, H_{n+1} \rangle.$$

Otherwise, we define \underline{a}_{α_n} and $\underline{a}_{\alpha_{n+1}}$ as in 3.4.11.3 by identification of roots.

Proof. — Let's prove the statement concerning α_n , the other case being analogous. Let $\gamma_n \in \Delta_{B_n} \setminus \Delta_{B_n}^{\tilde{S}_n}$ be its lift. We must have $\varpi_n \in (\hat{\Delta}_{B_n} \setminus \hat{\Delta}_{P_n}) \cup \hat{\Delta}_{Q_n}$. Then, as explained in the proof of Lemma 3.4.11.3, Lemmas A.2 of [Zyd20] and 6.1 of [Art78] provide constants \underline{a}_{α_n} such that

$$\lambda(\underline{a}_{\alpha_n}, Z, H_n, H_{n+1}) \gg_{T, T'} -\langle \gamma_n, H_n \rangle$$

Condition (3.4.11.13) and Lemma 3.4.6.1 prove that $\langle \alpha_n, H_n \rangle \gg_{T'} \langle \gamma_n, H_n \rangle$ which allows to conclude. \square

We are ready to define the desired $\lambda_1 = (\lambda_{1,n}, \lambda_{1,n+1}) \in \mathfrak{a}_{P_n}^* \times \mathfrak{a}_{P_{n+1}}^*$ and $\lambda_2 = (\lambda_{2,n}, \lambda_{2,n+1}) \in \mathfrak{a}_{R_n}^* \times \mathfrak{a}_{R_{n+1}}^*$ of Lemma 3.4.11.2. In Lemmas 3.4.11.3, 3.4.11.5 and 3.4.11.6 we have defined sets \underline{a}_α for all $\alpha \in \Delta_{P_{n+1}} \sqcup \Delta_{R_n}^{S_n} \sqcup \Delta_{R_{n+1}}^{S_{n+1}} \sqcup \Delta_{\tilde{S}_n} \sqcup \Delta_{\tilde{S}_{n+1}}$. Let $k \geq 0$. We set

$$\begin{aligned} \lambda_{1,n} &= \sum_{\varpi \in \hat{\Delta}_{P_n}} \left(\sum_{\alpha \in \Delta_{P_{n+1}}} k a_{\alpha, \varpi} + \sum_{\alpha \in \Delta_{R_n}^{S_n} \sqcup \Delta_{R_{n+1}}^{S_{n+1}} \sqcup \Delta_{\tilde{S}_n} \sqcup \Delta_{\tilde{S}_{n+1}}} a_{\alpha, \varpi} \right) \varpi, \\ \lambda_{1,n+1} &= \sum_{\varpi \in \hat{\Delta}_{P_{n+1}}} \left(\sum_{\alpha \in \Delta_{P_{n+1}}} k a_{\alpha, \varpi} + \sum_{\alpha \in \Delta_{R_n}^{S_n} \sqcup \Delta_{R_{n+1}}^{S_{n+1}} \sqcup \Delta_{\tilde{S}_n} \sqcup \Delta_{\tilde{S}_{n+1}}} a_{\alpha, \varpi} \right) \varpi, \\ \lambda_{2,n} &= \sum_{\varpi \in \hat{\Delta}_{R_n}} \left(\sum_{\alpha \in \Delta_{P_{n+1}}} k a_{\alpha, \varpi} + \sum_{\alpha \in \Delta_{R_n}^{S_n} \sqcup \Delta_{R_{n+1}}^{S_{n+1}} \sqcup \Delta_{\tilde{S}_n} \sqcup \Delta_{\tilde{S}_{n+1}}} a_{\alpha, \varpi} \right) \varpi, \\ \lambda_{2,n+1} &= \sum_{\varpi \in \hat{\Delta}_{R_{n+1}}} \left(\sum_{\alpha \in \Delta_{P_{n+1}}} k a_{\alpha, \varpi} + \sum_{\alpha \in \Delta_{R_n}^{S_n} \sqcup \Delta_{R_{n+1}}^{S_{n+1}} \sqcup \Delta_{\tilde{S}_n} \sqcup \Delta_{\tilde{S}_{n+1}}} a_{\alpha, \varpi} \right) \varpi. \end{aligned}$$

We have then for $Z \in \mathfrak{z}_P$ and $H = (H_1, H_2) \in \mathfrak{a}_{R_n}^{G_n} \times \mathfrak{a}_{R_{n+1}}^{G_{n+1}}$

$$(3.4.11.14) \quad \langle \lambda_1 - w' \lambda_2, Z \rangle + \langle \lambda_2 - w \lambda_1, H \rangle = \sum_{\alpha \in \Delta_{P_{n+1}}} k \lambda(\underline{a}_\alpha, Z, H_n, H_{n+1}) + \sum_{\alpha \in \Delta_{R_n}^{S_n} \sqcup \Delta_{R_{n+1}}^{S_{n+1}} \sqcup \Delta_{\tilde{S}_n} \sqcup \Delta_{\tilde{S}_{n+1}}} \lambda(\underline{a}_\alpha, Z, H_n, H_{n+1})$$

which, by Lemmas 3.4.11.3, 3.4.11.5 and 3.4.11.6 proves the desired inequality of Lemma 3.4.11.2 for k sufficiently large (the constant k is there to offset the $-\langle \alpha, Z \rangle$ terms of Lemma 3.4.11.5).

We have proven Lemma 3.4.11.2 which entails Lemma 3.4.11.1 which accomplishes the proof of assertion 1 of Theorem 3.4.3.2.

3.4.12. Proof of assertions 2 and 3 of Theorem 3.4.3.2. — If we replace κ_χ^T by K_χ^T and assume that f is compactly supported, statement 2 is proved in [Zyd20]. Practically the same method applies to κ_χ^T as well as Schwartz functions, since we have proven the convergence assertion of Theorem 3.4.3.2, which gives rise to the distribution i_χ . Its continuity follows immediately from the examination of the proof of assertion 1 and the analogue of Theorem 3.7 of [Zyd20]. What is not obvious however is that $i_\chi(f) = I_\chi(f)$. We will prove this for compactly supported f and the general case will follow by continuity.

As in §§3.4.11, unless otherwise stated, all sums of the type Σ_P or $\Sigma_{P \subset Q}$ are over elements of \mathcal{F}_{RS} .

Let $Q \in \mathcal{F}_{RS}$. Define

$$K_\chi^{T,Q}(x, y) = \sum_{Q \supset P} \epsilon_P^Q \sum_{\gamma \in (P \cap H)(F) \backslash (Q \cap H)(F)} \sum_{\delta \in P'(F) \backslash Q'(F)} \hat{\tau}_{P_{n+1}}^{Q_{n+1}}(H_{P_{n+1}}(\delta_n y_n) - T_{P_{n+1}}) K_{P, \chi}(\gamma x, \delta y).$$

For $P \subset Q$, elements of \mathcal{F}_{RS} , let

$$\mathfrak{a}_{P,Q}^{*, -} = \{\lambda \in \mathfrak{a}_{P_{n+1}}^* \mid \langle \lambda + 2\rho_Q, \varpi^\vee \rangle < 0, \forall \varpi \in \widehat{\Delta}_Q\}$$

where for $Q \in \mathcal{F}_{RS}$ we set

$$\rho_Q = \rho_{Q_{n+1}} - \rho_{Q_n} \in \mathfrak{a}_{0, n+1}.$$

Define for $\lambda \in \mathfrak{a}_{Q_{n+1}, \mathbb{C}}^*$

$$I_\chi^T(f, Q, \lambda) := \int_{(Q \cap H)(F) \backslash H(\mathbb{A})} \int_{Q'(F) \backslash G'(\mathbb{A})} e^{\langle \lambda, H_{Q_{n+1}}(x) \rangle} \tau_{Q_{n+1}}(H_{Q_{n+1}}(y_n) - T_{Q_{n+1}}) K_\chi^{T,Q}(x, y) \eta_{G'}(y) dy dx.$$

The integral converges absolutely for $\Re(\lambda) \in \mathfrak{a}_{Q,Q}^{*, -}$ and admits a meromorphic continuation, holomorphic at $\lambda = 0$ so that we have

$$(3.4.12.15) \quad I_\chi(f) = \sum_Q I_\chi^T(f, Q, 0)$$

for any T . The equality above is a formal consequence of Theorem 3.7 of [Zyd20].

Let $P \in \mathcal{F}_{RS}$. For a function $\phi : P_n(F) \backslash G_n(\mathbb{A}) \rightarrow \mathbb{C}$, we introduce the operator $\Lambda_{m'}^{T,P}$: it is a variant of the mixed operator $\Lambda_m^{T,P}$ of [JLR99]. It is defined in [Zyd20] §2.3 (where it is denoted by $\Lambda_m^{T,P}$). We have

$$\Lambda_{m'}^{T,P} \phi(y_n) = \sum_{P \supset Q} \epsilon_Q^P \sum_{\delta \in Q'_n(F) \backslash P'_n(F)} \hat{\tau}_{Q_{n+1}}^{P_{n+1}}(H_{Q_{n+1}}(\delta y_n) - T_{Q_{n+1}}) \phi_{Q_n}(\delta y_n), \quad y_n \in P'_n(F) \backslash G'_n(\mathbb{A}).$$

We will also use the operator $\Lambda_d^{T,P}$ as in (3.4.11.4).

Fix $R \subset R^1 \subset Q$ and $S \subset S^1 \subset Q$ all elements of \mathcal{F}_{RS} . Let $\lambda \in \mathfrak{a}_{R_{n+1}, \mathbb{C}}^*$. We define $I_\chi^T(f, R, R^1, S, S^1, Q, \lambda)$ as

$$\begin{aligned} & \int_{(R \cap H)(F) \backslash H(\mathbb{A})} \int_{(S'_n \times G'_{n+1})(F) \backslash G'(\mathbb{A})} e^{\langle \lambda, H_{R_{n+1}}(x) \rangle} \tau_{Q_{n+1}}(H_{Q_{n+1}}(x) - T'_{Q_{n+1}}) \tau_{Q_{n+1}}(H_{Q_{n+1}}(y_n) - T_{Q_{n+1}}) \\ & \sigma_{R_{n+1}}^{R_{n+1}^1, Q_{n+1}}(H_{R_{n+1}}(x) - T'_{R_{n+1}}) \sigma_{S_{n+1}}^{S_{n+1}^1, Q_{n+1}}(H_{S_{n+1}}(y_n) - T_{S_{n+1}}) \Lambda_d^{T', R} \Lambda_{m'}^{T, S} \\ & \left(\sum_{R \cup S \subset P \subset R^1 \cap S^1} \epsilon_{P_{n+1}}^{Q_{n+1}} \sum_{\delta_{n+1} \in P'_{n+1}(F) \backslash G'_{n+1}(F)} K_{P, \chi}(x, y_n, \delta_{n+1} y_{n+1}) \right) \eta_{G'}(y) dx dy \end{aligned}$$

where T' is a translate of T by an element of \mathfrak{a}_{n+1} that depends only on the support of f , $\Lambda_{m'}^{T, S}$ is applied with respect to y and $\Lambda_d^{T', R}$ with respect to x . The proof of Theorem 3.1 in [Zyd20] proves that the above integral converges absolutely for any $\Re(\lambda) \in \mathfrak{a}_{R,Q}^{*, -}$ and extends everywhere to a meromorphic function, holomorphic at 0. Moreover, we have the equality of meromorphic functions on $\mathfrak{a}_{Q_{n+1}, \mathbb{C}}^*$.

$$I_\chi^T(f, Q, \lambda) = \sum_{R \subset R^1 \subset Q} \sum_{S \subset S^1 \subset Q} I_\chi^T(f, R, R^1, S, S^1, Q, \lambda).$$

Fix $R \in \mathcal{F}_{RS}$ and a $\lambda \in \mathfrak{a}_{R,n+1,\mathbb{C}}^*$ such that $\Re(\lambda) \in \mathfrak{a}_{R,R}^{*,\bar{-}}$. Note that $\mathfrak{a}_{R,R}^{*,\bar{-}} \subset \mathfrak{a}_{R,Q}^{*,\bar{-}}$ for any $Q \supset R$. Fix also $S \in \mathcal{F}_{RS}$. We look at the sum

$$\sum_Q \sum_{R \subset R^1 \subset Q} \sum_{S \subset S^1 \subset Q} I_\chi^T(f, R, R^1, S, S^1, Q, \lambda).$$

We claim it equals $I_\chi^T(f, R, S, \lambda)$ defined by meromorphic continuation of

$$\int_{(R \cap H)(F) \backslash H(\mathbb{A})} \int_{(S'_n \times G'_{n+1})(F) \backslash G'(\mathbb{A})} e^{\langle \lambda, H_{R_{n+1}}(x) \rangle} \tau_{R_{n+1}}(H_{R_{n+1}}(x) - T'_{R_{n+1}}) \tau_{S_{n+1}}(H_{S_{n+1}}(y_n) - T_{S_{n+1}}) \Lambda_d^{T',R} \Lambda_{m'}^{T,S} K_\chi(x, y) \eta_{G'}(y) dx dy.$$

Fix $P \in \mathcal{F}_{RS}$ containing R and S . We see that we need to show that

$$K_{P,\chi}(x, y) \left(\sum_{Q \supset P} \epsilon_{P_{n+1}}^{Q_{n+1}} \tau_{Q_{n+1}}(X) \tau_{Q_{n+1}}(Y) \sum_{P \subset R^1, S^1 \subset Q} \sigma_{R_{n+1}}^{R^1, Q_{n+1}}(X) \sigma_{S_{n+1}}^{S^1, Q_{n+1}}(Y) \right) = \begin{cases} K_\chi(x, y) \tau_{R_{n+1}}(X) \tau_{S_{n+1}}(Y) & \text{if } P = G, \\ 0 & \text{if } P \neq G, \end{cases}$$

where $X = H_{R_{n+1}}(x) - T'_{R_{n+1}}$, $Y = H_{S_{n+1}}(y_n) - T_{S_{n+1}}$.

Since $\sum_{R^1 \supset P} \sigma_{R_{n+1}}^{R^1, Q_{n+1}} = \tau_{R_{n+1}}^{P_{n+1}} \hat{\tau}_{P_{n+1}}^{Q_{n+1}}$ we need to consider

$$K_{P,\chi}(x, y) \tau_{R_{n+1}}^{P_{n+1}}(X) \tau_{S_{n+1}}^{P_{n+1}}(Y) \left(\sum_{Q \supset P} \epsilon_{P_{n+1}}^{Q_{n+1}} \tau_{Q_{n+1}}(X) \hat{\tau}_{P_{n+1}}^{Q_{n+1}}(X) \tau_{Q_{n+1}}(Y) \hat{\tau}_{P_{n+1}}^{Q_{n+1}}(Y) \right).$$

We obtain the desired result if $P = G$. We can assume then $P \neq G$ and that $K_{P,\chi}(x, y) \neq 0$. We want to show the expression is zero. By the argument of the beginning of the proof of Theorem 3.1 in [Zyd20], and definition of T' , under the assumption $K_{P,\chi}(x, y) \neq 0$ we have that $\tau_{Q_{n+1}}(X) \hat{\tau}_{P_{n+1}}^{Q_{n+1}}(X) \tau_{Q_{n+1}}(Y) \hat{\tau}_{P_{n+1}}^{Q_{n+1}}(Y) = \tau_{Q_{n+1}}(Y) \hat{\tau}_{P_{n+1}}^{Q_{n+1}}(Y)$ which shows the desired vanishing by the Langlands Combinatorial Lemma.

We have shown thus that

$$I_\chi(f) = \sum_{R, S \in \mathcal{F}_{RS}} I_\chi^T(f, R, S, 0)$$

where T is arbitrary. Note however that if we fix R and set $I_\chi^T(f, R, \lambda)$ to be (3.4.12.16)

$$\int_{(R \cap H)(F) \backslash H(\mathbb{A})} \int_{G'(F) \backslash G'(\mathbb{A})} e^{\langle \lambda, H_{R_{n+1}}(x) \rangle} \tau_{R_{n+1}}(H_{R_{n+1}}(x) - T_{R_{n+1}}) \Lambda_d^{T,R} K_\chi(x, y) \eta_{G'}(y) dx dy$$

we get, using the inversion formula Lemme 2.4 [Zyd20]:

$$\sum_{S \in \mathcal{F}_{RS}} I_\chi^T(f, R, S, \lambda) = I_\chi^{T'}(f, R, \lambda).$$

This entails that

$$(3.4.12.17) \quad I_\chi(f) = \sum_{R \in \mathcal{F}_{RS}} I_\chi^{T'}(f, R, 0), \quad T' \in \mathfrak{a}_{n+1}.$$

Similar and more direct reasoning yields a formula for i_χ as follows. We define $i_\chi^T(f, R, S, Q, \lambda)$, for $R \subset S \subset Q$, elements of \mathcal{F}_{RS} , and $\lambda \in \mathfrak{a}_{R, \mathbb{C}}^*$, to be

$$\int_{(R \cap H)(F) \backslash H(\mathbb{A})} \int_{[G']} e^{\langle \lambda, H_{R_{n+1}}(x) \rangle} \tau_{Q_{n+1}}(H_{Q_{n+1}}(x) - T_{Q_{n+1}}) \sigma_{R_{n+1}}^{S_{n+1}, Q_{n+1}}(H_{R_{n+1}}(x) - T_{R_{n+1}}) \\ \Lambda_d^{T, R} \left(\sum_{R \subset P \subset S} \epsilon_{P_{n+1}}^{Q_{n+1}} \sum_{\delta \in P'(F) \backslash G'(F)} K_{P, \chi}(x, \delta y) \right) \eta_{G'}(y) dx dy$$

which converges absolutely for $\Re(\lambda) \in \mathfrak{a}_{R, Q}^{*-}$ and admits meromorphic continuation satisfying

$$\sum_{R \subset S \subset Q} i_\chi^T(f, R, S, Q, 0) = i_\chi(f), \quad T \in \mathfrak{a}_{n+1}.$$

Additionally, for a fixed R we have

$$\sum_{S \subset Q} i_\chi^T(f, R, S, Q, \lambda) = I_\chi^T(f, R, \lambda)$$

where $I_\chi^T(f, R, \lambda)$ is as in (3.4.12.16). Comparing the above two equalities with (3.4.12.17), taking $T' = T$, we get $i_\chi(f) = I_\chi(f)$ as desired.

3.5 Asymptotic formulas

3.5.1. Recall that we have defined integrals $I_\chi^T(f)$ and $i_\chi^T(f)$ (see Theorems 3.2.4.1 and 3.4.3.2).

Theorem 3.5.1.1. —

1. $i_\chi^T(f)$ is asymptotically equal to each of the integrals (3.3.3.2) (see Proposition 3.3.3.1) and (3.3.5.5) (see Proposition 3.3.5.1).
2. $I_\chi^T(f)$ is asymptotically equal to the integral (3.3.5.6) (see Proposition 3.3.5.1).

Remark 3.5.1.2. — The above theorem can be stated for the whole kernel K instead of K_χ . It can be proved in the same way.

Proof. — Let us revisit the proof of Theorem 3.4.3.2 keeping track of constants depending on the parameter T . It is not hard to see that Proposition 2.8 of [Zyd20], or any analogous result, including the original Lemma 1.4 of [Art80], gives a constant

$$e^{r\|T\|}$$

for r depending on r_1 and r'_1 . The expression (3.4.11.7) should be multiplied by $e^{r\|T\|}$ accordingly. We move on to the expression (3.4.11.12). The constants N_1 and N_2 are arbitrary and their choice is independent of T . They influence the constants N'_1 and N'_2 respectively, however. We take $N_2 = 0$ and choose N'_1 so that $N'_2 - N'_1 < 0$. To make sure that the integral over m_2 converges we choose the constant r'_2 in (3.4.11.7) small enough. This choice is independent of T as well. Choosing λ_1 (and λ_2) appropriately (choice not influencing dependency on T), the conclusion of the proof of Theorem 3.4.3.2 is that the integral (3.4.11.7) is bounded by

$$(3.5.1.1) \quad e^{r\|T\|} \int_{\mathfrak{z}_P} \sigma_P^Q(Z - T) e^{\langle -(\lambda_1 - w'\lambda_2), Z \rangle} dZ$$

However, looking at (3.4.11.14), we can make the constant k as big as we please, which, by Lemma 3.4.11.3, makes the exponent in the integral above as negative as we wish, without introducing

dependency on T by Remark 3.4.11.4. Reasoning as in the end of Theorem 2.2 of [Zyd16], having liberty with the exponent, we can make the integral (3.5.1.1) smaller than any power of $e^{-\|T\|}$ as long as $P \neq G$. We have showed thus that $i_\chi^T(f)$ asymptotically equals

$$(3.5.1.2) \quad \int_{[H]} \int_{[G']} \Lambda_d^T K_\chi(x, y) \eta_{G'}(y) dy dx.$$

Next, we show that (3.5.1.2) asymptotically equals $\int_{[H]} \int_{[G']} F^{G_{n+1}}(x, T) K_\chi(x, y) \eta_{G'}(y) dy dx$. Indeed, the argument is essentially identical to the one in [IY15], Proposition 3.8. One just needs to use the fact that for an element X in the universal enveloping algebra of $Lie(G) \otimes_{\mathbb{R}} \mathbb{C}$, the function $R(X)K_\chi(x, y)$, where $R(X)$ is the right action on $K_{f, \chi}(x, y)$ as a function of the variable x , is just $K_{R(X)f, \chi}(x, y)$. The uniform growth assumption of Proposition 3.8 in *loc. cit.* can then be replaced with the application of the bound (2.9.8.11).

Exactly the same reasoning works with Λ_d^T replaced with Λ_r^T which yield point 1. Point 2 follows exactly the same reasoning applied to the kernel K_χ^T . \square

4 Flicker-Rallis period of some spectral kernels

The goal of this chapter is to get the spectral expansion of the Flicker-Rallis integral of the automorphic kernel attached to a linear group and a specific cuspidal datum (called in §4.3.2 *-generic.) This is achieved in theorem 4.3.3.1. It turns out that the decomposition is discrete and is expressed in terms of some relative characters.

4.1 Flicker-Rallis intertwining periods and related distributions

4.1.1. Notations. — In all this section, we will fix an integer $n \geq 1$ and we will use notations of §§3.1.1 to 3.1.3. Since n will be fixed, we will drop the subscript n from the notation: $G = G_n$, $B = B_n$ etc. So we do not follow notations of §3.1.5: we hope that it will cause no confusion.

4.1.2. Flicker-Rallis periods. — Let π be a cuspidal automorphic representation of $G(\mathbb{A})$ with central character trivial on A_M^∞ . We shall denote by π^* the conjugate-dual representation of $G(\mathbb{A})$. We shall say that π is self conjugate-dual if $\pi \simeq \pi^*$ and that π is G' -distinguished, resp. (G', η) -distinguished, if the linear form (called the Flicker-Rallis period)

$$(4.1.2.1) \quad \varphi \mapsto \int_{[G']_0} \varphi(h) dh, \text{ resp. } \int_{[G']_0} \varphi(h) \eta(\det(h)) dh$$

does not vanish identically on $\mathcal{A}_\pi(G)$. Then π is self conjugate-dual if and only if π is either G' -distinguished or (G', η) -distinguished. However it cannot be both. This is related to the well-known factorisation of the Rankin-Selberg factorisation $L(s, \pi \times \pi \circ c)$ where c is the Galois involution of $G(\mathbb{A})$ in terms of Asai L -functions and to the fact that the residue at $s = 1$ of the Asai L -functions is expressed in terms of Flicker-Rallis periods (see [Fli88]).

4.1.3. In this chapter, we will focus on the period in (4.1.2.1) related to distinction. However it is clear that all the results hold *mutatis mutandis* for the period related to η -distinction.

4.1.4. Let $P = MN_P$ be a standard parabolic subgroup (with its standard decomposition). Let π be an irreducible cuspidal automorphic representation of M with central character trivial on A_M^∞ .

It will be convenient to write $M = G_{n_1} \times \dots \times G_{n_r}$ with $n_1 + \dots + n_r = n$. Accordingly we have $\pi = \sigma_1 \boxtimes \dots \boxtimes \sigma_r$ where σ_i is an irreducible cuspidal representation of G_{n_i} .

4.1.5. Let $\varphi \in \mathcal{A}_{P, \pi}(G)$. The parabolic subgroup $P' = P \cap G'$ of G' has the following Levi decomposition $M'N_{P'}$ where $M' = M \cap G'$. We then define the following integral which is a specific example of a Flicker-Rallis intertwining period introduced by Jacquet-Lapid-Rogawski (see [JLR99] section VII, note that our definition of $\mathcal{A}_{P, \pi}(G)$ is slightly different from theirs),

$$J(\varphi) = \int_{A_{M'}^\infty M'(F)N_{P'}(\mathbb{A}) \backslash G'(\mathbb{A})} \varphi(g) dg$$

Clearly we get a $G'(\mathbb{A})$ -invariant continuous linear form on $\mathcal{A}_{P, \pi}(G)$. Note that J does not vanish identically if and only if each component σ_i is $G'_{n_i}(F)$ -distinguished. In this case, we have $\pi = \pi^*$.

4.1.6. Let $Q \in \mathcal{P}(M)$. As recalled in §2.2.11, there is a unique pair (Q', w) such that the conditions are satisfied:

- $Q' = wQw^{-1}$ is the standard parabolic subgroup in the G -conjugacy class of Q ;
- $w \in W(P; Q')$.

Let $\lambda \in \mathfrak{a}_{P, \mathbb{C}}^{G, *}$. We have $M(w, \lambda)\varphi \in \mathcal{A}_{M_{Q'}, w\pi}(G)$ if λ is outside the singular hyperplanes of the intertwining operator. We shall define

$$(4.1.6.2) \quad J_Q(\varphi, \lambda) = J(M(w, \lambda)\varphi)$$

as a meromorphic function of λ .

4.1.7. Let $g \in G(\mathbb{A})$. Let's define for $\varphi, \psi \in \mathcal{A}_{P,\pi}(G)$

$$(4.1.7.3) \quad B_Q(g, \varphi, \psi, \lambda) = E(g, \varphi, \lambda) \cdot J_Q(\bar{\psi}, -\lambda).$$

as a meromorphic function of $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^{G,*}$. In fact, by the basic properties of Eisenstein series and intertwining operators, there exists an open subset $\omega_\pi \subset \mathfrak{a}_{P,\mathbb{C}}^{G,*}$ which is the complement of a union of hyperplanes of $\mathfrak{a}_{P,\mathbb{C}}^{G,*}$ such that :

- ω_π contains $i\mathfrak{a}_P^{G,*}$.
- for all $\varphi, \psi \in \mathcal{A}_{P,\pi}(G)$, the map $\lambda \mapsto B_Q(g, \varphi, \psi, \lambda)$ is holomorphic on ω_π and gives for each $\lambda \in \omega_\pi$ a continuous sesquilinear form in φ and ψ .

4.1.8. Let $f \in \mathcal{S}(G(\mathbb{A}))$, $g \in G(\mathbb{A})$ and $Q \in \mathcal{P}(M)$. Let's introduce the distribution

$$(4.1.8.4) \quad J_{Q,\pi}(g, \lambda, f) = \sum_{\varphi \in \mathcal{B}_{P,\pi}} B_Q(g, I_P(\lambda, f)\varphi, \varphi, \lambda)$$

where $\mathcal{B}_{P,\pi}$ is a K -basis of $\mathcal{A}_{P,\pi}(G)$ (see § 2.8.3 and $\lambda \in \omega_\pi$ (see § 4.1.7 for the notation ω_π). It follows from propositions 2.8.4.1 that $J_{Q,\pi}(g, \lambda)$ is a continuous distribution on $\mathcal{S}(G(\mathbb{A}))$.

4.1.9. A (G, M) -family. —

Proposition 4.1.9.1. — *The family $(J_{Q,\pi}(g, \lambda, f))_{Q \in \mathcal{P}(M)}$ is a (G, M) -family in the sense of Arthur (see [Art81]): namely each map*

$$\lambda \in \mathfrak{a}_P^{G,*} \mapsto J_{Q,\pi}(g, \lambda, f)$$

is smooth on $i\mathfrak{a}_P^{G,}$ (and even holomorphic on ω_π) and for adjacent elements $Q_1, Q_2 \in \mathcal{P}(M)$ we have*

$$(4.1.9.5) \quad J_{Q_1,\pi}(g, \lambda, f) = J_{Q_2,\pi}(g, \lambda, f)$$

on the hyperplane of $i\mathfrak{a}_L^{G,}$ defined by $\langle \lambda, \alpha^\vee \rangle = 0$ where α is the unique element in $\Delta_{Q_1} \cap (-\Delta_{Q_2})$.*

Proof. — The holomorphy on ω_π is obvious if f is K_∞ -finite (the sum in 4.1.8.4 is then finite). Let $C' \subset \omega_\pi$ be a compact subset. Using approximations of f by K_∞ -finite functions, one shows that $J_{Q,\pi}(g, \lambda, f)$ is a uniform limit on C' of holomorphic functions hence holomorphic.

Let $Q_1, Q_2 \in \mathcal{P}(M)$ be such that $\Delta_{Q_1} \cap (-\Delta_{Q_2})$ is a singleton $\{\alpha\}$. Let $\lambda \in i\mathfrak{a}_L^{G,*}$ such $\langle \lambda, \alpha^\vee \rangle = 0$. For $i = 1, 2$ let Q'_i be a standard parabolic subgroup and $w_i \in W(M, Q'_i)$ be such that $Q'_i = w_i Q_i w_i^{-1}$. Let $\beta = w_1 \alpha \in \Delta_{Q'_1}$ and Let s_β the simple reflection associated to β . Then we have $w_2 = s_\beta w_1$. Let $\varphi \in \mathcal{A}_{P,\pi}(G)$. Clearly it suffices to check the equality:

$$E(g, \varphi, \lambda) \cdot \overline{(J(M(w_1, \lambda)\varphi))} = E(g, \varphi, \lambda) \cdot \overline{J(M(w_2, \lambda)\varphi)}.$$

Using the functional equations of intertwining operators and Eisenstein series, we have $M(w_2, \lambda) = M(s_\beta, w_1 \lambda) M(w_1, \lambda)$ and $E(g, \varphi, \lambda) = E(g, M(w_1, \lambda)\varphi, w_1 \lambda)$. Thus up to a change of notations (replace P by Q'_1), we may assume that $Q_1 = P$ and thus $w_1 = 1$ and $\alpha = \beta$. We are reduced to prove

$$(4.1.9.6) \quad E(g, \varphi, \lambda) \cdot \overline{J(\varphi)} = E(g, \varphi, \lambda) \cdot \overline{J(M(s_\alpha, \lambda)\varphi)}$$

on the hyperplane $\langle \lambda, \alpha^\vee \rangle = 0$. The simple reflection s_α acts on M as a transposition of two consecutive blocks of M say G_{n_i} and $G_{n_{i+1}}$. Note that $M(s_\alpha, \lambda)\varphi = M(s_\alpha, 0)$. Then we have even a stronger property:

$$J(\varphi) = J(M(s_\alpha, 0)\varphi)$$

if $n_i \neq n_{i+1}$ or if $n_i = n_{i+1}$ but $\sigma_i \not\simeq \sigma_{i+1}^*$ (see lemma 8.1 case 1 of [Lap06]). Assume that $n_i = n_{i+1}$ and $\sigma_i \simeq \sigma_{i+1}^*$. The case where $\sigma_i \not\simeq \sigma_i^*$ is trivial (J is zero) so we shall also assume that $\sigma_i \simeq \sigma_i^*$. Then $M(s_\alpha, 0)\varphi = -\varphi$ ([KS88] proposition 6.3) and since $s_\alpha(\lambda) = \lambda$ we have $E(g, \varphi, \lambda) = 0$ so (4.1.9.6) is clear. \square

4.1.10. Majorization. — We will use the following proposition which results from Lapid's majorization of Eisenstein series (see [Lap06] proposition 6.1 and section 7). For the convenience of the reader, we sketch a proof.

Proposition 4.1.10.1. — *Let $f \in \mathcal{S}(G(\mathbb{A}))$. The map $\lambda \mapsto J_{Q,\pi}(g, \lambda, f)$ belongs to the Schwartz space $\mathcal{S}(\mathfrak{ia}_P^{G,*})$. Moreover*

$$f \mapsto J_{Q,\pi}(g, \cdot, f)$$

is a continuous map from $\mathcal{S}(G(\mathbb{A}))$ to $\mathcal{S}(\mathfrak{ia}_P^{G,})$ equipped with its usual topology.*

Proof. — let $C \in G(\mathbb{A}_f)$ be a compact subset and $K_0 \subset K^\infty$ be an open-compact subgroup such that $f \in \mathcal{S}(G(\mathbb{A}), C, K_0)$. For any $\alpha, \beta > 0$, we define an open subset $\omega_{\alpha,\beta}$ of $\mathfrak{a}_{P,\mathbb{C}}^{G,*}$ which contains $\mathfrak{ia}_{P,\mathbb{R}}^{G,*}$ by

$$\omega_{\alpha,\beta} = \{\lambda \in \mathfrak{a}_{P,\mathbb{C}}^{G,*} \mid \|\Re(\lambda)\| < \alpha(1 + \|\Im(\lambda)\|)^{-\beta}\}.$$

By the arguments in the proof of proposition 6.1 of [Lap06], one sees that there exist $\alpha, \beta > 0$ such that $\omega_{\alpha,\beta}$ is included in the open set ω_π of §4.1.7. In particular, $\lambda \mapsto J_{Q,\pi}(g, \lambda, f)$ is holomorphic on $\omega_{\alpha,\beta}$. Using Cauchy formula to control derivatives, it suffices to prove the following majorization: there exists a continuous semi-norm $\|\cdot\|$ on $\mathcal{S}(G(\mathbb{A}), C, K_0)$ and an open subset $\omega_{\alpha,\beta} \subset \omega_\pi$ such that for any integer $N \geq 1$ there exists $c > 0$ so that for all $f \in \mathcal{S}(G(\mathbb{A}), C, K_0)$ and all $\lambda \in \omega_{\alpha,\beta}$

$$(4.1.10.7) \quad |J_{Q,\pi}(g, \lambda, f)| \leq c \frac{\|f\|}{(1 + \|\lambda\|)^N}.$$

Let $m \geq 1$ be a large enough integer. Following the notations of proposition 2.8.4.3, we can write $f = f * g_1 + (f * Z) * g_2$; we get

$$\begin{aligned} J_{Q,\pi}(g, \lambda, f) &= \sum_{\varphi \in \mathcal{B}_{P,\pi}} E(g, I_P(\lambda, f)\varphi, \lambda) \overline{J_Q(I_P(-\lambda, g_1^\vee)\varphi, \lambda)} \\ &+ \sum_{\varphi \in \mathcal{B}_{P,\pi}} E(g, I_P(\lambda, f * Z)\varphi, \lambda) \overline{J_Q(I_P(-\lambda, g_2^\vee)\varphi, \lambda)} \end{aligned}$$

By a slight extension to Schwartz functions of Lapid's majorization (see [FLO12] remark C.2 about [Lap06] proposition 6.1), the expression

$$\left(\sum_{\varphi \in \mathcal{B}_{P,\pi}} |E(g, I_P(\lambda, f)\varphi, \lambda)|^2 \right)^{1/2}$$

and the same expression where f is replaced by $f * Z$ satisfy a bound like (4.1.10.7). Using Cauchy-Schwartz inequality, we are reduced to bound in λ (recall that g_i is independent of f)

$$(4.1.10.8) \quad \left(\sum_{\varphi \in \mathcal{B}_{P,\pi}} |J_Q(I_P(-\lambda, g_i^\vee)\varphi, \lambda)|^2 \right)^{1/2}.$$

Let w be such that wQw^{-1} is standard and $w \in W(P, wQw^{-1})$. At this point we will use the notations of the proof of proposition 2.8.4.1. There exists $c > 0$ and an integer r such that for all $\varphi' \in \mathcal{A}_{P,\pi}(G)^{K_0}$ we have

$$|J_Q(I_P(-\lambda, g_i^\vee)\varphi, \lambda)| = |J(M(w, \lambda)I_P(-\lambda, g_i^\vee)\varphi)| \leq c \|M(w, \lambda)I_P(-\lambda, g_i^\vee)\varphi\|_r$$

where $\|\varphi\|_r = \|R(1 + C_K)^r \varphi\|_{\text{Pet}}$. Then we need to bound the operator norm of the intertwining operator $M(w, \lambda)$. Using the normalization of intertwining operators, the bounds of normalizing factors [Lap06] lemma 5.1 and Müller-Speh's bound on the norm of normalized intertwining operators (see [MS04] proposition 4.2 and the proof of proposition 0.2), we get $c_1 > 0$, $N \in \mathbb{N}$ and $\alpha, \beta > 0$ such that for all $\tau \in \hat{K}_\infty$, $\lambda \in \omega_{\alpha, \beta}$ and $\varphi \in \mathcal{A}_{P, \pi}(G, K_0, \tau)$ we have

$$\|M(w, \lambda)I_P(-\lambda, g_i^\vee)\varphi\|_{K_0, r} \leq c_1(1 + \lambda_\tau)^N \|I_P(-\lambda, g_i^\vee)\varphi\|_r.$$

Using the same kind of arguments as in the proof of proposition 2.8.4.1 (see also remark (2.8.4.2)), one shows that there exist $\alpha, \beta > 0$ such that (4.1.10.8) is bounded independently of $\lambda \in \omega_{\alpha, \beta}$. \square

4.2 A spectral expansion of a truncated integral

4.2.1. Let $\chi \in \mathfrak{X}(G)$ be a cuspidal datum. We shall use the notation of §2.9.9. In particular, f is a function in $f \in \mathcal{S}(G(\mathbb{A}), C, K_0)$ and K_χ^0 is the attached kernel.

4.2.2. Let's consider a parameter T as in § 2.2.12. Following Jacquet-Lapid-Rogawski (see [JLR99]), we introduce the truncation operator Λ_m^T that associates to a function φ on $[G]$ the following function of the variable $h \in [G']$:

$$(4.2.2.1) \quad (\Lambda_m^T \varphi)(h) = \sum_P (-1)^{\dim(\mathfrak{a}_P^G)} \sum_{\delta \in P'(F) \backslash G'(F)} \hat{\tau}_P(H_P(\delta h) - T_P) \varphi_P(\delta h)$$

where the sum is over standard parabolic subgroup of G (those containing B) and φ_P is the constant term along P . Recall that $P' = G' \cap P$.

4.2.3. We shall define the mixed truncated kernel $K_\chi^0 \Lambda_m^T$: the notation means that the mixed truncation is applied to the second variable. This is a function on $G(\mathbb{A}) \times G'(\mathbb{A})$. To begin with we have:

Lemma 4.2.3.1. — *For $(x, y) \in G(\mathbb{A}) \times G'(\mathbb{A})$, we have:*

$$(K_\chi^0 \Lambda_m^T)(x, y) = \sum_{B \subset P} |\mathcal{P}(M_P)|^{-1} \int_{i\mathfrak{a}_P^{G, *}} \sum_{\varphi \in \mathcal{B}_{P, \chi}} E(x, I_P(\lambda, f)\varphi, \lambda) \overline{\Lambda_m^T E(y, \varphi, \lambda)} d\lambda.$$

Proof. — As $y \in G'(\mathbb{A})$, the mixed truncation is defined by a finite sum of constant terms of $K_\chi^0(x, \cdot)$ (in the second variable). The only point is to permute the sum over φ and the operator Λ_m^T . In fact using the continuity properties of Eisenstein series (see [Lap08]) and properties of mixed truncation operator (in particular a variant of lemma 1.4 of [Art80]), we can conclude as in the proof of proposition 2.8.4.1. \square

Lemma 4.2.3.2. — *For any integer N , there exists a continuous semi-norm $\|\cdot\|$ on $\mathcal{S}(G(\mathbb{A}), C, K_0)$ and an integer N' such that for all $X \in \mathcal{U}(\mathfrak{g}_C)$, all $x \in G(\mathbb{A})^1$, $y \in G'(\mathbb{A})^1$ and all $f \in \mathcal{S}(G(\mathbb{A}), C, K_0)$ we have*

$$(4.2.3.2) \quad \sum_{\chi \in \mathfrak{X}(G)} \sum_{B \subset P} |\mathcal{P}(M_P)|^{-1} \int_{i\mathfrak{a}_P^{G, *}} \sum_{\tau \in \hat{K}} \left| \sum_{\varphi \in \mathcal{B}_{P, \chi, \tau}} (R(X)E)(x, I_P(\lambda, f)\varphi, \lambda) \overline{\Lambda_m^T E(y, \varphi, \lambda)} d\lambda \right| \\ \leq \|f\| \|x\|_{[G]}^{N'} \|y\|_{[G]}^{-N}.$$

Proof. — By the basic properties of the mixed truncation operator (see lemma 1.4 of [Art80] and also [LR03] proof of lemma 8.2.1), for any N and N' there exists a finite family $(Y_i)_{i \in I}$ of

elements of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ such the expression (4.2.3.2) is majorized by the sum over $i \in I$ of $\|y\|_{[G]}^{-N}$ times the supremum over $g \in G'(\mathbb{A})^1$ of

$$\|g\|_{[G]}^{-N'} \sum_{\chi \in \mathfrak{X}} \sum_{B \subset P} |\mathcal{P}(M_P)|^{-1} \int_{\mathfrak{ia}_P^{G,*}} \sum_{\tau \in \hat{K}} \left| \sum_{\varphi \in \mathcal{B}_{P,\chi,\tau}} (R(X)E)(x, I_P(\lambda, f)\varphi, \lambda) \overline{R(Y_i)E(g, \varphi, \lambda)} \right| d\lambda.$$

Then the lemma is a straightforward consequence of lemma 2.9.9.1. \square

Proposition 4.2.3.3. — *For all $x \in G(\mathbb{A})$ and $\chi \in \mathfrak{X}(G)$, we have*

$$\int_{[G']_0} (K_{\chi}^0 \Lambda_m^T)(x, y) dy = \sum_{B \subset P} |\mathcal{P}(M_P)|^{-1} \int_{\mathfrak{ia}_P^{G,*}} \sum_{\varphi \in \mathcal{B}_{P,\chi}} E(x, I_P(\lambda, f)\varphi, \lambda) \overline{\int_{[G']_0} \Lambda_m^T E(y, \varphi, \lambda) dy} d\lambda.$$

Proof. — First one decomposes the sum over $\mathcal{B}_{P,\chi}$ as a sum over $\tau \in \hat{K}$ of finite sums over $\mathcal{B}_{P,\chi,\tau}$. Then, by the majorization of lemma 4.2.3.2 we can permute the integration over $[G']_0$ (which amounts to integrating over $[G']^1$) and the other sums or integrations in the expression we get in lemma 4.2.3.1. \square

4.3 The case of *-generic cuspidal data

4.3.1. We shall use the notations of section 4.2.

4.3.2. *-Generic cuspidal datum. — We shall say that a cuspidal datum $\chi \in \mathfrak{X}(G)$ is *-generic if for any representative (M, π) of χ and $w \in W(M)$ such that $w\pi$ is isomorphic to π or π^* we have $w = 1$. Let's denote by $\mathfrak{X}^*(G)$ the subset of *-generic cuspidal data.

With the notations of 4.1.4, we see that (M, π) is *-generic if and only if for all $1 \leq i, j \leq r$ such that $n_i = n_j$ one of the equalities $\sigma_i = \sigma_j$ or $\sigma_i = \sigma_j^*$ implies that $i = j$.

4.3.3. The next theorem is the main result of the section.

Theorem 4.3.3.1. — *Let $f \in \mathcal{S}(G(\mathbb{A}))$, let $\chi \in \mathfrak{X}(G)$ and let K_{χ} be the associated kernel. For any $g \in G(\mathbb{A})$, one has:*

1. *We have*

$$(4.3.3.1) \quad \int_{[G']} K_{\chi}(g, h) dh = \frac{1}{2} \int_{[G']_0} K_{\chi}^0(g, h) dh$$

where both integrals are absolutely convergent.

2. *If moreover $\chi \in \mathfrak{X}^*(G)$, we have, for any representative (M_P, π) of χ (where P is a standard parabolic subgroup of G),*

$$\int_{[G']} K_{\chi}(g, h) dh = 2^{-\dim(\mathfrak{a}_P)} J_{P,\pi}(g, f)$$

where one defines (see (4.1.8.4))

$$J_{P,\pi}(g, f) = J_{P,\pi}(g, 0, f).$$

In particular, the integral vanishes unless π is self conjugate dual and $M_{P'}$ -distinguished where $P' = G' \cap P$.

The assertion 1 follows readily from lemma 2.9.8.1, Fubini's theorem and the fact that the Haar measure on A_G^∞ is twice the Haar measure on $A_{G'}^\infty$ (see remark 3.1.3.1). The rest of the section is devoted to the proof of assertion 2 of theorem 4.3.3.1. The main steps are propositions 4.3.4.1 and 4.3.7.1.

4.3.4. A limit formula. — We shall use the notation $\lim_{T \rightarrow +\infty} f(T)$ to denote the limit of $f(T)$ when $\langle \alpha, T \rangle \rightarrow +\infty$ for all $\alpha \in \Delta_B$.

Proposition 4.3.4.1. — *Under the assumptions of theorem 4.3.3.1 (but with no genericity condition on χ), we have*

$$\lim_{T \rightarrow +\infty} \int_{[G']_0} (K_\chi^0 \Lambda_m^T)(g, h) dh = 2 \int_{[G']} K_\chi(g, h) dh.$$

Proof. — Let's denote $F^{G'}(\cdot, T)$ the function defined by Arthur relative to G' and its maximal compact subgroup K' (see [Art78] §6 and [Art85] lemma 2.1). It is the characteristic function of a compact of $[G']_0$. Using the fact that $h \mapsto K_\chi^0(g, h)$ is of uniform moderate growth (see lemma 2.9.8.1), we can conclude by a variant of [Art85] theorem 3.1 (see also in the same spirit [IY15] proposition 3.8) that

$$\lim_{T \rightarrow +\infty} \int_{[G']_0} (F^{G'}(h, T) K_\chi^0(g, h) - (K_\chi^0 \Lambda_m^T)(g, h)) dh = 0.$$

We have $\lim_{T \rightarrow +\infty} F^{G'}(h, T) = 1$. Thus we deduce by Lebesgue's theorem and the absolute convergence of the right-hand side of (4.3.3.1).

$$\lim_{T \rightarrow +\infty} \int_{[G']_0} F^{G'}(h, T) K_\chi^0(g, h) dh = \int_{[G']_0} K_\chi^0(g, h) dh$$

The proposition follows by (4.3.3.1). □

4.3.5. Let $\chi \in \mathfrak{X}^*(G)$. Let \mathcal{P}_χ be the set of standard parabolic subgroups such that there exists a cuspidal automorphic representation π of M_P such that (M_P, π) in the equivalence class defined by χ . Since $\chi \in \mathfrak{X}^*(G)$, the space $\mathcal{A}_{P, \chi}(G)$ is non-zero only if $P \in \mathcal{P}_\chi$. Let P be a standard parabolic subgroup and let (M_P, π) be a pair in χ . For any $P_1 \in \mathcal{P}_\chi$, by multiplicity-one theorem, we have

$$\mathcal{A}_{P_1, \chi}(G) = \bigoplus_{w \in W(P, P_1)} \mathcal{A}_{P, w\pi}.$$

In the following we set $M_1 = M_{P_1}$.

Let $P_1 \in \mathcal{P}_\chi$ and $g \in G(\mathbb{A})$. With the notations of section 4.1 (see eq. (4.1.8.4)), for all $Q \in \mathcal{P}(M_1)$, all $\lambda \in i\mathfrak{a}_{P_1, \mathbb{R}}^{G, *}$ we define

$$J_{Q, \chi}(g, \lambda, f) = \sum_{w \in W(P, P_1)} J_{Q, w\pi}(g, \lambda, f).$$

It's a continuous linear form on $\mathcal{S}(G(\mathbb{A}))$.

4.3.6. Proposition 4.3.6.1. — *For all $\chi \in \mathfrak{X}^*(G)$ and all $g \in G(\mathbb{A})$, we have*

$$(4.3.6.2) \quad \int_{[G']_0} (K_\chi^0 \Lambda_m^T)(g, h) dh = \frac{2^{-\dim(\mathfrak{a}_P^G)}}{|\mathcal{P}(M_P)|} \sum_{P_1 \in \mathcal{P}_\chi} \int_{i\mathfrak{a}_{P_1}^{G, *}} \sum_{Q \in \mathcal{P}(M_1)} J_{Q, \chi}(g, \lambda, f) \frac{\exp(-\langle \lambda, T_Q \rangle)}{\theta_Q(-\lambda)} d\lambda.$$

Proof. — This is an obvious consequence of the definitions, the proposition 4.2.3.3 and the lemma 4.3.6.2 below. \square

Lemma 4.3.6.2. — *Let $\chi \in \mathfrak{X}^*(G)$. Let $P_1 \in \mathcal{P}_\chi$ and $\varphi \in \mathcal{A}_{P_1, \chi}$. We have for all $\lambda \in i\mathfrak{a}_{P_1}^G$*

$$\int_{[G']_0} \Lambda_m^T E(y, \varphi, \lambda) dy = 2^{-\dim(\mathfrak{a}_{P_1}^G)} \sum_{Q \in \mathcal{P}(M_1)} J_Q(\varphi, \lambda) \frac{\exp(\langle \lambda, T_Q \rangle)}{\theta_Q(\lambda)}.$$

Proof. — This is simply a rephrasing in our particular situation of a key result of Jacquet-Lapid-Rogawski (see [JLR99] theorem 40). Indeed, because χ is $*$ -generic, theorem 40 of *ibid.* can be stated as:

$$\int_{[G']_0} \Lambda_m^T E(y, \varphi, \lambda) dy = 2^{-\dim(\mathfrak{a}_{P_1}^G)} \sum_{(Q, w)} J(M(w, \lambda)\varphi) \frac{\exp(\langle (w\lambda)_Q, T \rangle)}{\theta_Q(w\lambda)}$$

where the sum is over pair (Q, w) where Q is a standard parabolic subgroup and $w \in W(P_1, Q)$. \square

4.3.7. Proposition 4.3.7.1. — *Let $\chi \in \mathfrak{X}^*(G)$ and let (M_P, π) be a representative where P is a standard parabolic subgroup of G . We have:*

$$\lim_{T \rightarrow +\infty} \int_{[G']} (K_\chi^0 \Lambda_m^T)(g, h) dh = 2^{-\dim(\mathfrak{a}_P^G)} J_{P, \pi}(g, f)$$

where one defines

$$(4.3.7.3) \quad J_{P, \pi}(g, f) = J_{P, \pi}(g, 0, f).$$

Proof. — We start from the expansion (4.3.6.2) of proposition 4.3.6.1. For each $P_1 \in \mathcal{P}_\chi$, let $M_1 = M_{P_1}$. The family $(J_{Q, \chi}(g, \lambda, f))_{Q \in \mathcal{P}(M_1)}$ is a (G, M_1) -family of Schwartz functions on $i\mathfrak{a}_{P_1}^{G, *}$: this is a straightforward consequence of propositions 4.1.9.1 and 4.1.10.1. By [Lap11] Lemma 8, we have:

$$\lim_{T \rightarrow +\infty} \int_{i\mathfrak{a}_{P_1}^{G, *}} \sum_{Q \in \mathcal{P}(M_1)} J_{Q, \chi}(g, \lambda, f) \frac{\exp(-\langle \lambda, T_Q \rangle)}{\theta_Q(-\lambda)} d\lambda = J_{P_1, \chi}(g, 0, f)$$

By definition and lemma 4.3.7.2 below, one has:

$$\begin{aligned} J_{P_1, \chi}(g, 0, f) &= \sum_{w \in W(P, P_1)} J_{P_1, w\pi}(g, 0, f) \\ &= |W(P, P_1)| J_{P, \pi}(g, 0, f). \end{aligned}$$

Since $|\mathcal{P}(M_P)| = \sum_{P_1 \in \mathcal{P}_\chi} |W(P, P_1)|$ we get the expected limit. \square

Lemma 4.3.7.2. — *(Lapid) For any $w \in W(P, P_1)$, we have*

$$J_{P_1, w\pi}(g, 0, f) = J_{P, \pi}(g, 0, f).$$

Proof. — By definition, we have

$$J_{P_1, w\pi}(g, 0, f) = \sum_{\varphi \in \mathcal{B}_{P_1, w\pi}} E(g, \varphi, 0) \cdot J_1(\bar{\varphi})$$

where J_1 is the linear form on $\mathcal{A}_{P_1, w\pi}$ defined in §4.1.5 and $\mathcal{B}_{P_1, w\pi}$ is any K -basis of $\mathcal{A}_{P_1, w\pi}$. Now, the intertwining operator $M(w, 0)$ induces a unitary isomorphism from $\mathcal{A}_{P, \pi}$ to $\mathcal{A}_{P_1, w\pi}$, which sends K -bases to K -bases. Thus one has

$$\begin{aligned} J_{P_1, w\pi}(g, 0, f) &= \sum_{\varphi \in \mathcal{B}_{P, \pi}} E(g, M(w, 0)\varphi, 0) \cdot J_1(M(w, 0)\overline{\varphi}) \\ &= J_{P, \pi}(g, 0, f). \end{aligned}$$

The last equality results from the two equalities:

- $E(g, M(w, 0)\varphi, 0) = E(g, \varphi, 0)$;
- $J_1(M(w, 0)\overline{\varphi}) = J(\overline{\varphi})$ where J is the linear form on $\mathcal{A}_{P, \pi}$ defined in §4.1.5.

The first one is the functional equation of Eisenstein series and the second one is a consequence of case 1 of lemma 8.1 of [Lap06]. \square

5 The *-generic contribution in the Jacquet-Rallis trace formula

The goal of this chapter is to compute the contribution I_χ of the Jacquet-Rallis trace formula for *-generic cuspidal data χ . This is achieved in theorem 5.2.1.1 below. It turns out that for such χ the contribution I_χ is discrete and equal (up to an explicit constant) to a relative character defined in section 5.1 built upon Rankin-Selberg periods of Eisenstein series and Flicker-Rallis intertwining periods.

5.1 Relative characters

5.1.1. We will use the notations of section 3.1.

5.1.2. Let $\chi \in \mathfrak{X}(G)$ be a cuspidal datum and (M, π) be a representative where M is the standard Levi factor of the standard parabolic subgroup P of G . Recall that we have introduced a character $\eta_{G'}$ of $G'(\mathbb{A})$ (see §3.1.8). On $\mathcal{A}_{P, \pi}(G)$, we introduce the linear form J_η defined by

$$(5.1.2.1) \quad J_\eta(\varphi) = \int_{A_{M'} M'(F) N_{P'}(\mathbb{A}) \backslash G(\mathbb{A})} \varphi(g) \eta_{G'}(g) dg, \quad \forall \varphi \in \mathcal{A}_{P, \pi}(G)$$

where $M' = M \cap G'$ and $P' = P \cap G'$. This is a slight variation of that defined in §4.1.5.

We shall say that π is $(M', \eta_{G'})$ -distinguished if J_η does not vanish identically.

5.1.3. Relevant and generic cuspidal data. — We shall say that χ is *relevant* if π is $(M', \eta_{G'})$ -distinguished.

Let $\mathfrak{X}^*(G) = \mathfrak{X}^*(G_n) \times \mathfrak{X}^*(G_{n+1})$ (cf. §4.3.2). We shall say that χ is *-generic if it belongs to the subset $\mathfrak{X}^*(G)$. In particular, if χ is both relevant and generic (see §2.9.6) then it is *-generic.

5.1.4. Rankin-Selberg period of certain Eisenstein series. — Let $T \in \mathfrak{a}_{n+1}^+$. Recall that we have introduced in §3.3.2 the truncation operator Λ_r^T .

Proposition 5.1.4.1. — *Let Q be a parabolic subgroup of G and $Q' = Q \cap G'$. Let π be an irreducible cuspidal representation of M_Q which is $(M_{Q'}, \eta_{G'})$ -distinguished. Let $\varphi \in \mathcal{A}_{Q, \pi}(G)$. Then for a regular point $\lambda \in \mathfrak{a}_Q^{G, *}$ of the Eisenstein series $E(g, \varphi, \lambda)$ (see §2.7.3), the integral*

$$(5.1.4.2) \quad I(\varphi, \lambda) = \int_{[H]} \Lambda_r^T E(h, \varphi, \lambda) dh$$

is convergent and does not depend on T .

Remark 5.1.4.2. — The expression $I(\varphi, \lambda)$ is nothing else but the regularized Rankin-Selberg period of $E(\varphi, \lambda)$ as defined by Ichino-Yamana in [IY15].

Proof. — The convergence follows from proposition 3.3.2.1 and the fact that Eisenstein series are of moderate growth. It remains to prove that the integral does not depend on T . Recall that ι induces an isomorphism from G_n onto H . In the proof, it will be more convenient to work with G_n instead of H . However, by abuse of notations, for any $g \in G_n(\mathbb{A})$ and any function φ on $G(\mathbb{A})$ we shall write $\varphi(g)$ instead of $\varphi(\iota(g))$.

Let $T' \in \mathfrak{a}_{n+1}^+$. By lemma 2.2 of [IY15], we have

$$\Lambda_r^{T+T'} E(g, \varphi, \lambda) = \sum_{P \in \mathcal{F}_{RS}} \sum_{\delta \in (P \cap H)(F) \backslash H(F)} \Lambda_r^{T,P} E_{G_n \times P_{n+1}}(\delta g, \varphi, \lambda) \Gamma'_{P_{n+1}}(H_{P_{n+1}}(\delta g) - T_{P_{n+1}}, T')$$

where the notations are those of §§3.2.3 and 3.3.2. The other notations are borrowed from [Zyd18, eq. (4.4)]; the operator $\Lambda_r^{T,P}$ is the obvious variant of Λ_r^T and Γ'_P is an Arthur function whose precise definition is irrelevant here. We denote by $E_{G_n \times P_{n+1}}$ the constant term of E along $G_n \times P_{n+1}$. Thus, we have

$$\int_{[H]} \Lambda_r^{T+T'} E(g, \varphi, \lambda) dg = \sum_{P \in \mathcal{F}_{RS}} \int_{(P \cap H)(F) \backslash H(\mathbb{A})} (\Lambda_r^{T,P} E_{G_n \times P_{n+1}} P)(g, \varphi, \lambda) \Gamma'_{P_{n+1}}(H_{P_{n+1}}(g) - T_{P_{n+1}}, T') dg.$$

Let $P \in \mathcal{F}_{RS}$ be such that $P \subsetneq G$. It suffices to show that the terms corresponding to P vanish. We identify H with G_n . Then $P \cap H$ is identified with P_n . Let $M_n = M_{P_n}$. For an appropriate choice of a Haar measure on K_n , such a term can be written as

$$\int_{[M_n]} \int_{K_n} \exp(-\langle 2\rho_{P_n}, H_{P_n}(m) \rangle) (\Lambda_r^{T,P} E_P)(mk, \varphi, \lambda) \Gamma'_{P_{n+1}}(H_{P_{n+1}}(m) - T_{P_{n+1}}, T') dk dm,$$

where E_P denotes the constant term of E along $P = P_n \times P_{n+1}$. At this point, we may and shall assume that P is standard (if not, we may change B_n by a conjugate for the arguments). We have the usual formula for the constant term

$$E_P(m, \varphi, \lambda) = \sum_{w \in W(Q; P)} E^P(m, M(w, \lambda) \varphi, w \lambda).$$

where $W(Q; P)$ is the set of elements $w \in W$ that are of minimal length in double cosets $W^P w W^Q$. Let $w \in W(Q; P)$. Notice that the representation $w\pi$ is also $(wM_{Q'} w^{-1}, \eta_{G'})$ -distinguished. For the argument, we may and shall assume $w = 1$ (that is we assume that $Q \subset P$). Thus it suffices to show for all $k \in K$ the integral

$$(5.1.4.3) \quad \int_{[M_n]^1} \Lambda_r^{T,P} E^P(mk, \varphi, \lambda) dm$$

vanishes.

The group $M_{n+1} = M_{P_{n+1}}$ has a decomposition $G_{d_1} \times \dots \times G_{d_r}$ with $d_1 + \dots + d_r = n+1$. Each factor corresponds to a subset of the canonical basis (e_1, \dots, e_{n+1}) . We may assume that the factor G_{d_1} corresponds to a subset which does not contain e_{n+1} . As a consequence G_{d_1} is also a factor of M_n . We view $G_{d_1} \times G_{d_1}$ as a subgroup of $M_n \times M_{n+1}$. Let $Q_1 \times Q_2 = (G_{d_1} \times G_{d_1}) \cap Q \subset G_n \times G_{n+1}$. The representation π restricts to $M_{Q_1}(\mathbb{A})$ and $M_{Q_2}(\mathbb{A})$: this gives representations respectively denoted by π_1 and π_2 . As a factor of (5.1.4.3), we get

$$(5.1.4.4) \quad \int_{[G_{d_1}]^1} E(g, \varphi_1, \lambda_1) \Lambda^T E(g, \varphi_2, \lambda_2) dg$$

where $\varphi_i \in \mathcal{A}_{Q_i, \pi_i}(G_{d_1})$. Here the truncation is the usual Arthur's truncation operator on the group G_{d_1} . It is clear from Langlands' formula for the integral (5.1.4.4) (see [Art82]) that (5.1.4.4) vanishes unless there exists $w \in W(Q_1, Q_2)$ such that $\pi_2 \simeq w\pi_1$. But then π_2 would be $(M_{Q'_2}, (\eta_{d_1})^n)$ -distinguished and $(M_{Q'_2}, (\eta_{d_1})^{n+1})$ -distinguished with $\eta_{d_1} = \eta \circ \det_{d_1}$ and $M_{Q'_2} = M_{Q_2} \cap G'_{d_1}$. This is not possible. \square

5.1.5. Relative characters. — Let (P, π) be a pair for which P be a standard parabolic subgroup of G and π be a cuspidal automorphic representation of its standard Levi factor M_P . For any $\varphi \in \mathcal{A}_{P, \pi}(G)$, building upon the truncation operator Λ_r^T and the linear form J_η , we define the relative character $I_{P, \pi}^T$ for any $f \in \mathcal{S}(G(\mathbb{A}))$ by

$$I_{P, \pi}^T(f) = \sum_{\varphi \in \mathcal{B}_{P, \pi}} \int_{[H]} \Lambda_r^T E(h, I_P(0, f)\varphi, 0) dh \cdot \overline{J_\eta(\varphi)}$$

where the K -basis $\mathcal{B}_{P, \pi}$ is defined in §2.8.3. Using proposition 5.1.4.1, we have

$$I_{P, \pi}^T(f) = I_{P, \pi}(f)$$

where we define:

$$I_{P, \pi}(f) = \begin{cases} \sum_{\varphi \in \mathcal{B}_{P, \pi}} I(I_P(0, f)\varphi, 0) \cdot \overline{J_\eta(\varphi)} & \text{if } \pi \text{ is } (M_{P'}, \eta_{G'})\text{-distinguished;} \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 5.1.5.1. — *Let $\chi \in \mathfrak{X}^*(G)$. Let (P, π) be a representative. The map $f \mapsto I_{P, \pi}^T(f)$ (and thus $f \mapsto I_{P, \pi}(f)$) is well-defined and gives a continuous linear form on $\mathcal{S}(G(\mathbb{A}))$. It depends only on χ and not on the choice of (P, π) .*

Proof. — First we claim that $\varphi \mapsto \int_{[H]} \Lambda_r^T E(h, \varphi, 0) dh$ is a continuous map: this is an easy consequence of properties of Eisenstein series and the truncation operator Λ_r^T (see the proposition 3.3.2.1). On the other hand $\varphi \mapsto J_\eta(\varphi)$ is also continuous (see section 4.1). Thus the first assertion results from an application of proposition 2.8.4.1. The arguments of the proof of lemma 4.3.7.2 give the independence on the choice of (P, π) . \square

5.2 The *-generic contribution

5.2.1. Let $\chi \in \mathfrak{X}(G)$. Recall that we defined in theorem 3.2.4.1 a distribution I_χ on $\mathcal{S}(G(\mathbb{A}))$. Let (M, π) be a representative of χ where M is the standard Levi factor of the standard parabolic subgroup P of G . The following theorem is the main result of this chapter.

Theorem 5.2.1.1. — *Assume moreover $\chi \in \mathfrak{X}^*(G)$. We have*

$$I_\chi = 2^{-\dim(\mathfrak{a}_P)} I_{P, \pi}.$$

In particular, we have $I_\chi = 0$ unless χ is relevant.

The theorem is a direct consequence of the following proposition.

Proposition 5.2.1.2. — *Assume moreover $\chi \in \mathfrak{X}^*(G)$. We have for $T \in \mathfrak{a}_{n+1}^+$*

$$(5.2.1.1) \quad \int_{[H]} \int_{[G']} \Lambda_r^T K_\chi(x, y) \eta_{G'}(y) dx dy = 2^{-\dim(\mathfrak{a}_P)} I_{P, \pi}(f),$$

where the left-hand side is absolutely convergent (see proposition 3.3.3.1). In particular, the left-hand side does not depend on T .

Indeed, by theorem 3.3.7.1, I_χ is the constant term in the asymptotic expansion in T of the left-hand side of (5.2.1.1) hence $I_\chi = 2^{-\dim(\mathfrak{a}_P)} I_{P, \pi}$.

The rest of the section is devoted to the proof of proposition 5.2.1.2.

5.2.2. Proof of proposition 5.2.1.2. — We assume that $\chi \in \mathfrak{X}^*(G)$. The proof is a straightforward consequence of theorem 4.3.3.1 and some permutations between integrals, summations and the truncation. These permutations are provided by lemmas 5.2.2.1 and 5.2.2.3 below.

Lemma 5.2.2.1. — *For all $x \in [H]$, we have*

$$\int_{[G']} (\Lambda_r^T K_\chi)(x, y) \eta_{G'}(y) dx dy = \Lambda_r^T \left(\int_{[G']} K_\chi(\cdot, y) \eta_{G'}(y) dy \right) (x).$$

Remark 5.2.2.2. — On the left-hand side we apply the truncation operator Λ_r^T to the function $K_\chi(\cdot, y)$ (where y is fixed) and then we evaluate at x whereas on the right-hand side we apply the same operator to the function we get by integration of $K_\chi(\cdot, y) \eta_{G'}(y)$ over $y \in [G']$ and then we evaluate at x .

Proof. — Since x is fixed, the operator Λ_r^T is a finite sum of constant terms (see [Art78] lemma 5.1 for the finiteness). Then the lemma follows from Fubini's theorem which holds because we have

$$\int_{[N_Q]} \int_{[G']} |K_\chi(nx, y)| dndy < \infty$$

for all parabolic subgroups Q of G_{n+1} containing B_n . Here we identify N_Q with the subgroup $\{1\} \times N_Q$ of $G = G_n \times G_{n+1}$. The convergence of the integral results from the bound (3.3.3.3) above. \square

Lemma 5.2.2.3. — *We have*

$$\int_{[H]} \Lambda_r^T \left(\int_{[G']} K_\chi(\cdot, y) \eta_{G'}(y) dy \right) (h) dh = 2^{-\dim(\mathfrak{a}_P)} I_{P, \pi}(f).$$

Proof. — First, by theorem 4.3.3.1, we have for any $x \in [G]$:

$$\int_{[G']} K_\chi(x, y) \eta_{G'}(y) dy = 2^{-\dim(\mathfrak{a}_P)} \sum_{\varphi \in \mathcal{B}_{P, \pi}} E(x, I_P(0, f)\varphi, 0) dh \cdot \overline{J_\eta(\varphi)}$$

where the notations are borrowed from §5.1.5. Then we want to apply the truncation operator Λ_r^T and evaluate at $h \in [H]$. We want to show that this operation commutes with the summation over the orthonormal basis. As in the proof of lemma 5.2.2.1, it suffices to prove

$$\sum_{\varphi \in \mathcal{B}_{P, \pi}} \int_{[N_Q]} |E(ng, I_P(0, f)\varphi, 0)| dn \cdot |\overline{J_\eta(\varphi)}| < \infty$$

for any parabolic subgroups Q of G_{n+1} containing B_n , which is an easy consequence of continuity properties of Eisenstein series.

In this way, we get for $h \in [H]$:

$$\Lambda_r^T \left(\int_{[G']} K_\chi(\cdot, y) \eta_{G'}(y) dy \right) (h) = 2^{-\dim(\mathfrak{a}_P)} \sum_{\varphi \in \mathcal{B}_{P, \pi}} (\Lambda_r^T E)(h, I_P(0, f)\varphi, 0) \cdot \overline{J_\eta(\varphi)}.$$

By integration over $h \in [H]$, we have:

$$\int_{[H]} \Lambda_r^T \left(\int_{[G']} K_\chi(\cdot, y) \eta_{G'}(y) dy \right) (h) dh = 2^{-\dim(\mathfrak{a}_P)} \sum_{\varphi \in \mathcal{B}_{P, \pi}} \int_{[H]} (\Lambda_r^T E)(h, I_P(0, f)\varphi, 0) dh \cdot \overline{J_\eta(\varphi)}.$$

The right-hand side is nothing else but $2^{-\dim(\mathfrak{a}_P)} I_{P,\pi}(f)$. Still we have to justify the change of order of the integration and the summation. But it is easy to show that

$$\sum_{\varphi \in \mathcal{B}_{P,\pi}} \int_{[H]} |\Lambda_r^T E(h, I_P(0, f)\varphi, 0)| dh \cdot |\overline{J_\eta(\varphi)}| < \infty.$$

□

6 Spectral decomposition of the Flicker-Rallis period for certain cuspidal data

The goal of this chapter is to give another proof of the spectral decomposition of the Flicker-Rallis period for the same cuspidal data as in Section 4.3.2. The main result of this chapter (obtained as a combination of Theorem 6.2.5.1 and Theorem 6.2.6.1) can be used to get another version of Theorem 4.3.3.1 with a seemingly different relative character than $J_{P,\pi}$ (this will actually be done in §8.2.4). Of course, these two relative characters are the same. A direct proof of this fact will be given in Chapter 9.

6.1 Notation

6.1.1. In this chapter we adopt the set of notation introduced in Section 3.1: E/F is a quadratic extension of number fields, $G'_n = \mathrm{GL}_{n,F}$, $G_n = \mathrm{Res}_{E/F} \mathrm{GL}_{n,E}$, (B'_n, T'_n) , (B_n, T_n) are the standard Borel pairs of G'_n , G_n and K'_n , K_n the standard maximal compact subgroups of $G'_n(\mathbb{A})$, $G_n(\mathbb{A})$ respectively. Besides, we denote by N'_n , N_n the unipotent radicals of B'_n , B_n and we set

$$w_n = \begin{pmatrix} & & 1 \\ & \ddots & \\ & & \\ 1 & & \end{pmatrix} \in G'_n(F).$$

We write $e_n = (0, \dots, 0, 1)$ for the last element in the standard basis of F^n and we let $\mathcal{P}_n = \begin{pmatrix} \star & \star \\ 0 \dots 0 & 1 \end{pmatrix}$, $\mathcal{P}'_n = \mathcal{P}_n \cap G'_n$ be the mirabolic subgroups of G_n , G'_n respectively (that is the stabilizers of e_n for the natural right actions). The unipotent radicals of \mathcal{P}_n , \mathcal{P}'_n will be denoted by U_n and U'_n respectively. For nonnegative integers $m \leq n$, we embed G_m in G_n (resp. G'_m in G'_n) in the “upper left corner” by $g \mapsto \begin{pmatrix} g & \\ & I_{n-m} \end{pmatrix}$. Thus, in particular, we have $\mathcal{P}_n = G_{n-1}U_n$ and $\mathcal{P}'_n = G'_{n-1}U'_n$.

The entries of a matrix $g \in G_n(\mathbb{A})$ are written as $g_{i,j}$, $1 \leq i, j \leq n$, and the diagonal entries of an element $t \in T_n(\mathbb{A})$ as t_i , $1 \leq i \leq n$.

6.1.2. We fix a nontrivial additive character $\psi' : \mathbb{A}/F \rightarrow \mathbb{C}^\times$. For $\phi \in \mathcal{S}(\mathbb{A}^n)$, we define its Fourier transform $\hat{\phi} \in \mathcal{S}(\mathbb{A}^n)$ by

$$\hat{\phi}(x_1, \dots, x_n) = \int_{\mathbb{A}^n} \phi(y_1, \dots, y_n) \psi'(x_1 y_1 + \dots + x_n y_n) dy_1 \dots dy_n$$

the Haar measure on \mathbb{A}^n being chosen such that $\hat{\hat{\phi}}(x) = \phi(-x)$.

We denote by c the nontrivial Galois involution of E over F . Then, c acts naturally on $G_n(\mathbb{A})$ and thus on cuspidal automorphic representations of the latter. We denote this action by $\pi \mapsto \pi^c$. We fix $\tau \in E^\times$ such that $\tau^c = -\tau$ and we define $\psi : \mathbb{A}_E/E \rightarrow \mathbb{C}^\times$ by $\psi(z) = \psi'(\mathrm{Tr}_{E/F}(\tau z))$, $z \in \mathbb{A}_E$, where \mathbb{A}_E denotes the adèle ring of E and $\mathrm{Tr}_{E/F} : \mathbb{A}_E \rightarrow \mathbb{A}$ the trace map. We also define a generic character $\psi_n : [N_n] \rightarrow \mathbb{C}^\times$ by

$$\psi_n(u) = \psi \left((-1)^n \sum_{i=1}^{n-1} u_{i,i+1} \right), \quad u \in [N_n].$$

(The appearance of the sign $(-1)^n$ is only a convention that will be justified a posteriori in Chapter 7). Note that ψ is trivial on \mathbb{A} and therefore ψ_n is trivial on $N'_n(\mathbb{A})$. To any $f \in \mathcal{T}([G_n])$, we associate its *Whittaker function* W_f defined by

$$W_f(g) = \int_{[N_n]} f(ug) \psi_n(u)^{-1} du, \quad g \in G_n(\mathbb{A}).$$

6.2 Statements of the main results

6.2.1. Let $n \geq 1$ be a nonnegative integer. For $f \in \mathcal{T}([G_n])$, $\phi \in \mathcal{S}(\mathbb{A}^n)$ and $s \in \mathbb{C}$ we set

$$Z_{\psi}^{\text{FR}}(s, f, \phi) = \int_{N'_n(\mathbb{A}) \backslash G'_n(\mathbb{A})} W_f(h) \phi(e_n h) |\det h|^s dh$$

provided this expression converges absolutely.

6.2.2. Let $\chi \in \mathfrak{X}^*(G_n)$ be a $*$ -generic cuspidal datum (see §4.3.2 for the definition of $*$ -generic) represented by a pair (M_P, π) and set $\Pi = I_{P(\mathbb{A})}^{G_n(\mathbb{A})}(\pi)$. We can write

$$M_P = G_{n_1} \times \dots \times G_{n_k}$$

where n_1, \dots, n_k are positive integers such that $n_1 + \dots + n_k = n$. Then, π decomposes accordingly as a tensor product

$$\pi = \pi_1 \boxtimes \dots \boxtimes \pi_k$$

where for each $1 \leq i \leq k$, π_i is a cuspidal automorphic representation of $G_{n_i}(\mathbb{A})$.

6.2.3. Let $L(s, \Pi, \text{As})$ be the Shahidi's completed Asai L -function of Π [Sha90], [Gol94]. We have the decomposition

$$L(s, \Pi, \text{As}) = \prod_{i=1}^k L(s, \pi_i, \text{As}) \times \prod_{1 \leq i < j \leq k} L(s, \pi_i \times \pi_j^c).$$

As χ is $*$ -generic, the Rankin-Selberg L -functions $L(s, \pi_i \times \pi_j^c)$ are entire and non-vanishing at $s = 1$ [JS81b], [JS81a], [Sha81] whereas by [Fli88], $L(s, \pi_i, \text{As})$ has at most a simple pole at $s = 1$. Therefore, $L(s, \Pi, \text{As})$ has a pole of order at most k at $s = 1$ and this happens if and only if $L(s, \pi_i, \text{As})$ has a pole at $s = 1$ for every $1 \leq i \leq k$.

We say that the cuspidal datum χ is *distinguished* if $L(s, \Pi, \text{As})$ has a pole of order k at $s = 1$. By [Fli88], it is equivalent to ask π to be $M_{P'} = M_P \cap G'_n$ -distinguished.

6.2.4. For $f \in \mathcal{C}([G_n])$, we set $W_{f, \Pi} = W_{f_{\Pi}}$ where f_{Π} is defined as in Section 2.9.7. Then, $W_{f, \Pi}$ belongs to the Whittaker model $\mathcal{W}(\Pi, \psi_n)$ of Π with respect to ψ_n .

We define a continuous linear form β_n on $\mathcal{W}(\Pi, \psi_n)$ as follows. For S a finite set of places of F and $W \in \mathcal{W}(\Pi, \psi_n)$, we set

$$\beta_{n, S}(W) = \int_{N'_n(F_S) \backslash P'_n(F_S)} W(p_S) dp_S$$

the integral being convergent by (the same proof as) [BP18a, Proposition 2.6.1, Lemma 3.3.1] and the Jacquet-Shalika bound [JS81b]. By [Fli88, Proposition 3] and (2.3.2.3), for a given $W \in \mathcal{W}(\Pi, \psi_n)$, the quantity

$$\beta_n(W) = (\Delta_{G'}^{S, *})^{-1} L^{S, *}(1, \Pi, \text{As}) \beta_{n, S}(W)$$

is independent of S as long as it is sufficiently large (i.e. it contains all the Archimedean places as well as the non-Archimedean places where the situation is “ramified”). This defines the linear form β_n .

6.2.5. For every $f \in \mathcal{C}([G_n])$, we set

$${}^0 f(g) = \int_{A_{G_n}^{\infty}} f(ag) da, \quad g \in [G_n].$$

Theorem 6.2.5.1. —

1. Let $N \geq 0$. There exists $c_N > 0$ such that for every $f \in \mathcal{T}_N([G_n])$ and $\phi \in \mathcal{S}(\mathbb{A}^n)$, the expression defining $Z_\psi^{\text{FR}}(s, f, \phi)$ is absolutely convergent for $s \in \mathcal{H}_{>c_N}$ and the function $s \in \mathcal{H}_{>c_N} \mapsto Z_\psi^{\text{FR}}(s, f, \phi)$ is holomorphic and bounded in vertical strips. Moreover, for every $s \in \mathcal{H}_{>c_N}$, $(f, \phi) \mapsto Z_\psi^{\text{FR}}(s, f, \phi)$ is a (separately) continuous bilinear form on $\mathcal{T}_N([G_n]) \times \mathcal{S}(\mathbb{A}^n)$.
2. Let $\chi \in \mathfrak{X}^*(G_n)$. For every $f \in \mathcal{C}_\chi([G_n])$, the function $s \mapsto (s-1)Z_\psi^{\text{FR}}(s, {}^0f, \phi)$ admits an analytic continuation to $\mathcal{H}_{>1}$ with a limit at $s=1$. Moreover, we have

$$Z_\psi^{\text{FR},*}(1, {}^0f, \phi) := \lim_{s \rightarrow 1^+} (s-1)Z_\psi^{\text{FR}}(s, {}^0f, \phi) = \begin{cases} 2^{1-k} \widehat{\phi}(0) \beta_n(W_{f,\Pi}) & \text{if } \chi \text{ is distinguished,} \\ 0 & \text{otherwise.} \end{cases}$$

6.2.6. Theorem 6.2.6.1. — Let $\chi \in \mathfrak{X}^*(G_n)$. The linear form

$$P_{G'_n} : f \in \mathcal{C}([G_n]) \mapsto \int_{[G'_n]} f(h) dh$$

is well-defined (i.e. the integral converges) and continuous. Moreover, for every $f \in \mathcal{S}_\chi([G_n])$ and $\phi \in \mathcal{S}(\mathbb{A}^n)$ such that $\widehat{\phi}(0) = 1$ we have

$$(6.2.6.1) \quad P_{G'_n}(f) = \frac{1}{2} Z_\psi^{\text{FR},*}(1, {}^0f, \phi).$$

6.2.7. A direct consequence of Theorem 6.2.5.1 and Theorem 6.2.6.1 is the following corollary.

Corollary 6.2.7.1. — Let $\chi \in \mathfrak{X}^*(G_n)$ be represented by a pair (M_P, π) and set $\Pi = I_{P(\mathbb{A})}^{G_n(\mathbb{A})}(\pi)$. Then, for every $f \in \mathcal{S}_\chi([G_n])$ we have

$$P_{G'_n}(f) = \begin{cases} 2^{-\dim(A_P)} \beta_n(W_{f,\Pi}) & \text{if } \chi \text{ is distinguished,} \\ 0 & \text{otherwise.} \end{cases}$$

6.3 Proof of Theorem 6.2.5.1.2

Part 1. of Theorem 6.2.5.1 will be established in Section 6.5. Here, we give the proof of part 2. of this theorem. Let $f \in \mathcal{C}_\chi([G_n])$, $\phi \in \mathcal{S}(\mathbb{A}^n)$ and (M_P, π) be a pair representing the cuspidal datum χ as in Section 6.2. Set

$$\mathcal{A} := (i\mathbb{R})^k$$

and let \mathcal{A}_0 be the subspace of $\underline{x} = (x_1, \dots, x_k) \in \mathcal{A}$ such that $x_1 + \dots + x_k = 0$. We equip \mathcal{A} with the product of Lebesgue measures and \mathcal{A}_0 with the unique Haar measure such that the quotient measure on

$$\mathcal{A}/\mathcal{A}_0 \simeq i\mathbb{R}, (x_1, \dots, x_k) \mapsto x_1 + \dots + x_k$$

is again the Lebesgue measure.

There is an unique identification $\mathcal{A} \simeq i\mathfrak{a}_P^*$ which when it is composed with the map $\mu \in i\mathfrak{a}_P^* \mapsto \pi_\mu$ gives

$$(6.3.0.1) \quad \underline{x} = (x_1, \dots, x_k) \in \mathcal{A} \mapsto \pi_{\underline{x}} := \pi_1 |\det|_E^{x_1/n_1} \boxtimes \dots \boxtimes \pi_k |\det|_E^{x_k/n_k}.$$

For every $\underline{x} \in \mathcal{A}$, we set $\Pi_{\underline{x}} = I_{P(\mathbb{A})}^{G_n(\mathbb{A})}(\pi_{\underline{x}})$ and $f_{\underline{x}} = f_{\Pi_{\underline{x}}}$ following the definition of Section 2.9.7 (so that in particular $\Pi_0 = \Pi$ and $f_0 = f_{\Pi}$ with notation from the previous section).

The isomorphism (6.3.0.1) sends $\prod_{j=1}^k (in_j \mathbb{Z})$ onto $iX^*(P)$ hence, by (2.3.1.1), it also sends the measure on \mathcal{A} to $(n_1 \dots n_k)(2\pi)^k$ times the measure on $i\mathfrak{a}_P^*$. Therefore, by Theorem 2.9.7.1, we have

$${}^0f = \frac{(2\pi)^{-k}}{n_1 \dots n_k} \int_{A_{G_n}^\infty} \int_{\mathcal{A}} a \cdot f_{\underline{x}} d\underline{x} da = \frac{(2\pi)^{-k}}{n_1 \dots n_k} \int_{A_{G_n}^\infty} \int_{\mathcal{A}} |\det a|_E^{\frac{x_1 + \dots + x_k}{n}} f_{\underline{x}} d\underline{x} da.$$

We have an isomorphism $A_{G_n}^\infty \simeq \mathbb{R}_+^*$, $a \mapsto |\det a|_E$, sending the Haar measure on $A_{G_n}^\infty$ to $\frac{dt}{|t|}$ where dt is the Lebesgue measure. Thus, by Fourier inversion, the previous equality can be rewritten as

$$(6.3.0.2) \quad {}^0f = \frac{n}{n_1 \dots n_k} (2\pi)^{-k+1} \int_{\mathcal{A}_0} f_{\underline{x}} d\underline{x}$$

where the right-hand side is an absolutely convergent integral in $\mathcal{T}_N([G_n])$ for some $N > 0$. Therefore, by the first part of Theorem 6.2.5.1, there exists $c > 0$ such that for every $s \in \mathcal{H}_{>c}$ we have

$$(6.3.0.3) \quad Z_\psi^{\text{FR}}(s, {}^0f, \phi) = \frac{n}{n_1 \dots n_k} (2\pi)^{-k+1} \int_{\mathcal{A}_0} Z_\psi^{\text{FR}}(s, f_{\underline{x}}, \phi) d\underline{x}.$$

Let S_0 be a finite set of places of F including the Archimedean ones and outside of which π is unramified and let $S_{0,f} \subset S_0$ be the subset of finite places. Let $I \subseteq \{1, \dots, k\}$ be the subset of $1 \leq i \leq k$ such that $L(s, \pi_i, \text{As})$ has a pole at $s = 1$. We choose, for each $1 \leq i \leq k$ and $v \in S_{0,f}$, polynomials $Q_i(T), Q_{i,v}(T) \in \mathbb{C}[T]$ with roots in $\mathcal{H}_{|0,1[}$ and $\mathcal{H}_{|q_v^{-1},1[}$ respectively such that $s \mapsto Q_i(s)L_\infty(s, \pi_i, \text{As})$ and $s \mapsto Q_{i,v}(q_v^{-s})L_v(s, \pi_i, \text{As})$ have no pole in $\mathcal{H}_{|0,1[}$. Finally, we set

$$P(s, \underline{x}) = \prod_{i \in I} (s + \frac{2x_i}{n_i})(s - 1 + \frac{2x_i}{n_i}) \prod_{1 \leq i \leq k} Q_i(s + \frac{2x_i}{n_i}) \prod_{\substack{1 \leq i \leq k \\ v \in S_{0,f}}} Q_{i,v}(q_v^{-s - \frac{2x_i}{n_i}}) \quad \text{and} \quad \tilde{f}_{\underline{x}}(g) = f_{\underline{x}}({}^t g^{-1})$$

for every $\underline{x} \in \mathcal{A}_0$, $s \in \mathbb{C}$ and $g \in G_n(\mathbb{A})$. We will now check that the functions

$$(6.3.0.4) \quad (s, \underline{x}) \in \mathbb{C} \times \mathcal{A}_0 \mapsto P(s + \frac{1}{2}, \underline{x}) Z_\psi^{\text{FR}}(s + \frac{1}{2}, f_{\underline{x}}, \phi)$$

and

$$(6.3.0.5) \quad (s, \underline{x}) \in \mathbb{C} \times \mathcal{A}_0 \mapsto P(\frac{1}{2} - s, \underline{x}) Z_{\psi^{-1}}^{\text{FR}}(s + \frac{1}{2}, \tilde{f}_{\underline{x}}, \hat{\phi})$$

satisfy the conditions of Corollary A.0.10.1.

From the first part of Theorem 6.2.5.1, Theorem 2.9.7.1 and Lemma A.0.8.1, we deduce that these functions satisfy the first condition of Corollary A.0.10.1. To check that they also satisfy the second condition of Corollary A.0.10.1, we need to analyze more carefully the function $s \mapsto Z_\psi^{\text{FR}}(s, f_{\underline{x}}, \phi)$ for a fixed $\underline{x} \in \mathcal{A}_0$.

For S a sufficiently large finite set of places of F , that we assume to contain Archimedean places as well as the places where π , ψ' or ψ are ramified (thus $S_0 \subset S$), we have decompositions

$$\phi = \phi_S \phi^S \quad \text{and} \quad W_{f_{\underline{x}}} = W_{S, \underline{x}} W_{\underline{x}}^S$$

for every $\underline{x} \in \mathcal{A}_0$, where $\phi_S \in \mathcal{S}(F_S^n)$, ϕ^S is the characteristic function of $(\widehat{\mathcal{O}_F^S})^n$, $W_{S, \underline{x}} \in \mathcal{W}(\Pi_{\underline{x}, S}, \psi_{n, S})$ (that is the Whittaker model of the representation $\Pi_{\underline{x}, S}$ with respect to the character $\psi_{n, S} = \psi_{n|N(F_S)}$) and $W_{\underline{x}}^S \in \mathcal{W}(\Pi_{\underline{x}}^S, \psi_n^S)^{K_n^S}$ is such that $W_{\underline{x}}^S(1) = 1$. By [Fli88, Proposition 3] and (2.3.3.4), we then have

$$(6.3.0.6) \quad Z_\psi^{\text{FR}}(s, f_{\underline{x}}, \phi) = (\Delta_{G_n'}^{S,*})^{-1} L(s, \Pi_{\underline{x}}, \text{As}) \frac{Z_\psi^{\text{FR}}(s, W_{S, \underline{x}}, \phi_S)}{L_S(s, \Pi_{\underline{x}}, \text{As})}$$

for $s \in \mathcal{H}_{>c}$ where we have set

$$Z_\psi^{\text{FR}}(s, W_{S,\underline{x}}, \phi_S) = \int_{N'_n(F_S) \backslash G'_n(F_S)} W_{S,\underline{x}}(h_S) \phi_S(e_n h_S) |\det h_S|^s dh_S.$$

Moreover, by [BP18a, Theorem 3.5.1] the function $Z_\psi^{\text{FR}}(s, W_{S,\underline{x}}, \phi_S)$ extends meromorphically to the complex plane and satisfies the functional equation

$$(6.3.0.7) \quad \frac{Z_{\psi^{-1}}^{\text{FR}}(1-s, \widetilde{W}_{S,\underline{x}}, \widehat{\phi}_S)}{L_S(1-s, (\Pi_{\underline{x}})^\vee, \text{As})} = \epsilon(s, \Pi_{\underline{x}}, \text{As}) \frac{Z_\psi^{\text{FR}}(s, W_{S,\underline{x}}, \phi_S)}{L_S(s, \Pi_{\underline{x}}, \text{As})}$$

where $\widetilde{W}_{S,\underline{x}}(g) = W_{S,\underline{x}}(w_n {}^t g^{-1})$, $\widehat{\phi}_S$ is the (normalized) Fourier transform of ϕ_S with respect to the bicharacter $(\underline{u}, \underline{v}) \mapsto \psi'(u_1 v_1 + \dots + u_n v_n)$ and $\epsilon(s, \Pi_{\underline{x}}, \text{As})$ denotes the global epsilon factor of the Asai L -function $L(s, \Pi_{\underline{x}}, \text{As})$.

By (6.3.0.6), (6.3.0.7) as well as the meromorphic continuation and functional equation of $L(s, \Pi_{\underline{x}}, \text{As})$ [Sha90, Theorem 3.5(4)], we conclude that $Z_\psi^{\text{FR}}(s, f_{\underline{x}}, \phi)$ has a meromorphic continuation to \mathbb{C} satisfying the functional equation

$$(6.3.0.8) \quad Z_{\psi^{-1}}^{\text{FR}}(1-s, \widetilde{f}_{\underline{x}}, \widehat{\phi}) = Z_\psi^{\text{FR}}(s, f_{\underline{x}}, \phi).$$

On the other hand, we have the decomposition

$$L(s, \Pi_{\underline{x}}, \text{As}) = \prod_{i=1}^k L(s + \frac{2x_i}{n_i}, \pi_i, \text{As}) \times \prod_{1 \leq i < j \leq k} L(s + \frac{x_i}{n_i} + \frac{x_j}{n_j}, \pi_i \times \pi_j^c).$$

and, as $\chi \in \mathfrak{X}^*(G_n)$, the Rankin-Selberg L -functions $L(s, \pi_i \times \pi_j^c)$ are entire and bounded in vertical strips [Cog08, Theorem 4.1]. By the Jacquet-Shalika bound [JS81b] and the fact that the gamma function is of exponential decay in vertical strips, $Q_i(s) L_\infty(s, \pi_i, \text{As})$ and $Q_{i,v}(q_v^{-s}) L_v(s, \pi_i, \text{As})$ are holomorphic and bounded in vertical strips of $\mathcal{H}_{>0}$ for each $1 \leq i \leq k$ and $v \in S_{0,f}$. By [FL17, Lemma 5.2], $s \mapsto (s-1) L^{S_0}(s, \pi_i, \text{As})$, for $i \in I$, and $s \mapsto L^{S_0}(s, \pi_i, \text{As})$, for $i \notin I$, are also holomorphic and of finite order in vertical strips of $\mathcal{H}_{>0}$. Therefore, by the definition of P and the functional equation, $P(s, \underline{x}) L(s, \Pi_{\underline{x}}, \text{As})$ is entire and of finite order in vertical strips. By (6.3.0.6), (6.3.0.8) and [BP18a, Theorem 3.5.2], it follows that the functions (6.3.0.4), (6.3.0.5) are entire and of finite order in vertical strips in the first variable i.e. they also satisfy the second condition of Corollary A.0.10.1.

Thus, the conclusion of this corollary is valid and in particular the map

$$s \mapsto \left(\underline{x} \mapsto \prod_{i \in I} (s-1 + \frac{2x_i}{n_i}) Z_\psi^{\text{FR}}(s, f_{\underline{x}}, \phi) \right)$$

induces a holomorphic function $\mathcal{H}_{>1-\epsilon} \rightarrow \mathcal{S}(\mathcal{A}_0)$ for some $\epsilon > 0$. By (6.3.0.3) and [BP18b, Lemma 3.1.1, Proposition 3.1.2], it follows that $s \mapsto Z_\psi^{\text{FR}}(s, {}^0 f, \phi)$ extends analytically to $\mathcal{H}_{>1}$ and that

$$(6.3.0.9) \quad \lim_{s \rightarrow 1^+} (s-1) Z_\psi^{\text{FR}}(s, {}^0 f, \phi) = \begin{cases} 2^{1-k} \lim_{s \rightarrow 1} (s-1)^k Z_\psi^{\text{FR}}(s, f_0, \phi) & \text{if } I = \{1, \dots, k\}, \\ 0 & \text{otherwise} \end{cases}$$

Recall that $I = \{1, \dots, k\}$ if and only if $L(s, \Pi, \text{As})$ has a pole of order $k = rk(A_P)$ at $s = 1$. Moreover, by [BP18a, Lemma 3.3.1] and the Jacquet-Shalika bound [JS81b], the integral defining $Z_\psi^{\text{FR}}(s, W_{S,0}, \phi_S)$ is absolutely convergent in $\mathcal{H}_{>1-\epsilon}$ for some $\epsilon > 0$. Combining this with [BP18b, Lemma 2.16.3] and (6.3.0.6), in the case $I = \{1, \dots, k\}$ identity (6.3.0.9) can be rewritten as

$$\begin{aligned} \lim_{s \rightarrow 1^+} (s-1) Z_\psi^{\text{FR}}(s, {}^0 f, \phi) &= 2^{1-k} (\Delta_{G'_n}^{S,*})^{-1} L^{S,*}(1, \Pi, \text{As}) Z_\psi^{\text{FR}}(1, W_{S,0}, \phi_S) \\ &= 2^{1-k} (\Delta_{G'_n}^{S,*})^{-1} L^{S,*}(1, \Pi, \text{As}) \beta_{n,S}(W_{S,0}) \widehat{\phi}_S(0) \\ &= 2^{1-k} \widehat{\phi}(0) \beta_n(W_{f,\Pi}) \end{aligned}$$

and this ends the proof of Theorem 6.2.5.1.2.

6.4 Proof of Theorem 6.2.6.1

By (2.4.2.3), we have $\Xi^{[G_n]}(h) \ll \Xi^{[G'_n]}(h)^2$ for $h \in [G'_n]$. Hence, by (2.4.2.4), the linear form $P_{G'_n}$ is well-defined and continuous on $\mathcal{C}([G_n])$. This shows the first part of Theorem 6.2.6.1.

Let $f \in \mathcal{S}_\chi([G_n])$. Recall that $A_{G_n}^\infty = A_{G'_n}^\infty$ but the Haar measure on $A_{G_n}^\infty$ is twice the Haar measure on $A_{G'_n}^\infty$ (see Remark 3.1.3.1). Therefore, we have

$$P_{G'_n}(f) = \frac{1}{2} \int_{[G'_n]_0} {}^0f(h) dh.$$

Let $\phi \in \mathcal{S}(\mathbb{A}^n)$. We form the Epstein-Eisenstein series

$$E(h, \phi, s) = \int_{A_{G'_n}^\infty} \sum_{\gamma \in \mathcal{P}'_n(F) \backslash G'_n(F)} \phi(e_n \gamma a h) |\det(a h)|^s da, \quad h \in [G'_n], s \in \mathbb{C}.$$

This expression converges absolutely for $\Re(s) > 1$ and the map $s \mapsto E(\phi, s)$ extends to a meromorphic function valued in $\mathcal{T}([G'_n])$ with simple poles at $s = 0, 1$ of respective residues $\phi(0)$ and $\widehat{\phi}(0)$ (cf. [JS81b, Lemma 4.2]).

Consequently, the function

$$s \mapsto Z_n^{\text{FR}}(s, {}^0f, \phi) := \int_{[G'_n]_0} {}^0f(h) E(h, \phi, s) dh$$

is well-defined for $s \in \mathbb{C} \setminus \{0, 1\}$, meromorphic on \mathbb{C} with a simple pole at $s = 1$ whose residue is

$$(6.4.0.1) \quad \text{Res}_{s=1} Z_n^{\text{FR}}(s, {}^0f, \phi) = 2\widehat{\phi}(0) P_{G'_n}(f).$$

Unfolding the definition, formally we arrive at

$$(6.4.0.2) \quad Z_n^{\text{FR}}(s, {}^0f, \phi) = \int_{\mathcal{P}'_n(F) \backslash G'_n(\mathbb{A})} {}^0f(h) \phi(e_n h) |\det h|^s dh.$$

By Lemma 6.4.0.1 below, there exists $c_n > 0$ such that the last integral above is absolutely convergent for $s \in \mathcal{H}_{>c_n}$ and thus the equality above is justified for such s .

More generally, for every $1 \leq r \leq n$, let $N_{r,n}$ be the unipotent radical of the standard parabolic subgroup of G_n with Levi component $G_r \times (G_1)^{n-r}$, $N'_{r,n}$ be its intersection with G'_n and set

$${}^0f_{N_{r,n}, \psi}(g) = \int_{[N_{r,n}]} {}^0f(ug) \psi_n(u)^{-1} du, \quad g \in G(\mathbb{A}),$$

$$Z_r^{\text{FR}}(s, {}^0f, \phi) = \int_{\mathcal{P}'_r(F) N'_{r,n}(\mathbb{A}) \backslash G'_n(\mathbb{A})} {}^0f_{N_{r,n}, \psi}(h) \phi(e_n h) |\det h|^s dh, \quad s \in \mathbb{C},$$

provided the last expression above is convergent. The proof of the next lemma will be given in Section 6.5.

Lemma 6.4.0.1. — *For every $1 \leq r \leq n$, there exists $c_r > 0$ such that the expression defining $Z_r^{\text{FR}}(s, {}^0f, \phi)$ converges absolutely for $\Re(s) > c_r$.*

When $r = 1$, we have $N_{1,n} = N_n$ and ${}^0f_{N_{1,n}, \psi} = W_{0f}$ so that $Z_1^{\text{FR}}(s, {}^0f, \phi) = Z_\psi^{\text{FR}}(s, {}^0f, \phi)$. Therefore, by (6.4.0.1) and (6.4.0.2), the second part of Theorem 6.2.6.1 is a consequence of the following proposition.

Proposition 6.4.0.2. — *For every $1 \leq r \leq n$, the function $s \mapsto (s-1)Z_r^{\text{FR}}(s, {}^0f, \phi)$ extends to a holomorphic function on $\{s \in \mathbb{C} \mid \Re(s) > 1\}$ admitting a limit at $s = 1$. Moreover, we have*

$$\lim_{s \rightarrow 1+} (s-1)Z_r^{\text{FR}}(s, {}^0f, \phi) = \lim_{s \rightarrow 1+} (s-1)Z_r^{\text{FR}}(s, {}^0f, \phi).$$

Proof. — By descending induction on r , it suffices to establish the following:

(6.4.0.3) Let $1 \leq r \leq n-1$. There exists a function F_r holomorphic on $\mathcal{H}_{>1-\epsilon}$ for some $\epsilon > 0$ such that

$$Z_{r+1}^{\text{FR}}(s, {}^0f, \phi) = Z_r^{\text{FR}}(s, {}^0f, \phi) + F_r(s)$$

for all $s \in \mathbb{C}$ satisfying $\Re(s) > \max(c_r, c_{r+1})$.

Indeed, as $\mathcal{P}'_{r+1} = G'_r U'_r$, we have

$$(6.4.0.4) \quad Z_{r+1}^{\text{FR}}(s, {}^0f, \phi) = \int_{G'_r(F)N'_{r,n}(\mathbb{A}) \backslash G'_n(\mathbb{A})} \int_{[U'_{r+1}]} {}^0f_{N_{r+1,n},\psi}(uh) du \phi(e_n h) |\det h|^s dh.$$

By Fourier inversion on the locally compact abelian group $U_{r+1}(F)U'_{r+1}(\mathbb{A}) \backslash U_{r+1}(\mathbb{A})$, we have

$$(6.4.0.5) \quad \int_{[U'_{r+1}]} {}^0f_{N_{r+1,n},\psi}(uh) du = \sum_{\gamma \in \mathcal{P}'_r(F) \backslash G'_r(F)} ({}^0f_{N_{r+1,n},\psi})_{U_{r+1},\psi}(\gamma h) + ({}^0f_{N_{r+1,n},\psi})_{U_{r+1}}(h)$$

for all $h \in G'_n(\mathbb{A})$ where we have set

$$({}^0f_{N_{r+1,n},\psi})_{U_{r+1},\psi}(h) = \int_{[U_{r+1}]} {}^0f_{N_{r+1,n},\psi}(uh) \psi_n(u)^{-1} du = {}^0f_{N_{r,n},\psi}(h),$$

$$({}^0f_{N_{r+1,n},\psi})_{U_{r+1}}(h) = \int_{[U_{r+1}]} {}^0f_{N_{r+1,n},\psi}(uh) du.$$

By (6.4.0.4) and (6.4.0.5), we obtain

$$Z_{r+1}^{\text{FR}}(s, {}^0f, \phi) = Z_r^{\text{FR}}(s, {}^0f, \phi) + F_r(s)$$

for all $s \in \mathbb{C}$ such that $\Re(s) > \max(c_r, c_{r+1})$ and where we have set

$$F_r(s) = \int_{G'_r(F)N'_{r,n}(\mathbb{A}) \backslash G'_n(\mathbb{A})} ({}^0f_{N_{r+1,n},\psi})_{U_{r+1}}(h) \phi(e_n h) |\det h|^s dh.$$

It only remains to check that $F_r(s)$ extends to a holomorphic function on $\mathcal{H}_{>1-\epsilon}$ for some $\epsilon > 0$.

Let P_r be the standard parabolic subgroup of G_n with Levi component $M_r = G_r \times G_{n-r}$ and set $P'_r = P_r \cap G'_n$. We readily check that

$$({}^0f_{N_{r+1,n},\psi})_{U_{r+1}}(h) = \int_{[N_{n-r}]} {}^0f_{P_r} \left(\begin{pmatrix} I_r & \\ & u \end{pmatrix} h \right) \psi_n(u)^{-1} du = \int_{[N_{n-r}]} \int_{A_{G_n}^\infty} f_{P_r} \left(\begin{pmatrix} I_r & \\ & u \end{pmatrix} ah \right) da \psi_n(u)^{-1} du.$$

Therefore, by the Iwasawa decomposition $G'_n(\mathbb{A}) = P'_r(\mathbb{A})K'_n$ and since

$$\delta_{P'_r} \left(\begin{pmatrix} h_r & \\ & h_{n-r} \end{pmatrix} \right) = \delta_{P_r} \left(\begin{pmatrix} h_r & \\ & h_{n-r} \end{pmatrix} \right)^{1/2} = |\det h_r|^{n-r} |\det h_{n-r}|^{-r}$$

for all $h_r \in G'_r(\mathbb{A})$, $h_{n-r} \in G'_{n-r}(\mathbb{A})$, we have (for $\Re(s) > \max(c_r, c_{r+1})$ and a suitable choice of Haar measure on K'_n)

(6.4.0.6)

$$F_r(s) = \int_{K'_n \times [G'_r] \times N'_{n-r}(\mathbb{A}) \backslash G'_{n-r}(\mathbb{A})} \int_{[N_{n-r}]} \int_{A_{G_n}^\infty} f_{P_r,k,s} \left(a \begin{pmatrix} h_r & \\ & u h_{n-r} \end{pmatrix} \right) da \psi_n(u)^{-1} du |\det h_{n-r}|^{ns/(n-r)} \phi_{k,n-r}(e_{n-r} h_{n-r}) dh_{n-r} dh_r dk$$

where $f_{P_r,k,s} = \delta_{P_r}^{-1/2+s/2(n-r)}(R(k)f)_{P_r} |_{M_r(\mathbb{A})}$ and $\phi_{k,n-r}$ stands for the composition of $R(k)\phi$ with the inclusion $\mathbb{A}^{n-r} \rightarrow \mathbb{A}^n$, $x \mapsto (0, x)$. Let χ^M be the inverse image of χ in $\mathfrak{X}(M_r)$. By Corollary 2.9.6.2, we have $f_{P_r,k,s} \in \mathcal{C}_{\chi^M}([M_r])$ for every $(k, s) \in K_n \times \mathcal{H}_{>0}$ and the map

$$(k, s) \in K_n \times \mathcal{H}_{>0} \mapsto f_{P_r,k,s} \in \mathcal{C}_{\chi^M}([M_r])$$

is continuous, holomorphic in the second variable. In particular, for $\Re(s) > 0$ the integral

$$\int_{A_{G'_r}^\infty} \int_{[N_{n-r}]} \int_{A_{G_n}^\infty} f_{P_r, k, s} \left(a \begin{pmatrix} a'h_r & \\ & uh_{n-r} \end{pmatrix} \right) da da'$$

is absolutely convergent and equals, by the obvious change of variable, to

$$\frac{n}{n-r} \int_{A_{G'_r}^\infty} \int_{[N_{n-r}]} \int_{A_{G_{n-r}}^\infty} f_{P_r, k, s} \begin{pmatrix} a'h_r & \\ & uah_{n-r} \end{pmatrix} da da'.$$

It follows that (6.4.0.6) can be rewritten, for $\Re(s) \gg 1$, as

$$\begin{aligned} F_r(s) &= \frac{n}{n-r} \int_{K'_n} \int_{[G'_r] \times N'_{n-r}(\mathbb{A}) \backslash G'_{n-r}(\mathbb{A})} \int_{[N_{n-r}]} \int_{A_{G_r}^\infty} f_{P_r, k, s} \begin{pmatrix} h_r & \\ & uah_{n-r} \end{pmatrix} da \psi_n(u)^{-1} du \\ &\quad |\det h_{n-r}|^{ns/(n-r)} \phi_{k, n-r}(e_{n-r} h_{n-r}) dh_{n-r} dh_r dk \\ (6.4.0.7) \quad &= \frac{n}{n-r} \int_{K'_n} (P_{G'_r} \hat{\otimes} Z_{n-r}^{\text{FR}}(\frac{ns}{n-r})) (f_{P_r, k, s} \otimes \phi_{k, n-r}) dk \end{aligned}$$

where $Z_{n-r}^{\text{FR}}(s)$ stands for the bilinear form

$$(f', \phi') \in \mathcal{C}([G_{n-r}]) \times \mathcal{S}(\mathbb{A}^{n-r}) \mapsto Z_{\psi(-1)^r}^{\text{FR}}(s, {}^0 f', \phi').$$

On the other hand, by (2.9.5.9) we have

$$\mathcal{C}_{\chi^M}([M_r]) = \bigoplus_{(\chi_1, \chi_2) \in \mathfrak{X}(M_r) = \mathfrak{X}(G_r) \times \mathfrak{X}(G_{n-r}) \mapsto \chi} \mathcal{C}_{\chi_1}([G_r]) \hat{\otimes} \mathcal{C}_{\chi_2}([G_{n-r}])$$

and, as $\chi \in \mathfrak{X}^*(G_n)$, for every $(\chi_1, \chi_2) \in \mathfrak{X}(G_r) \times \mathfrak{X}(G_{n-r})$ mapping to $\chi \in \mathfrak{X}(G_n)$, we also have $\chi_2 \in \mathfrak{X}^*(G_{n-r})$. Therefore, by the first part of Theorem 6.2.6.1, Theorem 6.2.5.1 and (A.0.5.5), $s \mapsto P_{G'_r} \hat{\otimes} Z_{n-r}^{\text{FR}}(s)$ extends to an analytic family of (separately) continuous bilinear forms on $\mathcal{C}_\chi([M_r]) \times \mathcal{S}(\mathbb{A}^{n-r})$ for $s \in \mathcal{H}_{>1}$. Thus, by the first part of Theorem 6.2.6.1, (A.0.5.4) and the equality (6.4.0.7), $F_r(s)$ has an analytic continuation to $\{\Re(s) > 1 - r/n\}$. This ends the proof of the proposition and hence of Theorem 6.2.6.1. \square

6.5 Convergence of Zeta integrals

6.5.1. Proof. — (of Lemma 6.4.0.1) We only treat the case $1 \leq r \leq n-1$. The case $r = n$ can be dealt with in a similar manner, and is in fact easier.

Let Q_r be the standard parabolic subgroup of G_n with Levi component $G_r \times G_1^{n-r}$ and set $Q'_r = Q_r \cap G'_n$. Recall that $N_{r,n}$ is the unipotent radical of Q_r . Identifying $A_{G_n}^\infty \simeq \mathbb{R}_{>0}$, by the Iwasawa decomposition $G'_n(\mathbb{A}) = Q'_r(\mathbb{A})K'_n$, we need to show the convergence of

$$(6.5.1.1) \quad \int_{K'_n \times \mathcal{P}'_r(F) \backslash G'_r(\mathbb{A}) \times T'_{n-r}(\mathbb{A}) \times \mathbb{R}_{>0}} \left| (R(k)f)_{N_{r,n}, \psi} \begin{pmatrix} ah & \\ & at \end{pmatrix} \right| |R(k)\phi(t_{n-r}e_n)| \\ |\det h|^s |\det t|^s \delta_{Q'_r} \begin{pmatrix} h & \\ & t \end{pmatrix}^{-1} da dt dh dk$$

for $\Re(s) \gg 1$. We now apply Lemma 2.6.1.1. For this we note that $\psi_n|_{[N_{r,n}]} = \psi' \circ \ell$ where $\ell : N_{r,n} \rightarrow \mathbf{G}_a$ sends $u \in N_{r,n}$ to $\text{Tr}_{E/F}(\tau \sum_{i=r}^{n-1} u_{i,i+1})$ and $\tau \in E^\times$ is the unique trace-zero element such that $\psi(z) = \psi'(\text{Tr}_{E/F}(\tau z))$. We readily check that

$$\|\text{Ad}^*(m)\ell\|_{(N_{r,n})_{\text{ab}}^*(\mathbb{A})} \approx \|t_1^{-1}e_r h\|_{\mathbb{A}^r} \prod_{i=1}^{n-r-1} \|t_i t_{i+1}^{-1}\|_{\mathbb{A}}, \quad m = \begin{pmatrix} h & \\ & t \end{pmatrix} \in G'_r(\mathbb{A}) \times T'_{n-r}(\mathbb{A}).$$

Therefore, by Lemma 2.6.1.1.1, we can find $c > 0$ such that for every $N_1, N_2 > 0$ we have

$$(6.5.1.2) \quad \left| (R(k)f)_{N_{r,n},\psi} \begin{pmatrix} ah & \\ & at \end{pmatrix} \right| \ll \|ah\|_{[G'_r]}^{-N_2} \|t_1^{-1} e_r h\|_{\mathbb{A}^r}^{-N_1} \prod_{i=1}^{n-r-1} \|t_i t_{i+1}^{-1}\|_{\mathbb{A}}^{-N_1} \delta_{Q_r} \begin{pmatrix} h & \\ & t \end{pmatrix}^{-cN_2}$$

for $(k, h, t, a) \in K'_n \times G'_r(\mathbb{A}) \times T'_{n-r}(\mathbb{A}) \times \mathbb{R}_{>0}$. On the other hand, for every $N_1 > 0$, we have

$$|R(k)\phi(te_n)| \ll \|t\|_{\mathbb{A}}^{-N_1}, \quad (k, t) \in K'_n \times \mathbb{A}$$

and it is easy to check that for some $N_2 > 0$ we have

$$\|e_r h\|_{\mathbb{A}^r} \prod_{i=1}^{n-r} \|t_i\|_{\mathbb{A}} \ll \|t_{n-r}\|_{\mathbb{A}}^{N_2} \|t_1^{-1} e_r h\|_{\mathbb{A}^r}^{N_2} \prod_{i=1}^{n-r-1} \|t_i t_{i+1}^{-1}\|_{\mathbb{A}}^{N_2}, \quad (h, t) \in G'_r(\mathbb{A}) \times T'_{n-r}(\mathbb{A}).$$

As $\delta_{Q_r} \begin{pmatrix} h & \\ & t \end{pmatrix} = |\det h|^{n-r} \prod_{i=1}^{n-r} |t_i|^{n+1-2(r+i)}$ for every $(h, t) \in G'_r(\mathbb{A}) \times T'_{n-r}(\mathbb{A})$, combining this with (6.5.1.2), we deduce the existence of $c > 0$ such that for every $N_1, N_2 > 0$, (6.5.1.1) is essentially bounded by the product of

$$(6.5.1.3) \quad \int_{\mathcal{P}'_r(F) \setminus G'_r(\mathbb{A}) \times \mathbb{R}_{>0}} \|ah\|_{[G'_r]}^{-N_2} \|e_r h\|_{\mathbb{A}^r}^{-N_1} |\det h|^{s-(2cN_2+1)(n-r)} dadh$$

and

$$(6.5.1.4) \quad \int_{\mathbb{A}^\times} \|t\|_{\mathbb{A}}^{-N_1} |t|^{s-(2cN_2+1)(n+1-2(r+i))} dt$$

for $1 \leq i \leq n-r$.

Let $C_1, C_2 > 0$. By Lemma 2.6.2.1, for N_1 sufficiently large the integral (6.5.1.4) converges absolutely in the range

$$1 + (2cN_2 + 1)(n + 1 - 2(r + i)) < \Re(s) < C_1 + (2cN_2 + 1)(n + 1 - 2(r + i))$$

and for N_1, N_2 sufficiently large the integral (6.5.1.3) converges absolutely in the range

$$1 + (2cN_2 + 1)(n - r) < \Re(s) < C_2 + (2cN_2 + 1)(n - r).$$

Since $n + 1 - 2(r + i) < n - r$ for every $1 \leq i \leq n - r$, by taking $C_2 = 2$ and $C_1 \geq 2 + (2cN_2 + 1)(r + 2i - 1)$ for every $1 \leq i \leq n - r$, it follows that if $N_2 \gg 1$ and $N_1 \gg_{N_2} 1$ the integrals (6.5.1.3) and (6.5.1.4) are convergent in the range

$$1 + (2cN_2 + 1)(n - r) < \Re(s) < 2 + (2cN_2 + 1)(n - r).$$

The union of these open intervals for N_2 sufficiently large as above is of the form $]c_r, +\infty[$ which shows that $Z_r^{\text{FR}}(s, {}^0f, \phi)$ converges absolutely in the range $\Re(s) > c_r$ for a suitable $c_r > 0$. \square

6.5.2. Proof. — (of Theorem 6.2.5.1.1) Applying Lemma 2.6.1.1.2, the same manipulations as in the proof of Lemma 6.4.0.1 reduce us to showing the existence of $c_N > 0$ such that for every $C > c_N$ there exists $N' > 0$ satisfying that the integral

$$(6.5.2.5) \quad \int_{T'_n(\mathbb{A})} \prod_{i=1}^n \|t_i\|_{\mathbb{A}}^{-N'} \|t\|_{[T'_n]}^N \delta_{B'_n}(t)^{-1} |\det t|^s dt$$

converges in the range $s \in \mathcal{H}_{c_n, C}$ uniformly on compact subsets. But this follows again from Lemma 2.6.2.1 as there exists $M > 0$ such that

$$\|t\|_{[T'_n]}^N \delta_{B'_n}(t)^{-1} \ll \prod_{1 \leq i \leq n} \max(|t_i|, |t_i|^{-1})^M, \quad t \in [T'_n].$$

\square

7 Canonical extension of the Rankin-Selberg period for certain cuspidal data

This chapter is a continuation of Chapter 6 and we shall use the notation introduced there. The main goal is to show the existence of a canonical extension of corank one Rankin-Selberg periods to the space of uniform moderate growth functions for certain cuspidal data (see Theorem 7.1.4.1). Combining this with the results of Chapter 6, this will enable us to give an alternative proof of the spectral expansion of the Jacquet-Rallis trace formula for certain cuspidal data in Chapter 8.

7.1 Statements of the main results

7.1.1. Let $n \geq 1$ be a positive integer. We set $G = G_n \times G_{n+1}$ and $H = G_n$ that we consider as an algebraic subgroup of G via the diagonal inclusion $H \hookrightarrow G$. We also set $w = (w_n, w_{n+1}) \in G(F)$, $K = K_n \times K_{n+1}$, $N = N_n \times N_{n+1}$ and $N_H = N_n$. Put $\psi_N = \psi_n \boxtimes \psi_{n+1}$ (a generic character of $[N]$). We note that ψ_N is trivial on $[N_H]$ (see the convention in the definition of ψ_n in §6.1.2). To any function $f \in \mathcal{T}([G])$, we associate its *Whittaker function*

$$W_f(g) = \int_{[N]} f(ug) \psi_N(u)^{-1} du, \quad g \in G(\mathbb{A}).$$

For $f \in \mathcal{T}([G])$, we define

$$Z_\psi^{\text{RS}}(s, f) = \int_{N_H(\mathbb{A}) \backslash H(\mathbb{A})} W_f(h) |\det h|_E^s dh$$

for every $s \in \mathbb{C}$ for which the above expression converges absolutely.

7.1.2. H -generic cuspidal datum Let $\chi \in \mathfrak{X}(G)$ be a cuspidal datum represented by a pair (M_P, π) where $P = P_n \times P_{n+1}$ is a standard parabolic subgroup of G and $\pi = \pi_n \boxtimes \pi_{n+1}$ a cuspidal automorphic representation of $M_P(\mathbb{A})$ (with central character trivial on A_P^∞). We have decompositions

$$M_{P_n} = G_{n_1} \times \dots \times G_{n_k}, \quad M_{P_{n+1}} = G_{m_1} \times \dots \times G_{m_r}$$

and π_n, π_{n+1} decompose accordingly as tensor products

$$\pi_n = \pi_{n,1} \boxtimes \dots \boxtimes \pi_{n,k}, \quad \pi_{n+1} = \pi_{n+1,1} \boxtimes \dots \boxtimes \pi_{n+1,r}.$$

We will say that χ is H -generic if it satisfies the following condition:

(7.1.2.1) For every $1 \leq i \leq k$ and $1 \leq j \leq r$, we have $\pi_{n,i} \neq \pi_{n+1,j}$ or equivalently the Rankin-Selberg L -function $L(s, \pi_{n,i} \times \pi_{n+1,j})$ is entire.

7.1.3. Theorem 7.1.3.1. —

1. Let $N \geq 0$. There exists $c_N > 0$ such that for $f \in \mathcal{T}_N([G])$, the expression defining $Z_\psi^{\text{RS}}(s, f)$ converges absolutely for $s \in \mathcal{H}_{>c_N}$ and the map $s \in \mathcal{H}_{>c_N} \mapsto Z_\psi^{\text{RS}}(s, f)$ is holomorphic and bounded in vertical strips. Moreover, for every $s \in \mathcal{H}_{>c_N}$, $f \mapsto Z_\psi^{\text{RS}}(s, f)$ is a continuous functional on $\mathcal{T}_N([G])$.
2. Assume that $\chi \in \mathfrak{X}(G)$ is a H -generic cuspidal datum. Then, for every $f \in \mathcal{T}_\chi([G])$, the function $s \mapsto Z_\psi^{\text{RS}}(s, f)$ extends analytically to \mathbb{C} . Moreover, for every $s \in \mathbb{C}$ the linear form $f \in \mathcal{T}_\chi([G]) \mapsto Z_\psi^{\text{RS}}(s, f)$ is continuous.

7.1.4. Theorem 7.1.4.1. — Assume that $\chi \in \mathfrak{X}(G)$ is a H -generic cuspidal datum. The

restriction of the linear form

$$P_H : f \in \mathcal{S}([G]) \mapsto \int_{[H]} f(h) dh$$

to $\mathcal{S}_\chi([G])$ extends by continuity to $\mathcal{T}_\chi([G])$ and for every $f \in \mathcal{T}_\chi([G])$, we have

$$(7.1.4.2) \quad P_H(f) = Z_\psi^{\text{RS}}(0, f).$$

7.2 Proof of Theorem 7.1.3.1.2

Part 1. of Theorem 7.1.3.1 will be established in Section 7.4. Here, we give the proof of part 2. Let $f \in \mathcal{S}_\chi([G])$ and (M_P, π) be a cuspidal datum representing χ as in Section 7.1. We set $\Pi_{n,\mu} = I_{P_n(\mathbb{A})}^{G_n(\mathbb{A})}(\pi_{n,\lambda})$ and $\Pi_{n+1,\nu} = I_{P_{n+1}(\mathbb{A})}^{G_{n+1}(\mathbb{A})}(\pi_{n+1,\nu})$ for every $\mu \in \mathfrak{a}_{P_n, \mathbb{C}}^*$ and $\nu \in \mathfrak{a}_{P_{n+1}, \mathbb{C}}^*$. For $\lambda = (\mu, \nu) \in \mathfrak{a}_{P, \mathbb{C}}^*$, we also set $\Pi_\lambda = \Pi_{n,\mu} \boxtimes \Pi_{n+1,\nu}$ and $f_\lambda = f_{\Pi_\lambda}$ (following the notation from Section 2.9.7). Then, by Theorem 2.9.7.1 and the first part of Theorem 7.1.3.1, for $\Re(s) \gg 1$ we have

$$(7.2.0.1) \quad Z_\psi^{\text{RS}}(s, f) = \int_{i\mathfrak{a}_P^*} Z_\psi^{\text{RS}}(s, f_\lambda) d\lambda.$$

Set $\widetilde{f}_\lambda(g) = f_\lambda({}^t g^{-1})$ for every $\lambda \in i\mathfrak{a}_P^*$ and $g \in G(\mathbb{A})$. We will now check that the functions

$$(7.2.0.2) \quad (s, \lambda) \in \mathbb{C} \times i\mathfrak{a}_P^* \mapsto Z_\psi^{\text{RS}}(s, f_\lambda) \quad \text{and} \quad (s, \lambda) \in \mathbb{C} \times i\mathfrak{a}_P^* \mapsto Z_{\psi^{-1}}^{\text{RS}}(s, \widetilde{f}_\lambda)$$

satisfy the conditions of Corollary A.0.10.1.

For S a sufficiently large finite set of places of F , that we assume to contain Archimedean places as well as the places where π or ψ are ramified, we have, for every $\lambda \in i\mathfrak{a}_P^*$, a decomposition

$$W_{f_\lambda} = W_{\lambda, S} W_\lambda^S$$

where $W_{\lambda, S} \in \mathcal{W}(\Pi_{\lambda, S}, \psi_S)$ and $W_\lambda^S \in \mathcal{W}(\Pi_\lambda^S, \psi^S)^{K^S}$ is such that $W_\lambda^S(1) = 1$. By the unramified computation of local Rankin-Selberg integrals [JS81a, (3), p.781], [Cog08, Theorem 3.3] and (2.3.3.4), we have

$$(7.2.0.3) \quad Z_\psi^{\text{RS}}(s, f_\lambda) = (\Delta_H^{S,*})^{-1} L(s + \frac{1}{2}, \Pi_\lambda) \frac{Z_\psi^{\text{RS}}(s, W_{\lambda, S})}{L_S(s + \frac{1}{2}, \Pi_\lambda)}$$

for $\Re(s) \gg 1$ where we have set $L(s, \Pi_\lambda) = L(s, \Pi_{n,\mu} \times \Pi_{n+1,\nu})$ and

$$Z_\psi^{\text{RS}}(s, W_{\lambda, S}) = \int_{N_H(F_S) \backslash H(F_S)} W_{\lambda, S}(h_S) |\det h_S|_E^s dh_S$$

whenever $\lambda = (\mu, \nu) \in i\mathfrak{a}_P^*$. By [JPSS83], [Cog08, Theorem 4.1] and the condition that χ is H -generic (see (7.1.2.1)), the Rankin-Selberg L -function $L(s, \Pi_\lambda)$ is entire and bounded in vertical strips. On the other hand, by [JPSS83, Theorem 2.7] and [Jac09, Theorem 2.1], $s \mapsto \frac{Z_\psi^{\text{RS}}(s, W_{\lambda, S})}{L_S(s + \frac{1}{2}, \Pi_\lambda)}$ has a holomorphic continuation to \mathbb{C} which is of order at most 1 in vertical strips and satisfies the functional equation

$$(7.2.0.4) \quad \frac{Z_{\psi^{-1}}^{\text{RS}}(-s, \widetilde{W}_{\lambda, S})}{L_S(\frac{1}{2} - s, \Pi_\lambda^\vee)} = \epsilon(s + \frac{1}{2}, \Pi_\lambda) \frac{Z_\psi^{\text{RS}}(s, W_{\lambda, S})}{L_S(s + \frac{1}{2}, \Pi_\lambda)}$$

where $\widetilde{W}_{\lambda,S}(g_S) = W_{\lambda,S}(w^t g_S^{-1})$ and $\epsilon(s, \Pi_\lambda)$ denotes the global epsilon factor of the Rankin-Selberg L -function $L(s, \Pi_\lambda)$. Therefore, by (7.2.0.3) and the functional equation of the Rankin-Selberg L -function $L(s, \Pi_\lambda)$, we deduce that, for every $\lambda \in i\mathfrak{a}_P^*$, the function $s \mapsto Z_\psi^{\text{RS}}(s, f_\lambda)$ has a holomorphic continuation to \mathbb{C} which is of order at most 1 in vertical strips and satisfies the functional equation

$$(7.2.0.5) \quad Z_\psi^{\text{RS}}(s, f_\lambda) = Z_{\psi^{-1}}^{\text{RS}}(-s, \widetilde{f}_\lambda).$$

By the first part of Theorem 7.1.3.1, Theorem 2.9.7.1 and Lemma A.0.8.1, this shows that the functions in (7.2.0.2) satisfy the assumptions of Corollary A.0.10.1. Therefore, the map $s \mapsto (\lambda \in i\mathfrak{a}_P^* \mapsto Z_\psi^{\text{RS}}(s, f_\lambda))$ induces a holomorphic function $\mathbb{C} \rightarrow \mathcal{S}(i\mathfrak{a}_P^*)$ which is of finite order in vertical strips. By (7.2.0.1) and (7.2.0.5), this implies that $s \mapsto Z_\psi^{\text{RS}}(s, f)$ extends to a holomorphic function on \mathbb{C} of finite order in vertical strips satisfying the functional equation

$$(7.2.0.6) \quad Z_{\psi^{-1}}^{\text{RS}}(-s, \widetilde{f}) = Z_\psi^{\text{RS}}(s, f)$$

where $\widetilde{f}(g) = f({}^t g^{-1})$. As for $N > 0$, the closure of $\mathcal{S}_\chi([G])$ in $\mathcal{T}_{M,\chi}([G])$ contains $\mathcal{T}_{N,\chi}([G])$ for M sufficiently large (see (2.9.4.7)), Theorem 7.1.3.1.2 now follows from part 1. and Corollary A.0.10.2 (applied to the closure of $\mathcal{S}_\chi([G])$ in $\mathcal{T}_{M,\chi}([G])$).

7.3 Proof of Theorem 7.1.4.1

By Theorem 7.1.3.1.2, and since $\mathcal{S}_\chi([G])$ is dense in $\mathcal{T}_\chi([G])$ (see (2.9.4.7)), it suffices to check that identity (7.1.4.2) is valid for every $f \in \mathcal{S}_\chi([G])$. Therefore, let $f \in \mathcal{S}_\chi([G])$. For $1 \leq r \leq n$, we let $N_{r,n}$ and $N_{r,n+1}$ be the unipotent radicals of the standard parabolic subgroups of G_n and G_{n+1} with Levi components $G_r \times (G_1)^{n-r}$ and $G_r \times (G_1)^{n+1-r}$ respectively. Set $N_r^G = N_{r,n} \times N_{r,n+1}$ and $N_r^H = N_r^G \cap H = N_{r,n}$. For every $1 \leq r \leq n$, define

$$f_{N_r^G, \psi}(g) = \int_{[N_r^G]} f(ug) \psi_N(u)^{-1} du, \quad g \in G(\mathbb{A}).$$

Whenever the expression below converges absolutely, for every $1 \leq r \leq n$ and $s \in \mathbb{C}$ we set

$$Z_r^{\text{RS}}(s, f) = \int_{\mathcal{P}_r(F) N_r^H(\mathbb{A}) \backslash H(\mathbb{A})} f_{N_r^G, \psi}(h) |\det h|_E^s dh.$$

The proof of the next lemma will be given in Section 7.4.

Lemma 7.3.0.1. — *For every $1 \leq r \leq n$, there exists $c_r > 0$ such that the expression defining $Z_r^{\text{RS}}(s, f)$ converges absolutely for $s \in \mathcal{H}_{>c_r}$.*

To uniformize notation, for every $s \in \mathbb{C}$ we also set

$$Z_{n+1}^{\text{RS}}(s, f) = \int_{[H]} f(h) |\det h|_E^s dh.$$

Note that the above expression is absolutely convergent and defines an entire function of $s \in \mathbb{C}$ satisfying $P_H(f) = Z_{n+1}^{\text{RS}}(0, f)$. On the other hand, we have $Z_1^{\text{RS}}(s, f) = Z_\psi^{\text{RS}}(s, f)$. Hence, identity (7.1.4.2) is a consequence of the following proposition.

Proposition 7.3.0.2. — *For every $1 \leq r \leq n$, we have*

$$(7.3.0.1) \quad Z_{r+1}^{\text{RS}}(s, f) = Z_r^{\text{RS}}(s, f)$$

for $\Re(s) \gg 1$.

Proof. — Let $1 \leq r \leq n-1$. As $\mathcal{P}_{r+1} = G_r U_r$ and $N_{r,n} = U_r N_{r+1,n}$, for $s \in \mathcal{H}_{>c_{r+1}}$ we have

$$(7.3.0.2) \quad Z_{r+1}^{\text{RS}}(s, f) = \int_{G_r(F) N_r^H(\mathbb{A}) \backslash H(\mathbb{A})} \int_{[U_{r+1}^H]} f_{N_{r+1}^G, \psi}(uh) du |\det h|_E^s dh$$

where we have set $U_{r+1}^H = U_{r+1}$ viewed as a subgroup of $H = G_n$ (as always via the embedding in “the upper-left corner”). Similarly, we set $U_{r+1}^G = U_{r+1} \times U_{r+1}$ viewed as a subgroup of G . By Fourier inversion on the compact abelian group $U_{r+1}^G(F)U_{r+1}^H(\mathbb{A}) \backslash U_{r+1}^G(\mathbb{A})$, we have

$$(7.3.0.3) \quad \int_{[U_{r+1}^H]} f_{N_{r+1}^G, \psi}(uh) du = \sum_{\gamma \in \mathcal{P}_r(F) \backslash G_r(F)} (f_{N_{r+1}^G, \psi})_{U_{r+1}^G, \psi}(\gamma h) + (f_{N_{r+1}^G, \psi})_{U_{r+1}^G}(h)$$

for every $h \in H(\mathbb{A})$, where we have set

$$(f_{N_{r+1}^G, \psi})_{U_{r+1}^G, \psi}(h) = \int_{[U_{r+1}^G]} f_{N_{r+1}^G, \psi}(uh) \psi_N(u)^{-1} du = f_{N_r^G, \psi}(h),$$

$$(f_{N_{r+1}^G, \psi})_{U_{r+1}^G}(h) = \int_{[U_{r+1}^G]} f_{N_{r+1}^G, \psi}(uh) du.$$

By (7.3.0.2) and (7.3.0.3), we obtain

$$(7.3.0.4) \quad Z_{r+1}^{\text{RS}}(s, f) = Z_r^{\text{RS}}(s, f) + F_r(s)$$

for every $s \in \mathbb{C}$ such that $\Re(s) > \max(c_r, c_{r+1})$ where we have set

$$F_r(s) = \int_{G_r(F)N_r^H(\mathbb{A}) \backslash H(\mathbb{A})} (f_{N_{r+1}^G, \psi})_{U_{r+1}^G}(h) |\det h|_E^s dh.$$

By a similar argument, (7.3.0.4) still holds when $n = r$ if we set

$$F_n(s) = \int_{[H]} f_{U_{n+1}^G}(h) |\det h|_E^s dh$$

with $U_{n+1}^G = 1 \times U_{n+1}$ and

$$f_{U_{n+1}^G}(h) = \int_{[U_{n+1}^G]} f(uh) du.$$

From (7.3.0.4), we are reduced to showing that $F_r(s) = 0$ identically for every $1 \leq r \leq n$ and $\Re(s)$ sufficiently large. To uniformize notation, we set $f_{N_{n+1}^G, \psi} = f$. Let P_r be the standard parabolic subgroup of G with Levi component $L_r = (G_r \times G_{n-r}) \times (G_r \times G_{n+1-r})$ and set $P_r^H = P_r \cap H$. Then, we readily check that

$$(f_{N_{r+1}^G, \psi})_{U_{r+1}^G}(h) = \int_{[N_{n-r}] \times [N_{n+1-r}]} f_{P_r} \left(\left(\begin{pmatrix} I_r & \\ & u \end{pmatrix}, \begin{pmatrix} I_r & \\ & u' \end{pmatrix} \right) h \right) \psi_n(u)^{-1} \psi_{n+1}(u')^{-1} du' du$$

for every $h \in H(\mathbb{A})$ and $1 \leq r \leq n$. Therefore, by the Iwasawa decomposition $H(\mathbb{A}) = P_r^H(\mathbb{A})K_n$, we have

$$F_r(s) = \int_{K_n} \int_{[G_r] \times N_{n-r}(\mathbb{A}) \backslash G_{n-r}(\mathbb{A})} \int_{[N_{n-r}] \times [N_{n+1-r}]} f_{P_r} \left(\begin{pmatrix} h_r & \\ & u h_{n-r} \end{pmatrix} k, \begin{pmatrix} h_r & \\ & u' h_{n-r} \end{pmatrix} k \right) \\ \psi_n(u)^{-1} \psi_{n+1}(u')^{-1} du' du \delta_{Q_r^H} \left(\begin{pmatrix} h_r & \\ & h_{n-r} \end{pmatrix} \right)^{-1} |\det h_r|_E^s |\det h_{n-r}|_E^s dh_{n-r} dh_r dk.$$

By a painless calculation, left to the reader, we have

$$\delta_{P_r^H} \left(\begin{pmatrix} h_r & \\ & h_{n-r} \end{pmatrix} \right)^{-1} |\det h_r|_E^s |\det h_{n-r}|_E^s = \delta_{P_r} \left(\begin{pmatrix} h_r & \\ & h_{n-r} \end{pmatrix}, \begin{pmatrix} h_r & \\ & h_{n-r} \end{pmatrix} \right)^{-\frac{1}{2} + \alpha_r(s)} |\det h_{n-r}|_E^{s+2r\alpha_r(s)}$$

where $\alpha_r(s) = \frac{2s+1}{4n-4r+2}$. Let χ^L be the inverse image of χ in $\mathfrak{X}(L_r)$. By Corollary 2.9.6.2, for $\Re(\alpha_r(s)) > 0$ and every $k \in K_n$ the function $f_{P_r, k, s} := \delta_{P_r}^{-\frac{1}{2} + \alpha_r(s)} R(k) f_{P_r} \mid_{[L_r]}$ belongs to $\mathcal{C}_{\chi^L}([L_r])$. On the other hand, as $L_r = G_r \times G_{n-r} \times G_r \times G_{n+1-r}$, by (2.9.5.9) we have the decomposition

$$\mathcal{C}_{\chi^L}([L_r]) = \bigoplus_{(\chi_1, \chi_2) \in \mathfrak{X}(G_r^2) \times \mathfrak{X}(G_{n-r} \times G_{n+1-r}) \mapsto \chi} \mathcal{C}_{\chi_1}([G_r \times G_r]) \widehat{\otimes} \mathcal{C}_{\chi_2}([G_{n-r} \times G_{n+1-r}])$$

and the above equality can be rewritten as

$$(7.3.0.5) \quad F_r(s) = \int_{K_n} (P_{G_r^\Delta} \hat{\otimes} \mathcal{Z}_{n-r}^{\text{RS}}(s + 2r\alpha_r(s))) (f_{P_r, k, s}) dk$$

where $P_{G_r^\Delta}$ denotes the period integral over the diagonal subgroup of $G_r \times G_r$ and $\mathcal{Z}_{n-r}^{\text{RS}}(s)$ stands for the continuous linear form

$$f' \in \mathcal{C}([G_{n-r} \times G_{n+1-r}]) \mapsto Z_{\psi(-1)^r}^{\text{RS}}(s, f').$$

Since χ is H -generic, by (7.1.2.1) any preimage $(\chi_1, \chi_2) \in \mathfrak{X}(G_r^2) \times \mathfrak{X}(G_{n-r} \times G_{n+1-r})$ of χ with $\chi_1 = (\chi'_1, \chi''_1) \in \mathfrak{X}(G_r)^2$ we have $\chi'_1 \neq (\chi'_1)^\vee$. Hence, by definition of $\mathcal{C}_{\chi_1}([G_r \times G_r])$, $P_{G_r^\Delta}$ vanishes identically on $\mathcal{C}_{\chi_1}([G_r \times G_r])$. This implies that $F_r(s) = 0$ whenever $\Re(s) \gg 1$ and this ends the proof of the proposition. \square

7.4 Convergence of Zeta integrals

Proof. — (of Lemma 7.3.0.1) The argument is very similar to the proof of Lemma 6.4.0.1 so we only sketch it. Let $1 \leq r \leq n$ and Q_r^G be the standard parabolic subgroup of G with Levi component $(G_r \times (G_1)^{n-r}) \times (G_r \times (G_1)^{n+1-r})$ so that N_r^G is the unipotent radical of Q_r^G . Set $Q_r^H = Q_r^G \cap H$. By the Iwasawa decomposition $H(\mathbb{A}) = Q_r^H(\mathbb{A})K_n$, we need to show the convergence of

$$(7.4.0.1) \quad \int_{K_n \times \mathcal{P}_r(F) \backslash G_r(\mathbb{A}) \times T_{n-r}(\mathbb{A})} \left| (R(k)f)_{N_r^G, \psi} \begin{pmatrix} h & \\ & t \end{pmatrix} \right| |\det h|_E^s |\det t|_E^s \delta_{Q_r^H} \begin{pmatrix} h & \\ & t \end{pmatrix}^{-1} dt dh dk$$

for $\Re(s) \gg 1$. We apply Lemma 2.6.1.1.1 to $\psi_F = \psi'$ and

$$\ell : N_r^G \rightarrow \mathbf{G}_a,$$

$$(u, u') \mapsto \text{Tr}_{E/F} \left((-1)^n \tau \sum_{i=r}^{n-1} u_{i, i+1} + (-1)^{n+1} \tau \sum_{i=r}^n u'_{i, i+1} \right).$$

It is easy to see that there exists $N_0 > 0$ such that

$$(7.4.0.2) \quad \|e_r h\|_{\mathbb{A}_E^r} \prod_{i=1}^{n-r} \|t_i\|_{\mathbb{A}_E} \ll \|\text{Ad}^* \begin{pmatrix} h & \\ & t \end{pmatrix} \ell\|_{(N_r^G)^*_{\text{ab}}(\mathbb{A})}^{N_0}, \quad (h, t) \in G_r(\mathbb{A}) \times T_{n-r}(\mathbb{A}).$$

Therefore, from (7.4.0.2) and Lemma 2.6.1.1, there exists $c > 0$ such that for every $N_1, N_2 > 0$, (7.4.0.1) is essentially bounded by

$$\int_{\mathcal{P}_r(F) \backslash G_r(\mathbb{A}) \times T_{n-r}(\mathbb{A})} \|h\|_{[G_r]}^{-N_2} \|e_r h\|_{\mathbb{A}_E^r}^{-N_1} \prod_{i=1}^{n-r} \|t_i\|_{\mathbb{A}_E}^{-N_1} \delta_{Q_r^G} \begin{pmatrix} h & \\ & t \end{pmatrix}^{-cN_2} \delta_{Q_r^H} \begin{pmatrix} h & \\ & t \end{pmatrix}^{-1} |\det h|_E^s |\det t|_E^s dt dh.$$

Now, the convergence of the above expression for $\Re(s) \gg 1$, $N_2 \gg_s 1$ and $N_1 \gg_{s, N_2} 1$ can be shown as in the end of the proof of Lemma 6.4.0.1 using Lemma 2.6.2.1. \square

Proof. — (of Theorem 7.1.3.1.1) Applying Lemma 2.6.1.1.2 in a similar way, we are reduced to showing the existence of $c_N > 0$ such that for every $C > c_N$ there exists $N' > 0$ satisfying that the integral

$$\int_{T_n(\mathbb{A})} \prod_{i=1}^n \|t_i\|_{\mathbb{A}_E}^{-N'} \|t\|_{[T_n]}^N \delta_{B_n}(t)^{-1} |\det t|_E^s dt$$

converges in the range $s \in \mathcal{H}_{[c_N, C]}$ uniformly on any compact subsets. This is exactly what was established in the proof of Theorem 6.2.5.1.1 (up to replacing the base field F by E). \square

8 Contributions of certain cuspidal data to the Jacquet-Rallis trace formula: second proof

In this chapter, we adopt the set of notation introduced in Chapter 5. In particular, $n \geq 1$ is a positive integer, $G = G_n \times G_{n+1}$, $G' = G'_n \times G'_{n+1}$, $H = G_n$ with its diagonal embedding in G , $K = K_n \times K_{n+1}$ and $K' = K'_n \times K'_{n+1}$ are the standard maximal compact subgroups of $G(\mathbb{A})$ and $G'(\mathbb{A})$ respectively and $\eta_{G'} : [G'] \rightarrow \{\pm 1\}$ is the automorphic character defined in §3.1.8. We will also use notation from Chapters 6 and 7: $N = N_n \times N_{n+1}$ and $N_H = N_n$ are the standard maximal unipotent subgroups of G and H , $\psi_N = \psi_n \boxtimes \psi_{n+1}$ is a generic character of $[N]$ (where ψ_n and ψ_{n+1} are defined as in §6.1.2). We also set $\mathcal{P} = \mathcal{P}_n \times \mathcal{P}_{n+1}$ (resp. $\mathcal{P}' = \mathcal{P}'_n \times \mathcal{P}'_{n+1}$) where \mathcal{P}_n and \mathcal{P}_{n+1} (resp. \mathcal{P}'_n and \mathcal{P}'_{n+1}) stand for the mirabolic subgroups of G_n and G_{n+1} (resp. of G'_n and G'_{n+1}), $T = T_n \times T_{n+1}$ for the standard maximal torus of G and $N' = N'_n \times N'_{n+1}$ for the standard maximal unipotent subgroup of G' . Finally, as in §7.1.1, for every $f \in \mathcal{T}([G])$ we set

$$W_f(g) = \int_{[N]} f(ug) \psi_N(u)^{-1} du, \quad g \in G(\mathbb{A}).$$

8.1 Main result

8.1.1. Let $\chi \in \mathfrak{X}^*(G)$ be a *-generic cuspidal datum (see §5.1.3) represented by a pair (M_P, π) . We set $\Pi = I_{P(\mathbb{A})}^{G(\mathbb{A})}(\pi)$. We have decompositions $P = P_n \times P_{n+1}$, $\pi = \pi_n \boxtimes \pi_{n+1}$ and $\Pi = \Pi_n \boxtimes \Pi_{n+1}$ where: P_n, P_{n+1} are standard parabolic subgroups of G_n, G_{n+1} respectively with standard Levi components of the form

$$M_{P_n} = G_{n_1} \times \dots \times G_{n_k}, \quad M_{P_{n+1}} = G_{m_1} \times \dots \times G_{m_r},$$

π_n and π_{n+1} are cuspidal automorphic representations of $M_{P_n}(\mathbb{A}), M_{P_{n+1}}(\mathbb{A})$ decomposing into tensor products

$$\pi_n = \pi_{n,1} \boxtimes \dots \boxtimes \pi_{n,k}, \quad \pi_{n+1} = \pi_{n+1,1} \boxtimes \dots \boxtimes \pi_{n+1,r}$$

respectively and we have set $\Pi_n = I_{P_n(\mathbb{A})}^{G_n(\mathbb{A})}(\pi_n)$, $\Pi_{n+1} = I_{P_{n+1}(\mathbb{A})}^{G_{n+1}(\mathbb{A})}(\pi_{n+1})$. We write $\chi_n \in \mathfrak{X}^*(G_n)$ and $\chi_{n+1} \in \mathfrak{X}^*(G_{n+1})$ for the cuspidal data determined by the pairs (M_{P_n}, π_n) and $(M_{P_{n+1}}, \pi_{n+1})$ respectively.

The representation Π is generic and we denote by $\mathcal{W}(\Pi, \psi_N)$ its Whittaker model with respect to the character ψ_N . Also, for every $\phi \in \Pi$ we define

$$W_\phi(g) := W_{E(\phi)}(g) = \int_{[N]} E(ug, \phi) \psi_N(u)^{-1} du, \quad g \in G(\mathbb{A}).$$

Note that $W_\phi \in \mathcal{W}(\Pi, \psi_N)$.

8.1.2. We now define two continuous linear forms λ and β_η as well as a continuous invariant scalar product $\langle \cdot, \cdot \rangle_{\text{Whitt}}$ on $\mathcal{W}(\Pi, \psi_N)$. Let $W \in \mathcal{W}(\Pi, \psi_N)$.

- By [JPSS83] [Jac09], the Zeta integral (already encountered in Chapter 7)

$$Z^{\text{RS}}(s, W) = \int_{N_H(\mathbb{A}) \backslash H(\mathbb{A})} W(h) |\det h|_{\mathbb{A}_E}^s dh.$$

converges for $\Re(s) \gg 0$ and extends to a meromorphic function on \mathbb{C} with no pole at $s = 0$. We set

$$\lambda(W) = Z^{\text{RS}}(0, W).$$

- For S a sufficiently large finite set of places of F , we put

$$\beta_\eta(W) = (\Delta_{G'}^{S,*})^{-1} L^{S,*}(1, \Pi, \text{As}_G) \int_{N'(F_S) \backslash \mathcal{P}'(F_S)} W(p_S) \eta_{G'}(p_S) dp_S$$

where we have set $L(s, \Pi, \text{As}_G) = L(s, \Pi_n, \text{As}^{(-1)^{n+1}}) L(s, \Pi_{n+1}, \text{As}^{(-1)^n})$.

- Similarly, for S a sufficiently large finite set of places of F , we put

$$\langle W, W \rangle_{\text{Whitt}} = (\Delta_G^{S,*})^{-1} L^{S,*}(1, \Pi, \text{Ad}) \int_{N(F_S) \backslash \mathcal{P}(F_S)} |W(p_S)|^2 dp_S$$

where we have set $L(s, \Pi, \text{Ad}) = L(s, \Pi_n \times \Pi_n^\vee) L(s, \Pi_{n+1} \times \Pi_{n+1}^\vee)$.

That the above expressions converge and are independent of S as soon as it is chosen sufficiently large (depending on the level of W) follow from [Fli88] and [JS81b]. Moreover, the inner form $\langle \cdot, \cdot \rangle_{\text{Whitt}}$ is $G(\mathbb{A})$ -invariant by [Ber84] and [Bar03].

The next result follows from works of Jacquet-Shalika [JS81b], Shahidi [Sha81] and Lapid-Offen [FLO12, Appendix A]. For completeness, we explain the deduction (see §2.7.2 for our normalization of the Petersson inner product).

Theorem 8.1.2.1. — [Jacquet-Shalika, Shahidi, Lapid-Offen] *We have*

$$\langle \phi, \phi \rangle_{\text{Pet}} = \langle W_\phi, W_\phi \rangle_{\text{Whitt}}$$

for every $\phi \in \Pi$.

Proof. — Let $\phi \in \Pi$. By the Iwasawa decomposition, for a suitable Haar measure on K we have

$$\langle \phi, \phi \rangle_{\text{Pet}} = \int_K \int_{[M_P]_0} |\phi(mk)|^2 \delta_P(m)^{-1} dm dk.$$

Set $N^P = N \cap M_P$ and

$$\phi_{N^P, \psi}(g) = \int_{[N^P]} \phi(ug) \psi_N(u)^{-1} du, \quad g \in G(\mathbb{A}).$$

Let \mathcal{P}^P be the product of mirabolic groups $\prod_{i=1}^k \mathcal{P}_{n_i} \times \prod_{j=1}^r \mathcal{P}_{m_j}$. It is a subgroup of M_P . According to Jacquet-Shalika [JS81b, §4] (see also [FLO12, p.265] or [Zha14a, Proposition 3.1]⁵), for S a sufficiently large finite set of places of F we have

(8.1.2.1)

$$\begin{aligned} \int_{[M_P]_0} |\phi(mk)|^2 \delta_P(m)^{-1} dm &= (\Delta_{M_P}^{S,*})^{-1} \prod_{i=1}^k \text{Res}_{s=1} L^S(s, \pi_{n,i} \times \pi_{n,i}^\vee) \prod_{j=1}^r \text{Res}_{s=1} L^S(s, \pi_{n+1,j} \times \pi_{n+1,j}^\vee) \\ &\quad \times \int_{N^P(F_S) \backslash \mathcal{P}^P(F_S)} |\phi_{N^P, \psi}(p_S k)|^2 \delta_P(p_S)^{-1} dp_S \end{aligned}$$

for every $k \in K$. On the other hand, by [FLO12, Proposition A.2] we have

(8.1.2.2)

$$\begin{aligned} &\int_K \int_{N^P(F_S) \backslash \mathcal{P}^P(F_S)} |\phi_{N^P, \psi}(p_S k)|^2 \delta_P(p_S)^{-1} dp_S dk \\ &= \frac{\text{vol}_{G(\mathbb{A}^S)}(K^S)}{\text{vol}_{M_P(\mathbb{A}^S)}(K^S \cap M_P(\mathbb{A}^S))} \int_{P(F_S) \backslash G(F_S)} \int_{N^P(F_S) \backslash \mathcal{P}^P(F_S)} |\phi_{N^P, \psi}(p_S g_S)|^2 \delta_P(p_S)^{-1} dp_S dg_S \\ &= (\Delta_G^{S,*})^{-1} \Delta_{M_P}^{S,*} \int_{N(F_S) \backslash \mathcal{P}(F_S)} |\mathbf{W}_S(p_S, \phi_{N^P, \psi})|^2 dp_S. \end{aligned}$$

where $\mathbf{W}_S : I_{P(F_S)}^{G(F_S)}(\mathcal{W}(\pi_S, \psi_{N,S})) \rightarrow \mathcal{W}(\Pi_S, \psi_{N,S})$ stands for the *Jacquet functional*, defined as the value at $s = 0$ of the holomorphic continuation of

$$\mathbf{W}_s^S(g_S, \phi') = \int_{(w_P^G)^{-1} N^P(F_S) w_P^G \backslash N(F_S)} \phi'(w_P^G u_S g_S) \delta_P(w_P^G u_S g_S)^s \psi_N(u_S)^{-1} du_S, \quad \Re(s) \gg 1$$

⁵Note that our normalization of the Petterson inner product is different from *loc. cit.*

for $g_S \in G(F_S)$ and $\phi' \in I_{P(F_S)}^{G(F_S)}(\mathcal{W}(\pi_S, \psi_{N,S}))$ where $w_P^G = w^P w^G$ with w^P (resp. w^G) the permutation matrix representing the longest element in the Weyl group of T in M_P (resp. in G). Finally, by [Sha81, Sect. 4], we have

$$(8.1.2.3) \quad \mathbf{W}^S(\phi_{N^P, \psi}) = \prod_{1 \leq i < j \leq k} L^S(1, \pi_{n,i} \times \pi_{n,j}^\vee) \prod_{1 \leq i < j \leq r} L^S(1, \pi_{n+1,i} \times \pi_{n+1,j}^\vee) W_\phi.$$

(Note that, as χ is generic, the Rankin-Selberg L -functions $L(s, \pi_{n,i} \times \pi_{n,j}^\vee)$ and $L(s, \pi_{n+1,i} \times \pi_{n+1,j}^\vee)$ are all regular at $s = 1$.) As, for every $s \in \mathbb{R}$,

$$\begin{aligned} L^S(s, \Pi \times \Pi^\vee) &= \prod_{i=1}^k L^S(s, \pi_{n,i} \times \pi_{n,i}^\vee) \times \prod_{j=1}^r L^S(s, \pi_{n+1,j} \times \pi_{n+1,j}^\vee) \\ &\quad \times \left| \prod_{1 \leq i < j \leq k} L^S(s, \pi_{n,i} \times \pi_{n,j}^\vee) \right|^2 \times \left| \prod_{1 \leq i < j \leq r} L^S(s, \pi_{n+1,i} \times \pi_{n+1,j}^\vee) \right|^2, \end{aligned}$$

we deduce from (8.1.2.1), (8.1.2.2) and (8.1.2.3) the identity of the Theorem. \square

8.1.3. Relative characters. — Let $\mathcal{B}_{P,\pi}$ be a K -basis of Π as in §2.8.3. We define the *relative character* I_Π of Π as the following functional on $\mathcal{S}(G(\mathbb{A}))$:

$$I_\Pi(f) = \sum_{\phi \in \mathcal{B}_{P,\pi}} \frac{\lambda(R(f)W_\phi) \overline{\beta_\eta(W_\phi)}}{\langle W_\phi, W_\phi \rangle_{\text{Whitt}}}, \quad f \in \mathcal{S}(G(\mathbb{A})),$$

where the series converges, and does not depend on the choice of $\mathcal{B}_{P,\pi}$, by Proposition 2.8.4.1.

8.1.4. For every $f \in \mathcal{S}(G(\mathbb{A}))$, we set

$$K_{f,\chi}^1(g) = \int_{[H]} K_{f,\chi}(h, g) dh \quad \text{and} \quad K_{f,\chi}^2(g) = \int_{[G']} K_{f,\chi}(g, g') \eta_{G'}(g') dg', \quad g \in [G],$$

where the above expressions are absolutely convergent by Lemma 2.9.8.1.3.

Recall that the notion of relevant \ast -generic cuspidal datum has been defined in §5.1.3 and that we have defined for any $\chi \in \mathfrak{X}$ a distribution I_χ (see Theorem 3.2.4.1).

Theorem 8.1.4.1. — *Let $f \in \mathcal{S}(G(\mathbb{A}))$ and $\chi \in \mathfrak{X}^*(G)$. Then,*

1. *If χ is not relevant, we have $K_{f,\chi}^2(g) = 0$ for every $g \in [G]$ and moreover*

$$I_\chi(f) = 0.$$

2. *If χ is relevant, we have*

$$I_\chi(f) = \int_{[G']} K_{f,\chi}^1(g') \eta_{G'}(g') dg'$$

where the right-hand side converges absolutely and moreover

$$I_\chi(f) = 2^{-\dim(A_P)} I_\Pi(f).$$

The rest of this chapter is devoted to the proof of Theorem 8.1.4.1. Until the end, we fix a function $f \in \mathcal{S}_\chi(G(\mathbb{A}))$.

8.2 Proof of Theorem 8.1.4.1

8.2.1. We fix a character η_G of $[G]$ whose restriction to $[G']$ is equal to $\eta_{G'}$ (such a character exists as the idèle class group of F is a closed subgroup of the idèle class group of E) and we set $\tilde{\chi} = \eta_G \otimes \chi^\vee \in \mathfrak{X}^*(G)$. We can write $\tilde{\chi}$ as $(\tilde{\chi}_n, \tilde{\chi}_{n+1})$ where $\tilde{\chi}_k \in \mathfrak{X}^*(G_k)$ for $k = n, n+1$. For every $g \in [G]$, we denote by $\tilde{K}_{f,\chi}(g, \cdot)$ the function $\eta_G K_{f,\chi}(g, \cdot)$. By Lemma 2.9.8.1.2 and (2.9.5.9), we have

$$(8.2.1.1) \quad \tilde{K}_{f,\chi}(g, \cdot) \in \mathcal{S}_{\tilde{\chi}}([G]) = \mathcal{S}_{\tilde{\chi}_n}([G_n]) \hat{\otimes} \mathcal{S}_{\tilde{\chi}_{n+1}}([G_{n+1}])$$

for all $g \in [G]$. Moreover, with the notation of Theorem 6.2.6.1, we have

$$(8.2.1.2) \quad K_{f,\chi}^2(g) = P_{G'_n} \hat{\otimes} P_{G'_{n+1}}(\tilde{K}_{f,\chi}(g, \cdot)).$$

8.2.2. The non-relevant case Assume that χ is not relevant. By definition of a relevant cuspidal data (see §5.1.3), at least one of $\tilde{\chi}_n, \tilde{\chi}_{n+1}$ is not distinguished (see §6.2.3 for the definition of distinguished). Hence, by Theorem 6.2.5.1 and Theorem 6.2.6.1, $P_{G'_k}$ vanishes identically on $\mathcal{S}_{\tilde{\chi}_k}([G_k])$ for $k = n$ or $k = n+1$. Thus, by (8.2.1.1) and (8.2.1.2), the function $K_{f,\chi}^2$ vanishes identically. By Theorem 3.3.7.1 applied to the expression (3.3.5.5), this implies $I_\chi(f) = 0$. This proves part 1. of Theorem 8.1.4.1.

8.2.3. Regularized Rankin-Selberg period and convergence From now on, we assume that χ is relevant. By Lemma 2.9.8.1.2, for every $g \in [G]$ the function $K_{f,\chi}(\cdot, g)$ belongs to $\mathcal{S}_\chi([G])$. Since χ is relevant, it is H -generic in the sense of §7.1.2 (this follows from the dichotomy of §4.1.2). Therefore, by Theorem 7.1.4.1, P_H extends to a continuous linear form on $\mathcal{T}_\chi([G])$ that we shall denote by P_H^* . By definition of this extension and of the linear form λ (see §8.1.2), for every $\phi \in \Pi$ we have

$$(8.2.3.3) \quad P_H^*(E(\phi)) = \lambda(W_\phi).$$

By Lemma 2.9.8.1.3 there exists $N \geq 0$ such that the function

$$g' \in [G'] \mapsto K_{f,\chi}(\cdot, g') \in \mathcal{T}_N([G])$$

is absolutely integrable. As

$$K_{f,\chi}^1(g) = P_H(K_{f,\chi}(\cdot, g)) = P_H^*(K_{f,\chi}(\cdot, g)),$$

combined with Theorem 3.3.7.1 applied to the expression (3.3.5.6), this shows at once that the expression

$$\int_{[G']} K_{f,\chi}^1(g') \eta_{G'}(g') dg'$$

converges absolutely, is equal to $I_\chi(f)$ and that

$$(8.2.3.4) \quad I_\chi(f) = P_H^* \left(\int_{[G']} K_{f,\chi}(\cdot, g') \eta_{[G']}(g') dg' \right) = P_H^*(K_{f,\chi}^2).$$

8.2.4. Spectral expression of $K_{f,\chi}^2$ Set $\tilde{\Pi} = \Pi^\vee \otimes \eta_G$. We may write $\tilde{\Pi}$ as a tensor product $\tilde{\Pi}_n \boxtimes \tilde{\Pi}_{n+1}$ and we let

$$\beta = \beta_n \hat{\otimes} \beta_{n+1} : \mathcal{W}(\tilde{\Pi}, \psi_N) = \mathcal{W}(\tilde{\Pi}_n, \psi_n) \hat{\otimes} \mathcal{W}(\tilde{\Pi}_{n+1}, \psi_{n+1}) \rightarrow \mathbb{C}$$

be the (completed) tensor product of the linear forms β_n, β_{n+1} defined in §6.2.4. Fix $g \in [G]$ and set $\mathbf{f}_g = \tilde{K}_{f,\chi}(g, \cdot)$. Since χ is relevant, $\tilde{\chi}_n$ and $\tilde{\chi}_{n+1}$ are both distinguished. Note that the linear map

$$\mathbf{f} \in \mathcal{S}([G]) \mapsto W_{\mathbf{f}, \tilde{\Pi}} := W_{\mathbf{f}_\Pi} \in \mathcal{W}(\tilde{\Pi}, \psi_N)$$

is the (completed) tensor product of the continuous linear maps $\mathbf{f} \in \mathcal{S}([G_k]) \mapsto W_{\mathbf{f}, \tilde{\Pi}_k} \in \mathcal{W}(\tilde{\Pi}_k, \psi_k)$ for $k = n, n+1$ (as can be checked directly on pure tensors). Therefore, by (8.2.1.1), (8.2.1.2), Theorem 6.2.5.1 and Theorem 6.2.6.1 we have

$$(8.2.4.5) \quad K_{f, \chi}^2(g) = 2^{-\dim(A_P)} \beta(W_{\mathbf{f}_g, \tilde{\Pi}}).$$

Let $\mathcal{B}_{P, \pi}$ be a K -basis Π as in §2.8.3. Then, we have $\mathbf{f}_{g, \tilde{\Pi}} = \sum_{\phi \in \mathcal{B}_{P, \pi}} \langle \mathbf{f}_g, \eta_G E(\bar{\phi}) \rangle_{[G]} \eta_G E(\bar{\phi})$ where the sum converges absolutely in $\mathcal{T}_N([G])$ for some $N \geq 0$. Hence,

$$W_{\mathbf{f}_g, \tilde{\Pi}} = \sum_{\phi \in \mathcal{B}_{P, \pi}} \langle \mathbf{f}_g, \eta_G E(\bar{\phi}) \rangle_{[G]} \eta_G \overline{W_\phi}$$

in $\mathcal{W}(\tilde{\Pi}, \psi_N)$. On the other hand, we easily check that $\beta(\eta_G \overline{W_\phi}) = \overline{\beta_\eta(W_\phi)}$ and

$$\langle \mathbf{f}_g, \eta_G E(\bar{\phi}) \rangle_{[G]} = \langle K_{f, \chi}(g, \cdot), E(\bar{\phi}) \rangle_{[G]} = E(R(f)\phi)(g)$$

for every $\phi \in \mathcal{B}_{P, \pi}$. Therefore, by (8.2.4.5), we obtain

$$(8.2.4.6) \quad K_{f, \chi}^2(g) = 2^{-\dim(A_P)} \sum_{\phi \in \mathcal{B}_{P, \pi}} E(R(f)\phi)(g) \overline{\beta_\eta(W_\phi)}.$$

Note that by Proposition 2.8.4.1, the series above is actually absolutely convergent in $\mathcal{T}_N([G])$ for some $N \geq 0$ (and not just pointwise).

8.2.5. End of the proof By (8.2.3.4), (8.2.3.3) and (8.2.4.6), we obtain

$$I_\chi(f) = 2^{-\dim(A_P)} \sum_{\phi \in \mathcal{B}_{P, \pi}} \lambda(R(f)W_\phi) \overline{\beta_\eta(W_\phi)}.$$

Using Theorem 8.1.2.1 and since $\mathcal{B}_{P, \pi}$ is an orthonormal basis of Π , this can be rewritten as

$$I_\chi(f) = 2^{-\dim(A_P)} \sum_{\phi \in \mathcal{B}_{P, \pi}} \frac{\lambda(R(f)W_\phi) \overline{\beta_\eta(W_\phi)}}{\langle W_\phi, W_\phi \rangle_{\text{Whitt}}} = 2^{-\dim(A_P)} I_\Pi(f)$$

and this ends the proof of Theorem 8.1.4.1 in the relevant case.

9 Flicker-Rallis functional computation

The goal of this chapter is to prove Theorem 1.3.2.3 of the introduction that states that two natural functionals are equal. This is established in Theorem 9.2.5.1. The bulk of the work is in proving its local avatar. The case of split algebra E/F amounts to comparing scalar products which was done in Appendix A of [FLO12], which is an inspiration for this chapter.

9.1 Local comparison

9.1.1. Let E/F be an étale quadratic algebra over a local field F . Let $\text{Tr}_{E/F} : E \rightarrow F$ be the trace map. As in Paragraph 6.1.2, let $\psi' : F \rightarrow \mathbb{C}^\times$ be a non-trivial additive character, $\tau \in E^\times$ an element of trace 0 and we set $\psi : E \rightarrow \mathbb{C}^\times$ to be $\psi(x) = \psi'(\text{Tr}(\tau x))$. We use ψ' and ψ to define autodual Haar measures on F and E respectively. The duality $F \times E/F \rightarrow \mathbb{C}^\times$ given by $(x, y) \mapsto \psi(xy)$ defines a unique Haar measure on E/F dual to the one on F . This measure on E/F coincides with the quotient measure.

9.1.2. We employ the convention of Section 2.3.1 to define Haar measures (with ψ denoted here as ψ'). Let $k = E$ or F . We define the following measures on $GL_n(k)$ and its subgroups

- On $GL_n(k)$ we set

$$dx = \frac{dx_{ij}}{|\det x|_k^n}$$

where $x = (x_{ij})$.

- On standard Levi subgroups of $GL_n(k)$ we set the product measure using the measure defined above.
- On (semi) standard unipotent subgroups $N(k) \subset GL_n(k)$ we set the additive measure dn_{ij} where n_{ij} run through coordinates of N .
- If $P(k)$ is a standard Levi subgroup of $GL_n(k)$ with the standard Levi decomposition $N(k)M(k)$ we have the right-invariant measure $dp := dndm$ on $P(k)$ and the left invariant measure $\delta_{P_k}^{-1}dp$ where $\delta_{P_k} : P(k) \rightarrow \mathbb{R}_{>0}^\times$ is the Jacobian homomorphism for the adjoint action of $P(k)$ on $N(k)$.

With this normalization, we have for all $f \in C_c^\infty(GL_n(k))$

$$\int_{GL_n(k)} f dg = \int_{P(k)} \int_{\overline{N}(k)} f(pn) \delta_{P_k}(p)^{-1} dp dn.$$

where \overline{N} is the unipotent radical of the opposite parabolic to P .

9.1.3. We will use the notation introduced in Section 4 with some changes. All groups considered in this section are subgroups of $G_n = \text{Res}_{E/F} GL_n$. We write simply G for G_n , P_0 for the fixed minimal parabolic subgroup of G and N_0 for its unipotent radical. In order to be as compatible with Appendix A of [FLO12] as possible, instead of $G' = GL_n$ (defined over F) we write $G_F = GL_n$ and for any subgroup H of G we write H_F for $H \cap G_F$. We will often identify a group with its F points in this section.

9.1.4. We define the character $\psi : N_0 \rightarrow \mathbb{C}^\times$ as follows. Write $n \in N_0$ as

$$n = \begin{pmatrix} 1 & n_{12} & n_{13} & \dots & n_{1n} \\ 0 & 1 & n_{23} & \dots & n_{2n} \\ 0 & \ddots & \ddots & \ddots & n_{2n} \\ 0 & \ddots & \ddots & 1 & n_{n-1n} \\ 0 & \dots & 0 & 0 & 1 \end{pmatrix}, \quad n_{ij} \in E$$

and set $\psi(n) = \psi((-1)^n(n_{12} + n_{23} + \cdots + n_{n-1n}))$. This is the same character as the one from 6.1.2. By restriction, ψ defines a character of $N_0 \cap M$ for all standard Levi subgroups M .

9.1.5. We denote by $\mathcal{P} = \mathcal{P}_n$ the mirabolic subgroup of G defined as the stabilizer of the row vector $(0 \ \dots \ 0 \ 1)$.

Define the following functional on $C^\infty(N_0 \backslash \mathcal{P}, \psi) = \{f \in C^\infty(\mathcal{P}) \mid f(nx) = \psi(n)f(x), \ n \in N_0, \ x \in \mathcal{P}\}$

$$\beta(\varphi) = \beta_G(\varphi) = \int_{N_{0,F} \backslash \mathcal{P}_F} \varphi(p) \, dp.$$

Note that the integral is well defined as ψ is trivial on $N_{0,F}$. In the same way, we define β_M for all standard Levi subgroups M of G .

9.1.6. Let $\Pi_{gen}(G)$ be the set of irreducible generic complex representations of $G = GL_n(E)$. Let $\mathcal{W}(\pi) = \mathcal{W}^\psi(\pi)$ be the space of the Whittaker model of $\pi \in \Pi_{gen}(G)$ with respect to the character ψ . Let $\delta_g^\pi = \delta_g : \mathcal{W}(\pi) \rightarrow \mathbb{C}$ be the evaluation at $g \in G$. The group G acts on $\mathcal{W}(\pi)$ by right multiplication.

Fix $P = MN \in \mathcal{F}^G(P_0)$. Let w_M be the element in the Weyl group of G such that $w_M M w_M^{-1}$ is a standard Levi and the longest for this property. Let $P^w = N^w M^w \in \mathcal{F}^G(P_0)$ be the group whose Levi component is $M^w = w_M M w_M^{-1}$.

For $\sigma \in \Pi_{gen}(M)$ let $\text{Ind}_P^G(\mathcal{W}(\sigma))$ be the normalized (smooth) induction to G , from $\mathcal{W}(\sigma)$, seen as a representation of P via the natural map $P \rightarrow M$. For $\varphi \in \text{Ind}_P^G(\mathcal{W}(\sigma))$ let

$$\mathbf{W}(g, \varphi) = \int_{N^w} \delta_e^\sigma(\varphi(w_M^{-1} u' g)) \psi^{-1}(u') \, du'.$$

This is the so called Jacquet's integral. We have then that $\mathbf{W}_e(\varphi) := \mathbf{W}(e, \varphi)$ is a Whittaker functional on $\text{Ind}_P^G(\mathcal{W}(\sigma))$.

9.1.7. For $\sigma \in \Pi_{gen}(M)$ and $\varphi \in \text{Ind}_P^G(\mathcal{W}(\sigma))$ let

$$\beta'(\varphi) = \int_{P_F \backslash G_F} \beta_M(\varphi(g)) \, dg.$$

Theorem 9.1.7.1. — *Let $\sigma \in \Pi_{gen}(M)$. Suppose σ is unitary. We have then*

$$\beta'(\varphi) = \beta_G(\mathbf{W}(\varphi)).$$

Proof. — We follow very closely Appendix A of [FLO12].

We reduce the proof to the case P is maximal. Let $Q = LV \supset P$ be maximal and suppose the assertion holds for $M = M_P$. Then

$$\beta'(\varphi) = \int_{P_F \backslash G_F} \beta_M(\varphi(g)) \, dg = \int_{Q_F \backslash G_F} \int_{P_F \backslash Q_F} \delta_{Q_F}^{-1}(q) \beta_M(\varphi(qg)) \, dq \, dg.$$

The inner integral on the RHS by induction hypothesis equals

$$\beta_L(g \cdot \mathbf{W}^L(\varphi)).$$

If we let $\varphi'(g) = g \cdot \mathbf{W}^L(\varphi) \in \mathcal{W}(\text{Ind}_{L \cap P}^L(\mathcal{W}(\sigma)))$ then $\varphi' \in \text{Ind}_Q^G(\mathcal{W}(\text{Ind}_{L \cap P}^L(\mathcal{W}(\sigma))))$ and so by assumption and transitivity of Jacquet's integral we obtain

$$\int_{Q_F \backslash G_F} \beta_L(g \cdot \mathbf{W}^L(\varphi)) \, dg = \beta_G(\mathbf{W}(\varphi)).$$

Assume then that $P = MN$ is maximal of type (n_1, n_2) . In [FLO12], the authors use U instead of N . We will consequently use N in place of U here. Write $M = M_1 \times M_2$ with $M_i \cong \text{Res}_{E/F} GL_{n_i}$, M_1 being in the upper and M_2 in the lower diagonal.

Let

$$w = w_M^{-1} = \begin{pmatrix} 0 & I_{n_1} \\ I_{n_2} & 0 \end{pmatrix}.$$

Let $P' = M'N'$ be of type (n_2, n_1) so that $M' = M'_2 \times M'_1$ with $M'_i \cong \text{Res}_{E/F} GL_{n_i}$, M'_2 being in the upper and M'_1 in the lower diagonal. Let \mathcal{P}'_i be the mirabolic subgroup of M'_i . Let N'_i be the maximal upper triangular unipotent of M'_i , similarly without $'$. Note that $\mathcal{P}_i = w\mathcal{P}'_i w^{-1}$ etc.

We identify N' with the group of $n_2 \times n_1$ matrices. Let

$$C_i = \{I_n + \xi \mid \xi \text{ column vector of size } n_2 \text{ in the } i\text{-th column}\} \subset N', \quad i = n_2 + 1, \dots, n.$$

Let

$$R_i = \{I_n + \xi \mid \xi \text{ row vector of size } n_2 \text{ in the } i\text{-th row}\} \subset \overline{N}', \quad i = n_2 + 1, \dots, n.$$

We can identify C_i and R_j with E^n which induces a pairing between C_i and R_{i-1} that we will denote $\langle \cdot, \cdot \rangle_i$.

We note some obvious facts

- The groups R_i (resp. C_i) commute with each other and are normalized by M'_2 and M'_1 .
- The commutator set $[C_i, R_j]$ is contained in N'_1 for $j < i$.

We define the following groups

1. $X_i = C_{i+1} \cdots C_n$.
2. $Y_i = R_{n_2+1} \cdots R_{i-1}$.
3. $V_i = N'_1 X_i Y_i$. This is a unipotent group.
4. $V'_i = N'_1 X_{i-1} Y_i \supset V_i$. This is a unipotent group.
- 5.

$$S_i = \begin{cases} M'_2 V_i, & i > n_2, \\ \mathcal{P}'_2 N'_1 N', & i = n_2 \end{cases}$$

6. $S'_i = M'_2 V'_i$ for $i > n_2$.

Note that

- $S'_i = C_i S_i$ for $i > n_2$ as well as $S'_i = R_{i-1} S_{i-1}$ for $i > n_2 + 1$.
- Let δ_i and δ'_i be modular characters of S_i and S'_i respectively. It follows that $\delta'_i | S_i = |\det|_E \delta_i$ and $\delta'_i | S_{i-1} = |\det|_E^{-1} \delta_{i-1}$ in the above range.
- We have $\delta_i = |\det|_E^{n+n_2-2i+1}$ for $i \geq n_2$.

Let $\sigma = \sigma_1 \otimes \sigma_2$ be an irreducible representation of M , with $\sigma_i \in \Pi_{\text{gen}}(GL_{n_i}(E))$. We view σ_2 as a representation of M'_2 as well. Let us define

$$\begin{cases} \mathcal{A}_i = \text{Ind}_{S_i}^{\mathcal{P}_i} (\mathcal{W}(\sigma) \otimes \psi_i), & i = n_2 + 1, \dots, n, \\ \mathcal{A}_i = \text{Ind}_{N'_0}^{\mathcal{P}_i} \psi, & i = n_2. \end{cases}$$

Here, ψ_i is the character of V_i - the unipotent radical of S_i - whose restriction to $X_i Y_i$ is trivial and that coincides with ψ on N'_1 .

Explicitly, for $i > n_2$ we have

$$\mathcal{A}_i = \{\varphi : \mathcal{P} \rightarrow \mathcal{W}(\sigma_2) \mid \varphi(mvg) = \left(\frac{\delta_i(m)}{|\det m|} \right)^{1/2} \psi_i(v) \sigma(m) \varphi(g), \quad g \in \mathcal{P}, \quad m \in M'_2, \quad v \in V_i.\}$$

We also denote \mathcal{A}_i^2 the L^2 -induction version of the above as in [FLO12]. Note that

- $\mathcal{A}_i = \text{Ind}_{S'_i}^{\mathcal{P}}(\text{Ind}_{S_i}^{S'_i}((\mathcal{W}(\sigma) \otimes \psi_i)))$ for $i > n_2$.
- $\mathcal{A}_i = \text{Ind}_{S'_i}^{\mathcal{P}}(\text{Ind}_{S_{i-1}}^{S'_i}((\mathcal{W}(\sigma) \otimes \psi_{i-1})))$ for $i > n_2 + 1$.
- $\mathcal{A}_{n_2} = \text{Ind}_{S'_{n_2+1}}^{\mathcal{P}}(\text{Ind}_{N_0}^{S'_{n_2+1}} \psi)$.

For any $i > n_2$ the restriction map to C_i identifies $\text{Ind}_{S_i}^{S'_i}((\mathcal{W}(\sigma) \otimes \psi_i))$ with $C^\infty(C_i, \mathcal{W}(\sigma_2))$ because $S_i/S_{i-1} = C_i$. Let us denote $\varphi \mapsto \varphi|_{C_i}$ the restriction map and ι_{C_i} the map in the reverse order. Similarly, restriction to R_{i-1} identifies $\text{Ind}_{S_{i-1}}^{S'_i}((\mathcal{W}(\sigma) \otimes \psi_{i-1}))$ with $C^\infty(R_{i-1}, \mathcal{W}(\sigma_2))$. Let us denote $\varphi \mapsto \varphi|_{R_{i-1}}$ the restriction map and $\iota_{R_{i-1}}$ the map in the reverse order.

Given that C_i and R_i are in duality we have a Fourier transform

$$\mathcal{F}'_i : L^2(C_i, \overline{\mathcal{W}(\sigma_2)}) \rightarrow L^2(R_{i-1}, \overline{\mathcal{W}(\sigma_2)})$$

where $\overline{\mathcal{W}(\sigma_2)}$ is the L^2 completion of $\mathcal{W}(\sigma_2)$.

Lemma 9.1.7.2. — *For $i = n, \dots, n_2 + 2$, the above Fourier transform induces a map*

$$\mathcal{B}_i : \mathcal{A}_i^2 = \text{Ind}_{S'_i}^{\mathcal{P}}(\text{Ind}_{S_i}^{S'_i}((\mathcal{W}(\sigma) \otimes \psi_i))) \rightarrow \mathcal{A}_{i-1}^2 = \text{Ind}_{S'_i}^{\mathcal{P}}(\text{Ind}_{S_{i-1}}^{S'_i}((\mathcal{W}(\sigma) \otimes \psi_i)))$$

induced from the equivalence $\text{Ind}_{S_i}^{S'_i}((\mathcal{W}(\sigma) \otimes \psi_i)) \rightarrow \text{Ind}_{S_{i-1}}^{S'_i}((\mathcal{W}(\sigma) \otimes \psi_{i-1}))$ given by $\varphi \mapsto \iota_{R_{i-1}}(\mathcal{F}'_i(\varphi|_{C_i}))$. It is an equivalence of unitary representations.

Similarly, we have the map

$$\mathcal{F}_{n_2+1} : \text{Ind}_{S'_{n_2+1}}^{S'_{n_2+1}}((\mathcal{W}(\sigma) \otimes \psi_{n_2+1})) \rightarrow \text{Ind}_{N_0}^{S'_{n_2+1}} \psi$$

given by

$$\mathcal{F}_{n_2+1} \varphi(vm) = \psi(v) |\det m|_E^{1/2} \hat{\varphi}(\chi_m)(m), \quad m \in M'_2, \quad v \in V'_{n_2+1} = N'_1 N'$$

where

$$\chi_m : C_{n_2+1} \rightarrow \mathbb{C}^\times, \quad \chi_m(c) = \psi(mcm^{-1}), \quad \hat{\varphi}(\chi) = \int_{C_{n_2+1}} \varphi(c) \chi(c) dc.$$

Lemma 9.1.7.3. — *The above Fourier transform induces the equivalence of unitary representations*

$$\mathcal{B}_{n_2+1} : \mathcal{A}_{n_2+1}^2 = \text{Ind}_{S'_{n_2+1}}^{\mathcal{P}}(\text{Ind}_{S_{n_2+1}}^{S'_{n_2+1}}((\mathcal{W}(\sigma) \otimes \psi_{n_2+1}))) \rightarrow \mathcal{A}_{n_2}^2 = \text{Ind}_{S'_{n_2+1}}^{\mathcal{P}}(\text{Ind}_{N_0}^{S'_{n_2+1}}).$$

For $i = n, \dots, n_2 + 1$ let $\beta_i : \mathcal{A}_i \rightarrow \mathbb{C}$ be the following functional

$$\beta_i(\varphi) = \int_{S_{i,F} \setminus \mathcal{P}_F} \beta_{M'_2}(\varphi(p)) dp.$$

We also set $\beta_{n_2} = \beta_G$ on $\mathcal{A}_{n_2} = \text{Ind}_{N_0}^{\mathcal{P}} \psi$.

Lemma 9.1.7.4. — *For $i = n, \dots, n_2 + 2$ and $\varphi \in \mathcal{A}_i$ have*

$$\beta_i(\varphi) = \beta_{i-1}(\mathcal{B}_i(\varphi)).$$

Proof. — By equivariance property of \mathcal{B}_i it is enough to show the equality between $\int_{C_{i,F}} \beta_{M'_2}(\varphi(c)) dc$ and $\int_{R_{i-1,F}} \beta_{M'_2}(\mathcal{B}_i(\varphi)(r)) dr$. The duality $\langle \cdot, \cdot \rangle_i$ between C_i and R_{i-1} restricts to a duality between $C_i/C_{i,F}$ and $R_{i-1,F}$. We have thus

$$\begin{aligned} \int_{R_{i-1,F}} \beta_{M'_2}(\mathcal{B}_i(\varphi)(r)) dr &= \int_{R_{i-1,F}} \beta_{M'_2} \left(\int_{C_i} \varphi(c) \psi(\langle c, r \rangle_i) dc \right) dr = \\ &= \beta_{M'_2} \left(\int_{R_{i-1,F}} \int_{C_i/C_{i,F}} \left(\int_{C_i,F} \varphi(c^- + c) \psi(\langle c^-, r \rangle_i) dc \right) dc^- dr \right) = \int_{C_{i,F}} \beta_{M'_2}(\varphi(c)) dc. \end{aligned}$$

□

Lemma 9.1.7.5. — For $\varphi \in \mathcal{A}_{n_2+1}$ we have

$$\beta_{n_2+1}(\varphi) = \beta_{n_2}(\mathcal{B}_{n_2+1}(\varphi)).$$

Proof. — Again, it is enough to show the equality between $\int_{C_{n_2+1,F}} \beta_{M'_2}(\varphi(c)) dc$ and $\int_{N'_{2,F} \setminus M'_{2,F}} \mathcal{F}_{n_2+1}(\varphi)(m) dm$. We have

$$\begin{aligned} \int_{N'_{2,F} \setminus M'_{2,F}} \mathcal{F}_{n_2}(\varphi)(m) dm &= \int_{N'_{2,F} \setminus M'_{2,F}} \int_{C_{n_2+1}} \varphi(c)(m) \psi(mcm^{-1}) dc | \det m |_F dm = \\ &= \int_{\mathcal{P}'_{n_2,F} \setminus M'_{2,F}} \int_{C_{n_2+1}/C_{n_2+1,F}} \int_{C_{n_2+1,F}} \int_{N'_{2,F} \setminus \mathcal{P}'_{n_2,F}} \varphi(c+c^-)(pm) \psi(mc^-m^{-1}) | \det m |_F dp dc dc^- dm = \\ &= \int_{C_{n_2+1,F}} \int_{N'_{2,F} \setminus \mathcal{P}'_{n_2,F}} \varphi(c)(p) dp dc = \int_{C_{n_2+1,F}} \beta_{M'_2}(\varphi(c)) dc. \end{aligned}$$

□

Let $\varphi \in \text{Ind}_P^G(\mathcal{W}(\sigma_1 \otimes \sigma_2))$. For $m \in M_1$, let $\delta_m^1 : \mathcal{W}(\sigma_1 \otimes \sigma_2) \rightarrow \mathcal{W}(\sigma_2)$ be the evaluation map in the first variable. Define for $p \in \mathcal{P}$

$$\varphi_n(p) = \delta_e^1 \varphi(wp) \in \mathcal{A}_n.$$

We have then

$$\int_{P \setminus G} \|\varphi(g)\|_{L^2(\mathcal{W}(\sigma_1 \otimes \sigma_2))}^2 dg = \|\varphi_n\|_{\mathcal{A}_n}^2.$$

We set $\varphi_{i-1} = \mathcal{B}_i(\varphi_i)$ for $i = n, \dots, n_2 + 1$. As shown at the end of the Appendix A.3 of [FLO12], we have

$$(9.1.7.1) \quad \varphi_{n_2} = \mathbf{W}(\varphi).$$

Lemma 9.1.7.6. — For $\varphi \in \text{Ind}_P^G(\mathcal{W}(\sigma_1 \otimes \sigma_2))$, we have

$$\beta'(\varphi) = \beta_n(\varphi_n).$$

Proof. — Indeed

$$\begin{aligned} \beta'(\varphi) &= \int_{P_F \setminus G_F} \beta_M(\varphi(g)) dg = \int_{P_F \setminus G_F} \beta_M(\varphi(gw)) dg = \int_{\overline{N}_F} \beta_M(\varphi(uw)) du = \int_{N'_F} \beta_M(\varphi(wu')) du' = \\ &= \int_{N'_F} \int_{N_{1,F} \setminus \mathcal{P}_{1,F}} \beta_{M_2}(\delta_{m_1}^1(\varphi(wu'))) dm_1 du' = \int_{N'_F} \int_{N_{1,F} \setminus \mathcal{P}_{1,F}} \beta_{M_2}(\delta_e^1(\varphi(m_1 wu'))) \delta_P^{1/2}(m_1) dm_1 du' = \\ &= \int_{N'_F} \int_{N'_{1,F} \setminus \mathcal{P}'_{1,F}} \beta_{M'_2}(\delta_e^1(\varphi(wm'_1 u'))) \delta_{P'}^{-1}(m'_1) dm'_1 du' = \int_{S_{n,F} \setminus \mathcal{P}_F} \beta_{M'_2}(\delta_e^1(\varphi(wp))) dp. \end{aligned}$$

□

Now combining the equality (9.1.7.1) with Lemmas 9.1.7.4, 9.1.7.5, 9.1.7.6 we obtain the desired equality at least when computations in these Lemmas are justified. Taking $\varphi \in \text{Ind}_P^G(\mathcal{W}(\sigma_1 \otimes \sigma_2))$ supported on the big cell PwP' we can see that all integrals are absolutely convergent. By multiplicity (at most) one [Fli91, AG09], the Theorem 9.1.7.1 follows. □

9.2 Global comparison

9.2.1. We go back to the global setting and notation introduced in §3.1.

9.2.2. We normalize all local and global measures as in §2.3, with respect to a fixed character $\psi' : F \backslash \mathbb{A} \rightarrow \mathbb{C}^\times$. We have the quadratic character $\eta : F^\times \backslash \mathbb{A}^\times \rightarrow \mathbb{C}^\times$ associated to E/F and the associated character $\eta_{G'}$ of $G'(\mathbb{A})$ as defined in Paragraph 3.1.8.

9.2.3. As in §6.1.2, we also fix a non-trivial additive character $\psi : E \backslash \mathbb{A}_E \rightarrow \mathbb{C}^\times$, trivial on \mathbb{A} which is then used to define a non-degenerate character ψ_N of the maximal unipotent subgroup of $G(\mathbb{A})$ as in the beginning of 8.

9.2.4. Let $\chi \in \mathfrak{X}^*(G)$ (c.f. §5.1.3) and let (M, π) represent χ . Set $\Pi = \text{Ind}_{P(\mathbb{A})}^G(\mathbb{A})(\pi)$.

9.2.5. The comparison —

Theorem 9.2.5.1. — *For all $\phi \in \Pi$ we have*

$$J_\eta(\phi) = \beta_\eta(W_\phi)$$

where

- J_η is defined in 5.1.2.1;
- β_η is defined in 8.1.2.
- $W_\phi \in \mathcal{W}(\Pi, \psi_N)$ is defined in 8.1.1.

Proof. — The proof is essentially the same as of Theorem 8.1.2.1. The only difference is that the natural analogue of (8.1.2.1) is provided by Proposition 3.2 of [Zha14a] and the analogue of (8.1.2.2) is established invoking Theorem 9.1.7.1. □

Corollary 9.2.5.2. — *We have the equality of distributions on $\mathcal{S}(G(\mathbb{A}))$*

$$I_{P, \pi} = I_\Pi$$

where

1. $I_{P, \pi}$ is defined in §5.1.5.
2. I_Π is defined in §8.1.3.

Proof. — Looking at definitions of $I_{P, \pi}$ and I_Π , taking into consideration Theorem 8.1.2.1 and Theorem 9.2.5.1 above, we see that we need to establish for all $\phi \in \Pi$

$$\lambda(W_\phi) = I(\phi, 0)$$

where $\lambda = Z^{RS}(0, \cdot)$ is defined in §8.1.2 and $I(\phi, 0)$ is given by Proposition 5.1.4.1. This equality is precisely Theorem 1.1 of [IY15]. □

10 Proofs of the Gan-Gross-Prasad and Ichino-Ikeda conjectures

10.1 Identities among some global relative characters

10.1.1. Besides notation of Chapters 2 and 3, we shall use notation of Section 1. We fix an integer $n \geq 1$ and we will omit the subscript n : we will write \mathcal{H} for \mathcal{H}_n .

10.1.2. Relative characters for unitary groups. — Let $h \in \mathcal{H}$ be a Hermitian form. Let σ be an irreducible cuspidal automorphic subrepresentation of the group U_h . We define the relative character J_σ^h by

$$J_\sigma^h(f) = \sum_{\varphi} \mathcal{P}_h(\pi(f)\varphi) \overline{\mathcal{P}_h(\varphi)}, \quad \forall f \in \mathcal{S}(U_h(\mathbb{A}))$$

where φ runs over a K_h -basis (see 2.8.3) for some maximal compact subgroup $K_h \subset U_h(\mathbb{A})$. The period \mathcal{P}_h are those defined in 1.1.5. For any subset $\mathfrak{X}_0 \subset \mathfrak{X}(U_h)$ of cuspidal data which do not come from proper Levi subgroups (that is they are represented by pairs (U_h, τ) where τ is a cuspidal automorphic representation) we define more generally

$$(10.1.2.1) \quad J_{\mathfrak{X}_0}^h(f) = \sum_{\chi \in \mathfrak{X}_0} \sum_{\sigma} J_{\sigma}^h(f)$$

where the inner sum is over the set of the constituents σ of some decomposition of $L_{\chi}^2([U_h])$ (see §2.9.2.1) into irreducible subrepresentations. One can show that the double sum is absolutely convergent (see e.g. [Beu16, Proposition A.1.2]).

10.1.3. Let $V_{F,\infty} \subset S_0 \subset V_F$ be a finite set of places containing all the places that are ramified in E . For every $v \in V_F$, we set $E_v = E \otimes_F F_v$ and when $v \notin V_{F,\infty}$ we denote by $\mathcal{O}_{E_v} \subset E_v$ its ring of integers. Let $\mathcal{H}^\circ \subset \mathcal{H}$ be the (finite) subset of Hermitian spaces of rank n over E that admits a selfdual \mathcal{O}_{E_v} -lattice for every $v \notin S_0$.

For each $h \in \mathcal{H}^\circ$, the group U_h is naturally defined over $\mathcal{O}_F^{S_0}$ and we fix a choice of such a model. Since we are going to consider invariant distribution, this choice is irrelevant. We define the open compact subgroups $K_h^\circ = \prod_{v \notin S_0} U_h(\mathcal{O}_v)$ and $K^\circ = \prod_{v \notin S_0} G(\mathcal{O}_v)$ respectively of $U_h(\mathbb{A}^{S_0})$ and $G(\mathbb{A}^{S_0})$.

Let $v \notin S_0$. We denote by $\mathcal{S}^\circ(U_h(F_v))$, resp. $\mathcal{S}^\circ(G(F_v))$, the spherical Hecke algebra⁶ of complex functions on $U_h(F_v)$ (resp. $G(F_v)$) that are $U_h(\mathcal{O}_v)$ -bi-invariant (resp. $G(\mathcal{O}_v)$ -bi-invariant) and compactly supported.

We have the base change homomorphism

$$BC_{h,v} : \mathcal{S}^\circ(G(F_v)) \rightarrow \mathcal{S}^\circ(U_h(F_v)).$$

We denote by $\mathcal{S}^\circ(U_h(\mathbb{A}^{S_0}))$, resp. $\mathcal{S}^\circ(G(\mathbb{A}^{S_0}))$, the restricted tensor product of $\mathcal{S}^\circ(U_h(F_v))$, resp. $\mathcal{S}^\circ(G(F_v))$, for $v \notin S_0$. We have also a global base change homomorphism given by $BC_h^{S_0} = \otimes_{v \notin S_0} BC_{h,v}$.

We also denote by $\mathcal{S}^\circ(G(\mathbb{A})) \subset \mathcal{S}(G(\mathbb{A}))$ and $\mathcal{S}^\circ(U_h(\mathbb{A})) \subset \mathcal{S}(U_h(\mathbb{A}))$, for $h \in \mathcal{H}^\circ$, the subspaces of functions that are respectively bi- K° -invariant and bi- K_h° -invariant.

10.1.4. Transfer. — Let $h \in \mathcal{H}^\circ$. We shall say that $f_{S_0} \in \mathcal{S}(G(F_{S_0}))$ and $f_{S_0}^h \in \mathcal{S}(U_h(F_{S_0}))$ are transfers if the functions f_{S_0} and $f_{S_0}^h$ have matching regular orbital integrals in the sense of Definition 4.4 of [BLZZ19]. The Haar measures on the F_{S_0} -points of the involved groups are those defined in §2.3.3.

10.1.5. Let P be a standard parabolic subgroup of G and π be a cuspidal automorphic representation of M_P . Let $\chi \in \mathfrak{X}(G)$ be the class of the pair (M_P, π) . We assume henceforth that χ is a generic relevant cuspidal datum in the sense of §5.1.3.

⁶The product structure is given by the convolution where the Haar measure is normalized so that the characteristic functions of $U_h(\mathcal{O}_v)$ and $G(\mathcal{O}_v)$ are units.

Set $\Pi = \text{Ind}_P^G(\pi)$ for the corresponding parabolically induced representation. The assumption that χ is generic and relevant means exactly that Π is a Hermitian Arthur parameter (see §1.1.3). Moreover, we assume, as we may, that S_0 has been chosen such that Π admits K° -fixed vectors.

Attached to these data, we have three distributions denoted by I_χ , $I_{P,\pi}$ and I_Π . The first is constructed as a contribution of the Jacquet-Rallis trace formula and it is defined in Theorem 3.2.4.1. The second and third are relative characters built respectively in §5.1.5 and §8.1.3. The bulk of the paper was devoted to the proof of the following identities (see Theorem 5.2.1.1, Theorem 8.1.4.1 and Corollary 9.2.5.2)

$$(10.1.5.2) \quad I_\chi = 2^{-\dim(\mathfrak{a}_P)} I_{P,\pi} = 2^{-\dim(\mathfrak{a}_P)} I_\Pi.$$

10.1.6. Let S'_0 be the union of $S_0 \setminus V_{F,\infty}$ and the set of all finite places of F that are inert in E .

We define $\mathfrak{X}_0^h \subset \mathfrak{X}(U_h)$ as the set of equivalence classes of pairs (U_h, σ) where σ a cuspidal automorphic representation of $U_h(\mathbb{A})$ that satisfies the following conditions:

- σ is K_h° -unramified;
- for all $v \notin S'_0 \cup V_{F,\infty}$ the (split) base change of σ_v is Π_v .

Proposition 10.1.6.1. —

Let $f \in \mathcal{S}^\circ(G(\mathbb{A}))$ and $f^h \in \mathcal{S}^\circ(U_h(\mathbb{A}))$ for every $h \in \mathcal{H}^\circ$. Assume that the following properties are satisfied for every $h \in \mathcal{H}^\circ$:

1. $f = (\Delta_H^{S_0,*} \Delta_{G'}^{S_0,*}) f_{S_0} \otimes f^{S_0}$ with $f_{S_0} \in \mathcal{S}(G(F_{S_0}))$ and $f^{S_0} \in \mathcal{S}^\circ(G(\mathbb{A}^{S_0}))$.
2. $f^h = (\Delta_{U_h'}^{S_0})^2 f_{S_0}^h \otimes f^{h,S_0}$ with $f_{S_0}^h \in \mathcal{S}(U_h(F_{S_0}))$ and $f^{h,S_0} \in \mathcal{S}^\circ(U_h(\mathbb{A}^{S_0}))$.
3. The functions f_{S_0} and $f_{S_0}^h$ are transfers.
4. $f^{h,S_0} = BC_h^{S_0}(f^{S_0})$
5. The function f^{S_0} is a product of a smooth compactly supported function on the restricted product $\prod'_{v \notin S'_0} G(F_v)$ by the characteristic function of $\prod_{v \in S'_0 \setminus S_0} G(\mathcal{O}_v)$.

Then we have:

$$(10.1.6.3) \quad \sum_{h \in \mathcal{H}^\circ} J_{\mathfrak{X}_0^h}^h(f^h) = 2^{-\dim(\mathfrak{a}_P)} I_\Pi(f) = 2^{-\dim(\mathfrak{a}_P)} I_{P,\pi}(f).$$

Remark 10.1.6.2. — If the assumptions hold for the set S_0 , it also holds for any large enough finite set containing S_0 : this follows from the Jacquet-Rallis fundamental lemma (see [Yun11] and [BP]) and the simple expression of the transfer at split places (see [Zha14b, proposition 2.5]). We leave it to the reader to keep track of the different choices of Haar measures in these references.

Proof. —

The proof follows the same lines as the proof of [BLZZ19, Theorem 1.7]. For the convenience of the reader, we recall the main steps.

In Theorem 3.2.4.1 we defined a distribution I on $\mathcal{S}(G(\mathbb{A}))$: this is the “Jacquet-Rallis trace formula” for G . We have an analogous distribution J^h on $\mathcal{S}(U_h(\mathbb{A}))$ for each $h \in \mathcal{H}$: it is defined in [Zyd20, théorème 0.3] for compactly supported functions and extended to the Schwartz space in [CZ, §1.1.3 and théorème 15.2.3.1]. Note that, by the Jacquet-Rallis fundamental lemma [Yun11], [BP], for every $h \in \mathcal{H} \setminus \mathcal{H}^\circ$ there exists a place $v \in S'_0 \setminus S_0$ such that the characteristic function

$\mathbf{1}_{G(\mathcal{O}_v)}$ admits the zero function on $U_h(F_v)$ as a transfer. Therefore, by [CZ, théorème 1.6.1.1], the hypotheses of the proposition imply:

$$(10.1.6.4) \quad I(f) = \sum_{h \in \mathcal{H}^\circ} J^h(f^h).$$

We will denote by $\mathcal{M}^{S'_0}(G(\mathbb{A}))$, resp. $\mathcal{M}^{S'_0}(U_h(\mathbb{A}))$, the algebra of S'_0 -multipliers defined in [BLZZ19, definition 3.5] relatively to the subgroup $\prod_{v \notin S'_0} G(\mathcal{O}_v)$, resp. $\prod_{v \notin S'_0} U_h(\mathcal{O}_v)$. Any multiplier $\mu \in \mathcal{M}^{S'_0}(G(\mathbb{A}))$, resp. $\mu \in \mathcal{M}^{S'_0}(U_h(\mathbb{A}))$, gives rise to a linear operator $\mu*$ of the algebra $\mathcal{S}^\circ(G(\mathbb{A}))$, resp. $\mathcal{S}^\circ(U_h(\mathbb{A}))$ and for every admissible irreducible representation π of $G(\mathbb{A})$, resp. of $U_h(\mathbb{A})$, there exists a constant $\mu(\pi) \in \mathbb{C}$ such that $\pi(\mu * f) = \mu(\pi)\pi(f)$ for all $f \in \mathcal{S}^\circ(G(\mathbb{A}))$, resp. $f \in \mathcal{S}^\circ(U_h(\mathbb{A}))$.

Let ξ_Π be the infinitesimal character of Π . By [BLZZ19, Theorem 4.12 (4)], for every $h \in \mathcal{H}^\circ$ and $(U_h, \sigma) \in \mathfrak{X}_0^h$, the base-change of the infinitesimal character of σ is ξ_Π . However, the universal enveloping algebras of the complexified Lie algebras of U_h are all canonically identified for $h \in \mathcal{H}$ (since these are inner forms of each other) and base-change is injective at the level of infinitesimal characters. As, by [GRS11], there exists at least one $h \in \mathcal{H}^\circ$ such that the set \mathfrak{X}_0^h is nonempty (we may even take for h any quasi-split Hermitian form unramified outside S_0), there exists a common infinitesimal character ξ of all $(U_h, \sigma) \in \mathfrak{X}_0^h$, for $h \in \mathcal{H}^\circ$, whose base-change is ξ_Π .

By the strong multiplicity one theorem of Ramakrishnan (see [Ram18]) and Theorem 3.17 of [BLZZ19], one can find a multiplier $\mu \in \mathcal{M}^{S'_0}(G(\mathbb{A}))$ such that

- i. $\mu(\Pi) = 1$;
- ii. For all $\chi' \in \mathfrak{X}(G)$ such that $\chi' \neq \chi$, we have

$$K_{\mu*f, \chi'}^0 = 0$$

where the kernel $K_{\mu*f, \chi'}^0$ is defined as in §2.9.8.

By Theorem 3.6 and Theorem 4.12 (3) of [BLZZ19], for every $h \in \mathcal{H}^\circ$ there exists a multiplier $\mu^h \in \mathcal{M}^{S'_0}(U_h(\mathbb{A}))$ such that

- iii. $\mu^h(\sigma) = 1$ for all $(U_h, \sigma) \in \mathfrak{X}_0^h$;
- iv. For all $\chi' \in \mathfrak{X}(U_h)$ such that $\chi' \notin \mathfrak{X}_0^h$ and for all parabolic subgroups P of U_h , we have

$$K_{P, \mu^h*f^h, \chi'}^{U_h} = 0$$

where the left-hand side is the kernel of the operator given by the right convolution of $\mu^h * f^h$ on $L_\chi^2([U_h]_P)$ (see (2.9.2.1)).

Moreover, by [BLZZ19, Proposition 4.8, Lemma 4.10], we may choose μ and μ^h such that the functions $\mu * f$ and $\mu^h * f^h$, for $h \in \mathcal{H}^\circ$, still satisfy the assumptions of the proposition. So, in particular, from (10.1.6.4) applied to the functions $\mu * f$ and $(\mu^h * f^h)_{h \in \mathcal{H}^\circ}$ instead of f and $(f^h)_{h \in \mathcal{H}^\circ}$, we get

$$(10.1.6.5) \quad I(\mu * f) = \sum_{h \in \mathcal{H}^\circ} J^h(\mu^h * f^h).$$

Note that by conditions i. and iii. we have:

$$I_\Pi(\mu * f) = I_\Pi(f), \quad I_{P, \pi}(\mu * f) = I_{P, \pi}(f) \quad \text{and} \quad J_{\mathfrak{X}_0^h}^h(\mu^h * f^h) = J_{\mathfrak{X}_0^h}^h(f^h), \quad \text{for every } h \in \mathcal{H}^\circ.$$

Moreover, by ii. Theorem 3.3.7.1 applied to (3.3.3.2) and (10.1.5.2), we see that the left-hand side of (10.1.6.5) reduces to $I_\chi(\mu * f) = 2^{-\dim(\mathfrak{a}_P)} I_\Pi(\mu * f) = 2^{-\dim(\mathfrak{a}_P)} I_{P, \pi}(\mu * f)$. On the other hand, by iv. and the very definition of J^h given in [Zyd20], the right-hand side of (10.1.6.5) reduces to

$$\sum_{h \in \mathcal{H}^\circ} J_{\mathfrak{X}_0^h}^h(\mu^h * f^h) = \sum_{h \in \mathcal{H}^\circ} J_{\mathfrak{X}_0^h}^h(\mu^h * f^h).$$

Therefore, (10.1.6.5) gives precisely the identity of the proposition. \square

10.2 Proof of Theorem 1.1.5.1

10.2.1. Let $\Pi = \text{Ind}_P^G(\pi)$ be a Hermitian Arthur parameter of G . Note that by properties 1 and 2 of §1.1.3, the cuspidal datum χ associated to the pair (M_P, π) is generic and relevant in the sense of §4.3.2. For $h \in \mathcal{H}$ and σ a cuspidal automorphic representation of $U_h(\mathbb{A})$, it is readily seen that the linear form \mathcal{P}_h is nonzero on σ if and only if J_σ^h is not identically zero. On the other hand, the linear form J_η or β_η is always nonzero (this follows either from the fact that χ is relevant or is an easy consequence of [GK72], [Jac10, Proposition 5] and [Kem15]) whereas the linear form I , from Proposition 5.1.4.1, or λ , from §8.1.2, is nonzero if and only if $L(\frac{1}{2}, \Pi) \neq 0$ (as follows either from the work of Ichino and Yamana, see [IY15, corollary 5.7], or of Jacquet, Piatetski-Shapiro and Shalika [JPSS83], [Jac04]). Therefore, we similarly deduce that the distribution $I_{P,\pi}$ or I_Π is non-zero if and only if $L(\frac{1}{2}, \Pi) \neq 0$.

As a consequence, Theorem 1.1.5.1 amounts to the equivalence between the two assertions:

- (A) The distribution $I_{P,\pi}$ or I_Π is non-zero.
- (B) There exist $h \in \mathcal{H}$, $f \in \mathcal{S}(U_h(\mathbb{A}))$ and a cuspidal subrepresentation σ of U_h such that $BC(\sigma) = \Pi$ and $J_\sigma^h(f) \neq 0$.

10.2.2. Proof of (A) \Rightarrow (B). — We choose the S_0 of §10.1.3 such that I_Π is not identically zero on $f_1 \in \mathcal{S}^\circ(G(\mathbb{A}))$. Then Assertion (B) above is a consequence of Proposition 10.1.6.1: it suffices to take functions f and f^h for $h \in \mathcal{H}^\circ$ satisfying the hypotheses of that theorem and such that $I_\Pi(f) \neq 0$. That it is possible is implied by a combination of a result of [Xue19] and the existence of p -adic transfer [Zha14b].

10.2.3. Proof of (B) \Rightarrow (A). — We may choose the set S_0 so that there exist $h_0 \in \mathcal{H}^\circ$, $f_0^{h_0} \in \mathcal{S}^\circ(U_{h_0}(\mathbb{A}))$ and a cuspidal representation σ_0 of U_{h_0} such that for $v \in S'_0$ (see §10.1.6) $BC(\sigma_{0,v}) = \Pi_v$ and $J_{\sigma_0}^{h_0}(f_0^{h_0}) \neq 0$. For any other $h \in \mathcal{H}^\circ$ we set $f_0^h = 0$. Up to enlarging S_0 , we may assume that the family $(f_0^h)_{h \in \mathcal{H}^\circ}$ satisfies conditions 2. and 5. of Proposition 10.1.6.1. Moreover, we have (see [Zha14b, §2.5]) $J_{\sigma_0}^{h_0}(f_0^{h_0} * f_0^h) \geq 0$ for every $\sigma \in \mathfrak{X}_0^{h_0}$ and $J_{\sigma_0}^{h_0}(f_0^{h_0} * f_0^h) > 0$. In particular, the left hand side of (10.1.6.3) for the family $(f_0^h * f_0^h)_{h \in \mathcal{H}^\circ}$ is nonzero. Once again by [Xue19] and the existence of p -adic transfer [Zha14b], this implies that we can find test functions $f \in \mathcal{S}^\circ(G(\mathbb{A}))$ and $f^h \in \mathcal{S}^\circ(U_h(\mathbb{A}))$, for $h \in \mathcal{H}^\circ$, satisfying all the conditions of Proposition 10.1.6.1 and such that the left hand side of (10.1.6.3) is still nonzero. The conclusion of this proposition immediately gives Assertion (A).

10.3 Proof of Theorem 1.1.6.1

10.3.1. Let $h \in \mathcal{H}$ and σ be a cuspidal automorphic representation of $U_h(\mathbb{A})$ which is tempered everywhere. By [Mok15], [KMSW], σ admits a weak base-change Π to G . Moreover, by these references Π is also a strong base-change of σ : for every place v of F , the local base-change of σ_v (defined in [Mok15] and [KMSW]) coincides with Π_v . In particular, it follows that Π is also tempered everywhere.

We choose a finite set of places S_0 as in §10.1.3 such that $h \in \mathcal{H}^\circ$ and σ as well as the additive character ψ' used to normalize local Haar measures in Section 2.3 are unramified outside of S_0 .

For each place v of F , we define a distribution J_{σ_v} on $\mathcal{S}(U_h(F_v))$ by

$$J_{\sigma_v}(f_v^h) = \int_{U'_h(F_v)} \text{Trace}(\sigma_v(h_v)\sigma_v(f_v^h))dh_v, \quad f_v^h \in \mathcal{S}(U_h(F_v)),$$

where

$$\sigma_v(f_v^h) = \int_{U_h(F_v)} f_v^h(g_v)\sigma_v(g_v)dg_v$$

and the Haar measures are the one defined in §2.3.3. Moreover by [Har14], and since the representations σ_v are all tempered, the expression defining J_{σ_v} is absolutely convergent and for every

$v \notin S_0$ we have

$$J_{\sigma_v}(\mathbf{1}_{U_h(\mathcal{O}_v)}) = \Delta_{U'_h, v}^{-2} \frac{L(\frac{1}{2}, \Pi_v)}{L(1, \sigma_v, \text{Ad})}.$$

10.3.2. By [Zha14a, Lemma 1.7] and our choice of local Haar measures, Theorem 1.1.6.1 is equivalent to the following assertion: for all factorizable test function $f^h \in \mathcal{S}(U_{h_0}(\mathbb{A}))$ of the form $f^h = (\Delta_{U'_h}^{S_0})^2 \prod_{v \in S_0} f_v^h \times \prod_{v \notin S_0} \mathbf{1}_{U_h(\mathcal{O}_v)}$, we have

$$(10.3.2.1) \quad J_{\sigma}^h(f^h) = |S_{\Pi}|^{-1} \frac{L^{S_0}(\frac{1}{2}, \Pi)}{L^{S_0}(1, \sigma, \text{Ad})} \prod_{v \in S_0} J_{\sigma_v}(f_v^h).$$

10.3.3. For every place v of F , we define a local relative character I_{Π_v} on $G(F_v)$ by

$$I_{\Pi_v}(f_v) = \sum_{W_v \in \mathcal{W}(\Pi_v, \psi_{N, v})} \frac{\lambda_v(\Pi_v(f_v)W_v) \overline{\beta_{\eta, v}(W_v)}}{\langle W_v, W_v \rangle_{\text{Whitt}, v}}, \quad f_v \in \mathcal{S}(G(F_v)),$$

where the sum runs over a K_v -basis of the Whittaker model $\mathcal{W}(\Pi_v, \psi_{N, v})$ (in the sense of §2.8.3) and $\lambda_v, \beta_{\eta, v}, \langle \cdot, \cdot \rangle_{\text{Whitt}, v}$ are local analogs of the forms introduced in §8.1.2 given by

$$\begin{aligned} \lambda_v(W_v) &= \int_{N_H(F_v) \backslash H(F_v)} W_v(h_v) dh_v, \quad \beta_{\eta, v}(W_v) = \int_{N'(F_v) \backslash \mathcal{P}'(F_v)} W_v(p_v) \eta_{G', v}(p_v) dp_v, \\ \text{and } \langle W_v, W_v \rangle_{\text{Whitt}, v} &= \int_{N(F_v) \backslash \mathcal{P}(F_v)} |W_v(p_v)|^2 dp_v. \end{aligned}$$

Note that the above expressions, and in particular $\lambda_v(W_v)$, are all absolutely convergent due to the fact that Π_v is tempered (see [JPSS83, Proposition 8.4]). The above definition also implicitly depends on the choice of an additive character ψ of \mathbb{A}_E/E trivial on \mathbb{A} (through which the generic character ψ_N is defined, see beginning of Chapter 8 and §6.1.2) and up to enlarging S_0 , we may assume that ψ is unramified outside of S_0 . Then, it follows from the definition of I_{Π} that for every factorizable test function $f \in \mathcal{S}(G(\mathbb{A}))$ of the form $f = \Delta_H^{S_0, *} \Delta_{G'}^{S_0, *} \prod_{v \in S_0} f_v \times \prod_{v \notin S_0} \mathbf{1}_{G(\mathcal{O}_v)}$, we have

$$(10.3.3.2) \quad I_{\Pi}(f) = \frac{L^{S_0}(\frac{1}{2}, \Pi)}{L^{S_0}(1, \Pi, \text{As}_{G'})} \prod_{v \in S_0} I_{\Pi_v}(f_v).$$

10.3.4. Let f^h be a test function as in §10.3.2. Then, as both sides of (10.3.2.1) are continuous functionals in f_v^h for $v \in V_{F, \infty}$, by the main result of [Xue19] we may assume that for every $v \in V_{F, \infty}$ the function f_v^h admits a transfer $f_v \in \mathcal{S}(G(F_v))$. On the other hand, by [Zha14b], for every $v \in S_0 \setminus V_{F, \infty}$, the function f_v^h admits a transfer $f_v \in \mathcal{S}(G(F_v))$. Moreover, by the results of those references we may also choose the transfers such that for every $h' \in \mathcal{H}^{\circ}$ with $h' \neq h$, the zero function on $U_{h'}(F_{S_0})$ is a transfer of $f_{S_0} = \prod_{v \in S_0} f_v$. We set $f = \Delta_H^{S_0, *} \Delta_{G'}^{S_0, *} f_{S_0} \times \prod_{v \notin S_0} \mathbf{1}_{G(\mathcal{O}_v)}$. Then, setting $f^{h'} = 0$ for every $h' \in \mathcal{H}^{\circ} \setminus \{h\}$, the functions f and $(f^{h'})_{h' \in \mathcal{H}^{\circ}}$ satisfy the assumptions of Proposition 10.1.6.1. Therefore, we have

$$(10.3.4.3) \quad \sum_{\sigma \in \mathfrak{X}_0^h} J_{\sigma}^h(f^h) = 2^{-\dim(\mathfrak{a}_F)} I_{\Pi}(f).$$

10.3.5. If there exists a place $v \in S_0$ such that σ_v does not support any nonzero continuous $U'_h(F_v)$ -invariant functional, both sides of (10.3.2.1) are automatically zero.

Assume now that for every $v \in S_0$, the local representation σ_v supports a nonzero continuous $U'_h(F_v)$ -invariant functional. By the local Gan-Gross-Prasad conjecture [BP15], and the classification of cuspidal automorphic representations of U_h in terms of local L -packets [Mok15], [KMSW], it follows that all the terms except possibly $J_\sigma^h(f^h)$ in the left hand side of (10.3.4.3) are zero. Moreover, by [BP18b, Theorem 5.4.1] and since Π_v is the local base-change of σ_v , there are explicit constants $\kappa_v \in \mathbb{C}^\times$ for $v \in S_0$ satisfying $\prod_{v \in S_0} \kappa_v = 1$ and such that

$$(10.3.5.4) \quad I_{\Pi_v}(f_v) = \kappa_v J_{\sigma_v}(f_v^h)$$

for every $v \in S_0$. Combining this with (10.3.3.2), we get

$$\begin{aligned} J_\sigma^h(f^h) &= 2^{-\dim(\mathfrak{a}_P)} I_\Pi(f) = 2^{-\dim(\mathfrak{a}_P)} I_{P,\pi}(f) = 2^{-\dim(\mathfrak{a}_P)} \frac{L^{S_0}(\frac{1}{2}, \Pi)}{L^{S_0}(1, \Pi, \text{As}_{G'})} \prod_{v \in S_0} I_{\Pi_v}(f_v) \\ &= 2^{-\dim(\mathfrak{a}_P)} \frac{L^{S_0}(\frac{1}{2}, \Pi)}{L^{S_0}(1, \Pi, \text{As}_{G'})} \prod_{v \in S_0} J_{\sigma_v}(f_v^h). \end{aligned}$$

As $L^{S_0}(s, \Pi, \text{As}_{G'}) = L^{S_0}(s, \sigma, \text{Ad})$ and $|S_\Pi| = 2^{-\dim(\mathfrak{a}_P)}$, this exactly gives (10.3.2.1) and ends the proof of Theorem 1.1.6.1.

A Topological vector spaces

A.0.1. In this paper, by a *topological vector space* (TVS) we mean a complex locally convex separated vector spaces. Actually, most TVS encountered in this paper will be Fréchet or LF (that is a countable inductive limit of Fréchet spaces) or even strict LF (that is countable inductive limit $\varinjlim_n F_n$ of Fréchet spaces with closed embeddings $F_n \rightarrow F_{n+1}$ as connecting maps) spaces. Let E and F be TVS. We denote by E' the topological dual of E and by $\text{Hom}(F, E)$ the space of continuous linear mappings $F \rightarrow E$ both being equipped with their weak topologies (that is the topologies of pointwise convergence). Recall that a *total subspace* $H \subset E'$ is a subspace such that $\bigcap_{\lambda \in H} \text{Ker}(\lambda) = 0$. A *bounded* subset $B \subseteq E$ is one that is absorbed by any neighborhood of 0. If $B \subseteq E$ is bounded and absolutely convex, we define E_B to be the subspace generated by B equipped with the norm $\|e\|_B = \inf \{\lambda \geq 0 \mid e \in \lambda B\}$. Then, the natural inclusion $E_B \rightarrow E$ is continuous. The space E is said to be *quasi-complete* if every closed bounded subset of it is complete. Most TVS encountered in this paper will be quasi-complete (e.g. Fréchet or strict LF spaces).

A.0.2. We recall the notion of integral valued in a TVS in the form we use it in the core of the paper. Let (X, μ) be a measured space and $f : X \rightarrow E$ be a measurable function. When E is quasi-complete, we say that f is *absolutely integrable* if for every continuous semi-norm p on E the integral $\int_X p \circ f \mu$ converges. If this is the case, there exists a unique element

$$\int_X f \mu \in E$$

such that $\langle \lambda, \int_X f \mu \rangle = \int_X \langle \lambda, f \rangle \mu$ for every $\lambda \in E'$. This notion applies in particular to series $\sum_n f_n$ valued in a quasi-complete TVS E : the series is said to be *absolutely convergent* in E if for every continuous semi-norm p on E , the series $\sum_n p(f_n)$ converges, in which case $\sum_n f_n$ has a limit in E .

A.0.3. We will also freely use the notions of smooth or holomorphic functions valued in a TVS. For basic references on these subjects, we refer the reader to [Bou67, §2, §3], [Gro53, §2], [Gro73, Chap. 3, §8]. There are actually two ways to define smooth and holomorphic maps valued in E : either scalarly (that is after composition with any element of E') or by directly requiring the functions to be infinitely (complex) differentiable. These two definitions coincide when the space E is quasi-complete and, fortunately for us, we will only consider smooth/holomorphic functions valued in such spaces so that we don't have to distinguish.

Let M be a connected complex analytic manifold. A function $f : M \rightarrow E$ is holomorphic if and only if for every relatively compact open subset $\Omega \subseteq M$, there exists a bounded absolutely convex subset $B \subseteq E$ such that $f|_\Omega$ factorizes through a holomorphic map $\Omega \rightarrow E_B$ see [Gro53, §2, Remarque 2]. We also record the following convenient criterion of holomorphicity [Bou67, §3.3.1]:

(A.0.3.1) Assume that E is quasi-complete. A function $\varphi : M \rightarrow E$ is holomorphic if and only if it is continuous and for some total subspace $H \subseteq E'$, the functions $s \in M \mapsto \langle \varphi(s), \lambda \rangle$ are holomorphic for every $\lambda \in H$.

A.0.4. Assume that F is a LF space. As LF spaces are barreled [Trê67, Corollary 33.3] they satisfy the Banach-Steinhaus theorem [Trê67, Theorem 33.1] hence any bounded subset of $\text{Hom}(F, E)$ is equicontinuous (since $\text{Hom}(F, E)$ is equipped with the weak topology, that a subset $B \subseteq \text{Hom}(F, E)$ is bounded means that for every $f \in F$ the subset $\{T(f) \mid T \in B\}$ of E is itself bounded). This shows that for any bounded subset $B \subseteq \text{Hom}(F, E)$ the restriction of the canonical map $\text{Hom}(F, E) \times F \rightarrow E$ to $B \times F$ is continuous. Also, if E is quasi-complete then $\text{Hom}(F, E)$ is too [Trê67, §34.3 Corollary 2]. In particular, we get:

(A.0.4.2) Assume that F is LF and E is quasi-complete. Let $s \in M \mapsto T_s \in \text{Hom}(F, E)$ be holomorphic and $(s, k) \in M \times K \mapsto f_{s,k} \in F$ be a continuous map which is holomorphic in the first variable. Then, the map $(s, k) \in M \times K \mapsto T_s(f_{s,k}) \in E$ is continuous and holomorphic in the first variable.

Indeed, T has locally its image in a bounded set. Hence, by the above discussion, the map $(s, s', k) \in M \times M \times K \mapsto T_s(f_{s',k}) \in E$ is continuous. Moreover, this map is separately holomorphic in the variables s, s' . Thus, by Hartog's theorem, this map is holomorphic in the variables (s, s') which immediately implies the claim by “restriction to the diagonal”.

(A.0.4.3) Assume that F is LF and E is quasi-complete. Let $U \subseteq M$ be a nonempty open subset and $s \in U \mapsto T_s \in \text{Hom}(F, E)$ be a holomorphic map. If, for every $f \in F$ the map $s \mapsto T_s(f) \in E$ extends analytically to M then $T_s \in \text{Hom}(F, E)$ for every $s \in M$ and moreover $s \in M \mapsto T_s \in \text{Hom}(F, E)$ is holomorphic.

Indeed, $s \mapsto T_s$ induces a holomorphic map $M \rightarrow \text{Hom}(F, E)$ where $\text{Hom}(F, E)$ stands for the space of *all* linear maps $F \rightarrow E$ (not necessarily continuous) equipped with the topology of pointwise convergence. Hence, for every relatively compact connected open subset $\Omega \subseteq M$ such that $\Omega \cap U \neq \emptyset$ there exists a bounded subset $B \subseteq \text{Hom}(F, E)$ such that $s \mapsto T_s$ factorizes through a holomorphic map $\Omega \rightarrow \text{Hom}(F, E)_B$. By the Banach-Steinhaus theorem, $\text{Hom}(F, E) \cap \text{Hom}(F, E)_B$ is closed in $\text{Hom}(F, E)_B$ which immediately implies (by Hahn-Banach and the fact that Ω is connected) that $s \in \Omega \mapsto T_s$ factorizes through a holomorphic map $\Omega \rightarrow \text{Hom}(F, E) \cap \text{Hom}(F, E)_B$. The claim follows.

A.0.5. Let $\text{Bil}_s(E, F) = \text{Hom}(E, \text{Hom}(F, \mathbb{C}))$ be the space of separately continuous bilinear mappings $E \times F \rightarrow \mathbb{C}$ equipped with the topology of pointwise convergence. Applying (A.0.4.2) and (A.0.4.3) twice, we get:

(A.0.5.4) Assume that E and F are LF. Let $s \in M \mapsto B_s \in \text{Bil}_s(E, F)$ be holomorphic and $(s, k) \in M \times K \mapsto e_{s,k} \in E, (s, k) \in M \times K \mapsto f_{s,k} \in F$ be continuous maps which are holomorphic in the first variable. Then, the function $(s, k) \in M \times K \mapsto B_s(e_{s,k}, f_{s,k})$ is continuous and holomorphic in the first variable.

(A.0.5.5) Assume that both E and F are LF. Let $U \subseteq M$ be a nonempty open subset and $s \in U \mapsto B_s \in \text{Bil}_s(E, F)$ be a holomorphic map. If for every $(e, f) \in E \times F$ the function $s \mapsto B_s(e, f)$ extends analytically to M then $B_s \in \text{Bil}_s(E, F)$ for every $s \in M$ and moreover $s \in M \mapsto B_s \in \text{Bil}_s(E, F)$ is holomorphic.

A.0.6. We denote by $E \widehat{\otimes} F$ the completed projective tensor product [Trè67, Chap. 43]. It admits a canonical linear map $E \otimes F \rightarrow E \widehat{\otimes} F$ satisfying the following universal property: for every complete TVS G , precomposition yields an isomorphism

$$\text{Hom}(E \widehat{\otimes} F, G) \simeq \text{Bil}(E, F; G)$$

where $\text{Bil}(E, F; G)$ denotes the space of all *continuous* bilinear mappings $E \times F \rightarrow G$. In particular, if G and H are two other TVS and $T : E \rightarrow G, S : F \rightarrow H$ are continuous linear mapping, there is a unique continuous linear map $T \widehat{\otimes} S : E \widehat{\otimes} F \rightarrow G \widehat{\otimes} H$ which on $E \otimes F$ is given by $e \otimes f \mapsto T(e) \otimes S(f)$. Moreover, the topology induced from $E \widehat{\otimes} F$ on $E \otimes F$ is also associated to the family of semi-norms

$$(p \otimes q)(v) = \inf \left\{ \sum_i p(e_i) q(f_i) \mid v = \sum_i e_i \otimes f_i \right\}$$

where p (resp. q) runs over a family of semi-norms defining the topology on E (resp. F).

Assume now that E and F are spaces of (complex valued) functions on two sets X, Y and that their topologies are finer than the topology of pointwise convergence. When E is moreover a complete nuclear LF space, the following result of Grothendieck [Gro55, Théorème 13, Chap. II, §3 n.3] generally allows to describe $E \widehat{\otimes} F$ explicitly as a space of functions on $X \times Y$.

(A.0.6.6) Let $\mathcal{F}(X \times Y)$ be the space of all complex valued functions on $X \times Y$ equipped with the topology of pointwise convergence. Then the linear map $E \otimes F \rightarrow \mathcal{F}(X \times Y)$, $e \otimes f \mapsto ((x, y) \mapsto e(x)f(y))$, extends continuously to a linear embedding $E \widehat{\otimes} F \hookrightarrow \mathcal{F}(X \times Y)$ with image the space of functions $f : X \times Y \rightarrow \mathbb{C}$ satisfying the two conditions:

- For every $x \in X$, the function $y \in Y \mapsto f(x, y)$ belongs to the completion of F ;
- For every $\lambda \in F'$, the function $x \in X \mapsto \langle f(x, \cdot), \lambda \rangle$ belongs to E .

A.0.7. Let $C \in \mathbb{R} \cup \{-\infty\}$ and $f : \mathcal{H}_{>C} \rightarrow E$ be a holomorphic function. We say that f is of *order at most d in vertical strips* if for every $d' > d$ the function $z \mapsto e^{-|z|^{d'}} f(z)$ is *bounded in vertical strips* of $\mathcal{H}_{>C}$. We say that f is of *finite order in vertical strips* if it is of order at most d in vertical strips for some $d > 0$. Finally, we say that f is *rapidly decreasing in vertical strips* if for every $d > 0$ the function $z \mapsto |z|^d f(z)$ is bounded in vertical strips.

A.0.8. Let \mathcal{A} be a real vector space. Denote by $\text{Diff}(\mathcal{A})$ the space of complex polynomial differential operators on \mathcal{A} (which can be identified with $\text{Sym}(\mathcal{A}_{\mathbb{C}}^*) \otimes_{\mathbb{C}} \text{Sym}(\mathcal{A}_{\mathbb{C}})$). When E is quasi-complete, we define the space of *Schwartz functions on \mathcal{A} valued in E* , denoted by $\mathcal{S}(\mathcal{A}, E)$, as the space of smooth functions $f : \mathcal{A} \rightarrow E$ such that for every $D \in \text{Diff}(\mathcal{A})$, the function Df has bounded image. Note that if F is also quasi-complete and $T : E \rightarrow F$ is a continuous linear map then for every $f \in \mathcal{S}(\mathcal{A}, E)$, we have $T \circ f \in \mathcal{S}(\mathcal{A}, F)$. When $E = \mathbb{C}$, we simply set $\mathcal{S}(\mathcal{A}) = \mathcal{S}(\mathcal{A}, \mathbb{C})$ that we equip with its standard Fréchet topology.

Lemma A.0.8.1. — Assume that E is a strict LF space. Let $C > 0$, $d > 0$ and $s \in \mathcal{H}_{>C} \mapsto Z_s \in E'$ be a map such that for every $f \in E$, $s \in \mathcal{H}_{>C} \mapsto Z_s(f)$ is a holomorphic function of order at most d in vertical strips. Then, for every $f \in \mathcal{S}(\mathcal{A}, E)$, the map

$$(A.0.8.7) \quad s \in \mathcal{H}_{>C} \mapsto (\lambda \in \mathcal{A} \mapsto Z_s(f_\lambda)) \in \mathcal{S}(\mathcal{A})$$

is holomorphic and of finite order in vertical strips.

Proof. — Indeed, by the Banach-Steinhaus theorem, for every $d' > d$, every vertical strip $V \subseteq \mathcal{H}_{>C}$ and every bounded subset $B \subseteq E$ the set

$$\left\{ e^{-|s|^{d'}} Z_s(f) \mid s \in V, f \in B \right\} \subseteq \mathbb{C}$$

is bounded and, by [Trè67, Corollary 33.1], for every $s_0 \in \mathcal{H}_{>C}$, Z_s converges uniformly on compact subsets to Z_{s_0} as $s \rightarrow s_0$. Let $f \in \mathcal{S}(\mathcal{A}, E)$. Moreover, for every $D \in \text{Diff}(\mathcal{A})$ the set

$$\{Df_\lambda \mid \lambda \in \mathcal{A}\} \cup \{0\} \subseteq E$$

is compact. Therefore, for every $s_0 \in \mathcal{H}_{>C}$, $Z_s(Df_\lambda)$ converges to $Z_{s_0}(Df_\lambda)$ as $s \rightarrow s_0$ uniformly in $\lambda \in \mathcal{A}$ and $\left\{ e^{-|s|^{d'}} Z_s(Df_\lambda) \mid s \in V, \lambda \in \mathcal{A} \right\}$ is bounded for every $d' > d$ and every vertical strip $V \subseteq \mathcal{H}_{>C}$. This shows that the map (A.0.8.7) is continuous and of finite order in vertical strips. To conclude we apply the holomorphicity criterion (A.0.3.1) to $H \subseteq \mathcal{S}(\mathcal{A})'$ the subset of “evaluations at a point of \mathcal{A} ”. \square

A.0.9. Lemma A.0.9.1. — Assume that E is quasi-complete. Let $Z_+, Z_- : \mathcal{H}_{>C} \rightarrow E$ be holomorphic functions of finite order in vertical strips for some $C > 0$. Assume that there exists a total subspace $H \subset E'$ such that for every $\lambda \in H$, $Z_{+, \lambda} := \lambda \circ Z_+$ and $Z_{-, \lambda} := \lambda \circ Z_-$ extend to holomorphic functions on \mathbb{C} of finite order in vertical strips satisfying $Z_{+, \lambda}(z) = Z_{-, \lambda}(-z)$ for every $z \in \mathbb{C}$. Then, Z_+ and Z_- extend to holomorphic functions $\mathbb{C} \rightarrow E$ of finite order in vertical strips satisfying $Z_+(z) = Z_-(-z)$ for every $z \in \mathbb{C}$.

Proof. — Let $d > 0$ be such that Z_+ and Z_- are of order at most d in vertical strips of $\mathcal{H}_{>C}$. Then, by the Phragmen-Lindelöf principle, for every $\lambda \in H$, the holomorphic continuations of

$Z_{+,\lambda}$ and $Z_{-,\lambda}$ are also of order at most d in vertical strips. Therefore, up to multiplying Z_+ and Z_- by $z \mapsto e^{z^{4n+2}}$ for some $n \geq 0$, we may assume that all these functions are rapidly decreasing in vertical strips. Let $D > C$. Then, for every $z \in \mathcal{H}_{]-D,D[}$ and $\epsilon \in \{\pm\}$, we set

$$\Phi_\epsilon(z) = \frac{1}{2\pi} \left(\int_{-\infty}^{+\infty} \frac{Z_\epsilon(D+it)}{D+it-z} dt - \int_{-\infty}^{+\infty} \frac{Z_{-\epsilon}(D+it)}{D+it+z} dt \right).$$

Note that, since Z_+ and Z_- are rapidly decreasing in vertical strips and E is quasi-complete, the above integrals converge absolutely in E . By the usual holomorphicity criterion for parameter integrals, we readily check that the functions Φ_+ , Φ_- are holomorphic. Moreover, by the uniform boundedness principle, Φ_+ and Φ_- are bounded in vertical strips. Finally, by Cauchy's integration formula and the fact that the functions $Z_{+,\lambda}$, $Z_{-,\lambda}$ are rapidly decreasing in vertical strips, for every $\epsilon \in \{\pm\}$ and $\lambda \in H$ the functions $\lambda \circ \Phi_\epsilon$ and $Z_{\epsilon,\lambda}$ coincide on $\mathcal{H}_{]-D,D[}$. Therefore, as H is total, Φ_ϵ and Z_ϵ coincide on $\mathcal{H}_{]C,D[}$. This shows that Z_+ and Z_- admit holomorphic extensions bounded in vertical strips to $\mathcal{H}_{>-D}$ for every $D > C$ hence to \mathbb{C} . That the functional equation $Z_+(z) = Z_-(-z)$ holds for these extensions easily follows from the assumption. \square

A.0.10. Let \mathcal{A} be a real vector space. Specializing the previous lemma to $E = \mathcal{S}(\mathcal{A})$ and H the total subspace of E' given by “evaluations at a point of \mathcal{A} ” yields the following corollary.

Corollary A.0.10.1. — *Let $Z_+, Z_- : \mathcal{A} \times \mathbb{C} \rightarrow \mathbb{C}$ be two functions such that:*

1. *There exists $C > 0$ such that for every $s \in \mathcal{H}_{>C}$, the function $Z_+(\cdot, s)$, $Z_-(\cdot, s)$ belong to $\mathcal{S}(\mathcal{A})$ and the maps*

$$s \in \mathcal{H}_{>C} \mapsto Z_\epsilon(\cdot, s) \in \mathcal{S}(\mathcal{A}), \quad \epsilon \in \{\pm\},$$

are holomorphic functions of finite order in vertical strips;

2. *For every $\lambda \in \mathcal{A}$, $s \in \mathbb{C} \mapsto Z_+(\lambda, s)$ and $s \in \mathbb{C} \mapsto Z_-(\lambda, s)$ are holomorphic functions of finite order in vertical strips satisfying the functional equation*

$$Z_+(\lambda, s) = Z_-(\lambda, -s)$$

Then, for every $s \in \mathbb{C}$ the functions $Z_+(\cdot, s)$, $Z_-(\cdot, s)$ belong to $\mathcal{S}(\mathcal{A})$ the maps $s \in \mathbb{C} \mapsto Z_\epsilon(\cdot, s) \in \mathcal{S}(\mathcal{A})$, $\epsilon \in \{\pm\}$, are holomorphic.

Assume now that F is a LF space. As F is barreled, F' is quasi-complete [Trè67, §34.3 Corollary 2]. Specializing Lemma A.0.9.1 to $E = F'$ and H a dense subset of $E' = F$, we obtain the following.

Corollary A.0.10.2. — *Let F be a LF space, $C > 0$ and $Z_+, Z_- : \mathcal{H}_{>C} \times F \rightarrow \mathbb{C}$ be two functions. Assume that:*

1. *For every $s \in \mathcal{H}_{>C}$, $Z_+(s, \cdot)$ and $Z_-(s, \cdot)$ are continuous functionals on F ;*
2. *There exists $d > 0$ such that for every $f \in F$ and $\epsilon \in \{\pm\}$, $s \in \mathcal{H}_{>C} \mapsto Z_\epsilon(s, f)$ is a holomorphic function of order at most d in vertical strips;*
3. *For every $f \in H$ and $\epsilon \in \{\pm\}$, $s \mapsto Z_\epsilon(s, f)$ extends to a holomorphic function on \mathbb{C} of finite order in vertical strips satisfying*

$$Z_+(s, f) = Z_-(-s, f).$$

Then, Z_+ and Z_- extend to holomorphic functions $\mathbb{C} \rightarrow F'$ of finite order in vertical strips satisfying $Z_+(s, f) = Z_-(-s, f)$ for every $s \in \mathbb{C}$ and every $f \in F$.

References

- [AG09] A. Aizenbud and D. Gourevitch. Generalized Harish-Chandra descent, Gelfand pairs, and an Archimedean analog of Jacquet-Rallis’s theorem. *Duke Math. J.*, 149(3):509–567, 2009. With an appendix by the authors and Eitan Sayag.
- [Art78] J. Arthur. A trace formula for reductive groups I. Terms associated to classes in $G(\mathbb{Q})$. *Duke Math. J.*, 45:911–952, 1978.
- [Art80] J. Arthur. A trace formula for reductive groups II. *Comp. Math.*, 40:87–121, 1980.
- [Art81] J. Arthur. The trace formula in invariant form. *Ann. of Math. (2)*, 114(1):1–74, 1981.
- [Art82] J. Arthur. On the inner product of truncated Eisenstein series. *Duke Math. J.*, 49(1):35–70, 1982.
- [Art85] J. Arthur. A measure on the unipotent variety. *Canad. J. Math.*, 37(6):1237–1274, 1985.
- [Bar03] E. Baruch. A proof of Kirillov’s conjecture. *Ann. of Math. (2)*, 158(1):207–252, 2003.
- [Ber84] J. Bernstein. P -invariant distributions on $GL(N)$ and the classification of unitary representations of $GL(N)$ (non-Archimedean case). In *Lie group representations, II (College Park, Md., 1982/1983)*, volume 1041 of *Lecture Notes in Math.*, pages 50–102. Springer, Berlin, 1984.
- [Ber88] J. Bernstein. On the support of Plancherel measure. *J. Geom. Phys.*, 5(4):663–710 (1989), 1988.
- [Beu16] R. Beuzart-Plessis. Comparison of local spherical characters and the Ichino-Ikeda conjecture for unitary groups. *ArXiv e-prints*, February 2016.
- [BK14] J. Bernstein and B. Krötz. Smooth Fréchet globalizations of Harish-Chandra modules. *Israel J. Math.*, 199(1):45–111, 2014.
- [BL19] J. Bernstein and E. Lapid. On the meromorphic continuation of Eisenstein series. *arXiv:1911.02342*, 2019. prepublication.
- [BLZZ19] R. Beuzart-Plessis, Y. Liu, W. Zhang, and X. Zhu. Isolation of cuspidal spectrum, with application to the Gan–Gross–Prasad conjecture. *arXiv e-prints*, page arXiv:1912.07169, December 2019.
- [Bou67] N. Bourbaki. *Éléments de mathématique. Fasc. XXXIII. Variétés différentielles et analytiques. Fascicule de résultats (Paragraphes 1 à 7)*. Actualités Scientifiques et Industrielles, No. 1333. Hermann, Paris, 1967.
- [BP] R. Beuzart-Plessis. A new proof of Jacquet-Rallis’s fundamental lemma. *arXiv:1901.02653*.
- [BP15] R. Beuzart-Plessis. A local trace formula for the Gan-Gross-Prasad conjecture for unitary groups: the Archimedean case. *arXiv:1506.01452*, 2015. to appear in *Astérisque*.
- [BP18a] R. Beuzart-Plessis. Archimedean theory and ϵ -factors for the Asai Rankin-Selberg integrals. *arXiv:1812.00053*, 2018. prepublication.
- [BP18b] R. Beuzart-Plessis. Plancherel formula for $GL_n(F)\backslash GL_n(E)$ and applications to the Ichino-Ikeda and formal degree conjectures for unitary groups. *arXiv:1812.00047*, 2018. prepublication.
- [Cas89] W. Casselman. Introduction to the Schwartz space of $\Gamma\backslash G$. *Canad. J. Math.*, 41(2):285–320, 1989.

- [Cog08] J. Cogdell. Notes on L -functions for GL_n . In *School on Automorphic Forms on $GL(n)$* , volume 21 of *ICTP Lect. Notes*, pages 75–158. Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2008.
- [CZ] P.-H. Chaudouard and M. Zydor. Le transfert singulier pour la formule des traces de Jacquet-Rallis. *Comp. Math.*, accepted for publication.
- [FL17] T. Finis and E. Lapid. On the analytic properties of intertwining operators I: global normalizing factors. *Bull. Iranian Math. Soc.*, 43(4):235–277, 2017.
- [Fli88] Y. Flicker. Twisted tensors and Euler products. *Bull. Soc. Math. France*, 116(3):295–313, 1988.
- [Fli91] Y. Flicker. On distinguished representations. *J. Reine Angew. Math.*, 418:139–172, 1991.
- [FLO12] B. Feigon, E. Lapid, and O. Offen. On representations distinguished by unitary groups. *Publ. Math. Inst. Hautes Études Sci.*, 115:185–323, 2012.
- [GGP12] W. T. Gan, B. Gross, and D. Prasad. Symplectic local root numbers, central critical L values, and restriction problems in the representation theory of classical groups. Number 346, pages 1–109. 2012. Sur les conjectures de Gross et Prasad. I.
- [GJR09] D. Ginzburg, D. Jiang, and S. Rallis. Models for certain residual representations of unitary groups. In *Automorphic forms and L -functions I. Global aspects*, volume 488 of *Contemp. Math.*, pages 125–146. Amer. Math. Soc., Providence, RI, 2009.
- [GK72] I. M. Gelfand and D. A. Každan. Representations of the group $GL(n, K)$ where K is a local field. *Funkcional. Anal. i Priložen.*, 6(4):73–74, 1972.
- [GL06] S. Gelbart and E. Lapid. Lower bounds for L -functions at the edge of the critical strip. *Amer. J. Math.*, 128(3):619–638, 2006.
- [Gol94] D. Goldberg. Some results on reducibility for unitary groups and local Asai L -functions. *J. Reine Angew. Math.*, 448:65–95, 1994.
- [Gro53] A. Grothendieck. Sur certains espaces de fonctions holomorphes. I. *J. Reine Angew. Math.*, 192:35–64, 1953.
- [Gro55] A. Grothendieck. Produits tensoriels topologiques et espaces nucléaires. *Mem. Amer. Math. Soc.*, 16:Chapter 1: 196 pp.; Chapter 2: 140, 1955.
- [Gro73] A. Grothendieck. *Topological vector spaces*. Gordon and Breach Science Publishers, New York-London-Paris, 1973. Translated from the French by Orlando Chaljub, Notes on Mathematics and its Applications.
- [Gro97] B. Gross. On the motive of a reductive group. *Invent. Math.*, 130(2):287–313, 1997.
- [GRS11] D. Ginzburg, S. Rallis, and D. Soudry. *The descent map from automorphic representations of $GL(n)$ to classical groups*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2011.
- [Har14] R. N. Harris. The refined Gross-Prasad conjecture for unitary groups. *Int. Math. Res. Not. IMRN*, (2):303–389, 2014.
- [IY15] A. Ichino and S. Yamana. Periods of automorphic forms: the case of $(GL_{n+1} \times GL_n, GL_n)$. *Compos. Math.*, 151(4):665–712, 2015.
- [IY19] A. Ichino and S. Yamana. Periods of automorphic forms: the case of $(U_{n+1} \times U_n, U_n)$. *J. Reine Angew. Math.*, 746:1–38, 2019.

- [Jac04] H. Jacquet. Integral representation of Whittaker functions. In *Contributions to automorphic forms, geometry, and number theory*, pages 373–419. Johns Hopkins Univ. Press, Baltimore, MD, 2004.
- [Jac09] H. Jacquet. Archimedean Rankin-Selberg integrals. In *Automorphic forms and L-functions II. Local aspects*, volume 489 of *Contemp. Math.*, pages 57–172. Amer. Math. Soc., Providence, RI, 2009.
- [Jac10] H. Jacquet. Distinction by the quasi-split unitary group. *Israel J. Math.*, 178:269–324, 2010.
- [JLR99] H. Jacquet, E. Lapid, and J. Rogawski. Periods of automorphic forms. *J. Amer. Math. Soc.*, 12(1):173–240, 1999.
- [JPSS83] H. Jacquet, I. I. Piatetskii-Shapiro, and J. A. Shalika. Rankin-Selberg convolutions. *Amer. J. Math.*, 105(2):367–464, 1983.
- [JR11] H. Jacquet and S. Rallis. On the Gross-Prasad conjecture for unitary groups. In *On certain L-functions*, volume 13 of *Clay Math. Proc.*, pages 205–264. Amer. Math. Soc., Providence, RI, 2011.
- [JS81a] H. Jacquet and J. A. Shalika. On Euler products and the classification of automorphic forms. II. *Amer. J. Math.*, 103(4):777–815, 1981.
- [JS81b] H. Jacquet and J. A. Shalika. On Euler products and the classification of automorphic representations. I. *Amer. J. Math.*, 103(3):499–558, 1981.
- [JZ] D. Jiang and L. Zhang. Arthur parameters and cuspidal automorphic modules of classical groups. arXiv:1508.03205.
- [Kem15] A. Kemarsky. A note on the Kirillov model for representations of $GL_n(\mathbb{C})$. *C. R. Math. Acad. Sci. Paris*, 353(7):579–582, 2015.
- [KMSW] T. Kaletha, A. Mínguez, S.-W. Shin, and P.-J. White. Endoscopic classification of representations: Inner forms of unitary groups. arXiv:1409.3731.
- [KS88] C. Keys and F. Shahidi. Artin L -functions and normalization of intertwining operators. *Ann. Sci. École Norm. Sup. (4)*, 21(1):67–89, 1988.
- [Lap06] E. Lapid. On the fine spectral expansion of Jacquet’s relative trace formula. *J. Inst. Math. Jussieu*, 5(2):263–308, 2006.
- [Lap08] E. Lapid. A remark on Eisenstein series. In *Eisenstein series and applications*, volume 258 of *Progr. Math.*, pages 239–249. Birkhäuser Boston, Boston, MA, 2008.
- [Lap11] E. Lapid. On Arthur’s asymptotic inner product formula of truncated Eisenstein series. In *On certain L-functions*, volume 13 of *Clay Math. Proc.*, pages 309–331. Amer. Math. Soc., Providence, RI, 2011.
- [Lap13] E. Lapid. On the Harish-Chandra Schwartz space of $G(F)\backslash G(\mathbb{A})$. In *Automorphic representations and L-functions*, volume 22 of *Tata Inst. Fundam. Res. Stud. Math.*, pages 335–377. Tata Inst. Fund. Res., Mumbai, 2013. With an appendix by Farrell Brumley.
- [LR03] E. Lapid and J. Rogawski. Periods of Eisenstein series: the Galois case. *Duke Math. J.*, 120(1):153–226, 2003.
- [LW13] J.-P. Labesse and J.-L. Waldspurger. *La formule des traces tordue d’après le Friday Morning Seminar*, volume 31 of *CRM Monograph Series*. American Mathematical Society, Providence, RI, 2013. With a foreword by Robert Langlands [dual English/French text].

- [Mö0] W. Müller. On the singularities of residual intertwining operators. *Geom. Funct. Anal.*, 10(5):1118–1170, 2000.
- [Mok15] C. P. Mok. Endoscopic classification of representations of quasi-split unitary groups. *Mem. Amer. Math. Soc.*, 235(1108):vi+248, 2015.
- [MS04] W. Müller and B. Speh. Absolute convergence of the spectral side of the Arthur trace formula for GL_n . *Geom. Funct. Anal.*, 14(1):58–93, 2004. With an appendix by E. M. Lapid.
- [MW89] C. Mœglin and J.-L. Waldspurger. Le spectre résiduel de $GL(n)$. *Ann. Sci. École Norm. Sup. (4)*, 22(4):605–674, 1989.
- [MW94] C. Mœglin and J.-L. Waldspurger. *Décomposition spectrale et séries d'Eisenstein*, volume 113 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 1994. Une paraphrase de l'Écriture. [A paraphrase of Scripture].
- [Ram18] D. Ramakrishnan. A Theorem on $GL(n)$ à la Tchebotarev. *arXiv e-prints*, page arXiv:1806.08429, Jun 2018.
- [Sha81] F. Shahidi. On certain L -functions. *Amer. J. Math.*, 103(2):297–355, 1981.
- [Sha90] F. Shahidi. A proof of Langlands' conjecture on Plancherel measures; complementary series for p -adic groups. *Ann. of Math. (2)*, 132(2):273–330, 1990.
- [Trè67] F. Trèves. *Topological vector spaces, distributions and kernels*. Academic Press, New York-London, 1967.
- [Wal92] N. Wallach. *Real reductive groups. II*, volume 132 of *Pure and Applied Mathematics*. Academic Press, Inc., Boston, MA, 1992.
- [Wei82] A. Weil. *Adeles and algebraic groups*, volume 23 of *Progress in Mathematics*. Birkhäuser, Boston, Mass., 1982. With appendices by M. Demazure and Takashi Ono.
- [Xue19] H. Xue. On the global Gan-Gross-Prasad conjecture for unitary groups: approximating smooth transfer of Jacquet-Rallis. *J. Reine Angew. Math.*, 756:65–100, 2019.
- [Yun11] Z. Yun. The fundamental lemma of Jacquet and Rallis. *Duke Math. J.*, 156(2):167–227, 2011. With an appendix by Julia Gordon.
- [Zha14a] W. Zhang. Automorphic period and the central value of Rankin-Selberg L -function. *J. Amer. Math. Soc.*, 27:541–612, 2014.
- [Zha14b] W. Zhang. Fourier transform and the global Gan-Gross-Prasad conjecture for unitary groups. *Ann. of Math. (2)*, 180(3):971–1049, 2014.
- [Zyd16] M. Zydor. La variante infinitésimale de la formule des traces de Jacquet-Rallis pour les groupes unitaires. *Canad. J. Math.*, 68(6):1382–1435, 2016.
- [Zyd18] M. Zydor. La variante infinitésimale de la formule des traces de Jacquet-Rallis pour les groupes linéaires. *J. Inst. Math. Jussieu*, 17(4):735–783, 2018.
- [Zyd20] M. Zydor. Les formules des traces relatives de Jacquet-Rallis grossières. *J. Reine Angew. Math.*, 762:195–259, 2020.

Raphaël Beuzart-Plessis
Aix Marseille Univ
CNRS
Centrale Marseille
I2M

Marseille

France

email:

raphael.beuzart-plessis@univ-amu.fr

Pierre-Henri Chaudouard

Université de Paris

CNRS

Institut de Mathématiques de Jussieu-Paris Rive Gauche

F-75013 PARIS

France

email:

Pierre-Henri.Chaudouard@imj-prg.fr

Michał Zydor

University of Michigan

Ann Arbor, MI US

email:

zydor@umich.edu