



**HAL**  
open science

**Late Triassic (Tuvalian – Carnian, Tropites subbullatus/Anatropites spinosus zones) ostracods from Monte Gambanera (Castel di Iudica, Central-Eastern Sicily, Italy)**

S. Crasquin, Francesco Sciuto, Agatino Reitano, Rosa Maria Coco

► **To cite this version:**

S. Crasquin, Francesco Sciuto, Agatino Reitano, Rosa Maria Coco. Late Triassic (Tuvalian – Carnian, Tropites subbullatus/Anatropites spinosus zones) ostracods from Monte Gambanera (Castel di Iudica, Central-Eastern Sicily, Italy). *Bulletin de la Société Géologique de France*, 2020, <10.1051/bsgf/2020031>. <hal-02990524>

**HAL Id: hal-02990524**

**<https://hal.science/hal-02990524v1>**

Submitted on 9 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# Late Triassic (Tuvalian – Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones) ostracods from Monte Gambanera (Castel di Iudica, Central-Eastern Sicily, Italy)

Sylvie Crasquin<sup>1,\*</sup>, Francesco Sciuto<sup>2,\*</sup>, Agatino Reitano<sup>3</sup> and Rosa Maria Coco<sup>2</sup>

<sup>1</sup> CR2P – CNRS, MNHN, Sorbonne Université, 4, Place Jussieu, T. 46-56, E.5, F-75252 Paris Cedex 05, France

<sup>2</sup> Palaeoecological Research Group, Department of Biological, Geological and Environmental Science, Catania University, Corso Italia, 55, Catania 95129, Italy

<sup>3</sup> Natural History Museum, via degli Studi 9, 97013 Comiso, Ragusa, Italy

Received: 28 October 2019 / Accepted: 2 July 2020

**Abstract** – Ostracod associations coming from the Upper Triassic (*Tropites subbullatus*/*Anatropites spinosus* zone of the Tuvalian substage) clays and sandstones of the Mufara Formation outcropping along the west side of Monte Gambanera (Castel di Iudica, central-eastern Sicily) have been analysed for the first time. The specimens are relatively abundant, silicified, well preserved and often preserved as complete carapaces. Over 200 specimens have been determined. They belong to the families Healdiidae, Bairdiidae, Bythocypridae, Acratiidae, Cytheruridae, Limnocytheridae, Candonidae, Cavellinidae, Polycopidae and Thaumatoocypridae. Thirty-seven species are identified of which nine species are new: *Hungarella forelae* n.sp., *Hungarella siciliensis* n.sp., *Bairdia andreocrasquini* n.sp., *Bairdia gambaneraensis* n.sp., *Ptychobairdia iudicaensis* n.sp., *Ptychobairdia leonardi* n.sp., *Petasobairdia jeandercourti* n.sp., *Kerocythere dittainoensis* n.sp. and *Mockella barbroae* n.sp.

**Keywords:** Ostracods / systematic / Late Triassic / Neo-Tethys / Central-Eastern Sicily / Mount Gambanera / Mufara Formation

**Résumé** – Ostracodes du Trias supérieur (Tuvalien, Carnien, zones à *Tropites subbullatus*/*Anatropites spinosus*) de Monte Gambanera (Castel di Iudica, Sicile Centre Est, Italie). Pour la première fois est ici analysée une association d'ostracodes provenant du Trias supérieur (zones à *Tropites subbullatus*/*Anatropites spinosus* du sous étage Tuvalien) dans les argiles et grès de la Formation Mufara affleurant le flanc ouest du Mont Gambanera (Castel di Iudica, Sicile Centre Est). Les spécimens, silicifiés, sont relativement abondants, bien préservés et les plus souvent retrouvés sous formes de carapaces complètes. Plus des 200 spécimens ont été identifiés. Ils appartiennent aux familles Healdiidae, Bairdiidae, Bythocypridae, Acratiidae, Cytheruridae, Limnocytheridae, Candonidae, Cavellinidae, Polycopidae and Thaumatoocypridae. Trente-sept espèces sont reconnues dont sont nouvelles: *Hungarella forelae* n.sp., *Hungarella siciliensis* n.sp., *Bairdia andreocrasquini* n.sp., *Bairdia gambaneraensis* n.sp., *Ptychobairdia iudicaensis* n.sp., *Ptychobairdia leonardi* n.sp., *Petasobairdia jeandercourti* n.sp., *Kerocythere dittainoensis* n.sp. and *Mockella barbroae* n.sp.

**Mots clés :** ostracodes / systématique / Trias supérieur / Néo-Téthys / Sicile centre est / Mont Gambanera / Formation Mufara

## 1 Introduction

This is the second contribution on the Late Triassic ostracod fauna of the Mufara Formation outcropping in central eastern Sicily. Previously the ostracod fauna of the *Tropites dilleri*

zone of the Tuvalian substage outcropping at Monte Scalpello has been analysed (Crasquin *et al.*, 2018).

Now, a sedimentary level which is stratigraphically higher than the previous one and referable to the *Tropites subbullatus*/*Anatropites spinosus* zone of the Tuvalian substage, has been identified at the western side of the Monte Gambanera, nine kilometres south of Monte Scalpello (Fig. 1). The samples provided a rich and mostly well-preserved ostracod fauna.

\*Corresponding author:

[sylvie.crasquin@sorbonne-universite.fr](mailto:sylvie.crasquin@sorbonne-universite.fr)




**Fig. 1.** Geographical location of Monte Gambanera, Sicily, Italy and sample locality.

## 2 Geological setting and studied samples

Monte Gambanera is a modest relief located in central eastern Sicily (F 269 III NE of the Carta d'Italia alla scala 1:25 000) to the southeast of the town of Castel di Iudica (EN), about 40 kilometres west of Catania (Fig. 1). Structurally Monte Gambanera is part of the “Monte Judica Units” (Lentini *et al.*, 1987) and is inserted along the northern margin of the Gela Foredeep, in the geodynamic context of the southern end of the Maghrebian–Sicilian Southern Apennine nappes (Fig. 2) (Lentini *et al.*, 1987; Grasso, 2001 *inter alia*s).

In the Monte Gambanera area, the outcropping sediments are assigned to the Neo-Tethyan Mesozoic-Cenozoic complex which belongs to the so-called Imerese Succession (Lentini *et al.*, 1987; Montanari, 1987; *inter alia*s) or Imerese-Sicano Succession (Carrillat and Martini, 2009; Di Paolo *et al.*, 2012). The Imerese Basin, where these sedimentary successions were deposited, was delimited by the Panormide Carbonate Platform to the west and the Trapanese Carbonate Platform to the east and south (Catalano and D’Argenio, 1982; Montanari, 1987; Speranza and Minelli, 2014). In this basin, therefore, occurs a transitional facies between the Panormide and Trapanese shelf facies, on the one hand, and a deep marine facies of the Neo-Tethys, on the other. The sedimentary succession of Monte Gambanera (Fig. 2) starts with the “Carnian Flysch” (Auct.) or the Mufara Formation (Schmidt di Friedberg and Trovò, 1962). This unit, outcropping in the



**Fig. 2.** Stratigraphic series of Monte Gambanera, Sicily, Italy.

southern slopes of the mount, mainly consists of dark grey pelites, which locally contain rare ammonites, with rare interbeds of fossiliferous calcarenites and fibrous calcite with *Halobia* spp. imprints. The Mufara Fm. has been analysed since the beginning of the nineteenth century by Calcara (1840, 1845), Nelli (1899a, b) and subsequently by Gemmellaro (1904), Scalia (1907a, b, 1909, 1910–1914), Maugeri Patanè (1934) and Lentini (1974). These latter authors attributed these sediments to the Carnian (Late Triassic).

Stratigraphically, the Mufara Fm. outcropping at Monte Scalpello, can be referred to the *Tropites dilleri* zone of the Tuvalian substage (Crasquin *et al.*, 2018) due to the presence of Trachyceratidae (*Neoprotrachyceras*, *Trachysagenites*, *Pamphagosirenites*) and Tropitidae. A foraminifera, conodont and palynomorph biostratigraphical analysis, allowed to attribute the levels of the Mufara Fm. outcropping at Monte Gambanera to the *Tropites subbullatus/Anatropites spinosus* zone of the Tuvalian substage (Fig. 3) (Carrillat, 2001; Carrillat and Martini, 2009). In this stratigraphic horizon, cropping out near masseria Balconere at the west side of Mount Gambanera, two levels consisting of slightly silty clays have been sampled (Fig. 1). As they are stratigraphically very close and the number of specimens is quite low, we consider here the ostracod assemblages of both samples in all. They represent the ostracod association of the present study.

Twenty kilograms of sediments were collected from each of the two stratigraphic levels. Sediments were routinely washed, dried in oven and sieved. Then, ostracod specimens were picked out from the 63 µm fraction. The ostracod specimens were examined and measured under a stereomicroscope, then photographed under an LMU Tescan Vega II SEM. The material is housed in the Palaeontological Museum of the University of Catania. The repository number of the specimens are given in the systematic descriptions and/or in plate explanations. Over 200 specimens have been picked out from the two samples. The specimens are silicified, quite well preserved and often consist of complete carapaces.

## 3 Systematic palaeontology

**Abbreviations.** L: length; H: height; W: width; RV: right valve; LV: left valve; DB: dorsal border; VB: ventral border; AB: anterior border; PB: posterior border; PVB: postero-ventral border; AVB: antero-ventral border; PDB: postero-

STAGE	SUB STAGE	Ammonoid zones (Tethyan)	
CARNIAN	Tuvanian	Anatropites spinosus	
		Tropites subbullatus	
		Tropites dilleri	
	Julian	Austrotrachyceras austriacum	
		Trachyceras aonoides	
		Trachyceras aon	
		Daxatina canadensis	
	-227 ma		
-237 ma			

Fig. 3. Carnian ammonoid zones in Monte Scalpello (Crasquin *et al.*, 2018) and Monte Gambanera (present study) (after Lucas, 2010 modified).

dorsal border; ADB: antero-dorsal border. We follow here the general classification of Moore (1961) and Horne *et al.* (2002).

- Class Ostracoda Latreille, 1806
- Subclass Podocopa Müller, 1894
- Order Metacopida Sylvester-Bradley, 1961
- Suborder Metacopina Sylvester-Bradley, 1961
- Superfamily Healdiioidea Harlton, 1933
- Family Healdiidae Harlton, 1933

The systematics of Mesozoic Healdiidae is quite complicated and an important revision is necessary. We can't do it here because we have not enough material and most of the discrimination between the genera were based on muscles scars which are not preserved in the present material. However, we try to establish a way to distinguish the Triassic Healdiidae genera when only the external characters of carapaces are available. Therefore we follow the work of Kristan-Tollmann (1973, 1979). The Palaeozoic forms are considered to belong to the subfamily Healdiinae Harlton, 1933. The Triassic forms

belong to Hungarellinae Kristan-Tollmann, 1971 and Liassic ones to the subfamily Pseudohealdiinae Gründel, 1964 (Kristan-Tollmann, 1971).

Subfamily Hungarellinae Kristan-Tollmann, 1971

In the Subfamily Hungarellinae, seven genera have been described so far: *Triadiohealdia* Kristan-Tollmann, 1971; *Aneisohealdia* Kristan-Tollmann, 1971; *Labratella* Gramm, 1970; *Hungarella* Méhes, 1911; *Ogmoconchella* Gründel 1964 emend Michelsen, 1975; *Signohealdia* Kristan-Tollmann, 1971; *Torohealdia* Kristan-Tollmann, 1971.

A great confusion exists in the systematics of Late Permian – Triassic Healdiidae genera *Hungarella* – *Ogmoconcha* – *Ogmoconchella*. Some authors consider *Hungarella* Méhes, 1911 (which has no type material – Gerry and Kozur, 1973; but the Hungarian original material in under revision by E. Tóth, pers. com.) and *Ogmoconcha* Triebel, 1941 as synonyms (Moore, 1961; Anderson, 1964). Shaver (*in* Moore 1961), Sohn (1968) and Kristan-Tollmann (1971, 1977a, b) don't agree with this synonymy. In fact, the two genera are close but the valves are strongly dissymmetric in shape in *Hungarella*. The third genus, *Ogmoconchella* was introduced by Gründel (1964) and emended by Michelsen (1975) mainly due to the presence of a spine at PVB. In a recent revision, Forel and Crasquin (submitted) considered that until the relationship of *Ogmoconcha* and *Hungarella* is clarified, *Hungarella* should only been used for Triassic species to avoid artificially rooting *Ogmoconcha* down to the Triassic. Morphologically, the left and right valves of *Hungarella* are asymmetrical contrary to those of *Ogmoconcha* (Kristan-Tollmann, 1977a, b; Lord, 1982): in the absence of observable central muscle scars, all Triassic occurrences of *Ogmoconcha* and *Ogmoconchella* are re-attributed to the genus *Hungarella*.

Genus *Hungarella* Méhes, 1911

Type species *Bairdia problematica* Méhes, 1911

*Hungarella forelae* n.sp.

(Plate 1A)

2018 *Ogmoconchella felsooersensis* (Kozur, 1970); Crasquin *et al.* : 133, figs. 6A-B.

**Etymology.** Dedicated to Dr. Marie-Béatrice Forel, Museum national d'Histoire naturelle, Paris.

**Material.** Four complete carapaces.


**Holotype.** One complete carapace, collection number PMC O 21 H 13/10/2019, Plate 1A

**Paratype.** One complete carapace, collection number PMC O 77 P 13/10/2019, Crasquin *et al.*, 2018, fig. 6A.


**Diagnosis.** A species of *Hungarella* with triangular shape carapace, a posteroventral spine at RV, delicate flattening in blade shape at anterior border of RV.

**Description.** Massive carapace with a symmetric triangular shape; quite symmetric relative to H max; shape of left and right valves similar; LV is significantly larger than RV and radius of curvature of PB smaller than anterior one; LV overlaps RV all around the carapace with minimum at PVB; maximum of H located in front of or at mid L; maximum of L at mid H or a little below; VB quite straight; presence of a very fine flattening all around the AB of RV in blade shape; small spine more or less distinct at PVB of RV; dorsal view biconvex with valves almost symmetric in shape and W max at or just behind mid L; surface seems to be smooth.

**Dimensions.** L = 706–919 µm; H = 529–622 µm (see Fig. 4).



**Plate 1** Ostracods from Late Triassic (Tuvalian-Carnian) of Monte Gambanera, Sicily, Italy. All the specimens are stored in the Palaeontological Museum of the University of Catania. The repository numbers are given as PCM (Palaeontological Museum Catania) O (Ostracods) X H (number of holotype) or X P (number of paratype) or FS X (Figured Specimen number) registration date. Scale bars = 200  $\mu\text{m}$ . A: *Hungarella forelae* n. sp. A: holotype, right lateral view of a complete carapace, PMC O 21 H13/10/2019; paratype figured in Figure 6A (Crasquin et al., 2018). B-C: *Hungarella siciliensis* n. sp. B: holotype, right lateral view of a complete carapace, PMC O 22 H 13/10/2019; C: paratype, right lateral view of a complete carapace, PMC O 78 P 13/10/2019. D: *Hungarella* sp. A: right lateral view of a complete carapace, PCM O FS49. E-F: *Bairdia andreocrasquini* n. sp. E: holotype right lateral view of a complete carapace, PMC O 23 H 13/10/2019; F: paratype, right lateral view of a complete carapace, PMC O 79 P 13/10/2019. G-H: *Bairdia gambaneraensis* n. sp. G: holotype, right lateral view of a complete carapace, PMC O 24 H 13/10/2019; H: paratype, right lateral view of a complete carapace, PMC O 80 P 13/10/2019. I: *Bairdia cassiana* (Reuss, 1869). Right lateral view of a complete carapace, PCM O FS50. J: *Bairdia* cf. *monostorii* Forel and Grădinaru, 2018. Right lateral view of a complete carapace, PCM O FS51. K: *Bairdia* sp. 1 sensu Crasquin et al., 2018. Right lateral view of a complete carapace, PCM O FS52. L: *Bairdia* sp. A. Right lateral view of a complete carapace, PCM O FS53. M: *Bairdia* sp. B. Right lateral view of a complete carapace, PCM O FS54. N: *Bairdia* sp. C. Right lateral view of a complete carapace, PCM O FS55. O: *Bairdia* sp. D. Right lateral view of a complete carapace, PCM O FS56. P: *Bairdia* sp. E. Right lateral view of a complete carapace, PCM O FS57. Q: *Bairdia* sp. F. Right lateral view of a complete carapace, PCM O FS58. R: *Hiatabairdia subsymmetrica* Kristan-Tollmann, 1970. Lateral view of a left valve, PCM O FS59.



**Fig. 4.** Height (H)/length (L) diagram of figured specimens of the two new *Hungarella* species. In blue: *H. forelae* n. sp.; in orange: *H. siciliensis* n. sp.

**Occurrence.** Tuvalian–Carnian, *Tropites dilleri* zone (Crasquin et al., 2018) and *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

**Remarks.** The present specimens are close to *Ogmoconchella felsooersensis* (Kozur, 1970) from the early Anisian of Hungary (Kozur, 1970, Monostori, 1995) and Romania (Sebe et al., 2013). In a previous paper (Crasquin et al., 2018) two of the present specimens were attributed to this *O. felsooersensis*. Based on the new material this determination was revised and they are attributed to *Hungarella forelae*. Here the carapace is more triangular with a smaller radius of curvature of PB; the blade at AB is also more expressed here. The specimens described by Forel et al. (2019b; Plate 4, particularly fig. E, K) as *Hungarella gommerii* Forel, 2019 from the Carnian of Sichuan (South China) are very close to our specimens. However, the Sichuan specimens are smaller (biggest one  $\approx L = 600 \mu\text{m}$ ,  $H = 400 \mu\text{m}$ ) and show a smaller radius of curvature at both extremities. The largest specimen of *H. forelae* (Fig. 4) presents some morphological variability: overlap less important, at RV: the blade is located only at the ventral part of AB and occurrence of a small spine at the upper part of it, at LV: anteroventral blade seems to be also present. We consider that these morphological variations could be the expression of ontogenic variations but some doubt remains.

*Hungarella siciliensis* n. sp.

(Plate 1B and 1C)

**Etymology.** The species name refers to Sicily where the locus typicus is located.

**Material.** Four complete carapaces.

**Holotype.** One complete carapace, collection number PCM O 22 H 13/10/2019, Plate 1B.

**Paratype.** One complete carapace, collection number PCM O 78 P 13/10/2019, Plate 1C.

**Diagnosis.** A species of *Hungarella* with triangular shape carapace, two posteroventral spines at RV, flattening in blade shape plus a spine at anterior border of RV, spine at AB of LV.

**Description.** Massive stocky carapace with a symmetric triangular shape; quite symmetric relative to H max; general shape of valves similar, but LV is significantly larger than RV and radius of curvature of PB smaller than anterior one; LV overlaps RV all around the carapace with minimum at PVB;

maximum of H located at mid L or in front of mid L; maximum of L at mid H or a little below mid H; VB quite straight; presence of a very fine flattening at AB of RV in blade shape and a spine located near the maximum of convexity of AB; two more or less distinct spines at PVB of RV; one spine at AB of LV; dorsal view biconvex with valves almost symmetric in shape and W max at or just behind mid L; surface seems to be smooth.

**Dimensions.** L = 606–760  $\mu\text{m}$ ; H = 503–533  $\mu\text{m}$  (see Fig. 4).

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this work).

**Remarks.** *Hungarella siciliensis* n.sp. is very close to *H. forelae* n.sp.. The shapes of the valves are similar. The main differences between the two species is the presence of 2 spines at PVB of RV, presence of a spine at AB of booth valves and the less distinct “blade” at the AB of *H. siciliensis* n.sp.. We can’t exclude that these differences are due to morphological variability of *H. forelae* n.sp. (sexual or ontogenic). However, for the time being we have not enough specimens to settle this question.

*Hungarella* sp. A

(Plate 1D)

**Material.** One complete carapace.

**Dimensions.** H = 538  $\mu\text{m}$ ; L = 775  $\mu\text{m}$ .

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Order Podocopida Sars, 1860

Suborder Bairdiocopina Gründel, 1967

Superfamily Bairdiodea Sars, 1866

Family Bairdiidae, Sars 1866

Genus *Bairdia* McCoy, 1844

Type species *Bairdia curta* McCoy, 1844

*Bairdia andreocrasquini* n. sp.

(Plate 1E and 1F)

**Etymology.** In memory and honour of Dr. André Crasquin, father of the first author.

**Material.** Three complete carapaces and one broken carapace.

**Holotype.** One complete carapace, collection number PMC O 23 H 13/10/2019 (Plate 1E).

**Paratype.** One complete carapace, collection number PMC O 79 P 13/10/2019 (Plate 1F).

**Diagnosis.** A species of *Bairdia* with an elongated carapace, flattened AB and PB, and a ventral ridge along the posterior part of the VB and PB.

**Description.** Bairdioid carapace, quite elongated (H/L = 0.44); DB straight at RV and slightly convex at LV; ADB and PDB straight and quite symmetric with respect to DB; AB with large radius of curvature and maximum located above mid-H, AB strongly flattened laterally; VB slightly concave; PB slender and pointed, maximum of curvature located at lower 1/3 of H, strongly flattened laterally; presence of the ventral ridge which begins in posterior part of VB and runs along PB.

**Remarks.** This species is quite original and differs from all other ones by the specific characters (flattened AB and PB, and a ventral ridge along the posterior part of the VB and PB.). The shape of carapace is comparable to *Bairdia* sp. 4 *sensu* Forel

et al. (2019b) from the Carnian of China and to *Bairdia liviae* Forel and Grădinaru, 2018 from the Anisian of North Dobrogea, Romania (Forel and Grădinaru, 2018) but this last species does not show the specific characteristics.

**Dimensions.** H = 372–415  $\mu\text{m}$ ; L = 372–415  $\mu\text{m}$ .

**Occurrence.** Tuvalian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Bairdia gambaneraensis* n. sp.

(Plate 1G and 1H)

2018 *Bairdia* cf. *deformata* Kollmann, 1963; Crasquin et al.: 134, fig. 6K-L.

**Etymology.** Referring to the locus typicus Monte Gambanera, Sicily, Italy.

**Material.** Four complete carapaces.

**Holotype.** One carapace, collection number PMC O 24H 13/10/2019 (Plate 1G).

**Paratype.** One complete carapace, collection number PMC O 80 P 13/10/2019 (Plate 1H).

**Diagnosis.** A species of *Bairdia* with a very compact carapace, a continuously arched dorsal boarder and flattened and crenulated ventral parts of AB and PB

**Description.** Bairdioid carapace, quite short (H/L = 0.6–0.7), LV overlaps RV all around the carapace with minimum at AB and anterior part of VB; LV: all the dorsal part regularly arched; AB with large radius of curvature with maximum at mid-H, VB almost straight; BP with large radius of curvature with maximum at lower 1/3 of H; PDB arched; RV: DB straight, ADB straight with an angulation of 145°–150° against DB; AB with large radius of curvature; AVB and PVB flattened laterally in its very external part and with very fine crenulation; VB slightly concave; bairdioid beak quite absent; PDB straight with an angle of 125°–130° with DB; Presence of a shoulder on medio-dorsal part of LV; carapace biconvex and quite slender in dorsal view.

**Dimensions.** L = 886–910  $\mu\text{m}$ ; H = 600–643  $\mu\text{m}$ .

**Occurrence.** Tuvalian, *Tropites dilleri* zone (Crasquin et al., 2018), *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this work).

**Remarks.** *Bairdia gambaneraensis* n.sp. is similar to *Bairdia deformata* Kollmann, 1963 from the Rhaetian of Austria. The carapace of the present material is longer and presents a shoulder at LV. The new species has exactly the same general shape of valves as *B. penovoidea* Bolz, 1971 from Late Norian–Rhaetian of Austria and differs from this species only by a larger AB, a longer DB and the ventral crenulation of AB and PB.

*Bairdia cassiana* (Reuss, 1869)

(Plate 1I)

1869 *Cythere cassiana* n.sp.; Reuss: 108.

1869 *Bairdia cassiana* (Reuss, 1869); Gümbel: 180, pl. 5, figs. 18-19.

1958 *Bairdia cassiana* (Reuss, 1869); Styk: 171, fig. 3/1.


1970 *Bairdia cassiana* (Reuss, 1869); Ulrichs: 705-706, pl. 1, figs. 1-2.

1978 *Bairdia cassiana* (Reuss, 1869); Kristan-Tollmann: 81, pl. 1, fig. 4; pl. 6, fig. 6.

1991 *Bairdia cassiana* (Reuss, 1869); Kristan-Tollmann et al.: 200, pl. 1, fig. 5.

1995 *Bairdia cassiana rotundidorsata* n.ssp.; Monostori: 42, Pl. 2, figs. 4-5.

- 1996 *Bairdia (Rectobairdia) garciai* n.sp.; Crasquin-Soleau and Grădinaru: 77-78, pl. 2, figs. 5, 8.
- 2013 *Bairdia cassiana* (Reuss, 1869); Monostori and Tóth: 310, pl. 2, figs. 7, 8, 10.
- 2014 *Bairdia cassiana* (Reuss, 1869); Mette *et al.*: pl. 2, fig. 1.
- 2014 *Bairdia cassiana* (Reuss, 1869); Monostori and Tóth: 26, pl. 1, fig. 14.
- 2018 *Bairdia cassiana* (Reuss, 1869); Crasquin *et al.*: 134, fig. 6M.
- 2019a *Bairdia cassiana* (Reuss, 1869), Forel *et al.*, in press, figs. 4F–H.
- Material.** Four complete carapaces.
- Dimensions.** H = 486–533  $\mu\text{m}$ ; L = 840–948  $\mu\text{m}$ .
- Occurrence.** Early Carnian, Late Triassic, Southern Alps, Italy (Reuss 1869; Gümbel 1869; Ulrichs 1970; Kristan-Tollmann 1978); Carnian, Late Triassic, Święty Krzyż Mountain, Poland (Styk, 1958); Carnian, Late Triassic, Transdanubian Range, Hungary (Kristan-Tollmann, 1991); Late Anisian, Middle Triassic, Balaton Highland, Hungary (Monostori, 1995); Early Anisian, Middle Triassic, North Dobrogea, Romania (Crasquin-Soleau and Grădinaru, 1996); Ladinian, Middle Triassic, Balaton Highland, Hungary (Monostori and Tóth, 2013, 2014); Middle Anisian, Middle Triassic, Northern Calcareous Alps, Austria (Mette *et al.*, 2014); Carnian, Late Triassic, Karavanke Mountains, Slovenia (Forel *et al.*, 2019b); Tuvalian–Carnian, Tropites dilleri zone (Crasquin *et al.*, 2018) and Tropites subbullatus/Anatropites spinosus zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).
- Bairdia* cf. *monostorii* Forel and Grădinaru, 2018 (Plate 1J)
- 2018 *Bairdia* cf. *humilis* Monostori, 1995; Crasquin *et al.*: 134, figs. 6i–j.
- Material.** One complete carapace and one broken carapace.
- Dimensions.** (complete carapace) H = 311  $\mu\text{m}$ ; L = 806  $\mu\text{m}$ .
- Remarks.** Forel and Grădinaru (2018) renamed *Bairdia humilis* Monostori, 1995 in *Bairdia monostorii* nom. nov.
- Occurrence.** Tuvalian–Carnian, Tropites dilleri zone (Crasquin *et al.*, 2018) and Tropites dilleri zone (Crasquin *et al.*, 2018) and Tropites subbullatus/Anatropites spinosus zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).
- Bairata* sp.1 *sensu* Crasquin, Sciuto, Reitano, 2018 (Plate 1K)
- 2018 *Bairdia* sp. 1; Crasquin *et al.*: 134, fig. 6N–O
- Material.** One complete carapace and one right valve.
- Dimensions.** (complete carapace) H = 462  $\mu\text{m}$ ; L = 800  $\mu\text{m}$ .
- Remarks.** This species of *Bairdia* is characterized by the BD which is underlined by a blade, and by the reticulation of the carapace. The upper part of PB is quite horizontal and its radius of curvature is small. This could be a new species.
- Occurrence.** Tuvalian–Carnian, Tropites dilleri zone (Crasquin *et al.*, 2018) and Tropites subbullatus/Anatropites spinosus zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).
- Bairdia* sp. A (Plate 1L)
- Material.** Two complete carapaces.
- Dimensions.** H = 400–440  $\mu\text{m}$ ; L = 785–882  $\mu\text{m}$ .
- Occurrence.** Tuvalian–Carnian, Tropites subbullatus/Anatropites spinosus zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).
- Bairdia* sp. B (Plate 1M)
- Material.** One complete carapace.
- Dimensions.** H = 600  $\mu\text{m}$ ; L = 1000  $\mu\text{m}$ .
- Occurrence.** Tuvalian–Carnian, Tropites subbullatus/Anatropites spinosus zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).
- Bairdia* sp. C (Plate 1N)
- Material.** One complete carapace.
- Dimensions.** H = 643  $\mu\text{m}$ ; L = 1147  $\mu\text{m}$ .
- Remarks.** This species is characterized by its triangular shape, the flattening of the ventral borders and the reticulated surface. The PB has a very small radius of curvature.
- Occurrence.** Tuvalian–Carnian, Tropites subbullatus/Anatropites spinosus zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).
- Bairdia* sp. D (Plate 1O)
- Material.** One complete carapace.
- Dimensions.** H = 496  $\mu\text{m}$ ; L = 1014  $\mu\text{m}$ .
- Occurrence.** Tuvalian–Carnian, Tropites subbullatus/Anatropites spinosus zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).
- Bairdia* sp. E (Plate 1P)
- Material.** One complete carapace.
- Dimensions.** H = 476  $\mu\text{m}$ ; L = 953  $\mu\text{m}$ .
- Remarks.** This elongated species shows a blade underlying the BD.
- Occurrence.** Tuvalian–Carnian, Tropites subbullatus/Anatropites spinosus zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).
- Bairdia* sp. F (Plate 1Q)
- Material.** Two complete carapaces.
- Dimensions.** H = 600–615  $\mu\text{m}$ ; L = 885–953  $\mu\text{m}$ .
- Remarks.** This compact species (H/L = 0.64–0.67) has a flattened BP and AB and a shoulder at the dorsal part of the right valve.
- Occurrence.** Tuvalian–Carnian, Tropites subbullatus/Anatropites spinosus zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).
- Genus *Hiatobairdia* Kristan-Tollmann, 1970
- Type species: *Hiatobairdia subsymmetrica* Kristan-Tollmann, 1970
- Hiatobairdia subsymmetrica* Kristan-Tollmann, 1970 (Plate 1R)
- 1970 *Hiatobairdia subsymmetrica* n. gen. n. sp.; Kristan-Tollmann: 268, pl. 35, figs. 1–3.
- 1976 *Hiatobairdia subsymmetrica* Kristan-Tollmann, 1970; Tollmann: 276, pl. 163, fig. 14.
- 1978 *Hiatobairdia subsymmetrica deformis* n. sp.; Kristan-Tollmann: 83, pl. 4, figs. 1–7.
- 1979 *Hiatobairdia subsymmetrica* Kristan-Tollmann, 1970; Kristan-Tollmann *et al.*: 147, pl. 6, fig. 4.



**Plate 2.** Ostracods from Late Triassic (Tuvalian-Carnian) of Monte Gambanera, Sicily, Italy. All the specimens are stored in the Palaeontological Museum of the University of Catania. The repository numbers are given as PMC (Palaeontological Museum Catania) O (Ostracods) X H (number of holotype) or X P (number of paratype) or FS X (Figured Specimen number) registration date. Scale bars = 200  $\mu\text{m}$  except P-Q, R = 100  $\mu\text{m}$ . A: *Hiatobairdia* sp. A. Right lateral view of a complete carapace, PMC O FS60. B: *Mirabairdia pernodosa* Tollmann, 1963. Right lateral view of a complete carapace, PMC O FS61. C-D: *Ptychobairdia iudicaensis* n. sp. C: holotype, right lateral view of a complete carapace, PMC O25 H 13/10/2019; D: paratype, right lateral view of a complete carapace, PMC O 81 P 13/10/2019. E-F: *Ptychobairdia leonardoii* n. sp. E: holotype, lateral view of a right valve, PMC O 26 H 13/10/2019; F: paratype, lateral view of a left valve, PMC O 82 P 13/10/2019. G-H: *Petasobairdia jeandercourti* n. sp. G: holotype, right lateral view of a complete carapace, PMC O 27 H 13/10/2019; H: paratype, left lateral view of a complete carapace, PMC O 83 P 13/10/2019. I: *Urobairdia angusta* Kollmann, 1963. Right lateral view of a complete carapace, PCM O FS62. J: *Bairdiacypris triassica* Kozur, 1971c. Right lateral view of a complete carapace, PCM O FS63. K: *Bythocypris* sp. A. Left lateral view of a complete carapace, PCM O FS64. L: *Acratia maugerii* Crasquin *et al.*, 2018. Left lateral view of a complete carapace, PCM O FS65. M-N: *Kerocythere dittainoensis* n. sp. M: holotype, lateral view of a right valve, PMC O 28 H 13/10/2019; N: paratype, lateral view of a left valve, PMC O 84 P 13/10/2019. O: *Renngartenella sanctaecrusis* Kristan-Tollmann, 1973 ( $\delta$ ). Right lateral view of a complete carapace, PCM O FS66. P-Q: *Mockella barbroae* n. sp. P: holotype, right lateral view of a complete carapace, PMC O 29 H 13/10/2019; Q: paratype, right lateral view of a complete carapace, number PMC O 85 P13/10/2019. R: *Simeonella brotzenorum* Sohn, 1968. Lateral view of a right valve, PCM O FS67.

2018 *Hiatobairdia subsymmetrica* Kristan-Tollmann, 1970; Crasquin *et al.*: 134, figs. 6F–H.

**Material.** Three left valves.

**Dimensions.** H = 412–569  $\mu\text{m}$ ; L = 812–969  $\mu\text{m}$ .

**Occurrence.** Early Carnian of South Tyrol, Italy (Tollmann, 1976; Kristan-Tollmann, 1978); Rhaetian of Austrian Alps (Kristan-Tollmann, 1970) and Central Iran (Kristan-Tollmann *et al.*, 1979); Tuvalian–Carnian *Tropites dilleri* zone (Crasquin *et al.*, 2018) and *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study)

*Hiatobairdia* sp. A  
(Plate 2A)

**Material.** One complete carapace and one right valve.

**Dimensions.** (complete carapace) H = 533  $\mu\text{m}$ ;  
L = 948  $\mu\text{m}$ .

**Remarks.** This species has a straight DB and presents a ridge at the dorso-median part of the RV.


**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study)

Genus *Mirabairdia* Kollmann, 1963

Type species *Mirabairdia pernodosa* Kollmann, 1963

*Mirabairdia pernodosa* Kollmann, 1963

(Plate 2B)



**Fig. 5.** Height (H)/length (L) Diagram for *Ptychobairdia iudicaensis* n.sp. In red: right valves; in blue left valves.

1963 *Mirabairdia pernodosa* n. sp.; Kollmann: 177-178, pl. 1, figs. 1-2, pl. 8, figs. 1-6.

1971a *Triebelina (Mirabairdia) pernodosa illyrica* n.spp.; Kozur: 17, fig. 1G.

1971a *Triebelina (Mirabairdia) balatonica* n. sp.; Kozur: 15-16, figs. 2I, 3C.

1971 *Mirabairdia pernodosa* Kollm.; Kristan-Tollmann: text-fig. 1/8.

1984 *Triebelina (Mirabairdia) pernodosa illyrica* Kozur; Salaj and Jendrekova: pl. 2, figs. 1-4.

2014 *Triebelina (Mirabairdia) pernodosa* (Kollmann, 1963); Monostori and Tóth: 27-28, pl. 2, figs. 7-8.

**Material.** One complete carapace, 13 RV and 2 LV.

**Dimensions.** (complete carapace and LV) H (without spines) = 453–507  $\mu\text{m}$ ; L = 826–923  $\mu\text{m}$ .

**Occurrence.** Anisian, Western Carpathians, Slovakia (Salaj and Jendrekova, 1984; Kozur, 1971a); Anisian, Balaton Highland, Hungary (Kozur, 1971a); Ladinian, Dolomites, South Tirol, Italy, (Kristan-Tollmann, 1971); Ladinian, Northern Calcareous Alps, Cassian beds, Austria (Kollmann, 1963); Ladinian E-Bakony, Hungary (Monostori and Tóth, 2014); Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Genus *Ptychobairdia* Kollmann, 1960  
Type species *Ptychobairdia kuepperi* Kollmann, 1960

*Ptychobairdia iudicaensis* n. sp.  
(Plate 2C and 2D)

**Etymology.** From the locus typicus Castel di Iudica, Sicily, Italy.

**Material.** Four complete carapaces.

**Holotype.** One complete carapace, collection number PMC O 25 H 13/10/2019 (Plate 2C).

**Paratype.** One complete carapace, collection number PMC O 81 P 13/10/2019 (Plate 2D).

**Diagnosis.** A species of *Ptychobairdia* with a reticulated carapace which is flattened laterally all around except at the ventral part; LV significantly higher than RV, presence of vertical sulci at antero-dorsal part of the carapace.

**Description.** Carapace massive, high (H/L = 0.6); surface reticulated;

LV: Flattened laterally all around with maximum at DB and PB; BD strongly arched; AB with quite large radius of curvature and maximum at mid H; VB almost straight; BP with a very small curvature; two vertical sulci in dorsal part; LV strongly overlapping RV all around with maximum at BD.

RV: Strongly flattened all around except in ventral part; presence of a sulcus in AD part; BD long; AB with quite small radius of curvature; VB gently concave at its anterior part; BP very slender; DB, ADB, AVB, PVB, PDB straight.

**Remarks.** *Ptychobairdia iudicaensis* n. sp. is comparable with *P. oberhauseri* Kollmann, 1960 from the Rhaetian of Austria (Kollmann, 1960) and the Carnian–Norian of Queen Charlotte Island, Canada (Arias and Lord, 2000). The latter species is longer, has a smaller AB and shows a horizontal sulcus. *P. iudicaensis* n. sp. differs from *P. kristanae* Kollmann, 1960 from the Rhaetian–Early Jurassic of Austria (Kollmann, 1960, 1963) and the late Carnian of Sicily (Crasquin *et al.*, 2018) by its reticulated carapace and the RV being clearly smaller than LV.


**Dimensions.** L = 720–1083  $\mu\text{m}$ ; H = 480–667  $\mu\text{m}$  (see Fig. 5).

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Ptychobairdia leonardo* n. sp.  
(Plate 2E and 2F)

**Etymology.** Dedicated to Leonardo Reitano, son of Agatino Reitano.

**Material.** Six right valves, eight left valves.



**Fig. 6.** Height (H)/length (L) diagram for *Ptychobairdia leonardoï* n. sp. In blue: left valves; in red: right valves.

**Holotype.** One right valve, collection number PMC O 26 H 13/10/2019 (Plate 2E).

**Paratype.** One left valve, collection number PMC O 82 P 13/10/2019 (Plate 2F).

**Diagnosis.** A species of *Ptychobairdia* with a short coarse reticulated carapace, strongly compressed and finely reticulated AB and PB and a central node.

**Description.** Carapace with massive coarse reticulation, flattened laterally at AB and PB; DB straight at both valves and parallel to VB; ontogenic modifications of DB: at RV with nodules or blade at biggest specimens, at LV development of shoulders at each extremities in largest specimens; AB and PB with small radius of curvature, flattened laterally and covered by a fine reticulation; VB straight to slightly concave, with development of adventral structure; presence of a big node in median part of the carapace.

**Remarks.** *Ptychobairdia leonardoï* n. sp. differs from other species by the specific characters. *Margarobairdia zapfei* Kristan-Tollmann 1983 from the Anisian of South China (Kristan-Tollmann, 1983) has a similar valve shape but a different ornamentation.

**Dimensions.** L = 610–776  $\mu\text{m}$ ; H = 362–553  $\mu\text{m}$  (see Fig. 6).

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Genus *Petasobairdia* Chen and Shi, 1982  
Type species *Petasobairdia bicornuta* Chen and Shi, 1982  
*Petasobairdia jeandercourti* n.sp.  
(Plate 2G and 2H)

**Etymology.** Dedicated to past Pr. Jean Dercourt, who was the mentor of the first author.

**Material.** One broken carapace and three left valves.

**Holotype.** One broken carapace, collection number PMC O 27 H 13/10/2019 (Plate 2G).

**Paratype.** One left valve, collection number PMC O 83 P 13/10/2019 (Plate 2H).

**Diagnosis.** A species of *Petasobairdia* with a long reticulated carapace and elongated nodes at ADB and PDB of both valves.

**Description.** Carapace long (H/L  $\approx$  0.4), reticulated, strongly laterally compressed along AB, AVB, VB, PVB, PB; DB long and straight at both valves; presence of an elongated node at ADB and PDB of both valves; LV with two big horns with large base at each extremities of DB.

**Dimensions.** (holotype and paratype without spines) H = 348–373  $\mu\text{m}$ ; L = 826–853  $\mu\text{m}$ .

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

**Remarks.** *Petasobairdia jeandercourti* n. sp. is comparable to *P. bicornuta* Chen and Shi, 1982 from the Late Permian of Hubei Province (Chen and Shi, 1982) which has a much shorter DB. *P. longispinosa* (Kozur, 1971a, b, c) from Anisian of Slovakia (Kozur, 1971a), Anisian and Middle Triassic of Romania (Salaj and Jendrekova, 1984; Crasquin-Soleau and Grădinaru, 1996; Sebe *et al.*, 2013), Anisian of Austria (Mette *et al.*, 2014), Ladinian of Hungary (Monostori and Tóth, 2013) and Carnian of Southern Turkey (Forel *et al.*, 2017) has a shorter DB and doesn't have AD and PD nodes.

Genus *Urobairdia* Kollmann, 1963

Type species: *Urobairdia austriaca* Kollmann, 1963

*Urobairdia angusta* Kollmann, 1963

(Plate 2I)

1963 *Urobairdia angusta* n.g. n. sp.; Kollmann: 167, pl. 6, figs. 1–4.

1965 *Urobairdia angusta* Kollmann, 1963; Széles: 414, pl. 6, fig. 5.

1995 *Bairdia (Urobairdia) angusta recta* n. ssp.; Monostori: 43, pl.3, figs. 3–4.

2010 *Urobairdia angusta* Kollmann, 1963; Zorn: 271–272, pl. 6, figs. 12–15.

2013 *Bairdia (Urobairdia) angusta* Kollmann, 1963; Monostori and Tóth: 7, pl. 1, figs. 10–12.

2014 *Bairdia (Urobairdia) angusta* Kollmann, 1963; Monostori and Tóth: 25–26, Pl. 1, fig. 14.

**Material.** Two complete carapaces

**Dimensions.** H = 440–500  $\mu\text{m}$ ; L = 826–871  $\mu\text{m}$ .

**Occurrence.** Late Norian, Zlambach Formation, Austria (Kollmann, 1963; Zorn, 2010); Anisian, Felsöors, Hungary (Monostori, 1995); Carnian, Nosztori Valley, Hungary (Széles, 1965); Ladinian–Carnian, Balaton Highland, Hungary (Monostori and Tóth, 2013, 2014); Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Genus *Bairdiacypris* Bradfield 1935

Type species: *Bairdiacypris deloi* Bradfield, 1935.

*Bairdiacypris triassica* Kozur, 1971a, b, c

(Plate 2J)

1911 ?*Bairdia silicula* Jones; Méhés: 16–17, pl. 1, figs. 19–21

1971 *Bairdiacypris triassica* n.sp.; Kozur: 5–6, figs. 2H–L

2013 *Bairdiacypris triassica* Kozur, 1971a, b, c; Monostori and Tóth: 313–314, pl. 3, figs. 7–8

2014 *Bairdiacypris triassica* Kozur, 1971a, b, c; Monostori and Tóth: 25, pl. 1, fig. 12

2017 *Bairdiacypris triassica* Kozur, 1971c; Forel *et al.*: fig. 10U

**Material.** Two complete carapaces

**Dimensions.** H = 361–374  $\mu\text{m}$ ; L = 774–812  $\mu\text{m}$

**Occurrence.** Early Carnian, Balaton highland, Hungary (Méhés, 1911; Kozur, 1971c), Ladinian, Nosztori Valley, Hungary (Monostori and Tóth, 2013); Ladinian–Carnian, Balaton Highland (Monostori and Tóth, 2014), Carnian, Mersin, Turkey (Forel et al., 2017); Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study)

Family Bythocypridae Maddocks, 1969

Genus *Bythocypris* Brady, 1880

Type species: *Bythocypris reniformis* Brady, 1880

*Bythocypris* sp. A

(Plate 2K)

**Material.** Two complete carapaces and two LV.

**Dimensions.** H = 433–500  $\mu\text{m}$ ; L = 775–1090  $\mu\text{m}$ .

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study)

Family Acratidae Gründel, 1952

Genus *Acratia* Delo, 1930

Type species: *Acratia typica* Delo, 1930.

*Acratia maugeri* Crasquin et al., 2018

(Plate 2L)

1991 *Acratia* sp.; Kristan-Tollmann: 196, pl. 1, fig. 1.

2013 *Acratia goemoeryi* (Kozur, 1970); Monostori and Tóth: 6–7, pl. 4, only fig. 2.

2018 *Acratia maugeri* n.sp.; Crasquin et al.: 137–138, fig. 7H–J.

**Material.** One complete carapace and two LV.

**Dimensions.** H = 373–464  $\mu\text{m}$ ; L = 866–1066  $\mu\text{m}$ .

**Occurrence.** Late Ladinian of NE Iran (Kristan-Tollmann, 1991), Balaton Highland, Hungary (Monostori and Tóth, 2013); Tuvalian–Carnian, *Tropites dilleri* zone (Crasquin et al., 2018) and *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study)

Superfamily Cytheroidea Müller-Baird, 1850

Family Cytheruridae Müller, 1894

Subfamily Cytherurinae Müller, 1894

Genus *Kerocythere* Kozur and Nicklas, 1970

Type species *Cythere raibliana* Gümbel, 1869

*Kerocythere dittainoensis* n.sp.

(Plate 2M and 2N)

**Etymology.** Referring to the Dittaino river which flows near to the locus typicus.

**Material.** Four complete carapaces.

**Holotype.** One complete carapace, collection number PMC O 28 H 13/10/2019 (Plate 2M).

**Paratype.** One complete carapace, collection number PMC O 84 P 13/10/2019 (Plate 2N).

**Diagnosis.** A species of *Kerocythere* Kozur and Nicklas, 1970 with a subrectangular reticulate carapace, presence of a lateral thick ridge which ascends at PB and occurrence of ventral ridges, one thick and several thinner ones parallel to VB.

**Description.** Carapace subrectangular, almost equilateral; BD long and straight, presence of a ridge on each side of hinge; presence of an eye spot; AB with large radius of curvature with maximum located below mid-H, flattened laterally and smooth; VB almost straight; PB with small radius of curvature with maximum around mid H, upper and lower part quite straight; H max at anterior angle; L max at PB; sulcus more or

less developed in anterior 1/3 of L; surface reticulated and ornamented with possible pustules and ridges: one lateral, thick, reaching from antero-ventral part of the carapace up to PB, ascending in posterior part; group of ventral ridges, one thick parallel to VB and several (at least three) below.

**Dimensions.** L = 651–731  $\mu\text{m}$ ; H = 309–376  $\mu\text{m}$ .

**Occurrence.** *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

**Remarks.** *Kerocythere dittainoensis* n.sp. could be compared to *Kerocythere reticulata* Kristan-Tollmann, 1972 from early Carnian of the Dolomites (Italy). This species doesn't show the ventral group of ridges but has one ridge at the AD part of the carapace following the AB. The new species is also close to *Kerocythere tricostata* Forel, 2017 from the middle Carnian of southern Tauride-Anatolide platform (Turkey; Forel et al., 2017). At the present material the lateral ridge is longer, ascends at its posterior part and the surface is reticulated.

Family indet.

Genus *Renngartenella* Schneider 1957 (in Mandelstam et al., 1957)

*Renngartenella sanctaecrucis* Kristan-Tollmann, 1973

(Plate 2O)

1973 *Renngartenella sanctaecrucis* Kristan-Tollmann; Kristan-Tollmann and Hamedani: 215, 217–219, pl. 8, figs. 1–6; pl. 11, figs. 1, 3, 5, 6; pl. 12, fig. 10.

1979 *Renngartenella sanctaecrucis* Kristan-Tollmann, 1973; Liebermann: 215, pl. 5, fig. 2.

1982 *Renngartenella sanctaecrucis* Kristan-Tollmann, 1973; Basha: pl. 1, fig. 15.

1990 *Renngartenella sanctaecrucis* Kristan-Tollmann, 1973; Gerry et al.: 96, pl. 1, figs 11–13.

1994 *Renngartenella sanctaecrucis* Kristan-Tollmann, 1973; Monostori: 320, 322, figs 5/5–7.

2001 *Renngartenella sanctaecrucis* Kristan-Tollmann, 1973; Keim et al.: fig. 8C.

2014 *Renngartenella sanctaecrucis* Kristan-Tollmann, 1973; Monostori and Tóth: 29–30, pl.3, figs. 10–12.

2019a *Renngartenella sanctaecrucis* Kristan-Tollmann, 1973; Forel et al.: fig. 7T–U, in press.

**Material.** Ten carapaces.

**Dimensions.** H = 316–439  $\mu\text{m}$ ; L = 567–900  $\mu\text{m}$ .

**Occurrence.** Julian, early Carnian, Heiligkreuz Formation, Italy (Kristan-Tollmann and Hamedani, 1973); Carnian, Late Triassic, Italian Alps (Lieberman, 1979); Cordevolian, Carnian, Jordan (Basha, 1982); Carnian, Israel (Gerry et al., 1990); Carnian, Balaton Highland, Hungary (Monostori 1994; Monostori and Tóth, 2014); Carnian, Dolomites, Northern Italy (Keim et al., 2001); Carnian, Karavanke Mountains, Slovenia (Forel et al., 2019b); Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study)

Superfamily Cypridoidea Baird 1845

Family Limnocytheridae Klie, 1938

Genus *Mockella* Bunza and Kozur, 1971

Type species: *Mockella muelleri* Bunza and Kozur, 1971; subsequent designation (Kozur, 1973)

*Mockella barbroae* n. sp.

(Plate 2P and 2Q)

*Mockella barbroae* n.sp.




Fig. 7. Height (H)/length (L) diagram for *Mockella barbroae* n. sp.

2018 *Mockella muelleri* Bunza and Kozur, 1971, Crasquin *et al.*: 139, figs. 7O-P.

**Etymology.** Personal dedication of the first author to Mrs. Barbro Lamy, in token of friendship and affection.

**Material.** Seven complete carapaces and two carapaces from Crasquin *et al.*, 2018

**Holotype.** One complete carapace, collection number PMC O 29 H 13/10/2019 (Plate 2P).

**Paratype.** One complete carapace, collection number PMC O 85 P 13/10/2019 (Plate 2Q).

**Diagnosis.** A species of *Mockella* with a long subrectangular carapace and a well-developed rib all around the carapace.

**Description.** Carapace sub rectangular, long (H/L=0.535); eye spot well developed; AB with maximum of curvature located low between mid-H and lower 1/3 of H; VB straight; PB equivalent to AB in heteromorphs and smaller in tecnomorphs with maximum of curvature located above mid-H; anteromedian sulcus located in front of mid-L; posterior lobe well developed; anterior lobe less distinct; presence of a distinct ridge all around the carapace including BD; presence of additional ridges on lateral surface of the valves:

- a median ridge which begins at maximum of curvature of AB and precedes on the posterior lobe; this ridge is high and stands out on the surface;
- a lateral ridge below the lobes parallel to VB;
- a small ridge in upper part of AB and below the eye spot.

The valve surface is reticulated with 4 small pustules distributed parallel to AB; in dorsal view, the flanks are parallel. Sexual dimorphism present, expressed by the thickness of the posterior part of the carapace in heteromorphs.

**Dimensions.** L = 270–492 μm; H = 150–275 μm (see Fig. 7).

**Occurrence.** Late Carnian (*Tropites dilleri* zone), Murfa Formation, Sicily, Italy Tuvalian–Carnian (Crasquin *et al.*, 2018) and Tuvalian–Carnian, *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

**Remarks.** *Mockella barbroae* n.sp. is very close to *M. muelleri* Bunza and Kozur, 1971 from the late Carnian of Tyrol, Austria (Bunza and Kozur, 1971) and the Carnian of Monte Cammarata, Sicily (Cafiero and De Capoa Bonardi, 1982). In the study on the Mufara Formation (Crasquin *et al.*, 2018) we attributed the specimens to the latter species. The very well preserved present material enabled us to review our attribution. At the present specimens the BP is larger, the median ridge ends at the posterior lobe and doesn't reach the BP; an additional ridge is present below the lobes and the flanks are parallel in dorsal view

Genus *Simeonella* Sohn, 1968

Type species: *Simeonella brotzenorum* Sohn, 1968.

*Simeonella brotzenorum* Sohn, 1968

(Plate 2R)

1968 *Simeonella brotzenorum* n. sp.; Sohn: 23-24, pl. 2, figs. 1–4, 6–8, 12–22.

1971 *Simeonella brotzenorum alpina* n. ssp.; Bunza and Kozur: 4-5, pl. 1, figs. 5–7, 13.

1971 *Simeonella brotzenorum* n. ssp.; Bunza and Kozur: 5-6, pl. 1, fig. 3.

1973 *Simeonella brotzenorum* Sohn, 1968; Kristan-Tollmann and Hamedani: text-fig. 13/2.

1974 *Simeonella brotzenorum* Sohn, 1968; Hirsch and Gerry: pl. 2, figs. 1-2.

1979 *Simeonella brotzenorum* Sohn, 1968; Lieberman: 103, pl. 5, figs. 6-7.

1979 *Simeonella brotzenorum alpina* Bunza and Kozur, 1971; Styk: 119, pl. 28, figs. 9-10.



**3.** Ostracods from Late Triassic (Tuvalian-Carnian) of Monte Gambanera, Sicily, Italy. All the specimens are stored in the Palaeontological Museum of the University of Catania. The repository numbers are given as PCM (Palaeontological Museum Catania) O (Ostracods) FS X (Figured Specimen number) registration date. A, B, C, D: Scale bars = 200 μm; E, F, G, H, I: scale bars = 100 μm. A: *Paracypris?* cf. *redcarensis* (Blake, 1876). Lateral view of a right valve, PCM O FS68. B: *Paracypris?* sp. A. Right lateral view of a complete carapace, PCM O FS68. C: *Bektasia* sp. A. Right lateral view of a complete carapace, PCM O FS69. D: *Bektasia* sp. B. Right lateral view of a complete carapace, PCM O FS70. E: Podocopida gen. sp. indet. Lateral view of a right valve, PCM O FS71. F: *Polycoppe baudi* Crasquin-Soleau and Grădinaru 1996. Lateral view of a complete carapace, PCM O FS72. G: *?Polycoppe densoreticulata* Monostori and Tóth, 2013. Lateral view of a complete carapace, PCM O FS73. H: *Polycoppe* sp. A. Lateral view of a complete carapace, PCM O FS74. I: *Thaumatomma?* sp. A. Right lateral view of a complete carapace, PCM O FS75.

1982 *Simeonella brotzenorum* Sohn, 1968; Basha: pl. 1, fig. 11.

1990 *Simeonella brotzenorum* Sohn, 1968; Gerry et al.: 95, pl. 1, figs. 3–5.

1994 *Simeonella brotzenorum nostorica* n. ssp; Monostori: 324–325, text-fig. 6/1–6.

2001 *Simeonella brotzenorum nostorica* Monostori, 1994; Keim et al.: fig. 8B.

**Material.** Two carapaces.

**Dimensions.** H = 269–296 μm; L = 446–488 μm.

**Occurrence.** Ladinian to Carnian, Makhtesh Ramon, Israel (Sohn, 1968; Hirsch and Gerry, 1974; Gerry et al., 1990); Carnian, Northern Calcareous Alps, Austria (Bunza and Kozur, 1971; Kristan-Tollmann and Hamedani, 1973); Carnian, Julian Alps, Italy (Lieberman, 1979; Keim et al., 2001); Carnian, Poland (Styk, 1958), Carnian, Jordan Valley, Jordan (Basha, 1982); Carnian, Balaton Highland, Hungary (Monostori, 1994; Monostori and Tóth, 2014); Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this work)

Family Candonidae Kaufmann, 1900

Genus *Paracypris* Sars, 1866

*Paracypris?* cf. *redcarensis* (Blake, 1876) (Plate 3A)

**Material.** One complete carapace.

**Dimensions.** H = 257 μm; L = 561 μm.

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte

Gambanera, Central-Eastern Sicily, Italy (this study)

*Paracypris?* sp. A

(Plate 3B)

**Material.** One carapace.

**Dimensions.** H = 442 μm; L = 631 μm.

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-

Eastern Sicily, Italy (this study)

Order Platycopida Sars, 1866

Suborder Platycopina Sars, 1866

Superfamily Cavellinoidea Egorov, 1950

Family Cavellinidae Egorov, 1950

Genus *Bektasia* Özdikmen, 2010

Type species: *Reubenella avnimelechi* Sohn, 1968.

*Bektasia* sp. A

(Plate 3C)

**Material.** One carapace.

**Dimensions.** H = 477 μm; L = 787 μm.

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-

Eastern Sicily, Italy (this study)

*Bektasia* sp. B

(Plate 3D)

**Material.** One carapace.

**Dimensions.** H = 545 μm; L = 763 μm.

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-

Eastern Sicily, Italy (this study)

Order, genus and species *in det.*

Podocopida gen. sp. indet.

(Plate 3E)

**Material.** One carapace.

**Dimensions.** H = 188 μm; L = 364 μm.

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-

Eastern Sicily, Italy (this study)

Subclass Myodocopa Sars, 1866

Order Myodocopida Sars, 1866

Superfamily Polycopidea Sars, 1866

Family Polycopidae Sars, 1866

Genus *Polycoppe* Sars, 1866

*Polycoppe baudi* Crasquin-Soleau and Grădinaru 1996

(Plate 3F)

1996 *Polycoppe baudi* n.sp.; Crasquin and Grădinaru: 15–16, figs. 1–3.

2013 *Polycoppe baudi* Crasquin and Grădinaru 1996; Sebe et al.: pl. 1, fig. 1.

**Material.** Two carapaces.

**Dimensions.** H = 330–328 μm; L = 357–376 μm.

**Occurrences.** Early–middle Anisian, Uzum Bair, Dobrogea, Roumania (Crasquin-Soleau and Grădinaru, 1996; Sebe et al., 2013); Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern

Sicily, Italy (this study).

?*Polycope densoreticulata* Monostori and Tóth, 2013  
(Plate 3G)

2013 *Polycope densoreticulata* n.sp.; Monostori and Tóth:  
5, pl. 1, figs. 4–7.

**Material.** Three carapaces.

**Dimensions.** H = 204–293 µm; L = 231–306 µm.

**Occurrence.** Ladinian, Balaton Highland, Hungary  
(Monostori and Tóth, 2013); Tuvalian–Carnian, *Tropites*  
*subbullatus*/*Anatropites spinosus* zones, Monte Gambanera,  
Central-Eastern Sicily, Italy (this study)

*Polycope* sp. A

(Plate 3H)

**Material.** One carapace.

**Dimensions.** H = 289 µm; L = 278 µm.

**Occurrences.** Tuvalian–Carnian, *Tropites subbullatus*/  
*Anatropites spinosus* zones, Monte Gambanera, Central-  
Eastern Sicily, Italy (this study)

Order Halocyprida Dana, 1855

Suborder Halocypridina Dana, 1853

Superfamily Thaumatoocypridoidea Müller, 1906

Family Thaumatoocypridae Müller, 1906

Genus *Thaumatomma* Kornicker and Sohn, 1976

Type species: *Thaumatomma piscifrons* Kornicker and  
Sohn, 1976

*Thaumatomma* ? sp.A

(Plate 3I)

**Material.** Two carapaces.

**Dimensions.** H = 525–600 µm; L = 575–600 µm.

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/  
*Anatropites spinosus* zones, Monte Gambanera, Central-  
Eastern Sicily, Italy (this study).

#### 4 Assemblage analysis

The assemblage is composed of 200 specimens belonging to 10 families (plus two undetermined families), 19 genera and 37 species. The 10 determined families present are: Healdiidae, Bairdiidae, Bythocypridae, Acraatiidae, Cytheruridae, Limnocytheridae, Candonidae, Cavellinidae, Polycopidae, Thaumatoocypridae. The relative abundance of the different families expressed by the numbers of genera and species is given in Figure 8.

The Bairdiidae, the most abundant family in number of species (53%) and genera (37%) (Fig. 8), are present in marine environments ranging from very shallow waters up to deep seas. The morphology of the family changes from massive thick-shelled forms in nearshore environments to elongate thin-shelled forms beyond continental slope (particularly in genus *Bairdia*). In the same way, the carapaces of Acraatiidae lengthen with depth (as example: *Acratia goemeryi* Kozur, 1970 from Early-Smithian- to Late-Carnian-Triassic; see Forel et al., 2017). Here these two families present thick and ornamented shells (Plates 1E–1R and 2A–2L) which testify an open marine environment with moderate energy. The Healdiidae do also show a change of morphology with depth. The specimens with massive shells and small spines are neritic inhabitants (Plate 1A–1D) of relatively nearshore muddy conditions. In the deep sea, the specimens are thin shelled, elongate with long spines (e.g. Palaeozoic genus *Timorhealdia* Bless, 1987). In other families, some genera also show





Fig. 8. Diversity of ostracod families from the *Tropites subbullatus*/*Anatropites spinosus* zones represented by number of genera (A) and species (B) in the samples of Mount Gambanera.

different morphological adaptation from neritic to deep sea environments (*Healdia*, *Microcheilinella* etc. see synthesis Tab. 1 in Crasquin and Horne). The ratios between these different morphologies have been used to characterize the depth of at Palaeozoic and Early Mesozoic environments since quite a long time (model of Lethiers and Raymond, 1991). The present assemblage doesn't include any unequivocal deep water taxa such as those discovered recently in the Carnian of Southern Turkey for example (Forel et al., 2017) or of Sichuan, South China (Forel et al., 2019b). Although the number of specimens is very low, the diversity is quite high with 10 determined families (plus 2 undetermined), 17 genera and 37 species. This biodiversity testifies normal marine conditions and absence of environmental stress. The taxon *Simeonella brotzenorum* SOHN, 1968 which is characteristic of brackish–hypersaline conditions (Gerry et al., 1990; Monostori, 1994) is present but with only 2 carapaces. Quite all the specimens are preserved with the complete carapace. After the death of the specimens, the carapaces tend to open in a few hours (Guernet and Lethiers, 1989). This could suggest a rapid burial in the sediments due to a high sedimentation rate. Similar taphonomic characteristics were also found by Pokorny (1964) and Oertli (1971) for pelitic layer associations deposited in basins with extremely rapid distal sedimentation. We compare the results of the *Tropites subbullatus*/*Anatropites*



**Fig. 9.** Diversity of ostracod families from the *Tropites dilleri* zone represented by the number of genera (A) and species (B) in the samples of Mount Scalpello (data from Crasquin *et al.*, 2018).

*spinus* zones (this study), with the data obtained in levels just below in *Tropites dilleri* zone (Crasquin *et al.*, 2018) (Fig. 9). Through time, the assemblage became more diversified as recorded by the increasing number of families (8 to 12), genera (14 to 18) and species (23 to 36). The percentage of Bairdiidae decreases in favour of two families being absent before, Cytheruridae and Limnocytheridae which include typical genera of nearshore environments such as *Simoenella*, *Mockella* and *Kerocythere*. We observe also the presence of the brackish–hypersaline species *Renngartenella sanctaecrucis* Kristan-Tolmann, which was suggested by Gerry *et al.* (1990) to be a stenohaline ostracod.

For Monostori (1994), the dominance of three genera *Kerocythere*–*Renngartenella*–*Simeonella* seems to be a signal of salinity variability. Although these genera are not dominant here, their presence testifies a shallowing of environment from the *Tropites dilleri* zone to the *Tropites subbullatus*/*Anatropites spinus* zone. The genus *Acratia* is a typical Palaeozoic form present both in Eifelian (neritic) and Thuringian (deep) mega-assemblages (see synthesis in Crasquin and Horne, 2018). Nevertheless, this genus survived the Permian–Triassic extinction events. In 2013, Crasquin and Forel mentioned the last occurrence of neritic *Acratia* in the


Spathian and of deep marine *Acratia* in the Anisian (Crasquin-Soleau and Grădinaru, 1996). Since then, some deep marine forms were also found in the Ladinian of Balaton Highland (Monostori and Tóth, 2013), in the Carnian of Turkey (Forel *et al.*, 2017) and Slovenia (Forel *et al.*, 2019b). In 2013, Monostori and Tóth, described *Acratia goemeryi* from Ladinian neritic sediments of a borehole in Hungary. Three species of *Acratia* from the Carnian of Karavanke Mountains, Slovenia, are figured in Forel *et al.* (2019b). The occurrence of *Acratia maugerii* in the present material confirms that *Acratia* occurs in neritic environments of the Carnian.

## 5 Conclusion


The palaeoecological interpretation of the sedimentary facies of the Mufara Formation is extremely difficult due to the absence of intact outcrops. The original stratification and the sedimentary structures have been completely destroyed because of continuous agricultural processing of the pelitic soils and their very consistency which determines frequent drifts and landslides. In very few and limited locations parallel laminations and sandy levels were observed. The only useful palaeoecologic data are those obtained from the palaeontological analysis. The results of the ostracod fauna analysis allow the following conclusions:

- 1 The ostracod assemblage doesn't yield any evidence of deep marine taxa both at Mt. Scalpello (Crasquin *et al.*, 2018) and at Mt. Gambanera. The Mufara Basin, therefore, can be interpreted as a shallow marine basin (Fig. 10) within the deepest and most distal part of a vast continental shelf where the carbonate platforms Panormide, Trapanese, Saccense were located. The deep marine ostracod fauna discovered recently in the Carnian of Southern Turkey (Forel *et al.*, 2017) or in the South China (Forel *et al.*, 2019a) suggests a deepening of the Neo-Tethys basin towards the more eastern areas.
- 2 The Mufara Basin was a site of rapid and intense sedimentation probably linked to rapid bottom currents which, sometimes, displaced and transported (also *in vivo*) microfaunas from more superficial neighbouring environments. It is pointed out here that the sediments of the Mufara Basin at Monte Gambanera do not show vortex structures which were recognized in the Mufara Basin at Monte Scapello. This suggests, that the sediment environment of the Mufara Formation outcrops at Monte Gambanera (Fig. 11) was more distal and less turbulent than that of Mt which was effected by strong bottom traction and swirling currents (Crasquin *et al.*, 2018).
- 3 According to many authors, the Mufara basin is located in a transitional position between the bathyal Neotethys facies to the south and southeast and the carbonate platforms that surround it (Figs. 10 and 11).

**Acknowledgements.** The authors are grateful to the reviewers and the editors for detailed suggestions and comments to the manuscript. Thanks are due to Mr. Alfio Viola (Electronic Microscopy Laboratory, University of Catania) for SEM photos assistance. Catania Palaeoecological Research Group contribution n° 456.



**Fig. 10.** Palaeogeographic reconstruction of Tethyan (left) and central Mediterranean (right) areas during Late Triassic (after Di Stefano *et al.*, 2015, modified).



**Fig. 11.** Schematic palaeoenvironmental model for the late Carnian Mufara Formation basin (see also Fig. 10).

We thank the two reviewers Emőke Tóth from Eötvös Loránd University, Budapest, Hungary and Wolfgang Mette from Innsbruck University, Austria for their fruitful comments to improve our paper.

This research was supported by the University of Catania “Piano della Ricerca 2016/2018 (code no. 227221132118)” and “PiaCeRi–Piano Incentivi per la Ricerca di Ateneo 2020-22 linea di intervento 2”.

## References

Anderson FW. 1964. Rhaetic Ostracoda. *Bulletin of the Geological Survey of Great Britain* 21: 133–174.  
 Arias C, Lord A. 2000. Upper Triassic marine Ostracoda from the Queen Charlotte Islands British Columbia Canada. *Revista Española de Micropaleontología* 32: 175–192.  
 Baird W. 1845. Arrangement of the British Entomostraca, with a list of species, particularly noticing those which have as yet been discovered within the bounds of the club. *Transactions of Berwick's Naturalist Club* 2: 145–158.  
 Baird W. 1850. *The Natural History of the British Entomostraca*. London: Ray Society, 364 p.  
 Basha SHS. 1982. Microfauna from the Triassic rocks of Jordan. *Revue de Micropaléontologie* 25: 3–11.  
 Blake JE. 1876. Subclass Entomostraca. In: Late R, Blake JR, eds. *The Yorkshire Lias*, pp. 429–435.

Bless MJM. 1987. Lower Permian ostracodes from Timor (Indonesia). *Proceedings of Koninklijke Nederlandse Akademie van Wetenschappen* 90: 1–13.  
 Bolz H. 1971. Die Zlambach-Schichten (alpine Obertrias) unter besonderer Berücksichtigung der Ostrakoden, 1: Ostrakoden der Zlambach-Schichten, besonders Bairdiidae. *Senckenbergiana Lethaea* 52: 129–283.  
 Bradfield HH. 1935. Pennsylvanian ostracods of the Ardmore Basin, Oklahoma. *Bulletin of American Paleontology* 22: 1–145.  
 Brady GS. 1880 Report on the Ostracoda dredged by H.M.S. Challenger during the Years 1873–1876. Report on the Scientific Results of the Voyage of H.M.S. Challenger during the years 1873–76. *Zoology* 1(Part 3): 1–184.  
 Bunz G, Kozur H. 1971. Beiträge zur Ostracodenfauna der tethyalen Trias. *Geologisch Paläontologische Mitteilungen Innsbruck* 1: 1–76.  
 Cafiero B, De Capoa Bonardi P. 1982. Biostratigrafia del Trias pelagico della Sicilia. *Bollettino della Società Paleontologica Italiana* 21(1): 35–71.  
 Calcara P. 1840. Monografie dei generi Clausilia e Bulimo, coll’aggiunta di alcune nuove specie di conchiglie siciliane esistenti nella collezione della Sig. Teresa Gargotta in Salinas, 54 p.  
 Calcara P. 1845. Cenzo sui molluschi viventi e fossili della Sicilia da servire di supplemento ed insieme di critiche osservazioni all’opera di R.A. Philippi. Reale stamperia e Libreria Palermo, 65 p.  
 Carrillat A. 2001. Palaeoenvironmental reconstruction of the Mufara Formation (Upper Triassic, Sicily): biostratigraphy, organic facies,

- sedimentological and geochemical approach. Thèse Université de Genève, Terre et Environnement 27: 269 p.
- Carrillat A, Martini R. 2009. Palaeoenvironmental reconstruction of the Mufara Formation (Upper Triassic, Sicily): High resolution sedimentology, biostratigraphy and sea-level changes. *Palaeogeography, Palaeoclimatology, Palaeoecology* 283: 60–76.
- Catalano R, D’Argenio B. 1982. Schema geologico della Sicilia occidentale. In: Catalano R, D’Argenio B, eds. *Guida alla geologia della Sicilia occidentale. Guide geologiche regionali*. Memorie della Società Geologica Italiana suppl. A2: 9–41.
- Chen DQ, Shi CG. 1982. Latest Permian ostracoda from Nantong, Jiangsu and from Miannyang, Hubei. *Bulletin of the Nanjing Institute of Geology and Palaeontology. Academia Sinica* 4: 105–152 (in Chinese).
- Crasquin S, Horne DJ. 2018. The palaeopsychrosphere in the Devonian. *Lethaia* 51: 547–563.
- Crasquin S, Sciuto F, Reitano A. 2018. Late Carnian (Tuvalian, Tropites dilleri zone) ostracods (Crustacea) from the Mufara Formation (Monte Scalpello, Central-Eastern Sicily, Italy). *Annales de Paléontologie* 104: 129–142.
- Crasquin-Soleau S, Grădinaru E. 1996. Early Anisian ostracode fauna from the Tulcea unit (Cimmerian North Dobrogean orogeny, Romania). *Annales de Paléontologie* 82: 59–116.
- Dana J. 1853. Tribe III: Cyproidea, Ostracoda. In: *Crustacea of United States Exploring Expedition during the Years 1838, 1839, 1840, 1841, 1842, under the Command of Charles Wilkes, U.S.N., with Atlas of 96 plates*, 13(2), 1277–1304. Philadelphia: C. Sherman.
- Delo DM. 1930. Some Upper Carboniferous ostracods from the Shale Basin of Western Texas. *Journal of Paleontology* 4: 152–178.
- Di Paolo L, Aldega L, Corrado S, Mastalerz M. 2012. Maximum burial and unroofing of Mt. Judica recess area in Sicily: implication for the Apenninic–Maghrebic wedge dynamics. *Tectonophysics* 530–531: 193–207.
- Di Stefano P, Favara R, Luzio D, Renda P, Cacciatori MS, Calò M, et al. 2015. A regional-scale discontinuity in western Sicily revealed by a multidisciplinary approach: A new piece for understanding the geodynamic puzzle of the southern Mediterranean. *Tectonics* 34: 1–19.
- Drexler E. 1958. Foraminiferen und Ostracoden aus dem Lias von Siebeldingen/Pfalz. *Geologisches Jahrbuch* 75: 475–554.
- Egorov VG. 1950. Ostrakody franskogo iarusu Russkoi platformy, 1. Kloedenellidae: Moscow-Leningrad Vses. Neft. Nauchno-Issled. Geol. Razved. Inst., Moscow, Filial, 140 p.
- Forel M-B, Crasquin S. submitted. Bounded by crises: an updated view of the evolution of marine ostracods during the Triassic. *Marine Micropalaeontology*.
- Forel M-B, Grădinaru E. 2018. First report of Ostracods (Crustacea) associated with Bithynian (Anisian, Middle Triassic) Tubiphytes-microbial reef in the North Dobrogean Orogen (Romania). *Papers in Palaeontology* 4(2): 211–244.
- Forel M-B, Tekin UK, Okuyucu C, Bedi Y, Tuncer A, Crasquin S. 2017. Discovery of a long-term refuge for ostracods (Crustacea) after the end Permian extinction: a unique Carnian (Late Triassic) fauna from the Mersin Melange, southern Turkey. *Journal of Systematic Palaeontology* 17(1): 9–58.
- Forel M-B, Kolar-Jurkovsek T, Jurkovsek B. 2019a. Ostracods from the “Raibl Beds” (Carnian, Late Triassic) of Belca section in Karavanke Mountains, Northwestern Slovenia. *Geodiversitas*, in press.
- Forel M-B, Thuy B, Wisshak M. 2019b. Digging into the ancestral stocks of Jurassic lineages: ostracods (Crustacea) from Carnian (Late Triassic) sponge mounds from the Maantang Formation (South China). *Bulletin de la Société Géologique de France* 190(9): 29.
- Gemmellaro GG. 1904. I Cefalopodi del Trias Superiore della regione occidentale della Sicilia. *Giornale di Scienze Naturali ed Economiche di Palermo* 24: 319.
- Gerry E, Honigstein A, Rosenfeld A, Hirsch F, Eshet Y. 1990. The Carnian salinity crisis: ostracods and palynomorphs as indicators of palaeoenvironment. In: Whatley R, Maybury C, eds. *Ostracoda and global events*. London: Chapman & Hall, pp. 87–99.
- Gramm MN. 1970. Otpechatki adduktora Triasovykh cytherellid (Ostracoda) Primor’ya i nekotorye teorii filembriogeneza. *Paleontologicheskii Zhurnal* 1: 88–103.
- Grasso M. 2001. The apenninic-Maghrebic orogen in southern Italy, Sicily and adjacent areas. In: Vai GB, Martini IP, eds. *Anatomy of an Orogen: The Apennines and Adjacent Mediterranean Basins*. Kluwer Academic Publication GB, pp. 255–286.
- Gründel J. 1962. Zur Taxionomie der Ostracoden der Gattendorfia Stufe Thüringens. *Freiderger Forschungshilfe C* 151: 51–105.
- Gründel J. 1964. Zur Gattung *Healdia* (Ostracoda) und zu einigen verwandten Formen aus dem unter Jura. *Geologie* 13: 1–469.
- Gründel J. 1967. Zur Grossgliederung der Ordnung Podocopida G.W. Müller, 1894. *Neues Jahrbuch für Geologie and Palaontologie, Monatshefte* 6: 321–332.
- Guernet C, Lethiers F. 1989. Ostracodes et recherche des milieux anciens : possibilités et limites. *Bulletin de la Société Géologique de France* 8(5): 577–588.
- Gümbel CW. 1869. Über Foraminiferen, Ostracoden und mikroskopische Tier-Ueberreste in den St. Cassianer und Raibler Schichten. *Jahrbuch Geologischen Reichsanstalt* 19: 175–186.
- Harlton BH. 1933. Micropaleontology of the Pennsylvanian Johns Valley Shale of Ouachita Mountains, Oklahoma and its relationships to the Mississippian Caney Shale. *Journal of Paleontology* 7: 3–29.
- Hirsch F, Gerry E. 1974. Conodont and Ostracode-biostratigraphy of the Triassic in Israel. *Schriftenreihe der Erdwissenschaftlichen Kommissionen, Österreichische Akademie der Wissenschaften* 2: 107–112.
- Horne DJ, Cohen A, Martens K. 2002. Taxonomy, morphology and biology of Quaternary and living Ostracoda. In: Holmes JA, Chiva AR, eds. *The Ostracoda: Applications in Quaternary Research*. American Geophysical Union, Geophysical Monograph 131: 5–36.
- Kaufmann A. 1900. Über zwei neue Candona-Arten aus der Schweiz. *Zoologischer Anzeiger* 23: 108–110.
- Keim L, Brandner R, Krystyn L, Mette W. 2001. Termination of carbonate slope progradation: An example from the Carnian of the Dolomites, Northern Italy. *Sedimentary Geology* 143: 303–323.
- Klie W. 1938. Ostracoda, Muschelkrebse. In: *Dahl, Die Tierwelt Deutschlands*. Jena 34(3): 1–230.
- Kollmann K. 1960. Ostracoden aus der alpinen Trias Österreichs. I Parabairdia n. g. und Ptychobairdia n. g. (Bairdiidae). *Jahrbuch der Geologischen Bundesanstalt* 5: 79–105.
- Kollmann K. 1963. Ostracoden aus der alpinen Trias. II. –Weitere Bairdiidae. *Jahrbuch der Geologischen Bundesanstalt Wien* 106: 121–203.
- Kornicker LS, Sohn IG, 1976. Phylogeny, ontogeny, and morphology of living and fossil Thaumatozypridacea (Myodocopa: Ostracoda). *Smithsonian Contribution to Zoology* 219: 1–124.
- Kozur H. 1970. Neue Ostracoden-Arten aus dem Obersten Anis des Bakonyhochlandes (Ungarn). *Berichte des naturwissenschaftlich-medizinischen Vereins in Innsbruck* 58: 384–428.
- Kozur H. 1971a. Die Bairdiacea des Trias. Teil. 1: Skulpturierte Bairdiidae aus Mitteltriassischen Flachwasserablagerungen. *Geologisch Paläontologische Mitteilungen Innsbruck* 1: 1–27.

- 1289 Kozur H. 1971b. Die Bairdiacea der Trias. Teil 2: Skulpturierte  
1290 Bairdiidae aus Mitteltriassischen tiefschelfablagerungen. *Geo-*  
1291 *logisch Paläontologische Mitteilungen Innsbruck* 1(5): 1–21. 1352
- 1292 Kozur H. 1971c. Die Bairdiacea der Trias. Teil 3: Einige neue Arten  
1293 triassischer Bairdiacea und Bemerkungen zur Herkunft der  
1294 Macrocyprididae (Cypridacea). *Geologisch Paläontologische*  
1295 *Mitteilungen Innsbruck* 1: 1–18. 1356
- 1296 Kozur H. 1973. Beiträge zur ostracodenfauna der Trias. *Geologisch*  
1297 *Paläontologische Mitteilungen Innsbruck* 3(5): 1–41. 1358
- 1298 Kozur H, Nicklas L. 1970. Ostrakoden aus dem Plattenkalk-Niveau  
1299 des Haupt Dolomites (Rheatikum). In: Mostler H, ed. *Beiträge zur*  
1300 *Mikrofazies und Stratigraphie von Tirol und Vorarlberg*. Festband  
1301 des Geologischen Institutes, 300-Jahr-Feier Universität Innsbruck,  
1302 pp. 309–320. 1359
- 1303 Kristan-Tollmann E. 1970. Einige neue Bairdien (Ostracoda) aus der  
1304 alpinen Trias. *Neues Jahrbuch für Geologie und Paläontologie*  
1305 135: 268–310. 1360
- 1306 Kristan-Tollmann E. 1971. Weitere Beobachtungen an skulptierten  
1307 Bairdiidae (Ostracoda) der alpinen Trias. *Neues Jahrbuch für*  
1308 *Geologie und Paläontologie* 139: 57–87. 1362
- 1309 Kristan-Tollmann E. 1972. Die obertriadischen Arten der Ostraco-  
1310 dengattung *Kerocythere* Kozur and Nicklas 1970, und ihr  
1311 stratigraphischer weit. *Erdoel-Erdgas-Zeitschrift* 88: 43–49. 1363
- 1312 Kristan-Tollmann E. 1973. Zur phylogenetischen und stratigraphi-  
1313 schen Stellung der triadischen Healdiiden (Ostracoda) II. *Erdoel-*  
1314 *Erdgas-Zeitschrift* 89: 150–155. 1364
- 1315 Kristan-Tollmann E. 1977a. On the development of the muscle-scar  
1316 patterns in Triassic Ostracoda. In: Löffler H, Danielopol D, eds. In:  
1317 *Proceedings of the Sixth International Symposium on Ostracoda*  
1318 (Saalfelden, Salzburg), pp. 133–143. 1365
- 1319 Kristan-Tollmann E. 1977b. Zur Evolution des Schließmuskelfeldes  
1320 bei Healdiidae und Cytherellidae (Ostracoda). *Neues Jahrbuch für*  
1321 *Geologie und Paläontologie* 10: 621–639. 1366
- 1322 Kristan-Tollmann E. 1978. Bairdiidae (Ostracoda) aus den ober-  
1323 triadischen Cassianer Schichten der Ruones-Wiesen bei Corvara in  
1324 Südtirol. Schriftenreihe der Erdwissenschaftlichen Kommissionen.  
1325 *Österreichische Akademie der Wissenschaften* 4: 77–104. 1367
- 1326 Kristan-Tollmann E. 1983. Ostracoden aus dem Oberanis von  
1327 Leidapo bei Guiyang in Süd China. *Neue Beiträge zur*  
1328 *Biostratigraphie der Tethys-Trias* 121–176. 1368
- 1329 Kristan-Tollmann E. 1991. Ostracods from the Middle Triassic Sina  
1330 Formation (Aghdarband Group) in NE-Iran. *Abhandlungen der*  
1331 *Geologischen Bundesanstalt* 38: 195–200. 1369
- 1332 Kristan-Tollmann E, Hamedani A. 1973. Eine spezifische Mikro-  
1333 faunen-Vergesellschaftung aus den Opponitzer Schichten der  
1334 Oberkam der niederösterreichischen Kalkvoralpen. *Neues Jahr-*  
1335 *buch fuer Geologie und Palaeontologie* 143(2): 193–222. 1370
- 1336 Kristan-Tollmann E, Tollmann A, Hamedani A. 1979. Beiträge zur  
1337 Kenntnis der Trias von Persien. I. Revision der Triasgliederung,  
1338 Rhäfazies im Raum von Isfahan und Kössener Fazieseinschlag  
1339 bei Waliabad SE Abadeh. *Mitteilungen der Gesellschaft*  
1340 *der Geologie und Bergbaustudenten in Osterreich* 70:  
1341 119–186. 1371
- 1342 Latreille PA. 1806. Genera crustaceorum et insectorum secundum  
1343 ordinem naturalem in familias disposita, iconibus exemplisque  
1344 plurimis explicata. Parisiis et Argentorati, 303 p. 1372
- 1345 Lentini F. 1974. Caratteri stratigrafici e strutturali della zona di Monte  
1346 Judica (Sicilia Orientale). *Rivista Mineraria Siciliana* 145-147:  
1347 71–96. 1373
- 1348 Lentini F, Grasso M, Carbone S. 1987. *Introduzione alla geologia*  
1349 *della Sicilia e guida all'escursione*. *Convegno della Società*  
1350 *Geologica Italiana su sistemi avanfossa-avanpaese lungo la*  
*Catena Appenninico-Maghrebide, 22–25 aprile*, Naxos-Pergusa,  
Catania, 60 p. 1374
- Lethiers F, Raymond D. 1991. Les crises du Dévonien supérieur par  
l'étude des faunes d'ostracodes dans leur cadre paléogéographique.  
*Palaeogeography, Palaeoclimatology, Palaeoecology* 88: 133–  
146. 1375
- Lieberman HM. 1979. Die Bivalven- und Ostracodenfauna von Raibl  
und ihr stratigraphischer. *Verhandlungen der Geologischen*  
*Bundesanstalt* 2: 85–131. 1376
- Lord AR. 1982. Metacopine ostracods in the Lower Jurassic. In:  
Bannar ET, Lord AR, eds. *Aspects of Micropalaeontology*. London:  
Allen & Unwin, pp. 262–277. 1377
- Lucas SG. 2010. The Triassic timescale: an introduction. In: Lucas  
SG, ed. *The Triassic Timescale*, 334. Geological Society, London,  
Special Publications, pp. 1–16. 1378
- Maddocks RF. 1969. Revision of Recent Bairdiidae. *Bulletin of the*  
*United States National Museum* 295: 1–126. 1379
- Mandelstam NI, Schneider GF, Kuznetsova ZV, Kats FI. 1957. [New  
genera of Ostracoda in the families Cypridae and Cytheridae]. All  
–Union Palaeontological Society, Annual. *Palentologičesko-*  
*Obščestva Ezhegodnik* 16: 182–183 [in Russian]. 1380
- Maugeri Patané G. 1934. Sopra alcuni coralli del Trias superiore di M.  
Scalpello (Enna). *Relationes Consessuum Societatis Gioieniae*  
*Catinensis Natururarium Scientiarum* 67: 9. 1381
- McCoy F. 1844. A synopsis of the characters of the Carboniferous  
Limestone fossils of Ireland. Dublin: Dublin University Press, 207  
p. 1382
- Méhes G. 1911. Über Trias-Ostrakoden aus dem Bakony. Magyar  
Földrajzi Társaság, Budapest, Palaton-bizottsag, pp. 1–38. 1383
- Mette W, Mohtat-Aghai P. 1999. Ostracods and foraminifera from  
Upper Triassic intrashelf basin deposits in the Northern Calcareous  
Alps. *Geologisch Paläontologische Mitteilungen Innsbruck* (un-  
published manuscript). 1384
- Mette W, Honigstein A, Crasquin S. 2014. Deep-water ostracods from  
the Middle Anisian (Reifling Formation) of the Northern  
Calcareous Alps (Austria). *Journal of Micropalaeontology* 34:  
71–91. 1385
- Michelsen O. 1975. Lower Jurassic biostratigraphy and ostracods of  
the Danish Embayment. *Danmarks Geologiske Undersoegelse*,  
Raekke 2(104): 1–287. 1386
- Monostori M. 1994. Ostracod evidence of the Carnian salinity crisis  
in the Balaton Highland, Hungary. *Neues Jahrbuch für Geologie*  
*und Palaeontologie* 193: 311–331. 1387
- Monostori M. 1995. Environmental significance of the Anisian  
Ostracoda fauna from the Forras Hill near Felsoors (Balaton  
Highland, Transdanubia, Hungary). *Acta Geologica Hungarica* 39:  
37–56. 1388
- Monostori M, Tóth E. 2013. Ladinian (Middle Triassic) silicified  
ostracod faunas from the Balaton Highland (Hungary). *Rivista*  
*Italiana di Paleontologia e Stratigrafia* 119: 303–323. 1389
- Monostori M, Tóth E. 2014. Additional Middle to Upper Triassic  
ostracod faunas from the boreholes of Transdanubian Central  
Range (Hungary). *Hantkeniana* 9: 21–43. 1390
- Montanari L. 1987. Lineamenti stratigrafico-paleogeografici della  
Sicilia durante il Ciclo Alpino. *Memorie della Società Geologica*  
*Italiana* 38: 361–406. 1391
- Moore RC. 1961. Treatise of Invertebrate Paleontology. Part Q.  
Arthropoda 3, Crustacea, Ostracoda. Geological Society of  
America and University of Kansas Press, 442 p. 1392
- Müller GW. 1894. Die Ostracoden des Golfes von Neapel und der  
angrenzenden Meere Abschnilte. *Fauna und Flora des golfes von*  
*Neapel* 21: 1–404. 1393

- 1413 Müller GW. 1906. Die Ostracoden der Siboga-Expedition. In: Max  
1414 W, ed. *Siboga-Expedition, Uitkomsten op Zoologisch, Botanisch,*  
1415 *Oceanographisch en H.M. Siboga.* Monographien 30, EJ Brill,  
1416 Leiden, 40 p.
- 1417 Nelli B. 1899a. Il Raibliano del Monte Judica nella provincia di  
1418 Catania. *Atti della Reale Accademia Nazionale dei Lincei,*  
1419 *Rendiconti della Classe di Scienze fisiche, matematiche e naturali*  
1420 *s. 5, 8(1): 91–92.*
- 1421 Nelli B. 1899b. Il Raibl nei dintorni di M. Judica. *Bollettino della*  
1422 *Società Geologica Italiana* 18(2): 211–223.
- 1423 Oertli HJ. 1971. The aspect of ostracode faunas – A possible new tool  
1424 in petroleum sedimentology. *Bulletin du Centre de Recherches de*  
1425 *Pau* 5(suppl.): 137–151.
- 1426 Özdikmen H. 2010. Substitute names for three genera of Ostracoda  
1427 (Crustacea). *Munis Entomology and Zoology* 5: 315–316.
- 1428 Pokorný V. 1964. Some paleoecological problems in marine  
1429 ostracode faunas, demonstrated on the Upper Cretaceous ostracodes  
1430 of Bohemia, Czechoslovakia. *Pubblicazioni della Stazione*  
1431 *Zoologica di Napoli* 33(Suppl.): 462–479.
- 1432 Reuss AE. 1869. Zur fossilen Fauna der Oligocänschichten von Gaas.  
1433 Sitzungsberichte der österreichischen. *Akademie der Wissenschaften,*  
1434 *mathematisch-naturwissenschaftliche Klasse* Abt.1. 59(3):  
1435 446–488.
- 1436 Salaj J, Jendrekova O. 1984. Ecology and facial relation of some  
1437 groups of Triassic foraminifers and ostracods of stratigraphic  
1438 importance. *Geologický Zborník-Geologica Carpathica* 35(2):  
1439 231–240.
- 1440 Sars GO. 1866. Oversight af Norges marine Ostracoder. *Forhand-*  
1441 *linger i Videnskabs-Selskabet I Christiania* 1865, pp. 1–130.
- 1442 Scalia S. 1907a. I fossili del Trias superiore dell'Acquanova e di  
1443 Paraspora. *Bollettino dell'Accademia Gioenia di Scienze Naturali*  
1444 *in Catania* s. 2(1-2): 10–12.
- 1445 Scalia S. 1907b. La fauna del Trias superiore del gruppo montuoso di  
1446 Judica. *Bollettino dell'Accademia Gioenia di Scienze Naturali in*  
1447 *Catania* s. 2(3-4): 17–25.
- 1448 Scalia S. 1909. Il gruppo di M. Judica. *Bollettino della Società*  
1449 *Geologica Italiana* 28: 269–340.
- Scalia S. 1910–1914. La fauna del Trias superiore del gruppo del M.  
Judica: Parti 1–3. *Atti dell'Accademia Gioenia di Scienze Naturali*  
*in Catania* s. 5 (3-5-7): 51+58+25.
- Schmidt di Friedberg P, Trovò A. 1962. Contribution à l'étude  
structurale du groupe du Monte Judica (Sicile orientale). *Bulletin*  
*de la Société Géologique de France* 4(5): 754–759.
- Sebe O, Crasquin S, Grădinaru E. 2013. Early and Middle Anisian  
(Triassic) deep-water ostracods (Crustacea) from North Dobrogea  
(Romania). *Revue de Paléobiologie* 32: 509–529.
- Sohn IG. 1968. Triassic Ostracodes from Makhtesh Ramon, Israel.  
*Bulletin of the Geological Survey of Israel* 44: 1–71.
- Speranza F, Minelli L. 2014. Ultrathick Triassic dolomites control the  
rupture behavior of the central Apennine seismicity: Evidence from  
magnetic modeling of the L'Aquila fault zone. *Journal of*  
*Geophysical Research: Solid Earth* 119: 1–15.
- Styk O. 1958. Triassic microfauna in the neighbourhood of Chrzanow  
and the northwestern part of the Mesozoic periphery of the Swiety  
Krzysz Mts. *Bulletin Poland Instytut Geologiczny* 121, *Micro-*  
*palaeontological researches in Poland* 3: 163–176.
- Sylvester-Bradley PC. 1961. Suborder Metacopina Sylvester-Brad-  
ley, n. suborder. Q358–359. In: Moore RC, ed. *Treatise of*  
*invertebrate paleontology. Part Q, Arthropoda 3, Crustacea,*  
*Ostracoda.* Lawrence, KS: The Geological Society of America,  
Boulder, CO and University of Kansas Press, 442 p.
- Széles M. 1965. Ostracodák a Bakonyi Nosztori-Volgy felsokarni  
rétegeibol. *Földtani közlöny* 45: 412–417.
- Tollmann A. 1976. Analyse des klasschen nordalpinen Mesozoikums.  
*Monographie des Nördlichen Kalkalpen* 2: 1–580.
- Triebel E. 1941. Zur Morphologie und Okologie der fossilen  
Ostracoden. Mit Beschreibung einiger neuer Gattungen und Arten.  
*Senckenbergiana* 23: 294–400.
- Ulrichs M. 1970. Variability of some Ostracods from the Cassian beds  
(Alpine Triassic) depending on the ecology. *Bulletin du Centre*  
*Recherches de Pau – SNPA* 5: 695–715.
- Zorn I. 2010. Ostracodal Type Specimens Stored in the Paleontologi-  
cal Collection of the Geological Survey of Austria. *Jahrbuch der*  
*Geologischen Bundesanstalt* 150: 263–299.

**Cite this article as:** Crasquin S, Sciuto F, Reitano A, Coco RM. 2020. Late Triassic (Tuvanian – Carnian, *Tropites subbullatus/Anatropites spinosus* zones) ostracods from Monte Gambanera (Castel di Iudica, Central-Eastern Sicily, Italy), *BSGF - Earth Sciences Bulletin* Vol: 190069.