



**HAL**  
open science

# A glimpse at deep brain stimulation mechanisms using subthalamic nucleus optogenetic manipulations

Alix Tiran-Cappello, Yann Pelloux, Cécile Brocard, Mickaël Degoulet,  
Christelle Baunez

► **To cite this version:**

Alix Tiran-Cappello, Yann Pelloux, Cécile Brocard, Mickaël Degoulet, Christelle Baunez. A glimpse at deep brain stimulation mechanisms using subthalamic nucleus optogenetic manipulations. 2020. hal-02990849

**HAL Id: hal-02990849**

**<https://hal.science/hal-02990849v1>**

Preprint submitted on 4 Jan 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Title Page

**TITLE: A glimpse at deep brain stimulation mechanisms using subthalamic nucleus optogenetic manipulations**

**Authors:**

Alix Tiran-Cappello<sup>1</sup>, Yann Pelloux<sup>1†</sup>, Cécile Brocard<sup>1</sup>, Mickaël Degoulet<sup>1</sup>, Christelle Baunez<sup>1</sup>

**Affiliations :**

Institut de Neurosciences de la Timone, UMR7289 CNRS & Aix-Marseille Université, 27 Boulevard Jean Moulin - 13005 Marseille, France

† Present address: Department of Neuroscience and Physiology, Neuroscience Institute, New York University Medical Center, New York, NY 10016, USA

**Corresponding author:**

Christelle Baunez, Institut de Neurosciences de la Timone, UMR7289 CNRS & Aix-Marseille Université, 27 Boulevard Jean Moulin - 13005 Marseille, France

Phone: +33 491324062, email : [christelle.baunez@univ-amu.fr](mailto:christelle.baunez@univ-amu.fr)

**Short/running**

Probing subthalamic DBS on motivation with optogenetics

**title:**

**Keywords:** basal ganglia, high frequency stimulation, inactivation, food, cocaine, motivation.

**Number of words in the abstract: 212**

**Number of words in the main text: 4140**

**Number of figures: 5**

**Number of tables: 0**

**Supplemental information: 5**

## Abstract

Although deep brain stimulation (DBS) is now a widely used therapeutic strategy, its precise mechanism remains largely unclear. Since this approach is progressively extended to treat non-motor disorders such as depression, obsessive-compulsive disorders, the comprehension of its effects on motivated behaviors appears of the utmost importance for a possible application for addiction. In intact rats, we used inhibition and high frequency optogenetic activation of subthalamic nucleus (STN) neurons to test whether or not we could reproduce the effects of electric deep brain stimulation on rats' motivation for sweet food and cocaine. Rats' motivation was assessed using fixed-ratio 5 and progressive ratio schedules of reinforcement for both rewards and illumination was applied during behavioral testing. Efficiency of optogenetic manipulations has been validated using in-vitro electrophysiological recordings. Optogenetic inhibition of STN increased motivation for food and reduced motivation for cocaine. In contrast, optogenetic high frequency stimulation reduced the motivation for food without impacting motivation for cocaine. Optical inhibition mimics the effect of electric deep brain stimulation on food and cocaine motivation, confirming that the effects observed under electric DBS result from a specific inactivation of the STN. In contrast, optogenetic high frequency stimulation induces opposite effects to those of electric one, suggesting a stimulation of the STN that only seems to affect food motivation.

## 1 Main article text

### 2 Introduction

3           The subthalamic nucleus (STN) is a small homogenous glutamatergic structure of the  
4 basal ganglia. Historically known as a relay for motor information, STN is a well-  
5 established target for deep brain stimulation (DBS) treatment for motor symptoms of  
6 Parkinson's disease (PD) (Krack et al., 2010). STN functions have also been extended to  
7 decision-making, behavioral inhibition, reward encoding and control of motivation (Baunez  
8 et al., 2011; Hamani et al., 2017).

9           In rodents, the exploration of STN motivational functions has brought up interesting  
10 findings, such as the dissociative effect showing that both STN lesion and DBS increase the  
11 motivation for sweet food, while decreasing the motivation for cocaine (Baunez et al., 2005;  
12 Rouaud et al., 2010). Similarly STN pharmacological inactivation is able to reduce cocaine  
13 reinstatement (Bentzley and Aston-Jones, 2017). Finally, STN lesion and DBS have since  
14 been shown to prevent cocaine escalation and reescalation of cocaine or heroin intake  
15 (Pelloux et al., 2018; Wade et al., 2017).

16           In human, manipulations of the STN by DBS in PD patients have confirmed its  
17 involvement in motivated behavior such as hypersexuality (Doshi and Bhargava, 2008;  
18 Romito et al., 2002), or reduced dopaminergic dysregulation traits (Eusebio et al., 2013).  
19 STN DBS has progressively become a therapeutic strategy for the treatment of obsessive  
20 compulsive disorder (Mallet et al., 2008, 2002) and its potential to treat drug abuse disorder  
21 has been suggested (Pelloux and Baunez, 2013) and is being investigated. Despite its  
22 extensive use in these pathologies, the exact mechanism of STN DBS remains a matter of  
23 debate. For instance while being functionally similar to an inactivation in many cases,

24 electric STN DBS locally inhibits STN cell bodies (Benazzouz et al., 2000; Beurrier et al.,  
25 2001; Meissner et al., 2005; Tai et al., 2003; Welter et al., 2004), while increasing activity of  
26 STN output structures in some studies (Hashimoto et al., 2003; Maurice et al., 2003; Shehab  
27 et al., 2014), but has also been reported to inhibit the neurons of the substantia nigra  
28 reticulata (Benazzouz et al., 2000; Tai et al., 2003). It also impacts the activity of afferent  
29 structures and namely alters cortical activity through antidromic activation of the hyperdirect  
30 pathway (Li et al., 2012; McIntyre et al., 2004).

31 More recently, optogenetic studies have been developed and could allow a more  
32 specific way to alter the functioning of STN, while sparing the passing fibers and  
33 neighboring structures. Using this technique, it was shown that STN DBS effects in rodent  
34 PD models are highly dependent on the hyperdirect pathway (Gradinaru et al., 2009;  
35 Sanders and Jaeger, 2016). It was later demonstrated that optogenetic inhibition of the STN  
36 was sufficient to alleviate PD symptoms in rodents (Yoon et al., 2016, 2014), thus  
37 confirming historical pharmacological inactivation and lesion studies that had shown an  
38 improvement of motor functions in PD models after STN inhibition (Baunez et al., 1995;  
39 Bergman et al., 1990; Levy et al., 2001). To date, optogenetic experiments have been used to  
40 dissect circuits involved in the motor effects of STN DBS, however motivational aspects  
41 remain largely unexplored, even though a recent study used optogenetic control to  
42 demonstrate the causal role of the STN in pausing/interrupting initiated behavior (Fife et al.,  
43 2017). In the present work we used STN optogenetic modulation to compare its effects to  
44 those previously described of electric DBS on motivated behavior in rats (Rouaud et al.,  
45 2010) and then discuss the mechanisms involved. Thus the results presented here underline  
46 the specific contribution of STN neurons to DBS mechanisms.

47

## 48 Results

### 49 Motivation for sweet food

50 It has been previously shown that STN electric DBS increases the motivation for  
51 sweet food, while decreasing the motivation for cocaine (Rouaud et al., 2010). To test  
52 whether STN optogenetic inhibition could affect food-motivated behavior; we injected rats  
53 bilaterally with either ARCHT3.0 (inhibition, n=9) or EYFP (Control, n=7) expressed under  
54 the control of promoter CamKIIa (Fig. 1A) and subjected to fixed ratio 5 (FR5) and  
55 progressive ratio (PR) schedules of reinforcement. Due to relatively long experimental  
56 sessions, the light was applied intermittently by 5min of laser modulation in alternation with  
57 5 min OFF periods to reduce the risks of tissue damage (Fig. 1B). Under FR5 schedule of  
58 reinforcement, STN inactivation applied during eight consecutive sessions did not  
59 significantly change the number of food pellets obtained by the rats, although a trend  
60 towards an increase was observable at the end of the experiment and during the OFF period  
61 (Fig. 1C). Under PR schedule of reinforcement, the laser inducing inhibition transiently  
62 increased the motivation to work for sweet food reward in comparison with the level of  
63 motivation exhibited by the EYFP control group as shown by both the number of rewards  
64 obtained and the breakpoint reached (Fig. 1D). At the end of the OFF period, we performed  
65 an extra day of testing during which the inhibition induced by the laser modulation increased  
66 the breakpoint when compared to that of the EYFP control group (Fig. 1D). This also  
67 confirmed that the opsins were still functional at this stage of the experiment. Finally a  
68 detailed analysis over the first day of laser illumination and the laser challenged revealed an  
69 increased motivation for food in the inhibition group compared to the control group (Fig.  
70 1E).

71 We then tested the effects of high frequency optogenetic stimulation of the STN on  
72 food-motivated behavior, by injecting the fast kinetic channelrhodopsin CHETA-TC (n=7)  
73 or the EYFP (n=5) bilaterally in the STN (Fig. 2A). The light stimulation was applied at  
74 130Hz with a pulse-width of 2ms to mimic the electric DBS parameters in terms of  
75 frequency and using the same intermittent pattern described for inhibition experiment (Fig.  
76 2B). In the FR5 procedure, high frequency optogenetic stimulation reduced the number of  
77 pellets obtained by the rats but only when the power output was changed from 5mW to  
78 10mW (Fig. 2C & 2D). In the PR schedule of reinforcement, the 130Hz laser stimulation  
79 reduced the breakpoint of the CHETA-TC group compared to that reached by the EYFP  
80 control group (Fig. 2E). After a few consecutive sessions, the performance of the stimulated  
81 animals resumed to that of the control group. Interestingly when the stimulation was stopped  
82 we observed transient increase in the breakpoint of the animals that had been previously  
83 stimulated, thus indicating the presence of a rebound effect (Fig. 2E and 2F). We then tested  
84 various parameters to search for the optimal efficacy of the light stimulation. Variations of  
85 the frequencies of stimulation revealed that high frequencies (130Hz) are more efficient at  
86 reducing food-related motivation, in contrast with lower frequencies that did not modify the  
87 behavior (Fig. 2G). Finally, we tested the influence of the light pulse duration at 100Hz, by  
88 increasing the duration from 2ms to 3ms. This last manipulation further reduced motivation  
89 for sweet food (Fig. 2H).

90 Since the STN is primarily known for its motor functions, we performed a  
91 locomotion experiment to assess the effect of optogenetic modulation on such behavior.  
92 Laser beam activation increased the locomotor activity of the ‘inhibition’ group while it  
93 remained unaffected in the ‘stimulation’ group (Fig. S1A).

94

## 95 **Motivation for cocaine**

96           Because STN DBS reduces motivation for cocaine (Rouaud et al., 2010), we trained  
97 animals of control (n=5), inhibition (n=5) and stimulation (n=5) groups to nose-poke for  
98 cocaine under FR5 schedule of reinforcement. We applied the same intermittent laser pattern  
99 with 5min ON and OFF bins described previously during the food self-administration  
100 experiment. Optogenetic inhibition of the STN reduced the cocaine intake in ARCHT  
101 animals under FR5 schedule of reinforcement (Fig. 3A). Normalizing the rat intake relative  
102 to the last day of baseline further highlighted this decrease in cocaine intake (Fig. 3B & C).  
103 In contrast, the 130Hz stimulation at 10mW did not affect the cocaine intake (Fig. 3E-G). As  
104 optogenetic inhibition can impact locomotion, we calculated the percentage of active nose-  
105 pokes performed during laser bins, and none of the stimulation or inhibition groups  
106 displayed an altered pattern compared to the control group, thus ruling out the risk of non-  
107 specific effects related to motor activity (Fig. 3D & H). We also tested possible interactions  
108 of cocaine with laser stimulation on locomotor activity, and none of the three groups;  
109 control, inhibition and stimulation displayed any changes following the injection of a 5, 7.5  
110 or 10mg/kg dose of cocaine (Fig. S1B-D).

111           Under a PR schedule of reinforcement, laser inhibition of STN immediately reduced  
112 the number of injections obtained by the inhibition group and the maximum ratio reached by  
113 the rats, thus indicating a lower level of motivation for the cocaine (Fig. 4A & D). This  
114 effect persisted for a few days (Fig. 4B) and resulted in a global lower level of intake over  
115 the 8 days of laser modulation (Fig4. C). High frequency optogenetic stimulation at 130 Hz,  
116 unlike electrical 130Hz stimulation, had no effect on performance (Fig. 4E-D). Finally, we  
117 subjected the rats to acute 20Hz optogenetic stimulation, which had no impact on the



118 motivation to work for cocaine either (Fig. S3). As during FR5 conditions, laser illumination  
119 did not alter the pattern of nose pokes in the PR task (Fig. S4).

120

## 121 **Response of STN neurons to optogenetic modulation**

122 Ultimately, to confirm the effect of our optogenetic manipulations, we performed  
123 patch-clamp recordings on 19 adult rat brains infected with AAV virus to induce the  
124 expression of the opsins in the STN. We then confirmed that the presence of opsins did not  
125 deeply change the properties of the cells, as resting membrane potential, cell capacitance,  
126 membrane resistance, access resistance and firing rates were equivalent between stimulation,  
127 inhibition and control groups (Fig. S5A-E). To confirm the efficacy of the optogenetic  
128 modulation, we first applied discrete light pulses on STN neurons. Longer pulse-width  
129 duration and higher light intensity elicited higher inhibitory currents in ARCHT3.0 positive  
130 cells (Fig. S6A-C) and elicited higher depolarizing currents in CHETA-TC positive cells  
131 (Fig. S6D-F).

132 To further assess the effect of optogenetic inhibition, we electrically stimulated STN  
133 neurons thus forcing them to emit action potentials in a stable manner (10Hz, injected  
134 current: rheobase + 20-130pA) and then applied the laser stimulation. A 15s light-pulse on  
135 STN neurons expressing ARCHT3.0 successfully inhibited neuronal firing (Fig. 5A & B).  
136 We then applied the same light pattern that was used for the behavioral experiments. The  
137 cells were thus subjected to an alternation of 15s illumination followed by 5s of obscurity  
138 during 5min (Fig. 5C). Inhibition induced by the laser reduced the firing rate of STN  
139 neurons and when the laser was stopped the cells resumed to their basal firing rate (Fig. 5C  
140 & D). This confirmed that our manipulations were sufficient to inhibit STN neurons.

141 The effects of optogenetic stimulation were first assessed in STN neurons with 10s of  
142 laser illumination at various frequencies (Fig. 5E). Interestingly, higher frequencies of laser  
143 stimulation did not change the number of induced action potentials compared to lower  
144 frequencies (Fig. 5F). However higher frequencies significantly increased membrane  
145 potential, suggesting this pattern increased the excitability of STN neurons (Fig. 5G). To test  
146 this hypothesis we applied 10Hz electrical stimulation (injected current: Rheobase +0-50pA)  
147 to elicit a few action potentials and we applied 130Hz laser stimulation for 5min. Firing rate  
148 of STN neurons was significantly increased by the 130Hz laser stimulation, and once the  
149 laser was turned off firing rate was no longer different from the baseline period (Fig. 5H &  
150 I). This further enforced that 130Hz optogenetic stimulation is unlikely to induce action  
151 potentials on its own, but seems to increase STN neurons excitability.

152

## 153 Discussion

154 The data presented here show that STN optogenetic manipulations are sufficient to  
155 achieve bidirectional control on food motivation in rats. This positions the STN as a major  
156 contributor in the processes regulating sweet food reinforcement and motivation.  
157 Optogenetic inhibition mimics the effects obtained with either electric DBS or lesion of STN  
158 by increasing motivation for food and decreasing that for cocaine (Baunez et al., 2005;  
159 Rouaud et al., 2010). Optogenetic stimulation applied at 130Hz produces the mirror effect  
160 and reduces food motivation, revealing that optogenetic stimulation although applied at  
161 130Hz results in opposite effects to those of electric 130Hz stimulation. Interestingly, STN  
162 optogenetic manipulations have transitory effect on food and cocaine intake, while electric  
163 STN DBS produces long lasting effects (Rouaud et al., 2010). This indicates that the  
164 surrounding network and afferences strongly contribute to maintain these effects, in line

165 with previous studies showing that the hyperdirect pathway is critical in DBS treatment of  
166 some PD symptoms (Gradinaru et al., 2009; Li et al., 2012).

167 Because of our long lasting behavioral experiments, we used an intermittent laser  
168 modulation to lower the risk of tissue damage. Similar intensities have been used for  
169 inhibition and stimulation without any adverse consequences (Fife et al., 2017; Nieh et al.,  
170 2015). We did not observe any traces after histology that could be related to light induced  
171 damages, controls expressing EYFP were not impacted by the laser beam and animals of  
172 ‘inhibition’ and ‘stimulation’ groups were still responsive to light modulation even after  
173 several sessions, and the effect of optogenetic manipulation dissipated more or less quickly  
174 as the manipulation ceased, thus ruling out the risk of permanent alteration. Such  
175 intermittent pattern could hold a huge potential to achieve optogenetic control during  
176 prolonged experiments for future application.

177 We tested the influence of various parameters for optogenetic stimulation in the food  
178 experiment. Behavior was moderately impacted by the stimulation at 5mW, and significantly  
179 reduced by the stimulation with light intensity set at 10mW. This is explained by higher  
180 depolarizing currents in STN neurons (Fig. S6F) but also because light is transmitted  
181 through a larger volume of tissue and can stimulate a larger pool of neurons (Yizhar et al.,  
182 2011).

183 Our electrophysiological experiments show that high frequency optogenetic  
184 stimulation increases the activity of STN neurons at cell body level. Even if optogenetic  
185 stimulation of cell bodies can lead to a different pattern once the excitation wave reaches the  
186 axon (Yizhar et al., 2011), our results advocate in favor of an effective stimulation of STN  
187 output, as it produced opposite effects to those observed after optogenetic inhibition on  
188 motivation for food. Of note, optogenetic stimulation at high frequencies of glutamatergic

189 terminals has been used in vitro and in-vivo to induce LTD (Klavir et al., 2017), thus a  
190 similar effect could be accounted for the progressive loss of efficiency of the laser  
191 stimulation and for the rebound effect we observed during the food self-administration after  
192 several sessions of 130Hz optogenetic stimulation. In addition, our stimulation resulted in an  
193 increased activity of STN without locking it to a define rhythm, thus addressing one of the  
194 issues often raised by critics regarding optogenetic experiments, that is to determine if the  
195 activity resulting from the stimulation pattern possesses any physiological relevance  
196 (Deisseroth, 2015; Yizhar et al., 2011).

197         We reported increased locomotor activity consecutive to STN optogenetic inhibition  
198 (Fig. S1A). If this increased locomotor activity could explain the observed operant behavior  
199 for food reward, it is interesting to note that the laser inhibition did not change the number  
200 of responses in the inactive hole (Fig. S2C-D). Only the number of rewards and responses in  
201 the food magazine were increased by the laser inhibition (Fig. S2A-B). This suggests that  
202 even though the global activity of the animals is altered by the laser inhibition, it does not  
203 happen in an unspecific manner but remains goal-directed. Despite the fact that the first  
204 study of STN optogenetic modulation did not report any changes in locomotor activity in  
205 parkinsonian models, probably as a result of lower opsin efficiency (Gradinaru et al., 2009),  
206 more recent studies have shown that STN optogenetic inhibition could increase locomotion  
207 in PD rodent models (Yoon et al., 2016, 2014). It was also established that reducing STN  
208 glutamate transporter expression increases locomotor activity (Schweizer et al., 2014),  
209 although electric STN DBS has no effect on locomotor activity in naïve rats (Pelloux et al.,  
210 2018). Our results further extend these observations in naïve rats, and indicate that direct  
211 optogenetic inhibition of the STN is sufficient to increase locomotion. This confirms that  
212 specific inhibition of the STN leads to locomotor effect while the additional stimulation of

213 the afferences, efferences and neighboring structures by electric DBS can counteract this  
214 effect.

215 Finally, opsin expression was mainly observable in the limbic and associative  
216 territories of the STN, yet optogenetic modulation affects the motor activity, this further  
217 enforces the idea that in rat territories are highly intertwined and the STN represents a site of  
218 convergence for numerous cortical projections, as suggested by former studies (Janssen et  
219 al., 2017; Kita et al., 2014).

220 Although STN stimulation does not alter the motivation to work for cocaine nor the  
221 locomotor activity, STN inhibition impacts both of them, in line with previous studies  
222 (Baunez et al., 2005; Rouaud et al., 2010). Two elements can explain this finding. First, our  
223 optogenetic stimulation seems to increase STN excitability, and that increase might be  
224 insufficient to affect cocaine related motivation and locomotion. Second, the STN seems  
225 highly sensitive to specific ranges of oscillations. For instance in the context of PD, DBS at  
226 low frequency ( $< 20\text{Hz}$ ) has been reported to worsen the symptoms, while high frequency  
227 DBS exerts therapeutic effects (Eusebio et al., 2008; Timmermann et al., 2004). Very  
228 similarly, in the STN, very low frequencies (1-4Hz) have been closely tied to the severity of  
229 obsessive compulsive disorders (Welter et al., 2011). Finally, recent data from our group  
230 indicate that low frequency oscillations (Theta 8-12Hz and Beta 13-30Hz) develop in the  
231 STN during extended cocaine use, and that blocking these oscillations with DBS or lesion  
232 prevents escalation and re-escalation of cocaine intake (Pelloux et al., 2018). Thus, the key  
233 point to affect cocaine motivation and motor activity through STN stimulation probably lies  
234 within very specific ranges of oscillations or a specific driving rhythm rather than a general  
235 increase of activity.

236 Overall the present work emphasizes some of the specific contributions of STN in  
237 reward related processes: food and cocaine self-administration, as well as locomotor  
238 behavior. By providing data on three different behaviors that can be deeply impacted by  
239 DBS, we hope to offer a large comparative framework for the study of DBS effects. Since  
240 DBS is progressively applied for an increasing number of psychiatric pathologies such as  
241 depression, obsessive-compulsive disorders and addiction (Hamani et al., 2017; Mallet et al.,  
242 2008; Pelloux and Baunez, 2013), identifying the contribution of the structure itself and that  
243 of the networks supporting therapeutic effects appears critical to reduce the risks of side  
244 effects and refine the parameters and strategies used to target brain structures.

245

## 246 **Material and Methods**

### 247 **Animals**

248 Lister hooded male rats, 380-400g, were used in this study. They were housed in  
249 pairs; maintained in an inverted 12h dark/light cycle (light onset at 7 p.m.) in temperature-  
250 controlled room, with unlimited access to water. All experiments were carried out during the  
251 dark phase. For the food-rewarded experiments, rats were kept under moderate food  
252 restriction (15-16g/rat, 75-80% of normal daily intake). For the other experiments animals  
253 had no food restriction. All animal care and use conformed to the French regulation (Decree  
254 2010-118) and were approved by the local ethic committee and the University of Aix-  
255 Marseille (saisine #3129.01).

## 256 Viral vectors

257 We used AAV5 (UNC Vector core, Chapel Hill, USA) to transfect STN neurons,  
258 allowing the expression of recombinant protein under the CamKIIa promoter with the  
259 following constructs: AAV5-CaMKII-hChR2 (E123T/T159C)-p2A-EYFP-WPRE (CHETA-  
260 TC) for stimulation groups, AAV5-CaMKII-ArchT3.0-p2A-EYFP-WPRE for inhibition  
261 groups and AAV5-CaMKII-EYFP for the control groups.

262

## 263 Surgery

264 Animals were anesthetized with a combination of Ketamine (Imalgen, Merial,  
265 100mg/kg, s.c.) and medetomidine (Domitor, Janssen, 30mg/kg, s.c.).

266 *Optogenetic surgery:* Animals were placed in a Kopf stereotaxic apparatus to receive a  
267 bilateral 0.5  $\mu$ L injection of virus (0.16 $\mu$ L/min) using injectors placed at the following  
268 coordinates relative to bregma: AP: -3.7 mm; ML:  $\pm$ 2.4 mm; DV: -8.4 mm (from skull)  
269 according to the Paxinos and Watson atlas (Paxinos and Watson, 2005). Optic fibers were  
270 implanted 0.5mm above each injection site and secured within a head-cap made of dental  
271 cement on the skull.

272 *Catheter surgery:* A homemade silicone catheter was inserted 2.5-3.0cm in the right jugular  
273 vein and secured with ligatures and exited dorsally between the scapulae through a guide  
274 cannula secured with dental cement. Blood reflux in the tubing was checked to confirm  
275 correct placement of the catheter. Catheters were also daily flushed with heparin (Sanofi,  
276 3g/L) and antibiotic enrofloxacin (Baytril, Bayer, 8g/L) in 0.9% saline to prevent blood

277 clots and infection during the recovery period, and were also flushed before and after each  
278 self-administration session until the end of experiments.

279 After surgery, all rats were awakened with an injection of atipamezol (Antisedan, Janssen,  
280 0.15mg/kg, *i.m.*). Preventive long-acting amoxicillin treatment was applied (Duphamox LA,  
281 Pfizer, 100mg/kg, *s.c.*). They were allowed to recover for at least 7 days, during which they  
282 were daily monitored. To allow for satisfying level of expression of the opsins, a waiting  
283 period of 3 weeks was applied before any light delivery.

284

### 285 [In-vivo light delivery](#)

286         Implants were built with 230  $\mu\text{m}$  optic fibers (NA 0.22, Thorlabs) glued to 2.5mm  
287 ceramic ferrules (Thorlabs) using epoxy. Ferrules were slightly grinded with a Dremel to  
288 improve the contact with the dental cement. Light stimulation was performed during  
289 experiments using 200mW 532nm DPSS laser controlled with a signal generator. Optic  
290 rotary joints (DoricLenses) relieved the torsion due to animal movements and bilateral  
291 stimulation was achieved using an optic coupler (Thorlabs, FCMM625-50A). Light output  
292 was assessed before each experiment with a power meter (Thorlabs, PM20A) and  
293 parameters for stimulation were: 10mW, 2ms pulse, 130Hz; and for inhibition: 5mW, 15s  
294 pulse, 5s interval between pulses (resulting in a 0.2Hz pattern), unless specified otherwise.

### 295 [Food self-administration](#)

296         Rats were first trained during 10-15 days to perform the fixed-ratio 1 for sucrose  
297 pellets in standard rat operant chambers (MedAssociates) with an active hole paired with the  
298 delivery of the reward and illumination of an associated cue light and an inactive hole,



299 whose activation had no programmed consequences. The ratio was progressively increased  
300 until fixed ratio 5 (FR5), stability of performance was evaluated during 3 days before  
301 incrementing to the next ratio. Sessions of FR5 lasted 30 min. In the case of progressive  
302 ratio (PR) experiments, the ratios followed an arithmetically increasing schedule in steps of  
303 five, with three repetitions of each step (i.e., 1, 1, 1, 5, 5, 5, 10, 10, 10...). The sessions  
304 ended after 90min or after 5min without any response in the active hole and the last ratio  
305 completed was defined as the animal ‘breakpoint’.

### 306 Cocaine Self-Administration

307 The intravenous cocaine self-administration was conducted in the same operant  
308 boxes described above; the stainless steel guide cannula of the catheter was connected  
309 through steel-protected Tygon tubing to a swivel (Plastics One) and then a syringe filled  
310 with cocaine placed on an infusion pump (MedAssociates). Self-administration experiments  
311 used a dose of 250 $\mu$ g/90 $\mu$ L per cocaine injection (Coopérative pharmaceutique française).  
312 The daily sessions lasted for 2 h for FR5 or maximum 4 h for progressive ratio. In PR  
313 sessions, to limit the number of injections, the ratios for cocaine reward followed the  
314 modified equation of Roberts 1, 3, 6, 10, 15, 20, 25, 32, 40, 50, 62, 77, 95, 118, 145, 178,  
315 219, etc. (Depoortere et al., 1993). If the rats failed to complete a ratio within 60 min, the  
316 session ended.

### 317 Spontaneous- and cocaine-induced locomotor activity

318 Locomotor activity was measured as the distance traveled (in m) in a circular  
319 homemade Perspex open-field (60 cm diameter) with a video tracking performed with  
320 Bonsai (Open Ephys). Details of the procedure can be found in the supplementary material.

## 321 Electrophysiology

322 Full details regarding solutions and preparation of adult brain slices can be found in  
323 the supplementary materials and were adapted from other studies (McDevitt et al., 2014;  
324 Ting et al., 2014). Adult rats were anesthetized and perfused with ice-cold artificial  
325 cerebrospinal fluid. 200  $\mu\text{m}$  coronal slices containing the STN were prepared with a  
326 vibratome (1200S, Leica). Cells were visualized on an upright microscope with infrared  
327 differential interference contrast and fluorescence microscopy (Olympus, BX51WI).  
328 Recordings were made with pClamp 10.3 software using a MultiClamp 700B amplifier set  
329 with 4 kHz low-pass Bessel filter and 10 kHz digitization (Molecular Devices, Sunnydale,  
330 CA). Laser modulation was applied using a 532nm DPSS laser. The light beam was directed  
331 at the preparation using an optic fiber submerged into the bath. Laser intensities were  
332 adjusted to match those used in behavioral experiments.

333 To assess the influence of laser modulation on firing pattern, cells were imposed a  
334 given rhythm through current injection. Cells expressing inhibitory opsins ARCHT3.0 were  
335 driven at 10Hz, with injected current intensity corresponding to rheobase + 20-130pA in  
336 order to induce stable and robust firing pattern. Cells expressing excitatory opsins CHETA-  
337 TC were driven at 10Hz, intensity set at +0-50pA to elicit a few action potentials in a stable  
338 pattern. Stability of the excitation was assessed through a 2min baseline recording, then we  
339 applied the same laser pattern used for behavioral experiments for 5min and the cells were  
340 monitored for another 5min with the laser turned OFF.

341

342 Extended details for histological controls and statistic analyses can be found in the  
343 supplementary materials.

344

## 345 Acknowledgments

346 The authors thank Drs. F Brocard and R Bos for allowing use of their Patch clamp  
347 equipment and animal facility personal for technical support.

## 348 Funding

349 This work was funded by CNRS, Aix-Marseille Université; the MILDECA (Mission  
350 Interministérielle pour la Lutte contre les drogues et les conduits addictives; Programme  
351 Apprentis Chercheurs MAAD), the Fondation pour la Recherche Médicale  
352 (FRMDPA20140629789), the support of the A\*MIDEX project (ANR-11-IDEX-0001-02)  
353 funded by the « Investissements d’Avenir » French Government program, managed by the  
354 French National Research Agency (ANR). ATC was funded by the French Ministry of  
355 Higher Education and Research and by France Parkinson.

356

## 357 Disclosures

358 Drs. Baunez, Pelloux and Degoulet reported no biomedical financial interests or potential  
359 conflicts of interest. Alix Tiran-Cappello and Cécile Brocard reported no biomedical  
360 financial interests or potential conflicts of interest.

361

## References


- Baunez C, Dias C, Cador M, Amalric M. 2005. The subthalamic nucleus exerts opposite control on cocaine and “natural” rewards. *Nat Neurosci* **8**:484–489. doi:10.1038/nm1429
- Baunez C, Nieoullon A, Amalric M. 1995. In a rat model of parkinsonism, lesions of the subthalamic nucleus reverse increases of reaction time but induce a dramatic premature responding deficit. *J Neurosci Off J Soc Neurosci* **15**:6531–6541.
- Baunez C, Yelnik J, Mallet L. 2011. Six questions on the subthalamic nucleus: lessons from animal models and from stimulated patients. *Neuroscience, Function and Dysfunction of the Basal Ganglia* **198**:193–204. doi:10.1016/j.neuroscience.2011.09.059
- Benazzouz A, Gao DM, Ni ZG, Piallat B, Bouali-Benazzouz R, Benabid AL. 2000. Effect of high-frequency stimulation of the subthalamic nucleus on the neuronal activities of the substantia nigra pars reticulata and ventrolateral nucleus of the thalamus in the rat. *Neuroscience* **99**:289–295. doi:10.1016/S0306-4522(00)00199-8
- Bentzley BS, Aston-Jones G. 2017. Inhibiting subthalamic nucleus decreases cocaine demand and relapse: therapeutic potential. *Addict Biol* **22**:946–957. doi:10.1111/adb.12380
- Bergman H, Wichmann T, DeLong MR. 1990. Reversal of experimental parkinsonism by lesions of the subthalamic nucleus. *Science* **249**:1436–1438.
- Beurrier C, Bioulac B, Audin J, Hammond C. 2001. High-Frequency Stimulation Produces a Transient Blockade of Voltage-Gated Currents in Subthalamic Neurons. *J Neurophysiol* **85**:1351–1356.
- Deisseroth K. 2015. Optogenetics: 10 years of microbial opsins in neuroscience. *Nat Neurosci* **18**:1213–1225. doi:10.1038/nm.4091
- Depoortere RY, Li DH, Lane JD, Emmett-Oglesby MW. 1993. Parameters of self-administration of cocaine in rats under a progressive-ratio schedule. *Pharmacol Biochem Behav* **45**:539–548.
- Doshi P, Bhargava P. 2008. Hypersexuality following subthalamic nucleus stimulation for Parkinson’s disease. *Neurol India* **56**:474–476.
- Eusebio A, Chen CC, Lu CS, Lee ST, Tsai CH, Limousin P, Hariz M, Brown P. 2008. Effects of low-frequency stimulation of the subthalamic nucleus on movement in Parkinson’s disease. *Exp Neurol* **209**:125–130. doi:10.1016/j.expneurol.2007.09.007
- Eusebio A, Witjas T, Cohen J, Fluchère F, Jouve E, Régis J, Azulay J-P. 2013. Subthalamic nucleus stimulation and compulsive use of dopaminergic medication in Parkinson’s disease. *J Neurol Neurosurg Psychiatry* **84**:868–874. doi:10.1136/jnnp-2012-302387
- Fife KH, Gutierrez-Reed NA, Zell V, Bailly J, Lewis CM, Aron AR, Hnasko TS. 2017. Causal role for the subthalamic nucleus in interrupting behavior. *eLife* **6**. doi:10.7554/eLife.27689
- Gradinaru V, Mogri M, Thompson KR, Henderson JM, Deisseroth K. 2009. Optical Deconstruction of Parkinsonian Neural Circuitry. *Science* **324**:354–359. doi:10.1126/science.1167093
- Hamani C, Florence G, Heinsen H, Plantinga BR, Temel Y, Uludag K, Alho E, Teixeira MJ, Amaro E, Fonoff ET. 2017. Subthalamic Nucleus Deep Brain Stimulation: Basic Concepts and Novel Perspectives. *eNeuro* **4**:ENEURO.0140-17.2017. doi:10.1523/ENEURO.0140-17.2017

- Hashimoto T, Elder CM, Okun MS, Patrick SK, Vitek JL. 2003. Stimulation of the Subthalamic Nucleus Changes the Firing Pattern of Pallidal Neurons. *J Neurosci* **23**:1916–1923.
- Janssen MLF, Temel Y, Delaville C, Zwartjes DGM, Heida T, Deurwaerdère PD, Visser-Vandewalle V, Benazzouz A. 2017. Cortico-subthalamic inputs from the motor, limbic, and associative areas in normal and dopamine-depleted rats are not fully segregated. *Brain Struct Funct* **222**:2473–2485. doi:10.1007/s00429-016-1351-5
- Kita T, Osten P, Kita H. 2014. Rat subthalamic nucleus and zona incerta share extensively overlapped representations of cortical functional territories. *J Comp Neurol* **522**:4043–4056. doi:10.1002/cne.23655
- Klavir O, Prigge M, Sarel A, Paz R, Yizhar O. 2017. Manipulating fear associations via optogenetic modulation of amygdala inputs to prefrontal cortex. *Nat Neurosci* **20**:836–844. doi:10.1038/nn.4523
- Krack P, Hariz MI, Baunez C, Guridi J, Obeso JA. 2010. Deep brain stimulation: from neurology to psychiatry? *Trends Neurosci* **33**:474–484. doi:10.1016/j.tins.2010.07.002
- Levy R, Lang AE, Dostrovsky JO, Pahapill P, Romas J, Saint-Cyr J, Hutchison WD, Lozano AM. 2001. Lidocaine and muscimol microinjections in subthalamic nucleus reverse Parkinsonian symptoms. *Brain J Neurol* **124**:2105–2118.
- Li Q, Ke Y, Chan DCW, Qian Z-M, Yung KKL, Ko H, Arbuthnott GW, Yung W-H. 2012. Therapeutic Deep Brain Stimulation in Parkinsonian Rats Directly Influences Motor Cortex. *Neuron* **76**:1030–1041. doi:10.1016/j.neuron.2012.09.032
- Mallet L, Mesnage V, Houeto J-L, Pelissolo A, Yelnik J, Behar C, Gargiulo M, Welter M-L, Bonnet A-M, Pillon B, Cornu P, Dormont D, Pidoux B, Allilaire J-F, Agid Y. 2002. Compulsions, Parkinson's disease, and stimulation. *The Lancet* **360**:1302–1304. doi:10.1016/S0140-6736(02)11339-0
- Mallet L, Polosan M, Jaafari N, Baup N, Welter M-L, Fontaine D, du Montcel ST, Yelnik J, Chéreau I, Arbus C, Raoul S, Aouizerate B, Damier P, Chabardès S, Czernecki V, Ardouin C, Krebs M-O, Bardinnet E, Chaynes P, Burbaud P, Cornu P, Derost P, Bougerol T, Bataille B, Mattei V, Dormont D, Devaux B, Vérin M, Houeto J-L, Pollak P, Benabid A-L, Agid Y, Krack P, Millet B, Pelissolo A, STOC Study Group. 2008. Subthalamic nucleus stimulation in severe obsessive-compulsive disorder. *N Engl J Med* **359**:2121–2134. doi:10.1056/NEJMoa0708514
- Maurice N, Thierry A-M, Glowinski J, Deniau J-M. 2003. Spontaneous and Evoked Activity of Substantia Nigra Pars Reticulata Neurons during High-Frequency Stimulation of the Subthalamic Nucleus. *J Neurosci* **23**:9929–9936.
- McDevitt RA, Tiran-Cappello A, Shen H, Balderas I, Britt JP, Marino RAM, Chung SL, Richie CT, Harvey BK, Bonci A. 2014. Serotonergic versus Nonserotonergic Dorsal Raphe Projection Neurons: Differential Participation in Reward Circuitry. *Cell Rep* **8**:1857–1869. doi:10.1016/j.celrep.2014.08.037
- McIntyre CC, Savasta M, Kerkerian-Le Goff L, Vitek JL. 2004. Uncovering the mechanism(s) of action of deep brain stimulation: activation, inhibition, or both. *Clin Neurophysiol* **115**:1239–1248. doi:10.1016/j.clinph.2003.12.024
- Meissner W, Leblois A, Hansel D, Bioulac B, Gross CE, Benazzouz A, Boraud T. 2005. Subthalamic high frequency stimulation resets subthalamic firing and reduces abnormal oscillations. *Brain* **128**:2372–2382. doi:10.1093/brain/awh616
- Nieh EH, Matthews GA, Allsop SA, Presbrey KN, Leppla CA, Wichmann R, Neve R, Wildes CP, Tye KM. 2015. Decoding Neural Circuits that Control Compulsive Sucrose Seeking. *Cell* **160**:528–541. doi:10.1016/j.cell.2015.01.003

- Paxinos G, Watson C. 2005. *The Rat Brain in Stereotaxic Coordinates*. Elsevier Academic Press.
- Pelloux Y, Baunez C. 2013. Deep brain stimulation for addiction: why the subthalamic nucleus should be favored. *Curr Opin Neurobiol*, 23/4 *Addiction* **23**:713–720. doi:10.1016/j.conb.2013.02.016
- Pelloux Y, Degoulet M, Tiran-Cappello A, Cohen C, Lardeux S, George O, Koob GF, Ahmed SH, Baunez C. 2018. Subthalamic nucleus high frequency stimulation prevents and reverses escalated cocaine use. *Mol Psychiatry*. doi:10.1038/s41380-018-0080-y
- Romito LM, Raja M, Daniele A, Contarino MF, Bentivoglio AR, Barbier A, Scerrati M, Albanese A. 2002. Transient mania with hypersexuality after surgery for high frequency stimulation of the subthalamic nucleus in Parkinson's disease. *Mov Disord Off J Mov Disord Soc* **17**:1371–1374. doi:10.1002/mds.10265
- Rouaud T, Lardeux S, Panayotis N, Paleressompoulle D, Cador M, Baunez C. 2010. Reducing the desire for cocaine with subthalamic nucleus deep brain stimulation. *Proc Natl Acad Sci* **107**:1196–1200. doi:10.1073/pnas.0908189107
- Sanders TH, Jaeger D. 2016. Optogenetic stimulation of cortico-subthalamic projections is sufficient to ameliorate bradykinesia in 6-ohda lesioned mice. *Neurobiol Dis* **95**:225–237. doi:10.1016/j.nbd.2016.07.021
- Schweizer N, Pupe S, Arvidsson E, Nordenankar K, Smith-Anttila CJA, Mahmoudi S, André A, Dumas S, Rajagopalan A, Lévesque D, Leão RN, Wallén-Mackenzie Å. 2014. Limiting glutamate transmission in a Vglut2-expressing subpopulation of the subthalamic nucleus is sufficient to cause hyperlocomotion. *Proc Natl Acad Sci U S A* **111**:7837–7842. doi:10.1073/pnas.1323499111
- Shehab S, D'souza C, Ljubisavljevic M, Redgrave P. 2014. High-frequency electrical stimulation of the subthalamic nucleus excites target structures in a model using c-fos immunohistochemistry. *Neuroscience* **270**:212–225. doi:10.1016/j.neuroscience.2014.04.016
- Tai C-H, Boraud T, Bezard E, Bioulac B, Gross C, Benazzouz A. 2003. Electrophysiological and metabolic evidence that high-frequency stimulation of the subthalamic nucleus bridles neuronal activity in the subthalamic nucleus and the substantia nigra reticulata. *FASEB J* **17**:1820–1830. doi:10.1096/fj.03-0163com
- Timmermann L, Wojtecki L, Gross J, Lehrke R, Voges J, Maarouf M, Treuer H, Sturm V, Schnitzler A. 2004. Ten-Hertz stimulation of subthalamic nucleus deteriorates motor symptoms in Parkinson's disease. *Mov Disord* **19**:1328–1333. doi:10.1002/mds.20198
- Ting JT, Daigle TL, Chen Q, Feng G. 2014. Acute brain slice methods for adult and aging animals: application of targeted patch clamp analysis and optogenetics. *Methods Mol Biol Clifton NJ* **1183**:221–242. doi:10.1007/978-1-4939-1096-0\_14
- Wade CL, Kallupi M, Hernandez DO, Breyse E, Guglielmo G de, Crawford E, Koob GF, Schweitzer P, Baunez C, George O. 2017. High-Frequency Stimulation of the Subthalamic Nucleus Blocks Compulsive-Like Re-Escalation of Heroin Taking in Rats. *Neuropsychopharmacology* **42**:1850–1859. doi:10.1038/npp.2016.270
- Welter M-L, Burbaud P, Fernandez-Vidal S, Bardinet E, Coste J, Piallat B, Borg M, Besnard S, Sauleau P, Devaux B, Pidoux B, Chaynes P, Tézenas du Montcel S, Bastian A, Langbour N, Teillant A, Haynes W, Yelnik J, Karachi C, Mallet L. 2011. Basal ganglia dysfunction in OCD: subthalamic neuronal activity correlates with symptoms severity and predicts high-frequency stimulation efficacy. *Transl Psychiatry* **1**:e5. doi:10.1038/tp.2011.5

- Welter M-L, Houeto J-L, Bonnet A-M, Bejjani P-B, Mesnage V, Dormont D, Navarro S, Cornu P, Agid Y, Pidoux B. 2004. Effects of High-Frequency Stimulation on Subthalamic Neuronal Activity in Parkinsonian Patients. *Arch Neurol* **61**:89–96. doi:10.1001/archneur.61.1.89
- Yizhar O, Fenno LE, Davidson TJ, Mogri M, Deisseroth K. 2011. Optogenetics in Neural Systems. *Neuron* **71**:9–34. doi:10.1016/j.neuron.2011.06.004
- Yoon HH, Min J, Hwang E, Lee CJ, Suh J-KF, Hwang O, Jeon SR. 2016. Optogenetic Inhibition of the Subthalamic Nucleus Reduces Levodopa-Induced Dyskinesias in a Rat Model of Parkinson's Disease. *Stereotact Funct Neurosurg* **94**:41–53. doi:10.1159/000442891
- Yoon HH, Park JH, Kim YH, Min J, Hwang E, Lee CJ, Suh J-KF, Hwang O, Jeon SR. 2014. Optogenetic inactivation of the subthalamic nucleus improves forelimb akinesia in a rat model of Parkinson disease. *Neurosurgery* **74**:533–541. doi:10.1227/NEU.0000000000000297


## Figure legends





**Fig. 1:** STN optogenetic inactivation increases the motivation of rats to work for sweet food.


**A.** Representative image showing the fluorescence in the STN (red lines) for an animal expressing ARCHT. Dotted rectangles indicate the position of the optic fibers. **B.** Schematic representation of laser pattern used for STN inhibition, which consisted in an alternation of 5min ON (15s illumination followed by 5s of obscurity) and OFF bins, and during ON bins laser was applied using a repeating pattern of 15s illumination followed by 5s of obscurity. **C.** Under FR5 schedule of reinforcement, laser inhibition of the STN applied during 8 consecutive sessions, did not significantly alter the number of pellets obtained by the rats (Mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}}(19, 247) = 1.550$ ;  $P = 0.0699$ ). **D.** Under PR schedule of reinforcement, laser inhibition transiently increased the breakpoint (i.e. last ratio completed) of rat working for sugar pellets (Mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}}(7, 91) = 2.574$ ,  $P = 0.0182$ , @ interaction effect). One day after the OFF period, another day of laser inhibition (laser challenge) was performed, which resulted in an increased breakpoint in the rats subjected to optogenetic inhibition (Mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}}(1, 13) = 8.174$ ,  $P = 0.0134$ , Fisher's post-hoc test: \*  $P < 0.05$ , vs control, ###  $P < 0.001$ , vs OFF). **E.** Percentage of changes in breakpoint when the laser illumination was turned ON, relative to previous day without laser modulation. Data correspond to the baseline-laser transition and the laser challenge (Mann-Whitney test,  $P = 0.0324$ )



**Fig. 2:** STN optogenetic stimulation at 130Hz reduces motivation for sweet food. **A.**

Example of fluorescence observed in the STN (red lines) in an animal transfected with


CHETA-TC. **B.** Schematic representation of laser pattern used for STN stimulation, which consisted in an alternation of 5min ON and OFF bins, during the former, the laser was applied at 130Hz without interruptions. **C.** In the FR5 task, the influence of 130Hz STN optogenetic stimulation on the number of pellets obtained by the rats was assessed over multiple sessions, application of laser stimulation with an intensity of 5mW and 10mW reduced the food intake. (Mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}} (23, 230) = 1.794, P = 0.0168, *$  interaction effect). **D.** Comparison of the influence of laser intensity on rat performance calculated on 4-day blocks relative to the last block with no laser stimulation (Mixed ANOVA,  $F_{\text{intensity} \times \text{subjects}} (2, 20) = 4.610, P = 0.0226$ ; Sidak's post-hoc test vs. control group: \*\*,  $P < 0.01$ ). **E.** Under PR schedule of reinforcement, the break point of stimulation group was reduced by the 130Hz optogenetic stimulation (Mixed ANOVA, Interaction  $F_{\text{sessions} \times \text{subjects}} (17, 153) = 2.736, P = 0.0006$ ; Sidak's post-hoc test vs. control group: \*\*,  $P < 0.01$ ). **F.** The performances of the 2 first OFF sessions were normalized using the last 2 sessions under optogenetic stimulation: after 8 consecutive sessions of laser stimulation, animals exhibited a transient increase in breakpoint when the laser was turned OFF (Mann Whitney test,  $P = 0.0476$ ). **G.** Efficiency of laser stimulation was assessed at various frequencies revealing higher impact for high frequencies. Data were averaged over 3-day blocks (Mixed ANOVA,  $F_{\text{frequency} \times \text{subjects}} (4, 36) = 3.001, P = 0.0310$ ; Sidak's post-hoc test vs. control group: \*\*,  $P < 0.01$ ). **H.** The influence of the pulse duration on the breakpoint was assessed at 100Hz. 3ms pulses were more effective than 2ms pulses to impact food motivation in the PR task, data were averaged over 3-day blocks (Mixed ANOVA,  $F_{\text{pulse duration} \times \text{subjects}} (2, 18) = 4.670, P = 0.0232$ ; Sidak's post-hoc test vs. control group: \*,  $P < 0.05$ ).



**Fig. 3:** STN optogenetic inhibition but not stimulation reduces cocaine intake in FR5 task.


**A.** Example of STN fluorescence in the STN (red lines) of an animal expressing inhibitory opsins ARCHT. **B.** Laser inhibition reduced the number of cocaine injections obtained by the rats in the FR5 (Mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}} (19, 152) = 1.958, P = 0.0137$ ). **C.** Normalization using the performance during the last day of baseline further highlighted this decrease (Mixed ANOVA,  $F_{\text{Group}} (1, 8) = 5.772, P = 0.0430$ ; Sidak post-hoc test: \*,  $P <$

0.05). **D.** Averaged performance across the eight sessions of laser inhibition revealed a lower level of intake compared to controls (Mann Whitney test,  $P = 0.0317$ , ###,  $P < 0.001$  for inhibition group). **E.** The percentage of active nose-poke when the laser stimulation was turned ON (i.e. during laser bins) was calculated and was not different between stimulation and control group (Mann Whitney test,  $P = 0.2222$ ). **F.** Typical image of fluorescence in the STN (red lines) of an animal expressing excitatory opsins CHETA-TC. **G.** 130Hz laser stimulation did not change the number of cocaine injection taken by the stimulated animals compared to the control group (Mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}}(19, 152) = 0.5815$ ;  $P = 0.9151$ ). **H.** Normalization to the last day of baseline revealed a similar level of performance when controlling for inter- individual variability (Mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}}(11, 88) = 0.7318$ ,  $P = 0.7053$ ). **I.** Average variation of performance over the 8 laser sessions did not emphasize any effect of 130Hz laser stimulation (Mann Whitney test,  $P = 0.8889$ ). **J.** The percentage of active nose-pokes when laser stimulation was turned ON did not differ between ‘stimulation’ and ‘control’ groups (Mann Whitney test,  $P = 0.9999$ ).



**Fig. 4:** STN optogenetic inhibition reduces cocaine intake in PR task and optogenetic stimulation produces a moderate trend toward a decrease. **A.** Laser inhibition reduced the number of cocaine injections obtained by the rats in the PR task (Mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}} (19, 133) = 1.972$ ,  $P = 0.0138$ , ####,  $P < 0.0001$  for inhibition group). **B.** Normalization of data by the last day of baseline allowed controlling for inter-individual variability and thus highlighted changes in cocaine intake during laser inhibition (Mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}} (11, 77) = 2.144$ ,  $P = 0.0265$ ; Sidak post-hoc test: \*,  $P < 0.05$ , vs control). **C.** Average variations in performance over the 8 laser sessions of inhibition further accentuated this

decrease (Mann Whitney test,  $P = 0.0159$ ). **D.** Variations in breakpoint (ratio) at baseline-laser transition for inhibition and control groups (Mann Whitney test,  $P = 0.0079$ ). **E.** 130Hz laser stimulation did not change cocaine intake of stimulated animals compared to the control group (Mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}} (19, 133) = 1.185, P = 0.2792$ ). **F.** Normalization to the last day of baseline did not highlight further variations in cocaine intake (Mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}} (11, 77) = 1.620, P = 0.1097$ ). **G.** Average performance over laser sessions revealed a similar level of intake (Mann Whitney test,  $P = 0.2540$ ). **H.** Variations in breakpoint (ratio) at baseline-laser transition for stimulation and control groups (Mann Whitney test,  $P = 0.9999$ ).



**Fig. 5:** Effects of optogenetic modulation on STN neurons. **A.** Typical response recorded in a STN neuron expressing ARCHT driven at a 10Hz frequency undergoing 15s of laser inhibition. **B.** Firing rate in STN neurons expressing inhibitory opsin ARCHT was reduced by a single 15s inhibitory light pulse (Friedman test,  $P < 0.0001$ ; Dunn's post-hoc test: \*\*\*,  $P < 0.001$ , vs baseline,  $n = 10$  cells). **C.** Example of response for a STN neuron during 5 min



of laser inhibition applied with the same pattern as during behavioral experiments. **D.** Firing rate in STN neurons expressing ARCHT was reduced by 5min of application of laser pattern used during behavioral testing (Friedman test,  $P = 0.0207$ ; Dunn's post-hoc test: \*,  $P < 0.05$ , vs baseline,  $n = 7$  cells). **E.** Typical excitatory response recorded in current-clamp mode for a STN neuron expressing excitatory opsin CHETA-TC and facing 10s of laser stimulation at various frequencies: 20 (light grey), 50 (purple) and 130Hz (blue). **F.** Number of action potentials elicited during 10s of optogenetic stimulation (Friedman test,  $P = 0.6145$ ,  $n = 10$  cells). **G.** Variation of membrane potential caused by the 10s optogenetic stimulation at various frequencies (Friedman test,  $P < 0.0001$ ; Dunn's post-hoc test: \*\*\*,  $P < 0.001$ , vs 10Hz,  $n = 10$  cells). **H.** Example of response in CHETA-TC positive STN neuron at onset (left panel) and offset (right panel) of 5min laser stimulation at 130Hz used during behavioral testing. **I.** Average changes in firing rate induced by 5 min of 130Hz laser stimulation (Friedman test,  $P = 0.0289$ ; Dunn's post-hoc test: \*,  $P < 0.05$ , vs baseline,  $n = 8$  cells).

## Supplemental information

### Electrophysiology

Animals were anesthetized and perfused intra-cardiacally with ice-cold artificial cerebrospinal fluid (ACSF). 200  $\mu\text{m}$  coronal slices containing the STN were prepared in ice-cold ACSF. After being cut, the slices were maintained for 10 minutes at 33°C and then transferred to holding ACSF at room temperature. ACSF used for perfusion, cutting, and recovery contained NMDG as a sodium substitute and contained, in mM: 92 NMDG, 2.5 KCl, 1.25  $\text{NaH}_2\text{PO}_4$ , 30  $\text{NaHCO}_3$ , 20 HEPES, 25 glucose, 2 thiourea, 5 Na-ascorbate, 3 Na-pyruvate, 0.5  $\text{CaCl}_2$  and 10  $\text{MgSO}_4\cdot 7\text{H}_2\text{O}$  (pH: 7.35). ACSF used for holding slices prior to recording was identical, but contained 92 mM NaCl instead of NMDG and contained 1 mM MgCl and 2 mM  $\text{CaCl}_2$ . ACSF used to perfuse slices during recording was maintained at 31°C and contained, in mM, 125 NaCl, 2.5 KCl, 1.25  $\text{NaHPO}_4$ , 1 MgCl, 2.4 CaCl, 26  $\text{NaHCO}_3$  and 11 glucose. All ACSF preparations were saturated with 95%  $\text{O}_2$  and 5%  $\text{CO}_2$ .

Cells were patched using glass pipets with resistance 3.5-5.0M $\Omega$ , filled with internal solution containing, in mM, 140 potassium gluconate, 5 NaCl, 2 MgCl<sub>2</sub>, 10 HEPES, 0.5 EGTA, 2 ATP, 0.4 GTP (pH: 7.35). Series resistance was monitored during experiment with 10mV hyperpolarizing pulses and maintained below 20M $\Omega$ , or cells were discarded. Firing pattern of STN neurons was determined by injecting depolarizing currents during 500ms with 20pA increment steps.

Cells were optically stimulated through a 200 $\mu\text{m}$  fiber placed in the bath with light beam aimed at the STN. Intensity was set at 10mW for stimulation and 5mW for inhibition experiment, except when the influence of the light intensity was assessed. For experiments

testing the influence of light intensity and pulse width, cells were held at -60mV in current-clamp configuration and discrete light pulses were applied. When testing the influence of light intensity, pulse duration was set to 15s and 1s for inhibition and stimulation groups, respectively. Testing the influence of the pulse duration was performed at constant light intensity: 10mW (Stimulation) and 5mW (Inhibition).

### Spontaneous- and cocaine-induced locomotor activity

Locomotor activity was measured as the distance traveled (in m) in a circular homemade Perspex open-field (60 cm diameter) with a video tracking performed with Bonsai (Open Ephys), recorded on a PC and analyzed offline. Rats were placed in the open-field for 30 min of habituation and were connected to the optic fiber cable and further recorded for 30 min during which they underwent laser stimulation for 2 bins of 5 min each. Their spontaneous locomotion was measured. For the measure of cocaine-induced locomotion, the animals were injected at the end of the habituation with either 0.9% NaCl (1 ml/kg) or one of the various doses of cocaine (5; 7.5 and 10 mg/kg, *s.c.*) whose order was counterbalanced in a pseudo-latin square manner, with a five-day interval between two injections to allow for washout. The doses were chosen according to previous studies in line with their reinforcing effect associated with limited effect on locomotion to avoid stereotyped behavior in case of potentiation by the STN manipulation (Baunez et al., 2005).

## Histology

At the end of the experiment, the rats were deeply anesthetized with pentobarbital (Dolethal, *i.p.*) and perfused intracardially with 4% paraformaldehyde dissolved in PBS. The brains were extracted and after cryo-protection in 30% sucrose, they were frozen into liquid isopentane (-80°C) to be further cut in 40µm coronal slices with a cryostat

For the optogenetic experiments, the brain sections were examined for optic fibers location and for native fluorescence expression with an epifluorescence microscope (Zeiss, Imager.z2) immediately after being cut, except for the CHETA-TC animals, in which native fluorescence was present but required immuno-staining to control the exact boundaries of the viral expression.

Sections underwent a 90 min permeation step (PBS, 1% bovine serum albumin (BSA) 2% normal goat serum (NGS), 0.4% TritonX-100), 3x5min PBS washes, incubation with primary antibody (mouse anti-GFP, A11120, Life technologies; 1:200, in PBS 1% BSA, 2% NGS, 0.2% TritonX-100) at 4°C overnight. The sections were then washed 3x5min with PBS followed by 2h incubation at room temperature with secondary antibody (Goat anti-mouse Alexa 488, A11011, Life technologies, 1:400 in PBS 1% BSA, 2% NGS), and finally washed 3x5min with PBS before being mounted on to glass slides with homemade mounting medium.


67 animals were used in the optogenetic procedure, among which 2 were excluded after loss of optic fiber implants, 3 for absence of fluorescence in either one or both STN, and 1 due to optic fiber misplacement.

## Statistical analyses


Data are expressed as mean  $\pm$  SEM with the sample size indicated for each group. Analyses were performed with Prism (GraphPad) and Matlab (Mathworks) software. Behavioral data were analyzed with mixed-model design ANOVA (mixed ANOVA) with groups as between factor and sessions, time bins, or light parameters as a repeated within factor. Follow-up analysis was conducted using Sidak's post-hoc test for multiple comparisons or Fisher's post-hoc test for single comparison. Data regarding cocaine-induced locomotion were analyzed with 2-way Repeated Measure ANOVA with dose and time bins as repeated within factors. Two samples comparisons were analyzed with non-parametric Mann-Whitney test. Electrophysiological data were analyzed with Friedman test followed by Dunn's post-hoc test, variations in cells properties were analyzed with Kruskal-Wallis test or mixed ANOVA. Only  $P$ -values  $< 0.05$  were considered significant.

## Supplementary figures


**A**




**B**



**C**




**D**



**Supplementary Fig. 1:** Effect of STN optogenetic modulation on rat spontaneous and cocaine-induced locomotion. **A.** Locomotor activity expressed in distance travelled (in m) per 5 min bins during the 30min of habituation (left) and 30min of test (right) with 2 bins of


laser activation (yellow) for the control (black dots), stimulation (blue dots) and inhibition (green dots) groups. There was a significant interaction effect when comparing ‘inhibition’ group to ‘control’ (mixed ANOVA,  $F_{\text{Time bins} \times \text{subjects}} (5, 105) = 8.576, P < 0.0001$ ; Sidak’s post-hoc test: \*  $p < 0.05$ , \*\*\*  $p < 0.001$ ) while no effect was observable when comparing ‘stimulation’ and ‘control’ groups (mixed ANOVA,  $F_{\text{Time bins} \times \text{subjects}} (5, 85) = 1.253, P = 0.2920$ ). **B – D.** Effect of STN optogenetic modulation on cocaine-induced locomotion tested with various doses of cocaine: 5mg/kg (orange), 7.5mg/kg (brown), 10mg/kg (red) and compared to the saline control (black) in control (**B.**), inhibition (**C.**) and stimulation (**D.**) groups. There was no effect of the cocaine dose on the locomotor activity in any of the 3 groups: ‘control’, ‘inhibition’ and ‘stimulation’ (Repeated Measure 2-way ANOVA,  $F_{\text{Time bins} \times \text{dose}} (15, 120) = 1.288, P = 0.2202$ ;  $F_{\text{Time bins} \times \text{dose}} (15, 120) = 0.4931, P = 0.9402$ ;  $F_{\text{Time bins} \times \text{dose}} (15, 40) = 0.7500, P = 0.7207$ , respectively). The laser beam activation produced an increase in locomotor activity in the ‘inhibition’ group regardless of the dose of cocaine (Repeated Measure 2-way ANOVA,  $F_{\text{laser bins}} (5, 120) = 17.26, P < 0.0001$ ).




**Supplementary Fig. 2:** Effects of STN optogenetic modulation on food magazine visits and inactive hole responses recorded in the progressive ratio (PR) task illustrated for each session (8 sessions of baseline, 8 sessions with laser ON (i.e. Laser), 4 sessions with laser OFF (i.e. OFF) and 1 session with laser ON again (i.e. challenge)). **A.** Influence of STN optogenetic inhibition on the number of responses made in the food magazine. When the laser was turned on, there was a significant increase in food magazine responses at baseline-laser transition (1<sup>st</sup> session of the laser ON period)(mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}}(1, 13) = 8.809, P = 0.0109$ ) and at off-challenge transition (mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}}(1, 13) = 7.968, P = 0.0144$ ). **B.** Influence of STN 130Hz optogenetic stimulation on the responses made in the food magazine. The 130 Hz stimulation induced a strong trend towards decreased number of responses in the stimulated animals (mixed ANOVA,  $F_{\text{Group}}(1, 9) = 4.864, P = 0.0549$ ). **C.** Effect of STN optogenetic inhibition on the number of responses made in the inactive hole. The number of erroneous responses remained unaffected either during baseline-laser transition (mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}}(1, 13) = 0.008801, P = 0.9267$ ) or during the laser challenge (mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}}(1, 13) = 1.267, P =$




0.2807). **D.** Effect of STN optogenetic 130 Hz stimulation on the number of responses made in the inactive hole. The number of erroneous responses was not affected by the 130Hz optogenetic stimulation (mixed ANOVA,  $F_{\text{sessions} \times \text{subjects}} (1, 9) = 3.286, P = 0.1033$ ). Fisher's post-hoc test: \*  $p < 0.05$ , \*\*  $p < 0.01$ , vs control.



**Supplementary Fig. 3:** Effect of acute 20 Hz optogenetic stimulation applied during one session of progressive ratio for cocaine. The motivation of the animals (controls in black and stimulated in blue) is expressed in number of injections obtained during one session with laser OFF, one session with laser ON (20 Hz) and one last session with laser OFF. The acute 20Hz optogenetic stimulation did not alter motivation for cocaine (mixed ANOVA, F sessions x subjects (2, 14) = 0.3486, P = 0.7116).




**Supplementary Fig. 4:** Evaluation of the percentage of active nose-pokes while the laser was activated. **A.** The percentage of active nose-poke when the laser beam was turned ON (i.e. during laser bins) was not modified compared to controls (Mann Whitney test,  $P = 0.2857$ ). **B.** Laser stimulation did not alter the pattern of nose-poke of stimulated animals compared to the control group (Mann Whitney test,  $P = 0.4127$ )



**Supplementary Fig. 5:** Cells properties of STN neurons after viral infection recorded in whole cell configuration for controls (EYFP, n=8 cells, black bars), stimulation (CHETA-

TC, n=10 cells, blue bars) and inhibition (ARCT, n=11 cells, green bars). Cell resting membrane potential (**A.**, Kruskal-Wallis test,  $P = 0.8214$ ), cell capacitance (**B.**, Kruskal-Wallis test,  $P = 0.0644$ ), membrane resistance (**C.**, Kruskal-Wallis test,  $P = 0.7397$ ), and access resistance (**D.**, Kruskal-Wallis test,  $P = 0.3158$ ) were unaffected. **E.** The average number of action potentials (APs) elicited by a 500ms step of hyperpolarization was not modified due to the presence of the opsins (**E.**, mixed ANOVA,  $F_{\text{Frequency} \times \text{Group}}(30, 390) = 0.2480, P > 0.9999$ ).



**Supplementary Fig. 6:** Evaluation of light stimulation parameters in cells expressing inhibitory opsins (ARCHT, n=11 cells) or excitatory opsins (CHETA-TC, n=10 cells). **A.** Example of light induced inhibitory currents in one STN neuron expressing ARCHT by a 5s (brown) and 15s (blue) pulse. **B.** Average area under the curve (AUC) calculated for different durations of light pulses with intensity set at 5mW (Friedman test,  $\chi^2(5) = 51.88$ ,  $p < 0.0001$ , Dunn's post-hoc test: \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$ ; vs 1s pulse). **C.** Average current induced by a single 15s pulse applied with different light intensities (Friedman test,  $\chi^2(2) = 22.00$ ,  $p < 0.0001$ ; Dunn's post-hoc test: \*,  $p < 0.05$ ; \*\*\*,  $p < 0.001$ ). **D.** Example of induced depolarizing current in CHETA-TC cells induced by a 0.5ms (pink), 2ms (brown)

or 5ms (purple) light pulse. **E.** Average induced current in CHETA-TC cells depending on the duration of the light pulse with intensity set at 10mW (Friedman test,  $\chi^2 (5) = 46.00$ ,  $p < 0.0001$ ; Dunn's post-hoc test: \*\*\*,  $p < 0.001$ ; vs 0.5ms pulse). **F.** Induced current by a 1s light pulse as a function of light intensity (Friedman test,  $\chi^2 (3) = 28.92$ ,  $p < 0.0001$ ; Dunn's post-hoc test: \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$ ; vs 2.5mW)