

Contribution to the reappraisal of the mid Paleogene ichtyofauna of Western Africa with three new enigmatical elasmobranchs from Thanetian–Lutetian of Senegal

Bernard Siguendibo Sambou, Lionel Hautier, Raphael Sarr, Rodolphe Tabuce, Fabrice Lihoreau, Moustapha Thiam, Renaud Lebrun, Jérémy Martin, Henri Cappetta, Sylvain Adnet

To cite this version:

Bernard Siguendibo Sambou, Lionel Hautier, Raphael Sarr, Rodolphe Tabuce, Fabrice Lihoreau, et al.. Contribution to the reappraisal of the mid Paleogene ichtyofauna of Western Africa with three new enigmatical elasmobranchs from Thanetian–Lutetian of Senegal. Annales de Paléontologie, 2020, 106 (3), pp.102400. $10.1016/j.annpal.2020.102400$. hal-02992155

HAL Id: hal-02992155 <https://hal.science/hal-02992155v1>

Submitted on 23 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

 Contribution to the reappraisal of mid Paleogene ichtyofauna of Western Africa with three new enigmatical elasmobranchs from Thanetian-Lutetian of Senegal. Contribution à la réévaluation de l'ichtyofaune du Paléogène moyen en Afrique de l'Ouest avec trois nouveaux élasmobranches énigmatiques du Thanétien-Lutétien du Sénégal. 7 Bernard Siguendibo Sambou², Lionel Hautier¹, Raphael Sarr², Rodolphe Tabuce¹, Fabrice 8 Lihoreau¹, Moustapha Thiam², Renaud Lebrun¹, Jeremy E. Martin³, Henri Cappetta¹, Sylvain Adnet^{1*} 11 ¹Institut des Sciences de l'Evolution, UMR5554, CNRS, IRD, EPHE, Université de Montpellier, Montpellier, France ² Département de Géologie, Faculté des Sciences et Techniques, Université Cheikh Anta Diop de Dakar, B. P. 5005 Dakar, Sénégal ³Univ. Lyon, ENS de Lyon, Université Claude Bernard Lyon 1, CNRS, UMR 5276 Laboratoire de Géologie de Lyon: Terre, Planètes, Environnement, F-69342 46 Allée d'Italie, Lyon, France *Corresponding author: sylvain.adnet@umontpellier.fr

-
-
-
-

Abstract.

 We report here three new elasmobranch fossil taxa from Thanetian-Lower Lutetian nearshore marine deposits of northeastern (Matam region) and central-western (Sine-Saloum region) Senegal. These three new taxa represent the oldest species of the enigmatical elasmobranch *Odontorhytis*, the oldest putative representatives of marine potamotrygonid, and an uncertain dasyatoid genus with the smallest grinding dentition ever described. These new taxa, representing the second Cenozoic elasmobranch remains formally described from Senegal, broaden our understanding of their evolutionary and biogeographic history in the equatorial Eastern Atlantic during this time period. Their occurrences confirm hypothesised stratigraphically correlations between the top of Matam Fm. and the base of Lam Lam Fm., refine the marine connections between these two Senegalese regions, and suggest that these genera were broadly distributed within the shallow marine settings of the equatorial Eastern Atlantic during latest Paleocene – early Middle Eocene.

Keywords: Elasmobranch, new taxa, Paleogene, Senegal.

Résumé.

 Nous rapportons ici trois nouveaux taxa de fossiles d'élasmobranches des dépôts marins littoraux du Thanétien-Lutétien inférieur du Nord-Est (région de Matam) et du Centre-Ouest (région du Sine-Saloum) du Sénégal. Ces trois nouveaux taxa représentent les espèces les plus anciennes d'élasmobranches énigmatiques appartenant au genre *Odontorhytis*, aux potamotrygonides marins et à un dasyatoïde avec la plus petite denture broyeuse jamais décrite. Ces nouveaux taxa, complétant l'unique description d'élasmobranches cénozoïques nouveaux au Sénégal, élargissent notre compréhension de leur histoire évolutive et biogéographique de l'Atlantique Est équatorial au cours de cette période. Leurs occurrences confirment des corrélations stratigraphiques hypothétiques entre le sommet de la Formation de Matam et la base de la Formation de Lam Lam. Celles-ci affinent aussi les liens marins entre ces deux régions sénégalaises et leur comparaison avec des occurrences reconnues en Afrique du Nord et de l'Ouest suggèrent que ces genres étaient largement répartis et diversifiés dans les environnements marins peu profonds de l'Atlantique Est équatorial au cours du Paléocène terminal et du début de l'Éocène moyen.

Mots-clés: Elasmobranche, nouveaux taxa, Paléogène, Sénégal.

-
-
-
-

Introduction

 Biogeographic origins of African modern vertebrates have been the focus of considerable research since last decade in warm Paleogene period of West Africa. Among them, elasmobranch fossil record devoted to mid Paleogene was very partial with a unique description of taxa such as *Lophobatis* Cappetta, 1986, preliminary reports (Pascal, 1987; Cappetta and Traverse, 1988; Stevens et al., 2011), and/or ancient works that (e.g. Strömer, 1910, White, 1926, 1934, 1955). Biogeographic interpretations of past elasmobranch diversity in Central Western Atlantic during Paleogene have been mainly evidenced based on the richer northwestern (e.g. Morocco) and southwestern (e.g. Angola) African deposits (e.g. Arambourg, 1952; Noubhani and Cappetta, 1997; Dartevelle and Casier, 1943, 1959; Solé et al., 2018). However, a corresponding body of research is largely lacking for the Paleogene equatorial Eastern Atlantic elasmobranchs, despite the fact that nearly all of the known Cenozoic rocks along the West African coast are marine, especially near the Guinean Gulf. Indeed, reappraisal of the mid Paleogene Western African faunas is essential to better understand the elasmobranch evolution in the equatorial paleo-Atlantic. Currently, the Senegalese elasmobranch Cenozoic fossil record is only represented by mid Paleogene taxa from the phosphate ores of Matam (Pascal, 1987; Sambou et al., 2018; Sambou 2018) and Taïba (Hautier et al., 2012, 2014). Other associated marine vertebrate, including bony fish like amiid or scombrid (O-Leary et al. 2012, Cleason et al. 2015), sea cows and whales (Hautier et al., 2012) are recovered in Early-Middle Eocene from same or different localities (e.g. Gossas, Bargny). We report here, three new taxa of elasmobranchs with unambiguous dental morphologies that support biostratigraphical correlations between Formations from two

Geological setting, material and methods.

Most of the specimens were found in the phosphate ore of Ndendouri-Ouali Diala near Matam

(sub-prefecture of Orkadiéré, Matam region, approximatively 15°18'N, 13°01'W, see

Sambou et al., 2017 for details) located on the left riverbank of the Senegal River,

northeasternstern part of Senegal (Fig. 1). The studied material was collected in water-well

cuttings during ancient campaigns of M. Pascal (see Pascal, 1987) that allowed to collect an

amount of 450 kg of phosphatic sediment and 50 kg of sandy sediment in a series of wells

(collecting H.C.) and from recent wells drilled manually in the subsoil of Ndendouri-Ouali

Diala (collecting B.S.) and following a NW direction (see Sambou, 2018). Sambou et al.

(2017) recently revised the faunas from the Gorgol and Matam Fm. outcropping in the

phosphate ore of Matam as previously reported in Pascal (1987). 38 species of elasmobranch

have been recognized in the different levels of the 20 well-drilling and 33 different samples

along the Matam series (see Sambou et al., 2017 : Tab.1 for details). These faunas include

several determined and undetermined species of *Brachycarcharias, "Cretolamna",*

Protoginglymostoma, Ginglymostoma, Nebrius, Rhizoprionodon, Physogaleus, Triakis,

Foumtizia, Rhynchobatus, Pristis, Anoxypristis, Eotorpedo, Lophobatis, Myliobatis,

Rhinoptera, Leidybatis, Garabatis, Coupatezia, Ouledia and the three new taxa reported here.

From these, five fossiliferous associations were identified along the available

sedimentological successions (see Fig. 1), and confirmed the late Thanetian age of the top of

the Gorgol Fm. (association 1 – lithological Unit 0), the Ypresian age of the phosphate

 deposits belonging to Matam Fm (rich associations 2-4, lithological Units 1-3), and the Lutetian age of the higher clay-limestone unit of Matam Fm. (association 5, lithological Unit 4).

 The other specimens come from the water-well cuttings located at Sanghaye Bele village (14°16'55.5"N, 16°6'14.01"W, Fig. 1), north of Kaolack in the Sine-Saloum region (central- western part of Senegal, see Tabuce et al., 2019 for details). The fossiliferous deposits attributed to the Lam Lam Fm. correspond to coquina limestone and yellow to white clayey limestone alternating with marls. Material was recovered by soft acid attack (acetic 8%) of 30 kg of sediments (see also Tabuce et al. 2019 for details). If the base of the Lam-Lam Formation is E8 in age (lower Lutetian), the top of the formation (Lam-Lam marls) yielded a rich planktonic foraminifera association that indicates the E9 biozone (top of lower Lutetian) (Flicoteaux 1974, Roger et al. 2009). These regional correlations indicate a confident lower Lutetian age (E8-E9 biozones, 49 to 45,8 Ma ~44 M.a., Vandenberghe et al. 2012) for the fossiliferous level in water-well of Sanghaye Bele (see Tabuce et al. 2019 for correlations). Some elasmobranch remains were collected in the 30 kg of limestone and include for instance some teeth of *Chiloscyllium* sp., *Nebrius* cf. *obliquus*, ?*Rhizoprionodon* sp., *Physogaleus* sp., *Rhynchobatus* sp., *Rhinobatos* sp., *Ouledia* sp., *?Jacquhermania attiai*, *Arechia* sp., Dasyatoid indet., *Coupatezia* sp. as well as two of the three new taxa described here. The elasmobranch assemblage from the coquina limestone of the Lam Lam Fm. in Sanghaye Bele would then be contemporaneous with the coquina limestone of the Ngolothie borehole near Kaolack (Tabuce et al. 2019) and most probably sub contemporaneous with the elasmobranch assemblage of the clay-limestone unit ending the Matam Fm. in Sambou et al. (2017).

FIG.1 here

Systematic Palaeontology

- *Institutional Abbreviations* Fossil material, including type series, is housed at the
- University of Montpellier, France (abbreviations SES, MAM) or at the University Cheikh
- Anta Diop of Dakar, Senegal (abbreviations SNSAN/2011/). Tooth terminology follows
- Cappetta (2012)
-
- Class CHONDRICHTHYES Huxley, 1880
- Subclass ELASMOBRANCHII Bonaparte, 1838
- SubCohort NEOSELACHII Compagno, 1977
- Superorder BATOMORPHII Cappetta, 1980
- Order MYLIOBATIFORMES Compagno, 1973
- Superfamily DASYATOIDEA Compagno, 1973
- ?Potamotrygonidae Garman, 1877
- ?Styracurinae Carvalho et al., 2016
-
- *Atlantitrygon nov.gen.*
- *Type species***:** *Dasyatis sudrei* Cappetta, 1972
- *Type material***:** Hototype: SES64, Sessao, Thanetian of Niger (Cappetta, 1972 : Plate 6 fig. 3)
- refigured in figure 2 A ; paratypes SES62-63, 65-67 (Cappetta, 1972 : Plate 6 fig. 1-2,4-6)
- partially refigured in figure 2 B-D.

 *Etymology***:** simplified Greek *"Atlanti"* refers to the past distribution of representatives, currently limited to tropical Atlantic sea along the western African coast, and Greek "*trygon"* meaning stingray.

 *Diagnosis***:** Dasyatoid genus characterized by gracile medium sized teeth with high crown compared to root, a development in height of the lingual zone and the lingual marginal face of crown (including the transversal keel); a long, slender cusp mesiodistally flattened and highlighted by an occlusal median crest, a junction between the undulating transverse keel and the median crest of cusp forming a three-pointed star in occlusale view ; a narrow lingual face and a concave labial face of crown with finely ornamented enameloid; a root with two sub-triangular lobes elongated transversally.

 Occurrence: from Thanetian (Niger, Togo and Senegal) with *A. sudrei* to lower Lutetian (Senegal) with *A. senegalensis* nov. sp.

 Discussion: Cappetta (1972) described *Dasyatis sudrei* from Sessao, Niger. At that time, diagnosis characters, e.g. the long slender cusp, were evaluated as enough peculiar compared to other species of *Dasyatis* s.l. to justify the creation of a new species (Cappetta, 1972 : p. 207). The author suggested that this taxon could be close to genus *Himantura*, unknown at the fossil state in 1972. In fact, a slender long cusp is known in many male teeth belonging to some dasyatidspecies (e.g. *Dasyatis, Hypanus*) and many Urotrygonidae (e.g. *Urobatis*) and Urolophidae (e.g. *Urolophus*) but this long cusp is usually rooted on the edge of lingual zone and its top is delimited by the transversal keel that follows the upper borders of cusp. The upper face of cusp, often concave, belongs in fact to the labial face, contrary to what is observed in *Atlantitrygon*. The unique dasyatoid teeth that display a comparable "displaced" cusp are observable in living western Atlantic "*Himantura" schmardae* (Werner, 1904 : see Cappetta, 2012, fig. 410) recently attributed to genus *Styracura* Carvalho et al., 2016wich also includes *S. pacifica* (Beebe and Tee-Van, 1941). Both these marine species, recently included in Potamotrygonidae, are considered more closely related to freshwater potamotrygonins (Carvalho et al., 2016). The inclusion of *Atlantitrygon* in the marine Styracurinae thus remains hypothetic and is only based on global resemblances of the tooth shape and the presence of a displaced main cusp. However, the teeth of *S. schmardae* are easily distinguishable from *Atlantitrygon* nov. gen*.* by having a straight transversal keel, a second transverse keel, running more or less parallel to main one, a cusp without crest and a ornamentations absent, contrary to the condition seen in teeth of *Atlantitrygon* nov. gen. Finally, the unique combination of characters in *Atlantitrygon* nov. gen. does support the creation of this new fossil genus but its systematic position among the Potamotrygonidae remains nevertheless uncertain, partly because our limited knowledge of the dental variation in living batoid (Guinot et al., 2018).

Atlantitrygon senegalensis nov. sp.

*Etymology***:** Refers to the country where the type series was found

Type material : Holotype Fig. 2 K, MAM-09 (well F0 9.18-9.3m, Unit 2 in Sambou et al.,

2017) Ndendouri-Ouali Diala, ?Early Ypresian, Matam Fm.; Paratypes Fig. 2 F-J, MAM-04

to MAM-08 and MAM-10 to MAM-11 (well F0 9.18-9.30m, Unit 2 in Sambou et al., 2017)

Ndendouri-Ouali Diala, ?Early Ypresian, Matam Fm

*Additional material***:** a fifteen teeth from Lower Lutetian (E8-E9 biozones, 49 to 45,8 My) of

Sine-Saloum region, Senegal and also partially figured Fig. 2 N-Q

Diagnosis: The new species distinguishes from the unique type species of genus by having

teeth with higher transversal keel that looks as more protruding, undulated and often marked

by three rounded pseudo cusplets in lingual view. Two of these cusplets are located at the

 margin while the third one, the highest, is situated in the cusp axis. Compared to the type species, ornamentation of the enameloid of the lingual and labial faces is more finely developed, especially under the transversal keel with an alveolar network shape. *Occurrence***:** Ypresian to Lower Lutetian of Matam and Lower Lutetian of Sine-Saloum regions, Senegal, possibly in the Ypresian of Mali (Tamaguilet) and Morocco (Ouled Abdoun)

 Description: The tooth morphology is consistent with that of dasyatoids, a paraphyletic group 217 that includes many different families of stingrays (e.g. DasyatidaeUrobatidae, Urolophidae, Urotrygonidae, Potamotrygonidae, Plesiobatidae) with no pavement-like arrangement. The dentition is monognathic heterodont with high-crowned teeth (e.g. Fig. 2 I), which decrease in size toward the commissure (e.g. Fig. 2 M, P). The holotype (Fig. 2 K) is a tooth probably located medially along the jaw. It displays a high crown (Fig. 2 K2) with robust cusp (Fig. 2 222 K2). In occlusal view (Fig. 2 K1), the crown is semi-oval in shape, broader than long, with a long, slender cusp with median crest on its occlusal part. This median crest does not reach the apex of cusp (Fig. 2K1). The crown has a high, transverse keel, which divides the crown into distinct labial and lingual parts (Fig. 2K1). The junction between the undulating transverse keel and the median crest of cusp forms a three-pointed star (Fig. 2 K1). The transverse keel is quite salient and forms an erected cutting edge. The upper part of the lingual face (below the transverse keel) and the labial face are fully ornamented. This ornamentation is quite delicate and organized as an alveolar network with larger alveolus on concave basin of the labial face (Fig. 2 K3). The upper part of the ornamented labial visor (labial border of labial face) is reduced to a straight crest that overhangs the lower part of the unornamented labial visor. The lower part of lingual face, also unornamented below the cusp axis, is relatively expanded marginally. The root is narrow in profile (Fig. 2 K2), of holaulacorhizous type with two short lobes separated by a low groove. A large central foramen opens in labial part of the

 nutritive groove. Tooth variability, in part due to heterodonty, mainly involves the general shape of the crown, which is sometimes subtriangular in shape (e.g. Fig. 2 I) with often a long slender cusp. This cusp is always present in all specimens suggesting no sexual heterodonty. In more anterior files, the crown sometimes displays a very high, undulating transverse keel 239 that forms three pseudo rounded cusplets in lingual view (e.g. Fig. 2 L1, O1, N1). One of them is located at the junction between the transverse keel and the median crest of the cusp. Lingual ornamentations seemingly are absent on the lower part of lingual face and below the cusp axis of all specimens, conversely to the condition seen in the whole labial face and in the narrow band of lingual marginal face just under the transversal keel. In anterior teeth, this ornamented upper part of lingual face is sometimes set apart from the unornamented lower part by rounded transverse keels in lingual view (Fig. 2 L, N). Enameloid ornamentation is quite delicate in the upper part of the lingual face (Fig. 2 H3, L2) and could be organized as an alveolar network with larger alveolus on concave basin of the labial face (Fig. 2 K3, M2). This ornamentation reduces gradually near the keel and crests where it takes the form of lace when not abraded. The labial visor is merely variable, being often reduced to a salient and straight crest delimiting the labial border of the labial face. The basal view of the crown (Fig. 251 2 H2, I1, O2) shows a broad and slightly convex crown rim at the outer part, which gradually narrows down to half of its width at the inner part. The crown–root junction is located in a shallow depression at the centre of the basal surface of the crown. The root is lower than the crown, of holaulacorhizous type with two lobes, which are triangular (fig. 2 I1, H2) to semi- circular in shape (fig. 2 O2) in basal view. The root base has a well-developed and deep median nutritive groove that encloses a single large central foramen (fig. 2 M1, O2). No inner or outer foramina are discernible on the specimens. The lower Lutetian material from Sine- Saloum (fig. 2 N-P) is quite similar to the type series from Matam region (fig. 2 F-M) despite its poorer state of conservation.

 Discussion: Compared to *A. sudrei*, the new species *A. senegalensis* nov. sp. displays teeth with similar shape, but its transversal keel is higher, more protruding, undulated and often marked by three rounded pseudo cusplets in lingual view. Two of these cusplets are located at the margin while the third one, the most developed, is situated in the cusp axis at the junction of the three-pointed star. Compared to *A. sudrei*, less salient crests are present on the concave labial face of the crown in *A. senegalensis* nov. sp., as observable in type series (e.g. fig. 2 B). The ornamentation of the enameloid in the band of the lingual marginal face under transversal keel (as those of the labial face) is more finely developed compared to *A. sudrei* with alveolar network shape, except near the keel and crests where they reduce gradually. Only a few specimens of *A. sudrei* have been recovered in the Thanetian of Matam, Senegal (fig. 2 E). It seems more abundant in the Nigerian type locality (Cappetta, 1972) and in Togolese Thanetian deposits (Guinot et al., 2020). Although badly preserved (partial crown with broken cusp), the tooth crown shows that the cusp was mesiodistally flattened and that the junction between the low transverse keel and the cusp formed a three-pointed star in occlusal view. The enameloid of the crown concave labial face is also finely ornamented (fig. 2 E2). Some teeth from Ypresian deposits of Matam (e.g. fig. 2 F, H) show a morphology very close to that of *A. sudrei* teeth with a weak transverse keel. This probably indicates a close filiation between the two chronospecies of the new genus. Similar teeth to *A. senegalensis* were also observed (H.C. pers. observations) in the Ypresian of Morocco (Ouled Abdoun) and Mali (Tamaguilet), which might indicate that *Atlantitrygon* representatives had a wider distribution along the tropical western Atlantic coasts, from Niger to Morocco, during the Thanetian-Ypresian period.

FIG.2 here

 occlusal view with depressed visor suggesting a light teeth imbrication or at least a direct contact between alternating teeth. The lingual face of the crown is straight and mainly unornamented, excepted just under the transverse keel where it is slightly concave with an enameloid marked by vertical folders. The labial face is flat, strongly ornamented with sorted micrometric holes piercing the whole surface. The labial visor is medially concave and its ornamentation is less homogeneous compared to labial face. The teeth roots are bilobate, vertically directed, narrower than the crown and relatively high with two pinched lobes separated by large but soft nutritive groove where a unique large foramen opens medially. *Description*: The teeth are very small, not exceeding 1.3 mm in size. They exhibit reduced sexual dimorphism in crown morphology, the crown teeth of presumed breeding males bearing a short cusp lingually directed (fig 3 A-B). The base of cusp is sometimes as wide as the transverse keel or medially centered (fig. 3 A-B). The non-cuspidate teeth of presumed females (conceivably teeth of non-breeding male or teeth from the lateral files of breeding males) display diamond-shaped occlusal outline. The holotype (fig. 3 D) probably belongs to a latero-anterior file of a female jaw. The crown is high with developed lingual face, which is sub-trapezoid in lingual view (fig. 3 D1). The lingual extremity is medially convex under the vertical bulge and the marginal borders are straight to slightly concave near the transverse keel. The enameloid is mainly unornamented except near the transverse keel where several enameloid folders are easily distinguishable (fig. 3 D1). The labial face is flat, its contour is diamond-shaped in occlusal view (fig. 3 D2) and displays many circular holes deeply piercing the enameloid surface (fig. 3 D3). The hole size is relatively homogenous except near the labial visor where the ornamentation is more unstructured. The labial visor is medially concave, which suggests a light imbrication with the lingual face of tooth directly anterior to it. The lower part of labial visor is thick and unornamented. The crown/root junction is located in a shallow depression in the centre of the basal crown surface. The root is narrower

 than the crown, of holaulacorhizous type with two lobes, which are semi-circular in shape in basal view. The root lobes are slightly lingually directed and almost vertical in profile. The root base displays a weak median nutritive groove that encloses a large central foramen. Male teeth (fig. 3 A-B) are similar in shape with a development of the medial cusp that could be high and as large than transversal keel, small (fig. 3 A) or reduced to a slight protuberanceof the transverse keel (fig. 3 B2). Such variability is probably due to different file position of the teeth, the higher teeth being possibly located near the symphysis. The female teeth are less variable, their crown being more or less narrow (fig. 3 C-D) or extended mesio-distally (fig.3 E). The lateral teeth have a mesio-distally extended labial face, a developed labial visor, a transverse keel slightly worn by functional wearing, and an oblique root lobe as in some extant myliobatoid with pavement-like dentition. Despite the limited sample of this minute species, teeth of female (fig. C-H) seem twice as large as those identified as belonging to male, conformably with expected size difference between sexes in most batoid species. No clear dental difference is obvious between specimens from the type series (Ypresian) and those from the Lower Lutetian (fig. 3 F-H).

 *Discussion***:** Currently, *Microtriftis* nov. gen. is only affiliated to dasyatoid of *incertae familiae*. If the morphology of isolated teeth (e.g. high crown with narrower root vertically oriented, flat ornamented labial face with concave visor in all files) suggests that each tooth is in contact with its neighbouring teeth to form a pavement-like dentition as observable in living myliobatoid, there are significant difference with their living and fossil representatives. In *Microtriftis* nov. gen., the root is always of holaulacorhizous type, like many dasyatoid but unlike myliobatoid that present a root of polyhaulacorhizous type. The imbrication of teeth (assumed here on alternating gridlines) seems light in *Microtriftis* nov. gen., the lingual face of tooth and the labial visor of previous tooth file could not be interlocked, in contrast to the condition observed in myliobatoid jaws. Other Paleogene dasyatoid *incertae sedis* developed

 a grinding dentition with joined alternating teeth (e.g. *Heterobatis* Noubhani and Cappetta 1997, *Hypolophodon* Cappetta 1980; *Ishaquia* Noubhani and Cappetta 1997; *Myliodasyatis* Noubhani and Cappetta 1997; *Phosphatodon* Cappetta 2012; *Meridiana* Case 1994). Among them, only *Heterobatis* can be reasonably compared to new genus, the other genera largely differing both in size and tooth morphology. The teeth of *Heterobatis* are also relatively small (until 3mm tooth size) with a flat labial face sharing ornamentation of enameloid. The anterior teeth display a clear dasyatid pattern with a crown face and a root directed lingually whereas the lateral teeth are sometimes very large, of polyhaulacorhizous type, and remind myliobatids.

 The teeth of *Microtriftis* nov. gen. recall those of another Paleocene dasyatoid *incertae sedis*, *Delpitia* Noubhani and Cappetta 1997, with similar sexual heterodonty and ornamentation of the labial face. However, *Delpitia* is larger and possess a typical dasyatid root, directed lingually with lobes enlarged basally. No evidence of interconnecting/joining teeth is visible on type species *D. reticulata* (Noubhani and Cappetta 1997: Pl. 68, fig. 1-6). The minute new taxon is only recorded in Ypresian to Lower Lutetian marine deposits of Central and Northeastern Senegal (Sine-Saloum region and Matam) even if some early-middle Eocene specimens collected in well drilling from Guinea Bissau (Prian, 1986, 1987) are very similar (HC pers. observ.).

- FIG.3 here
-

NEOSELACHII Compagno, 1977 *incertae ordinis*

Odontorhytis Böhm 1926

*Type species***:** *O. pappenheimi* Böhm, 1926

 This enigmatical taxon is commonly recovered with other elasmobranchs in Eocene nearshore marine deposits of Africa and central Asia. Considering its uncertain systematic position among sharks, authors generally report its occurrence with some caution. Whatever the final outcome on systematic issues, this taxon could be a significant part of the Eocene neashore elasmobranch assemblages. Inappropriately attributed to a percoid fish or to a rajid by Priem (1905; 1915 respectively) based on middle-late Eocene Egyptian specimens, Böhm (1926) described similar teeth from Namibia as *Odontorhytis pappenheimi*, but also considered that it belongs to bony fish of family Lophiidae. Cappetta (1981) definitively attributed this taxon to an undetermined elasmobranch and figurated late Ypresian specimens from Ouled Abdoun (Morocco) that he attributed to the unique species *O. pappenheimi* of Böhm. However, and as noted Cappetta (2012 : 326), clear differences can be noted between Thanetian-Ypresian teeth on one side and middle-late Eocene teeth (e.g. *O. pappenheimi* Böhm, 1926) on the other side, asserting than late Ypresian material from Morocco and several other ones, must be likely considered as representatives of a distinct and unnamed species (Cappetta, 1987, 2012)

- *Odontorhytis priemi* nov. sp.
- Rajidae or Scylliidae Priem (1909: fig. 31-35)
- *Odontorhytis pappenheimi* Cappetta (1981 plate 1 fig. 2-4)
- *Odontorhytis* sp*.* Noubhani and Cappetta (1997: text only)
- *Odontorhytis* sp. Gheerbrant et al. (1993: text only)
- *Odontorhytis* sp. Cappetta (2012: fig. 313: C-E)
- *Odontorhytis* sp. Sambou et al. (2017: text only, in part)

 *Type material***:** Holotype: fig. 4 C, MAM-19 (well F0, 9.18-9.30 m, Unit 2 in Sambou et al., 2017) – Ndendouri-Ouali Diala, ?Early Ypresian, Matam Fm. ; Paratypes: MAM-18, MAM- 20 (well F3, Unit 2 in Sambou et al., 2017) Ndendouri-Ouali Diala, ?Early Ypresian, Matam Fm., Senegal.

 *Additional material***:** Sixteen teeth, partially broken, were mainly found in Thanetian (MAM- 22 to 24, well P53, Unit 0 in Sambou et al., 2017) and Ypresian levels of Ndendouri-Ouali Diala (Units 1-3 in Sambou et al., 2017). We can include those from Morocco: MYB8-9, DYD1-2 from Mera-el-arech (sillon A-B) and Recette 4 (Sillon D), Ouled Abdoun, Morocco – Late Ypresian, figured in Cappetta (1981 plate 1 fig. 2-4) and Cappetta (2012: fig. 313: C-E).

 *Etymology***:** in honours to the French paleoichtyologist F. Priem (1857-1919) who firstly signalled material attributed to this new species.

 Diagnosis: New species of *Odontorhytis* known by isolated teeth exhibiting reduced heterodonty. The new species is mainly characterized by teeth compressed mesio-distally, a fine cusp of sigmoidal profile with lateral cutting edges and marked enameloid folds on both face, a pair of very low lateral cusplets sometimes reduced to minute bulges in small specimens or simply lacking in the largest ones, the lack of lingual barb near the apex, and the lack of a sharp median crest on the crown labial face. The root is of cordiform outline in lingual or labial view, ever larger than crown with large median groove entirely developed on the root depth.

 *Occurrences***:** from Thanetian to Ypresian of North and Western Africa (e.g. Mali, Senegal, Morocco and Tunisia).

 *Description***:** The variation of the tooth morphology in this elasmobranch genus is weak and only involved the presence/development of lateral cusplets compared to the overall/absolute size, which prevents any inference on possible developmental heterodonty. The teeth are small to medium in size and laterally compressed. The holotype is of medium size, probably from one of the anterior files (Fig. 4 C). The weakly enamelled cusp is high, pointed, and of sigmoid profile (fig. 4 C2). The crown labial face is rather flat towards the apex of cusp but convex at its base where several fine folds run on its lower half. It extends downward, overlapping the root by an expansion, high and narrow, more or less pinched laterally (fig. 4 C1) that may bear a sagittal keel on the probably more anterior teeth (fig. 4 B1, H1). The lingual face of crown is strongly concave and bears parallel and well-marked vertical folds (fig. 4 C3). These folds rarely reach the apex of cusp but a short, sharp medio-sagittal crest is sometimes observable near the apex in some teeth (fig. 4 E1). The lateral cutting edges are well marked on the cusp but do not reach the base of the crown on many teeth, holotype included (fig. 4 C2). At the base of the cusp, a pair of very low lateral denticles is usually visible near the onset of the cutting edges (fig. 4 F-H), one of the denticles can however disappear in some teeth (fig. 4 D) and could be reduced to tiny or inconspicuous bulges on more largest teeth, holotype included (fig. 4 B-C). The root is of cordiform outline in lingual view (fig. 4 B2, C3, E1), without well-differentiated lobes. In labial view, it extends well beyond the crown. Its basal face is very concave in profile (fig. 4 B3, C2, E2); it has a well- marked lingual protuberance below the crown and displays an axial nutritive groove on all surface (fig. 4 C3). This groove is relatively deep at the level of the protuberance where a foramen opens (fig. 4 C3). Usually, the basal face lifts slightly on the edges of the groove (fig. 4 E1). Narrow small elliptical foramina open laterally on the labial surface of the root, near 449 the crown-root boundary (fig. 4 C1). Unfortunately, the Thanetian material (Fig 4 F-H) is poorly preserved in comparison to the Ypresian sample (fig. 4 B-E). However, the root

 morphology of the Thanetian teeth seemly resemble those of Ypresian teeth, but they are smaller, more often cuspidate, and their root outline seems sharper and less thick in profile. *Discussion***:** Compared to the middle-late Eocene representatives of genus (including *O. pappenheimi, O. bahariensis* Salame and Asan, 2019 and possible other unnamed species in Tabuce et al., 2005), the cusp is thinner, strongly bent lingually and of sigmoidal profile in *O. priemi* nov. sp. The lateral cutting edges are thin but distinct and always present in *O. priemi* nov. sp. They do not reach the crown base. A pair of very low and sharp lateral cusplets is sometimes present at the base of the cusp, near the beginning of cutting edges. These cusplets can be weak and sometimes be reduced to minute bulges or even absent in some teeth. The crown lingual face bears numerous fine folds that do not reach the apex. Compared to *O. priemi* nov. sp., a barb may be present on the lingual side of the apex in the middle-late Eocene specimens (e.g. *O. pappeheimi, O. bahariensis*). In the latter, the crown is more strongly mesio-distally compressed and slightly curved distally in labial view. No lateral cutting edge is visible on the cusp nor lateral cusplets at the base of crown. On the other side, an unusual distinct and sharp median crest is present on the cusp labial face, no crest –but similar fine folds– is present on the lingual face, which is transversally convex as observed in *O. priemi* nov. sp.

 In both group, the root is high with a heart-like outline (particularly in *O. priemi* nov. sp.) but without differentiated lobes. The profile of its basal face is very concave; it shows a well- marked lingual protuberance and bears an axial furrow on its whole surface. This furrow is much deeper at the end of protuberance where some foramina can be observed. The two groups of species differ from each other only by the morphology of the root (as the crown), which is mesio-distally compressed in *O. pappeheimi* and *O. bahariensis* compared to *O. priemi* nov. sp. There is an overlap in tooth size between *O. priemi* nov. sp. and *O. pappenheimi/ O. bahariensis* but the Thanetian-Ypresian species rarely reach more than 7mm in height whereas the middle-late Eocene specimens can easily measure more than 1cm in height (especially in *O. pappenheimi)*. Distinction between *O. pappenheimi* and *O. bahariensis* remains unclear. Without diagnosis, the latter seems to have smaller gracile teeth than in *O. pappenheimi*, with crown relatively higher with shorter labial cutting edge (Salame and Asan, 2019). Ignoring the ontogenetic heterodonty of this ambiguous neoselachian, *O. bahariensis* could in fact represent the youngest specimens of *O. pappenheimi*, usually recovered in the same areas and ages. The new species *O. priemi* nov. sp. is also present in the Ypresian of Mali (Cappetta 1987), Thanetian-Ypresian of Ouarzazate and phosphate of Ouled Abdoun, Morocco (Cappetta, 1981; Gheerbrant et al., 1993; Noubhani and Cappetta, 1997), and in phosphates of Redeyef, Tunisia (Priem, 1909). The second morphotype (including *O. pappenheimi* and ?*O. bahariensis*) also occurs in Senegalese deposits (Fig. 4 A) and was recovered in the top level of the Matam phosphate deposits (wells P1A, P1B, R4, I1 et G0 see Sambou et al. 2017) and in Sanghaye Bele water well, Sine-Saloum region. This morphotype is recovered elsewhere in the middle Lutetian of Morocco (Tabuce et al., 2005), in the late Lutetian-early Bartonian of Egypt (Strougo et al., 2007, Underwood et al., 2011, Salame and Asan, 2019) but seems more frequent in the late middle Eocene–late Eocene from southwestern Morocco (Adnet et al., 2010), Tunisia (Merzeraux et al., 2017; Sweydan et al., 2018), Egypt (Priem, 1905 ; Case and Cappetta, 1990 ; Murray et al., 2010; Adnet et al., 2011 ; Underwood et al., 2011 ; Murray et al., 2014), Namibia (Böhm 1926), and Pakistan (Case and West, 1991). Its occurrence in the early Rupelian of Jebel Qatrani–Quarry E, Fayum (Murray et al., 2014) is based on a unique broken tooth and must be confirmed by additional samples. These two species do not overlap in the fossil record, suggesting a replacement of taxa during the late Ypresian or near the Ypresian/Lutetian boundary. However, the youngest representatives of *O. priemi* nov. sp. from Matam (e.g. fig. 4 B) show a seemly transitional morphology toward those of "*O. pappenheimi* group" in having larger

 teeth, more compressed mesio-distally with no lateral cusplets. Murray et al. (2010) suggested that *Odontorythis pappenheimi* as the other elasmobranch elements from BQ-2 locality (Fayum, Egypt) probably represent freshwater-tolerant species. The co-occurrence of *Odontorhytis* teeth with mammals remains in different localities (e.g. Gheerbrant et al., 1993; Tabuce et al., 2005; Merzeraux et al., 2016; Murray et al., 2010, 2014; Tabuce et al., 2019) seems to confirm this ecological abilities even if the other listed occurrences, including those of Matam or Morocco, testify that it is firstly a marine elasmobranch.

FIG.4 here

Paleobiological and Paleoecological inferences.

 Among Potamotrygonidae, which also include the marine Styracurinae (Carvalho et al., 2016), the sister subfamily Potamotrygoninae (South American river stingrays) is the most diverse and famous group of obligate freshwater stingrays. South American fossils of this subfamily drew particular attention this last decade (e.g. Brito and Deynat, 2004; Adnet et al., 2014; Chabain et al., 2017) with evidences that they were abundant in middle Eocene deposits of Peruvian Amazonia. In contrast, the living styracurines (amphi-American whipray) are less abundant. They are currently frequent in the Central eastern Pacific (*Styracura pacifica*) and Central western Atlantic (*S. schmardae*) shallow waters until the Amazon river mount southward (Last et al., 2016; Bloom and Lovejoy 2017). No fossil representatives of Styracurine (or formally *Himantura*) are known in fossiliferous deposits along the Central Eastern Pacific or Central Western Atlantic coasts (e.g. Landini et al., 2017; Carrillo-Bricenio et al., 2014, 2015, 2016, 2018). However, the styracurines likely inhabited the Eastern Pacific–Caribbean zone since the middle Eocene, which corresponds to the oldest occurrence of their sister taxa, the potamotrygonins (Adnet et al., 2014). The attribution of *Atlantitrygon*

 to Styracurinae is based on the similar overall shape of the teeth and the presence of a displaced main cusp. Further evidences (as coeval records in Caribbean zone) are needed to confirm such a systematic assumption but *Atlantitrygon* (and mainly the oldest species *A. sudrei*) might represent one of the marine–brackish pool species of potamotrygonid. Subsequently, the multiple occurrence of *Atlantitrygon* in Western Africa deposits may indicate that past styracurines were widespread in the Central Atlantic from western to eastern coasts during the warming period of Paleogene.

 Solving the systematic affinities of *Microtriftis* among dasyatoid is out of scope of this study. The minute teeth seem partially interlocked forming the pavement-like teeth suited for crushing or grinding preys. Ornamented enameloid of labial surface in *Microtriftis* could probably strengthen the grater function of pavement-like teeth as observed in many other living and fossil taxa with such a dentition (e.g. *Myliobatis, Rhombodus, Coupatezia, Dasyatis sensu lato, Pastinachus*). Specialization for extreme durophagy, including an extremely robust dentition with pavement-like teeth, is redundant in myliobatiform evolution (e.g. Underwood et al. 2015, 2017). Their teeth can be only partially interlocked to crush preys, or can be strongly interfitted and jointed together thanks to complementary faces of crown to grind hard mollusc as observed in Myliobatidae and Rhinopteridae (grinding type). *Microtriftis* share the same interlocking pattern than many other fossil dasyatoid (e.g. *Aturobatis, Heterobatis, Hypolophodon, Ishaquia, Meridiana, Myliodasyatis, Phosphatodon*), which could also be considered as grinders (high crown, polygonal outline). However, the dentition of *Microtriftis* is unique in its extreme small size. The number of tooth files in *Microtriftis* jaws, as in all grinding taxa with holaulacorhize root (Cappetta 2012), was likely important. If we cannot surely estimate the size of occlusal surface of pavement of both jaws, it unlikely exceeded two centimetres in width considering the individual tooth size (0.5 mm). Living stingrays with interlocking plates of flat, pavement-like teeth are known to grind

 molluscs and other hard-shelled prey. *Microtriftis* then represents one of the smallest stingrays with grinding teeth, which raises the question on the type of preys it could predate on.

 The Thanetian-Ypresian occurrence of the enigmatical *Odontorythis* completes the fossil record of the genus usually known by the middle-late Eocene. The tooth morphology of its oldest species *Odontorythis priemi* nov. sp*.* does not clarify its systematic position among Neoselachii. The presence of lateral cutting edges and additional cusplets is more common in shark teeth compared to the unusual morphology observed in Lutetian-Priabonian representatives, including its sister species *O. pappenheimi.* Observations of transitional forms between these two chronospecies *O. priemi* nov. sp*.* and *O. pappenheimi* and their relatives suggest that they progressively replaced each other during the latest early Eocene in Senegal and elsewhere.

Conclusions.

 The Senegalese Paleogene fauna is significantly rich (see Pascal, 1987 and Sambou et al., 2017 for listing) with numerous other taxa recovered elsewhere. The focus on three new taxa described herein confirms the dating of fossiliferous levels of the phosphate series of Ndendouri-Ouali Diala (Matam) and improves the stratigraphical correlations between the phosphate deposits of Matam area and those of Sine-Saloum region during the Paleogene. The presence of *Odontorhytis* cf. *pappenheimi* and absence of *O. priemi* in the top level of the Matam Fm. (Matam area) and in Lam Lam Fm. in Sine-Saloum regions also confirm that both deposits are penecontemporaneous (see Tabuce et al., 2019). A previous study confirmed that the coquina limestone of the Lam Lam Fm. in Sanghaye Bele is subcontemporaneous with

 those of the clay-limestone unit ending the Matam Fm., and that it is indicative of a littoral environment (Sambou et al. 2017).

 The Gorgol Fm. (Thanetian) in Matam area probably corresponds to neritic deposal as suggested by analysis of the elasmobranch fauna (Sambou et al., 2017) and lithology (Pascal, 1987). The presence of *Atlantitrygon sudrei* in Thanetian deposits of Senegal, Niger and Togo suggests the establishment of an extended marine realm with faunal connection along the past Guinea Gulf. During the lower Eocene period, the Senegal basins underwent a succession of transgressions and regressions that install the mesobathyal and neritic deposits in the southwestern region (Thiam, 2014; Thiam and Sarr, 2014). The Ypresian marine deposits in the Matam area (lower part of Matam Fm.) are mainly dominated by limestones, clays, and phosphates deposited during the transgression phases that largely extended the marine area to the eastern edge of the Senegal basin (Monciardini and Pascal, 1966). A preliminary study of the elasmobranch fauna (see Sambou et al., 2017) enabled to reconstruct a continuous sea water communication with the large and deeper southermost gulf (Slansky, 1980) to the North Gambia (Miyouna et al., 2009). Ostracods also testify that this inner platform was flooded by relative warm to temperate sea waters (Thiam and Sarr, 2018). Presence of *Atlantitrygon senegalensis* nov. sp. and *Odontorhytis priemi* nov. sp. in subcontemporaneous deposits of Mali and Morocco also suggests wider faunal connection along the Central Western Atlantic coasts during the Ypresian period. Coupled with some Ypresian tectonics (Miyouna et al 2009), the decrease of sea level began at the end of Ypresian as previously reported in Casamance area (Ly AND Anglada, 1991) and in the Mbour-Joal area (Western Senegal; Sarr 1999). The strong decrease in selachian diversity recovered in the higher clay-limestone unit (top of Matam Fm.) coupled with the persistence of only small coastal species are clearly indicative of a more littoral environment in Matam (Sambou et al., 2017) with contemporaneous development of anoxic conditions in southwestern plateform (Thiam and

Sarr, 2018). However, the co-occurrence of *Atlantitrygon senegalensis* nov. sp. and

Odontorhytis cf. *pappenheimi* in the early Lutetian limestones of both localities also indicates

the persistence of a relictual Est-West marine connection in central Senegal (between Matam

and Sine-Saloum areas) during the early Lutetian regional regression (Thiam, 2014 ; Thiam

and Sarr, 2014, 2018)

Acknowledgements

We thank the Department of Geology of the University Cheikh-Anta-Diop of Dakar,

especially Professor El Hadji Sow (Head of Department). We are very grateful to Eugène N.

Faye (Director of the SERPM, Société d'Etudes et de Réalisation des Phosphates de Matam)

and El Hadji M. Fall (Director of MINTECH international) for providing full access to the

quarry of Matam. . We are also grateful to Guillaume Guinot (University of Montpellier) for

the data concerning the Paleocene record of Togo, Elisabeth Cabirou (University of

Montpellier) for treatment of tooth illustrations, Giuseppe Marramà and anonymous reviewer

for improving this work. This research is part of the PaleoSen project (www.paleosen.com),

and was supported by the French ANR-PALASIAFRICA Program (ANR-08-JCJC-0017),

PICS-CNRS, the International Exchange Scheme of the Royal Society, and the National

Geographic Society's Global Exploration Fund (Northern Europe), and the Cooperation and

Cultural Action Service of French Embassy in Senegal (N°2019-0371853/SCAC/ESRI –

Programme d'Appui à la Recherche France Sénégal). Lionel Hautier also gratefully

acknowledges Sidney Sussex College (Cambridge, UK).

References

-
- Adnet, S., Salas Gismondi, R., Antoine, P. O., 2014. Comparisons of Dental Morphology in
- River Stingrays (Chondrichthyes: Potamotrygonidae) with New Fossils from the Middle
- Eocene of Peruvian Amazonia Rekindle Debate on Their Evolution. Naturwissenschaften 101 (1), 33–45.
-
- Arambourg, C., 1952. Les Vertébrés Fossiles Des Gisements de Phosphates (Maroc-Algérie-

Tunisie). Notes et Mémoires du Service géologique du Maroc 92, 1–372.

-
- Beebe, W., Tee-Van, J., 1941. Eastern Pacific expeditions of the New York Zoological
- Society. XXVIII. Fishes from the tropical eastern Pacific. Part 3. Rays, Mantas and
- Chimaeras. Zoologica, Scientific Contributions of the New York Zoological Society 26 (3), 245–280.
-
- Bloom, D. D., Lovejo, N. R., 2017. On the Origins of Marine-Derived Freshwater Fishes in South America, Journal of Biogeography 44 (9), 1927–1938.
-
- Böhm, J., 1926. Über tertiäre Versteinerungen von den Bogenfelsen diamantfeldern, In E.
- Kaiser (Edt.), *Die Diamantwüste Südwestafrikas*. Berlin. pp. 55–87.
-
- Brito, P. M., Deynat, P. P., 2004. Freshwater Stingrays from the Miocene of South America
- with Comments on the Rise of Potamotrygonids (Batoids, Myliobatiformes), In *Recent*

 advances in the Origin and Early Radiation of Vertebrates, verlag Dr. Friedrich Pfeil: München, Germany, pp 575–582.

- Cappetta, H., 1972. Les Poissons Crétacés et Tertiaires Du Bassin Des Iullemmeden
- (République du Niger). Palaeovertebrata 5 (5), 179–251.
-
- Cappetta, H., 1980. Modification Du Statut Générique de Quelques Espèces de Sélaciens Crétacés et Tertiaires. Palaeovertebrata 10 (1), 29–42.
-
- Cappetta, H., 1981. Additions a La Faune de Sélaciens Fossiles Du Maroc. 1: Sur La Présence
- Des Genres Heptranchias, Alopias et Odontorhytis Dans l'Yprésien Des Ouled Abdoun.
- Géobios 14 (5), 563–575.
-
- Cappetta, H., 1987. Chondrichthyes II Mesozoic and Cenozoic Elasmobranchii. Schultze, H.-
- P. (Edt.) *Handbook of Paleoichthyology,* Gustav Fischer Verlag: Stuttgart-New York. 193 p.

- Cappetta, H., 2012. Chondrichthyes II Mesozoic and Cenozoic Elasmobranchii: Teeth.
- Schultze, H.-P., (Edt.) *Handbook of Paleoichthyology*; verlag Dr. Friedrich Pfeil: Stuttgart-
- New York. 512 p.
-
- Cappetta, H., Traverse, M., 1988. Une Riche Faune de Sélaciens Dans Le Bassin à Phosphate de Kpogamé-Hahotoé (Eocène Moyen Du Togo): Note Préliminaire et Précisions Sur la
- Structure et l'âge du Gisement. Géobios 21 (3), 359–365.
-

- Carrillo-Briceño, J. D., Argyriou, T., Zapata, V., Kindlimann, R., Jaramillo, C. A., 2016. New
- Early Miocene (Aquitanian) Elasmobranchii Assemblage from the La Guajira Peninsula,
- Colombia. Ameghiniana 53 (2), 77–99.
-
- Carrillo-Briceño, J. D., Carrillo, J. D., Aguilera, O. A., Sanchez-Villagra, M. R., 2018. Shark and Ray Diversity in the Tropical America (Neotropics)—an Examination of Environmental
- and Historical Factors Affecting Diversity. PeerJ 6, e5313.
-
- Carvalho, M. R. D., Loboda, T. S., Da Silva, J. P. C. B., 2016. New Subfamily, Styracurinae,
- and New Genus, *Styracura*, for *Himantura Schmardae* (Werner, 1904) and *Himantura*
- *Pacifica* (Beebe and Tee-Van, 1941) (Chondrichthyes: Myliobatiformes). Zootaxa 4175 (3), 201-221.
-
- Case, G. R., 1994. Fossil Fish Remains from the Late Paleocene Tuscahoma and Early
- Eocene Bashi Formations of Meridian, Lauderdale County, Mississippi. Part I. Selachians.
- Palaeontographica Abteilung serie A 230 (4–6), 97–138.
-
- Case, G. R., Cappetta, H., 1990. The Eocene Selachians Fauna from the Fayum Depression in Egypt. Palaeontographica Abteilung Serie A 212, 1–30.
-
- Case, G. R., West, R. M., 1991. Geology and Paleontology of the Eocene Drazinda Shale
- Member of the Khirthar Formation, Central Western Pakistan, Part II Late Eocene Fishes.
- Tertiary Research 12 (3–4), 105–120.
-
- Chabain, J., Antoine, P.-O., Altamirano-Sierra, A. J., Marivaux, L., Pujos, F., Salas Gismondi,
- R., Adnet, S., 2017. Cenozoic Batoids from Contamana (Peruvian Amazonia) with Focus on
- Freshwater Potamotrygonins and Their Paleoenvironmental Significance. Géobios 50 (5),
-
- Claeson, K. M., Sarr, R., Hill, R. V., Sow, E. H., Malou, R., O'Leary, M. A., 2015. New
- Fossil Scombrid (Pelagia: Scombridae) Fishes Preserved as Predator and Prey from the
- Eocene of Senegal. Proceedings of the Academy of Natural Sciences of Philadelphia 164 (1),
- 133–147.

389–400.

- Dartevelle, E., Casier, E., 1943. Les Poissons Fossiles Du Bas-Congo et Des Région Voisines. Annales du Musée du Congo Belge, Série A (Minéralogie Géologie, Paléontologie) 3- 2 (1), 1–200.
-
- Dartevelle, E., Casier, E., 1959. Les Poissons Fossiles Du Bas Congo et Des Région Voisines. Annales du Musée du Congo Belge, Série A (Minéralogie Géologie, Paléontologie) 3- 2 (3), 257–568.
-
- Flicoteaux, R., 1974. Biozonation des formations paléogènes de la retombée orientale du dôme de Ndias (Sénégal occidental). Vth African Colloquium on Micropalaeontology, Addis-Abeba, Ethiopia, 1972, Madrid, Enadimsa p. 349-362.
-
- Guinot, G., Adnet, S., Shimada, K., Underwood, C. J., Siversson, M., Ward, D. J., Kriwet, J.,
- Cappetta, H., 2018. On the Need of Providing Tooth Morphology in Descriptions of Extant
- Elasmobranch Species. Zootaxa 4461 (1), 118-126.
-
- Guinot, G., Amoudji, Y.Z., Kassegne, K.E., Sarr, R., Johnson, A.K.C., Da Costa, P.Y.D.,
- Hautier, L., 2020. Elasmobranchs from the upper Palaeocene of Togo. Géobios, in press.
- https://doi.org/10.1016/j.geobios.2019.12.003.
-
- Gheerbrant, E., Cappetta, H., Feist, M., Jaeger, J. J., Sudre, J., Vianey-Liaud, M., Sige, B.,
- 1993. La Succession Des Faunes de Vertébres d'age Paléocène Supérieur et Éocène Inférieur
- Dans Le Bassin d'Ouarzazate, Maroc. Contexte Géologique, Portée Biogéographique et
- Paléogéographique. Newsletters on Stratigraphy 28 (1), 35–58.
-
- Hautier, L., Sarr, R., Tabuce, R., Lihoreau, F., Adnet, S., Domning, D. P., Samb, M., Hameh,
- P. M., 2012. First Prorastomid Sirenian from Senegal (Western Africa) and the Old World
- Origin of Sea Cows. Journal of Vertebrate Paleontology 32 (5), 1218–1222.
-
- Hautier, L., Sarr, R., Lihoreau, F., Tabuce, R., Marwan Hameh, P., 2014. First record of the family Protocetidae in the Lutetian of Senegal (West Africa). Palaeovertebrata 1-7, 10.18563/pv.38.2.e2
-
- Landini, W., Altamirano-Sierra, A., Collareta, A., Di Celma, C., Urbina, M., Bianucci, G.,
- 2017. The Late Miocene Elasmobranch Assemblage from Cerro Colorado (Pisco Formation,
- Peru). Journal of South American Earth Sciences 73, 168–190.
-
- Last, P. R., De Carvalho, M. R., Corrigan, S., Naylor, G. J., Séret, B., Yang, L., 2016. *The*
- *Rays of the World*, CSIRO Publishing, Melbourne. 40, 790 pp.
-
- Ly, A., Anglada, R., 1991. Le bassin sénégalo-mauritanien dans l'évolution des marges périatlantiques au Tertiaire. Cahier de Micropaléontologie. Nouvelle Série 6 (2), 23-47.

- Merzeraud, G., Essid, E. M., Marzougui, W., Ammar, H. K., Adnet, S., Marivaux, L., Tabuce,
- R., Vianey-Liaud, M., 2016. Stratigraphie et Sédimentologie Des Dépôts Marins et
- Continentaux d'âge Éocène Moyen à Miocène en Tunisie Centrale (Région Du Djebel El

Kébar). Bulletin de la Société Géologique de France 187 (1), 11–25.

-
- Miyouna, T., Sow, E.H., Diop, A., Ndiaye, P.M., Niang Diop, I., 2009. Evolution Morpho- structurale et bathymétrique de la partie Septentrionale du bassin Sénégalais au Paléogène. 763 Journal des Sciences 9 (1), $1 - 16$.
-
- Monciardini, C., Pascal, M., 1966. Recherche de phosphates. Rapport complémentaire. Le phosphate de chaux au Sénégal. Synthèse des résultats acquis, étude paléogéographique et essai prévisionnel. Rapport du BRGM, DAK 66 A8, 31p (1)
-

- Noubhani, A., Cappetta, H., 1995. Batoïdes Nouveaux Ou Peu Connus (Neoselachii:
- Rajiformes, Myliobatiformes) des Phosphates Maastrichtiens et Paléocènes du Maroc.

Belgian Geological Survey, Professional Paper 278*,* 157–183.

- Noubhani, A., Cappetta, H., 1997. Les Orectolobiformes, Carcharhiniformes et
- Myliobatiformes (Elasmobranchii, Neoselachii) Des Bassins à Phosphate Du Maroc
- (Maastrichtien-Lutétien Basal). Systématique, Biostratigraphie, Évolution et Dynamique Des

Faunes. Palaeo Ichthyologica 8, 1–327.

 O'Leary, M.A., Sarr, R., Malou, R., Sow, E.H., Lepre, C., Hill, R.V., 2012. A new fossil amiid from the Eocene of Senegal and the persistence of extinct marine amiids after the Cretaceous–Paleogene boundary. Copeia 4, 603-608.

 Pascal, M. 1987. Nouvelles découvertes de minéralisations phosphatées au Sénégal : les gisements de N'Diendouri-Ouali Diala (Département de Matam, Région du Fleuve), Chronique de la recherche minière 486 : 3-24.

 Prian, J.P., 1986. Géologie de la bordure méridionale du golfe tertiaire de Casamance (Sénégal, Guinée Bissau). Document du BRGM n°101, ISSN 0221-2536, 5-32.

- Prian, J. P., 1987. The Farim-Saliquinhe Eocene phosphate deposits Guinea Bissau, West Africa. In : *Phosphate deposits of the World* Vol. 2 Phosphate rock resources. Notholt, A.J.G,
- Sheldon, R.P., Davidson, D.F. (Edt.), Cambridge University press, pp. 277-283.

- Priem, M. F., 1905. Sur les poissons fossiles de l'Eocène moyen d'Egypte. Extrait du Bulletin de la Société Géologique de France 4 V, 633–641.
-
- Priem, M. F., 1909. Note Sur Les Poissons Fossiles Des Phosphates de Tunisie et d'Algérie.

Extrait du Bulletin de la Société Géologique de France 4 IX*,* 315–324.

-
- Priem, M. F., 1915. Sur Les Vertébrés Du Crétacé et de l'Eocène d'Egypte. Extrait du
- Bulletin de la Société Géologique de France 4 XIV*,* 366–382.

 Roger, J., Noël, B.J., Barusseau, J. P., Serrano, O., Nehlig, P., Duvail, C., 2009. Notice explicative de la carte géologique du Sénégal à 1/500 000, feuilles nord-ouest, nord-est et sud- ouest. Ministère des Mines, de l'Industrie et des PME, Direction des Mines et de la Géologie, Dakar, 61 p.

- Salame, I., Asan, A., 2019. A new *Odontorhytis* species (Chondrichtyes) from the middle Eocene of Elgedida Mine, Bahariya Oasis, Egypt. Egyptian Journal of Geology 63, 407-415.
-

 Sarr, R., 1999. Le Paléogène de la région de Mbour-Joal (Sénégal occidental) : biostratigraphie, étude systématique des ostracodes, paléoenvironnement. Revue de Paléobiologie, Genève 18 (1), 1-29.

Solé, F., Noiret, C., Desmares, D., Adnet, S., Taverne, L., De Putter, T., Mees, F., Yans, J.,

Steeman, T., Louwye, S., Folie, A., Stevens, N.J, Gunnel, G.F., Baudet, D., Kitambala Yaya,

833 N ., Smith, T., 2018. Reassessment of Historical Sections from the Paleogene Marine Margin

of the Congo Basin Reveals an Almost Complete Absence of Danian Deposits. Geoscience

- Frontiers 10 (3), 1039-1063.
-
- Stevens, N. J., Eastman, J. T., Odunze, S. O., Cooper, L. N., Obi, G. C., 2011. Paleocene

Ichthyofauna and Paleoenvironmental Setting, Imo Formation, Southeastern Nigeria. Neues

Jahrbuch für Geologie und Paläontologie Abhteilung 260 (3), 289–296.

- Strömer, E., 1910. Reptilien-Und Fischreste Aus Dem Marinen Alttertiär von Südtogo
- (Westafrika). Monatsbericht der deutschen geologischen Gesellschaft 62 (7), 478–505.

- Underwood, C. J., Ward ,D. J., King, C., Antar, S. M., Zalmout, I. S., Gingerich, P. D., 2011.
- Shark and Ray Faunas in the Middle and Late Eocene of the Fayum Area, Egypt. Proceedings
- of the Geologists' Association 122 (1), 47–66.
-
- Underwood, C. J., Johanson, Z., Welten, M., Metscher, B., Rasch, L. J., Fraser, G. J., Smith,
- M. M., 2015. Development and Evolution of Dentition Pattern and Tooth Order in the Skates
- And Rays (Batoidea; Chondrichthyes). Plos One 10 (4), e0122553.
-
- Underwood, C. J., Kolmann, M. A., Ward, D. J., 2017. Paleogene Origin of Planktivory in the
- Batoidea. Journal of Vertebrate Paleontology 37 (3), e1293068.
-
- Vandenberghe, N., Hilgen, F.J.,Speijer, R.P., 2012. The Paleogene Period. In: Gradstein, F.
- M., Ogg, J. G., Schmitz, M. D., and Ogg, G. M., (Edts.) *The Geological Time Scale* 2012,
- Oxford, Elsevier Science, p. 855-921
-
- Werner, F., 1904. Die Fische der zoologisch-vergleichend-anatomischen Sammlung der
- Wiener Universität. I. Teil. Cyclostomen, Chondropterygier, Ganoiden, Dipnoer. Zoologische
- Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere (Jena) 21 (3), 263– 302.
-
- White, E. I., 1926. Eocene Fishes from Nigeria. Bulletin of the Geological Survey Nigeria 10, 1–82.
-
- White, E. I., 1934. Fossil Fishes of Sokoto Province. Bulletin of the Geological Survey Nigeria 14, 1–78.

 Fig. 1. Location and hypothetical stratigraphic correlation of Senegal Paleogene deposits between studied fossil localities of Sanghaye Bele and Ndendouri-Ouali Diala. Logs and details from Tabuce et *al.* (2019) and Sambou et *al.* (2017), respectively. Black teeth outline indicating levels with elasmobranch remains.

 Fig. 1. Localisation et hypothèse de corrélation stratigraphique des gisements paléogènes du Sénégal entre les localités fossilifères étudiées de Sanghaye Bélé et Ndendouri-Ouali Diala. Logs et détails des coupes issus de Tabuce et *al.* 2019 et Sambou et *al.* 2017, respectivement. Les silhouettes des dents noires indiquent les niveaux avec les restes d'élasmobranches.

 Fig. 2. *Atlantitrygon sudrei* nov. sp.: Thanetian of Sessao, Niger - Fig. A. SES64 (holotype); B. SES65; C. SES62; D. SES63; E. MAM-03 (well P53, Unit 0 in Sambou et *al.*, 2017) ;

- *Atlantitrygon senegalensis* nov. sp.: Fig F-J: MAM-04 to MAM-08; Fig. K: MAM-09
- (holotype); Fig. L-M: MAM-10 to MAM-11 (well F0 9.18-9.3m, Unit 2 in Sambou et *al.*,
- 2017) ?Early Ypresian, Matam Fm. ; Fig. N-Q. SNSAN/2011/01 to SNSAN/2011/04 Lower
- Lutetian, Lam Lam Fm.
- **Fig. 2.** *Atlantitrygon sudrei* nov. sp.: Thanétien de Sessao, Niger Fig. A. SES64 (holotype);
- B. SES65; C. SES62; D. SES63; E. MAM-03 (puits P53, Unité 0 dans Sambou et *al.*, 2017) ;
- *Atlantitrygon senegalensis* nov. sp.: Fig F-J: MAM-04 à MAM-08; Fig. K: MAM-09
- (holotype); Fig. L-M: MAM-10 à MAM-11 (puits F0 9.18-9.3m, Unité 2 dans Sambou et *al.*,
- 2017) ?Yprésien inférieur, Formation de Matam; Fig. N-Q. SNSAN/2011/01 à
- SNSAN/2011/04 Lutétien inférieur, Formation de Lam Lam.

 Fig. 3. *Microtriftis matami* nov. gen. nov. sp. A MAM-12 (well F0 9.18-9.3m, Unit 2 in Sambou et al., 2017) ?Early Ypresian, Matam Fm.; B-C MAM-13 and MAM-14 (well P1, Unit 2 in Sambou et al., 2017) ?Early Ypresian, Matam Fm.; D-E MAM-15 (holotype) and 931 MAM-16 (well F0 9.18-9.3m, Unit 2 in Sambou et al., 2017) ?Early Ypresian, Matam Fm.; F-H SNSAN/2011/05 to SNSAN/2011/07 Lower Lutetian, Lam Lam Fm.

- **Fig. 3.** *Microtriftis matami* nov. gen. nov. sp. A MAM-12 (puits F0 9.18-9.3m, Unité 2 dans
- Sambou et *al.*, 2017) ?Yprésien inférieur, Formation de Matam; B-C MAM-13 et MAM-14
- (puits P1, Unité 2 dans Sambou et *al.*, 2017) ?Yprésien inférieur, Formation de Matam; D-E
- MAM-15 (holotype) et MAM-16 (puits F0 9.18-9.3m, Unité 2 dans Sambou et *al.*, 2017)
- ?Yprésien inférieur, Formation de Matam ; F-H SNSAN/2011/04 à SNSAN/2011/07 Lutétien
- inférieur, Formation de Lam Lam.

Fig. 4. A. *O. cf. pappenheimi*: Lower Lutetian, Lam Lam. Fm., SNSAN/2011/08; B-F.

Odontorhytis priemi nov.sp., : B: MAM-18 from (well F3, Unit 2 in Sambou et al., 2017)

?Early Ypresian, Matam Fm., C: MAM-19 (well F0 9.18-9.3m, Unit 2 in Sambou et al., 2017)

(holotype) ?Early Ypresian, Matam Fm, D: MAM-20 (well F3, Unit 2 in Sambou et al., 2017)

?Early Ypresian, Matam Fm; E: MAM-21 (well P1A, Unit 2 in Sambou et al., 2017) ?Early

Ypresian, Matam Fm ; F-H MAM-22 to MAM-24 (well P53, Unit 0 in Sambou et al., 2017);

Thanetian, Gorgol Fm.

Fig. 4. A. *O. cf. pappenheimi*: Lutétien inférieur, Formation de Lam Lam, SNSAN/2011/08;

B-F. *Odontorhytis priemi* nov.sp., : B: MAM-18 de (puits F3, Unité 2 dans Sambou et *al.*,

2017) ?Yprésien inférieur, Formation de Matam, C: MAM-19 (puits F0 9.18-9.3m, Unité 2

- dans Sambou et *al.*, 2017) (holotype) ?Yprésien inférieur, Formation de Matam, D: MAM-20
- (puits F3, Unité 2 dans Sambou et *al.*, 2017) ?Yprésien inférieur, Formation de Matam; E:
- MAM-21 (puits P1A, Unité 2 dans Sambou et *al.*, 2017) ?Yprésien inférieur, Formation de
- Matam; F-H MAM-22 à MAM-24 (puits P53, Unité 0 dans Sambou et *al.*, 2017); Thanétien,
- Formation de Gorgol.