



HAL
open science

From neglected to dissected: How technological advances are leading the way to the study of *Coxiella burnetii* pathogenesis

Melanie Burette, Matteo Bonazzi

► To cite this version:

Melanie Burette, Matteo Bonazzi. From neglected to dissected: How technological advances are leading the way to the study of *Coxiella burnetii* pathogenesis. *Cellular Microbiology*, 2020, Celebrating the Career and Legacy of Professor Pascale Cossart, 22 (4), pp.e13180. 10.1111/cmi.13180. hal-02993239

HAL Id: hal-02993239

<https://hal.science/hal-02993239>

Submitted on 9 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

From neglected to dissected: how technological advances are leading the way to the study of *Coxiella burnetii* pathogenesis

Melanie Burette¹ & Matteo Bonazzi¹

1. IRIM, UMR 9004 CNRS, Université de Montpellier, Montpellier, France.

For correspondence: matteo.bonazzi@irim.cnrs.fr

Abstract

Coxiella burnetii is an obligate intracellular bacterial pathogen responsible for severe worldwide outbreaks of the zoonosis Q fever. The remarkable resistance to environmental stress, extremely low infectious dose and ease of dissemination, contributed to the classification of *C. burnetii* as a class B biothreat. Unique among intracellular pathogens, *C. burnetii* escapes immune surveillance and replicates within large autophagolysosome-like compartments called *Coxiella*-containing vacuoles (CCVs). The biogenesis of these compartments depends on the subversion of several host signaling pathways. For years, the obligate intracellular nature of *C. burnetii* imposed significant experimental obstacles to the study of its pathogenic traits. With the development of an axenic culture medium in 2009, *C. burnetii* became genetically tractable, thus allowing the implementation of mutagenesis tools and screening approaches to identify its virulence determinants and investigate its complex interaction with host cells. Here, we review the key advances that have contributed to our knowledge of *C. burnetii* pathogenesis, leading to the rise of this once-neglected pathogen to an exceptional organism to study the intravacuolar lifestyle.

Introducing *Coxiella burnetii*

Coxiella burnetii is a Gram negative obligate intracellular pathogen, and the causative agent of Q fever, a worldwide zoonotic disease (Eldin *et al.*, 2017). Animal coxiellosis is mostly associated with abortion, stillbirths, and weak offspring (Fig. 1). Desiccation of placental materials, excretions of birthing fluids, urine, feces and milk from infected animals contribute to shed *C. burnetii* into the environment (Eldin *et al.*, 2017). Transmission to humans results from exposure to contaminated aerosols and dust (Fig. 1). In humans, *C. burnetii* infections are often asymptomatic and self-limiting, however, 40% of individuals infected with *C. burnetii* develop an acute disease which is associated with a flu-like syndrome, pneumonia, hepatitis and chronic fatigue (Eldin *et al.*, 2017) (Fig. 1). Acute disease may convert into a chronic illness with severe complications, including endocarditis (Eldin *et al.*, 2017). An important risk factor for Q fever outbreaks is the multiple zoonotic reservoirs of *C. burnetii*, which include domestic livestock and wild animals (mammals, reptiles, birds and ticks) (Eldin *et al.*, 2017). *C. burnetii* is extremely infectious, with 1-to-10 bacteria being sufficient to cause disease (Brooke *et al.*, 2013). The low infectious dose coupled to remarkable environmental stability contribute to the significant spreading of *C. burnetii* infections well away from the outbreak source and led to the classification of *C. burnetii* as a category B biothreat (Madariaga *et al.*, 2003).

C. burnetii is a stealth pathogen that actively escapes innate immune recognition by inhibiting the NF- κ B pathway (Mahapatra *et al.*, 2016) and inflammasome activation (Cunha *et al.*, 2015). Infected cells are also protected from apoptosis, thereby preserving the bacterial replicative niche over long periods (Lührmann *et al.*, 2017). *C. burnetii* enters macrophages by phagocytosis through the interaction with α V β 3 integrins (Capo *et al.*, 1999). In contrast, in non-phagocytic cells,

internalization is facilitated by the invasin OmpA (Outer membrane protein A) (Martinez *et al.*, 2014) (Fig. 1). Following internalization, bacteria reside within early endosomal compartments, also called early CCVs, that passively mature along the endocytic pathway by successive fusion events with early and late endosomes and lysosomes (Fig. 1). Maturation is accompanied by the acidification of the CCV lumen (Heinzen *et al.*, 1996), which is required to activate bacterial metabolism (Hackstadt and Williams, 1981) and the translocation of bacterial effector proteins by a Dot/Icm Type 4b Secretion System (T4SS) (Newton *et al.*, 2013). Thus, by 48 hours post-infection, cells display a single, large, mature CCV, where markers of multiple membrane compartments co-exist (Dragan and Voth, 2019), which is indicative of the capacity of *C. burnetii* to hijack multiple host membrane trafficking pathways (Fig. 1). Importantly, the biogenesis of these remarkable compartments is a two-step process requiring both cellular and bacterial factors. This review will focus on the recent technological advances that fostered remarkable progress in our understanding of the complex host/pathogen interplay that controls the generation of the *C. burnetii* replicative niche.

The long and winding road (to genetic manipulation)

Together with its high infectivity, the obligate intracellular nature of *C. burnetii* has imposed severe constraints on the study of its pathogenesis. An important step towards the development of tools to investigate *C. burnetii* infection was the isolation of a Phase II variant (NMII) of the Nine Mile Phase I strain (NMI), presenting a single deletion of 21 genes involved in lipopolysaccharide biosynthesis and displaying loss-of-virulence phenotype in animal models (Moos and Hackstadt, 1987; Hoover *et al.*, 2002). Thus, NMII was allowed for manipulation in biosafety level-2 (BSL-2) environments (Fig. 2D), as opposed to other *C. burnetii* strains that require

manipulation in biosafety level-3 (BSL-3) confinement. Importantly, loss-of-virulence of NMII has no impact either on intracellular growth kinetics or development of CCVs in cultured cells, making NMII a relevant model to study host-pathogen interactions (Howe *et al.*, 2010).

Nevertheless, bacterial amplification of *C. burnetii* in embryonated eggs and/or cultured cells represented a real obstacle for its genetic manipulation. Indeed, transposon mutagenesis allowed the generation of *C. burnetii* mutants (Beare *et al.*, 2009); however, amplification and isolation of these from infected cells ruled out the possibility of isolating mutations in virulence genes. Thus, until 2011, *ftsZ* was the only *C. burnetii* gene characterized using Himar1 transposon mutagenesis (Beare *et al.*, 2009) (Fig. 2F).

Whole-genome sequencing of *C. burnetii* NMI RSA493 in 2003 allowed the identification of typical features, including a Dot/Icm secretion system, which is highly homologous to that of *L. pneumophila* (Seshadri *et al.*, 2003) (Fig. 2C). This finding was pivotal to develop bioinformatics-based approaches to predict effector proteins-coding genes. These revealed that similar to *L. pneumophila*, effector proteins translocated by *C. burnetii* encode a C-terminal secretion signal called the E-block motif and are mostly under the control of a PmrA response regulator (Zusman *et al.*, 2007). Due to the genetic intractability of *C. burnetii* at that time, candidate effector proteins were tested for secretion using *L. pneumophila* as a surrogate model, exploiting the homologies between the Dot/Icm secretion systems of the two pathogens. This approach allowed the Roy laboratory to validate the translocation of four *C. burnetii* effector proteins encoding ankyrin repeat homology domains (ARHDs) (Pan *et al.*, 2008) (Fig. 2E). Later work from the Samuel laboratory reported the first

large-scale, bioinformatics-based identification of *C. burnetii* effector proteins and used the *L. pneumophila* system to validate the secretion of 32 new T4SS substrates (Chen *et al.*, 2010) (Fig. 2H). Moreover, using a shuttle plasmid system for the expression of recombinant proteins in *C. burnetii*, Chen and colleagues also demonstrated that the functionality of the Dot/Icm secretion system (Chen *et al.*, 2010).

One year later, a genetic screen of *C. burnetii* proteins carrying C-terminal secretion signals led to the identification of additional effector proteins (Carey *et al.*, 2011) (Fig. 2I). For this study, wild type (wt) *L. pneumophila* or the T4SS-defective $\Delta dotA$ mutant were transformed with a library containing adenylate cyclase enzyme (CyaA)-tagged random fragments of *C. burnetii* genome, leading to the validation of 7 additional *C. burnetii* effectors (Carey *et al.*, 2011). These were further tested for their intracellular localization and function, which indicated a possible implication in the manipulation of host membrane trafficking. Of note, this is the first study to report a replication phenotype associated with a transposon insertion in the *C. burnetii* Dot/Icm gene *icmL* (*icmL*::Tn), thus demonstrating the importance of *C. burnetii* effector protein translocation for infection (Carey *et al.*, 2011) (Fig. 2I). Today, the development of tailored bioinformatics algorithms for the identification of T4SS effector proteins led to the identification of over 140 candidate *C. burnetii* T4SS effectors (Voth *et al.*, 2009; Chen *et al.*, 2010; Carey *et al.*, 2011; Maturana *et al.*, 2013; Weber *et al.*, 2013). Recent genome comparison of the *C. burnetii* strains NMII RSA493, Henzerling RSA331, G Q212, K Q154 and Dugway 5J108-111, revealed that many genes encoding candidate effector proteins are either pseudogenized or missing altogether, leaving only 44 out of the 143 identified effector protein-coding genes intact across all strains (Larson *et al.*, 2016). Despite these significant advances, functional

analysis of *C. burnetii* genes involved in virulence was still limited by the genetic intractability of this pathogen.

A game-changer in the study of *C. burnetii* infections has been the development of a synthetic medium (ACCM for Acidified Citrate Cysteine Medium), followed by the development of ACCM-2, allowing the extracellular culture of this obligate intracellular bacterium (Omsland *et al.*, 2009; Omsland *et al.*, 2011) (Fig. 2G). ACCM-2 development stems from metabolic requirement studies using microarrays, genomic reconstruction of metabolic pathways and metabolite typing (Omsland *et al.*, 2009; Omsland *et al.*, 2011). These studies highlighted that *C. burnetii* requires acid activation buffer (pH 4.75), to reproduce the CCV microenvironment (Heinzen *et al.*, 1996) and increase its metabolic potential as well as low oxygen levels (2.5%) to facilitate the bacterial micro-aerophilic respiration (Omsland *et al.*, 2009; Omsland *et al.*, 2011). This scientific milestone finally enabled genetic manipulation of *C. burnetii*, leading to a new era in our understanding of its pathogenesis (Fig. 2).

From genes to function: characterization of *C. burnetii* effectors involved in vacuole biogenesis

With the development of axenic culture, targeted deletion of *C. burnetii* was formally achievable (albeit remaining extremely challenging), and the role of *C. burnetii* effector proteins in infection could be tested directly. The Heinzen laboratory first exploited axenic culture with the generation of a *dotA* deletion mutant (Beare *et al.*, 2012) (Fig. 2J) and with the identification and characterization of CvpA (for *Coxiella* vacuolar protein A). This effector protein localizes at CCVs and reroutes recycling endosomes to this compartment by interacting with the clathrin adaptor protein AP2 (Larson *et al.*, 2013) (Fig. 1). In a follow-up study, the Heinzen laboratory identified four

additional members of what constitutes today the Cvp sub-class of *C. burnetii* effector proteins (CvpA, CvpB/Cig2, and CvpC-to-E, Fig. 1) (Larson *et al.*, 2015). Accordingly, targeted deletion of *C. burnetii* vacuolar proteins severely affects the biogenesis of CCVs (Larson *et al.*, 2013; Larson *et al.*, 2015).

If targeted deletion of *C. burnetii* genes remains challenging, *Himar1*-based transposon mutagenesis has been extensively applied to the generation of libraries of *C. burnetii* mutants. Thus, by 2013, the Samuel laboratory reported the first study that combined bioinformatics-mediated identification of candidate effector proteins with transposon mutagenesis (Weber *et al.*, 2013) (Fig. 2N). Searching the *C. burnetii* genome for T4SS features (PmrA consensus sequences, E-block motifs, and homologies with known effectors), led to the identification of 234 genes encoding putative *C. burnetii* effector proteins. β -lactamase translocation assay in *L. pneumophila* validated 53 T4SS substrates, most of which were never reported before. Transposon mutagenesis showed that ten effector proteins were involved in the biogenesis of CCVs and bacterial replication (Weber *et al.*, 2013) (Fig. 2N).

The following year, two independent studies reported the large-scale identification of *C. burnetii* virulence determinants based on the generation of two libraries of *C. burnetii* transposon mutants (Martinez *et al.*, 2014; Newton *et al.*, 2014) (Fig. 2O, R). The Bonazzi library consisted of 3000 GFP-tagged mutants, among which over 1000 were sequenced, annotated, and screened using quantitative, multiparametric image analysis to identify bacterial factors involved in host cell invasion, intracellular replication and persistence (Martinez *et al.*, 2014; Martinez *et al.*, 2015). This approach allowed to validate the function of 16 out of the 22 genes constituting the *C. burnetii* Dot/Icm secretion system, characterize the phenotype associated with transposon insertions in 31 genes encoding effector proteins and

identify the first *C. burnetii* invasin OmpA (for Outer membrane protein A), which is necessary and sufficient to trigger internalization by non-phagocytic cells (Martinez *et al.*, 2014) (Fig. 1 & 2P). This study was also the first to report the use of the insect model *Galleria mellonella* to investigate *C. burnetii* virulence *in vivo* (Martinez *et al.*, 2014).

The Roy laboratory used a modified-Himar1 expressing mCherry fluorescent protein to generate over 3200 mutants. These were visually screened in HeLa cells to identify genes important in the biogenesis of CCVs and bacterial replication using the lysosomal marker LAMP-1 to visualize the membranes of the *C. burnetii* replicative compartment (Newton *et al.*, 2014) (Fig. 2O). This approach allowed the isolation of mutants characterized by different intracellular phenotypes, including defects in intracellular replication and homotypic fusion of independent CCVs or the appearance of filamentous bacteria. Of note, mutants carrying a transposon insertion in gene encoding *CBU1751 (cig57)* displayed a severe vacuole biogenesis defect (Newton *et al.*, 2014). The Newton laboratory has further characterized the role of this effector protein in the development of CCVs and intracellular replication, showing that Cig57 subverts clathrin-mediated traffic by interacting with FCHO2, an accessory protein of clathrin-coated pits (Latomanski *et al.*, 2016) (Fig. 1). More recently, CTLC (clathrin heavy chain) has been observed at CCVs, where it plays an essential role in vacuole expansion (Latomanski and Newton, 2018). Importantly, CTLC recruitment to the CCV is related to autophagy, and conversely, the fusion of autophagosomes with CCVs is dependent on CTLC (Latomanski and Newton, 2018). The Roy visual screen also highlighted a multivacuolar phenotype associated with transposon insertions in the gene *CBU0021 (cig2)*, identical to that previously observed following the knockdown of autophagy-related genes (McDonough *et al.*, 2013), suggesting a functional link

between a bacterial effector protein and the manipulation of a specific host cell function (Fig. 2O). This hypothesis was further investigated, demonstrating that Cig2 contributes to the recruitment of the autophagy machinery to CCVs (Fig. 1), thus facilitating their homotypic fusion and contributing to an enhanced bacterial virulence in the *Galleria mellonella* infection model (Kohler *et al.*, 2016).

It takes two to tango: role of host cell pathways in *C. burnetii* vacuole biogenesis

If on one hand effector proteins translocation is critical for pathogenesis, host cell proteins, lipids, and membrane trafficking pathways also play a significant role in *C. burnetii* infections. As mentioned above, the maturation of CCVs along the endocytic pathway is a pre-requisite for effector proteins translocation (Newton *et al.*, 2013)(Fig. 2L); thus, Rab GTPases of the endocytic pathway play a key role in *C. burnetii* intracellular replication (Fig. 1) (Beron *et al.*, 2002; Romano *et al.*, 2007; Campoy *et al.*, 2011). Indeed, silencing of either Rab5 or Rab7 correlates with effector protein translocation defects (Newton *et al.*, 2016) (Fig. 2R).

A comprehensive characterization of the host cell components required for the biogenesis of CCVs was provided for the first time by a genome-wide screen using siRNA targeting eukaryotic genes in *C. burnetii*-infected HeLa cells (McDonough *et al.*, 2013) (Fig. 2M). Host determinants required for *C. burnetii* infection were identified by the analysis of the number and size of CCVs, revealing the importance of several eukaryotic pathways in critical infection events. As expected, the silencing of genes encoding pH-regulating proteins CLN3 and CLCN5, as well as components of the vacuolar ATPase resulted in a defect in the biogenesis of CCVs (McDonough *et al.*, 2013). Besides, the depletion of the retromer cargo complex VPS26-VPS29-VPS35 leads to defective in bacterial replication, revealing a role for retrograde membrane

trafficking in *C. burnetii* infections (McDonough *et al.*, 2013). Interestingly, seminal work from the Colombo laboratory in the past decade demonstrated that the induction of autophagy favors the biogenesis of CCVs and that *C. burnetii* actively manipulates autophagy during infections (Gutierrez *et al.*, 2005; Romano *et al.*, 2007). However, the precise role of autophagy in CCVs' development remained to be characterized. The genome-wide screening approach of the Roy laboratory revealed that knockdown of autophagy proteins syntaxin-17, ATG5 and ATG12 results in the formation of multiple CCVs of smaller size as compared to control cells (McDonough *et al.*, 2013). It was later demonstrated that this multivacuolar phenotype is the result of the defective homotypic fusion of CCVs (Newton *et al.*, 2014; Martinez *et al.*, 2016).

The Voth and Heinzen laboratories have further explored the complex interplay between *C. burnetii* and autophagy by. The autophagy-associated cargo receptor p62 is actively recruited at CCVs, independently of LC3-interacting domains (Winchell *et al.*, 2018). Interestingly, *C. burnetii* infections seem to preserve p62 from degradation upon induction of autophagy (Winchell *et al.*, 2018). Accordingly, *C. burnetii* infections inhibit mTORC1, a master regulator of autophagy, by a non-canonical mechanism that does not result in accelerated autophagy, nor a block of the autophagic flux (Larson *et al.*, 2019). Finally, autophagy is also involved in repairing the membranes of CCVs during expansion, which are subject to transitory damage and loss-of-acidification (Mansilla Pareja *et al.*, 2017) (Fig. 2T).

Besides host cell proteins, lipids also play a significant role in the development of CCVs, which are rich in sterols (Gilk *et al.*, 2010). Indeed, cholesterol homeostasis regulates CCVs biogenesis and intracellular survival of *C. burnetii* (Gilk *et al.*, 2013; Mulye *et al.*, 2017) (Fig. 2K). Accordingly, *C. burnetii* actively manipulates cholesterol metabolism via a eukaryotic-like $\Delta 24$ sterol reductase (Gilk *et al.*, 2010). Furthermore,

an image-based supervised machine learning approach led to the identification of CvpB/Cig2 as the first *C. burnetii* effector protein to bind phosphoinositides and manipulate their metabolism (Martinez *et al.*, 2016) (Fig. 2S). Indeed, CvpB/Cig2 binds phosphatidylinositol-3-phosphate (PI(3)P) and phosphatidylserine (PS) and perturbs the activity of the PI3-kinase PIKFYVE. This inhibition results in an enrichment of PI(3)P at CCVs, which is essential for their homotypic fusion (Martinez *et al.*, 2016). Importantly, despite the apparent defect in the biogenesis of CCVs, *cvpB* transposon mutants were unaffected in their capacity of replicating within infected cells. However, *cvpB* mutants are attenuated in the *in vivo* models *Galleria mellonella* (Martinez *et al.*, 2016) and SCID mice (van Schaik *et al.*, 2017), demonstrating that the biogenesis of CCVs can modulate *C. burnetii* virulence, independently of bacterial replication.

To infinity and beyond (conclusions)

As of today, *C. burnetii* stands as the sole example of an obligate intracellular bacterial pathogen for which a specific axenic culture medium has been developed. Consequently, research on this dangerous and complex pathogen has bloomed during this decade, with the development of genetic tools and phenotypic screening approaches to better understand the complex interactions established between *C. burnetii* and its host. The generation of libraries of *C. burnetii* transposon mutants combined with the development of multi-parametric screening approaches have allowed the rapid and unbiased identification of microbial genes involved in specific steps of the infectious process. This has been facilitated by the fact that the majority of *C. burnetii* transposon mutants isolated to date display a significant phenotype during infection (Martinez *et al.*, 2014), suggesting a milder functional gene redundancy as compared to the closely related pathogen *L. pneumophila*. Accordingly,

bioinformatics predicted ~ 150 effector proteins compared to the 300 translocated by *Legionella*.

Nevertheless, understanding how these newly identified virulence determinants manipulate host cell functions remains exceptionally challenging. Probably due to the complex nature of this compartment, a large proportion of the *C. burnetii* effector proteins identified and characterized to date are involved in the biogenesis of the CCV; however, proteins involved in the manipulation of other signaling pathways including apoptosis and inflammation have been identified and partially characterized. With the development of advanced bioinformatics approaches such as unsupervised machine learning, hierarchical clustering and Bayesian network analysis, we can now compare microbial-targeted and host-targeted phenotypic screens, to identify sets of bacterial and eukaryotic genes predicted to be involved in the same biological process during infection.

Despite the giant leaps taken since the development of axenic culture, many technical barriers still exist today. Directed mutagenesis remains challenging, limiting our studies to the mutants available in the transposon libraries hosted in the laboratories that undertook this endeavor. Nevertheless, our knowledge of this once-neglected pathogen is ever increasing. This will allow, in the near future, to address burning questions on *C. burnetii* pathogenesis, including a characterization of the strategies used to evade immune recognition, cell-to-cell spread, and, consequently, dissemination to distant organs (heart and liver) following infection of alveolar macrophages.

Acknowledgements

Work in our laboratory is supported by the Agence Nationale de la Recherche (ANR; ANR-14-CE14-0012-01, project AttaQ and ANR- 17-CE15-0021, project QPID) and by the ERA-NET Infect-ERA (ANR-13-IFEC-0003, project EUGENPATH). We apologize to all colleagues whose work was not discussed in this review.

Conflict of interest

The authors declare no conflict of interest.

References

- Beare, P. A, Larson, C.L., Gilk, S.D., and Heinzen, R. A (2012) Two systems for targeted gene deletion in *Coxiella burnetii*. *Appl Environ Microbiol* **78**: 4580–9.
- Beare, P.A., Howe, D., Cockrell, D.C., Omsland, A., Hansen, B., and Heinzen, R.A. (2009) Characterization of a *Coxiella burnetii* *ftsZ* mutant generated by Himar1 transposon mutagenesis. *J Bacteriol* **191**: 1369–1381
- Beron, W., Gutierrez, M.G., Rabinovitch, M., and Colombo, M.I. (2002) *Coxiella burnetii* Localizes in a Rab7-Labeled Compartment with Autophagic Characteristics. *Infect Immun* **70**: 5816–5821
- Brooke, R.J., Kretzschmar, M.E.E., Mutters, N.T., and Teunis, P.F. (2013) Human dose response relation for airborne exposure to *Coxiella burnetii*. *BMC Infect Dis* **13**: 488 BMC Infectious Diseases.
- Campoy, E.M., Zoppino, F.C.M., and Colombo, M.I. (2011) The Early Secretory Pathway Contributes to the Growth of the *Coxiella* -Replicative Niche. *Infect Immun* **79**: 402–413
- Capo, C., Lindberg, F.P., Meconi, S., Zaffran, Y., Tardei, G., Brown, E.J., *et al.*

(1999) Subversion of monocyte functions by coxiella burnetii: impairment of the cross-talk between alphavbeta3 integrin and CR3. *J Immunol* **163**: 6078–85

Carey, K.L., Newton, H.J., Lührmann, A., and Roy, C.R. (2011) The Coxiella burnetii Dot/Icm System Delivers a Unique Repertoire of Type IV Effectors into Host Cells and Is Required for Intracellular Replication. *PLoS Pathog* **7**: e1002056

Chen, C., Banga, S., Mertens, K., Weber, M.M., Gorbasljeva, I., Tan, Y., *et al.* (2010) Large-scale identification and translocation of type IV secretion substrates by Coxiella burnetii. *Proc Natl Acad Sci* **107**: 21755–21760.

Cunha, L.D., Ribeiro, J.M., Fernandes, T.D., Massis, L.M., Khoo, C.A., Moffatt, J.H., *et al.* (2015) Inhibition of inflammasome activation by Coxiella burnetii type IV secretion system effector IcaA. *Nat Commun* **6**: 10205

Dragan, A.L., and Voth, D.E. (2019) Coxiella burnetii: international pathogen of mystery. *Microbes Infect* 1–11

Eldin, C., Mélenotte, C., Mediannikov, O., Ghigo, E., Million, M., Edouard, S., *et al.* (2017) From Q Fever to Coxiella burnetii Infection: a Paradigm Change. *Clin Microbiol Rev* **30**: 115–190

Gilk, S.D., Beare, P.A., and Heinzen, R.A. (2010) Coxiella burnetii expresses a functional $\Delta 24$ sterol reductase. *J Bacteriol* **192**: 6154–6159.

Gilk, S.D., Cockrell, D.C., Luterbach, C., Hansen, B., Knodler, L.A., Ibarra, J.A., *et al.* (2013) Bacterial Colonization of Host Cells in the Absence of Cholesterol. *PLoS Pathog* **9**.

Gutierrez, M.G., Vázquez, C.L., Munafó, D.B., Zoppino, F.C.M., Berón, W., Rabinovitch, M., and Colombo, M.I. (2005) Autophagy induction favours the generation and maturation of the Coxiella-replicative vacuoles. *Cell Microbiol* **7**: 981–

Hackstadt, T., and Williams, J.C. (1981) Biochemical stratagem for obligate parasitism of eukaryotic cells by *Coxiella burnetii*. *Proc Natl Acad Sci* **78**: 3240–3244

Heinzen, R.A., Scidmore, M.A., Rockey, D.D., and Hackstadt, T. (1996) Differential interaction with endocytic and exocytic pathways distinguish parasitophorous vacuoles of *Coxiella burnetii* and *Chlamydia trachomatis*. *Infect Immun* **64**: 796–809.

Hoover, T.A., Culp, D.W., Vodkin, M.H., Williams, J.C., and Thompson, H.A. (2002) Chromosomal DNA Deletions Explain Phenotypic Characteristics of Two Antigenic Variants, Phase II and RSA 514 (Crazy), of the *Coxiella burnetii* Nine Mile Strain. *Infect Immun* **70**: 6726–6733

Howe, D., Shannon, J.G., Winfree, S., Dorward, D.W., and Heinzen, R.A. (2010) *Coxiella burnetii* phase I and II variants replicate with similar kinetics in degradative phagolysosome-like compartments of human macrophages. *Infect Immun* **78**: 3465–3474.

Kohler, L.J., Reed, S.R., Sarraf, S.A., Arteaga, D.D., Newton, H.J., and Roy, C.R. (2016) Effector Protein Cig2 Decreases Host Tolerance of Infection by Directing Constitutive Fusion of Autophagosomes with the *Coxiella* -Containing Vacuole. *MBio* **7**

Larson, C.L., Beare, P.A., Howe, D., and Heinzen, R.A. (2013) *Coxiella burnetii* effector protein subverts clathrin-mediated vesicular trafficking for pathogen vacuole biogenesis. *Proc Natl Acad Sci* **110**: E4770–E4779

Larson, C.L., Beare, P.A., Voth, D.E., Howe, D., Cockrell, D.C., Bastidas, R.J., *et al.* (2015) *Coxiella burnetii* Effector Proteins That Localize to the Parasitophorous Vacuole Membrane Promote Intracellular Replication. *Infect Immun* **83**: 661–670.

Larson, C.L., Martinez, E., Beare, P.A., Jeffrey, B., Heinzen, R.A., and Bonazzi, M. (2016) Right on Q: Genetics begin to unravel *Coxiella burnetii* host cell interactions.

Future Microbiol **11**: 919–939.

Larson, C.L., Sandoz, K.M., Cockrell, D.C., and Heinzen, R.A. (2019) Noncanonical Inhibition of mTORC1 by *Coxiella burnetii* Promotes Replication within a Phagolysosome-Like Vacuole. **10**: 1–16.

Latomanski, E.A., and Newton, H.J. (2018) Interaction between autophagic vesicles and the *Coxiella*-containing vacuole requires CLTC (clathrin heavy chain). *Autophagy* **14**: 1710–1725

Latomanski, E.A., Newton, P., Khoo, C.A., and Newton, H.J. (2016) The Effector Cig57 Hijacks FCHO-Mediated Vesicular Trafficking to Facilitate Intracellular Replication of *Coxiella burnetii*. *PLoS Pathog* **12**: 1–24.

Lührmann, A., Newton, H.J., and Bonazzi, M. (2017) Beginning to understand the role of the Type IV secretion system effector proteins in *coxiella burnetii* pathogenesis. *Curr Top Microbiol Immunol* **413**: 243–268.

Madariaga, M.G., Rezai, K., Trenholme, G.M., and Weinstein, R.A. (2003) Q fever: a biological weapon in your backyard. *Lancet Infect Dis* **3**: 709–721

Mahapatra, S., Gallaher, B., Smith, S.C., Graham, J.G., Voth, D.E., and Shaw, E.I. (2016) *Coxiella burnetii* Employs the Dot/Icm Type IV Secretion System to Modulate Host NF- κ B/RelA Activation. *Front Cell Infect Microbiol* **6**: 1–13.

Mansilla Pareja, M.E., Bongiovanni, A., Lafont, F., and Colombo, M.I. (2017) Alterations of the *Coxiella burnetii* Replicative Vacuole Membrane Integrity and Interplay with the Autophagy Pathway. *Front Cell Infect Microbiol* **7**: 1–17.

Martinez, E., Allombert, J., Cantet, F., Lakhani, A., Yandrapalli, N., Neyret, A., *et al.* (2016) *Coxiella burnetii* effector CvpB modulates phosphoinositide metabolism for optimal vacuole development. *Proc Natl Acad Sci* **113**: E3260–E3269

Martinez, E., Cantet, F., and Bonazzi, M. (2015) Generation and multi-phenotypic

high-content screening of *Coxiella burnetii* transposon mutants. *J Vis Exp* **2015**

Martinez, E., Cantet, F., Fava, L., Norville, I., and Bonazzi, M. (2014) Identification of OmpA, a *Coxiella burnetii* Protein Involved in Host Cell Invasion, by Multi-Phenotypic High-Content Screening. *PLoS Pathog* **10**: e1004013

Maturana, P., Graham, J.G., Sharma, U.M., and Voth, D.E. (2013) Refining the plasmid-encoded type IV secretion system substrate repertoire of *Coxiella burnetii*. *J Bacteriol* .

McDonough, J.A., Newton, H.J., Klum, S., Swiss, R., Agaisse, H., and Roy, C.R. (2013) Host Pathways Important for *Coxiella burnetii* Infection Revealed by Genome-Wide RNA Interference Screening. *MBio* **4**: 1–13

Moos, A., and Hackstadt, T. (1987) Comparative virulence of intra- and interstrain lipopolysaccharide variants of *Coxiella burnetii* in the guinea pig model. *Infect Immun* **55**: 1144–1150.

Mulye, M., Samanta, D., Winfree, S., Heinzen, R.A., and Gilk, S.D. (2017) Elevated cholesterol in the *Coxiella burnetii* intracellular niche is bacteriolytic. *MBio* **8**.

Newton, H.J., Kohler, L.J., McDonough, J.A., Temoche-Diaz, M., Crabill, E., Hartland, E.L., and Roy, C.R. (2014) A Screen of *Coxiella burnetii* Mutants Reveals Important Roles for Dot/Icm Effectors and Host Autophagy in Vacuole Biogenesis. *PLoS Pathog* **10**: e1004286

Newton, H.J., McDonough, J.A., and Roy, C.R. (2013) Effector Protein Translocation by the *Coxiella burnetii* Dot/Icm Type IV Secretion System Requires Endocytic Maturation of the Pathogen-Occupied Vacuole. *PLoS One* **8**: e54566

Newton, P., Latomanski, E.A., and Newton, H.J. (2016) Applying Fluorescence Resonance Energy Transfer (FRET) to Examine Effector Translocation Efficiency by *Coxiella burnetii* during siRNA Silencing. *J Vis Exp*

Omsland, A., Beare, P.A., Hill, J., Cockrell, D.C., Howe, D., Hansen, B., *et al.* (2011) Isolation from Animal Tissue and Genetic Transformation of *Coxiella burnetii* Are Facilitated by an Improved Axenic Growth Medium. *Appl Environ Microbiol* **77**: 3720–3725

Omsland, A., Cockrell, D.C., Howe, D., Fischer, E.R., Virtaneva, K., Sturdevant, D.E., *et al.* (2009) Host cell-free growth of the Q fever bacterium *Coxiella burnetii*. *Proc Natl Acad Sci* **106**: 4430–4434

Pan, X., Lührmann, A., Satoh, A., Laskowski-Arce, M.A., and Roy, C.R. (2008) Ankyrin repeat proteins comprise a diverse family of bacterial type IV effectors. *Science (80-)* **320**: 1651–1654.

Romano, P.S., Gutierrez, M.G., Berón, W., Rabinovitch, M., and Colombo, M.I. (2007) The autophagic pathway is actively modulated by phase II *Coxiella burnetii* to efficiently replicate in the host cell. *Cell Microbiol* **9**: 891–909

Schaik, E.J. van, Case, E.D., Martinez, E., Bonazzi, M., and Samuel, J.E. (2017) The SCID mouse model for identifying virulence determinants in *Coxiella burnetii*. *Front Cell Infect Microbiol* **7**: 1–10.

Seshadri, R., Paulsen, I.T., Eisen, J.A., Read, T.D., Nelson, K.E., Nelson, W.C., *et al.* (2003) Complete genome sequence of the Q-fever pathogen *Coxiella burnetii*. *Proc Natl Acad Sci* **100**: 5455–5460

Voth, D.E., Howe, D., Beare, P.A., Vogel, J.P., Unsworth, N., Samuel, J.E., and Heinzen, R.A. (2009) The *Coxiella burnetii* ankyrin repeat domain-containing protein family is heterogeneous, with C-terminal truncations that influence Dot/Icm-mediated secretion. *J Bacteriol* **191**: 4232–4242.

Weber, M.M., Chen, C., Rowin, K., Mertens, K., Galvan, G., Zhi, H., *et al.* (2013) Identification of *Coxiella burnetii* type IV secretion substrates required for intracellular

replication and Coxiella-containing vacuole formation. *J Bacteriol* **195**: 3914–3924.

Winchell, C.G., Dragan, A.L., Brann, K.R., Onyilagha, F.I., Kurten, R.C., and Voth,

D.E. (2018) Coxiella burnetii Subverts p62/Sequestosome 1 and Activates Nrf2

Signaling in Human Macrophages . *Infect Immun* **86**.

Zusman, T., Aloni, G., Halperin, E., Kotzer, H., Degtyar, E., Feldman, M., and Segal,

G. (2007) The response regulator PmrA is a major regulator of the icm/dot type IV

secretion system in Legionella pneumophila and Coxiella burnetii. *Mol Microbiol* **63**:

1508–1523

Figure legends

Figure 1. Overview of *C. burnetii* infections. *C. burnetii* is an obligate intracellular pathogen which infects wild and farm animals. Bacteria are shed in the environment with birth products and excretions leading to the contamination of hay and dust. Human infection occurs 2-3 weeks after the inhalation of contaminated particles, followed by the internalization of *C. burnetii* by alveolar macrophages. Human Q fever remain asymptomatic in 60% of infected individuals, whereas 40% develop an acute, febrile disease, which can turn into a chronic disease with more severe symptoms including endocarditis. *C. burnetii* invades eukaryotic cell through phagocytosis, which is facilitated by the bacterial invasin OmpA. Early CCVs mature along the endocytic pathway by successive fusion events with early endosomes (EEs), late endosomes (LEs) and lysosomes. Acidification of the CCV activates bacterial metabolism and the translocation of bacterial effector proteins (purple circles and ovals) by the Dot/Icm secretion system. Several effector proteins are collectively beneficial for the biogenesis of the mature CCV where bacterial replication occurs. These include the Cvp family (for *Coxiella* vacuolar proteins) of effectors that localize at CCV membranes and manipulate host membrane trafficking pathways. CvpA interacts with the clathrin adaptor AP2 at recycling endosomes (REs), re-routing these compartments to the forming CCV. CvpB binds PI(3)P at early endosomes and enriches this phospholipid at CCVs. This is required to favor the autophagy-mediated homotypic fusion between CCVs. Although not a Cvp, Cig57 interacts with the clathrin adaptor FCHO2 at clathrin-coated pits and re-routes clathrin-mediated membrane traffic to the CCV.

Figure 2. Milestone discoveries that have contributed to the knowledge of *C. burnetii* pathogenesis.



