

# Palmitoylated Cysteines in Chikungunya Virus nsP1 Are Critical for Targeting to Cholesterol-Rich Plasma Membrane Microdomains with Functional Consequences for Viral Genome Replication

William Bakhache, Aymeric Neyret, Eric Bernard, Andres Merits, Laurence

Briant

### ► To cite this version:

William Bakhache, Aymeric Neyret, Eric Bernard, Andres Merits, Laurence Briant. Palmitoylated Cysteines in Chikungunya Virus nsP1 Are Critical for Targeting to Cholesterol-Rich Plasma Membrane Microdomains with Functional Consequences for Viral Genome Replication. Journal of Virology, 2020, 94 (10), 10.1128/JVI.02183-19. hal-02994629

### HAL Id: hal-02994629 https://hal.science/hal-02994629

Submitted on 8 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Palmitoylated cysteines in Chikungunya virus nsP1 are critical for targeting to cholesterol-rich
2	plasma membrane microdomains with functional consequences for viral genome replication
3	
4	William Bakhache <sup>a</sup> , Aymeric Neyret <sup>a</sup> , Eric Bernard <sup>a</sup> , Andres Merits <sup>b</sup> and Laurence Briant <sup>a#</sup>
5	
6	<sup>a</sup> IRIM, Univ. Montpellier - CNRS UMR9004, Montpellier, France.
7	<sup>b</sup> Institute of Technology, University of Tartu, Tartu, Estonia.
8	
9	Running head: CHIKV nsP1 affinity for cholesterol enriched membranes
10	
11 12 13	<sup>#</sup> Address correspondence to Laurence Briant, IRIM - Institut de recherche en Infectiologie de Montpellier, UMR9004 CNRS-UM. 1919 route de Mende 34293 Montpellier Cedex 5 <u>laurence.briant@irim.cnrs.fr</u>
14	
15 16	<b>Key words</b> : Chikungunya virus, Methyl/guanylyltransferase, Cholesterol, Plasma membrane, Replication complexes, Alphavirus,

#### 18 Abstract

19 In mammalian cells, alphavirus replication complexes are anchored to the plasma membrane. This 20 interaction with lipid bilayers is mediated through the viral methyl/guanylyltransferase nsP1 and 21 reinforced by palmitoylation of cysteine residue(s) in the C-terminal region of this protein. Lipid 22 content of membranes supporting nsP1 anchoring remains poorly studied. Here, we explore the 23 membrane binding capacity of nsP1 with regards to cholesterol. Using the medically important 24 Chikungunya virus (CHIKV) as a model, we report that nsP1 co-segregates with cholesterol-rich 25 detergent-resistant membrane microdomains (DRMs), also called lipid rafts. In search for critical 26 factor for cholesterol partitioning, we identify nsP1 palmitoylated cysteines as major players in this 27 process. In cells infected with CHIKV or transfected with CHIKV trans-replicase plasmids, nsP1, 28 together with the other nonstructural proteins, are detected in DRMs. While the functional 29 importance of CHIKV nsP1 preference for cholesterol-rich membrane domains remains to be 30 determined, we observed that U18666A- and imipramine-induced sequestration of cholesterol in 31 late endosomes redirected nsP1 to these compartments and simultaneously dramatically decreased 32 CHIKV genome replication. A parallel study of Sindbis virus (SINV) revealed that nsP1 from this 33 divergent alphavirus displays a low affinity for cholesterol and only moderately segregates with 34 DRMs. Behaviors of CHIKV and SINV with regards to cholesterol therefore match with the previously 35 reported differences in requirement for nsP1 palmitoylation, that is dispensable for SINV but strictly 36 required for CHIKV replication. Altogether, this study highlights the functional importance of nsP1 37 segregation with DRMs and provides new insight into the functional role of nsP1 palmitoylated 38 cysteines during alphavirus replication.

#### 39 Importance

40 Functional alphavirus replication complexes are anchored to the host cell membranes through the41 interaction of nsP1 with the lipid bilayers. In this work, we investigate the importance of cholesterol

42 for such association. We show that nsP1 has affinity for cholesterol-rich membrane microdomains 43 formed at the plasma membrane and identify conserved palmitoylated cysteine(s) in nsP1 as the key 44 determinant for cholesterol affinity. We demonstrate that drug-induced cholesterol sequestration in 45 late endosomes not only redirects nsP1 to this compartment but also dramatically decreases genome 46 replication, suggesting the functional importance of nsP1 targeting to cholesterol-rich plasma 47 membrane microdomains. Finally, we evidence that nsP1 from Chikungunya and Sindbis viruses 48 display different sensitivity to cholesterol sequestering agents, that parallel with their difference in 49 the requirement for nsP1 palmitoylation for replication. This research, therefore, gives new insight 50 into the functional role of palmitoylated cysteines in nsP1 for the assembly of functional alphavirus 51 replication complexes in their mammalian host.

52

#### 54 Introduction

55 In the last decade, evidence has pointed toward the intricate relationship between host lipid 56 metabolism and the replication of viral pathogens. Indeed, viruses can co-opt or reprogram lipid 57 signaling, synthesis and metabolism either to generate ATP, to extend cellular membranes or to 58 remodel membrane lipid content. These modifications will serve to create an environment that is 59 optimal for viral replication. This need is dictated by the pivotal role played by membranes in almost 60 all steps of the virus life cycle. Indeed, the importance of cellular lipids during the binding/entry 61 process and assembly/budding of new infectious progeny into the extracellular space has long been 62 appreciated (for review see (1)). However, the discovery that viruses with a positive-strand RNA 63 genome ((+)RNA viruses) replicate in association with host cell membranes has expanded the 64 regulatory function of lipids to genome replication. It is now well established that such viruses create 65 membranous compartments, also called virus replication organelles, originating from the 66 endoplasmic reticulum (ER), Golgi apparatus, mitochondria, peroxisomes, endosomes/lysosomes or 67 from the plasma membrane (PM) (2). In these compartments, viral replication proteins bind cell 68 membranes with an affinity for determined lipid species (3, 4).

69 Cell membranes are composed of phospholipids, glycolipids, and cholesterol. Among other lipids, 70 cholesterol constitutes a unique type of cellular membrane building block. It is responsible for 71 regulating fluidity and impermeability to lipid bilayers. In vertebrate cells, cholesterol homeostasis is 72 maintained through de novo synthesis in the ER, with 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-73 CoA) reductase being the rate-limiting enzyme, and through receptor-mediated intake from the 74 extracellular medium in the form of low-density lipoproteins (LDLs) (5). In this scenario, LDLs are 75 delivered to and hydrolyzed in late endosomes/lysosomes (LE/Ls) where free cholesterol is released. 76 Then, cholesterol requires proper intracellular transport to exit the LE/Ls and reach its final 77 destination mainly at the PM, in the Golgi apparatus and the ER. This is ensured by Niemann-Pick C 1 78 and 2 (NPC1 and NPC2) proteins localized at the limiting membrane of lysosomes and in the

79 lysosomal lumen respectively (6). Once at the PM, cholesterol together with glycosphingolipids, 80 glycophosphatidylinositol (GPI)-anchored proteins and transmembrane proteins can cluster into 81 discrete domains. These cholesterol-enriched detergent-resistant membrane microdomains, referred 82 to as DRMs or lipid rafts, have been identified as platforms for both endocytosis of penetrating viral 83 particles and for progeny assembly and budding (7). More recently, cholesterol regulatory function 84 was extended to the replication step of (+)RNA viruses. The local accumulation of cholesterol was 85 proposed to contribute to the creation of a membrane microenvironment conducive to assembly and 86 optimal functionning of replication complexes formed by members of the Flaviviridae family, 87 including hepatitis C virus (HCV) (8-10) and West Nile virus (11), by members of Picornaviridae 88 family, including Coxsackievirus and poliovirus (12) and by plant viruses from the Bromoviridae family 89 (Brome mosaic virus; BMV) (13). Consistent with the idea that cholesterol accumulation may be 90 required for optimal activity of viral replication machinery, manipulation of cholesterol metabolism 91 was found to impair genome replication of taxonomically divergent (+)RNA viruses (8, 12, 14–18).

92 Alphaviruses are (+)RNA viruses, which are predominantly transmitted to vertebrates by mosquito 93 vectors. Chikungunya virus (CHIKV) is an Old World alphavirus causing millions of infections in 94 tropical and subtropical geographical areas with a potential risk of spreading to regions with a 95 temperate climate. It has recently received significant attention as a consequence of its re-96 emergence in the Indian Ocean and Caribbean Islands before spreading worldwide (19). CHIKV, like 97 other alphaviruses, replicates its genome in membranous niches derived from the host PM (20–22). 98 The replication complex confined in these organelles, termed as spherules, contains the four non-99 structural proteins nsP1, nsP2, nsP3 and nsP4 encoded by the 5' ORF in the viral genome which are 100 expressed in the form of P123 and P1234 polyprotein precursors. In this complex, membrane binding 101 is mediated by nsP1, the viral methyltransferase (MTase) and guanylyltransferase (GTase), which 102 catalyzes the formation of the cap structure at the 5' end of nascent positive stand viral RNAs (23, 103 24). A putative  $\alpha$ -helix central to nsP1 was proposed to play a crucial role for membrane attachment 104 (25, 26). S-acylated cysteines located in the C-terminal region of this protein were then proposed to stabilize this interaction (27). The functional requirement for nsP1 interaction with membranes varies among alphaviruses. It is strictly required for Semliki Forest virus (SFV) nsP1 enzymatic activity (28) while it is dispensable for that of Sindbis virus (SINV) nsP1 (29). More specifically, alanine substitution of palmitoylated cysteines has only a marginal inhibitory effect on SINV replication (30, 31) while leading to the acquisition of compensatory mutations for SFV (31, 32). Recently, this mutation was reported to completely abolish the replication of CHIKV (33, 34).

111 In vitro, SFV nsP1 has affinity for anionic phospholipids especially phosphatidylserine, 112 phosphatidylglycerol, and cardiolipin; these lipid species significantly improve its capping activity (25, 113 28). In cells, however, nsP1 affinity for specific lipids remains almost uninvestigated. In recent years, 114 cholesterol metabolism was reported to be critical for the replication of alphavirus genome. Indeed, 115 SINV RNA replication and protein synthesis is significantly decreased in fibroblasts from patients with 116 type A Niemann-Pick disease (NPD-A) which induces cholesterol and sphingolipid storage in LE/Ls 117 (35). More recently, we reported that U18666A, a class II cationic amphipathic steroid  $3-\beta$ -[2-118 (diethylamine)ethoxy]androst-5-en-17-one, and the anti-depressant drug imipramine, which both 119 phenocopy NPD-A, are CHIKV inhibitors with potential activity against RNA replication steps (16). 120 Here, we further question the importance of cholesterol metabolism in the alphavirus life cycle. We 121 especially explore the outcome of cholesterol manipulation on nsP1 subcellular distribution and 122 membrane anchoring. We show that nsP1 has affinity for cholesterol-rich membranes. Indeed, 123 U18666A or imigramine redirect CHIKV nsP1 to Lamp2-positive endosomes where unesterified 124 cholesterol accumulates. At the plasma membrane, nsP1 co-segregates with cholesterol-containing 125 DRMs as attested by membrane flotation assays. Investigating the molecular basis of nsP1 targeting 126 to DRMs revealed that the cysteines, previously reported to be palmitoylated (34), are the main 127 determinants for association with these domains. Interestingly, when expressed transiently in the 128 context of a P1234 polyprotein precursor or by an infectious CHIKV, nsP2, nsP3, and nsP4 were found 129 to co-segregate with nsP1 in cholesterol-rich membrane fractions, a property that was abolished 130 when nsP1 cysteines were mutated. In a parallel study we found that cysteine-mediated nsP1 association with cholesterol-rich membranes is conserved for SINV despite being less marked than for CHIKV. Moreover, SINV nsP1 was also less sensitive to U18666A-induced cholesterol sequestration. This phenotype with regard to nsP1 cholesterol partitioning parallels the reduced requirement for cysteine palmitoylation previously reported for SINV replication (30, 31). Altogether this study provides clues on the proviral role of cholesterol in alphavirus replication, suggesting its regulatory function in nsPs association with the PM and presumably in the formation of functional replication complexes.

138 Results

#### 139 Cholesterol is pivotal for alphavirus genome replication.

140 First, we set out to study the involvement of cholesterol homeostasis in CHIKV replication. 141 Cholesterogenesis was inhibited using lovastatin, an FDA-approved inhibitor of HMG-CoA reductase, 142 that catalyzes the conversion of HMG-CoA to mevalonate in the cholesterol biosynthesis pathway 143 (Fig 1A) (36). Cholesterol availability at the PM was reduced using U18666A or imipramine. By 144 targeting NPC1 transporter, both drugs block the transfer of endocytosed cholesterol from late 145 endosomes to different organelles, including the PM, without significant effect on other lipid species 146 (37). Additionally, U18666A inhibits enzymes of the cholesterol synthesis pathway (38). Each drug, 147 used in a concentration range that was controlled to have limited toxicity (Fig 1B-b,d,f), was added to 148 HEK293T cells 30 mins before infection with CHIKV-LR-5'GFP at a multiplicity of infection (MOI) of 149 0.5. Cells were maintained in culture with the drugs for 24 hrs and then infection was monitored 150 measuring levels of GFP reporter expressed by this recombinant virus. In each case, CHIKV infection 151 was significantly reduced as compared with the mock-treated condition (Fig 1B-a,c,e).

We and others have demonstrated that the depletion of membrane cholesterol is deleterious for fusion of the virion and host membranes (39, 40). Because host membranes are also pivotal for replication of alphavirus genome through the creation of membranous replication organelles, we

155 investigated whether cholesterol biosynthesis is also required for the post-entry step of CHIKV 156 infection cycle. To this end, we performed experiments in which cells were treated with cholesterol 157 metabolism and transport inhibitors 1 hr after CHIKV infection. In these conditions, lovastatin, 158 U18666A and imipramine decreased CHIKV genome replication (Fig 2A). To definitively omit drug 159 effects on viral entry, we finally took advantage of a CHIKV trans-replication system that 160 recapitulates post-entry events of CHIKV life cycle (33). Plasmids CMV-P1234 and HSPoll-Fluc-Gluc, 161 encoding the P1234 polyprotein and a replication-competent template RNA containing firefly and 162 Gaussia luciferase reporter genes under the control of genomic and subgenomic viral promoters 163 respectively (Fig 2B), were cotransfected in HEK293T cells. Consequences of cholesterol transport 164 inhibition were assayed by adding increasing concentrations of lovastatin, U18666A, and imipramine 165 to transfected cells and quantification of reporter activities in cell lysates. As depicted in Figure 2C, 166 reporter expression directed by both genomic and subgenomic promoters was decreased by the 167 drugs when compared to control conditions. Altogether, these results indicate that cholesterol 168 homeostasis, including ongoing biosynthesis and transport of unesterified cholesterol to the host 169 membranes, is pivotal for CHIKV genome replication.

# 170 α-helix and putative palmitoylated cysteines cooperate for CHIKV nsP1 binding to host 171 membranes.

172 Alphavirus replication complexes are anchored to the host membranes thanks to nsP1 membrane 173 binding capacity. While extensively reported for nsP1 encoded by SFV and SINV (25, 30), this feature 174 has not yet been studied functionally for CHIKV. To investigate CHIKV nsP1 behavior with regard to 175 host cell membranes, a plasmid encoding a GFP-fused CHIKV nsP1 protein (GFP-nsP1; Fig 3A) was 176 generated and used to transfect HEK293T cells. NsP1 membrane association was investigated by 177 fractionation of transfected cells. Post-nuclear extract prepared from GFP-nsP1-expressing cells was 178 separated into membranous (P25) and cytosolic (S25) samples by differential centrifugation as 179 previously reported (41). Each fraction was resolved using SDS-PAGE and probed with anti-GFP

180 antibodies and with antibodies against  $Na^+/K^+$  ATPase or GAPDH, that respectively associate with 181 membrane and cytosolic compartments. In these conditions, GFP alone was detected in the cytosolic 182 sample together with GAPDH (Fig 3B and 3C). By contrast, GFP-nsP1 was detected in the membrane 183 fraction also containing Na<sup>+</sup>/K<sup>+</sup> ATPase. In parallel HeLa cells expressing these proteins were analyzed 184 using confocal microscopy. As expected, the fluorescence of individual GFP was diffuse in the 185 cytoplasm and nucleus. However, GFP-nsP1 fluorescence overlapped with PM stained using wheat 186 germ hemagglutinin (WGA)-conjugated with Alexa Fluor 647 (Fig 3D). As previously reported for 187 related alphaviruses (42), expression of GFP-nsP1 generated huge membrane reshaping creating 188 filopodia- and lamellipodia-like structures covering the entire cell surface, that stained positive for 189 the green fluorescence. Of note, this profile was also observed for a C-terminally GFP-tagged nsP1-190 GFP protein as well as for an untagged nsP1 detected by mean of anti-nsP1 serum (Fig 3E), thereby 191 supporting that GFP-nsP1 behaves as native nsP1 regarding localization and association with 192 membranes. According to the focal plane chosen, a fraction of each of these proteins was also 193 detected as small cytosolic aggregates as illustrated in Figure 3E.

194 Because alphavirus nsP1 was previously proposed to be trafficked to endosomes (43) and in light of 195 our confocal microscopy analysis, we further questioned nsP1 subcellular distribution by performing 196 fractionation assays that separated the PM from other cell membranes. The post-nuclear extract was 197 prepared from GFP-nsP1 expressing cells and separated by isopycnic centrifugation in a self-forming 198 linear 10-20-30% iodixanol density gradient. Twenty-four samples were collected from top to bottom 199 and assayed by western blotting for GFP-nsP1 content. A roughly equal proportion of GFP-nsP1 was 200 detected in the top fractions 1 and 2 and also in fractions 9 to 12 that all stained positive for the 201  $Na^{+}/K^{+}$  ATPase membrane marker (Fig 3F). These fractions also contained flotillin-1 (FLOT1) that is 202 known to localize predominantly to the PM and endosomal compartments, i.e. late endosomes and 203 recycling endosomes (44). By contrast, individual GFP segregated with fractions 17 to 24 at the 204 bottom of the gradient that corresponded to the cytosolic compartment. Altogether, these results 205 show that CHIKV nsP1 is a membrane-associated protein, that cofractionates equally with the PM

and some internal membranes, probably endosomal in nature, suggesting that this protein maytraffic between the two compartments.

208 We next questioned whether the functions of membrane binding determinants previously identified 209 in alphavirus nsP1 proteins are also conserved for CHIKV nsP1. Extensive analysis performed using 210 SFV and SINV as models established that membrane association of nsP1 relies on a central conserved 211 sequence that folds as an amphipathic  $\alpha$ -helix when studied as a synthetic peptide in solution (25). In 212 this sequence, a pivotal role in membrane anchoring was attributed to tryptophan at position 259 213  $(W_{259})$  that sinks into the phospholipid bilayer (25). This interaction was proposed to be reinforced by 214 the presence of cysteine residue at position 420 (C420) in SINV or cysteine residues 418-420 in SFV 215 (C418-420) which are covalently palmitoylated and render the protein highly hydrophobic (27, 30). 216 These motifs are conserved in the CHIKV nsP1 sequence where W258 and C417-419 of residues 217 correspond to W259 and C418-420 of SFV nsP1, respectively (Fig 3A). Recently, C417-419 in CHIKV 218 nsP1 were reported to be palmitoylated (34), therefore further attesting that nsP1 palmitoylation is a 219 conserved feature among Old World alphaviruses. To study the contribution of each motif in CHIKV 220 nsP1 membrane binding, we generated GFP-nsP1 mutants in which W258 (GFP-nsP1<sub>W258A</sub>) or  $C_{417-419}$ 221 (GFP-nsP1<sub>3A</sub>) residues in nsP1 sequence were replaced by alanine as well as a double nsP1 mutant 222 bearing a combination of W<sub>258</sub>A or C<sub>417-419</sub>A mutations (GFP-nsP1<sub>DM</sub>). Fluorescence microscopy of cells 223 transfected with the corresponding plasmids evidenced a diffuse green fluorescence that 224 predominated in the cytoplasm by contrast with cells expressing GFP-nsP1 that was mainly detected 225 at the plasma membrane (Fig 3D). Fractionation experiments confirmed that GFP-nsP1<sub>W258A</sub> and GFP-226 nsP1<sub>3A</sub> were more abundant in the cytosolic fraction than GFP-nsP1, with a significant amount of 227 each mutant protein remaining associated with cell membranes (Fig 3B and 3C). The same cell 228 extracts were then subjected to membrane flotation assay in iodixanol gradient to appreciate their 229 capacity to associate with the internal/plasma membrane compartments. GFP-nsP1<sub>W258A</sub> and GFP-230 nsP1<sub>3A</sub> proteins were both detected in fractions 18-24 of the gradient thereby confirming that these 231 proteins have a decreased membrane affinity (Fig 3F). Each mutant protein was also present at the 232 top of the gradient in fractions 1-2 corresponding to the PM and in fractions 9-12 corresponding to 233 internal membranes. Distribution of mutant proteins in these fractions generally mimicked that of 234 GFP-nsP1, except that GFP-nsP1<sub>W258A</sub> was somewhat more abundant in internal membrane fractions. 235 This was also reflected in more diffuse localization of GFP fluorescence in cells expressing GFP-236 nsP1<sub>W258A</sub> (compare panels b and d of Fig. 3D) indicating possible role of W258 for PM affinity of CHIKV nsP1. Nevertheless, the observed differences were small suggesting that  $W_{258}A$  and  $C_{417-419}A$ 237 238 mutations despite decreasing membrane affinity do not prominently modify nsP1 distribution 239 between intracellular and plasma membranes or nsP1 trafficking capacity. Interestingly, membrane 240 association was further decreased when  $W_{258}A$  and  $C_{417-419}A$  mutations were combined in GFP-nsP1<sub>DM</sub> 241 (Fig 3B and 3D). Altogether these results establish that CHIKV nsP1 associates both with the plasma 242 and internal membranes, an association dictated by the cooperation of motifs that contain C417-419 243 and W258 amino acids in this protein.

#### 244 Cholesterol storage defect redirects CHIKV nsP1 to late endosomes.

245 Based on the observation that cholesterol synthesis and transport is required for CHIKV RNA 246 replication and on the intimate relationship of CHIKV nsP1 with membranes, we next wondered 247 whether cholesterol homeostasis has an impact on nsP1 behavior. We first investigated nsP1 248 subcellular localization with respect to cholesterol distribution. Cells transfected with plasmid 249 expressing GFP-nsP1 were incubated with U18666A or imipramine to generate a cholesterol 250 entrapment in LE/Ls (45). This capacity was controlled using the antifungal antibiotic filipin III that 251 forms a fluorescent complex upon association with unesterified cholesterol (46). In the presence of 252 U18666A or imipramine, filipin III evidenced the formation of large fluorescent cytosolic aggregates 253 that contrasted with the presence of cholesterol at the PM and more evenly distributed in the 254 cytoplasm of mock-treated cells (Fig 4A and data not shown). For U18666A, these clusters colocalized 255 with Lamp2 (Fig 4B), a LE/Ls marker, thereby confirming that this pharmacological agent stimulates 256 the accumulation of unesterified cholesterol in late endosomes as previously reported (47). Using

257 these experimental conditions, we next assessed the impact of cholesterol storage defect on GFP-258 nsP1 subcellular localization and membrane affinity. Fractionation assays established that drug 259 treatment did not increase amount of cytosolic nsP1 (Fig 4C; S25 fraction) indicating that it had no 260 consequence on nsP1 membrane affinity. However, microscopy imaging revealed that, in the 261 presence of U18666A, a significant part of GFP-nsP1 fluorescence was redirected from the PM to 262 cytosolic aggregates. This signal overlapped with filipin III staining as attested by cross-sectional 263 analysis of the fluorescent signals (Fig 4D and 4E). In similar experimental conditions, no 264 redistribution of individual GFP fluorescence was observed. Accordingly, upon cholesterol storage 265 condition, GFP-nsP1 colocalized with unesterified cholesterol stored in late endosomes as quantified 266 by calculating Mander's coefficient (Fig 4F). This situation contrasted with that of mock-treated cells 267 in which the GFP-nsP1 fluorescence colocalized with filipin III at the PM and did not overlap with 268 Lamp 2 staining. Altogether, our results argue that the inhibition of NPC1-mediated cholesterol 269 transport by U18666A redirects GFP-nsP1 to LE/Ls where unesterified cholesterol accumulates. Re-270 targeting GFP-nsP1 to these compartments has no significant impact on nsP1 membrane binding 271 capacity.

#### 272 C417-419 but not W258 residue determine CHIKV nsP1 sensitivity to cholesterol distribution.

273 Palmitoylation governs protein trafficking and association with membranes. A major focus of studies 274 on protein palmitoylation has been the role of this modification in promoting interaction with 275 gangliosides and cholesterol, leading at certain conditions, to protein translocation to raft/caveolae 276 membrane domains (48). Given that the C417-429 in CHIKV nsP1 required for optimal membrane 277 association have recently been reported to be palmitoylated (34), we investigated their contribution 278 to GFP-nsP1 cholesterol affinity. The above described experiments were repeated using cells 279 transfected with GFP-nsP1<sub>3A</sub> plasmid. In the presence of U18666A, GFP-nsP1<sub>3A</sub> fluorescence 280 remained diffuse in the cytoplasm whether the cells were maintained in the presence of U18666A or 281 with an appropriate concentration of vehicle (Fig 5C). GFP-nsP1 $_{W258A}$  fluorescence was concentrated

in cytosolic foci colocalized with filipin III fluorescence (Fig 5A) as observed for GFP-nsP1. These phenotypes were confirmed by cross-sectional analysis of the fluorescent signals (Fig 5B and D) and by calculation of Mander's coefficients (Fig 5E). Accordingly, W258, even if required for strengthening nsP1 membrane affinity, is not critical for cholesterol affinity. In contrast, cysteine-toalanine substitution in mutant GFP- nsP1<sub>3A</sub> abolished sensitivity to U18666A, suggesting a critical role of palmitoylated cysteines in nsP1 cholesterol affinity.

288 To further explore GFP-nsP1<sub>3A</sub> behavior with regard to cholesterol, we took advantage of CD81, a 289 heavily palmitoylated tetraspanin, segregating mainly with DRMs, as a lipid raft biomarker (49). 290 Indeed, in our hands, this protein was mainly detected at the PM of untreated U2OS cells (Fig 6A). It 291 was redirected to cytosolic aggregates in cells cultured with U18666A or imipramine (Fig 6B) where it 292 colocalized with GFP-nsP1 as attested by Mander's and Pearson's coefficient calculation (Fig 6D and 293 6E). CD81 therefore appears as a sensitive cholesterol sensor. Using this property, we explored the 294 importance of nsP1 palmitoylated cysteines for cholesterol dependency. Cells transfected to express 295 GFP-nsP1<sub>3A</sub> were incubated in the presence of U18666A. Despite inducing the clustering of CD81 into 296 intracellular compartments, this treatment did not affect GFP-nsP1<sub>3A</sub> distribution (Fig 6C). Indeed, 297 GFP-nsP1<sub>3A</sub> remained detected as a diffuse cytoplasmic protein as observed in mock-treated cells 298 thereby contrasting with the phenotype observed for GFP-nsP1. No colocalization of CD81 and GFP-299 nsP1<sub>3A</sub> fluorescence was observed (Fig 6D). Optical sectioning (z-stack) and three-dimensional (3D) 300 volume reconstruction from image stacks confirmed that conversely to GFP-nsP1, GFP-nsP1<sub>3A</sub> poorly 301 colocalized with CD81 in drug treated cells (Fig 6E and 6F). These phenotypes were confirmed from 302 cells cultured with imipramine. According to these results, nsP1 behaves similarly to CD81 with 303 regard to cholesterol storage, a property that requires the presence of palmitoylated cysteines in 304 nsP1 C-terminus.

305 CHIKV nsP1 partitions with DRMs.

306 Given that nsP1 subcellular localization is sensitive to cholesterol redistribution to the PM, we 307 questioned its capacity to segregate with cholesterol-enriched membrane microdomains. Cholesterol 308 is not uniformly distributed in membranes. In living cells, it concentrates in nanoscale assemblies, 309 also enriched in sphingolipids and glycosylphosphatidylinositol (GPI)-anchored proteins, referred to 310 as lipid rafts. These compartments are characterized biochemically by their insolubility in non-ionic 311 detergents, a property reflected in their name (DRMs - detergent-resistant membrane 312 microdomains), and by their light density on sucrose gradients. Therefore, they can be separated 313 from non-raft membranes by centrifugation methods. Samples prepared from cells transfected with 314 GFP-nsP1 expression plasmid were treated with Triton X-100 at 4°C, separated on a 10-80% sucrose 315 density gradient, and then analyzed by western blot. Detergent-resistant fractions corresponding to 316 DRMs were identified utilizing an antibody against FLOT1, a well-known raft-associated protein (50). 317 In our experimental conditions, FLOT1 fractionated into light density fractions 1 and 2 at the top of 318 the gradient thereby identifying DRMs (Fig 7A). In contrast, the non-raft marker Na<sup>+</sup>/K<sup>+</sup> ATPase 319 remained associated with fractions 7-9 of heavier density corresponding to non-raft membranes and 320 cytosolic compartment (detergent sensitive; DS). Using this protocol, more than 85% of the GFP-nsP1 321 protein was detected in FLOT1-positive fractions at the top of the gradient supporting its capacity to 322 associate with cholesterol-enriched DRMs (Fig 7A and 7B). To confirm GFP-nsP1 affinity for 323 cholesterol-enriched microdomains, this experiment was repeated starting from cells cultured in the 324 presence of the cholesterol depleting agent methyl- $\beta$ -cyclodextrin ( $\beta$ MCD). Due to its ability to 325 sequester cholesterol in its hydrophobic pocket,  $\beta$ MCD extracts cholesterol from the lipid bilayer and 326 disrupts DRMs (51). Under these cholesterol depletion conditions, FLOT1 was redistributed from 327 DRMs to DS fractions and sedimented at the bottom of the density gradient (Fig 7C and 7D). Analysis 328 of the same fractions with anti-nsP1 antibodies revealed that GFP-nsP1 was barely detectable in top 329 fractions while it accumulated in fractions of heavier density together with FLOT1. Therefore, GFP-330 nsP1 fractionation with DRMs is sensitive to cholesterol extraction from cell membranes. Finally, this 331 experiment was repeated starting from cells expressing GFP-nsP1<sub>W258A</sub> or GFP-nsP1<sub>3A</sub> mutant proteins that displayed different sensitivity to U18666A. As observed for GFP-nsP1, the GFP-nsP1<sub>W258A</sub> mainly segregated with DRMs fractions attesting that raft association was not disrupted by mutation of W258 residue in the putative  $\alpha$ -helix of nsP1 (Fig 7E and 7F). Analyzing the behavior of the GFPnsP1<sub>3A</sub> mutant protein revealed, in contrast, that most of this protein was absent from the top fraction and co-segregated with the DS fractions. These experiments demonstrate that the presence of GFP-nsP1 in DRMs is mainly dictated by acylated cysteines while mutation in the putative  $\alpha$ -helical peptide has only marginal impact on this phenotype.

#### 339 nsPs associate with DRM fractions in cells with CHIKV RNA replication

340 Unlike SFV, CHIKV replication is completely abolished by nsP1<sub>3A</sub> mutation regardless of the type and 341 growth temperature of cells (33, 34). In contrast, nsP1<sub>W258A</sub> mutation allows virus to grow efficiently 342 in insect cells or in mammalian cells cultivated at 28°C (33, 52). Combined with our observations, this 343 strongly indicates that the presence of nsP1 in DRMs is an absolute requirement for CHIKV genome 344 replication. Therefore, we questioned whether nsP1 affinity for DRMs was conserved in cells 345 containing functional CHIKV replicase complexes. To this end, HEK293T cells either transfected with a 346 trans-replication system that reproduces CHIKV RNA replication (Fig 2B) or infected with the CHIKV-347 LR-5'GFP were used. DRMs isolation followed by immunoblot analysis revealed that a significant 348 fraction of nsP1 expressed in transfected or infected cells (~85 and ~40%, respectively) was detected 349 in cholesterol-rich fractions (Fig 8D and 8E). NsP1 is the only alphavirus nonstructural protein with 350 membrane affinity. In the replication complex, nsP1 co-localizes with nsP2, nsP3, and nsP4 (53–56). 351 Thus, it is suspected to play a critical role in replication complex anchoring to the PM. Therefore, we 352 investigated whether nsP1 capacity to segregate with DRMs has an impact on the association of 353 other nsPs with specialized membrane microdomains. Probing gradient fractions prepared from 354 transfected and infected cells with antibodies against nsPs established that a significant part of each 355 of them was detected in DS fractions thereby agreeing with the previously reported capacity of nsPs 356 to be present in different cytosolic compartments (57). However, approximately 10-30 % of total

357 nsP3 and nsP4 levels were also present in raft fractions. In some, but not in all experiments, nsP2 was 358 also detectable in DRMs; most probably reflecting the weaker interaction of nsP2 with other 359 components of replicase complexes. In order to further assess the relevance for the presence of 360 nsP2, nsP3 and nsP4 in DRMs fractions, these experiments were reproduced starting from cells 361 transfected with a CHIKV trans-replicase system in which the nsP1 protein contained the C418-420A 362 mutation. As expected, this mutation prevented nsP1 association with DRMs (Fig 8C). Concomitantly, 363 nsP2, nsP3 and nsP4 were also excluded from these fractions. Altogether these results confirm the 364 capacity of nsP1, expressed in the context of trans-replicase or by infectious CHIKV, to associate with 365 DRMs, an association that dictates targeting of other nsPs, albeit at low levels, to cholesterol-366 enriched membrane microdomain. Furthermore, the C<sub>418-420</sub>A mutation that completely inactivates 367 CHIKV trans-replicase (33) completely prevented association of CHIKV nsPs with DRMs.

#### 368 Conservation of nsP1-directed cholesterol affinity among divergent Old World alphaviruses.

369 Despite the presence of palmitoylated cysteine residues in the C-terminal region of nsP1 is a 370 conserved feature of distantly related Old World alphaviruses (30, 34), functional studies have 371 highlighted differences with regard to cysteine requirement for genome replication. Cysteine-to-372 alanine mutation are lethal for CHIKV (33, 34) but is well tolerated by SINV (30). In the light of such 373 differences and of the herein evidenced role of the CHIKV nsP1 cysteines in lipid raft association, we 374 next questioned whether SINV nsP1 behaves similarly with respect to cholesterol. To achieve this, a 375 plasmid encoding a GFP-fused SINV nsP1 protein was generated (Fig 9A) and used for the 376 transfection of HEK293T cells. Then, we tested SINV nsP1 partitioning to DRMs by membrane 377 flotation assays. As previously observed for CHIKV nsP1, SINV nsP1 was also detected in the top 378 fractions of the density gradient supporting its ability to partition with cholesterol-rich membrane 379 microdomains. However, contrasting with CHIKV for which almost 85% of nsP1 associated to DRM 380 fractions, a roughly equal proportion of the SINV nsP1 was associated with DRMs and DS (Fig 9B). We 381 further investigated SINV GFP-nsP1 affinity for cholesterol by testing its sensitivity to U18666A in

382 confocal microscopy experiments. In control conditions, SINV GFP-nsP1 was detected at the PM, 383 including in filopodia-like membrane protrusions that were abundantly observed as reported before 384 for the untagged protein (42) and for CHIKV GFP-nsP1 (Fig 9C). In the presence of U18666A, SINV 385 GFP-nsP1 was still mainly detected at the PM, while cholesterol stained with filipin was concentrated 386 in intracellular storage compartments as expected. In these conditions, co-localization of GFP-nsP1 387 SINV with cholesterol-enriched endosomes was unfrequently detected. This is in contrast with the 388 results obtained for CHIKV nsP1 (Fig 4D) (Mander's coefficient of 0.394 ± 0.032 and 0.217 ± 0.001 for 389 CHIKV and SINV nsP1 respectively). Next, we investigated the role of palmitoylated cysteine by 390 repeating these experiments starting from cells expressing a cysteine-to-alanine SINV nsP1 391 palmitoylation-negative mutant protein (GFP-nsP1<sub>C420A</sub>). As shown in Figure 9B, the DRMs association 392 of GFP-nsP1<sub>C420A</sub> was significantly reduced when compared with that of SINV GFP-nsP1 (Fig 8A). 393 Analyzing SINV GFP-nsP1 C420A subcellular localization in U18666A treated cells confirmed that this 394 mutant did not colocalize with filipin-labelled cholesterol (Fig 9D). Altogether these results suggest 395 that SINV nsP1 is targeted to DRMs, a property that depends of palmitoylated cysteine as previously 396 observed for CHIKV nsP1. However, compared with CHIKV counterpart, SINV nsP1 is equally 397 abundant in DS compartments and displays only modest sensitivity to U18666A suggesting a reduced 398 affinity for cholesterol.

#### 399 Discussion

The present study identifies CHIKV nsP1 as a lipid-raft co-segregating protein with an affinity for cholesterol. We defined cysteine residues that can be palmitoylated as the molecular determinant important for this targeting. In the context of cells with ongoing CHIKV RNA replication, nsP1, together with a fraction of other nsPs, partitions with cholesterol-rich DRMs. Together with evidence that drugs reducing cholesterol availability at the PM impair CHIKV RNA replication, our results support that nsP1 targeting to cholesterol-rich PM microdomains may have a functional importance for viral genome replication.

407 Cholesterol is a main component of membranes. Together with sphingolipids, it segregates into 408 discrete microdomains, referred to as lipid rafts or DRMs, present both on the inner and the outer 409 leaflet of the PM (58, 59). These membrane domains with a size on the nm scale are highly dynamic. 410 They accumulate a subset of membrane proteins, mainly GPI-anchored proteins, transmembrane 411 proteins, and acylated cell components (60, 61). Based on these properties, rafts were seen as 412 platforms that compartmentalize cellular processes with an important function in receptor-ligand 413 interaction, signal transduction and endocytosis (60). Herein, we establish that CHIKV nsP1 associates 414 with the PM. In this compartment, a pool of nsP1 is targeted to cholesterol-containing DRMs (Fig 415 10a), an association that was reversed by  $\beta$ MCD cholesterol-removal agent. Affinity for cholesterol 416 was further supported by investigating nsP1 behavior with regard to U18666A- or imipramine-417 mediated cholesterol sequestration in LE/Ls. We found that intracellular cholesterol storage resulted 418 in nsP1 accumulation in endosomes without consequence on overall nsP1 membrane binding ability 419 (Fig 10b). Altogether these results indicate that availability of cholesterol at the PM is required for 420 appropriate targeting of CHIKV nsP1 to this compartment.

421 In the last decade, the biochemical or biophysical underpinnings that govern nsP1 association with 422 membranes have been the focus of an intense attention. For SFV and SINV, the central  $\alpha$ -helical 423 motif in nsP1 spanning amino acid residues 245 to 264 was proposed as the main determinant for 424 membrane anchoring to lipid bilayers with W259 residue being critical for hydrophobic interactions 425 with the phospholipid acyl chains (25). Conserved acylated cysteine(s) in nsP1 were proposed to 426 tighten this membrane interaction (26, 27, 30, 34). Using cell fractionation assays, we show that both 427  $W_{258}A$  substitution in the putative  $\alpha$ -helix of CHIKV nsP1 and  $C_{417-419}A$  mutations indeed decrease 428 nsP1 affinity for cell membranes. Interestingly, combining W258A and C417-419A mutations further 429 reduced nsP1 membrane association thereby suggesting that in the CHIKV nsP1 the two domains 430 may synergize for membrane interaction, a situation that was not described for other alphaviruses. 431 Analyzing the contribution of these membrane interaction determinants in nsP1 targeting to lipid

432 rafts revealed that the W258A mutation had only a slight effect on nsP1 co-fractionation with 433 cholesterol-rich domains. This mutation also slightly reduced PM association of nsP1 and facilitated 434 its association with internal membranes. This may indicate that W258 residue is important for 435 membrane/plasma membrane targeting of nsP1 but not for its palmitoylation, DRMs association and 436 enzymatic activity. How these properties correlate with the proposed role of W258 residue as one of 437 membrane anchors of nsP1 is currently unclear. For the proper understanding of the somewhat 438 controversial data regarding the importance of the W258 residue, the structure of the membrane 439 bound  $\alpha$ -helical peptide of nsP1 (25) should be compared with the structure of membrane-bound 440 enzymatically active nsP1 which, to this date, is not available. In contrast to the W<sub>258</sub>A mutation the 441 effects of the C<sub>417-419</sub>A mutation on CHIKV nsP1 were unambiguous. This mutation, previously 442 reported to prevent CHIKV nsP1 palmitoylation and replication (33, 34), dramatically reduced DRMs 443 co-fractionation (Fig 10c). Moreover, by contrast with wild-type nsP1, nsP1<sub>C417-419A</sub> sequestration with 444 Lamp2 in endosomes could not be observed under U18666A or imipramine treatment. According to 445 these experiments, acylation appears as critical to direct nsP1 to cholesterol-enriched membrane 446 microdomains. This result parallels previous evidence regarding the role of palmitoylation in cellular 447 (p59<sup>fyn</sup> and p60<sup>src</sup>) (62) or viral (influenza hemagglutinin) (63) proteins association with rafts. This 448 observation raised the question of the functional outcome of nsP1 association with cholesterol-rich 449 DRMs.

450 Because other nonstructural proteins of alphaviruses cannot directly associate with membranes, 451 nsP1 plays a decisive role in proper targeting and membrane binding of other nsPs involved in the 452 formation and functioning of alphavirus replication complex (24). Starting from cells infected with 453 CHIKV, we found that a fraction of each of the four nsPs was associated with DRMs. These results 454 were confirmed in cells transfected with plasmids encoding fora CHIKV trans-replication system, in 455 which other non-structural proteins co-sedimented with DRMs depending on the integrity of nsP1 456 C417-419 residues. Currently the functional importance of nsPs targeting to rafts is unknown. 457 Nevertheless, we observed that in addition to nsP1 sequestration into late endosomes, U18666A and

imipramine also generated a significant drop in CHIKV genome replication. This raises the question ofthe functional consequences of nsP1 mistargeting and of its impact on replication complex assembly.

460 Over two decades, alphaviruses encoding nsP1 mutants with reduced membrane binding ability have 461 been in the focus of different studies. They established that nsP1 palmitoylation had only a mild 462 impact on SINV or SFV infectivity, highlighting the essential role of the W259 residue in nsP1's central 463  $\alpha$ -helix for both membrane anchoring and genome replication (30, 31). For CHIKV, the W<sub>258</sub>A mutant 464 is viable, albeit having a temperature-sensitive phenotype (33). By contrast, the C<sub>417-419</sub>A mutation 465 results in complete inactivation of the CHIKV replicase, leading to non-functional enzymes unable to 466 synthesize any viral RNAs both in mammalian and insect cells (33, 34, 52). Here, the direct 467 comparison of SINV and CHIKV revealed that SINV nsP1 partitions with DRMs but to a lesser extent 468 than observed for CHIKV nsP1. This phenotype was equally dependent upon conserved cysteine in 469 nsP1. Moreover, SINV nsP1 targeting to the PM was less sensitive to cholesterol manipulation by 470 U18666A. These discrepancies therefore parallel the differences in nsP1 palmitoylation requirement 471 previously reported for CHIKV and SINV replication (33). However, counterintuitively, SINV genome 472 replication was also sensitive to U18666A and cholesterol sequestration (this study and (35)) albeit 473 less than observed for CHIKV (data not shown). Conversely to CHIKV, this might not reflect 474 dependency on PM rafts, but instead cholesterol-induced alteration of endosomes where SINV 475 preferentially replicates as proposed by others (35, 64).

In the last decade, cholesterol-rich membrane microdomains, beyond their role in virus entry and exit, have also been identified as platforms for the assembly and anchoring of replication complexes produced by a broad range of RNA viruses including HCV, picornaviruses or flaviviruses (9, 10, 12, 18, 65). A dramatic reduction of viral RNA replication was observed upon membrane cholesterol extraction or in conditions reducing cholesterol availability (12, 14, 66, 67) establishing functional importance for this association. Besides changing membrane composition and fluidity, which may affect different interactions between virus-encoded replicase subunits, cholesterol partitioning in

483 membranes also attract cellular proteins, with functions in cell signaling and intracellular trafficking.
484 We can assume that targeting CHIKV nsP1 and possibly other nsPs at these sites may favor
485 interaction with raft-associated cellular factors indispensable for viral replication. The exact necessity
486 for nsP1 partitioning to DRMs therefore will deserve further investigations.

487

#### 488 Materials and Methods

489 Antibodies and reagents: The following antibodies/reagents and respective dilutions were used in 490 this study: rabbit polyclonal antisera against CHIKV nsP1, nsP2, nsP3, nsP4 (all in-house, 1:1,000), 491 monoclonal antibodies against GAPDH (1:1,000) (Santa Cruz Biotechnologies Inc.), Na<sup>+</sup> K<sup>+</sup> ATPase 492 (1:50,000) (Ab76020, Abcam), GFP (1:1,000) (Chromotek), Lamp2 (1:1,000), CD81 (1:500) (Clone JS-493 81, BD Biosciences) and flotillin-1 (1:1,000) (BD Biosciences). Secondary antibodies conjugated to 494 horseradish peroxidase or Alexa Fluor were purchased from Jackson Immunoresearch and Thermo 495 Fisher Scientific, respectively. Filipin III, U18666A, Lovastatin, methyl-β-cyclodextrin were purchased 496 from Sigma-Aldrich. Imipramine was obtained from Abcam and WGA-647 from Thermo Fisher 497 Scientific.

498 **Cells:** HEK293T cells (ATCC # ACS-4500), HeLa cells (ATCC # CRM-CCL2), BHK21 cells (ATCC # CCL-10) 499 and U2OS cells (ATCC # HTB-96) used for propagation and Vero cells (ATCC # CCL-81) used for 500 titration of the CHIKV were cultured in Dulbecco's modified Eagle's medium (DMEM, Thermo Fischer 501 Scientific) supplemented with penicillin and 10% fetal calf serum (FCS, Lonza) and grown at 37°C in a 502 5% CO<sub>2</sub> atmosphere. Viability of cells incubated with drugs for 24 hrs was measured using Cell Titer 503 96 Aqueous one solution cell proliferation assay (Promega) according to the manufacturer's protocol.

504 **Viruses:** The pCHIKV-LR-5'GFP, full-length molecular clone of CHIKV (LR2006OPY1 strain) with GFP 505 reporter (68), was linearized and transcribed *in vitro* using the mMESSAGE mMACHINE kit (Ambion-

Life Technologies). 1  $\mu$ g of RNA was then transfected with lipofectamine 2000 (Thermo Fisher Scientific) into 10<sup>5</sup> HEK293T cells and the cells were incubated at 37°C for 24 hrs. Culture medium was collected, and virus stock was amplified on BHK-21 cells. After 48 hrs at 37°C, the supernatant was collected, filtered through a 0.45  $\mu$ m membrane, aliquoted and stored at -80°C. Viral stocks were tittered using plaque assay as previously reported (39).

Infection with CHIKV-reporter viruses: The cells (70-80% confluence) were rinsed once with PBS before infection with CHIKV-LR-5'GFP diluted to achieve the desired MOI. For pre-infection experiments, the cells were pre-incubated with the virus for 1 hr or 2 hrs before drug addition. For post-infection experiments, the cells were incubated with drugs for 30 min before infection. After 24 hrs in culture, the cells were lysed with RIPA buffer. GFP reporter fluorescence was measured directly from the cell lysate using an Infinite F200PRO fluorometer (Tecan). Values were normalized to the protein content in the sample determined using the BCA Assay (Pierce).

Plasmids and transfection: Sequence encoding for CHIKV nsP1 was amplified by PCR using pCHIKV-LR-5'GFP as a template; obtained fragment was cloned into the pEGFP-C1 plasmid as previously described (69). Sequences encoding for a  $nsP1_{W258A}$ ,  $nsP1_{C417-419A}$  and  $nsP1_{DM}$  were generated using the Quikchange site-directed mutagenesis kit by Agilent. SINV nsP1 and  $nsP1_{420A}$  were obtained using PCR and pTOTO1101 (70) or its derivative pSINV-C<sub>420</sub>A1 (30) as templates. These inserts were cloned in frame with eGFP in the peGFP-C1 plasmid. Cells were transfected with obtained plasmids using JetPei reagent (Polyplus Transfection) according to manufacturer recommendations.

Trans-replication assay: For trans-replication assays, CMV-P1234 (71) or CMV-SINV-P1234 (72) plasmids encoding nonstructural polyprotein from CHIKV or SINV respectively were cotransfected together with the HSPolI-Fluc-Gluc (33) or HSPolI-SINV-tFluc-Gluc (73) template plasmids encoding for replication-competent template RNA of CHIKV or SINV containing firefly and *Gaussia* luciferase reporter sequences placed under control of genomic and subgenomic promoters respectively. Equal amounts of plasmids were transfected into HEK293T cells using JetPei transfection reagent. After 24

hrs in culture, cells were washed in PBS and lysed using Passive Lysis Buffer (Promega). Expression of firefly and *Gaussia* luciferase was determined using the Dual-Glo Luciferase Assay system (Promega) and a Spark luminometer (Tecan). Reporter activities were normalized to the protein content in the sample determined using the BCA Assay (Pierce).

Immunoblotting: Samples were separated by SDS-PAGE and then transferred to PVDF membrane (Hybond, Amersham). Membranes were blocked using 5% non-fat dry milk in PBS and probed with appropriate primary antibodies. After wash steps with PBS – 0.1% Tween 20, the membranes were probed with HRP-conjugated secondary antibodies. The revelation was performed by incubating the membranes with either Luminata Forte (Merck) or Clarity Max (BioRad) and then image acquisition was done using a Chemidoc (Bio-Rad).

541 Cell fractionation and membrane flotation assays: Cells were incubated in hypotonic buffer (10mM 542 Tris/HCI [pH 7.4], 10 mM NaCl supplemented with protease inhibitors) for 10 min on ice and then 543 lysed with a Dounce homogenizer (30-40 strokes). The lysates were clarified by low-speed 544 centrifugation at 1,000 g for 10 min. Obtained post-nuclear supernatants (PNS) were then adjusted 545 to a final concentration of 500 mM NaCl and incubated for 30 min on ice. After ultracentrifugation at 546 25,000 g for 20 min, the cytosolic (supernatant, S25) and membrane fraction (pellet, P25) were 547 collected. P25 samples were solubilized in lysis buffer composed of 1% Brij 96 in 20 mM Tris/HCI [pH 548 7.5] before analysis. For membrane flotation experiments, cells were resuspended in 250 mM 549 sucrose in PBS supplemented with protease inhibitors and then lysed with a Dounce homogenizer 550 (30-40 strokes). Cell lysates were spun at 1,000 g for 10 min to pellet the nuclei. The supernatant 551 referred to as crude lysate (CL) was then adjusted to 30% iodixanol concentration by mixing Optiprep 552 (Axis-Shield). CL (4 mL) was loaded at the bottom of a centrifuge tube, and then overlaid with 4 mL 553 20% iodixanol and then 4 mL 10% iodixanol. The gradient was then spun 200,000 g at 4°C for 16 hrs 554 in a Beckman SW41 rotor. Finally, 24 fractions were collected from top to bottom.

555 Detergent-resistant membrane isolation: Cells were lysed on ice in TNE buffer (10 mM Tris/HCI [pH 556 7.5], 100 mM NaCl, 10 mM EDTA) containing 0.5% Triton X-100 for 30 min. Lysates were then further 557 treated with the Dounce homogenizer, and then clarified by low-speed centrifugation at 1,000 g for 558 10 min to obtain the PNS. PNS (0.5 mL) was adjusted to 60% sucrose by adding 1.5 mL of 80% sucrose 559 TNE (w/v). The lysates were layered over 500  $\mu$ L of 80% sucrose TNE, then covered with 2 mL of 50% 560 sucrose TNE, 6 mL of 38% sucrose TNE, and 1.5 mL of 10% sucrose TNE. The sucrose gradients were 561 centrifuged at 100,000 g at 4°C for 18 hrs in a Beckman SW41 rotor (Beckman Coulter). Nine fractions 562 were then collected and analyzed by immunoblotting.

563 Immunofluorescence microscopy and image analysis: Cells grown on glass coverslips were washed 564 with PBS and then fixed with 4% paraformaldehyde/PBS (Sigma Aldrich) for 10 min. For intracellular 565 labeling, the cells were permeabilized with 0.1% Triton-X100 in PBS and blocked for 30 min with PBS 566 containing 0.2% Bovine Serum Albumin. Incubation with primary antibody was performed for 1 hr. 567 After washes with PBS, secondary reagents were added for 30 min. DAPI (Sigma-Aldrich) was used to 568 stain the nuclei. Filipin or WGA-647 staining was performed by incubation at room temperature for 569 either 1 hr or 10 min respectively. After final washes, coverslips were mounted with Prolong Gold 570 antifade mounting media (Thermo Fisher Scientific). Images were acquired using a Leica SP5-SMD 571 scanning confocal microscope equipped with a 63×, 1.4 numerical aperture Leica Apochromat oil 572 lense at the Montpellier Resources Imaging platform. Image analysis was performed utilizing Fiji 573 ImageJ and the JACoP plugin. For Mander's analysis, the threshold and region of interest were 574 maintained consistent between the different conditions. 3D reconstruction was performed by Imaris 575 software.

576 Statistical analysis: All of the analyses (unpaired Student's t-test) were performed using GraphPad
577 Prism version 6 (GraphPad Software Inc.). A p-value of <0.05 was considered statistically significant.</li>
578 Following designations are used on figures: \* p<0.05; \*\* p<0.001; \*\*\*\* p<0.0001; \*\*\*\* p<0.0001.</li>

579 Acknowledgments

- 580 We are indebted to Fabien Blanchet for helpful discussions and Joe McKellar for manuscript edition
- 581 (IRIM, Montpellier, France), to Cécile Gauthier-Rouvière (CRBM, Montpellier, France) for reagents
- 582 and to the Montpellier RIO Imaging platform staff for technical assistance.

#### 583 Authors contribution

584 Conceptualization: LB, AM; Methodology: LB; Funding acquisition: LB; Investigation-main 585 experiments: WB. Investigations - additional experiments: AN, EB; Supervision: LB; Writing original 586 draft: LB, AM, WB.

#### 587 Financial disclosure statement

- 588 The funders had no role in study design, data collection and interpretation, or the decision to submit
- the work for publication.

#### 590 Funding

- 591 This work was funded by the Agence Nationale de la Recherche (ANR-18-CE11-0026-01). WB is a
- 592 fellow of Montpellier Méditerranée Infections Infectiopôle Sud foundation

#### 593 Competing interests

- 594 The authors have declared that no competing interests exist.
- 595

#### 596 References

- Lorizate M, Kräusslich H-G. 2011. Role of Lipids in Virus Replication. Cold Spring Harb Perspect
   Biol 3:a004820.
- Paul D, Bartenschlager R. 2013. Architecture and biogenesis of plus-strand RNA virus replication
   factories. World J Virol 2:32–48.
- 601 3. Sasvari Z, Nagy PD. 2010. Making of viral replication organelles by remodeling interior 602 membranes. Viruses 2:2436–42.
- 603 4. de Castro IF, Volonte L, Risco C. 2013. Virus factories: biogenesis and structural design. Cell
  604 Microbiol 15:24–34.

- 6055.Goldstein JL, Brown MS. 1984. Progress in understanding the LDL receptor and HMG-CoA606reductase, two membrane proteins that regulate the plasma cholesterol. J Lipid Res 25:1450–6071461.
- 608 6. Infante RE, Wang ML, Radhakrishnan A, Kwon HJ, Brown MS, Goldstein JL. 2008. NPC2 facilitates
  609 bidirectional transfer of cholesterol between NPC1 and lipid bilayers, a step in cholesterol egress
  610 from lysosomes. Proc Natl Acad Sci USA 105:15287–15292.
- 611 7. Mercer J, Schelhaas M, Helenius A. 2010. Virus entry by endocytosis. Annu Rev Biochem 79:803–
  612 33.
- 8. Stoeck IK, Lee JY, Tabata K, Romero-Brey I, Paul D, Schult P, Lohmann V, Kaderali L,
  Bartenschlager R. 2018. Hepatitis C Virus Replication Depends on Endosomal Cholesterol
  Homeostasis. J Virol 92.
- 616 9. Aizaki H, Lee K-J, Sung VM-H, Ishiko H, Lai MMC. 2004. Characterization of the hepatitis C virus
   617 RNA replication complex associated with lipid rafts. Virology 324:450–461.
- 618 10. Shi ST, Lee K-J, Aizaki H, Hwang SB, Lai MMC. 2003. Hepatitis C Virus RNA Replication Occurs on
  619 a Detergent-Resistant Membrane That Cofractionates with Caveolin-2. J Virol 77:4160–4168.
- Mackenzie JM, Khromykh AA, Parton RG. 2007. Cholesterol manipulation by West Nile virus
   perturbs the cellular immune response. Cell Host Microbe 2:229–39.
- 12. Ilnytska O, Santiana M, Hsu NY, Du WL, Chen YH, Viktorova EG, Belov G, Brinker A, Storch J,
  Moore C, Dixon JL, Altan-Bonnet N. 2013. Enteroviruses harness the cellular endocytic
  machinery to remodel the host cell cholesterol landscape for effective viral replication. Cell Host
  Microbe 14:281–93.
- Ku K, Nagy PD. 2015. RNA virus replication depends on enrichment of
   phosphatidylethanolamine at replication sites in subcellular membranes. P Natl Acad Sci USA
   112:E1782–E1791.
- Albulescu L, Wubbolts R, van Kuppeveld FJ, Strating JR. 2015. Cholesterol shuttling is important
   for RNA replication of coxsackievirus B3 and encephalomyocarditis virus. Cell Microbiol
   17:1144–56.
- Rothwell C, LeBreton A, Young Ng C, Lim JYH, Liu W, Vasudevan S, Labow M, Gu F, Gaither LA.
   2009. Cholesterol biosynthesis modulation regulates dengue viral replication. Virology 389:8–19.
- 634 16. Wichit S, Hamel R, Bernard E, Talignani L, Diop F, Ferraris P, Liegeois F, Ekchariyawat P,
  635 Luplertlop N, Surasombatpattana P, Thomas F, Merits A, Choumet V, Roques P, Yssel H, Briant L,
  636 Misse D. 2017. Imipramine Inhibits Chikungunya Virus Replication in Human Skin Fibroblasts
  637 through Interference with Intracellular Cholesterol Trafficking. Sci Rep 7:3145.
- Takano T, Endoh M, Fukatsu H, Sakurada H, Doki T, Hohdatsu T. 2017. The cholesterol transport
  inhibitor U18666A inhibits type I feline coronavirus infection. Antiviral Research 145:96–102.
- 640 18. Sagan SM, Rouleau Y, Leggiadro C, Supekova L, Schultz PG, Su AI, Pezacki JP. 2006. The influence
  641 of cholesterol and lipid metabolism on host cell structure and hepatitis C virus replication.
  642 Biochem Cell Biol 84:67–79.
- 643 19. Weaver SC, Lecuit M. 2015. Chikungunya Virus and the Global Spread of a Mosquito-Borne
  644 Disease. N Engl J Med 372:1231–1239.
- 645 20. Froshauer S, Kartenbeck J, Helenius A. 1988. Alphavirus RNA replicase is located on the cytoplasmic surface of endosomes and lysosomes. J Cell Biol 107:2075–86.
- 647 21. Kujala P, Ikaheimonen A, Ehsani N, Vihinen H, Auvinen P, Kaariainen L. 2001. Biogenesis of the
  648 Semliki Forest virus RNA replication complex. J Virol 75:3873–84.
- 649 22. Spuul P, Balistreri G, Hellstrom K, Golubtsov AV, Jokitalo E, Ahola T. 2011. Assembly of 650 alphavirus replication complexes from RNA and protein components in a novel trans-replication

- 651 system in mammalian cells. J Virol 85:4739–51.
- Ahola T, Kaariainen L. 1995. Reaction in alphavirus mRNA capping: formation of a covalent
   complex of nonstructural protein nsP1 with 7-methyl-GMP. P Natl Acad Sci USA 92:507–11.
- Salonen A, Vasiljeva L, Merits A, Magden J, Jokitalo E, Kaariainen L. 2003. Properly folded
   nonstructural polyprotein directs the semliki forest virus replication complex to the endosomal
   compartment. J Virol 77:1691–702.
- 657 25. Lampio A, Kilpelainen I, Pesonen S, Karhi K, Auvinen P, Somerharju P, Kaariainen L. 2000.
  658 Membrane binding mechanism of an RNA virus-capping enzyme. J Biol Chem 275:37853–9.
- 659 26. Spuul P, Salonen A, Merits A, Jokitalo E, Kaariainen L, Ahola T. 2007. Role of the amphipathic
  660 peptide of Semliki forest virus replicase protein nsP1 in membrane association and virus
  661 replication. J Virol 81:872–83.
- 27. Laakkonen P, Ahola T, Kaariainen L. 1996. The effects of palmitoylation on membrane
   association of Semliki forest virus RNA capping enzyme. J Biol Chem 271:28567–71.
- Ahola T, Lampio A, Auvinen P, Kaariainen L. 1999. Semliki Forest virus mRNA capping enzyme
   requires association with anionic membrane phospholipids for activity. EMBO J 18:3164–72.
- Tomar S, Narwal M, Harms E, Smith JL, Kuhn RJ. 2011. Heterologous production, purification and
   characterization of enzymatically active Sindbis virus nonstructural protein nsP1. Protein Expr
   Purif 79:277–84.
- 30. Ahola T, Kujala P, Tuittila M, Blom T, Laakkonen P, Hinkkanen A, Auvinen P. 2000. Effects of
  Palmitoylation of Replicase Protein nsP1 on Alphavirus Infection. Journal of Virology 74:6725–
  671 6733.
- 31. Zusinaite E, Tints K, Kiiver K, Spuul P, Karo-Astover L, Merits A, Sarand I. 2007. Mutations at the
  palmitoylation site of non-structural protein nsP1 of Semliki Forest virus attenuate virus
  replication and cause accumulation of compensatory mutations. J Gen Virol 88:1977–85.
- 675 32. Karo-Astover L, Šarova O, Merits A, Žusinaite E. 2010. The infection of mammalian and insect 676 cells with SFV bearing nsP1 palmitoylation mutations. Virus Research 153:277–287.
- Age Utt, Rausalu K, Jakobson M, Männik A, Alphey L, Fragkoudis R, Merits A. 2019. Design and
  use of Chikungunya virus replication templates utilizing mammalian and mosquito RNA
  polymerase I mediated transcription. Journal of Virology JVI.00794-19.
- 680 34. Zhang N, Zhao H, Zhang L. 2019. Fatty Acid Synthase Promotes the Palmitoylation of 681 Chikungunya Virus nsP1. Journal of Virology 93:e01747-18.
- Solution
  Sol
- 685 36. Alberts AW. 1990. Lovastatin and Simvastatin Inhibitors of HMG CoA Reductase and
   686 Cholesterol Biosynthesis. CRD 77:14–21.
- Sobo K, Blanc IL, Luyet P-P, Fivaz M, Ferguson C, Parton RG, Gruenberg J, Goot FG van der. 2007.
  Late Endosomal Cholesterol Accumulation Leads to Impaired Intra-Endosomal Trafficking. PLOS
  ONE 2:e851.
- 690 38. Cenedella RJ. 2009. Cholesterol Synthesis Inhibitor U18666A and the Role of Sterol Metabolism
  691 and Trafficking in Numerous Pathophysiological Processes. Lipids 44:477–487.
- 692 39. Bernard E, Solignat M, Gay B, Chazal N, Higgs S, Devaux C, Briant L. 2010. Endocytosis of
  693 Chikungunya Virus into Mammalian Cells: Role of Clathrin and Early Endosomal Compartments.
  694 PLoS ONE 5.
- 695 40. Lu YE, Cassese T, Kielian M. 1999. The cholesterol requirement for sindbis virus entry and exit

- and characterization of a spike protein region involved in cholesterol dependence. J Virol73:4272-8.
- 698 41. Barton DJ, Sawicki SG, Sawicki DL. 1991. Solubilization and immunoprecipitation of alphavirus
   699 replication complexes. J Virol 65:1496–506.
- 42. Laakkonen P, Auvinen P, Kujala P, Kaariainen L. 1998. Alphavirus replicase protein NSP1 induces
   filopodia and rearrangement of actin filaments. J Virol 72:10265–9.
- Peranen J, Laakkonen P, Hyvonen M, Kaariainen L. 1995. The alphavirus replicase protein nsP1 is
   membrane-associated and has affinity to endocytic organelles. Virology 208:610–20.
- 44. Dermine J-F, Duclos S, Garin J, St-Louis F, Rea S, Parton RG, Desjardins M. 2001. Flotillin-1enriched Lipid Raft Domains Accumulate on Maturing Phagosomes. J Biol Chem 276:18507–
  18512.
- 45. Lu F, Liang Q, Abi-Mosleh L, Das A, De Brabander JK, Goldstein JL, Brown MS. 2015. Identification
   of NPC1 as the target of U18666A, an inhibitor of lysosomal cholesterol export and Ebola
   infection. eLife 4:e12177.
- 46. Schroeder F, Holland JF, Bieber LL. 1971. Fluorometric evidence for the binding of cholesterol to
  the filipin complex. J Antibiot 24:846–849.
- 47. Liscum L, Faust JR. 1989. The intracellular transport of low density lipoprotein-derived
  cholesterol is inhibited in Chinese hamster ovary cells cultured with 3-beta-[2(diethylamino)ethoxy]androst-5-en-17-one. J Biol Chem 264:11796–11806.
- 715 48. Charollais J, Goot FGVD. 2009. Palmitoylation of membrane proteins (Review). Molecular
   716 Membrane Biology 26:55–66.
- 49. Charrin S, Manié S, Thiele C, Billard M, Gerlier D, Boucheix C, Rubinstein E. 2003. A physical and
  functional link between cholesterol and tetraspanins. European Journal of Immunology
  33:2479–2489.
- 50. Eckert GP, Igbavboa U, Müller WE, Wood WG. 2003. Lipid rafts of purified mouse brain
  synaptosomes prepared with or without detergent reveal different lipid and protein domains.
  Brain Research 962:144–150.
- 51. Christian AE, Haynes MP, Phillips MC, Rothblat GH. 1997. Use of cyclodextrins for manipulating
   cellular cholesterol content. J Lipid Res 38:2264–2272.
- 52. Bartholomeeusen K, Utt A, Coppens S, Rausalu K, Vereecken K, Arien KK, Merits A. 2018. A
   Chikungunya Virus trans-Replicase System Reveals the Importance of Delayed Nonstructural
   Polyprotein Processing for Efficient Replication Complex Formation in Mosquito Cells. J Virol 92.
- Rana J, Rajasekharan S, Gulati S, Dudha N, Gupta A, Chaudhary VK, Gupta S. 2014. Network
   mapping among the functional domains of Chikungunya virus nonstructural proteins. Proteins
   82:2403–11.
- 54. Sreejith R, Rana J, Dudha N, Kumar K, Gabrani R, Sharma SK, Gupta A, Vrati S, Chaudhary VK,
  Gupta S. 2012. Mapping interactions of Chikungunya virus nonstructural proteins. Virus Res
  169:231–6.
- 55. Kumar S, Kumar A, Mamidi P, Tiwari A, Kumar S, Mayavannan A, Mudulli S, Singh AK, Subudhi
  BB, Chattopadhyay S. 2018. Chikungunya virus nsP1 interacts directly with nsP2 and modulates
  its ATPase activity. Scientific Reports 8:1045.
- 737 56. Remenyi R, Gao Y, Hughes RE, Curd A, Zothner C, Peckham M, Merits A, Harris M. 2018.
  738 Persistent Replication of a Chikungunya Virus Replicon in Human Cells Is Associated with
  739 Presence of Stable Cytoplasmic Granules Containing Nonstructural Protein 3. J Virol 92.
- 740 57. Pietila MK, Albulescu IC, Hemert MJV, Ahola T. 2017. Polyprotein Processing as a Determinant
   741 for in Vitro Activity of Semliki Forest Virus Replicase. Viruses 9.

- 58. Lingwood D, Simons K. 2010. Lipid Rafts As a Membrane-Organizing Principle. Science 327:46–
  50.
- 59. Hancock JF, Parton RG. 2005. Ras plasma membrane signalling platforms. Biochemical Journal
  389:1–11.
- 746 60. Simons K, Ikonen E. 1997. Functional rafts in cell membranes. Nature 387:569–572.
- 61. Levental I, Lingwood D, Grzybek M, Coskun Ü, Simons K. 2010. Palmitoylation regulates raft
  affinity for the majority of integral raft proteins. PNAS 107:22050–22054.
- Shenoy-Scaria AM, Dietzen DJ, Kwong J, Link DC, Lublin DM. 1994. Cysteine3 of Src family
   protein tyrosine kinase determines palmitoylation and localization in caveolae. The Journal of
   Cell Biology 126:353–363.
- Sobocińska J, Roszczenko-Jasińska P, Ciesielska A, Kwiatkowska K. 2018. Protein Palmitoylation
   and Its Role in Bacterial and Viral Infections. Front Immunol 8.
- 64. Thaa B, Biasiotto R, Eng K, Neuvonen M, Gotte B, Rheinemann L, Mutso M, Utt A, Varghese F,
  Balistreri G, Merits A, Ahola T, McInerney GM. 2015. Differential Phosphatidylinositol-3-KinaseAkt-mTOR Activation by Semliki Forest and Chikungunya Viruses Is Dependent on nsP3 and
  Connected to Replication Complex Internalization. J Virol 89:11420–37.
- 65. Cordero JG, Juárez ML, González-Y-Merchand JA, Barrón LC, Castañeda BG. 2014. Caveolin-1 in
   Lipid Rafts Interacts with Dengue Virus NS3 during Polyprotein Processing and Replication in
   HMEC-1 Cells. PLOS ONE 9:e90704.
- 66. Paul D, Hoppe S, Saher G, Krijnse-Locker J, Bartenschlager R. 2013. Morphological and
   biochemical characterization of the membranous hepatitis C virus replication compartment. J
   Virol 87:10612–27.
- 764 67. Ye J, Wang C, Sumpter R, Brown MS, Goldstein JL, Gale M. 2003. Disruption of hepatitis C virus
   765 RNA replication through inhibition of host protein geranylgeranylation. PNAS 100:15865–15870.
- 766 68. Vanlandingham DL, Tsetsarkin K, Hong C, Klingler K, McElroy KL, Lehane MJ, Higgs S. 2005.
  767 Development and characterization of a double subgenomic chikungunya virus infectious clone
  768 to express heterologous genes in Aedes aegypti mosquitoes. Insect Biochem Mol Biol 35:1162–
  769 70.
- 69. Matkovic R, Bernard E, Fontanel S, Eldin P, Chazal N, Hassan Hersi D, Merits A, Peloponese JM,
  Briant L. 2018. The host DHX9 DExH Box helicase is recruited to Chikungunya virus replication
  complexes for optimal genomic RNA translation. J Virol.
- 70. Polo JM, Davis NL, Rice CM, Huang HV, Johnston RE. 1988. Molecular analysis of Sindbis virus pathogenesis in neonatal mice by using virus recombinants constructed in vitro. Journal of Virology 62:2124–2133.
- 776 71. Utt A, Quirin T, Saul S, Hellstrom K, Ahola T, Merits A. 2016. Versatile Trans-Replication Systems
   777 for Chikungunya Virus Allow Functional Analysis and Tagging of Every Replicase Protein. PLoS
   778 ONE 11:e0151616.
- 779
   72. Götte B, Utt A, Fragkoudis R, Merits A, McInerney GM. 2020. Sensitivity of alphaviruses to G3BP
   780 deletion correlates with efficiency of replicase polyprotein processing. Journal of Virology.
- 781 73. Götte B, Utt, Age, Fragkoudis , Rennos, Merits, Andres, McInerney, Gerald M. Submitted to
   782 Journal of Virology JVI01681-19R1. Sensitivity of alphaviruses to G3BP deletion correlates with
   783 efficiency of replicase polyprotein processing.
- 784

785 Figure legends

786

#### 787 Figure 1: Cholesterol homeostasis regulates CHIKV life cycle

788 (A) Cholesterol metabolism and biosynthesis inhibitors. (B) Drugs affecting cholesterol metabolism 789 inhibit CHIKV infection. HEK293T cells treated for 30 min with the indicated concentrations of (a) 790 lovastatin, (c) U18666A, (e) imipramine were infected with CHIKV-LR-5'GFP at a MOI of 0.5. Infection 791 was monitored after 24 hrs by quantification of GFP reporter in the cell lysates. For each drug 792 concentration, cell viability was determined after 24 hrs using Cell Titer Glo assay (b, d, f). Values are 793 expressed as a percentage of control condition that was taken as 100%. A representative experiment 794 is shown. Values are means of triplicates ± SEM. p-values are calculated by comparing each treated-795 condition with the mock.

796

#### 797 Figure 2: Cholesterol transport inhibitors regulate intracellular CHIKV replication

798 (A) Post-infection addition of lovastatin, U18666A or imipramine reduces CHIKV replication. 799 Increasing concentrations of drugs were added to HEK293T cells 1 hr after CHIKV infection at a MOI 800 of 1. After 24 hrs, CHIKV replication was monitored by quantification of GFP fluorescence in the cell 801 lysates. Values, normalized to the protein content in the samples, are expressed as a percentage of 802 the non-infected (NI) condition. (B) Schematics of CMV-P1234 and HSPoll-Fluc-Gluc constructs, 803 plasmid backbones are shown. (C) HEK293T cells were transfected with plasmids depicted in (B) and 804 treated with the indicated concentrations of lovastatin, U18666A or imipramine at 2 hrs post-805 transfection. Firefly and Gaussia luciferase activities were determined after 24 hrs in culture. A 806 representative experiment is shown. Values are expressed as a percentage of the untreated 807 condition and are means of triplicates ± SEM. NT designates non-transfected control cells. p-values 808 are calculated by comparing each treated-condition with the mock.

#### 809 Figure 3: CHIKV nsP1 associates with the PM through conserved sequence motifs

810 (A) Organization of the CHIKV nsP1. Conserved amino-acids involved in the central  $\alpha$ -helix (aa 244-811 263) and those required for nsP1 acylation (417-419) are indicated. Mutants used in this study are 812 depicted. (B) Membrane affinity of GFP-nP1, GFP-nsP1<sub>3A</sub>, GFP-nsP1<sub>W258A</sub> and GFP-nsP1<sub>DM</sub> was 813 determined by cell fractionation assays. HEK293T cells transfected to express each GFP-fused protein 814 were fractionated to produce cytosolic (S25) and membranous (P25) fractions. Equal amounts of 815 fractions were analyzed by SDS-PAGE and western blotting with anti-nsP1 or anti-GFP antibodies. 816 Antibodies against Na<sup>+</sup>/K<sup>+</sup> ATPase (ATPase) and GAPDH were used to control the separation of 817 membranes and cytosolic compartments respectively. For each experiment a sample of the 818 unfractionated cell lysate (CL) was run in parallel to determine the relative amount of each nsP1 819 mutant expressed. Cells expressing the GFP protein alone are shown as a control. (C) The intensity of 820 bands in S25 and P25 samples from panel B was determined using Image J software and plotted as a 821 histogram. (D) Subcellular localization of GFP-nsP1 and its mutant was determined by confocal 822 imaging of transfected HeLa cells. Cell membranes are labeled with Alexa Fluor 647-conjugated 823 wheat germ hemagglutinin (WGA). Scale bars are 5 µm. (E) Confocal microscopy imaging of HeLa 824 cells transfected with expression plasmids for nsP1-GFP or for an untagged nsP1. The untagged nsP1 825 protein was detected with rabbit antiserum and Alexa-647-conjugated secondary reagents. Nuclei 826 were stained with DAPI before analysis by confocal imaging. GFP-nsP1 at the plasma membrane and 827 filopodia-like structures formed at the surface of transfected cells are indicated by white arrows. 828 Cytosolic aggregates are indicated by blue arrows. Bars: 5 µm. (F) Lysates of cells expressing GFP-829 nP1, GFP-nsP1<sub>3A</sub>, GFP-nsP1<sub>W258A</sub>, GFP were subjected to membrane flotation assay in iodixanol 830 gradient. Fractions collected from top to bottom were separated on SDS-PAGE and probed with 831 antibodies specific for nsP1. Lysates from GFP expressing cells were probed with antibodies specific 832 for GFP, flotillin-1 (FLOT1) or ATPase. Data are representative of three separate experiments.

#### 833 Figure 4: Cholesterol trafficking inhibitors redirect nsP1 to endo/lysosomal compartments

834 (A) HeLa cells cultured in the presence of 12  $\mu$ M U18666A were stained with filipin III to visualize cholesterol distribution. Cells cultured in medium containing an appropriate amount of vehicle are

836 shown as control (Mock). (B) Cells expressing the GFP-nsP1 protein were maintained for 24 hrs in the 837 presence of 12  $\mu$ M U18666A and then co-labeled with filipin III and anti-Lamp2 antibodies to 838 visualize cholesterol and endo/lysosomes respectively. Cells maintained with vehicle alone are 839 shown as controls (Mock). (C) Consequences of U18666A treatment on GFP-nsP1 membrane binding 840 capacity was determined by cell fractionation assays. Crude lysates (CL) and cytosolic fractions (S25) 841 prepared from mock-treated or U18666A-treated HEK293T cells were separated on SDS-PAGE and 842 probed with antibodies against GFP or GAPDH. (D) HeLa cells transfected with GFP or GFP-nsP1 expression plasmid and maintained with 12  $\mu$ M U18666A or in medium alone (Mock) for 24 hrs were 843 844 stained with filipin III and DAPI and then processed for confocal imaging. (E) Histograms indicate the 845 signal intensity profile of filipin III and green fluorescence along the white lines in (D). Bars: 5 µm. (F) 846 Colocalization of filipin and GFP signals in U18666A-treated cells was determined by calculation of 847 Mander's overlap coefficients ( $n \ge 20$  cells).

## 848 Figure 5: GFP-nsP1<sub>W258A</sub> and GFP-nsP1<sub>3A</sub> display different sensitivity to U18666A-induced 849 cholesterol storage defect.

HeLa cells expressing either GFP-nsP1<sub>3A</sub> (A) or GFP-nsP1<sub>W258A</sub> (C) were incubated with 12  $\mu$ M U18666A for 24 hrs. Unesterified cholesterol was stained with filipin-III and cells were processed for confocal microscopy. (B) and (D) Histograms showing filipin III and GFP fluorescence intensity along the white lines indicated in (A) and (B) respectively. Bars: 5  $\mu$ m. (E) Colocalization of filipin and GFP signals in U18666A-treated cells was determined by calculation of Mander's overlap coefficients (n ≥ 20 cells).

# Figure 6: C417-419 are key determinants for CHIKV nsP1 targeting to CD81-positive cholesterol-rich compartments

 $U_{2OS}$  cells expressing (A) the GFP, (B) GFP-nsP1, (C) GFP-nsP1<sub>3A</sub> were cultured in control condition or in medium supplemented with 12  $\mu$ M U18666A or 75  $\mu$ M imipramine for 24 hrs. Then, the cells were probed with anti-CD81 mAbs and Alexa Fluor-594-conjugated secondary antibodies. Nuclei were stained with DAPI and the cells were processed for confocal microscopy. Scale bars are 5  $\mu$ m. (D) Quantification of the degrees of colocalization of GFP fluorescence and CD81 staining in treated cells was determined by calculation of Mander's and Pearson's overlap coefficients (n  $\ge$  20 cells). (E and F) 3D image reconstruction was performed from z-stacks acquired from treated cells expressing either GFP-nsP1 or GFP-nsP1<sub>3A</sub>.

#### 866 Figure 7: CHIKV nsP1 partitioning with lipid rafts in human cells requires C417-419

867 (A) HEK293T cells were transfected with plasmids encoding GFP-nsP1 and processed for raft 868 isolation. Nine fractions of the density gradients, numbered from top to bottom, were collected and 869 analyzed using SDS-PAGE. Proteins were probed with antibodies against nsP1, FLOT1 and Na+/K+ 870 ATPase. Fractions containing detergent-resistant membranes (DRMs) and detergent soluble 871 membranes (DS) are indicated. (B) The amount of GFP-nsP1 and FLOT1 associated with each fraction 872 was determined by densitometry scanning of immunoblots and expressed as a percentage of total 873 protein expression level. The diagram is representative of three experiments. (C, D) A similar 874 experiment was performed starting from cells incubated for 30 minutes in the presence of 10 mM 875  $\beta$ MCD. (E) Cells expressing the GFP-nsP1<sub>3A</sub> or GFP-nsP1<sub>W258A</sub> mutants were processed as in (A). (F) 876 Amounts of GFP-nsP1, GFP-nsP1<sub>3A</sub> or GFP-nsP1<sub>W258A</sub> in DRM and DS fractions were determined by 877 densitometry scanning of the immunoblots shown in (A) and (E) and expressed as a percentage of 878 total protein signal.

# Figure 8: nsPs expressed by a CHIKV trans-replicase or an infectious CHIKV co-segregate with DRMs depending of nsP1 C417-419. DRMs were isolated from (A) HEK293T cells transfected with CMV-P1234 and HSPolI-Fluc-Gluc plasmids (depicted in Fig 2B); (B) cells infected with CHIKV-LR-5'GFP (MOI 0.1); (C) cells transfected with CMV-P1<sub>3A</sub>234 and HSPolI-Fluc-Gluc plasmids. After 24 hrs, fractions were prepared and analyzed as described for Fig 7A, immunoblots were probed with antibodies against each CHIKV nonstructural protein as indicated. (D and E) Amounts of nsPs in DRM

and DS fractions were determined by densitometry scanning of the immunoblots shown in (A) and(B) respectively and expressed as a percentage of total protein signal.

887 Figure 9: SINV nsP1 affinity for cholesterol. (A) Organization of the SINV nsP1 and mutant studied. 888 Conserved amino-acids involved in the central  $\alpha$ -helix (aa 245-264) and palmitoylated cysteine (C420) 889 are indicated. Mutant used in this study is depicted. (B) HEK293T cells transfected with either SINV 890 GFP-nsP1 or GFP-nsP1<sub>C420A</sub> plasmids were processed for DRMs isolation as described for Fig 7. 891 Fractions (numbered from top to bottom) were probed in immunoblot with antibodies against GFP. 892 (C) HeLa cells were transfected with a plasmid encoding a SINV GFP-nsP1 protein for 4 hrs and 893 cultured for an additional 24 hrs in the presence of 12  $\mu$ M U18666A. Cells were labeled with filipin III 894 to visualize cholesterol. Controls cells (Mock) were maintained with vehicle. (D) HeLa cells 895 transfected to express the GFP-nsP1<sub>C420A</sub> mutant were processed as in (C). Bars: 5  $\mu$ m.

896 Figure 10: Model for nsP1/cholesterol interplay in transfected and CHIKV-infected cells. a) 897 Palmitoylated cysteines in wild type nsP1 are critical for targeting to cholesterol-rich membranes 898 microdomains. In infected cells, the preferential association of nsP1 with DRMs increases its local 899 concentration which may facilitate its oligomerization and dictates the localization of a fraction of 900 other nsPs to lipid rafts. In this scenario CHIKV efficiently replicates its genome. b) Imipramine and 901 U18666A, by inhibiting cholesterol trafficking to the PM, generate nsP1 accumulation in Lamp2-902 positive late endosomes/lysosomes. Association of nsP2, nsP3 and nsP4 to cholesterol rich 903 microdomains is no more observed. Under these conditions, CHIKV genome replication is abolished 904 by a mechanism that remains to be fully elucidated. c) nsP1<sub>3A</sub> that weakly interacts with the plasma 905 membrane is not targeted to DRMs. In this situation CHIKV genomes replication is impaired. For 906 simplicity, only mature nsPs are shown in this model. In natural infection, the membrane association 907 begins by modification and membrane targeting of polyprotein precursors.



Β





















F

С

Ε



Α







D









**Distance (Pixels)** 





















