



Geometric categories in cognition

Moira Dillon, Marianne Duyck, Stanislas Dehaene, Marie Amalric, Véronique Izard

► To cite this version:

Moira Dillon, Marianne Duyck, Stanislas Dehaene, Marie Amalric, Véronique Izard. Geometric categories in cognition. *Journal of Experimental Psychology. Human Perception and Performance*, 2019, 45 (9), pp.1236-1247. <10.1037/xhp0000663>. <hal-02997511>

HAL Id: hal-02997511

<https://hal.science/hal-02997511v1>

Submitted on 19 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



AMERICAN
PSYCHOLOGICAL
ASSOCIATION

Journal of Experimental Psychology: Human Perception and Performance

Manuscript version of

Geometric Categories in Cognition

Moira R. Dillon, Marianne Duyck, Stanislas Dehaene, Marie Amalric, Véronique Izard

Funded by:

- European Research Council
- National Science Foundation
- Norman Henry Anderson Graduate Psychology Fund

© 2019, American Psychological Association. This manuscript is not the copy of record and may not exactly replicate the final, authoritative version of the article. Please do not copy or cite without authors' permission. The final version of record is available via its DOI: <https://dx.doi.org/10.1037/xhp0000663>

This article is intended solely for the personal use of the individual user and is not to be disseminated broadly.



CHORUS *Advancing Public Access to Research*

Title Geometric categories in cognition

Authors Moira R. Dillon^{1*†}, Marianne Duyck^{2,3,4*}, Stanislas Dehaene^{5,6}, Marie Amalric^{6,7},
Véronique Izard^{2,3}

Affiliations

¹Department of Psychology, New York University, New York, NY, USA

²CNRS (Integrative Neuroscience and Cognition Center, UMR 8002), Paris, France

³Université Paris Descartes, Sorbonne Paris Cité, Paris, France

⁴Laboratory of Sensorimotor Research, NEI/NIH, Bethesda, USA.

⁵Collège de France, Paris, France

⁶Cognitive Neuroimaging Unit, CEA DSV/I2BM, INSERM, Université Paris-Sud, Université
Paris-Saclay, NeuroSpin Center, Gif/Yvette, France

⁷Sorbonne Universités, UPMC Université Paris 6, IFD, 4 Place Jussieu, Paris, France

*These authors contributed equally and are listed in alphabetical order.

†To whom correspondence should be addressed:

Moira Dillon

Department of Psychology, New York University

6 Washington Place

New York, NY 10003

USA

Abstract

At the scale in which we live, space is continuous. Nevertheless, our perception and cognition parse the world into categories, whether physical, like *scene* or *object*, or abstract, like *infinitesimal point* or *seven*. The present study focuses on two categories of special angles in planar geometry, *parallels* and *perpendiculars*, and we evaluate how these categories might be reflected in adults' basic angle discrimination. In the first experiment, participants were most precise when detecting two parallel or perpendicular lines among other pairs of lines at different relative orientations. Detection was also enhanced for two connected lines whose angle approached 90°, with precision peaking at 90°. These patterns emerged despite large variations in the scales and orientations of the angle exemplars. In the second experiment, the enhanced detection of perpendiculars persisted when stimuli were rotated in depth, indicating a capacity to discriminate shapes based on perpendicularity in three dimensions despite large variation in angles' two-dimensional projections. The results suggest that two categorical concepts which lie at the foundation of Euclidean geometry, parallelism and perpendicularity, are reflected in our discrimination of simple visual forms, and they pave the way for future studies exploring the developmental and evolutionary origins of these cognitive categories.

Keywords: categories; shape discrimination; spatial cognition; angle; geometry; parallel; perpendicular; **Significance:** In this paper, we discover that two categories of special angles in formal geometry, *parallels* and *perpendiculars*, are robustly reflected in adults' basic angle discrimination. Our results both characterize the psychophysical properties of angle discrimination, which has been debated in prior literature, and also link vision research to the conceptual, formal, and school-relevant spatial understanding that supports abstract mathematics.

Introduction

Categories of all kinds pervade human cognition and organize our physical and mental worlds. Some categories are present early in human development and are rooted in perception. For example, very young infants show more precise discrimination (indicated by an increase in sucking frequency) of two synthetic speech sounds separated by a fixed distance in voice onset time when that distance distinguishes voiced and unvoiced stop consonants (e.g. /b/ and /p/) compared to when it does not (Eimas et al., 1971; Liberman et al., 1961). Moreover, even early brain responses are modulated by categories of visual stimuli: Localized regions of the visual cortex of infants as young as four months respond preferentially to exemplar pictures belonging to categories like *faces* and *scenes* (Deen et al., 2017). Nevertheless, such early emerging categorical processing is reshaped by experience. For example, in the first year of life, infants' auditory discrimination becomes specialized to the phonemic categories present in their native language (Kuhl et al., 1992; Werker & Tees, 1984), and children's explicit discrimination of phonemes develops reciprocally with culturally constructed reading and writing systems (see Anthony & Francis, 2005 for a review). Such complex developmental stories raise intriguing questions about the origins of our explicit categorical knowledge in adulthood: To what extent is such knowledge innate, rooted in our perceptual experiences, or acquired by explicit education or immersion in a specific language and culture? And what of those categories with no apparent perceptual origins or constraints, like the concept of *irrational number* or other such abstract concepts often found in formal mathematics? In the present study, we investigate whether two categories of special angles that lie at the foundation of formal, Euclidean geometry, *parallelism* and *perpendicularity* (Euclid, 1990/300 B.C.E.), are reflected in adults' basic angle

discrimination. We do so by evaluating whether adults' discrimination acuity is enhanced around these category boundaries in a variety of perceptual contexts.

At the turn of the 20th century, Wilhelm Wundt and Hermann von Helmholtz independently suggested that angle discrimination may not be a wholly continuous process. They observed that small angles are judged to be somewhat bigger than their actual size and that big angles are judged to be somewhat smaller (von Helmholtz, 1897; Wundt, 1897). More recent work on angle discrimination has aimed to explain and quantify this observation by suggesting, for example, that errors in discrimination reflect orientation selectivity in the visual cortex that leads to orientation distortions (Carpenter & Blakemore, 1973) or that the errors are rooted in an inference about how an angle's appearance reflects its real-world size (Howe & Purves, 2005; Nundy, Lotto, Coppola, Shimpf, & Purves, 2000).

A number of studies have focused on comparing the threshold at which we are able to discriminate various angles, but the results have been mixed. In addition, such studies often incompletely controlled for a variety of visual cues in the angle stimuli, such as orientation or size, and so conclusions about angle discrimination specifically have been difficult to make (Werkhoven & Koenderink, 1993; Wenderoth & Johnson, 1984; Snippe & Koenderink 1994; Regan & Hamstra 1992). Chen & Levi (1996) suggested, for example, that there are different detection thresholds for angles of specific sizes, reporting more fine-grained discrimination of 90° angles in a discrimination space that otherwise follows Weber's Law. While such thresholds were measured for 12 different reference angles (ranging from 15° -180°), the angles were presented at only two different orientations, vertical or oblique, so it remains unclear whether the thresholds were generalizable to more variable orientations. Heeley & Buchanan-Smith (1996) found a similar pattern of discrimination thresholds with angles presented at random orientations,

but they did not simultaneously vary the lengths of the lines that formed the angles, allowing for other, global shape cues to drive participants' performance. Regan, Gray, & Hamstra (1996), in contrast, found fairly constant thresholds of angle discrimination between 20° and 160° when angles were presented at a wide range of orientations. The authors suggest, however, that their discrepant findings might have been due to their unique methodology: Participants compared each stimulus to a reference angle size internalized through many practice trials, rather than to a physical display. The existing literature thus suggests that discrimination may be most precise around 90° angles, but it remains inconclusive.

In the present study, we measured the discrimination thresholds for many different angles using an intruder task (after Dehaene, Izard, Pica, & Spelke, 2006). We evaluated whether participants could locate an intruder that differed in angle size among five other angle exemplars of the same angle size as each other. The lines forming the angles varied considerably in their lengths and orientations to ensure that responses were made on the basis of angle alone within an individual trial. In addition, no individual lines were oriented within 10° of the horizontal or vertical to avoid any specialized angle discrimination that might occur when individual lines are orientated at the horizontal or vertical (Xu, Chen, & Kuai, 2018).

We hypothesized that angle discrimination may reflect categories of parallelism and perpendicularity in two ways. First, detection thresholds may be more precise when parallels or perpendiculars serve as reference angles, compared to intruder angles of other sizes. Second, the detection of angle intruders may be asymmetric such that, for reference angles near 0°/180° and 90°, intruders whose angle size moves towards versus away from the parallel or perpendicular category boundaries will be easier to detect. This second prediction should hold when intruders cross the category boundary and also, perhaps, when intruders approach but do not cross the

boundary. In Experiment 1, we find evidence for such categories. In Experiment 2, we demonstrate the robustness of these categories by imposing an additional rotation in depth in the experimental displays, dissociating 3D perpendicularity from 2D angle.

Experiment 1

Methods

Participants

Eight adults (four women; $M_{age} = 25$ years; range 19-28 years) participated in this experiment. The sample size was set in advance based on the maximum sample size (8 participants) used in several other studies investigating angle discrimination and presenting large numbers of trials to individual participants (Chen & Levi, 1996; Regan, Gray, & Hamstra, 1996; Heeley & Buchanan-Smith, 1996; Snippe & Koenderink 1994; Regan & Hamstra 1992, Xu et al., 2018). In addition, stimuli were piloted in advance on three of the study's authors, and effects were robust enough to emerge in each pilot participant. As such, we also illustrate individual participant results in the figures and **SM**. All participants had normal or corrected-to-normal vision and had completed high school; most of them had also received a college or advanced degree. None were informed of the purpose and hypotheses of the study until after it was completed. This study was part of a research program approved by the Paris Descartes Ethics Committee (Conseil d'Évaluation Éthique pour les Recherches En Santé, CERES) and each participant provided informed consent prior to the experiment. Participants were paid 10€ per hour plus an additional sum, which depended on their performance (see below). On average, participants earned 77.38€ (range 75.60€-80.07€).

Design, Apparatus, Procedure, and Stimuli

For each trial, participants were presented with six angles, five of which were identical in their angle size (hereafter referred to as the “reference” angle) and one of which differed in its angle size (hereafter referred to as the “deviant” angle). Participants were asked to identify the deviant angle. In the *connected lines* condition (**Figure 1A**), angles were composed of two lines that met at one end. Participants saw trials in seven different blocks, each block with a different reference angle. The angle measures of the stimuli are summarized **Table 1**. The reference angles were chosen to evaluate discrimination thresholds symmetrically around 90°, and the angle differences between the reference angles and each deviant were chosen to capture potentially large differences in discrimination thresholds across the different references (if, e.g., discrimination followed Weber’s Law). To probe the detection of 90° deviants specifically, +/-10° were presented in the 80° and 100° reference blocks. In the *non-connected lines* condition (**Figure 1B**), two lines were oriented relative to one another, but did not meet, and participants saw eight different reference angles in separate blocks (**Table 1**). We again added 90° deviants in the 80° and 100° blocks. We also added 0° (parallel) deviants in the 10° block and removed from that block the -12° and -18° deviants since these values were not geometrically possible (i.e., they would have resulted in -2° and -8° deviants, which would be equivalent to 2° and 8° angles). Each type of deviant was presented 18 times in a random order in a block, three times at each of six possible target locations. The number of trials per block thus ranged from 126 to 162. Finally, despite the addition of 0° and 90° deviants in some reference blocks, those angles were not over-represented in the experiment. For example, in the connected lines condition, a total of 756 90° exemplars were presented, compared to 810 exemplars of 80° and 100° angles and 720 exemplars of the angles in the other reference blocks (**Table 1**).

		Reference Angles								
		0	10	25	55	80	90	100	125	155
Deviant Angle Difference	-18	18		7	37	62	72	82	107	137
	-12	12		13	43	68	78	88	113	143
	-10		0					90		
	-7	7	3	18	48	73	83	93	118	148
	-3	3	7	22	52	77	87	97	122	152
	+3	3	13	28	58	83	93	103	128	158
	+7	7	17	32	62	87	97	107	132	162
	+10					90				
	+12	12	22	37	67	92	102	112	137	167
	+18	18	28	43	73	98	108	118	143	173

Table 1. Summary of the stimuli presented in Experiments 1 & 2. The 25° and 155° reference angles were presented only in Experiment 1 and the 0° and 10° reference angles were presented only in the non-connected lines condition of Experiment 1. All angle measures are in degrees.




Figure 1. A. Exemplar trials from the connected lines condition. Participants showed smaller discrimination thresholds with 90° reference angles (top) than with 80° reference angles

(bottom). **B.** Exemplar trials from the non-connected lines condition. Participants easily distinguished non-parallel lines among parallel lines (top) but had difficulty with larger deviants when reference angles were 10° (bottom). Here, deviants differ from the reference angles by 24° (the difference used during training trials). For illustration purposes, the correct responses are circled in red.

Participants were seated in a lit room (62 cd.m^{-2}) at eye level and 48 cm away from the center of an LCD monitor (60 Hz) subtending 44×32 degrees of visual angle (dva). On every trial, six angles appeared at the same time in one of the six white circular placeholders (124 cd.m^{-2} , 5 dva radius). These placeholders were equally distributed around a 11 dva radius circle centered on a central black fixation dot (0.15 cd.m^{-2} , 0.4 dva radius) on a light grey screen (92 cd.m^{-2}). While maintaining their head position with a chin rest, participants were given 5 s to look at all of the angles before the angles disappeared. Participants could respond 250 ms after the presentation onset or up to 30 s after the angles disappeared. At the beginning of each trial, the cursor was positioned at the center of the screen. Participants clicked on the location of the deviant angle and received informative auditory feedback. The next trial started 500 ms after the response. After every quarter of a block, the percentage of correct responding was displayed as well as the sum earned during that block. Participants earned 0.014€ per correct response.

In both conditions, the lines that formed each angle were the same length, but these lengths varied across angles in the same display (chosen randomly from a uniform distribution between 2 and 4.4 dva). Lines were never within 10° of the vertical or horizontal axes of the screen. In the non-connected lines condition, one of the lines was displaced relative to the other line both along it (up to its midpoint) and orthogonally (0.6-2.5 dva) to it. Finally, to ensure that all angles in each display were presented at sufficiently different orientations, each was initially assigned to either 0° , 60° , 120° , 180° , 240° , and 300° relative to the screen and then each was

jittered randomly between $\pm 30^\circ$. Each angle was then translated to a random location within 0.4 dva of its placeholder's edge.

The experiment was divided into two sessions corresponding to the two conditions, which took place on different days within a two-week period. The order of conditions was counterbalanced across participants. Each session started with a short training phase consisting of two trials per reference angle displaying $\pm 24^\circ$ angle deviants. These trials were first presented with unlimited viewing time and then with the 5 s viewing time used in the actual experiment. Each block started with an introductory screen instructing participants to click on the shape that had a different angle size from the rest. This introductory screen also displayed one example of the reference angle for that block, oriented such that a vertical line would bisect the angle. The order of the blocks was random for each participant. Each session lasted approximately 3 hours, with a 15-30 min break after the fourth block in each condition.

Analyses

Participants' performance for each reference angle and for both the connected and non-connected line conditions was fit with down-pointing Gaussian curves constrained to chance performance (chance = 0.167) at a difference of zero degrees and perfect performance at a difference of infinity. The fitted Gaussians were used to estimate individual participants' thresholds. These thresholds corresponded to the difference in degrees between the reference and the deviant such that the participant could detect the deviant on half of the trials and was guessing on the other half (i.e., performance of 0.583, halfway between chance and perfect performance). To capture possible asymmetries in participants' responses, performance was fit separately for smaller and larger deviants. Overall thresholds for each reference were obtained by

averaging these two values. Because we did not present the same number and measure of angle deviants across reference angles (e.g., we included 10° deviants in the 80° block, but not in the 55° block), such thresholds were more appropriate than raw accuracies to compare performance across reference angles. This measure was decided on after pilot testing and in advance of any data collection.

In the connected lines condition, all eight participants performed above chance on all seven reference angle conditions with both smaller and larger deviants (binomial tests, two-tailed, all $ps < 0.05$), yielding 112 data sets with above-chance performance. In the non-connected lines condition, however, 17 of the 128 data sets did not significantly differ from chance, thus yielding unreliable detection thresholds (see **Figure 2** for the accuracy curves of a representative participant and **Figure S1** for all individual accuracy curves). To analyze the results of the non-connected lines condition, we thus used non-parametric, rank-order tests instead of parametric tests, which replaced estimated threshold values with ranks based on thresholds' relative magnitudes. As confirmation of our results for this condition using this method, we also analyzed participants' accuracy using parametric tests, and we obtained the same results (see **SM, Table S1, Figure S2**).

First, we evaluated participants' performance in the connected lines condition. Using planned one-tailed t -tests, we examined whether participants' detection thresholds were more precise for the perpendicular (90°) reference angles compared to the 80° and 100° reference angles (the closest references to 90°, a 10° difference) and to the 55° and 125° reference angles (references further from 90°, a 35° difference). In those same reference blocks, we then examined whether deviant detection was more precise when deviants approached or crossed the 90° boundary compared to when they did not. For example, we evaluated whether thresholds in

detecting larger deviants in 80° reference blocks were more precise than detecting smaller deviants in 80° reference blocks. Since all four comparisons tested whether angle discrimination was influenced by the category of right angles, p -values were adjusted for multiple comparisons using Holm's method.

For the non-connected lines condition, we investigated whether detection thresholds were more precise for the perpendicular (90°) and parallel (0°) reference angles using planned Holm-corrected, one-tailed Wilcoxon signed-ranks tests. As in the connected lines condition, we compared the thresholds of detecting deviants with a 90° reference both to those with an 80°/100° reference and also to those with a 55°/125° reference. In addition, we also compared the thresholds of detecting deviants with a 0° reference to the two closest reference blocks (10° and 25°). Finally, we tested for asymmetries in detection thresholds around these categories by examining whether deviant detection was more precise when deviants approached or crossed the 90° or 0° boundaries compared to when they did not.




Figure 2. Model-fit curves for one participant (S2) for all reference angles in the connected lines condition of Experiment 1. Individual data points are additionally represented for the 80°, 90°, and 100° reference blocks. These curves illustrate lower thresholds in the 90° reference block (i.e., a more narrow curve, in yellow) and lower thresholds for deviants towards versus away from 90° (i.e., an asymmetry in the curves around 0° of deviation) in the 80° (in green) and 100° (in orange) reference blocks.

Results

Connected Lines Condition

Figure 3A displays the individual and average detection thresholds for each angle reference in the connected lines condition. All participants had similarly shaped curves, in which detection thresholds were smaller as reference angles approached 0°/180°. Strikingly, all participants also showed steep drops in their detection thresholds as angle references approached 90°. Group-wise analyses, summarized in **Table 2**, corroborated these results, finding significantly more precise detection thresholds for 90° reference angles compared to 80°/100°

reference angles ($t(7) = 5.47, p < .001$, Cohen's $d = 1.94$) and $55^\circ/125^\circ$ reference angles ($t(7) = 10.05, p < .001$, Cohen's $d = 3.55$). Such differences are characteristic of a categorical effect at 90° .

Moreover, we observed significant asymmetries in the blocks where the deviants crossed the 90° boundary (80° and 100° references). In these blocks, deviant detection was more precise towards 90° compared to away from it (**Table 2**; $t(7) = -6.29, p < .001$, Cohen's $d = 2.23$). However, no asymmetries were observed in the $55^\circ/125^\circ$ reference blocks ($t(7) = 0.21, p = .419$, Cohen's $d = 0.07$). Thus, as deviants crossed 90° , their discrimination became more precise, again signaling an influence of the perpendicular category on performance (see **Figure 3B**).

Perpendicular			
	90	80/100	55/125
Connected Lines – thresholds	5	8.9***	11.2***
Connected Lines – asymmetry		toward: 6.9; away: 10.8**	toward: 11.1; away: 11.2
Non-Connected Lines – thresholds	8	13.6*	20.7*
Non-Connected Lines – asymmetry		toward: 10.1; away: 16.6*	toward: 19.5; away: 19.0
Parallel			
	0	10	25
Non-Connected Lines – thresholds	3	13.3*	20.0*
Non-Connected Lines – asymmetry		toward: 5.1; away: 21.2*	toward: 12.1; away: 28.4*

Table 2. Group-wise mean (for the connected lines condition) or median (for the non-connected lines) detection thresholds for perpendicular and parallel reference angle blocks compared to other reference blocks and detection asymmetries. Planned, Holm-corrected one-tailed t-tests (for the connected lines condition) or Wilcoxon signed-ranks tests (for the non-connected lines condition) compare the 90° or 0° reference blocks to the other reference blocks and the magnitude of the asymmetry effects. All angle measures are in degrees. * $p < .05$, ** $p < .01$, *** $p < .001$.




Figure 3. A. Estimated detection thresholds for each participant (grey lines) and on average (black line) at different reference angles for the connected lines condition. All curves are characterized by threshold drops towards 0°/180° as well as a sharp drop at 90°. The dashed curve corresponds to the participant shown in **Fig. 2. B.** Asymmetries in the thresholds for detecting smaller versus larger deviants at different reference angles. Positive values on the y-axis indicate greater success in detecting larger versus smaller deviants, and negative values indicate the opposite. The dashed curve corresponds to the participant shown in **Fig. 2.** For 80° and 100° references, acuity is better (smaller thresholds) when the deviant is in the direction of 90°. The asymmetry in detection thresholds crosses zero at almost exactly 90°, and there appears to be an approximately identical advantage on both sides of 90°.

Non-Connected Lines Condition

Figure 4A displays the individual and median detection thresholds for each reference angle in the non-connected lines condition. All participants had similarly shaped curves, in which detection thresholds were smallest at 0° compared to other reference angles (see **Figure S1** for individual participant accuracy curves). Group-wise analyses corroborated these results (see **Table 2**), finding more precise detection thresholds for 0° reference angles compared to 10° and 25° reference angles (10°: $Z = 2.15$, $p = .016$, 8/8 participants showed the effect; 25°: $Z = 2.15$, $p = .016$, 8/8 participants showed the effect). Moreover, participants performed better with angles deviating towards versus away from 0° with both 10° and 25° reference angles (10°: $Z =$

2.15, $p = .016$, 8/8 participants showed the effect; 25°: $Z = 2.15$, $p = .016$, 8/8 participants showed the effect; see **Figure 4B**) suggesting that parallel lines might serve as an anchor for judgments of discrimination.

Detection thresholds were quite variable across participants between 10° and 80°, though they exhibit a similar shape, first increasing at smaller reference angles and then decreasing at larger reference angles. Evidently, acute angles are difficult to differentiate with non-connected lines. Nevertheless, as in the connected lines condition, participants' precision increased around 90° compared to 80°/100° reference angles ($Z = 2.15$, $p = .016$; 8/8 participants showed the effect) and to 55°/125° reference angles ($Z = 2.15$, $p = .016$; 8/8 participants showed the effect; see **Table 2**). We also observed significant asymmetries in detection thresholds, with better discrimination as deviants approached or crossed the 90° boundary in the 80°/100° reference blocks ($Z = 2.15$, $p = .016$, 8/8 participants showed the effect), but not in the 55°/125° reference blocks ($Z = 0.07$, $p = .473$, 4/8 participants showed the effect; see **Fig. 4B**).




Figure 4. A. Estimated detection thresholds for each participant (grey lines) and the group medians (black line) at different reference angles for the non-connected lines condition. Thresholds at some reference angles for some individuals exceed the graph limit. All curves are characterized by low thresholds for the 0° reference angle as well as a sharp drop at the 90° reference angle. **B.** Asymmetries in the detection thresholds for larger versus smaller deviants at

different reference angles. The detection threshold asymmetry crosses zero at around 90°, with an approximately equal advantage on both sides.

Discussion

This experiment presents two main findings. First, acuity in angle discrimination varies massively across angles. In particular, thresholds are smallest for parallels and perpendiculars. When angles are acute or obtuse and far from these two categories, thresholds of discrimination are high, though when angles deviate from such acute or obtuse angles and approach the parallel or perpendicular categories, discrimination becomes more precise. These effects persist over variations in the orientation and scale of the angle exemplars. While other studies reviewed above have found similar categorical responses to parallels and perpendiculars under some conditions, our results suggest that angle discrimination in general and the more precise discrimination of these angle categories in particular are invariant to scale and orientation.

How robust, then, are the categories of parallelism and perpendicularity in the face of more drastic variation in the angle stimuli, such as rotations in 3D depth, which are prevalent in everyday viewing conditions? In Experiment 1, where stimuli were presented in the frontoparallel plane, the higher precision for parallel and perpendicular lines could have arisen from a specific sensitivity at the retinal or retinotopic level. If, however, the effect arises at a more abstract representational level, i.e., that parallelism and perpendicularity are properties that apply to the arrangement of lines in 3D space, then the effect should remain even if the stimuli are rotated in 3D depth, off the frontoparallel plane. This manipulation applies specifically to perpendicular lines: While parallel lines remain parallel even with rotation in depth (save a few, “accidental” viewpoints where their projections coincide: Amir, Biederman & Hayworth, 2011; 2012; Biederman, Yue, & Davidoff, 2009), perpendicular lines vary greatly in their projected

angle measure when the lines are rotated in depth (Nundy et al., 2000). Experiment 2 was therefore conducted both to evaluate how well the detection of perpendicular lines is preserved under viewing conditions that include additional rotations in 3D depth and to examine whether perpendicularity as a category persists under these conditions.

Experiment 2

In this experiment, we presented new participants with stimuli derived from the connected lines condition of Experiment 1. In the *slanted screen* condition (**Figure 5A**), participants viewed the same stimuli as Experiment 1, but on a screen that was rotated in depth by 45°. In the *normal viewing* condition (**Figure 5B**), participants were presented with the same stimuli as Experiment 1 with an un-rotated, frontoparallel screen (i.e., a direct replication of **Experiment 1**). By comparing participants' performance in these two conditions, we could evaluate whether both angle discrimination in general and the perpendicular category in particular persist with the rotation of our visual stimuli in depth. In the *projected stimuli* condition (**Figure 5C**), participants viewed the projection of the stimuli presented in the slanted screen condition on a frontoparallel screen. With this condition, we could evaluate whether the 3D context, presenting consistent cues to the reference frame transformation (like would be present in our everyday object recognition, see Nakayama, He, & Shimojo, 1995) or some general property of the stimuli's projection (which has a less clear connection to our everyday perceptual experience) might explain participants' responses in the slanted screen condition. If participants showed the same pattern of performance on the projected stimuli condition and the slanted screen condition, then there may be some geometric information at the level of the retinal projection that is guiding responses on the slanted screen condition.




Figure 5. The top panel illustrates on overhead perspective of the setup for the screen and participant for the three conditions: (A) *slanted screen*; (B) *normal viewing*; (C) *projected stimuli*. The lower panel illustrates the same trial (100° referent angle and 124° deviant) as it would be seen on the screen in each condition. The angles in the slanted screen condition are identical to the angles in the projected stimuli condition when projected on to the frontoparallel plane, as illustrated here. For illustration purposes, the correct response is circled in red.

Methods

Participants

Twelve adults (10 women; $M_{age} = 23$ years; range 19-38 years) participated in this experiment. The sample size was set in advance of data collection. With the sample size and smallest effect size from Experiment 1 ($N = 8$, Cohen's $d = 1.94$), our power to detect group-wise categorical effects within condition would be .999. We decided to increase our sample size by four participants compared to Experiment 1 since we also planned to compare results across conditions in this experiment and investigate effects by condition. All participants had normal or corrected-to-normal vision and had completed high school; most of them had also received a college or advanced degree. None were informed of the purpose and hypotheses of the study

until after it was completed. Each provided informed consent prior to the experiment and were paid 10€ per hour plus an additional sum, which depended on their performance. On average, participants earned 72.95€ (range 70.47€-75.56€).

Design, Apparatus, Procedure, and Stimuli

The distance of the participant to the center of the screen, the lighting conditions, the placeholder sizes, the screen luminance, the presentation duration, and the stimuli parameters were identical in all conditions of Experiment 2 to the connected lines condition of Experiment 1, except that the lines that formed each angle were slightly shorter (uniform distribution from 1.6 to 3.5 dva). In the projected stimuli condition, the angles were transformed by reducing all horizontal coordinates of each angle by a factor of $\cos(\frac{\pi}{4})$, while keeping all vertical coordinates constant. This transformation was applied to the angles only so that the external reference frame in the slanted and projected stimuli conditions (placeholders' shapes and screen frame) provided cues to enforce the perception of a slanted or frontoparallel presentation of the stimuli. As in Experiment 1, trials were blocked by reference angle but fewer reference angles were presented to focus on the comparisons that were most relevant to exploring the categorical effect (see **Table 1**). Each type of deviant was presented 18 times in a random order in a block, three times at each of six possible target locations. The total number of trials per block thus ranged from 144 to 162.

The experiment was divided into three sessions, one for each of the three experimental conditions. The sessions took place on different days, one to five days apart, and the order of the condition presented at each session was counterbalanced across participants. Sessions started with a short training phase consisting of two trials per reference angle displaying +/- 24° angle deviants. Five blocks corresponding to the five reference angles followed the training phase. The

order of the blocks within a session was chosen randomly for each participant. Each session lasted approximately 1.5 hours. As in Experiment 1, at the beginning of each block, participants were given a slide with instructions, showing an example of the reference angle in the vertical orientation. This slide was identical in the slanted screen, normal viewing, and projected stimuli conditions.

Analyses

As in Experiment 1, we fit performance with asymmetric down-pointing Gaussian curves. However, in several cases participants responded at chance (across participants: 10/120 blocks in the slanted screen condition; 3/120 blocks in the normal viewing condition; and 45/120 blocks in the projected condition; see **Figure S3** for individual accuracy curves). Such at-chance performance resulted in very high and unreliable estimated threshold values. We thus again used non-parametric rank tests to minimize the impact of these extreme values in our comparisons, and we also conducted parametric analyses on accuracy data, which yielded identical results except where indicated (see **SM, Table S2**).

As in Experiment 1, for all conditions of Experiment 2, we examined whether angle detection was more precise for the 90° reference angle compared to the other reference angles that were close to and far from 90° (i.e., 80°/100° reference angles and 55°/125° reference angles). We also evaluated whether deviant detection was more precise when deviants approached or crossed the 90° boundary compared to when they did not (all using Holm-corrected, one-tailed Wilcoxon signed-ranks tests). We only evaluated the asymmetries in the 80°/100° reference angles since the asymmetries in the 55°/125° reference angles were not significant in Experiment 1. By comparing the findings across the three conditions, first we asked whether categorical effects were present in the slanted screen condition, as in the normal

viewing condition (we compared the thresholds and asymmetries in these two conditions using Scheirer-Ray-Hare [SRH] tests, a non-parametric equivalent of a two-way repeated measures ANOVA). Second, we asked whether the effects observed with the slanted screen were due to information present in the stimuli's projection rather than based on shape information invariant to 3D rotations (by comparing the slanted screen condition to the projected stimuli condition using a second SRH test).

Results

Figure 6A shows the median detection thresholds for each reference angle in all three conditions, and **Table 3** shows tests for within-condition categorical effects. First, in the slanted screen condition, we observed more precise detection thresholds for 90° reference angles compared to both the 80°/100° reference angles ($Z = 3.18$, $p < .001$, 12/12 participants showed the effect), and also the 55°/125° reference angles ($Z = 3.18$, $p < .001$, 12/12 participants showed the effect). Second, this categorical effect did not differ across the slanted screen and normal viewing conditions. An SRH test comparing these two conditions revealed a significant main effect of reference angle ($H = 8.37$, $p = .004$), a significant main effect of condition ($H = 5.87$, $p = .015$), but, crucially, no interaction between reference angle and condition ($H = 0.20$, $p = .653$). As such, our normal viewing condition also replicated the results of Experiment 1, finding smaller detection thresholds with the 90° reference angle compared to the other reference angles (80°/100°: $Z = 2.47$, $p = .007$, 11/12 participants showed the effect; 55°/125°: $Z = 2.98$, $p = .001$, 11/12 participants showed the effect).

Perpendicular			
	90	80/100	55/125

Slanted Screen - thresholds	9.4	13.2***	15.9***
Slanted Screen - asymmetry		toward: 12.8; away: 14.8*	
Normal Viewing - thresholds	7.1	10.6**	12.2**
Normal Viewing - asymmetry		toward: 9.5; away: 11.3*	
Projected Stimuli - thresholds	27.3	26.5	25.9
Projected Stimuli - asymmetry		toward: 22.1; away: 28.1	

Table 3. Group-wise median detection thresholds for perpendicular reference angle blocks compared to other reference blocks. Planned, Holm-corrected one-tailed Wilcoxon signed-ranks tests compare the 90° reference blocks to the other reference blocks. All angle measures are in degrees. ** $p < .01$ *** $p < .001$.




Figure 6. A. Median thresholds for different reference angles in the three conditions of Experiment 2. While the slanted screen and normal viewing conditions show a sharp drop at the 90° reference angle, the projected stimuli condition does not. **B.** Asymmetries in the threshold of detection for larger versus smaller deviants at different reference angles.

Does the increased performance with the 90° reference angle in the slanted screen condition result from an analysis of the stimuli as projected or from an invariance of the detection of perpendicularity over rotations in depth? If the former, then the same categorical effects should be present in the projected stimuli condition. An SRH test comparing the slanted

screen and projected stimuli conditions revealed a significant main effect of reference angle ($H = 5.89, p = .015$), a significant main effect of condition ($H = 11.67, p < .001$), and an interaction between reference angle and condition ($H = 8.56, p = .003$). Indeed, contrary to the slanted screen condition, there was no sign of categorical effects in the projected stimuli condition, as performance with the 90° reference angle was no different from the other reference angles ($ps = 1.00$).

Next, we assessed whether performance for reference angles $80^\circ/100^\circ$ was better when the deviant approached or crossed the 90° boundary (see **Figure 6B**). Performance was significantly asymmetric in the slanted screen condition ($Z = 1.85, p = 0.032$, 9/12 participants showed the effect). An SRH test comparing the thresholds towards and away from 90° in the slanted screen and normal viewing conditions revealed a significant main effect of deviant direction ($H = 5.00, p = .025$), condition ($H = 7.29, p = .007$), but no interaction between these two factors ($H = 0.02, p = .896$). In the normal viewing condition too, there was a significant asymmetry in detection thresholds ($Z = 1.85, p = .032$, 8/12 participants showed the effect). Thus, the asymmetry effects were comparable across the slanted screen and normal viewing conditions and were comparable to Experiment 1.

Finally, an SRH test comparing the slanted screen and projected stimuli conditions on the asymmetry of their deviant detection thresholds revealed a main effect of deviant direction ($H = 6.88, p = .009$), condition ($H = 11.38, p < .001$), and no interaction between the two factors ($H = 0.09, p = .770$). Indeed, asymmetries followed the same pattern in the projected stimuli condition as in the other conditions, with slightly better performance for angles deviating towards 90° versus away from 90° , though this difference was not significant ($Z = 1.20, p = .116$, 8/12 participants showed the effect; see **SM, Table S2**). These weak asymmetry effects in the

projected stimuli condition may have been due to a residual property of the angle transformation, rather than to participants' categorical discrimination of projected 90° angles. In particular, when the stimuli were transformed, the range of angle sizes presented on the screen differed based on the reference angle, with the greatest range occurring at 90°. Moreover, projected deviant angles fell outside the range of projected reference angles around 90° more often for deviants towards 90° than for deviants away from 90°, making deviants towards 90° easier to detect.

Discussion

This experiment builds on the findings of Experiment 1 by suggesting that both angle discrimination in general, and the perpendicular category in particular, are largely invariant to the rotation of shape stimuli in depth. A 45° rotation of the screen in the slanted screen condition degraded discrimination performance uniformly, yet left the relative discriminability of angles of different sizes intact, including preserving peak discriminability at 90°. Because the 90° angles in that condition were defined only in three dimensions with widely varying 2D projections, these results suggest that participants detected the angle intruder by monitoring the 3D angle sizes rather than their retinotopic, 2D projections. This conclusion is further supported by comparisons between the slanted screen condition and the projected stimuli condition. Here, the 2D angles that were projected on the retina were identical between conditions, yet accuracy varied greatly, suggesting that the preserved shape discrimination in the slanted screen condition was not due to properties of the 2D projection of the stimuli. Indeed, participants performed so poorly in the projected stimuli condition that the data were difficult to model, and it is thus hard to make specific conclusions about the characteristics of our shape discrimination with stimuli

transformed in this way. By contrast, rotating shapes in depth, with all of the real-world depth cues intact, led to a largely preserved ability to discriminate angles.

General Discussion

Across two experiments presenting angle-intruder detection tasks, we demonstrated that the discriminability of both connected and non-connected lines forming different angles varies greatly depending on the size of the angles being discriminated. Most notably, thresholds of discrimination were significantly smaller when lines formed parallels and perpendiculars. When angles were acute or obtuse and far from these two special angle categories, thresholds of discrimination were high, though when angles deviated from nearer acute or obtuse angles to approach the parallel or perpendicular categories, discrimination became more precise. This pattern of results persisted not only over variations in the orientations and scale of the angle exemplars, but also over their rotations in depth. In particular, when the very same angle stimuli were presented on a screen rotated 45° in depth, the relative discriminability of angles of different sizes persisted. Most remarkably, the significantly smaller threshold for detecting perpendicular angles persisted, despite the variability in the 2D projections of these angles, and this result was not due to properties of the stimuli's 2D projection. Our findings thus suggest that angle discrimination in general and the special angle categories of parallelism and perpendicularity in particular affect participants' shape judgments by their real-world angle information.

The prior literature had outlined different models of angle discrimination to which we can compare our results. In two cases, more precise discrimination for horizontal and vertical lines was invoked to explain better acuity for 90° and $0/180^\circ$ angles: Chen & Levi (1996) suggested

an orientation-independent, Weber-like discrimination space except at 90° , where discrimination is heightened and rooted in the presence of vertical and horizontal lines; Xu et al. (2018) proposed that angles which include either a vertical or horizontal line are discriminated most precisely. In contrast, Heely & Buchanan-Smith (1996) proposed that, unlike the discrimination of individual line orientations, angle discrimination operates over a reference frame that is object-centered, like the discrimination of other, more complex objects. Our findings thus support and extend this last model: Not only did our stimuli specifically avoid using lines at or near the horizontal or vertical, but also more precise discrimination persisted when the angles were rotated in 3D depth, suggesting that angle discrimination unfolds at the level of the 3D angle size, as it might for the shape of complex 3D objects.

What then is the status of the geometric categories of *parallelism* and *perpendicularity*? Some research has suggested that sensitivity to parallelism, at least, arises early in child development. Even four-year-old children indicate that a pair of parallel lines is the “most different” from five other pairs of lines that present continuous differences in angle at varied absolute orientations and scales (Izard et al., 2011a; 2011b). Children’s knowledge of the word “parallel,” moreover, has no relation to this choice (Izard et al., 2011a; 2011b). As such, children’s judgments of parallelism may be rooted in the recognition of more basic shape properties, for example, that two lines have the same orientation or that they maintain a constant distance from each other — and perhaps these very properties contribute to the categorical effects for parallels documented here. Future research should explore whether children display the same categorical effects around the discrimination of parallels as adults and whether their specialized treatment of parallel lines changes with more visual experience or with explicit learning of formal geometry in school.

What about perpendicularity, whose angle measure changes greatly with rotations in 3D depth? Unlike their performance with parallel lines, not until age seven or older do children pick out a pair of perpendicular lines from other pairs of lines that vary continuously in angle, and individual children's knowledge of the lexical terms "right angle" and "perpendicular" correlates with this choice (Izard et al., 2011a). The present task with adults did not rule out the possibility that explicitly learned conceptual representations were supporting participants' performance. All the adults in this study had benefitted from a formal education at least through high-school, which likely included a geometry class where the concepts of parallelism and perpendicularity were taught. In particular, since participants were introduced to each block of trials with an image that depicted the reference angle, it was possible that, for the parallel and perpendicular categories, participants labeled the reference and used that label to access a stored representation of an exemplar from that category from memory, making its recognition more accurate (Firestone & Scholl, 2016). Like other studies examining advantages in discrimination for complex spatial stimuli, defined, e.g., by topological relations, it is unknown the extent to which the present findings reflect verbal coding or pure visual processing (Lovett & Franconeri, 2017). Indeed, a suite of studies using search tasks and deviant detection tasks show enhanced detection of visual stimuli including colors, simple shapes, and more complex objects when category boundaries line up with linguistic labels (see Goldstone & Hendrickson, 2010; Lupyan, 2012). Future studies using the same angle stimuli with adults from other cultures, children, and non-human animals might shed further light on whether the effects observed in this study were perceptual or cognitive.

Nevertheless, the detection of perpendicular angles may indeed be universal. A group of adults from the Mundurucu tribe in the Amazon, who, unlike the adults in the present sample,

have no specialized geometric training or vocabulary, pick out a pair of perpendicular lines as the “most different” from other pairs of lines that vary continuously in angle (Izard et al., 2011a). While categorization of perpendiculars (as in this study with the Mundurucu adults, Izard et al., 2011a) and categorical effects in discrimination (as measured with the adults in this paper) may dissociate, it is possible that a perceptually based perpendicular category arises spontaneously in development and that education in formal geometry refines it or heightens attention to it (see Piazza, Pica, Izard, Spelke, & Dehaene, 2013 for further exploration of this suggestion in the numerical domain). If this is the case, it remains a challenge to identify a perceptual learning mechanism that might operate over such a long period of development and create such a robust category.

Might visual experience through development bolster our recognition of 3D shape information from which a heightened sensitivity to perpendicular angles could emerge? Previous work has shown that high-level perceptual learning can occur over extended periods of time, for example, to create or refine specific object categories like letters and faces (Dehaene, Cohen, Sigman, & Vinckier, 2005; Maurer, Le Grand, & Mondloch, 2002). This kind of perceptual learning is nevertheless limited, in that categorical effects disappear when, for example, face stimuli are inverted (Maurer et al., 2002). Such limits may be adaptive to the context: While face recognition is not invariant to large 2D orientation changes, it is invariant over large changes in viewpoint (i.e., in full view or in profile), expression, and lighting (Anselmi & Poggio, 2014).

The perceptual learning associated with 3D object recognition may accumulate 2D and 3D rotational invariance through development since objects are often seen rotated in 2D or 3D space. While newborns differentiate among 2D angle shapes given long periods of habituation (Slater, Mattock, Brown, & Bremner, 1991), seven-month-old infants fail to differentiate shapes

differing in angle during brief exposures to 2D forms with simultaneous variations in scale, direction, and orientation (Dillon, Izard, & Spelke, in preparation). Research with toddlers has suggested that 3D object recognition undergoes protracted development, with a spurt in the ability to recognize objects by their 3D geometric shapes between the ages of 18-24 months (Augustine, Smith, & Jones, 2011; Smith, 2009), and studies with older children have shown that the period during which we become better able to recognize unfamiliar viewpoints of 3D objects captured in 2D drawings extends to adulthood (Jüttner, Müller, & Rentschler, 2006; Jüttner, Wakui, Petters, Kaur, & Davidoff, 2013; Landau, Hoffman, & Kurz, 2006). Further developmental work, investigating invariances in angle detection may begin to shed light on how such shape representations become more robust to every-day 3D viewing conditions.

Even if angle detection in general becomes more robust to rotations in 3D depth through development, such development alone does not explain the emergence of a perpendicular category. A rotationally invariant representation of perpendicularity could be singled out because a perpendicular line is the most symmetrical position for a line relative to another line (i.e., splitting a line in half) or to a plane, and both infants and young children show some sensitivity to symmetry (Bornstein, Ferdinandsen, & Gross, 1981; Huang, Xue, Spelke, Huang, Zheng, Peng, 2018). A rotationally invariant representation of perpendicularity could also derive from representations of the absolute vertical and horizontal. These orientations are robustly represented early in development and across animal species (Appelle, 1972). For example, 5-month-old infants take longer to look towards a deviant oblique line among other oblique lines versus among vertically oriented lines the same angle measure away (Franklin, Catherwood, Alvarez, & Axelsson, 2010).

Finally, the present study raises questions about the relations between our perception of angles and our conception of formal Euclidean geometry. Euclid famously did not explain what he means by “equal” angles when, at the beginning of his *Elements* (Book 1, Definition 10), he defines a *perpendicular* as a line with equal angles to either side as it stands up from a baseline. This definition is strikingly perceptual: We see what he means. Do we need to see the abstractions of geometry to conceive of them? Are they driven by mental imagery of specific spatial exemplars, abstract idealized concepts, or linguistic representations? Indeed, what are the mental representations guiding our geometric reasoning? Individuals differ in their ability to discriminate angles: Might individual differences in our angle discrimination predict our ability to reason about the properties of points, lines, and figures on the Euclidean plane? Could training in angle discrimination cause short- or long-term benefits to geometric reasoning or judgment? We are only just beginning to probe how the human mind navigates the perceptual and conceptual worlds of geometry.

Author Contributions

All authors conceived of and designed the study. M. Duyck programmed the stimuli. M. R. Dillon, M. Duyck, and V. Izard collected the data. M. R. Dillon, M. Duyck, and V. Izard performed the data analysis. M. R. Dillon, M. Duyck, and V. Izard drafted the manuscript, and all authors provided critical revisions. All authors approved the final version of the manuscript.

Acknowledgements

This work was supported by a Starting Grant of the European Research Council to V.I. (FP7 Project MathConstruction 263179), by a grant from the National Science Foundation to M.R.D. (DGE-1144152), and by the Norman Henry Anderson Graduate Psychology Fund to M.R.D.

Authors' Note

The full data set and analysis code are available on the Open Science Framework at: <https://osf.io/wa64q/>. Code for the generation of the experimental stimuli are available upon request.

References

- Amir, O., Biederman, I., & Hayworth, K. J. (2011). The neural basis for shape preferences. *Vision Research*, 51(20), 2198-2206.
- Amir, O., Biederman, I., & Hayworth, K. J. (2012). Sensitivity to nonaccidental properties across various shape dimensions. *Vision Research*, 62, 35-43.
- Anselmi, F., & Poggio, T. (2014). *Representation learning in sensory cortex: a theory*. Center for Brains, Minds and Machines (CBMM).
- Anthony, J. L., & Francis, D. J. (2005). Development of phonological awareness. *Current Directions in Psychological Science*, 14(5), 255-259.
- Appelle, S. (1972). Perception and discrimination as a function of stimulus orientation: the "oblique effect" in man and animals. *Psychological Bulletin*, 78(4), 266.
- Augustine, E., Smith, L. B., & Jones, S. S. (2011). Parts and relations in young children's shape-based object recognition. *Journal of Cognition and Development*, 12(4), 556-572.
- Biederman, I., Yue, X., & Davidoff, J. (2009). Representation of shape in individuals from a culture with minimal exposure to regular, simple artifacts: Sensitivity to nonaccidental versus metric properties. *Psychological Science*, 20(12), 1437-1442.
- Bornstein, M. H., Ferdinandsen, K., & Gross, C. G. (1981). Perception of symmetry in infancy. *Developmental Psychology*, 17(1), 82.
- Cantlon, J. F., Pined, P., Dehaene, S., & Pelphrey, K. A. (2010). Cortical representations of symbols, objects, and faces are pruned back during early childhood. *Cerebral Cortex*, 21(1), 191-199.
- Carpenter, R. H. S., & Blakemore, C. (1973). Interactions between orientations in human vision. *Experimental Brain Research*, 18(3), 287-303.

- 699 Chen, S., & Levi, D. M. (1996). Angle judgment: Is the whole the sum of its parts?. *Vision*
700 *Research*, 36(12), 1721-1735.
- 701 Deen, B., Richardson, H., Dilks, D. D., Takahashi, A., Keil, B., Wald, L. L., ... & Saxe, R.
702 (2017). Organization of high-level visual cortex in human infants. *Nature*
703 *Communications*, 8, 13995.
- 704 Dehaene, S., Cohen, L., Sigman, M., & Vinckier, F. (2005). The neural code for written words: a
705 proposal. *Trends in Cognitive Sciences*, 9(7), 335-341.
- 706 Dehaene, S., Izard, V., Pica, P., & Spelke, E. (2006). Core knowledge of geometry in an
707 Amazonian indigene group. *Science*, 311(5759), 381-384.
- 708 Dillon, M. R., Izard, V., & Spelke, E. S. (in preparation). Infants' sensitivity to shape changes in
709 2D visual forms.
- 710 Eimas, P. D., Siqueland, E. R., Jusczyk, P., & Vigorito, J. (1971). Speech perception in
711 infants. *Science*, 171(3968), 303-306.
- 712 Euclid. (1990). *Great Books of the Western World: The thirteen books of Euclid's Elements. The*
713 *works of Archimedes, including The Method. Introduction to Arithmetic by Nicomachus*
714 (2 ed.). Chicago: Encyclopedia Britannica. (Original work written c. 300 B.C.E.).
- 715 Firestone, C., & Scholl, B. J. (2016). Cognition does not affect perception: Evaluating the
716 evidence for "top-down" effects. *Behavioral and Brain Sciences*, 39, 1-19.
- 717 Franklin, A., Catherwood, D., Alvarez, J., & Axelsson, E. (2010). Hemispheric asymmetries in
718 categorical perception of orientation in infants and adults. *Neuropsychologia*, 48(9),
719 2648-2657.
- 720 Goldstone, R. L., & Hendrickson, A. T. (2010). Categorical perception. *Wiley Interdisciplinary*
721 *Reviews: Cognitive Science*, 1(1), 69-78.

- 722 Heeley, D. W., & Buchanan-Smith, H. M. (1996). Mechanisms specialized for the perception of
723 image geometry. *Vision Research*, 36(22), 3607-3627.
- 724 Howe, C. Q., & Purves, D. (2005). Natural-scene geometry predicts the perception of angles and
725 line orientation. *Proceedings of the National Academy of Sciences of the United States of*
726 *America*, 102(4), 1228-1233.
- 727 Izard, V., Pica, P., Dehaene, S., Hinchey, D., & Spelke, E. S. (2011a). Geometry as a universal
728 mental construction. *Space, Time and Number in the Brain*, 19, 319-332.
- 729 Izard, V., Pica, P., Spelke, E. S., & Dehaene, S. (2011b). Flexible intuitions of Euclidean
730 geometry in an Amazonian indigene group. *Proceedings of the National Academy of*
731 *Sciences*, 108(24), 9782-9787.
- 732 Huang, Y., Xue, X., Spelke, E., Huang, L., Zheng, W., & Peng, K. (2018). The aesthetic
733 preference for symmetry dissociates from early-emerging attention to
734 symmetry. *Scientific Reports*, 8(1), 6263.
- 735 Jüttner, M., Müller, A., & Rentschler, I. (2006). A developmental dissociation of view-dependent
736 and view-invariant object recognition in adolescence. *Behavioural Brain Research*,
737 175(2), 420-424.
- 738 Jüttner, M., Wakui, E., Petters, D., Kaur, S., & Davidoff, J. (2013). Developmental trajectories of
739 part-based and configural object recognition in adolescence. *Developmental Psychology*,
740 49(1), 161.
- 741 Kuhl, P. K., Williams, K. A., Lacerda, F., Stevens, K. N., & Lindblom, B. (1992). Linguistic
742 experience alters phonetic perception in infants by 6 months of age. *Science*, 255, 606-
743 608.

- 744 Landau, B., Hoffman, J. E., & Kurz, N. (2006). Object recognition with severe spatial deficits in
745 Williams syndrome: sparing and breakdown. *Cognition*, 100(3), 483-510.
- 746 Liberman, A. M., Harris, K. S., Kinney, J. A., & Lane, H. (1961). The discrimination of relative
747 onset-time of the components of certain speech and nonspeech patterns. *Journal of*
748 *Experimental Psychology*, 61(5), 379.
- 749 Lovett, A., & Franconeri, S. L. (2017). Topological relations between objects are categorically
750 coded. *Psychological Science*, 28(10), 1408-1418.
- 751 Lupyan, G. (2012). Linguistically modulated perception and cognition: the label-feedback
752 hypothesis. *Frontiers in Psychology*, 3, 54.
- 753 Maurer, D., Le Grand, R., & Mondloch, C. J. (2002). The many faces of configural
754 processing. *Trends in Cognitive Sciences*, 6(6), 255-260.
- 755 Nakayama, K., He, Z. J., & Shimojo, S. (1995). Visual surface representation: A critical link
756 between lower-level and higher-level vision. In S. M. Kosslyn & D. N. Osherson (Eds.),
757 *Visual cognition: An invitation to cognitive science* (pp. 1-70). Cambridge, MA: MIT
758 Press.
- 759 Nundy, S., Lotto, B., Coppola, D., Shimpi, A., & Purves, D. (2000). Why are angles
760 misperceived? *Proceedings of the National Academy of Sciences*, 97(10), 5592-5597.
- 761 Piazza, M., Pica, P., Izard, V., Spelke, E. S., & Dehaene, S. (2013). Education enhances the
762 acuity of the nonverbal approximate number system. *Psychological Science*, 24(6), 1037-
763 1043.
- 764 Regan, D., Gray, R., & Hamstra, S. J. (1996). Evidence for a neural mechanism that encodes
765 angles. *Vision Research*, 36(2), 323-IN3.

- 766 Slater, A., Mattock, A., Brown, E., & Bremner, J. G. (1991). Form perception at birth:
767 Revisited. *Journal of Experimental Child Psychology*, 51(3), 395-406.
- 768 Smith, L. B. (2009). From fragments to geometric shape: Changes in visual object recognition
769 between 18 and 24 months. *Current Directions in Psychological Science*, 18(5), 290-294.
- 770 von Helmholtz, H. (1909). *Outlines of Psychology*. (J. P. C. Southall, Trans.). The Optical
771 Society of America. (Original work published 1897).
- 772 Wenderoth, P., & Johnson, M. (1984). The effects of angle-arm length on judgments of angle
773 magnitude and orientation contrast. *Perception & Psychophysics*, 36(6), 538-544.
- 774 Werker, J. F., & Tees, R. C. (1984). Cross-language speech perception: Evidence for perceptual
775 reorganization during the first year of life. *Infant Behavior and Development*, 7(1), 49-63.
- 776 Werkhoven, P., & Koenderink, J. J. (1993). Visual size invariance does not apply to geometric
777 angle and speed of rotation. *Perception*, 22(2), 177-184.
- 778 Wundt, W. M. (1925). *Helmholtz's Treatise on Physiological Optics*. (C. H. Judd, Trans.).
779 Leipzig: Wilhelm Engelmann. (Original work published 1896).
- 780 Xu, Z. X., Chen, Y., & Kuai, S. G. (2018). The human visual system estimates angle features in
781 an internal reference frame: A computational and psychophysical study. *Journal of*
782 *Vision*, 18(13):10, 1-11.