

764 **Variance of the effective optimum**

765 By definition, the variance in the per-generation effective optimum is

$$766 \text{Var}[\theta_{tot}] = \text{Var}[\sum_{k=1}^n e_k \theta_k] = \sigma_{\theta}^2 \sum_{k=1}^n e_k^2 + 2 \sum_{i=1}^{n-1} \sum_{j=i+1}^n \text{cov}[e_i \theta_i, e_j \theta_j], \quad (\text{A1})$$

767 where  $\sigma_{\theta}^2$  is the stationary variance of the per-episode optimum. Rewriting (A1) using the  
768 properties of the autocorrelation of the environment (eq. 11) leads to

$$769 \text{Var}[\theta_{tot}] = \sigma_{\theta}^2 (\sum_{k=1}^n e_k^2 + 2 \sum_{i=1}^{n-1} \sum_{j=i+1}^n e_i e_j \rho^{(j-i)\alpha T/n}), \quad (\text{A2})$$

770 with  $e_i = \omega_{tot}^2 / \omega_i^2$ . The first term in the right-hand side of equation (A2) corresponds to the direct  
771 contribution of the variance of the environment. The second term on the right-hand side results  
772 from the autocorrelation of the environment between selection episodes within a generation,  
773 generating additional variation to the effective optimum. With the further assumption that the  
774 strength of selection is the same at all episodes of selection,  $\omega_i^2 = \omega^2$ , Equation A2 further  
775 simplifies in

$$776 \text{Var}[\theta_{tot}] = \sigma_{\theta}^2 \left( \frac{1}{n} + \frac{2}{n} \sum_{i=1}^{n-1} \sum_{j=i+1}^n \rho^{(j-i)\alpha T/n} \right) = \sigma_{\theta}^2 \frac{n - n\rho^{2\alpha T/n} + 2\rho^{\alpha T/n} (\rho^{\alpha T} - 1)}{n^2 (\rho^{\alpha T/n} - 1)^2} \quad (\text{A3})$$

777 The variance of the effective optimum  $\theta_{tot}$  is always lower than the variance of the environment  
778 (set to 1 in figure A1), and decreases with increasing number of selection episodes (fig A1-A).


779 However,  $\text{Var}[\theta_{tot}]$  does not decrease without bounds, instead reaching a limit value for a large

780 number of episodes of selection,  $\lim_{n \rightarrow \infty} (\text{Var}[\theta_{tot}]) = 2 \frac{\psi - \ln \psi - 1}{(\ln \psi)^2}$ , which only depend on the

781 autocorrelation of the environment over the time window for selection  $\psi = \rho^{\alpha T}$ . In addition, for

782 a given number of selection episodes, the variance of the effective optimum increases with the

783 autocorrelation of the environment (fig. A1-B).



785

786 Figure A1: Variance of the effective optimum as a function of the number of selection episodes (A) and as a function  
 787 of the autocorrelation of the environment (B). Crosses represent the results from simulations where the strength of  
 788 selection is drawn from a probability distribution as described in fig. 2 caption. Full lines represent the expectation  
 789 from eq. A3 and dashed lines represent the limit for a large number of selection episodes. Panel A: from light grey to  
 790 dark  $\rho_{\theta, T} = 0.2, 0.5$  and  $0.9$  respectively. Panel B: the gray scale represents different number of selection episode per  
 791 generation (see within the panel). For both panels,  $\bar{\omega}_{tot}^2 = 25$ ,  $\sigma_{\theta}^2 = 1$ ,  $\alpha = 1$ .

## 792 Variance of the selection gradient

793 The variance of  $\beta_{tot}$  is proportional to the variance of the distance of the mean phenotype to the  
 794 optimum

$$795 \text{Var}[\beta_{tot}] = (S_{tot})^2 \text{Var}[(\bar{z} - \theta_{tot})] = (S_{tot})^2 \text{E}[(\bar{z} - \theta_{tot})^2]. \quad (\text{A4})$$

796 By rearranging equation 10 in the main text (Charlesworth 1993), the mean phenotypic value at  
 797 generation  $g$  is

$$798 \quad \bar{z}_g = (1 - S_{tot}G_0)^g \bar{z}_0 + S_{tot}G_0 \sum_{i=1}^g (1 - S_{tot}G_0)^{i-1} \theta_{tot,g-i}. \quad (A5)$$

799 After a sufficiently large number of generations, the first term in equation (A5), corresponding to  
 800 the initial conditions, can be neglected. We thus obtain

$$801 \quad E[(\bar{z} - \theta_{tot})^2] = E \left[ \left( S_{tot}G_0 \sum_{i=1}^g (1 - S_{tot}G_0)^{i-1} \theta_{tot,g-i} - \theta_{tot,g} \right)^2 \right], \quad (A6)$$

802 By developing terms, using the autocorrelation of the effective optimum (eq. 12) and assuming  
 803 that  $g \rightarrow \infty$ , we obtain equation (16) in the main text (see also Charlesworth 1993; eq. 19a)

$$804 \quad \text{Var}[\beta_{tot}] = 2\text{Var}[\theta_{tot}] S_{tot}^2 \frac{1 - \rho(1 + S_{tot}G_0(\varphi - 1))}{(2 - S_{tot}G_0)(1 - \rho(1 - S_{tot}G_0))}.$$

### 805 **Autocorrelation of the selection gradient**

806 The autocovariance of the selection gradient over  $\tau$  generations is

$$807 \quad \text{Cov}[\beta_{tot,g}, \beta_{tot,g-\tau}] = S_{tot}^2 (E[\theta_{tot,g} \theta_{tot,g-\tau}] + E[\bar{z}_g \bar{z}_{g-\tau}] - E[\bar{z}_{g-\tau} \theta_{tot,g}] - E[\theta_{tot,g-\tau} \bar{z}_g]). \quad (A7)$$

### 808 *Continuous time approach*

809 Following Lande and Shannon (1996), the dynamics can be approximated in continuous time. The  
 810 continuous time approximation eases the analysis. After long enough time so that the we can  
 811 neglect the initial conditions, the mean phenotype at time  $t$  is

$$812 \quad \bar{z}_t = S_{tot}G_0 \int_0^\infty e^{-S_{tot}G_0 x} \theta_{tot,t-x} dx. \quad (A8)$$

813 Equation (A8) is the continuous time equivalent to equation (A6) once the effect of initial  
 814 conditions has been erased. From equations (A7) and (A8), the method to derive the

815 autocorrelation of the selection gradient is described in Chevin and Haller (2014, appendix for the  
816 autoregressive optimum). This approach cannot a priori be used in the context of the present model  
817 because the inflation factor is 1 for  $\tau = 0$  and is not defined for  $0 < \tau < 1$ . However, we found  
818 that the naïve approach to use equation (12) in the main text directly in the derivations from Chevin  
819 and Haller (2014) matches closely the exact discrete time approach that we develop in the next  
820 section. As we could not demonstrate this observation formally, we chose not to present this result  
821 in our main text. We nevertheless provide this result here as a heuristic:

$$822 \quad \rho_{\beta_{tot},\tau} = \varphi \frac{\rho^{\tau T - S_{tot}G_0 \frac{T_\theta}{T}} e^{-S_{tot}G_0\tau}}{1 - S_{tot}G_0 \frac{T_\theta}{T}}, \quad (A9)$$

823 where  $\frac{T_\theta}{T}$  measures the characteristic timescale of the autocorrelation of the environment in units  
824 of generation time.

### 825 *Discrete time approach*

826 In contrast with previous studies (Lande and Shannon 1996, Chevin and Haller 2014), we calculate  
827 the autocorrelation of the selection gradient in discrete time. Similarly to the calculation for the  
828 variance of the selection gradient, we use equation (A5) assuming that many generations occurred  
829 so that the first term can be neglected. We also use equation (12) in the main text for the  
830 autocorrelation of the effective optimum  $\rho_{\theta_{tot},\tau} = \rho^{\tau T} \varphi$ , reminding that  $\varphi = 1$  for  $\tau = 0$ . The  
831 general procedure follows Chevin and Haller (2014, appendix for the autoregressive optimum) by  
832 calculating each expectation from equation (A7). Even though we kept notations concise, each  
833 sum in the expectations below can be resolved exactly. The first expectation is straightforward

$$834 \quad E[\theta_{tot,g} \theta_{tot,g-\tau}] = \varphi \text{Var}[\theta_{tot}] \rho^{\tau T}. \quad (A10)$$

835 The expectation for the first cross product is

836  $E[\bar{z}_{g-\tau}\theta_{tot,g}] = S_{tot}G_0 \sum_{i=1}^{g-\tau} (1 - S_{tot}G_0)^{i-1} E[\theta_{tot,g}\theta_{tot,g-i-\tau}]$ , taking into account the

837 discontinuity of the inflation factor  $I_f$  in  $i = g - \tau$ , it results in

$$838 \quad E[\bar{z}_{g-\tau}\theta_{tot,g}] = \varphi \text{Var}[\theta_{tot}] S_{tot}G_0 \sum_{i=1}^{g-\tau-1} (1 - S_{tot}G_0)^{i-1} \rho^{(\tau+i)T} + S_{tot}G_0(1 -$$

$$839 \quad S_{tot}G_0)^{g-\tau-1} \text{Var}[\theta_{tot}]. \quad (\text{A11})$$

840 The expectation for second cross product is

$$841 \quad E[\theta_{tot,g-\tau}\bar{z}_g] = S_{tot}G_0 \sum_{i=1}^g (1 - S_{tot}G_0)^{i-1} E(\theta_{tot,g-i}\theta_{tot,g-\tau}).$$

842 We need to distinguish 3 cases,  $i < \tau$ ,  $i > \tau$  and  $i = \tau$  :

$$843 \quad E[\theta_{tot,g-\tau}\bar{z}_g] = \varphi \text{Var}[\theta_{tot}] S_{tot}G_0 \left( \sum_{i=1}^{\tau-1} (1 - S_{tot}G_0)^{i-1} \rho^{(\tau-i)T} + \sum_{i=\tau+1}^g (1 -$$

$$844 \quad S_{tot}G_0)^{i-1} \rho^{(i-\tau)T} + \varphi^{-1} (1 - S_{tot}G_0)^{\tau-1} \right). \quad (\text{A12})$$

845 Lastly,

$$846 \quad E[\bar{z}_g\bar{z}_{g-\tau}] = (S_{tot}G_0)^2 \sum_{i=1}^g \sum_{k=1}^{g-\tau} (1 - S_{tot}G_0)^{i+k-2} E[\theta_{tot,g-i}\theta_{tot,g-\tau-k}].$$

847 To calculate the double sum, it is convenient to perform a change of variables

$$848 \quad \sum_{i=1}^g \sum_{k=1}^{g-\tau} (1 - S_{tot}G_0)^{i+k-2} E[\theta_{tot,g-i}\theta_{tot,g-\tau-k}] = \sum_{i=1}^g \sum_{j=\tau+1}^g (1 -$$

$$849 \quad S_{tot}G_0)^{i+j-\tau-2} E[\theta_{tot,g-i}\theta_{tot,g-j}].$$

850 We then need to distinguish 4 cases,

$$851 \quad \text{for } i < \tau+1, A = \varphi \text{Var}[\theta_{tot}] \sum_{i=1}^{\tau} \sum_{j=\tau+1}^g (1 - S_{tot}G_0)^{i+j-\tau-2} \rho^{(j-i)T}.$$

$$852 \quad \text{for } i > \tau \text{ and } i = j, B = \text{Var}[\theta_{tot}] \sum_{i=\tau+1}^g (1 - S_{tot}G_0)^{2i-\tau-2}.$$

853 for  $i > \tau$  and  $j > i$ ,  $C = \varphi \text{Var}[\theta_{tot}] \sum_{j=\tau+2}^g \sum_{i=\tau+1}^{j-1} (1 - S_{tot} G_0)^{i+j-\tau-2} \rho^{(j-i)T}$ .

854 for  $i > \tau$  and  $i > j$ ,  $D = \varphi \text{Var}[\theta_{tot}] \sum_{i=\tau+2}^g \sum_{j=\tau+1}^{i-1} (1 - S_{tot} G_0)^{i+j-\tau-2} \rho^{(i-j)T}$ ,

855 with  $C = D$ , and resulting in

$$856 \quad E[\bar{z}_g \bar{z}_{g-\tau}] = (S_{tot} G_0)^2 (A + B + 2C). \quad (\text{A13})$$

857 Adding equations (A10) to (A13), finding the limit of this sum for large  $g$ , dividing by the

858 variance of the selection gradient (eq. 16) and rearranging leads to the autocorrelation of the

859 selection gradient

$$860 \quad \rho_{\beta_{tot}, \tau} = \frac{\varphi \varepsilon \rho^{\tau T} + \gamma (1 - S_{tot} G_0)^{\tau}}{\vartheta}, \quad (\text{A14})$$

861 where,

$$862 \quad \varepsilon = (\rho^T - 1)^2 (2 - S_{tot} G_0),$$

$$863 \quad \gamma = \frac{S_{tot} G_0}{(1 - S_{tot} G_0)} \left( \rho^T (1 - \varphi) (S_{tot} G_0)^2 - (1 - \rho^T) (1 + \rho^T (2\varphi - 1)) (1 - S_{tot} G_0) \right),$$

$$864 \quad \vartheta = 2(\rho^T - 1 + S_{tot} G_0)(\rho^T - 1 + \rho^T(\varphi - 1)S_{tot} G_0).$$

865 Equation (A14) is written in a form similar to equation (A9), decomposing the effects of the

866 autocorrelation of the environment and of the deviations of the mean phenotype to the optimum.

867 An interpretation for equation (A14) is proposed in the main text.

868 In the special case where there is a unique episode of selection per generation, i.e.  $\varphi = 1$ , equation

869 (A14) is

$$870 \quad \rho_{\beta_{tot}, \tau} = \frac{\rho_{\theta}^{\tau T} - \kappa (1 - S_{tot} G_0)^{\tau}}{1 - \kappa}, \quad (\text{A15})$$

871 where  $\kappa = \frac{(1+\rho^T)S_{tot} G_0}{(1-\rho^T)(2-S_{tot} G_0)}$ .

872 Equation (A15) is very similar to equation 5b in Chevin and Haller (2014). Comparing equation  
873 (A14) and (A15) shows that the inflation factor, i.e. the autocorrelation of selection within  
874 generation, complicates the analysis by introducing new terms at the interface between the  
875 autocorrelation of the environment and the dynamics of selection within and across generations.  
876 A Taylor expansion performed on the reciprocals of equation (A15) and 5b in Chevin and Haller  
877 (2014) shows that both equations are identical to at least the second order. This result can be  
878 confirmed by representing both functions for different autocorrelation of the environment. Both  
879 functions match exactly on most of the autocorrelation range with minor deviation at very weak  
880 autocorrelation (not shown).

### 881 **Changes in (co)variances under selection**

882 Here we derive the changes in the variances and covariances of phenotypic components after each  
883 selection episode. Following classical quantitative genetics, we assume that the phenotype  $z$  of  
884 any individual prior to any selection in a generation is the sum of two normally distributed  
885 components: the breeding value  $x$  and environmental effect  $e$ . At the beginning of a generation,  
886 these phenotypic components have means  $\bar{x}$  and  $\bar{e} = 0$ , variances  $G$  and  $E$ , and are uncorrelated,  
887 such that their covariance is  $C_{x,e} = 0$  and the total phenotypic variance is  $P = G + E$ . Changes in  
888 the mean breeding value under selection are obtained from changes in mean phenotype using the  
889 regression of breeding values on phenotypes, which at the beginning of a generation has slope  
890  $\frac{\text{Cov}(x,z)}{P} = \frac{G}{G+E} = h^2$ , following the usual notation for heritability.

891 We consider episodes of selection caused by a Gaussian fitness peak, such that individuals with  
892 phenotype  $z$  have fitness  $(z) \propto \exp(-\frac{(z-\theta)^2}{2\omega^2})$  (where we omit indices for selection episodes and  
893 generation for simplicity). It is informative to start by investigating the first selection episode in a  
894 generation, where  $x$  and  $e$  are independent and  $\bar{e} = 0$ ,  $\bar{z} = \bar{x}$ . The fitness function on breeding  
895 values is obtained by integrating over the distribution of environmental effects,  $\tilde{W}(x) \propto$   
896  $\exp(-\frac{(x-\theta)^2}{2(\omega^2+E)})$ , and the fitness function on environmental effects is obtained similarly by  
897 integrating over breeding values,  $W_e(e) \propto \exp(-\frac{(\bar{x}+e-\theta)^2}{2(\omega^2+G)})$ . The fitness function on breeding  
898 values (respectively, on environmental effects) is thus also Gaussian, with squared width  $\omega^2 + V_e$   
899 (respectively  $\omega^2 + G$ ) and optimum  $\theta$  (respectively  $\theta - \bar{x}$ ). With a Gaussian fitness peak, the  
900 distribution of phenotypes, breeding values and environmental effects after selection remain  
901 Gaussian. Denoting the per-episode strength of stabilizing selection as  $S = \frac{1}{\omega^2+P}$ , the means after  
902 selection are

$$903 \quad \bar{z}^* = \theta + \frac{\omega^2}{\omega^2+P}(\bar{z} - \theta) = \theta + (1 - SP)(\bar{z} - \theta) = \bar{z} - SP(\bar{z} - \theta) \quad (\text{A16a})$$

$$904 \quad \bar{x}^* = \theta + \frac{\omega^2+V_e}{\omega^2+P}(\bar{x} - \theta) = \theta + (1 - SG)(\bar{x} - \theta) = \bar{x} - SG(\bar{x} - \theta) \quad (\text{A16b})$$

$$905 \quad \bar{e}^* = \theta - \bar{x} + \frac{\omega^2+G}{\omega^2+P}(\bar{x} - \theta) = \theta - \bar{x} + (1 - SE)(\bar{x} - \theta) = -SE(\bar{z} - \theta), \quad (\text{A16c})$$

906 where we have used the fact that  $\bar{e} = 0$  and thus  $\bar{z} = \bar{x}$  prior to any selection. Note that  $\bar{x}^* - \bar{x} =$   
907  $h^2(\bar{z}^* - \bar{z})$  and  $\bar{e}^* - \bar{e} = (1 - h^2)(\bar{z}^* - \bar{z})$ , as required. The variances after selection are

$$908 \quad P^* = \left(\frac{1}{\omega^2} + \frac{1}{P}\right)^{-1} = (1 - SP)P \quad (\text{A17a})$$

$$909 \quad G^* = \left(\frac{1}{\omega^2+E} + \frac{1}{G}\right)^{-1} = (1 - SG)G \quad (\text{A17b})$$



910  $E^* = \left( \frac{1}{\omega^2 + G} + \frac{1}{E} \right)^{-1} = (1 - SE)E,$  (A17c)

911 which satisfy the relationships  $G^* - G = h^4(P^* - P)$  and  $E^* - E = (1 - h^2)^2(P^* - P)$ . The  
 912 phenotypic variance after selection must also satisfy

913  $P^* = G^* + E^* + 2C_{x,e}^*$  (A18)

914 so the covariance of breeding values and environmental effects after selection is

915  $C_{x,e}^* = \frac{P^* - G^* - E^*}{2}.$  (A19)

916 Using  $P = G + E$  this yields

917  $C_{x,e}^* = \frac{(1-SP)P - (1-SG)G - (1-SE)E}{2} = -S \frac{P^2 - G^2 - E^2}{2} = -SGE,$  (A20)

918 which satisfies  $C_{x,e}^* = C_{x,e}^* - C_{x,e} = h^2(1 - h^2)(P^* - P)$ .

919 These formulas apply to the first selection episode, but in further episodes the change in the mean  
 920 breeding value and environmental effects also depend on their covariance. More general formulas  
 921 for change at any selection episode can be derived through a multivariate approach. Let us denote  
 922 the vector of breeding values and environmental effects as  $\mathbf{y} = (x, e)^T$  (where superscript T  
 923 denotes a transposition), with mean  $\bar{\mathbf{y}}$  and covariance matrix  $\mathbf{Y}$ . Substituting  $x + e$  for  $z$  in the  
 924 fitness function  $W(z)$ , and integrating over the joint distribution of  $x$  and  $e$ , the mean fitness is

925  $\bar{W} \propto \exp\left(-\frac{S_C(\bar{x} + \bar{e} - \theta)^2}{2}\right)$  (A21)

926  $S_C = \frac{1}{\omega^2 + G + E + 2C_{x,e}}$  (A22)

927 Then using standard results from evolutionary quantitative genetics (Lande 1979, 1980, Lande and  
 928 Arnold 1983 ) and properties of the Gaussian function, the change in the vector of mean breeding  
 929 values and environmental effects under selection within a generation is

$$930 \quad \Delta^* \bar{\mathbf{y}} = \mathbf{Y} \partial_{\bar{\mathbf{y}}} \ln \bar{W} \quad (\text{A23})$$

931 where  $\partial_{\bar{\mathbf{y}}}$  is the vector of partial derivatives with respect to each element of  $\bar{\mathbf{y}}$ , yielding

$$932 \quad \Delta^* \bar{x} = -S_C (\bar{x} + \bar{e} - \theta) (G + C_{x,e}) \quad (\text{A24a})$$

$$933 \quad \Delta^* \bar{e} = -S_C (\bar{x} + \bar{e} - \theta) (E + C_{x,e}) \quad (\text{A24b})$$

$$934 \quad \Delta^* \bar{z} = \Delta^* \bar{x} + \Delta^* \bar{e} = -S_C (\bar{x} + \bar{e} - \theta) (G + E + 2C_{x,e}) \quad (\text{A24c})$$

935 When  $\bar{e} = C_{x,e} = 0$ , these simplify to  $\Delta^* \bar{x} = -SG(\bar{x} - \theta)$  and  $\Delta^* \bar{e} = -SE(\bar{x} - \theta)$ , consistent  
 936 with the formulas above for the first episode of selection. The recursion for the covariance matrix  
 937 is

$$938 \quad \Delta^* \mathbf{Y} = \mathbf{Y} (\partial_{\bar{\mathbf{y}}}^2 \ln \bar{W}) \mathbf{Y} \quad (\text{A25})$$

939 where  $\partial_{\bar{\mathbf{y}}}^2$  is the Hessian matrix of second partial derivatives with respect to each element of  $\bar{\mathbf{y}}$ ,  
 940 yielding

$$941 \quad \Delta^* G = -S_C (G + C_{x,e})^2 \quad (\text{A26})$$

$$942 \quad \Delta^* E = -S_C (E + C_{x,e})^2 \quad (\text{A27})$$

$$943 \quad \Delta^* C_{x,e} = -S_C (G + C_{x,e}) (E + C_{x,e}) \quad (\text{A28})$$

944 When  $\bar{e} = C_{x,e} = 0$ , these simplify to  $\Delta^* G = -SG^2$ ,  $\Delta^* E = -SE^2$ , and  $C_{x,e} = -SGE$ , again  
 945 consistent with the formulas above for the first episode of selection.

946 In the general case where  $\bar{e}$  and  $C_{x,e}$  are non-zero, as is expected if the population has previously  
947 undergone any episode of selection in the ongoing generation, then the relationship between  
948 phenotypic change and change in breeding values cannot be inferred directly from current  
949 components of phenotypic variance. Indeed we have

$$950 \frac{\Delta^* \bar{x}}{\Delta^* \bar{z}} = \frac{G + C_{x,e}}{G + E + 2C_{x,e}} = \frac{\text{Cov}(x, x+e)}{\text{Var}(x+e)} \quad (\text{A29})$$

951 which is still a regression slope of breeding values on phenotypes, but this no longer translates into  
952 a ratio of additive genetic variance on phenotypic variance, because of the covariance between  
953 breeding values and environmental values. However, it can be shown that this ratio always equals  
954 the ratio  $\frac{G}{G+E} = h^2$  at the beginning of a generation.

## 955 **References**

- 956
- 957 Charlesworth, B. 1993. Directional selection and the evolution of sex and recombination. *Genetics*  
958 *Research* **61**:205-224.
- 959 Chevin, L. M., and B. C. Haller. 2014. The temporal distribution of directional gradients under  
960 selection for an optimum. *Evolution* **68**:3381-3394.
- 961 Lande, R. 1979. Quantitative genetic analysis of multivariate evolution, applied to brain: body size  
962 allometry. *Evolution* **33**:402-416.
- 963 Lande, R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters.  
964 *Evolution* **34**:292-305.
- 965 Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters.  
966 *Evolution* **37**:1210-1226.

967 Lande, R., and S. Shannon. 1996. The role of genetic variation in adaptation and population  
968 persistence in a changing environment. *Evolution* **50**:434-437.

969

970