



HAL
open science

**MONOTONICITY AND COMPLETE
MONOTONICITY OF TWO FUNCTIONS DEFINED
BY THREE DERIVATIVES OF A FUNCTION
INVOLVING TRIGAMMA FUNCTION**

Feng Qi, Ling-Xiong Han, Hong-Ping Yin

► **To cite this version:**

Feng Qi, Ling-Xiong Han, Hong-Ping Yin. MONOTONICITY AND COMPLETE MONOTONICITY OF TWO FUNCTIONS DEFINED BY THREE DERIVATIVES OF A FUNCTION INVOLVING TRIGAMMA FUNCTION. 2020. hal-02998203

HAL Id: hal-02998203

<https://hal.science/hal-02998203>

Preprint submitted on 10 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MONOTONICITY AND COMPLETE MONOTONICITY OF TWO FUNCTIONS DEFINED BY THREE DERIVATIVES OF A FUNCTION INVOLVING TRIGAMMA FUNCTION

FENG QI, LING-XIONG HAN, AND HONG-PING YIN

Dedicated to people facing and battling COVID-19

ABSTRACT. In the paper, by convolution theorem of the Laplace transforms, a monotonicity rule for the ratio of two Laplace transforms, Bernstein's theorem for completely monotonic functions, and other analytic techniques, the authors

- (1) verify decreasing monotonicity of a ratio between three derivatives of a function involving trigamma function;
- (2) find necessary and sufficient conditions for a function defined by three derivatives of a function involving trigamma function to be completely monotonic.

These results confirm previous guesses posed by the first author and generalize corresponding known conclusions.

CONTENTS

1.	Introduction	1
2.	Lemmas	3
3.	Decreasing monotonicity	6
4.	Necessary and sufficient conditions of complete monotonicity	7
5.	Remarks	8
	References	9

1. INTRODUCTION

In the literature [1, Section 6.4], the function

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt, \quad \Re(z) > 0$$

and its logarithmic derivative $\psi(z) = [\ln \Gamma(z)]' = \frac{\Gamma'(z)}{\Gamma(z)}$ are called Euler's gamma function and digamma function respectively. Further, the functions $\psi'(z)$, $\psi''(z)$, $\psi'''(z)$, and $\psi^{(4)}(z)$ are known as the trigamma, tetragamma, pentagamma, and

2010 *Mathematics Subject Classification*. Primary 33B15; Secondary 26A48, 26D07, 33B10, 44A10, 44A35.

Key words and phrases. monotonicity; complete monotonicity; completely monotonic function; trigamma function; derivative; ratio; convolution theorem; inequality; monotonicity rule; Laplace transform; Bernstein's theorem; exponential function; guess.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\text{A}\text{T}\text{E}\mathcal{X}$.

hexagamma functions respectively. All the derivatives $\psi^{(k)}(z)$ for $k \geq 0$ are known as polygamma functions.

Recall from Chapter XIII in [4], Chapter 1 in [11], and Chapter IV in [12] that, if a function $f(t)$ on an interval I has derivatives of all orders on I and satisfies inequalities $(-1)^n f^{(n)}(t) \geq 0$ for $t \in I$ and $n \in \{0\} \cup \mathbb{N}$, then we call $f(t)$ a completely monotonic function on I .

Let $\Phi(x) = x\psi'(x) - 1 = x[\psi'(x) - \frac{1}{x}]$ on $(0, \infty)$. In [7, Theorem 4.1] and [9, Theorem 4], the first author turned out the following necessary and sufficient conditions and double inequality:

- (1) if and only if $\alpha \geq 2$, the function $\mathfrak{H}_\alpha(x) = \Phi'(x) + \alpha\Phi^2(x)$ is completely monotonic on $(0, \infty)$;
- (2) if and only if $\alpha \leq 1$, the function $-\mathfrak{H}_\alpha(x)$ is completely monotonic on $(0, \infty)$;
- (3) the double inequality $-2 < \frac{\Phi'(x)}{\Phi^2(x)} < -1$ is valid and sharp in the sense that the lower and upper bounds -2 and -1 cannot be replaced by any bigger and smaller ones respectively.

In [7, Theorem 1.1], the first author found the following necessary and sufficient conditions and limits:

- (1) if and only if $\beta \geq 2$, the function $H_\beta(x) = \frac{\Phi'(x)}{\Phi^\beta(x)}$ is decreasing on $(0, \infty)$, with the limits

$$\lim_{x \rightarrow 0^+} H_\beta(x) = \begin{cases} -1, & \beta = 2 \\ 0, & \beta > 2 \end{cases} \quad \text{and} \quad \lim_{x \rightarrow \infty} H_\beta(x) = \begin{cases} -2, & \beta = 2 \\ -\infty, & \beta > 2; \end{cases}$$

- (2) if $\beta \leq 1$, the function $H_\beta(x)$ is increasing on $(0, \infty)$, with the limits

$$H_\beta(x) \rightarrow \begin{cases} -\infty, & x \rightarrow 0^+ \\ 0, & x \rightarrow \infty. \end{cases}$$

For $k \in \{0\} \cup \mathbb{N}$ and $\lambda_k, \mu_k \in \mathbb{R}$, let

$$\mathfrak{J}_{k, \lambda_k}(x) = \Phi^{(2k+1)}(x) + \lambda_k [\Phi^{(k)}(x)]^2 \quad \text{and} \quad J_{k, \mu_k}(x) = \frac{\Phi^{(2k+1)}(x)}{[(-1)^k \Phi^{(k)}(x)]^{\mu_k}}$$

on $(0, \infty)$. In [5, Theorems 3.1 and 4.1], the first author presented the following necessary and sufficient conditions, limits, and double inequality:

- (1) if and only if $\lambda_k \geq \frac{(2k+2)!}{k!(k+1)!}$, the function $\mathfrak{J}_{k, \lambda_k}(x)$ is completely monotonic on $(0, \infty)$;
- (2) if and only if $\lambda_k \leq \frac{1}{2} \frac{(2k+2)!}{k!(k+1)!}$, the function $-\mathfrak{J}_{k, \lambda_k}(x)$ is completely monotonic on $(0, \infty)$;
- (3) if and only if $\mu_k \geq 2$, the function $J_{k, \mu_k}(x)$ is decreasing on $(0, \infty)$, with the limits

$$\lim_{x \rightarrow 0^+} J_{k, \mu_k}(x) = \begin{cases} -\frac{1}{2} \frac{(2k+2)!}{k!(k+1)!}, & \mu_k = 2 \\ 0, & \mu_k > 2 \end{cases}$$

and

$$\lim_{x \rightarrow \infty} J_{k, \mu_k}(x) = \begin{cases} -\frac{(2k+2)!}{k!(k+1)!}, & \mu_k = 2 \\ -\infty, & \mu_k > 2; \end{cases}$$

(4) if $\mu_k \leq 1$, the function $J_{k,\mu_k}(x)$ is increasing on $(0, \infty)$, with the limits

$$J_{k,\mu_k}(x) \rightarrow \begin{cases} -\infty, & x \rightarrow 0^+ \\ 0, & x \rightarrow \infty; \end{cases}$$

(5) the double inequality

$$-\frac{(2k+2)!}{k!(k+1)!} < \frac{\Phi^{(2k+1)}(x)}{[\Phi^{(k)}(x)]^2} < -\frac{1}{2} \frac{(2k+2)!}{k!(k+1)!}$$

is valid on $(0, \infty)$ and sharp in the sense that the lower and upper bounds cannot be replaced by any larger and smaller numbers respectively.

For $k \geq m \geq 0$, let

$$\mathcal{J}_{k,m}(x) = \frac{\Phi^{(2k+2)}(x)}{\Phi^{(k-m)}(x)\Phi^{(k+m+1)}(x)}$$

on $(0, \infty)$. In [5, Remark 5.3], the first author guessed that the function $\mathcal{J}_{k,m}(x)$ for $k \geq m \geq 0$ should be decreasing on $(0, \infty)$ and that the double inequality

$$-\frac{2(2k+2)!}{k!(k+1)!} < \mathcal{J}_{k,0}(x) < -\frac{(2k+2)!}{k!(k+1)!} \quad (1.1)$$

for $k \geq 0$ should be valid on $(0, \infty)$ and sharp in the sense that the lower and upper bounds cannot be replaced by any larger and smaller numbers respectively.

For $m, n \in \{0\} \cup \mathbb{N}$ and $\omega_{m,n} \in \mathbb{R}$, let

$$Y_{m,n}(x) = \frac{\Phi^{(m+n+1)}(x)}{\Phi^{(m)}(x)\Phi^{(n)}(x)} \quad (1.2)$$

and

$$\mathcal{Y}_{m,n;\omega_{m,n}}(x) = \Phi^{(m+n+1)}(x) + \omega_{m,n}\Phi^{(m)}(x)\Phi^{(n)}(x). \quad (1.3)$$

It is clear that

$$Y_{m,n}(x) = Y_{n,m}(x), \quad \mathcal{Y}_{m,n;\omega_{m,n}}(x) = \mathcal{Y}_{n,m;\omega_{n,m}}(x), \\ Y_{k-m,k+m+1}(x) = \mathcal{J}_{k,m}(x), \quad \mathcal{Y}_{k,k;\omega_{k,k}}(x) = \mathfrak{J}_{\omega_{k,k}}(x), \quad \mathcal{Y}_{0,0;\omega_{0,0}}(x) = \mathfrak{H}_{\omega_{0,0}}(x).$$

In this paper, we will prove decreasing monotonicity of the function $Y_{m,n}(x)$ and find necessary and sufficient conditions on $\omega_{m,n}$ for $\pm(-1)^{m+n+1}\mathcal{Y}_{m,n;\omega_{m,n}}(x)$ to be completely monotonic on $(0, \infty)$. These results confirm the above guesses and generalize corresponding ones in [5, 7, 9] mentioned above.

2. LEMMAS

The following lemmas are necessary in this paper.

Lemma 2.1 (Convolution theorem for the Laplace transforms [12, pp. 91–92]). *Let $f_k(t)$ for $k = 1, 2$ be piecewise continuous in arbitrary finite intervals included in $(0, \infty)$. If there exist some constants $M_k > 0$ and $c_k \geq 0$ such that $|f_k(t)| \leq M_k e^{c_k t}$ for $k = 1, 2$, then*

$$\int_0^\infty \left[\int_0^t f_1(u)f_2(t-u)du \right] e^{-st} dt = \int_0^\infty f_1(u)e^{-su} du \int_0^\infty f_2(v)e^{-sv} dv.$$

Lemma 2.2 ([13, Lemma 4]). *Let the functions $A(t)$ and $B(t) \neq 0$ be defined on $(0, \infty)$ such that their Laplace transforms $\int_0^\infty A(t)e^{-xt} dt$ and $\int_0^\infty B(t)e^{-xt} dt$ exist. If the ratio $\frac{A(t)}{B(t)}$ is increasing, then the ratio $\frac{\int_0^\infty A(t)e^{-xt} dt}{\int_0^\infty B(t)e^{-xt} dt}$ is decreasing on $(0, \infty)$.*

Lemma 2.3. *Let $x, y \in \mathbb{R}$ such that $0 < 2x < y$.*

(1) *When $y > 2x > 2(2 + \frac{1}{\ln 2}) = 6.885390\dots$, the function*

$$F(x, y) = 2\left(\frac{1}{x} - \frac{1}{y-x}\right) + \frac{1}{2}\left(\frac{2^{y-x}}{y-x} - \frac{2^x}{x}\right) - \frac{2^{y-x} - 2^x}{(y-x)x}$$

is positive.

(2) *For $k, m \in \mathbb{N}$ such that $6 \leq 2m < k$, the sequence $F(m, k)$ is positive.*

Proof. The function $F(x, y)$ can be rearranged as

$$F(x, y) = \frac{2(y-2x) + 2^{x-1}[2-y+x+(x-2)2^{y-2x}]}{x(y-x)}.$$

Therefore, it suffices to prove $2-y+x+(x-2)2^{y-2x} > 0$, that is,

$$2^{y-2x} > \frac{y-x-2}{x-2}. \quad (2.1)$$

Replacing $y-2x$ by t in (2.1) leads to

$$2^t > \frac{t+x-2}{x-2} = 1 + \frac{t}{x-2} \quad (2.2)$$

for $t > 0$ and $x > 2$. The inequality (2.2) can be reformulated as $x > 2 + \frac{t}{2^t-1}$. Since the function $\frac{t}{2^t-1}$ is decreasing from $(0, \infty)$ onto $(0, \frac{1}{\ln 2})$, it is sufficient for $x > 2 + \frac{1}{\ln 2} = 3.442695\dots$

Repeating those arguments before the inequality (2.1) hints us that, for proving $F(m, k) > 0$, it is sufficient to show

$$2^{k-2m} > \frac{k-m-2}{m-2} = 1 + \frac{k-2m}{m-2}$$

which can be rewritten as

$$\frac{k-2m}{2^{k-2m}-1} < m-2 \quad (2.3)$$

Since $\frac{t}{2^t-1}$ is decreasing in $t \in (0, \infty)$ and $k-2m \geq 1$, the largest value of the left hand side in the inequality (2.3) is $\frac{1}{2^1-1} = 1$ which means that the strict inequality (2.3) is valid for all $m \geq 4$. As a result, the sequence $F(m, k)$ is positive for all $m \geq 4$.

When $m = 3$, the sequence $F(3, k)$ is

$$F(3, k) = \frac{2^k - 32k + 128}{48(k-3)} = \frac{2^5 [2^{k-5} - (k-4)]}{48(k-3)}$$

which is positive for all $k > 2 \cdot 3 = 6$. The proof of Lemma 2.3 is complete. \square

Lemma 2.4. *Let*

$$h(t) = \begin{cases} \frac{e^t(e^t - 1 - t)}{(e^t - 1)^2}, & t \neq 0 \\ \frac{1}{2}, & t = 0 \end{cases}$$

on $(-\infty, \infty)$. Then, for any fixed $s \in (0, 1)$, the ratio $\frac{h(st)}{h^s(t)}$ is increasing in t from $(0, \infty)$ onto $(\frac{1}{2^{1-s}}, 1)$.

Proof. It is easy to see that

$$\lim_{t \rightarrow 0} \frac{h(st)}{h^s(t)} = \frac{\lim_{t \rightarrow 0} h(st)}{\lim_{t \rightarrow 0} h^s(t)} = \frac{\frac{1}{2}}{\frac{1}{2^s}} = \frac{1}{2^{1-s}}$$

and

$$\lim_{t \rightarrow \infty} H_s(t) = \frac{\lim_{t \rightarrow \infty} h(st)}{\lim_{t \rightarrow 0} h^s(t)} = \frac{1}{1^s} = 1.$$

Direct differentiating and expanding to power series give

$$\begin{aligned} \frac{d}{dt} \left[\frac{h(st)}{h^s(t)} \right] &= - \frac{se^{(1+s)t} \left[\begin{array}{l} (t-2)e^{(1+2s)t} + (t+2)e^{2st} + (2-st)e^{(2+s)t} \\ + 4(s-1)te^{(1+s)t} - (2st^2 + 3st + 2)e^{st} \\ - (st+2)e^{2t} + (2st^2 + 3t + 2)e^t + (s-1)t \end{array} \right]}{(e^t - 1)^3 (e^{st} - 1)^3 h^{s+1}(t)} \\ &= \frac{se^{(1+s)t} \sum_{k=7}^{\infty} \left(\begin{array}{l} (3k+2)s^k + 2(2s+1)^k + ks(s+2)^{k-1} \\ + 2k^2 s^{k-1} + 4k(s+1)^{k-1} \\ + 2k(1+2^{k-2})s + 2^{k+1} \\ - [2^{k+1}s^k + 2(s+2)^k \\ + 4ks(s+1)^{k-1} + 2k(1+2^{k-2})s^{k-1} \\ + k(2s+1)^{k-1} + 2k^2s + 3k + 2] \end{array} \right) \frac{t^k}{k!}}{(e^t - 1)^3 (e^{st} - 1)^3 h^{s+1}(t)} \\ &= \frac{se^{(1+s)t} \sum_{k=7}^{\infty} \left(\begin{array}{l} \sum_{m=1}^{k-1} [k2^{k-m} \binom{k-1}{m-1} + 4k \binom{k-1}{m} + 2^{m+1} \binom{k}{m}] s^m \\ + 2k(1-k+2^{k-2})(s-s^{k-1}) \\ - \sum_{m=1}^{k-1} [k2^m \binom{k-1}{m} + 4k \binom{k-1}{m-1} + 2^{k-m+1} \binom{k}{m}] s^m \end{array} \right) \frac{t^k}{k!}}{(e^t - 1)^3 (e^{st} - 1)^3 h^{s+1}(t)} \\ &= \frac{se^{(1+s)t} \sum_{k=7}^{\infty} \left(\begin{array}{l} \sum_{m=1}^{k-1} [k2^{k-m} \binom{k-1}{m-1} + 4k \binom{k-1}{m} + 2^{m+1} \binom{k}{m}] (s^m - s^{k-m}) \\ + 2k(1-k+2^{k-2})(s-s^{k-1}) \end{array} \right) \frac{t^k}{k!}}{(e^t - 1)^3 (e^{st} - 1)^3 h^{s+1}(t)} \\ &= \frac{se^{(1+s)t} \sum_{k=7}^{\infty} \left(\begin{array}{l} \sum_{3 \leq m < \frac{k}{2}} [k2^{k-m} \binom{k-1}{m-1} - k2^m \binom{k-1}{k-m-1}] \\ + 4k \binom{k-1}{m} - 4k \binom{k-1}{k-m} \\ + 2^{m+1} \binom{k}{m} - 2^{k-m+1} \binom{k}{k-m} \end{array} \right) (s^m - s^{k-m}) \frac{t^k}{k!}}{(e^t - 1)^3 (e^{st} - 1)^3 h^{s+1}(t)} \\ &= \frac{se^{(1+s)t} \sum_{k=7}^{\infty} \left(\begin{array}{l} 2 \sum_{3 \leq m < \frac{k}{2}} \frac{k!}{(m-1)!(k-m-1)!} \left[2 \left(\frac{1}{m} - \frac{1}{k-m} \right) \right. \\ \left. + \frac{1}{2} \left(\frac{2^{k-m}}{k-m} - \frac{2^m}{m} \right) - \frac{2^{k-m-2m}}{(k-m)m} \right] (s^m - s^{k-m}) \end{array} \right) \frac{t^k}{k!}}{(e^t - 1)^3 (e^{st} - 1)^3 h^{s+1}(t)} \\ &= \frac{s^4 e^{(1+s)t} \left(\begin{array}{l} \frac{1-s}{36} t^7 + \frac{1-s^2}{45} t^8 + \frac{22(1-s^3)+15(s-s^2)}{2160} t^9 \\ + \frac{52(1-s^4)+63(s-s^3)}{15120} t^{10} \\ + \frac{285(1-s^5)+470(s-s^4)+238(s^2-s^3)}{302400} t^{11} + \dots \end{array} \right)}{(e^t - 1)^3 (e^{st} - 1)^3 h^{s+1}(t)}. \end{aligned}$$

Utilizing Lemma 2.3 reveals that the derivative $\frac{d}{dt} \left[\frac{h(st)}{h^s(t)} \right]$ is positive for $s \in (0, 1)$ and $t > 0$. Consequently, for $s \in (0, 1)$, the ratio $\frac{h(st)}{h^s(t)}$ is increasing in $t > 0$. The proof of Lemma 2.4 is complete. \square

Lemma 2.5 ([5, Lemma 2.2]). *For $k \geq 0$, the function $(-1)^k \Phi^{(k)}(x)$ is completely monotonic on $(0, \infty)$, with the limits*

$$(-1)^k x^{k+1} \Phi^{(k)}(x) \rightarrow \begin{cases} k!, & x \rightarrow 0^+; \\ \frac{k!}{2}, & x \rightarrow \infty. \end{cases} \quad (2.4)$$

Lemma 2.6 (Bernstein's theorem [12, p. 161, Theorem 12b]). *A function $f(x)$ is completely monotonic on $(0, \infty)$ if and only if*

$$f(x) = \int_0^\infty e^{-xt} d\sigma(t), \quad x \in (0, \infty), \quad (2.5)$$

where $\sigma(s)$ is non-decreasing and the integral in (2.5) converges for $x \in (0, \infty)$.

3. DECREASING MONOTONICITY

In this section, we prove that the function $Y_{m,n}(x)$ defined in (1.2) is decreasing.

Theorem 3.1. *For $m, n \in \{0\} \cup \mathbb{N}$, the function $Y_{m,n}(x)$ defined in (1.2) is decreasing in x from $(0, \infty)$ onto the interval $\left(-\frac{2(m+n+1)!}{m!n!}, -\frac{(m+n+1)!}{m!n!}\right)$. Consequently, for $m, n \in \{0\} \cup \mathbb{N}$, the double inequality*

$$-\frac{2(m+n+1)!}{m!n!} < Y_{m,n}(x) < -\frac{(m+n+1)!}{m!n!} \quad (3.1)$$

is valid on $(0, \infty)$ and sharp in the sense that the lower and upper bounds cannot be replaced by any larger and smaller numbers respectively.

Proof. In the proof of [9, Theorem 4], the first author established that

$$\Phi(x) = \int_0^\infty h(t) e^{-xt} dt. \quad (3.2)$$

Then the ratio $Y_{m,n}(x)$ can be rewritten as

$$\begin{aligned} Y_{m,n}(x) &= -\frac{\int_0^\infty t^{m+n+1} h(t) e^{-xt} dt}{\int_0^\infty t^m h(t) e^{-xt} dt \int_0^\infty t^n h(t) e^{-xt} dt} \\ &= -\frac{\int_0^\infty t^{m+n+1} h(t) e^{-xt} dt}{\int_0^\infty \left[\int_0^t u^m (t-u)^n h(u) h(t-u) du \right] e^{-xt} dt}, \end{aligned}$$

where we used Lemma 2.1. Basing on Lemma 2.2, in order to prove decreasing monotonicity of $Y_{m,n}(x)$, it suffices to show that the ratio

$$\mathfrak{Y}_{m,n}(t) = \frac{t^{m+n+1} h(t)}{\int_0^t u^m (t-u)^n h(u) h(t-u) du} \quad (3.3)$$

is decreasing in $t \in (0, \infty)$. By changing the variable $u = \frac{1+v}{2}t$, the denominator of $\mathfrak{Y}_{m,n}(t)$ becomes

$$\left(\frac{t}{2}\right)^{m+n+1} \int_{-1}^1 (1+v)^m (1-v)^n h\left(\frac{1+v}{2}t\right) h\left(\frac{1-v}{2}t\right) dv.$$

Accordingly, we obtain

$$\begin{aligned} \frac{1}{\mathfrak{Y}_{m,n}(t)} &= \frac{\int_{-1}^1 (1+v)^m (1-v)^n h\left(\frac{1+v}{2}t\right) h\left(\frac{1-v}{2}t\right) dv}{2^{m+n+1} h(t)} \\ &= \frac{1}{2^{m+n+1}} \int_{-1}^1 (1+v)^m (1-v)^n \frac{h\left(\frac{1+v}{2}t\right) h\left(\frac{1-v}{2}t\right)}{h(t)} dv \quad (3.4) \\ &= \frac{1}{2^{m+n+1}} \int_{-1}^1 (1+v)^m (1-v)^n \frac{h(st)}{h^s(t)} \frac{h((1-s)t)}{h^{1-s}(t)} dv, \end{aligned}$$

where $s = \frac{1+v}{2} \in (0, 1)$. From Lemma 2.4, we find that the function $\frac{h(st)}{h^s(t)} \frac{h((1-s)t)}{h^{1-s}(t)}$ is increasing in $t \in (0, \infty)$ for any fixed $s \in (0, 1)$. Hence, the function $\mathfrak{Y}_{m,n}(t)$ is decreasing on $(0, \infty)$. Therefore, the function $Y_{m,n}(x)$ for $m, n \in \{0\} \cup \mathbb{N}$ is decreasing on $(0, \infty)$.

Making use of the limits in (2.4) in Lemma 2.5 yields

$$\begin{aligned} Y_{m,n}(x) &= -\frac{(-1)^{m+n+1} x^{m+n+2} \Phi^{(m+n+1)}(x)}{[(-1)^m x^{m+1} \Phi^{(m)}(x)][(-1)^n x^{k+m+2} \Phi^{(n)}(x)]} \\ &\rightarrow \begin{cases} -\frac{(m+n+1)!}{m!n!}, & x \rightarrow 0^+; \\ -\frac{2(m+n+1)!}{m!n!}, & x \rightarrow \infty. \end{cases} \end{aligned}$$

The proof of Theorem 3.1 is complete. \square

4. NECESSARY AND SUFFICIENT CONDITIONS OF COMPLETE MONOTONICITY

In this section, we discover necessary and sufficient conditions on $\omega_{m,n}$ for the function $\pm(-1)^{m+n+1} \mathcal{Y}_{m,n;\omega_{m,n}}(x)$ defined in (1.3) to be completely monotonic.

Theorem 4.1. For $m, n \in \{0\} \cup \mathbb{N}$ and $\omega_{m,n} \in \mathbb{R}$,

- (1) if and only if $\omega_{m,n} \leq \frac{(m+n+1)!}{m!n!}$, the function $(-1)^{m+n+1} \mathcal{Y}_{m,n;\omega_{m,n}}(x)$ is completely monotonic on $(0, \infty)$;
- (2) if and only if $\omega_{m,n} \geq \frac{2(m+n+1)!}{m!n!}$, the function $(-1)^{m+n} \mathcal{Y}_{m,n;\omega_{m,n}}(x)$ is completely monotonic on $(0, \infty)$;
- (3) the double inequality (3.1) is valid on $(0, \infty)$ and sharp in the sense that the lower and upper bounds cannot be replaced by any larger and smaller numbers respectively.

Proof. As did in the proof of Theorem 3.1, by virtue of the integral representation (3.2) and Lemma 2.1, we acquire

$$\begin{aligned} (-1)^{m+n+1} \mathcal{Y}_{m,n;\omega_{m,n}}(x) &= \left[\int_0^\infty t^{m+n+1} h(t) e^{-xt} dt \right. \\ &\quad \left. - \omega_{m,n} \int_0^\infty t^m h(t) e^{-xt} dt \int_0^\infty t^n h(t) e^{-xt} dt \right] \\ &= \int_0^\infty \left[t^{m+n+1} h(t) - \omega_{m,n} \int_0^t u^m (t-u)^n h(u) h(t-u) du \right] e^{-xt} dt \\ &= \int_0^\infty \left[1 - \frac{\omega_{m,n}}{\mathfrak{Y}_{m,n}(t)} \right] t^{m+n+1} h(t) e^{-xt} dt, \end{aligned}$$

where $\mathfrak{Y}_{m,n}(t)$ is defined by (3.3) and it has been proved in the proof of Theorem 3.1 to be decreasing on $(0, \infty)$. From Lemma 2.4, we conclude that the function $\frac{h(st)}{h^s(t)} \frac{h((1-s)t)}{h^{1-s}(t)}$ is increasing in t from $(0, \infty)$ onto $(\frac{1}{2}, 1)$. Accordingly, by virtue of (3.4), we arrive at the sharp inequalities

$$\frac{1}{2^{m+n+2}} \int_{-1}^1 (1+v)^m (1-v)^n dv < \frac{1}{\mathfrak{Y}_{m,n}(t)} < \frac{1}{2^{m+n+1}} \int_{-1}^1 (1+v)^m (1-v)^n dv.$$

Since

$$\begin{aligned} \int_{-1}^1 (1+v)^m (1-v)^n dv &= \int_0^1 [(1+v)^m (1-v)^n + (1-v)^m (1+v)^n] dv \\ &= 2^{m+n+1} B(m+1, n+1) \\ &= 2^{m+n+1} \frac{m!n!}{(m+n+1)!}, \end{aligned}$$

where we used the formula

$$\begin{aligned} \int_0^1 [(1+x)^{\mu-1} (1-x)^{\nu-1} + (1+x)^{\nu-1} (1-x)^{\mu-1}] dx &= 2^{\mu+\nu-1} B(\mu, \nu) \\ &= 2^{\mu+\nu-1} \frac{\Gamma(\mu)\Gamma(\nu)}{\Gamma(\mu+\nu)} \end{aligned}$$

for $\Re(\mu), \Re(\nu) > 0$ in [3, p. 321, 3.214], the double inequality

$$\frac{1}{2} \frac{m!n!}{(m+n+1)!} < \frac{1}{\mathfrak{Y}_{m,n}(t)} < \frac{m!n!}{(m+n+1)!}$$

is valid and sharp on $(0, \infty)$. Consequently, by virtue of Lemma 2.6, if and only if $\omega_{m,n} \leq \frac{(m+n+1)!}{m!n!}$, the function $(-1)^{m+n+1} \mathcal{Y}_{m,n;\omega_{m,n}}(x)$ is completely monotonic on $(0, \infty)$; if and only if $\omega_{m,n} \geq \frac{2(m+n+1)!}{m!n!}$, the function $(-1)^{m+n} \mathcal{Y}_{m,n;\omega_{m,n}}(x)$ is completely monotonic on $(0, \infty)$.

The double inequality (3.1) follows from complete monotonicity of the functions $\pm(-1)^{m+n+1} \mathcal{Y}_{m,n;\omega_{m,n}}(x)$. The proof of the sharpness of the double inequality (3.1) is the same as did in the proof of Theorem 3.1. The proof of Theorem 4.1 is complete. \square

5. REMARKS

In this section, we list several remarks related to our main results and their proofs in this paper.

Remark 5.1. Lemma 2.4 in this paper generalizes a conclusion in [7, Lemma 2.3]: the function $\frac{h(2t)}{h^2(t)}$ is decreasing from $(0, \infty)$ onto $(1, 2)$.

Remark 5.2. The function $F(x, y)$ discussed in Lemma 2.3 can be reformulated as

$$F(x, y) = \left(\frac{1}{x} - \frac{1}{y-x} \right) \left[2 - \frac{1}{2} \frac{\frac{2^{y-x}}{y-x} - \frac{2^x}{x}}{\frac{1}{y-x} - \frac{1}{x}} - \frac{2^{y-x} - 2^x}{(y-x) - x} \right]$$

in which the functions

$$\frac{\frac{2^{y-x}}{y-x} - \frac{2^x}{x}}{\frac{1}{y-x} - \frac{1}{x}} \quad \text{and} \quad \frac{2^{y-x} - 2^x}{(y-x) - x}$$

can be regarded as special means [2, 10].

Let $x, y \in \mathbb{R}$ such that $0 < x < \frac{y}{2}$. Motivated by Lemma 2.3, we guess that,

- (1) when $2 < x < \frac{y}{2}$, the function $F(x, y)$ is positive;
- (2) when $y > 4$ and $0 < x < 2$, the function $F(x, y)$ is negative.

Furthermore, one can discuss positivity and negativity of the function $F(x, y)$ for all x, y satisfying $0 < x < \frac{y}{2}$.

Remark 5.3. When taking $m = k$ and $n = k + 1$, the double inequality (3.1) in Theorem 3.1 becomes the double inequality (1.1) guessed by the first author in [5, Remark 5.3].

Remark 5.4. For $m, n \in \{0\} \cup \mathbb{N}$, direct differentiation gives

$$Y'_{m,n}(x) = \frac{\Phi^{(m+n+2)}(x)[\Phi^{(m)}(x)\Phi^{(n)}(x)] - \Phi^{(m+n+1)}(x)[\Phi^{(m)}(x)\Phi^{(n)}(x)]'}{[\Phi^{(m)}(x)\Phi^{(n)}(x)]^2}$$

on $(0, \infty)$. The decreasing monotonicity of $Y_{m,n}(x)$ in Theorem 3.1 implies that, for $m, n \in \{0\} \cup \mathbb{N}$, the inequality

$$\Phi^{(m+n+1)}(x)[\Phi^{(m)}(x)\Phi^{(n)}(x)]' > \Phi^{(m+n+2)}(x)[\Phi^{(m)}(x)\Phi^{(n)}(x)],$$

equivalently,

$$\frac{[\Phi^{(m)}(x)\Phi^{(n)}(x)]'}{\Phi^{(m)}(x)\Phi^{(n)}(x)} < \frac{\Phi^{(m+n+2)}(x)}{\Phi^{(m+n+1)}(x)},$$

is valid on $(0, \infty)$.

We guess that, for $m, n \in \{0\} \cup \mathbb{N}$, the function

$$\Phi^{(m+n+1)}(x)[\Phi^{(m)}(x)\Phi^{(n)}(x)]' - \Phi^{(m+n+2)}(x)[\Phi^{(m)}(x)\Phi^{(n)}(x)]$$

is completely monotonic in $x \in (0, \infty)$.

One can also consider necessary and sufficient conditions on $\Lambda_{m,n} \in \mathbb{R}$ for $m, n \in \{0\} \cup \mathbb{N}$ such that the function

$$\Phi^{(m+n+1)}(x)[\Phi^{(m)}(x)\Phi^{(n)}(x)]' - \Lambda_{m,n}\Phi^{(m+n+2)}(x)[\Phi^{(m)}(x)\Phi^{(n)}(x)]$$

and its negativity are respectively completely monotonic on $(0, \infty)$.

Remark 5.5. This paper is the sixth one in a series of articles including [5, 6, 7, 8, 9].

REFERENCES

- [1] M. Abramowitz and I. A. Stegun (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series **55**, Reprint of the 1972 edition, Dover Publications, Inc., New York, 1992.
- [2] P. S. Bullen, *Handbook of Means and Their Inequalities*, Revised from the 1988 original [P. S. Bullen, D. S. Mitrinović and P. M. Vasić, *Means and Their Inequalities*, Reidel, Dordrecht], Mathematics and its Applications **560**, Kluwer Academic Publishers Group, Dordrecht, 2003; available online at <https://doi.org/10.1007/978-94-017-0399-4>.
- [3] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, Translated from the Russian, Translation edited and with a preface by Daniel Zwillinger and Victor Moll, Eighth edition, Revised from the seventh edition, Elsevier/Academic Press, Amsterdam, 2015; available online at <https://doi.org/10.1016/B978-0-12-384933-5.00013-8>.
- [4] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht-Boston-London, 1993; available online at <https://doi.org/10.1007/978-94-017-1043-5>.
- [5] F. Qi, *Complete monotonicity and monotonicity of two functions defined by two derivatives of a function involving trigamma function*, HAL preprint (2020), available online at <https://hal.archives-ouvertes.fr/hal-02993723>.

- [6] F. Qi, *Lower bound of sectional curvature of manifold of beta distributions and complete monotonicity of functions involving polygamma functions*, MDPI Preprints **2020**, 2020110315, 22 pages; available online at <https://doi.org/10.20944/preprints202011.0315.v1>.
- [7] F. Qi, *Monotonicity of a ratio involving trigamma and tetragamma functions*, OSF Preprints (2020), available online at <https://doi.org/10.31219/osf.io/5rfb8>.
- [8] F. Qi, *Necessary and sufficient conditions for two functions defined by two derivatives of a function involving trigamma function to be completely monotonic or monotonic*, OSF Preprints (2020), available online at <https://doi.org/10.31219/osf.io/6ar4p>.
- [9] F. Qi, *Some properties of several functions involving polygamma functions and originating from the sectional curvature of the beta manifold*, São Paulo J. Math. Sci. **15** (2021), in press; available online at <https://doi.org/10.1007/s40863-020-00193-1>.
- [10] F. Qi and D. Lim, *Integral representations of bivariate complex geometric mean and their applications*, J. Comput. Appl. Math. **330** (2018), 41–58; available online at <https://doi.org/10.1016/j.cam.2017.08.005>.
- [11] R. L. Schilling, R. Song, and Z. Vondraček, *Bernstein Functions*, 2nd ed., de Gruyter Studies in Mathematics **37**, Walter de Gruyter, Berlin, Germany, 2012; available online at <https://doi.org/10.1515/9783110269338>.
- [12] D. V. Widder, *The Laplace Transform*, Princeton University Press, Princeton, 1946.
- [13] Z.-H. Yang and J.-F. Tian, *Monotonicity and inequalities for the gamma function*, J. Inequal. Appl. **2017**, Paper No. 317, 15 pages; available online at <https://doi.org/10.1186/s13660-017-1591-9>.

INSTITUTE OF MATHEMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO 454010, HENAN, CHINA; COLLEGE OF MATHEMATICS, INNER MONGOLIA UNIVERSITY FOR NATIONALITIES, TONGLIAO 028043, INNER MONGOLIA, CHINA; SCHOOL OF MATHEMATICAL SCIENCES, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN 300387, CHINA

Email address: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com
URL: <https://qifeng618.wordpress.com>, <https://orcid.org/0000-0001-6239-2968>

COLLEGE OF MATHEMATICS AND PHYSICS, INNER MONGOLIA UNIVERSITY FOR NATIONALITIES, TONGLIAO 028043, INNER MONGOLIA, CHINA

Email address: hlx2980@163.com, hanlingxiong@outlook.com
URL: <https://orcid.org/0000-0003-1346-9943>

COLLEGE OF MATHEMATICS AND PHYSICS, INNER MONGOLIA UNIVERSITY FOR NATIONALITIES, TONGLIAO 028043, INNER MONGOLIA, CHINA

Email address: hongpingyin@qq.com
URL: <https://orcid.org/0000-0001-7481-0194>