

IL-7 as an adjuvant for mucosal vaccine development

Magali Rancez, S Logerot, S. Figueiredo, B Charmeteau-De-Muylder, I Bourgault-Villada, A Couëdel-Courteille, R Cheynier, Carolina Moraes-Cabe

▶ To cite this version:

Magali Rancez, S Logerot, S. Figueiredo, B Charmeteau-De-Muylder, I Bourgault-Villada, et al.: IL-7 as an adjuvant for mucosal vaccine development. 22nd International AIDS Conference (AIDS 2018), Jul 2018, Amsterdam, Netherlands. 2018. hal-03000425

HAL Id: hal-03000425 https://hal.science/hal-03000425

Submitted on 11 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.









IL-7 AS AN ADJUVANT FOR MUCOSAL VACCINE DEVELOPMENT

M. Rancez¹, S. Logerot¹, S. Figueiredo¹, B. Charmeteau-de-Muylder¹, I. Bourgault-Villada¹, A.Couëdel-Courteille², R. Cheynier¹ 1- Institut Cochin / IINSERM U 1016 / CNRS UMR 8104 / Université Paris Descartes - Sorbonne Paris Cité, Paris, France. 2- Institut Cochin / IINSERM U 1016 / CNRS UMR 8104 / Université Paris Descartes - Sorbonne Paris Cité / Université Paris Diderot, Paris, France.

INTRODUCTION and OBJECTIVES

Development of mucosal immune responses, while certainly essential to allow protection against numerous sexually transmitted infections, is still hard to induce in most models. Despite some progresses in the development of adjuvants possibly helping for the setting of this kind of immunity, results in terms of protection as for induction of the local antibody production and specific T-cells remain largely disappointing.

We have showed that systemic injection of IL-7 stimulates chemokine-induced recruitment of circulating T-cells into the mucosa (Blood 2009, 114(4):816-25), and evidenced local IL-7 expression in acutely infected mucosae (Front Immunol 2017, 19;8:588).

In this study, we aimed at using recombinant glycosylated simian IL-7 (sIL-7) as an immunostimulating molecule for vaginal immune responses in vaccine protocols. We administrated sIL-7 by spray directly on rhesus macague's vaginal mucosa surface and guantified local transcription of 19 chemokines by gRT-PCR and cell infiltration by immunochemistry plus image analysis to define the optimal dose of sIL-7 to be sprayed on mucosal surface. Then, six macaques were immunized with antigens (Diphtheria Toxoid (DT) and the HIV-1 gp41-P1 peptide) applied directly on the vaginal mucosa, two days after either IL-7 or PBS administration. The immunizations were repeated thrice, four months apart, and the macaques were euthanized 2 weeks after the last immunization. Antigen-specific IgA and IgG productions were quantified in vaginal secretions by ELISA. Antigen-specific plasma cells were detected by reverse immunohistochemistry and by B-cell ELISPOT.



ACKNOWLEDGMENT

The authors thank the staffs of the Institut Pasteur (Paris, France) and Infectious Disease Models and Innovative Therapies (IDMIT, Fontenay-aux-Roses, France) Primate Centers. The authors greatly acknowledge Maryline Favier, Franck Letourneur and Pierre Bourdoncle, respectively heads of HistIM, GENOM'IC and IMAG'IC core facilities of the Institut Cochin. The authors thank Thomas Guilbert from IMAG'IC platform at the Institut Cochin for writing the routine for ImageJ image analysis software. The authors acknowledge Cytheris S.A. (now Revimmune Inc.), for R-sIL-7gly (sIL-7). This work was supported by ANRS, INSERM and Université Paris Descartes.

Presented at the 22nd International AIDS Conference – Amsterdam, the Netherlands





2 ш

RIS DID

Fig 1: Impact of vaginal sIL-7 administration on chemokine transcription

(A) mRNAs coding for CXCL12 and CCL7, were quantified and normalized to HPRT mRNAs in each vaginal biopsies sampled one month before (PRE) and 48^H after the administration of $1\mu g$ (n=2), $5\mu g$ (n=2), $10\mu g$ (n=3) or 15µg (n=2) of sIL-7 (POST), by vaginal spray. Each point represents the median of a triplicate quantification in a given biopsy. (B) mRNAs coding for 19 chemokines, were guantified in triplicate and normalized to HPRT mRNAs in each vaginal biopsies sampled one month before (PRE) and 48^{H} after the administration of 10µg (n=3) and 15µg (n=2) of sIL-7 (POST), by vaginal spray. Each point represents the mean of all the quantifications performed per animal PRE and POST administration of the sIL-7. Four biopsies per animal PRE and POST administration. *: p<0.05 by the Wilcoxon Signed-Rank Test.

Fig 2: Immune cell recruitment into vaginal chorion of sIL-7 treated macaques

(A) Sections of monkey vaginal mucosa biopsies sampled one month before (Ctrl), or 48^{H} after (IL-7) the administration of 10µg of sIL-7 by vaginal spray were labeled for CD3, CD11c, PM-2K and CD20 (Alexa-Fluor® 546) in combinaison with CD4 or CD8, DC-SIGN, CD20 and MHC-II MamuLa-DR (Alexa-Fluor[®] 488), respectively. Nuclei were stained with DAPI. EP= Pluristratified Epithelium. (B, C) Cell quantifications by image analysis of IHF staining (expressed as number of cells per 500,000 µm2 of chorion \pm SEM) were realized from 20 to 32 vaginal mucosa macagues, except for PM-2K+ cells piopsies (8 guantification: 5 macagues) sampled one month before (Ctrl - white bars) and 48^{H} after (IL-7 48^{H} - black bars) the administration of 10µg of sIL-7. (Mann-Whitney U test).



maintenance

Specific anti-DT IgGs (A, D) and IgAs (B, E), and anti-P1 IgAs (C, F), were quantified by ELISA in vaginal secretions of 6 rhesus macaques that received vaginal administration of either 10µg of sIL-7 (black bars, n=3) or PBS (white bars, n=3), followed at day 2 by antigen administration (DT and P1). Similar immunizations were repeated thrice, every 4 months. Results are expressed as optical density over IgG or IgA concentration at any time-point for the 3 animals from the same group in any given sample for specific anti-DT (A-B, D-E) or as ng of specific anti P1 per mg of total IgAs (C, F). Bars and error bars represent means \pm SEM. Non interpretable data due to menstruations were excluded. (MANOVA with Fisher test). D0: Administration of sIL-7 or PBS; W: Week post-antigens administration.

> sIL-7-dependent immunization protocol induces ectopic lymphoid follicles genesis in vaginal mucosa IL-7+Aqs



We here evidenced that 10µg of sIL-7, administered by spray on rhesus macaques vaginal mucosa, stimulates local transcription of CCL2, CCL5, CCL7, CCL17, CCL19, CCL20, CCL22, CXCL10, CXCL12, CXCL13, and CX₃CL1 by 48^H after administration. Local expression of these chemokines leads to massive mDC, macrophages, NK, CD4⁺ T-, CD8⁺ T-, and infiltration into the vaginal mucosa.

Healthy rhesus macagues immunized with a cocktail of antigens (Diphtheria Toxoid + the HIV-1 gp41-P1 peptide) directly applied on the vaginal mucosa two days after vagin (10µg) administration developed an earlier and stronger mucosal antigen-specific immune response as compared to animals receiving the antigenic cocktail alone. Indeed, dur immunization protocol, sIL-7 pre-treatment promoted then sustained a robust production of antigen-specific IgAs and IgGs in vaginal secretions, as detected by ELISA.

By attracting immune cells, local sIL-7 administration prepares the mucosal immune system, gathering conditions that result in enhanced antigen-specific vaginal immune res Moreover, throughout the immunization protocol, CXCL12, CXCL13, CCL19 and CCL21 known to contribute to tertiary lymphoid organ formation and organization (Nat Immunol 344-53), which could support the induction of immune responses, in particular by sustaining productive immunoglobulin class switching in germinal centers, were increased onl vaginal mucosa of sIL-7 treated macaques, and sIL-7 pretreatment favored the formation of tertiary lymphoid organs genesis in the vaginal mucosa. Besides, high levels of IgAs vaginal secretions are produced by mucosally localized plasma cells as evidenced by reverse IHF. CXCL12 is known as mucosal chemoattractant molecule for IgA- and IgG-secreting cells (J Exp Med 2002, 196(1):65-75), suggesting its role in mucosal infiltration of DT- and P1-specific plasma cells. Finally, we evidenced an Increased frequency of B-cell resp peripheral blood.

Altogether, these data demonstrate the capacity of sIL-7 to be used as a mucosal adjuvant leading to enhanced mucosal antibody response, a very promising strategy to protection to numerous sexually transmitted infections.

RESULTS **MUCOSAL ADJUVANT POTENTIAL OF sIL-7 in DT and P1 IMMUNIZATION**

Preferential localization of DT-specific lgAs pl cells in the vaginal mucosa of sIL-7 treated macagu



Figure 4: DT specific IgGs plasma cells are more abundant in draining lymph nodes while DT specific IgAs plasma cells a abundant in the vaginal mucosa of sIL-7 treated macagues at necro IgG and IgA anti-DT-specific ASC were quantified by B-cells ELISPOT assay on cells from iliac LN from 3 macaques per group (A) and on isolated cells from chorion from 1 animal per group (C), sampled at necropsy (group IL-7+Ags, black PBS+Ags, white bars). Results are expressed as IgG or IgA anti-DT-specific ASG cells. Bars and error bars represent means \pm SEM. LN: Lymph nodes; Ags: Antie antibody secreting cells. (Mann-Whitney U test). Sections of monkey iliac Li vaginal fornix (D), sampled at necropsy, were labeled after incubation with DT. an anti-DT antibody. Representative examples of the draining LN of sIL-7-tre 7+Ags) or ctrl (PBS+Ags) macaques co-labeled with DT (green) and anti-IgG (r anti-IgA (red, D) are shown. Nuclei were stained with DAPI. (B, D) Arrowheads single-stained cells. (D) Arrows identify IgA anti-DT-specific ASC.

Fig 5: Ectopic lymphoid follicles in vaginal mucosa treated macaques at necropsy

Sections of monkey vaginal mucosa, lower part (A1, B1), upper part and vaginal fornix (A3, B3), sampled at necropsy (2 weeks after the from sIL-7-treated (A, IL-7+Ags) or control (B, PBS+Ags) macag labeled for CD3 (Alexa-Fluor[®] 546), CD20 (Alexa-Fluor[®] 488) and CD3 Fluor® 633), respectively. Nuclei were stained with DAPI. EP= Plur Epithelium; Ch= Chorion.

CONCLUSION

<u>IgAs plasma</u> <u>macaques</u>
DT lgG
s <u>100μm</u> DT IgA
POT assay on isolated ted cells from vaginal 2-7+Ags, black bars; or 7-specific ASC per 10 ⁶ <i>es; Ags: Antigens; ASC:</i> nkey iliac LN (B) and tion with DT, thanks to N of sIL-7-treated (IL- id anti-IgG (red, B) or) Arrowheads indicate
mucosa of sIL-7
), upper part (A2, B2), eks after the 3 rd boost) +Ags) macaques were 488) and CD31 (Alexa- API. EP= Pluristratified
19, CCL20, CCL21, 08⁺ T-, and B-cells
after vaginal sIL-7 ndeed, during the
mmune responses. <i>Immunol 2006, 7</i> : reased only in the vels of IgAs in the S-secreting plasma B-cell response in
strategy to confer