




ANALYSIS OF THE KUROSHIO EXTENSION DECadal VARIABILITY : Internal Mechanisms and External Forcings

^{1,2} Fedele G. (giusy.fedele@unive.it),

³ Penduff T., ⁴ Pierini S., ² Bellucci A., ² Masina S.



¹ Ca' Foscari University of Venice ² Centro Euro-Mediterraneo sui Cambiamenti Climatici, CMCC, Bologna

³ Institut des Géosciences de l'Environnement, IGE, Grenoble ⁴ Parthenope University of Naples



Università
Ca' Foscari
Venezia




PRIMAVERA

Credit: C. Henze, NASA Advanced Supercomputing

Why study the Kuroshio Extension (KE) from a Climatic Point of View?


- Highest eddy kinetic energy level (EKE) in the North Pacific Ocean. EKE changes lead to high temperature anomalies that are capable of enhancing the variability of the midlatitude coupled ocean-atmosphere system.
- The KE is known to undergo relevant variations on the decadal time scale. The mechanisms that generate the mean structure of the KE and its low frequency variability (LFV) are still under debate.




(Figures Adapted from Qiu and Chen, 2005)

Introduction

Frontal Variability (FV)



Broad-Scale Variability (BV)



- Synchronization with external forcings (Qiu and Chen, 2005, 2010).
- Many authors have suggested that nonlinear intrinsic oceanic mechanisms can play a fundamental role in the phenomenon (Pierini, 2006).
- The KE LFV is interpreted as a case of intrinsic climate variability paced by external forcing through the coherence resonance mechanism (Pierini, 2014).

QUESTION

What is the role of the intrinsic variability and external forcing in driving the KE LFV?

QUESTION

What is the role of the intrinsic variability and external forcing in driving the KE LFV?

METHOD

To explore a *large* ensemble of ocean-only eddy-permitting model simulations: OCCIPUT


OCCIPUT*

SIMULATION	EXTENT AND RESOLUTION	MEMBERS AND DURATION	FORCING	SIMULATED VARIABILITY	TIME SAMPLING
OCCITENS	Global 1/4°	50-members ENSEMBLE 1980-2015	Full variability (ERA-interim)	Forced and intrinsic variability	5 days
OCCICLIM	Global 1/4°	1 member 300 years	Climatological Seasonal Cycle	Pure intrinsic variability	5 days

Reference SSH isoline of the jet 0.7 meters

Average in the box (141-153°E, 32-38°N)

Max total geost. velocity in the box
(141-153°E, 30-40°) integrated zonally



Gaussian metrics are appropriate since the climatological PDFs for these variables are pretty gaussian

We take advantage of some definitions and notations (Leroux et al., 2018):

$$f_i(t) = \langle f_i(t) \rangle + f'_i(t)$$

We define $f_i(t, f)$ as every Index after nonlinear detrending, where t stands for the time (yr) and i is the index of the ensemble member.

We use overbars for the time-mean operator and angled brackets for the ensemble-mean operator.

$$\left\{ \begin{array}{l} A_{intrinsic} = \sqrt{\varepsilon^2} \\ A_{forced} = \sigma(f) \\ A_{tot, i} = \sigma_{fi} \\ R = \frac{A_{intrinsic}}{A_{forced}} \end{array} \right.$$

$\varepsilon^2(t)$: the time variance operator applied to the ensemble mean


$\varepsilon^2(t)$: ensemble variance operator applied to $f'_i(t)$

$A_{intrinsic}$: Amplitude of the intrinsic variability

A_{forced} : Amplitude of the forced variability

A_{tot} : Amplitude of the total variability in each member

R : Amplitude Ratio



Gaussian metrics are appropriate since the climatological PDFs for these variables are pretty gaussian

We take advantage of some definitions and notations (Leroux et al., 2018):

$$f_i(t) = \langle f_i(t) \rangle + f'_i(t)$$

We define $f_i(t, j)$ as every Index after nonlinear detrending, where t stands for the time (yr) and i is the index of the ensemble member.

We use overbars for the time-mean operator and angled brackets for the ensemble-mean operator.

$$\left\{ \begin{array}{l} A_{intrinsic} = \sqrt{\varepsilon^2} \\ A_{forced} = \sigma(f) \\ A_{tot, i} = \sigma_{fi} \\ R = \frac{A_{intrinsic}}{A_{forced}} \end{array} \right.$$

$\varepsilon^2(t)$: the time variance operator applied to the ensemble mean


$\varepsilon^2(t)$: ensemble variance operator applied to $f'(t)$

$A_{intrinsic}$: Amplitude of the intrinsic variability

A_{forced} : Amplitude of the forced variability

A_{tot} : Amplitude of the total variability in each member

R : Amplitude Ratio



Pure intrinsic variability


EXTENT AND RESOLUTION	MEMBERS AND DURATION	FORCING	TIME SAMPLING
Global 1/4°	1 member 300 yrs	Climatological Seasonal Cycle	5 days

STRATEGY:


TO CREATE A PSEUDO-ENSEMBLE FROM OCCICLIM

50 members x 36 yrs have been built randomly from the indices already computed, in order to compare the pure intrinsic variability with the total one given by OCCITENS.


OCCITENS




OCCITENS: 10% isoline



OCCICLIM




OCCICLIM: 10% isoline




Φ, Ψ : Latitude and Velocity Indices. Changing the Y-axis variable we always get the same footprint of the attractor.

OCCITENS


OCCICLIM





OCCITENS: 10% isoline




TWO
MODES




OCCICLIM: 10% isoline




OCCITENS




OCCITENS: 10% isoline



OCCICLIM




OCCICLIM: 10% isoline






OCCITENS

OCCICLIM





Center of Gravity






OCCITENS

OCCICLIM



INTRINSIC VARIABILITY PACE BY AN EXTERNAL FORCING





The external forcing acts reducing the entropy of the dynamical system

CONCLUSIONS


- OCCIPUT ensemble simulation : disentangle the intrinsic and forced variabilities
- The KE LFV is a case of intrinsic variability paced by an external forcing
- The external forcing acts reducing the entropy of the dynamical system

WORK IN PROGRESS

What is the ability of state of the art models in reproducing the observed KE LFV: the role of horizontal resolution -PRIMAVERA project- Model data from **HIGHRESMIP-CMIP6**

HighResMIP -CMIP6	Coupled Model	Configuration	Resolution	Forcing	Output Frequency	Vertical Levels
OCEAN ATMOSPHERE	NEMO v3.6 CAM 4	CMCC-CM2- HR4**	0.25° 1°	Control-1950	Daily (100 yrs)	50
OCEAN ATMOSPHERE	NEMO v3.6 CAM 4	CMCC-CM2- VHR4**	0.25° 0.25°	Control-1950	Daily (100 yrs)	50

** Model data are from the H2020 PRIMAVERA Project



PRIMAVERA

Fedele et al., in preparation



THANKS FOR YOUR ATTENTION


Fedele G. (giusy.fedele@unive.it)



Università
Ca' Foscari
Venezia



PRIMAVERA



Credit: C. Henze, NASA Advanced Supercomputing