



HAL
open science

Access to a stabilized i -motif DNA structure through four successive ligation reactions on a cyclopeptide scaffold

Alexandre Devaux, Laureen Bonnat, Thomas Lavergne, Eric Defrancq

► **To cite this version:**

Alexandre Devaux, Laureen Bonnat, Thomas Lavergne, Eric Defrancq. Access to a stabilized i -motif DNA structure through four successive ligation reactions on a cyclopeptide scaffold. *Organic & Biomolecular Chemistry*, 2020, 18 (32), pp.6394-6406. 10.1039/D0OB01311K . hal-03001772

HAL Id: hal-03001772

<https://hal.science/hal-03001772v1>

Submitted on 23 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Access to a stabilized *i*-motif DNA structure through four successive ligation reactions on a cyclopeptide scaffold

Alexandre Devaux, Laureen Bonnat, Thomas Lavergne* and Eric Defrancq*

Univ. Grenoble Alpes, Département de Chimie Moléculaire, CNRS, 570 rue de la chimie, CS 40700, Grenoble 38000, France

ABSTRACT. I-motifs are largely underexplored tetraplex nucleic acid structures which have been suggested to play essential biological functions and might constitute future therapeutic targets. I-motifs generally require acidic conditions to fold *in vitro*, a particularity which significantly complicates the use of native *i*-motif forming sequences for interactions studies with potential ligands and biological components (*e.g.* proteins). In this context, we report herein on the assembly of a peptide-DNA conjugate capable to fold at room temperature into a stable *i*-motif structure at neutral pH. To achieve the controlled assembly of the *i*-motif forming conjugate, we developed a new synthetic pathway of four successive orthogonal ligation reactions between bifunctional C-rich DNA strands and a tetrafunctional cyclopeptide scaffold.

Introduction

Over half a century ago, the seminal contributions of Watson, Crick, Wilkins and Franklin established the double-helical structure of DNA in which two antiparallel strands are held together through canonical A/T and G/C base pairing. However, the past decades have brought accumulating evidence of the existence of a wide variety of alternative nucleic acid topologies including hairpin, triplex, cruciform and tetraplex structures. In particular, G- and C-rich sequences have been shown to form tetraplex architectures, respectively G-quadruplex (G4) and *i*-motif structures.^{1, 2} Unlike duplex DNA, G4 and *i*-motif formation does not involve canonical Watson-Crick base pairing: indeed a G4 DNA is formed from the stacking of Hoogsteen hydrogen bonded guanine tetrads whereas for *i*-motif formation two anti-parallel duplexes are held together by hemi-protonated C-C⁺ base pairs intercalated with each other (Fig. 1).




Figure 1. A/ G-quartet with Hoogsteen hydrogen bonded guanine and schematic drawing of a G4 structure and B/ hemi-protonated C-C⁺ base pairs and schematic representation of an *i*-motif structure.

G4 DNA structures have been the subject of intensive investigations. Sequencing and bioinformatics analyses of the human genome indicate that it contains as much as 700,000 sequences having the potential to form stable G-quadruplex structures³ and a large number of studies reported on their central biological functions within physiological and pathological processes.⁴ Although *i*-motif DNA structures have attracted less attention, several recent studies suggest that they may also hold essential biological functions. By using Quadparser algorithm, more than 5,000

putative *i*-motif forming sequences have been found in the human genome. Interestingly more than 600 were found to be located in promotor sequences of genes such as *c-myc*, *bcl-2* or *c-kit*.⁵ Recent studies have proposed that *i*-motifs are implicated in the regulation of transcription⁶ and the integrity of telomeric DNA.⁷ Proteins such as hnRNP LL have been identified to interact with *i*-motif structures acting as an activating transcription factor.⁸ More recently, two independent studies have demonstrated the presence of *i*-motif structures in human cells through *in cellulo* NMR spectroscopy⁹ and immunofluorescence by using a specific antibody against *i*-motif.¹⁰ In addition to the regulation of cellular processes in human cells, an *i*-motif forming sequence has been identified in the long terminal repeat promoter of the HIV-1 proviral genome where it has been shown to modulate viral transcription.¹¹

Encouraged by those inspiring reports, several groups have engaged in successful efforts to identify small molecule ligands that interact with C-rich sequences, stabilizing or destabilizing the corresponding *i*-motif structures.¹² Some of those ligands have been shown to modulate gene regulation and expression and they now constitute promising probes and drug candidates targeting *i*-motif sequences within cancer-related genes or viral genomes.⁶

Beyond their biological occurrences, *i*-motifs have also attracted a huge interest in the field of functional nanostructures. *I*-motif structures which can fold and unfold reversibly depending on their environments, have been manipulated by chemists to create nanotools such as in cell pH sensors¹³ or pH responsive shape-memory hydrogels.¹⁴ Indeed, the formation and the stability of *i*-motif DNA structures are affected *in vitro* by a number of parameters such as the length and the nature of the sequence, the solute concentration and the pH. In particular, *i*-motifs are generally formed *in vitro* under acidic pH, a particularity that complicates the use of native *i*-motif sequences to study the interactions of potential ligands.

As a consequence, numerous strategies have been explored to stabilize the *i*-motif structure and to promote its formation at physiological pH. Extending the number of C-C+ base pairs and adjusting the sequence of the loop region can help to stabilize the folded structure.¹⁵ For a given sequence of interest, modifications of the sugar,¹⁶ base¹⁷ and phosphate moieties¹⁸ can promote the formation of the *i*-motif. However, all of those modifications are susceptible to affect the recognition of the native structure of interest by putative targeting ligands.

In this context, the design of biomolecular conjugates capable to fold into well-defined and stable native *i*-motif structures would be of interest as they would find applications for the development of structure-specific *i*-motif binding ligands as well as for the identification of their structure-specific binding partners (*e.g.*, proteins) *via* pull-down strategies.

In the past, our group, along with others have proposed the use of peptide scaffolds to assemble stable mimics of tetraplex DNA structures (as in Fig. 2).¹⁹ Our strategy is based on the use of regioselectively addressable cyclopeptide platforms for directing the intramolecular assembly of the attached oligonucleotides into the desired tetraplex topology. By site-specifically fixing DNA and/or RNA G4-forming sequences onto cyclopeptide scaffolds, we have been able to trigger the formation of topologically controlled and thermally stabilized G4 architectures such as parallel tetramolecular DNA and RNA G4, antiparallel dimeric DNA G4 and DNA:RNA hybrid G4 structures.^{19a, 20, 21} For the preparation of the parallel DNA (or RNA) tetrameric systems (Fig. 2A), four oxime ligation reactions were simultaneously performed between an aminoxy-containing cyclopeptide platform and DNA (or RNA) strands that bear aldehyde complementary functions. Whereas the controlled anchoring of nucleic acid strands to form antiparallel structures such as the *i*-motif structure (Fig. 2B) required two sequential ligations and for that purpose oxime ligation (OL) and copper-catalyzed azide-alkyne cycloaddition (CuAAC) reactions were performed between an aminoxy- and azide-containing cyclopeptide and DNA strands that bore either aldehyde or alkyne functions.²² More recently, a more sophisticated system consisting in a biologically relevant viral G4 (Fig. 2C) has been elaborated with the site-specific attachment of two distinct oligonucleotides in three successive ligations involving the OL, CuAAC and a third compatible ligation, namely the thiol-iodoacetamide (TC) SN2 reaction. This latter ligation was found to fulfil the criteria of chemical orthogonality associated with the synthesis of such bioconjugate and the resulting thioether linkage had no perturbative effects on the folding of the G4 structure.²³ All the resulting tetraplex mimics obtained so far displayed increased stabilities and reduced structural polymorphism. Owing to those properties, the different G4 motifs have found applications in the selection and characterization of structure-specific ligands.²⁴

To further expand this approach and to access more sophisticated biomolecular systems mimicking nucleic acid structures of high biological interest, we explored strategies to fully control the

assembly of DNA-peptide conjugates *via* four successive ligations steps. As a proof of concept, we report herein on our unsuccessful (**1a**, Fig. 2D) and successful (**1b**, Fig. 2E) optimization efforts to assemble and stabilize a biologically relevant *i*-motif structure derived from the human telomeric sequence (CCCTAA)₄. Unlike the *i*-motif structure in Fig. 2B, which consisted of four tracts of four cytosine forming a tetramolecular *i*-motif structure without biological relevance, the *i*-motif structure within **1b** was designed to form *via* the association of four tracts of three cytosine forming a bimolecular *i*-motif structure displaying two 5'-TAA^{3'} loops. This new conjugate therefore precisely mimics the biologically relevant structure of the human telomeric *i*-motif motif and in particular the loops organization, a feature which is crucial for the subsequent selection and characterization of structure-specific ligands.

To assemble the *i*-motif conjugates **1a** and **1b** *via* four successive orthogonal reactions, we have chosen to make use of the three previously exploited ligations (*vide supra*) *i.e.* the oxime ligation (OL), the copper-catalyzed azide–alkyne cycloaddition (CuAAC) and the thiol-iodoacetamide SN2 reaction (TC) with the objective to use one of them twice. Oxime ligation has already been used for the successive incorporation of reporters for the preparation of ODN bis-conjugates.²⁵ However both partners for the OL (*i.e.* aldehyde and aminoxy functions) are not very stable and thus, if one would want to perform two OL stepwise for the synthesis of sophisticated systems such as **1a/b**, non-trivial protection strategies should be implemented (*i.e.* use of different protecting groups for the aldehyde or aminoxy groups). In our hand, TC reactions have been found to be more sluggish than OL and CuAAC during the formation of DNA-peptide conjugates and therefore repeating twice this reaction in the course of the assembly of those conjugates may prove problematic. On the other hand, CuAAC reactions have been repeatedly used for the successive incorporation of reporters at internal positions or at 3', 5' extremities of nucleic acids.²⁶ Furthermore, protected alkynes can be used along with a free alkyne to allow for sequential CuAAC reactions after mild deprotections.^{26a} CuAAC can also be performed in presence of free amines which can later be converted into azides *via* mild diazotransfer reaction for subsequent CuAAC reactions.²⁷ Finally, both the alkyne and the azide are stable in a wide range of non-reductive experimental conditions. Based on the robustness and flexibility of the CuAAC reaction, we decided to explore the formation of the *i*-motif structural mimic *via* the stepwise assembly of

peptide-DNA conjugates through four successive ligations with one OL, one TC and two CuAAC reactions.

Two strategies were explored and involved the use of two diversely functionalized cyclopeptides and three distinct bifunctional C-rich oligonucleotides. The first strategy (**Route A**, Scheme 1) consists in starting with the oxime ligation following by a first CuAAC reaction to attach bifunctional oligonucleotide **2**. Subsequent thiol-iodoacetamide SN2 (TC) and CuAAC reactions are performed to incorporate the second bifunctional oligonucleotide **3** toward the final conjugate (OL → CuAAC → TC → CuAAC). Unfortunately, the route A was found inefficient to afford the expected target **1a** (Fig. 2D). A second route (**Route B**, Scheme 2) was thus explored consisting in starting the assembly with the least yield-efficient TC chemistry using oligonucleotide **10** and achieving the assembly by the most efficient OL reaction. The following synthetic pathway was therefore used: TC → CuAAC → CuAAC → OL. This route allowed us to access the conjugate **1b** (Fig. 2E). The resulting structural mimic was found to fold into a stable *i*-motif structure at physiological pH and room temperature paving the way to functional applications.





Figure 2. Use of cyclopeptide scaffold for the design of stabilized tetraplex DNA structures: **A/** Parallel G4-DNA,^{19a} **B/** I-motif DNA structure without loops,²² **C/** Antiparallel G4 structure from HIV virus sequence,²³ **D/** Telomeric *i*-motif mimic **1a** and **E/** Telomeric *i*-motif mimic **1b**. For the sake of clarity, the positive charges shared by the two cytosines involved in the C-C base pairs of structures B, D and E are not displayed. Dotted lines represent the base pairing.

Results and discussion

Route A (OL → CuAAC → TC → CuAAC)

This strategy requires the attachment of oligonucleotides **2** ($5'Y\text{-ACCCTAACCCCT-X}^{3'}$) and **3** ($5'Y\text{-ACCCTAACCCCT-Z}^{3'}$) in which X represents a 3'-aldehyde linker, Y a 5'-alkynyl linker and Z a 3'-thiol linker through OL, CuAAC and TC reactions onto corresponding peptide scaffold **4a** bearing the complementary functional groups (Scheme 1).

Cyclopeptide scaffold **4a** was prepared using standard fluorenylmethoxycarbonyl (Fmoc)/tBu solid-phase peptide synthesis (SPPS) on a 2-chlorotrityl resin followed by a cyclization reaction in solution according to reported procedures (Scheme S1).²⁸ Briefly, the biotin, aminoxy and azido moieties were incorporated during SPPS by using a *N* ϵ -biotin-functionalized lysine building block, a *N* ϵ -ethoxyethylidene (Eei) protected aminoxy functionalized lysine building block and the *L*-azidonorleucine building block, respectively. To equip the cyclopeptide with the chloroacetamide function and the second azido group, two other orthogonally protected lysine (*i.e.* *N* ϵ -Boc and *N* ϵ -Dde protected lysine) were introduced during SPPS. After the cleavage from the resin in slight acidic conditions, to prevent the cleavage of ethoxyethylidene (Eei) protecting group, and cyclization of the linear peptide in diluted conditions, the chloroacetamide group was anchored on the lysine side chain using chloroacetic anhydride after selective removal of bis-*N*-[1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl] (Dde) lysine protecting group with hydrazine. Final treatment in acidic conditions allowed the removal of tert-butoxycarbonyl (Boc) as well as the ethoxyethylidene protections and afforded desired cyclopeptide scaffold **4a**. To synthesize oligonucleotides **2** and **3**, solid-phase oligonucleotide syntheses (SPOS), using phosphoramidite chemistry, were performed on commercially available glyceryl CPG solid support and 3'-thiol modifier C3 S-S CPG solid support, respectively. Alkynyl function was introduced at the 5'-terminus of oligonucleotides **2** and **3** by using commercially available 5'-hexynyl (β -cyanoethyl) phosphoramidite. Following the cleavage from the solid support and deprotection steps, oligonucleotide **2** was obtained after oxidative cleavage of the 3'-diol group. To obtain oligonucleotide **3**, a 100 mM DTT treatment was performed to reduce the disulfide linkage generated upon the cleavage from the solid support.



Scheme 1. Synthesis of *i*-motif **1a** through *Route A*. a) Oligonucleotide **2** in 0.4 M ammonium acetate buffer (pH 4.5) at 55°C for 45 min; b) 100 mM HEPES buffer (pH 7.4), CuSO_4 (6 equiv), THPTA (30 equiv), sodium ascorbate (30 equiv) at 37°C for 2 h; c) Oligonucleotide **3** in $\text{H}_2\text{O}/\text{CH}_3\text{CN}$ (9:1, v/v), TCEP (2 equiv), 500 mM KCl, DIEA (45 equiv), KI (60 equiv) at RT for 3 h; d) 50 mM NaHCO_3 buffer/MeOH, CuSO_4 (2.8 equiv), imidazole-1-sulfonyl azide hydrochloride ISAHC (20 equiv), 30 min. at 60°C.

Toward the incorporation of the first oligonucleotide, **2** was grafted on cyclopeptide scaffold **4a** through oxime bond formation to generate intermediate **5**. The coupling reaction was carried

out in ammonium acetate buffer (pH 4.5) at 55°C using a slight excess of aminoxy-containing peptide **4a**. The reaction was monitored by RP-HPLC and proceeded to completion within 45 min. Purification through RP-HPLC followed by desalting using size-exclusion chromatography (SEC) afforded pure conjugate **5** with an isolated yield of 65%. A first intramolecular CuAAC reaction was then performed in HEPES buffer (pH 7.4) in the presence of CuSO₄, tris(benzyltriazolylmethyl)amine (THPTA) ligand and sodium ascorbate at 37°C for 2 h to afford, following desalting using size-exclusion chromatography (SEC) and RP-HPLC purification, conjugate **6** in 72% yield. Conjugates **5** and **6** were characterized by ESI MS analysis, which showed an excellent agreement between the experimentally determined molecular weight and the calculated value (Figures S15 and S17).

From intermediate **6**, we envisioned to perform first the TC reaction and to finish the assembly with a CuAAC reaction. This decision was mainly guided by the instability of the thiol function during CuAAC reaction. Toward **1a** from **6**, two alternative pathways have been investigated that consist in performing the diazo transfer reaction either after (*a*) or before (*b*) the TC reaction (Scheme 1). Previous studies have shown a poor reactivity for the TC reaction when a first oligonucleotide was already anchored on the cyclopeptide scaffold and consequently an optimization of TC reaction had to be performed.²³ The assembly of conjugate **6** with thiol containing oligonucleotide **3** was thus carried out in water at room temperature in presence of tris(2-carboxyethyl)phosphine (TCEP) to prevent the formation of disulfide species, potassium iodide (KI) for *in situ* activation of the reaction through the transient formation of a reactive iodoacetamide group and DIEA to obtain a pH around 8.5. Furthermore, under those optimized conditions, the addition of KCl is absolutely required for cations shielding to neutralize unfavorable charge repulsions between conjugate **6** and non-complementary oligonucleotide **3**. Intermediate **7** was obtained in 74% yield after RP-HPLC purification and then subjected to diazo transfer reaction with ISAHC (imidazole-1-sulfonylazide hydrochloride) reagent by using previously reported conditions²⁷ to afford conjugate **8** in 80% yield. The other alternative (*way b*) has also been explored by performing first the diazo transfer reaction on conjugate **6** to give intermediate **9** and subsequent TC reaction with oligonucleotide **3** in the same optimized conditions as above to afford conjugate **8** in

30% overall yield. Conjugate **8** obtained by both pathways was characterized by ESI-MS analysis which showed a good agreement between the experimentally determined molecular weight and the calculated value (Fig. S21). It is noteworthy that the diazo transfer reaction was found efficient and compatible with the sophisticated conjugates **6** and **7**.

Toward **1a**, a second CuAAC reaction had to be achieved through reaction of the alkyne moiety on the 5'-extremity of the grafted oligonucleotide and the azide function carried by the cyclopeptide. By using the standard and above depicted CuAAC conditions as well as in presence of 500 mM KCl with the aim to neutralize unfavorable charge repulsions between the oligonucleotide strands, no transformation was observed as evidenced by RP-HPLC monitoring (data not shown). We only observed the recovery of the starting material **8**. Variation of reactant equivalents and temperatures did not prove beneficial to promote the formation of **1a**. As *i*-motifs are known to form at acidic pH, we thus envisioned to perform CuAAC reaction in slightly acidic conditions compatible with CuAAC. At slight acidic pH, the DNA strands of conjugate **8** were expected to adopt an antiparallel topology (pre-organization to an *i*-motif structure) and thus to hold the azide and alkyne groups close to one another thereby lowering the entropic cost of the CuAAC reaction. CD analysis of starting conjugate **8** indeed revealed the formation of antiparallel *i*-motif topology at pH 4, 5 and 6 whereas at pH 7 no structuring was noticed (Fig. S39). CuAAC reactions were thus carried out at pH 4, 5, 6 and 7 in the above conditions (with and without KCl) and again no conversion toward **1a** could be observed by RP-HPLC monitoring regardless of the conditions: only the starting material **8** was observed (data not shown). The absence of conversion could be explained by the ineffectiveness of the CuAAC reaction in acidic conditions.

Although *Route A* was found unsuitable to afford the desired constrained *i*-motif **1a** it provided precious information to design the second strategy (*Route B*). Indeed, a major drawback with *Route A* is that during the last synthetic step *i.e.* CuAAC reaction, the molecular weight of starting material **8** and final product **1a** are the same thus precluding the use of mass spectrometry to monitor the transformation and identify the final conjugate. Also, the TC reaction was again shown to be the least efficient and hard to carry out when an oligonucleotide is already


grafted on the molecular scaffold. We thus decided for *Route B* to use the least efficient TC reaction in an early stage of the synthetic process while the most efficient OL reaction would be used at the last step. Furthermore, intramolecular OL reaction results in the formation of water by-product leading to a final product whose molecular weight is distinct from its precursor thus allowing the use of MS for the monitoring of the reaction and identification of final product **1b**. Moreover, OL reaction is known to be more efficient in acidic conditions and the formation of i-motif is also known to be favored in acidic pH. We thus anticipated a pre-organization of the i-motif leading to a spatial proximity of the two reactive function (*i.e.* aldehyde and aminoxy moieties) to promote the last intramolecular reaction.

Route B (TC → CuAAC → CuAAC → OL)

This strategy requires the successive attachment of oligonucleotide **10** ($5'Y\text{-ACCCTAACCCCT-X}3'$) in which *X* represents a 3'-azide linker and *Y* a 5'-thiol linker and above described oligonucleotide **2** through TC, CuAAC and OL reactions onto corresponding peptide scaffold **4b** bearing the complementary functional groups (Scheme 2). Cyclopeptide **4b** thus contains a protected aminoxy group for OL reaction, an alkyne function for the first CuAAC reaction, a chloroacetamide moiety for TC reaction and an amino function to be transformed into an azide for the second CuAAC reaction. It was prepared as for peptide **4a** using standard fluorenylmethoxycarbonyl (Fmoc)/tBu solid-phase peptide synthesis (SPPS) on a 2-chlorotrityl resin and cyclization reaction in solution (see Scheme S2). Again the biotin and aminoxy moieties were incorporated during SPPS by using a *N* ϵ -biotin-functionalized lysine building block and a *N* ϵ -ethoxyethylidene (Eei) protected aminoxy functionalized lysine building block and two other orthogonally protected lysine (*i.e.* *N* ϵ -Alloc and *N* ϵ -Dde protected lysine) were introduced during SPPS for the introduction of chloroacetamide and azide moieties as above. The alkyne group was introduced as a building block during SPPS by using a *N* ϵ -pentyne functionalized lysine building block. After the cleavage from the resin in slight acidic conditions (*i.e.* to prevent the removal of Eei aminoxy protecting group) and cyclization of the linear peptide, the chloroacetamide group was anchored on the lysine side chain using chloroacetic anhydride after selective removal of Dde protecting group with hydrazine. It

should be noticed that allylic alcohol should be used during the Dde cleavage to prevent the reduction of Alloc protection. Alloc protecting group was next removed through treatment with $\text{Pd}^0(\text{PPh}_3)_4$ in presence of phenylsilane and afforded desired cyclopeptide **4b**. Oligonucleotide **10** ($5'$ -Y-ACCCTAACCCCT-X $3'$) was prepared by solid-phase oligonucleotide syntheses (SPOS) using phosphoramidite chemistry, on commercially available 3'-PT-amino-modifier C3 CPG. Thiol function was introduced at the 5'-terminus by using commercially available 5'-S-Trityl modifier C6 phosphoramidite. The introduction of azide function was achieved through diazo transfer reaction as described above on the S-trityl protected thiol oligonucleotide following by thiol deprotection in presence of silver nitrate as reported.²⁹

With *Route B*, the first step consists in the use of the TC reaction which was carried out in the above depicted conditions *i.e.* in presence of tris(2-carboxyethyl)phosphine (TCEP), potassium iodide (KI) and DIEA. In this case, the addition of KCl for cations shielding is not required. It should be noted that due to the presence of an azide group on oligonucleotide **10**, TCEP had to be added during the reaction and not prior to the addition of all other components in order to avoid partial reduction of the azide group. Desalting using size-exclusion chromatography (SEC) followed by RP-HPLC purification afforded pure conjugate **11** with an isolated yield of 53%. A first intramolecular CuAAC reaction was then performed in PBS buffer (pH 7.4) in the presence of CuSO_4 , THPTA ligand and sodium ascorbate at 37°C for 2 h. As previously observed, the addition of KCl was required in order to neutralize unfavorable charge repulsions. After addition of EDTA, desalting using size-exclusion chromatography (SEC) and RP-HPLC purification, conjugate **12** was obtained in 40% yield. The free amino function on the peptide moiety was then transformed into an azide by using a diazo transfer reaction which was carried out as described above to afford, in quantitative yield, conjugate **13** after purification by size-exclusion chromatography. Conjugates **11-13** were characterized by ESI-MS analysis which showed a good agreement between the experimentally determined molecular weight and the calculated value (Figures S26, S29 and S31).



Scheme 2. Synthesis of *i*-motif **1b** through *Route B*: a) Oligonucleotide **10** in water, TCEP (1 equiv), DIEA (60 equiv), KI (60 equiv), pH 8.5, 90 min. at 37°C; b) 50 mM PBS buffer, 10% DMF, $CuSO_4$ (3 equiv), THPTA (30 equiv), sodium ascorbate (30 equiv), 500 mM KCl pH 7.4, 2h at 37°C; c) 50 mM $NaHCO_3$ buffer/MeOH, $CuSO_4$ (3 equiv), imidazole-1-sulfonyl azide hydrochloride ISAHC (30 equiv), 1h at 60°C; d) Oligonucleotide **2** in 50 mM PBS buffer, 10% DMF, $CuSO_4$ (3 equiv), THPTA (30 equiv), sodium ascorbate (30 equiv), 500 mM KCl pH 7.4, 1h at 60°C; e) 1% TFA in water, 1h at room temperature.

From intermediate **13**, two pathways can be considered to access the expected product **1b**. Indeed, a first OL could be performed to attach oligonucleotide **2** followed by a second CuAAC reaction or *vice-versa*. As mentioned above, the major drawback with the CuAAC reaction is that there is no molecular weight difference between the starting material and the reaction product and a retention time difference is expected to analyze and purify the crude reaction mixture.

Furthermore, the CuAAC reaction was found to be inefficient in the last step of the assembly of **1a**. For those reasons, we envisioned to perform the CuAAC reaction first followed by the most reactive oxime reaction for the last step. CuAAC reaction was carried out in the optimized conditions as described above *i.e.* in presence of 500 mM KCl for neutralizing the unfavorable charge repulsions between **13** and oligonucleotide **2**. However, a low conversion rate was obtained at room temperature (around 20-30% according to LC-MS monitoring). Increasing the temperature to 60°C was found beneficial to afford intermediate **14** with a satisfactory 48% yield after desalting using size-exclusion chromatography (SEC) and RP-HPLC purification.

The last step consisted in the one-pot deprotection of the aminoxy function and the oxime reaction. Treatment of conjugate **14** in mild acidic conditions (1% TFA) allowed the cleavage of Eei protecting group and the simultaneous oxime bond formation. As anticipated the monitoring of the reaction by LC-MS analysis showed the loss of the Eei protection as well as the loss of a water molecule thus demonstrating the formation of the oxime bond to afford the *i*-motif forming conjugate **1b** (Figures 3 and S37, S38). It should be mentioned that the intermediate free aminoxy compound was not observed which emphasizes the high reactivity of the oxime reaction. As expected, the efficiency of the oxime formation might have been promoted by the preorganization of the DNA strands into an *i*-motif structure under the acidic pH required to perform the oxime reaction.




Figure 3. UPLC-MS monitoring of OL for the crude transformation of **14** to **1b** corresponding to all peak. Insert: UPLC profile of crude mixture, column heated at 60°C, detection at 260 nm.

Characterization of *i*-motif formation

To confirm the formation of an *i*-motif structure within **1b** and to gage its stability, CD studies were performed at different pH at a concentration of 2 μ M at 20°C. Previous investigations have indeed shown that *i*-motif DNA can be associated with a specific CD signature.³⁰ CD spectrum of the biomolecular assembly **1b** recorded at pH 4 displayed a positive peak at 286 nm and a negative peak at 262 nm, characteristic of the formation of *i*-motif topology. For comparison CD spectrum of the native telomeric sequence **15** d(5'(CCCCTAA)₃CCCT3') was recorded and showed similar signals (Fig. 4A displays the CD spectra of **1a** and **15** at pH=4) that confirms the formation of *i*-motif structure in **1b**.

The influence of pH on the formation of the *i*-motif structure was then investigated with both compounds **1b** and oligonucleotide **15**. By varying the pH from 4 to 7 the shape of the signal changed with the appearance of a positive peak at around 270 nm at pH = 7 (Fig. 4B) for native telomeric sequence **15** which could be attributed to the “unstructured” DNA motif in agreement with reported studies.^{15a} Whereas for constrained system **1b**, the characteristic signal of the ~~antiparallel~~ *i*-motif structure is still observed at pH 7 (Fig. 4C). This data is of interest as it suggests that interaction studies with potential ligands as well as with biological components (*e.g.* proteins) might be performed at physiological pH with the mimic structure **1b**.




Figure 4. **A/** CD spectra of biomolecular assembly **1b** (black line) and control sequence **15** (red line) at pH = 4; **B/** CD spectra of native telomeric sequence **15** at different pH; **C/** CD spectra of biomolecular assembly **1b** at different pH. Measurements were carried out at 2 μ M in 10 mM PBS buffer at 20°C.

The thermal stability of the *i*-motif structure formed within **1b** was further evaluated by CD denaturation studies at pH 6 and at neutral pH 7 (Figures S40 and S41). At pH 6 the melting temperature was 40°C, a temperature which is on par with the reported melting temperature of the native telomeric *i*-motif sequence at pH 5.5.^{15a} On the other hand, we found that the telomeric *i*-motif structural mimic within **1b** possess a melting temperature of 31°C at pH 7. This demonstrates a significant stabilization from the native structure, which was shown to unfold into a random coil at pH 6.5 with no determinable *T_m* at physiological pH,^{15a} and further confirms that **1b** might be exploited to study the interaction of *i*-motif-targeting ligands at neutral pH and room temperature.

Experimental section

General remarks

ESI mass spectra were realized using an Esquire 3000 spectrometer from Bruker or an Acquity UPLC/MS system from Waters equipped with an SQ Detector 2. MALDI-TOF mass spectra were carried out by using an Autoflex Bruker with hydropiccolinic acid (HPA) (45 mg, ammonium citrate (4 mg) in 500 mL H₂O/CH₃CN) as the matrix. Peptides were analyzed in the positive mode and oligonucleotides and conjugates in the negative mode.

Peptide synthesis

General details: The peptide synthesis was performed on a Syro II synthesizer using Fmoc/tBu strategy on a 2-chlorotrityl resin. The course of reactions were monitored by using UPLC-MS system Waters including reverse phase chromatography using a Nucleosil C18 column (130 Å, 2.1 x 50 mm, 1.7 µm) within a column heater set at 60°C and with detection by UV at 214 nm and 250 nm and combined electron spray ionization mass spectrometry. RP-HPLC purifications were performed on a Gilson or Waters system with Nucleosil C18 column (100 Å, 250 x 21 mm, 7 µm) with UV monitoring at 214 nm and 250 nm. 20 mL min⁻¹ flow linear gradient was applied from 95%

solvent A1 (0.1% formic acid in water) and 5% solvent B1 (0.1% formic acid in acetonitrile) to 100% B1 for 20 min.

Synthesis of peptide 4a: The linear peptide was synthesized using 2-chlorotrityl (loading of 0.83 mmol/g) commercial resin. Fmoc-Gly-OH (3 equiv) was coupled on the resin in anhydrous DCM in a glass reaction vessel fitted with a sintered glass. pH was adjusted to 8 by using DIEA. The mixture was stirred for 1h at room temperature. A DCM/MeOH/DIEA solution (17/2/1, v/v/v) was then added. Fmoc protecting group was removed using three washes with a 20% piperidine solution in DMF (40 mL). The resin loading was assessed by quantification of free dibenzofulvene using UV absorbance at 299 nm (loading of 0.40 mmol g⁻¹, yield: 48%). The elongation was then performed on a Syro II using Fmoc/tBu strategy on the above prepared resin using HBTU as activator. Fmoc-Ala-OH, Fmoc-Gly-OH, Fmoc-Lys(Boc)-OH, Fmoc-Pro-OH, Fmoc-Lys(Dde)-OH, and Fmoc-Lys(biotin)-OH were commercially available. Fmoc-azidonorleucine and Fmoc-Lys(Eei-Aoa)-OH were synthesized according to reported protocols.^{31, 32} The elongated peptide was then cleaved from the resin by using DCM/TFE/AcOH solution (70/20/10, v/v/v). The resulting solution was evaporated under vacuum and the peptide was precipitated in ether to give a white powder. The crude product was used without any further purification in the next step. The cyclization of the linear peptide was performed in DMF at 10⁻³ M concentration in the presence of PyBOP (1.2 equiv). The pH was adjusted to 8-9 by using DIEA and the solution was stirred at room temperature until complete peptide cyclization occurred (UPLC monitoring). The solvent was evaporated under vacuum then the crude peptide was precipitated in ether and purified on RP-HPLC and freeze-dried to give a white powder. ESI-MS (+): *m/z* calcd for C₇₈H₁₂₇N₂₀O₁₉S: 1679.9, *m/z* found: 1680.5 [M+H]⁺. The Dde protecting group was then removed using a 2% hydrazine solution in DMF. The mixture was stirred at room temperature until complete deprotection (UPLC monitoring). The solvent was evaporated under vacuum then the crude peptide was precipitated in ether to give a white powder that was used without any further purification. The peptide was next dissolved in DMF to reach a 3.10⁻³ M concentration. The pH was adjusted to 8-9 using DIEA and chloroacetic anhydride (2 equiv) was added. The reaction was stirred for 1h30 at room temperature. The solvent was evaporated under vacuum then the crude peptides were precipitated in ether. The crude product was purified on RP-HPLC and freeze-dried to obtain a white powder. ESI-MS (+): *m/z* calcd for

$C_{70}H_{116}N_{20}O_{18}S$: 1591.8, m/z found: 1613.9 $[M+Na]^+$, HR-ESI-MS (+): m/z calcd for $C_{70}H_{116}N_{20}O_{18}S$: 1591.8186, m/z found: 1591.8196. The Boc protection on the lysine side chain and ethoxyethylidene on the aminoxy moiety were then cleaved through treatment with a TFA/TIS/ H_2O (80/16/4, v/v/v) solution and the mixture was stirred for 1h at room temperature. The solvent was then evaporated under vacuum and the crude peptide was precipitated in ether to afford **4a** as a white powder. ESI-MS (+): m/z calcd for $C_{61}H_{102}N_{20}O_{15}S$: 1421.7, m/z found: 1421.9 $[M+H]^+$.

Synthesis of peptide 4b: Peptide **4b** was prepared using the same procedure as for peptide **4a**. The following protected amino acids were used: Fmoc-Ala-OH, Fmoc-Gly-OH, Fmoc-Lys(Boc)-OH, Fmoc-Pro-OH, Fmoc-Lys(Dde)-OH, Fmoc-Lys(Alloc)-OH and Fmoc-Lys(biotin)-OH which are commercially available and Fmoc-Lys (N-pentynoic acid)-OH and Fmoc-Lys(Eei-Aoa)-OH building blocks which were prepared as previously reported.^{28, 32} After cleavage from the resin with DCM/TFE/AcOH solution (70/20/10, v/v/v) and cyclization as above, the cyclized intermediate peptide was purified by RP-HPLC and freeze-dried to give a white powder. ESI-MS (+): m/z calcd for $C_{82}H_{128}N_{18}O_{20}S$: 1716.92, m/z found: 1716.97 $[M+H]^+$. Dde removal and introduction of chloroacetamide group were carried out as above. ESI-MS (+): m/z calcd for $C_{74}H_{118}N_{18}O_{19}S$: 1629.34, m/z found: 1629.95 $[M+H]^+$. The peptide was then dissolved in anhydrous CH_2Cl_2 / anhydrous DMF (3/1, v/v) and $Pd[P(Ph)_3]_4$ (0.2 equiv) and $PhSiH_3$ (100 equiv) were added. The reaction was stirred at room temperature for 30 minutes and quenching with MeOH. The solution was evaporated under vacuum and finally purified on RP-HPLC to afford peptide **4b**. ESI-MS (+): m/z calcd for $C_{70}H_{115}N_{18}O_{17}S$: 1546.27, m/z found: 1546.93 $[M+H]^+$; ESI-HRMS (+): m/z calcd for $C_{70}H_{114}N_{18}O_{17}S$: 1545.8018, m/z found: 1545.8003 $[M]^+$.

Oligonucleotide synthesis

General details. The oligonucleotides were prepared using β -cyanoethylphosphoramidite chemistry on a 3400 DNA synthesizer at 1 μ mol scale. RP-HPLC analyses were performed on a Waters HPLC system using C18 Nucleosil column (Macherey-Nagel, 100 Å, 250 X 4.6 mm, 5 μ m) with UV-monitoring at 260 and 280 nm. The RP-HPLC purifications of oligonucleotides were performed on

a Gilson system with Nucleosil C-18 column (Macherey-Nagel, 100 Å, 250 X 10 mm, 7 µm) with UV-monitoring at 260 and 280 nm using a 4 mL/min flow linear gradient. Solvent A2 (50 mM triethylammonium acetate buffer with 5% acetonitrile) and solvent B2 (acetonitrile with 5% water) were used.

Synthesis of oligonucleotide 2: Oligonucleotide **2** was prepared according to a previously reported procedure.²² ESI-MS (-): m/z calcd for $C_{112}H_{148}N_{37}O_{70}P_{12}$: 3502.60, m/z found: 3502.68 [M-H]⁻.

Synthesis of oligonucleotide 3: The 5'-hexynyl, 3' S-protected intermediate oligonucleotide precursor of oligonucleotide **3** was obtained from automated synthesis using commercially available 5'-hexynyl (β-cyanoethyl) phosphoramidite and 3'-thiol-modifier C3 S-S CPG resin at 1 µmol scale. The crude product was purified on RP-HPLC. ESI-MS (-): m/z calcd for $C_{116}H_{156}N_{37}O_{70}P_{12}S_2$: 3624.5, m/z found: 3625.5 [M-H]⁻. Dithiothreitol (100 equiv) was then added to a solution of 3' S-protected oligonucleotide in Tris.HCl buffer 1M, pH 8.5 to obtain a final concentration of 10^{-3} M. The mixture was stirred for 1h at room temperature and the crude was purified on RP-HPLC then desalted to afford oligonucleotide **3**. ESI-MS (-): m/z calcd for $C_{113}H_{151}N_{37}O_{69}P_{12}S$: 3534.4, m/z found: 3534.9 [M-H]⁻.

Synthesis of oligonucleotide 10: Precursor oligonucleotide bearing a 5'S-trityl group and a 3'-amino group was obtained from automated synthesis using commercially available 5'-thiol modifier C6 phosphoramidite and 3'-PT-amino-modifier C3 CPG resin at 1 µmol scale. The crude product was purified on RP-HPLC. ESI-MS (-): m/z calcd for $C_{132}H_{171}N_{38}O_{69}P_{12}S$: 3797.73, m/z found: 3797.11 [M-H]⁻. The intermediate oligonucleotide was then suspended in NaHCO₃ buffer (50 mM, pH 8.5)/MeOH (3/1, v/v) in a closed screw-cap plastic vial. CuSO₄ (3 equiv) and imidazole-1-sulfonyl azide hydrochloride (ISAHC, 30 equiv) were added. Afterwards, the solution was heated at 60°C. The reaction was followed by RP-HPLC and UPLC-MS. Finally, the solution was cooled at room temperature and desalted by SEC. ESI-MS (-): m/z calcd for $C_{132}H_{169}N_{40}O_{69}P_{12}S$: 3821.7, m/z found: 3821.5 [M-H]⁻. The azide containing oligonucleotide was next suspended in triethylammonium acetate (TEAA) buffer (0.1M, pH 6.5). 0.15 volume of 1M aqueous silver nitrate solution and 0.5 volume of acetic acid were added. The mixture was vortexed and left to react for 45 minutes

at room temperature. Afterwards, 0.2 volume of 1M aqueous DTT solution was added. The mixture was vortexed and left to react for 15 minutes at room temperature. After, the solution was centrifuged in order to remove the silver-DTT complex and the supernatant was recovered. The precipitate was washed with 3 volumes of TEAA buffer. Finally, the solution was desalted by SEC. ESI-MS (-): m/z calcd for $C_{113}H_{155}N_{40}O_{69}P_{12}S$: 3579.63, m/z found: 3579.58 [M-H]⁻.

Conjugate synthesis

Synthesis of conjugate 5 (oxime ligation): Oligonucleotide **2** (1 equiv, 500 nmol) was dissolved in 0.4 M ammonium acetate buffer (pH 4.5, concentration 10^{-3} M) and aminoxy peptide **4a** (2 equiv, 1 μ mol) was added. The solution was stirred at 55°C for 45 min. The crude mixture was purified on RP-HPLC. Conjugate **5** was desalted on a NAP-25 column and freeze dried (325 nmol, 65% yield). MALDI-Tof-MS (-): m/z calcd for $C_{173}H_{245}N_{57}O_{84}P_{12}S$: 4906.3, m/z found: 4906.9 [M-H]⁻.

Synthesis of conjugate 6 (CuAAC reaction): Conjugate **5** (300 nmol) was dissolved in 100 mM HEPES buffer (pH 7.4, concentration 10^{-4} M) and $CuSO_4$ (6 equiv), THPTA (30 equiv), and sodium ascorbate (30 equiv) were added to this solution. The reaction was stirred at 37°C for 2h and stopped with the addition of 0.5 M ethylenediaminetetraacetic acid (EDTA) solution (50 equiv). The resulting reaction mixture was desalted using a NAP-25 column and the crude was purified on RP-HPLC. Conjugate **6** was desalted on a NAP-25 column and freeze dried (216 nmol, 72% yield). ESI-MS (-): m/z calcd for $C_{173}H_{245}N_{57}O_{84}P_{12}S$: 4906.3, m/z found: 4907.2 [M-H]⁻.

Synthesis of conjugate 7 (thioether formation): Conjugate **6** (1 equiv, 100 nmol) and oligonucleotide **3** (2 equiv, 200 nmol) were dissolved in a solution of H_2O/CH_3CN (9:1, v/v) at $5 \cdot 10^{-4}$ M. TCEP (2 equiv), 500 mM KCl, DIEA (45 equiv), and KI (60 equiv) were added to this solution. The reaction was stirred at room temperature for 3h and the crude was purified on RP-HPLC. Conjugate **7** was desalted on a NAP-25 column and freeze dried (74 nmol, 74% yield). ESI-MS (-): m/z calcd for $C_{286}H_{395}N_{94}O_{153}P_{24}S_2$: 8405.2, m/z found: 8403.5 [M-H]⁻.

Synthesis of conjugate 8 (diazo transfer reaction), way (a): To a solution conjugate **7** (40 nmol, concentration 3.10^{-4} M) in a solution of 50 mM NaHCO₃ / (MeOH/H₂O, 1/1) 75/25 at pH 8.5 were added CuSO₄ (2.8 equiv) and ISAHC (20 equiv). The reaction was stirred at 60°C for 30 min and quenching with 0.5 M EDTA solution (50 equiv). The product was purified on RP-HPLC then desalted and freeze-dried (31.6 nmol,; 79 % yield). ESI-MS (-): *m/z* calcd for C₂₈₆H₃₉₃N₉₆O₁₅₃P₂₄S₂: 8430.8, *m/z* found: 8433.8 [M-H]⁻.

Synthesis of conjugate 9 (diazo transfer reaction): The same diazo transfer protocol was used from conjugate **6** (100 nmol) to give conjugate **9** (50 nmol, 50% yield). MALDI-Tof-MS (-): *m/z* calcd for C₁₇₃H₂₄₃N₅₉O₈₄P₁₂SCl: 4932.3, *m/z* found: 4934.4 [M-H]⁻.

Synthesis of conjugate 8 (thioether formation), way (b): The same thiol chloroacetamide reaction was carried out from conjugate **9** (40 nmol) and oligonucleotide **3** (80 nmol) to afford conjugate **8** (24 nmol, 60% yield).

Synthesis of conjugate 11 (thioether formation): From the starting material oligonucleotide **10** (1 eq, 220 nmol) and peptide **4b** (440 nmol, 2 eq), the same protocol as for the synthesis of conjugate **7** was carried out excepted that TCEP was added during the course of the reaction and no KCl was added (116 nmol, yield: 53%). ESI-MS (-) *m/z* calcd C₁₈₃H₂₆₈N₅₈O₈₆P₁₂S₂: 5089.46, *m/z* found: 5089.68 [M-H]⁻.

Synthesis of conjugate 12 (CuAAC reaction): To a solution of conjugate **11** (1 eq, 150 nmol) in 50 mM PBS buffer (pH 7.4) were added CuSO₄ (3 equiv, THPTA (30 equiv), sodium ascorbate (30 equiv), 500 mM KCl and DMF (10% vol). The reaction was stirred at 37°C for 2h and quenching with 0.5 M EDTA solution (50 equiv). The product was desalted on a NAP-25 column then purified on RP-HPLC and freeze-dried (60 nmol, yield: 40%). ESI-MS (-) *m/z* calcd C₁₈₃H₂₆₈N₅₈O₈₆P₁₂S₂: 5089.46, *m/z* found: 5089.62 [M-H]⁻.

Synthesis of conjugate 13 (diazo transfer reaction): From conjugate **12** (1 eq, 90 nmol), diazo transfer reaction was carried out as for conjugate **6** to afford conjugate **13** in quantitative yield. ESI-MS (-) *m/z* calcd C₁₈₃H₂₆₇N₆₀O₈₆P₁₂S₂: 5116.45, *m/z* found: 5116.21 [M-H]⁻.

Synthesis of conjugate 14 (CuAAC reaction): From the starting conjugate **13** (1 eq, 70 nmol) and oligonucleotide **2** (1.2 eq, 84 nmol), the same protocol as for the preparation of conjugate **12** was applied (42 nmol, yield: 48%). MALDI ToF-MS (-) m/z calcd $C_{295}H_{413}N_{97}O_{156}P_{24}S_2$: 8617.05, m/z found: 8613.61 [M-H]⁻.

Synthesis of conjugate 1b (oxime ligation): The conjugate **14** (1 equiv, 9 nmol) was suspended in water (350 μ L) in a closed screw-cap glass vial. A 2% aqueous TFA solution (350 μ L) was added in order to have a final solution of TFA 1%vol. The reaction was stirred at room temperature for 1h. The product was desalted by SEC then freeze-dried. The yield was considered as quantitative. MALDI ToF-MS (-) m/z calcd $C_{291}H_{405}N_{97}O_{154}P_{24}S_2$: 8528.99, m/z found: 8526.68 [M-H]⁻.

Circular dichroism studies

Analyses were recorded on a Jasco J-810 spectropolarimeter using 1 cm length quartz cuvette. Spectra were recorded at 20 °C or every 5 °C in a range of 5 to 90°C with a wavelength range of 220 to 330 nm. For each temperature, the spectrum was an average of three scans with a 0.5 s response time, a 1 nm data pitch, a 4 nm bandwidth and a 200 nm min⁻¹ scanning speed. Melting temperatures were obtained using Boltzmann fit on Origin software. Each curve fit was only accepted with a $r_{value} > 0.99$.

Conclusion

In summary, we have designed and assembled a peptide-DNA conjugate capable of folding into a stable *i*-motif structure. The assembly was performed *via* the stepwise connection of two *i*-motif forming DNA strands through four consecutive orthogonal ligation reactions. The sequential synthetic pathway was optimized to provide a full control in the assembly of such sophisticated *i*-motif forming conjugate. This detailed optimization effort shades some new light on the concrete hurdles associated with the assembly of such advanced molecular objects. For instance, we found that TC and CuACC reactions were less fit than the OL reaction to achieve the challenging formation of intramolecular covalent bonds within bulky molecular conjugates. We also further

demonstrate than a mild acidic treatment can be used for the one-pot aminoxy deprotection and oxime bound formation in presence of acid sensitive heteropolymer DNA sequence.

The resulting conjugate was found by CD to fold, at room temperature, into a *i*-motif structure at acidic and neutral pH. This stabilization of the secondary structure was obtained without major modifications of a large portion of the native structure. The stabilized *i*-motif mimic may therefore be used to study, at physiologically relevant pH, the interaction of the *i*-motif with putative *i*-motif targeting ligands by using SPR. This unique molecular object might also be used to pull-down and further characterize proteins that interact with *i*-motif structures. In both cases, one could take advantage of the biotin moiety built-in the *i*-motif conjugate for surface immobilization. Finally, such novel *i*-motif forming conjugate with shifted pH sensitivity might be exploited to design original DNA nanodevices with new sets of characteristics and performances.

Supporting Information

RP-HPLC profiles of products and ESI-MS and MALDI Tof-MS spectra of purified ODNs, CD spectra.

Acknowledgment

This work was supported by Agence Nationale de la Recherche (ANR-16-CE11-0006-01). The Nanobio-ICMG platform (FR2607) is acknowledged for providing facilities for synthesis and purification of oligonucleotides (R. Lartia) as well as for mass spectrometry analyses (A. Durand, L. Fort and R. Guéret).

References

- 1 J. Spiegel, S. Adhikari and S. Balasubramanian, The Structure and Function of DNA G-Quadruplexes. *Trends in Chemistry*, 2020, **2**, 123-136.
- 2 (a) K. Gehring, J.-L. Leroy and M. Guéron, A Tetrameric DNA Structure with Protonated Cytosine-Cytosine Base Pairs, *Nature*, 1993, **363** (6429), 561-565; (b) S. Benabou, A. Avino, R. Eritja, C. Gonzalez

-
- and R. Gargallo, Fundamental Aspects of the Nucleic Acid i-Motif Structures, *RSC Adv.*, 2014, **4**, 26956-26980.
- 3 V. S. Chambers, G. Marsico, J. M. Boutell, M. Di Antonio, G. P. Smith and S. Balasubramanian, High-throughput Sequencing of DNA G-Quadruplex Structures in the Human Genome. *Nat. Biotechnol.*, 2015, **33** (8), 877-881.
- 4 T. Tian, Y.-Q. Chen, S.-R. Wang and X. Zhou, G-Quadruplex: A Regulator of Gene Expression and Its Chemical Targeting. *Chem*, 2018, **4** (6), 1314-1344.
- 5 E. P. Wright, J. L. Huppert and Z. A. E. Waller, Identification of Multiple Genomic DNA Sequences Which Form i-Motif Structures at Neutral pH. *Nucleic Acids Research*, 2017, **45** (6), 2951-2959.
- 6 (a) C. E. Kaiser, N. A. Van Ert, P. Agrawal, R. Chawla, D. Yang and L. H. Hurley, Insight into the Complexity of the i-Motif and G-Quadruplex DNA Structures Formed in the KRAS Promoter and Subsequent Drug-Induced Gene Repression, *J. Am. Chem. Soc.* 2017, **139** (25), 8522-8536; (b) S. Takahashi, J. A. Brazier and N. Sugimoto, Topological Impact of Noncanonical DNA Structures on Klenow Fragment of DNA Polymerase *Proc. Natl. Acad. Sci. USA* 2017, **114** (36), 9605-9610; (c) S. Kendrick, A. Muranyi, V. Gokhale, L. H. Hurley and L. M. Rimsza, Simultaneous Drug Targeting of the Promoter MYC G-Quadruplex and BCL2 i-Motif in Diffuse Large B-Cell Lymphoma Delays Tumor Growth, *J. Med. Chem.* 2017, **60** (15), 6587-6597.
- 7 (a) Y. Chen, K. Qu, C. Zhao, L. Wu, J. Ren, J. Wang and X. Qu, Insights into the Biomedical Effects of Carboxylated Single-Wall Carbon Nanotubes on Telomerase and Telomeres, *Nat. Commun.* 2012, **3**, 1074; (b) P. Skolakova, S. Foldynova-Trantirkova, K. Bednarova, R. Fiala, M. Vorlickova and L. Trantirek, Unique C. Elegans Telomeric Overhang Structures reveal the Evolutionarily Conserved Properties of Telomeric DNA, *Nucleic Acids Res.* 2015, **43** (9), 4733-4745.
- 8 (a) B. Roy, P. Talukder, H. J. Kang, S. S. Tsuen, M. P. Alam, L. H. Hurley and S. M. Hecht, Interaction of Individual Structural Domains of hnRNP LL with the BCL2 Promoter i-Motif DNA, *J. Am. Chem. Soc.* 2016, **138** (34), 10950-10962; (b) L. Lannes, P. Young, C. Richter, N. Morgner and H. Schwalbe, Interaction of the N-Terminal Tandem Domains of hnRNP LL with the BCL2 Promoter i-Motif DNA Sequence, *ChemBioChem* 2017, **18** (20), 2033-2044.
- 9 S. Dzatko, M. Krafcikova, R. Hänsel-Hertsch, T. Fessl, R. Fiala, T. Loja, D. Krafcik, J.-L. Mergny, S. Foldynova-Trantirkova and L. Trantirek, Evaluation of the Stability of DNA I-Motifs in the Nuclei of Living Mammalian Cells. *Angew. Chem. Int. Ed.*, 2018, **57** (8), 2165-2169.

-
- 10 M. Zeraati, D. B. Langley, P. Schofield, A. L. Moye, R. Rouet, W. E. Hughes, T. M. Bryan, M. E. Dinger and D. Christ, I-Motif DNA Structures Are Formed in the Nuclei of Human Cells. *Nature Chem.*, 2018, **10** (6), 631-637.
- 11 E. Ruggiero, S. Lago, P. Šket, M. Nadai, I. Frasson, J. Plavec and S. N Richter, A Dynamic i-Motif with a Duplex Stem-Loop in the Long Terminal Repeat Promoter of the HIV-1 Proviral Genome Modulates Viral Transcription. *Nucleic Acids Res.*, 2019, **47** (21), 11057-11068.
- 12 S. S. Masoud and K. Nagasawa, I-Motif-Binding Ligands and Their Effects on the Structure and Biological Functions of i-Motif. *Chem. Pharm. Bull.*, 2018, **66** (12), 1091-1103.
- 13 J. Huang, Y. He, X. Yang, K. Wang, L. Ying, K. Quan, Y. Yang and B. Yin, I-motif-based Nano-Flares for Sensing pH Changes in Live Cells. *Chem. Commun.*, 2014, **50**, 15768-15771.
- 14 W. Guo, C.-H. Lu, R. Orbach, F. Wang, X.-J. Qi, A. Ceconello, D. Seliktar and I. Willner, pH-Stimulated DNA Hydrogels Exhibiting Shape-Memory Properties. *Adv. Mater.*, 2015, **27** (1), 73-78.
- 15 (a) E. P. Wright, J. L. Huppert and Z.A. E. Waller, Identification of Multiple Genomic DNA Sequences Which Form i-Motif Structures at Neutral pH. *Nucleic Acids Res.* 2017, **45** (22), 2951–2959; (b) A. M. Fleming, Y. Ding, R. A. Rogers, J. Zhu, J. Zhu, A. D. Burton, C. B. Carlisle and C. J. Burrows, C. J. 4n–1 Is a “Sweet Spot” in DNA i-Motif Folding of 2'-Deoxycytidine Homopolymers. *J. Am. Chem. Soc.* 2017, **139** (13), 4682-4689.
- 16 (a) L. Lacroix, J.-L. Mergny, J.-L. Leroy and C. Hélène, Inability of RNA To Form the I-Motif: Implications for Triplex Formation. *Biochemistry* 1996, **35** (26), 8715-8722; (b) S. Robidoux and M. J. Damha, D-2-Deoxyribose and D-Arabinose, but Not D-Ribose, Stabilize the Cytosine Tetrad (i-DNA) Structure. *J. Biomol. Struct. Dyn.* 1997, **15** (3), 529-535; (c) H. A. Assi, R. El-Khoury, C. González and M. J. Damha, 2'-Fluoroarabinonucleic Acid Modification Traps G-quadruplex and I-Motif Structures in Human Telomeric DNA. *Nucleic Acids Res.*, 2017, **45** (20), 11535-11546; (d) A. Aviñó, M. Dellafiore, R. Gargallo, C. González, A. M. Iribarren, J. Montserrat and R. Eritja, Stabilization of Telomeric I-Motif Structures by (2'S)-2'-Deoxy-2'-C-Methylcytidine Residues. *ChemBioChem*, 2017, **18** (12), 1123-1128.
- 17 (a) V. B. Tsvetkov, T. S. Zatsepin, E. S. Belyaev, Y. I. Kostyukevich, G. V. Shpakovski, V. V. Podgorsky, G. E. Pozmogova, A. M. Varizhuk and A. V. Aralov, i-Clamp Phenoxazine for the Fine Tuning of DNA I-Motif Stability. *Nucleic Acids Res.*, 2018, **46** (6), 2751-2764; (b) P. Školáková, D. Renčiuk, J. Palacký, D. Krafčík, Z. Dvořáková, I. Kejnovská, K. Bednářová and M. Vorlíčková, Systematic Investigation of Sequence Requirements for DNA I-Motif Formation. *Nucleic Acids Res.* 2019, **47** (5), 2177-2189; (c) P. Perlíková, K. K. Karlsen, E. B. Pedersen and J. Wengel, Unlocked Nucleic Acids with a Pyrene-Modified

-
- Uracil: Synthesis, Hybridization Studies, Fluorescent Properties and i-Motif Stability. *ChemBioChem*, 2014, **15** (1), 146-156; (d) B. Xu, G. Devi and F. Shao, Regulation of Telomeric i-Motif Stability by 5-Methylcytosine and 5-Hydroxymethylcytosine Modification. *Org. Biomol. Chem.*, 2015, **13** (20), 5646-5651; (d) L. Lannes, S. Halder, Y. Krishnan and H. Schwalbe, Tuning the pH Response of i-Motif DNA Oligonucleotides. *ChemBioChem*, 2015, **16** (11), 1647-1656; (e) F. Seela, S. Budow and P. Leonard, Oligonucleotides Forming an i-Motif: the pH-Dependent Assembly of Individual Strands and Branched Structures Containing 2'-Deoxy-5-propynylcytidine. *Org. Biomol. Chem.*, 2007, **5** (12), 1858-1872; (f) B. Yang and M. T. Rodgers, Base-Pairing Energies of Proton-Bound Heterodimers of Cytosine and Modified Cytosines: Implications for the Stability of DNA i-Motif Conformations. *J. Amer. Chem. Soc.*, 2014, **136** (1), 282-290.
- 18 (a) J.-L. Mergny and L. Lacroix, Kinetics and Thermodynamics of I-DNA Formation: Phosphodiester versus Modified Oligodeoxynucleotides. *Nucleic Acids Res.*, 1998, **26** (21), 4797-4803; (b) J. A. Brazier, J. Fisher and R. Cosstick, Stabilization of the DNA I-Motif Structure by Incorporation of 3'-S-Phosphorothiolate Linkages. *Angew. Chem. Int. Ed.*, 2006, **45** (1), 114-117.
- 19 (a) P. Murat, D. Cressend, N. Spinelli, A. Van der Heyden, P. Labbe, P. Dumy and E. Defrancq, A Novel Conformationally Constrained Parallel G Quadruplex, *ChemBioChem*, 2008, **9** (16), 2588-2591; (b) V. Romanucci, M. Gaglione, A. Messere, N. Potenza, A. Zarrelli, S. Noppen, S. Liekens, J. Balzarini and G. Di Fabio, Hairpin Oligonucleotides forming G-Quadruplexes: New Aptamers with anti-HIV Activity, *Eur. J. Med. Chem.*, 2015, **89**, 51-58; (c) M. Nikan, B. O. Patrick and J. C. Sherman, Crystal Structure of a Template-Assembled Synthetic G-Quadruplex, *ChemBioChem*, 2012, **13** (10), 1413-1415; (d) R. Ferreira, M. Alvira, A. Avino, I. Gomez-Pinto, C. Gonzalez, V. Gabelica and R. Eritja, Synthesis and Structural Characterization of Stable Branched DNA G-Quadruplexes using the Trebler Phosphoramidite, *ChemistryOpen*, 2012, **1**, 106-114; (e) G. Oliviero, N. Borbone, J. Amato, S. D'Errico, A. Galeone, G. Piccialli, M. Varra and L. Mayol, Synthesis of Quadruplex-Forming Tetra-end-Linked Oligonucleotides: Effects of the Linker Size on Quadruplex Topology and Stability, *Biopolymers*, 2009, **91** (6), 466-477.
- 20 L. Bonnat, J. Dejeu, H. Bonnet, B. Gennaro, O. Jarjayes, F. Thomas, T. Lavergne and E. Defrancq, Templated Formation of Discrete RNA and DNA: RNA Hybrid G-Quadruplexes and Their Interactions with Targeting Ligands, *Chem. Eur. J.*, 2016, **22** (9), 3139-3147.
- 21 R. Bonnet, T. Lavergne, B. Gennaro, N. Spinelli and E. Defrancq, Construction of anti-Parallel G-Quadruplexes through Sequential Templated Click, *Chem. Commun.*, 2015, **51** (23), 4850-4853.

-
- 22 R. Bonnet, P. Murat, N. Spinelli and E. Defrancq, Click–click Chemistry on a Peptidic Scaffold for easy Access to Tetrameric DNA Structures, *Chem. Commun.*, 2012, **48** (48), 5992-5994.
- 23 L. Bonnat, L. Bar, B. Gennaro, H. Bonnet, O. Jarjayes, F. Thomas, J. Dejeu, E. Defrancq and T. Lavergne, Template-Mediated Stabilization of a DNA G-Quadruplex formed in the HIV-1 Promoter and Comparative Binding Studies, *Chem. Eur. J.*, **2017**, *23* (23), 5602-5613.
- 24 (a) P. Murat, R. Bonnet, A. Van der Heyden, N. Spinelli, P. Labbe, D. Monchaud, M. P. Teulade-Fichou, P. Dumy and E. Defrancq, Template-Assembled Synthetic G-Quadruplex (TASQ): a Useful System for Investigating the Interactions of Ligands with Constrained Quadruplex Topologies, *Chem. Eur. J.*, 2010, **16** (20), 6106-6114; (b) G. Piraux, L. Bar, M. Abraham, T. Lavergne, H. Jamet, J. Dejeu, L. Marcelis, E. Defrancq and B. Elias, New Ruthenium-Based Probes for Selective G-Quadruplex Targeting, *Chem. Eur. J.*, 2017, **23** (49), 11872-11880; (c) L. Lecarme, E. Prado, A. De Rache, M. L. Nicolau-Travers, R. Bonnet, A. van Der Heyden, C. Philouze, D. Gomez, J. L. Mergny, H. Jamet, E. Defrancq, O. Jarjayes and F. Thomas, Interaction of Polycationic Ni(II)-Salophen Complexes with G-Quadruplex DNA, *Inorg. Chem.*, 2014, **53** (23), 12519-12531.
- 25 (a) O. P. Edupuganti, Y. Singh, E. Defrancq and Pascal Dumy, New Strategy for the Synthesis of 3',5'-Bifunctionalized Oligonucleotide Conjugates through Sequential Formation of Chemoselective Oxime Bonds, *Chem. Eur. J.*, 2004, **10** (23), 5988-5995; (b) O. P. Edupuganti, O. Renaudet, E. Defrancq and P. Dumy, The Oxime Bond Formation as an Efficient Chemical Tool for the Preparation of 3',5'-Bifunctionalised Oligodeoxyribonucleotides, *Bioorg. Med. Chem. Lett.*, 2004, **14** (11), 2839-2842.
- 26 (a) P. M. E. Gramlich, S. Warncke, J. Gierlich and T. Carell, Click-click-click: Single to Triple Modification of DNA, *Angew. Chem. Int. Ed.* 2008, **47** (18), 3442-3444; (b) S. De Tito, F. Morvan, A. Meyer, J. J. Vasseur, A. Cummaro, L. Petraccone, B. Pagano, E. Novellino, A. Randazzo, C. Giancola and D. Montesarchio, Fluorescence Enhancement upon G-Quadruplex Folding : Synthesis, Structure, and Biophysical Characterization of a Dansyl/Cyclodextrin-Tagged Thrombin Binding Aptamer, *Bioconjugate Chem.* 2013, **24** (11), 1917-1927.
- 27 R. Lartia, P. Murat, P. Dumy and Eric Defrancq, Versatile Introduction of Azido Moiety into Oligonucleotides through Diazo Transfer Reaction, *Org. Lett.*, 2011, **13** (20), 5672-5675.
- 28 M. Galibert, P. Dumy and D. Boturyn, One-Pot Approach to Well-Defined Biomolecular Assemblies by Orthogonal Chemoselective Ligations. *Angew. Chem. Int. Ed.* 2009, **48** (14), 2576-2579.

-
- 29 B. A. Connolly and P. Rider, Chemical Synthesis of Oligonucleotides Containing a Free Sulphydryl Group and Subsequent Attachment of Thiol Specific Probes. *Nucleic Acids Res.*, 1985, **13** (12), 4485-4502.
- 30 J. Kypr, I. Kejnovska, D. Renciuik and M. Vorlickova, Circular Dichroism and Conformational Polymorphism of DNA, *Nucleic Acids Res.*, 2009, **37** (6), 1713-1725.
- 31 E. D. Goddard-Borger and R. V. Stick, An efficient, Inexpensive, and Shelf-Stable Diazotransfer Reagent: Imidazole-1-sulfonyl zide Hydrochloride. *Org. Lett.* 2007, **9**, 3797-3800.
- 32 S. Foillard, M. O. Rasmussen, J. Razkin, D. Boturyn and P. Dumy, 1-Ethoxyethylidene, a New Group for the Stepwise SPPS of Aminoxyacetic Acid Containing Peptides. *J. Org. Chem.* 2008, **73** (3), 983-991.