



HAL
open science

Hitting topological minor models in planar graphs is fixed parameter tractable

Petr A. Golovach, Giannos Stamoulis, Dimitrios M. Thilikos

► **To cite this version:**

Petr A. Golovach, Giannos Stamoulis, Dimitrios M. Thilikos. Hitting topological minor models in planar graphs is fixed parameter tractable. SODA 2020 - 31st ACM/SIAM Symposium on Discrete Algorithms, Jan 2020, Salt Lake City, UT, United States. pp.931-950, 10.1137/1.9781611975994.56 . hal-03003167

HAL Id: hal-03003167

<https://hal.science/hal-03003167v1>

Submitted on 20 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hitting Topological Minor Models in Planar Graphs is Fixed Parameter Tractable¹

Petr A. Golovach^{2,3,4} Giannos Stamoulis^{5,6} Dimitrios M. Thilikos^{4,7,8}

Abstract

For a finite collection of graphs \mathcal{F} , the \mathcal{F} -TM-DELETION problem has as input an n -vertex graph G and an integer k and asks whether there exists a set $S \subseteq V(G)$ with $|S| \leq k$ such that $G \setminus S$ does not contain any of the graphs in \mathcal{F} as a topological minor. We prove that for every such \mathcal{F} , \mathcal{F} -TM-DELETION is fixed parameter tractable on planar graphs. In particular, we provide an $f(h, k) \cdot n^2$ algorithm where h is an upper bound to the vertices of the graphs in \mathcal{F} .

Keywords: Topological minors, irrelevant vertex technique, treewidth, vertex deletion problems

¹Emails: Petr.Golovach@uib.no, giannos95@gmail.com, sedthilk@thilikos.info.

²Department of Informatics, University of Bergen, Norway.

³Supported by the Research Council of Norway via the projects “CLASSIS” and “MULTIVAL”.

⁴Supported by the Research Council of Norway and the French Ministry of Europe and Foreign Affairs, via the Franco-Norwegian project PHC AURORA 2019.

⁵Department of Mathematics, National and Kapodistrian University of Athens, Athens, Greece.

⁶Inter-university Postgraduate Programme “Algorithms, Logic, and Discrete Mathematics” (ALMA).

⁷AIGCo project-team, LIRMM, CNRS, Université de Montpellier, Montpellier, France.

⁸Supported by projects DEMOGRAPH (ANR-16-CE40-0028) and ESIGMA (ANR-17-CE23-0010).

Contents

1	Introduction	1
1.1	The \mathcal{P} -deletion problem and its variants	1
1.2	Our contribution.	2
1.3	High level description of our algorithm	2
2	Definitions and preliminaries	5
2.1	Basic concepts on Graphs	6
2.2	Railed annuli	7
2.3	Rerouting topological minors	8
2.4	Boundaried graphs and folios	9
3	The two main subroutines of the algorithm	10
3.1	Reducing the solution space	11
3.2	Finding an irrelevant area	13
4	Proof of the main result	17
5	Proof of the Model Taming theorem	18
5.1	Linkages in railed annuli	18
5.2	Taming a Linkage	19
5.3	Rerouting a linkage	26
6	Conclusions	30

1 Introduction

1.1 The \mathcal{P} -deletion problem and its variants

In general, a \mathcal{P} -DELETION problem is determined by some graph class \mathcal{P} and asks, given an n -vertex graph G and an integer k , whether G can be transformed to a graph in \mathcal{P} after the deletion of k vertices. In other words, the class \mathcal{P} represents some desired property that we want to impose to the input graph after deleting k vertices. This is a general graph modification problem with great expressive power as it encompasses many problems, depending on the choice of the property \mathcal{P} . Unfortunately for most instantiations of \mathcal{P} , this problem is not expected to admit a polynomial time algorithm. Lewis and Yannakakis showed in [21] that for any non-trivial and hereditary graph class \mathcal{P} , the \mathcal{P} -vertex deletion problem is NP-complete. Given this hardness result, an attractive alternative is to consider the standard parameterized version of the problem, called p - \mathcal{P} -DELETION where the parameter is the number k of vertex deletions. In this case the challenge is to investigate for which instantiations of \mathcal{P} , p - \mathcal{P} -DELETION is fixed parameter tractable (or, in short, is FPT), i.e., it can be solved by an $O_k(n^c)$ -time algorithm¹ (or FPT-algorithm), for some constant c . There is a long line of research on this general question. In many cases, this concerns particular properties and possible optimizations of the contribution of k in the function hidden in the “ O_k ” notation (see e.g. [3]). However, it is interesting to notice that FPT-algorithms exist for general families of properties. In this direction the more general (and compact) results concern properties \mathcal{P} that can be characterized by the exclusion of some finite set \mathcal{F} of graphs (i.e., of size bounded by some constant h) with respect to some partial ordering relation \leq . We define

$$\mathcal{P}_{\mathcal{F},\leq} = \{G \mid \forall H \in \mathcal{F} H \not\leq G\}$$

and ask whether p - $\mathcal{P}_{\mathcal{F},\leq}$ -DELETION is FPT. Let us now consider the general status of this problem for the main known instantiations of the partial ordering relation \leq .

(1) \leq is the *contraction*² relation: then there are graphs H such that $\mathcal{P}_{\{H\},\leq}$ -DELETION is NP-complete even for the case where $k = 0$. For instance one may take H to be the path on 4 vertices, as indicated in [5]. Using the terminology of fixed parameter complexity, this implies that there are choices of \mathcal{F} such that p - $\mathcal{P}_{\mathcal{F},\leq}$ -DELETION is para-NP-complete.

(2) \leq is the *induced minor*³ relation: as on the previous case there are choices of \mathcal{F} such that p - $\mathcal{P}_{\mathcal{F},\leq}$ -DELETION is para-NP-complete. For instance, one may consider \mathcal{F} to contain the graph in [10, Theorem 4.3].

(3) \leq is the *subgraph* or the *induced subgraph* relation: because of the result of Cai in [6], p - $\mathcal{P}_{\mathcal{F},\leq}$ -DELETION is FPT, for every \mathcal{F} . In particular, the result in [6] implies an $O(h^k n^{h+1})$ -time algorithm for both these problems. However, if instead we parameterize $\mathcal{P}_{\mathcal{F},\leq}$ -DELETION by h ,

¹ Let $(x_1, \dots, x_l) \in \mathbb{N}^l$ and $\chi, \psi : \mathbb{N} \rightarrow \mathbb{N}$. We use the notation $\chi(n) = O_{x_1, \dots, x_l}(\psi(n))$ to denote that there exists a computable function $\phi : \mathbb{N}^l \rightarrow \mathbb{N}$ such that $\chi(n) = O(\phi(x_1, \dots, x_l) \cdot \psi(n))$.

² A graph G is a *contraction* of a graph G' if G can be obtained from G' by applying edge contractions.

³ A graph G is an *induced minor* of a graph G' if G can be obtained from some contraction of G' after removing vertices.

then there are instantiations of \mathcal{F} for which the problem is W[1]-hard even for $k = 0$: just take $\mathcal{F} = \{K_h\}$ to generate the p -CLIQUE problem.

(4) \leq is the *minor*⁴ relation: again p - $\mathcal{P}_{\mathcal{F}, \leq}$ -DELETION is FPT, for every \mathcal{F} . To see this, observe that, for every k , the set of YES-instances of this problem is closed under taking of minors. On the other hand, Robertson and Seymour [25] proved that graphs are well-quasi-ordered with respect to the minor relation. These two facts together imply that there is a finite set \mathcal{B}_k (whose size depends on k and h) such that (G, k) is a YES-instance if and only if G contains a graph in \mathcal{B}_k . As minor checking for a graph on c vertices can be done in $O_c(n^3)$ -steps [24], we derive the existence of an $O_{k,h}(n^3)$ -step algorithm.

1.2 Our contribution.

Interestingly, we are not aware of other partial ordering relations where p - $\mathcal{P}_{\mathcal{F}, \leq}$ -DELETION is FPT, for every \mathcal{F} . Among the possible candidates, the most relevant one is the topological minor relation, denoted by \preceq : a graph H is a *topological minor* of a graph G if G contains as a subgraph some subdivision⁵ of H .

In this paper we make a first step on the study of the p - $\mathcal{P}_{\mathcal{F}, \preceq}$ -DELETION problem, also called \mathcal{F} -TM-DELETION, and we conjecture that it is FPT. Unfortunately, there are no known meta-algorithmic results, similar to those of the case of minors, that permit a straightforward resolution of this conjecture, as graphs are not well-quasi-ordered under topological minors. On the positive side, there is an algorithm that checks topological minor containment in $O_h(n^3)$ -time [13] and this result would be a special case of our conjecture for the case where $k = 0$. In this paper we prove that this conjecture is true, when we are restricted to planar graphs. Moreover, we develop results and techniques that may serve as the base of its full resolution.

Given a finite set \mathcal{F} of graphs we use $h(\mathcal{F})$ for the maximum size of a graph in \mathcal{F} . We also write $\mathcal{F} \not\preceq G$ to denote the fact that some of the graphs in \mathcal{F} is a topological minor of G . We define $\mathbf{p}_{\mathcal{F}}(G) = \min\{k \mid \exists S \subseteq V(G) : |S| \leq k \wedge \mathcal{F} \not\preceq G \setminus S\}$. The main result of this paper is the following:

Theorem 1. *There exists an algorithm that given a finite set of graphs \mathcal{F} , a $k \in \mathbb{N}$, and a planar graph G , outputs whether $\mathbf{p}_{\mathcal{F}}(G) \leq k$ in $O_{h,k}(n^2)$ steps, where $h = h(\mathcal{F})$.*

1.3 High level description of our algorithm

Our main approach towards proving [Theorem 1](#) is the application of the so-called *irrelevant vertex technique*. This technique was introduced for the first time by Robertson and Seymour in [24] for the design of an FPT-algorithm for the DISJOINT PATHS problem, parameterized by the number of terminals. Subsequently, it was applied, in diverse ways, for the design of FPT-algorithms for several graph-theoretical problems and is now considered as a powerful technique of parameterized algorithm design [1, 9, 11, 12, 14–19, 22, 23]. We also refer to [8, Chapter 7] for

⁴A graph G is an *minor* of a graph G' if G is the contraction of some subgraph of G' .

⁵A graph G is a *subdivision* of a graph G' if G can be obtained by G' if we replace its edges by paths with the same endpoints.

a high-level overview of the irrelevant vertex technique. The general algorithmic paradigm of the irrelevant vertex technique takes profit of some structural characteristic of the input graph in order to detect, in FPT-time, some vertex, called *irrelevant*, whose removal from G generates an equivalent instance of the problem. By recursing on the produced equivalent instance we end up with a graph where the structural parameter is bounded (by some function of k), a fact that permits the resolution of the problem with other techniques – typically by dynamic programming. In most of the times, this structural parameter is treewidth (see [Section 2](#) for the formal definition) and this is the one that we use in this paper. Towards proving [Theorem 1](#), the application of the irrelevant vertex technique is based to the following theorem.

Theorem 2. *There exists a function $f_1 : \mathbb{N} \rightarrow \mathbb{N}$, and an algorithm with the following specifications:*

Find_Irrelevant_Vertex(k, h, G)

Input: $k, h \in \mathbb{N}_{\geq 0}$ and an n -vertex planar graph G

Output:

1. an (irrelevant) vertex $v \in V(G)$ such that, for every graph class \mathcal{F} where $h(\mathcal{F}) \leq h$, it holds that $\mathbf{p}_{\mathcal{F}}(G) \leq k \iff \mathbf{p}_{\mathcal{F}}(G \setminus v) \leq k$ or
2. a tree decomposition of G of width at most $f_1(h) \cdot k$.

Moreover, this algorithm runs in $O_{k,h}(n)$ steps.

After applying the algorithm of [Theorem 2](#) at most n times, the problem is reduced to instances of bounded treewidth. As topological minor containment can be expressed by a MSOL formula and vertex deletion to some MSOL definable property can also be expressed in MSOL, it follows from the Theorem of Courcelle [7] (see also [2, 4, 28]) that the problem for reduced instances can be solved in $O_{k,h}(n)$ steps. [Theorem 1](#) follows.

In the rest of this section we give an outline on how [Theorem 2](#) is proved. All combinatorial concepts used in this description are presented in an intuitive way; formal definitions can be found in [Section 2](#). Given a tuple of variables $\mathbf{x} = (x_1, \dots, x_q)$ by the term \mathbf{x} -big/small we refer to a quantity that is lower/upper bounded by some function of \mathbf{x} . Alternatively, we use the term \mathbf{x} -many/few that is defined analogously. We work on some embedding of G in the plane.

Walls and annuli. An important combinatorial object is the one of a r -wall, as the one in [Figure 1](#), that can be seen the union of r horizontal paths intersected by r vertical paths. The layers of a wall W are defined as indicated in [Figure 1](#).

We call the outermost layer *perimeter* of the W . Using a result of [11] we know that if the treewidth of a planar graph is (k, h) -big, then G contains a (k, h) -big wall such that the subgraph of G , called *the compass* of W , inside the closed disk defined by the perimeter of W has (k, h) -small treewidth (see [Proposition 1](#)). This additional property will permit us to answer queries expressed by MSOL sentences on subgraphs of the compass of W .

The next step is to detect some more structure in the wall W that is intuitively depicted in the left side of [Figure 2](#). We first distinguish the collection \mathcal{C} of the (k, h) -many outmost layers,




Figure 1: A 17-wall and its 8 layers.

drawn in yellow, and then we consider in the rest of W a packing of (k, h) -many (h) -big walls, drawn in green. This is done in [Lemma 1](#).




Figure 2: Left: The partition of a wall into a yellow annulus and several green subwalls. Right: An example of a $(5, 8)$ -railed annulus depicted in yellow; its inner disk D_5 is depicted in green.

We now work on the “annulus” of the (k, h) -many outer layers of W . For this, it is convenient to see those cycles as “crossed” by a collection \mathcal{P} of disjoint paths called *rails*. We call this system of cycles and rails *railed annulus*, denoted by $\mathcal{A} = (\mathcal{C}, \mathcal{P})$. (see the right side of [Figure 2](#) for an example of a railed annulus with 5 cycles and 8 rails).

Taming topological minor models. Notice that if H is a topological minor of a graph G , then this is materialized by a pair (M, T) where M is a subgraph of G and B is a set of vertices of M , called *branches*, such that all vertices of $V(G) \setminus B$ have degree 2. We say that (M, T) is a *topological minor model* of H in G if a graph isomorphic to H is created after dissolving in M all vertices in $V(G) \setminus B$. For simplicity, assume that $\mathcal{F} = \{H\}$ and recall that $P_{\mathcal{F}}(G) \leq k$ if there is a set $S \subseteq V(G)$, $|S| \leq k$, called from now on *solution set*, that intersects all topological minor models of H in G .

Our next aim is to analyze how topological minor models of H may cross the cycles and the rails of a railed annulus $\mathcal{A} = (\mathcal{C}, \mathcal{P})$. For this we dedicate [Section 5](#) to the proof of a general theorem stating that if the branches of (M, A) are situated outside the annulus and the annulus is (h) -big then it is possible to find an alternative “rail-tamed” model (M', T') of G , whose intersection with the “middle cycle” of \mathcal{A} consists only of (h) -few rail vertices.

We refer to this theorem as the “*model taming theorem*” ([Theorem 3](#)). As it has independent combinatorial interest, we present it in a slightly more general form that will appear useful on further algorithmic applications. The proof of this theorem is technical and it is based on the so-called *unique linkage theorem* by Robertson and Seymour in [26, 27] (also appeared in an alternative form as the *unique-linkage theorem* in [20]).

Representations of topological minor models. Using the model-taming theorem, we can pick a (h) -small collection \mathcal{P}' of the rails of \mathcal{A} for which the following holds: for every topological minor model (M, T) of H that crosses \mathcal{A} , there is a disk Δ bounded by some cycle C of \mathcal{A} and a “tamed” version (M', T') of (M, T) that *represents* (M, T) in the sense that a set of vertices that are “not so close” to C , intersects $M \cap \Delta$ iff the same set intersects $M' \cap \Delta$. From now on we refer to the instantiations of $M' \cap \Delta$ as the *inner tamed models* of \mathcal{A} and we can see them as models representing the “inner part” of all annulus-crossing models.

Reducing the solution space. The next step is to compute, for every cycle C of \mathcal{A} , a set S_C of at (k, h) -many vertices intersecting each possible inner tamed model. This computation can be done in $O_{k,h}(n)$ -time as this question can be expressed in MSOL and concerns subgraphs of the compass of W that has (k, h) -small treewidth. Let Δ_{in} be the disk bounded by the innermost cycle of \mathcal{C} (cycle C_5 in [Figure 2](#)). We then compute $S_{\text{in}} = \Delta_{\text{in}} \cap (\bigcup_{C \in \mathcal{C}} S_C)$ and observe that S_{in} has (k, h) -small size. Based on the fact that the inner tamed models represent the inner part of all models crossing \mathcal{A} and the fact that all these models are intersected by subsets of at most k vertices whose restriction in Δ_{in} is in S_{in} , we prove that if $G \setminus S$ does not contain any topological minor model of H , then the same holds if we restrict S to contains only members of S_{in} ([Lemma 3](#)). This is an important restriction of the solution space of the problem in what concerns its intersection with Δ_{in} . As the (h) -big sub-walls packed inside Δ_{in} are (k, h) -many, there is a sub-wall whose compass can be avoided by all possible solution sets.

Finding an irrelevant vertex. We now fix our attention to the solution-free compass of some (h) -big subwall of W . Once again, we see this wall as a railed annulus \mathcal{A}' and use the model taming theorem in order to represent all ways topological minor models of H can “invade” the compass of W by tamed topological models going through the rails of \mathcal{A}' . This in turn permit us to detect a vertex v of the solution-free compass of W such that if a solution set S intersects a topological minor model that contains v , then it should also intersect some representation of it that avoids v , therefore v is irrelevant ([Lemma 4](#)).

2 Definitions and preliminaries

We denote by \mathbb{N} the set of all non-negative integers. Given an $n \in \mathbb{N}$, we denote by $\mathbb{N}_{\geq n}$ the set containing all integers equal or greater than n . Given two integers x and y we define by $[x, y] = \{x, x + 1, \dots, y - 1, y\}$. Given an $n \in \mathbb{N}_{\geq 1}$, we also define $[n] = \{1, \dots, n\}$. Let U be a set, $r \in \mathbb{N}_{\geq 1}$, and $\mathcal{A} = [A_1, \dots, A_r] \subseteq (2^U)^r$, $\mathcal{B} = [B_1, \dots, B_r] \subseteq (2^U)^r$. We say that $\mathcal{A} \subseteq \mathcal{B}$ if for all $i \in [r]$, $A_i \subseteq B_i$. Also, if $S \subseteq U$ we denote $\mathcal{A} \cap S = [A_1 \cap S, \dots, A_r \cap S]$.

2.1 Basic concepts on Graphs

All graphs in this paper are undirected, finite, and they do not have loops or multiple edges. If $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are graphs, then we denote $G_1 \cap G_2 = (V_1 \cap V_2, E_1 \cap E_2)$ and $G_1 \cup G_2 = (V_1 \cup V_2, E_1 \cup E_2)$. Also, given a graph G and a set $S \subseteq V(G)$, we denote by $G \setminus S$ the graph obtained if we remove from G the vertices in S , along with their incident edges. Given a graph G , we say that the pair (A, B) is a *separation* of G if $A \cup B = V(G)$ and there is no edge in G with one endpoint in $A \setminus B$ and the other in $B \setminus A$. A *path (cycle)* in a graph G is a connected subgraph with all vertices of degree at most (exactly) 2. A path is *trivial* if it has only one vertex and is empty if it is the empty graph (i.e., the graph with empty vertex set).

Partially disk-embedded graphs. A *closed disk* (resp. *open disk*) of Δ is a subset of Δ that is homeomorphic to the set $\{(x, y) \mid x^2 + y^2 \leq 1\}$ (resp. $\{(x, y) \mid x^2 + y^2 < 1\}$). A *disk* of Δ is a closed or an open disk of Δ . We use $\mathbf{bor}(\Delta)$ to denote the boundary of Δ and $\mathbf{int}(\Delta)$ to denote the open disk $\Delta \setminus \mathbf{bor}(\Delta)$. We say that a graph G is *partially disk-embedded in some closed disk* Δ , if there is some subgraph K of G that is embedded in Δ such that $\mathbf{bor}(\Delta)$ is a cycle of K and $(V(G) \cap \Delta, V(G) \setminus \mathbf{int}(\Delta))$ is a separation of G . From now on, we use the term *partially Δ -embedded graph* G to denote that a graph G is partially disk-embedded in some closed disk Δ . We also call the graph K *compass* of the partially Δ -embedded graph G and we always assume that we accompany a partially Δ -embedded graph G together with an embedding of its compass in Δ that is the set $G \cap \Delta$.

Let G be a partially Δ -embedded graph and let $\mathcal{C} = [C_1, \dots, C_r]$, $r \geq 2$, be a collection of vertex-disjoint cycles of the compass of G . We say that the sequence \mathcal{C} is a *Δ -nested sequence of cycles* of G if every C_i is the boundary of an open disk D_i of Δ such that $\Delta \supseteq D_1 \supseteq \dots \supseteq D_r$. From now on, each Δ -nested sequence \mathcal{C} will be accompanied with the sequence $[D_1, \dots, D_r]$ of the corresponding open disks as well as the sequence $[\overline{D}_1, \dots, \overline{D}_r]$ of their closures. Given $x, y \in [r]$ where $x \leq y$, we call the set $\overline{D}_x \setminus D_y$ *(x, y) -annulus* of \mathcal{C} and we denote it by $\mathbf{ann}(\mathcal{C}, x, y)$. Finally, we say that $\mathbf{ann}(\mathcal{C}, 1, r)$ is the *annulus* of \mathcal{C} and we denote it by $\mathbf{ann}(\mathcal{C})$.

Grids and Walls. Let $k, r \in \mathbb{N}$. The $(k \times r)$ -*grid* is the Cartesian product of two paths on k and r vertices respectively. An *elementary r -wall*, for some odd $r \geq 3$, is the graph obtained from a $(2r \times r)$ -grid with vertices (x, y) , $x \in [2r] \times [r]$, after the removal of the “vertical” edges $\{(x, y), (x, y + 1)\}$ for odd $x + y$, and then the removal of all vertices of degree one. Notice that, as $r \geq 3$, an elementary r -wall is a planar graph that has a unique (up to topological isomorphism) embedding in the plane such that all its finite faces are incident to exactly six edges. The *perimeter* of an elementary r -wall is the cycle bounding its infinite face. Given an elementary wall \overline{W} , a *vertical path* of \overline{W} is one whose vertices, in ordering of appearance, are $(i, 1), (i, 2), (i + 1, 2), (i + 1, 3), (i, 3), (i, 4), (i + 1, 4), (i + 1, 5), (i, 5), \dots, (i, r - 2), (i, r - 1), (i + 1, r - 1), (i + 1, r)$, for some $i \in \{1, 3, \dots, 2r - 1\}$. Also an *horizontal path* of \overline{W} is the one whose vertices, in ordering of appearance, are $(1, j), (2, j), \dots, (2r, j)$, for some $j \in [2, r - 1]$, or $(1, 1), (2, 1), \dots, (2r - 1, 1)$ or $(2, r), (2, r), \dots, (2r, r)$. (see Figure 1).

An *r -wall* is any graph W obtained from an elementary r -wall \overline{W} after subdividing edges. We call the vertices that were added after the subdivision operations *subdivision vertices*. The

perimeter of W is the cycle of W whose non-subdivision vertices are the vertices of the perimeter of \overline{W} . Also, a vertical (resp. horizontal) path of W is a subdivided vertical (resp. horizontal) path of \overline{W} . An r' -*subwall* W' of a wall W is any r' -wall that is a subgraph of W and whose horizontal/vertical paths are subpaths of the horizontal/vertical paths of W .

A subgraph W of a graph G is called a *wall* of G if W is an r -wall for some odd $r \geq 3$ and we refer to r as the *height* of the wall W . The *layers* of an r -wall W are recursively defined as follows. The first layer of W is its perimeter. For $i = 2, \dots, (r-1)/2$, the i -th layer of W is the $(i-1)$ -th layer of the subwall W' obtained from W after removing from W its perimeter and all occurring vertices of degree one. Notice that each $(2r+1)$ -wall has r layers.

Treewidth. Given a $k \in \mathbb{N}_{\geq 0}$, we say that a graph G is a k -*tree* if G is isomorphic to K_{k+1} or (recursively) there is a vertex v in G where $N_G[\{v\}]$ isomorphic to K_{k+1} and $G \setminus \{v\}$ is a k -tree. The *treewidth* of a G is the minimum k for which G is a subgraph of some k -tree.

The following result from [11] intuitively states that given a $q \in \mathbb{N}$ and a graph G with “big” enough treewidth, we can find a q -wall of G whose compass has “small” enough treewidth.

Proposition 1 ([11]). *There exists a constant c_1 and an algorithm with the following specifications:*

Find_Wall(G, q)

Input: a planar graph G and a $q \in \mathbb{N}$.

Output:

1. *A q -wall W of G whose compass has treewidth at most $c_1 \cdot q$ or*
2. *A tree decomposition of G of width at most $c_1 \cdot q$.*

Moreover, this algorithm runs in $O_q(n)$ steps.

2.2 Railed annuli

Let $r \in \mathbb{N}_{\geq 3}$ and $q \in \mathbb{N}_{\geq 3}$. Assume also that r is an odd number. An (r, q) -*railed annulus* of a partially Δ -embedded graph G is a pair $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ where $\mathcal{C} = [C_1, \dots, C_r]$ is a Δ -nested collection of cycles of G and $\mathcal{P} = [P_1, \dots, P_q]$ is a collection of pairwise vertex-disjoint paths in G such that

- For every $j \in [q]$, $P_j \subseteq \text{ann}(\mathcal{C})$.
- For every $(i, j) \in [r] \times [q]$, $C_i \cap P_j$ is a non-empty path, that we denote $P_{i,j}$.

We refer to the paths of \mathcal{P} as the *rails* of \mathcal{A} and to the cycles of \mathcal{C} as the *cycles* of \mathcal{A} .

Let $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ be an (r, q) -railed annulus of a partially Δ -embedded graph G . We call \overline{D}_r (resp. \overline{D}_1) the *inner* (resp. *outer*) *disk* of \mathcal{A} . We also extend the notion of an annulus and we say that the *annulus* of $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ is the annulus of \mathcal{C} .

We now prove the following lemma which intuitively states that there is an algorithm that given a “big enough” wall, it outputs a collection of railed annuli whose number and size will be useful in the proof of **Theorem 2**.

Lemma 1. *There exists a function $f_2 : \mathbb{N}^3 \rightarrow \mathbb{N}$ and an algorithm with the following specifications:*

Find_Collection_of_Annuli(x, y, z, Δ, G, W)

Input: two odd integers $x, y \in \mathbb{N}_{\geq 3}$, an integer $z \in \mathbb{N}$, a partially Δ -embedded graph G and a q -wall W of the compass of G whose perimeter is the boundary of Δ and such that $q \geq f_2(x, y, z)$.

Output: a collection $\mathfrak{A} = \{\mathcal{A}_0, \mathcal{A}_1, \dots, \mathcal{A}_z\}$ of railed annuli of the compass of G such that

- \mathcal{A}_0 is a (x, x) -railed annulus whose outer disk is Δ and whose inner disk is Δ' ,
- for $i \in [z]$, \mathcal{A}_i is a (y, y) -railed annulus of $G \cap \Delta'$, and
- for every $i, j \in [z]$ where $i \neq j$, the outer disk of \mathcal{A}_i and the outer disk of \mathcal{A}_j are disjoint.

Moreover, this algorithm runs in $O(n)$ steps and $f_2(x, y, z) = O(x + y\sqrt{z})$.

Proof. Let $y' := y + \lceil (y - 2)/4 \rceil$ and assume that y' is an odd integer (otherwise, make it odd by adding 1) and let $f_2(x, y, z) = x + \max\{\lceil (x - 2)/4 \rceil, \lceil \sqrt{z}/2 \rceil \cdot y'\} + 1$. We argue that the following holds:

Claim: Let $p \in \mathbb{Z}_{\geq 3}$. If H is an h -wall of G , where $h \geq p + \lceil (p - 2)/4 \rceil$, then H contains a (p, p) -railed annulus $\mathcal{A} = (\mathcal{C}, \mathcal{P})$, where $\mathcal{C} = [C_1, \dots, C_p]$ and for every $i \in [p]$, C_i is the i -th layer of H .

Proof of Claim: Let H be an h -wall of G , where $h \geq p + \lceil (p - 2)/4 \rceil$. We define the Δ -nested collection $\mathcal{C} = [C_1, \dots, C_p]$ of cycles of G , where, for every $i \in [p]$, C_i is the i -th layer of H . Let $\hat{\mathcal{P}}$ be the collection of the vertical and horizontal paths of H that intersect C_p . Observe that for every $i \in [p - 1]$, every path in $\hat{\mathcal{P}}$ also intersects C_i and that $\hat{\mathcal{P}} \cap \text{ann}(\mathcal{C})$ is a collection of pairwise-vertex disjoint paths of G . Also, notice that since $h - p \geq \lceil (p - 2)/4 \rceil$ then $\hat{\mathcal{P}} \cap \text{ann}(\mathcal{C})$ contains at least p paths. Let $\mathcal{P} := [P_1, \dots, P_p]$ be a subset of $\hat{\mathcal{P}} \cap \text{ann}(\mathcal{C})$. Then, \mathcal{P} is a collection of pairwise vertex-disjoint paths of G and it holds that for every $j \in [p]$, $P_j \subseteq \text{ann}(\mathcal{C})$ and for every $(i, j) \in [p] \times [p]$, $C_i \cap P_j$ is a non-empty path. Therefore, H contains a (p, p) -railed annulus $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ of G and the claim follows.

Following the claim above, for $H := W$, $h := q$, and $p := x$, since $q \geq x + \lceil (x - 2)/4 \rceil$, we deduce the existence of a (x, x) -railed annulus \mathcal{A}_0 whose inner disk is \overline{D}_x and whose outer disk is \overline{D}_1 - that is Δ . Observe that since $q - x \geq \lceil \sqrt{z}/2 \rceil \cdot y' + 1$, then there exists an r -wall \hat{W} of G for some odd $r \in \mathbb{Z}_{\geq 3}$ such that $r \geq \lceil \sqrt{z}/2 \rceil \cdot y'$ and $\hat{W} \subseteq G \cap \overline{D}_x$.

Now, notice that \hat{W} contains a collection $\mathcal{W} = \{W'_1, \dots, W'_z\}$ of z y' -subwalls of W such that, for every $i, j \in [z], i \neq j$, $K(W'_i) \cap K(W'_j) = \emptyset$. Therefore, for every $i \in [z]$, applying again the claim above for $H := W'_i$, $h := y'$ and $p := y$, we deduce the existence of a (y, y) -railed annulus \mathcal{A}_i of W'_i . Furthermore, for every $i, j \in [z], i \neq j$, recall that $K(W'_i) \cap K(W'_j) = \emptyset$ which implies that the outer disk of \mathcal{A}_i and the outer disk of \mathcal{A}_j are disjoint. The proof concludes by setting $\mathfrak{A} = \{\mathcal{A}_0, \mathcal{A}_1, \dots, \mathcal{A}_z\}$. \square

2.3 Rerouting topological minors

We say that (M, T) is a *tm-pair* if M is a graph, $T \subseteq V(M)$, and all vertices in $V(M) \setminus T$ have degree two. We denote by $\text{diss}(M, T)$ the graph obtained from M by dissolving all vertices in

$V(M) \setminus T$. A *tm-pair* of a graph G is a *tm-pair* (M, T) where M is a subgraph of G . Given two graphs H and G , we say that a *tm-pair* (M, T) of G , is a *topological minor model of H in G* if H is isomorphic to $\text{diss}(M, T)$. We call the vertices in T *branch vertices* of (M, T) .

Topological minor models in railed annuli. Let G be a partially Δ -embedded graph, let H be a graph, $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ be a (r, q) -railed annulus of G . Let $r = 2t + 1$. Let also $s \in [r]$ where $s = 2t' + 1$. Given some $I \subseteq [q]$, we say that a topological minor model (M, T) of H in G is (s, I) -*confined in \mathcal{A}* if

$$M \cap \text{ann}(\mathcal{C}, t - t', t + t') \subseteq \bigcup_{i \in I} P_i.$$

Intuitively, the above definition demands that M traverses the “middle” (s, q) -annulus by intersecting it only the rails of \mathcal{A} .

Our algorithms are based on the following combinatorial result whose proof is postponed in [Section 5](#).

Theorem 3 (Model Taming). *There exist two functions $f_3, f_4 : \mathbb{N}_{\geq 0} \rightarrow \mathbb{N}_{\geq 0}$ such that if*

- s is a positive odd integer,
- H is a graph on g edges,
- G is a partially Δ -embedded graph,
- $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ is a (r, q) -railed annulus of G , where $r \geq f_4(g) + 2 + s$ and $q \geq 5/2 \cdot f_3(g)$,
- (M, T) is a topological minor model of H in G such that $T \cap \text{ann}(\mathcal{A}) = \emptyset$, and
- $I \subseteq [q]$ where $|I| > f_3(g)$,

then G contains an topological minor model (\tilde{M}, \tilde{T}) of H in G such that

1. $\tilde{T} = T$,
2. \tilde{M} is (s, I) -confined in \mathcal{A} and
3. $\tilde{M} \setminus \text{ann}(\mathcal{A}) \subseteq M \setminus \text{ann}(\mathcal{A})$.

Apart from being the combinatorial base of our results, the Model Taming theorem will appear useful in future results using the irrelevant vertex technique.

2.4 Boundaried graphs and folios

Let $t \in \mathbb{N}$. A t -*boundaried graph* is a triple $\mathbf{G} = (G, B, \rho)$ where G is a graph, $B \subseteq V(G)$, $|B| \leq t$, and $\rho : B \rightarrow [t]$ is an injective function. We call B the *boundary* of \mathbf{G} and we call the vertices of B the *boundary vertices* of \mathbf{G} . We also call G the *underlying graph* of \mathbf{G} . Moreover, we call $|B|$ the *boundary size* of \mathbf{G} . We say that the t -boundaried $\mathbf{G}' = (G', B', \rho')$ is a *subgraph* of \mathbf{G} if G' is a subgraph of G , $B' = B \cap V(G')$, and $\rho' = \rho|_{B'}$. Also, for $S \subseteq V(G)$, we define

$\mathbf{G} \setminus S$ to be the t -boundaried graph (G', B', ρ') where $G' = G \setminus S$, $B' = B \setminus S$ and $\rho' = \rho|_{B'}$. Two t -boundaried graphs $\mathbf{G}_1 = (G_1, B_1, \rho_1)$ and $\mathbf{G}_2 = (G_2, B_2, \rho_2)$ are isomorphic if G_1 is isomorphic to G_2 via a bijection $\phi : V(G_1) \rightarrow V(G_2)$ such that $\rho_1 = \rho_2 \circ \phi|_{B_1}$, i.e., the vertices of B_1 are mapped via ϕ to equally indexed vertices of B_2 . A *boundaried graph* is any t -boundaried graph for some $t \in \mathbb{N}$.

We also define the *treewidth* of a boundaried graph $\mathbf{G} = (G, B, \rho)$, denoted by $\mathbf{tw}(\mathbf{G})$ as the minimum width of a tree decomposition (T, χ) of G for which there is some $t \in V(T)$ such that $B \subseteq \chi(t)$. Notice that the treewidth of a t -boundaried graph is always lower bounded by its boundary size.

Topological minors of boundaried graphs. If $\mathbf{M} = (M, B, \rho)$ is a boundaried graph and $T \subseteq V(M)$ with $B \subseteq T$, we call (\mathbf{M}, T) a *btm-pair* and we define $\mathbf{diss}(\mathbf{M}, T) = (\mathbf{diss}(M, T), B, \rho)$ (notice that we consider all boundary vertices to be branch vertices, therefore we do not permit their dissolution). If $\mathbf{G} = (G, B, \rho)$ is a boundaried graph and (M, T) is a *tm-pair* of G where $B \subseteq T$, then we say that (\mathbf{M}, T) , where $\mathbf{M} = (M, B, \rho)$, is a *btm-pair* of $\mathbf{G} = (G, B, \rho)$. Let $\mathbf{G}_i = (G_i, B_i, \rho_i), i \in [2]$. We say that \mathbf{G}_1 is a *topological minor* of \mathbf{G}_2 , denoted by $\mathbf{G}_1 \preceq \mathbf{G}_2$, if there is a *btm-pair* (\mathbf{M}, T) of \mathbf{G}_2 such that $\mathbf{diss}(\mathbf{M}, T)$ is isomorphic to \mathbf{G}_1 . We call $\mathbf{diss}(\mathbf{M}, T)$ *representation* of the *btm-pair* (\mathbf{M}, T) of \mathbf{G} .

Folios. Let $h, t \in \mathbb{N}$ where $h \geq t$. We denote by $\mathcal{B}_h^{(t)}$ the set of all (pairwise non-isomorphic) t -boundaried graphs with at most h vertices. We set the function $f_5 : \mathbb{N}^2 \rightarrow \mathbb{N}$ such that $f_5(t, h) = |\mathcal{B}_h^{(t)}|$. Given a t -boundaried graph \mathbf{G} and an integer $h \in \mathbb{N}$, we define the *h-folio* of \mathbf{G} , denoted by $\mathcal{F}_h^{(t)}(\mathbf{G})$, as the set containing all t -boundaried graphs in $\mathcal{B}_h^{(t)}$ that are representations of the *btm-pairs* of \mathbf{G} . Notice that $|\mathcal{F}_h^{(t)}(\mathbf{G})| \leq f_5(t, h)$.

Given that topological minor containment can be expressed in Monadic Second Order logic, the next lemma follows from Courcelle's theorem.

Lemma 2. *There is an algorithm with the following specifications:*

Compute_Folio (h, w, t, \mathbf{G})

Input: $h, w, t \in \mathbb{N}$, where $h \geq t$ and a t -boundaried graph \mathbf{G} of treewidth at most w .

Output: the set $\mathcal{F}_h^{(t)}(\mathbf{G})$.

Moreover, this algorithm runs in $O_{h,w}(n)$ steps.

3 The two main subroutines of the algorithm

In this section, we provide two main subroutines that will be useful in the proof of [Theorem 2](#). For now on, functions f_3, f_4 will always denote the functions of [Theorem 3](#).

Boundaried graphs in railed annuli. Let $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ be a (r, q) -railed annulus of a partially Δ -embedded graph G . We can see each path P_j in \mathcal{P} as being oriented towards the ‘‘inner’’ part of Δ , i.e., starting from an endpoint of $P_{1,j}$ and finishing to an endpoint of $P_{r,j}$. For every $(i, j) \in [r] \times [q]$, we define $r_{i,j}$ as the first vertex of P_j that appears in $P_{i,j}$ while traversing P_j according to this orientation.




Figure 3: Visualization of the partition of the cycles of \mathcal{A} in sets $A_i, i \in [k+1]$ and of the partition in sets $B_{i,j}, i, j \in [k+1] \times [h+1]$.

Given an $i \in [r]$ and a $t \in [q]$, we define the t -boundaried graph $\mathbf{G}_{i,t} = (G_i, R_{i,t}, \rho_{i,t})$ where $G_i = G \cap \overline{D}_i$, $R_{i,t} = \{r_{i,1} \dots, r_{i,t}\}$ and, for $j \in [t]$, $\rho(t_{i,j}) = j$.

3.1 Reducing the solution space

We now prove the following lemma that intuitively states that there is an algorithm that given a graph G and a “big enough” railed annulus \mathcal{A} of G , it “reduces” the set of vertices that are candidates to the hitting set S .

Lemma 3. *There exists an algorithm with the following specifications:*

Reduce_Solution_Space $(k, h, g, \Delta, G, w, \mathcal{C}, \mathcal{P})$

Input: three integers $k, h, g \in \mathbb{N}_{\geq 0}$, a partially Δ -embedded graph G whose compass has treewidth $\leq w$ and an (r, q) -railed annulus $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ of G , where $r = (k+1)(h+1)(f_4(g) + 3)$ and $q \geq 5/2 \cdot f_3(g)$.

Output: a set $R \subseteq D_r \cap V(G)$ such that

- $|R| \leq f_5(f_3(g) + 1, h + f_3(g) + 1)^{h+1} \cdot k(k+1)$ and
- for every h -vertex and g -edge graph H and every $S \subseteq V(G)$, if $|S| \leq k$ and $H \not\subseteq G \setminus S$, then there is some $S' \subseteq (V(G) \setminus D_r) \cup R$ such that $|S'| \leq k$ and $H \not\subseteq G \setminus S'$.

Moreover, this algorithm runs in $O_{h,w}(n)$ steps.

Proof. We set $\mu := f_4(g) + 3$ and $\lambda := f_3(g)$. Given an $i \in [k+1]$ we use notation A_i as a shortcut to $\text{ann}(\mathcal{C}, (i-1)(h+1)\mu + 1, i(h+1)\mu)$ and for every $j \in [h+1]$ we define $B_{i,j} = \text{ann}(\mathcal{C}, (i-1)(h+1)\mu + (j-1)\mu + 1, (i-1)(h+1)\mu + j\mu)$. Intuitively, we partition \mathcal{C} in $k+1$ sets of consecutive cycles (i.e., the cycles of $A_i, i \in [k+1]$) and then, for every $i \in [k+1]$ we partition the set of cycles of A_i into $h+1$ sets of consecutive cycles (i.e., the cycles of $B_{i,j}, j \in [h+1]$). Notice that for every $i, j \in [k+1] \times [h+1]$, $|B_{i,j} \cap \mathcal{C}| = \mu$.

Also, we define for every $(i, j) \in [k + 1] \times [h + 1]$ the $(\lambda + 1)$ -boundaried graph $\hat{\mathbf{G}}_{i,j} = \mathbf{G}_{(i-1)(h+1)\mu+(j-1)\mu+\lceil\mu/2\rceil,\lambda+1}$ (to get some intuition, notice that the boundary vertices of $\hat{\mathbf{G}}_{i,j}$ lie on the “middle” cycle of $B_{i,j}$ – see [Figure 3](#)). Let $i \in [k + 1]$ and for every $j \in [h + 1]$, let $\mathcal{F}_j := \mathcal{F}_{h+\lambda+1}^{(\lambda+1)}(\hat{\mathbf{G}}_{i,j})$. Then we define the collection $\mathfrak{F}_i = [\mathcal{F}_1, \dots, \mathcal{F}_{h+1}] \in (\mathcal{B}_{h+\lambda+1}^{(\lambda+1)})^{h+1}$. Let S_{i,\mathfrak{F}_i} be a subset of $V(G) \cap D_{(i-1)(h+1)\mu+1}$ of at most k vertices such that, for every $j \in [h + 1]$, it holds that $\mathcal{F}_{h+\lambda+1}^{(\lambda+1)}(\hat{\mathbf{G}}_{i,j} \setminus S) \subseteq \mathcal{F}_j$. If such a set does not exist, then we set $S_{i,\mathfrak{F}_i} = \emptyset$. We define

$$R = \left(\bigcup_{\substack{i \in [k+1] \\ \mathfrak{F}_i \in (\mathcal{B}_{h+\lambda+1}^{(\lambda+1)})^{h+1}}} S_{i,\mathfrak{F}_i} \right) \cap D_r$$

Notice that as each $\hat{\mathbf{G}}_{i,h+1}$ is a subgraph of the compass of G , it has treewidth at most w . Moreover, the set S_{i,\mathfrak{F}_i} can be expressed in MSOL and, again from Courcelle’s theorem, each S_{i,\mathfrak{F}_i} , and therefore R as well, can be computed in $O_{k,g}(|G|)$ steps.

Let H be h -vertex graph and g -edge graph and $S \subseteq V(G)$ such that $|S| \leq k$ and $H \not\leq G \setminus S$. As $r = (k + 1)(h + 1)\mu$ and $|S| \leq k$, then by the pigeonhole’s principle there is some $\ell \in [k + 1]$ such that $S \cap A_\ell = \emptyset$. (In case there are many such ℓ ’s, we take the minimum one.) Let $S_{\text{in}} = S \cap D_{\ell(h+1)\mu}$ and $S_{\text{out}} = S \setminus \bar{D}_{(\ell-1)(h+1)\mu+1}$. Let also $k_{\text{in}} := |S_{\text{in}}|$ and $k_{\text{out}} := |S_{\text{out}}|$ and keep in mind that $k_{\text{in}} + k_{\text{out}} = |S| \leq k$. Let \mathcal{H} be the set of all topological minor models of H in G and notice that for every $(M, T) \in \mathcal{H}$, $S \cap V(M) \neq \emptyset$, i.e., S intersects at least one vertex of each graph in \mathcal{H} . Let \mathcal{H}_ℓ be the members of \mathcal{H} that are intersected *only* by vertices in S_{in} .

The next claim shows that there is a collection of cycles of \mathcal{A} such that for each tm-pair $(M, T) \in \mathcal{H}_\ell$ there exists a cycle C of this collection and a tm-pair $(\tilde{M}, \tilde{T}) \in \mathcal{H}_\ell$ that is equivalent to (M, T) and is “tamed in C ” in the sense that $M \cap C$ is a subgraph of the rails of \mathcal{A} .

Claim: For every $(M, T) \in \mathcal{H}_\ell$, there is an $j_M \in [h + 1]$ and a topological minor model (\tilde{M}, \tilde{T}) of H in $G \setminus S$ such that $\tilde{M} \setminus A_\ell \subseteq M \setminus A_\ell$ and whose intersection with C_{y_M} is the union of the paths $\{P_{y_M, c_1^M}, \dots, P_{y_M, c_{z_M}^M}\}$ where $y_M = (\ell - 1)(h + 1)\mu + (j_M - 1)\mu + \lceil\mu/2\rceil$ and $\{c_1^M, \dots, c_{z_M}^M\} \subseteq [\lambda + 1]$ (see [Figure 4](#)).




Figure 4: Visualization of the statement of the Claim. (M, T) is depicted in the left figure, while (\tilde{M}, \tilde{T}) is depicted in the right figure.

Proof of Claim: Let $(M, T) \in \mathcal{H}_\ell$ and notice that $S_{\text{in}} \cap V(M) \neq \emptyset$. As $|T| = h$, there is some $j_M \in [h + 1]$ such that $T \cap B_{\ell,j_M} = \emptyset$ (if many such j_M ’s exist, take the minimum one). We

use notation $\mathcal{A}^{(M)} = (\mathcal{C}^{(M)}, \mathcal{P}^{(M)})$ instead of $\mathcal{A} \cap B_{\ell, j_M}$. We can now apply [Theorem 3](#) for $s = 1$, $\mathcal{A} := \mathcal{A}^{(M)}$, and $I = [\lambda + 1]$ and obtain a topological minor model (\tilde{M}, \tilde{T}) of H in G such that $\tilde{T} = T$, \tilde{M} is $(1, I)$ -confined in $\mathcal{A}^{(M)}$ and $\tilde{M} \setminus B_{\ell, j_M} \subseteq M \setminus B_{\ell, j_M}$, which implies that $\tilde{M} \setminus A_\ell \subseteq M \setminus A_\ell$. Let $y_M = (\ell - 1)(h + 1)\mu + (i_M - 1)\mu + \lceil \mu/2 \rceil$. Notice that (\tilde{M}, \tilde{T}) is a topological minor model of H in $G \setminus S$ whose intersection with C_{y_M} is the union of some of the paths in $\{P_{y_M, 1}, \dots, P_{y_M, \lambda+1}\}$. Suppose that these paths are $\{P_{y_M, c_1^M}, \dots, P_{y_M, c_{z_M}^M}\}$ where $\{c_1^M, \dots, c_{z_M}^M\} \subseteq [\lambda + 1]$. The claim follows.

Following the above claim, for every $(M, T) \in \mathcal{H}_\ell$ we define the $(\lambda + 1)$ -boundaried graph $\mathbf{G}_M = (G_M, B_M, \rho_M)$ where $G_M = (\tilde{M} \cap \bar{D}_{y_M}) \cup (B_M, \emptyset)$ (i.e. the graph $\tilde{M} \cap \bar{D}_{y_M}$ together with the isolated vertices B_M), $B_M = \{r_{y_M, 1}, \dots, r_{y_M, \lambda+1}\}$, and for $d \in [\lambda + 1]$, $\rho_M(r_{y_M, d}) = d$.

We now define, for every $j \in [h + 1]$, the set $\mathcal{F}_j = \{\text{diss}(\mathbf{G}_M, T \cup B_M) \mid j_M = j \text{ and } (M, T) \in \mathcal{H}_\ell\}$ and we set $\mathfrak{F}_S = [\mathcal{F}_1, \dots, \mathcal{F}_{h+1}]$.

Notice that S_{in} is a subset of $V(G) \cap D_{(\ell-1)(h+1)\mu+1}$ of k_{in} vertices such that, for every $j \in [h + 1]$, it holds that $\mathcal{F}_{h+\lambda+1}^{(\lambda+1)}(\hat{\mathbf{G}}_{\ell, j} \setminus S_{\text{in}}) = \mathcal{F}_j$. Clearly, $|S_{\ell, \mathfrak{F}_S}| \leq S_{\text{in}}$. We now set $S' = S_{\ell, \mathfrak{F}_S} \cup S_{\text{out}}$ and we observe that $S_{\ell, \mathfrak{F}_S} \subseteq R$, $|S'| \leq k$, and that $S' \subseteq (V(G) \setminus D_r) \cup R$. It remains to prove that $H \not\leq G \setminus S'$.

Suppose to the contrary that the graph $G \setminus S'$ contains some topological minor model (M, T) of H as a subgraph. According to the claim above, there is an $j_M \in [h + 1]$ and a topological minor model (\tilde{M}, \tilde{T}) of H in $G \setminus S$ such that $\tilde{M} \setminus A_\ell \subseteq M \setminus A_\ell$ and whose intersection with C_{y_M} is the union of the paths $\{P_{y_M, c_1^M}, \dots, P_{y_M, c_{z_M}^M}\}$ where $y_M = (\ell - 1)(h + 1)\mu + (i_M - 1)\mu + \lceil \mu/2 \rceil + 1$ and $\{c_1^M, \dots, c_{z_M}^M\} \subseteq [\lambda + 1]$. Therefore $\text{diss}(\mathbf{G}_M, T \cup B_M) \in \mathcal{F}_{j_M}$, which contradicts the definition of S_{ℓ, \mathfrak{F}_S} . \square

3.2 Finding an irrelevant area

Before we proceed with the proof of the second result of this section we need some more definitions. Let $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ be an (r, q) -railed annulus of a partially Δ -embedded graph G .

For every $i \in [r]$, we define $F_{\mathcal{A}}^{(i)}$ as the edge set of the unique $(P_{i, q}, P_{i, 1})$ -path that does not contain any vertex from P_2 . We also set $F_{\mathcal{A}} = \bigcup_{i \in [r]} F_{\mathcal{A}}^{(i)}$. Let $(i, j, j') \in [r] \times [q]^2$ where $j \neq j'$. We denote by $L_{i, j \rightarrow j'}$ the shortest path in C_i starting from a vertex of $P_{i, j}$ and finishing to a vertex of $P_{i, j'}$ and that does not contain any edge from $F_{\mathcal{A}}$. Let $(i, i', j) \in [r]^2 \times [q]$ where $i \neq i'$ (see [Figure 5](#)). We denote by $R_{i \rightarrow i', j}$ the shortest path in P_j starting from a vertex of $P_{i, j}$ and finishing to a vertex of $P_{i', j}$. Let $(i, i', j, j') \in [r]^2 \times [q]^2$ such that $i < i'$ and $j < j'$. We define $\Delta_{i, i', j, j'}$ as the closed disk bounded by the unique cycle in the graph

$$P_{i, j} \cup L_{i, j \rightarrow j'} \cup P_{i, j'} \cup R_{i \rightarrow i', j'} \cup P_{i', j'} \cup L_{i', j' \rightarrow j} \cup P_{i', j} \cup R_{i' \rightarrow i, j}.$$

The next lemma intuitively states that there exists an algorithm that given a partially Δ -embedded graph G and a “big enough” railed annulus of G , then there exists a bidimensional area $\Delta' \subseteq \Delta$ such that for every hitting set S outside Δ , $\Delta' \cap V(G)$ is an irrelevant part of the instance.




Figure 5: An example of a $(5,8)$ -railed annulus \mathcal{A} , the set $F_{\mathcal{A}}$ (depicted in green), and the graphs $L_{2,5 \rightarrow 7}$ (depicted in red), $R_{2 \rightarrow 4,1}$ (depicted in yellow), and $\Delta_{3,5,2,5}$ (depicted in blue).

Lemma 4. *There exists an algorithm with the following specifications:*

Find_irrelevant_area $(h, g, b, \Delta, G, w, \mathcal{C}, \mathcal{P})$

Input: three integers $h, g \in \mathbb{N}_{\geq 1}$ and $b \in \mathbb{N}_{\geq 2}$, a partially Δ -embedded graph G whose compass has treewidth at most w , and an (r, q) -railed annulus $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ of G , where $r = f_5(f_3(g) + 1, h + f_3(g) + 1) \cdot ((h + 1)(f_4(g) + 3) + b + 1)$ and $q = \max\{5/2 \cdot f_3(g), f_3(g) + b\}$.

Output: a closed disk $\Delta' \subseteq \Delta$ such that

- $\mathbf{tw}(G \cap \Delta') \geq b$ and
- for every h -vertex and g -edge graph H and for every $S \subseteq V(G) \setminus \Delta$, if $H \not\subseteq (G \setminus (\Delta' \cap V(G))) \setminus S$ then $H \not\subseteq G \setminus S$.

Moreover, this algorithm runs in $O_{h,w}(b \cdot |G|)$ steps.

Proof of Lemma 4. Let $t := f_3(g) + 1$, $\mu := f_4(g) + 3$, $\ell := (h + 1)\mu + b + 1$. Using this notation we have that $r = f_5(t, h + t) \cdot \ell$.

We consider the t -boundaried graphs $\mathbf{G}_{i,t}$, $i \in [r]$. As the underlying graph of each $\mathbf{G}_{i,t}$ is a subgraph of the compass of G , we have that $\mathbf{tw}(\mathbf{G}_{i,t}) \leq w + t = O_h(w)$. For each $i \in [r]$, we call the algorithm **Compute_Folio** $(h + t, t, \mathbf{G}_{i,t}, w + t)$ and compute the $(h + t)$ -folio of $\mathbf{G}_{i,t}$ which, from now on, we denote by \mathcal{F}_i . According to **Lemma 2** \mathcal{F}_i , for all $i \in [r]$ can be computed in $O_{h,w}(r \cdot |G|) = O_{h,w}(b \cdot |G|)$ steps. We observe that if $1 \leq i \leq i' \leq r$, then $\mathcal{F}_{i'} \subseteq \mathcal{F}_i$. This, together with the fact that $|\mathcal{B}_h^{(t)}| = f_5(t, h + t)$, implies that there is an $i' \in [r - \ell + 1]$ such that $\mathcal{F}_{i'} = \mathcal{F}_{i'+1} = \dots = \mathcal{F}_{i'+\ell-1}$. We define

$$\Delta' = \Delta_{i'+\mu(h+1), i'+\ell-2, t+1, t+b}$$

and notice that $G \cap \Delta'$ contains a $(b \times b)$ -grid as a minor, therefore $\mathbf{tw}(G \cap \Delta') \geq b$ (see **Figure 6**). Also keep in mind that Δ' does not intersect the cycle $C_{i'+\ell-1}$.




Figure 6: An example showing the disk Δ' .

Let now H be a h -vertex and g -edge graph and $S \subseteq V(G) \setminus \Delta$ such that $H \not\leq (G \setminus \Delta) \setminus S$. It remains to prove that $H \not\leq G \setminus S$. Suppose to the contrary that the graph $G \setminus S$ contains some topological minor model (M, T) of H as a subgraph. As $|T| = h$ and $\ell = \mu \cdot (h + 1) + b + 1$ there is some $i'' \in [i', i' + \ell - \mu]$ such that $T \cap \text{ann}(\mathcal{C}, i'', i'' + \mu - 1) = \emptyset$.

We consider the (μ, q) -railed annulus $\mathcal{A}' = (\mathcal{C}', \mathcal{P})$ of $G \setminus S$ where

- $\mathcal{C}' = [C'_1, \dots, C'_\mu] := [C_{i''}, \dots, C_{i'' + \mu - 1}]$ and
- $\mathcal{P}' = [P'_1, \dots, P'_q] := [P_1 \cap \text{ann}(\mathcal{A}'), \dots, P_q \cap \text{ann}(\mathcal{A}')]$. (See Figure 7.)




Figure 7: An example showing the (μ, q) -railed annulus \mathcal{A}' .

We are now in position to apply [Theorem 3](#) for $s := 1$, $H, G := G \setminus S$, $\mathcal{A} := \mathcal{A}'$, $r := \mu$, M , and $I = [t]$. We deduce the existence of a topological minor model (\tilde{M}, \tilde{T}) of H in $G \setminus S$ such that $\tilde{T} = T$, \tilde{M} is $(1, I)$ -confined in \mathcal{A}' , and $\tilde{M} \setminus \text{ann}(\mathcal{A}') \subseteq M \setminus \text{ann}(\mathcal{A}')$ (see [Figure 7](#)).




Figure 8: An example of (\tilde{M}, \tilde{T}) , the result of applying [Theorem 3](#) in the railed annulus \mathcal{A}' .

Let $y = i'' + \lfloor \mu/2 \rfloor$. Notice now that $\tilde{M} \cap C_y$ is the union of some of the paths in $\{P_{y,1}, \dots, P_{y,t}\}$. Suppose that these paths are $\{P_{y,c_1}, \dots, P_{y,c_\mu}\}$ where $\{c_1, \dots, c_\mu\} \subseteq [t]$. We consider the boundaried graph $\mathbf{M}_y = (M_y, B_y, \rho_y)$ where $M_y = (\tilde{M} \cap \overline{D}_y) \cup (B_y, \emptyset)$ (i.e. the graph $\tilde{M} \cap \overline{D}_y$ together with the isolated vertices B_y), $B_y = \{r_{y,1}, \dots, r_{y,t}\}$, and for every $d \in [t]$, $\rho_y(r_{y,d}) = d$. We also define $\hat{M}_y = \tilde{M} \setminus D_y \setminus \bigcup_{d \in [t]} (V(P_{y,c_d}) \setminus r_{y,c_d})$. Keep in mind that \hat{M}_y does not intersect the disk Δ' (see [Figure 7](#)).




Figure 9: The graphs M_y (depicted in red) and \hat{M}_y (depicted in blue).

Now consider the t -boundaried graph $\text{diss}(\mathbf{M}_y, T \cup B_y)$ and notice that it is isomorphic to a member $\mathbf{F} \in \mathcal{F}_y$. We set $y' = i' + \ell - 1$. Recall that $\mathbf{F} \in \mathcal{F}_{y'}$, as $\mathcal{F}_y = \mathcal{F}_{y'}$. This means that $\mathbf{G}_{y'}$ contains as a subgraph a model $\mathbf{M}_{y'} = (M_{y'}, B_{y'}, \rho_{y'})$ of \mathbf{F} where $B_{y'} = \{r_{y',1}, \dots, r_{y',t}\}$, and for every $d \in [t]$, $\rho_{y'}(r_{y',d}) = d$. Notice that $M_{y'}$ does not intersect Δ' . Let $\hat{M}_{y'}$ be the graph obtained from $M_{y'}$ after removing the vertices $r_{y',j}, j \in [t] \setminus \{c_1, \dots, c_\mu\}$. For every $d \in [\mu]$, we define $P_d^* = P_{y,c_d} \cup R_{y \rightarrow y', c_d}$ and observe that none of the paths in $\mathcal{P}^* = \{P_d^* \mid d \in [\mu]\}$ intersects Δ' . Consider now the graph $M_0 := \hat{M}_y \cup \hat{M}_{y'} \cup \mathbf{UP}^*$ and observe that (M_0, T) is a topological minor model of H in $G \setminus S$ such that $V(M_0) \cap \Delta' = \emptyset$. Therefore $H \preceq (G \setminus \Delta') \setminus S$, a contradiction (see [Figure 10](#)).




Figure 10: Visualization of the last part of the proof.

□

4 Proof of the main result

Now we have all necessary results in order to prove [Theorem 2](#).

Proof of [Theorem 2](#). Let $g := \binom{h}{2}$. Also, let

$$\begin{aligned} x &:= (k+1)(h+1)(f_4(g)+3), \\ y &:= f_5(f_3(g)+1, h+f_3(g)+1) \cdot ((h+1)(f_4(g)+3)+1), \text{ and} \\ z &:= f_5(f_3(g)+1, h+f_3(g)+1)^{h+1} \cdot k(k+1)+3. \end{aligned}$$

For $q := f_2(x, y, z)$, we call the algorithm **Find_Wall**(G, q) of [Proposition 1](#) which outputs either a q -wall W of G whose compass has treewidth at most $c_1 \cdot q$ or a tree decomposition of G of width at most $c_1 \cdot q$. We consider the first case.

Let Δ be the closed disk defined by the compass of W . Then, we call the algorithm **Find_Collection_of_Annuli**(x, y, z, Δ, G, W) of [Lemma 1](#) which outputs a collection $\mathfrak{A} = \{\mathcal{A}_0, \mathcal{A}_1, \dots, \mathcal{A}_z\}$ of railed annuli of the compass of G such that

- \mathcal{A}_0 is a (x, x) -railed annulus whose outer disk is Δ and whose inner disk is Δ' ,
- for $i \in [z]$, \mathcal{A}_i is a (y, y) -railed annulus of $G \cap \Delta'$, and
- for every $i, j \in [z]$ where $i \neq j$, the outer disk of \mathcal{A}_i and the outer disk of \mathcal{A}_j are disjoint.

We call the algorithm **Reduce_Solution_Space**($k, h, g, \Delta, G, w, \mathcal{C}, \mathcal{P}$) of [Lemma 3](#) for $(\mathcal{C}, \mathcal{P}) := \mathcal{A}_0$ and $w := c_1 \cdot q$ which outputs a set $R \subseteq \Delta' \cap V(G)$ such that

- $|R| \leq f_5(f_3(g)+1, h+f_3(g)+1)^{h+1} \cdot k(k+1) = z - 1$ and
- for every graph H on at most h vertices and g edges and every $|S| \leq k$ and $H \not\subseteq G \setminus S$, then there is some $S' \subseteq (V(G) \setminus \Delta') \cup R$ such that $|S'| \leq k$ and $H \not\subseteq G \setminus S'$.

Since $|R| < z$ then there exists a $j \in [z]$ such that $R \cap \text{ann}(\mathcal{A}_j) = \emptyset$. Let $(\mathcal{C}^{(j)}, \mathcal{P}^{(j)}) := \mathcal{A}_j$. Now, for $b := 2$, the algorithm **Find_irrelevant_area** $(h, g, b, \Delta, G, w, \mathcal{C}^{(j)}, \mathcal{P}^{(j)})$ of [Lemma 4](#) computes a closed disk $\Delta' \subseteq \Delta$ such that

- $\text{tw}(G \cap \Delta') \geq b$, and
- for every graph H on at most h vertices and g edges and every $S \subseteq V(G) \setminus \Delta$, if $H \not\subseteq (G \setminus (\Delta' \cap V(G))) \setminus S$ then $H \not\subseteq G \setminus S$.

Therefore, there exists a vertex $v \in V(G) \cap \Delta'$ such that $\mathbf{p}_H(G) \leq k \iff \mathbf{p}_H(G \setminus v) \leq k$. \square

5 Proof of the Model Taming theorem

This section is devoted to the proof of [Theorem 3](#) that is the base of the correctness of both algorithmic results of the previous section.

5.1 Linkages in railed annuli

A *linkage* in a graph G is a subgraph L of G whose connected components are all non-trivial paths. The *paths* of a linkage are its connected components and we denote them by $\mathcal{P}(L)$. The *size* of L is the number of its paths and is denoted by $|L|$. The *terminals* of a linkage L , denoted by $T(L)$, are the endpoints of the paths in $\mathcal{P}(L)$, and the *pattern* of L is the set

$$\{\{s, t\} \mid \mathcal{P}(L) \text{ contains some } (s, t)\text{-path}\}.$$

Two linkages L_1, L_2 of G are *equivalent* if they have the same pattern and we denote this fact by $L_1 \equiv L_2$. We say that a linkage L of a graph G is *vital* if $V(L) = V(G)$ and there is no other linkage of G that is equivalent to L .

Let G be a partially Δ -embedded graph, let $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ be a (r, q) -railed annulus of G and L be a linkage of G . If L is a linkage of a partially Δ -embedded graph, and $D \subseteq \Delta$, then we say that L is *D-avoiding* if $T(L) \cap D = \emptyset$. We also say that L is *D-free* if $D \cap L = \emptyset$. We also say that L is *A-avoiding* if it is $\text{ann}(\mathcal{C})$ -avoiding (see [Figure 11](#)).




Figure 11: An example of a railed annulus \mathcal{A} , a closed disk D (depicted in blue) and a linkage L (depicted in red) that is D -free and \mathcal{A} -avoiding.

Let $r = 2t + 1$. Let also $s \in [r]$ where $s = 2t' + 1$. Given some $I \subseteq [q]$, we say that a linkage L is (s, I) -confined in \mathcal{A} if

$$L \cap \text{ann}(\mathcal{C}, t - t', t + t') \subseteq \bigcup_{i \in I} P_i.$$

Our purpose is to prove the following.

Theorem 4. *There exist two functions $f_3, f_4 : \mathbb{N}_{\geq 0} \rightarrow \mathbb{N}_{\geq 0}$ such that for every odd $s \in \mathbb{N}_{\geq 1}$ and every $k \in \mathbb{N}_{\geq 0}$, if G is a partially Δ -embedded graph, $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ is a (r, q) -railed annulus of G , where $r \geq f_4(k) + s$ and $q \geq 5/2 \cdot f_3(k)$, L is an \mathcal{A} -avoiding linkage of size at most k , and $I \subseteq [q]$, where $|I| > f_3(k)$, then G contains a linkage \tilde{L} where $\tilde{L} \equiv L$, \tilde{L} is \mathcal{A} -avoiding, $\tilde{L} \setminus \text{ann}(\mathcal{C}) \subseteq L \setminus \text{ann}(\mathcal{C})$, and \tilde{L} is (s, I) -confined in \mathcal{A} .*

We say that a function is *even* if its images are even numbers. We state the following result.

Proposition 2 ([20, 26]). *There exists an even function $f_3 : \mathbb{N}_{\geq 0} \times \mathbb{N}_{\geq 0} \rightarrow \mathbb{N}_{\geq 0}$ such that if G is a graph and L is a vital linkage of G , then $\text{tw}(G) \leq f_3(|L|)$.*

In the above proposition, f_3 is a non-decreasing function that is important for the statement of many of the results of this paper. For this reason, for now on, f_3 will always denote the function of [Proposition 2](#).

5.2 Taming a Linkage

LB-pairs. Given a graph G , a *LB-pair* of G is a pair (L, B) where B is a subgraph of G with maximum degree 2 and L is a linkage of G . We define $\text{cae}(L, B) = |E(L) \setminus E(B)|$ (i.e., the number of linkage edges that are not edges of B).

Lemma 5. *Let (L, B) be an LB-pair of a G . If $\text{tw}(L \cup B) > f_3(|L|)$, then G contains a linkage L' where*

1. $\text{cae}(L', B) < \text{cae}(L, B)$,
2. $L' \equiv L$,
3. $L' \subseteq L \cup B$.

Proof. Let $H = L \cup B$. From [Proposition 2](#), L is not a vital linkage of H , therefore, H contains a linkage L' such that $L \neq L'$ and $L' \equiv L$. Notice that $E(L') \setminus E(B) \subseteq E(L) \setminus E(B)$. It remains to prove that this inclusion is proper.

Let $\{x, y\}$ be a member of the common pattern of L and L' such that the (x, y) -path P of L is different than the (x, y) -path P' of L' . Clearly, P and P' , when oriented from x to y , have a common part P^* . Formally, this is the connected component of $P \cap P'$ that contains x . Let e be the $(m + 1)$ th edge of P , starting from x , where m is the length of P^* . Notice that $e \in E(L) \setminus E(B)$, while $e \notin E(L') \setminus E(B)$.

We conclude that $E(L') \setminus E(B) \subsetneq E(L) \setminus E(B)$, therefore $|E(L') \setminus E(B)| < |E(L) \setminus E(B)|$, as required. \square

Minimal linkages. Let G be a partially Δ -embedded graph, \mathcal{C} be a Δ -nested cycle collection of G , $D \subseteq \Delta$, L be a $\text{ann}(\mathcal{C})$ -avoiding and D -free linkage of G . We say that a linkage L' of G is (\mathcal{C}, D, L) -minimal if, among all the $\text{ann}(\mathcal{C})$ -avoiding linkages of G that are equivalent to L and are subgraphs of $L \cup (\mathbf{UC} \setminus D)$, L' is one where the quantity $\text{cae}(L', \mathbf{UC} \setminus D)$ is minimized.

Lemma 6. *Let G be a partially Δ -embedded graph, \mathcal{C} be a Δ -nested cycle collection of G , $D \subseteq \Delta$, L be an $\text{ann}(\mathcal{C})$ -avoiding and D -free linkage of G , and L' be a (\mathcal{C}, D, L) -minimal linkage of G , then $\text{tw}(L' \cup (\mathbf{UC} \setminus D)) \leq f_3(|L'|)$.*

Proof. Let $B = \mathbf{UC} \setminus D$ and observe that (L', B) is an LB-pair of G . Assume, towards a contradiction, that $\text{tw}(L' \cup B) > f_3(|L'|)$. From Lemma 5, G contains a linkage L'' that is equivalent to L' where $\text{cae}(L'', B) < \text{cae}(L', B)$ and $L'' \subseteq L' \cup B$. This contradicts the choice of L' as a (\mathcal{C}, D, L) -minimal linkage of G . \square

Streams, rivers, mountains, and valleys. Let G be a partially Δ -embedded graph, $\mathcal{C} = [C_1, \dots, C_r]$ be Δ -nested cycle collection of G , and L be a $\text{ann}(\mathcal{C})$ -avoiding linkage of G . A (\mathcal{C}, L) -stream of G is a subpath of L that is a subset P of $\text{ann}(\mathcal{C})$ and such that $V(P \cap C_1)$ consists of the one endpoint of P and $V(P \cap C_r)$ consists of the other. A disjoint collection of (\mathcal{C}, L) -streams of G is a collection \mathcal{R} of (\mathcal{C}, L) -streams such that \mathbf{UR} is a linkage of G . A (\mathcal{C}, L) -river of G is a (\mathcal{C}, L) -stream that is a subpath of a connected component of $L \cap \text{ann}(\mathcal{C})$ that has one of its endpoints in C_1 and the other in C_r . Notice that not each (\mathcal{C}, L) -stream of G is a (\mathcal{C}, L) -river and any collection of (\mathcal{C}, L) -rivers is a disjoint collection of (\mathcal{C}, L) -streams (see Figure 12).




Figure 12: An example of a (\mathcal{C}, L) -stream (depicted in solid red) and a (\mathcal{C}, L) -river (depicted in solid blue).

Let $i \in [r]$ and $D \subseteq \Delta$. An (\mathcal{C}, D, L) -mountain (resp. (\mathcal{C}, D, L) -valley) of G based on C_i is a non-trivial subpath P of some path of L where

1. $P \subseteq \overline{D}_i$ (resp. $P \subseteq \Delta \setminus D_i$),
2. $P \cap D_r = \emptyset$ (resp. $P \cap (\Delta \setminus \overline{D}_1) = \emptyset$),
3. $P \cap C_i$ has two connected components, each containing exactly one of the endpoints of P ,
4. if D' is the closure of the connected component of $D_i \setminus P$ (resp. $(\Delta \setminus \overline{D}_i) \setminus P$) that does not contain D_r (resp. $\Delta \setminus \overline{D}_1$), then $D' \cap T(L) = \emptyset$ and $D' \cap D = \emptyset$.




Figure 13: An example of a (\mathcal{C}, D, L) -valley (depicted in solid red), and some (\mathcal{C}, D, L) -mountains (depicted in solid colors). Notice that the (\mathcal{C}, D, L) -mountain depicted in green is tight.

Clearly, in (4), D' is a closed disk. We call it, the *disk* of the (\mathcal{C}, D, L) -mountain (resp. valley) P and we denote it by $\text{disk}(P)$. Notice that there is no (\mathcal{C}, D, L) -mountain based on C_r and there is no (\mathcal{C}, D, L) -valley based on C_1 .

A (\mathcal{C}, D, L) -mountain (resp. (\mathcal{C}, D, L) -valley) of G is any (\mathcal{C}, D, L) -mountain (resp. (\mathcal{C}, D, L) -valley) of G based on some of the cycles of \mathcal{C} .

The *height* (resp. *depth*) of a (\mathcal{C}, D, L) -mountain (resp. (\mathcal{C}, D, L) -valley) P that is based on C_i is the maximum j such that C_{i+j-1} (resp. C_{i-j+1}) intersects P and, in both cases, we denote it by $\text{dehe}(P)$. Moreover, the height (resp. depth) of P is at least 1 and at most r .

Notice that if a (\mathcal{C}, L) -stream P of G is a subpath of a (\mathcal{C}, D, L) -mountain P' or a (\mathcal{C}, D, L) -valley P' of G then $\text{dehe}(P') = r$. Moreover, if a (\mathcal{C}, L) -stream P of G is not a subpath of some (\mathcal{C}, D, L) -mountain or some (\mathcal{C}, D, L) -valley of G , then P is a (\mathcal{C}, L) -river of G .

We say that a (\mathcal{C}, D, L) -mountain (resp. (\mathcal{C}, D, L) -valley) P based on C_i , is *tight* if $\text{dehe}(P) = d \geq 2$ and there is a sequence $[P_2, \dots, P_d]$ of (\mathcal{C}, D, L) -mountains (resp. (\mathcal{C}, D, L) -valleys) based on C_i such that

- $P = P_d$,
- $\forall j \in [2, d]$, $\text{dehe}(P_j) = j$, and
- $\forall j \in [2, d - 1]$, $P_j \subseteq \text{disk}(P_{j+1})$ (see Figure 13).

Lemma 7. *Let G be a partially Δ -embedded graph, \mathcal{C} be a Δ -nested cycle collection of G , $D \subseteq \Delta$, L be a $\text{ann}(\mathcal{C})$ -avoiding and D -free linkage of G . Let also L' be a (\mathcal{C}, D, L) -minimal linkage of G . Then all (\mathcal{C}, D, L') -mountains (resp. (\mathcal{C}, D, L') -valleys) of G are tight.*

Proof. Let $B = \mathbf{UC} \setminus D$. We present the proof for the case of (\mathcal{C}, D, L') -mountains as the case of (\mathcal{C}, D, L') -valleys is symmetric.

Claim: Let $i \in \mathbb{N}_{\geq 1}, j \in \mathbb{N}_{\geq 2}$. If P_j is a (\mathcal{C}, D, L') -mountain of G based on C_i such that $\text{dehe}(P_j) = j$, then there exists a (\mathcal{C}, D, L') -mountain P' based on C_i such that $\text{dehe}(P') = j - 1$ and $P' \subseteq \text{disk}(P_j)$.

Proof of Claim: Suppose to the contrary that there does not exist a (\mathcal{C}, D, L') -mountain P' based on C_i such that $\text{dehe}(P') = j - 1$ and $P' \subseteq \text{disk}(P_j)$. Let $P_j^{(j-1)} = (P_j \setminus D_{i+(j-1)-1}) \cup (C_{i+(j-1)-1} \cap \text{disk}(P_j))$ and notice that $\text{dehe}(P_j^{(j-1)}) = j - 1$ (see Figure 14).




Figure 14: An example of a (\mathcal{C}, D, L') -mountain P_j of G based on C_i such that $\text{dehe}(P_j) = j$ (depicted in red) and the (\mathcal{C}, D, L') -mountain $P_j^{(j-1)}$ (depicted in green).

Then G contains a linkage $L'' = (L' \setminus P_j) \cup (P_j^{(j-1)})$ that is equivalent to L where $\text{cae}(L'', B) < \text{cae}(L', B)$ and $L'' \subseteq L' \cup B$. This contradicts the choice of L' as a (\mathcal{C}, D, L) -minimal linkage of G . The claim follows.

Let P be a (\mathcal{C}, D, L') -mountain of G based on C_i such that $\text{dehe}(P) = d \geq 2$. The fact that P is tight follows by recursively applying the Claim above. \square

Orderings of streams. Let G be a partially Δ -embedded graph, \mathcal{C} be a Δ -nested cycle collection of G , D be an open disk where $D \subseteq \text{ann}(\mathcal{C})$, L be an $\text{ann}(\mathcal{C})$ -avoiding and D -free linkage of G .

If \mathcal{Z} is a disjoint collection of (\mathcal{C}, L) -streams of G we define its D -ordering as follows: Consider the sequence $[Z_1, \dots, Z_d]$ such that for each $i \in [d]$, one, say D_i , of the two connected components of $\text{ann}(\mathcal{C}) \setminus (Z_i \cup Z_{i+1})$ does not intersect $\mathbf{U}\mathcal{Z}$ (here Z_{d+1} denotes Z_1). Among all $(d - 1)!$ such sequences we insist that $[Z_1, \dots, Z_d]$ is the unique one where $D \subseteq D_q$ and that the order of \mathcal{Z} is the counter-clockwise order that its elements appear around $\text{ann}(\mathcal{C})$ (see Figure 15). We call $[Z_1, \dots, Z_d]$ the D -ordering of \mathcal{Z} .




Figure 15: An example of an Δ -nested cycle collection \mathcal{C} , an open disk $D \subseteq \text{ann}(\mathcal{C})$ (depicted in blue), a linkage L (depicted in red) that is D -free and $\text{ann}(\mathcal{C})$ -avoiding, a disjoint collection \mathcal{Z} of (\mathcal{C}, L) -streams, and the D -ordering $[Z_1, \dots, Z_5]$ of \mathcal{Z} .

Brambles. Given a graph G , we say that a subset S of $V(G)$ is *connected* if $G[S]$ is connected. Given $S_1, S_2 \subseteq V(G)$, we say that S_1 and S_2 touch if either $S_1 \cap S_2 \neq \emptyset$ or there is an edge $e \in E(G)$ where $e \cap S_1 \neq \emptyset$ and $e \cap S_2 \neq \emptyset$. A *bramble* in G is a collection \mathcal{B} of pairwise touching connected subsets of $V(G)$. The *order* of a bramble \mathcal{B} is the minimum number of vertices that intersect all of its elements.

Proposition 3 ([29]). *Let $k \in \mathbb{N}_{\geq 0}$. A graph G has a bramble of order $k + 1$ if and only if $\text{tw}(G) \geq k$.*

We now use the notions of ordering of streams and brambles to prove the following result.

Lemma 8. *Let G be a partially Δ -embedded graph, \mathcal{C} be Δ -nested cycle collection of G , D be an open disk where $D \subseteq \text{ann}(\mathcal{C})$, L be an $\text{ann}(\mathcal{C})$ -avoiding and D -free linkage of G , and \mathcal{Z} be a disjoint collection of (\mathcal{C}, L) -streams of G . Then $\text{tw}(L \cup (\mathbf{UC} \setminus D)) \geq \min\{|\mathcal{C}|, |\mathcal{Z}|\}$.*

Proof. Let $[Z_1, \dots, Z_d]$ be the D -ordering of \mathcal{Z} and let D' be the connected component of $\text{ann}(\mathcal{C}) \setminus (Z_d \cup Z_1)$ that contains D . Let $r = \min\{|\mathcal{C}|, |\mathcal{Z}|\}$, and let $[Z_1, \dots, Z_r]$ be the sequence consisting of the first r elements of the D -ordering of \mathcal{Z} . Let also \mathcal{C}' be the sequence consisting of the first r elements of \mathcal{C} . Notice that there is a disjoint collection $\mathcal{Z}' = [Z'_1, \dots, Z'_r]$ of (\mathcal{C}', L) -streams of G such that for each $i \in [r]$, $Z'_i \subseteq Z_i$.

We now set $\mathcal{B} = \mathcal{C}' \setminus D'$, denote $\mathcal{B} = [B_1, \dots, B_r]$, and notice that both \mathcal{B} and \mathcal{Z}' are sequences of paths in G , such that both \mathbf{UB} and \mathbf{UZ}' are linkages of G . Consider now the graph $Q = \mathbf{UB} \cup \mathbf{UZ}'$ and notice that $C = B_1 \cup Z'_1 \cup B_r \cup Z'_r$ is a cycle of G .

As $Q \subseteq L \cup (\mathbf{UC} \setminus D)$, it remains to prove that $\text{tw}(Q) \geq r$. For this, because of **Proposition 3**, it suffices to give a bramble of Q of order $r + 1$. For each $(i, j) \in [2, r - 1]^2$ we define $X^{(i,j)} = (B_i \cup Z'_j) \setminus V(C)$. It is easy to check that $\mathcal{X} = \{X^{(i,j)} \mid (i, j) \in [2, r - 1]^2\}$ is a bramble of Q of order $\geq r - 2$. Let also $X^{(1)} = Z_1 \setminus B_1$, $X^{(2)} = B_1$, and $X^{(3)} = Z'_r \cup B_r$. Notice that $\mathcal{X} \cup \{X^{(1)}, X^{(2)}, X^{(3)}\}$ is also a bramble of Q and its order is the order of \mathcal{X} incremented by 3. Therefore Q contains a bramble of order at least $r + 1$, as required (see **Figure 5.2**).




Figure 16: An example of the construction of a bramble of Q , where $|\mathcal{B}| = 5$ and $|\mathcal{Z}'| = 5$. Here, $X^{(2,2)}, X^{(3,3)}, X^{(4,4)}$ are depicted in red, green, and yellow, respectively, while $X^{(1)}, X^{(2)}, X^{(3)}$ are depicted in orange, brown, and blue, respectively.

□

Lemma 9. Let G be a partially Δ -embedded graph, \mathcal{C} be a Δ -nested cycle collection of G , D be a connected subset of Δ , and L be a $\text{ann}(\mathcal{C})$ -avoiding and D -free linkage of G . If P is a tight (\mathcal{C}, D, L) -mountain (resp. (\mathcal{C}, D, L) -valley) of G , then $\text{tw}(L \cup (\mathbf{UC} \setminus D)) \geq \frac{2}{3} \cdot \text{dehe}(P)$.

Proof. Let $d = \text{dehe}(P)$. We examine the non-trivial case where $d \geq 3$. We present the proof for the case where P is a (\mathcal{C}, D, L) -mountain as the case where P is a (\mathcal{C}, D, L) -valley is symmetric.

We assume that P is based on C_i , for some $i \in [r]$. By the definition of tightness, there is a sequence $\mathcal{P} = [P_2, \dots, P_d = P]$ of (\mathcal{C}, D, L) -mountains (resp. (\mathcal{C}, D, L) -valleys) based on C_i such that

- $P = P_d$,
- $\forall i \in [2, d]$, $\text{dehe}(P_i) = i$, and
- $\forall i \in [2, d-1]$, $P_i \subseteq \text{disk}(P_{i+1})$.

For every $j \in [2, d]$, we denote $\mathcal{C}^{(j)} = [C_i, \dots, C_{i+j-1}]$. Notice that for every $j \in [2, d]$, L is an $\text{ann}(\mathcal{C}^{(j)})$ -avoiding and D -free linkage of G .

Claim: For every $j \in [2, d-1]$ there exists a disjoint collection \mathcal{Z}_j of $(\mathcal{C}^{(j)}, L)$ -streams of G where $|\mathcal{Z}_j| \geq 2(d-j) + 1$.

Proof of Claim: Let $j \in [2, d-1]$. Observe that for each $h \in [j+1, d]$ exactly two of the connected components of $\text{ann}(\mathcal{C}^{(j)}) \cap P_h$ are $(\mathcal{C}^{(j)}, L)$ -rivers in G . This implies that there is a collection \mathcal{R}_j of at least $2(d-j)$ $(\mathcal{C}^{(j)}, L)$ -rivers in G . Recall that \mathcal{R}_j is a disjoint collection of $(\mathcal{C}^{(j)}, L)$ -streams of G . Observe also that we can pick some sub-path of $\text{ann}(\mathcal{C}^{(j)}) \cap P_j$ that has one endpoint in C_i and the other in C_{i+j-1} . As this path does not share vertices with any of the paths in \mathcal{R}_j we can add it in \mathcal{R}_j and obtain a disjoint collection \mathcal{Z}_j of $(\mathcal{C}^{(j)}, L)$ -streams of G where $|\mathcal{Z}_j| \geq 2(d-j) + 1$. Claim follows (see [Figure 17](#)).




Figure 17: An example of a tight (\mathcal{C}, D, L) -mountain P based on C_i of height d and the respective sequence of (\mathcal{C}, D, L) -mountains based on C_i (depicted in red), an annulus $\mathcal{C}^{(j)}$ (depicted in blue), for some $j \in [2, d]$, and a disjoint collection \mathcal{Z}_j (depicted in green) of $(\mathcal{C}^{(j)}, L)$ -streams of G .

We now set $j' = \lfloor (2d+1)/3 \rfloor$ and observe that $2 \leq j' \leq d-1$. The above claim implies that there exists a disjoint collection $\mathcal{Z}_{j'}$ of $(\mathcal{C}^{(j')}, L)$ -streams of G such that $|\mathcal{Z}_{j'}| \geq 2(d-j') + 1 \geq j' = |\mathcal{C}^{(j')}|$. Therefore, we can apply [Lemma 8](#) on $\mathcal{C}^{(j')}$ and deduce that $\text{tw}(L \cup (\mathbf{UC}^{(j')} \setminus D)) \geq j'$. The Lemma follows as $L \cup (\mathbf{UC}^{(j')} \setminus D) \subseteq L \cup (\mathbf{UC} \setminus D)$ and $\lfloor (2d+1)/3 \rfloor \geq 2d/3$. \square

Lemma 10. *Let G be a partially Δ -embedded graph, \mathcal{C} be a Δ -nested cycle collection of G , D be a connected subset of Δ , L be a $\text{ann}(\mathcal{C})$ -avoiding and D -free linkage of G , and L' be a (\mathcal{C}, D, L) -minimal linkage of G . Then all (\mathcal{C}, D, L') -mountains (resp. (\mathcal{C}, D, L') -valleys) of G have height (resp. depth) at most $\frac{3}{2} \cdot f_3(|L'|)$.*

Proof. We set $B = \mathbf{UC} \setminus D$. By Lemma 6, $\text{tw}(L' \cup B) \leq f_3(|L'|)$. Let P be a (\mathcal{C}, D, L') -mountain (resp. (\mathcal{C}, D, L') -valley) of G based on C_i , for some $i \in [r-1]$ (resp. $i \in [2, r]$). From Lemma 7, P should be tight and, from Lemma 9, $\text{tw}(L' \cup B) \geq \frac{2}{3} \cdot \text{dehe}(P)$. Therefore, $\text{dehe}(P) \leq \frac{3}{2} \cdot f_3(|L'|)$. \square

Lemma 11. *Let G be a partially Δ -embedded graph, $\mathcal{C} = [C_1, \dots, C_r]$ be a Δ -nested cycle collection of G , and L be a \overline{D}_1 -avoiding linkage. Then there is a linkage L' of G such that*

1. L' is \overline{D}_1 -avoiding,
2. $L' \equiv L$,
3. L' is $\overline{D}_{3m/2+1}$ -free, where $m = f_3(|L'|)$.

Proof. Let G^+ be the graph obtained if we take its disjoint union with a cycle $C_{r+1} \subseteq D_r$ and we set $\mathcal{C}^+ = [C_1, \dots, C_r, C_{r+1}]$. Observe that L is an $\text{ann}(\mathcal{C}^+)$ -avoiding linkage of G^+ . Let L' be a $(\mathcal{C}^+, \emptyset, L)$ -minimal linkage of G^+ .

As $L' \equiv L$, L' is a \overline{D}_1 -avoiding linkage of both G and G^+ . Therefore L' satisfies conditions 1 and 2. For condition 3, assume to the contrary that L' is a linkage of G that is intersecting $\overline{D}_{3m/2+1}$. As L' is a \overline{D}_1 -avoiding linkage of G^+ we obtain that G^+ contains some $(\mathcal{C}, \emptyset, L')$ -mountain P , based on C_1 where $\text{dehe}(P) > 3m/2$. On the other side, as L is an $\text{ann}(\mathcal{C}^+)$ -avoiding linkage of G^+ we can apply Lemma 10, on G^+ , \mathcal{C}^+ , \emptyset , L , and L' and obtain that $\text{dehe}(P) \leq 3m/2$, a contradiction. \square

Lemma 12. *Let G be a partially Δ -embedded graph, \mathcal{C} be a Δ -nested cycle collection of G , $D \subseteq \Delta$, and L be an \mathcal{A} -avoiding and D -free linkage. If $|\mathcal{C}| > m = f_3(|L|)$, then G contains a linkage L' of G such that*

1. $L' \equiv L$,
2. $L' \cap D = \emptyset$,
3. All (\mathcal{C}, D, L') -mountains of G have height at most $\frac{3}{2}m$,
4. All (\mathcal{C}, D, L') -valleys of G have depth at most $\frac{3}{2}m$, and
5. L' has at most m \mathcal{A} -rivers.

Proof. Let L' be a (\mathcal{C}, D, L) -minimal linkage. (1) and (2) follow by the definition of a (\mathcal{C}, D, L) -minimal linkage. (3) and (4) follow directly from Lemma 10. To prove (5), assume that G contains a collection \mathcal{Z} of (\mathcal{C}, L') -rivers where $|\mathcal{Z}| > m$. Recall that \mathcal{Z} is a disjoint collection of (\mathcal{C}, L') -streams of G . From Lemma 8, $\text{tw}(L' \cup (\mathbf{UC} \setminus D)) \geq \min\{|\mathcal{C}|, |\mathcal{Z}|\} > m$. We arrive to a contradiction as, from Lemma 6, $\text{tw}(L' \cup (\mathbf{UC} \setminus D)) \leq m$. \square

5.3 Rerouting a linkage

The following proposition is a direct consequence of [1, Lemma 7].

Proposition 4. *Let k, k', d be integers such that $0 \leq d \leq k' \leq k$. Let Γ be a $(k \times k')$ -grid and let $\{p_1^{\text{up}}, \dots, p_\rho^{\text{up}}\}$ (resp. $\{p_1^{\text{down}}, \dots, p_\rho^{\text{down}}\}$) be vertices of the higher (resp. lower) horizontal line arranged as they appear in it from left to right. Then the grid Γ contains ρ pairwise disjoint paths P_1, \dots, P_ρ such that, for every $h \in [\rho]$, the endpoints of P_h are p_h^{up} and p_h^{down} .*

Given two vertex disjoint paths P_1 and P_2 of G , we say that an (P_1, P_2) -path of G is a path that whose one endpoint is a vertex of P_1 the other endpoint is a vertex of H_2 and contains all edges of $P_1 \cup P_2$. We now prove the following:

Lemma 13. *Let $r, q, s \in \mathbb{N}_{\geq 3}$, $b, d \in \mathbb{N}_{\geq 0}$, such that $r \geq s + 2b$ and $q \geq b + d$, where r and s are odd numbers. If G is a partially Δ -embedded graph, \mathcal{A} is a (r, q) -railed annulus of G , $I \subseteq [q]$ where $|I| \geq d$, then there is a linkage K of G such that,*

(a) *there is an ordering $\mathcal{P}(K) = [K_1, \dots, K_d]$, where for $i \in [d]$, K_i is a $(P_{1,b+i}, P_{r,b+i})$ -path of G .*

(b) *K is (s, I) -confined in \mathcal{A} .*

Proof. Let $\mathcal{A} = (\mathcal{C}, \mathcal{P})$, let $t = \lfloor r/2 \rfloor$ and $t' = \lfloor s/2 \rfloor$. Also, let $\{i_1, \dots, i_d\} \subseteq I$ such that $\forall j \in [d-1], i_j < i_{j+1}$.

Claim: There is a collection of pairwise disjoint paths $\mathcal{P}^{\text{down}} = \{P_1^{\text{down}}, \dots, P_d^{\text{down}}\}$ such that, for every $h \in [d]$, P_h^{down} is a $(P_{1,b+h}, P_{b,i_h})$ -path.

Proof of Claim: For $i \in [b], j \in [q]$ let $p_{i,j}$ be the vertex obtained after contracting all edges in $P_{i,j}$. We also define:

- $E_1 = \{e = \{p_{i,j}, p_{i,j+1}\} \mid e \text{ is the edge obtained after contracting all but one of the edges of } L_{i,j \rightarrow j+1}, i \in [b], j \in [q-1]\}$ and
- $E_2 = \{e = \{p_{i,j}, p_{i+1,j}\} \mid e \text{ is the edge obtained after contracting all but one of the edges of } R_{i \rightarrow i+1,j}, i \in [b-1], j \in [q]\}$.

Let H be the graph where $V(H) = \{p_{i,j} \mid (i, j) \in [b] \times [q]\}$ and $E(H) = E_1 \cup E_2$. Observe that H is isomorphic to a $(q \times b)$ -grid (see Figure 18). For $h \in [d]$, let p_h^{low} (resp. p_h^{high}) be the vertex $p_{1,b+h}$ (resp. p_{b,i_h}).




Figure 18: An example showing the construction of the graph H . For every $h \in [d]$, the resulting vertices p_h^{low} and p_h^{high} (corresponding to the vertices of the paths $P_{1,b+h}$ and P_{b,i_h} , respectively) are depicted in white.

Due to [Proposition 4](#), H contains d pairwise disjoint paths P_1, \dots, P_d such that, for every $h \in [d]$, the endpoints of P_h are p_h^{low} and p_h^{high} . Therefore, if we substitute every vertex of each P_i with the the edges that where contracted in G in order to obtain it in H , we obtain the claimed result.

By applying the previous claim symmetrically, we can find a collection of pairwise disjoint paths $\mathcal{P}^{\text{up}} = \{P_1^{\text{up}}, \dots, P_d^{\text{up}}\}$ such that, for every $h \in [d]$, P_h^{up} is a $(P_{r-b, i_h}, P_{r, b+h})$ path.

Now, for every $h \in [d]$, let $P_h^{\text{mid}} = \text{ann}(\mathcal{C}, b, r-b) \cap P_{i_h}$ and let $K_h = P_h^{\text{down}} \cup P_h^{\text{mid}} \cup P_h^{\text{up}}$. Since $r \geq s + 2b$ and $s = 2t' + 1$ then $\text{ann}(\mathcal{C}, t-t', t+t') \subseteq \text{ann}(\mathcal{C}, b, r-b)$ and therefore $K = \{K_1, \dots, K_d\}$ is the desired linkage. This concludes the proof. \square

Let $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ be an (r, q) -railed annulus of a partially Δ -embedded graph G . We set $z = \lfloor \min\{r, q\}/2 \rfloor$. For each $i \in [z]$, we define $C_i^{\mathcal{A}}$ as the unique cycle of the graph

$$L_{i, i \rightarrow q-i+1} \cup L_{r-i+1, i \rightarrow q-i+1} \cup R_{i \rightarrow r-i+1, i} \cup R_{i \rightarrow r-i+1, q-i+1}.$$

Notice that if $r, q \geq 5$, then $[C_1^{\mathcal{A}}, \dots, C_z^{\mathcal{A}}]$ is a Δ -nested collection of cycles of G and we denote it by $\mathcal{C}_{\mathcal{A}}$ (see [Figure 19](#)).




Figure 19: An example of an $(8, 8)$ -railed annulus $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ and the sequence $\mathcal{C}_{\mathcal{A}}$ (depicted in red).

Proof of [Theorem 4](#). Recall that f_3 is the function of [Proposition 2](#). We define $f_4(k) := 3 \cdot (f_3(k))^2 + 6 \cdot f_3(k) + 2$. For simplicity, we use $m = f_3(k)$. Let also $b = 3m/2$, and keep in mind that $r \geq f_4(k) + s = 3m^2 + 6m + 2 + s = 2(m+1) \cdot b + 2 + s + 2b$ and that $|I| \geq m + 1$.

Recall that $\mathcal{C}_{\mathcal{A}} = [C'_1, \dots, C'_z]$, where $z = \lfloor \min\{q, r\}/2 \rfloor$, is a Δ -nested collection of cycles of G . For each $i \in [z]$, we denote by D'_i (resp. \overline{D}'_i) the open (closed) disk corresponding to C'_i . Let also $\check{D} := D'_{b+1}$ and $D := \overline{D}'_{b+1}$. Keep in mind that $\overline{D}'_1 = \Delta_{1, r, 1, q}$ and $D = \Delta_{b+1, r-b, b+1, q-b}$.

Observe now that L is a \overline{D}'_1 -avoiding linkage. By applying [Lemma 11](#) on G , $\mathcal{C}_{\mathcal{A}}$, L , and \overline{D}'_1 , we obtain that G has a \overline{D}'_1 -avoiding and D -free linkage L' such that $L' \equiv L$.

It is easy to verify that L' is \mathcal{A} -avoiding, $r \geq a > m$, $D \subseteq \text{ann}(\mathcal{C})$, and $|L'| = |L| \leq k$. Therefore, we may apply [Lemma 12](#) on k , G , \mathcal{A} , D , and L' . We obtain a D -free linkage L'' of G that is equivalent to L' (and therefore to L as well) and such that

- (a) All (\mathcal{C}, D, L'') -mountains/valleys of G have height/depth at most b .
- (b) L'' has at most m \mathcal{A} -rivers of G ,

Let $\mathcal{Z} = [Z_1, \dots, Z_d]$ the D -ordering of the \mathcal{A} -rivers of L'' in G and keep in mind that, from (b), $d \leq m$.




Figure 20: Visualization of an (r, q) -railed annulus and the notations introduced above.

For every $i \in [d]$, we define x_i^{down} (resp. x_i^{up}) as the vertex in the path $C_{(i+1) \cdot b+1} \setminus \check{D}$ (resp. $C_{r-(i+1) \cdot b} \setminus \check{D}$) that belongs in Z_i and has the minimum possible distance to the vertices of the path $P_{(i+1) \cdot b+1, q-b}$ (resp. $P_{r-(i+1) \cdot b, q-b}$). We also denote by Q_i^{down} (resp. Q_i^{up}) the path certifying this minimum distance.

For $i \in [d]$, let Z_i^{down} and Z_i^{up} be the two connected components of the graph obtained from Z_i if we remove the edges of its $(x_i^{\text{down}}, x_i^{\text{up}})$ -subpath (see Figure 20). We choose Z_i^{down} (resp. Z_i^{up}) so that it intersects C_1 (resp. C_r).

Claim: For $i \in [d]$, Z_{i-1}^{down} (resp. Z_{i-1}^{up}) does not intersect Q_i^{down} (resp. Q_i^{up}) — where Z_0^{down} (resp. Z_0^{up}) denotes Z_q .

Proof of claim: If $Z_{i-1}^{\text{down}} \cap Q_i^{\text{down}} \neq \emptyset$ (resp. $Z_{i-1}^{\text{up}} \cap Q_i^{\text{up}} \neq \emptyset$) for some $i \in [d]$, then some of the connected components of $Z_{i-1}^{\text{down}} \cap \bar{D}_{i \cdot b+1}$ (resp. $Z_{i-1}^{\text{up}} \cap (\Delta \setminus D_{r-i \cdot b})$) whose endpoints are in $C_{i \cdot b+1}$ (resp. $C_{r-i \cdot b}$) should be a (\mathcal{C}, D, L'') -mountain (resp. (\mathcal{C}, D, L'') -valley) of G of height (resp. depth) $> b$, a contradiction to (a). Claim follows.

Because of the above claim, it follows that the paths $Q_i^{\text{down}} \cup Z_i^{\text{down}}$ (resp. $Q_i^{\text{up}} \cup Z_i^{\text{up}}$), $i \in [d]$ are pairwise vertex-disjoint $(Z_i \cap C_1, P_{(i+1) \cdot b+1, q-b})$ -paths (resp. $(Z_i \cap C_r, P_{r-(i+1) \cdot b, q-b})$ -paths) in G that do not intersect the open disk \check{D} .

Let $w = (m+1) \cdot b + 2$ and $w' = r - (m+1) \cdot b - 1$. For $i \in [d]$, we now define (see Figure 21)

$$Y_i^{\text{down}} = \text{the } (P_{q-b}, P_{b+i})\text{-path in } L_{(i+1) \cdot b+1, q-b \rightarrow b+i} \cup P_{(i+1) \cdot b+1, b+i} \cup R_{(i+1) \cdot b+1 \rightarrow w, b+i},$$

$$Y_i^{\text{up}} = \text{the } (P_{q-b}, P_{b+i})\text{-path in } L_{r-(i+1) \cdot b, q-b \rightarrow b+i} \cup P_{r-(i+1) \cdot b+1, b+i} \cup R_{r-(i+1) \cdot b \rightarrow w', b+i}.$$

By the definition of Y_i^{down} and Y_i^{up} , the graphs $X_i^{\text{down}} = Z_i^{\text{down}} \cup Q_i^{\text{down}} \cup Y_i^{\text{down}}$ and $X_i^{\text{up}} = Z_i^{\text{up}} \cup Q_i^{\text{up}} \cup Y_i^{\text{up}}$, $i \in [d]$, are pairwise vertex-disjoint paths. In particular, taking into account the definition of Y_i^{down} and Y_i^{up} , we have that

$$X_i^{\text{down}} \text{ is a } (Z_i \cap C_1, P_{w, b+i})\text{-path} \quad \text{and} \quad (1)$$

$$X_i^{\text{up}} \text{ is a } (Z_i \cap C_r, P_{w', b+i})\text{-path} \quad (2)$$

Let $\Omega = \text{ann}(\mathcal{C}, w, w')$ and $K' = \bigcup X_i^{\text{down}} \cup \bigcup X_i^{\text{up}}$. Observe that

$$K' \cap \Omega = \{P_{w, b+i}, P_{w', b+i} \mid i \in [d]\}. \quad (3)$$




Figure 21: Visualization of the definition of Y_i^{up} and Y_i^{down} , $i \in [d]$.

Let $\bar{\mathcal{A}} = (\bar{\mathcal{C}}, \bar{\mathcal{P}})$, where $\bar{\mathcal{C}} = [C_w, \dots, C_{w'}]$ and $\bar{\mathcal{P}} = \mathcal{P} \cap \Omega$.

Notice that $|\bar{\mathcal{C}}| = w' - w + 1 = r - 2(m+1) \cdot b - 2 \geq s + 2b$. Notice also that $d \leq |I|$, $I \subseteq [q]$. Finally, $b = 3/2m$ and $d \leq m$ imply that $d + b \leq 5/2m \leq q$. We can now apply [Lemma 13](#) for $r, q, s, b, d, \bar{\mathcal{A}}$, and I and obtain a linkage K of $\bar{\mathcal{A}}$ satisfying properties (a), and (b) of [Lemma 13](#).

From Property (a) we can write $\mathcal{P}(K) = [K_1, \dots, K_d]$ and, using [\(3\)](#), we deduce that that, for $i \in [d]$, K_i is a $(P_{w,b+i}, P_{w',b+i})$ -path of G . This, together with [\(1\)](#), [\(2\)](#), and [\(3\)](#), implies that $K \cup K'$ is a linkage of G where $K \cup K' \subseteq \text{ann}(\mathcal{C})$. From Property (b), K is (s, I) -confined in $\bar{\mathcal{A}}$, therefore, from [\(3\)](#), $K \cup K'$ is (s, I) -confined in \mathcal{A} . Observe also that each of the d paths of $\mathcal{P}(K \cup K')$ is a $(Z_i \cap C_1, Z_i \cap C_r)$ -path of G for some $i \in [d]$. We define

$$\tilde{L} = (L \setminus A') \cup K \cup K'$$

where $A' = \text{ann}(\mathcal{C}) \setminus (C_1 \cup C_r)$. By definition \tilde{L} is a linkage of G where $\tilde{L} \equiv L$ and $\tilde{L} \setminus \text{ann}(\mathcal{C}) \subseteq L \setminus \text{ann}(\mathcal{C})$. Finally, as $K \cup K'$ is (s, I) -confined in \mathcal{A} , then \tilde{L} is (s, I) -confined in \mathcal{A} as well. \square

Now, since we proved [Theorem 4](#), we will use it to in order to prove [Theorem 3](#).

Proof of Theorem 3. Let s be a positive odd integer, H be a graph on g edges, G be a partially Δ -embedded graph, $\mathcal{A} = (\mathcal{C}, \mathcal{P})$ be a (r, q) -railed annulus of G , where $r \geq f_4(g) + 2 + s$ and $q \geq 5/2 \cdot f_3(g)$, (M, T) be a topological minor model of H in G such that $T \cap \text{ann}(\mathcal{A}) = \emptyset$.

Let $\mathcal{A}' = ([C_2, \dots, C_{r-1}], \mathcal{P} \cap \text{ann}(\mathcal{C}, 2, r-1))$ and keep in mind that \mathcal{A}' is a (r, q) -railed annulus of G , where $r \geq f_4(g) + s$ and $q \geq 5/2 \cdot f_3(g)$. Notice that it also holds that $T \cap \text{ann}(\mathcal{A}') = \emptyset$ (see [Figure 22](#)).

Let $\tilde{M}^{(1)} = \tilde{M}[N_{\tilde{M}}[\tilde{T}]]$. Notice that all the connected components of $M \setminus T$ are paths of G . Let L be the linkage of $G \setminus T$ created if we take the union of all non-trivial connected components of $M \setminus T$. Observe that $\mathcal{P}(L)$ is the set of all paths of G connecting neighbors of branch vertices of M and consisting only of subdividing vertices of M . Also, notice that since $T \cap \text{ann}(\mathcal{A}') = \emptyset$, then L is \mathcal{A}' -avoiding and there is a one-to-one correspondence of $\mathcal{P}(L)$ with $E(H)$ and thus $|L| \leq g$.




Figure 22: An example of a topological minor model (M, T) of H in G . Vertices of T are depicted in blue while the neighbors of vertices of T that are also subdividing vertices are depicted in red. Also, $\text{ann}(\mathcal{A}')$ is depicted in green.

Let $I \subseteq [q]$, where $|I| > f_3(h)$. By applying [Theorem 4](#) for s, g, G, \mathcal{A}', L , and I we obtain a linkage \tilde{L} of G such that $\tilde{L} \equiv L$, \tilde{L} is \mathcal{A}' -avoiding, $\tilde{L} \setminus \text{ann}(\mathcal{A}') \subseteq L \setminus \text{ann}(\mathcal{A}')$, and \tilde{L} (s, I) -confined in \mathcal{A}' . We define

$$\tilde{M} = (M \setminus L) \cup \tilde{L}.$$

By definition, (\tilde{M}, T) is a topological minor model of H in G . Also, since $L, \tilde{L} \subseteq \text{ann}(\mathcal{A})$, then $\tilde{M} \setminus \text{ann}(\mathcal{A}) \subseteq M \setminus \text{ann}(\mathcal{A})$. Finally, as \tilde{L} is (s, I) -confined in \mathcal{A}' then \tilde{M} is (s, I) -confined in \mathcal{A} as well. \square

6 Conclusions

In this paper we prove that \mathcal{F} -TM-DELETION is Fixed Parameter Tractable on planar graphs by designing an $O_{k,h}(n^2)$ -time algorithm for his problem. The remaining question is whether the same result can be derived for *all* graphs, as we conjectured in the introduction. Towards this, we chose to state all combinatorial theorems of this paper in more general forms. Based on them, a straightforward generalization is possible for the class of surface embeddable graphs, that is graphs with Euler genus at most γ . Indeed, the only piece of the proof that needs extension is the starting point of the proof, that is the algorithm of [Proposition 1](#), that can easily be extended to work on graphs of Euler genus γ . Using this, we can directly derive a $O_{k,h,\gamma}(n^2)$ -time algorithm for the version of the problem on surfaces. It follows that will much more effort it is possible to extend the result to every class that excludes some fixed graph as a minor. However, for a complete resolution of our conjecture one has to deal with the case where the input graph contains a big clique minor. We believe that the techniques of the algorithm of [\[13\]](#) can be a good starting point in this direction. However, the technical challenges of such an extension are cumbersome.

References

- [1] Isolde Adler, Stavros G. Kolliopoulos, Philipp Klaus Krause, Daniel Lokshtanov, Saket Saurabh, and Dimitrios M. Thilikos. Irrelevant vertices for the planar disjoint paths problem. *J. Comb. Theory, Ser. B*, 122:815–843, 2017.
- [2] Stefan Arnborg, Jens Lagergren, and Detlef Seese. Easy problems for tree-decomposable graphs. *Journal of Algorithms*, 12:308–340, 1991.
- [3] Hans L. Bodlaender, Pinar Heggernes, and Daniel Lokshtanov. Graph modification problems (dagstuhl seminar 14071). *Dagstuhl Reports*, 4(2):38–59, 2014.
- [4] Richard B. Borie, R. Gary Parker, and Craig A. Tovey. Automatic generation of linear-time algorithms from predicate calculus descriptions of problems on recursively constructed graph families. *Algorithmica*, 7:555–581, 1992.
- [5] A. E. Brouwer and H. J. Veldman. Contractibility and NP-completeness. *Journal of Graph Theory*, 11(1):71–79, 1987.
- [6] Leizhen Cai. Fixed-parameter tractability of graph modification problems for hereditary properties. *Information Processing Letters*, 58:171–176, 1996.
- [7] Bruno Courcelle and M. Mosbah. Monadic second-order evaluations on tree-decomposable graphs. *Theoretical Computer Science*, 109:49–82, 1993.
- [8] Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michał Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015.
- [9] Marek Cygan, Dániel Marx, Marcin Pilipczuk, and Michał Pilipczuk. The planar directed k -vertex-disjoint paths problem is fixed-parameter tractable. In *54th Annual IEEE Symposium on Foundations of Computer Science, FOCS 2013, 26-29 October, 2013, Berkeley, CA, USA*, pages 197–206, 2013.
- [10] M. R. Fellows, J. Kratochvíl, M. Middendorf, and F. Pfeiffer. The complexity of induced minors and related problems. *Algorithmica*, 13(3):266–282, 1995.
- [11] Petr A. Golovach, Marcin Kaminski, Spyridon Maniatis, and Dimitrios M. Thilikos. The parameterized complexity of graph cyclability. *SIAM J. Discrete Math.*, 31(1):511–541, 2017.
- [12] Petr A. Golovach, Pim van ’t Hof, and Daniël Paulusma. Obtaining planarity by contracting few edges. *Theor. Comput. Sci.*, 476:38–46, 2013.
- [13] Martin Grohe, Ken-ichi Kawarabayashi, Dániel Marx, and Paul Wollan. Finding topological subgraphs is fixed-parameter tractable. In *Proceedings of the 43rd ACM Symposium on Theory of Computing, (STOC 2011)*, pages 479–488, 2011.
- [14] Martin Grohe, Ken-ichi Kawarabayashi, Dániel Marx, and Paul Wollan. Finding topological subgraphs is fixed-parameter tractable. In *STOC’11*, pages 479–488. ACM, 2011.

- [15] Bart M. P. Jansen, Daniel Lokshantov, and Saket Saurabh. A near-optimal planarization algorithm. In *Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '14, pages 1802–1811. SIAM, 2014.
- [16] Ken-ichi Kawarabayashi and Yusuke Kobayashi. The induced disjoint path problem. In *13th Conference on Integer Programming and Combinatorial Optimization, IPCO 2008*, volume 5035 of *Lecture Notes in Computer Science*, pages 47–61. Springer, Berlin, 2008.
- [17] Ken-ichi Kawarabayashi and Bojan Mohar. Graph and map isomorphism and all polyhedral embeddings in linear time. In *STOC'08*, pages 471–480. ACM, 2008.
- [18] Ken-ichi Kawarabayashi, Bojan Mohar, and Bruce A. Reed. A simpler linear time algorithm for embedding graphs into an arbitrary surface and the genus of graphs of bounded tree-width. In *FOCS'08*, pages 771–780. IEEE Computer Society, 2008.
- [19] Ken-ichi Kawarabayashi and Bruce A. Reed. Odd cycle packing. In *Proceedings of the 42nd ACM Symposium on Theory of Computing, STOC 2010*, pages 695–704, 2010.
- [20] Ken-ichi Kawarabayashi and Paul Wollan. A shorter proof of the graph minor algorithm: The unique linkage theorem. In *Proceedings of the Forty-second ACM Symposium on Theory of Computing*, STOC '10, pages 687–694, New York, NY, USA, 2010. ACM.
- [21] J.M. Lewis and C.H. Papadimitriou. The node-deletion problem for hereditary properties is np-complete. *J. Comput. System Sci.*, 20:219–230, 1980.
- [22] Dániel Marx. Chordal deletion is fixed-parameter tractable. *Algorithmica*, 57(4):747–768, 2010.
- [23] Dániel Marx and Ildikó Schlotter. Obtaining a planar graph by vertex deletion. *Algorithmica*, 62(3-4):807–822, 2012.
- [24] N. Robertson and P.D. Seymour. Graph minors .xiii. the disjoint paths problem. *Journal of Combinatorial Theory, Series B*, 63(1):65 – 110, 1995.
- [25] Neil Robertson and P. D. Seymour. Graph Minors. XX. Wagner’s conjecture. *Journal of Combinatorial Theory, Series B*, 92(2):325–357, 2004.
- [26] Neil Robertson and Paul D. Seymour. Graph Minors. XXI. Graphs with unique linkages. *J. Comb. Theory, Ser. B*, 99(3):583–616, 2009.
- [27] Neil Robertson and Paul D. Seymour. Graph minors. XXII. Irrelevant vertices in linkage problems. *J. Comb. Theory, Ser. B*, 102(2):530–563, 2012.
- [28] Detlef Seese. Linear time computable problems and first-order descriptions. *Mathematical Structures in Computer Science*, 6(6):505–526, 1996.
- [29] Paul D. Seymour and Robin Thomas. Graph searching and a min-max theorem for tree-width. *J. Comb. Theory Ser. B*, 58(1):22–33, 1993.