



HAL
open science

Proteomic and metabolomic profiling underlines the stage- and time-dependent effects of high temperature on grape berry metabolism

David Lecourieux, Christian Kappel, Stéphane Claverol, Philippe Pieri, Regina Feil, John E Lunn, Marc Bonneau, Lijun Wang, Eric Gomès, Serge Delrot, et al.

► To cite this version:

David Lecourieux, Christian Kappel, Stéphane Claverol, Philippe Pieri, Regina Feil, et al.. Proteomic and metabolomic profiling underlines the stage- and time-dependent effects of high temperature on grape berry metabolism. *Journal of Integrative Plant Biology*, 2020, 62 (8), pp.1132-1158. 10.1111/jipb.12894 . hal-03003936

HAL Id: hal-03003936

<https://hal.science/hal-03003936v1>

Submitted on 1 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Running Title:** Heat stress impacts grape berry proteome and metabolism

2

3 **Proteomic and metabolomic profiling underlines the stage- and time-**
4 **dependent effects of high temperature on grape berry metabolism**

5

6

7 **David Lecourieux^{1*}, Christian Kappel², Stéphane Claverol³, Philippe Pieri¹, Regina Feil⁴, John**
8 **E. Lunn⁴, Marc Bonneau³, Lijun Wang⁵, Eric Gomès¹, Serge Delrot¹ and Fatma Lecourieux⁶**

9

10

11 ¹ UMR1287 EGFV, Université de Bordeaux, INRA, Bordeaux Sciences Agro, ISVV, 33140 Villenave
12 d'Ornon, France.

13 ² Institut für Biochemie und Biologie, Universität Potsdam, D-14476 Potsdam, Germany

14 ³ Université de Bordeaux, Plateforme Protéome, Centre de Génomique Fonctionnelle Bordeaux, 33076
15 Bordeaux, France.

16 ⁴ Max Planck Institute of Molecular Plant Physiology, Wissenschaftspark Golm, Am Mühlberg 1,
17 14476 Potsdam-Golm, Germany

18 ⁵ Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China

19 ⁶ UMR1287 EGFV, CNRS, Université de Bordeaux, INRA, Bordeaux Sciences Agro, ISVV, 33140
20 Villenave d'Ornon, France.

21

22

23 *Corresponding author: david.lecourieux@inra.fr

24

25

26 **ABSTRACT**

27 Climate change scenarios predict an increase in mean air temperatures and in the frequency, intensity,
28 and length of extreme temperature events in many wine-growing regions worldwide. Because elevated
29 temperature has detrimental effects on the berry growth and composition, it threatens the economic
30 and environmental sustainability of wine production. Using Cabernet Sauvignon fruit-bearing cuttings,
31 we investigated the effects of high temperature (HT) on grapevine berries through a label-free shotgun
32 proteomic analysis coupled to a complementary metabolomic study. Among the 2279 proteins
33 identified, 592 differentially abundant proteins were found in berries exposed to HT. The gene
34 ontology categories “Stress”, “Protein”, “Secondary metabolism” and “Cell wall” were predominantly
35 altered under HT. High temperatures strongly impaired carbohydrate and energy metabolism, and the
36 effects depended on the stage of development and duration of treatment. Transcript amounts correlated
37 poorly with protein expression levels in HT berries, highlighting the value of proteomic studies in the
38 context of heat stress. Furthermore, this work reveals that HT alters key proteins driving berry
39 development and ripening. Finally, we provide a list of differentially abundant proteins that can be
40 considered as potential markers for developing or selecting grape varieties that are better adapted to
41 warmer climates or extreme heat waves.

42

43

44 **Keywords:** *Vitis vinifera*, berry, heat stress, proteomic, primary and secondary metabolisms

45

46

47

48 INTRODUCTION

49 Grapevine is economically the most important fruit crop in the world, providing dried fruits, table
50 grapes and the basis for wine making (www.oiv.int; www.fao.org). The combination of climate, soil
51 and viticultural practices in a given region constitutes a highly interactive system known as “terroir”
52 (Deloire et al. 2005). The interaction between “terroir” and the plant’s genotype (cultivar) significantly
53 affects vine development and berry composition, and thereafter determines wine quality and character
54 (Van Leeuwen and Seguin 2006). Grapevine requires suitable temperatures, radiation intensities and
55 duration, and water availability during the growth cycle for optimal fruit production and quality (Kuhn
56 et al. 2014). Because of on-going climate change, the environmental parameters in most wine
57 producing regions around the world are expected to change, potentially lowering productivity and
58 altering the traditional characteristics (“typicity”) of the wine from a given region (Schultz and Jones
59 2010). While some of these climatic constraints can be overcome, at least partially, by viticultural
60 practices, temperature remains more difficult to control under field conditions (Van Leeuwen et al.
61 2013). Climate change scenarios predict an increase in global mean temperatures between 1.5 and
62 5°C, during the 21st century (IPCC, 2014), in addition to an increase in the frequency, intensity, and
63 duration of extreme temperature events (Lobell et al. 2008). Considering that air temperature is a key
64 climate variable affecting grapevine phenology, berry metabolism and composition (Jones et al. 2005;
65 Martínez-Lüscher et al. 2016), global warming represents a serious threat for wine production and/or
66 typicity in many wine-growing regions worldwide (Fraga et al. 2016).

67 Temperatures above the optimum are sensed as heat stress by living organisms. Heat stress disturbs
68 cellular homeostasis and can lead to severe retardation in growth and development and even death
69 (Wahid et al. 2007). In grapevine, warming promotes vegetative growth and disturbs the plant’s
70 carbon balance (Greer and Weedon 2012), adversely affecting flower set and young berry
71 development (Greer and Weston 2010). HS also impacts fruit primary and secondary metabolism,
72 desynchronizing sugar and organic acid metabolism and delaying sugar and polyphenol accumulation
73 during ripening (Torregrosa et al. 2017; Gouot et al. 2018). As a consequence, heat waves occurring
74 during key berry development stages can dramatically affect the organoleptic properties of wine even
75 when more favourable weather conditions prevail for the remaining of the season.

76 In this context, the common viticultural practice of removing leaves around the fruit zone to
77 enhance aroma development and avoid fungal infections (Alem et al. 2018), might be questioned since
78 it affects light exposure and temperature of the berries. Because of the direct irradiative effects,
79 temperature within the fruit may differ very significantly from that of the whole plant or surrounding
80 air (Cola et al., 2009). The microclimate (mostly temperature and light conditions) strongly influences
81 the metabolite composition of berries, affecting flavonols, anthocyanins, amino acids and aroma
82 precursors (Downey et al. 2006; Pereira et al. 2006; Asproudi et al. 2016; Young et al. 2016; Reshef et
83 al. 2017). These effects can be explained, at least in part, by the substantial remodelling of the fruit
84 transcriptome observed under these conditions (du Plessis et al. 2017; Lecourieux et al. 2017).

85 In a previous report, we showed dramatic biochemical and transcriptomic changes in heat-exposed
86 berries, depending on both the developmental stage and the stress duration (Lecourieux et al. 2017).
87 Although transcriptomic analyses have offered comprehensive insights into the mechanisms
88 underlying the HS responses in grape (Liu et al. 2012; Carbonell-Bejerano et al. 2013; Rienth et al.
89 2014b; Jiang et al. 2017; Lecourieux et al. 2017), it is recognised that changes in transcript abundance
90 do not always correlate with the levels of the encoded proteins (Keller and Simm 2018). Furthermore,
91 proteins represent a more functional level of gene expression than transcripts because they have a
92 direct impact on metabolism and other cellular processes. Therefore, state of the art mass spectrometry
93 analysis of the proteome is a powerful tool to study molecular mechanisms and biological traits in
94 plants, including responses to abiotic stresses (Kosova et al. 2018). Using shotgun proteomic tools
95 such as iTRAQ (isobaric Tags for Relative and Absolute Quantitation) and label-free quantification
96 techniques, several proteomic studies have recently investigated the heat responses of grapevine leaves
97 (Liu et al. 2014; Jiang et al. 2017) or grape cell cultures (George et al. 2015).

98 To better understand the consequences of high temperature (HT), inherent to leaf removal, on berry
99 development and to improve our knowledge about the molecular mechanisms involved in grapevine
100 fruit response to HT, the present work combines transcriptomic, proteomic and metabolite analysis
101 using Cabernet Sauvignon fruit-bearing cuttings. A label-free proteomic approach was used to
102 investigate changes in the grape berry proteome resulting from direct exposure to elevated
103 temperature. The collected proteomic data were compared with our previous transcriptomic analysis
104 performed on the same berry samples (Lecourieux et al., 2017). These data showed a strong and
105 negative effect of HT on proteins related to carbohydrate and energy metabolism, which led us to
106 analyse the impact of HT on primary metabolites using tandem mass spectrometry (LC-MS/MS).
107 Finally, we focused on proteins accumulating in berries exposed to HT that are potentially involved in
108 thermotolerance. Altogether, this work helps to better understand the consequences of heat stress on
109 developing grape berries and provides a list of potential heat-tolerance molecular markers that could
110 be useful for breeders and viticulturists.

111

112 **RESULTS AND DISCUSSION**

113

114 **Quantitative analysis of heat stress responsive proteins in grape berries**

115 Experiments were performed on fruit-bearing cuttings of *V. vinifera* Cabernet Sauvignon as described
116 in Lecourieux et al. (2017). In brief, berries at three different stages of development were exposed to
117 elevated temperatures for 12 h (7:00 am to 7:00 pm) during each day of treatment and harvested for
118 analysis after 1 and 7 days of treatment. The average pulp temperature of the heat-treated berries was
119 about 8°C higher than in control berries. A label-free proteomic approach was performed to identify
120 proteins with differential abundance between control and heat-treated berry samples.

121 Using the parameters and filters described in the Material and Methods, the conducted label-free

122 experiment led to the identification of 3972 protein groups (from here on referred to as proteins) in the
123 combined dataset from the six conditions (three stages x two treatment durations), of which 2279
124 showed two or more unique peptides (**Table 1**). Protein abundance changes between control and heat-
125 exposed fruits were considered to be significant when a minimum fold change of 1.5 ($\log_2FC > 0.58$ or
126 < -0.58) was reached, with a significance threshold of $p < 0.05$. Only proteins identified in all three
127 biological replicates were considered for quantification. Among the quantified proteins with a
128 minimum of two unique peptides, 592 DAPs (differentially abundant proteins) displayed significant
129 differences in abundance between control and HT berries in at least one of the six conditions. The
130 entire dataset is available via ProteomeXchange (identifier PXD014693) and in Supplementary **Table**
131 **S1**. Further information on the 592 DAPs is provided in Supplementary **Table S2**. Among these 592
132 DAPs, 501 proteins were identified in only one condition (139 up-regulated and 362 down-regulated)
133 while the 91 remaining proteins were retrieved in more than one condition (16 up-regulated, 61 down-
134 regulated, and 14 proteins that were up-regulated in some conditions but down-regulated in others)
135 (**Table 1 and Table S2**). More DAPs were identified at the ripening stage (304 DAPs) than at the
136 green (146 DAPs) or veraison stages (222 DAPs). Whereas the numbers of up- and down-regulated
137 proteins were similar at veraison (115 up, 107 down), more down-regulated proteins were found when
138 the HT was applied at the green stage (24 up, 122 down) or ripening (49 up, 255 down) stages (**Figure**
139 **1A**). The DAPs identified after short (1 day) treatments were mostly different from those observed
140 after long (7 days) treatments. Indeed, only 1 (up), 9 (4 up, 5 down) and 18 (2 up, 16 down) common
141 DAPs between 1D or 7D of HT were observed at the green, veraison and middle ripening stages,
142 respectively (**Figure 1B and Tables S2, S3, S4 & S5**). Similarly, at all three stages, only one common
143 down-regulated DAP was identified after 1 day of HT (VIT_12s0057g01020, fasciclin-like protein),
144 with none being detected after 7 days of treatment (**Table S2**). The number of common DAPs per
145 stage increased when looking for proteins with modified abundance after 1 and/or 7 days treatment. In
146 this case, 7 DAPs (3 up, 4 down) were found in common for the three stages, and up to 31 DAPs were
147 obtained when comparing heat effects between green and veraison stages (**Figure 1C and**
148 **Supplementary Table 2**).

149 To determine the functional categories affected by HT, the identified DAPs were functionally
150 classified using MapMan BIN codes (Usadel et al. 2005) and a *V. vinifera* mapping file (Lecourieux et
151 al., 2017). The predominant proteins changed in berries in response to HT were assigned to “protein
152 metabolism” (114/592 DAPs, 19.3%), “stress” (54/592 DAPs, 9.1%), “secondary metabolism”
153 (31/592 DAPs, 5.2%) and “cell wall” (25/592 DAPs, 4.2%) (**Figure 2**). Except for the categories
154 “stress” (36 up, 18 down) and “RNA” (10 up, 9 down), all the others categories contained more down-
155 regulated proteins than up-regulated ones, with particularly asymmetrical changes in some categories,
156 such as “lipid metabolism” (0 up, 16 down) and “transport” (2 up, 17 down). No particular effect of
157 HT according to the developmental stage was observed on these functional categories, whereas
158 categories linked to carbohydrate and energy metabolism (glycolysis, tricarboxylic acid (TCA) cycle,

159 fermentation, gluconeogenesis, minor and major carbohydrate (CHO) metabolism) or related to
160 “photosynthesis”, “cell wall” and “redox” were mostly impacted during veraison and/or ripening
161 stages (**Supplementary Table 2**).

162

163 **Poor correlation between the transcriptome and proteome in heat-stressed berries**

164 Global analyses based on gene expression and protein profiles provide a powerful tool for
165 understanding how organisms respond to environment changes. To integrate proteomic data with
166 transcriptomic results, the present protein abundance data were compared to each corresponding
167 differentially expressed gene (DEG) identified in the same grape berry samples exposed to HT
168 (Lecourieux et al., 2017). Whereas 592 DAPs were retrieved in at least one of the six HT conditions
169 (**Table 1, Supplementary Table S1**), 6848 DEGs were identified using the same berry samples
170 (Lecourieux et al. 2017). A linear correlation between differentially expressed transcripts and the
171 corresponding proteins was established for 64 out of 592 DAPs (**Figure 4, Supplementary Table**
172 **S11**). Two types of correlation profiles were found. The first one corresponds to proteins and
173 transcripts that exhibit profiles with the same tendency (up- or down-regulation in response to HT) and
174 the second one includes proteins and transcripts that are inversely correlated for a same gene. The best
175 relationship between mRNA and protein levels (39 DAPs) was observed when HT was applied for one
176 day at veraison, and for proteins belonging to functional BINs associated to “heat stress” (25
177 members), “hormone” (4), and “secondary metabolism” (5) (**Figure 4, Supplementary Table S11**).

178 Our results underline the poor correlation between the proteome and the transcriptome in HT
179 berries, with only 11% (64/592 DAPs) of the differentially abundant proteins that were also impacted
180 at the mRNA expression level. This is not unexpected since HT leads to denaturation and aggregation
181 of proteins, and because translation, especially under elevated temperature, is governed by a variety of
182 regulatory mechanisms independent of transcription rate (Bokszczanin and Fragkostefanakis 2013).
183 These include the impacts of miRNAs on mRNA stability and translatability, mRNA half-life, global
184 and transcript specific translational rates, as well as protein turnover. Any lag between the synthesis of
185 a gene transcript and its translation into a protein could also give rise to an apparent discrepancy
186 between transcript and protein abundance at a given sampling time (Keller and Simm 2018). Last but
187 not least, the poor correlation between transcript and protein levels in heat-treated berry samples may
188 also be due to alternative splicing (AS) events. Increasing evidence has shown that AS is a critical
189 post-transcriptional event and plays an important role in plant stress responses (Baurle 2016).
190 Particularly, many genes are under transcriptional regulation by the circadian clock, and several
191 circadian clock genes show AS in response to even small changes in temperature (James et al. 2018).
192 In grapevine, AS changes were observed for nearly 70% of the genes in leaves exposed to
193 temperatures above 35°C. In the same study, only 64 out of the 808 identified DAPs were retrieved as
194 differentially expressed at the transcript level (Jiang et al. 2017), that fits well with the weak
195 transcript/protein correlation observed in our work. Jiang et al. (2017) suggested that AS changes

196 under high temperature could provide molecular plasticity allowing the plants to adapt to stress
197 conditions. Using *Arabidopsis* (*Arabidopsis thaliana*) plants exposed or not to a non-lethal HT,
198 referred to as priming, Ling et al. (2018) have recently demonstrated that AS functions as a novel
199 component of heat-shock memory.

200 Despite the generally poor correlation between individual transcripts and proteins levels in berries
201 exposed to HT, there was a much better correspondence at the functional category level, with
202 transcripts and proteins assigned to the “stress”, “protein metabolism” and “secondary metabolism”
203 MapMan BINs showing consistent responses as a group (**Supplementary Figure S1**). We also
204 noticed that, at the transcript level, most of the functional categories were differentially affected
205 according to the developmental stage and the stress duration whereas it was less obvious at the protein
206 level. The best relationship between transcripts and proteins levels was observed with genes related to
207 the category “stress”, and especially to the “heat” sub-category. Among the 30 HSPs whose protein
208 abundance increased in response to HT, 25 were also up-regulated at the transcript level
209 (**Supplementary Table S12**). Most proteins are inherently thermo-labile, with some becoming
210 denatured at the temperatures that grapevines experience during a heat wave. In addition, high
211 temperature is often associated with oxidative stress leading to production of reactive oxygen species
212 that irreversibly damage proteins. Maintaining proteins in stable and functional conformations and
213 preventing their aggregation are essential for survival of cells under high temperature. HSPs function
214 as molecular chaperones in maintaining protein quality and folding, and are required for the
215 acquisition of plant thermotolerance (Bokszczanin and Fragkostefanakis 2013). It is well established
216 that transcription factors from the HSF family regulate the heat stress response in plants, through a
217 massive up-regulation of *HSP* genes (Scharf et al. 2012). However, as transcription factors are low
218 abundance proteins they largely escaped proteomic detection, and no HSF was retrieved in the present
219 study despite the observed transcriptional up-regulation of six *VviHSF* genes in heat-exposed berries
220 (Lecourieux et al. 2017).

221 In good agreement with our transcriptomic data, the protein abundance of the functional category
222 “Protein metabolism” was strongly impacted with nearly 20% of the identified DAPs that belong to
223 this category being affected, especially those belonging to the sub-categories “protein synthesis” (34
224 DAPs) and “protein degradation” (55 DAPs) (**Supplementary Table S6**). However, the HT triggered
225 a strong down-accumulation effect on the “protein metabolism” category (92 out of the 114 DAPs), in
226 contrast with what was observed on transcripts (333 up- and 307 down-regulated genes). A number of
227 studies demonstrated that different abiotic stresses, including heat, result in a general inhibition of
228 protein translation (Yanguéz et al. 2013). Accordingly, we observed a significant decrease in
229 abundance of many proteins related to protein synthesis including various ribosome structural proteins
230 and several translation initiation factors (**Supplementary Table S6**). Whereas inhibiting translation
231 under stress helps to reduce energy consumption, certain specific mRNAs are selectively translated to
232 produce relevant proteins involved in the proper establishment of the stress adaptation process. By

233 studying the translome of Arabidopsis seedling exposed to HT, Yanguéz et al. (2013) observed that
234 regulation of gene expression at the translational level is superimposed on regulation at the
235 transcriptional level, and often reinforces the response of individual genes/proteins to heat stress.
236 Protein translation is accomplished by the combined actions of the ribosomes and ancillary proteins
237 such as initiation, elongation, and termination factors. Yanguéz et al. (2013) showed that translation
238 could be regulated at the initiation phase in Arabidopsis seedlings exposed to HT, even if the
239 mechanisms involved in this regulation remain unclear. To date, there is some evidence for the role of
240 different plant translation initiation factors (eIFs) in various abiotic stresses responses (Dutt et al.
241 2015). Our proteomic data highlighted the increase in abundance of four eIFs upon HS, namely eIF1A
242 (VIT_15s0046g01500), two isoforms of eIF3 (VIT_14s0060g01630, VIT_00s0880g00020) and eIF5
243 (VIT_14s0006g01990), whereas others eIF isoforms were decreased (**Supplementary Table S6**).
244 eIF5 (VIT_14s0006g01990) showed the highest accumulation score (Log₂FC 5.22) within this
245 functional category. The importance of eIF5 proteins in thermotolerance was highlighted in pea
246 (Suragani et al. 2011) and Arabidopsis (Xu et al. 2011; Zhang et al. 2017).

247

248 **Heat stress alters key proteins driving berry development and ripening**

249

250 *Heat stress impacts the secondary metabolism in developing berry*

251 The strong effects of HT observed on transcripts assigned to the “secondary metabolism” functional
252 category (291 DEGs across the six conditions, (Lecourieux et al. 2017)) were confirmed at the protein
253 level (31 out of the 592 DAPs), albeit to a lesser extent (**Supplementary Table S8**). These 31 DAPs
254 (11 up, 20 down) mainly belonged to the subcategories: “isoprenoids” (7 DAPs), “phenylpropanoid-
255 lignin” (12) and “flavonoids” (7).

256 HT altered expression of numerous genes related to aromatic potential, including genes encoding
257 biosynthetic enzymes for terpenes and methoxypyrazines (Lecourieux et al. 2017), but the proteomic
258 data from the current study suggested that these changes in transcript abundance were not translated
259 through to the protein level. A few proteins related to isoprenoid subfamily were down-accumulated
260 under HT including a 3-ketoacyl-CoA synthase (KCS10, VIT_04s0008g02250), which is involved in
261 the synthesis of very long chain fatty acids that are incorporated into plant lipids (precursors of
262 cuticular waxes), and several enzymes related to the synthesis of precursors of terpenoids
263 (**Supplementary Table S8**). By contrast, a carotenoid cleavage dioxygenase (CCD1,
264 VIT_13s0064g00810) was found more abundant in stressed berries at veraison. Expression of various
265 *VvCCD* genes was reported to increase towards berry ripening and enzymes in this family were
266 demonstrated to promote desirable flavor and aroma compounds in grapes and wine (Mathieu et al.
267 2005; Rienth et al. 2014a; Alem et al. 2018). The putative role of CCD proteins in heat responses is
268 still unknown but these proteins were suggested to enhance tolerance to various abiotic stresses
269 through their enzymatic products (e.g. apocarotenoids) acting as stress signals (Havaux 2014). For

270 instance, some members of the CCD family are involved in generating carotenoid-derived
271 phytohormones, such as strigolactones known to be involved in various plant developmental and
272 adaptation processes (Jia et al. 2018).

273 It is well known that extreme high temperatures significantly affect the phenylpropanoid content of
274 grape berries, with the impact being dependent on the intensity and duration of the HS as well as the
275 developmental stage of the fruit (Gouot et al. 2018). Consistent with changes observed at the transcript
276 level (Lecourieux et al. 2017), HT triggered an enhanced abundance of proteins involved in lignin
277 synthesis, including shikimate *O*-hydroxycinnamoyltransferase (HCT; VIT_09s0018g01190),
278 cinnamyl alcohol dehydrogenase (CAD; VIT_03s0180g00260, VIT_02s0025g03100,
279 VIT_00s0615g00030) and caffeic acid *O*-methyltransferase (CCoAOMT1; VIT_16s0098g00850). A
280 significant increase in lignin biosynthesis would confer additional strength to lignified secondary cell
281 wall, helping berries to protect against shriveling mediated-HT. For instance, the activation of the
282 lignin pathway is a reaction to advanced stages of dehydration in grape berry fruit (Zamboni et al.
283 2010). Key enzymes of the general phenylpropanoid pathway (phenylalanine ammonia-lyase PAL,
284 VIT_06s0004g02620, VIT_08s0040g01710, VIT_13s0019g04460; cinnamate 4-hydroxylase C4H,
285 VIT_06s0004g08150; 4-coumarate-CoA ligase 4CL; VIT_11s0052g01090), enzymes from the
286 flavonoid biosynthesis pathway (dihydroflavonol 4-reductase DFR, VIT_18s0001g12800; flavonoid
287 3'-hydroxylase F3'H, VIT_17s0000g07210; flavonoid 3',5'-hydroxylase F3'5'H,
288 VIT_06s0009g02860; flavanone 3'-hydroxylase F3H, VIT_18s0001g14310; isoflavone reductase,
289 VIT_03s0038g04700) as well as enzymes mediating the glycosylation of flavonoids (UDP-glucosyl
290 transferases, VIT_12s0055g00290, VIT_02s0033g00130, VIT_17s0000g04750; UDP-rhamnosyl
291 transferase, VIT_00s0218g00150) were also affected at the protein level upon HT. Most of these
292 proteins were less abundant in heated berries, suggesting a sharp decline of the corresponding
293 enzymatic activities (**Supplementary Tables S2 & S8**). This lower amount of key enzymes probably
294 contributes to the reduction of flavonoid synthesis in grape berries exposed to heat. However, other
295 concomitant processes may explain the decrease of flavonoid contents in warmed berries, including
296 reduced biosynthetic enzyme activities due to non-optimal temperature, enzymatic oxidation or
297 chemical degradation after ROS scavenging (Gouot et al. 2018). Upon their biosynthesis in the
298 cytosol, flavonoids are rapidly accumulated into vacuoles or other cellular compartments through
299 vesicle trafficking, membrane transporters or glutathione S-transferase (GST)-mediated transport
300 (Zhao 2015). In addition to transport of flavonoids into vacuoles mediated by ATP-binding cassette
301 (ABC) transporters (Francisco et al. 2013), there is also H⁺-dependent transport mechanism in
302 grapevine (Gomez et al. 2011; Kuang et al. 2019). Vacuolar transport of anthocyanins and other
303 metabolites, such as malate, directly or indirectly depends on the transmembrane pH gradient
304 generated by both tonoplast H⁺-ATPases and H⁺-pyrophosphatases (V-PPase) (Shiratake and
305 Martinoia 2007). Three V-ATPase subunits (VIT_03s0038g00790, VIT_18s0001g01020,
306 VIT_04s0008g02460) and two V-PPases (VIT_11s0118g00350, VIT_09s0002g07880) showed strong

307 decrease in protein abundance in HT berries, and this may impair the efficient transport of
308 anthocyanins into the vacuole. Finally, the translational repression of several GST proteins
309 (VIT_01s0026g01340, VIT_01s0011g01900, VIT_08s0040g03100, VIT_19s0093g00310,
310 VIT_07s0104g01800) observed in heat-stressed berries during ripening could have an impact on fruit
311 secondary metabolism as well (Perez-Diaz et al. 2016).

312

313 *Harmful effects of heat stress on carbohydrate and energy berry metabolism*

314 A drastic negative effect of HT was observed for proteins assigned to primary carbohydrate
315 metabolism (**Supplementary Table S9**). Almost all the DAPs related to the following processes were
316 under-accumulated after HT: major CHO metabolism (6 out of 8 DAPs), glycolysis (8 out of 9
317 DAPs), fermentation (4 out of 5 DAPs), gluconeogenesis (3 out of 4 DAPs), oxidative pentose
318 phosphate pathway (6 DAPs) and TCA cycle (13 out of 14 DAPs). To better understand the
319 consequences of this HT repressive effect, and to expand beyond the previously reported metabolite
320 changes (Lecourieux et al., 2017), we measured an additional 31 metabolites that are involved in sugar
321 or organic acid metabolism, or respiration. To put the measured metabolites into a metabolic context, a
322 schematic representation of plant central carbon metabolism was created to display the metabolite
323 profiles under HT compared to control conditions, for each developmental stage and each stress
324 duration (**Figure 3, Supplementary Table S10**). A significant effect of the developmental stage (two-
325 way ANOVA, $p < 0.05$) was observed for most quantified metabolites (23 out of 31), in agreement
326 with the previously reported metabolite profiles in developing berries from Cabernet Sauvignon fruit-
327 bearing cuttings (Dai et al. 2013). HT significantly influenced the content of 14 metabolites including
328 shikimate, glycerate, glycerol-3-P (G-3-P), several glycolytic intermediates (fructose 6-phosphate
329 (Fru6P), 3-phosphoglycerate (3-PGA) and pyruvate), fumarate, and several sugar phosphates (glucose
330 6-P (G6P), trehalose 6-P (T6P), sucrose 6'-P (S6P) and mannose 6-P (M6P)). The well-known
331 accelerated decrease of malate under elevated temperature was also observed although the changes
332 were below the level of significance in an ANOVA test. However, when considered over the 14-day
333 time frame of the experiments, the decrease in malate was statistically significant according to a t-test
334 (**Figure 3**). This agrees with the significantly lower malate content detected previously in HT berries
335 at harvest (Lecourieux et al., 2017).

336 Metabolism undergoes profound reprogramming during fleshy fruit development (Beauvoit et al.
337 2018). The contents of primary and secondary metabolites have been investigated in developing grape
338 berries (Dai et al. 2013; Degu et al. 2014; Cuadros-Inostroza et al. 2016; Wang et al. 2017). Using
339 Cabernet Sauvignon fruit-bearing cuttings as in the present work, Dai et al. (2013) conducted a deep
340 analysis of metabolites from central carbon metabolism using an LC-MS/MS approach. They reported
341 the timing of important switches in primary carbohydrate metabolism during grape berry development
342 and discussed the relationship between these metabolite changes and transcriptional and translational
343 modifications within the fruit. Similarly, Wang et al. (2017) reported close correlations between the

344 metabolome and the proteome at the interface of primary and secondary metabolism in developing *V.*
345 *vinifera* Early Campbell fruits. Down-regulation of carbohydrate production and protein synthesis is a
346 conserved response of plants to abiotic stresses (Cramer et al. 2011). This may be a way for plants to
347 save energy, and reflects a shift from plant growth to mechanisms of protection. However, with the
348 exception of organic acids, the effects of high temperature on berry primary metabolism in general are
349 still poorly described (Serrano et al. 2017). It is well known that warm climatic conditions enhance the
350 degradation of malate in grapevine fruits, leading to a decrease in total acidity (Sweetman et al. 2014).
351 Sugar metabolism is desynchronized from organic acid metabolism in ripening berries at high
352 temperatures (Rienth et al. 2016). The micro-environment plays an active role in controlling the
353 dynamics of primary metabolite accumulation that affect fruit composition. For instance, heating
354 strongly alters the berry concentration of various amino acids, including phenylalanine and γ -
355 aminobutyric acid, and decreases malate content (Lecourieux et al. 2017). Integrating microclimatic
356 parameters and metabolomic analysis, Reshef et al. (2017) reported that solar irradiance affected the
357 overall levels and patterns of accumulation of sugars, organic acids, amino acids and
358 phenylpropanoids, across the grape cluster. The relationship between the diurnal microclimatic
359 changes (radiation and temperature) and the berry metabolic dynamics was recently described (Reshef
360 et al. 2019). The present study provides additional evidence about the harmful effects of heat on berry
361 primary metabolism through the quantification of 31 metabolite intermediates related to sugar
362 accumulation, glycolysis, and the TCA cycle, as well as through the identification of numerous
363 proteins related to the corresponding metabolic pathways whose abundance is affected by HS. Among
364 the 31 metabolites quantified, 14 were significantly impacted upon HT, with reduced contents in
365 warmed berries for all of these (**Figure 3, Supplementary Table S10**). Some intermediates were
366 more affected when the HT was applied during the green stage (glycerate, fumarate), and others
367 during ripening (T6P and S6P), whereas the remaining ones were negatively impacted at all three
368 developmental stages (pyruvate, shikimate, G3P, 3-PGA, G6P, Man6P, F6P). Reduced malate
369 contents were also observed under HS, including the green stage (**Figure 3**). These primary metabolite
370 profiles coincide with a dramatic down-regulation of numerous enzymes related to carbohydrate and
371 energy metabolism in heat-treated berries (48 out of 56 DAPs, **Supplementary Table S9**), and more
372 than 70% of these DAPs were repressed during the ripening period. Using a Cabernet Sauvignon cell
373 suspension culture subjected to HS (42°C), George et al. (2015) observed a similar decline in proteins
374 related to sugar metabolism. Our results highlighted changes in many metabolic pathways in grape
375 berries exposed to heat, possibly due to the inhibitory effect of elevated temperature on metabolic
376 enzyme activities from these pathways.

377 Several enzymes from the pathways of sugar and sugar-phosphate metabolism were less abundant
378 in heated berries, including a fructokinase (FRK, VIT_14s0006g01410) and two hexokinases (HXK,
379 VIT_18s0001g14230, VIT_11s0016g03070). The decrease in these enzymes may contribute to the
380 lower concentration in sugar phosphates (T6P, S6P, G6P and F6P) under heat, since the hexoses

381 (glucose and fructose) resulting from cleavage of imported sucrose can be converted to hexose
382 phosphates by FRK and HXK in the cytosol (Stein and Granot 2018). In plants, trehalose 6-phosphate
383 (T6P) acts as a signal of sucrose availability connecting plant growth and development to its metabolic
384 status, and it was reported that T6P levels can exhibit dynamic responses to environmental cues
385 (Figueroa and Lunn 2016). An increase in T6P content was observed in berries from veraison on, but
386 this increase was lower under heat (**Figure 3, Supplementary Table S10**). In some plants, T6P
387 inhibits the SnRK1 Ser/Thr protein kinase, in turn contributing to a reduction in anthocyanin content
388 (Baena-Gonzalez et al. 2007). It is thus tempting to speculate that the increase in T6P concentration
389 from veraison onwards contributes to anthocyanin accumulation through repression of SnRK1, and
390 that this regulatory process is less effective under heat. However, the roles of T6P as a signalling
391 molecule and the SnRK1/T6P pathway during the berry ripening process and in HT response remain
392 to be demonstrated. In sink organs such as developing berries, sucrose cleavage is carried out either by
393 invertase (INV) to yield glucose and fructose, or by sucrose synthase (SUS) to yield UDPG and
394 fructose. Our proteomic analysis revealed a strong accumulation of a SUS isoform after HT at
395 veraison (SUS2, VIT_05s0077g01930). However, glucose, fructose and sucrose contents were not
396 differentially altered by heat treatment (**Figure 3, Supplementary Table S10**), whereas a delay in the
397 increase in total soluble solids (TSS) content was observed in developing berries after heat exposure
398 during the green stage (Lecourieux et al. 2017). Accumulation of SUS may be a sign of hypoxia
399 resulting from increased respiration rate under high temperature conditions, thus consuming oxygen
400 faster than it can diffuse into the bulky grape berry tissue. Such as *AtSUS1* and *AtSUS4* in
401 Arabidopsis, *SUS* genes are described as hypoxic responsive genes and a role for sucrose synthase as
402 part of the acclimation mechanism to anoxia in dicots was proposed (Santaniello et al., 2014).

403 The present study also revealed the lower abundance of many glycolytic enzymes (9 DAPs) in
404 warmed berries, mostly during ripening, together with the reduced content of some glycolytic
405 intermediates (Fru6P, 3-PGA, pyruvate). A similar decrease in abundance of glycolytic enzymes or
406 glycolytic intermediates was also reported for some grape varieties including Cabernet Sauvignon,
407 when berries ripen under normal growth conditions (Giribaldi et al. 2007; Dai et al. 2013; Martinez-
408 Estes et al. 2013). Thus, the consequence of heat stress on the glycolytic pathway remains unclear
409 since the glycolytic flux might have been maintained or even increased at high temperature although
410 the decrease of some glycolytic-related DAPs. Disturbing the glycolytic pathway may impair the
411 generation of energy (ATP), reducing equivalents (NADH) and various intermediates required for the
412 production of amino acid, lipids and secondary metabolites. For example, it has been reported that cell
413 energy, reducing power and α -ketoglutarate availability are potential drivers for flavonoid
414 biosynthesis in grape cells (Soubeyrand et al. 2018).

415 The TCA cycle is a central metabolic hub since it generates energy, reducing power and carbon
416 skeletons. Several enzymes from the TCA cycle were repressed in heated berries but consequences on
417 the corresponding TCA intermediates were not obvious within the time frame of the analysis. The

418 transcript levels of MDH, which catalyzes a reversible reaction between oxaloacetate and malate, and
419 ME which oxidizes malate to pyruvate, were reported to increase during grape berry ripening. This
420 higher expression might contribute to the decline of malate concentration after veraison (Martinez-
421 Estes et al., 2011; Wang et al., 2017). In our study, however, heat exposure reduced protein amount of
422 both enzymes in ripening fruits (MDH: VIT_10s0003g01000, VIT_19s0014g01640,
423 VIT_00s0373g00040; ME: VIT_15s0046g03670, VIT_11s0016g03210), therefore questioning their
424 precise role in hastening the malate decrease under heat. The amount of a phosphoenolpyruvate
425 carboxylase (PEPC, VIT_19s0014g01390) was also reduced by HT at the green stage. This may
426 contribute to lower the amount of malate under heat before veraison (**Figure 3**), since the first step of
427 anaplerotic malate synthesis involves the carboxylation of PEP by PEPC (Sweetman et al. 2009).
428 However, the loss of malate during ripening has been attributed to increased degradation rather than
429 decreased synthesis pre-veraison (Ruffner et al. 1976; Sweetman et al. 2014). Accordingly, PEPC
430 activity decreases in pre-veraison berries with day heating, and malate content did not correlate with
431 changes in PEPC activity (Sweetman et al. 2014). The PEPC activity is highly regulated by reversible
432 protein phosphorylation and mono-ubiquitination, which affects its sensitivity to feedback inhibition
433 by malate (Ruiz-Ballesta et al. 2016). Thus, changes in the phosphorylation and/or the ubiquitination
434 status of PEPC might be more important than changes in total PEPC protein abundance. Energy-
435 consuming processes such as gluconeogenesis may also be impacted in ripening berries exposed to
436 HT, with three down-regulated enzymes (MDH, PEPCK and isocitrate lyase). Lower MDH amounts
437 would reduce the accumulation of oxaloacetate (OAA), required for PEP formation through
438 phosphoenolpyruvate carboxykinase activity (PEPCK). In normal growth conditions, both PEPCK
439 activity and transcript abundance are increased in berries after veraison (Walker et al. 2015). In
440 ripening grapes, PEPCK may function in gluconeogenesis when malate released from the vacuole
441 exceeds the demand from other malate-consuming processes (Walker et al. 2015).

442

443 *Heat stress promotes changes in lipid metabolism and cell wall remodelling*

444 The functional category “Lipid metabolism” was particularly affected in heat-exposed berries since all
445 the lipid-related proteins (20 DAPs) identified in our work were negatively regulated after HT
446 (**Supplementary Table S2**). Moreover, we also observed that warmed berries contain lower amount
447 of glycerol 3-phosphate, precursor of the membrane glycerolipids (**Figure 3**). These lipid-related
448 DAPs are involved in fatty acid (FA) synthesis, FA elongation or FA degradation, among which were
449 the following enzymes: acetyl-CoA carboxylase (ACC, VIT_18s0001g04980, VIT_11s0065g00360),
450 glycerol-3-phosphate acyltransferase (GPAT, VIT_15s0046g02400), beta-ketoacyl-CoA synthase
451 (KCS, VIT_13s0067g03890) and phosphoinositide phosphatase (SAC, VIT_04s0044g00030). ACC
452 proteins catalyse the carboxylation of acetyl-CoA to form malonyl-CoA during fatty acid synthesis
453 (Sasaki and Nagano 2004). The cytosolic ACCase contributes to the synthesis of very-long chain fatty
454 acids, with a role in membrane stability and biosynthesis of flavonoids. Its lower abundance in

455 warmed berries may negatively impact the flavonoid accumulation upon veraison. The heat-repressive
456 effects observed at the protein level may also lead to a deep remodelling of the berry protection
457 system. Particularly, the down-regulation of some proteins such as GPAT and KCS may disturb the
458 berry protective barrier, namely the cuticle, since these two proteins belong to families involved in
459 cutin and wax biosynthesis, respectively (Guo et al. 2016; Petit et al. 2016). The strongest translational
460 repression was observed for a protein from the SAC (suppressor of actin) family (\log_2FC : -6.22) when
461 the HS was applied during the green stage. SAC proteins are phosphatases that hydrolyze
462 polyphosphoinositides, and roles in stress responses, cell wall formation, vacuolar trafficking and
463 vacuolar morphology and function have been demonstrated in plants (Novakova et al. 2014). Among
464 the known effects of high temperatures, one of the most relevant is the alteration of membrane-linked
465 processes due to modifications in membrane fluidity and permeability (Niu and Xiang 2018). Heat
466 stress decreases the total content of lipids and glycolipids (Escandon et al. 2018). To preserve
467 membrane integrity under elevated temperatures, plants can regulate the ratio of bilayer to non-bilayer-
468 forming lipids (Welti et al. 2007). Plant thermotolerance might also be improved by reducing the
469 membrane content of unsaturated fatty acids or by accumulating more saturated fatty acids prior to
470 HS, thus helping to maintain optimal membrane fluidity (Higashi et al. 2015; Escandon et al. 2018).
471 Accordingly, we found that two desaturase proteins (VIT_05s0094g00750, VIT_00s0499g00030)
472 were down-regulated in warmed berries. Membrane components may also be involved in initiating
473 key signalling pathways that lead to the appropriate plant responses to heat (Niu and Xiang 2018).
474 Consequences of the heat-induced lipid metabolism changes for berry development and quality remain
475 to be determined.

476 The cell wall forms a dynamic boundary for plant cells and plays important roles during grape
477 berry development, governing the potential for cell expansion during fruit growth and modulating the
478 texture of mature berries through the depolymerisation and the solubilisation of cell wall polymers
479 (Goulao et al. 2012b). Cell wall remodelling also constitutes an important component of plant
480 responses to HS that maintain overall function and growth, although the cell wall factors contributing
481 to the acquisition of plant thermotolerance remain largely unknown (Wu et al. 2018). Our previous
482 transcriptomic analysis highlighted the strong effect of HT on genes involved in cell wall homeostasis,
483 and especially on enzymes contributing to cell wall expansion and loosening such as xyloglucan endo-
484 transglycosylase and expansins (Lecourieux et al. 2017). At the protein level, most of the 25 DAPs
485 related to the category “cell wall” displayed lower amounts in heated berries, except for a few proteins
486 among which were two pectin methylesterases (PME, VIT_07s0005g00730, VIT_11s0016g00290)
487 and two polygalacturonases (PG, VIT_02s0025g01330, VIT_08s0007g08330) that accumulate in
488 ripening berries exposed to heat. Both PME and PG are implicated in pectin disassembly and fruit
489 softening process, but it is difficult to assess the impact of changes in these four proteins, since both
490 enzymes belong to large families, with 36 PMEs and 60 PGs being annotated in the reference grape
491 genome (Goulao et al. 2012a). However, there is increasing evidence that cell wall-modifying

492 enzymes are involved in plant responses to HS, and PMEs appear to be key players in the process (Wu
493 et al. 2018). Similar observations were made in fruits of other species exposed to high temperature,
494 such as strawberry (Langer et al. 2018).

495

496 **Identification of proteins with a potential role in berry responses to heat**

497 Global warming represents a substantial risk for the sustainability of wine production in many regions.
498 Plant breeding combined with molecular plant biotechnology has the potential to deliver stable yields
499 and quality under warmer climate conditions, but this requires the identification of key traits of
500 tolerance to heat (Torregrosa et al. 2017). Our proteomic study revealed 54 DAPs related to the
501 functional category “Stress” and mainly belonging to the “abiotic/heat stress” cluster (**Supplementary**
502 **Table S7**; 35 out of the 54 “stress” DAPs). Most of the proteins associated with the “heat stress”
503 category were over-accumulated, mainly at veraison stage, and all were members of the heat shock
504 protein (HSP) family. Looking for potential protective mechanisms activated in berries exposed to HT,
505 we paid particular attention to up-regulated proteins previously described in the HS responses and
506 thermotolerance in other plant species (**Table 2**; **Supplementary Tables S7 and S12**). We also listed
507 some proteins that increased in abundance under HT and thus have a putative role in stress tolerance.
508 These proteins (**Table 2**) are potential markers for developing or selecting grape varieties that are
509 better adapted to warmer climate or the extreme heat waves that are expected due to climate change.

510 Cabernet Sauvignon berry can improve its HS response abilities by elevating the levels of
511 particular proteins associated with the maintenance of protein homeostasis (synthesis, folding and
512 degradation). HSP/chaperones are considered as powerful buffers against environmental stress
513 including HS. Their chaperone functions include not only protein folding, but also signalling and
514 protein targeting and degradation (Bokszczanin and Fragkostefanakis 2013). Data collected over the
515 last decades suggest that members of the HSP/chaperone network are valuable targets for engineering
516 multiple stress resistance in crops (Jacob et al. 2017). Among the 30 up-regulated HSPs in heat-
517 exposed berries, four of these were increased irrespective of the fruit developmental stage (HSP70,
518 HSP90, HSP101, HSP17.6II), and all of these were described as required for thermotolerance in
519 various plant species (**Table 2**). Another up-regulated protein under HS corresponds to a Bcl-2-
520 associated athanogene isoform (BAG). Plant BAGs are multifunctional proteins that regulate cell
521 development and contribute to cytoprotection from pathogen attack and abiotic stress (Doukhanina et
522 al. 2006). In particular, AtBAG7 was described to be a key player of the heat-induced unfolded protein
523 response (UPR) pathway that alleviates the detrimental effects of accumulation of misfolded proteins
524 (Li et al. 2017). VvROF1 abundance was also increased in HS conditions. In Arabidopsis, AtROF1
525 encodes a peptidyl prolyl isomerase catalysing the *cis/trans* isomerization of peptide bonds N-terminal
526 to proline residues within polypeptide chains. AtROF1 is transcriptionally induced by HT and is
527 involved in acquired thermotolerance through the interaction with HSP90.1 and HSFA2 (Meiri and

528 Breiman 2009). We also observed that two thaumatin-like proteins (VvTLPs) from the osmotin sub-
529 family strongly accumulated in heat stressed berries. The grape genome contains 33 VvTLPs (Yan et
530 al. 2017), and this family is known to be involved in biotic and abiotic stress responses including
531 grapevine (Liu et al. 2010). Osmotins have been shown to have multiple functions in enhancing stress
532 tolerance in many plant species, although the mechanisms by which osmotins mediate plant response
533 to abiotic stress are not well established (de Jesus-Pires et al. 2019). Finally, a strong increase in the
534 translation initiation factor VveIF5 was detected in heated berries (**Table 2**). The role of eIF5 in
535 thermotolerance was described in Arabidopsis and pea (Suragani et al., 2011; Zhang et al., 2017).

536 How plants sense and transduce the HT signal is still an important topic to be addressed. It was
537 reported that phytochromes function as plant thermosensors (Jung et al. 2016). Transduction of the HT
538 signal involves various players (Li et al. 2018), including members of the plasma membrane cyclic
539 nucleotide gated calcium channel family (CNGC). In Arabidopsis, CNGC2 and CNGC6 were
540 described as heat sensors, contributing to the Ca²⁺ signalling induced by HT and leading to the onset
541 of plant acquired thermotolerance (Finka et al. 2012). Whereas *CNGC6* gene is ubiquitously expressed
542 in Arabidopsis (Gao et al. 2012), our results revealed that the protein abundance of the grape ortholog
543 VvCNGC2 increased after berry exposure to heat, underlining the need for *de novo* synthesis of this
544 protein to achieve its role in thermotolerance.

545 Integration of elevated temperature, signal transduction and the appropriate stress response is
546 mediated, at least partially, by plant hormones (Ku et al. 2018). Several proteins related to hormone
547 homeostasis were seen to be more abundant in heat-exposed berries (**Table 2**). Four abscisic acid
548 (ABA)-related DAPs were up-regulated in the present study. The first DAP is a 9-cis-epoxycarotenoid
549 dioxygenase (NCED). This enzyme catalyses the first committed step in ABA biosynthesis (Tan et al.
550 1997) and NCED overexpression improves drought tolerance in grape (He et al. 2018). At the
551 transcript level, a mixed picture has emerged for the *NCED* isogenes in HS grape berries (Carbonell-
552 Bejerano et al. 2013; Rienth et al. 2014a; Lecourieux et al. 2017). The second ABA-related DAP was
553 GER5/GRE5, a protein closely related to GEM (GLABRA2 Expression Modulator) and belonging to
554 a subfamily of the GRAM domain-containing proteins (Mauri et al. 2016). In heat-stressed berries,
555 GER5 was up-regulated at both mRNA and protein levels (**Supplementary Table S11**). GER5 and
556 closely related GRAM domain genes (GER1, GEM) were proposed to be required in the reproductive
557 development of Arabidopsis whereas a putative role in abiotic stress responses remains unclear
558 (Bhatta et al. 2018). The third ABA-related DAP is an ABA-, stress-, and ripening- (ASR) induced
559 protein known to play prominent roles in the protection of plants against abiotic stress (Gonzalez and
560 Iusem 2014). For instance, the grape ASR VvMSA was reported to be involved in responses to water
561 deficit (Medici et al. 2014). The last ABA-related DAP is RACK1 (receptor for activated C kinase 1).
562 In rice, this protein positively regulates seed germination by controlling endogenous levels of ABA
563 and H₂O₂ (Zhang et al. 2014). More broadly, plant RACK1 proteins regulate various signalling
564 pathways ranging from developmental processes such as seed germination, flowering and leaf

565 initiation, to immune and stress responses against pathogens and environmental stimuli (Islas-Flores et
566 al. 2015). Another phytohormone-related DAP was a brassinosteroid-related protein phosphatase
567 which accumulated in heated berries. In *Arabidopsis*, this protein is one of a four-member gene family
568 (BSL, for BRI1-SUPPRESSOR1 Like) shown to participate in brassinosteroid signalling (Maselli et
569 al. 2014), and potentially involved in stress tolerance (Gruszka 2018).

570 Heat stress is accompanied by oxidative stress, which may trigger adverse effects through oxidative
571 damage of cell membranes and important macromolecules (Bokszczanin and Fragkostefanakis 2013).
572 Therefore, redox regulation and signalling have been recognised as crucial mechanisms for acute
573 responses to HT (Katano et al. 2018). Our proteomic analysis of HT berries highlighted several up-
574 regulated DAPs with antioxidant properties, including two glutathione S-transferases (GSTs), a
575 glutathione reductase (GR, VIT_07s0141g0039), one catalase, a protein-disulfide isomerase (PDI), an
576 aldo-keto reductase (AKR), two mannitol dehydrogenases (MTD) and an *S*-nitrosoglutathione
577 reductase (GSNOR). Glutathione S-transferases belong to a well-characterized detoxification enzyme
578 family, catalyzing the conjugation of the reduced glutathione tripeptide (GSH) to electrophilic
579 substrates (Kumar and Trivedi 2018). GSTs quench reactive molecules through the addition of GSH
580 and protect the cell from oxidative damage. Accordingly, plant GSTs play an important role in abiotic
581 and biotic stress responses (Nianiou-Obeidat et al. 2017). The two up-regulated GSTs were observed
582 at veraison but an opposite pattern was seen at ripening with five GST isoforms being strongly
583 repressed at the protein level. This GST down-regulation observed under HT may negatively impact
584 the accumulation efficiency of flavonoids into the vacuolar compartment of ripening berries. Indeed,
585 GSTs function as non-enzymatic carriers in intracellular transport, allowing the accumulation of
586 flavonoid-GSH conjugates into vacuoles (Petruzza et al. 2013). The GST-mediated transport of
587 anthocyanins and proanthocyanidins has been established in grape as well (Perez-Diaz et al. 2016).
588 Among the four other heat-affected proteins, GR catalyses the reduction of glutathione disulfide
589 (GSSG) to GSH, and its role in heat tolerance was described in different plant species (Jegadeesan et
590 al. 2018). The ROS detoxifying enzyme catalase is considered as an acclimation protein required for
591 basal thermotolerance (Vanderauwera et al. 2011), whereas *PDI* encodes an oxidoreductase enzyme
592 belonging to the thioredoxin superfamily. PDI can be considered as a folding enzyme since it catalyses
593 the formation of correct disulfide bridges between the cysteine residues in proteins to maintain ER-
594 homeostasis (Park and Seo 2015). In potato, StPDI1 was found to interact with a sucrose transporter
595 (StSUT1) (Krugel et al. 2012), and a role in abiotic stress tolerance and H₂O₂ detoxification was
596 suggested (Eggert et al. 2016). Recently, the heterologous over-expression of a *Methanothermobacter*
597 *thermautotrophicus* PDI with a chaperone function and disulfide isomerase activity could confer
598 thermotolerance to transgenic rice (Wang et al. 2018). The aldo-keto reductase (AKR) superfamily
599 contains proteins able to reduce a broad spectrum of substrates, ranging from simple sugars to
600 potentially toxic aldehydes. Plant AKRs are involved in diverse metabolic reactions including reactive
601 aldehyde detoxification, biosynthesis of osmolytes, secondary metabolism and membrane transport.

602 These proteins may also confer multiple stress tolerance (Sengupta et al. 2015), including heat stress
603 tolerance as reported in rice (Turoczy et al. 2011). Two mannitol dehydrogenases (MTD) increased in
604 abundance in grape berries exposed to HT. The role of mannitol and by extension, its catabolic
605 enzyme MTD, is well established in resistance to both biotic and abiotic stresses. Several studies
606 suggest that mannitol protects plants from various abiotic stresses by acting as an osmoprotectant and
607 by quenching the damaging ROS (Patel and Williamson 2016). In grapevine, VvMTD1 was described
608 as involved in water-deficit stress tolerance (Conde et al. 2015). Finally, GSNOR has an important
609 function in the homeostasis of NO (nitric oxide), which is considered as a broad-spectrum anti-stress
610 molecule when acting at low concentration in plants. Whereas NO was described to accumulate in
611 tobacco exposed to HT (Gould et al. 2003), the key role of GSNOR1-mediated de-nitrosylation in
612 thermotolerance was revealed by the characterization of *Arabidopsis hot5* (sensitive to hot
613 temperatures) knock-out mutants (Lee et al. 2008). Taken together, these up-regulated and redox-
614 related proteins may play a role in signalling as well as in preventing harmful effects of ROS and
615 reactive nitrogen species in berries under HS conditions.

616 Finally, our proteomic work highlighted the heat-induced accumulation of two proteins related to
617 vitamin metabolism, namely VvPDX1.2 and VvPANK (**Table 2**). The pseudoenzyme PDX1.2
618 (Pyridoxine biosynthesis 1) is a noncatalytic homolog of the PDX1 subunit of the vitamin B₆
619 biosynthesis protein machinery. Transcriptionally regulated by HsfA1, PDX1.2 serves to stabilize the
620 catalytic PDX1s under HT, thereby maintaining vitamin B₆ homeostasis in plants (Dell'Aglio et al.
621 2017). AtPDX1.2 was reported as essential for plant growth under heat and oxidative stress conditions
622 (Moccand et al. 2014). The second up-regulated protein corresponds to a pantothenate kinase (PANK).
623 Pantothenate (known as vitamin B₅) is the key precursor of CoA and acyl carrier protein, which are
624 essential co-factors for many metabolic enzymes (Raman and Rathinasabapathi 2004). The PANK
625 converts pantothenate to phosphopantothenate, using ATP as phosphate donor. PANK activity plays a
626 critical role in regulating intracellular coenzyme A (CoA) levels in bacteria and animals but little is
627 known about the role of this enzyme in plants (Ottenhof et al. 2004). It was suggested that an
628 increased level of CoA, through an increased PANK activity, may be responsible for improved plant
629 growth and stress resistance (Rubio et al. 2008).

630

631 **Conclusion**

632 Global warming represents a substantial risk for the sustainability of wine production in many regions.
633 Plant breeding combined with molecular plant biotechnology have the potential to deliver stable yields
634 and quality under warmer climate conditions, but this requires the identification of key traits of
635 tolerance to heat. To this end, we combined omics approaches to decipher the molecular components
636 and processes affected in developing Cabernet Sauvignon berries exposed to heat. In the present study,
637 592 differentially abundant proteins were identified in heated berries among which some may play a

638 substantial role in heat tolerance. Our data also show that there is not a straightforward correlation
639 between the heat-modulated berry transcriptome and proteome, indicating the major involvement of
640 post-transcriptional regulation under heat stress. These pronounced changes contribute on the one
641 hand to generate fruit damaging effects, including disruption of carbohydrate and secondary
642 metabolism, and on the other hand to mobilize key heat-signalling players as well as protective
643 molecules such as antioxidants and chaperones (**Figure 5**). In summary, these findings help elucidate
644 heat-triggering effects on developing grape berries and identify genes and proteins that potentially
645 control heat acclimation in grapevine. The functional characterization of some putative candidates is in
646 progress.

647

648 **MATERIAL AND METHODS**

649 **Plant material, temperature and sampling**

650 Elevated temperature (+ 8°C) was directed on clusters of *V. vinifera* Cabernet Sauvignon fruiting
651 cuttings. The HT was applied at three different berry developmental stages and maintained throughout
652 the light period (7:00 am to 7:00 pm) until harvesting. To separate short- and long-term responses,
653 berries were collected after 1 and 7 days of treatment. Details of the experimental protocol and
654 sampling were provided in Lecourieux et al. (2017). The same pools of deseeded berries were used for
655 both the transcriptomic analysis reported in Lecourieux et al. (2017) and the proteomic analysis
656 reported here.

657

658 **Tissue preparation and total protein extraction**

659 Each biological replicate contained 25 pooled berries underwent independent protein extraction using
660 a protocol adapted from Lücker et al. (2009) and Martinez-Estezo et al. (2011). Briefly, deseeded
661 berries were ground to a fine powder in a mortar with liquid nitrogen and 3 g of frozen ground tissue
662 were used to prepare a total protein extract. Pectins and pigments were removed after incubation of the
663 frozen tissues in 25 mL of a cold (-20°C) ethyl acetate: ethanol (1 : 2 (v/v) for 30 min at -20°C with
664 repeated vortexing. The protein extract was precipitated by centrifugation for 3 min at 21000 × g at
665 4°C, and a second ethyl acetate: ethanol wash was performed in a similar way. The pellet was next
666 extracted twice with cold (-20°C) 100% acetone by vortexing and centrifuging, as before. The pellet
667 was transferred to a mortar and dried in a fume hood until complete acetone evaporation. The dried
668 pellet was finely ground after addition of 1/3 vol of sand, and washed with 1.5 mL chilled 10% (w/v)
669 trichloroacetic acid (TCA) in acetone. This washing step was repeated seven times. This was followed
670 by two washes with aqueous 10% (v/v) TCA, two with chilled 80% (v/v) acetone, and drying at 4 °C.
671 Each washing step included incubation at -20°C for 20 min followed by centrifugation for 3 min at
672 21000 × g at 4°C. Subsequently the dried pellet was resuspended in 500 µL protein extraction buffer
673 containing: 0.7 M sucrose, 0.1 M KCl, 0.5 M Tris-HCl (pH 7.5), 50 mM EDTA, 1% PVPP, 1%

674 DTT and protease inhibitor cocktail (Roche, Germany), and incubated at room temperature (RT) for
675 30 min with intermittent vortexing. Then, an equal volume of Tris-saturated phenol (pH 7.5) was
676 added, the sample was incubated at RT for 30 min with vortexing every 5 min. After centrifugation at
677 $15,000 \times g$ at RT for 40 min, the phenol phase in the upper layer was transferred into a new 2-mL
678 Eppendorf tubes and the remaining aqueous lower phase was re-extracted with another 500 μ L Tris-
679 saturated phenol (pH 7.5). The two phenol phases were combined and washed twice with an equal
680 volume of phenol washing buffer (0.7 M sucrose, 0.1 M KCl, 0.5 M Tris (pH 7.0, 50 mM EDTA, 1%
681 DTT, and protease inhibitor cocktail). Proteins contained in the organic phase were precipitated
682 overnight at 4°C with 5 volumes of 0.1 M ammonium acetate in methanol. Precipitated proteins were
683 washed three times in 0.1 M ammonium acetate in methanol and twice in ice-cold 80% (v/v) acetone.
684 All these precipitating and washing steps included incubation for 30 min followed by centrifugation at
685 $15,000 \times g$ at 4°C for 40 min. Finally, proteins were dissolved in 200 μ L freshly prepared
686 solubilization solution (7 M urea, 70 mM SDS, 20 mM DTT) and quantified according to the Bradford
687 method (Bradford, 1976) with bovine serum albumin as standard.

688

689 **Sample preparation and protein digestion**

690 Thirty μ g of each protein sample (three biological replicates for each condition; 12 conditions in total:
691 three stages x two treatments x two treatment durations) were solubilized in Laemmli buffer and were
692 separated by SDS-PAGE in a 10% (w/v) acrylamide gel. After colloidal blue staining, each lane was
693 horizontally divided into four parts of equal length and each part was subsequently cut in 1 mm x 1
694 mm gel pieces. Gel pieces were destained in 25 mM ammonium bicarbonate dissolved in 50% (v/v)
695 acetonitrile (ACN), rinsed twice in ultrapure water and dehydrated in 100% ACN for 10 min. After
696 ACN removal, gel pieces were dried at room temperature, covered with proteolysis solution
697 (containing 10 ng/ μ l trypsin dissolved in 40 mM NH_4HCO_3 and 10% (v/v) ACN), rehydrated at 4 °C
698 for 10 min, and finally incubated overnight at 37 °C. The gel pieces were then incubated for 15 min in
699 40 mM NH_4HCO_3 and 10% (v/v) ACN at room temperature with rotary shaking. The supernatant was
700 collected, and an $\text{H}_2\text{O}/\text{ACN}/\text{formic acid}$ (47.5:47.5:5 by volume) extraction solution was added and
701 gel slices incubated for 15 min. The extraction step was repeated twice. Supernatants were pooled and
702 concentrated in a vacuum centrifuge to a final volume of 40 μ L. Digests were finally acidified by
703 addition of 2.4 μ L of 5% (v/v) formic acid and the resulting peptide mixtures stored at -20 °C until
704 analysis.

705

706 **nLC-MS/MS analysis**

707 Peptide mixtures were analyzed on an Ultimate 3000 nanoLC system (Dionex, Amsterdam, The
708 Netherlands) coupled to a nanospray LTQ-Orbitrap XL mass spectrometer (ThermoFinnigan, San
709 Jose, USA). Ten microliters of peptide mixtures were loaded onto a 300- μ m-inner diameter x 5-mm
710 C_{18} PepMapTM trap column (LC Packings) at a flow rate of 30 μ L/min. The peptides were eluted from

711 the trap column onto an analytical 75-mm id x 15-cm C18 Pep-Map column (LC Packings) with a 5–
712 40% linear gradient of solvent B in 105 min (solvent A was 0.1% formic acid in 5% ACN, and solvent
713 B was 0.1% formic acid in 80% ACN). The separation flow rate was set at 200 nL/min. The mass
714 spectrometer operated in positive ion mode at a 1.8-kV needle voltage and a 39-V capillary voltage.
715 Data were acquired in a data-dependent mode alternating a Fourier transform mass spectrometry
716 (FTMS) scan survey over the range m/z 300–1700 and six ion trap MS/MS scans with CID (Collision
717 Induced Dissociation) as activation mode. MS/MS spectra were acquired using a 3- m/z unit ion
718 isolation window and a normalized collision energy of 35%. Only +2 and +3 charge-state ions were
719 selected for fragmentation. Dynamic exclusion duration was set to 30s.

720

721 **Database search and MS data processing**

722 Data were searched using the SEQUEST algorithm (supported by the Proteome Discoverer 1.4
723 software; Thermo Fisher Scientific Inc.) against a subset of the 2012.01 version of UniProt database
724 restricted to *Vitis vinifera* Reference Proteome Set (29,817 entries). Spectra from peptides higher than
725 5000 Da or lower than 350 Da were rejected. The search parameters were as follows: mass accuracy of
726 the monoisotopic peptide precursor and peptide fragments was set to 10 ppm and 0.8 Da respectively.
727 Only *b*- and *g*-ions were considered for mass calculation. Oxidation of methionines (+16 Da) was
728 considered as a variable modification. Two missed trypsin cleavages were allowed. Peptide validation
729 was performed using the Percolator algorithm (Käll et al. 2007), and only “high confidence” peptides
730 were retained corresponding to a 1% false positive rate at the peptide level.

731

732 **Label-Free Quantitative Data Analysis**

733 Raw LC-MS/MS data were imported into the Progenesis LC-MS 4.0 (Non Linear Dynamics)
734 software. Data processing includes the following steps: (i) Features detection, (ii) Features alignment
735 across the six samples for each part, (iii) Volume integration for 2-6 charge-state ions, (iv)
736 Normalization on total protein abundance, (v) Import of sequence information, (vi) ANOVA test at the
737 peptide level and filtering for features $p < 0.05$, (vii) Calculation of protein abundance (sum of the
738 volume of corresponding peptides), (viii) ANOVA test at protein level and filtering for features p
739 < 0.05 . Only non-conflicting features and unique peptides were considered for calculation at the
740 protein level. Quantitative data were considered for proteins quantified by a minimum of two peptides.
741 The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via
742 the PRIDE (Perez-Riverol et al. 2019) partner repository with the dataset identifier PXD014693.

743 Protein abundance changes between control and heat-exposed fruits were considered to be
744 significant when a minimum fold change of 1.5 ($\log_2FC > 0.58$ or < -0.58) was reached, with a
745 significance threshold of $p < 0.05$. Only proteins identified in all three biological replicates were
746 considered for quantification. Significantly affected protein categories based on the MapMan
747 Ontology (Usadel et al., 2005) were identified using a Chi-square test. MapMan mappings for the

748 Cribi 12X grapevine genome were based on closest homologs regarding to the *Arabidopsis thaliana*
749 genome. Protein annotations were taken from Grimplet et al. (2012).

750

751 **Primary metabolism quantification**

752 Deseeded berries were ground to a fine powder in liquid nitrogen and aliquots (15–20 mg) of frozen
753 powder were extracted with chloroform/methanol as described by Lunn et al. (2006). Primary
754 metabolites were measured by LC-MS/MS as described by Lunn et al. (2006), with modifications as
755 described in Figueroa et al. (2016) and Dai et al. (2013).

756

757 **ACKNOWLEDGMENTS**

758 The authors would like to thank Christelle Renaud and Ghislaine Hilbert for metabolite
759 quantifications. For the production of the fruiting cuttings and technical assistance during greenhouse
760 experiments, we express our gratitude to Jean Pierre Petit, Nicolas Hoquard and Guillaume Pacreau.

761

762 **FUNDING**

763 This research received funding from the Agence Nationale de la Recherche for the project
764 “DURAVITIS” (grant no. ANR-2010-GENM-004-01).

765

766 **AUTHOR CONTRIBUTIONS**

767 DL, FL and PP designed the research; DL oversaw the research; PP and DL performed the greenhouse
768 experiments; FL, JC, and DL carried out the protein extraction; CK performed the bioinformatics
769 analysis; RF, DL and JEL did the metabolic analysis; SC and MB did the metabolic analysis, FL, CK,
770 and DL analysed and interpreted the data; DL and FL drafted the manuscript; SD, JEL, LW and EG
771 critically revised the manuscript. All authors read and approved the final manuscript.

772

773 **CONFLICTS OF INTEREST**

774 The authors declare no conflicts of interest.

775 **REFERENCES**

- 776
- 777 Alem H, Rigou P, Schneider R, Ojeda H, Torregrosa L (2018) Impact of agronomic
778 practices on grape aroma composition: a review. **J Sci Food Agric** 99: 975-985
- 779 Asproudi A, Petrozziello M, Cavalletto S, Guidoni S (2016) Grape aroma precursors in cv.
780 Nebbiolo as affected by vine microclimate. **Food chemistry** 211: 947-956
- 781 Baena-Gonzalez E, Rolland F, Thevelein JM, Sheen J (2007) A central integrator of
782 transcription networks in plant stress and energy signalling. **Nature** 448: 938-942
- 783 Baurle I (2016) Plant Heat Adaptation: priming in response to heat stress.
784 **F1000Research** 5:
- 785 Beauvoit B, Belouah I, Bertin N, Cakpo CB, Colombié S, Dai Z, Gautier H, Génard M, Moing
786 A, Roch L, Vercambre G, Gibon Y (2018) Putting primary metabolism into perspective to
787 obtain better fruits. **Annals of botany** 122: 1-21
- 788 Bhatta M, Morgounov A, Belamkar V, Baenziger PS (2018) Genome-Wide Association
789 Study Reveals Novel Genomic Regions for Grain Yield and Yield-Related Traits in
790 Drought-Stressed Synthetic Hexaploid Wheat. **Int J Mol Sci** 19:
- 791 Bokszczanin KL, Fragkostefanakis S (2013) Perspectives on deciphering mechanisms
792 underlying plant heat stress response and thermotolerance. **Front Plant Sci** 4: 315
- 793 Carbonell-Bejerano P, Santa Maria E, Torres-Perez R, Royo C, Lijavetzky D, Bravo G,
794 Aguirreolea J, Sanchez-Diaz M, Antolin MC, Martinez-Zapater JM (2013)
795 Thermotolerance responses in ripening berries of *Vitis vinifera* L. cv Muscat Hamburg.
796 **Plant Cell Physiol** 54: 1200-1216
- 797 Conde A, Regalado A, Rodrigues D, Costa JM, Blumwald E, Chaves MM, Gerós H (2015)
798 Polyols in grape berry: transport and metabolic adjustments as a physiological strategy
799 for water-deficit stress tolerance in grapevine. **Journal of Experimental Botany** 66:
800 889-906
- 801 Cramer GR, Urano K, Delrot S, Pezzotti M, Shinozaki K (2011) Effects of abiotic stress on
802 plants: a systems biology perspective. **BMC Plant Biol** 11: 163
- 803 Cuadros-Inostroza A, Ruiz-Lara S, Gonzalez E, Eckardt A, Willmitzer L, Pena-Cortes H
804 (2016) GC-MS metabolic profiling of Cabernet Sauvignon and Merlot cultivars during
805 grapevine berry development and network analysis reveals a stage- and cultivar-
806 dependent connectivity of primary metabolites. **Metabolomics : Official journal of the**
807 **Metabolomic Society** 12: 39
- 808 Dai ZW, Leon C, Feil R, Lunn JE, Delrot S, Gomes E (2013) Metabolic profiling reveals
809 coordinated switches in primary carbohydrate metabolism in grape berry (*Vitis vinifera*
810 L.), a non-climacteric fleshy fruit. **J Exp Bot** 64: 1345-1355
- 811 de Jesus-Pires C, Ferreira-Neto JRC, Pacifico Bezerra-Neto J, Kido EA, de Oliveira Silva RL,
812 Pandolfi V, Wanderley-Nogueira AC, Binneck E, da Costa AF, Pio-Ribeiro G, Pereira-
813 Andrade G, Sittolin IM, Freire-Filho F, Benko-Iseppon AM (2019) Plant Thaumatin-like
814 Proteins: Function, Evolution and Biotechnological Applications. **Current protein &**
815 **peptide science**:
- 816 Degu A, Hochberg U, Sikron N, Venturini L, Buson G, Ghan R, Plaschkes I, Batushansky A,
817 Chalifa-Caspi V, Mattivi F, Delledonne M, Pezzotti M, Rachmilevitch S, Cramer GR, Fait A
818 (2014) Metabolite and transcript profiling of berry skin during fruit development
819 elucidates differential regulation between Cabernet Sauvignon and Shiraz cultivars at
820 branching points in the polyphenol pathway. **BMC Plant Biol** 14: 188
- 821 Dell'Aglio E, Boycheva S, Fitzpatrick TB (2017) The Pseudoenzyme PDX1.2 Sustains
822 Vitamin B₆ Biosynthesis as a Function of Heat Stress. **Plant Physiology**
823 174: 2098-2112

- 824 Deloire A, Vaudour E, Carey VA, Bonnardot V, Van Leeuwen C (2005) Grapevine
825 responses to terroir: a global approach. **OENO One** 39: 149-162
- 826 Doukhanina EV, Chen S, van der Zalm E, Godzik A, Reed J, Dickman MB (2006)
827 Identification and functional characterization of the BAG protein family in Arabidopsis
828 thaliana. **Journal of Biological Chemistry** 281: 18793-18801
- 829 Downey MO, Dokoozlian NK, Krstic MP (2006) Cultural practice and environmental
830 impacts on the flavonoid composition of grapes and wine: a review of recent research.
831 **American Journal of Enology and Viticulture** 57: 257-268
- 832 du Plessis K, Young PR, Eyeghe-Bickong HA, Vivier MA (2017) The Transcriptional
833 Responses and Metabolic Consequences of Acclimation to Elevated Light Exposure in
834 Grapevine Berries. **Front Plant Sci** 8: 1261
- 835 Dutt S, Parkash J, Mehra R, Sharma N, Singh B, Raigond P, Joshi A, Chopra S, Singh BP
836 (2015) Translation initiation in plants: roles and implications beyond protein synthesis.
837 **Biologia Plantarum** 59: 401-412
- 838 Eggert E, Obata T, Gerstenberger A, Gier K, Brandt T, Fernie AR, Schulze W, Kuhn C
839 (2016) A sucrose transporter-interacting protein disulphide isomerase affects redox
840 homeostasis and links sucrose partitioning with abiotic stress tolerance. **Plant, cell &**
841 **environment** 39: 1366-1380
- 842 Escandon M, Meijon M, Valledor L, Pascual J, Pinto G, Canal MJ (2018) Metabolome
843 Integrated Analysis of High-Temperature Response in Pinus radiata. **Front Plant Sci** 9:
844 485
- 845 Figueroa CM, Feil R, Ishihara H, Watanabe M, Kolling K, Krause U, Hohne M, Encke B,
846 Plaxton WC, Zeeman SC, Li Z, Schulze WX, Hoefgen R, Stitt M, Lunn JE (2016) Trehalose
847 6-phosphate coordinates organic and amino acid metabolism with carbon availability.
848 **The Plant journal : for cell and molecular biology** 85: 410-423
- 849 Figueroa CM, Lunn JE (2016) A Tale of Two Sugars: Trehalose 6-Phosphate and Sucrose.
850 **Plant Physiology** 172: 7-27
- 851 Finka A, Cuendet AF, Maathuis FJ, Saidi Y, Goloubinoff P (2012) Plasma membrane cyclic
852 nucleotide gated calcium channels control land plant thermal sensing and acquired
853 thermotolerance. **The Plant cell** 24: 3333-3348
- 854 Fraga H, Garcia de Cortazar Atauri I, Malheiro AC, Santos JA (2016) Modelling climate
855 change impacts on viticultural yield, phenology and stress conditions in Europe. **Global**
856 **change biology** 22: 3774-3788
- 857 Francisco RM, Regalado A, Ageorges A, Burla BJ, Bassin B, Eisenach C, Zarrouk O, Vialet S,
858 Marlin T, Chaves MM, Martinoia E, Nagy R (2013) ABCC1, an ATP binding cassette
859 protein from grape berry, transports anthocyanidin 3-O-Glucosides. **The Plant cell** 25:
860 1840-1854
- 861 Gao F, Han X, Wu J, Zheng S, Shang Z, Sun D, Zhou R, Li B (2012) A heat-activated
862 calcium-permeable channel--Arabidopsis cyclic nucleotide-gated ion channel 6--is
863 involved in heat shock responses. **The Plant journal : for cell and molecular biology**
864 70: 1056-1069
- 865 George IS, Pascovici D, Mirzaei M, Haynes PA (2015) Quantitative proteomic analysis of
866 cabernet sauvignon grape cells exposed to thermal stresses reveals alterations in sugar
867 and phenylpropanoid metabolism. **Proteomics**:
- 868 Giribaldi M, Perugini I, Sauvage FX, Schubert A (2007) Analysis of protein changes
869 during grape berry ripening by 2 - DE and MALDI - TOF. **Proteomics** 7: 3154-3170
- 870 Gomez C, Conejero G, Torregrosa L, Cheynier V, Terrier N, Ageorges A (2011) In vivo
871 grapevine anthocyanin transport involves vesicle-mediated trafficking and the
872 contribution of anthoMATE transporters and GST. **The Plant Journal** 67: 960-970

- 873 Gonzalez RM, Iusem ND (2014) Twenty years of research on Asr (ABA-stress-ripening)
874 genes and proteins. **Planta** 239: 941-949
- 875 Goulao L, Fernandes J, Lopes P, Amâncio S (2012a) *Tackling the cell wall of the grape*
876 *berry*. Bentham Science Publishers Sharjah, UAE,
- 877 Goulao L, Fernandes J, Lopes P, Amancio S (2012b) Tackling the cell wall of the grape
878 berry. **The biochemistry of the grape berry**: 172
- 879 Gould K, Lamotte O, Klinguer A, Pugin A, Wendehenne D (2003) Nitric oxide production
880 in tobacco leaf cells: a generalized stress response? **Plant, cell & environment** 26:
881 1851-1862
- 882 Gouot JC, Smith JP, Holzapfel BP, Walker AR, Barril C (2018) Grape berry flavonoids: A
883 review of their biochemical responses to high and extreme high temperatures. **J Exp**
884 **Bot**:
- 885 Greer DH, Weedon MM (2012) Modelling photosynthetic responses to temperature of
886 grapevine (*Vitis vinifera* cv. Semillon) leaves on vines grown in a hot climate. **Plant, cell**
887 **& environment** 35: 1050-1064
- 888 Greer DH, Weston C (2010) Heat stress affects flowering, berry growth, sugar
889 accumulation and photosynthesis of *Vitis vinifera* cv. Semillon grapevines grown in a
890 controlled environment. **Functional Plant Biology** 37: 206-214
- 891 Gruszka D (2018) Crosstalk of the Brassinosteroid Signalosome with Phytohormonal
892 and Stress Signaling Components Maintains a Balance between the Processes of Growth
893 and Stress Tolerance. **Int J Mol Sci** 19:
- 894 Guo HS, Zhang YM, Sun XQ, Li MM, Hang YY, Xue JY (2016) Evolution of the KCS gene
895 family in plants: the history of gene duplication, sub/neofunctionalization and
896 redundancy. **Molecular genetics and genomics : MGG** 291: 739-752
- 897 Havaux M (2014) Carotenoid oxidation products as stress signals in plants. **The Plant**
898 **Journal** 79: 597-606
- 899 He R, Zhuang Y, Cai Y, Aguero CB, Liu S, Wu J, Deng S, Walker MA, Lu J, Zhang Y (2018)
900 Overexpression of 9-cis-Epoxycarotenoid Dioxygenase Cisgene in Grapevine Increases
901 Drought Tolerance and Results in Pleiotropic Effects. **Front Plant Sci** 9: 970
- 902 Higashi Y, Okazaki Y, Myouga F, Shinozaki K, Saito K (2015) Landscape of the lipidome
903 and transcriptome under heat stress in *Arabidopsis thaliana*. **Scientific reports** 5:
904 10533
- 905 Islas-Flores T, Rahman A, Ullah H, Villanueva MA (2015) The Receptor for Activated C
906 Kinase in Plant Signaling: Tale of a Promiscuous Little Molecule. **Front Plant Sci** 6: 1090
- 907 Jacob P, Hirt H, Bendahmane A (2017) The heat-shock protein/chaperone network and
908 multiple stress resistance. **Plant Biotechnol J** 15: 405-414
- 909 James AB, Calixto CPG, Tzioutziou NA, Guo W, Zhang R, Simpson CG, Jiang W, Nimmo GA,
910 Brown JWS, Nimmo HG (2018) How does temperature affect splicing events? Isoform
911 switching of splicing factors regulates splicing of LATE ELONGATED HYPOCOTYL (LHY).
912 **Plant, cell & environment** 41: 1539-1550
- 913 Jegadeesan S, Chaturvedi P, Ghatak A, Pressman E, Meir S, Faigenboim A, Rutley N, Beery
914 A, Harel A, Weckwerth W, Firon N (2018) Proteomics of Heat-Stress and Ethylene-
915 Mediated Thermotolerance Mechanisms in Tomato Pollen Grains. **Front Plant Sci** 9:
916 1558
- 917 Jia KP, Baz L, Al-Babili S (2018) From carotenoids to strigolactones. **J Exp Bot** 69: 2189-
918 2204
- 919 Jiang J, Liu X, Liu C, Liu G, Li S, Wang L (2017) Integrating Omics and Alternative Splicing
920 Reveals Insights into Grape Response to High Temperature. **Plant Physiol** 173: 1502-
921 1518

- 922 Jones GV, White MA, Cooper OR, Storchmann K (2005) Climate change and global wine
923 quality. **Climatic change** 73: 319-343
- 924 Jung JH, Domijan M, Klose C, Biswas S, Ezer D, Gao M, Khattak AK, Box MS,
925 Charoensawan V, Cortijo S, Kumar M, Grant A, Locke JC, Schafer E, Jaeger KE, Wigge PA
926 (2016) Phytochromes function as thermosensors in Arabidopsis. **Science** 354: 886-889
- 927 Käll L, Canterbury JD, Weston J, Noble WS, MacCoss MJ (2007) Semi-supervised learning
928 for peptide identification from shotgun proteomics datasets. **Nature methods** 4: 923
- 929 Katano K, Honda K, Suzuki N (2018) Integration between ROS Regulatory Systems and
930 Other Signals in the Regulation of Various Types of Heat Responses in Plants. **Int J Mol**
931 **Sci** 19:
- 932 Keller M, Simm S (2018) The coupling of transcriptome and proteome adaptation during
933 development and heat stress response of tomato pollen. **BMC Genomics** 19: 447
- 934 Kosova K, Vitamvas P, Urban MO, Prasil IT, Renaut J (2018) Plant Abiotic Stress
935 Proteomics: The Major Factors Determining Alterations in Cellular Proteome. **Front**
936 **Plant Sci** 9: 122
- 937 Krugel U, He HX, Gier K, Reins J, Chincinska I, Grimm B, Schulze WX, Kuhn C (2012) The
938 potato sucrose transporter StSUT1 interacts with a DRM-associated protein disulfide
939 isomerase. **Mol Plant** 5: 43-62
- 940 Ku YS, Sintaha M, Cheung MY, Lam HM (2018) Plant Hormone Signaling Crosstalks
941 between Biotic and Abiotic Stress Responses. **Int J Mol Sci** 19:
- 942 Kuang L, Chen S, Guo Y, Ma H (2019) Quantitative Proteome Analysis Reveals Changes in
943 the Protein Landscape During Grape Berry Development With a Focus on Vacuolar
944 Transport Proteins. **Front Plant Sci** 10: 641
- 945 Kuhn N, Guan L, Dai ZW, Wu B-H, Lauvergeat V, Gomès E, Li S-H, Godoy F, Arce-Johnson
946 P, Delrot S (2014) Berry ripening: recently heard through the grapevine. **Journal of**
947 **experimental botany** 65: 4543-4559
- 948 Kumar S, Trivedi PK (2018) Glutathione S-Transferases: Role in Combating Abiotic
949 Stresses Including Arsenic Detoxification in Plants. **Front Plant Sci** 9: 751
- 950 Langer SE, Oviedo NC, Marina M, Burgos JL, Martinez GA, Civello PM, Villarreal NM
951 (2018) Effects of heat treatment on enzyme activity and expression of key genes
952 controlling cell wall remodeling in strawberry fruit. **Plant physiology and**
953 **biochemistry : PPB / Societe francaise de physiologie vegetale** 130: 334-344
- 954 Lecourieux F, Kappel C, Pieri P, Charon J, Pillet J, Hilbert G, Renaud C, Gomes E, Delrot S,
955 Lecourieux D (2017) Dissecting the Biochemical and Transcriptomic Effects of a Locally
956 Applied Heat Treatment on Developing Cabernet Sauvignon Grape Berries. **Front Plant**
957 **Sci** 8: 53
- 958 Lee U, Wie C, Fernandez BO, Feelisch M, Vierling E (2008) Modulation of nitrosative
959 stress by S-nitrosoglutathione reductase is critical for thermotolerance and plant growth
960 in Arabidopsis. **The Plant cell** 20: 786-802
- 961 Li B, Gao K, Ren H, Tang W (2018) Molecular mechanisms governing plant responses to
962 high temperatures. **Journal of integrative plant biology** 60: 757-779
- 963 Li Y, Williams B, Dickman M (2017) Arabidopsis B-cell lymphoma2 (Bcl-2)-associated
964 athanogene 7 (BAG7)-mediated heat tolerance requires translocation, sumoylation and
965 binding to WRKY29. **New Phytol** 214: 695-705
- 966 Ling Y, Serrano N, Gao G, Atia M, Mokhtar M, Woo YH, Bazin J, Veluchamy A, Benhamed
967 M, Crespi M, Gehring C, Reddy ASN, Mahfouz MM (2018) Thermoprimering triggers
968 splicing memory in Arabidopsis. **J Exp Bot** 69: 2659-2675

- 969 Liu GT, Ma L, Duan W, Wang BC, Li JH, Xu HG, Yan XQ, Yan BF, Li SH, Wang LJ (2014)
970 Differential proteomic analysis of grapevine leaves by iTRAQ reveals responses to heat
971 stress and subsequent recovery. **BMC Plant Biol** 14: 110
- 972 Liu GT, Wang JF, Cramer G, Dai ZW, Duan W, Xu HG, Wu BH, Fan PG, Wang LJ, Li SH
973 (2012) Transcriptomic analysis of grape (*Vitis vinifera* L.) leaves during and after
974 recovery from heat stress. **BMC Plant Biol** 12: 174
- 975 Liu JJ, Sturrock R, Ekramoddoullah AK (2010) The superfamily of thaumatin-like
976 proteins: its origin, evolution, and expression towards biological function. **Plant Cell**
977 **Rep** 29: 419-436
- 978 Lobell DB, Burke MB, Tebaldi C, Mastrandrea MD, Falcon WP, Naylor RL (2008)
979 Prioritizing climate change adaptation needs for food security in 2030. **Science** 319:
980 607-610
- 981 Lunn JE, Feil R, Hendriks JH, Gibon Y, Morcuende R, Osuna D, Scheible WR, Carillo P,
982 Hajirezaei MR, Stitt M (2006) Sugar-induced increases in trehalose 6-phosphate are
983 correlated with redox activation of ADPglucose pyrophosphorylase and higher rates of
984 starch synthesis in *Arabidopsis thaliana*. **The Biochemical journal** 397: 139-148
- 985 Martinez-Esteso MJ, Vilella-Anton MT, Pedreno MA, Valero ML, Bru-Martinez R (2013)
986 iTRAQ-based protein profiling provides insights into the central metabolism changes
987 driving grape berry development and ripening. **BMC Plant Biol** 13: 167
- 988 Martínez-Lüscher J, Kizildeniz T, Vučetić V, Dai Z, Luedeling E, van Leeuwen C, Gomès E,
989 Pascual I, Irigoyen JJ, Morales F (2016) Sensitivity of grapevine phenology to water
990 availability, temperature and CO₂ concentration. **Frontiers in Environmental Science**
991 4: 48
- 992 Maselli GA, Slamovits CH, Bianchi JJ, Vilarrasa-Blasi J, Caño-Delgado AI, Mora-García S
993 (2014) Revisiting the Evolutionary History and Roles of Protein Phosphatases with
994 Kelch-Like Domains in Plants. **Plant Physiology** 164: 1527-1541
- 995 Mathieu S, Terrier N, Procureur J, Bigey F, Gunata Z (2005) A carotenoid cleavage
996 dioxygenase from *Vitis vinifera* L.: functional characterization and expression during
997 grape berry development in relation to C₁₃-norisoprenoid accumulation. **J Exp Bot** 56:
998 2721-2731
- 999 Mauri N, Fernández-Marcos M, Costas C, Desvoyes B, Pichel A, Caro E, Gutierrez C (2016)
1000 GEM, a member of the GRAM domain family of proteins, is part of the ABA signaling
1001 pathway. **Scientific reports** 6: 22660
- 1002 Medici A, Laloi M, Atanassova R (2014) Profiling of sugar transporter genes in grapevine
1003 coping with water deficit. **FEBS letters** 588: 3989-3997
- 1004 Meiri D, Breiman A (2009) *Arabidopsis* ROF1 (FKBP62) modulates thermotolerance by
1005 interacting with HSP90.1 and affecting the accumulation of HsfA2-regulated sHSPs. **The**
1006 **Plant journal : for cell and molecular biology** 59: 387-399
- 1007 Moccand C, Boycheva S, Surriabre P, Tambasco-Studart M, Raschke M, Kaufmann M,
1008 Fitzpatrick TB (2014) The pseudoenzyme PDX1. 2 boosts vitamin B6 biosynthesis under
1009 heat and oxidative stress in *Arabidopsis*. **Journal of Biological Chemistry**: jbc. M113.
1010 540526
- 1011 Naniou-Obeidat I, Madesis P, Kissoudis C, Voulgari G, Chronopoulou E, Tsaftaris A,
1012 Labrou NE (2017) Plant glutathione transferase-mediated stress tolerance: functions
1013 and biotechnological applications. **Plant Cell Rep** 36: 791-805
- 1014 Niu Y, Xiang Y (2018) An Overview of Biomembrane Functions in Plant Responses to
1015 High-Temperature Stress. **Front Plant Sci** 9: 915
- 1016 Novakova P, Hirsch S, Feraru E, Tejos R, van Wijk R, Viaene T, Heilmann M, Lerche J, De
1017 Rycke R, Feraru MI, Grones P, Van Montagu M, Heilmann I, Munnik T, Friml J (2014) SAC

- 1018 phosphoinositide phosphatases at the tonoplast mediate vacuolar function in
1019 Arabidopsis. **Proceedings of the National Academy of Sciences of the United States**
1020 **of America** 111: 2818-2823
- 1021 Ottenhof HH, Ashurst JL, Whitney HM, Saldanha SA, Schmitzberger F, Gweon HS,
1022 Blundell TL, Abell C, Smith AG (2004) Organisation of the pantothenate (vitamin B5)
1023 biosynthesis pathway in higher plants. **The Plant Journal** 37: 61-72
- 1024 Park C-J, Seo Y-S (2015) Heat shock proteins: a review of the molecular chaperones for
1025 plant immunity. **The plant pathology journal** 31: 323
- 1026 Patel TK, Williamson JD (2016) Mannitol in Plants, Fungi, and Plant-Fungal Interactions.
1027 **Trends in plant science** 21: 486-497
- 1028 Pereira G, Gaudillere J-P, Pieri P, Hilbert G, Maucourt M, Deborde C, Moing A, Rolin D
1029 (2006) Microclimate influence on mineral and metabolic profiles of grape berries.
1030 **Journal of Agricultural and Food Chemistry** 54: 6765-6775
- 1031 Perez-Diaz R, Madrid-Espinoza J, Salinas-Cornejo J, Gonzalez-Villanueva E, Ruiz-Lara S
1032 (2016) Differential Roles for VviGST1, VviGST3, and VviGST4 in Proanthocyanidin and
1033 Anthocyanin Transport in *Vitis vinifera*. **Front Plant Sci** 7: 1166
- 1034 Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ,
1035 Inuganti A, Griss J, Mayer G, Eisenacher M, Perez E, Uszkoreit J, Pfeuffer J, Sachsenberg T,
1036 Yilmaz S, Tiwary S, Cox J, Audain E, Walzer M, Jarnuczak AF, Ternent T, Brazma A,
1037 Vizcaino JA (2019) The PRIDE database and related tools and resources in 2019:
1038 improving support for quantification data. **Nucleic acids research** 47: D442-d450
- 1039 Petit J, Bres C, Mauxion JP, Tai FW, Martin LB, Fich EA, Joubes J, Rose JK, Domergue F,
1040 Rothan C (2016) The Glycerol-3-Phosphate Acyltransferase GPAT6 from Tomato Plays a
1041 Central Role in Fruit Cutin Biosynthesis. **Plant Physiol** 171: 894-913
- 1042 Petruzza E, Braidot E, Zancani M, Peresson C, Bertolini A, Patui S, Vianello A (2013) Plant
1043 flavonoids--biosynthesis, transport and involvement in stress responses. **Int J Mol Sci**
1044 14: 14950-14973
- 1045 Raman SB, Rathinasabapathi B (2004) Pantothenate synthesis in plants. **Plant Science**
1046 167: 961-968
- 1047 Reshef N, Fait A, Agam N (2019) Grape berry position affects the diurnal dynamics of its
1048 metabolic profile. **Plant, cell & environment**:
- 1049 Reshef N, Walbaum N, Agam N, Fait A (2017) Sunlight Modulates Fruit Metabolic Profile
1050 and Shapes the Spatial Pattern of Compound Accumulation within the Grape Cluster.
1051 **Front Plant Sci** 8: 70
- 1052 Rienth M, Torregrosa L, Kelly MT, Luchaire N, Pellegrino A, Grimplet J, Romieu C (2014a)
1053 Is transcriptomic regulation of berry development more important at night than during
1054 the day? **PLoS One** 9: e88844
- 1055 Rienth M, Torregrosa L, Luchaire N, Chatbanyong R, Lecourieux D, Kelly MT, Romieu C
1056 (2014b) Day and night heat stress trigger different transcriptomic responses in green
1057 and ripening grapevine (*vitis vinifera*) fruit. **BMC Plant Biol** 14: 108
- 1058 Rienth M, Torregrosa L, Sarah G, Ardisson M, Brillouet JM, Romieu C (2016)
1059 Temperature desynchronizes sugar and organic acid metabolism in ripening grapevine
1060 fruits and remodels their transcriptome. **BMC Plant Biol** 16: 164
- 1061 Rubio S, Whitehead L, Larson TR, Graham IA, Rodriguez PL (2008) The coenzyme a
1062 biosynthetic enzyme phosphopantetheine adenyltransferase plays a crucial role in
1063 plant growth, salt/osmotic stress resistance, and seed lipid storage. **Plant Physiol** 148:
1064 546-556
- 1065 Ruffner HP, Hawker JS, Hale CR (1976) Temperature and enzymic control of malate
1066 metabolism in berries of *Vitis vinifera*. **Phytochemistry** 15: 1877-1880

- 1067 Ruiz-Ballesta I, Baena G, Gandullo J, Wang L, She YM, Plaxton WC, Echevarria C (2016)
1068 New insights into the post-translational modification of multiple phosphoenolpyruvate
1069 carboxylase isoenzymes by phosphorylation and monoubiquitination during sorghum
1070 seed development and germination. **J Exp Bot** 67: 3523-3536
- 1071 Sasaki Y, Nagano Y (2004) Plant acetyl-CoA carboxylase: structure, biosynthesis,
1072 regulation, and gene manipulation for plant breeding. **Bioscience, biotechnology, and**
1073 **biochemistry** 68: 1175-1184
- 1074 Scharf KD, Berberich T, Ebersberger I, Nover L (2012) The plant heat stress
1075 transcription factor (Hsf) family: structure, function and evolution. **Biochimica et**
1076 **biophysica acta** 1819: 104-119
- 1077 Schultz HR, Jones GV (2010) Climate induced historic and future changes in viticulture.
1078 **Journal of Wine Research** 21: 137-145
- 1079 Sengupta D, Naik D, Reddy AR (2015) Plant aldo-keto reductases (AKRs) as multi-
1080 tasking soldiers involved in diverse plant metabolic processes and stress defense: A
1081 structure-function update. **Journal of plant physiology** 179: 40-55
- 1082 Serrano A, Espinoza C, Armijo G, Inostroza-Blancheteau C, Poblete E, Meyer-Regueiro C,
1083 Arce A, Parada F, Santibanez C, Arce-Johnson P (2017) Omics Approaches for
1084 Understanding Grapevine Berry Development: Regulatory Networks Associated with
1085 Endogenous Processes and Environmental Responses. **Front Plant Sci** 8: 1486
- 1086 Shiratake K, Martinoia E (2007) Transporters in fruit vacuoles. **Plant Biotechnology** 24:
1087 127-133
- 1088 Soubeyrand E, Colombie S, Beauvoit B, Dai Z, Cluzet S, Hilbert G, Renaud C, Maneta-
1089 Peyret L, Dieuaide-Noubhani M, Merillon JM, Gibon Y, Delrot S, Gomes E (2018)
1090 Constraint-Based Modeling Highlights Cell Energy, Redox Status and alpha-
1091 Ketoglutarate Availability as Metabolic Drivers for Anthocyanin Accumulation in Grape
1092 Cells Under Nitrogen Limitation. **Front Plant Sci** 9: 421
- 1093 Stein O, Granot D (2018) Plant Fructokinases: Evolutionary, Developmental, and
1094 Metabolic Aspects in Sink Tissues. **Front Plant Sci** 9: 339
- 1095 Suragani M, Rasheedi S, Hasnain SE, Ehtesham NZ (2011) The translation initiation
1096 factor, PelF5B, from *Pisum sativum* displays chaperone activity. **Biochemical and**
1097 **biophysical research communications** 414: 390-396
- 1098 Sweetman C, Deluc LG, Cramer GR, Ford CM, Soole KL (2009) Regulation of malate
1099 metabolism in grape berry and other developing fruits. **Phytochemistry** 70: 1329-1344
- 1100 Sweetman C, Sadras VO, Hancock RD, Soole KL, Ford CM (2014) Metabolic effects of
1101 elevated temperature on organic acid degradation in ripening *Vitis vinifera* fruit. **J Exp**
1102 **Bot**:
- 1103 Tan BC, Schwartz SH, Zeevaart JA, McCarty DR (1997) Genetic control of abscisic acid
1104 biosynthesis in maize. **Proceedings of the National Academy of Sciences** 94: 12235-
1105 12240
- 1106 Torregrosa L, Bigard A, Doligez A, Lecourieux D, Rienth M, Luchaire N, Pieri P,
1107 Chatbanyong R, Shahood R, Farnos M, Roux C, Adiveze A, Pillet J, Sire Y, Zumstein E,
1108 Veyret M, Le Cunff L, Lecourieux F, Saurin N, Muller B, Ojeda H, Houel C, Péros J-P, This P,
1109 Pellegrino A, Romieu C (2017) Developmental, molecular and genetic studies on
1110 grapevine response to temperature open breeding strategies for adaptation to warming.
1111 **OENO One** 51: 155-165
- 1112 Turoczy Z, Kis P, Torok K, Cserhati M, Lendvai A, Dudits D, Horvath GV (2011)
1113 Overproduction of a rice aldo-keto reductase increases oxidative and heat stress
1114 tolerance by malondialdehyde and methylglyoxal detoxification. **Plant Mol Biol** 75: 399-
1115 412

- 1116 Usadel B, Nagel A, Thimm O, Redestig H, Blaesing OE, Palacios-Rojas N, Selbig J,
1117 Hannemann J, Piques MC, Steinhauser D, Scheible WR, Gibon Y, Morcuende R, Weicht D,
1118 Meyer S, Stitt M (2005) Extension of the visualization tool MapMan to allow statistical
1119 analysis of arrays, display of corresponding genes, and comparison with known
1120 responses. **Plant Physiol** 138: 1195-1204
- 1121 Van Leeuwen C, Schultz HR, de Cortazar-Atauri IG, Duchêne E, Ollat N, Pieri P, Bois B,
1122 Goutouly J-P, Quénot H, Touzard J-M (2013) Why climate change will not dramatically
1123 decrease viticultural suitability in main wine-producing areas by 2050. **Proceedings of**
1124 **the National Academy of Sciences** 110: E3051-E3052
- 1125 Van Leeuwen C, Seguin G (2006) The concept of terroir in viticulture. **Journal of wine**
1126 **research** 17: 1-10
- 1127 Vanderauwera S, Suzuki N, Miller G, van de Cotte B, Morsa S, Ravanat JL, Hegie A,
1128 Triantaphylides C, Shulaev V, Van Montagu MC, Van Breusegem F, Mittler R (2011)
1129 Extranuclear protection of chromosomal DNA from oxidative stress. **Proceedings of the**
1130 **National Academy of Sciences of the United States of America** 108: 1711-1716
- 1131 Wahid A, Gelani S, Ashraf M, Foolad MR (2007) Heat tolerance in plants: an overview.
1132 **Environmental and experimental botany** 61: 199-223
- 1133 Walker RP, Battistelli A, Moscatello S, Tecsi L, Leegood RC, Famiani F (2015)
1134 Phosphoenolpyruvate carboxykinase and gluconeogenesis in grape pericarp. **Plant**
1135 **physiology and biochemistry : PPB / Societe francaise de physiologie vegetale** 97:
1136 62-69
- 1137 Wang L, Sun X, Weiszmann J, Weckwerth W (2017) System-Level and Granger Network
1138 Analysis of Integrated Proteomic and Metabolomic Dynamics Identifies Key Points of
1139 Grape Berry Development at the Interface of Primary and Secondary Metabolism. **Front**
1140 **Plant Sci** 8: 1066
- 1141 Wang X, Chen J, Liu C, Luo J, Yan X, Aihua A, Cai Y, Xie H, Ding X, Peng X (2018) Over-
1142 expression of a protein disulfide isomerase gene from *Methanothermobacter*
1143 *thermautotrophicus*, enhances heat stress tolerance in rice. **Gene** 684: 124-130
- 1144 Welti R, Shah J, Li W, Li M, Chen J, Burke J, Fauconnier M-L, Chapman K, Chye M, Wang X
1145 (2007) Plant lipidomics: discerning biological function by profiling plant complex lipids
1146 using mass spectrometry. **Frontiers in bioscience: a journal and virtual library** 12:
1147 Wu HC, Bulgakov VP, Jinn TL (2018) Pectin Methylsterases: Cell Wall Remodeling
1148 Proteins Are Required for Plant Response to Heat Stress. **Front Plant Sci** 9: 1612
- 1149 Xu J, Zhang B, Jiang C, Ming F (2011) RceIF5A, encoding an eukaryotic translation
1150 initiation factor 5A in *Rosa chinensis*, can enhance thermotolerance, oxidative and
1151 osmotic stress resistance of *Arabidopsis thaliana*. **Plant Mol Biol** 75: 167-178
- 1152 Yan X, Qiao H, Zhang X, Guo C, Wang M, Wang Y, Wang X (2017) Analysis of the grape
1153 (*Vitis vinifera* L.) thaumatin-like protein (TLP) gene family and demonstration that
1154 TLP29 contributes to disease resistance. **Scientific reports** 7: 4269
- 1155 Yanguez E, Castro-Sanz AB, Fernandez-Bautista N, Oliveros JC, Castellano MM (2013)
1156 Analysis of genome-wide changes in the transcriptome of *Arabidopsis* seedlings subjected
1157 to heat stress. **PLoS One** 8: e71425
- 1158 Young PR, Eyeghe-Bickong HA, du Plessis K, Alexandersson E, Jacobson DA, Coetzee Z,
1159 Deloire A, Vivier MA (2016) Grapevine plasticity in response to an altered microclimate:
1160 Sauvignon Blanc modulates specific metabolites in response to increased berry
1161 exposure. **Plant physiology** 170: 1235-1254
- 1162 Zamboni A, Di Carli M, Guzzo F, Stocchero M, Zenoni S, Ferrarini A, Tononi P, Toffali K,
1163 Desiderio A, Lilley KS, Pe ME, Benvenuto E, Delledonne M, Pezzotti M (2010)

1164 Identification of putative stage-specific grapevine berry biomarkers and omics data
1165 integration into networks. **Plant Physiol** 154: 1439-1459
1166 Zhang D, Chen L, Li D, Lv B, Chen Y, Chen J, XuejiaoYan, Liang J (2014) OsRACK1 is
1167 involved in abscisic acid- and H₂O₂-mediated signaling to regulate seed germination in
1168 rice (*Oryza sativa*, L.). **PLoS One** 9: e97120
1169 Zhang L, Liu X, Gaikwad K, Kou X, Wang F, Tian X, Xin M, Ni Z, Sun Q, Peng H, Vierling E
1170 (2017) Mutations in eIF5B Confer Thermosensitive and Pleiotropic Phenotypes via
1171 Translation Defects in *Arabidopsis thaliana*. **The Plant cell** 29: 1952-1969
1172 Zhao J (2015) Flavonoid transport mechanisms: how to go, and with whom. **Trends in**
1173 **plant science** 20: 576-585
1174
1175

1176 **FIGURE LEGENDS**

1177

1178 **Figure 1. Venn diagrams displaying the numbers of DAPs in heat-treated berries, according to**
1179 **the developmental stage (A) or the treatment duration (B).** DAPs were selected using a 1.5 fold
1180 expression change and an adjusted $p < 0.05$ (with FDR correction). Only proteins identified in all three
1181 biological replicates and with a minimum of two unique peptides were considered for quantification.
1182 The detailed information of the common DAPs upon HT for each stage or for each exposure time was
1183 listed as in Supplementary Tables 1 and 2.

1184

1185 **Figure 2. Total number of differentially abundant proteins for each MapMan main category.**
1186 Category description is given on the y-axis. Numbers of down- and up-regulated proteins across the 6
1187 experimental conditions are provided on the x-axis for each category.

1188

1189 **Figure 3. Metabolite profiles of Cabernet Sauvignon developing berries exposed (red line) or not**
1190 **(blue line) to heat stress.** Metabolites are shown in their respective metabolic pathways (sugar
1191 metabolism, glycolysis, and the TCA cycle) and their temporal profiles during berry treatment
1192 ($\mu\text{mol.g}^{-1}$ FW for glucose, and fructose; and nmol.g^{-1} FW for the others) are presented alongside. For
1193 each profile, the x-axis shows the developing stage (G: middle-green, V: véraison and R: middle-
1194 ripening) and the stress duration (1, 7 and 14 days).

1195

1196 **Figure 4. Correlation between the expression of proteins and mRNAs in developing control and**
1197 **HT berries.** DAPs and DEGs were selected using a 1.5 and 2 fold expression change, respectively,
1198 and an adjusted $p < 0.05$ (with FDR correction). Venn diagrams display the number of DAPs and
1199 DEGs (red: up-regulated and blue: down-regulated) according to the developing stage (A: middle-
1200 green, B; véraison and C: middle-ripening) and the stress duration (1 and 7 days).

1201

1202 **Figure 5. Schematic overview of heat stress effects on developing Cabernet Sauvignon berries.**
1203 This illustration summarizes the main findings deduced from transcriptomic, proteomic and
1204 metabolomics analyses (this work and (Lecourieux et al. 2017)). Up and down arrows reflect the
1205 accumulation levels of the corresponding transcripts, proteins or metabolites in heated berries.

1206

1207 **SUPPORTING INFORMATION**

1208 **Figure S1:** MapMan GO categories impacted at the transcriptomic and proteomic levels in heated
1209 grape berries

1210

1211 **Table S1:** All identified proteins with calculated differentials and corresponding statistics

1212

1213 **Table S2:** List and information about the differentially abundant proteins between control and heat
1214 stress conditions

1215

1216 **Table S3:** Differentially abundant proteins between control and heat stress condition at green stage

1217

1218 **Table S4:** Differentially abundant proteins between control and heat stress condition at veraison stage

1219

1220 **Table S5:** Differentially abundant proteins between control and heat stress condition at ripening stage

1221

1222 **Table S6:** List of "Protein metabolism" related DAPs

1223

1224 **Table S7:** List of "Stress" related DAPs

1225

1226 **Table S8:** List of "Secondary metabolism" related DAPs

1227

1228 **Table S9:** List of "carbohydrate and energy metabolism" related DAPs

1229

1230 **Table S10:** Effects of the HT and developmental stages on individual metabolites,

1231

1232 **Table S11:** List of overlapping DEGs and DAPs in Cabernet Sauvignon berries exposed or not to HT

1233

1234 **Table S12:** DAPs with potential role in thermotolerance.

1235

1236 **Table 1: Amount of identified proteins and differentially abundant proteins in control**
 1237 **and heat berries according to the developmental stage and the stress duration**

Samples	Total protein identified	Total protein quantified (≥ 2 peptides)	Total DAPs	Number of DAPs up-regulated	Number of DAPs down-regulated	Number of DAPs up- or down-regulated
All 6 conditions	3972	2279	592			
			→ DAPs in 1 condition (501)	139	362	--
			→ DAPs in > 1 condition (91)	16	61	14
G1D		1633	39	11	28	
G7D		1263	108	14	94	
V1D		1580	148	70	78	
V7D		1054	83	49	34	
R1D		1244	238	38	200	
R7D		857	84	13	71	


1238
 1239
 1240
 1241

1242 **Table 2. Deregulated proteins with potential role in heat-induced berry responses**
1243

Functional category	Grape accession number	Putative Name	Function	Related proteins - role described in the literature	References		
Stress	VIT_16s0050g01150	VvHSP90	Chaperone	AtHSP90	Role in thermotolerance	Hong & Vierling, 2001; Meiri and Breiman, 2009; Jacob et al., 2017	
	VIT_17s0000g07190	VvHSP101		AtHSP101			
	VIT_18s0041g01230	VvHSP70		AtHSP70			
	VIT_04s0008g01570	VvHSP17.6II		AtHSP17.6II			
	VIT_02s0025g04330	VvTLP	Osmotin	Thaumatococcus-like proteins	Enhance abiotic stress tolerance in many plant species	Le et al., 2018 de Jesus-Pires et al., 2019	
	VIT_02s0025g04290						
	VIT_15s0046g01560	VvBAG	Chaperone	AtBAG7 (BCL-2-ASSOCIATED ATHANOGENE 7)	Role in the unfolded protein response (UPR) to mitigate heat stress	Williams et al., 2010 Li et al., 2017	
	VIT_00s0769g00010	VvROF1	Peptidyl propyl isomerase	AtROF1 (AtFKBP62)	Role in thermotolerance	Meiri & Breiman, 2009	
	Protein synthesis	VIT_14s0006g01990	VveIF5	Translation initiation factor	AteIF5 PseIF5	Role in thermotolerance	Suragami et al., 2011; Zhang et al., 2017
		VIT_14s0108g01420	VvCNGC2	Cyclic nucleotide gated calcium channel	AtCNGC2 AtCNGC6 PpCNGCb	Primary heat sensor Role in thermotolerance	Saidi et al., 2009 Finka et al., 2012 Gao et al., 2012
Hormone homeostasis	VIT_13s0064g00810	VvNCED	9-cis-epoxycarotenoid dioxygenase		ABA biosynthesis Strigolactone biosynthesis Role in drought tolerance in grape	He et al., 2018 Jia et al., 2018	
	VIT_01s0010g01840	VvGER5		GEM (Glabra2 Expression Modulator)	ABA signalling pathway	Baron et al., 2014; Bhatta et al., 2018	
	VIT_18s0072g00380	VvASR		VvMSA	Responses to water deficit	Medici et al., 2014	
	VIT_17s0000g02750	VvRACK1	Receptor for activated kinase 1		Role in various signalling pathways including drought and salt stress	Islas-Flores et al., 2015	
	VIT_06s0009g00260	VvBSL	Protein phosphatase	BRI1-SUPPRESSOR1	Role in brassinosteroid signalling Role in stress tolerance	Maselli et al., 2014; Gruszka, 2018	
Redox regulation	VIT_05s0049g01090	VvGST	Glutathione S-transferase		Role in abiotic and biotic responses	Nianiou-Obeidat et al., 2017	
	VIT_19s0015g02610	VvGR	Glutathione reductase		Role in heat tolerance	Khanna-Chopra & Chauhan, 2015; Jegadeesan et al., 2018	
	VIT_07s0141g00039						
	VIT_18s0122g01320	VvCAT	Catalase		Role in basal thermotolerance	Vanderauwera et al., 2011	
	VIT_00s0615g00030	VvMTD	Mannitol dehydrogenase	VvMTD1	Resistance to abiotic and biotic stress	Patel & Williamson, 2016	
	VIT_00s0371g00050						
	VIT_06s0004g02890	VvPDI	Protein-disulfide isomerase		Role in abiotic stress tolerance and H ₂ O ₂ detoxification	Eggert et al., 2016	
	VIT_07s0005g04600	VvGSNOR	Chaperone S-nitrosoglutathione reductase	AtHOT5	Role in thermotolerance	Gould et al., 2003; Lee et al., 2008	
Co-factor and vitamin metabolism	VIT_09s0002g04490	VvPDX1.2	Pseudoenzyme	AtPDX1.2	Role under heat and oxidative stress conditions	Mocceand et al., 2014; Dell'Aglio et al., 2017	
	VIT_01s0127g00490	VvPANK	Pantothenate kinase		Induced in response to salt stress	Liu et al., 2013; Aloisi et al., 2016	

1244
1245

1246 **Figures:**
1247




1248

1249

1250 **Figure 1. Venn diagrams displaying the numbers of DAPs in heat-treated berries, according to**
1251 **the developmental stage (A) or the treatment duration (B).** DAPs were selected using a 1.5 fold
1252 expression change and an adjusted $p < 0.05$ (with FDR correction). Only proteins identified in all three
1253 biological replicates and with a minimum of two unique peptides were considered for quantification.
1254 The detailed information of the common DAPs upon HT for each stage or for each exposure time was
1255 listed as in Supplementary Tables 1 and 2.

1256



1257


1258

1259 **Figure 2. Total number of differentially abundant proteins for each MapMan main category.**

1260 Category description is given on the y-axis. Numbers of down- and up-regulated proteins across the 6

1261 experimental conditions are provided on the x-axis for each category.

1262



1263

1264

1265


1266

1267

1268


1269

Figure 3. Metabolite profiles of Cabernet Sauvignon developing berries exposed (red line) or not (blue line) to heat stress. Metabolites are shown in their respective metabolic pathways (sugar metabolism, glycolysis, and the TCA cycle) and their temporal profiles during berry treatment ($\mu\text{mol.g}^{-1}$ FW for glucose, and fructose; and nmol.g^{-1} FW for the others) are presented alongside. For each profile, the x-axis shows the developing stage (G: middle-green, V: véraison and R: middle-ripening) and the stress duration (1, 7 and 14 days).



1270
1271
1272
1273
1274
1275
1276
1277

Figure 4. Correlation between the expression of proteins and mRNAs in developing control and HT berries. DAPs and DEGs were selected using a 1.5 and 2 fold expression change, respectively, and an adjusted $p < 0.05$ (with FDR correction). Venn diagrams display the number of DAPs and DEGs (red: up-regulated and blue: down-regulated) according to the developing stage (A: middle-green, B; véraison and C: middle-ripening) and the stress duration (1 and 7 days).



1278

1279

1280 **Figure 5. Schematic overview of heat stress effects on developing Cabernet Sauvignon berries.**

1281 This illustration summarizes the main findings deduced from transcriptomic, proteomic and
 1282 metabolomics analyses (this work and (Lecourieux et al. 2017)). Up and down arrows reflect the
 1283 accumulation levels of the corresponding transcripts, proteins or metabolites in heated berries.

1284