



**HAL**  
open science

## Bioactive trace metals and their isotopes as paleoproductivity proxies: An assessment using GEOTRACES-era data

Tristan Horner, Susan Little, Tim Conway, Jesse Farmer, Jennifer Hertzberg,  
Alastair Lough, Jennifer Mckay, Allyson Tessin, Stephen Galer, Sam Jaccard,  
et al.

### ► To cite this version:

Tristan Horner, Susan Little, Tim Conway, Jesse Farmer, Jennifer Hertzberg, et al.. Bioactive trace metals and their isotopes as paleoproductivity proxies: An assessment using GEOTRACES-era data. *Global Biogeochemical Cycles*, In press, pp.e2020GB006814. 10.1029/2020GB006814 . hal-03003951v1

**HAL Id: hal-03003951**

**<https://hal.science/hal-03003951v1>**

Submitted on 19 Nov 2020 (v1), last revised 23 Nov 2021 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Bioactive trace metals and their isotopes as paleoproductivity proxies: An assessment using GEOTRACES-era data

T.J. Horner<sup>1,2,\*</sup>, S.H. Little<sup>3,\*</sup>, T.M. Conway<sup>4,\*</sup>, J.R. Farmer<sup>5,6,\*</sup>, J.E. Hertzberg<sup>7</sup>, A.J.M. Lough<sup>8,9</sup>, J. McKay<sup>10</sup>, A. Tessin<sup>11</sup>, S.J.G. Galer<sup>6</sup>, S.L. Jaccard<sup>12</sup>, F. Lacan<sup>13</sup>, A. Paytan<sup>14</sup>, K. Wuttig<sup>15</sup>, and GEOTRACES–PAGES Biological Productivity Working Group Members<sup>16</sup>.

<sup>1</sup>NIRVANA Labs, <sup>2</sup>Department of Marine Chemistry & Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA, USA; <sup>3</sup>Department of Earth Sciences, University College London, London, GBR; <sup>4</sup>College of Marine Science, University of South Florida, FL, USA; <sup>5</sup>Department of Geosciences, Princeton University, Princeton, NJ, USA ; <sup>6</sup>Max-Planck Institute for Chemistry, Mainz, DEU; <sup>7</sup>Department of Ocean, Earth & Atmospheric Sciences, Old Dominion University, Norfolk, VA, USA; <sup>8</sup>National Oceanography Centre, University of Southampton, Southampton, GBR; <sup>9</sup>Now at: School of Geography, University of Leeds, Leeds, GBR; <sup>10</sup>College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, OR, USA; <sup>11</sup>Department of Geology, Kent State University, Kent, OH, USA; <sup>12</sup>Institute of Geological Sciences and Oeschger Center for Climate Change Research, University of Bern, Bern, CHE; <sup>13</sup>LEGOS, University of Toulouse, CNRS, CNES, IRD, UPS, Toulouse, FRA; <sup>14</sup>Institute of Marine Sciences, University of California Santa Cruz, Santa Cruz, CA, USA; <sup>15</sup>Antarctic Climate and Ecosystems Cooperative Research Centre, University of Tasmania, Hobart, AUS; <sup>16</sup>A full list of working group members and their affiliations appears at the end of the manuscript.

\*To whom correspondence should be addressed: T.J.H. ([Tristan.Horner@whoi.edu](mailto:Tristan.Horner@whoi.edu)), S.H.L. ([susan.little@ucl.ac.uk](mailto:susan.little@ucl.ac.uk)), T.M.C. ([tmconway@usf.edu](mailto:tmconway@usf.edu)), or J.R.F. ([jesse.farmer@princeton.edu](mailto:jesse.farmer@princeton.edu)).

## Key Points

- This review assesses the potential of a number of bioactive trace elements and their isotopes to inform on past ocean productivity
- Distributions, drivers, and depositional archives are described for iron, zinc, copper, cadmium, molybdenum, barium, nickel, and silver
- Future priorities include quantification of ‘missing’ flux terms, constraining circulation influences, and identifying sedimentary archives

**Keywords:** marine chemistry; iron; zinc; copper; cadmium; molybdenum; barium; nickel; silver.

## Abstract

1  
2 The ocean's biological carbon pump redistributes climatically-significant quantities of carbon from the  
3 atmosphere to the ocean interior and seafloor. How the biological pump operated in the past is therefore  
4 important for understanding past atmospheric carbon dioxide concentrations and Earth's climate history.  
5 Due to their intimate association with biological processes, several bioactive trace metals and their isotopes  
6 are thought to be promising proxies for productivity, including: iron, zinc, copper, cadmium, molybdenum,  
7 barium, nickel, and silver. Here we review the oceanic distributions, driving processes, and depositional  
8 archives for these eight elements and their isotopes based on GEOTRACES-era datasets. We offer an  
9 assessment of the overall maturity of each isotope system to serve as a proxy for diagnosing aspects of past  
10 ocean export productivity, and identify priorities for future research. Despite many of the elements reviewed  
11 here sharing a common biological driving processes, we show that key aspects of the biogeochemical cycle  
12 of each element are often unique. Rather than being a source of confusion, it is our hope that combining the  
13 unique perspectives afforded by each bioactive trace element will enable painting a more complete picture  
14 of marine paleoproductivity, biogeochemical cycles, and Earth's climate history.

## 15 1. Introduction

16 The ocean plays host to three carbon ‘pumps’ that redistribute climatically-significant quantities of carbon  
 17 dioxide (CO<sub>2</sub>) from the atmosphere to the ocean interior and seafloor (Volk & Hoffert, 1985). These ocean  
 18 carbon pumps—biological, carbonate, and solubility— influence Earth’s climate over timescales ranging  
 19 from decades to millions of years (e.g., Volk & Hoffert, 1985; Sigman et al., 2010; Khatiwala et al., 2019).  
 20 The biological pump is of particular interest as it connects the cycles of C to those of O<sub>2</sub>, (micro)nutrients,  
 21 and marine biology, and today accounts for as much as 70 % of the ‘contribution ’ of all three carbon pumps  
 22 (Sarmiento & Gruber, 2006). The biological pump redistributes atmospheric carbon in two steps. First,  
 23 phytoplankton, photoautotrophic microbes, use sunlight to transform ambient DIC (dissolved inorganic  
 24 carbon) into POC (particulate organic carbon), represented here by CO<sub>2</sub> and glucose, respectively, by the  
 25 simplified reaction:



27 The second step requires that some fraction of the newly-formed POC sinks into the ocean interior through  
 28 a combination of biological and physical aggregation processes (e.g., Alldredge & Silver 1988). The  
 29 resulting surface ocean DIC deficit promotes the invasion of atmospheric CO<sub>2</sub> into seawater to maintain  
 30 air–sea CO<sub>2</sub> equilibrium, driving an overall reduction in atmospheric *p*CO<sub>2</sub>. (This definition of the  
 31 biological pump neglects dissolved organic carbon export, which is comparatively understudied, though  
 32 may account for as much as one-third of C export; e.g., Carlson et al., 2010; Giering et al., 2014.)  
 33 Importantly, Reaction [1] requires sunlight and can only occur in the euphotic layer of the ocean. In contrast,  
 34 aerobic heterotrophic respiration can occur wherever POC and O<sub>2</sub> are present:



36 (There are a number of O<sub>2</sub>-independent respiration pathways that are reviewed in detail elsewhere; e.g.,  
 37 Froelich et al., 1979.)

38


39 While the representation of all POC as glucose (CH<sub>2</sub>O) is instructive for illustrating an important biotic  
 40 transformation in the ocean, it is also simplistic; microbial biomass consists of dozens of bioactive elements  
 41 that serve many essential functions (e.g., da Silva & Williams, 1991). The elemental stoichiometry of POC  
 42 can thus be expanded to include a number of major and micronutrient elements, as illustrated by the  
 43 extended Redfield ratio reported by Ho et al. (2003):



45 With this extended stoichiometry in mind, it is clear that Reactions [1] and [2]—the production and  
46 regeneration of organic matter, respectively—will not only generate gradients in the dissolved  
47 concentration of DIC and O<sub>2</sub>, but also for many other bioactive elements associated with POC cycling.  
48 These gradients will be steepest for those elements possessing shorter residence times and where biological  
49 uptake and regeneration are the most important processes driving their vertical distributions. Likewise, such  
50 gradients may be almost absent for elements that possess long residence times or are primarily cycled by  
51 processes disconnected from productivity.

52

53 For those bioactive metals where biological processes are important, the implication of Reactions [1] and  
54 [2] is that many of the metals listed in [3] may, in turn, be used as proxies of POC cycling and hence  
55 paleoproductivity. A key motivation for using these elements as tracers of (past) POC cycling is that  
56 bioactive metal distributions are often set over significant spatiotemporal scales. For example, temporal  
57 changes in POC fluxes from a single sediment core would, at most, reflect local changes in export  
58 productivity, though such reconstructions are oftentimes unreliable indicators of productivity owing to  
59 significant preservation biases (e.g., Rühlemann et al., 1999). Biases aside, building a regional picture of  
60 paleoproductivity in this manner would require sampling many regions of the seafloor (e.g., Cartapanis et  
61 al., 2016) and conducting many more analyses. In contrast, the flux and residence time of nutrients in the  
62 euphotic ocean can be diagnostic of the productivity of entire ecosystems (e.g., Dugdale & Goering, 1967).  
63 Indeed, large-scale features of past ocean productivity are routinely reconstructed using the abundance and  
64 stable isotopic compositions of macronutrient elements (C, N, and Si; see Farmer et al., *this issue*). It thus  
65 follows that the abundance and isotopic compositions of several bioactive trace elements and their isotopes  
66 (TEI's) cycled along with POC could also serve as valuable proxies for past export productivity. However,  
67 in order to use systems of trace elements and their isotopes as proxies for productivity, it is necessary to  
68 develop a comprehensive understanding of the marine behavior of these elements, including: mapping their  
69 distribution in the ocean; elucidating the drivers of the distribution; characterizing sources, sinks, and  
70 transformations associated with biological, physical and chemical (notably redox) reactions; and,  
71 recognizing if (and how) a given element is incorporated and preserved in marine sediments.



72  
 73 **Figure 1 | Overview of bioactive trace metal distributions discussed in this review.** **A** Map showing locations of  
 74 the representative depth profiles shown in both the lower panels and discussed throughout this review (Sections. 3–  
 75 10). The specific station (**bold**) and cruise identifier are given for each location; circles, squares, and triangles denote  
 76 stations in the Pacific, Atlantic, and Southern Oceans, respectively. The lack of exemplar stations from the Indian Ocean  
 77 reflects the current paucity of GEOTRACES-compliant campaigns in this basin. Two stations sampled from the Atlantic  
 78 Sector of the Southern Ocean are designated by a single symbol: St. S5 (GIPY4), situated at 57.55 S, 0.04 W, and  
 79 PS71-138 (GIPY5) at 61.00 S on the Zero Meridian. **B–I** Depth profiles of bioactive trace metal concentrations for Fe  
 80 (B), Zn (C), Cu (D), Cd (E), Mo (F), Ba (G), Ni (H), and Ag (I). Profiles are shown in semi-log space to illustrate variations  
 81 in the epi- and mesopelagic realm. Originators are cited in the Data Sources section. Map created using Ocean Data  
 82 View (Schlitzer, R., <https://odv.awi.de>, 2018).

83

84 The wealth of new TEI data from the GEOTRACES program (Fig. 1) now permits an assessment of whether  
85 certain bioactive metals are promising proxies for past ocean productivity. This contribution represents such  
86 an assessment. Our study synthesizes what is known about the processes governing the cycling of several  
87 bioactive TEIs in seawater, explores the level of development and readiness of each TEI system to inform  
88 on aspects of past ocean productivity, and identifies areas where further research is most needed to improve  
89 our understanding of the geochemistry of TEIs in the past and present ocean. We base our assessment on  
90 publicly-available results from the international GEOTRACES programme (e.g., Mawji et al. 2015;  
91 Schlitzer et al., 2018). Our study is not intended to be a thorough review of all available techniques used to  
92 reconstruct paleoproductivity. Instead, we focus on bioactive TEI systems that are either recognized as  
93 micronutrients (such as those in Eq. 3) or exhibit nutrient-like dissolved profiles in seawater, regardless of  
94 their nutritional status to phytoplankton (e.g., Ba, Ag). This will not be the last word on the topic; our  
95 understanding of many of these TEIs has rapidly evolved in recent years, and will continue to evolve as  
96 new data are generated.

97

98 This contribution is structured such that each bioactive TEI system is reviewed similarly and systematically.  
99 The order in which TEI's are assessed follows the extended Redfield ratio—iron (Fe; Sec. 3.), zinc (Zn;  
100 Sec. 4), copper (Cu; Sec. 5), cadmium (Cd; Sec. 6), and molybdenum (Mo; Sec. 7)—before describing the  
101 nonessential elements barium (Ba; Sec. 8), nickel (Ni; Sec. 9.), and silver (Ag, Sec. 10) in order of  
102 decreasing dissolved concentration in seawater. The assessment of each TEI system is organized around  
103 four questions:

- 104 1. What is the modern marine distribution of this TEI system?
- 105 2. Which biological, chemical, and physical processes are most important for maintaining this  
106 distribution?
- 107 3. In what form is this TEI system incorporated into sediments?
- 108 4. Are there clear priorities for improving the utility of this system to track paleoproductivity?

109 This structure results in some repetition of the main distributions, drivers, and sedimentary archives  
110 between individual TEI systems. This redundancy is deliberate: each section can be read independently  
111 without reference to other TEIs. We close our review by assessing the 'maturity' of each system based on  
112 a comparison to more established productivity proxies, offer suggestions for future studies, and discuss  
113 prospects for paleoproductivity reconstructions using bioactive TEI isotope systems.



## 114 2. Definitions

### 115 2.1. Paleoproductivity

116 What is (paleo)productivity? This definition is significant since the biological productivity of an ecosystem  
117 can be described by a number of nested C cycles, whereby only some sub-cycles contribute to the  
118 sequestration of atmospheric CO<sub>2</sub> as POC in the ocean interior and the geological record. Moreover,  
119 productivity estimates can diverge depending on the tracer (e.g., Bender et al., 1987) or reference depths  
120 (e.g., Buesseler et al., 2020) used as the basis of any calculation. Individual proxies for POC cycling may  
121 be sensitive to only certain components of the C cycle and could thus yield conflicting insights into  
122 paleoproductivity. For consistency with the macronutrient literature, we use the definitions described by  
123 Sigman & Hain (2012): GPP, NPP, and NEP, which correspond to gross primary production, net primary  
124 production, and net ecosystem (or ‘export’) production, respectively. Gross PP refers to total autotrophic  
125 production of POC (or O<sub>2</sub>; i.e., Eq. [1]). Net PP is defined as GPP minus autotrophic respiration—the  
126 metabolic O<sub>2</sub> requirements of primary producers—and is effectively the rate at which phytoplankton  
127 produce new biomass. Lastly, NEP refers to GPP minus all autotrophic *and* heterotrophic respiration in an  
128 ecosystem (i.e., NPP minus heterotrophic respiration). If considering only the sunlit surface ocean, NEP is  
129 equivalent to export production; the flux of POC from the surface ocean must, over sufficiently long  
130 timescales, balance the vertical supply of dissolved nutrients. Likewise, integrated over the entire euphotic  
131 ocean, NEP represents the upper limit of the CO<sub>2</sub> sequestering capacity of the biological pump: the amount  
132 of POC buried in marine sediments must be less than, or equivalent to, annual NEP (minus any contributions  
133 from terrestrial OC). Accordingly, NEP is perhaps the most relevant term to understanding how the  
134 biological pump draws down and stores atmospheric CO<sub>2</sub>. Indeed, variations in NEP are implicated as a  
135 key driver of glacial–interglacial variations in *p*CO<sub>2</sub> and hence climate (e.g., Broecker, 1982; Boyle, 1988a;  
136 Berger et al., 1989; Paytan, 2009). Thus, when describing and assessing the utility of each TEI system to  
137 inform on past productivity, we are specifically concerned with whether and how that system is related to  
138 NEP.

139

### 140 2.2. Key processes

141 In addition to only being sensitive to certain POC sub-cycles, the TEI systems reviewed here exhibit a  
142 number of instances where their cycling may be decoupled from those of POC. Broadly, these processes  
143 can be categorized as affecting the sources, internal cycling, or sinks of TEIs, but not always the  
144 macronutrients or C. These additional processes may lead to differences between, or even a complete



145 decoupling of (paleo)productivity estimates using TEI- and macronutrient-based productivity measures.  
 146 These three areas are introduced here and described in detail in the sections that follow.

147  
 148 First, many elements, including bioactive TEIs, can be decoupled from DIC cycling through ‘external’  
 149 sources that mask any POC-related drawdown. Though rarely significant for DIC, these sources—rivers,  
 150 dust, desorption from particles, sediments, and hydrothermalism—may be significant terms in local and  
 151 regional TEI budgets if the magnitude is comparable to the dissolved supply from ocean circulation.  
 152 Examples include dust-derived Cu to surface planktonic communities (e.g., Paytan et al., 2009) and  
 153 hydrothermal-derived Fe to the deep ocean (e.g., Resing et al., 2015). Consideration of these external  
 154 sources is particularly important when close to such point sources and for elements with residence times  
 155 that are less than or similar to the mixing time of the ocean (~1 kyr).

156  
 157 Second, the internal cycle of TEIs can be complex; TEI distributions can be decoupled from those of the  
 158 macronutrients through myriad interactions with particles, be they biological (uptake and remineralization),  
 159 physical (adsorption, desorption), or chemical (e.g., redox transformation leading to precipitation or  
 160 dissolution; e.g., Boyd et al., 2017).

161 Biological uptake itself may decouple TEIs from macronutrients since organisms appear to have wider  
 162 tolerances for the intracellular proportions of certain trace elements compared to those of C, N, and P. The  
 163 physiological mechanisms enabling this plasticity, and the feedback interactions that drive it, are beyond  
 164 the scope of this review, and are discussed in detail elsewhere (e.g., Sunda, 2012; Morel et al., 2020). From  
 165 a proxy perspective, this flexibility may cause uncertainty in paleoproductivity estimates; the more variable  
 166 the C:TEI stoichiometry of organisms within an ecosystem, the more uncertain the paleoproductivity  
 167 estimate derived from that trace element. (The corollary being that the more rigid the stoichiometry, the  
 168 more robust the paleoproductivity estimate.) An extreme example concerns nonessential elements (e.g.,  
 169 Ba), or metals that are only essential for certain groups of organisms within an ecosystem (e.g., Ni for  
 170 methanotrophs). Productivity estimates derived from the export of these nonessential elements are  
 171 potentially susceptible to decoupling from productivity cycles as their export is not intrinsically tied to the  
 172 overall functioning of an ecosystem.

173 Remineralization—the regeneration of POC to inorganic dissolved nutrients—can similarly decouple the  
 174 internal cycles of TEIs and macronutrients. In the case of a scarce nutrient, such as Fe, individual organisms  
 175 (e.g., Saito et al., 2011) and even entire ecosystems (e.g., Rafter et al., 2017) may have evolved mechanisms  
 176 to retain scarce resources. Likewise, macro- and micronutrient TEIs may be regenerated by heterotrophic

177 organisms at different rates (e.g., Twining et al., 2014; Ohnemus et al., 2019). Thus, the number of  
178 productivity cycles that must occur before a trace element is exported out of the euphotic zone may differ  
179 from that of C or other macronutrients.

180 The processes of adsorption and desorption can also fractionate TEI:macronutrient relationships. Originally  
181 developed in the context of dissolved and particulate thorium isotopes (Bacon & Anderson, 1982),  
182 reversible scavenging is now suggested to play a role in the vertical cycling of other metals, including Cu  
183 (e.g., Little et al., 2013) and Fe (e.g., Abadie et al., 2017). Reversible scavenging is a continuous process  
184 that occurs between particle surfaces and dissolved species. While this process can occur at any depth,  
185 scavenging intensity is positively correlated with the quantity of particles, and so is most important in the  
186 upper water column. Likewise, while dissolved TEIs may be scavenged by any class of particle (e.g., opal,  
187 lithogenics), recent modeling efforts indicate that certain particle types may preferentially scavenge certain  
188 elements (e.g., Lerner et al., 2018). Scavenging may also affect TEIs primarily cycled by organic matter;  
189 secondary phases may scavenge metals during remineralization, which could affect TEI distributions in the  
190 upper water column (e.g., Zn, John & Conway, 2014; Co, Hawco et al., 2018; Fe, Tagliabue et al., 2019).

191 Changes in the ambient redox environment, such as in an oxygen minimum zone (OMZ), may also enhance  
192 dissolved–particulate transformations for certain TEIs. Whether OMZs act as a source or a sink depends on  
193 the TEI; whereas some elements may be released during reductive dissolution of particulate Fe–Mn oxides,  
194 others may exhibit enhanced scavenging onto particles in strongly reducing environments.

195

196 Third, many TEIs have significant output fluxes that are not associated with organic matter. The processes  
197 mediating these burial fluxes are diverse, ranging from scavenging (e.g., Cu, Mo) to precipitation into  
198 (organo)minerals (e.g., Ba; Défarge, 2011). Consequently, changes in the non POC-associated burial flux  
199 term(s) may drive the ocean to a new TEI:C stoichiometry that could be difficult to recognize or interpret  
200 in the sedimentary record.

201

202 While these three categories of processes may appear insurmountable obstacles to the reliable application  
203 of TEIs to reconstruct paleoproductivity, there are reasons to be optimistic. Indeed, the fact that so many  
204 confounding processes are now recognized highlights just how far our understanding of TEI geochemistry  
205 has evolved in recent years. Moreover, many of these processes primarily concern the use of trace element  
206 abundances to reconstruct past productivity. As we show in the following sections, the processes controlling  
207 the abundance of an element may be distinct from that controlling its isotopic composition. These distinct  
208 controls may enable isotopic analyses to reduce, or even eliminate, certain ambiguities in

209 paleoceanographic data, highlighting a potentially valuable role for TEI systems in the reconstruction of  
 210 paleoproductivity.

211

### 212 **2.3. Isotope notation**

213 The bioactive TEI literature abounds with isotope notations (e.g.,  $\epsilon$ ,  $\delta$ ), reference materials (e.g., JMC,  
 214 NIST), and isotope ratio pairs (e.g.,  $^{57}\text{Fe}/^{54}\text{Fe}$  versus  $^{56}\text{Fe}/^{54}\text{Fe}$ ;  $^{137}\text{Ba}/^{134}\text{Ba}$  versus  $^{138}\text{Ba}/^{134}\text{Ba}$ ). There are  
 215 merits to each choice and it is not our intention to review these here. However, we believe that the sheer  
 216 number of ways in which TEI data have been reported can be confusing to scientists in other disciplines  
 217 and this confusion ultimately diminishes the reach and utility of TEI-based research. To avoid furthering  
 218 this confusion, we have adopted a number of conventions that apply throughout this review, regardless of  
 219 how literature data were originally reported. First, we use a single isotope notation throughout (' $\delta$ '; i.e., the  
 220 delta notation). Second, we express all data relative to the most widely accepted standard for each isotope  
 221 system. For many isotope systems, the most widely-accepted standard may have since been exhausted (e.g.,  
 222 JMC Lyon for Zn). In those cases, there are usually cross-calibrated secondary materials that can be used  
 223 to report new isotope data in terms of 'legacy' materials (e.g. AA-ETH for Zn, Archer et al., 2017). Third,  
 224 we report TEI data using the same isotope ratio pairs as used in the GEOTRACES data products (e.g.,  
 225 Mawji et al., 2015; Schlitzer et al., 2018). We note that isotope data are a unitless ratio quantity (Coplen,  
 226 2011), though are commonly reported with 'units' of ‰ (i.e., parts per one-thousand):

$$227 \quad \delta^x\text{TE} = R_{\text{sample}} / R_{\text{standard}} - 1 \quad [4]$$

228 where  $\delta^x\text{TE}$  represents  $\delta^{56}\text{Fe}$ ,  $\delta^{66}\text{Zn}$ ,  $\delta^{65}\text{Cu}$ ,  $\delta^{114}\text{Cd}$ ,  $\delta^{98}\text{Mo}$ ,  $\delta^{138}\text{Ba}$ ,  $\delta^{60}\text{Ni}$ , or  $\delta^{109}\text{Ag}$  and  $R$  represents  
 229  $^{56}\text{Fe}/^{54}\text{Fe}$ ,  $^{66}\text{Zn}/^{64}\text{Zn}$ ,  $^{65}\text{Cu}/^{63}\text{Cu}$ ,  $^{114}\text{Cd}/^{110}\text{Cd}$ ,  $^{98}\text{Mo}/^{95}\text{Mo}$ ,  $^{138}\text{Ba}/^{134}\text{Ba}$ ,  $^{60}\text{Ni}/^{58}\text{Ni}$ , or  $^{109}\text{Ag}/^{107}\text{Ag}$  in either a  
 230 sample or standard. For clarity, all isotopic data reviewed here have been calculated using notation in Eq.  
 231 4, and renormalized to the following standards, regardless of how the data originators reported their results:  
 232 IRMM-014 for Fe (e.g., Dauphas et al., 2017), JMC-Lyon for Zn (e.g., Moynier et al., 2017), NIST SRM  
 233 976 for Cu (*ibid.*), NIST SRM 3108 for Cd (e.g., Abouchami et al., 2013), NIST SRM 3134 +0.25 ‰ for  
 234 Mo (Nägler et al., 2014), NIST SRM 3104a for Ba (e.g., Horner et al., 2015a), NIST SRM 986 for Ni (e.g.,  
 235 Elliott & Steele, 2017), and NIST SRM 978a for Ag (e.g., Woodland et al., 2005).

236

237 We summarize salient features of each TEI in Table 1. The review of each TEI system follows.

238

239

240  
241

**Table 1 | Summary of oceanic concentrations, isotopic compositions, and mean ocean residence times for bioactive TEIs discussed in this review. Superscripts denote references listed in the Data Sources section.**

<b>Element</b>	<b>Dissolved concentration range (nmol kg<sup>-1</sup>)</b>	<b>Mean Upper Continental Crust composition (‰)</b>	<b>Mean deep ocean isotopic composition (‰)</b>	<b>Range of deep ocean isotope compositions (‰)</b>	<b>Residence time estimates (kyr)</b>
<b>Fe</b>	0.01–100 <sup>a</sup>	+0.1 <sup>d</sup>	Variable <sup>k</sup>	-2.4 to +1.5 <sup>k</sup>	0.004–0.6 <sup>r</sup>
<b>Zn</b>	0.01–10 <sup>a</sup>	+0.3 <sup>e</sup>	≈+0.5 <sup>l</sup>	-0.2 to +0.6 <sup>l</sup>	1–11 <sup>r</sup>
<b>Cu</b>	0.5–4 <sup>a</sup>	+0.1 <sup>e</sup>	≈+0.7 <sup>m</sup>	+0.6 to +0.8 <sup>m</sup>	2–5 <sup>r</sup>
<b>Cd</b>	0.00003–1.2 <sup>a</sup>	0.0 <sup>f</sup>	≈+0.3 <sup>n</sup>	+0.2 to +0.4 <sup>n</sup>	22–105 <sup>r</sup>
<b>Mo</b>	100 <sup>b</sup>	+0.4 <sup>g</sup>	≈+2.3 <sup>o</sup>	Homogeneous <sup>o</sup>	440 <sup>s</sup>
<b>Ba</b>	35–160 <sup>a</sup>	0.0 <sup>h</sup>	≈+0.3 <sup>p</sup>	+0.2 to +0.4 <sup>p</sup>	8 <sup>t</sup>
<b>Ni</b>	1.5–11 <sup>a</sup>	+0.1 <sup>i</sup>	≈+1.3 <sup>q</sup>	Homogeneous <sup>q</sup>	10–30 <sup>u</sup>
<b>Ag</b>	0.0002–0.1 <sup>c</sup>	0.0 <sup>j</sup>	Unknown	Unknown	0.4 <sup>v</sup>

### 242 3. Iron

243 Iron (Fe) plays a key role within phytoplankton as an electron carrier for photosynthesis and respiration  
244 processes, as well as within enzymes necessary for photosynthesis and nitrogen fixation (Morel & Price,  
245 2003). However, in oxygenated seawater, Fe(II) is rapidly oxidized to Fe(III), which is highly insoluble  
246 (Liu & Millero, 2002). Intense biological demand coupled to low solubility results in generally sub-  
247 nanomolar concentrations for dissolved Fe throughout the oceans, and Fe concentrations approach the low  
248 picomolar range in some surface regions far from Fe sources, such as the vast Southern Ocean (Chever et  
249 al., 2010; Klunder et al., 2011; Schlitzer et al., 2018). Consequently, biological production in about 30 %  
250 of the modern surface ocean is thought to be limited primarily by Fe (Moore et al., 2013), principally in  
251 upwelling regions where deep water is depleted in Fe relative to the macronutrients nitrate and phosphate  
252 (e.g., Moore, 2016; Boyd et al., 2017). In these regions, termed ‘High Nutrient Low Chlorophyll’ (HNLC),  
253 Fe supply can limit primary productivity (e.g., Martin & Fitzwater, 1988) and potentially the exchange of  
254 carbon between the ocean and atmosphere. Furthermore, a changing supply of Fe from sources such as  
255 atmospheric dust, hydrothermal venting, or sedimentary release to the surface oceans through geological  
256 time can exert a significant control on both the distribution of primary productivity in the oceans and,  
257 through this, the global carbon cycle. Changes in supply of dust to the Fe-limited Southern Oceans has been  
258 shown to correlate with climate variability on millennial time scales and has also been invoked to explain  
259 the dramatic sawtooth glacial–interglacial shifts in atmospheric carbon dioxide (Martin, 1990; Sigman &  
260 Boyle, 2000; Martínez-García et al., 2011; 2014). The strong link between Fe supply and primary  
261 productivity means that ratios of Fe to other elements and/or Fe isotope ratios ( $\delta^{56}\text{Fe}$ ) might be useful  
262 proxies for investigating changes in paleoproductivity, provided the processes that fractionate them are  
263 dominantly linked to primary production, can be constrained, and suitable archives identified.

264


#### 265 3.1. Marine distribution

266 The distribution of dissolved Fe in seawater is driven by a mixture of competing processes, including  
267 biological uptake and (deeper) regeneration; distinct sources of Fe at shallow, intermediate, and deep  
268 depths; adsorption/desorption processes onto organic and lithogenic particles; dissolution and precipitation  
269 processes; and complexation to organic ligands (Fig. 2; e.g., Boyd & Ellwood, 2010; Labatut et al. 2014;  
270 Tagliabue et al., 2017). As a result, dissolved Fe displays what has historically been termed as hybrid-type  
271 depth profile in the open ocean, which exhibits a number of similarities between different ocean basins  
272 (Fig. 3). Surface Fe is drawn down in surface waters by biological uptake, and can even be as low as 0.01  
273 nmol kg<sup>-1</sup> in Fe limited regions. However, some areas of the oceans, such as the ‘dusty’ North Atlantic,

274 surface Fe concentrations can be driven as high as  $2 \text{ nmol kg}^{-1}$  as a result of dust events (Sedwick et al.,  
275 2005). Below the surface mixed layers, regeneration of biogenic material, reversible particle scavenging,  
276 and complexation by organic ligands act to keep the background deep ocean Fe concentrations at around  
277  $0.4\text{--}0.6 \text{ nmol kg}^{-1}$  (e.g., Lauderdale et al., 2020). Against this background, it has been known for decades  
278 that Fe concentrations in deeper waters are elevated near point sources such as sedimentary margins  
279 (Johnson et al., 1999). Recently, a range of studies including those conducted as part of the GEOTRACES  
280 program, have illuminated this picture, showing that deep sources of Fe—such as sedimentary and  
281 hydrothermal release—are widespread, may have distinct traceable  $\delta^{56}\text{Fe}$  source signatures, and that this  
282 Fe can be transported over large distances through the ocean interior (e.g., Radic et al. 2011; Saito et al.,  
283 2013; Conway and John, 2014; Resing et al., 2015; Nishioka et al., 2020). Despite exhibiting well-defined  
284 deep maxima close to point sources, Fe distributions at shallow and intermediate depths are much more  
285 variable and models have struggled to reproduce these variations (e.g., Tagliabue et al., 2016). The extent  
286 to which deeply sourced Fe is supplied to surface seawater is thus equivocal (c.f., Tagliabue et al., 2010;  
287 Roshan et al., 2020).

288  
289 The origin of Fe variability in the shallow and intermediate ocean is thought to reflect local differences in  
290 the competition between uptake, regeneration, sources, and scavenging. The same processes influence  
291  $\delta^{56}\text{Fe}$ ; however, unlike variations in Fe concentrations,  $\delta^{56}\text{Fe}$  exhibits dramatic variability between—and  
292 even within—ocean basins (Fig. 2; Schlitzer et al., 2017). Such water column variability in  $\delta^{56}\text{Fe}$ , from  
293 isotope compositions as light as  $-3 \text{ ‰}$  (John et al., 2012), to as heavy as  $\sim +1.5 \text{ ‰}$  (Ellwood et al., 2020),  
294 is thought to be driven by Fe source signatures and therefore also oceanic circulation, and a combination of  
295 biological uptake, Fe complexation to organic ligands in surface waters, and non-reductive release of  
296 dissolved Fe from particles (notably lithogenic particles) during desorption and/or ligand-promoted  
297 dissolution (e.g. Conway and John, 2014; Abadie et al. 2017; Ellwood et al., 2020; Fig. 3). As such,  $\delta^{56}\text{Fe}$   
298 may provide insight to the contribution of Fe sources at the basin scale (e.g. Conway & John, 2014), as well  
299 as for Fe cycling processes such as uptake and regeneration.

300




301  
 302 **Figure 2 | Representative profiles of dissolved Fe concentrations ( $[Fe]$ ) and Fe-isotopic compositions ( $\delta^{56}Fe$ )**  
 303 **in the Eastern Tropical North Atlantic (squares, dashed line) and Northeast Pacific (circles, solid line) Oceans.**  
 304 This comparison illustrates that despite possessing similar dissolved concentration profiles, the isotopic behavior of Fe  
 305 is markedly different between basins.  
 306

307 Despite numerous studies of both small- (e.g., Martin et al., 1990; Coale, 1991; Coale et al., 2003) and  
 308 large-scale iron fertilization (e.g., de Baar et al., 2005; Boyd et al., 2007; Boyd & Ellwood, 2010), there are  
 309 still uncertainties in how changing Fe supply to the surface ocean may affect phytoplankton growth in Fe-  
 310 limited regions, and/or nitrogen fixers under nitrate limitation, leading to gaps in our understanding of  
 311 linking Fe cycling directly to climate change (Misumi et al., 2014). One major challenge is assessing what  
 312 portion of the dissolved Fe pool is available for uptake by different microbes, termed ‘bioavailable’ Fe.  
 313 Such uncertainties on Fe supply, speciation, and bioavailability in the surface ocean are compounded by  
 314 limitations in our ability to constrain the supply of dissolved Fe in upwelled deep waters. For example,  
 315 while the ratio of C to macronutrients such N and P in the deep ocean is well known and the residence time  
 316 and distribution can be accurately reproduced using apparent oxygen utilization (AOU), this is not the case  
 317 for Fe. Only around 10–15 % of cellular Fe ( $Fe:C = 18\text{--}33 \mu\text{mol}:\text{mol}$ ) appears to be regenerated in the deep  
 318 ocean ( $Fe:C = 4\text{--}6$ ; Twining & Baines, 2013), leading to a weaker correlation between Fe and AOU, even in  
 319 regions away from Fe sources (Rijkenberg et al., 2014) and upwelling deep waters that are depleted in



320 dissolved Fe relative to macronutrients. Since incubation experiments show that Fe associated with sinking  
 321 organic matter from the subsurface is effectively regenerated (Velasquez et al., 2016), much of the released  
 322 Fe must be rapidly scavenged (Tagliabue et al., 2019). Despite this scavenging, vertical transport is still  
 323 thought to be the major source of dissolved Fe for phytoplankton in most Fe-limited regions, indicating that  
 324 additional research into the interplay between Fe supply (sources, stabilization, and transport) and demand  
 325 (biological uptake, scavenging) are required.  
 326  
 327



328  
 329 **Figure 3 | Processes driving iron isotope variations in modern seawater.** The oceans' internal cycle of Fe is  
 330 perhaps the most complex of the TEIs discussed here, exhibiting several significant sources, sinks, and transformations  
 331 not directly associated with biological productivity.

332  
 333 **3.2. Driving processes**

334 **3.2.1. Biological**

335 Initial studies of processes that fractionate Fe isotopes were optimistic that  $\delta^{56}\text{Fe}$  would make for a powerful  
 336 proxy of physiological 'biosignatures' (Beard et al., 1999), especially once measurement of  $\delta^{56}\text{Fe}$  was

337 extended to seawater by Lacan et al. (2008). However, much of this early optimism faded once it was found  
338 that other factors were also important in setting dissolved  $\delta^{56}\text{Fe}$ , such as external Fe sources (e.g., Radic et  
339 al., 2011; John et al., 2012; Sec. 3.2.3.). Moreover, organisms exhibit significant variability in Fe:C ratios  
340 (e.g., Twining & Baines, 2013), suggesting that Fe cycling may be partially decoupled from ecosystem  
341 productivity. Despite the more nuanced picture, there is a growing body of evidence suggesting that  
342 phytoplankton probably preferentially incorporate light Fe isotopes from seawater and, in some  
343 circumstances, can render detectable changes in dissolved  $\delta^{56}\text{Fe}$ . For example, studies from isolated eddies,  
344 the Mertz Polynya, and the open Southern Ocean showed that surface  $\delta^{56}\text{Fe}$  at picomolar dissolved Fe  
345 concentrations are isotopically heavy ( $>+1\text{‰}$ ), which has been attributed to the combination of surface  
346 uptake, regeneration, and organic complexation (Lacan et al. 2008; Ellwood et al., 2015; 2020). Estimates  
347 for the magnitude of fractionation due to biological uptake range between  $-0.1$  (Radic et al., 2011) and  $-1.0$   
348  $\text{‰}$  (Ellwood et al., 2020). The magnitude and direction of any Fe isotope fractionation during uptake may  
349 depend on the phytoplankton species, uptake mechanism, and Fe species consumed. Additional research is  
350 required on all three fronts. Below the surface mixed layer,  $\delta^{56}\text{Fe}$  of dissolved Fe appears to be primarily a  
351 reflection of the isotope signature of Fe sources, rather than reflecting a dominant influence from the  
352 biological processes of remineralization (e.g. Abadie et al., 2017; Conway & John, 2014a; John et al.,  
353 2018a; Labatut et al., 2014). Accordingly, there remains the possibility for  $\delta^{56}\text{Fe}$  of surface seawater to be  
354 linked to productivity in some oceanographic settings, though the relative role of other Fe-cycling processes  
355 and the degree to which they erase any diagnostic productivity signatures requires further investigation.

356

### 357 3.2.2. *Chemical*

358 The chemical behavior of Fe in seawater is complex and has the potential to decouple Fe cycling from  
359 macronutrients and thus productivity. Unlike the macronutrients, which are present as aqueous ions in  
360 solution, Fe is scarcely soluble in seawater, and much of what constitutes ‘dissolved’ Fe—operationally  
361 defined as that which can pass through a 0.2 or 0.4  $\mu\text{m}$  filter—is in actuality a soup containing organic  
362 complexes, nanoparticles, colloids, and a small fraction of truly ionic Fe. The controls governing, and the  
363 extent to which exchange occurs between these forms of dissolved Fe, are areas of focused interest (e.g.,  
364 Fitzsimmons & Boyle, 2014; Fitzsimmons et al., 2015). Additionally, dissolved Fe is subject to strong  
365 removal via scavenging, which lowers the Fe:macronutrient ratio of waters returned to the surface via  
366 upwelling (Moore, 2016). Chemical processes can also exert a significant influence over the isotopic

367 composition of Fe in seawater, such as through redox transformations, exchange reactions (e.g.,  
368 complexation, particle interactions), and by authigenic precipitation, discussed below.

369 Redox transformations drive large Fe isotope effects (e.g., Johnson et al., 2002; Skulan et al., 2002; Welch  
370 et al., 2003; Anbar et al., 2005). Indeed, much of the Fe isotope variation in Earth's ancient, more reducing  
371 past likely derives from fractionations associated with redox transformations (e.g., Johnson et al., 2008).  
372 While the role of redox relative to other processes is diminished in today's largely-oxygenated ocean, it is  
373 nonetheless an important mediator of Fe isotope source compositions, particularly within the ocean interior.  
374 This is neatly illustrated using the example of sediment dissolution, which can occur with or without a  
375 change in the redox state of Fe. Bulk marine sediments typically possess a composition similar to the crustal  
376 composition of +0.1 ‰ (Beard et al., 2003; Poitrasson, 2006). However, dissolved Fe(II) derived from  
377 bacterially-mediated reductive dissolution in sediments has been characterized by  $\delta^{56}\text{Fe}$  between -1 and -4  
378 ‰ (Berquist & Boyle, 2006; Severmann et al., 2006; 2010; Homoky et al., 2009; 2013; Klar et al., 2017a;  
379 Henkel et al., 2018), whereas Fe derived from non-reductive dissolution processes is thought to be  
380 considerably heavier, between +0.1 to +0.3‰ (Homoky et al., 2009; 2013; Radic et al., 2011). Further  
381 modification of reductive end-member compositions is possible upon contact with oxidizing seawater,  
382 potentially masking true source signatures. Oceanic water column dissolved  $\delta^{56}\text{Fe}$  compositions attributed  
383 to either non-reductive (+0.1 to +0.4 ‰) or reductive (-0.3 to -3.3 ‰) release of Fe from sediments have  
384 now been observed globally (Radic et al., 2011; John et al., 2012; Staubwasser et al., 2013; Conway &  
385 John, 2014a; Labutut et al., 2014; Conway & John 2015a; Chever et al., 2015; Fitzsimmons et al., 2016;  
386 Klar et al., 2017a; Abadie et al., 2017; Klar et al., 2018; Rolison et al., 2018; John et al., 2018a; Charette et  
387 al., 2020).

388 Exchange reactions can also fractionate primary Fe isotope compositions. For example, natural lithogenic  
389 dust is thought to possess a relatively narrow range of Fe isotope compositions (i.e.,  $\delta^{56}\text{Fe} \approx +0.1 \pm 0.2\text{‰}$ ;  
390 Waeles et al., 2007; Mead et al., 2013; Conway et al., 2019; Chen et al., 2020), reflecting the overall  
391 homogeneity of the upper continental crust. However, dissolved Fe in seawater attributed to dissolving dust  
392 particles is isotopically heavy, around +0.7 ‰ (Conway & John, 2014a). This fractionation is thought to  
393 reflect dissolution in concert with, and complexation by, strong (organic) ligands (Fishwick et al., 2014),  
394 which have been experimentally shown to preferentially bind heavy Fe isotopes (Dideriksen et al., 2008;  
395 Morgan et al., 2010). Fractionation effects may also arise during exchange of Fe between dissolved and  
396 particulate forms, though the magnitude of the effect depends on whether the exchange is primarily physical

397 (negligible fractionation; e.g. Fitzsimmons et al., 2017) or chemical (from  $\approx+0.3$  up to  $+1$  ‰; Labutut et  
398 al., 2014; Fitzsimmons et al., 2015).

399 Lastly, authigenic precipitation can control the isotopic composition of Fe released by large point sources,  
400 such as hydrothermal vents and margin sediments. Iron in hydrothermal vent fluids possesses end-member  
401 compositions ranging from  $-0.7$  to  $+0.1$ ‰ (Sharma et al., 2001; Beard et al., 2003; Severmann et al., 2004;  
402 Rouxel et al., 2008; Bennett et al., 2009; Rouxel et al., 2016; Nasemann et al., 2018; Rouxel et al., 2018).  
403 However, precipitation of Fe into authigenic minerals can render significant changes in dissolved  $\delta^{56}\text{Fe}$   
404 (e.g., Severmann et al., 2004; Bennett et al., 2009). The direction of fractionation depends on (e.g., Lough  
405 et al., 2017)—and may thus be diagnostic of (e.g., Horner et al., 2015b)—the specific transformations  
406 occurring: Fe sulfides and oxides preferentially incorporate isotopically light and heavy Fe, respectively  
407 (e.g., Skulan et al. 2002; Rouxel et al., 2008). Mineral precipitation can drive  $\delta^{56}\text{Fe}$  of residual Fe stabilized  
408 in seawater to values ranging between  $-2.4$  to  $+1.5$  ‰, depending on the authigenic mineral produced  
409 (Conway & John, 2014a; Ellwood et al., 2015; Fitzsimmons et al., 2016; 2017; Klar et al., 2017b; Lough  
410 et al., 2017; Rouxel et al., 2018). Analogous processes appear to operate along continental margins,  
411 whereby ‘light’ Fe, mobilized by reductive dissolution, encounters oxidizing seawater and forms  
412 precipitates that are heavier than the source Fe (though overall still considerably lighter than background  
413 seawater; e.g., Marsay et al., 2018).


414 Additional nuances relating to  $\delta^{56}\text{Fe}$  signatures of local point sources such as Fe released from  
415 anthropogenic aerosol dust, glaciers, icebergs, and rivers are also emerging, with such sources varying  
416 dramatically over short spatial and temporal scales and with *in situ* chemical reactions (e.g. Fantle &  
417 DePaolo, 2004; Berquist & Boyle, 2006; Ingri et al., 2006; Escoube et al., 2009; Poitrasson et al., 2014;  
418 Mead et al., 2013; Ilina et al., 2013; Akerman 2014; Chen et al., 2014; Escoube et al., 2015; Mullholland  
419 2015; Zhang et al., 2015; Kurisu et al., 2016a;b; Stevenson et al., 2017; Conway et al., 2019). The net effect  
420 of the dominance of Fe source on dissolved  $\delta^{56}\text{Fe}$  in seawater means that any signal from biological uptake  
421 in the Fe-depleted mixed layer, even if could be recorded in an archive, may be easily overprinted by even  
422 a small addition of new Fe from either above or below.

423

### 424 3.2.3. Physical

425 The residence time of Fe in seawater is substantially less than the mixing time of the global ocean (Table  
426 1). Local and regional Fe sources can thus drive large differences in dissolved [Fe] and  $\delta^{56}\text{Fe}$  between ocean  
427 basins (Fig. 3). Local source signatures—[Fe],  $\delta^{56}\text{Fe}$ , and perhaps Fe speciation—can be transported and

428 retained over the scale of individual ocean basins (Fig. 4; Conway & John, 2014a; Abadie et al., 2017).  
 429 This is arguably one of the marquee findings of the GEOTRACES program, and underpins the utility of  
 430 sedimentary  $\delta^{56}\text{Fe}$  to reconstruct past marine Fe sources, discussed next.



431  
 432 **Figure 4 | Three-dimensional scene depicting dissolved iron concentrations in the Atlantic Ocean** (Schlitzer,  
 433 2017). This perspective illustrates the density of GEOTRACES data in the region as well as the influence of multiple  
 434 Fe 'hot spots', such as mid-ocean ridges and continental margins. Data are available in the GEOTRACES Intermediate  
 435 Data Product 2017 (Schlitzer et al., 2018); names of data originators appear in the figure.

436

437

### 438 3.3. Marine archives

#### 439 3.3.1. Surface ocean

440 A requirement of the application of any paleoproxy is the availability of suitable sedimentary archives.

441 These archives must have both a high fidelity for the signal of interest and be robust to post-depositional



442 alteration. Archives most relevant to reconstructing paleoproductivity should capture surface ocean  $\delta^{56}\text{Fe}$ ;  
443 however, there are few—if any—reliable archives. The lack of surface-water archives reflects two related  
444 challenges. First, most archives for surface seawater are derived from biominerals, such as foraminifera,  
445 diatoms, sponges, and corals. These archives possess vanishingly low Fe concentrations, such that Fe  
446 isotopic analysis of these substrates has proven difficult. Second, following burial, many biominerals will  
447 act as substrates for authigenic mineral formation. These authigenic minerals, such as clays (e.g., Badaut &  
448 Risacher, 1983) and Fe–Mn oxides (e.g., Boyle, 1981), possess Fe concentrations far in excess of those in  
449 the underlying biomineral, necessitating significant physical and chemical cleaning (e.g., Cheng et al.,  
450 2000).

451 Despite these obstacles, there are three positive signs that reconstructing past surface seawater  $\delta^{56}\text{Fe}$  is  
452 possible. First, biogenic opal may contain Fe at concentrations in the  $\mu\text{g g}^{-1}$  range (Ellwood & Hunter, 2000;  
453 Lal et al., 2006; Shemesh et al., 1988; Sun et al., 2016), which is tractable for  $\delta^{56}\text{Fe}$  analysis. Second, the  
454 Fe content of diatoms is correlated with ambient dissolved [Fe] (Twining & Baines, 2013). Lastly, the  
455 positive relationship between the Fe content of diatoms and corresponding seawater appears to hold through  
456 sinking and sedimentation (Pichevin et al., 2014), indicating that diatoms are a potential window into past  
457 surface ocean Fe chemistry. Whether these relationships also extend to  $\delta^{56}\text{Fe}$  remains to be seen, and will  
458 require additional core-top calibrations, incubation experiments, and detailed assessment of the efficacy of  
459 chemical cleaning.

460

### 461 3.3.2. *Deep ocean*

462 In contrast to surface seawater, there are a number of studies examining Fe sources and fluxes in the meso-  
463 and bathypelagic ocean. To constrain Fe sources, researchers have examined the Fe isotope composition of  
464 Fe-rich sediments, including red clays (Tegler et al., 2018), polymetallic nodules (Marcus et al., 2015), and  
465 Fe–Mn crusts (Zhu et al., 2000; Levasseur et al., 2004; Chu et al., 2006; Horner et al., 2015b; Liu et al.,  
466 2020). Ferromanganese crusts are currently the best-studied for Fe isotopes (Fig. 5); Fe–Mn crusts are  
467 slowly-accumulating deposits ( $\sim\text{mm Myr}^{-1}$ ) that record ambient seawater  $\delta^{56}\text{Fe}$  with a spatially-invariant  
468 offset of  $-0.77\pm 0.06\text{‰}$  (Horner et al., 2015b). The constancy of the offset implies that the Fe isotope  
469 composition of individual Fe–Mn crust layers can be interpreted in terms of past dissolved  $\delta^{56}\text{Fe}$ , and thus  
470 past Fe sources. Curiously, however, the Fe-isotopic variability of Fe–Mn deposits recovered from the  
471 central Pacific—particularly layers formed before  $\sim 20\text{ Ma}$ —exceed the range of modern deep-ocean  $\delta^{56}\text{Fe}$   
472 (though only in the positive direction; Fig. 5). The reasons for elevated  $\delta^{56}\text{Fe}$  in the past are debated. Horner  
473 et al. (2015b) report that heavy  $\delta^{56}\text{Fe}$  could arise through widespread secondary modification of large Fe

474 sources through authigenic reactions, such as sulfide precipitation. In contrast, Johnson et al. (2020) contend  
475 that the elevated  $\delta^{56}\text{Fe}$  reflects extensive biological modification of dissolved Fe, driven by large-scale Fe  
476 fertilization. Regardless, the variation in these records points to a dynamic and enigmatic Fe cycle in Earth's  
477 past, and indicates that Fe–Mn crusts have a largely untapped potential to reconstruct spatiotemporal  
478 variations in this cycle.

479


480 Other approaches are also showing promise to study Fe fluxes to the deep ocean through time. For example,  
481 researchers have constrained the rate of sedimentary accumulation of hydrothermally-derived Fe and Cu  
482 using constant flux proxies such as extraterrestrial helium-3 (e.g., Middleton et al., 2016) or thorium-230  
483 (Costa et al., 2017). These studies report that hydrothermal activity may be coherent with sea-level changes  
484 on Quaternary glacial–interglacial cycles, suggesting a potentially remarkable set of connections between  
485 the solid Earth, ocean chemistry, and global climate (e.g., Cullen & Coogan, 2017). Reconstruction of  
486 hydrothermal metal fluxes over million-year timescales may also be possible using the geochemistry of  
487 pelagic clays (e.g., Dunlea et al., 2015), though such approaches are still in their infancy.

488

489 Although existing Fe-isotopic and Fe flux records cannot be directly interpreted in terms of past  
490 productivity, the supply and cycling of Fe in seawater is an important control over the biological  
491 productivity of the ocean. Thus, records of past oceanic Fe sources may prove most valuable as context for



492 interpreting other proxy records, or for generating novel hypotheses regarding the connections between the  
 493 Fe cycle and global climate.  
 494



495  
 496 **Figure 5 | Ferromanganese crust records of central Pacific  $\delta^{56}\text{Fe}$  since the Late Cretaceous.** Records from CD29-  
 497 2, 28DSR9, and METG-03 from Horner et al. (2015b), Chu et al. (2006) and Liu et al. (2020), respectively. The record  
 498 from CD29-2 was interpreted by Horner et al. (2015b) as evidencing the importance of deep (non-eolian) Fe sources  
 499 to central Pacific Fe budgets throughout much of the Cenozoic and Late Cretaceous. In contrast, the  $\delta^{56}\text{Fe}$  recorded  
 500 by three central Pacific Fe–Mn crusts have been similar since the late Miocene, consistent with a regional Fe source  
 501 derived largely from non-reductive sediment dissolution and/or eolian deposition.

### 502 3.4. Prospects

503 While the marine Fe cycle is complex, Fe isotopes are proving to be a valuable tool for studying the modern  
504 Fe cycle. Detailed study of this cycle reveals that it is driven by a multitude of biological, physical and  
505 chemical processes, amongst which productivity is but one small part. In our view, this means that there  
506 are considerable obstacles to using  $\delta^{56}\text{Fe}$  as a paleoproductivity tracer. However, this does not preclude  
507  $\delta^{56}\text{Fe}$  from emerging as a powerful tracer for studying the dynamics of the Fe cycle in the (paleo)oceans.  
508 Such a tracer would be especially powerful given the proximal connection between Fe supply and the  
509 biological productivity of the ocean.


510 Exploiting  $\delta^{56}\text{Fe}$  will require resolving and refining several ambiguities. First, there is a clear need to better  
511 constrain the Fe-isotopic fractionation factor associated with biological uptake (in variable conditions and  
512 from different species) and to diagnose locations where dissolved  $\delta^{56}\text{Fe}$  is most affected by productivity.  
513 Second, the fractionation factors for remineralization and scavenging are essentially unknown, though field  
514 data suggests that the net result of these effects is relatively small (e.g. Radic et al., 2011; Labutut et al.,  
515 2104). Constraining these fractionation factors will be particularly important for developing novel archives  
516 of the paleo Fe cycle, such as pelagic clays (e.g., Tegler et al., 2018). Third, any sedimentary reconstruction  
517 of past Fe-isotopic chemistry will need to consider the high degree of spatial variability in modern  $\delta^{56}\text{Fe}$ .  
518 This will necessitate spatially-distributed core sampling, similar to the approach used to constrain basin-  
519 scale patterns of dust deposition over glacial–interglacial timescales (e.g., Costa et al., 2016; Winckler et  
520 al., 2016). Addressing these priorities will provide valuable constraints on the extent to which the Fe cycle  
521 has influenced primary productivity over recent geological history, and provide key insights into the  
522 potential sensitivity of Earth’s climate to perturbations in marine trace element cycles.

523

524 **4. Zinc**

525 **4.1. Marine distribution**

526 Consistent with its importance as a micronutrient, dissolved Zn has a nutrient-type distribution in the ocean  
 527 (e.g., Bruland, 1980). Typical surface ocean Zn concentrations are 0.01 to 0.5 nmol kg<sup>-1</sup>, compared to deep  
 528 water concentrations of ~2.5 nmol kg<sup>-1</sup> in the north Atlantic and ~10 nmol kg<sup>-1</sup> in the north Pacific (Schlitzer  
 529 et al., 2018; Fig. 6). The dissolved Zn distribution in the ocean closely follows that of the macronutrient Si  
 530 (Bruland, 1980), at least partially due to the similar behaviour of both elements in the Southern Ocean  
 531 (Vance et al., 2017; de Souza et al., 2018; Weber et al., 2018; Roshan et al., 2018; Middag et al., 2019).  
 532 Decoupling of Zn from Si is observed in regions remote from Southern Ocean influence (Janssen & Cullen,  
 533 2015; Vance et al., 2019).



534  
 535 **Figure 6 | Representative profiles of dissolved Zn concentrations ([Zn]) and Zn-isotopic compositions ( $\delta^{66}\text{Zn}$ )**  
 536 **in the Eastern Tropical North Atlantic (squares, dashed line), Northeast Pacific (circles, solid line) and Southern**  
 537 **(triangles, dotted line) Oceans.** This comparison illustrates that despite possessing distinct dissolved concentration  
 538 profiles, the isotopic behavior of Zn is similar between basins.


539

540 It was hoped that Zn isotope ratios would provide a means of tracing biological carbon cycling in the past  
541 and present ocean (e.g., Maréchal et al., 2000; Pichat et al., 2003; Bermin et al., 2006). Zinc isotope data  
542 are reported relative to JMC-Lyon, which has now expired. New  $\delta^{66}\text{Zn}$  data are typically measured relative  
543 to AA-ETH or IRMM-3702 Zn standards and corrected by the offset between these and JMC-Lyon ( $\approx 0.3\%$ ;  
544 Moeller et al., 2006; Archer et al., 2017). The apparent preferential uptake of light Zn isotopes by  
545 phytoplankton observed in some culture experiments (John et al., 2007; Köbberich and Vance, 2017;  
546 Samanta et al., 2018) predicted an isotopically heavy residual pool in surface seawater, analogous to that  
547 observed for  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ,  $\delta^{30}\text{Si}$  and  $\delta^{114}\text{Cd}$  (Cd reviewed in Sec. 6., C, N and Si in Farmer et al., *this issue*).  
548 However, the test of this hypothesis, through analyses of  $\delta^{66}\text{Zn}$  in seawater, awaited a GEOTRACES-driven  
549 revolution in large volume trace metal clean sampling (Cutter et al., 2017), alongside analytical advances  
550 including new chemical separation procedures and MC-ICP-MS (multiple-collector inductively-coupled  
551 plasma mass spectrometry; Bermin et al., 2006; Conway et al., 2013; Takano et al., 2017).

552

553 A consistent picture of the distribution of  $\delta^{66}\text{Zn}$  in the ocean has since emerged. The deep ocean is  
554 isotopically homogeneous, with a  $\delta^{66}\text{Zn}$  signature of about  $+0.45\%$  (Fig. 6.; Conway & John, 2014b; Zhao  
555 et al., 2014; Conway & John, 2015a; Samanta et al., 2017; Takano et al., 2017; John et al., 2018b; Wang et  
556 al., 2019a; Vance et al., 2019; Sieber et al., 2020; Liao et al., 2020; Lemaitre et al., 2020), and isotopically  
557 heavier than the upper continental crust (UCC;  $\delta^{66}\text{Zn} +0.3\%$ ; Moynier et al., 2017). Deviations to deep  
558 ocean  $\delta^{66}\text{Zn}$  compositions as light as  $-0.2\%$  have been observed near sediments or hydrothermal Zn sources  
559 (Conway and John, 2014b; Lemaitre et al., 2020). In the surface, while heavy  $\delta^{66}\text{Zn}$  isotope compositions  
560 have been observed, the upper water column for Zn is often isotopically light or, in the Southern Ocean and  
561 North Pacific, only very slightly fractionated towards heavier values (Conway & John, 2014b; 2015a; Zhao  
562 et al., 2014; Samanta et al., 2017; Takano et al., 2017; Wang et al., 2019a; Sieber et al., 2020; Vance et al.,  
563 2019; Liao et al., 2020; Lemaitre et al., 2020). Possible reasons for the discrepancy between expected and  
564 observed upper water column  $\delta^{66}\text{Zn}$  are discussed in detail below. They include preferential scavenging of  
565 isotopically heavy Zn on particle surfaces (e.g., John & Conway, 2014; Weber et al., 2018), an isotopically  
566 light source of Zn, possibly from shallow remineralization of organic material (Samanta et al., 2017; Vance  
567 et al., 2019) or from anthropogenic aerosols (Liao et al., 2020; Lemaitre et al., 2020). We note here that  
568 these complexities will likely make it challenging to use Zn isotopes as a paleoproductivity proxy.

569



570  
 571 **Figure 7 | Processes driving zinc isotope variations in modern seawater.** Though biological processes are capable  
 572 of influencing dissolved  $\delta^{66}\text{Zn}$ , they do not appear to be the dominant driver of Zn isotope variations in the marine realm.

573

## 574 4.2. Driving processes

### 575 4.2.1 Biological

576 Zinc is a metal center in two key enzymes: carbonic anhydrase, necessary for carbon fixation, and alkaline  
 577 phosphatase, necessary for dissolved organic phosphorus uptake by marine organisms (Morel et al., 1994;  
 578 Shaked et al., 2006; reviewed by Sinoir et al., 2012). Zinc also has an array of other physiological roles in  
 579 marine organisms, as exemplified by the observation that Zn contents of phytoplankton cells are of similar  
 580 magnitude to the micronutrient Fe (Twining & Baines, 2013).

581 Zinc can be growth-limiting for phytoplankton grown in culture (Anderson et al., 1978; Brand et al., 1983;  
 582 Morel et al., 1994), but Zn (co)-limitation (with Fe, Co) has only rarely been observed in the open ocean  
 583 (e.g., Coale, 1991; Coale et al., 2003; Franck et al., 2003; Ellwood, 2004; Lohan et al., 2005). This  
 584 difference between culture and field may reflect the ability of some phytoplankton to substitute Cd or Co

585 for Zn in some enzyme systems when ambient Zn concentrations are low (e.g., Morel et al., 1994; Lee &  
586 Morel, 1995; Yee & Morel, 1996; Kellogg et al., 2020). Nevertheless, Zn availability has been shown to  
587 influence species composition and phytoplankton growth, including rates of calcification and alkaline  
588 phosphatase activity (Sunda & Huntsman, 1995; Crawford et al., 2003; Schulz et al., 2004; Shaked et al.,  
589 2006; Mahaffey et al., 2014). In culture, phytoplankton biomass is typically enriched in the light isotopes  
590 of Zn (John et al., 2007; Köbberich and Vance, 2017; 2019; Samanta et al., 2018). Rather than reflecting a  
591 kinetic isotope effect on uptake, it has been suggested that Zn speciation in the media controls cellular  $\delta^{66}\text{Zn}$   
592 values; specifically, strong organic ligands (e.g., EDTA) present in the media preferentially complex heavy  
593 Zn isotopes, rendering the bioavailable Zn pool isotopically light (John et al., 2007; Köbberich and Vance,  
594 2017; 2019; Fig. 7).

595

#### 596 4.2.2. Chemical

597 Like most bioessential metals, Zn bioavailability is dictated by its chemical speciation (e.g., Anderson et  
598 al., 1978), which is dominated in the ocean by complexation to strong ( $K' \sim 10^9 - 10^{11}$ ) organic ligands  
599 (Bruland, 1989; Donat & Bruland, 1990; Ellwood & Van Den Berg, 2000; Jakuba et al., 2012; Kim et al.,  
600 2015). Inorganic Zn is considered to make up <5% of the total Zn pool in most ocean regions, with the  
601 exception of the Southern Ocean, where strong upwelling of nutrient-rich deep waters leads to Zn  
602 concentrations in excess of complexing ligands (Baars & Croot, 2011). Note that while strongly-complexed  
603 Zn is unlikely to be bioavailable, the presence of ‘weak’ ligands (or more labile ligands) can enhance Zn  
604 uptake (Aristilde & Xu, 2012). As noted above, the role of the diversity of organic (and inorganic) Zn-  
605 binding ligands in determining dissolved and particulate  $\delta^{66}\text{Zn}$  values remains to be fully evaluated.

606 A role for scavenging in the marine cycling of Zn and Zn isotopes has been widely cited (e.g., John &  
607 Conway, 2014; Weber et al., 2018; Roshan et al., 2018; Liao et al., 2020; Fig. 7). It is argued that scavenging  
608 can a) explain the widespread observation of isotopically light Zn in the upper ocean, via preferential  
609 removal of heavy Zn isotopes on particles, and b) explain elevated concentration of Zn in the deep Pacific  
610 compared to that supplied in southern-sourced deep waters (Weber et al., 2018; Roshan et al., 2018).  
611 Elevated Zn in the deep Pacific has also been attributed to additional Zn input via, e.g., hydrothermalism  
612 (Roshan et al., 2018), with a lighter than deep ocean  $\delta^{66}\text{Zn}$  signature (John et al., 2018b). Local and basin  
613 scale deviations towards lighter deep ocean  $\delta^{66}\text{Zn}$  compositions in both the Atlantic and Pacific have been  
614 attributed to sedimentary input (Conway & John, 2014b; John et al., 2017; Liao et al., 2020; Lemaitre et  
615 al., 2020), and could also reflect hydrothermalism (Conway & John, 2014b; John et al., 2018b; Lemaitre et  
616 al., 2020). Anthropogenic aerosol deposition is thought to supply significant Zn to regions of the surface



617 ocean (e.g., Liao & Ho, 2018), with possible direct and indirect (via scavenging) regional impacts on upper  
618 ocean  $\delta^{66}\text{Zn}$  values (Liao et al., 2020; Lemaitre et al., 2020).

619 Lastly, Janssen & Cullen (2015) suggest that decoupling of Zn and Si in the northeast Pacific reflects the  
620 formation of Zn sulfides in the oxygen deficient zone (ODZ), directly equivalent to proposed Cd sulfide  
621 precipitation; this hypothesis is discussed in more detail in Sec. 6.2.3. To date, however, there remains scant  
622 evidence for water column Zn-sulfide precipitation in ODZs (e.g., Conway & John, 2014b; John et al.,  
623 2018b; Vance et al., 2019). That said, Zn-sulfide precipitation is undoubtedly important in euxinic basins  
624 such as the Black Sea and Cariaco Basin (Vance et al., 2016; Isson et al., 2018), and has been postulated to  
625 occur within the porewaters of oxygen deficient, organic-rich sediments (Sec. 4.3.3.; Little et al., 2016).

626

#### 627 4.2.3. Physical

628 In common with many of the TEIs discussed here, the physical ocean circulation exerts a first order control  
629 on the distribution of dissolved Zn and  $\delta^{66}\text{Zn}$  compositions (Vance et al., 2017; de Souza et al., 2018; Weber  
630 et al., 2018; Sieber et al., 2020). Subantarctic water masses have distinct low Zn:PO<sub>4</sub> and Si:PO<sub>4</sub> ratios, due  
631 to the elevated uptake of Zn and Si by diatoms in the surface Southern Ocean (Sarmiento et al., 2004; Vance  
632 et al., 2017). Remineralization of these Zn- and Si-rich diatoms at depth imprints a correspondingly high  
633 Zn:PO<sub>4</sub> and high Si:PO<sub>4</sub> fingerprint on Antarctic bottom waters. This coupling of Zn and Si in the Southern  
634 Ocean forms the basis of the global Zn:Si correlation via the advection of southern sourced water masses  
635 towards the low latitudes, where they fill much of the ocean interior (de Souza et al., 2012; Holzer et al.,  
636 2014). The homogeneity of deep ocean Zn isotope compositions reflects the limited degree of Zn isotope  
637 fractionation on uptake by Southern Ocean diatoms (Zhao et al., 2014; Wang et al., 2019a), which results  
638 in intermediate and deep southern-sourced water masses with limited or no isotopic contrast (Sieber et al.,  
639 2020). The oceanic Zn:Si correlation persists despite shallower remineralization of Zn relative to Si  
640 (Twining et al., 2014), and is especially clear in the Atlantic, underlining that the mixing of water masses  
641 is the dominant control on dissolved Zn in this basin (Vance et al., 2017; Weber et al., 2018; Middag et al.,  
642 2019). Decoupling of Zn and Si is observed in the upper water column of the northeast Pacific, far from the  
643 Southern Ocean influence (Janssen & Cullen, 2015; Vance et al., 2019), likely resulting from the shallower  
644 remineralisation of Zn (from organic ‘soft parts’) relative to Si (from opal; Vance et al., 2019).




### 645 **4.3. Sedimentary archives**

#### 646 *4.3.1. Ferromanganese sediments*

647 Manganese oxides are strong sorbents of positively charged, divalent trace metals, due to their negative  
648 layer charge at the pH of natural waters (e.g., Koschinsky & Halbach, 1995). The phylломanganate  
649 birnessite is the main Mn- and TE-bearing phase in oxic marine sediments (e.g., Koschinsky and Hein,  
650 2003), as well as for several elements described here, including Zn, Cu, Ni, Cd, and Mo (though Mo exists  
651 in seawater as the molybdate anion). Ferromanganese crusts and nodules incorporate TEs during growth,  
652 leading to  $10^6$ -fold enrichments compared to seawater concentrations (e.g., Arrhenius, 1963; Aplin and  
653 Cronan, 1985). As a result, Fe–Mn sediments are often one of the first marine sediment types to be targeted  
654 in the development of a new isotopic tracer.

655 In the case of Zn, Fe–Mn crusts and nodules are isotopically heavy compared to seawater ( $\approx +0.45\text{‰}$ ),  
656 exhibiting compositions between  $+0.9$  to  $+1 \text{‰}$  (Maréchal et al., 2000; Little et al., 2014a; Fig. 8). The  
657 heavy signature is broadly consistent with inorganic sorption experiments of Zn on birnessite, with  
658 experiments at high ionic strength yielding  $\Delta^{66}\text{Zn}_{\text{sorbed-aqueous}}$  (where  $\Delta^{66}\text{Zn}_{\text{sorbed-aqueous}} = \delta^{66}\text{Zn}_{\text{sorbed}} -$   
659  $\delta^{66}\text{Zn}_{\text{aqueous}}$ ) of  $+0.16$  to  $+2.7\text{‰}$  (Bryan et al., 2015). The magnitude of fractionation observed in experiments  
660 is typically considerably larger than in natural Fe–Mn crusts and nodules, which Bryan et al. (2015) suggest  
661 may reflect sorption of Zn on birnessite via two different molecular mechanisms, each associated with  
662 different isotopic fractionation factors, as is the case for thallium (e.g., Nielsen et al., 2013). In addition,  
663 the influence of organic and inorganic speciation of Zn in seawater may play a role, though this remains to  
664 be fully evaluated (e.g., Little et al., 2014b).

665 Little et al. (2014a) observed no marked temporal changes in  $\delta^{66}\text{Zn}$  values for three Fe–Mn crusts from  
666 each of the major ocean basins over the last  $\sim 20$  Ma. Assuming no significant diffusion or resetting of crust  
667 Zn isotope compositions on this timescale, this observation suggests that, on a global basis, the marine Zn  
668 cycle has been in isotopic steady state for at least 20 Myr (Fig. 8)



669

670 **Figure 8 | Deep ocean Zn isotope constancy over the past 20 Myr.** These records are derived from Fe–Mn crusts  
 671 recovered from the Atlantic (square, dashed line), Indian (triangle, solid line), and Pacific (circle, dotted line) Oceans.  
 672 Assuming the isotopic offset between dissolved Zn in seawater and Fe–Mn crusts has remained unchanged at  $\approx 0.55$   
 673 ‰ over this time, these records imply only minimal Zn-isotopic variations in the deep ocean over the past  $\sim 20$  Myr. Two  
 674 anomalous measurements from 109D-C (in parentheses) possess low levels of authigenic Zn enrichment, indicative of  
 675 detrital contamination (Little et al., 2014a).

676

### 677 4.3.2. Biogenic silica

678 The correlation of Zn with Si in the modern ocean led to the suggestion that Zn/Si measured in diatom opal  
679 may be a proxy for Zn/Si (and thus [Zn]) of past seawater. However, culturing and  $\mu$ -XRF analyses revealed  
680 that only a small fraction (1–3%) of the diatom Zn quota is incorporated into the opal frustules, with the  
681 remainder present in the organic ‘soft parts’ of the diatom cells (Ellwood & Hunter, 2000; Twining et al.,  
682 2004). The Zn concentration in opal ( $Zn_{\text{opal}}$ ) also more closely reflects the bioavailable Zn in seawater than  
683 ambient Zn:Si (Ellwood & Hunter, 2000). Nevertheless, if the mechanisms of Zn incorporation into  
684 biogenic opal can be understood, and calibrated, a Zn:Si proxy of Zn bioavailability could help shed light  
685 on micronutrient limitation of the biological pump.

686 Andersen et al. (2011) analyzed Zn isotopes ( $\delta^{66}Zn_{\text{opal}}$ ) and  $Zn/Si_{\text{opal}}$  in diatom opal isolated from core top  
687 sediments from the Southern Ocean. They observed isotopically heavy Zn in opal (at +0.7 to +1.5‰), and  
688 an inverse relationship of  $\delta^{66}Zn_{\text{opal}}$  with  $Zn/Si_{\text{opal}}$ . Consistent with culturing studies, core top  $Zn/Si_{\text{opal}}$   
689 appears to be linked to bioavailable Zn concentrations in ambient surface seawater. The authors suggested  
690 that  $\delta^{66}Zn_{\text{opal}}$  should also reflect the isotope composition of bioavailable Zn in surface seawater, which, they  
691 predicted, should be isotopically heavy due to the predicted preferential incorporation of light isotopes into  
692 phytoplankton organic material. In this view, the extent of uptake—nutrient utilization—would be recorded  
693 by the systematics of  $Zn/Si_{\text{opal}}$  and  $\delta^{66}Zn_{\text{opal}}$ . However, this study predated the recent surge in seawater  $\delta^{66}Zn$   
694 measurements. As discussed above, surface water  $\delta^{66}Zn$  analyses in the Southern Ocean have not borne out  
695 the prediction of isotopically-heavy residual surface waters, with little to no fractionation observed (Zhao  
696 et al., 2014; Wang et al., 2019a; Sieber et al., 2020).

697 More recently, Hendry and Andersen (2013) have shown that some sponge spicules faithfully record  
698 seawater  $\delta^{66}Zn$ . Sponges are primarily deep-sea organisms. Hence, if we can deconvolve the controls on  
699 diatom  $Zn/Si$  and  $\delta^{66}Zn_{\text{opal}}$ , a combination of  $Zn/Si$  and  $\delta^{66}Zn$  measurements in diatoms and sponges (as  
700 used for Si isotopes, Farmer et al., *this issue*) could provide a strong basis for unravelling the past ocean  
701 global Zn cycle, including the role of Southern Ocean processes in the biological carbon pump.

702

### 703 4.3.3. Carbonates

704 Carbonates may provide an alternative archive for Zn (as Zn/Ca) and  $\delta^{66}Zn$ . For example, Marchitto et al.  
705 (2000) showed that Zn/Ca ratios in benthic foraminifera reflect bottom water dissolved Zn concentrations.  
706 However, Zn concentrations in individual microfossil shells are extremely low (of the order  $\sim 0.1$  ng); at

707 this level, with current analytical capabilities and blank contributions, more than one hundred benthic  
 708 foraminifera would be required for a single Zn isotope measurement.

709 To circumvent the issue of low Zn contents of individual shells, Pichat et al. (2003) utilized a selective  
 710 carbonate dissolution procedure on bulk sediment from the equatorial Pacific, mostly consisting of  
 711 coccoliths. These authors argued that isotopically heavy  $\delta^{66}\text{Zn}_{\text{carb}}$  reflected ambient surface seawater, with  
 712 values modulated by changes in biological productivity due to varying seasonal insolation. Similarly  
 713 isotopically heavy Zn in ancient carbonates has also been argued to reflect strong biological utilization in  
 714 surface waters (Kunzmann et al., 2013; cf. John et al., 2017; Liu et al., 2017). However, as discussed, the  
 715 interpretation of these data should be revisited in the light of recent seawater  $\delta^{66}\text{Zn}$  data, where isotopically  
 716 heavy surface ocean Zn isotope compositions are rarely observed.

717 Selective carbonate dissolution procedures present two problems. First, Zn is present at high concentrations  
 718 in potential contaminating material, e.g., lithogenic or authigenic (e.g. Fe-Mn oxide) phases, and second,  
 719 non-quantitative leaching introduces the potential for stable isotope fractionation (Revels et al., 2015).  
 720 Nevertheless, several lab groups are now investing considerable effort to systematically assess these  
 721 procedures, with promising results (e.g., Clarkson et al., 2018).

722 Finally, deep-sea coral skeletons also provide promise as a potential intermediate-deep water archive of  
 723  $\delta^{66}\text{Zn}$  (e.g., Little et al., 2017a). Their size and global distribution, combined with the ability to assign  
 724 precise ages to individual specimens, gives corals some distinct advantages over traditional sedimentary  
 725 paleoclimate archives (Robinson et al., 2014).

726

#### 727 4.3.4. Organic-rich sediments

728 Quantitative removal of Zn into Zn sulfides in the euxinic Black Sea water column leads to sedimentary  
 729  $\delta^{66}\text{Zn}$  values that preserve the global seawater signature (Vance et al., 2016). This behaviour is analogous  
 730 to the Mo isotope system (Sec. 7; e.g., Neubert et al., 2008; Nägler et al., 2011), and suggests that euxinic  
 731 black shales represent an archive for seawater  $\delta^{66}\text{Zn}$  values. By extension, Isson et al. (2018) propose that  
 732 a chemically isolated sulfide-bound Zn fraction also preserves seawater  $\delta^{66}\text{Zn}$  values, demonstrating the  
 733 effectiveness of their approach for core top sediments from the euxinic Cariaco Basin. However, it is not  
 734 yet clear if the sulfide fraction of organic shales from non-euxinic sites will necessarily record seawater  
 735  $\delta^{66}\text{Zn}$ , as (non-quantitative) precipitation of Zn sulfides is associated with light isotope fractionation (Fujii  
 736 et al., 2011a; Vance et al., 2016).

737 Authigenic Zn in organic rich sediments deposited in productive continental margin settings is isotopically  
738 light (on average, about +0.1‰) compared to seawater (at about +0.45 ‰; Little et al., 2016). One possible  
739 explanation for this signature is the preferential biological uptake of light isotopes into organic matter (Little  
740 et al., 2016; John et al., 2017). The second possible explanation, touched upon above, is light Zn isotope  
741 fractionation during Zn sulfide precipitation, either in sediments or, possibly, in reducing  
742 microenvironments in the water column (Janssen & Cullen, 2015; Little et al., 2016; Vance et al., 2016;  
743 Bianchi et al., 2018). While a biologically-driven signature would suggest utility for  $\delta^{66}\text{Zn}$  in organic-rich  
744 sediments as a paleoproductivity proxy, a role for sulfidation highlights the importance of redox conditions  
745 during diagenesis, particularly sulphate reduction, in driving the  $\delta^{66}\text{Zn}$  values preserved in organic-rich  
746 sediments. We note that sedimentary redox is linked to overlying productivity (e.g., Hartnett et al., 1998),  
747 but in a complex manner that would make a purely productivity based interpretation of preserved  $\delta^{66}\text{Zn}$   
748 values in black shales challenging.

749

#### 750 **4.4. Prospects**

751 To summarize, recent seawater, experimental, and sedimentary Zn isotope research implies that the  
752 relationship between Zn isotopes and (paleo)productivity is not straightforward. This adds uncertainty to  
753 the application of using Zn isotopes as a tracer of paleoproductivity. Fundamental to the application of Zn  
754 isotopes as a paleoproductivity proxy (or other type of paleoenvironmental tracer) is a better understanding  
755 of the mechanisms driving the modern upper water column distribution of  $\delta^{66}\text{Zn}$ . Future work should target  
756 coupled dissolved and particulate phase  $\delta^{66}\text{Zn}$  data, in concert with detailed biological and chemical  
757 speciation data. To date, particulate phase  $\delta^{66}\text{Zn}$  data has proved particularly challenging due to Zn being  
758 extremely contamination prone.

759 We recommend revisiting existing archive data in the context of modern seawater  $\delta^{66}\text{Zn}$  data. In general,  
760 we encourage that the future interpretation of Zn/Si, Zn/Ca, Zn-TOC and  $\delta^{66}\text{Zn}$  data be made within a  
761 mechanistic framework, via careful sequential extraction procedures, application of appropriate micro-  
762 analytical techniques (e.g., LA-ICP-MS, nano-SIMS,  $\mu$ -XAFS), and further experimental and theoretical  
763 work.

764 We note that there are systematic variations in sedimentary Zn isotope compositions on Gyr timescales. For  
765 example, Isson et al. (2018) reported  $\delta^{66}\text{Zn}$  of black-shale associated Zn spanning  $\sim 3.5$  Gyr, as well as  
766 bitumen and kerogen spanning the last 1.5 Gyr. The black shale record shows a shift from UCC-like values  
767 (centered on  $\approx +0.4$  ‰) to more marine-like  $\delta^{66}\text{Zn}$  (centered on  $\approx +0.6$  ‰) around 800 Ma, which the authors

768 interpret as evidencing a transition to a more eukaryote-dominated marine ecosystem. While this feature is  
769 consistent with the higher affinity of eukaryotes for Zn compared to prokaryotes (Dupont et al., 2006; 2010),  
770 there remains considerable uncertainty regarding the influence of Earth's overall redox state on marine Zn  
771 isotope systematics.


772 In carbonates, Yan et al. (2019) compile three  $\delta^{66}\text{Zn}$  datasets from contemporaneous Ediacaran (635 Ma,  
773 Marinoan) post-glacial cap carbonates (Kunzmann et al., 2013; John et al., 2017; Lv et al., 2018), which  
774 show systematic changes in Zn isotope compositions over this period of marked global change. Similarly,  
775 Sweere et al. (2018) present data showing marked shifts in carbonate-bound Zn isotope compositions in  
776 several geological sections spanning a Cretaceous ocean anoxic event (OAE 2). As yet, there is no  
777 consensus on the drivers of these intriguing isotopic shifts, but these and similar records serve to motivate  
778 the future of Zn isotope research.

779

780 **5. Copper**

781 **5.1. Marine distribution**

782 The distribution of Cu in the ocean has been described as ‘hybrid-type’, because it is intermediate between  
 783 nutrient- and scavenged-type elements (Bruland & Lohan, 2003). Depth profiles of dissolved Cu typically  
 784 show approximately linear increases in concentration with depth (e.g., Boyle et al., 1977). Surface dissolved  
 785 Cu concentrations are typically about 0.5 to 1 nmol kg<sup>-1</sup>, compared to deep Atlantic concentrations of ~2.5  
 786 nmol kg<sup>-1</sup> and deep Pacific concentrations of ~4 nmol kg<sup>-1</sup> (Fig. 9). The distribution of dissolved Cu has  
 787 been attributed to a combination of biological uptake and remineralization, benthic flux from sediments  
 788 (e.g., Boyle et al., 1977; Roshan & Wu, 2015a; Little et al., 2018), and/or reversible scavenging (Little et  
 789 al., 2013; Richon et al., 2019). These processes are superimposed on the first-order distribution established  
 790 via the physical ocean circulation (Roshan & Wu, 2015a).




791 **Figure 9 | Representative profiles of dissolved Cu concentrations ([Cu]) and Cu-isotopic compositions (δ<sup>65</sup>Cu)**  
 792 **in the Southwest Atlantic (squares, dashed line), Subtropical South Pacific (circles, solid line) and Southern**  
 793 **(triangles, dotted line) Oceans. This comparison illustrates that the isotopic behavior of Cu is similar between basins,**  
 794 **reflecting the importance of complexation by strong organic ligands.**  
 795

796



797 Copper isotopes are reported as relative to the NIST SRM 976 standard, though due to a shortage of this  
 798 material, two other certified reference standards are now available: ERM-AE633 ( $\delta^{65}\text{Cu}_{\text{SRM976}} = -0.01 \pm 0.05$   
 799 ‰) and ERM-AE647 ( $\delta^{65}\text{Cu}_{\text{SRM976}} = +0.21 \pm 0.05$  ‰; Moeller et al., 2012; Moynier et al., 2017). For  
 800 consistency and ease of comparison, Moynier et al. (2017) recommend that future data be reported relative  
 801 to NIST SRM 976. We adopt this convention. The analysis of Cu isotopes in seawater is challenging, due  
 802 to both Cu's strong organic complexation, and only two isotopes precluding the use of a double spike  
 803 technique (reviewed in Little et al., 2018). The data presented to date indicate that deep seawater  $\delta^{65}\text{Cu}$   
 804 values are isotopically heavy (at about +0.7 ‰) compared to the upper continental crust (UCC; at about  
 805 +0.1 ‰; Table 1; Moynier et al., 2017). Lighter Cu isotope compositions in the upper water column and  
 806 along margins are thought to reflect local sources of isotopically light Cu (e.g., aerosols, riverine  
 807 particulates, sediments; Takano et al., 2014; Little et al., 2018).

808



809

810 **Figure 10 | Processes driving copper isotope variations in modern seawater.** Biological productivity exerts only a  
 811 modest impact on the marine Cu isotope cycle and thus there is no obvious route to developing Cu isotopes (or Cu  
 812 concentrations) as a paleoproductivity proxy.

## 813 **5.2. Driving processes**

### 814 *5.2.1. Biological*

815 Copper is bioessential, but cellular Cu contents are approximately 2–10 fold lower than the micronutrients  
816 Fe and Zn (Twining & Baines, 2013). The redox-active behaviour of Cu (existing as  $\text{Cu}^{2+}$  or  $\text{Cu}^+$  in  
817 biological systems) enables its role in electron transport, for example in the Cu-containing proteins  
818 plastocyanin and cytochrome c oxidase (Ridge et al., 2008). Copper uptake of some Fe-limited  
819 phytoplankton may increase, either due to the replacement of Fe-containing with Cu-containing enzymes  
820 (e.g., Peers and Price, 2006), or the involvement of Cu in the high-affinity Fe uptake systems (Annett et al.,  
821 2008; Maldonado et al., 2006; Guo et al., 2012). Despite its biological function, Cu is also extremely toxic  
822 due to the formation of reactive oxygen species, which pose a threat to DNA, lipids, and proteins (Ridge et  
823 al., 2008). Copper toxicity thresholds vary by phytoplankton group, with smaller organisms generally more  
824 sensitive than larger ones (e.g., cyanobacteria cf. diatoms; Brand et al., 1986), and coastal strains more  
825 resistant than open ocean strains (e.g., Peers et al., 2005).

826 A small number of studies have investigated Cu isotope fractionation during cellular uptake or cell surface  
827 adsorption by microorganisms (Pokrovsky et al., 2008; Navarrete et al., 2011; Cadiou et al., 2017; Coutaud  
828 et al., 2018; 2019). The results of these experiments are somewhat variable, with enrichment of either light  
829 or heavy Cu isotopes observed during assimilation and adsorption. However, assimilation in culture  
830 generally favours light Cu isotopes (Navarrete et al., 2011; Cadiou et al., 2017; Fig. 10). The complexity in  
831 Cu isotopic behaviour has been attributed to small changes in Cu speciation or redox during uptake and/or  
832 release of Cu (Coutaud et al., 2018; 2019).

833

### 834 *5.2.2. Chemical*

835 In seawater, the vast majority of Cu is complexed to strong organic ligands (more than 99.8% complexed  
836 in surface northeast Pacific), which lower free  $\text{Cu}^{2+}$  concentrations to below toxic levels (e.g., Coale &  
837 Bruland, 1988; 1990; Moffett & Dupont, 2007). It is thought that ligands are primarily produced by biota  
838 for the purpose of detoxification (e.g., Moffett et al., 1990; Moffett & Brand, 1996), albeit recent work  
839 suggesting that strongly complexed Cu is bioavailable to some eukaryotes, which appear to have a higher  
840 cellular Cu requirement (and higher thresholds of Cu toxicity) than prokaryotes (Semeniuk et al., 2009;  
841 2015).

842 Both theory and experiments predict preferential complexation of heavy isotopes by strong organic ligands  
843 (Sherman, 2013; Fujii et al., 2013; Ryan et al., 2014; Sherman et al., 2015), and organic complexation is  
844 thought to play a key role in the modern oceanic budget and distribution of Cu isotopes (Vance et al., 2008;  
845 Little et al., 2014a; Thompson & Ellwood, 2014; Takano et al., 2014; Little et al., 2018). The small pool of  
846 non-complexed  $\text{Cu}^{2+}$  in seawater is thus expected to be isotopically light (e.g., Little et al., 2014b, 2018).

847 Based on a surface complexation model with the phyllo-manganate birnessite, the principal scavenging  
848 phase of divalent trace metals in oxic sediments, Sherman & Peacock (2010) calculated that the  
849 concentration of dissolved Cu in deep waters should be orders of magnitude lower than is actually observed.  
850 They attribute this difference to the chelation of “*essentially all dissolved Cu*” by organic ligands (Sherman  
851 & Peacock, 2010), consistent with observations (e.g., Moffett & Dupont, 2007; Heller & Croot, 2015;  
852 Jacquot & Moffett, 2015). Nevertheless, some form or forms of scavenging are also thought to play a role  
853 in the oceanic Cu distribution.

854 Reversible scavenging, a term used to describe the equilibrium between a scavenged and dissolved metal  
855 pool, has been proposed as the driving process behind the generally monotonic, linear increases in dissolved  
856 [Cu] depth (Little et al., 2013; Richon et al., 2019; Fig. 9). In some regions with high particulate loads (e.g.,  
857 some hydrothermal plumes, benthic nepheloid layers), scavenging removal of Cu has been observed  
858 (Jacquot and Moffett, 2015; Roshan & Wu, 2015a). Preferential scavenging of light Cu isotopes by  
859 particulate (e.g., oxyhydroxide) phases has also been proposed as an explanation for isotopically heavy  
860 seawater Cu isotope compositions (e.g., Takano et al., 2014), though the driving mechanisms leading to  
861 isotopically light particulate Cu remain to be fully established (see section 5.3; Fig. 10).

862 The shorter residence time of Cu (2-3.3 kyr; Little et al., 2017b), compared to metals with longer residence  
863 times (e.g., Cd, Zn, Ni), mean that, as for Fe, regional and local sources of Cu to the ocean play a relatively  
864 larger role in determining dissolved Cu distributions than for some other TEIs. Sources of Cu include  
865 aerosols (both natural and anthropogenic; e.g., Takano et al., 2014; Yang et al., 2019), benthic fluxes from  
866 sediments (e.g., Boyle et al., 1977; Heller & Croot, 2015; Roshan & Wu, 2015a; Little et al., 2018), and  
867 dissolved or particulate riverine sources (e.g., Vance et al., 2008; Little et al., 2018; Richon et al., 2019). A  
868 possible small hydrothermal source has been identified in the South Pacific from the East Pacific Rise  
869 (Roshan & Wu, 2018), which is in contrast to the scavenging removal of Cu observed around hydrothermal  
870 vents elsewhere (e.g., Jacquot and Moffett, 2015).

871

872 *5.2.3. Physical*

873 Copper's shorter residence time relative to Cd, Zn and Ni also means that the Southern Ocean and wider  
874 physical ocean circulation play a somewhat smaller role in oceanic Cu and  $\delta^{65}\text{Cu}$  distributions compared to  
875 many of the other TEIs discussed herein. Nonetheless, the imprint of circulation is evident in certain  
876 circumstances, such as in the Atlantic and the upper 2 km of the South Pacific where dissolved Cu is  
877 correlated with dissolved silicate (Roshan & Wu, 2015a; 2018; Little et al., 2018).


878

879 **5.3. Marine archives**880 *5.3.1. Ferromanganese sediments*

881 Ferromanganese sediments (crusts and nodules) exhibit Cu isotope compositions of +0.3 to +0.5 ‰  
882 (Albarède, 2004; Little et al., 2014). This means that, on average, Fe–Mn sediments are approximately 0.35  
883 ‰ lighter than deep seawater, which averages +0.7 ‰ (Fig. 11). The explanation for this offset is uncertain,  
884 but may reflect either strong organic complexation of Cu in seawater (Little et al., 2014b), or the enrichment  
885 of light Cu isotopes on the birnessite mineral surface, as recently observed in inorganic experiments  
886 whereby sorbed Cu exhibited  $\delta^{65}\text{Cu}$  that was  $0.45 \pm 0.18$  ‰ lighter than Cu in solution (Ijichi et al., 2018).  
887 At equilibrium in seawater, however, the strong complexation and mineral sorption effects should be  
888 additive, leading to a much larger isotopic offset than the  $\sim 0.35$  ‰ observed, suggesting that one of the two  
889 effects is not expressed in nature for reasons as yet unclear.

890 Little et al. (2014a) observed no marked changes in deep-ocean  $\delta^{65}\text{Cu}$  recovered from three Fe–Mn crusts  
891 from each of the major ocean basins over the last  $\sim 20$  Ma. Assuming Cu diffusion is slower than crust  
892 growth rate, the lack of Cu isotope variation indicates that, on a global basis, the marine Cu cycle has been  
893 in isotopic steady state for at least 20 Myr (Fig. 10).

894



895

896 **Figure 11 | Deep ocean Cu isotope constancy over the past 20 Myr.** These records are derived from Fe–Mn crusts  
 897 recovered from the Atlantic (square, dashed line), Indian (triangle, solid line), and Pacific (circle, dotted line) Oceans.  
 898 Assuming the isotopic offset between dissolved Cu in seawater and Fe–Mn crusts has remained unchanged at ≈0.35  
 899 ‰ over this time, these records imply that the inter-basin Cu-isotopic variations observed in modern crusts and the Cu  
 900 isotope cycle itself have remained relatively stable for at least 20 Myr. Two samples from 109D-C possess low levels  
 901 of authigenic Cu enrichment indicating detrital contamination (parentheses, arrow; Little et al., 2014a).  
 902

### 903 5.3.2. *Organic-rich sediments*

904 Qualitative arguments for high organic matter fluxes (i.e., higher paleoproductivity) have been made based  
905 on elevated Cu and Ni concentrations in ancient organic-rich sediments (e.g., Tribovillard et al., 2006). This  
906 approach is supported by positive correlations between Cu and Ni with TOC in modern continental margin  
907 sediments (Sec. 9.3). Applying this [metal]–TOC approach to study past productivity has three  
908 prerequisites:

- 909 1. Organic matter is the primary vector supplying the TE to sediments;
- 910 2. Sediments are reducing, and must remain reducing during diagenesis and burial (note that this  
911 differs for Ba, since BaSO<sub>4</sub> preservation requires more oxidizing conditions; Sec. 8);
- 912 3. The system is open marine

913 A significant challenge to this approach for Cu (as well as Zn, Cd, Mo, and other chalcophile elements) is  
914 its reactivity towards sulfide, as illustrated by the quantitative removal of Cu from the euxinic Black Sea  
915 water column and resultant enrichment in underlying sediments (Tankéré et al., 2001; Little et al., 2015).  
916 Thus, for Cu, prerequisite number (1), that the metal is primarily supplied to sediment associated with  
917 organic matter, is not fulfilled. We return to this approach for Ni in Sec. 9.3.2.

918 Authigenic Cu in modern organic-rich sediments is generally similar in isotopic composition to Fe–Mn  
919 sediments and suspended particulate material collected from the South Atlantic, all at about +0.3‰ (Little  
920 et al., 2017b; 2018; Ciscato et al., 2019). This homogeneity in authigenic Cu isotope compositions has been  
921 suggested to reflect an equilibrium isotope fractionation in the aqueous phase between organically  
922 complexed Cu and inorganic Cu<sup>2+</sup>, with the latter approximately 0.4‰ lighter than ligand-bound Cu,  
923 followed by near quantitative scavenging of inorganic Cu<sup>2+</sup> by (any type of) particulate material (Little et  
924 al., 2017b; 2018). If correct, this hypothesis suggests that authigenic Cu isotope compositions in marine  
925 sediments may reflect the evolution of organic complexation on geological timescales.

926 Ciscato et al. (2018; 2019) developed a new approach to isolate TEIs associated with two different fractions  
927 in organic-rich sediments, the ‘organic-pyrite fraction’ (OPF) and ‘HF-digestible fraction’ (HFD). They  
928 find that the OPF of modern Peru margin sediments typically contains >50 % of total Cu and is variably  
929 isotopically light compared to bulk authigenic Cu. They suggest this signature reflects incomplete  
930 sulfidation under variable water column and sedimentary redox conditions (e.g., Bianchi et al., 2018).  
931 Unlike in modern sediments, in ancient shales (ranging in age from 0.4 to 3.4 Ga) bulk Cu concentrations  
932 do not correlate with TOC and >80% of Cu is hosted in the HFD fraction. In addition, the OPF fraction in  
933 ancient shales is markedly isotopically heavier than it is in modern sediments (Ciscato et al., 2019). This



934 difference between modern and ancient Cu partitioning may reflect diagenetic or metamorphic processing,  
935 or it may be a primary feature relating to differences in the Cu isotope composition of seawater, or  
936 differences in the mechanism(s) of Cu sequestration into sediments.

937

#### 938 **5.4. Prospects**

939 Despite the biological importance of Cu, the modern biogeochemical cycle of Cu suggests there is no clear  
940 route to developing Cu isotopes (or Cu concentrations) as a paleoproductivity proxy. However, organic  
941 ligands play a key role in the cycling of Cu and Cu isotopes, suggesting potential for the use of Cu isotopes  
942 in tracing the evolution of organic complexation on geological timescales. Additionally, careful sequential  
943 extraction procedures, such as those described by Ciscato et al. (2019), may allow for the direct probing of  
944 past seawater properties, such as redox state.

945 At present, there are few applications of Cu isotopes to study ancient biogeochemical cycles. Interestingly  
946 however, there are two black shale records that indicate a shift from UCC-like to heavy Cu isotope values  
947 across the Great Oxidation Event (GOE, ~2.4 Ga; Chi Fru et al., 2016; Ciscato et al., 2019). Chi Fru et al.  
948 (2016) interpreted this shift as reflecting the onset of oxidative weathering and waning of iron formation  
949 deposition, with the latter process driving pre-GOE seawater isotopically light due to the preferential  
950 scavenging of heavy Cu isotopes to Fe oxides. However, a recent analysis of two classic pre-GOE sequences  
951 containing iron formations do not support this earlier hypothesis, with  $\delta^{65}\text{Cu}$  remaining close to 0 ‰  
952 (Thibon et al., 2019). Thus, while the limited available data preclude confident interpretations, there are  
953 tantalizing tastes of future research directions in Cu isotope geochemistry.

954

955

956

957

958

959

## 960 **6. Cadmium**

961 Dissolved cadmium (Cd) concentrations closely correlate with the macronutrient phosphate in the oceans  
962 ( $\text{PO}_4^{3-}$ ; e.g., Boyle et al., 1976; Bruland, 1980). The nutrient-like properties of Cd—and attendant  
963 correlations with  $\text{PO}_4^{3-}$ —have been documented in multiple ocean basins and in multiple dimensions (i.e.,  
964 vertically, spatially, and temporally). This overall behavior implies that Cd somehow participates in marine  
965 biological processes and that the resulting distribution captured by sediments can be connected to  
966 underlying patterns of primary productivity. Recently, results from the GEOTRACES program have  
967 highlighted new nuances to this cycle that act to decouple phosphate and Cd in certain situations, which  
968 affect the interpretation of Cd distributions in the modern ocean as well as in the sedimentary record.  
969 Accordingly, this section will review the first-order features of marine Cd (and Cd isotope) distributions,  
970 discuss known driving mechanisms, highlight recent progress on identifying suitable sedimentary archives,  
971 and offer recommendations for future studies.


972

### 973 **6.1. Marine distribution**

974 Away from major upwelling regions, surface water dissolved Cd concentrations are typically between 1–  
975  $100 \text{ pmol kg}^{-1}$ , but can reach as low as  $30 \text{ fmol kg}^{-1}$  (Schlitzer et al., 2017; Fig. 12). The majority of this  
976 small surface inventory is thought to be complexed by strong organic ligands (e.g., Bruland, 1992; Ellwood,  
977 2004). In intermediate and deep waters, dissolved Cd concentrations are significantly elevated relative to  
978 surface waters, ranging from  $0.5 \text{ nmol kg}^{-1}$  in intermediate and deep waters in the north Atlantic to  $\approx 1.2$   
979  $\text{nmol kg}^{-1}$  in the oldest deep waters of the north Pacific (Schlitzer et al., 2018). As with Zn (Sec. 4.), the  
980 overall distribution of dissolved Cd throughout the oceans is driven principally by biological and physical  
981 processes in the Southern Ocean, and the lateral circulation of Southern Ocean water masses (e.g., Baars et  
982 al., 2014; Middag et al., 2018; Sieber et al., 2019a). Thus, the shape of vertical Cd concentration profiles at  
983 lower latitudes arises largely from horizontal transport and mixing of high-Cd Southern Ocean-sourced  
984 water masses, with a modest contribution from regeneration of sinking particles (i.e., 5–40 %; Middag et  
985 al., 2018). These processes result in Cd having ‘nutrient-type’ one dimensional water column profiles, with

986 a progressive increase in intermediate- and deep-water [Cd] along the pathways of meridional overturning  
 987 circulation (e.g., de Baar et al., 1994; Middag et al., 2018).

988



989  
 990 **Figure 12 | Representative profiles of dissolved Cd concentrations ([Cd]) and Cd-isotopic compositions**  
 991 **( $\delta^{114}\text{Cd}$ ) in the Eastern Tropical North Atlantic (squares, dashed line), Northeast Pacific (circles, solid line), and**  
 992 **Southern (triangles, dotted) Oceans.** Note the break in scale in  $\delta^{114}\text{Cd}$  above 1 ‰, illustrating the extreme isotopic  
 993 compositions observed in the most Cd-depleted surface samples. Notably, such extreme compositions are not  
 994 observed in the similarly Cd-depleted surface waters of the southern hemisphere gyres (e.g., Gault-Ringold et al., 2011;  
 995 Xie et al., 2017; George et al., 2019), nor in the surface of the high-latitude Southern Ocean, where dissolved [Cd] is  
 996 elevated (e.g., Abouchami et al., 2011; 2014). This comparison illustrates that the processes leading to distinct  
 997 dissolved concentration profiles effect only modest changes in dissolved  $\delta^{114}\text{Cd}$  between basins.

998


999 The past decade has seen an explosion in the number of studies employing Cd stable isotopes to investigate  
 1000 marine Cd cycling. The majority of extant Cd isotope data are reported relative to the NIST SRM 3108  
 1001 standard, though several earlier studies, reviewed by Rehkämper et al. (2012), were reported relative to  
 1002 other in-house materials. Cross-calibration of these materials is described in detail by Abouchami et al.  
 1003 (2013). While the earliest study of Cd isotope variations in seawater was unable to unambiguously identify  
 1004 systematic patterns in the water column, the authors noted that cultures of phytoplankton preferentially

1005 incorporated isotopically-light Cd relative to the media (Lacan et al., 2006). Assuming biological processes  
1006 were responsible for Cd uptake, this observation led to two key predictions for marine Cd isotope  
1007 systematics: that Cd-depleted surface waters should exhibit isotopically ‘heavier’ compositions than Cd-  
1008 replete deep waters; and, that the degree of isotopic fractionation should be proportional to the extent of Cd  
1009 removal into particles. Indeed, this is precisely the pattern that was first reported by Ripperger et al. (2007).

1010 Subsequent studies have corroborated this general one-dimensional pattern in the Southern (e.g.,  
1011 Abouchami et al., 2011; 2014; Xue et al., 2013; Sieber et al. 2019a), Atlantic (e.g., Xue et al., 2012; Conway  
1012 & John, 2015a; Xie et al., 2017; Xie et al., 2019a), Arctic (Zhang et al., 2019), and Pacific Oceans (Yang  
1013 et al., 2012; Yang et al., 2014; Conway & John, 2015b; Janssen et al., 2017; John et al., 2018; Yang et al.,  
1014 2018; Xie et al., 2019b; Sieber et al., 2019b). These studies have shown that the deep ocean (>500–1,000  
1015 m) is largely homogenous in Cd isotope composition ( $\delta^{114}\text{Cd}$  of +0.2 to +0.3 ‰). This deep water  $\delta^{114}\text{Cd}$   
1016 value is heavier than the upper continental crust  $\delta^{114}\text{Cd}$  signature of ~0‰ (Schmitt et al., 2009a), similar to  
1017 that observed for other metals such as Zn, Ba, and Ni (Sections 4, 8, and 9.). Intermediate-depth waters  
1018 relating to water masses such as Antarctic Intermediate Water (AAIW) exhibit slightly heavier  $\delta^{114}\text{Cd}$   
1019 values (+0.4 to +0.5 ‰), while surface waters generally exhibit heavier isotopic compositions (up to  
1020 ~+1‰). As for Cd concentrations, these one dimensional  $\delta^{114}\text{Cd}$  profile shapes arise largely from the  
1021 combination of Southern Ocean biological processes and lateral circulation of water masses, as well as  
1022 some contribution from local surface uptake and regeneration (Abouchami et al., 2014; Sieber et al., 2019b).  
1023 Some studies have reported  $\delta^{114}\text{Cd}$  values of up to +5 ‰ in surface waters of the northern hemisphere gyres  
1024 (Ripperger et al., 2007; Xue et al., 2012; Conway & John, 2015a; 2015b), while others have reported more  
1025 muted fractionation or even a switch to lighter than deep-ocean compositions in surface waters (Gault-  
1026 Ringold et al., 2012; Xie et al., 2017; Janssen et al., 2017; Xie et al., 2019a; George et al., 2019; Sieber et  
1027 al., 2019b; Xie et al., 2019). The origin of these extreme  $\delta^{114}\text{Cd}$  values remains ambiguous.

1028 Overall, although the first-order distributions of dissolved Cd concentration and isotope composition in  
1029 surface waters are consistent with intense cycling by biological processes, this range of observations  
1030 suggests that the processes influencing surface Cd isotope compositions may be more complex. For  
1031 example, interactions with ligands, effects arising from recycling of organic matter, differences in  
1032 fractionation factor between different organisms, and supply of Cd from external Cd sources may all  
1033 influence surface water dissolved  $\delta^{114}\text{Cd}$ . This raises three related questions to be addressed in the remainder  
1034 of this section: What are the candidate biological processes that drive these patterns? How do other physical

1035 or chemical processes play a role in setting marine Cd distributions? Can Cd-based reconstructions of these  
 1036 processes be used to inform on past productivity?  
 1037



1038  
 1039 **Figure 13 | Processes driving cadmium isotope variations in modern seawater.** Biological processes exert a  
 1040 significant control on surface water Cd cycling, implying that  $\delta^{114}\text{Cd}$  is broadly responsive to productivity. However,  
 1041 important redox-dependent processes remain to be fully elucidated, particularly those occurring around OMZs.

1042  
 1043 **6.2. Driving processes**

1044 *6.2.1. Biological*

1045 As noted above, the nutrient-like distribution of Cd implies intense biological cycling in seawater, even  
 1046 though Cd is considered toxic (e.g., Waldron & Robinson, 2009). This dichotomy has inspired a significant  
 1047 body of research investigating the role that Cd plays in microbial physiology. These studies showed that  
 1048 Cd uptake by marine microbes exhibits three noteworthy dependencies. First, cellular Cd quotas are  
 1049 strongly positively correlated with the Cd content of their environment, both in culture (see compilation by  
 1050 Twining & Baines, 2013) and from oceanographic data (Middag et al., 2018). Second, microbial Cd uptake

1051 is diminished when the concentration of other divalent cations increases, particularly so for Fe, Mn, and  
1052 Zn. Likewise, lower concentrations of these divalent cations cause increases in Cd uptake (e.g., Sunda &  
1053 Huntsman, 2000; Cullen et al., 2003). Some diatoms have even shown capacity to substitute much of their  
1054 metabolic Zn requirements with Cd (Price & Morel, 1990). Third, Cd uptake is also influenced by speciation  
1055 of inorganic carbon, with low  $p\text{CO}_2$  promoting higher cellular Cd quotas (e.g., Cullen et al., 1999; Cullen  
1056 & Sherrell, 2005; de Baar et al., 2017). The connection between Cd and carbon speciation is particularly  
1057 intriguing given the discovery of the  $\zeta$ -class of carbonic anhydrase that can utilize Cd (or Zn) as the catalytic  
1058 metal (e.g., Lane et al., 2005; Xu et al., 2008).

1059 Despite these dependencies, the extent to which active physiological utilization of Cd controls global  
1060 patterns of Cd uptake is unclear. For example, genes encoding the  $\zeta$  carbonic anhydrase metalloenzyme  
1061 were not found in green algae nor coccolithophores, and were similarly absent from many species of diatom  
1062 (Park et al., 2007). Thus, it is similarly plausible that some part of the biological Cd cycle is driven by  
1063 organisms inadvertently removing Cd from seawater while attempting to source other metals (e.g., Boyle,  
1064 1988b; Horner et al., 2013), or that microbes require Cd to populate other Cd-centered metalloenzymes that  
1065 await discovery.

1066 The role of microbial physiology in mediating Cd isotope fractionation is comparatively understudied.  
1067 Despite this, it appears that biological fractionation of Cd isotopes is ubiquitous; fresh- (Lacan et al., 2006)  
1068 and saltwater (John & Conway, 2014) green algae, incubations of unfiltered seawater (Xue et al., 2012),  
1069 heterotrophic bacteria (Horner et al., 2013), and shallow marine particles (e.g., Yang et al., 2015; Janssen  
1070 et al., 2019) all indicate that biological particles accumulate isotopically-light Cd from their environment  
1071 with a fractionation between  $-0.3$  to  $-0.8$  ‰ (Fig. 13). How—and perhaps even whether—this range of  
1072 offsets is related to environmental context, experimental setup, species, or specific microbial functions is  
1073 unknown.

1074

### 1075 *6.2.2. Chemical*

1076 The role of chemical processes in mediating global Cd distributions is the most under-constrained. Recent  
1077 studies suggested that pelagic partitioning of Cd into sulfides in oxygen-minimum zones may be a  
1078 significant loss term (Janssen et al., 2014; Bianchi et al., 2018; Plass et al., 2020). Moreover, the Cd isotope  
1079 effect associated with sulfide precipitation identified by Guoinseau et al. (2018) is consistent with field  
1080 data, and particulate Cd is known to accumulate more rapidly in sediments that are bathed by bottom waters  
1081 containing low dissolved oxygen (e.g., van Geen et al., 1995; Sec. 6.3; Fig. 13). Collectively, these



1082 observations suggest a potential redox sensitivity in sedimentary Cd isotope distributions that deserves  
1083 additional scrutiny. Indeed, the influence of sediments as the most important marine Cd sink can be seen in  
1084 some water column profiles of  $\delta^{114}\text{Cd}$  (Xie et al., 2019b). Similarly, data from the hydrothermal TAG site  
1085 in the North Atlantic suggests that hydrothermal plumes may scavenge Cd from seawater, constituting a  
1086 small sink of isotopically-light Cd, though this does not have an observable effect on deep ocean  $\delta^{114}\text{Cd}$   
1087 values outside of the plume itself (Conway & John, 2015b; Fig. 13).

1088 Other potential sources and sinks include rivers, atmospheric deposition, and sediments; however, none of  
1089 these interfaces exhibit significant anomalies in Cd concentration or  $\delta^{114}\text{Cd}$  in GEOTRACES-era datasets.  
1090 This finding is in accord with earlier research by Martin & Thomas (1994), though there exist two possible  
1091 exceptions. The first concerns the role of atmospheric aerosols, which have been invoked to explain the Cd  
1092 isotope composition of surface waters in the Southwest Pacific (e.g., George et al., 2019) and South China  
1093 Sea (e.g., Yang et al., 2012). Modern aerosol inputs may be largely anthropogenic in origin. Anthropogenic  
1094 forms of Cd exhibit a relatively narrow range of isotopic compositions that are typically—though not  
1095 always (e.g., Shiel et al., 2010)—lighter than dust-derived Cd (e.g., Bridgestock et al., 2017). Second,  
1096 interactions with organic ligands have also been invoked to explain the relatively muted pattern of Cd  
1097 isotope fractionation in the surface of the south Atlantic Ocean (e.g. Xie et al., 2017; Guinoiseau et al.,  
1098 2018), but there are as yet no corroborating field or experimental data examining the role of organic ligands  
1099 in mediating Cd isotope fractionation in seawater.

1100

### 1101 6.2.3. *Physical*

1102 Physical processes are similarly influential in mediating the global distribution and isotopic composition of  
1103 Cd throughout the global ocean, particularly those processes occurring in the Southern Ocean. Antarctic  
1104 Intermediate- and Bottom Waters masses possess higher dissolved  $[\text{Cd}]:[\text{PO}_4^{3-}]$  than North Atlantic Deep  
1105 Water (e.g., de Baar et al., 1994; Middag et al., 2018). Mixing between these southern- and northern-sourced  
1106 water masses likely contributes to the well-known ‘kink’ in the Cd– $\text{PO}_4^{3-}$  relationship (e.g., Frew & Hunter,  
1107 1992; Elderfield & Rickaby, 2000; Quay & Wu, 2015). Why northern- and southern-sourced intermediate  
1108 and deep waters possess different dissolved  $[\text{Cd}]:[\text{PO}_4^{3-}]$  is debated, and likely reflects regionally-distinct  
1109 fractionation of Cd and P during biological uptake (e.g., Sunda & Huntsman, 2000; Cullen et al., 2003) and  
1110 during remineralization (e.g., Baars et al., 2014; Roshan & Wu, 2015). In the Atlantic, however, the  
1111 importance of remineralization to deep water Cd budgets is of secondary significance: the ratio of  
1112 regenerated-to-preformed  $[\text{Cd}]$  is ~30 % in the mesopelagic, and generally <10 % in the deep ocean (Roshan  
1113 & Wu, 2015b; Middag et al., 2018). Given the low proportion of regenerated Cd in the deep Atlantic, the

1114 ratio of dissolved  $[Cd]:[PO_4^{3-}]$  and the overall distribution of  $[Cd]$  are essentially governed by the  
1115  $[Cd]:[PO_4^{3-}]$  of the source waters (Middag et al., 2018) and the prevailing geometry of ocean circulation,  
1116 respectively (Boyle, 1988b).

1117 Recent Cd isotope data from the South Atlantic (Xie et al., 2017), South Pacific (George et al., 2019; Sieber  
1118 et al., 2019a), and Arctic (Zhang et al., 2019) also support the importance of mixing in mediating deep-  
1119 ocean Cd distributions, though it should be noted that the isotopic contrast between mixing end-members  
1120 is small, relative to measurement precision (e.g., Janssen et al., 2017; Fig. 12). For example, biological  
1121 uptake of light Cd in the source regions of intermediate waters in the surface Southern Ocean results in  
1122 isotopically-heavy pre-formed  $\delta^{114}Cd$  signatures being imparted to Cd-depleted intermediate water masses  
1123 (e.g., +0.45 ‰ in AAIW; +0.65 ‰ in Subantarctic Mode Water, SAMW; Xue et al., 2013; Abouchami et  
1124 al., 2014; Sieber et al., 2019b). Lateral circulation of these southern-sourced water masses then transfer this  
1125 signature northward to intermediate depths in the Atlantic and Pacific Oceans (e.g. Xue et al., 2012;  
1126 Abouchami et al., 2014; Conway & John, 2015a; Sieber et al., 2019b). This effect is more pronounced in  
1127 the North Atlantic than in the Pacific, where southward flowing NADW also carries isotopically-heavy Cd  
1128 southward at depths of 1,000–3,000m (Xue et al., 2012; Conway & John, 2015a).

1129

### 1130 **6.3. Marine archives**

#### 1131 *6.3.1. Carbonates*

1132 There is a long history of Cd concentration measurements in marine carbonates, particularly corals and  
1133 foraminifera, most commonly reported as Cd:Ca molar ratios. In principle, carbonates are an appealing  
1134 archive of ambient Cd chemistry since inorganic partition coefficients are  $\gg 1$  (e.g., Tesoriero & Pankow,  
1135 1996) and the Cd:Ca of many types of carbonate exhibit a strong proportionality with ambient dissolved  
1136  $[Cd]$ . In practice, however, most biogenic carbonates exhibit partition coefficients closer to unity (Boyle,  
1137 1988b), and resultant Cd:Ca is also sensitive to the species (Boyle, 1992) and temperature of calcification  
1138 (e.g., Rickaby & Elderfield, 1999). As such, Cd:Ca in carbonates has found the most utility where ambient  
1139 dissolved  $[Cd]$ —and attendant carbonate Cd:Ca—exceeds several  $100 \text{ pmol kg}^{-1}$ , such as in tracing  
1140 industrial fallout (e.g., Shen et al., 1987) or in studies of Quaternary deep ocean circulation (e.g., Boyle &  
1141 Keigwin, 1985; van Geen et al., 1992; Adkins et al., 1998; Farmer et al., 2019).

1142 There are far fewer studies examining the Cd isotope composition of marine carbonates as tracers of  
1143 historical Cd isotope chemistry. Inorganic partitioning experiments indicate that Cd isotopes are  
1144 fractionated during incorporation into calcite by  $\approx -0.5 \text{ ‰}$  (Horner et al., 2011). The isotopic effect is


1145 temperature- and Mg-independent, but vanishes at low salinity. This inorganic calibration has been used to  
1146 interpret patterns of Cd isotope fractionation preserved in bulk carbonates from the Neoproterozoic Eon  
1147 (1,000–541 Ma). The variations in these sediments are interpreted as evidencing changes in biological  
1148 productivity (e.g., Hohl et al., 2017) and Cd sinks (e.g., John et al., 2017) through time. Applications of Cd  
1149 isotopes to foraminifera to study problems in Quaternary paleoceanography are precluded by the large  
1150 sample requirements; obtaining 1 ng of Cd—the minimum quantity needed for a reasonable isotope  
1151 measurement (Ripperger & Rehkämper, 2007; Schmitt et al., 2009b)—requires picking (and cleaning) of  
1152 10's of mg of foraminiferal tests. Alleviation of such limitations awaits development of automated picking  
1153 and screening systems, or vast improvements in ion transmission efficiency for isotope ratio mass  
1154 spectrometry.

1155

### 1156 6.3.2. *Ferromanganese sediments*

1157 Ferromanganese sediments have shown the most promise for recording deep-ocean Cd isotope chemistry.  
1158 Both ferromanganese nodules (Schmitt et al., 2009a) and crusts (Horner et al., 2010) reflect ambient  
1159 seawater Cd isotope compositions with negligible fractionation (Fig. 14A), consistent with Cd–Mn-  
1160 oxyhydroxide partitioning experiments conducted at high ionic strength that show only minor Cd isotope  
1161 fractionation (Wasylenki et al., 2014). Cadmium is predicted to have a similar diffusivity in Fe–Mn crusts  
1162 to Hf (Horner et al., 2010), implying that long-term records are likely to exhibit some diffusive smoothing  
1163 while preserving larger variations (e.g., Henderson & Burton, 1999). As with Zn and Cu isotopes however,  
1164 time-resolved records of  $\delta^{114}\text{Cd}$  recovered from Fe–Mn crusts indicate minimal variation over the last 20  
1165 Myr, and perhaps even the Cenozoic (Fig. 14B); modeling conducted by Murphy (2016) suggests that Cd  
1166 diffusion may occur much faster than predicted by the diffusion model of Henderson & Burton (1999). If

1167 so, the lack of variation in  $\delta^{114}\text{Cd}$  over the Cenozoic may simply reflect a ‘resetting’ of all crust layers  
 1168 toward modern deep-ocean Cd isotope compositions.




1169  
 1170 **Figure 14 | Ferromanganese crusts records of deep-water Cd isotope compositions. (A)** Compilation of ‘core-  
 1171 top’ Fe–Mn crust  $\delta^{114}\text{Cd}$ ; data from Schmitt et al. (2009a; open symbols) and Horner et al. (2010; closed symbols). In  
 1172 general, Southern Ocean samples exhibit lighter  $\delta^{114}\text{Cd}$  than other basins, consistent with profiles of  $\delta^{114}\text{Cd}$  (e.g., Fig.  
 1173 12). **(B)** Cenozoic records of deep-water  $\delta^{114}\text{Cd}$  recovered from four Fe–Mn crusts; data from Murphy, 2016 (closed  
 1174 symbols) and Schmitt et al. (2009a; open symbols). These records have been plotted using the authors’ preferred age  
 1175 models, meaning that there are some differences between the chronology of CD29-2 shown here compared to Fig. 5.  
 1176 Such differences may be immaterial however, given that Cd isotopes in Fe–Mn crusts are potentially subject to diffusive  
 1177 ‘resetting’ over time (Murphy, 2016).

1178  
 1179 **6.3.3. Organic-rich sediments**

1180 Organic-rich sediments are the principal sink of dissolved Cd from the modern oceans (e.g., Rosenthal et  
 1181 al., 1995; van Geen et al., 1995; Little et al., 2015). A significant fraction of the total Cd in organic-rich  
 1182 sediments is derived from sinking organic matter. Accordingly, the Cd content of organic-rich sediments  
 1183 has been applied as a proxy for past productivity (e.g., Brumsack, 2006). However, there is also a redox  
 1184 sensitivity: Cd contents are generally elevated in organic-rich sediments that are bathed by low-oxygen  
 1185 waters (Fig. 15). This enrichment likely derives from three processes. First, low-oxygen environments may  
 1186 limit oxidation—and thus favor preservation—of settling Cd sulfide particles formed in the water column.  
 1187 Second, the chalcophile nature of Cd means that even trace levels of hydrogen sulfide may cause Cd to (co-

1188 )precipitate. Thus, as organic matter is remineralized within the sediment column, any Cd liberated to  
 1189 porewaters is proportionally more likely to re-precipitate into sulfide minerals, relative to harder metals  
 1190 (and carbon), effectively ‘trapping’ remineralized Cd in sediments (e.g., Rosenthal et al., 1995). Third,  
 1191 recent evidence suggests that Cd may also directly (co-)precipitate from seawater and into sediments when  
 1192 plumes of hydrogen sulfide interact with bottom waters (e.g., Xie et al., 2019b; Plass et al., 2020). The  
 1193 relative importance of these three processes to the O<sub>2</sub>-dependent pattern of Cd accumulation remains to be  
 1194 fully elucidated.

1195 While the contribution of organic-rich sediments to the isotopic mass balance of Cd is presently unknown,  
 1196 their significance to the marine Cd budget suggests that, globally, the Cd isotope composition of these  
 1197 sediments should balance the riverine flux  $\delta^{114}\text{Cd} \approx +0.1 \pm 0.1\%$  (Lambelet et al., 2013). Alternatively, the  
 1198 other minor sinks of Cd—carbonates, ferromanganese oxides, clays—must possess large isotopic offsets  
 1199 relative to seawater, which seems unlikely given existing field and experimental data. Obtaining modern  
 1200 calibrations of Cd isotope partitioning into organic-rich sediments should be considered a priority.



1201  
 1202 **Figure 15 | Cadmium concentrations in California Margin sediments.** Left: Regionally-representative dissolved  
 1203 [O<sub>2</sub>] profile from the northeast Pacific showing broad minimum between 600–800 m. Profile from 35.5°N, 122.5°W (from  
 1204 World Ocean Atlas; Garcia et al., 2013). Right: Solid-phase Cd concentrations for northeast Pacific core-top (closed  
 1205 symbols) and “slightly deeper” (8–10 cm; open symbols) sediment samples; dashed line indicates arithmetic mean.  
 1206 These samples evidence a maximum in authigenic Cd deposition at the top of the OMZ, which may originate from  
 1207 processes occurring in the water or sediment column.

1208

1209 **6.4 Prospects**

1210 The overview provided above indicates that Cd participates in marine biological processes and that its  
 1211 distribution is sensitive to the biological productivity of the oceans. How this sensitivity is transcribed into  
 1212 marine sediments remains uncertain, however. Additionally, there are several other processes that have the  
 1213 potential to render isotope effects that require further exploration before Cd isotopes can be solely  
 1214 interpreted as a productivity proxy, such as: biological fractionation effects, authigenic transformations,  
 1215 and redox sensitivities. Relatedly, the fidelity of many types of marine sediment to record ambient Cd  
 1216 isotope chemistry remain inadequately constrained. With these two themes in mind, we suggest several  
 1217 areas for additional research that may help to address the overarching question as to whether Cd isotope-  
 1218 based reconstructions can be used to inform on past ocean productivity.

1219

1220 **6.4.1. Modern**

1221 Several questions persist regarding the modern Cd isotope cycle. We list five of the most pressing below  
 1222 and offer possible remedies to each.

- 1223 ● Are the ‘extreme’ (i.e.,  $\delta^{114}\text{Cd}$  of + 5 ‰; Fig. 12) values seen in surface waters of the northern-  
 1224 hemisphere gyres real? That is, are these heavy compositions true oceanographic features that are  
 1225 absent from the southern hemisphere, or do they represent analytical artifacts unique to MC-ICP-  
 1226 MS? This issue will require intercalibration between the two main approaches—the other being  
 1227 thermal ionization mass spectrometry (TIMS)—that various groups use to measure Cd isotopes in  
 1228 seawater.
- 1229 ● To what extent do local Cd isotope compositions in surface waters reflect larger-scale processes?  
 1230 Addressing this issue will require elucidating the role of organismal uptake, ligands, and mixing,  
 1231 which would benefit from conducting additional experimentation with plankton, coupling of  
 1232 isotope methods with electrochemistry, and numerical modeling, respectively.
- 1233 ● Is there a ‘Redfield’ stoichiometry for Cd in organic matter? If so, what controls it? Existing culture  
 1234 (Ho et al., 2003) and field (Ohnemus et al., 2017) data suggest a wide range of cellular Cd:P, which

1235 have been suggested as reflecting species and local supply ratios, respectively (in addition to the  
 1236 aforementioned feedback interactions). Further experimentation with model organisms is needed.  
 1237 ● Do all organisms contribute to Cd isotope fractionation in seawater? Uptake of Cd into cells is  
 1238 widespread, whereas the genes encoding the  $\zeta$ -class of carbonic anhydrases are not. Again, further  
 1239 experimentation with marine-relevant organisms is necessary.  
 1240 ● How are the ‘light’ Cd isotope compositions seen in suspended particles above oxygen-minimum  
 1241 zones related to those putatively accumulating in sediments? Do these values relate to sulfide  
 1242 precipitation and what is their influence on the global mass balance? What controls the Cd isotope  
 1243 composition of these particles? Addressing these questions will require examining the Cd isotope  
 1244 composition of particles from oxygen-rich oceanographic regions, identifying whether processes  
 1245 associated with particle regeneration affect Cd isotope compositions, and surveying core-top  
 1246 sediments.

1247

1248 *6.4.2. Paleo*

1249 As with the modern cycle, several ambiguities persist, though the most pressing relates to archives. Indeed,  
 1250 it appears that a major obstacle preventing the widespread application of Cd isotope-based proxies in  
 1251 paleoceanography has been the lack of suitable archives.

- 1252 ● Concerning carbonates, the main question is still whether it is possible to isolate sufficient  
 1253 quantities of foraminiferal-bound Cd to reconstruct past seawater  $\delta^{114}\text{Cd}$ . Does species matter, or  
 1254 can mixed assemblages be used?
- 1255 ● For organic-rich sediments, the role of biology itself must be considered. That is, how important  
 1256 might it be that different organisms exhibit different magnitudes of Cd isotope fractionation (and  
 1257 Cd:C stoichiometry)? If important, how best to interpret Cd isotope records—species, evolutionary  
 1258 innovations, productivity?

1259


1260



1261 **7. Molybdenum**

1262 **7.1. Marine distribution**

1263 Today, Mo is one of the most abundant trace metals in seawater (Morris, 1975; Bruland 1983; Collier 1985).  
 1264 Molybdenum possesses a long residence time (~440,000 yr; Miller et al., 2011), is conservative with respect  
 1265 to salinity, and exhibits a uniform isotopic composition in oxygenated seawater (e.g., Barling et al., 2001;  
 1266 Siebert et al., 2003). Based on 168 seawater samples from the Atlantic, Pacific, and Southern Oceans  
 1267 analyzed by Nakagawa et al. (2012), the average salinity-normalized dissolved [Mo] and  $\delta^{98}\text{Mo}$  of the ocean  
 1268 are  $107 \pm 6 \text{ nmol kg}^{-1}$  and  $+2.36 \pm 0.10 \text{ ‰}$ , respectively (both values  $\pm 2 \text{ SD}$ ; Fig. 16).




1269 **Figure 16 | Representative profiles of dissolved Mo concentrations ([Mo]) and Mo-isotopic compositions**  
 1270 **( $\delta^{98}\text{Mo}$ ) in the Northwest Atlantic (squares, dashed line), Southern (triangles, dotted line), and Northwest**  
 1271 **Pacific (circles, solid line) Oceans. This comparison illustrates that the dissolved behavior of Mo is essentially**  
 1272 **invariant throughout the global oceans.**  
 1273

1274  
 1275 Given that no significant gradients in dissolved [Mo] are expected, there are few new open-ocean  $\delta^{98}\text{Mo}$   
 1276 data measured as part of GEOTRACES. However, dissolved [Mo] has been measured on two  
 1277 GEOTRACES transects (GP16 and GA02), demonstrating four intriguing instances where [Mo] deviated

1278 from an otherwise conservative distribution. These instances are briefly described below and covered in  
1279 detail by Ho et al. (2018). First, the most significant dissolved [Mo] anomalies (~5 % drawdown) are  
1280 associated with intense scavenging by particulate Fe (oxy)hydroxides close to the Peruvian OMZ. Second,  
1281 dissolved [Mo] is slightly drawn down (<5 %) in some samples directly above the East Pacific Rise  
1282 hydrothermal ridge crest, again mostly likely driven by scavenging onto hydrothermally-derived Fe–Mn  
1283 oxides. Some Mo drawdown is also observed in some far-field hydrothermal samples, though the  
1284 mechanism is unclear. Third, following normalization to a salinity of 35, surface seawater shows a minor  
1285 dissolved [Mo] drawdown, implying either biological uptake or adsorption to biotic particles. Though not  
1286 a true oceanographic feature, a fourth type of dissolved Mo anomaly is also noteworthy: bottle storage  
1287 artefacts. Ho et al. (2018) found that many samples with initially low values showed an increase in dissolved  
1288 [Mo] with increasing storage time, implying a change in Mo speciation to a form that is detectable by ICP-  
1289 MS. In contrast to the relative constancy of dissolved [Mo] in open ocean settings, there are a number of  
1290 studies of showing striking dissolved Mo variations in coastal and estuarine systems (e.g., Dalai et al., 2005;  
1291 Dellwig et al., 2007; Joung and Shiller, 2016; Wang et al., 2016) as well as in modern restricted settings,  
1292 such as the Black and Baltic Seas (Nägler et al., 2011).

1293



1294

1295 **Figure 17 | Processes driving molybdenum isotope variations in modern seawater.** While biological processes  
 1296 may exert a slight influence on surface water Mo distributions, the main drivers of marine Mo cycling are related to the  
 1297 balance between scavenging pathways, which are redox dependent.

1298

## 1299 7.2. Driving processes

### 1300 7.2.1. Biological

1301 Molybdenum is a bio-essential micronutrient required by enzymes that catalyze key reactions in the global  
 1302 C, N, and S cycles (Mendel & Bittner, 2006). Importantly, Mo is a co-factor of the primary nitrogenase  
 1303 enzyme complex, meaning that Mo is required for energy-efficient nitrogen fixation. Additionally, Mo is  
 1304 required for over 30 other enzymes that control biologically essential redox processes (Kendall et al., 2017).  
 1305 Despite its biological importance, biological activity does not appear to be a dominant process in setting  
 1306 the distribution of Mo in seawater (Fig. 17), likely due to the high abundance of Mo in the modern ocean.  
 1307 Results from the East Pacific Zonal Transect do, however, suggest some Mo removal by biological uptake  
 1308 and/or adsorption onto biogenic particles within regions of elevated chlorophyll (Ho et al., 2018).  
 1309 Experimental data indicate that biological uptake of Mo imparts a small negative isotope fractionation on  
 1310 the order of 0.3 ‰ (Wasylenki et al., 2007).

1311 Enhanced removal of Mo from seawater in regions with high export of organic carbon likely explains some  
1312 of the non-conservative behavior observed in modern coastal regions. A significant relationship between  
1313 Mo and total organic carbon has also been observed in marine euxinic sediments (e.g. Helz et al., 1996;  
1314 Algeo & Lyons, 2006; McManus et al., 2006; Lyons et al., 2009). Furthermore, it has been shown that Mo  
1315 interactions with organic matter can control Mo cycling in both soils (Wichard et al., 2009; King et al.,  
1316 2014, 2016; Marks et al., 2015; Siebert et al., 2015) and marine sediments (Wagner et al., 2017; Tessin et  
1317 al., 2019). The formation of Mo complexes containing organic ligands has been proposed as an explanation  
1318 for the correlation between sedimentary Mo and organic carbon, suggesting that organic matter may play  
1319 an important role in both the delivery and burial of Mo in sediments (Wagner et al., 2017). Accordingly, it  
1320 is possible that Mo accumulation in marine sediments is linked to export production and, more specifically,  
1321 delivery and burial of organic carbon in sediments.

1322

### 1323 *7.2.2. Chemical*

1324 Due to the long residence time and abundance of Mo, non-conservative Mo behavior is typically localized  
1325 to areas with high particle concentrations, such as close to the continental margin, in the euphotic ocean, or  
1326 around hydrothermal vents (e.g., Goto et al., 2020). In the present oxic water column, Mo is present  
1327 primarily as the oxyanion molybdate ( $\text{MoO}_4^{2-}$ ) and Mo sorption onto Mn oxyhydroxides represents the  
1328 most significant modern Mo sink (Fig. 17; Bertine & Turekian, 1973; Scott & Lyons, 2012). Since the Fe–  
1329 Mn oxide sink preferentially incorporates isotopically light Mo from seawater with an isotopic effect of  
1330  $\approx 3.0$  ‰, the fraction of Mo that is buried in Fe–Mn oxides exerts a major control over the Mo isotope  
1331 composition of seawater.

1332 In the presence of sulfide, the oxygen atoms in molybdate are progressively substituted for sulfur, producing  
1333 particle reactive thiomolybdate species ( $\text{MoO}_{4-x}\text{S}_x^{2-}$ ; Erickson & Helz, 2000; Vorlicek et al., 2015).  
1334 Dissolved Mo can thus be strongly drawn down in sulfidic environments, such as the Black Sea.  
1335 Importantly, this drawdown occurs with a small, but non-zero isotope effect of  $\approx 0.5 \pm 0.3$  ‰, whereby  
1336 isotopically light Mo is preferentially scavenged (e.g., Nägler et al., 2011). Thiomolybdate is also  
1337 scavenged in from sulfidic sediment porewaters leading to significant Mo accumulations within sediments  
1338 deposited in anoxic and euxinic environments (e.g., Emerson & Husted, 1991; Crusius et al., 1996; Scott  
1339 & Lyons, 2012). Long-term Mo burial is associated with Fe–S minerals (Chappaz et al., 2014; Vorlicek et  
1340 al., 2018) and/or organic matter (Dahl et al., 2017; Tessin et al., 2019), depending on the biogeochemical  
1341 conditions prevailing at a given basin.

1342

1343 **7.2.3. Physical**

1344 The residence time of dissolved Mo is significantly longer than the mixing time of the ocean (Table 1).  
1345 Thus, variations in dissolved [Mo] and  $\delta^{98}\text{Mo}$  are not influenced by the geometry of modern overturning  
1346 circulation. Whole-ocean changes in dissolved [Mo] and  $\delta^{98}\text{Mo}$  are, however, possible, and much of what  
1347 we know about such changes is gleaned from studies of the geological past, particularly during periods of  
1348 ocean anoxia. Based on this research, it has been suggested that during intervals of lower ocean  
1349 oxygenation, the ocean Mo inventory may have been low enough to limit marine primary productivity  
1350 (Anbar & Knoll, 2002; Algeo, 2004; Glass et al., 2009; Reinhard et al., 2013). This limitation has been  
1351 demonstrated in certain lake ecosystems (Goldman, 1960; Glass et al., 2012). Under strongly euxinic  
1352 conditions ( $[\text{H}_2\text{S}]_{\text{aq}} > 11 \mu\text{M}$ ), thermodynamic calculations predict that tetrathiomolybdate ( $\text{MoS}_4^{2-}$ ) becomes  
1353 the most predominant species (Erickson and Helz, 2000), which can result in quantitative Mo drawdown  
1354 from seawater into seawater with negligible isotope fractionation. Evidence from Lake Cadagno, the Black  
1355 and Baltic Seas, and seasonally anoxic fjords supports the occurrence of near-quantitative drawdown of Mo  
1356 from seawater in these environments (Neubert et al., 2008; Dahl et al., 2010; Nägler et al., 2011;  
1357 Noordmann et al., 2015). Indeed, this hypothesis is the foundation of several paleoceanographic studies that  
1358 assume that the sedimentary Mo isotopic signatures, deposited in euxinic settings, faithfully capture the  
1359  $\delta^{98}\text{Mo}$  of oxygenated (surface) seawater and can be further interpreted in terms of the fraction of the seafloor  
1360 that is oxygenated (e.g., Kendall et al., 2015; Dickson, 2017).

1361

1362 **7.3. Sedimentary archives**

1363 The majority of Mo paleoceanographic studies focus on the measurement of Mo concentrations and  
1364 isotopes within bulk sediments. While many applications of Mo within the sedimentary record focus on  
1365 early Earth, Mo proxies have been used to constrain environmental conditions in the Holocene (e.g.  
1366 Hardisty et al., 2016; van Helmond et al., 2018) and Pleistocene (e.g. Dean et al., 2006; Scholz et al., 2017).  
1367 However, Mo and its isotopes are predominantly used as tracers of redox conditions and not productivity.

1368 Research on biological archives of Mo concentrations and isotopes has met with mixed results. Research  
1369 examining  $\delta^{98}\text{Mo}$  in corals indicates that corals may accurately record seawater Mo isotope concentrations  
1370 (Voegelin et al., 2009). However, follow-up studies suggested a temperature-dependent fractionation  
1371 between seawater and corals that is related to the activity of symbiotic zooxanthellae (Wang et al., 2019b).  
1372 Moreover, bivalve shell Mo:Ca ratios have been determined to have no relationship to oceanographic

1373 conditions (Vihtakari et al., 2017), whereas Tabouret et al. (2012) suggest a mechanism related to trophic  
1374 uptake, but not to ambient dissolved [Mo]. A third study proposed that Mo:Ca peaks in bivalve carbonate  
1375 are controlled by ingestion of phytoplankton grown on nitrate due to high concentrations of Mo associated  
1376 with nitrate reductase, indicating that bivalves may provide an archive for surface water nitrate uptake and  
1377 a potential proxy for the balance between new and regenerated productivity (Thebault et al., 2009).

1378

#### 1379 **7.4. Prospects**

1380 At present, it does not appear that bulk sediment  $\delta^{98}\text{Mo}$  will be useful for reconstructing biological  
1381 productivity, even though Mo is cycled by organic matter both actively (e.g., Liermann et al., 2005;  
1382 Wasylenki et al., 2007) and passively (e.g., Kowalski et al., 2013; King et al., 2018). That we see no  
1383 pathway to using Mo as a productivity tracer reflects, in part, the difficulty in disentangling biological  
1384 processes that exert relatively modest Mo isotope fractionations from those associated with thiomolybdate  
1385 transformations (Tossell, 2005), or scavenging processes that possess large fractionation effects (e.g., Mn-  
1386 or Fe-oxide scavenging; Siebert et al., 2003; Barling & Anbar, 2004; Wasylenki et al., 2008; Brucker et al.,  
1387 2009; Goldberg et al., 2009; 2012). Additionally, within the modern ocean and likely within the recent  
1388 geologic past, the global ocean reservoir of Mo is too large and well mixed for biological associated  
1389 fractionations to significantly impact the global Mo isotopic composition.

1390 The use of bulk sedimentary Mo concentrations as a proxy for export of organic carbon to the seafloor is  
1391 more promising, but numerous caveats exist. Specifically, other mechanisms for enhanced delivery,  
1392 sequestration, and burial complicate any efforts to quantitatively relate Mo enrichments to increased export  
1393 productivity (e.g., Scholz et al., 2017). Redox conditions and, in particular, the presence of sulfide in the  
1394 water column and sediment pore waters will be a primary control on Mo accumulation (e.g., Hardisty et al.,  
1395 2018). Sedimentary Mo enrichments can also be produced through shuttling of Mo adsorbed to the surface  
1396 of Fe and Mn oxides to the seafloor (Algeo & Lyons, 2006; Algeo & Tribovillard, 2009; Dellwig et al.,  
1397 2010; Scholz et al., 2013). At a minimum, independent constraints on water column and pore water redox  
1398 conditions using Fe speciation, other trace metals and/or fossil redox proxies are required before an  
1399 argument can be made relating Mo concentrations to export productivity. Additionally, the quantitative  
1400 relationship between Mo and organic carbon may be impacted by aqueous Mo concentrations, which may  
1401 have varied over Earth's history, or if depositional environments become restricted (i.e., Mo drawdown  
1402 leads to a lower Mo:TOC ratio; Algeo & Lyons, 2006). Conversely, the utility of bulk sediment Mo

1403 concentrations and isotopes may lie in constraining redox conditions to improve the interpretation of other  
1404 trace metal proxies that are more strongly controlled by primary and/or export productivity.

1405 The most promising future avenue for Mo-based productivity proxies may emerge from fossil-specific  
1406 measurements of Mo concentrations (and perhaps  $\delta^{98}\text{Mo}$ ). Currently, research has focused on large fossils  
1407 (corals and bivalves) and has led to mixed results on the utility of Mo in reconstructing productivity-related  
1408 parameters. However, analytical progress may allow for measurement of smaller sample sizes, which may  
1409 provide opportunities to explore new archives of past marine Mo geochemistry.

1410


1411

1412



1413 **8. Barium**

1414 Barium is a widely-used tracer of marine export production, past and present. This section provides an  
 1415 overview of marine Ba geochemistry, focusing on driving mechanisms, palaeoceanographic applications,  
 1416 and emerging insights from Ba stable isotopes. We close by suggesting several areas for additional research.



1417  
 1418 **Figure 18 | Representative profiles of dissolved Ba concentrations ( $[Ba]$ ) and Ba-isotopic compositions ( $\delta^{138}Ba$ )**  
 1419 **from the Eastern Tropical North Atlantic (squares, dashed line), northeast Pacific (circles, solid line), and**  
 1420 **Southern (triangles, dotted lines) Oceans.** This comparison illustrates that the oceanographic processes leading to  
 1421 distinct dissolved concentration profiles render significant changes in dissolved  $\delta^{138}Ba$  between basins.

1422

1423 **8.1. Marine distribution**

1424 The nutrient-like distribution of dissolved Ba has been documented in the literature since the 1970's (e.g.,  
 1425 Wolgemuth & Broecker, 1970). However, it was not until later in the decade that the GEOSECS Program  
 1426 fully revealed the three-dimensional marine distribution of Ba (e.g., Chan et al., 1976; 1977). These  
 1427 geochemical ocean sections highlighted vertical, zonal, and meridional variations in dissolved  $[Ba]$  related  
 1428 to the major biogeochemical and hydrographic features of the ocean. In nutrient-depleted surface waters,  
 1429 dissolved  $[Ba]$  exhibits low concentrations between 35–45 nM. Similarly, nutrient-replete deep waters are

1430 enriched in [Ba], though typically by no more than a factor of four above surface values. The spatial  
1431 resolution of GEOSECS illustrated the importance of hydrography; [Ba] increases along the meridional  
1432 overturning circulation from  $\approx 50$  nM in deep waters of the North Atlantic, to  $\approx 90$  nM in the Southern Ocean,  
1433 up to  $\approx 180$  nM in the deep northeast Pacific (Fig. 18; Chan et al., 1976; 1977).


1434 Though it was long suspected that the major dissolved–particulate transformation of Ba was related to the  
1435 mineral barite ( $\text{BaSO}_4$ ; e.g., Chow & Goldberg, 1960; Turekian, 1968), this was not confirmed until the  
1436 1980's (e.g., Dehairs et al., 1980; Bishop, 1988). Barite crystals are now recognized as an ubiquitous  
1437 component of marine particulate matter, with up to  $10^4$  discrete, micron-sized crystals present per L of  
1438 seawater (Dehairs et al., 1980). However, the distribution of particulate  $\text{BaSO}_4$  is distinct from primary  
1439 biogenic phases that exhibit Martin-like distributions with maxima in the euphotic zone. The minimum  
1440 particulate Ba concentration is typically situated in the euphotic zone and the maximum slightly below,  
1441 usually in the upper mesopelagic (e.g., Ohnemus & Lam, 2015; Ohnemus et al., 2019). This distribution  
1442 likely reflects the fact that neither Ba nor  $\text{BaSO}_4$  are utilized for physiological processes by any of the major  
1443 marine primary producers. Particulate Ba fluxes are nevertheless strongly correlated with export  
1444 productivity in well-oxygenated environments (e.g., Bishop, 1988; Dymond et al., 1992; Francois et al.,  
1445 1995; Dymond & Collier, 1996; McManus et al., 2002) and therefore sedimentary Ba abundances have  
1446 been widely used as a proxy to reconstruct past changes in ocean export production (e.g., Francois et al.,  
1447 1995; Paytan et al., 1996; Paytan & Griffith, 2007 and references therein; Costa et al., 2016, Winckler et  
1448 al., 2016).

1449 Recent studies of Ba stable isotope geochemistry have added a new dimension with which to study marine  
1450 Ba cycling. Von Allmen et al. (2010) first reported that isotopically light Ba is preferentially incorporated  
1451 into  $\text{BaSO}_4$ , with a particulate–dissolved Ba-isotopic offset of  $\approx -0.3$  ‰. The direction of this offset implies  
1452 that residual solutions, such as seawater, should exhibit Ba-isotopic compositions heavier than those of  
1453 sedimented  $\text{BaSO}_4$ . This was corroborated for Atlantic seawater by Horner et al. (2015a), showing that  
1454 dissolved  $\delta^{138}\text{Ba}$  displays a mirror image of dissolved [Ba]: Ba-depleted surface water masses exhibited  
1455 ‘heavy’ Ba-isotopic compositions ( $\approx +0.6$  ‰), whereas Ba-replete deep waters possessed ‘lighter’ values  
1456  $\approx +0.3$  ‰. (Notably, all values are considerably heavier than the upper continental crust, which possesses  
1457  $\delta^{138}\text{Ba} \approx 0.0 \pm 0.1$  ‰; Nan et al., 2018.) Similar patterns have since been corroborated in other ocean basins  
1458 (Bates et al., 2017; Hsieh & Henderson, 2017; Bridgestock et al., 2018; Geyman et al., 2019; Cao et al.,  
1459 2020; Fig. 18).

1460 These isotopic studies have yielded a number of novel insights into the marine Ba cycle. First, these data  
1461 underscore the importance of physical mixing (i.e., ocean circulation) in mediating patterns of dissolved

1462  $\delta^{138}\text{Ba}$  and, by extension, [Ba]. Second, both the regression of dissolved Ba-isotopic data (Bates et al., 2017;  
1463 Hsieh & Henderson, 2017) and comparison of co-located seawater and particulates (Horner et al., 2017)  
1464 imply an average particulate–dissolved Ba-isotopic offset  $\approx -0.5$  ‰, somewhat larger than the experimental  
1465 results reported by Von Allmen et al. (2010). Third, marine sediments—both bulk (Bridgestock et al., 2018)  
1466 and  $\text{BaSO}_4$  isolates therefrom (Crockford et al., 2019)—faithfully reflect the  $\approx -0.5$  ‰ Ba-isotopic offset  
1467 from surface seawater. Consequently, the mean  $\delta^{138}\text{Ba}$  of globally-sedimented  $\text{BaSO}_4$  is  $\approx +0.1 \pm 0.1$  ‰  
1468 (Crockford et al., 2019). Since  $\text{BaSO}_4$  is the dominant oceanic output (e.g., Paytan & Kastner, 1996), these  
1469 data imply that mean isotopic composition of Ba delivered to the ocean should possess an average  
1470 composition  $\approx +0.1 \pm 0.1$  ‰. At present however, the main Ba inputs are unable to close the marine Ba-  
1471 isotopic budget: rivers, the principal Ba source to seawater, rivers, are too heavy, exhibiting compositions  
1472 generally  $\geq +0.2$  ‰ (e.g., Cao et al., 2020; Gou et al., 2020); and, groundwater discharge, while possessing  
1473 the necessary ‘light’ composition of  $\approx +0.1 \pm 0.1$  ‰, is too small a Ba flux to balance the budget (Mayfield

1474 et al., in press). Thus, either the marine Ba-isotopic budget is currently out of steady state, or an additional  
 1475 Ba source possessing a light isotopic composition remains to be identified.  
 1476



1477  
 1478 **Figure 19 | Processes driving barium isotope variations in modern seawater.** Though biological processes exert  
 1479 only a minor direct influence on  $\delta^{138}\text{Ba}$ , the biologically-mediated cycling of  $\text{BaSO}_4$  drives large variations in marine Ba  
 1480 isotope cycling, thereby connecting  $\delta^{138}\text{Ba}$  to (paleo)productivity.

1481  
 1482 **8.2. Driving processes**

1483 Seawater is undersaturated with respect to  $\text{BaSO}_4$  almost everywhere (Monnin et al., 1999; Rushdi et al.,  
 1484 2000; Monnin & Cividini, 2006). However, micro-crystalline  $\text{BaSO}_4$  is ubiquitous in the ocean. This ‘barite

1485 paradox' remains an area of active research. Proposed driving mechanisms broadly fall into two categories:  
1486 'active' biological and 'passive' chemical precipitation.

1487

### 1488 *8.2.1. Biological*

1489 Several organisms are known to precipitate BaSO<sub>4</sub> intracellularly, possibly for the purposes of gravitropism  
1490 (e.g., Gooday & Nott, 1982; Finlay et al., 1983). However, the organisms known to actively precipitate  
1491 BaSO<sub>4</sub> are not abundant in seawater, nor do they constitute a significant fraction of marine primary  
1492 productivity. Likewise, acantharea—organisms that precipitate SrSO<sub>4</sub> (celestite) tests that can contain  
1493 considerable quantities of Ba (e.g., Bernstein & Byrne, 2004)—are not necessary for driving significant  
1494 dissolved [Ba] drawdown in the ocean (Esser & Volpe, 2002) or for barite precipitation (Ganeshram et al.,  
1495 2003). Thus, existing evidence does not support a significant role for active biological processes in driving  
1496 marine Ba cycling (Fig. 19).

1497

### 1498 *8.2.2. Chemical*

1499 Passive chemical precipitation is likely the major contributor to particulate BaSO<sub>4</sub> stocks and  
1500 sedimentation. Given that seawater is largely undersaturated with respect to BaSO<sub>4</sub>, passive precipitation  
1501 is thought to occur within particle-associated microenvironments that are supersaturated with respect to  
1502 BaSO<sub>4</sub> (e.g., Chow & Goldberg, 1960; Dehairs et al., 1987). The development of BaSO<sub>4</sub>-supersaturated  
1503 microenvironments is hypothesized to relate to the heterotrophic oxidation of organic matter (Chow &  
1504 Goldberg, 1960), whereby Ba and sulfate ions are concentrated in chemically-isolated micro-zones during  
1505 bacterially-mediated mineralization of organic matter. Once sufficient quantities of Ba and sulfate ions have  
1506 accumulated and the microenvironment becomes supersaturated, BaSO<sub>4</sub> precipitation occurs. Thus, passive  
1507 precipitation of BaSO<sub>4</sub> is possible even in strongly undersaturated environments with low ambient sulfate  
1508 (e.g., Horner et al., 2017). Continued mineralization destroys the microenvironment, ceasing precipitation  
1509 and exposing BaSO<sub>4</sub> precipitates to undersaturated seawater where they begin to dissolve.

1510 The widespread association between pelagic BaSO<sub>4</sub> and aggregates of decaying organic matter provides  
1511 indirect support for this process (Dehairs et al., 1980; Bishop, 1988). Indeed, the peak in particulate Ba—  
1512 and presumably BaSO<sub>4</sub>—abundance is found in the upper mesopelagic zone where most organic matter  
1513 mineralization occurs (e.g., Sternberg et al., 2008). Microscale mechanisms remain unresolved, however;  
1514 recent studies indicate that biofilms may play an important role accumulating Ba (e.g., Martinez-Ruiz et

1515 al., 2018; 2019), which can promote precipitation of BaSO<sub>4</sub> nanoparticles from undersaturated solutions  
1516 (Deng et al., 2019).

1517 Regardless of the precise microscale mechanism, precipitation of particulate BaSO<sub>4</sub> is ubiquitous in the  
1518 marine realm. Given that BaSO<sub>4</sub> precipitation renders a substantial negative isotope fractionation of  $\approx 0.5$   
1519 ‰, it is highly likely that BaSO<sub>4</sub> cycling drives much of the Ba isotope variability in the ocean (Fig. 19).  
1520 From a paleoproxy perspective, this is ideal; BaSO<sub>4</sub> formation is related to the decay of organic matter and  
1521 not by the presence of any specific organism (e.g., Jacquet et al., 2007; Dehairs et al., 2008). Downward  
1522 transport of particulate BaSO<sub>4</sub> is driven by aggregation with larger particles (Lam & Marchal, 2014). The  
1523 efficiency of this downward transport depends on the same biophysical processes that export organic matter,  
1524 thus connecting the export flux of BaSO<sub>4</sub> to that of organic carbon (Fig. 19).

1525 Barites formed in the ocean through this passive chemical pathway are commonly termed marine, pelagic,  
1526 authigenic, or biogenic. Though none of these terms are perfect descriptors of the chemical processes  
1527 involved, ‘marine’ and ‘biogenic’ are the most ambiguous and their use is discouraged; the former  
1528 encompasses all BaSO<sub>4</sub> formed in the marine realm—including diagenetic, cold seep, and hydrothermal—  
1529 whereas the latter could be taken to imply only those precipitates brought about by active biological  
1530 processes. While ‘authigenic’ is an informative descriptor, it has also been used to describe sedimentary  
1531 BaSO<sub>4</sub> that formed via diagenetic redistribution on or below the seafloor (e.g., Torres et al., 1996). Thus,  
1532 we recommend use of the term ‘pelagic’ when describing chemically-precipitated microcrystalline BaSO<sub>4</sub>,  
1533 and encourage authors to include this definition in their works.

1534

### 1535 *8.2.3. Physical*

1536 Barium exhibits a nutrient-like profile in the oceans, similar to that of Si (silicic acid) and alkalinity (Fig.  
1537 18). However, the extent to which this pattern arises from non-conservative biogeochemical processes  
1538 versus physical mixing remains unresolved. Results from the GEOTRACES program are facilitating  
1539 renewed interest into this topic, which is being investigated using two complementary approaches. In the  
1540 first, biogeochemical contributions to basin-scale Ba distributions are isolated using statistical methods,  
1541 such as OMPA (Optimum Multiparameter water mass Analysis). These statistical methods showed that  
1542 mixing is dominant in the Mediterranean (Jullion et al., 2017) and North Atlantic (Le Roy et al., 2018), but  
1543 that sea ice-related processes may be important in the Arctic (Hendry et al., 2018). Second, the influence  
1544 of mixing is evidenced from emerging Ba stable isotope data. Indeed, the importance of mixing has been  
1545 documented vertically (Horner et al., 2015a), zonally (Bates et al., 2017; Bridgestock et al., 2018), and

1546 meridionally (Bates et al., 2017; Hsieh & Henderson, 2017). Together, these new approaches imply that *in*  
1547 *situ* biogeochemical processes exert a relatively minor influence on basin-scale Ba distributions.

1548

### 1549 **8.3. Marine archives**

1550 Given the connections between export production and BaSO<sub>4</sub> fluxes, the major archive of historical changes  
1551 in Ba cycling is BaSO<sub>4</sub> itself. Indeed, the sedimentary accumulation of BaSO<sub>4</sub>—most commonly  
1552 determined as the fraction of Ba in excess of the detrital Ba background or the deposition rate of BaSO<sub>4</sub>  
1553 itself—has been extensively used to reconstruct past changes in export production (e.g., Schmitz, 1987;  
1554 Francois et al., 1995; Paytan et al., 1996; Frank et al., 2000; Jaccard et al., 2005; Jaccard et al., 2013; Ma  
1555 et al., 2014; 2015, Costa et al. 2016, Winckler et al., 2016). An estimated 30 % of the BaSO<sub>4</sub> microcrystals  
1556 formed in seawater are buried in oxygenated sediments (e.g. Dymond et al., 1992), a considerably higher  
1557 fraction than for organic carbon (Paytan & Kastner, 1996). However, in oligotrophic regimes where both  
1558 BaSO<sub>4</sub> fluxes and sedimentation rates are low, prolonged exposure to undersaturated seawater results in  
1559 poor preservation (Eagle et al., 2003; Serno et al., 2014). Similarly, in high-productivity coastal upwelling  
1560 settings, sedimentary mineralization of organic matter consumes oxygen present in pore waters, driving  
1561 conditions down the redox tower and toward sulfate reduction, hampering BaSO<sub>4</sub> preservation (McManus  
1562 et al., 1998; Paytan & Griffith, 2007; Carter et al., 2020).

1563 Recently, the isotopic composition of Ba in bulk sediments has been investigated as a proxy for the Ba  
1564 isotope composition of the Ba source (i.e., dissolved Ba in epipelagic and upper-mesopelagic seawater).  
1565 Applications to date have explored the recovery of the biological carbon pump following the PETM (~56  
1566 Ma; Bridgestock et al., 2019) and the origin of enigmatic sedimentary BaSO<sub>4</sub> deposited in the aftermath of  
1567 the Marinoan glaciation (~635 Ma; Crockford et al., 2019) and Great Oxidation Event (~2,000 Ma;  
1568 Hodgskiss et al., 2019). Proceedings from recent geochemistry conferences indicate that many more  
1569 investigations are underway.

1570 Lastly, the amount—and, more recently the isotopic composition (e.g., Hemsing et al., 2018; Geyman et  
1571 al., 2019; Liu et al., 2019)—of Ba in marine carbonates has been extensively used to reconstruct the Ba  
1572 content of past environments. These studies are not discussed here as they only indirectly relate to  
1573 productivity, and instead pertain primarily to reconstructing ocean circulation and/or terrestrial runoff.

1574



1575 **8.4. Prospects**

1576 Barium exhibits several nutrient-like properties. Dissolved Ba distributions resemble those of other  
 1577 nutrients and particulate abundances are intimately associated with the processes of organic carbon  
 1578 remineralization and export. Despite these connections, several aspects of Ba cycling—both in the modern  
 1579 and past oceans—remain unresolved. We suggest several areas for additional research that will help widen  
 1580 the applicability of Ba-based proxies in paleoceanography.

1581

1582 *8.4.1. Modern*

1583 Several questions persist regarding the modern Ba cycle. Below we list three and offer possible remedies  
 1584 to each.

- 1585 • To what extent do the similar distributions in [Ba], [Si], alkalinity, and [<sup>226</sup>Ra] reflect true  
 1586 biogeochemical coupling versus passive physical mixing? As noted above, statistical analysis of  
 1587 Ba (and Si, Ra) distributions in regions will offer valuable insights, particularly if conducted in  
 1588 regions with weaker overturning circulation. Likewise, additional profiles of dissolved Ba isotope  
 1589 distributions will enable quantification of mixing relationships.
- 1590 • Closing the Ba-isotopic mass balance of the ocean. This will require detailed evaluation of other  
 1591 putative Ba sources, such as hydrothermal vents, cold seeps and other benthic sources (e.g.,  
 1592 Hoppema et al., 2010), atmospheric deposition, and the importance of estuarine enhancement of  
 1593 riverine Ba fluxes (e.g., Hanor & Chan, 1977; Edmond et al., 1978) as well as Ba fractionation  
 1594 associated with other phases like the adsorption on Fe-Mn oxyhydroxides.
- 1595 • What is the mechanism of BaSO<sub>4</sub> precipitation? While the microenvironment-mediated model  
 1596 appears most likely, the (bio)chemical mechanisms of precipitation remain ambiguous: How and  
 1597 from what are Ba and sulfate ions liberated during bacterial mineralization? How are they  
 1598 accumulated? Do different substrate organisms and/or heterotrophic communities influence the  
 1599 amount of BaSO<sub>4</sub> precipitated? Addressing these questions will require additional field and  
 1600 laboratory studies, which can then be compared against distributions of particulate BaSO<sub>4</sub> in the  
 1601 ocean interior. Depending on their importance, these nuances may require ecological  
 1602 parameterizations in numerical models of Ba cycling.
- 1603 • Narrowing estimates of the fractionation factor between BaSO<sub>4</sub> and seawater. Existing laboratory  
 1604 studies place this estimate  $\approx -0.3$  ‰, whereas a wide-range of field data suggest that it is  
 1605 considerably larger at  $\approx -0.5$  ‰. Accounting for this  $\approx 0.2$  ‰ difference is both important and

1606 justifies additional experimentation, as it indicates incomplete understanding of the processes that  
 1607 fractionate Ba isotopes in the marine realm.

1608

1609 *8.4.2. Paleo*

1610 As with the modern cycle, several ambiguities persist. Assuming that BaSO<sub>4</sub> remains the preferred archive  
 1611 of past Ba cycle variations, the most pressing relates to diagenesis.


1612 • What is the effect of early diagenesis on the Ba isotope composition of sedimentary BaSO<sub>4</sub>?  
 1613 Likewise, do diagenetic BaSO<sub>4</sub> retain any primary Ba isotope information? Assessing these  
 1614 questions will require studies of co-located BaSO<sub>4</sub> and porewaters from environments at various  
 1615 stages of early diagenesis. Answering these questions is critical in establishing the validity of Ba  
 1616 isotopes in barite as a paleoceanographic proxy.

1617 • Are BaSO<sub>4</sub> accumulation rates impacted by ambient [Ba]? Related to the question concerning  
 1618 BaSO<sub>4</sub> precipitation mechanisms above, it is presently unknown if the quantity of BaSO<sub>4</sub> yielded  
 1619 by remineralization is impacted by ambient [Ba]—does more BaSO<sub>4</sub> precipitation occur at higher  
 1620 ambient [Ba]? Likewise, to what extent does BaSO<sub>4</sub> preservation depend on ambient [Ba]? These  
 1621 considerations are significant when considering longer-term records, particularly when marine  
 1622 sulfate levels were lower-than-modern (and [Ba] presumably higher; e.g., Hanor, 2000; Wei &  
 1623 Algeo, in press). Finally, does the seawater temperature at the depth of POC mineralization impact  
 1624 the relationship between POC and barite formation? These questions are best addressed through a  
 1625 combination of experimentation (e.g., cultures, precipitation), field studies in basins with large  
 1626 gradients in [Ba], and numerical experiments incorporating saturation state modeling.

1627 **9. Nickel**

1628 **9.1. Marine distribution**

1629 Nickel has a classic nutrient-type distribution in the oceans with one unusual feature; surface dissolved Ni  
 1630 concentrations never drop below around 1.8–2 nmol kg<sup>-1</sup> (e.g., Sclater et al., 1976; Bruland, 1980; Fig. 20).  
 1631 There is evidence, however, that this residual pool in surface waters is not bioavailable, as summarized  
 1632 recently by Archer et al. (2020). Deep water dissolved Ni concentrations are 4–5 nmol kg<sup>-1</sup> in the Atlantic,  
 1633 and ~9 nmol kg<sup>-1</sup> in the north Pacific (Fig. 20).




1634 **Figure 20 | Representative profiles of dissolved Ni concentrations ([Ni]) and Ni-isotopic compositions (δ<sup>60</sup>Ni)**  
 1635 **from the South Atlantic (squares, dashed line), Southern (triangles, dotted line), and Subtropical South Pacific**  
 1636 **(circles, solid line) Oceans. This comparison illustrates that the oceanographic processes leading to distinct dissolved**  
 1637 **concentration profiles render only small changes in dissolved δ<sup>60</sup>Ni between basins.**  
 1638

1639  
 1640 Developing robust analytical protocols for analyzing Ni isotopes for a range of sample matrices has proven  
 1641 somewhat more challenging than for some of the other TEIs discussed here due to the difficulty in  
 1642 separating Ni from interfering elements (e.g., Fe, a major isobaric interference on <sup>58</sup>Ni). Chemical  
 1643 purification protocols now use a sequence of resins including anion exchange (e.g., AG MP-1M or AG1-

1644 X8, Bio-Rad) and either a Ni-specific resin (containing dimethylglyoxime) or Nobias PA-1 (Hitachi High  
 1645 Technologies; e.g., Cameron et al., 2009; Gueguen et al., 2013; Wang et al., 2019a; Yang et al., 2020). The  
 1646 limited data subsequently reported for the oceanic dissolved pool of Ni is isotopically homogeneous at  
 1647 depth ( $\delta^{60}\text{Ni} \approx +1.3 \text{ ‰}$ ), with a small shift towards heavier values (up to +1.7 ‰) in the upper water column  
 1648 (Takano et al., 2017; Wang et al., 2019a; Yang et al., 2020; Archer et al., 2020). The latter is proposed to  
 1649 reflect a small kinetic isotope fractionation during biological uptake. Similar to Zn, Cu, Cd, Mo, and Ba,  
 1650 the isotopic composition of Ni in seawater is isotopically heavy compared to the UCC (Table 1).

1651



1652 **Figure 21 | Processes driving nickel isotope variations in modern seawater.** Biological processes are recognized  
 1653 to drive a small, but systematic increase in dissolved  $\delta^{60}\text{Ni}$  and marine Ni isotope values are preserved in certain  
 1654 sediments. Other abiotic Ni cycling processes may be important however, such as sorption to Mn-oxide minerals, and  
 1655 the significance of these fractionations to global Ni cycling remains to be fully elucidated.  
 1656

1657

1658

1659

1660

1661 **9.2. Driving processes**

1662 *9.2.1 Biological*

1663 To date, eight Ni-based enzyme systems have been identified (Ragsdale, 2009), including urease, which is  
 1664 key to the global nitrogen cycle, and methyl-CoM reductase, which catalyses the production of all  
 1665 biologically generated methane on Earth. The obligate requirement of methanogens for Ni has led to interest  
 1666 in developing Ni and Ni isotopes as a tracer of methane production on the early Earth (e.g. Cameron et al.,  
 1667 2009; Konhauser et al., 2009; Wang et al., 2019c). In the modern ocean the highest Ni quotas are found in  
 1668 diazotrophs (N-fixers), thought to reflect the presence of Ni-Fe hydrogenases (which catalyse H<sub>2</sub> produced  
 1669 during N fixation), Ni-superoxide dismutase (Ni-SOD) and urease (Tamagnini et al., 2002; Dupont et al.,  
 1670 2008a; 2008b; Nuester et al., 2012). Nickel limitation of phytoplankton grown on urea has been shown in  
 1671 culture and in natural assemblages, suggesting the Ni-N co-limitation of phytoplankton growth may be  
 1672 relevant in the ocean (Price & Morel, 1991; Dupont et al. 2008a; 2010). Significant Ni is also found in  
 1673 diatom frustules (about 50 % of diatom cellular quotas; Twining et al., 2012). The latter observation is  
 1674 thought to play a role in the similarity of Ni and Si oceanic distributions (Twining et al., 2012).

1675 No culture data are available to determine the degree of Ni isotope fractionation during biological uptake.  
 1676 Upper ocean data suggest no fractionation or a small preference for the light isotope (up to  $\alpha \approx 0.9997$ ;  
 1677 Takano et al., 2017; Archer et al., 2020; Yang et al., 2020; Fig. 21), consistent with the Ni isotope  
 1678 systematics observed in organic-rich sediments (Ciscato et al., 2018) and water column particulates (Takano  
 1679 et al., 2020). Interestingly, new water column data from the South Atlantic suggest distinct ecological  
 1680 differences in Ni drawdown and Ni isotope fractionation compared to other TEIs (e.g., Zn, Cd). Limited  
 1681 drawdown and Ni isotope fractionation is observed in the diatom-dominated regime south of the Polar Front  
 1682 in the Southern Ocean. In contrast, more marked drawdown and significant Ni isotope fractionation is  
 1683 observed north of the Polar Front, which has been attributed to the predominance of nitrate-limited, Ni-  
 1684 requiring cyanobacteria (Archer et al., 2020).

1685

1686 *9.2.2 Chemical*

1687 Nickel is partially complexed by strong organic ligands in coastal and open ocean environments (5–70 %;  
 1688 e.g., Donat et al., 1994; Saito et al., 2004; Boiteau et al., 2016), though slow water exchange kinetics of Ni  
 1689 make these complexation measurements particularly challenging (Hudson & Morel, 1993). Slow exchange  
 1690 kinetics may also explain the residual pool of non-bioavailable Ni in the surface ocean (e.g., Mackay et al.,

1691 2002; Dupont et al., 2010). Speciation models suggest that the remainder of the Ni is present as free  $\text{Ni}^{2+}$   
 1692 and  $\text{NiCl}^+$ , with a small fraction present as  $\text{NiCO}_3^0$  (Zirino and Yamamoto, 1972; Turner et al., 1981).

1693 Nickel cycling is tightly coupled to that of Mn, in sediments, pore waters, and across the redoxcline of the  
 1694 Black Sea (e.g., Klinkhammer, 1980; Koschinsky and Hein, 2003; Vance et al., 2016). In the Black Sea,  
 1695 for example, Mn redox cycling is associated with preferential sorption of light Ni isotopes, with a large  
 1696 negative fractionation of  $\approx -4\text{‰}$  (Vance et al., 2016). This large fractionation is consistent with experimental  
 1697 sorption of Ni on birnessite (Wasylenki et al., 2014; Sorensen et al., 2020).

1698 Unlike more strongly chalcophile elements like Cd, Cu, and Zn, dissolved Ni is not strongly drawn down  
 1699 in the euxinic portion of the Black Sea water column (Tankéré et al., 2001; Vance et al., 2016). However,  
 1700 Ni is enriched in Black Sea sediments (Little et al., 2015), with  $\delta^{60}\text{Ni}$  compositions notably lighter (at +0.3  
 1701 to +0.6‰) than Ni sources to the basin (at about +1.3‰; Vance et al., 2016; Fig. 21). Vance et al. (2016)  
 1702 attributed these light isotope compositions to the scavenging of sulfidized Ni species, which are predicted  
 1703 to be isotopically light (Fujii et al., 2011b).

1704

### 1705 *9.2.3 Physical*

1706 The most recent estimate for residence time of Ni in the ocean is approximately  $\sim 20$  kyr (Little et al., 2020),  
 1707 considerably longer than the mixing time of the ocean. As a result, in parallel with the other TEIs discussed  
 1708 herein, Ni and Ni isotope distributions are modulated at first order by the geometry of physical ocean  
 1709 circulation. The importance of diatom uptake in the Southern Ocean in partially coupling oceanic Ni and  
 1710 Si (Twining et al., 2012) was introduced above, and the homogeneity of deep ocean Ni isotope compositions  
 1711 supports an important role for southern-sourced water masses in the Ni distribution (Takano et al., 2017;  
 1712 Wang et al., 2019a; Archer et al., 2020).

1713


## 1714 **9.3. Marine archives**

### 1715 *9.3.1. Ferromanganese sediments*

1716 Ferromanganese crusts exhibit variable Ni isotope compositions (Fig. 22; Gall et al., 2013; Gueguen et al.,  
 1717 2016), which are, on average, slightly isotopically heavier (at  $\delta^{60}\text{Ni}$  of +1.6‰) than seawater ( $\delta^{60}\text{Ni}$  at about  
 1718 +1.3‰). However, experiments suggest that sorption of Ni to birnessite (the primary Ni-hosting phase in  
 1719 Fe–Mn sediments) should be associated with a large negative isotope effect (of about 3 to 4 ‰; Wasylenki



1720 et al., 2014; Sorensen et al., 2020). It remains unclear why the full isotope effect is not expressed in Fe–Mn  
 1721 crusts. Intense Mn cycling occurs across the redoxcline of the Black Sea; this cycling is associated with  
 1722 large variations in Ni and Ni isotopes, consistent with the experimentally observed light isotope effect on  
 1723 sorption to birnessite (Vance et al., 2016). Recent data from Mn-rich sediments that have undergone  
 1724 diagenesis also point to the preservation of a large negative Ni isotope effect (Little et al., 2020).



1725  
 1726 **Figure 22 | Four Fe–Mn records of Ni isotope compositions from the Pacific Ocean.** Assuming that Fe–Mn crusts  
 1727 have always formed with an offset  $\approx +0.5$  ‰ with respect to ambient dissolved Ni in seawater, these records illustrate  
 1728 that the Ni isotope composition of seawater has varied by only  $\pm 0.2$  ‰ over the Cenozoic. Data for J2-480, J2-480-  
 1729 R14, and ZEP2-DR06-03 are from Gueguen et al. (2016). Data from CD29-2 are from Gall et al. (2013). All four records  
 1730 have been plotted using the authors' preferred age model, meaning that there are some differences between the  
 1731 chronology of CD29-2 shown here compared to Figs. 5, 14.




1732 Though promising, the development of Fe–Mn crusts as a tracer of past oceanic Ni cycling awaits a  
1733 mechanistic understanding of the processes driving their variably isotopically heavy signature, as well as  
1734 an awareness of the likely complicating role of diagenetic remobilisation of Ni (e.g., Atkins et al., 2016;  
1735 Little et al., 2020).

1736

### 1737 9.3.2. *Organic-rich sediments*

1738 As introduced in section 5.3.2 (Cu), qualitative arguments for high organic matter fluxes (i.e., increased  
1739 paleoproductivity) have been made based on elevated Cu and Ni concentrations in ancient organic-rich  
1740 sediments (e.g., Tribovillard et al., 2006). For Ni, this approach is supported by positive correlations with  
1741 TOC in modern continental margin sediments (Fig. 23).

1742 Nickel does not precipitate in the presence of water column dissolved sulfide and is therefore a better  
1743 candidate for the Metal–TOC approach than the more strongly chalcophile elements, like Cu. Nevertheless,  
1744 Ni cycling is strongly linked to the redox cycling of Mn, so prerequisite (2)—that depositional environments  
1745 remain reducing—is critical. In addition, sedimentary and water column data from the Black Sea indicate  
1746 that the Fe–Mn redox shuttle provides an alternative supply route for Ni to sediments in these settings (Little  
1747 et al., 2015; Vance et al., 2016). Finally, an open marine setting (prerequisite 3) would be required to make  
1748 estimates of the relative productivity of two different sites based on their absolute measured Ni:TOC ratios.  
1749 Otherwise the degree of basin restriction will exert the primary control on nutrient supply, and therefore  
1750 the degree of trace metal enrichment (Algeo & Maynard, 2008; Little et al., 2015).



1751  
 1752 **Figure 23 | Correlation of Ni and TOC content in organic-rich sediments from the Peru Margin.** Data from Ciscato  
 1753 et al. (2018). The best-fit regression of these data yields a Ni–TOC slope of  $9.6 \pm 1.1$  (mean  $\pm$  2 SD), similar to the “ $\approx 9$ ”  
 1754 reported by Böning et al. (2015).

1755  
 1756 Ciscato et al. (2018) investigated the distribution of Ni and its isotopes in two fractions isolated from Peru  
 1757 margin organic-rich sediments. The HF-HCl digestible fraction (usually containing >80% of total Ni)  
 1758 exhibited  $\delta^{60}\text{Ni}$  values similar to modern deep seawater (at about +1.2‰). Meanwhile, these authors found  
 1759 variable Ni isotope compositions in the organic-pyrite fraction (OPF), which they suggested record the  
 1760 fractionation imparted by biological uptake in the photic zone. Systematic relationships between Ni–TOC,

1761  $\delta^{60}\text{Ni}_{\text{OPF}}$ , and  $\delta^{13}\text{C}$  indicate that there is merit in continuing to investigate Ni and Ni isotopes as a  
1762 paleoproductivity tracer (Ciscato et al., 2018).

1763

#### 1764 **9.4. Prospects**

1765 To date, Ni and its isotopes have been under-developed as a potential paleoproductivity proxy. Recent data,  
1766 both from the dissolved phase in seawater and in two different fractions isolated from anoxic organic-rich  
1767 sediments, suggest promise in the coupling of Ni and C and their isotopes. However, in oxic-to-suboxic  
1768 settings, Ni and its isotopes in sediments are strongly influenced by the diagenetic redox cycling of Mn. In  
1769 these settings, we recommend exploratory Ni and  $\delta^{60}\text{Ni}$  work focusing on carbonate and siliceous  
1770 biominerals, work that is increasingly tractable with new and improved chemical separation and analytical  
1771 techniques.


1772 Finally, in a completely different approach, Wang et al. (2019c) present  $\delta^{60}\text{Ni}$  values of Precambrian glacial  
1773 diamictites, which are suggested to represent the chemical weathering residues of the UCC. They find a  
1774 small shift towards heavier Ni isotope compositions across the Great Oxidation Event, which they relate to  
1775 the onset of oxidative weathering of crustal sulfides (Wang et al., 2019c). Combined with the proposed  
1776 importance of Ni to the maintenance of methanogenesis during this time period (e.g. Konhauser et al.,  
1777 2009), it is hoped that future Ni stable isotope analyses will shed further light on the paleoenvironmental  
1778 conditions on the early Earth.

1779

1780 **10. Silver**

1781 **10.1. Marine distribution**

1782 Silver is the scarcest of the TEIs described here, both within Earth’s crust and in the modern ocean.  
 1783 Moreover, Ag is an usual candidate for a paleoproductivity proxy given that it possesses no known  
 1784 biological function and is most widely known for its antimicrobial properties. Despite this, Ag exhibits a  
 1785 characteristic nutrient-like profile in seawater, most similar to that of Si (and Zn, Ba; Fig. 24).




1786 **Figure 24 | Representative profiles of dissolved Ag concentrations ([Ag]) from the Northeast Pacific (circles,**  
 1787 **solid line) and Southern (triangles, dotted line) Ocean. No Ag-isotopic data for seawater are available at this time.**  
 1788

1789  
 1790 Dissolved Ag occurs primarily as chloride complexes in seawater (Cowan et al., 1985, Miller & Bruland,  
 1791 1995) and exhibits a nutrient-type depth profile. Dissolved [Ag] in surface waters is typically <5 pmol kg<sup>-1</sup>,  
 1792 and ranges from 5–30 pmol kg<sup>-1</sup> in deep waters of the Atlantic Ocean to 50–114 pmol kg<sup>-1</sup> in the Pacific

1793 Ocean (Fig. 24; Flegal et al., 1995; Rivera-Duarte et al., 1999; Ndung'u et al., 2001; Zhang et al., 2001;  
 1794 2004; Ranville and Flegal, 2005; Boye et al., 2012). The total observed range in seawater is 0.2–115 pmol  
 1795 kg<sup>-1</sup> (Gallon and Flegal, 2014). Despite its potential as a biogeochemical proxy, no Ag isotope data yet  
 1796 exist for dissolved or particulate phases in the water column, nor in marine sedimentary archives.  
 1797 Accordingly, we cannot directly assess the utility of Ag isotopes to infer paleoproductivity, though we can  
 1798 deduce a number of processes that are likely to influence dissolved  $\delta^{109}\text{Ag}$  based on the processes known  
 1799 to cycle Ag (Fig. 25).

1800



1801  
 1802 **Figure 25 | Processes likely to drive silver isotope variations in modern seawater.** Despite there being no Ag  
 1803 isotope data for seawater at this time, we can infer a number of processes that are most likely to influence dissolved  
 1804  $\delta^{109}\text{Ag}$  from measurements of [Ag] in seawater and sediments.

1805

1806

1807 **10.2. Driving processes**

1808 The 1D depth profile for dissolved Ag is similar to that of Si (and Zn, Ba), resulting in a strong positive  
 1809 correlation between these elements (Fig. 1). This has led researchers to suggest that Ag is taken up by  
 1810 diatoms, incorporated into their frustules, and then released as the frustules dissolve (e.g., Flegal et al.,

1811 1995). Silver might then be delivered to the seafloor with opal, potentially making it useful as a  
1812 paleoproductivity proxy (Friedl & Pedersen, 2002). However, the correlation between dissolved Ag and Si  
1813 is non-linear, indicating that other factors are at play (Zhang et al., 2001; 2004). Furthermore, while Ag is  
1814 taken up by various types of phytoplankton, including diatoms (Fisher & Wentz, 1993), experiments  
1815 conducted using the marine diatom *Thalassiosira pseudonana* show that most of the Ag is associated with  
1816 the organic fraction rather than the opal (Wagner, 2013; Fig. 25). Martin et al. (1983) also hypothesized  
1817 that high particulate Ag concentration within the euphotic zone (40–70 m) off the west coast of Mexico  
1818 were due to the formation of Ag-organic complexes. Interestingly, particulate Ag concentrations are even  
1819 higher well below the euphotic zone, at a depth corresponding to the upper portion of the oxygen minimum  
1820 zone (Martin et al., 1983). It could also be that a global Ag:Si correlation arises at least in part from  
1821 biological processes occurring in the surface of the Southern Ocean, whereby intermediate and mode water  
1822 masses with low preformed [Ag] are advected to lower latitudes, analogous to the mechanism proposed for  
1823 Zn–Si (Vance et al., 2017) and Ba–Si coupling (Horner et al., 2015a). If correct, Ag is unlikely to be coupled  
1824 directly to opal with a simple relationship that can be used to reconstruct past diatom productivity.

1825

1826 Dissolved Ag cycling may become decoupled from productivity by possessing different source and sink  
1827 terms relative to carbon and the macronutrients (Gallon and Flegal, 2014 and references therein). For  
1828 example, Ranville & Flegal (2005) and Ranville et al. (2010) invoked an anthropogenic aerosol source of  
1829 Ag to surface and intermediate waters to explain north Pacific water column data. Other complications may  
1830 arise in low-oxygen environments—waters from the northeast Pacific (Kramer et al., 2011) and  
1831 southeastern Atlantic (Boye et al., 2012) exhibit a deficit in dissolved Ag relative to Si within their  
1832 respective OMZs. These deficits imply preferential removal of Ag over Si, which may occur locally or  
1833 ‘upstream’. If occurring locally, a putative mechanism is (co-)precipitation with other chalcophile elements,  
1834 analogous to the sulfide-mediated mechanism proposed for Cd in OMZs (Janssen et al., 2014; Sec. 6).  
1835 Alternatively, the deficit may reflect low preformed Ag:Si in intermediate waters, which is inherited from  
1836 preferential drawdown of Ag over Si in regions upstream where these intermediate waters were last  
1837 ventilated. Both interpretations have implications for the use of Ag (isotopes) as a paleoproxy: the former  
1838 implies a redox sensitivity that depends on the changing location, spatial extent, and intensity of low-oxygen  
1839 regions in the oceans over time. The latter implies a sensitivity to ecology and the geometry of ocean  
1840 circulation. Both warrant additional scrutiny.

1841


### 1842 **10.3. Marine archives**

1843 Bulk sediments are the main archive that has been investigated for their potential to record information  
1844 about the marine Ag cycle. Silver concentrations in bulk sediments from open-marine environments range  
1845 from  $\leq 100 \text{ ng g}^{-1}$  (i.e., typical lithogenic values) up to 100s of  $\text{ng g}^{-1}$  (Koide et al., 1986). In general, low  
1846 concentrations are typical of well-oxygenated sediments, while higher concentrations are found in oxygen-  
1847 poor and sulfidic environments. The general consensus has been that Ag enrichment in anoxic sediments is  
1848 the result of post-depositional precipitation of  $\text{Ag}_2\text{S}$  (Koide et al., 1986) or possibly Ag selenide (Crusius  
1849 & Thomson, 2003; Böning et al., 2005). However, high Ag concentrations are also documented in marine  
1850 sediments that are only weakly reducing (Böning et al., 2004, 2005; McKay & Pedersen, 2008; Morford et  
1851 al., 2008). Furthermore, even in anoxic sediments, the degree of Ag enrichment exceeds what would be  
1852 expected from post-depositional Ag precipitation alone (Borchers et al., 2005; McKay and Pedersen, 2008;  
1853 Böning et al., 2009). Thus, redox-controlled, post-depositional precipitation is not the primary control on  
1854 Ag accumulation in marine sediments; there must also be a flux of non-lithogenic, particulate Ag to the  
1855 seafloor (McKay & Pedersen, 2008).

1856 McKay & Pedersen (2008) hypothesized that Ag, like Ba, accumulates in organic-rich settling particles.  
1857 However, in contrast to Ba, Ag precipitation requires a reduced microenvironment within the particle,  
1858 which is only favored in oxygen-poor waters. The analysis of sediment trap samples from the northeast  
1859 Pacific show that the fluxes of particulate Ag and particulate organic carbon flux positively correlate  
1860 (Martin et al., 1983). This correlation is also seen in surface sediments (McKay & Pedersen, 2008) and  
1861 sediment cores from the northeastern Pacific (Fig. 26). These data broadly support the use of Ag  
1862 concentrations as a paleoproductivity proxy, with an important caveat: the post-depositional preservation  
1863 of particulate Ag requires that sediments remain reducing (prerequisite 2.), as settling particulate Ag formed  
1864 below the euphotic zone is not preserved if sediments are oxidizing (McKay and Pedersen, 2008; Morford  
1865 et al., 2008). Thus, while the delivery of Ag to sediments appears related to productivity (Wagner et al.,  
1866 2013), Ag preservation will likely depend on the depositional environment. Regardless, these observations  
1867 are promising from the point of view of tracing past productivity.

1868





1869

1870 **Figure 26 | Concentrations of silver and total organic carbon in sediment cores from the northeast Pacific**  
 1871 **Ocean.** These data illustrate that Ag is broadly correlated with organic matter in bulk sediments, which is highly  
 1872 promising from the perspective of developing Ag as a paleoproductivity proxy.

1873

1874

1875 **10.4. Prospects**

1876 The apparent linkages between dissolved and particulate [Ag], macronutrients, and organic matter provides  
 1877 tantalizing evidence that Ag may be coupled to surface productivity (Fig. 24). Moreover, core-top studies  
 1878 indicate that the geochemical signature of this coupling is preserved under certain environmental conditions

1879 (Fig. 26). Despite this progress, the study of Ag in marine biogeochemical cycles remains in its infancy,  
1880 particularly compared to many of the other elements described here. Additional constraints are needed in  
1881 several areas, including: the role of biogeochemical processes in mediating Ag distributions in the water  
1882 column, the dominant controls on the downward transport of Ag through the oceans, and on the controls on  
1883 Ag preservation in sediments. Given what has been learned from the application of the other TEI systems  
1884 described here, new analytical developments in Ag isotope geochemistry could help place valuable  
1885 constraints in several of these areas.

1886

1887

## 1888 **11. Synthesis**

1889 We now assess the overall readiness of each TEI system to reconstruct productivity, summarized in Table  
1890 2. This exercise is analogous to the assessment of analytical techniques used in chemical oceanography  
1891 described by Fassbender et al. (2017). Our assessment is similarly conducted in two dimensions. First, we  
1892 identify five objectives toward the development of a reliable productivity proxy, ranging from development  
1893 of the analytical capabilities necessary to measure that species, to constraining diagenetic effects, and  
1894 ending with the goal of using that species to reliably reconstruct paleoproductivity itself. Second, we assess  
1895 the level of development within each objective. Our reasoning behind the assignments is described above  
1896 in Sections 3–10.

1897

1898 Reliable application of a proxy requires that five objectives be serially met (e.g., Hillaire-Marcel & Vernal,  
1899 2007; Table 2). In practical terms, however, the final stage (proxy application) is often realized before many  
1900 of the supporting objectives; variations in TEI ratios in the sedimentary record commonly provide the  
1901 motivation for developing a more holistic understanding of that TEI system in the modern environment. (A  
1902 common critique of this approach being that subsequent studies often invalidate earlier interpretations.) The  
1903 five objectives that follow are common to the development of almost any proxy. First, it is essential to  
1904 develop the analytical capability to measure the species of interest commensurate with the quantities  
1905 typically encountered in the environment. In the case of TEI abundance proxies, the development and  
1906 widespread adoption of ICP-MS instrumentation coupled to automated sample preparation systems has  
1907 enabled low-blank–high-throughput–high-sensitivity analyses of multiple TE's in both seawater (e.g.,  
1908 Wuttig et al., 2019) and sediments alike (e.g., Wefing et al., 2017). In contrast, the techniques required to  
1909 measure many TEI systems have only been developed within the past decade—or are still in development—  
1910 and generally remain labor intensive and time consuming. Second, it is important to map the broad vertical  
1911 and spatial patterns of a TEI system in the modern ocean. The GEOTRACES program has provided a  
1912 coordinated opportunity to study the basin-scale distributions of multiple TEIs. Third, the utility of a proxy  
1913 is significantly increased if the driving processes are understood. These processes may be isolated through  
1914 a number of approaches, including: lab-based analogue experimentation, numerical modeling, and high-  
1915 resolution spatiotemporal environmental studies. Fourth, paleoceanographic proxies require sedimentary  
1916 archives—a substrate from which to reconstruct the variable of interest. The latter necessitates knowledge  
1917 of how a TEI system partitions between seawater and sediment, such as through a core-top study, ideally  
1918 conducted across large environmental gradients. Since many marine sedimentary archives are biogenic in  
1919 origin, additional experimentation isolating ‘vital effects’ may be necessary. Fifth, proxies are only as

1920 reliable as their archives are hardy. Diagenetic processes may alter primary environmental signatures, and  
 1921 recognizing these effects is imperative for reliably reconstructing past environmental conditions.

1922

1923 We assess the level of development within each objective on a four-point scale: unknown, developing,  
 1924 applied, and unlikely. The levels define a continuum from least to most understood, and are a useful  
 1925 shorthand for illustrating where additional work is most needed. Assignment of ‘unknown,’ implies that  
 1926 too little is presently known to reliably assess progress towards that objective. These TEI systems may or  
 1927 may not ultimately be useful in reconstructing ocean productivity. ‘Developing’ objectives are those where  
 1928 there are pilot studies on that topic, but overall there are an insufficient number to define general rules for  
 1929 that TEI system. If a TEI system is widely recognized to be useful towards some objective, it is given a  
 1930 score of ‘applied’. Though these scores are subjective, they are roughly equivalent to the number of  
 1931 spatiotemporal and lab studies of that TEI system, and the number of inter-calibrated laboratories. If the  
 1932 evidence indicates that a TEI system is not suitable for reconstruction of past productivity, a score of  
 1933 ‘unlikely’ is given. This does not rule out future developments, such as identification of environmental  
 1934 control variables or new sedimentary archives, only that current data (and archives) do not support use of  
 1935 the TEI system towards this goal. Lastly, we recognize that there are continual refinements to analytical  
 1936 protocols, environmental distributions, etc. and thus, at some level, all five objectives could reasonably be  
 1937 described as ‘developing’. Rather, our assignments are intended to give a relative sense of understanding  
 1938 between different TEI systems toward the overarching goal of reconstructing past ocean productivity.

1939

1940

1941

1942 **Table 2 | Proxy development assessment.** The progress toward five objectives is assessed for each TE proxy system  
 1943 and assigned a development level ranging unknown to unlikely, corresponding to the least and most certain  
 1944 assignments, respectively. Some TEI systems possess split designations. Definitions of each development level are  
 1945 described in Sec. 11. and discussed in detail for each TEI system in the corresponding TEI section.

Proxy	1. Analytical protocols?	2. Distribution in environment?	3. Driving process isolated?	4. Sedimentary archives?	5. Effect of diagenesis?	Goal: Productivity reconstruction?
[Fe] $\delta^{56}\text{Fe}$	Applied	Applied	Developing	Developing	Developing	Unknown
[Zn] $\delta^{66}\text{Zn}$	Applied	Applied	Developing	Developing	Developing	Unknown
[Cu] $\delta^{65}\text{Cu}$	Applied	Applied	Developing	Developing	Developing	Unknown
[Ni] $\delta^{60}\text{Ni}$	Applied	Applied	Developing	Developing	Developing	Unknown
[Cd] $\delta^{114}\text{Cd}$	Applied	Applied	Developing	Developing	Developing	Unknown
[Ba] $\delta^{138}\text{Ba}$	Applied	Applied	Developing	Developing	Unknown	Unknown
[Mo] $\delta^{98}\text{Mo}$	Applied	Applied	Applied	Developing	Unknown	Unknown
[Ag] $\delta^{109}\text{Ag}$	Developing	Unknown	Unknown	Unknown	Unknown	Unknown

**Key**

- Unknown
- Developing
- Applied
- Unlikely

1946  
 1947  
 1948

## 1949 **12. Conclusions and outlook**

1950

1951 This review and synthesis allows us to draw a number of inferences regarding the overall suitability,  
1952 similarities, and research priorities for using Fe, Zn, Cu, Cd, Mo, Ba, Ni, and Ag and their isotopes as  
1953 paleoproductivity proxies. Regarding suitability, our proxy development assessment highlights that our  
1954 knowledge of bioactive trace metal isotope systematics lags far behind those of trace-element abundances.  
1955 This field is nascent; excepting Mo, the very first isotope data for seawater for all of the elements reviewed  
1956 here were published within the last 15 year, or are yet to be reported, as is the case for Ag. The field is thus  
1957 decades behind the trace element concentration community. Despite the lack of detail however, we can  
1958 classify the elements reviewed here into three broad categories: those where the preponderance of evidence  
1959 indicates that a particular bioactive metal isotope system is unlikely to inform on paleoproductivity, such  
1960 as  $\delta^{56}\text{Fe}$ ,  $\delta^{65}\text{Cu}$ , and  $\delta^{98}\text{Mo}$ ; those where there are promising signs, but significant calibration and validation  
1961 remains to be completed, such as for  $\delta^{66}\text{Zn}$  and  $\delta^{114}\text{Cd}$ ; and, those elements where simply too little is  
1962 presently known to confidently assign utility in diagnosing paleoproductivity such as for  $\delta^{138}\text{Ba}$ ,  $\delta^{60}\text{Ni}$ , and  
1963  $\delta^{109}\text{Ag}$ .

1964

1965 Our review highlights a number of instances where the bioactive metals and their isotopes share similarities  
1966 or—perhaps more importantly—exhibit divergent behaviors, despite sharing a common biological driver.  
1967 The most significant similarity is ocean circulation; trace element isotope distributions reflect a mixture of  
1968 local (i.e., *in situ*) processes, and regional (or *ex situ*) conditions that are set ‘upstream’ of any given locality.  
1969 Indeed, researchers are recognizing that, much like the macronutrients (see Farmer et al., *this issue*), the  
1970 first-order features of marine  $\delta^{66}\text{Zn}$ ,  $\delta^{65}\text{Cu}$ ,  $\delta^{114}\text{Cd}$ ,  $\delta^{138}\text{Ba}$ , and  $\delta^{60}\text{Ni}$  distributions are not controlled locally  
1971 by dissolved–particulate transformations, but reflect a regionally-integrated history of vertical cycling and  
1972 mixing that is imparted over the scale of an ocean basin. Today, the Southern Ocean represents the common  
1973 starting point in controlling nutrient and bioactive metal inventories for much of the low latitude  
1974 thermocline; whether, when, and how this pattern might have changed in Earth’s past will be an important  
1975 consideration for interpreting paleoproductivity records derived from these proxies. Where the bioactive  
1976 metals diverge in their response is equally instructive from the point of view of proxy development. For  
1977 example, if experiencing a common forcing, such as a temporary shift in the loci of most intense primary  
1978 productivity, subtle differences in the response of the various bioactive metals reviewed here may hold the  
1979 key to understanding whether the underlying change was truly productivity related. For example,  
1980 sedimentary accumulation and preservation of Cd and Ba exhibit opposing redox sensitivities; Cd burial is  
1981 enhanced at low  $[\text{O}_2]$ , whereas Ba is oftentimes diminished. Recognizing these opposing controls may

1982 enable reconciling inconsistent proxy records, thus building a more holistic picture of marine  
1983 biogeochemistry at key points in Earth's history.

1984

1985 Lastly, we outline three priorities for further study. First, a number of modern ambiguities require  
1986 addressing. Most notable is the apparent 'missing' source and/or sink terms for the whole-ocean isotopic  
1987 budgets of several elements, including Cu, Cd, Ba, and Ni. Though it is possible that these missing source  
1988 and sink terms represent actual short-term imbalances in the flux of these elements to and from the ocean,  
1989 we view it more likely that there are major fluxes that remain uncharacterized. The origin of these  
1990 imbalances is most pressing for elements where the output flux associated with organic matter constitutes  
1991 a secondary sink (e.g., Zn, Ni), as the isotopic budget of these elements is most susceptible to decoupling  
1992 from productivity. Second, the significance of ocean circulation in mediating basin-scale bioactive metal  
1993 isotope distributions implies that paleoceanographic interpretations made from a single site cannot be  
1994 uniquely interpreted in terms of either changes in productivity or ocean circulation without additional  
1995 constraints. Such constraints could take the form of independent circulation estimates—from numerical  
1996 models or canonical circulation proxies measured in the same samples—or by measuring bioactive metal  
1997 isotope distributions in spatially-distributed sediment samples. Third, the lack of suitable archives with  
1998 which to reconstruct surface water bioactive metal isotope compositions afflicts almost every element  
1999 reviewed here. Overcoming this limitation will require the most creativity; we suggest more studies testing  
2000 the fidelity of non-traditional substrates (e.g., mixed foraminiferal assemblages, coccolith calcite, diatom  
2001 opal), periodic re-assessment of the feasibility of traditional substrates following analytical advancements  
2002 (e.g., improvements in ion transmission efficiency, large-scale [automated and/or crowd-sourced] picking  
2003 of monospecific foraminiferal assemblages), and development and validation of selective extraction  
2004 protocols that can be used to isolate phases of interest from complex matrices. Though attempting to  
2005 overcome these limitations may be considered high risk, we believe that this risk is more than justified by  
2006 the reward of developing a more complete understanding of Earth's biogeochemical history.

2007

2008

2009

2010



2011 **Postscripts**

2012

2013 **Acknowledgments**

2014 This contribution grew (and grew) out of the joint GEOTRACES–PAGES workshop held in Aix-en-  
 2015 Provence in December 2018. The authors extend tremendous thanks to the workshop organizers, and in  
 2016 particular Bob Anderson, Catherine Jeandel, Laurence Vidal, and Kazuyo Tachikawa. T.J.H. acknowledges  
 2017 support from the U.S. National Science Foundation; S.H.L. from the U.K. Natural Environment Research  
 2018 Council (NE/P018181/1); T.M.C. from the University of South Florida; and, J.R.F. from the Max Planck  
 2019 Society, the Tuttle Fund of the Department of Geosciences of Princeton University, the Grand Challenges  
 2020 Program of the Princeton Environmental Institute, and the Andlinger Center for Energy and the  
 2021 Environment of Princeton University. Most importantly, we acknowledge the hundreds of scientists who  
 2022 have generated GEOTRACES-compliant TEI data that made this synthesis possible.

2023

2024 **Data sources**

2025 Dissolved TEI data are sourced from the GEOTRACES Intermediate Data Products in 2014 (Mawji et al.,  
 2026 2015) and 2017 (Schlitzer et al., 2018). Citations to the primary data sources used in the figures and Table  
 2027 1 are given below.

2028

2029 *Figures*

2030 Iron: Conway & John, 2014a (Atlantic); Conway & John, 2015a (Pacific); Horner et al., 2015b (CD29-2  
 2031 record); Chu et al., 2006 (28DSR9 record); Liu et al., 2020 (METG-03 record). Zinc: Conway & John,  
 2032 2014b (Atlantic); Conway & John, 2015a (Pacific); Wang et al., 2019a (Southern); Little et al., 2014 (Fe–  
 2033 Mn crust records). Copper: Little et al., 2018 (Atlantic); Takano et al., 2017 (Pacific); Boye et al., 2012  
 2034 (Southern); Little et al., 2014 (Fe–Mn crust records). Cadmium: Conway and John, 2015b (Atlantic);  
 2035 Conway & John, 2015a (Pacific); Abouchami et al., 2014 (Southern); Schmitt et al., 2009a (Fe–Mn crust  
 2036 data); Horner et al., 2010 (Fe–Mn crust ‘core-top’ data); Murphy, 2016 (‘down core’ Fe–Mn crust records);  
 2037 Schmitt et al., 2009a (‘down core’ Fe–Mn crust record); van Geen et al., 1995 (California Margin core-top  
 2038 data). Molybdenum: Nakagawa et al., 2012 (all basins). Barium: Bates et al., 2017 (Atlantic); Geyman et  
 2039 al., 2019 (Pacific); Hsieh & Henderson, 2017 (Southern). Nickel: Archer et al., 2020 (Atlantic); Takano et  
 2040 al., 2017 (Pacific); Wang et al., 2019a (Southern); Gall et al., 2013 (CD29-2 record); Gueguen et al., 2016  
 2041 (Fe–Mn crust records). Silver: Fischer et al., 2018 (Pacific); Boye et al., 2012 (Southern). Core MD02-2496  
 2042 (Vancouver Island Margin; 48.97°N, 127.04°W, 1,243 m water depth) from Chang et al., 2014; Core

2043 W8709-13PC (Oregon Margin; 42.116°N, 125.75°W; 2,712 m water depth) from Kienast, 2003; ODP  
 2044 1017E (Southern California; 34.53°N, 121.1°W; 955 m water depth) from Hendy & Pedersen, 2005; ODP  
 2045 893A (Santa Barbara Basin; 34.287°N, 120.036°W; 576.5 m water depth) from Ivanochko, 2001.

2046 *Table 1*

2047 <sup>a</sup>Schlitzer (2017); <sup>b</sup>Morris et al. (1975); <sup>c</sup>Gallon and Flegal, 2014; <sup>d</sup>Beard et al. (2003); <sup>e</sup>Moynier et al.  
 2048 (2017) and references therein; <sup>f</sup>Schmitt et al. (2009a); <sup>g</sup>Willbold & Elliott (2017); <sup>h</sup>Nan et al., 2018;  
 2049 <sup>i</sup>Cameron et al. (2009); <sup>j</sup>USGS reference material SCO-1 (Cody Shale) is used given paucity of  
 2050 representative data; Schönbächler et al. (2007); <sup>k</sup>Section 3 and references therein; <sup>l</sup>Section 4 and references  
 2051 therein; <sup>m</sup>Section 5 and references therein; <sup>n</sup>Section 6 and references therein; <sup>o</sup>Data from Nakagawa et al.  
 2052 (2012) renormalized to NIST SRM 3134 + 0.25 ‰ based on Goldberg et al. (2013); <sup>p</sup>Section 8 and  
 2053 references therein; <sup>q</sup>Section 9 and references therein; <sup>r</sup>Hayes et al. (2018) and references therein; <sup>s</sup>Miller et  
 2054 al. (2011); <sup>t</sup>Dickens et al. (2003); <sup>u</sup>Cameron & Vance (2014) and references therein; Little et al. (2015;  
 2055 2020); <sup>v</sup>Broecker & Peng (1982).

2056

2057 **Competing interests**

2058 The authors declare that they are not aware of any financial conflicts of interest, real or perceived.

2059

2060 **GEOTRACES–PAGES Biological Productivity Working Group Members**

2061 C. Bolton (CEREGE, Univ. Aix-Marseille, CNRS, IRD, Collège de France, INRAE, Aix-en-Provence,  
 2062 France); E. Calvo (Institut de Ciències del Mar, CSIC, Barcelona, Spain); D. Cardinal (LOCEAN  
 2063 (UMR7159), Sorbonne Université, IRD, CNRS, MNHN, Paris, France); T. de Garidel-Thoron  
 2064 (CEREGE, Univ. Aix-Marseille, CNRS, IRD, Collège de France, INRAE Aix-en-Provence, France); S.  
 2065 Fietz (Department of Earth Sciences, Stellenbosch University, Stellenbosch South Africa); K. R.  
 2066 Hendry (School of Earth Sciences, University of Bristol, Bristol, UK); F. Marcantonio (Department  
 2067 of Geology and Geophysics, Texas A&M University), P. A. Rafter (Department of Earth System  
 2068 Science, University of California, Irvine, CA, USA); H. Ren (Department of Geosciences, National  
 2069 Taiwan University, Taipei, Taiwan); C. J. Somes (GEOMAR Helmholtz Centre for Ocean Research  
 2070 Kiel, 24105 Kiel, Germany); J. N. Sutton (Univ Brest, CNRS, IRD, Ifremer, Institut Universitaire  
 2071 Européen de la Mer, LEMAR, Plouzané, France); A. Torfstein (Institute of Earth Sciences, Hebrew  
 2072 University, Jerusalem, & InterUniversity Institute for Marine Sciences, Eilat, Israel); G. Winckler (Lamont-  
 2073 Doherty Earth Observatory of Columbia University, Palisades, NY, USA).

2074 **References**

- 2075 Abadie, C., Lacan, F., Radic, A., Pradoux, C., & Poitrasson, F. (2017). Iron isotopes reveal distinct  
 2076 dissolved iron sources and pathways in the intermediate versus deep Southern Ocean. *Proc. Natl.*  
 2077 *Acad. Sci.*, **114**, 858-863.
- 2078 Abouchami, W., Galer, S. J. G., de Baar, H. J. W., Alderkamp, A. C., Middag, R., Laan, P., Feldmann, H.,  
 2079 & Andreae, M. O. (2011). Modulation of the Southern Ocean cadmium isotope signature by ocean  
 2080 circulation and primary productivity. *Earth Planet. Sci. Lett.*, **305**(1-2), 83-91.
- 2081 Abouchami, W., Galer, S. J. G., Horner, T. J., Rehkämper, M., Wombacher, F., Xue, Z., Lambelet, M.,  
 2082 Gault-Ringold, M., Stirling, C. H., Schönbächler, M., Shiel, A. E., Weis, D., & Holdship, P. F.  
 2083 (2013). A common reference material for cadmium isotope studies—NIST SRM 3108.  
 2084 *Geostandards and Geoanalytical Research*, **37**(1), 5-17.
- 2085 Abouchami, W., Galer, S. J. G., de Baar, H. J. W., Middag, R., Vance, D., Zhao, Y., Klunder, M., Mezger,  
 2086 K., Feldmann, H., & Andreae, M. O. (2014). Biogeochemical cycling of cadmium isotopes in the  
 2087 Southern Ocean along the Zero Meridian. *Geochim. Cosmochim. Acta*, **127**, 348-367.
- 2088 Adkins, J. F., Cheng, H., Boyle, E. A., Druffel, E. R., & Edwards, R. L. (1998). Deep-sea coral evidence  
 2089 for rapid change in ventilation of the deep North Atlantic 15,400 years ago. *Science*, **280**(5364),  
 2090 725-728.
- 2091 Akerman, A., Poitrasson, F., Oliva, P., Audry, S., Prunier, J., & Braun, J. J. (2014). The isotopic fingerprint  
 2092 of Fe cycling in an equatorial soil-plant-water system: The Nsimi watershed, South Cameroon.  
 2093 *Chem. Geol.*, **385**, 104–116.
- 2094 Albarède, F. (2004). The stable isotope geochemistry of copper and zinc. *Reviews in Mineralogy and*  
 2095 *Geochemistry*, **55**(1), 409-427.
- 2096 Algeo, T. J. (2004). Can marine anoxic events draw down the trace element inventory of seawater? *Geology*,  
 2097 **32** (12), 1057-1060, doi:10.1130/G20896.1.
- 2098 Algeo, T. J., & Lyons, T. W. (2006). Mo-total organic carbon covariation in modern anoxic marine  
 2099 environments: Implications for analysis of paleoredox and paleohydrographic conditions:  
 2100 *Paleoceanography*, **21**, doi:10.1029/2004PA001112.
- 2101 Algeo T. J., & Maynard J. B. (2008). Trace-metal covariation as a guide to water-mass conditions in ancient  
 2102 anoxic marine environments. *Geosphere* **4**, 872.
- 2103 Algeo, T. J., & Tribovillard, N. (2009). Environmental analysis of paleoceanographic systems based on  
 2104 molybdenum-uranium covariation: *Chem. Geol.*, **268**, 211–225,  
 2105 doi:10.1016/j.chemgeo.2009.09.001.
- 2106 Alldredge, A. L., & Silver, M. W. (1988). Characteristics, dynamics and significance of marine snow. *Prog.*  
 2107 *Oceanogr.*, **20**(1), 41-82.
- 2108 Anbar, A. D., & Knoll, A. H. (2002). Proterozoic Ocean Chemistry and Evolution: A Bioinorganic Bridge?  
 2109 *Science*, **297**, 1137–1142, doi:10.1126/science.1069651.
- 2110 Anbar, A. D., Jarzecki, A. A., & Spiro, T. G. (2005) Theoretical investigation of iron isotope fractionation  
 2111 between  $\text{Fe}(\text{H}_2\text{O})_6^{3+}$  and  $\text{Fe}(\text{H}_2\text{O})_6^{2+}$ : Implications for iron stable isotope geochemistry. *Geochim.*  
 2112 *Cosmochim. Acta.* **69**, 825–837. <https://doi.org/10.1016/j.gca.2004.06.012>.
- 2113 Andersen, M. B., Vance, D., Archer, C., Anderson, R. F., Ellwood, M. J., & Allen, C. S. (2011). The Zn  
 2114 abundance and isotopic composition of diatom frustules, a proxy for Zn availability in ocean  
 2115 surface seawater. *Earth Planet. Sci. Lett.*, **301**, 137–145.  
 2116 <http://dx.doi.org/10.1016/j.epsl.2010.10.032>.

- 2117 Anderson, M. A., Morel, F. M. M., & Guillard, R. R. L. (1978). Growth limitation of a coastal diatom by  
2118 low zinc ion activity. *Nature*, **276**, 70-71.
- 2119 Annett, A. L., Lapi, S., Ruth, T. J., & Maldonado, M. T. (2008). The effects of Cu and Fe availability on  
2120 the growth and Cu:C ratios of marine diatoms. *Limnol. Oceanogr.* **53**, 2451–2461.
- 2121 Aplin, A. C., & Cronan, D. S. (1985). Ferromanganese oxide deposits from the central Pacific Ocean, I.  
2122 Encrustations from the Line Islands Archipelago. *Geochim. Cosmochim. Acta*, **49**(2), 427-436.
- 2123 Archer, C., Andersen, M. B., Cloquet, C., Conway, T. M., Dong, S., Ellwood, M., Moore, R., Nelson, J.,  
2124 Rehkämper, M., Rouxel, O., Samanta, M., Shin, K.-C., Sohrin, Y., Takano, S., & Wasylenki, L.  
2125 (2017). Inter-calibration of a proposed new primary reference standard AA-ETH Zn for zinc  
2126 isotopic analysis. *Journal of Analytical Atomic Spectrometry*, **32**(2), 415-419.
- 2127 Archer, C., Vance, D., Milne, A., & Lohan, M. C. (2020). The oceanic biogeochemistry of nickel and its  
2128 isotopes: New data from the South Atlantic and the Southern Ocean biogeochemical divide. *Earth  
2129 Planet. Sci. Lett.*, **535**, 116118.
- 2130 Aristilde, L., & Xu, Y. (2012). Weak organic ligands enhance Zinc uptake in Marine phytoplankton.  
2131 *Environ. Sci. Technol.*, **46**, 5438-5445.
- 2132 Arrhenius, G. (1963). *Pelagic sediments*. Scripps Institution of Oceanography, University of California.
- 2133 Atkins, A. L., Shaw, S., & Peacock, C. L. (2016). Release of Ni from birnessite during transformation of  
2134 birnessite to todorokite: Implications for Ni cycling in marine sediments. *Geochim. Cosmochim.  
2135 Acta*, **189**, 158-183.
- 2136 Baars, O., & Croot, P. L. (2011). The speciation of dissolved zinc in the Atlantic sector of the Southern  
2137 Ocean. *Deep. Res. Part II Top. Stud. Oceanogr.*, **58**, 2707–2719.  
2138 <http://dx.doi.org/10.1016/j.dsr2.2010.10.041>.
- 2139 Baars, O., Abouchami, W., Galer, S. J., Boye, M., & Croot, P. L. (2014). Dissolved cadmium in the  
2140 Southern Ocean: distribution, speciation, and relation to phosphate. *Limnol. Oceanogr.*, **59**(2), 385-  
2141 399.
- 2142 Bacon, M.P. and Anderson, R.F. (1982). Distribution of thorium isotopes between dissolved and particulate  
2143 forms in the deep sea. *J. Geophys. Res. Oceans*, **87**(C3), 2045-2056.
- 2144 Badaut, D., & Risacher, F. (1983). Authigenic smectite on diatom frustules in Bolivian saline lakes.  
2145 *Geochim. Cosmochim. Acta*, **47**(3), 363-375.
- 2146 Barling, J., Arnold, G., & Anbar, A. D., (2001). Natural mass-dependent variations in the isotopic  
2147 composition of molybdenum. *Earth Planet. Sci. Lett.*, **193**, 447–457, doi:10.1016/S0012-  
2148 821X(01)00514-3.
- 2149 Barling, J., & Anbar, A. D. (2004). Molybdenum isotope fractionation during adsorption by manganese  
2150 oxides. *Earth Planet. Sci. Lett.*, **217**, 315-329, doi:10.1016/S0012-821X(03)00608-3.
- 2151 Bates, S. L., Hendry, K. R., Pryer, H. V., Kinsley, C. W., Pyle, K. M., Woodward, E. M. S., & Horner, T.  
2152 J. (2017). Barium isotopes reveal role of ocean circulation on barium cycling in the Atlantic.  
2153 *Geochim. Cosmochim. Acta*, **204**, 286-299.
- 2154 Beard, B.L., Johnson, C.M., Cox, L., Sun, H., Neelson, K.H., & Aguilar, C. (1999). Iron Isotope  
2155 Biosignatures. *Science* **285**, 1889-1892.
- 2156 Beard, B. L., Johnson, C. M., Skulan, J. L., Neelson, K. H., Cox, L., & Sun, H. (2003). Application of Fe  
2157 isotopes to tracing the geochemical and biological cycling of Fe. *Chem. Geol.*, **195**(1-4), 87-117.

- 2158 Bender, M., Grande, K., Johnson, K., Marra, J., Williams, P. J. L., Sieburth, J., Pilson, M., Langdon, C.,  
 2159 Hitchcock, G., Orchoado, J., Hunt, C., Donaghy, P., & Heinemann, K. (1987). A comparison of four  
 2160 methods for determining planktonic community production. *Limnol. Oceanogr.*, **32**(5), 1085-1098.
- 2161 Bennett, S. A., Rouxel, O., Schmidt, K., Garbe-Schönberg, D., Statham, P. J., & German, C. R. (2009).  
 2162 Iron isotope fractionation in a buoyant hydrothermal plume, 5 S Mid-Atlantic Ridge. *Geochim.*  
 2163 *Cosmochim. Acta*, **73**(19), 5619-5634.
- 2164 Bergquist, B. A., & Boyle, E. A. (2006). Iron isotopes in the Amazon River system: Weathering and  
 2165 transport signatures. *Earth Planet. Sci. Lett.*, **248**(1-2), 54-68.  
 2166 <https://doi.org/10.1016/j.epsl.2006.05.004>.
- 2167 Berger, W. H., Smetacek, V., & Wefer, G. (1989). Ocean productivity and paleoproductivity - an overview.  
 2168 In W. H. Berger, V. S. Smetacek, G. Wefer (Eds) *Productivity of the Oceans present and past:*  
 2169 *Report of the Dahlem Workshop on Productivity of the Ocean, Berlin, 1988*, Life Sciences Research  
 2170 Reports 44, Wiley & Sons, Chichester, pp. 1-34.
- 2171 Bermin, J., Vance, D., Archer, C., & Statham, P. J. (2006). The determination of the isotopic composition  
 2172 of Cu and Zn in seawater. *Chem. Geol.*, **226**, 280-297.
- 2173 Bernstein, R. E., & Byrne, R. H. (2004). Acantharians and marine barite. *Mar. Chem.*, **86** (1-2), 45-50.
- 2174 Bertine, K. K., & Turekian, K. K. (1973). Molybdenum in marine deposits. *Geochim. Cosmochim. Acta*,  
 2175 **37** (6), 1415-1434.
- 2176 Bianchi, D., Weber, T. S., Kiko, R., & Deutsch, C. (2018). Global niche of marine anaerobic metabolisms  
 2177 expanded by particle microenvironments. *Nat. Geosci.*, **11**, 1-6.
- 2178 Bishop, J. K. (1988). The barite-opal-organic carbon association in oceanic particulate matter. *Nature*, **332**  
 2179 (6162), 341.
- 2180 Böning, P., Brumsack, H.-J., Böttcher, M.E., Schnetger, B., Kriete, C., Kallmeyer, J., & Borchers, S.L.  
 2181 (2004). Geochemistry of Peruvian near-surface sediments. *Geochim. Cosmochim. Acta*, **68**, 4429-  
 2182 4451.
- 2183 Böning, P., Cuypers, S., Grunwald, M., Schnetger, B., & Brumsack, H.-J. (2005). Geochemical  
 2184 characteristics of Chilean upwelling sediments at ~36°S. *Mar. Geol.*, **220**, 1-21.
- 2185 Böning, P., Brumsack, H.-J., Schnetger, B., & Grunwald, M. (2009). Trace element signatures of Chilean  
 2186 upwelling sediments at ~36°S. *Mar. Geol.*, **259**, 112-121.
- 2187 Böning, P., Shaw, T., Pahnke, K., & Brumsack, H.-J. (2015). Nickel as an indicator of fresh organic matter  
 2188 in upwelling sediments. *Geochim. Cosmochim. Acta*, **162**, 99-108, doi:10.1016/j.gca.2015.04.027.
- 2189 Borchers, S.L., Schnetger, B., Boning, P., & Brumsack, H.-J. (2005). Geochemical signatures of the  
 2190 Namibian diatom belt: Perennial upwelling and intermittent anoxia. *Geochemistry, Geophysics,*  
 2191 *Geosystems*, **6**(4), Q06006, doi:10.1029/2004GC000886.
- 2192 Boyd, P. W., Jickells, T., Law, C. S., Blain, S., Boyle, E. A., Buesseler, K. O., Coale, K. H., Cullen, J. J.,  
 2193 de Baar, H. J. W., Follows, M., Harvey, M., Lancelot, C., Levasseur, M., Owens, N. P. J., Pollard,  
 2194 R., Rivkin, R. B., Sarimento, J., Schoemann, V., Smetacek, V., Takeda, S., Tsuda, A., Turner, S.,  
 2195 & Watson, A. J.. (2007). Mesoscale iron enrichment experiments 1993-2005: synthesis and future  
 2196 directions. *Science*, **315**(5812), 612-617.
- 2197 Boyd, P. W., & Ellwood, M. J. (2010). The biogeochemical cycle of iron in the ocean. *Nat. Geosci.* **3**(10),  
 2198 675-682.
- 2199 Boyd, P. W., Ellwood, M. J., Tagliabue, A., & Twining, B. S. (2017). Biotic and abiotic retention, recycling  
 2200 and remineralization of metals in the ocean. *Nat. Geosci.*, **10**(3), 167-173.



- 2201 Boye, M., Wake, D. B., Lopez Gracia, P., Bown, J., Baker, A. R., & Achterberg, E. P. (2012). Distributions  
 2202 of dissolved trace metals (Cd, Cu, Mn, Pb, Ag) in the southeastern Atlantic and the Southern Ocean.  
 2203 *Biogeosciences*, **9**, 3231-3246, doi:10.5194/bg-9-3231-2012.
- 2204 Boyle, E. A. (1981). Cadmium, zinc, copper, and barium in foraminifera tests. *Earth Planet. Sci. Lett.*,  
 2205 **53**(1), 11-35.
- 2206 Boyle, E. A. (1988a). Vertical oceanic nutrient fractionation and glacial/interglacial CO<sub>2</sub> cycles. *Nature*,  
 2207 **331**(6151), 55-56.
- 2208 Boyle, E. A. (1988b). Cadmium: Chemical tracer of deepwater paleoceanography. *Paleoceanography*, **3**(4),  
 2209 471-489.
- 2210 Boyle, E. A. (1992). Cadmium and  $\delta^{13}\text{C}$  paleochemical ocean distributions during the stage 2 glacial  
 2211 maximum. *Annual Review of Earth and Planetary Sciences*, **20**(1), 245-287.
- 2212 Boyle, E. A., & Keigwin, L. D. (1985). Comparison of Atlantic and Pacific paleochemical records for the  
 2213 last 215,000 years: Changes in deep ocean circulation and chemical inventories. *Earth Planet. Sci.*  
 2214 *Lett.*, **76**(1-2), 135-150.
- 2215 Boyle, E. A., Sclater, F., & Edmond, J. M. (1976). On the marine geochemistry of cadmium. *Nature*,  
 2216 **263**(5572), 42.
- 2217 Boyle, E. A., Sclater, F. R., & Edmond, J. M. (1977). The distribution of dissolved copper in the Pacific.  
 2218 *Earth Planet. Sci. Lett.*, **37**, 38-54.
- 2219 Brand, L. E., Sunda, W. G., & Guillard, R. R. L. (1983). Limitation of marine phytoplankton by zinc,  
 2220 manganese, and iron. *Limnol. Oceanogr.*, **28**, 1182-1198.
- 2221 Brand L. E., Sunda, W. G., & Guillard, R. R. L. (1986). Reduction of marine phytoplankton reproduction  
 2222 rates by copper and cadmium. *J. Exp. Mar. Bio. Ecol.* **96**, 225-250.
- 2223 Bridgestock, L., Rehkämper, M., van de Flierdt, T., Murphy, K., Khondoker, R., Baker, A. R., Chance, R.,  
 2224 Strekopytov, S., Humphreys-Williams, E., Achterberg, E. P. (2017). The Cd isotope composition  
 2225 of atmospheric aerosols from the Tropical Atlantic Ocean. *Geophys. Res. Lett.* **44**(6), 2932-2940.  
 2226 <https://doi.org/10.1002/2017GL072748>.
- 2227 Bridgestock, L., Hsieh, Y. T., Porcelli, D., Homoky, W. B., Bryan, A., & Henderson, G. M. (2018). Controls  
 2228 on the barium isotope compositions of marine sediments. *Earth Planet. Sci. Lett.*, **481**, 101-110.
- 2229 Bridgestock, L., Hsieh, Y. T., Porcelli, D., & Henderson, G. M. (2019). Increased export production during  
 2230 recovery from the Paleocene–Eocene thermal maximum constrained by sedimentary Ba isotopes.  
 2231 *Earth Planet. Sci. Lett.*, **510**, 53-63.
- 2232 Broecker, W.S. (1982). Glacial to interglacial changes in ocean chemistry. *Prog. Oceanogr.* **11**, 151-197.
- 2233 Broecker, W.S., & Peng, T. H. (1982). *Tracers in the Sea*. Eldigo Press, Palisades, NY.
- 2234 Brucker, R. L. P., McManus, J., Severmann, S., & Berelson, W. M. (2009). Molybdenum behavior during  
 2235 early diagenesis: Insights from Mo isotopes. *Geochemistry, Geophysics, Geosystems*, **10**,  
 2236 doi:10.1029/2008GC002180.
- 2237 Bruland, K. W. (1980). Oceanographic distributions of cadmium, zinc, nickel, and copper in the North  
 2238 Pacific. *Earth Planet. Sci. Lett.*, **47**(2), 176-198.
- 2239 Bruland, K. W. (1989). Complexation of zinc by natural organic ligands in the central North Pacific. *Limnol.*  
 2240 *Oceanogr.*, **34**, 269-285.
- 2241 Bruland, K. W. (1992). Complexation of cadmium by natural organic ligands in the central North Pacific.  
 2242 *Limnol. Oceanogr.*, **37**(5), 1008-1017.

- 2243 Bruland, K. W., & Lohan, M. C. (2003). Controls of Trace Metals in Seawater. In *Treatise on Geochemistry*,  
 2244 Elsevier, 23-47.
- 2245 Brumsack, H. J. (2006). The trace metal content of recent organic carbon-rich sediments: implications for  
 2246 Cretaceous black shale formation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **232**(2-  
 2247 4), 344-361.
- 2248 Bryan, A. L., Dong, S., Wilkes, E. B., & Wasylenki, L. E. (2015). Zinc isotope fractionation during  
 2249 adsorption onto Mn oxyhydroxide at low and high ionic strength. *Geochim. Cosmochim. Acta*, **157**,  
 2250 182-197.
- 2251 Buesseler, K. O., Boyd, P. W., Black, E. E., & Siegel, D. A. (2020). Metrics that matter for assessing the  
 2252 ocean biological carbon pump. *Proc. Natl. Acad. Sci.*, **117**(18), 9679-9687.
- 2253 Cadiou, J. L., Pichat, S., Bondanese, V. P., Soulard, A., Fujii, T., Albarède, F., & Oger, P. (2017). Copper  
 2254 transporters are responsible for copper isotopic fractionation in eukaryotic cells. *Sci. Rep.*, **7**(1), 1-  
 2255 10.
- 2256 Cameron, V., Vance, D., Archer, C., & House, C. H. (2009). A biomarker based on the stable isotopes of  
 2257 nickel. *Proceedings of the National Academy of Sciences*, **106**(27), 10944-10948.
- 2258 Carlson, C. A., Hansell, D. A., Nelson, N. B., Siegel, D. A., Smethie, W. M., Khatiwala, S., Meyers, M.  
 2259 M., & Halewood, E. (2010). Dissolved organic carbon export and subsequent remineralization in  
 2260 the mesopelagic and bathypelagic realms of the North Atlantic basin. *Deep Sea Res. Pt. II*, **57**(16),  
 2261 1433-1445.
- 2262 Cao, Z., Siebert, C., Hathorne, E. C., Dai, M., & Frank, M. (2020). Corrigendum to "Constraining the  
 2263 oceanic barium cycle with stable barium isotopes" [*Earth Planet. Sci. Lett.*, **434** (2016) 1-9], doi:  
 2264 10.1016/j.epsl.2019.116003/
- 2265 Cartapanis, O., Bianchi, D., Jaccard, S. L., & Galbraith, E. D. (2016). Global pulses of organic carbon burial  
 2266 in deep-sea sediments during glacial maxima. *Nat. Commun.*, **7**, 10796,  
 2267 doi:10.1038/ncomms10796.
- 2268 Carter, S. C., Paytan, A., & Griffith, E. M. (2020). Toward an Improved Understanding of the Marine  
 2269 Barium Cycle and the Application of Marine Barite as a Paleoproductivity Proxy. *Minerals*, **10**(5),  
 2270 421.
- 2271 Chan, L. H., Edmond, J. M., Stallard, R. F., Broecker, W. S., Chung, Y. C., Weiss, R. F., & Ku, T. L.  
 2272 (1976). Radium and barium at GEOSECS stations in the Atlantic and Pacific. *Earth Planet. Sci.*  
 2273 *Lett.*, **32**(2), 258-267.
- 2274 Chan, L. H., Drummond, D., Edmond, J. M., & Grant, B. (1977). On the barium data from the Atlantic  
 2275 GEOSECS expedition. *Deep Sea Res.*, **24**(7), 613-649.
- 2276 Chang, A. S., Pedersen, T. F., & Hendy, I. L. (2014). Effects of productivity, glaciation, and ventilation on  
 2277 late Quaternary sedimentary redox and trace element accumulation on the Vancouver Island  
 2278 margin, western Canada. *Paleoceanogr.*, **29**(7), 730-746.
- 2279 Chappaz, A., Lyons, T. W., Gregory, D. D., Reinhard, C. T., Gill, B. C., Li, C., & Large, R. R. (2014).  
 2280 Does pyrite act as an important host for molybdenum in modern and ancient euxinic sediments?  
 2281 *Geochim. Cosmochim. Acta*, **126**, 112–122, doi:10.1016/j.gca.2013.10.028.
- 2282 Charette, M. A., et al. (2020). The Transpolar Drift as a Source of Riverine and Shelf-Derived Trace  
 2283 Elements to the Central Arctic Ocean. *J. Geophys. Res. Oceans*, **125**(5), e2019JC015920.  
 2284 <https://doi.org/10.1029/2019JC015920>.
- 2285 Chen, J.-B., Busigny, V., Gaillardet, J., Louvat, P., & Wang, Y.-N. (2014). Iron isotopes in the Seine River  
 2286 (France): Natural vs anthropogenic sources. *Geochim. Cosmochim. Acta*, **128**, 128-143.



- 2287 Chen., T., Wenqiang, L., Gao, B., Liu, R., Li, G., Zhao, L., and Ji, J. (2020). Reactive iron isotope signatures  
 2288 of the East Asian dust particles: Implications for iron cycling the deep North Pacific. *Chem. Geol.*  
 2289 **531**, 119342. <https://doi.org/10.1016/j.chemgeo.2019.119342>.
- 2290 Cheng, H., Adkins, J., Edwards, R. L., & Boyle, E. A. (2000). U-Th dating of deep-sea corals. *Geochim.*  
 2291 *Cosmochim. Acta*, **64**(14), 2401-2416.
- 2292 Chever, F., Bucciarelli, E., Sarthou, G., Speich, S., Arhan, M., Penven, P., & Tagliabue, A. (2010). Physical  
 2293 speciation of iron in the Atlantic sector of the Southern Ocean along a transect from the subtropical  
 2294 domain to the Weddell Sea Gyre. *J. Geophys. Res. Oceans*, **115**, C10059,  
 2295 doi:10.1029/2009JC005880.
- 2296 Chever, F., Rouxel, O. J., Croot, P. L., Ponzevera, E., Wuttig, K., & Auro, M. (2015). Total dissolvable and  
 2297 dissolved iron isotopes in the water column of the Peru upwelling regime. *Geochim. Cosmochim.*  
 2298 *Acta*, **162**, 66-82. <https://doi.org/10.1016/j.gca.2015.04.031>.
- 2299 Chow, T. J., & Goldberg, E. D. (1960). On the marine geochemistry of barium. *Geochim. Cosmochim.*  
 2300 *Acta*, **20**(3-4), 192-198.
- 2301 Chu, N., Johnson, C., Beard, B., German, C., Nesbitt, R., Frank, M., Bohn, M., Kubik, P., Usui, A., &  
 2302 Graham, I. (2006). Evidence for hydrothermal venting in Fe isotope compositions of the deep  
 2303 Pacific Ocean through time. *Earth Planet. Sci. Lett.*, **245**, 202-217.
- 2304 Ciscato E. R., Bontognali T. R. R., & Vance, D. (2018). Nickel and its isotopes in organic-rich sediments:  
 2305 implications for oceanic budgets and a potential record of ancient seawater. *Earth Planet. Sci. Lett.*  
 2306 **494**, 239–250. <https://doi.org/10.1016/j.epsl.2018.04.061>.
- 2307 Ciscato, E. R., Bontognali, T. R., Poulton, S. W., & Vance, D. (2019). Copper and its Isotopes in Organic-  
 2308 Rich Sediments: From the Modern Peru Margin to Archean Shales. *Geosciences*, **9**(8), 325.
- 2309 Clarkson, M., Müsing, K., Andersen, M. B., & Vance, D. (2018). Reducing the role of contaminant phases  
 2310 for metal isotopes in carbonates. In *Goldschmidt Abstracts* p. 441.
- 2311 Coale, K. H. (1991). Effects of iron, manganese, copper and zinc enrichments on productivity and biomass  
 2312 in the subarctic Pacific. *Limnol. Oceanogr.*, **36**, 1851–1864.
- 2313 Coale K. H., & Bruland K. W. (1988). Copper complexation in the Northeast Pacific. *Limnol. Oceanogr.*  
 2314 **33**, 1084–1101.
- 2315 Coale K. H., & Bruland K. W. (1990). Spatial and temporal variability in copper complexation in the North  
 2316 Pacific. *Deep Sea Res. Part A, Oceanogr. Res. Pap.* **37**, 317–336.
- 2317 Coale, K. H., Wang, X., Tanner, S. J., & Johnson, K. S. (2003). Phytoplankton growth and biological  
 2318 response to iron and zinc addition in the Ross Sea and Antarctic Circumpolar Current along 170°W.  
 2319 *Deep. Res. Part II Top. Stud. Oceanogr.*, **50**, 635–653.
- 2320 Collier, R. W. (1985). Molybdenum in the Northeast Pacific Ocean. *Limnol. Oceanogr.*, **30**, 1351–1354,  
 2321 doi:10.4319/lo.1985.30.6.1351
- 2322 Conway, T. M., & John, S. G. (2014a). Quantification of dissolved iron sources to the North Atlantic Ocean.  
 2323 *Nature*, 511, 212-215.
- 2324 Conway, T. M., & John, S. G. (2014b). The biogeochemical cycling of zinc and zinc isotopes in the North  
 2325 Atlantic Ocean. *Global Biogeochem. Cycles*, 1111–1128.
- 2326 Conway, T. M., & John, S. G. (2015a). The cycling of iron, zinc and cadmium in the North East Pacific  
 2327 Ocean—Insights from stable isotopes. *Geochim. Cosmochim. Acta*, **164**, 262-283.
- 2328 Conway, T. M., & John, S. G. (2015b). Biogeochemical cycling of cadmium isotopes along a high-  
 2329 resolution section through the North Atlantic Ocean. *Geochim. Cosmochim. Acta*, **148**, 269-283.

- 2330 Conway, T. M., Rosenberg, A. D., Adkins, J. F., & John, S. G. (2013). A new method for precise  
 2331 determination of iron, zinc and cadmium stable isotope ratios in seawater by double-spike mass  
 2332 spectrometry. *Anal. Chim. Acta*, **793**, 44–52.
- 2333 Conway, T. M., Hamilton, D. S., Shelley, R. U., Aguilar-Islas, A. M., Landing, W. M., Mahowald, N. M.,  
 2334 & John, S. G. (2019). Tracing and constraining anthropogenic aerosol iron fluxes to the North  
 2335 Atlantic Ocean using iron isotopes. *Nat. Commun.*, **10**, 2628, doi:10.1038/s41467-019-10457-w.
- 2336 Coplen, T. B. (2011). Guidelines and recommended terms for expression of stable-isotope-ratio and gas-  
 2337 ratio measurement results. *Rapid communications in mass spectrometry*, **25**(17), 2538-2560.
- 2338 Costa, K. M., McManus, J. F., Anderson, R. F., Ren, H., Sigman, D. M., Winckler, G., Fleisher, M. Q.,  
 2339 Marcantonio, F., & Ravelo, A. C. (2016). No iron fertilization in the equatorial Pacific Ocean  
 2340 during the last ice age. *Nature*, **529**(7587), 519-522.
- 2341 Costa, K. M., McManus, J. F., Middleton, J. L., Langmuir, C. H., Huybers, P. J., Winckler, G., &  
 2342 Mukhopadhyay, S. (2017). Hydrothermal deposition on the Juan de Fuca Ridge over multiple  
 2343 glacial–interglacial cycles. *Earth Planet. Sci. Lett.*, **479**, 120-132.
- 2344 Coutaud, M., Méheut, M., Glatzel, P., Pokrovski, G. S., Viers, J., Rols, J. L., & Pokrovsky, O. S. (2018).  
 2345 Small changes in Cu redox state and speciation generate large isotope fractionation during  
 2346 adsorption and incorporation of Cu by a phototrophic biofilm. *Geochim. Cosmochim. Acta*, **220**, 1-  
 2347 18.
- 2348 Coutaud, M., Méheut, M., Viers, J., Rols, J. L., & Pokrovsky, O. S. (2019). Copper isotope fractionation  
 2349 during excretion from a phototrophic biofilm. *Chem. Geol.*, **513**, 88-100.
- 2350 Cowan, C. E., Jenne, E. A., & Crecelius, E. A. (1985). Silver speciation in seawater: The importance of  
 2351 sulfide and organic complexes. In: A.C. Sigleo and A. Hattori (Eds.), *Marine and Estuarine*  
 2352 *Geochemistry*, Lewis Publishers Inc., Chelsea, MI, USA.
- 2353 Crawford, D. W. W., Lipsen, M. S. S., Purdie, D. A. A., Lohan, M. C. C., Statham, P. J. J., Whitney, F. A.  
 2354 A., Putland, J. N. N., Johnson, W. K. K., Sutherland N., Peterson, T. D. D., Harrison, P. J. J., &  
 2355 Wong C. S. S. (2003). Influence of zinc and iron enrichments on phytoplankton growth in the  
 2356 northeastern subarctic Pacific. *Limnol. Oceanogr.*, **48**, 1583–1600.
- 2357 Crockford, P. W., Wing, B. A., Paytan, A., Hodgskiss, M. S., Mayfield, K. K., Hayles, J. A., Middleton, J.  
 2358 E., Ahm, A.-S. C., Johnston, D. T., Caxito, F., Uhlein, G., Halverson, G. P., Eickmann, B., Torres,  
 2359 M., & Horner, T. J. (2019). Barium-isotopic constraints on the origin of post-Marinoan barites.  
 2360 *Earth Planet. Sci. Lett.*, **519**, 234-244.
- 2361 Crusius, J., & Thomson, J. (2003). Mobility of authigenic rhenium, silver, and selenium during  
 2362 postdepositional oxidation in marine sediments. *Geochim. Cosmochim. Acta*, **67**(2), 265-273.
- 2363 Crusius, J., Calvert, S., Pedersen, T., & Sage, D. (1996). Rhenium and molybdenum enrichments in  
 2364 sediments as indicators of oxic, suboxic and sulfidic conditions of deposition. *Earth Planet. Sci.*  
 2365 *Lett.*, **145**, 65–78, doi:10.1016/S0012-821X(96)00204-X.
- 2366 Cullen, J. T. (2006). On the nonlinear relationship between dissolved cadmium and phosphate in the modern  
 2367 global ocean: could chronic iron limitation of phytoplankton growth cause the kink? *Limnol.*  
 2368 *Oceanogr.*, **51**(3), 1369-1380.
- 2369 Cullen, J. T., & Sherrell, R. M. (2005). Effects of dissolved carbon dioxide, zinc, and manganese on the  
 2370 cadmium to phosphorus ratio in natural phytoplankton assemblages. *Limnol. Oceanogr.*, **50**(4),  
 2371 1193-1204.

- 2372 Cullen, J. T., Chase, Z., Coale, K. H., Fitzwater, S. E., & Sherrell, R. M. (2003). Effect of iron limitation  
 2373 on the cadmium to phosphorus ratio of natural phytoplankton assemblages from the Southern  
 2374 Ocean. *Limnology and Oceanography*, **48**(3), 1079-1087.
- 2375 Cullen, J. T., Lane, T. W., Morel, F. M., & Sherrell, R. M. (1999). Modulation of cadmium uptake in  
 2376 phytoplankton by seawater CO<sub>2</sub> concentration. *Nature*, **402**(6758), 165.
- 2377 Cullen, J. T., Chase, Z., Coale, K. H., Fitzwater, S. E., & Sherrell, R. M. (2003). Effect of iron limitation  
 2378 on the cadmium to phosphorus ratio of natural phytoplankton assemblages from the Southern  
 2379 Ocean. *Limnol. Oceanogr.*, **48**(3), 1079-1087.
- 2380 Cullen, J. T., & Coogan, L. A. (2017). Changes in Fe Oxidation Rate in Hydrothermal Plumes as a Potential  
 2381 Driver of Enhanced Hydrothermal Input to Near-Ridge Sediments During Glacial Terminations.  
 2382 *Geophys. Res. Lett.*, **44**(23), 11-951.
- 2383 Cutter, G., Casciotti, K., Croot, P., Geibert, W., Heimbürger, L.-E., Lohan, M., & van de Flierdt, T. (2017).  
 2384 *Sampling and Sample-handling Protocols for GEOTRACES Cruises*, Version 3.  
 2385 <http://www.geotraces.org/images/stories/documents/intercalibration/Cookbook.pdf> [accessed  
 2386 2020-06-23].
- 2387 da Silva, J. F., & Williams, R. J. P. (1991). *The biological chemistry of the elements: the inorganic*  
 2388 *chemistry of life*. Oxford University Press.
- 2389 Dahl, T. W., Anbar, A. D., Gordon, G. W., Rosing, M. T., Frei, R., & Canfield, D. E. (2010). The behavior  
 2390 of molybdenum and its isotopes across the chemocline and in the sediments of sulfidic Lake  
 2391 Cadagno, Switzerland. *Geochim. Cosmochim. Acta*, **74**, 144–163, doi:10.1016/j.gca.2009.09.018.
- 2392 Dahl, T. W., Chappaz, A., Hoek, J., McKenzie, C. J., Svane, S., and Canfield, D. E. (2017). Evidence of  
 2393 molybdenum association with particulate organic matter under sulfidic conditions. *Geobiology*, **15**,  
 2394 311–323, doi:10.1111/gbi.12220.
- 2395 Dalai, T.K., Nishimura, K., & Nozaki, Y. (2005). Geochemistry of molybdenum in the Chao Phraya River  
 2396 estuary, Thailand: Role of suboxic diagenesis and porewater transport. *Chem. Geol.*, **218**, 189–202,  
 2397 doi:10.1016/j.chemgeo.2005.01.002.
- 2398 de Baar, H. J. W., Saager, P. M., Nolting, R. F., & van der Meer, J. (1994). Cadmium versus phosphate in  
 2399 the world ocean. *Mar. Chem.*, **46**(3), 261-281.
- 2400 de Baar, H. J., Boyd, P. W., Coale, K. H., Landry, M. R., Tsuda, A., Assmy, P., ... & Buesseler, K. O.  
 2401 (2005). Synthesis of iron fertilization experiments: from the iron age in the age of enlightenment.  
 2402 *Journal of Geophysical Research: Oceans*, **110**(C9).
- 2403 de Baar, H. J. W., van Heuven, S. M., Abouchami, W., Xue, Z., Galer, S. J. G., Rehkämper, M., Middag,  
 2404 R. & van Ooijen, J. (2017). Interactions of dissolved CO<sub>2</sub> with cadmium isotopes in the Southern  
 2405 Ocean. *Mar. Chem.* **195**, 105-121.
- 2406 de Souza, G. F., Reynolds, B. C., Rickli, J., Frank, M., Saito, M. A., Gerringa, L. J., & Bourdon, B. (2012).  
 2407 Southern Ocean control of silicon stable isotope distribution in the deep Atlantic Ocean. *Global*  
 2408 *Biogeochem. Cycles*, **26**(2).
- 2409 de Souza, G. F., Khatiwala, S. P., Hain, M. P., Little, S. H., & Vance, D. (2018). On the origin of the marine  
 2410 zinc–silicon correlation. *Earth Planet. Sci. Lett.*, **492**, 22-34.
- 2411 Dean, W. E., Zheng, Y., Ortiz, J. D., & van Geen, A. (2006). Sediment Cd and Mo accumulation in the  
 2412 oxygen-minimum zone off western Baja California linked to global climate over the past 52 kyr.  
 2413 *Paleoceanography*, **21**, PA4209, doi:10.1029/2005PA001239.
- 2414 Défarge, C. (2011). Organomineralization. Reitner, J., & Thiel, V. (eds.). *Encyclopedia of Geobiology*,  
 2415 Springer Verlag, pp.697-701, 2011, doi:10.1007/978-1-4020-9212-1\_159.

- 2416 Dehairs, F., Chesselet, R., & Jedwab, J. (1980). Discrete suspended particles of barite and the barium cycle  
 2417 in the open ocean. *Earth Planet. Sci. Lett.*, **49**(2), 528-550.
- 2418 Dehairs, F., Lambert, C. E., Chesselet, R., & Risler, N. (1987). The biological production of marine  
 2419 suspended barite and the barium cycle in the Western Mediterranean Sea. *Biogeochemistry*, **4**(2),  
 2420 119-140.
- 2421 Dehairs, F., Jacquet, S., Savoye, N., Van Mooy, B. A., Buesseler, K. O., Bishop, J. K., Lamborg, C. H.,  
 2422 Elskens, M., Baeyens, W., Boyd, P. W., Casciotti, K. L., & Monnin, C. (2008). Barium in twilight  
 2423 zone suspended matter as a potential proxy for particulate organic carbon remineralization: Results  
 2424 for the North Pacific. *Deep Sea Res. Pt. II*, **55**(14-15), 1673-1683.
- 2425 Dellwig, O., Beck, M., Lemke, A., Lunau, M., Kolditz, K., Schnetger, B., & Brumsack, H.-J. (2007). Non-  
 2426 conservative behaviour of molybdenum in coastal waters: Coupling geochemical, biological, and  
 2427 sedimentological processes. *Geochim. Cosmochim. Acta*, **71**, 2745–2761,  
 2428 doi:10.1016/j.gca.2007.03.014.
- 2429 Dellwig, O., Leipe, T., März, C., Glockzin, M., Pollehne, F., Schnetger, B., Yakushev, E. V., Böttcher, M.  
 2430 E., & Brumsack, H. J. (2010). A new particulate Mn-Fe-P-shuttle at the redoxcline of anoxic basins.  
 2431 *Geochim. Cosmochim. Acta*, **74**, 7100–7115, doi:10.1016/j.gca.2010.09.017.
- 2432 Deng, N., Stack, A. G., Weber, J., Cao, B., De Yoreo, J. J., & Hu, Y. (2019). Organic–mineral interfacial  
 2433 chemistry drives heterogeneous nucleation of Sr-rich (Ba<sub>x</sub>, Sr<sub>1-x</sub>)SO<sub>4</sub> from undersaturated solution.  
 2434 *Proc. Natl. Acad. Sci.*, 201821065.
- 2435 Dickens, G.R., Fewless, T., Thomas, E. & Bralower, T.J. (2003). Excess barite accumulation during the  
 2436 Paleocene–Eocene Thermal Maximum: Massive input of dissolved barium from seafloor gas  
 2437 hydrate reservoirs. In Wing, S. L, Gingerich, P. D., Schmitz, B., & Thomas, E. (Eds.) *Causes and  
 2438 consequences of globally warm climates in the early Paleogene*, **369**, 11-23, Geological Society of  
 2439 America Special Papers; doi:10.1130/0-8137-2369-8.11.
- 2440 Dickson, A. J. (2017). A molybdenum-isotope perspective on Phanerozoic deoxygenation events. *Nat.*  
 2441 *Geosci.*, **10**, 721–726, doi:10.1038/NGEO3028.
- 2442 Dideriksen, K., Baker, J. A., & Stipp, S. L. S. (2008). Equilibrium Fe isotope fractionation between  
 2443 inorganic aqueous Fe (III) and the siderophore complex, Fe (III)-desferrioxamine B. *Earth Planet.  
 2444 Sci. Lett.*, **269**(1-2), 280-290.
- 2445 Donat, J. R., & Bruland, K. W. (1990). A comparison of two voltammetric techniques for determining zinc  
 2446 speciation in Northeast Pacific Ocean waters. *Mar. Chem.*, **28**, 301–323.
- 2447 Donat J. R., Lao, K. A., & Bruland, K. W. (1994) Speciation of dissolved copper and nickel in South San  
 2448 Francisco Bay: a multi-method approach. *Anal. Chim. Acta* **284**, 547–571.
- 2449 Dunlea, A. G., Murray, R. W., Sauvage, J., Spivack, A. J., Harris, R. N., & D'Hondt, S. (2015). Dust,  
 2450 volcanic ash, and the evolution of the South Pacific Gyre through the Cenozoic. *Paleoceanogr.*,  
 2451 **30**(8), 1078-1099.
- 2452 Dugdale, R. C., & Goering, J. J. (1967). Uptake of new and regenerated forms of nitrogen in primary  
 2453 production. *Limnol. Oceanogr.*, **12**(2), 196–206.
- 2454 Dupont, C. L., Moffett, J. W., Bidigare, R. R., & Ahner, B. A. (2006). Distributions of dissolved and  
 2455 particulate biogenic thiols in the subarctic Pacific Ocean. *Deep. Res. Part I Oceanogr. Res. Pap.*,  
 2456 **53**, 1961–1974.
- 2457 Dupont, C. L., Butcher, A., Valas, R. E., Bourne, P. E., & Caetano-Anolles, G. (2010). History of biological  
 2458 metal utilization inferred through phylogenomic analysis of protein structures. *Proc. Natl. Acad.  
 2459 Sci.*, **107**, 10567–10572.



- 2460 Dymond, J., Suess, E., & Lyle, M. (1992). Barium in deep-sea sediment: A geochemical proxy for  
2461 paleoproductivity. *Paleoceanography*, **7**(2), 163-181.
- 2462 Dymond, J., & Collier, R. (1996). Particulate barium fluxes and their relationships to biological  
2463 productivity. *Deep Sea Research Part II: Topical Studies in Oceanography*, **43**(4-6), 1283-1308.
- 2464 Eagle, M., Paytan, A., Arrigo, K. R., van Dijken, G., & Murray, R. W. (2003). A comparison between  
2465 excess barium and barite as indicators of carbon export. *Paleoceanography*, **18**(1), 1021,  
2466 <https://doi.org/10.1029/2002PA000793>.
- 2467 Edmond, J. M., Boyle, E. D., Drummond, D., Grant, B., & Mislick, T. (1978). Desorption of barium in the  
2468 plume of the Zaire (Congo) River. *Netherlands Journal of Sea Research*, **12**(3-4), 324-328.
- 2469 Elderfield, H., & Rickaby, R. E. M. (2000). Oceanic Cd/P ratio and nutrient utilization in the glacial  
2470 Southern Ocean. *Nature*, **405**(6784), 305-310.
- 2471 Elliott, T., & Steele, R. C. (2017). The isotope geochemistry of Ni. *Reviews in Mineralogy and  
2472 Geochemistry*, **82**(1), 511-542.
- 2473 Ellwood, M. J. (2004). Zinc and cadmium speciation in subantarctic waters east of New Zealand. *Mar.  
2474 Chem.*, **87**(1-2), 37-58.
- 2475 Ellwood, M. J., & Hunter, K. A. (2000). The Incorporation of Zinc and Iron into the Frustule of the Marine  
2476 Diatom *Thalassiosira pseudonana*. *Limnol. Oceanogr.*, **45**, 1517-1524.
- 2477 Ellwood, M. J., & Van Den Berg, C. M. G. (2000). Zinc speciation in the Northeastern Atlantic Ocean.  
2478 *Mar. Chem.*, **68**, 295-306.
- 2479 Ellwood, M.J., Hutchins, D.A., Lohan, M.C., Milne, A., Nasemann, P., Nodder, S.D., Sander, S.G.,  
2480 Strzepek, R., Wilhelm, S.W., & Boyd, P.W. (2015). Iron stable isotopes track pelagic iron cycling  
2481 during a subtropical phytoplankton bloom. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E15-E20.
- 2482 Ellwood, M. J., Strzepek, R. F., Strutton, P. G., Trull, T. W., Fourquez, M., & Boyd, P. W. (2020). Distinct  
2483 iron cycling in a Southern Ocean eddy. *Nature Communications*, **11**(1), 1-8.
- 2484 Emerson, S. R., & Husted, S. S. (1991). Ocean anoxia and the concentrations of molybdenum and  
2485 vanadium in seawater. *Mar. Chem.* **34**, 177-196, doi:10.1016/0304-4203(91)90002-E.
- 2486 Erickson, B. E., & Helz, G. R. (2000). Molybdenum(VI) speciation in sulfidic waters: Stability and lability  
2487 of thiomolybdates. *Geochim. Cosmochim. Acta*, **64**, 1149-1158, doi:10.1016/S0016-  
2488 7037(99)00423-8.
- 2489 Escoube, R., Rouxel, O. J., Pokrovsky, O. S., Schroth, A., Max Holmes, R., & Donard, O. F. X. (2015).  
2490 Iron isotope systematics in Arctic rivers. *Comptes Rendus Geosci.*, **347**(7-8), 377-385.
- 2491 Escoube, R., Rouxel, O. J., Sholkovitz, E., & Donard, O. F. X. (2009). Iron isotope systematics in estuaries:  
2492 The case of North River, Massachusetts (USA). *Geochim. Cosmochim. Acta*, **73**(14), 4045-4059.
- 2493 Esser, B. K., & Volpe, A. M. (2002). At-sea high-resolution chemical mapping: extreme barium depletion  
2494 in North Pacific surface water. *Mar. Chem.*, **79**(2), 67-79.
- 2495 Fantle, M. S., & DePaolo, D. J., (2004). Iron isotopic fractionation during continental weathering. *Earth  
2496 Planet. Sci. Lett.*, **228**(3-4), 547-562.
- 2497 Farmer, J. R., Hönisch, B., Haynes, L. L., Kroon, D., Jung, S., Ford, H. L., Raymo, M. E., Jaume-Seguí,  
2498 M., Bell, D. B., Goldstein, S. L., Pena, L. D., Yehudai, M., & Kim, J. (2019). Deep Atlantic Ocean  
2499 carbon storage and the rise of 100,000-year glacial cycles. *Nat. Geosci.* **12**, 355-360,  
2500 <https://doi.org/10.1038/s41561-019-0334-6>.

- 2501 Fassbender, A. J., Palevsky, H. I., Martz, T. R., Ingalls, A. E., Gledhill, M., Fawcett, S. E., Brandes, J. A.,  
 2502 Aluwihare, L. I., & the participants of COME ABOARD and DISCO XXV (2017). Perspectives  
 2503 on Chemical Oceanography in the 21st century: Participants of the COME ABOARD Meeting  
 2504 examine aspects of the field in the context of 40 years of DISCO. *Mar. Chem.*, **196**, 181-190.
- 2505 Finlay, B. J., Hetherington, N. B., & Da Vison, W. (1983). Active biological participation in lacustrine  
 2506 barium chemistry. *Geochim. Cosmochim. Acta*, **47**(7), 1325-1329.
- 2507 Fischer, L., Smith, G., Hann, S., & Bruland, K. W. (2018). Ultra-trace analysis of silver and platinum in  
 2508 seawater by ICP-SFMS after off-line matrix separation and pre-concentration. *Mar. Chem.*, **199**,  
 2509 44-52.
- 2510 Fisher, N.S., & Wente, M. (1993). The release of trace elements by dying marine phytoplankton. *Deep-Sea  
 2511 Research I*, **40**(4), 671-694.
- 2512 Fishwick, M. P., Sedwick, P. N., Lohan, M. C., Worsfeld, P. J., Buck, K. N., Church, T. M., Ussher, S. J.  
 2513 (2014). The impact of changing surface ocean conditions on the dissolution of aerosol iron. *Global  
 2514 Biogeochem. Cycles*, **28**(11), 1235-1250. <https://doi.org/10.1002/2014GB004921>.
- 2515 Fitzsimmons, J. N., & Boyle, E. A. (2014). Both soluble and colloidal iron phases control dissolved iron  
 2516 variability in the tropical North Atlantic Ocean. *Geochim. Cosmochim. Acta*, **125**, 539-550.
- 2517 Fitzsimmons, J. N., Carrasco, G. G., Wu, J., Roshan, S., Hatta, M., Measures, C. I., Conway, T. M., John,  
 2518 S.G., & Boyle, E. A. (2015). Partitioning of dissolved iron and iron isotopes into soluble and  
 2519 colloidal phases along the GA03 GEOTRACES North Atlantic Transect. *Deep Sea Res. Pt. II*, **116**,  
 2520 130-151.
- 2521 Fitzsimmons, J. N., Conway, T. M., Lee, J.-M., Kayser, R., Thyng, K. M., John, S. G. and Boyle, E. A.  
 2522 (2016). Dissolved iron and iron isotopes in the Southeastern Pacific Ocean. *Global Biogeochem.  
 2523 Cycles*, **30**(10), 1372-1395. <https://doi.org/10.1002/2015GB005357>.
- 2524 Fitzsimmons, J. N., John, S. G., Marsay, C. M., Hoffman, C. L., Nicholas, S. L., Toner, B. M., German, C.  
 2525 R., & Sherrell, R. M. (2017). Iron persistence in a distal hydrothermal plume supported by  
 2526 dissolved-particulate exchange. *Nature Geoscience*, **10**(3), 195-201.
- 2527 Flegal, A.R., Sanudo-Wilhelmy, S.A., & Scelfo, G.M. (1995). Silver in the eastern Atlantic Ocean. *Mar.  
 2528 Chem.*, **49**, 315-320.
- 2529 Franck, V. M., Bruland, K. W., Hutchins, D. A., & Brzezinski, M. A. (2003). Iron and zinc effects on silicic  
 2530 acid and nitrate uptake kinetics in three high-nutrient, low-chlorophyll (HNLC) regions. *Mar. Ecol.  
 2531 Prog. Ser.*, **252**, 15-33.
- 2532 Francois, R., Honjo, S., Manganini, S. J., & Ravizza, G. E. (1995). Biogenic barium fluxes to the deep sea:  
 2533 Implications for paleoproductivity reconstruction. *Global Biogeochem. Cycles*, **9**(2), 289-303.
- 2534 Frew, R. D., & Hunter, K. A. (1992). Influence of Southern Ocean waters on the cadmium-phosphate  
 2535 properties of the global ocean. *Nature*, **360**(6400), 144.
- 2536 Friedl, G., & Pedersen, T. (2002). Silver as a new tracer for diatom production. *Eawag News*, **52**, 14-15.
- 2537 Froelich, P., Klinkhammer, G. P., Bender, M. L., Luedtke, N. A., Heath, G. R., Cullen, D., Dauphin, P.,  
 2538 Hammond D., Hartman, B., & Maynard, V. (1979). Early oxidation of organic matter in pelagic  
 2539 sediments of the eastern equatorial Atlantic: suboxic diagenesis. *Geochimica et Cosmochimica  
 2540 Acta*, **43**(7), 1075-1090.
- 2541 Fru, E. C., Rodríguez, N. P., Partin, C. A., Lalonde, S. V., Andersson, P., Weiss, D. J., El Albani, A.,  
 2542 Rodushkin, I., & Konhauser, K. O. (2016). Cu isotopes in marine black shales record the Great  
 2543 Oxidation Event. *Proceedings of the National Academy of Sciences*, **113**(18), 4941-4946.

- 2544 Fujii, T., Moynier, F., Pons, M. L., & Albarède, F. (2011a). The origin of Zn isotope fractionation in  
2545 sulfides. *Geochim. Cosmochim. Acta*, **75**(23), 7632-7643.
- 2546 Fujii, T., Moynier, F., Dauphas, N., & Abe, M. (2011b). Theoretical and experimental investigation of  
2547 nickel isotopic fractionation in species relevant to modern and ancient oceans. *Geochim.*  
2548 *Cosmochim. Acta*, **75**(2), 469-482.
- 2549 Fujii, T., Moynier, F., Abe, M., Nemoto, K., & Albarède, F. (2013). Copper isotope fractionation between  
2550 aqueous compounds relevant to low temperature geochemistry and biology. *Geochim. Cosmochim.*  
2551 *Acta*, **110**, 29-44.
- 2552 Gall, L., Williams, H. M., Siebert, C., Halliday, A. N., Herrington, R. J., & Hein, J. R. (2013). Nickel  
2553 isotopic compositions of ferromanganese crusts and the constancy of deep ocean inputs and  
2554 continental weathering effects over the Cenozoic. *Earth Planet. Sci. Lett.*, **375**, 148-155.
- 2555 Gallon, C., & Flegal, A. R. (2015). Sources, Fluxes, and Biogeochemical Cycling of Silver in the Oceans.  
2556 In: Whitacre, D. (eds) *Rev. Environ. Contamin. Toxicol.* **235**. Springer, Cham.  
2557 [https://doi.org/10.1007/978-3-319-10861-2\\_2](https://doi.org/10.1007/978-3-319-10861-2_2).
- 2558 Ganeshram, R. S., François, R., Commeau, J., & Brown-Leger, S. L. (2003). An experimental investigation  
2559 of barite formation in seawater. *Geochim. Cosmochim. Acta*, **67**(14), 2599-2605.
- 2560 Garcia, H. E., R. A. Locarnini, T. P. Boyer, J. I. Antonov, O.K. Baranova, M.M. Zweng, J.R. Reagan, D.R.  
2561 Johnson, 2014. World Ocean Atlas 2013, Volume 3: Dissolved Oxygen, Apparent Oxygen  
2562 Utilization, and Oxygen Saturation. S. Levitus, Ed., A. Mishonov Technical Ed.; *NOAA Atlas*  
2563 *NESDIS 75*, 27 pp.
- 2564 Gault-Ringold, M., Adu, T., Stirling, C. H., Frew, R. D., & Hunter, K. A. (2012). Anomalous  
2565 biogeochemical behavior of cadmium in Subantarctic surface waters: mechanistic constraints from  
2566 cadmium isotopes. *Earth Planet. Sci. Lett.*, **341**, 94-103.
- 2567 George, E., Stirling, C. H., Gault-Ringold, M., Ellwood, M. J., & Middag, R. (2019). Marine  
2568 biogeochemical cycling of cadmium and cadmium isotopes in the extreme nutrient-depleted  
2569 subtropical gyre of the South West Pacific Ocean. *Earth Planet. Sci. Lett.*, **514**, 84-95.
- 2570 Georgiev, S. V., Horner, T. J., Stein, H. J., Hannah, J. L., Bingen, B., & Rehkämper, M. (2015). Cadmium-  
2571 isotopic evidence for increasing primary productivity during the Late Permian anoxic event. *Earth*  
2572 *Planet. Sci. Lett.*, **410**, 84-96.
- 2573 Geyman, B. M., Ptacek, J. L., LaVigne, M., & Horner, T. J. (2019). Barium in deep-sea bamboo corals:  
2574 Phase associations, barium stable isotopes, & prospects for paleoceanography. *Earth Planet. Sci.*  
2575 *Lett.*, **525**, 115751.
- 2576 Giering, S. L., Sanders, R., Lampitt, R. S., Anderson, T. R., Tamburini, C., Boutrif, M., Zubkov, M. V.,  
2577 Marsay, C. M., Henson, S. A., Saw, K., Cook, K., & Mayor, D. J.(2014). Reconciliation of the  
2578 carbon budget in the ocean's twilight zone. *Nature*, **507**(7493), 480-483.
- 2579 Glass, J. B., Axler, R. P., Chandra, S., & Goldman, C. R. (2012). Molybdenum limitation of microbial  
2580 nitrogen assimilation in aquatic ecosystems and pure cultures. *Frontiers in Microbiology*, **3**, 1–11,  
2581 doi:10.3389/fmicb.2012.00331.
- 2582 Glass, J. B., Wolfe-Simon, F., & Anbar, A. D. (2009). Coevolution of metal availability and nitrogen  
2583 assimilation in cyanobacteria and algae. *Geobiology*, **7**, 100–123, doi:10.1111/j.1472-  
2584 4669.2009.00190.x.
- 2585 Goldberg, T., Archer, C., Vance, D., & Poulton, S. W. (2009). Mo isotope fractionation during adsorption  
2586 to Fe (oxyhydr)oxides. *Geochim. Cosmochim. Acta*, **73**, 6502–6516,  
2587 doi:10.1016/j.gca.2009.08.004.



- 2588 Goldberg, T., Archer, C., Vance, D., Thamdrup, B., McAnena, A., & Poulton, S. W. (2012). Controls on  
 2589 Mo isotope fractionations in a Mn-rich anoxic marine sediment, Gullmar Fjord, Sweden. *Chem.*  
 2590 *Geol.*, **296-297**, 73-82, doi:10.1016/j.chemgeo.2011.12.020.
- 2591 Goldberg, T., Gordon, G., Izon, G., Archer, C., Pearce, C. R., McManus, J., Anbar, A. D., & Rehkämper,  
 2592 M. (2013). Resolution of inter-laboratory discrepancies in Mo isotope data: an intercalibration.  
 2593 *Journal of Analytical Atomic Spectrometry*, **28**(5), 724-735.
- 2594 Goldman, C. R. (1960). Molybdenum as a Factor Limiting Primary Productivity in Castle Lake, California.  
 2595 *Science*, **132**, 1016–1017, doi:10.1126/science.132.3433.1016.
- 2596 Gooday, A. J., & Nott, J. A. (1982). Intracellular barite crystals in two xenophyophores, *Aschemonella*  
 2597 *ramuliformis* and *Galatheammima* sp. (Protozoa: Rhizopoda) with comments on the taxonomy of  
 2598 *A. ramuliformis*. *Journal of the Marine Biological Association of the United Kingdom*, **62**(3), 595-  
 2599 605.
- 2600 Goto, K. T., Sekine, Y., Shimoda, G., Hein, J. R., Aoki, S., Ishikawa, A., Suzuki, K., Gordon, G. W., &  
 2601 Anbar, A. D. (2020). A framework for understanding Mo isotope records of Archean and  
 2602 Paleoproterozoic Fe-and Mn-rich sedimentary rocks: Insights from modern marine hydrothermal  
 2603 Fe-Mn oxides. *Geochim. Cosmochim. Acta*, **280**, 221-236.
- 2604 Gou, L. F., Jin, Z., Galy, A., Gong, Y. Z., Nan, X. Y., Jin, C., Wang, X.-D., Bouchez, J., Cai, H.-M., Chen,  
 2605 J.-B., Yu, H.-M., & Huang, F. (2020). Seasonal riverine barium isotopic variation in the middle  
 2606 Yellow River: Sources and fractionation. *Earth and Planetary Science Letters*, **531**, 115990.
- 2607 Gueguen, B., Rouxel, O., Ponzevera, E., Bekker, A., & Fouquet, Y. (2013). Nickel isotope variations in  
 2608 terrestrial silicate rocks and geological reference materials measured by MC-ICP-MS.  
 2609 *Geostandards and Geoanalytical Research*, **37**(3), 297-317.
- 2610 Gueguen, B., Rouxel, O., Rouget, M. L., Bollinger, C., Ponzevera, E., Germain, Y., & Fouquet, Y. (2016).  
 2611 Comparative geochemistry of four ferromanganese crusts from the Pacific Ocean and significance  
 2612 for the use of Ni isotopes as paleoceanographic tracers. *Geochim. Cosmochim. Acta*, **189**, 214-235.
- 2613 Guinoiseau, D., Galer, S. J., & Abouchami, W. (2018). Effect of cadmium sulphide precipitation on the  
 2614 partitioning of Cd isotopes: Implications for the oceanic Cd cycle. *Earth Planet. Sci. Lett.*, **498**,  
 2615 300-308.
- 2616 Guo, J., Lapi, S., Ruth, T. J., & Maldonado, M. T. (2012). The effects of iron and copper availability on the  
 2617 copper stoichiometry of marine phytoplankton. *J. Phycol.* **48**, 312–325.
- 2618 Hanor, J. S., & Chan, L. H. (1977). Non-conservative behavior of barium during mixing of Mississippi  
 2619 River and Gulf of Mexico waters. *Earth and Planetary Science Letters*, **37**(2), 242-250.
- 2620 Hanor, J. S. (2000). Barite–celestine geochemistry and environments of formation. *Reviews in Mineralogy*  
 2621 *and Geochemistry*, **40**(1), 193-275.
- 2622 Hardisty, D. S., Riedinger, N., Planavsky, N. J., Asael, D., Andrén, T., et al. (2016). A Holocene history of  
 2623 dynamic water column redox conditions in the Landsort Deep, Baltic Sea. *Am. J. Sci.*, **316**, 713-  
 2624 745, doi:10.2475/08.2016.01.
- 2625 Hardisty, D. S., Lyons, T. W., Riedinger, N., Isson, T. T., Owens, J. D., Aller, R. C., Rye, D. M., Planavsky,  
 2626 N. J., Reinhard, C. T., Gill, B. C., Masterson, A. L., Asael, D., & Johnston, D. T. (2018). An  
 2627 evaluation of sedimentary molybdenum and iron as proxies for pore fluid paleoredox conditions.  
 2628 *Am. J. Sci.*, **318**(5), 527-556.
- 2629 Hartnett, H. E., Keil, R. G., Hedges, J. I., & Devol, A. H. (1998). Influence of oxygen exposure time on  
 2630 organic carbon preservation in continental margin sediments. *Nature*, **391**(6667), 572-575.

- 2631 Hawco, N. J., Lam, P. J., Lee, J. M., Ohnemus, D. C., Noble, A. E., Wyatt, N. J., Lohan, M. C., & Saito,  
 2632 M. A. (2018). Cobalt scavenging in the mesopelagic ocean and its influence on global mass  
 2633 balance: Synthesizing water column and sedimentary fluxes. *Marine Chemistry*, **201**, 151-166.
- 2634 Hayes, C. T., Anderson, R. F., Cheng, H., Conway, T. M., Edwards, R. L., Fleisher, M. Q., Ho, P., Huang,  
 2635 K.-F., John, S. G., Landing, W. M., Little, S. H., Lu, Y., Morton, P. L., Moran, S. B., Robinson, L.  
 2636 F., Shelley, R. U., Shiller, A. M., & Zheng, X.-Y. (2018). Replacement times of a spectrum of  
 2637 elements in the North Atlantic based on thorium supply. *Global Biogeochem. Cycles*, **32**(9), 1294-  
 2638 1311.
- 2639 Heller, M. I., & Croot, P. L. (2015). Copper speciation and distribution in the Atlantic sector of the Southern  
 2640 Ocean. *Mar. Chem.*, **173**, 253-268.
- 2641 Helz, G. R., Miller, C. V., Charnock, J. M., Mosselmans, J. F. W., Patrick, R. A. D., Garner, C. D., &  
 2642 Vaughan, D. J. (1996). Mechanism of molybdenum removal from the sea and its concentration in  
 2643 black shales: EXAFS evidence. *Geochim. Cosmochim. Acta*, **60**, 3631–3642, doi:10.1016/0016-  
 2644 7037(96)00195-0.
- 2645 Hemsing, F., Hsieh, Y. T., Bridgestock, L., Spooner, P. T., Robinson, L. F., Frank, N., & Henderson, G.  
 2646 M. (2018). Barium isotopes in cold-water corals. *Earth and Planetary Science Letters*, **491**, 183-  
 2647 192.
- 2648 Henderson, G. M., & Burton, K. W. (1999). Using (234U/238U) to assess diffusion rates of isotope tracers  
 2649 in ferromanganese crusts. *Earth Planet. Sci. Lett.*, **170**(3), 169-179.
- 2650 Hendry, K. R., & Andersen, M. B. (2013). The zinc isotopic composition of siliceous marine sponges:  
 2651 Investigating nature's sediment traps. *Chem. Geol.*, **354**, 33–41. Available at:  
 2652 <http://dx.doi.org/10.1016/j.chemgeo.2013.06.025>.
- 2653 Hendry, K. R., Pyle, K. M., Barney Butler, G., Cooper, A., Fransson, A., Chierici, M., Leng, M. J., Meyer,  
 2654 A., & Dodd, P.A. (2018). Spatiotemporal Variability of Barium in Arctic Sea-Ice and Seawater. *J.*  
 2655 *Geophys. Res. Oceans*, **123**(5), 3507-3522.
- 2656 Hendy, I. L., & Pedersen, T. F. (2005). Is pore water oxygen content decoupled from productivity on the  
 2657 California Margin? Trace element results from Ocean Drilling Program Hole 1017E, San Lucia  
 2658 slope, California. *Paleoceanography*, **20**(4).
- 2659 Henkel, S., Kasten, S., Hartmann, J. F., Silva-Busso, A., & Staubwasser, M. (2018). Iron cycling and stable  
 2660 Fe isotope fractionation in Antarctic shelf sediments, King George Island. *Geochim. Cosmochim.*  
 2661 *Acta*, **237**, 320-338. <https://doi.org/10.1016/j.gca.2018.06.042>.
- 2662 Hillaire-Marcel, C., & De Vernal, A. (Eds.). (2007). *Proxies in Late Cenozoic Paleooceanography*. Elsevier.
- 2663 Ho, P., Lee, J. M., Heller, M. I., Lam, P. J., and Shiller, A. M. (2018). The distribution of dissolved and  
 2664 particulate Mo and V along the U.S. GEOTRACES East Pacific Zonal Transect (GP16): The roles  
 2665 of oxides and biogenic particles in their distributions in the oxygen deficient zone and the  
 2666 hydrothermal plume. *Mar. Chem.*, **201**, 242–255, doi:10.1016/j.marchem.2017.12.003.
- 2667 Ho, T. Y., Quigg, A., Finkel, Z. V., Milligan, A. J., Wyman, K., Falkowski, P. G., & Morel, F. M. (2003).  
 2668 The elemental composition of some marine phytoplankton 1. *Journal of Phycology*, **39**(6), 1145-  
 2669 1159.
- 2670 Hodgskiss, M. S., Crockford, P. W., Peng, Y., Wing, B. A., & Horner, T. J. (2019). A productivity collapse  
 2671 to end Earth's Great Oxidation. *Proc. Natl. Acad. Sci.*, **116**(35), 17207-17212.
- 2672 Hohl, S. V., Galer, S. J. G., Gamper, A., & Becker, H. (2017). Cadmium isotope variations in  
 2673 Neoproterozoic carbonates—A tracer of biologic production?. *Geochemical Perspective Letters*, **3**,  
 2674 32-44.

- 2675 Hohl, S. V., Jiang, S. Y., Wei, H. Z., Pi, D. H., Liu, Q., Viehmann, S., & Galer, S. J. (2019). Cd Isotopes  
 2676 trace periodic (bio)geochemical metal cycling at the verge of the Cambrian animal evolution.  
 2677 *Geochim. Cosmochim. Acta*, **263**, 195-214, doi:10.1016/j.gca.2019.07.036.
- 2678 Holzer, M., Primeau, F. W., DeVries, T., & Matear, R. (2014). The Southern Ocean silicon trap: Data-  
 2679 constrained estimates of regenerated silicic acid, trapping efficiencies, and global transport paths.  
 2680 *J. Geophys. Res. Oceans*, **119**(1), 313-331.
- 2681 Homoky, W. B., Severmann, S., Mills, R. A., & Fones, G. R. (2009). Pore-fluid Fe isotopes reflect the  
 2682 extent of benthic Fe redox recycling: evidence from continental shelf and deep-sea sediments.  
 2683 *Geology*, **37**, 751-754. <https://doi.org/10.1130/G25731A.1>.
- 2684 Homoky, W. B., John, S. G., Conway T. M., & Mills, R. A. (2013). Distinct iron isotopic signatures and  
 2685 supply from marine sediment dissolution. *Nat. Comms*, **4**, 2143.  
 2686 <https://doi.org/10.1038/ncomms3143>.
- 2687 Hoppema, M., Dehairs, F., Navez, J., Monnin, C., Jeandel, C., Fahrbach, E., & De Baar, H. J. W. (2010).  
 2688 Distribution of barium in the Weddell Gyre: Impact of circulation and biogeochemical processes.  
 2689 *Marine Chemistry*, **122**(1-4), 118-129.
- 2690 Horner, T. J., Schönbacher, M., Rehkämper, M., Nielsen, S. G., Williams, H., Halliday, A. N., Xue, Z., &  
 2691 Hein, J. R. (2010). Ferromanganese crusts as archives of deep water Cd isotope compositions.  
 2692 *Geochemistry, Geophysics, Geosystems*, **11**(4), Q04001, <https://doi.org/10.1029/2009GC002987>.
- 2693 Horner, T. J., Rickaby, R. E., & Henderson, G. M. (2011). Isotopic fractionation of cadmium into calcite.  
 2694 *Earth Planet. Sci. Lett.*, **312**(1-2), 243-253.
- 2695 Horner, T. J., Lee, R. B., Henderson, G. M., & Rickaby, R. E. (2013). Nonspecific uptake and homeostasis  
 2696 drive the oceanic cadmium cycle. *Proc. Natl. Acad. Sci.*, **110**(7), 2500-2505.
- 2697 Horner, T. J., Kinsley, C. W., & Nielsen, S. G. (2015a). Barium-isotopic fractionation in seawater mediated  
 2698 by barite cycling and oceanic circulation. *Earth Planet. Sci. Lett.*, **430**, 511-522.
- 2699 Horner, T. J., Williams, H. M., Hein, J. R., Saito, M. A., Burton, K. W., Halliday, A. N., & Nielsen, S. G.  
 2700 (2015b). Persistence of deeply sourced iron in the Pacific Ocean. *Proc. Natl. Acad. Sci.*, **112**, 1292-  
 2701 1297.
- 2702 Horner, T. J., Pryer, H. V., Nielsen, S. G., Crockford, P. W., Gauglitz, J. M., Wing, B. A., & Ricketts, R.  
 2703 D. (2017). Pelagic barite precipitation at micromolar ambient sulfate. *Nat. Commun.*, **8**(1), 1342.
- 2704 Hsieh, Y. T., & Henderson, G. M. (2017). Barium stable isotopes in the global ocean: tracer of Ba inputs  
 2705 and utilization. *Earth Planet. Sci. Lett.*, **473**, 269-278.
- 2706 Hudson, R. J. M., & Morel, F. M. M. (1993). Trace metal transport by marine microorganisms: Implications  
 2707 of metal coordination kinetics. *Deep Sea Res.* **40**, 129-150.
- 2708 Ijichi, Y., Ohno, T., & Sakata, S. (2018). Copper isotopic fractionation during adsorption on manganese  
 2709 oxide: Effects of pH and desorption. *Geochemical Journal*, **52**(2), e1-e6.
- 2710 Ilina, S. M., Poitrasson, F., Lapitskiy, S. A., Alekhin, Y. V., Viers, J., & Pokrovsky, O. S. (2013). Extreme  
 2711 iron isotope fractionation between colloids and particles of boreal and temperate organic-rich  
 2712 waters. *Geochim. et Cosmochim. Acta*, **101**, 96-111.
- 2713 Ingri, J., Malinovsky, D., Rodushkin, I., Baxter, D.C., Widerlund, A., Andersson, P., Gustafsson, Ö.,  
 2714 Forsling, W. & Öhlander, B. (2006). Iron isotope fractionation in river colloidal matter. *Earth  
 2715 Planet. Sci. Lett.*, **245**(3-4), 792-798.
- 2716 Isson, T. T., Love, G. D., Dupont, C. L., Reinhard, C. T., Zumberge, A. J., Asael, D., Gueguen, B., McCrow,  
 2717 J., Gill, B. C., Owens, J., Rainbird, R. H., Rooney, A. D., Zhao, M. Y., Stueeken, E. E., Konhauser,

- 2718 K. O., John, S. G., Lyons, T. W., & Planavsky, N. J. (2018). Tracking the rise of eukaryotes to  
2719 ecological dominance with zinc isotopes. *Geobiology*, **16**, 341–352.
- 2720 Ivanochko, T. S. (2001). Productivity influences on oxygenation of the Santa Barbara Basin, California,  
2721 during the Late Quaternary (Doctoral dissertation, University of British Columbia).
- 2722 Jaccard, S. L., Haug, G. H., Sigman, D. M., Pedersen, T. F., Thierstein, H. R., & Röhl, U. (2005).  
2723 Glacial/interglacial changes in subarctic North Pacific stratification. *Science*, **308**(5724), 1003-  
2724 1006.
- 2725 Jaccard, S. L., Hayes, C. T., Martínez-García, A., Hodell, D. A., Anderson, R. F., Sigman, D. M., & Haug,  
2726 G. H. (2013). Two modes of change in Southern Ocean productivity over the past million years.  
2727 *Science*, **339**(6126), 1419-1423.
- 2728 Jacquet, S. H., Dehairs, F., Elskens, M., Savoye, N., & Cardinal, D. (2007). Barium cycling along WOCE  
2729 SR3 line in the Southern Ocean. *Mar. Chem.*, **106**(1-2), 33-45.
- 2730 Jacquot, J. E., & Moffett, J. W. (2015). Copper distribution and speciation across the International  
2731 GEOTRACES Section GA03. *Deep Sea Res. Pt. II*, **116**, 187-207, doi:10.1016/j.dsr2.2014.11.013.
- 2732 Jakuba, R. W., Saito, M. A., Moffett, J. W., & Xu, Y. (2012). Dissolved zinc in the subarctic North Pacific  
2733 and Bering Sea: Its distribution, speciation, and importance to primary producers. *Global  
2734 Biogeochem. Cycles*, **26**, 1–15.
- 2735 Janssen, D. J., & Cullen, J. T. (2015). Decoupling of zinc and silicic acid in the subarctic northeast Pacific  
2736 interior. *Mar. Chem.*, **177**, 124–133. <http://dx.doi.org/10.1016/j.marchem.2015.03.014>.
- 2737 Janssen, D. J., Conway, T. M., John, S. G., Christian, J. R., Kramer, D. I., Pedersen, T. F., & Cullen, J. T.  
2738 (2014). Undocumented water column sink for cadmium in open ocean oxygen-deficient zones.  
2739 *Proc. Natl. Acad. Sci.*, **111**(19), 6888-6893.
- 2740 Janssen, D. J., Abouchami, W., Galer, S. J. G., & Cullen, J. T. (2017). Fine-scale spatial and interannual  
2741 cadmium isotope variability in the subarctic northeast Pacific. *Earth Planet. Sci. Lett.*, **472**, 241-  
2742 252.
- 2743 Janssen, D. J., Abouchami, W., Galer, S. J., Purdon, K. B., & Cullen, J. T. (2019). Particulate cadmium  
2744 stable isotopes in the subarctic northeast Pacific reveal dynamic Cd cycling and a new isotopically  
2745 light Cd sink. *Earth Planet. Sci. Lett.*, **515**, 67-78.
- 2746 John, S. G., & Conway, T. M. (2014). A role for scavenging in the marine biogeochemical cycling of zinc  
2747 and zinc isotopes. *Earth Planet. Sci. Lett.*, **394**, 159-167.
- 2748 John, S. G., Geis R. W., Saito M. A., & Boyle E. A. (2007). Zinc isotope fractionation during high affinity  
2749 and low-affinity zinc transport by the marine diatom *Thalassiosira oceanica*. *Limnol. Oceanogr.*,  
2750 **52**, 2710–2714.
- 2751 John, S. G., Mendez, J., Moffett, J., & Adkins, J. (2012). The flux of iron and iron isotopes from San Pedro  
2752 Basin sediments. *Geochim. Cosmochim. Acta*, **93**, 14-29.
- 2753 John, S. G., Kunzmann, M., Townsend, E. J., & Rosenberg, A. D. (2017). Zinc and cadmium stable isotopes  
2754 in the geological record: A case study from the post-snowball Earth Nuccaleena cap dolostone.  
2755 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **466**, 202-208.
- 2756 John, S. G., Helgoe, J., Townsend, E., Weber, T., DeVries, T., Tagliabue, A., Moore, K., Lam, P., Marsay,  
2757 C.M., & Till, C. (2018a). Biogeochemical cycling of Fe and Fe stable isotopes in the Eastern  
2758 Tropical South Pacific. *Mar. Chem.* **201**, 66-76.
- 2759 John, S. G., Helgoe, J., & Townsend, E. (2018b). Biogeochemical cycling of Zn and Cd and their stable  
2760 isotopes in the Eastern Tropical South Pacific. *Mar. Chem.*, **201**, 256-262.



- 2761 Johnson, C.M., Skulan, J.L., Beard, B. L., Sun, H., Neelson, K. H., & Braterman, P. S. (2002). Isotopic  
 2762 fractionation between Fe(III) and Fe(II) in aqueous solutions. *Earth. Planet. Sci. Lett.*, **195**, 141–  
 2763 153. [https://doi.org/10.1016/S0012-821X\(01\)00581-7](https://doi.org/10.1016/S0012-821X(01)00581-7).
- 2764 Johnson, C. M., Beard, B. L., & Roden, E. E. (2008). The iron isotope fingerprints of redox and  
 2765 biogeochemical cycling in modern and ancient Earth. *Annu. Rev. Earth Planet. Sci.*, **36**, 457–493.
- 2766 Johnson, C. M., Beard, B. L., & Weyer, S. (2020). *Iron Geochemistry: An Isotopic Perspective*. Springer.  
 2767 doi:10.1007/978-3-030-33828-2.
- 2768 Joung, D. J., and Shiller, A. M. (2016). Temporal and spatial variations of dissolved and colloidal trace  
 2769 elements in Louisiana Shelf waters. *Mar. Chem.* **181**, 25–43, doi:10.1016/j.marchem.2016.03.003.
- 2770 Jullion, L., Jacquet, S. H. M., & Tanhua, T. (2017). Untangling biogeochemical processes from the impact  
 2771 of ocean circulation: First insight on the Mediterranean dissolved barium dynamics. *Global  
 2772 Biogeochem. Cycles*, **31**(8), 1256–1270.
- 2773 Kellogg, M. M., McIlvin, M. R., Vedamati, J., Twining, B. S., Moffett, J. W., Marchetti, A., Moran, D. M.,  
 2774 & Saito, M. A. (2020). Efficient zinc/cobalt inter-replacement in northeast Pacific diatoms and  
 2775 relationship to high surface dissolved Co: Zn ratios. *Limnol. Oceanogr.* doi:10.1002/lno.11471.
- 2776 Kendall, B., Komiya, T., Lyons, T. W., Bates, S. M., Gordon, G. W., Romaniello, S. J., Jiang, G., Creaser,  
 2777 R. A., Xiao, S., McFadden, K., Sawaki, Y., Tahata, M., Shu, D., Han, J., Li, Y., Chu, X., & Anbar,  
 2778 A. D. (2015). Uranium and molybdenum isotope evidence for an episode of widespread ocean  
 2779 oxygenation during the late ediacaran period. *Geochim. Cosmochim. Acta*, **156**, 173–193,  
 2780 doi:10.1016/j.gca.2015.02.025.
- 2781 Kendall, B., Dahl, T. W., & Anbar, A. D. (2017). The Stable Isotope Geochemistry of Molybdenum. *Rev.  
 2782 Min. Geochem.*, **82**, 683–732, doi:10.2138/rmg.2017.82.16.
- 2783 Khatiwala, S., Schmittner, A., & Muglia, J. (2019). Air-sea disequilibrium enhances ocean carbon storage  
 2784 during glacial periods. *Science Advances*, **5**(6), eaaw4981.
- 2785 Kienast, S. (2003). Palaeoceanography of the mid-latitude North East Pacific: during the late Pleistocene  
 2786 (Doctoral dissertation, University of British Columbia).
- 2787 Kim, T., Obata, H., Kondo, Y., Ogawa, H., & Gamo, T. (2015). Distribution and speciation of dissolved  
 2788 zinc in the western North Pacific and its adjacent seas. *Mar. Chem.*, **173**, 330–341.  
 2789 <http://dx.doi.org/10.1016/j.marchem.2014.10.016>.
- 2790 King, E. K., Thompson, A., Hodges, C., & Pett-Ridge, J. C. (2014). Towards Understanding Temporal and  
 2791 Spatial Patterns of Molybdenum in the Critical Zone. *Procedia Earth and Planetary Science*, **10**,  
 2792 56–62, doi:10.1016/j.proeps.2014.08.011.
- 2793 King, E. K., Thompson, A., Chadwick, O. A., & Pett-Ridge, J. C. (2016). Molybdenum sources and isotopic  
 2794 composition during early stages of pedogenesis along a basaltic climate transect. *Chem. Geol.*, **445**,  
 2795 54–67, doi:10.1016/j.chemgeo.2016.01.024.
- 2796 King, E. K., Perakis, S. S., & Pett-Ridge, J. C. (2018). Molybdenum isotope fractionation during adsorption  
 2797 to organic matter. *Geochim. Cosmochim. Acta*, **222**, 584–598, doi:10.1016/j.gca.2017.11.014.
- 2798 Klar, J. K., James, R. H., Gibbs, D., Lough, A., Parkinson, I., Milton, J. A., Hawkes, J. A., & Connelly, D.  
 2799 P. (2017a). Isotopic signature of dissolved iron delivered to the Southern Ocean from hydrothermal  
 2800 vents in the East Scotia Sea. *Geology*, **45**(4), 351–354. <https://doi.org/10.1130/G38432.1>.
- 2801 Klar, J. K., Homoky, W. B., Statham, P. J., Birchill, A. J., Harris, E. L., Woodward, E. M. S., Silburn, B.,  
 2802 Cooper, M. J., James, R. H., Connelly, D. P., Chever, F., Lichtschlag, A., & Graves, C. (2017b).  
 2803 Stability of dissolved and soluble Fe(II) in shelf sediment pore waters and release to an oxic water  
 2804 column. *Biogeochem.*, **135**, 1–19. <https://doi.org/10.1007/s10533-017-0309-x>.

- 2805 Klar, J. K., Schlosser, C., Milton, J. A., Woodward, E. M. S., Lacan, F., Parkinson, I. J., Achterberg, E., &  
 2806 James, R. H. (2018). Sources of dissolved iron to oxygen minimum zone waters on the Senegalese  
 2807 continental margin in the tropical North Atlantic Ocean: Insights from iron isotopes. *Geochim.*  
 2808 *Cosmochim. Acta*, **236**, 60-78. <https://doi.org/10.1016/j.gca.2018.02.031>.
- 2809 Klinkhammer, G. P. (1980). Early diagenesis in sediments from the eastern equatorial Pacific, II. Pore water  
 2810 metal results. *Earth Planet. Sci. Lett.*, **49**(1), 81-101.
- 2811 Klunder, M. B., Laan, P., Middag, R., De Baar, H. J. W., & van Ooijen, J. C. (2011). Dissolved iron in the  
 2812 Southern Ocean (Atlantic sector). *Deep Sea Res. Pt. II*, **58**, 2678-2694,  
 2813 doi:10.1016/j.dsr2.2010.10.042.
- 2814 Köbberich, M., & Vance, D. (2017). Kinetic control on Zn isotope signatures recorded in marine diatoms.  
 2815 *Geochim. Cosmochim. Acta*, **210**, 97–113. <http://dx.doi.org/10.1016/j.gca.2017.04.014>.
- 2816 Köbberich, M., & Vance, D. (2019). Zn isotope fractionation during uptake into marine phytoplankton:  
 2817 Implications for oceanic zinc isotopes. *Chem. Geol.*, **523**, 154-161,  
 2818 doi:10.1016/j.chemgeo.2019.04.004.
- 2819 Koide, M., Hodge, V.F., Yang, J.S., Stallard, M., & Goldberg, E.G. (1986). Some comparative marine  
 2820 chemistries of rhenium, gold, silver and molybdenum. *Appl. Geochem.*, **1**, 705-714.
- 2821 Konhauser, K. O., Pecoits, E., Lalonde, S. V., Papineau, D., Nisbet, E. G., Barley, M. E., Arndt, N. T.,  
 2822 Zahnle, K., & Kamber, B. S. (2009). Oceanic nickel depletion and a methanogen famine before the  
 2823 Great Oxidation Event. *Nature*, **458**(7239), 750-753.
- 2824 Koschinsky, A., & Halbach, P. (1995). Sequential leaching of marine ferromanganese precipitates: Genetic  
 2825 implications. *Geochim. Cosmochim. Acta*, **59**(24), 5113-5132.
- 2826 Koschinsky, A., & Hein, J. R. (2003). Uptake of elements from seawater by ferromanganese crusts: solid-  
 2827 phase associations and seawater speciation. *Mar. Geol.*, **198**(3-4), 331-351.
- 2828 Kowalski, N., Dellwig, O., Beck, M., Gräwe, U., Neubert, N., Nägler, T. F., Badewien, T. H., Brumsack,  
 2829 H. J., van Beusekom, J. E. E., & Böttcher, M. E. (2013). Pelagic molybdenum concentration  
 2830 anomalies and the impact of sediment resuspension on the molybdenum budget in two tidal systems  
 2831 of the North Sea. *Geochim. Cosmochim. Acta*, **119**, 198–211, doi:10.1016/j.gca.2013.05.046.
- 2832 Kramer, D., Cullen, J. T., Christian, J. R., Johnson, W. K., & Pedersen, T. F. (2011). Silver in the subarctic  
 2833 northeast Pacific Ocean: Explaining the basin scale distribution of silver. *Mar. Chem.*, **123**, 133-  
 2834 142.
- 2835 Kurisu, M., Takahashi, Y., Iizuka, T. & Uematsu, M. (2016a). Very low isotope ratio of iron in fine aerosols  
 2836 related to its contribution to the surface ocean. *J. Geophys. Res. Atmos.* **121**, 11,119–11,136.  
 2837 <https://doi.org/10.1002/2016JD024957>.
- 2838 Kurisu, M., Sakata, K., Miyamoto, C., Takaku, Y., Iizuka, T., & Takahashi, Y. (2016b). Variation of iron  
 2839 isotope ratios in anthropogenic materials emitted through combustion processes. *Chem. Lett.* **45**,  
 2840 970–972 (2016). <https://doi.org/10.1246/cl.160451>.
- 2841 Kunzmann, M., Halverson, G. P., Sossi, P. A., Raub, T. D., Payne, J. L., & Kirby, J. (2013). Zn isotope  
 2842 evidence for immediate resumption of primary productivity after snowball Earth. *Geology*, **41**, 27–  
 2843 30.
- 2844 Labatut, M., Lacan, F., Pradoux, C., Chmeleff, J., Radic, A., Murray, J. W., Poitrasson, F., Johansen, A.  
 2845 M., & Thil, F. (2014). Iron sources and dissolved-particulate interactions in the seawater of the  
 2846 Western Equatorial Pacific, iron isotope perspectives. *Global Biogeochem. Cycles* **28**, 1044-1065.
- 2847 Lacan, F., Francois, R., Ji, Y., & Sherrell, R. M. (2006). Cadmium isotopic composition in the ocean.  
 2848 *Geochim. Cosmochim. Acta*, **70**(20), 5104-5118.

- 2849 Lacan, F., Radic, A., Jeandel, C., Poitrasson, F., Sarthou, G., Pradoux, C., & Freydier, R. (2008).  
 2850 Measurement of the isotopic composition of dissolved iron in the open ocean. *Geophys. Res. Lett.*,  
 2851 **35**, L24610, doi:10.1029/2008GL035841.
- 2852 Lal, D., Charles, C., Vacher, L., Goswami, J. N., Jull, A. J. T., McHargue, L., & Finkel, R. C. (2006). Paleo-  
 2853 ocean chemistry records in marine opal: Implications for fluxes of trace elements, cosmogenic  
 2854 nuclides (Be-10 and Al-26), and biological productivity. *Geochim. Cosmochim. Acta* **70**, 3275-  
 2855 3289.
- 2856 Lam, P. J., & Marchal, O. (2015). Insights into particle cycling from thorium and particle data. *Annual*  
 2857 *Review of Marine Science*, **7**, 159-184.
- 2858 Lambelet, M., Rehkämper, M., van de Flierdt, T., Xue, Z., Kreissig, K., Coles, B., Porcelli, D., &  
 2859 Andersson, P. (2013). Isotopic analysis of Cd in the mixing zone of Siberian rivers with the Arctic  
 2860 Ocean—New constraints on marine Cd cycling and the isotope composition of riverine Cd. *Earth*  
 2861 *Planet. Sci. Lett.*, **361**, 64-73.
- 2862 Lane, T. W., Saito, M. A., George, G. N., Pickering, I. J., Prince, R. C., & Morel, F. M. (2005).  
 2863 Biochemistry: a cadmium enzyme from a marine diatom. *Nature*, **435**(7038), 42.
- 2864 Lauderdale, J. M., Braakman, R., Forget, G., Dutkiewicz, S., & Follows, M. J. (2020). Microbial feedbacks  
 2865 optimize ocean iron availability. *Proc. Natl. Acad. Sci.*, **117**(9), 4842-4849.
- 2866 Lee, J. G., & Morel, F. M. M. (1995). Replacement of zinc by cadmium in marine phytoplankton. *Mar.*  
 2867 *Ecol. Prog. Ser.*, **127**, 305–309.
- 2868 Lemaitre, N., de Souza, G. F., Archer, C., Wang, R. M., Planquette, H., Sarthou, G., & Vance, D. (2020).  
 2869 Pervasive sources of isotopically light zinc in the North Atlantic Ocean. *Earth Planet. Sci. Lett.*,  
 2870 **539**, doi:10.1016/j.epsl.2020.116216.
- 2871 Lerner, P., Marchal, O., Lam, P. J., & Solow, A. (2018). Effects of particle composition on thorium  
 2872 scavenging in the North Atlantic. *Geochimica et Cosmochimica Acta*, **233**, 115-134.
- 2873 Liao, W. H., & Ho, T. Y. (2018). Particulate trace metal composition and sources in the Kuroshio adjacent  
 2874 to the East China Sea: The importance of aerosol deposition. *Journal of Geophysical Research:*  
 2875 *Oceans*, **123**(9), 6207-6223.
- 2876 Liao, W. H., Takano, S., Yang, S. C., Huang, K. F., Sohrin, Y., & Ho, T. Y. (2020). Zn isotope composition  
 2877 in the water column of the Northwestern Pacific Ocean: the importance of external sources. *Global*  
 2878 *Biogeochem. Cycles*, **34**(1), e2019GB006379.
- 2879 Liermann, L. J., Guynn, R. L., Anbar, A., & Brantley, S. L. (2005). Production of a molybdophore during  
 2880 metal-targeted dissolution of silicates by soil bacteria. *Chem. Geol.*, **220**, 285–302,  
 2881 doi:10.1016/j.chemgeo.2005.04.013.
- 2882 Little, S.H., Vance, D., Siddall, M. and Gasson, E. (2013). A modeling assessment of the role of reversible  
 2883 scavenging in controlling oceanic dissolved Cu and Zn distributions. *Global Biogeochem. Cycles*  
 2884 **27**(3), 780-791.
- 2885 Little, S.H., Sherman, D.M., Vance, D. and Hein, J.R. (2014a). Molecular controls on Cu and Zn isotopic  
 2886 fractionation in Fe–Mn crusts. *Earth Planet. Sci. Lett.* **396**, 213-222.
- 2887 Little, S. H., Vance, D., Walker-Brown, C., & Landing, W. M. (2014b). The oceanic mass balance of copper  
 2888 and zinc isotopes, investigated by analysis of their inputs, and outputs to ferromanganese oxide  
 2889 sediments. *Geochim. Cosmochim. Acta* **125**, 673–693.
- 2890 Little, S. H., Vance, D., Lyons, T. W., & McManus, J. (2015). Controls on trace metal authigenic  
 2891 enrichment in reducing sediments: insights from modern oxygen-deficient settings. *Am. J. Sci.*,  
 2892 **315**(2), 77-119.



- 2893 Little, S. H., Vance, D., McManus, J., & Severmann, S. (2016). Key role of continental margin sediments  
2894 in the oceanic mass balance of Zn and Zn isotopes. *Geology*, **44**, 207–210.
- 2895 Little, S. H., van de Flierdt, T., Rehkämper, M., Wilson, D. J., Adkins, J. F., & Robinson, L. F. (2017a).  
2896 Deep sea corals as archives of seawater Zn isotopes. In *Goldschmidt Abstracts*, p. 2387.
- 2897 Little, S.H., Vance, D., McManus, J., Severmann, S. and Lyons, T.W. (2017b). Copper isotope signatures  
2898 in modern marine sediments. *Geochim. Cosmochim. Acta*, **212**, pp.253-273.
- 2899 Little, S. H., Archer, C., Milne, A., Schlosser, C., Achterberg, E. P., Lohan, M. C., & Vance, D. (2018).  
2900 Paired dissolved and particulate phase Cu isotope distributions in the South Atlantic. *Chem. Geol.*  
2901 **502**, 29-43.
- 2902 Little, S.H., Archer, C., McManus, J., Najorka, J., Wegorzewski, A.V. and Vance, D. (2020). Towards  
2903 balancing the oceanic Ni budget. *Earth Planet. Sci. Lett.* **547**, p.116461.
- 2904 Liu, R., Guo, B., Wang, M., Li, W., Yang, T., Ling, H., & Chen, T. (2020). Isotopic fingerprinting of  
2905 dissolved iron sources in the deep western Pacific since the late Miocene. *Science China Earth*  
2906 *Sciences*, **63**, doi:10.1007/s11430-020-9648-6.
- 2907 Liu, S. A., Wu, H., Shen, S. Z., Jiang, G., Zhang, S., Lv, Y., Zhang, H., & Li, S. (2017). Zinc isotope  
2908 evidence for intensive magmatism immediately before the end-Permian mass extinction. *Geology*,  
2909 **45**, 343–346.
- 2910 Liu, X., & Millero, F. J. (2002). The solubility of iron in seawater. *Mar. Chem.*, **77**(1), 43-54.
- 2911 Le Roy, E. L., Sanial, V., Charette, M. A., van Beek, P., Lacan, F., Jacquet, S. H., Henderson, P. B.,  
2912 Souhaut, M., García-Ibáñez, M. I., Jeandel, C., Pérez, F. F., & Sarthou, G. (2018). The <sup>226</sup>Ra–Ba  
2913 relationship in the North Atlantic during GEOTRACES-GA01. *Biogeosciences*, **15**(9), 3027-3048.
- 2914 Levasseur, S., Frank, M., Hein, J. R., & Halliday, A. N. (2004). The global variation in the iron isotope  
2915 composition of marine hydrogenetic ferromanganese deposits: implications for seawater  
2916 chemistry? *Earth Planet. Sci. Lett.*, **224**(1-2), 91-105.
- 2917 Liu, Y., Li, X., Zeng, Z., Yu, H. M., Huang, F., Felis, T., & Shen, C. C. (2019). Annually-resolved coral  
2918 skeletal  $\delta^{138/134}\text{Ba}$  records: A new proxy for oceanic Ba cycling. *Geochim. Cosmochim. Acta*, **247**,  
2919 27-39.
- 2920 Lohan, M. C., Crawford, D. W., Purdie, D. A., & Statham, P. J. (2005). Iron and Zinc Enrichments in the  
2921 Northeastern Subarctic Pacific: Ligand Production and Zinc Availability in Response to  
2922 Phytoplankton Growth. *Limnol. Oceanogr.*, **50**, 1427–1437.
- 2923 Lough, A. J. M., Klar, J. K., Homoky, W. B., Comer-Warner, S. A., Milton, J. A., Connelly, D. P., James,  
2924 R. H., & Mills, R. A. (2017). Opposing authigenic controls on the isotopic signature of dissolved  
2925 iron in hydrothermal plumes. *Geochim. Cosmochim. Acta*, **202**, 1-20.
- 2926 Lv, Y., Liu, S. A., Wu, H., Hohl, S. V., Chen, S., & Li, S. (2018). Zn-Sr isotope records of the Ediacaran  
2927 Doushantuo Formation in South China: diagenesis assessment and implications. *Geochim.*  
2928 *Cosmochim. Acta*, **239**, 330-345.
- 2929 Lyons, T. W., Anbar, A. D., Severmann, S., Scott, C., & Gill, B. C. (2009). Tracking Euxinia in the Ancient  
2930 Ocean: A Multiproxy Perspective and Proterozoic Case Study. *Annu. Rev. Earth Planet. Sci.*, **37**,  
2931 507–534, doi:10.1146/annurev.earth.36.031207.124233.
- 2932 Ma, Z., Gray, E., Thomas, E., Murphy, B., Zachos, J., & Paytan, A. (2014). Carbon sequestration during  
2933 the Palaeocene–Eocene Thermal Maximum by an efficient biological pump. *Nat. Geosci.*, **7**(5),  
2934 382.

- 2935 Ma, Z., Ravelo, A. C., Liu, Z., Zhou, L., & Paytan, A. (2015). Export production fluctuations in the eastern  
 2936 equatorial Pacific during the Pliocene-Pleistocene: Reconstruction using barite accumulation rates.  
 2937 *Paleoceanography*, **30**(11), 1455-1469.
- 2938 Mahaffey, C., Reynolds, S., Davis, C. E., & Lohan, M. C. (2014). Alkaline phosphatase activity in the  
 2939 subtropical ocean: insights from nutrient, dust and trace metal addition experiments. *Front. Mar.*  
 2940 *Sci.*, **1**, 1–13.
- 2941 Maldonado, M. T., Allen, A. E., Chong, J. S., Lin, K., Leus, D., Karpenko, N., & Harris, S. L. (2006).  
 2942 Copper-dependent iron transport in coastal and oceanic diatoms. *Limnol. Oceanogr.* **51**, 1729–  
 2943 1743.
- 2944 Marchitto, T. M., Curry, W. B., & Oppo, D. W. (2000) Zinc concentrations in benthic foraminifera reflect  
 2945 seawater chemistry. *Paleoceanography*, **15**, 299–306.
- 2946 Marcus, M. A., Edwards, K. J., Gueguen, B., Fakra, S. C., Horn, G., Jelinski, N. A., Rouxel, O., Sorensen,  
 2947 J., & Toner, B. M. (2015). Iron mineral structure, reactivity, and isotopic composition in a South  
 2948 Pacific Gyre ferromanganese nodule over 4 Ma. *Geochim. Cosmochim. Acta*, **171**, 61-79.
- 2949 Maréchal, C. N., Nicolas, E., Douchet, C., & Albarède, F. (2000). Abundance of zinc isotopes as a marine  
 2950 biogeochemical tracer. *Geochem. Geophys. Geosyst.*, **1**, 1015,  
 2951 <http://doi.wiley.com/10.1029/1999GC000029>.
- 2952 Marks, J. A., Perakis, S. S., King, E. K., & Pett-Ridge, J. (2015). Soil organic matter regulates molybdenum  
 2953 storage and mobility in forests. *Biogeochemistry*, **125**, 167–183, doi:10.1007/s10533-015-0121-4.
- 2954 Marsay, C. M., Lam, P. J., Heller, M. I., Lee, J. M., & John, S. G. (2018). Distribution and isotopic signature  
 2955 of ligand-leachable particulate iron along the GEOTRACES GP16 East Pacific Zonal Transect.  
 2956 *Mar. Chem.*, **201**, 198-211.
- 2957 Martin, J. H. (1990). Glacial-interglacial CO<sub>2</sub> change: the iron hypothesis. *Paleoceanography* **5**, 1-13.
- 2958 Martin, J. H., & Fitzwater, S. E. (1988). Iron deficiency limits phytoplankton growth in the north-east  
 2959 Pacific subarctic. *Nature*, **331**(6154), 341-343.
- 2960 Martin, J. H., Fitzwater, S. E., & Gordon, R. M. (1990). Iron deficiency limits phytoplankton growth in  
 2961 Antarctic waters. *Global Biogeochem. Cycles*, **4**(1), 5-12.
- 2962 Martin, J. H., Knauer, G. A. and Gordon, R. M. (1983) Silver distributions and fluxes in north-east Pacific  
 2963 waters. *Nature*, **305**, 306-309.
- 2964 Martin, J. M., & Thomas, A. J. (1994). The global insignificance of telluric input of dissolved trace metals  
 2965 (Cd, Cu, Ni and Zn) to ocean margins. *Mar. Chem.*, **46**(1-2), 165-178.
- 2966 Martínez-García, A., Rosell-Melé, A., Jaccard, S. L., Geibert, W., Sigman, D. M., & Haug, G. H. (2011).  
 2967 Southern Ocean dust–climate coupling over the past four million years. *Nature*, **476**(7360), 312-  
 2968 315.
- 2969 Martínez-García, A., Sigman, D. M., Ren, H., Anderson, R. F., Straub, M., Hodell, D. A., et al. (2014). Iron  
 2970 fertilization of the Subantarctic Ocean during the last ice age. *Science*, **343**, 1347-1350,  
 2971 doi:10.1126/science.1246848.
- 2972 Martínez-Ruiz, F., Jroundi, F., Paytan, A., Guerra-Tschuschke, I., del Mar Abad, M., & González-Muñoz,  
 2973 M. T. (2018). Barium bioaccumulation by bacterial biofilms and implications for Ba cycling and  
 2974 use of Ba proxies. *Nat. Commun.*, **9**.
- 2975 Martínez-Ruiz, F., Paytan, A., González-Muñoz, M. T., Jroundi, F., Abad, M. M., Lam, P. J., Bishop, J. K.  
 2976 B., Horner, T. J., Morton, P. L., & Kastner, M. (2019). Barite formation in the ocean: Origin of  
 2977 amorphous and crystalline precipitates. *Chem. Geol.*, **511**, 441-451.

- 2978 Mathur, R., Arribas, A., Megaw, P., Wilson, M., Stroup, S., Meyer-Arrivillaga, D., & Arribas, I. (2018).  
 2979 Fractionation of silver isotopes in native silver explained by redox reactions. *Geochim. Cosmochim.*  
 2980 *Acta*, **224**, 313-326.
- 2981 Mawji, E. et al. (2015). The GEOTRACES Intermediate Data Product 2014. *Mar. Chem.*, **177**, 1-8,  
 2982 doi:10.1016/j.marchem.2015.04.005.
- 2983 Mayfield, K. K., A. Eisenhauer, D. Santiago Ramos, J. A. Higgins, T. J. Horner, M. E. Auro, T. Magna, N.  
 2984 Moosdorf, M. A. Charette, M. E. Gonnea, C. E. Brady, N. Komar, B. Peucker-Ehrenbrink, and A.  
 2985 Paytan (in press) The Importance of Groundwater Discharge in Marine Isotope Budgets, *Nature*  
 2986 *Communications*.
- 2987 McKay, J. L., & Pedersen, T. F. (2008). The accumulation of silver in marine sediments: A link to biogenic  
 2988 Ba and marine productivity. *Global Biogeochem. Cycles*, **22**, GB4010,  
 2989 doi:10.1029/2007GB003136
- 2990 McManus, J., Berelson, W. M., Klinkhammer, G. P., Johnson, K. S., Coale, K. H., Anderson, R. F., Kumar,  
 2991 N., Burdige, D. J., Hammond, D. E., Brumsack, H. J., McCorkle, D. C., & Rushdi, A. (1998).  
 2992 Geochemistry of barium in marine sediments: Implications for its use as a paleoproxy. *Geochim.*  
 2993 *Cosmochim. Acta*, **62**(21-22), 3453-3473.
- 2994 McManus, J., Dymond, J., Dunbar, R. B., & Collier, R. W. (2002). Particulate barium fluxes in the Ross  
 2995 Sea. *Mar. Geol.*, **184**(1-2), 1-15.
- 2996 McManus, J., Berelson, W. M., Severmann, S., Poulson, R. L., Hammond, D. E., Klinkhammer, G. P., &  
 2997 Holm, C. (2006). Molybdenum and uranium geochemistry in continental margin sediments:  
 2998 Paleoproxy potential. *Geochim. Cosmochim. Acta*, **70**, 4643-4662,  
 2999 doi:10.1016/j.gca.2006.06.1564.
- 3000 Mead, C., Herckes, P., Majestic, B. J., & Anbar, A. D. (2013). Source apportionment of aerosol iron in the  
 3001 marine environment using iron isotope analysis. *Geophys. Res. Lett.*, **40**(21), 5722-5727.
- 3002 Mendel, R. R., & Bittner, F. (2006) Cell biology of molybdenum. *Biochim. Biophys. Acta*, **1763**(7), 621-  
 3003 635, doi:10.1016/j.bbamcr.2006.03.013.
- 3004 Middag, R., van Heuven, S. M., Bruland, K. W., & de Baar, H. J. (2018). The relationship between cadmium  
 3005 and phosphate in the Atlantic Ocean unraveled. *Earth Planet. Sci. Lett.*, **492**, 79-88.
- 3006 Middag, R., de Baar, H. J. W., & Bruland, K. W. (2019). The relationships between dissolved zinc and  
 3007 major nutrients phosphate and silicate along the GEOTRACES GA02 transect in the west Atlantic  
 3008 Ocean. *Global Biogeochem. Cycles*, **33**, 63-84, doi:10.1029/2018GB006034.
- 3009 Middleton, J. L., Langmuir, C. H., Mukhopadhyay, S., McManus, J. F., & Mitrovica, J. X. (2016).  
 3010 Hydrothermal iron flux variability following rapid sea level changes. *Geophys. Res. Lett.*, **43**, 3848-  
 3011 3856.
- 3012 Miller, C. A., Peucker-Ehrenbrink, B., Walker, B. D., & Marcantonio, F. (2011). Re-assessing the surface  
 3013 cycling of molybdenum and rhenium. *Geochim. Cosmochim. Acta*, **75**, 7146-7179,  
 3014 doi:10.1016/j.gca.2011.09.005.
- 3015 Miller, L. A., & Bruland, K. W. (1995). Organic speciation of silver in marine waters. *Environ. Sci.*  
 3016 *Technol.*, **29**, 2616-2621.
- 3017 Misumi, K., Lindsay, K., Moore, J. K., Doney, S. C., Bryan, F. O., Tsumune, D., & Yoshida, Y. (2014).  
 3018 The iron budget in ocean surface waters in the 20th and 21st centuries: projections by the  
 3019 Community Earth System Model version. *Biogeosciences*, **11**, 33-55.

- 3020 Moeller, K., Schoenberg, R., Pedersen, R. B., Weiss, D., & Dong, S. (2012). Calibration of the new certified  
 3021 reference materials ERM-AE633 and ERM-AE647 for copper and IRMM-3702 for zinc isotope  
 3022 amount ratio determinations. *Geostan. Geoanal. Res.* **36**(2), 177-199.
- 3023 Moffett, J. W., & Brand, L. E. (1996). Production of strong, extracellular Cu chelators by marine  
 3024 cyanobacteria in response to Cu stress. *Limnol. Oceanogr.* **41**, 388–395.
- 3025 Moffett, J. W., & Dupont, C. (2007). Cu complexation by organic ligands in the sub-arctic NW Pacific and  
 3026 Bering Sea. *Deep. Res. Part I Oceanogr. Res. Pap.* **54**, 586–595.
- 3027 Moffett, J. W., Zika, R. G., & Brand, L. E. (1990). Distribution and potential sources and sinks of copper  
 3028 chelators in the Sargasso Sea. *Deep Sea Res. Part A, Oceanogr. Res. Pap.* **37**, 27–36.
- 3029 Monnin, C., Jeandel, C., Cattaldo, T., & Dehairs, F. (1999). The marine barite saturation state of the world's  
 3030 oceans. *Mar. Chem.*, **65**(3-4), 253-261.
- 3031 Monnin, C., & Cividini, D. (2006). The saturation state of the world's ocean with respect to (Ba, Sr)SO<sub>4</sub>  
 3032 solid solutions. *Geochim. Cosmochim. Acta*, **70**(13), 3290-3298.
- 3033 Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., Galbraith, E. D.,  
 3034 Geider, R. J., Guieu, C., Jaccard, S. L., Jickells, T. D., La Roche, J., Lenton, T. M., Mahowald, N.  
 3035 M., Marañon, E., Marinov, I., Moore, J.K., Nakatsuka, T., Oschlies, A., Saito, M. A., Thingstad,  
 3036 T. F., Tsuda, A., & Ulloa, O. (2013). Processes and patterns of oceanic nutrient limitation. *Nat.*  
 3037 *Geosci.* **6**, 701-710.
- 3038 Moore, C. M. (2016). Diagnosing oceanic nutrient deficiency. *Philosophical Transactions of the Royal*  
 3039 *Society A: Mathematical, Physical and Engineering Sciences*, **374**(2081), 20150290.
- 3040 Morel, F. M. M., & Price, N. M. (2003). The biogeochemical cycles of trace metals in the oceans. *Science*,  
 3041 **300**, 944-947, doi:10.1126/science.1083545.
- 3042 Morel, F. M. M., Reinfelder, J. R., Roberts, S. B., Chamberlain, C. P., Lee, J. G., & Yee, D. (1994) Zinc  
 3043 and carbon co-limitation of marine phytoplankton. *Nature*, **369**, 740-742.
- 3044 Morel, F. M., Lam, P. J., & Saito, M. A. (2020). Trace Metal Substitution in Marine Phytoplankton. *Annual*  
 3045 *Review of Earth and Planetary Sciences*, **48**.
- 3046 Morford, J. L., Kalnejais, L. H., Helman, P., Yen, G., & Reinard, M. (2008). Geochemical cycling of silver  
 3047 in marine sediments along an offshore transect. *Mar. Chem.*, **110**, 77-88.
- 3048 Morgan, J. L., Wasylenki, L. E., Nuester, J., & Anbar, A. D. (2010). Fe isotope fractionation during  
 3049 equilibration of Fe–organic complexes. *Environ. Sci. Technol.*, **44**(16), 6095-6101.
- 3050 Morris, A. W. (1975). Dissolved molybdenum and vanadium in the northeast Atlantic Ocean. *Deep Sea*  
 3051 *Research and Oceanographic Abstracts*, **22**, 49–54, doi:10.1016/0011-7471(75)90018-2.
- 3052 Moynier, F., Vance, D., Fujii, T., & Savage, P. (2017). The Isotope Geochemistry of Zinc and Copper. *Rev.*  
 3053 *Mineral. Geochemistry* **82**, 543–600.
- 3054 Murphy, K. L. (2016). *Isotopic studies in marine geochemistry* (Doctoral dissertation, Imperial College  
 3055 London), doi:10.25560/68270.
- 3056 Mulholland, D.S., Poitrasson, F., Boaventura, G.R., Allard, T., Vieira, L.C., Santos, R.V., Mancini, L. &  
 3057 Seyler, P. (2015). Insights into iron sources and pathways in the Amazon River provided by isotopic  
 3058 and spectroscopic studies. *Geochim. Cosmochim. Acta*, **150**, 142–159.
- 3059 Nögler, T. F., Neubert, N., Böttcher, M. E., Dellwig, O., & Schnetger, B. (2011). Molybdenum isotope  
 3060 fractionation in pelagic euxinia: Evidence from the modern Black and Baltic Seas. *Chem. Geol.*,  
 3061 **289**, 1–11, doi:10.1016/j.chemgeo.2011.07.001.



- 3062 Nagler, T. F., Anbar, A. D., Archer, C., Goldberg, T., Gordon, G. W., Greber, N. D., Seibert, C., Sohrin,  
 3063 Y., & Vance, D. (2014). Proposal for an international molybdenum isotope measurement standard  
 3064 and data representation. *Geostand. Geoanal. Res.*, **38**(2), 149-151.
- 3065 Nakagawa, Y., Takano, S., Firdaus, M. L., Norisuye, K., Hirata, T., Vance, D., & Sohrin, Y. (2012). The  
 3066 molybdenum isotopic composition of the modern ocean. *Geochemical Journal*, **46**, 131–141,  
 3067 doi:10.2343/geochemj.1.0158.
- 3068 Nan, X. Y., Yu, H. M., Rudnick, R. L., Gaschnig, R. M., Xu, J., Li, W.-Y., Zhang, Q., Jin, Z.-D., Li, X.-  
 3069 H., & Huang, F. (2018). Barium isotopic composition of the upper continental crust. *Geochim.  
 3070 Cosmochim. Acta*, **233**, 33-49.
- 3071 Nasemann, P., Gault-Ringold, M., Stirling, C. H., Koschinsky, A., & Sander, S. G. (2018). Processes  
 3072 affecting the isotopic composition of dissolved iron in hydrothermal plumes: A case study from the  
 3073 Vanuatu back-arc. *Chem. Geol.* **476**, 70-84. <https://doi.org/10.1016/j.chemgeo.2017.11.005>.
- 3074 Navarrete, J. U., Borrok, D. M., Viveros, M., & Ellzey, J. T. (2011). Copper isotope fractionation during  
 3075 surface adsorption and intracellular incorporation by bacteria. *Geochim. Cosmochim. Acta*, **75**(3),  
 3076 784-799.
- 3077 Nishioka, J., Obata, H., Ogawa, H., Ono, K., Yamashita, Y., Lee, K., Takeda, S., & Yasuda, I. (2020).  
 3078 Subpolar marginal seas fuel the North Pacific through the intermediate water at the termination of  
 3079 the global ocean circulation. *Proc. Natl Acad. Sci. U. S. A.* **117**(23), 12665-12673  
 3080 <https://doi.org/10.1073/pnas.2000658117>.
- 3081 Ndung'u, K., Thomas, M.A., & Flegal, A.R. (2001). Silver in the western equatorial and South Atlantic  
 3082 Ocean. *Deep-Sea Research II*, **48**, 2933-2945.
- 3083 Neubert, N., Nagler, T. F., and Bottcher, M. E. (2008). Sulfidity controls molybdenum isotope fractionation  
 3084 into euxinic sediments: Evidence from the modern Black Sea. *Geology*, **36**, 775–778,  
 3085 doi:10.1130/G24959A.1.
- 3086 Nielsen, S. G., Wasylenki, L. E., Rehkamper, M., Peacock, C. L., Xue, Z., & Moon, E. M. (2013). Towards  
 3087 an understanding of thallium isotope fractionation during adsorption to manganese oxides.  
 3088 *Geochim. Cosmochim. Acta*, **117**, 252-265.
- 3089 Noordmann, J., Weyer, S., Montoya-Pino, C., Dellwig, O., Neubert, N., Eckert, S., Paetzel, M., & Bottcher,  
 3090 M. E. (2015). Uranium and molybdenum isotope systematics in modern euxinic basins: Case  
 3091 studies from the central Baltic Sea and the Kyllaren fjord (Norway). *Chem. Geol.*, **396**, 182–195,  
 3092 doi:10.1016/j.chemgeo.2014.12.012.
- 3093 Ohnemus, D. C., & Lam, P. J. (2015). Cycling of lithogenic marine particles in the US GEOTRACES North  
 3094 Atlantic transect. *Deep Sea Res. Pt. II*, **116**, 283-302.
- 3095 Ohnemus, D. C., Rauschenberg, S., Cutter, G. A., Fitzsimmons, J. N., Sherrell, R. M., & Twining, B. S.  
 3096 (2017). Elevated trace metal content of prokaryotic communities associated with marine oxygen  
 3097 deficient zones. *Limnol. Oceanog.*, **62**(1), 3-25.
- 3098 Ohnemus, D. C., Torrie, R., & Twining, B. S. (2019). Exposing the Distributions and Elemental  
 3099 Associations of Scavenged Particulate Phases in the Ocean Using Basin-Scale Multi-Element Data  
 3100 Sets. *Global Biogeochem. Cycles*, **33**(6), 725-748.
- 3101 Park, H., Song, B., & Morel, F. M. (2007). Diversity of the cadmium-containing carbonic anhydrase in  
 3102 marine diatoms and natural waters. *Environmental Microbiology*, **9**(2), 403-413.
- 3103 Paytan, A. (2009). Ocean Paleoproductivity. In: Gornitz V. (eds) *Encyclopedia of Paleoclimatology and  
 3104 Ancient Environments*. Encyclopedia of Earth Sciences Series, Springer, Dordrecht.

- 3105 Paytan, A., & Kastner, M. (1996). Benthic Ba fluxes in the central Equatorial Pacific, implications for the  
3106 oceanic Ba cycle. *Earth Planet. Sci. Lett.*, **142**(3-4), 439-450.
- 3107 Paytan, A., Kastner, M., & Chavez, F. P. (1996). Glacial to interglacial fluctuations in productivity in the  
3108 equatorial Pacific as indicated by marine barite. *Science*, **274**(5291), 1355-1357.
- 3109 Paytan, A., & Griffith, E. M. (2007). Marine barite: Recorder of variations in ocean export productivity.  
3110 *Deep Sea Res. Pt. II*, **54**(5-7), 687-705.
- 3111 Paytan, A., Mackey, K. R., Chen, Y., Lima, I. D., Doney, S. C., Mahowald, N., Labiosa, R., & Post, A. F.  
3112 (2009). Toxicity of atmospheric aerosols on marine phytoplankton. *Proc. Natl. Acad. Sci.*, **106**(12),  
3113 4601-4605.
- 3114 Peers, G., & Price, N. M. (2006). Copper-containing plastocyanin used for electron transport by an oceanic  
3115 diatom. *Nature* **441**, 341–344.
- 3116 Peers, G., Quesnel, S., & Price, N. M. (2005). Copper requirements for iron acquisition and growth of  
3117 coastal and oceanic diatoms. *Limnol. Oceanogr.* **50**, 1149–1158.
- 3118 Pichat, S., Douchet, C., & Albarède, F. (2003). Zinc isotope variations in deep-sea carbonates from the  
3119 eastern equatorial Pacific over the last 175 ka. *Earth Planet. Sci. Lett.*, **210**, 167–178.
- 3120 Pichevin, L. E., Ganeshram, R. S., Geibert, W., Thunell, R., & Hinton, R. (2014). Silica burial enhanced  
3121 by iron limitation in oceanic upwelling margins. *Nat. Geosci.*, **7**(7), 541-546.
- 3122 Plass, A., Schlosser, C., Sommer, S., Dale, A. W., Achterberg, E. P., & Scholz, F. (2020). The control of  
3123 hydrogen sulfide on benthic iron and cadmium fluxes in the oxygen minimum zone off Peru.  
3124 *Biogeosciences*, **17**(13), 3685-3704.
- 3125 Poitrasson, F. (2006). On the iron isotope homogeneity level of the continental crust. *Chem. Geol.*, **235**(1-  
3126 2), 195-200.
- 3127 Poitrasson, F., Vieira, L. C., Seyler, P., dos Santos Pinheiro, G. M., Mulholland, D. S., Bonnet, M. P.,  
3128 Martinez, J.-M., Lima, B. A., Boaventura, G. R., Chmeleff, J., Dantas, E. L., Guyot, J.-L., Mancini,  
3129 L., Pimentel, M. M., Santos, R. V., Sondag, F., & Vauchel, P. (2014). Iron isotope composition of  
3130 the bulk waters and sediments from the Amazon River Basin. *Chem. Geol.*, **377**, 1-11.
- 3131 Pokrovsky, O. S., Viers, J., Emnova, E. E., Kompantseva, E. I., & Freydier, R. (2008). Copper isotope  
3132 fractionation during its interaction with soil and aquatic microorganisms and metal oxy (hydr)  
3133 oxides: Possible structural control. *Geochim. Cosmochim. Acta*, **72**(7), 1742-1757.
- 3134 Price, N. M., & Morel, F. M. M. (1990). Cadmium and cobalt substitution for zinc in a marine diatom.  
3135 *Nature*, **344**(6267), 658.
- 3136 Price, N. M., & Morel, F. M. M. (1991). Colimitation of phytoplankton growth by nickel and nitrogen.  
3137 *Limnol. Oceanogr.* **50**, 1071-1077.
- 3138 Quay, P., & Wu, J. (2015). Impact of end-member mixing on depth distributions of  $\delta^{13}\text{C}$ , cadmium and  
3139 nutrients in the N. Atlantic Ocean. *Deep Sea Res. Pt. II*, **116**, 107-116.
- 3140 Radic, A., Lacan, F., & Murray, J. W. (2011). Iron isotopes in the seawater of the equatorial Pacific Ocean:  
3141 New constraints for the oceanic iron cycle. *Earth Planet. Sci. Lett.*, **306**, 1-10.
- 3142 Rafter, P. A., Sigman, D. M., & Mackey, K. R. (2017). Recycled iron fuels new production in the eastern  
3143 equatorial Pacific Ocean. *Nat. Commun.*, **8**(1), 1100.
- 3144 Ragsdale, S. W. (2009). Nickel-based enzyme systems. *J. Biol. Chem.* **284**, 18571–18575.
- 3145 Ranville, M. A., & Flegal, A.R. (2005). Silver in the North Pacific Ocean. *Geochemistry, Geophysics,*  
3146 *Geosystems*, **6**(3), doi:10.1029/2004GC000770

- 3147 Ranville, M. A., Cutter, G. A., Buck, C. S., Landing, W. M., Cutter, L. S., Resing, J. A., & Flegal, A. R.  
 3148 (2010). Aeolian contamination of Se and Ag in the North Pacific from Asian fossil fuel combustion.  
 3149 *Environ. Sci. Tech.*, **44**, 1587–1593.
- 3150 Rehkämper, M., Wombacher, F., Horner, T. J., & Xue, Z. (2012). Natural and anthropogenic Cd isotope  
 3151 variations. In Baskaran, M. (Ed.) *Handbook of Environmental Isotope Geochemistry* (pp. 125-154).  
 3152 Springer, Berlin, Heidelberg, doi:10.1007/978-3-642-10637-8\_8.
- 3153 Reinhard, C. T., Planavsky, N. J., Robbins, L. J., Partin, C. A., Gill, B. C., Lalonde, S. V., Bekker, A.,  
 3154 Konhauser, K. O., & Lyons, T. W. (2013). Proterozoic ocean redox and biogeochemical stasis.  
 3155 *Proc. Natl. Acad. Sci.*, **110**, 5357–5362, doi:10.1073/pnas.1208622110.
- 3156 Resing, J. A., Sedwick, P. N., German, C. R., Jenkins, W. J., Moffett, J. W., Sohst, B. M., & Tagliabue, A.  
 3157 (2015). Basin-scale transport of hydrothermal dissolved metals across the South Pacific Ocean.  
 3158 *Nature*, **523**(7559), 200-203.
- 3159 Revels, B. N., Zhang, R., Adkins, J. F., & John, S. G. (2015). Fractionation of iron isotopes during leaching  
 3160 of natural particles by acidic and circumneutral leaches and development of an optimal leach for  
 3161 marine particulate iron isotopes. *Geochim. Cosmochim. Acta*, **166**, 92–104.
- 3162 Richon, C., & Tagliabue, A. (2019). Insights into the major processes driving the global distribution of  
 3163 copper in the ocean from a global model. *Global Biogeochem. Cycles*, **33**(12), 1594-1610.
- 3164 Rickaby, R. E. M., & Elderfield, H. (1999). Planktonic foraminiferal Cd/Ca: paleonutrients or  
 3165 paleotemperature?. *Paleoceanography*, **14**(3), 293-303.
- 3166 Ridge, P. G., Zhang, Y., & Gladyshev, V. N. (2008). Comparative genomic analyses of copper transporters  
 3167 and cuproproteomes reveal evolutionary dynamics of copper utilization and its link to oxygen.  
 3168 *PLoS One* **3**(1), <https://doi.org/10.1371/journal.pone.0001378>.
- 3169 Rijkenberg, M. J. A., Middag, R., Laan, P., Gerringa, L. J., van Aken, H. M., Schoemann, V., de Jong, J.  
 3170 T. M., & de Baar, H. J. W. (2014). The distribution of dissolved iron in the West Atlantic Ocean.  
 3171 *PloS One*, **9**(6).
- 3172 Ripperger, S., & Rehkämper, M. (2007). Precise determination of cadmium isotope fractionation in  
 3173 seawater by double spike MC-ICPMS. *Geochim. Cosmochim. Acta*, **71**(3), 631-642.
- 3174 Ripperger, S., Rehkämper, M., Porcelli, D., & Halliday, A. N. (2007). Cadmium isotope fractionation in  
 3175 seawater—A signature of biological activity. *Earth Planet. Sci. Lett.*, **261**(3-4), 670-684.
- 3176 Rivera-Duarte, I., Flegal, A. R., Sañudo-Wilhelmy, S. A., & Véron, A. J. (1999). Silver in the far North  
 3177 Atlantic Ocean. *Deep-Sea Research II*, **46**, 979-990.
- 3178 Robinson, L. F., Adkins, J. F., Frank, N., Gagnon, A. C., Prouty, N. G., Roark, E. B., & van de Fliedrt, T.  
 3179 (2014). The geochemistry of deep-sea coral skeletons: A review of vital effects and applications  
 3180 for palaeoceanography. *Deep. Res. Part II Top. Stud. Oceanogr.*, **99**, 184–198.  
 3181 <http://dx.doi.org/10.1016/j.dsr2.2013.06.005>.
- 3182 Rohatgi, A. (2019) WebPlotDigitizer (version 4.2; April 2019). <https://automeris.io/WebPlotDigitizer>  
 3183 [accessed 2020-06-22].
- 3184 Rolison, J. M., Stirling, C. H., Middag, R., Gault-Ringold, M., George, E., & Rijkenberg, M. J. A. (2018).  
 3185 Iron isotope fractionation during pyrite formation in a sulfidic Precambrian ocean analogue. *Earth*  
 3186 *Planet. Sci. Lett.*, **488**, 1-13. <https://doi.org/10.1016/j.epsl.2018.02.006>.
- 3187 Roshan, S., & Wu, J. (2015a). The distribution of dissolved copper in the tropical-subtropical north Atlantic  
 3188 across the GEOTRACES GA03 transect. *Mar. Chem.*, **176**, 189-198,  
 3189 doi:10.1016/j.marchem.2015.09.006.



- 3190 Roshan, S., & Wu, J. (2015b). Cadmium regeneration within the North Atlantic. *Global Biogeochem.*  
 3191 *Cycles*, **29**(12), 2082-2094.
- 3192 Roshan, S., & Wu, J. (2018). Dissolved and colloidal copper in the tropical South Pacific. *Geochim.*  
 3193 *Cosmochim. Acta*, **233**, 81-94, doi:10.1016/j.gca.2018.05.008.
- 3194 Roshan, S., DeVries, T., Wu, J., & Chen, G. (2018). The internal cycling of zinc in the ocean. *Global*  
 3195 *Biogeochem. Cycles*, **32**(12), 1833-1849, doi:10.1029/2018GB006045.
- 3196 Roshan, S., DeVries, T., Wu, J., John, S., & Weber, T. (2020). Reversible scavenging traps hydrothermal  
 3197 iron in the deep ocean. *Earth and Planetary Science Letters*, 542, 116297.
- 3198 Rühlemann, C., Müller, P. J., & Schneider, R. R. (1999). Organic carbon and
- 3199 Rouxel, O., Shanks III, W. C., Bach, W., & Edwards, K. J. (2008). Integrated Fe-and S-isotope study of  
 3200 seafloor hydrothermal vents at East Pacific Rise 9–10 N. *Chem. Geol.*, **252**(3-4), 214-227.
- 3201 Rouxel, O., Toner, B. M., Manganini, S. J., & German, C. R. (2016). Geochemistry and iron isotope  
 3202 systematics of hydrothermal plume fall-out at East Pacific Rise 9°50'N. *Chem. Geol.* **441**, 212-234.  
 3203 <https://doi.org/10.1016/j.chemgeo.2016.08.027>.
- 3204 Rouxel, O., Toner, B., Germain, Y., & Glazer, B. (2018). Geochemical and iron isotopic insights into  
 3205 hydrothermal iron oxyhydroxide deposit formation at Loihi Seamount. *Geochim. Cosmochim.*  
 3206 *Acta*, **220**, 449-482. <https://doi.org/10.1016/j.gca.2017.09.050>.
- 3207 Rosenthal, Y., Lam, P., Boyle, E. A., & Thomson, J. (1995). Authigenic cadmium enrichments in suboxic  
 3208 sediments: Precipitation and postdepositional mobility. *Earth Planet. Sci. Lett.*, **132**(1-4), 99-111.
- 3209 carbonate as paleoproductivity proxies: Examples from high and low productivity areas of the tropical  
 3210 Atlantic. In *Use of Proxies in Paleoceanography* (pp. 315-344). Springer, Berlin, Heidelberg.
- 3211 Rushdi, A. I., McManus, J., & Collier, R. W. (2000). Marine barite and celestite saturation in seawater.  
 3212 *Mar. Chem.*, **69**(1-2), 19-31.
- 3213 Ryan, B. M., Kirby, J. K., Degryse, F., Scheiderich, K., & McLaughlin, M. J. (2014). Copper isotope  
 3214 fractionation during equilibration with natural and synthetic ligands. *Environ. Sci. Technol.*, **48**(15),  
 3215 8620-8626.
- 3216 Saito, M. A., Moffett, J. W., & DiTullio, G. R. (2004). Cobalt and nickel in the Peru upwelling region: A  
 3217 major flux of labile cobalt utilized as a micronutrient. *Global Biogeochem. Cycles* **18**, 1–14.
- 3218 Saito, M. A., Bertrand, E. M., Dutkiewicz, S., Bulygin, V. V., Moran, D. M., Monteiro, F. M., Follows, M.  
 3219 J., Valois, F. W., & Waterbury, J. B. (2011). Iron conservation by reduction of metalloenzyme  
 3220 inventories in the marine diazotroph *Crocospaera watsonii*. *Proc. Natl. Acad. Sci.*, **108**(6), 2184-  
 3221 2189.
- 3222 Saito, M. A., Noble, A. E., Tagliabue, A., Goepfert, T. J., Lamborg, C. H., & Jenkins, W. J. (2013). Slow-  
 3223 spreading submarine ridges in the South Atlantic as a significant oceanic iron source. *Nat. Geosci.*,  
 3224 **6**(9), 775-779.
- 3225 Samanta, M., Ellwood, M. J., Sinoir, M., & Hassler, C. S. (2017). Dissolved zinc isotope cycling in the  
 3226 Tasman Sea, SW Pacific Ocean. *Mar. Chem.*, **192**, 1–12.  
 3227 <http://dx.doi.org/10.1016/j.marchem.2017.03.004>.
- 3228 Samanta, M., Ellwood, M. J., & Strzepek, R. F. (2018). Zinc isotope fractionation by *Emiliania huxleyi*  
 3229 cultured across a range of free zinc ion concentrations. *Limnol. Oceanogr.*, **63**, 660–671.
- 3230 Sarmiento, J. L., & Gruber, M. (2006). Carbon Cycle. In *Ocean Biogeochemical Dynamics* (pp. 318-358).  
 3231 Princeton, NJ: Princeton University Press.

- 3232 Sarmiento, J. L., Gruber, N., Brzezinski, M. A., & Dunne, J. P. (2004). High-latitude controls of thermocline  
3233 nutrients and low latitude biological productivity. *Nature*, **427**(6969), 56-60.
- 3234 Sclater, F. R., Boyle, E., & Edmond, J. M. (1976). On the marine geochemistry of nickel. *Earth Planet. Sci.*  
3235 *Lett.*, **31**(1), 119-128.
- 3236 Schönbacher, M., Carlson, R. W., Horan, M. F., Mock, T. D., & Hauri, E. H. (2007). High precision Ag  
3237 isotope measurements in geologic materials by multiple-collector ICPMS: An evaluation of dry  
3238 versus wet plasma. *Intl J. Mass Spectrom.*, **261**(2-3), 183-191.
- 3239 Sedwick, P. N., Church, T. M., Bowie, A.R., Marsay, C. M., Ussher, S. J., Achilles, K. M., Lethaby, P. J.,  
3240 Johnson, R. J., Sarin, M. M., McGillicuddy, D. J. (2005). Iron in the Sargasso Sea (Bermuda  
3241 Atlantic Time-series Study region) during summer: Eolian imprint, spatiotemporal variability, and  
3242 ecological implications. *Global Biogeochem. Cycles*, **19**, GB4006.  
3243 <https://doi.org/10.1029/2004GB002445>.
- 3244 Sharma, M., Polizzotto, M., & Anbar, A. D. (2001). Iron isotopes in hot springs along the Juan de Fuca  
3245 ridge. *Earth Planet. Sci. Lett.* **194**(1-2), 39-51. [https://doi.org/10.1016/S0012-821X\(01\)00538-6](https://doi.org/10.1016/S0012-821X(01)00538-6).
- 3246 Sherman, D. M. (2013). Equilibrium isotopic fractionation of copper during oxidation/reduction, aqueous  
3247 complexation and ore-forming processes: Predictions from hybrid density functional theory.  
3248 *Geochim. Cosmochim. Acta*, **118**, 85-97.
- 3249 Sherman, D. M., Little, S. H., & Vance, D. (2015). Reply to comment on "Molecular controls on Cu and  
3250 Zn isotopic fractionation in Fe-Mn crusts". *Earth Planet. Sci. Lett.*, **411**, 313-315.
- 3251 Sherman, D. M., & Peacock, C. L. (2010). Surface complexation of Cu on birnessite ( $\delta$ -MnO<sub>2</sub>): Controls  
3252 on Cu in the deep ocean. *Geochim. Cosmochim. Acta*, **74**(23), 6721-6730.
- 3253 Schlitzer, R., et al. (2018). The GEOTRACES intermediate data product 2017. *Chem. Geol.*, **493**, 210-223.
- 3254 Schlitzer, R. (2019) eGEOTRACES - Electronic Atlas of GEOTRACES Sections and Animated 3D Scenes,  
3255 <http://egeotraces.org> [accessed 2020-08-31].
- 3256 Schmitt, A. D., Galer, S. J., & Abouchami, W. (2009a). Mass-dependent cadmium isotopic variations in  
3257 nature with emphasis on the marine environment. *Earth Planet. Sci. Lett.*, **277**(1-2), 262-272.
- 3258 Schmitt, A. D., Galer, S. J., & Abouchami, W. (2009b). High-precision cadmium stable isotope  
3259 measurements by double spike thermal ionisation mass spectrometry. *Journal of Analytical Atomic*  
3260 *Spectrometry*, **24**(8), 1079-1088.
- 3261 Schmitz, B. (1987). Barium, equatorial high productivity, and the northward wandering of the Indian  
3262 continent. *Paleoceanography*, **2**(1), 63-77.
- 3263 Scholz, F., McManus, J., & Sommer, S. (2013). The manganese and iron shuttle in a modern euxinic basin  
3264 and implications for molybdenum cycling at euxinic ocean margins. *Chem. Geol.*, **355**, 56-68,  
3265 doi:10.1016/j.chemgeo.2013.07.006.
- 3266 Scholz, F., Siebert, C., Dale, A. W., & Frank, M. (2017). Intense molybdenum accumulation in sediments  
3267 unearth a nitrogenous water column and implications for the reconstruction of paleo-redox  
3268 conditions based on molybdenum isotopes. *Geochim. Cosmochim. Acta*, **213**, 400-417.
- 3269 Schulz, K. G., Zondervan, I., Gerringa, L. J. A., Timmermans, K. R., Veldhuis, M. J. W., & Riebesell, U.  
3270 (2004). Effect of trace metal availability on coccolithophorid calcification. *Nature*, **430**, 673-676.
- 3271 Scott, C., & Lyons, T. W. (2012). Contrasting molybdenum cycling and isotopic properties in euxinic  
3272 versus non-euxinic sediments and sedimentary rocks: Refining the paleoproxies. *Chem. Geol.*,  
3273 **324-325**, 19-27, doi:10.1016/j.chemgeo.2012.05.012.

- 3274 Semeniuk, D. M., Cullen, J. T., Johnson, W. K., Gagnon, K., Ruth, T. J., & Maldonado, M. T. (2009).  
 3275 Plankton copper requirements and uptake in the subarctic Northeast Pacific Ocean. *Deep Sea*  
 3276 *Research Part I: Oceanographic Research Papers*, **56**(7), 1130-1142.
- 3277 Semeniuk, D. M., Bundy, R. M., Payne, C. D., Barbeau, K. A., & Maldonado, M. T. (2015). Acquisition  
 3278 of organically complexed copper by marine phytoplankton and bacteria in the northeast subarctic  
 3279 Pacific Ocean. *Mar. Chem.* **173**, 222–233.
- 3280 Serno, S., Winckler, G., Anderson, R. F., Hayes, C. T., Ren, H., Gersonde, R., & Haug, G. H. (2014). Using  
 3281 the natural spatial pattern of marine productivity in the Subarctic North Pacific to evaluate  
 3282 paleoproductivity proxies. *Paleoceanography*, **29**(5), 438-453.
- 3283 Severmann, S., Johnson, C. M., Beard, B. L., German, C. R., Edmonds, H. N., Chiba, H., & Green, D. R.  
 3284 H. (2004). The effect of plume processes on the Fe isotope composition of hydrothermally derived  
 3285 Fe in the deep ocean as inferred from the Rainbow vent site, Mid-Atlantic Ridge, 36°14' N. *Earth*  
 3286 *Planet. Sci. Lett.*, **225**(1-2), 63-76. <https://doi.org/10.1016/j.epsl.2004.06.001>.
- 3287 Severmann, S., Johnson, C. M., Beard, B. L., & McManus, J. (2006). The effect of early diagenesis on the  
 3288 Fe isotope compositions of porewaters and authigenic minerals in continental margin sediments.  
 3289 *Geochim. Cosmochim. Acta*, **70**(8). 2006-2022. <https://doi.org/10.1016/j.gca.2006.01.007>.
- 3290 Severmann, S., McManus, J., Berelson, W. M., & Hammond, D. E. (2010). The continental shelf benthic  
 3291 iron flux and its isotope composition. *Geochim. Cosmochim. Acta*, **74**(14), 3984-4004.  
 3292 <https://doi.org/10.1016/j.gca.2010.04.022>.
- 3293 Shaked, Y., Xu, Y., Leblanc, K., & Morel, F. M. M. (2006) Zinc availability and alkaline phosphatase  
 3294 activity in *Emiliania huxleyi*: Implications for Zn-P co-limitation in the ocean. *Limnol. Oceanogr.*,  
 3295 **51**, 299–309.
- 3296 Shemesh, A., Mortlock, R. A., Smith, R. J., & Froelich, P. N. (1988). Determination of Ge/Si in marine  
 3297 siliceous microfossils: Separation, cleaning and dissolution of diatoms and radiolaria. *Mar. Chem.*  
 3298 **25**, 305-323.
- 3299 Shen, G. T., Boyle, E. A., & Lea, D. W. (1987). Cadmium in corals as a tracer of historical upwelling and  
 3300 industrial fallout. *Nature*, **328**(6133), 794.
- 3301 Shiel, A. E., Weis, D., & Orians, K. J. (2010). Evaluation of zinc, cadmium and lead isotope fractionation  
 3302 during smelting and refining. *Science of the Total Environment*, **408**(11), 2357-2368.
- 3303 Sieber, M., Conway, T. M., de Souza, G. F., Hassler, C. S., Ellwood, M. J., & Vance, D. (2019a). High-  
 3304 resolution Cd isotope systematics in multiple zones of the Southern Ocean from the Antarctic  
 3305 Circumnavigation Expedition. *Earth Planet. Sci. Lett.*, **527**, doi:10.1016/j.epsl.2019.115799.
- 3306 Sieber, M., Conway, T. M., de Souza, G. F., Obata, H., Takano, S., Sohrin, Y., & Vance, D. (2019b).  
 3307 Physical and biogeochemical controls on the distribution of dissolved cadmium and its isotopes in  
 3308 the Southwest Pacific Ocean. *Chem. Geol.*, **511**, 494-509.
- 3309 Sieber, M., Conway, T. M., de Souza, G. F., Hassler, C. S., Ellwood, M. J., & Vance, D. (2020). Cycling  
 3310 of zinc and its isotopes across multiple zones of the Southern Ocean: Insights from the Antarctic  
 3311 Circumnavigation Expedition. *Geochim. Cosmochim. Acta*, **268**, 310-324,  
 3312 doi:10.1016/j.gca.2019.09.039.
- 3313 Siebert, C., Nägler, T. F., von Blanckenburg, F., and Kramers, J. D. (2003). Molybdenum isotope records  
 3314 as a potential new proxy for paleoceanography. *Earth Planet. Sci. Lett.*, **211**, 159–171,  
 3315 doi:10.1016/S0012-821X(03)00189-4.

- 3316 Siebert, C., Pett-Ridge, J. C., Opfergelt, S., Guicharnaud, R. A., Halliday, A. N., & Burton, K. W. (2015).  
 3317 Molybdenum isotope fractionation in soils: Influence of redox conditions, organic matter, and  
 3318 atmospheric inputs. *Geochim. Cosmochim. Acta*, **162**, 1–24, doi:10.1016/j.gca.2015.04.007.
- 3319 Sigman, D. M., & Boyle, E. A. (2000). Glacial/interglacial variations in atmospheric carbon dioxide.  
 3320 *Nature*, **407**(6806), 859-869.
- 3321 Sigman, D. M., & Hain, M. P. (2012). The Biological Productivity of the Ocean: Section 1, *Nature*  
 3322 *Education Knowledge*, **3**(10):21.
- 3323 Sigman, D. M., Hain, M. P., & Haug, G. H. (2010). The polar ocean and glacial cycles in atmospheric CO<sub>2</sub>  
 3324 concentration, *Nature*, **466**, 47-55, doi:10.1038/nature09149.
- 3325 Skulan, J. L., Beard, B. L., & Johnson, C. M. (2002). Kinetic and equilibrium Fe isotope fractionation  
 3326 between aqueous Fe (III) and hematite. *Geochim. Cosmochim. Acta*, **66**(17), 2995-3015.
- 3327 Sinoir, M., Butler, E. C. V., Bowie, A. R., Mongin, M., Nesterenko, P. N., & Hassler, C. S. (2012). Zinc  
 3328 marine biogeochemistry in seawater: A review. *Mar. Freshw. Res.*, **63**, 644–657.
- 3329 Sorensen, J. V., Gueguen, B., Stewart, B. D., Peña, J., Rouxel, O., & Toner, B. M. (2020). Large nickel  
 3330 isotope fractionation caused by surface complexation reactions with hexagonal birnessite. *Chem.*  
 3331 *Geol.*, **537**, 119481.
- 3332 Staubwasser, M., Schoenberg, R., von Blanckenburg, F., Krüger, S., & Pohl, C. (2013) Isotope fractionation  
 3333 between dissolved and suspended particulate Fe in the oxic and anoxic water column of the Baltic  
 3334 Sea. *Biogeosciences*, **10**, 233-245. doi:10.5194/bg-10-233-2013.
- 3335 Sternberg, E., Jeandel, C., Robin, E., & Souhaut, M. (2008). Seasonal cycle of suspended barite in the  
 3336 Mediterranean sea. *Geochim. Cosmochim. Acta*, **72**(16), 4020-4034.
- 3337 Stevenson, E. I., Fantle, M. S., Das, S. B., Williams, H. M., & Aciego, S. M. (2017). The iron isotopic  
 3338 composition of subglacial streams draining the Greenland ice sheet. *Geochim. Cosmochim. Acta*,  
 3339 **213**, 237-254.
- 3340 Sun, W.-P., Hu, C.-Y., Han, Z.-B., Shen, C., & Pan, J.-M. (2016). Zn/Si records in diatom opal from Prydz  
 3341 Bay, East Antarctica. *Marine Geology* **381**, 34-41.
- 3342 Sunda, W. G. (2012). Feedback interactions between trace metal nutrients and phytoplankton in the ocean.  
 3343 *Front. Microbiol.*, **3**(204), doi:10.3389/fmicb.2012.00204.
- 3344 Sunda, W. G., & Huntsman, S. A. (1995). Iron uptake and growth limitation in oceanic and coastal  
 3345 phytoplankton. *Mar. Chem.*, **50**, 189–206.
- 3346 Sunda, W. G., & Huntsman, S. A. (2000). Effect of Zn, Mn, and Fe on Cd accumulation in phytoplankton:  
 3347 implications for oceanic Cd cycling. *Limnol. Oceanogr.*, **45**(7), 1501-1516.
- 3348 Sweere, T. C., Dickson, A. J., Jenkyns, H. C., Porcelli, D., Elrick, M., van den Boorn, S. H., & Henderson,  
 3349 G. M. (2018). Isotopic evidence for changes in the zinc cycle during Oceanic Anoxic Event 2 (Late  
 3350 Cretaceous). *Geology*, **46**(5), 463-466.
- 3351 Tabouret, H., Pomerleau, S., Jolivet, A., Pécheyran, C., Riso, R., et al. (2012). Specific pathways for the  
 3352 incorporation of dissolved barium and molybdenum into the bivalve shell: An isotopic tracer  
 3353 approach in the juvenile Great Scallop (*Pecten maximus*). *Mar. Environ. Res.*, **78**, 15-25,  
 3354 doi:10.1016/j.marenvres.2012.03.006.
- 3355 Tagliabue, A., Aumont, O., DeAth, R., Dunne, J. P., Dutkiewicz, S., Galbraith, E., Misumi, K., Moore, J.  
 3356 K., Ridgwell, A., Sherman, E., Stock, C., Vichi, M., Völker, C., & Yool, A. (2016). How well do  
 3357 global ocean biogeochemistry models simulate dissolved iron distributions?. *Global Biogeochem.*  
 3358 *Cycles*, **30**(2), 149-174.



- 3359 Tagliabue, A., Bowie, A. R., Boyd, P. W., Buck, K. N., Johnson, K. S., & Saito, M. A. (2017). The integral  
3360 role of iron in ocean biogeochemistry. *Nature*, **543**(7643), 51-59.
- 3361 Tagliabue, A., Bowie, A. R., DeVries, T., Ellwood, M. J., Landing, W. M., Milne, A., Ohnemus, D. C.,  
3362 Twining, B. S., & Boyd, P. W. (2019). The interplay between regeneration and scavenging fluxes  
3363 drives ocean iron cycling. *Nat. Commun.*, 10(1), 1-8.
- 3364 Takano, S., Tanimizu, M., Hirata, T., & Sohrin, Y. (2014). Isotopic constraints on biogeochemical cycling  
3365 of copper in the ocean. *Nat. Commun.* **5**, 5663.
- 3366 Takano, S., Tanimizu, M., Hirata, T., Shin, K. C., Fukami, Y., Suzuki, K., & Sohrin, Y. (2017). A simple  
3367 and rapid method for isotopic analysis of nickel, copper, and zinc in seawater using chelating  
3368 extraction and anion exchange. *Anal. Chim. Acta* **967**, 1–11.
- 3369 Tankéré, S. P. C., Muller, F. L. L., Burton, J. D., Statham, P. J., Guieu, C., & Martin, J. M. (2001). Trace  
3370 metal distributions in shelf waters of the northwestern Black Sea. *Cont. Shelf Res.* **21**, 1501–1532.
- 3371 Tegler, L. A., Dunlea, A. G., Romaniello, S. J., Anbar, A. D., & Horner, T. J. (2018). More Clear than Mud:  
3372 Dating Deep-Sea Clays with Osmium Isotopes to Unravel Seawater Metal Cycling through the  
3373 Cenozoic. In *2018 Ocean Sciences Meeting*. AGU.
- 3374 Tesoriero, A. J., & Pankow, J. F. (1996). Solid solution partitioning of Sr<sup>2+</sup>, Ba<sup>2+</sup>, and Cd<sup>2+</sup> to calcite.  
3375 *Geochim. Cosmochim. Acta*, **60**(6), 1053-1063.
- 3376 Tessin, A., Chappaz, A., Hendy, I., & Sheldon, N. (2019). Molybdenum speciation as a paleo-redox proxy:  
3377 A case study from Late Cretaceous Western Interior Seaway black shales. *Geology*, **47**, 59–62,  
3378 doi:10.1130/G45785.1.
- 3379 Thébault, J., Chauvaud, L., L'Helguen, S., Clavier, J., Barats, A., et al. (2009). Barium and molybdenum  
3380 records in bivalve shells: Geochemical proxies for phytoplankton dynamics in coastal  
3381 environments? *Limnol. Oceanogr.*, **54**(3), 1002-1014, doi:10.4319/lo.2009.54.3.1002.
- 3382 Thibon, F., Blichert-Toft, J., Albarede, F., Foden, J., & Tsikos, H. (2019). A critical evaluation of copper  
3383 isotopes in Precambrian Iron Formations as a paleoceanographic proxy. *Geochim. Cosmochim.*  
3384 *Acta*, **264**, 130-140.
- 3385 Thompson, C. M., & Ellwood, M. J. (2014). Dissolved copper isotope biogeochemistry in the Tasman Sea,  
3386 SW Pacific Ocean. *Mar. Chem.* **165**, 1–9.
- 3387 Torres, M. E., Brumsack, H. J., Bohrmann, G., & Emeis, K. C. (1996). Barite fronts in continental margin  
3388 sediments: a new look at barium remobilization in the zone of sulfate reduction and formation of  
3389 heavy barites in diagenetic fronts. *Chem. Geol.*, **127**(1-3), 125-139.
- 3390 Tossell, J. A. (2005). Calculating the partitioning of the isotopes of Mo between oxidic and sulfidic species  
3391 in aqueous solution. *Geochim. Cosmochim. Acta*, **69**, 2981–2993, doi:10.1016/j.gca.2005.01.016.
- 3392 Tribouillard, N., Algeo, T. J., Lyons, T., & Riboulleau, A. (2006). Trace metals as paleoredox and  
3393 paleoproductivity proxies: an update. *Chem. Geol.*, **232**(1-2), 12-32.
- 3394 Turekian, K. K. (1968). Deep-sea deposition of barium, cobalt and silver. *Geochim. Cosmochim. Acta*,  
3395 **32**(6), 603-612.
- 3396 Turner, D. R., Whitfield, M., & Dickson, A. G. (1981). The equilibrium speciation of dissolved components  
3397 in freshwater and sea water at 25°C and 1 atm pressure. *Geochim. Cosmochim. Acta*, **45**, 855–881.
- 3398 Twining, B. S., & Baines, S. B. (2013). The trace metal composition of marine phytoplankton. *Annual*  
3399 *Review of Marine Science*, **5**, 191-215.

- 3400 Twining, B. S., Baines, S. B., & Fisher, N. S. (2004) Element stoichiometries of individual plankton cells  
 3401 collected during the Southern Ocean Iron Experiment (SOFEX). *Limnol. Oceanogr.*, **49**, 2115–  
 3402 2128.
- 3403 Twining, B. S., Baines, S. B., Vogt, S., & Nelson, D. M. (2012) Role of diatoms in nickel biogeochemistry  
 3404 in the ocean. *Glob. Biogeochem. Cycles*, **26** (4), GB4001, doi:10.1029/2011GB004233.
- 3405 Twining, B. S., Nodder, S. D., King, A. L., Hutchins, D. A., LeClerc, G. R., DeBruyn, J. M., Maas, E. W.,  
 3406 Vogt, S., Wilhelm S. W., & Boyd, P. W. (2014). Differential remineralization of major and trace  
 3407 elements in sinking diatoms. *Limnol. Oceanogr.*, **59**(3), 689-704.
- 3408 van Geen, A., Luoma, S. N., Fuller, C. C., Anima, R., Clifton, H. E., & Trumbore, S. (1992). Evidence  
 3409 from Cd/Ca ratios in foraminifera for greater upwelling off California 4,000 years ago. *Nature*,  
 3410 **358**(6381), 54.
- 3411 van Geen, A., McCorkle, D. C., & Klinkhammer, G. P. (1995). Sensitivity of the phosphate-cadmium-  
 3412 carbon isotope relation in the ocean to cadmium removal by suboxic sediments. *Paleoceanography*,  
 3413 **10**(2), 159-169.
- 3414 van Helmond, N. A. G. M., Jilbert, T., & Slomp, C. P. (2018). Hypoxia in the Holocene Baltic Sea:  
 3415 Comparing modern versus past intervals using sedimentary trace metals. *Chem. Geol.*, **493**, 478-  
 3416 490, doi:10.1016/j.chemgeo.2018.06.028.
- 3417 Vance, D., Archer, C., Bermin, J., Perkins, J., Statham, P. J., Lohan, M. C., Ellwood, M. J., & Mills, R. A.  
 3418 (2008). The copper isotope geochemistry of rivers and the oceans. *Earth Planet. Sci. Lett.* **274**,  
 3419 204–213.
- 3420 Vance, D., Little, S. H., Archer, C., Cameron, V., Andersen, M. B., Rijkenberg, M. J. A., & Lyons, T. W.  
 3421 (2016). The oceanic budgets of nickel and zinc isotopes: the importance of sulfidic environments  
 3422 as illustrated by the Black Sea. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* **374**, 20150294.
- 3423 Vance, D., Little, S. H., de Souza, G. F., Khatiwala, S., Lohan, M. C., & Middag, R. (2017). Silicon and  
 3424 zinc biogeochemical cycles coupled through the Southern Ocean. *Nat. Geosci.*, **10**, 202–206.
- 3425 Vance, D., de Souza, G. F., Zhao, Y., Cullen, J. T., & Lohan, M. C. (2019). The relationship between zinc,  
 3426 its isotopes, and the major nutrients in the North-East Pacific. *Earth Planet. Sci. Lett.*, **525**,  
 3427 doi:10.1016/j.epsl.2019.115748.
- 3428 Velasquez, I. B., Ibisani, E., Maas, E. W., Boyd, P. W., Nodder, S., & Sander, S. G. (2016). Ferrioxamine  
 3429 Siderophores Detected amongst Iron Binding Ligands Produced during the Remineralization of  
 3430 Marine Particles. *Front. Mar. Sci.* **3**, 172. <https://doi.org/10.3389/fmars.2016.00172>.
- 3431 Vihtakari, M., Ambrose Jr., W. G., Renaud, P. E., Locke V, W. L., Carroll, M. L., et al. (2017). A key to  
 3432 the past? Element ratios as environmental proxies in two Arctic bivalves. *Palaeogeogr.*  
 3433 *Palaeoclimatol. Palaeoecol.*, **465**, 316-332, doi:10.1016/j.palaeo.2016.10.020.
- 3434 Voegelin, A. R., Nägler, T. F., Samankassou, E., & Villa, I. M. (2009). Molybdenum isotopic composition  
 3435 of modern and Carboniferous carbonates. *Chem. Geol.*, **265**, 488-498,  
 3436 doi:10.1016/j.chemgeo.2009.05.015.
- 3437 Volk, T., & Hoffert, M. I. (1985). Ocean carbon pumps: Analysis of relative strengths and efficiencies in  
 3438 ocean-driven atmospheric CO<sub>2</sub> changes. *The carbon cycle and atmospheric CO<sub>2</sub>: natural variations*  
 3439 *Archean to present*, **32**, 99-110.
- 3440 Von Allmen, K., Böttcher, M. E., Samankassou, E., & Nägler, T. F. (2010). Barium isotope fractionation  
 3441 in the global barium cycle: First evidence from barium minerals and precipitation experiments.  
 3442 *Chem. Geol.*, **277**(1-2), 70-77.



- 3443 Vorliceck, T. P., Chappaz, A., Groskreutz, L. M., Young, N., & Lyons, T. W. (2015). A new analytical  
 3444 approach to determining Mo and Re speciation in sulfidic waters. *Chem. Geol.*, **403**, 52–57,  
 3445 doi:10.1016/j.chemgeo.2015.03.003.
- 3446 Vorliceck, T. P., Helz, G. R., Chappaz, A., Vue, P., Vezina, A., & Hunter, W. (2018). Molybdenum Burial  
 3447 Mechanism in Sulfidic Sediments: Iron-Sulfide Pathway. *ACS Earth and Space Chemistry*, **2**, 565–  
 3448 576, doi:10.1021/acsearthspacechem.8b00016.
- 3449 Waeles, M., Baker, A. R., Jickells, T., & Hoogewerff, J. (2007). Global dust teleconnections: aerosol iron  
 3450 solubility and stable isotope composition. *Environmental Chemistry*, **4**(4), 233-237.
- 3451 Wagner, M. E. (2013). *Silver as a novel tracer for Late Quaternary Southern Ocean biological and*  
 3452 *geophysical processes*. PhD thesis, University of Michigan, 189p.
- 3453 Wagner, M., Hendy, I. L., McKay, J. L., & Pedersen, T.F. (2013). Influence of biological productivity on  
 3454 silver and redox-sensitive trace metal accumulation in Southern Ocean surface sediments, Pacific  
 3455 sector. *Earth Planet. Sci. Lett.*, **380**, 31-40.
- 3456 Wagner, M., Chappaz, A., & Lyons, T.W. (2017). Molybdenum speciation and burial pathway in weakly  
 3457 sulfidic environments: Insights from XAFS. *Geochim. Cosmochim. Acta*, **206**, 18–29,  
 3458 doi:10.1016/j.gca.2017.02.018.
- 3459 Waldron, K. J., & Robinson, N. J. (2009). How do bacterial cells ensure that metalloproteins get the correct  
 3460 metal? *Nature Reviews Microbiology*, **7**(1), 25.
- 3461 Wang, D., Xia, W., Lu, S., Wang, G., Liu, Q., Moore, W. S., and Arthur Chen, C. T. (2016). The  
 3462 nonconservative property of dissolved molybdenum in the western Taiwan Strait: Relevance of  
 3463 submarine groundwater discharges and biological utilization. *Geochemistry, Geophysics,*  
 3464 *Geosystems*, **17**, 28–43, doi:10.1002/2014GC005708.
- 3465 Wang, R. M., Archer, C., Bowie, A. R., & Vance, D. (2019a). Zinc and nickel isotopes in seawater from  
 3466 the Indian Sector of the Southern Ocean: The impact of natural iron fertilization versus Southern  
 3467 Ocean hydrography and biogeochemistry. *Chem. Geol.*, **511**, 452-464.
- 3468 Wang, Z., Li, J., Wei, G., Deng, W., Chen, X., Zeng, T., Wang, X., Ma, J., Zhang, L., Tu, X., Wang, Q., &  
 3469 McCulloch, M. (2019b). Biologically controlled Mo isotope fractionation in coral reef systems.  
 3470 *Geochim. Cosmochim. Acta*, **262**, 128-142, doi:10.1016/j.gca.2019.07.037.
- 3471 Wang, S.-J., Rudnick, R. L., Gaschnig, R. M., Wang, H., & Wasylenki, L. E. (2019c). Methanogenesis  
 3472 sustained by sulfide weathering during the Great Oxidation Event. *Nat. Geosci.*, **12**, 296-300,  
 3473 doi:10.1038/s41561-019-0320-z.
- 3474 Wasylenki, L. E., Anbar, A. D., Liermann, L. J., Mathur, R., Gordon, G. W., & Brantley, S. L. (2007).  
 3475 Isotope fractionation during microbial metal uptake measured by MC-ICP-MS. *J. Anal. At.*  
 3476 *Spectrom.*, **22**, 905-910, doi:10.1039/b705476a.
- 3477 Wasylenki, L. E., Rolfé, B. A., Weeks, C. L., Spiro, T. G., and Anbar, A. D. (2008). Experimental  
 3478 investigation of the effects of temperature and ionic strength on Mo isotope fractionation during  
 3479 adsorption to manganese oxides. *Geochim. Cosmochim. Acta*, **72**, 5997–6005,  
 3480 doi:10.1016/j.gca.2008.08.027.
- 3481 Wasylenki, L. E., Swihart, J. W., & Romaniello, S. J. (2014). Cadmium isotope fractionation during  
 3482 adsorption to Mn oxyhydroxide at low and high ionic strength. *Geochim. Cosmochim. Acta*, **140**,  
 3483 212-226.
- 3484 Weber, T., John, S. G., Tagliabue, A., & DeVries, T. (2018). Biological uptake and reversible scavenging  
 3485 of zinc in the global ocean. *Science*, **361**, 72–76.

- 3486 Wefing, A. M., Arps, J., Blaser, P., Wienberg, C., Hebbeln, D., & Frank, N. (2017). High precision U-series  
 3487 dating of scleractinian cold-water corals using an automated chromatographic U and Th extraction.  
 3488 *Chem. Geol.*, **475**, 140-148.
- 3489 Wei, W., & Algeo, T. J. (in press). Secular variation in the elemental composition of marine shales since  
 3490 840 Ma: Tectonic and seawater influences. *Geochim. Cosmochim. Acta*, doi:  
 3491 10.1016/j.gca.2020.01.033.
- 3492 Welch, S. A., Beard, B. L., Johnson, C. M., & Braterman, P. S. (2003). Kinetic and equilibrium Fe isotope  
 3493 fractionation between aqueous Fe(II) and Fe(III). *Geochim. Cosmochim. Acta*, **67**, 4231-4250.  
 3494 [https://doi.org/10.1016/S0016-7037\(03\)00266-7](https://doi.org/10.1016/S0016-7037(03)00266-7).
- 3495 Wichard, T., Mishra, B., Myneni, S. C. B., Bellenger, J. P., & Kraepiel, A. M. L. (2009). Storage and  
 3496 bioavailability of molybdenum in soils increased by organic matter complexation. *Nat. Geosci.*, **2**,  
 3497 625–630, doi:10.1038/ngeo589.
- 3498 Willbold, M., & Elliott, T. (2017). Molybdenum isotope variations in magmatic rocks. *Chem. Geol.*, **449**,  
 3499 253-268.
- 3500 Winckler, G., Anderson, R. F., Jaccard, S. L., & Marcantonio, F. (2016). Ocean dynamics, not dust, have  
 3501 controlled equatorial Pacific productivity over the past 500,000 years. *Proceedings of the National  
 3502 Academy of Sciences*, **113**(22), 6119-6124.
- 3503 Wolgemuth, K., & Broecker, W. S. (1970). Barium in sea water. *Earth Planet. Sci. Lett.*, **8**(5), 372-378.
- 3504 Woodland, S. J., Rehkämper, M., Halliday, A. N., Lee, D.-C., Hatterndorf, B., & Günther, D. (2005)  
 3505 Accurate measurement of silver isotopic compositions in geological materials including lowPd/Ag  
 3506 meteorites. *Geochim. Cosmochim. Acta*, **69**(8). doi:10.1016/j.gca.2004.10.012.
- 3507 Wuttig, K., Townsend, A. T., van der Merwe, P., Gault-Ringold, M., Holmes, T., Schallenberg, C., Latour,  
 3508 P., Tonnard, M., Rijkenberg, M. J. A., Lannuzel, D., & Bowie, A. R. (2019). Critical evaluation of  
 3509 a seaFAST system for the analysis of trace metals in marine samples. *Talanta*, **197**, 653-668.
- 3510 Xie, R. C., Galer, S. J., Abouchami, W., Rijkenberg, M. J., De Baar, H. J., De Jong, J., & Andreae, M. O.  
 3511 (2017). Non-Rayleigh control of upper-ocean Cd isotope fractionation in the western South  
 3512 Atlantic. *Earth Planet. Sci. Lett.*, **471**, 94-103.
- 3513 Xie, R. C., Galer, S. J. G., Abouchami, W., & Frank, M. (2019a) Limited impact of eolian and riverine  
 3514 sources on the biogeochemical cycling of Cd in the tropical Atlantic. *Chem. Geol.* **511**, 371-379.  
 3515 <https://doi.org/10.1016/j.chemgeo.2018.10.018>.
- 3516 Xie, R., C., Rehkämper, M., Grasse, P., van de Flierdt, T., Frank, M., & Xue, Z. (2019b). Isotopic evidence  
 3517 for complex biogeochemical cycling of Cd in the eastern tropical South Pacific. *Earth Planet. Sci.  
 3518 Lett.* **512**, 134-146. <https://doi.org/10.1016/j.epsl.2019.02.001>.
- 3519 Xu, Y., Feng, L., Jeffrey, P. D., Shi, Y., & Morel, F. M. (2008). Structure and metal exchange in the  
 3520 cadmium carbonic anhydrase of marine diatoms. *Nature*, **452**(7183), 56.
- 3521 Xue, Z., Rehkämper, M., Schönbacher, M., Statham, P. J., & Coles, B. J. (2012). A new methodology for  
 3522 precise cadmium isotope analyses of seawater. *Analytical and Bioanalytical Chemistry*, **402**(2),  
 3523 883-893.
- 3524 Xue, Z., Rehkämper, M., Horner, T. J., Abouchami, W., Middag, R., van de Flierdt, T., & de Baar, H. J.  
 3525 (2013). Cadmium isotope variations in the Southern Ocean. *Earth Planet. Sci. Lett.*, **382**, 161-172.
- 3526 Yang, S.-C., Lee, D.-L., & Ho, T.-Y. (2012). The isotopic composition of Cadmium in the water column  
 3527 of the South China Sea. *Geochim. Cosmochim. Acta*, **98**, 66-77.  
 3528 <https://doi.org/10.1016/j.gca.2012.09.022>.

- 3529 Yang, S.-C., Lee, D.-L., Ho, T.-Y., Wen, L.-S., & Yang, H.-H. (2014). The isotopic composition of  
 3530 dissolved cadmium in the water column of the West Philippine Sea. *Front. Mar. Sci.* **1**(61).  
 3531 <https://doi.org/10.3389/fmars.2014.00061>
- 3532 Yang, S.-C., Lee, D.-C., & Ho, T.-Y. (2015). Cd isotopic composition in the suspended and sinking particles  
 3533 of the surface water of the South China Sea: The effects of biotic activities. *Earth Planet. Sci. Lett.*,  
 3534 **428**, 63-72.
- 3535 Yang, S.-C., Zhang, J., Sohrin, Y., & Ho, T.-Y. (2018). Cadmium cycling in the water column of the  
 3536 Kuroshio-Oyashio Extension region: Insights from dissolved and particulate isotopic composition.  
 3537 *Geochim. Cosmochim. Acta*, **233**, 66-80.
- 3538 Yang, T., Chen, Y. Zhou, S., & Li, H. (2019). Impacts of aerosol copper on marine phytoplankton: A  
 3539 review. *Atmosphere*, **10**(7), 414, doi:10.3390/atmos10070414.
- 3540 Yang, S.-C., Hawco, N. J., Pinedo-González, P., Bian, X., Huang, K.-F., & John, S. G. (2020). A new  
 3541 purification method for Ni and Cu stable isotopes in seawater provides evidence for widespread Ni  
 3542 isotope fractionation by phytoplankton in the North Pacific. *Chem. Geol.* **547**, 119662.  
 3543 <https://doi.org/10.1016/j.chemgeo.2020.119662>.
- 3544 Yee, D., & Morel, F. M. M. (1996). In vivo substitution in carbonic of zinc by cobalt of a marine anhydrase  
 3545 diatom. *Limnol. Oceanogr.*, **41**, 573–577.
- 3546 Zhang, R., John, S. G., Zhang, J., Ren, J., Wu, y., Zhu, Z., Liu, S., Zhu, X., Marsay, C. M., & Wenger, F.  
 3547 (2015). Transport and reaction of iron and iron stable isotopes in glacial meltwaters on Svalbard  
 3548 near Kongsfjorden: From rivers to estuary to ocean. *Earth Planet. Sci. Lett.*, **424**, 201-211.  
 3549 <https://doi.org/10.1016/j.epsl.2015.05.031>.
- 3550 Zhang, R., Jensen, L. T., Fitzsimmons, J. N., Sherrell, R. M., & John, S. (2019). Dissolved cadmium and  
 3551 cadmium stable isotopes in the western Arctic Ocean. *Geochim. Cosmochim. Acta*, **258**, 258-273,  
 3552 <https://doi.org/10.1016/j.gca.2019.05.028>.
- 3553 Zhang, Y., Amakawa, H., & Nozaki, Y. (2001). Oceanic profiles of dissolved silver: precise measurements  
 3554 in the basins of western North Pacific, Sea of Okhotsk, and the Japan Sea. *Marine Chemistry*, **75**,  
 3555 151-163.
- 3556 Zhang, Y., Obata, H., & Nozaki, Y. (2004). Silver in the Pacific Ocean and the Bering Sea. *Geochemical  
 3557 Journal*, **38**, 623-633.
- 3558 Zhao, Y., Vance, D., Abouchami, W., & de Baar, H. J. W. (2014). Biogeochemical cycling of zinc and its  
 3559 isotopes in the Southern Ocean. *Geochim. Cosmochim. Acta*, **125**, 653–672.  
 3560 <http://dx.doi.org/10.1016/j.gca.2013.07.045>.
- 3561 Zhu, X. K., O'Nions, R. K., Guo, Y., & Reynolds, B. C. (2000). Secular variation of iron isotopes in North  
 3562 Atlantic deep water. *Science*, **287**(5460).
- 3563 Zirino, A., & Yamamoto, S. (1972). A pH-dependent model for the chemical speciation of copper, zinc,  
 3564 cadmium, and lead in seawater. *Limnol. Oceanogr.* **17**, 661–671.