



HAL
open science

Diel cycles of carbon, nutrient and metal in humic lakes of permafrost peatlands

L.S. Shirokova, D. Payandi-Rolland, A.G. Lim, R.M. Manasypov, J. Allen, J.-L. Rols, P. Bénézet, J. Karlsson, O.S. Pokrovsky

► **To cite this version:**

L.S. Shirokova, D. Payandi-Rolland, A.G. Lim, R.M. Manasypov, J. Allen, et al.. Diel cycles of carbon, nutrient and metal in humic lakes of permafrost peatlands. *Science of the Total Environment*, 2020, 737, pp.139671. 10.1016/j.scitotenv.2020.139671 . hal-03003987

HAL Id: hal-03003987

<https://hal.science/hal-03003987>

Submitted on 25 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Diel cycles of carbon, nutrient and metal in humic lakes of permafrost peatlands

L.S. Shirokova^{a, b}, D. Payandi-Rolland^a, A.G. Lim^c, R.M. Manasyrov^c, J. Allen^d, J.-L. Rols^d, P. Benezeth^a, J. Karlsson^e, O.S. Pokrovsky^{a, *}

^a Géosciences Environnement Toulouse, GET – CNRS – IRD – OMP – Université de Toulouse, 14, Avenue Edouard Belin, 31400 Toulouse, France

^b Institute of Ecological Problems of the North, Russian Academy of Science, 23, Naberezhnaya Sev. Dviny, Arkhangelsk, Russia

^c BIO-GEO-CLIM Laboratory, Tomsk State University, 36 Lenina Pr., Tomsk, Russia

^d EcoLab, Université de Toulouse, CNRS, Toulouse, France

^e Climate Impacts Research Centre (CIRC), Department of Ecology and Environmental Science, Umeå University, SE-901 87 Umeå, Sweden

ARTICLE INFO

Article history:

Received 14 April 2020

Received in revised form 21 May 2020

Accepted 22 May 2020

Available online xxx

Editor: Fernando A.L. Pacheco

Keywords:

Carbon dioxide
Methane
Emission
Thermokarst
Pond
Trace metal
Organic carbon
Peat
Warming

ABSTRACT

Despite the importance of surface waters of permafrost landscapes in carbon (C) emission and dissolved C and metal storage and export, the majority of available observations in high latitude aquatic systems deal with punctual or seasonal sampling without accounting for diurnal variations in temperature and primary productivity-respiration cycles. Towards providing comprehensive understanding of diel variations in CO₂ emission, organic C and element concentrations in lakes of frozen peatlands, we monitored, each 2h over 2 days, the water temperature, pH, CO₂ fluxes, CO₂, CH₄, dissolved organic and inorganic carbon (DOC and DIC, respectively), nutrients, carboxylic acids, bacterial number, and major and trace elements in two acidic (pH=3.6 and 4.0) and humic (DOC=15 and 35 mg L⁻¹) thermokarst lakes of discontinuous permafrost zone in Western Siberia. We discovered a factor of 2 to 3 higher CO₂ concentrations and fluxes during the night compared to daytime in the high-DOC lake. The emission fluxes in the low-DOC lake increased from zero to negative values during the day to highly positive values during the end of night and early morning. The methane concentration varied within a factor of 5 without any link to the diurnal cycle. The bulk of dissolved (< 0.45 μm) hydrochemical parameters remained highly stable with ±10% variation in concentration over 2 days of observation (DOC, DIC, SUVA_{254 nm}, carboxylates (formate, oxalate, pyruvate and glutarate), Mn, Fe, Al, other trace elements). Concentrations of Si, P, K, Cu varied within ±20% whereas those of Zn and Ni ranged by a factor of 2 to 4 without any link to diurnal pattern. Overall, the impact of diel cycle on CH₄, DOC, nutrient and metal concentration was below 10%. However, neglecting night-time period may underestimate net CO₂ emission by ca. 30 to 50% in small organic-rich thaw ponds and switch the CO₂ exchange from uptake/zero to net emission in larger thermokarst lakes. Given the dominance of large lakes in permafrost regions, the global underestimation of the emission flux may be quite high. As such, monitoring CO₂ concentrations and fluxes in thermokarst lakes during months of extended night time (August to October) is mandatory for assessing the net emissions from lentic waters of frozen peatlands.

© 2020.

1. Introduction

The emission of greenhouse gases (GHG), particularly CO₂ and CH₄, from inland waters at high latitudes is a critical issue of these aquatic system behaviors under on-going climate warming. Due to sizeable amount of GHG and organic carbon (OC) stored in frozen organic-rich soils and produced during soil interaction with surface waters, high-latitude regions, notably peatlands, are at the forefront of environmental research as they can provide crucial information for modeling of terrestrial C cycle modification in response to permafrost thaw, water temperature rise and aquatic biotic community change (Gorham, 1991; Roehm et al., 2009; Peura et al., 2019). A major issue of lake biogeochemistry is the variation of GHG dynamics in re-

sponse to photosynthesis/respiration cycle of phyto- and bacterio-plankton (Szyper et al., 1992; Pollard, 2013; Hanson et al., 2006; Sadro et al., 2014) and benthic macrophytes and periphyton (Yvon-Durocher et al., 2014). The diel CO₂ concentration and emission patterns linked to variations in water temperatures and sunlight availability are fairly well documented in temperate rivers and ponds (Baehr and DeGrandpre, 2004; Forget et al., 2009; Lynch et al., 2010; Xiao et al., 2013; Yang et al., 2015; Liu et al., 2016; Spafford and Risk, 2018; DelVecchia et al., 2019). However, due to logistical constraints, the majorities of studies in high latitude, permafrost-affected peatland lakes dealt with one punctual sampling, usually during summer period (e.g., Shirokova et al., 2013; Pokrovsky et al., 2011, 2013), with the exception of several seasonally-rather than daily-resolved measurements (Jonsson et al., 2008; Laurion et al., 2010; Marushchak et al., 2013; Podgrajsek et al., 2014; Manasyrov et al., 2015; Serikova et al., 2019). Although the latter studies addressed the seasonality of GHG emission, the diurnal variability of concentration

* Corresponding author.

Email address: oleg.pokrovsky@get.omp.eu (O.S. Pokrovsky)

and fluxes in these aquatic settings is rarely considered. In fact, there are various factors controlling DOC concentration and gas exchange of the lake with atmosphere and most of these factors are expected to vary diurnally. The 'external' factors are DOC and DIC (CO_2) delivery to the lake via shallow surface flow, subsurface (supra-permafrost) flow and possibly underground discharge via taliks (unfrozen water paths through permafrost). The 'internal' factors are determined by metabolism of plankton and periphyton, photo-oxidation of DOC, GHG generation or uptake in sediments, and gas exchange velocity (k) that could vary depending on the wind speed and day/night temperature variation via convective mixing (i.e., Eugster et al., 2003). Because the superposition of these external and internal gas sources likely varies across the systems and the time of the day, their overall impact on net C exchange has to be resolved via diurnal measurements on contrasting but representative lakes of the permafrost zone. Thus, recent in situ observations of CO_2 dynamics in tundra and alpine streams indicate net daytime uptake of CO_2 by photosynthesis (Peter et al., 2014; Crawford et al., 2017; Rocher-Ros et al., 2019) whereas Arctic lakes showed sizeable diurnal variability in CO_2 concentration and fluxes (Lundin et al., 2016).

In addition to GHG gases that are sensitive to diurnal pattern, dissolved organic carbon and trace elements bound to organic complexes are also likely to be affected by variations of light and temperature. However, the majority of trace metal studies in lakes dealt with one sampling during the day (i.e., Hamilton-Taylor and Willis, 1990; Nriagu et al., 1996; Falkner et al., 1997; Viollier et al., 1997; Albéric et al., 2000) and although strong diel variations of solutes were reported in temperate rivers (Brick and Moore, 1996; Neal et al., 2002; Jones et al., 2004; Gammons et al., 2005; Parker et al., 2007; Carling et al., 2011; Nimick et al., 2011; Vorobyev et al., 2019), subarctic organic-rich lakes remained very poorly investigated with the exception of humic lakes in boreal non-permafrost zone (i.e., Shirokova et al., 2010; Pokrovsky and Shirokova, 2013). At the same time, diurnal patterns of trace metals, involved in redox cycles and linked to photo-reduction and photo-oxidation (McMahon, 1969; Barry et al., 1994; Emmenegger et al., 2001), including colloidal forms of trace elements (Pokrovsky and Shirokova, 2013) are fairly well established in temperate lake waters but totally unknown in permafrost-affected humic lakes.

The majority of dystrophic to mesotrophic aquatic systems in frozen peatlands drastically differ from oligotrophic northern rivers and lakes in the sense that the peatland thermokarst lakes and ponds are *i*) highly humic, and typically contain between 10 and 100 mg L^{-1} of DOC and *ii*) quite acidic, with pH ranging between 3.5 and 5.5 (Pokrovsky et al., 2014). As a result, the GHG and DOC regime in humic lake may be strongly controlled by a balance between the productivity of plankton, periphyton and macrophytes, and the heterotrophic bacteria mineralization of dissolved and particulate OC in the water column and sediments. Indeed, the heterotrophic bacterioplankton is capable to mineralize dissolved organic matter (DOM) of both allochthonous and autochthonous origin (Tranvik, 1988, 1989; Kritzberg et al., 2004) and thus can potentially liberate or consume associated trace metals. At the same time, limited photic layer and generally small abundance of phytoplankton in thermokarst peatland lakes (Pavlova et al., 2016) might preclude the cycle of CO_2 consumption during the day and its production during the night, unlike it is established in oligotrophic lakes of the subarctic (Ask et al., 2012). Further, considering that photo-oxidation of DOM can be the dominant pathway of C processing in arctic waters (Cory et al., 2014), this may increase CO_2 concentration from night to day (Bertilsson and Tranvik, 2000) thus counteracting the GPP effect, especially in shallow humic waters.

Therefore, the working hypotheses of this study is that the variability in external factors like temperature and light will exert primary control on DOC and related trace elements, and this will produce a diel cycle in C and elements concentrations and GHG emissions in thermokarst lakes. Further, we anticipate that the higher the DOC in a peatland lake, the smaller will be the response of water column hydrochemistry to diel variations in light and temperature. This response may be essentially controlled by external vs internal sources and thus likely to vary depending on lake size. In order to test these hypotheses, we studied two thaw ponds with different size and DOM content and we quantified (1) the variation of dissolved C and metal concentration between day and night and (2) the degree of diel variation in CO_2 flux, in order to reveal the control of hydrochemical parameters by external factors such as temperature and light. For this, we monitored CO_2 , CH_4 , organic carbon, major and trace element concentrations and CO_2 fluxes in thermokarst lakes during 2 summer days of anticyclone weather. Obtained results allowed to improve monitoring practices and demonstrated the need of revising the existing estimations of CO_2 emission from high latitude inland waters, which are mainly based on daytime period.

2. Study site, material and methods

We studied typical thermokarst lake (Lake Trisino, N $63^\circ 47' 09.35''$, E $75^\circ 38' 54.57''$, $S_{\text{area}} = 16,920 \text{ m}^2$, $S_{\text{watershed}} = 0.108 \text{ km}^2$, average depth 0.5 m) and thaw pond (Chernoe, N $63^\circ 46' 55.72''$, E $75^\circ 39' 11.73''$, $S_{\text{area}} = 1830 \text{ m}^2$, $S_{\text{watershed}} \leq 500 \text{ m}^2$, average depth 0.3 m) located at the INTERACT Research Station (Khanymey), discontinuous permafrost zone of the Western Siberian Lowland (WSL) as shown in Fig. S1 of Supplement. The permafrost here is abundant within flat-mound peat bogs on the watershed divides. The depth of the active (seasonally unfrozen) layer ranges from $145 \pm 20 \text{ cm}$ within the depressions to $41 \pm 5 \text{ cm}$ at the mounds (Raudina et al., 2018). Within the area of the Khanymey Research station, there are numerous thermokarst lakes and thaw ponds formed due to thawing of histosols (0.1 to 1.4 m peat thickness) over mineral (essentially Pleistocene sands) substrates (Shirokova et al., 2013). The feeding of small thaw ponds and large thermokarst lakes occurs either by supra-permafrost flow from adjacent mounds or by surface runoff from the elevated parts of the lake watershed.

The observations were performed during 48 h on July 26–28th 2018, at stable anticyclone conditions without any precipitation event. The July was chosen as the month of maximal surface water temperature and ecosystem productivity. Although the nighttime is much longer in August and September, possible rain events can affect the biotic cycle, thus preventing us from stable and continuous daily measurements. Samples were collected in the middle of the lake, at the depth of 20 cm and 10 cm (Trisino and Chernoe, respectively), from a PVC boat, which was gently moved over the lake surface using a rope held by a person standing on the shore to avoid disturbance of sediments. Every 2 h, the dissolved oxygen (Cellox 325; accuracy of $\pm 5\%$), specific conductivity (TetraCon 325; $\pm 1.5\%$), pH (SenTix 41; $\pm 0.02 \text{ pH}$) and water temperature ($\pm 0.2^\circ \text{C}$) were measured in-situ at 20 cm depth using a WTW 3320 Multimeter. Air temperature and atmospheric pressure were measured using an ADC Summit handheld weather station (Silva). The partial pressure of CO_2 ($p\text{CO}_2$) in water was measured in-situ using a hand-held infrared gas analyzer (IRGA, GMT222 CARBOCAP 3000 ppm probe, Vaisala; $\pm 1.5\%$) enclosed within a waterproof and gas-permeable membrane and connected to a logger (MI70, Vaisala; $\pm 0.2\%$, see details in Serikova et al., 2019).

Water samples for DOC, DIC, and dissolved CO_2 and CH_4 were collected and analyzed following methods described elsewhere (Manasypov et al., 2015; Polishchuk et al., 2018; Serikova et al., 2018). In brief, for DOC analyses, collected waters were filtered on-site in pre-washed sterile 30-mL polypropylene Nalgene® bottles through single-use Minisart filter units (0.45 μm pore size, Sartorius, acetate cellulose filter). The first 20 mL of the filtrate was always discarded. The DOC and DIC were analyzed using a Carbon Total Analyzer (Shimadzu TOC VSCN) with an uncertainty better than 3% (see Prokushkin et al., 2011 for methodology). For calculating SUVA_{254} , we measured ultraviolet absorbance at 254 nm (UV_{254}) using a 10-mm quartz cuvette on a Bruker CARY-50 UV-VIS spectrophotometer following Abbt-Braun and Frimmel (1999). Filtered solutions for cations and trace element analyses were acidified ($\text{pH}=2$) with ultrapure double-distilled HNO_3 and stored in HDPE bottles previously washed with 0.1 M HCl and rinsed with ultrapure MilliQ deionized water. The preparation of sampling bottles was performed in a clean bench room class A 10000. Major cations (Ca, Mg, Na, K), Si and trace elements (TE) were determined without pre-concentration by ICP-MS Agilent 7500ce, routinely used in our laboratory for analysis of organic-rich lakes from frozen peatlands (cf. Pokrovsky et al., 2016). We obtained a good agreement between replicated measurements of SLRS-5 (River water reference material for trace metals) and the certified values (relative difference < 10% SD on repeated measurements), for all major and trace elements, except B and P (30%). Chloride, sulfide, fluoride and carboxylic organic ligands (lactate, acetate, propionate, formate, butyrate, pyruvate, galacturonate, glutarate, malate, tartrate, oxalate, phthalate, quinate, and citrate) were measured using a high-performance ion chromatography (Dionex ICS-5000+, analytical column: AS11-HC 4 μm), with an uncertainty of 5% and detection limits of $2 \mu\text{g L}^{-1}$. Phosphate was also measured by an IC with an uncertainty of 2% and a detection limit of $1 \mu\text{g L}^{-1}$. Total microbial cell concentration was measured after sample fixation in glutaraldehyde, by a flow cytometry (Guava® EasyCyte™ systems, Merck). Cells were stained using 1 μL of a 10 times diluted SYBR GREEN solution (10,000 \times , Merck), added to 250 μL of each sample before analysis. Particles were identified as cells based on green fluorescence and forward scatter (Marie et al., 1999).

For GHG analyses, unfiltered water was sampled in 25-mL Serum bottles that were closed without air bubbles using vinyl stoppers and aluminum caps and immediately poisoned by adding 0.2 mL of saturated HgCl_2 using a two-way needle system. The CH_4 concentrations were measured using a Bruker GC-456 gas chromatograph (GC) equipped with flame ionization detector. The CO_2 fluxes (FCO_2) were measured in both lakes using small (30 cm diameter) floating chambers equipped with non-dispersive infrared CO_2 logger (ELG, SenseAir, Bastviken et al., 2015; Alin et al., 2011) and calculated following methods described elsewhere (Vachon et al., 2010; Serikova et al., 2019). The CO_2 loggers were calibrated in the lab against pure N_2 - CO_2 mixtures before each sampling campaign. We placed chambers along a transect from the shore to the center of the lakes, with 6 chambers for the Lake Trisino and 4 chambers for the smaller Chernoe. The CO_2 accumulation rate inside each chamber was recorded continuously at 5 min interval. The chambers were ventilated each 2 h and the CO_2 accumulation rate inside each chamber was calculated by linear regression for 50 to 100 min accumulation time. Examples of CO_2 concentration vs. time in both lakes are shown in Fig. S2. Note that, although linear regression can underestimate CO_2 flux in the static floating chamber method (i.e., Xiao et al., 2016), the use of exponential model of Xiao et al. (2016) instead of linear regression

yields minor corrections to the final flux values (within ± 5 –10%) which is below the uncertainty of replicates.

The normality of DOC, major and trace elements, GHG concentration and fluxes was assessed via the Shapiro-Wilk test. Because the data were not normally distributed, we used non-parametric statistics. A non-parametric Mann-Whitney U test was taken for treating the data due to the short series of observation, non-normal distribution of number of parameters and some amount of outliers. Regressions were used to examine the relationships between pCO_2 , FCO_2 and water temperature. We used the median, 1st and 3rd quartiles to trace the dependence of all hydrochemical components on the period of observation (day vs. night). Differences in concentrations and fluxes between day and night time and between two lakes were tested using Mann-Whitney U test for paired data sets with significance level at 0.05, given the limited number of data (short series of observation and some amount of outliers).

3. Results

3.1. Lack of diel variations in major lake water parameters, nutrients, trace elements and bacteria

Concentrations of GHG and dissolved components (DOC, DIC, major cations and anions, and 30 trace elements), pH and specific conductivity during the two days observation are provided in the Mendeleev Data Repository (Pokrovsky et al., 2019). The two chosen lakes are representative for the region in terms of DOC, GHG and metal concentration as illustrated in plots of lake water hydrochemical parameters as a function of lake surface area (Figs. S3 and S4 of the Supplement). The DOC-rich thaw pond Chernoe had about 2 times higher DOC concentration (significant at $p < 0.001$) than thermokarst lake Trisino and exhibited respectively, 2 to 3 times higher concentrations of divalent metal cations (Mg, Ca, Sr, Ba, Mn, Co, Ni, Cu, Pb, Cd), trivalent and tetravalent TE (Al, Ti, Fe, Zr, REEs, Th), Cr, As, Cs, Nb, and U, which are usually bound to organic colloids in thermokarst waters (Pokrovsky et al., 2016).

The summer-time period of lake sampling corresponds to most stable chemical composition of lake waters (Manasypov et al., 2015 and Fig. S5). This period is often used for C and element pool inventory (Polishchuk et al., 2018) because the lake volume and shoreline are most stable. Moreover, our two day measurements are fully consistent with previous observations under normal summer conditions, as illustrated in Table S1 which lists the mean (\pm SD) values of pH, DOC, DIC and trace element in both lakes during summer 2013, 2015, 2016 and this study. The Mann-Whitney U test did not show significant difference between different years of both lakes.

For convenience, we listed the 48-h median (\pm IQR) values of all measured parameters at the end of the day and end of the night, corresponding to minimum and maximum of CO_2 concentration and emission (see Section 3.2. below) in Table 1. The corresponding median values integrated over two days and nights are given in Table S2. The water temperature of the two lakes followed a similar diurnal cycle with a minimum temperature attained approximately 2 h after sunrise (Fig. 1 A). The difference between day and night was significant, as confirmed by Mann-Whitney U test ($p=0.006$ and 0.008 for Trisino and Chernoe, respectively). The pH exhibited rather weak (± 0.2 to 0.3 units) fluctuations with generally higher values during the day compared to night time (Fig. 1 B). The water column remained O_2 -saturated (90%) for both lakes and the O_2 concentration increased by ca. 20–30% from the middle of the night until noon, generally following the pattern of pH (not shown). The DOC remained fairly constant with non-systematic variations around the mean value (\pm SD) of 35.2 ± 2.0 and $15.5 \pm 0.9 \text{ mg L}^{-1}$ in Chernoe and Trisino, respectively

Table 1

Median (\pm IQR) concentration in the Trisino Lake and the Chernoe Pond during the period of low and high emission CO_2 . The minimum is calculated at the end of the day (late afternoon) and the maximum corresponds to the end of night/early morning.

Variable	Trisino Lake		Chernoe Pond	
	Day min	Night max	Day min	Night max
T_{waters} , °C	22.2 \pm 1.6	18.9 \pm 3.4	22.1 \pm 1	18.3 \pm 3.05
T_{airs} , °C	21.3 \pm 3.9	15.2 \pm 4.5	21.3 \pm 3.9	15 \pm 6.2
pH	3.91 \pm 0.03	3.81 \pm 0.04	3.68 \pm 0.19	3.73 \pm 0.04
SC ^a , $\mu\text{S cm}^{-1}$	19 \pm 1	19 \pm 1	28 \pm 1	29 \pm 0.5
flux CO_2 , $\text{g C m}^{-2} \text{d}^{-1}$	-0.03 \pm 0.02	0.2 \pm 0.14	0.08 \pm 0.06	0.22 \pm 0.08
CH_4 , $\mu\text{mol L}^{-1}$	0.23 \pm 0.06	0.42 \pm 0.36	3.74 \pm 2.54	4.84 \pm 1.51
CO_2 , ppmv	360 \pm 90	740 \pm 148	580 \pm 390	1210 \pm 155
DIC, mg L^{-1}	0.6 \pm 0.05	0.6 \pm 0.03	0.6 \pm 0.11	0.7 \pm 0.21
DOC, mg L^{-1}	15.7 \pm 1.12	14.9 \pm 0.44	35.9 \pm 3.09	33.5 \pm 4.77
SUVA ₂₅₄	4.54 \pm 0.29	4.67 \pm 0.06	4.92 \pm 0.32	5.15 \pm 0.55
Li, $\mu\text{g L}^{-1}$	0.65 \pm 0.04	0.64 \pm 0.01	0.61 \pm 0.01	0.58 \pm 0.01
B, $\mu\text{g L}^{-1}$	5.62 \pm 3.66	3.51 \pm 1.06	8.66 \pm 2.69	5.37 \pm 0.54
Na, $\mu\text{g L}^{-1}$	369 \pm 13.4	367 \pm 8.1	373 \pm 37.1	339 \pm 24.8
Mg, $\mu\text{g L}^{-1}$	243 \pm 1.9	243 \pm 1.3	437 \pm 2.9	431 \pm 18.3
Al, $\mu\text{g L}^{-1}$	64.1 \pm 2.23	62.9 \pm 3.94	137 \pm 3	136 \pm 7.9
Si, $\mu\text{g L}^{-1}$	138 \pm 19	143 \pm 31.6	227 \pm 37.9	191 \pm 57.2
K, $\mu\text{g L}^{-1}$	98.7 \pm 26.3	85.4 \pm 21.9	93.5 \pm 11.6	79.8 \pm 32.9
Ca, $\mu\text{g L}^{-1}$	44.9 \pm 4.24	42.5 \pm 21.2	538 \pm 18.7	533 \pm 77.6
Mn, $\mu\text{g L}^{-1}$	8.9 \pm 0.06	8.99 \pm 0.24	20.3 \pm 0.61	20.2 \pm 0.61
Fe, $\mu\text{g L}^{-1}$	90 \pm 2.1	88.8 \pm 2.71	274 \pm 27.6	272 \pm 18
Co, $\mu\text{g L}^{-1}$	0.08 \pm 0.01	0.06 \pm 0.01	0.12 \pm 0.01	0.11 \pm 0.004
Ni, $\mu\text{g L}^{-1}$	0.22 \pm 0.04	0.24 \pm 0.1	0.42 \pm 0.06	0.45 \pm 0.3
Cu, $\mu\text{g L}^{-1}$	0.16 \pm 0.06	0.17 \pm 0.04	0.47 \pm 0.01	0.45 \pm 0.05
Zn, $\mu\text{g L}^{-1}$	6.1 \pm 8.57	3.84 \pm 1.73	12.2 \pm 8.12	6.41 \pm 0.85
As, $\mu\text{g L}^{-1}$	0.32 \pm 0.002	0.33 \pm 0.01	0.52 \pm 0.01	0.52 \pm 0.02
Rb, $\mu\text{g L}^{-1}$	0.02 \pm 0.03	0.04 \pm 0.03	0.07 \pm 0.01	0.08 \pm 0.07
Sr, $\mu\text{g L}^{-1}$	1.92 \pm 0.06	1.93 \pm 0.12	4.38 \pm 0.05	4.37 \pm 0.38
Y, $\mu\text{g L}^{-1}$	0.03 \pm 0.002	0.03 \pm 0.001	0.04 \pm 0.002	0.03 \pm 0.002
Pb, $\mu\text{g L}^{-1}$	0.11 \pm 0.04	0.15 \pm 0.05	0.35 \pm 0.02	0.36 \pm 0.01
Ti, $\mu\text{g L}^{-1}$	0.42 \pm 0.02	0.42 \pm 0.02	1.4 \pm 0.06	1.35 \pm 0.07
V, $\mu\text{g L}^{-1}$	0.92 \pm 0.01	0.87 \pm 0.02	1.02 \pm 0.05	0.98 \pm 0.02
Cr, $\mu\text{g L}^{-1}$	0.39 \pm 0.09	0.33 \pm 0.05	0.71 \pm 0.05	0.6 \pm 0.04
Zr, $\mu\text{g L}^{-1}$	0.06 \pm 0.002	0.05 \pm 0.002	0.11 \pm 0.003	0.1 \pm 0.01
Mo, $\mu\text{g L}^{-1}$	0.004 \pm 0.004	0.005 \pm 0.003	0.01 \pm 0.002	0.01 \pm 0.004
Cd, $\mu\text{g L}^{-1}$	0.01 \pm 0.002	0.01 \pm 0.001	0.03 \pm 0.01	0.03 \pm 0.004

Table 1 (Continued)

Variable	Trisino Lake		Chernoe Pond	
	Day min	Night max	Day min	Night max
Sb, $\mu\text{g L}^{-1}$	0.04 \pm 0.002	0.04 \pm 0.002	0.05 \pm 0.003	0.04 \pm 0.002
Cs, $\mu\text{g L}^{-1}$	0.001 \pm 0.0001	0.001 \pm 0.001	0.01 \pm 0.001	0.01 \pm 0.002
Ba, $\mu\text{g L}^{-1}$	2.14 \pm 0.22	2.13 \pm 0.28	4.01 \pm 0.21	4.06 \pm 0.5
La, $\mu\text{g L}^{-1}$	0.01 \pm 0.001	0.01 \pm 0.001	0.02 \pm 0.003	0.02 \pm 0.001
Ce, $\mu\text{g L}^{-1}$	0.02 \pm 0.002	0.02 \pm 0.002	0.04 \pm 0.004	0.05 \pm 0.002
Th, $\mu\text{g L}^{-1}$	0.01 \pm 0.0004	0.01 \pm 0.001	0.02 \pm 0.0001	0.02 \pm 0.001
U, $\mu\text{g L}^{-1}$	0.002 \pm 0.001	0.002 \pm 0.0005	0.01 \pm 0.0003	0.01 \pm 0.0004

^a SC – specific conductivity.


(Fig. 1 C). The SUVA₂₅₄ was also constant and fluctuated non-systematically between 4 and 5 $\text{L mg C}^{-1} \text{m}^{-1}$ for both lakes (Fig. 1 D).

Phosphate and silica concentrations did not exhibit discernable diel variation in both lakes; PO_4 remained essentially constant at 17 to 20 $\mu\text{g P-PO}_4 \text{L}^{-1}$ (Fig. S6 A), whereas Si fluctuated between 150 and 250 $\mu\text{g L}^{-1}$ without distinct trend (Fig. S6 B). The concentration of carboxylates remained constant (formate, oxalate, pyruvate) or fluctuated without any diel pattern (acetate and lactate) as illustrated in Fig. S7. The concentration of formate and oxalate was ca. 30 to 60% higher in organic-rich Chernoe compared to organic-poor Trisino, and glutarate was detected only in Chernoe, also without diurnal pattern. The total cell number and biomass of bacteria were similar between the two lakes and varied within 10–15% throughout the observations without any systematic difference ($p > 0.05$ of Mann-Whitney U test) between the day and the night period (512,000 \pm 41,000 cell mL^{-1} in Trisino and 531,000 \pm 80,000 cell mL^{-1} in Chernoe).

The DIC in both lakes was very low, 0.6 to 0.7 mg L^{-1} and did not respond to day/night cycle (Fig. S8 A). The major cations (K, Na, Ca, Mg) and anions (Cl, SO_4 , F) concentrations and specific conductivity (S.C.) also remained constant within $\pm 10\%$ as illustrated in Fig. S8 B for specific conductivity. Most trace elements did not exhibit any sizeable variations over diurnal cycle as their concentrations remained stable within $\pm 10\%$ (Li, B, Al, Ti, V, Cr, Mn, Fe, Co, As, Rb, Sr, Y, Zr, Nb, Cd, Cs, Ba, REEs, Pb, Th, U) as illustrated in Figs. S8–S9. The exceptions were Cu, Zn and Ni whose concentrations varied by a factor of 1.5 to 3.0, respectively, however, without any discernable diel cycle (Fig. S9 D and E). These trace elements did not exhibit significant difference between their day-time and night-time concentrations over 48 h of monitoring as confirmed by Mann-Whitney U test ($p > 0.05$ for both lakes).

3.2. Dissolved CO_2 and CH_4 and emission flux of CO_2

The pCO_2 ranged from 300 to 800 ppmv in Trisino and from 400 to 1400 ppmv in Chernoe. The pCO_2 demonstrated clear diurnal pattern with a factor of 2 to 4 higher concentrations achieved at the end of the night and early morning (Fig. 2 A) compared to the middle of the day. The difference in CO_2 concentration between the end of night/early morning peak and the end of day minimum averaged over 48 h of observation was significant for both lakes ($Z = -2.48$, $p = 0.013$ for Trisino; $Z = -2.80$, $p = 0.005$ for Chernoe). The diurnal pattern of pCO_2 significantly correlated with water temperature (T_{water}): $R_{\text{Spearman}} = -0.52$, $p < 0.01$ and $R_{\text{Spearman}} = -0.57$, $p < 0.005$



for Trisino and Chernoe, respectively (Fig. 2 C). The FCO_2 was close to zero or negative during the day time in Trisino ($-0.02 \pm 0.02 \text{ g C m}^{-2} \text{ d}^{-1}$, mean \pm SD) but positive in Chernoe ($0.12 \pm 0.07 \text{ g C m}^{-2} \text{ d}^{-1}$, mean \pm SD). During the night, the FCO_2 strongly increased and became positive in Trisino ($0.12 \pm 0.1 \text{ g C m}^{-2} \text{ d}^{-1}$, mean \pm SD), whereas its increase in Chernoe was a factor of 2 to 5 between the day minimum and the night maximum (Fig. 2 B). Similar to pCO_2 , the FCO_2 followed the diurnal pattern which was shifted with respect to solar radiation intensity and significantly ($R_{\text{Spearman}} = -0.56$, $p < 0.01$ and $R_{\text{Spearman}} = -0.57$, $p < 0.01$ for Trisino and Chernoe, respectively) negatively correlated with water temperature (Fig. 2 D). The difference in FCO_2 between early morning maximum and late afternoon minimum was significant ($Z = -2.65$, $p = 0.008$ for Trisino; $Z = -2.19$, $p = 0.028$ for Chernoe). Furthermore, to check the difference in the degree of pCO_2 and FCO_2 response to diel cycles of temperature and light between two lakes, we used a Mann-Whitney U test. The lakes were significantly different ($Z = -2.00$, $p = 0.045$ for the end of the day minimum; $Z = -2.80$, $p = 0.005$ for the early morning maximum).

The CH_4 concentration ranged between 0.15 and $0.5 \mu\text{mol L}^{-1}$ in DOC-poor Trisino and between 2 and $7 \mu\text{mol L}^{-1}$ in organic-rich Chernoe (Fig. 3). Possible impact of bubble emissions was taken into account via assessing lateral heterogeneity of CH_4 concentrations across the lakes, and reflected in the uncertainty attached to the data at each time of the day. These uncertainties typically ranged from 10 to 20% . There was no systematic trend of CH_4 with time and diurnal pattern in large Lake Trisino; however, the CH_4 concentrations in small and shallow Chernoe pond increased up to ca. 5-fold after the beginning of sampling without following the diurnal pattern. We therefore cannot exclude sediment disturbance and bubble release due to boat movement, which interfered with methane measurements in the Chernoe Lake given its low depth.

4. Discussion

4.1. High stability of pH, nutrients, organic acids and dissolved ($<0.45 \mu\text{m}$) components of thermokarst lakes

Diurnal variations of major hydrochemical and biological parameters (Eiler et al., 2006) and trace metals, involved in redox cycles (McMahon, 1969; Barry et al., 1994; Emmenegger et al., 2001; Pokrovsky and Shirokova, 2013) in lake waters are fairly well known. The present study adds a novel dimension to the existing knowledge via reporting the chemical composition of previously unknown humic and acidic thermokarst lakes, which are widely distributed in the permafrost zone of Northern Eurasia (Polishchuk et al., 2018). Over 2 days and nights monitoring in the Trisino Lake, the pH decreased by 0.1 to 0.2 unit during the night compared to day time period, presumably reflecting the primary production – mineralization processes (Wright and Mills, 1967; Zirino et al., 1986), as it is known in non-permafrost humic lakes of circum-neutral pH (Shirokova et al., 2010). The low pH of studied lakes (3.86 and 3.71 of Trisino and Chernoe, respectively) stems from a combination of high DOC, acidic nature of surrounding peat and ground and submerged vegetation (mostly mosses). Note that the diel pH variations were much lower (ca., 0.1 ± 0.1 unit) in organic-rich Chernoe which had two times higher DOC concentration. Overall, the obtained re-

sults dismiss the hypothesis of temperature and light control on DOC and trace element concentration in lakes and ponds of permafrost peatlands. Moreover, the degree of response in solute concentrations (except pH and CO_2) to diel variation in external parameters was not related to the DOC level in humic lakes.

The constant concentration of DOC and lack of discernable diel pattern in SUVA in both lakes suggest that the pool of dissolved organic carbon was not affected, at the diel scale, by the five main processes controlling DOM level in thermokarst lake waters. These are: *i*) soil water input via supra-permafrost flow (Raudina et al., 2018); *ii*) benthic respiration of DOM (Demars, 2019), *iii*) photo-degradation (Cory et al., 2014), *iv*) heterotrophic consumption of DOM in the water column (Kritzberg et al., 2006; Shirokova et al., 2009; Peura et al., 2019) and *v*) phytoplankton and macrophytes exometabolite production (Demarty and Prairie, 2009). First of all, it is possible that the main processes controlling DOC input and removal in the water column, such as phytoplankton metabolite production and photo-oxidation, compensate each other at the diurnal scale. Similarly, the DOC delivery via supra-permafrost flow may accelerate during the day due to enhanced soil heating and ice thaw, but this freshly-delivered allochthonous DOC can be quickly, on the scale of first hours, processed by photolysis (Shirokova et al., 2019). Note that low impact of the phytoplankton and heterotrophs on DOC and SUVA is consistent with lack of diurnal variation in limiting nutrients (Si, PO_4), simple organic acids that can be used for heterotrophic metabolism (acetate, lactate, formate, pyruvate, oxalate), and the total bacterial number. A lack of Si concentration change between day and night is also consistent with low abundance of diatoms in thermokarst lakes of Western Siberia, given that the phytoplankton of acidic and humic waters is dominated by cyanobacteria and green algae (Pavlova et al., 2016). Finally, in agreement with results on some temperate rivers (Spencer et al., 2007) and boreal lakes (Shirokova et al., 2010; Pokrovsky and Shirokova, 2013), it is possible that the changes in bulk DOC concentration simply could not be quantified because the biogeochemically-processed fraction of DOC was small and not detectable at overall high DOC level in humic waters. Therefore, the present study corroborates recent results on humic (24 mg DOC L^{-1}) and circumneutral ($\text{pH} = 7.5\text{--}9.7$) subarctic lake which demonstrated limited variability in DOC and SUVA on a diel scale compared to a seasonal scale (Johnston et al., 2019). The latter was essentially linked to the change in specific organic compounds (decrease of aromatic and polyphenolic classes) and increase in aliphatic compounds, following pulses of allochthonous and autochthonous DOM. Presumably, these pulses are not detectable over hours to daily scale observation, especially in high-DOC aquatic environments. Similar deduction can be applied for the lack of DIC pattern: at the acidic pH of Trisino and Chernoe water, the DIC represents quite small fraction of CO_2 and thus the DIC variations linked to photosynthesis cycle could not be resolved due to high pCO_2 variations.

For other dissolved ($<0.45 \mu\text{m}$) components of the water column, we observed quite high stability of concentrations in both lakes, which signifies that the role of diel processes in the water column, the water-sediment exchange and short-term input of supra-permafrost waters were negligible during 2 days of observation. This result is also important from methodological point of view as it suggests that for most dissolved major and trace elements of thermokarst lakes in discontinuous permafrost zone, a single one-time sampling at any pe-

Fig. 1. Air and water temperature and direct solar radiation (DSR) (A), pH (B), DOC concentration (C), and SUVA (D) variations in surface water measured in the two lakes of Western Siberia (Trisino and Chernoe). The shadowed area represents night time observations (same for Figs. 2 and 3).




Fig. 2. Diurnal pattern of pCO₂ (A) and CO₂ flux (FCO₂) (B) measured in the two lakes of Western Siberia. The negative correlations between pCO₂ and water temperature (C) and FCO₂ and water temperature (D) are significant at p < 0.01. The error bars represent the s.d. of six (Trisino) and four (Chernoe) chamber measurements across the lakes.

riod of the day or night is sufficiently representative for a given lake during specific season of the year.

Another important result of our observations is that there was a factor of 2 to 3 higher concentration of DOC and ~30 elements in DOC-rich Chernoe thaw pond compared to the DOC-poor Trisino Lake. It is known that the majority of cations are strongly bound to organic and organo-ferric colloids in thermokarst lakes and ponds of discontinuous permafrost zone of WSL (typically from 80 to 99%,

Pokrovsky et al., 2016). Given that both DOC and Fe concentrations are 2 to 3 times higher in the Chernoe Lake compared to the Trisino Lake, the concentration of organo-ferric colloids that bind many trace elements is also higher in the Chernoe Lake. The exceptions are Li, B, Na, K, Si, V, Mo and Sb that are not bound to organic colloids and thus exhibit similar (within ±20%) concentrations in both lakes. Note that while total dissolved phosphorus (P_{tot}) was 5 to 10 times lower in the low-DOC Trisino compared to the high-DOC Chernoe, the phos-




Fig. 3. Diurnal pattern concentration of dissolved CH_4 measured in the two lakes of Western Siberia. The uncertainties of CH_4 concentration are within the symbols size. They stem from 2 to 3 individual measurements in various part of the lake at each time of the day.

phate anion concentrations were rather similar between lakes. As such, the difference in P_{tot} behavior is due to its limitation by organo-ferric colloids rather than due to phytoplankton uptake of P in the Trisino Lake.

4.2. Diurnal pattern of CO_2 concentration and flux

Although the present study is probably the first that documents significant diel variations of $p\text{CO}_2$ flux using direct chamber measurements in humic thermokarst lakes of permafrost peatlands, our findings are in general agreement with large body of studies in non-permafrost rivers and lakes showing similar pattern of decreased CO_2 during daylight hours (Naiman, 1983; Guasch et al., 1998; Bott et al., 2006; Vesala et al., 2006; Parker et al., 2010; Vale et al., 2018; Rocher-Ros et al., 2019) and highest values in the early morning (Dawson et al., 2001; Reiman and Xu, 2019; Yang et al., 2019). The $p\text{CO}_2$ and FCO_2 diel patterns in permafrost peatland waters confirm the hypothesis that the high-DOC lake exhibits weaker response to temperature and light variations compared to the low-DOC lake.

The diel variabilities of net ecosystem production and related CO_2 effluxes are mostly pronounced in oligotrophic lakes (Spafford and Risk, 2018; Liu et al., 2016). In such DOC-poor lakes, the community respiration rates can vary by an order of magnitude, yielding up to 3 mg L^{-1} diel changes in DOC concentration and being strongly linked to diel oscillations in water temperature (i.e., Bogard et al., 2019). In other cases, the typical variations of DOC concentrations linked to bacterio-plankton respiration are much lower (for example, 0.1 to 0.3 mg L^{-1} over 5 h of night time, Sadro et al., 2011). In contrast, at the high level of DOC in humic thermokarst lakes of peatlands, the 0.1 to 3 mg L^{-1} diel variations in DOC cannot be resolved with sufficient precision. Indeed, the $0.3 \text{ g C m}^{-2} \text{ d}^{-1}$ variation in CO_2 flux observed in this work (Fig. 2 b) corresponds to a 0.3 mg L^{-1} variation in DOC concentration (due to bacterial metabolism or photolysis) in a lake of 1 m depth. In shallower, 30-cm deep thaw ponds like Chernoe, the effect of DOM processing the water column capable to produce the observed CO_2 flux is 3 times higher, but still does not exceed 1 mg L^{-1} of DOC. Such low effects can be hardly resolved at high DOC level of studied ponds (15 to 35 mg L^{-1}). At the same time, the role of DOM photolysis as a cause of CO_2 emission in subarctic waters (i.e. Cory et al., 2014; Bowen et al., 2019) can be weakly pronounced in permafrost peatlands because the acidic humic waters of thermokarst lakes from frozen peatlands are highly resistant to photo-degradation (Shirokova et al., 2019). This is further confirmed by the

fact that the concentrations of photo-sensitive components (SUVA_{254} , Fe, Mn) remained conservative with $<10\%$ variations during full period of observations (Fig. 1 D, Figs. S9 A and B). It is also possible that the photo-oxidation of DOC to CO_2 occurring during the day (Cory et al., 2014; Bertilsson and Tranvik, 2000) is masked by CO_2 consumption due to benthic photosynthesis, as it was suggested in some Arctic streams (Rocher-Ros, 2019).

We therefore hypothesize that the respiration of organic matter in sediments (benthic microbial metabolism), rather than DOM processing in the water column are mainly responsible for high FCO_2 in general, as it was recently shown in a small catchment on peat soils in boreal zone (Demars, 2019). The CO_2 uptake during day time may be linked to macrophytes and submerged mosses that partially colonize the bottom of shallow thermokarst lakes and thaw ponds in this region (Pokrovsky et al., 2014), as it is also known in other Arctic lakes (Squires et al., 2009; Tank et al., 2011; Bouchard et al., 2014). In this regard, the diurnal pattern of $p\text{CO}_2$ and FCO_2 can be interpreted as a result of a balance between sediment respiration and GPP of periphyton, aquatic plants and submerged mosses. As for the external supply of CO_2 from surrounding soils to the lake water (via suprapermafrost flow, Raudina et al., 2018), the variation of this factor counteracts the observed $p\text{CO}_2$ pattern because the maximal soil water input due to ice thaw is likely to occur during the highest air temperature (late afternoon) which corresponds to minimal $p\text{CO}_2$ level in the lake waters. Similarly, convective mixing is unlikely to contribute to the CO_2 emission pattern because, during period of observation at the anticyclonic weather, the $p\text{CO}_2$ and FCO_2 were strongly correlated ($r^2=0.90$, $p<0.01$) and the wind speed during the anticyclone remained quite low and did not vary between day and night.

4.3. Consequences for global CO_2 emission, and C and metal pools assessment in thermokarst lakes of Western Siberia lowland

The observed stability of most dissolved component concentrations in thermokarst lakes of permafrost peatlands has important implication for the evaluation of pools and fluxes of C and related elements. Highly conservative behavior of DOC, DIC, CH_4 , nutrients, major and trace elements, notably micro-nutrients (except Zn), toxicants and geochemical traces during two days and nights of monitoring demonstrates the representability of one single sampling during the day or night for assessing seasonal (week, month - averaged) behavior of solutes in thermokarst lakes of frozen peatlands. In particular, the pools of DOC and element calculated in thaw ponds and thermokarst lakes of the whole territory of Western Siberia, based on

single-point sampling (Polishchuk et al., 2018), provides reasonable approximation for the summer time period. This conclusion corroborates results on diel and seasonal variability of dissolved components in N. American subarctic lake (Johnston et al., 2019).

In contrast, systematically higher FCO₂ during the night compared to the day may require some corrections on global estimations of CO₂ emissions from lakes and ponds of frozen peatlands. The majority of available measurements on thermokarst lakes and thaw ponds were performed during day-time, without accounting for possible diurnal cycle (see Serikova et al., 2019 and references therein). Considering full duration of unfrozen period on lakes of discontinuous permafrost zone of WSL (mean multi-annual of 137 days, from May 25 to October 7), the night time represents 30% of the ice-free period. The enhanced FCO₂ during the night from small thaw ponds (< 2000 m²) such as Lake Chernoe thus increases the average emission from 0.10 to 0.20 g C m⁻² d⁻¹. However, because the contribution of such ponds to overall water area is negligibly small (< 1%, Polishchuk et al., 2018), this does not affect the regional C emission balance. In contrast, medium and large thermokarst lakes (> 1–10 ha) provide the main water surfaces and CO₂ aquatic pools in WSL (Polishchuk et al., 2018). In these lakes, the day-time FCO₂ in summer could fluctuate around zero (± 0.02 g C m⁻² d⁻¹) but rise to 0.25 ± 0.05 g C m⁻² d⁻¹ at the end of the night as it is observed in the Trisino Lake.

The median day-time CO₂ flux from lakes of discontinuous permafrost zone of the WSL in summer is 0.74 g C m⁻² d⁻¹ but ranges from -0.3 to 3.13 g C m⁻² d⁻¹ during day time (calculated based on data of 21 lakes in the region, Serikova et al., 2019). Although the variations of CO₂ flux between individual lakes may strongly exceed the effect of diurnal cycle, a 0.2 g C m⁻² d⁻¹ increase between day and night may produce sizeable increase in seasonal emission fluxes. Therefore, given the scarcity of diurnal observations on large lakes and the absence of data on night-time emission during ice-off and ice-on periods (when the emission is sizably higher than in summer, Serikova et al., 2019), the quantification of the night-time CO₂ emission at the regional level over full open water period of the year is of high priority.

5. Conclusions

We observed stable concentrations of pH, DOC, SUVA₂₅₄, CH₄, nutrients, carboxylic acids, most major and trace elements in waters of acidic and humic thermokarst lake and thaw pond during a continuous 2-days observation at generally anticyclone weather. In contrast, the CO₂ concentration and emission negatively correlated with water temperature and exhibited clear diurnal pattern with a maximum after the sunrise (minimal surface water temperature) and a minimum during late afternoon day-time (maximum surface water temperature). Among major drivers of observed CO₂ variation, we exclude photosynthetic/heterotrophic activity of the plankton in the water column and photolysis of DOM. Rather, the balance between benthic respiration of dissolved and particulate organic matter and primary productivity of periphyton, aquatic macrophytes and submerged mosses was responsible for enhanced CO₂ emission during the night and CO₂ decrease or even uptake during the day. Neglecting night-time CO₂ flux, especially in large thermokarst lakes, can sizably underestimate the region-upscaled global C emission. This calls for a need to study diurnal variations in C emission in large (10–500 ha) thermokarst lakes because they mostly contribute to overall inland water surface of the region. These measurements should be focused on late summer and autumn periods when the heterotrophic respiration dominates over primary productivity and the diurnal cycles are most pronounced.

Uncited reference

Mortimer, 1959

CRedit authorship contribution statement

L.S. Shirokova: Conceptualization, Investigation, Writing - original draft. **D. Payandi-Rolland:** Conceptualization, Investigation, Writing - original draft. **A.G. Lim:** Investigation. **R.M. Manasypov:** Investigation. **J. Allen:** Formal analysis. **J.-L. Rols:** Formal analysis. **P. Benezech:** Formal analysis. **J. Karlsson:** Formal analysis. **O.S. Pokrovsky:** Conceptualization, Writing - original draft.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This work was supported by RSCF grants No. 19-77-00073 (30%, sampling) and 18-77-10045 (20%, data analysis), the Ministry of Education and Science of the Russian Federation no. 1.8195.2017/9.10, RFBR grants No 17-55-16008, No 19-05-50096, and No 20-05-00729, and RSF grant No 18-77-10045 for data analysis. The fieldwork was partially supported by the international program INTERACT for the 2018 campaign. We thank Frédéric Julien (EcoLab) for Dionex HPLC measurements of carboxylic organic ligands.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2020.139671>.

References

- Abbt-Braun, G., Frimmel, F.H., 1999. Basic characterization of Norwegian NOM samples — similarities and differences. *Environ. Int.* 25, 161–180. [https://doi.org/10.1016/S0160-4120\(98\)00118-4](https://doi.org/10.1016/S0160-4120(98)00118-4).
- Albéric, P., Voillier, E., Jézéquel, D., Grosbois, C., Michard, G., 2000. Interactions between trace elements and dissolved organic matter in the stagnant anoxic deep layer of a meromictic lake. *Limnol. Oceanogr.* 45, 1088–1096. <https://doi.org/10.4319/lo.2000.45.5.1088>.
- Alin, S.R., Rasera, M. de F.F.L., Salimon, C.I., Richey, J.E., Holtgrieve, G.W., Krusche, A.V., Snidvongs, A., 2011. Physical controls on carbon dioxide transfer velocity and flux in low-gradient river systems and implications for regional carbon budgets. *Journal of Geophysical Research: Biogeosciences* 116 (G1), G01009. <https://doi.org/10.1029/2010JG001398>.
- Ask, J., Karlsson, J., Jansson, M., 2012. Net ecosystem production in clear-water and brown-water lakes. *Glob. Biogeochem. Cycles* 26, GB1017. <https://doi.org/10.1029/2010GB003951>.
- Baehr, M.M., DeGrandpre, M.D., 2004. In situ pCO₂ and O₂ measurements in a lake during turnover and stratification: observations and modeling. *Limnol. Oceanogr.* 49 (2), 330–340.
- Barry, R.C., Schnoor, J.L., Sulzberger, B., Sigg, L., Stumm, W., 1994. Iron oxidation kinetics in an acidic alpine lake. *Water Res.* 28, 323–333. [https://doi.org/10.1016/0043-1354\(94\)90270-4](https://doi.org/10.1016/0043-1354(94)90270-4).
- Bastviken, D., Sundgren, I., Natchimuthu, S., Reyier, H., Gålfalk, M., 2015. Technical note: cost-efficient approaches to measure carbon dioxide (CO₂) fluxes and concentrations in terrestrial and aquatic environments using mini loggers. *Biogeosciences* 12, 3849–3859. <https://doi.org/10.5194/bg-12-3849-2015>.
- Bertilsson, S., Tranvik, L.J., 2000. Photochemical transformation of dissolved organic matter in lakes. *Limnol. Oceanogr.* 45, 753–762. <https://doi.org/10.4319/lo.2000.45.4.0753>.
- Bogard, M.J., Johnston, S.E., Dornblaser, M., Spencer, R.G.M., Striegler, R.G., Butman, D.E., 2019. Extreme rates and diel variability of planktonic respiration in

- a shallow sub-arctic lake. *Aquat. Sci.* 81, 60. <https://doi.org/10.1007/s00027-019-0657-9>.
- Bott, T.L., Montgomery, D.S., Newbold, J.D., Arscott, D.B., Dow, C.L., Aufdenkampe, A.K., Jackson, J.K., Kaplan, L.A., 2006. Ecosystem metabolism in streams of the Catskill Mountains (Delaware and Hudson River watersheds) and Lower Hudson Valley. *J. N. Am. Benthol. Soc.* 25, 1018–1044.
- Bouchard, F., Francus, P., Pienitz, R., Laurion, I., Feyte, S., 2014. Subarctic thermokarst ponds: investigating recent landscape evolution and sediment dynamics in thawed permafrost of northern Québec (Canada). *Arctic, Antarctic, Alpine Res.* 46, 251–271. <https://doi.org/10.1657/1938-4246-46.1.251>.
- Bowen, J.C., Kaplan, L.A., Cory, R.M., 2019. Photodegradation disproportionately impacts biodegradation of semi-labile DOM in streams. *Limnology and Oceanography* 1–9. <https://doi.org/10.1002/lno.11244>, (press).
- Brick, C.M., Moore, J.N., 1996. Diel variation of trace metals in the upper Clark Fork River, Montana. *Environ. Sci. Technol.* 30, 1953–1960. <https://doi.org/10.1021/es9506465>.
- Carling, G.T., Fernandez, D.P., Rudd, A., Pazmino, E., Johnson, W.P., 2011. Trace element diel variations and particulate pulses in perimeter freshwater wetlands of Great Salt Lake, Utah. *Chem. Geol.* 283, 87–98. <https://doi.org/10.1016/j.chemgeo.2011.01.001>.
- Cory, R.M., Ward, C.P., Crump, B.C., Kling, G.W., 2014. Sunlight controls water column processing of carbon in arctic fresh waters. *Science* 345, 925–928. <https://doi.org/10.1126/science.1253119>.
- Crawford, J.T., Stanley, E.H., Dornblaser, M.M., Striegl, R.G., 2017. CO₂ time series patterns in contrasting headwater streams of North America. *Aquat. Sci.* 79, 473–486. <https://doi.org/10.1007/s00027-016-0511-2>.
- Dawson, J.J.C., Billett, M.F., Hope, D., 2001. Diurnal variations in the carbon chemistry of two acidic peatland streams in north-east Scotland. *Freshw. Biol.* 46, 1309–1322. <https://doi.org/10.1046/j.1365-2427.2001.00751.x>.
- DelVecchia, A.G., Balik, J.A., Campbell, S.K., Taylor, B.W., West, D.C., Wissinger, S.A., 2019. Carbon dioxide concentrations and efflux from permanent, semi-permanent, and temporary subalpine ponds. *Wetlands* 1–15. <https://doi.org/10.1007/s13157-019-01140-3>, (press).
- Demars, B.O.L., 2019. Hydrological pulses and burning of dissolved organic carbon by stream respiration. *Limnol. Oceanogr.* 64, 406–421. <https://doi.org/10.1002/lno.11048>.
- Demarty, M., Prairie, Y.T., 2009. In situ dissolved organic carbon (DOC) release by submerged macrophyte–epiphyte communities in southern Quebec lakes. *Can. J. Fish. Aquat. Sci.* 66, 1522–1531. <https://doi.org/10.1139/F09-099>.
- Eiler, A., Olsson, J.A., Bertilsson, S., 2006. Diurnal variations in the auto- and heterotrophic activity of cyanobacterial phycospheres (*Gloeotrichia echinulata*) and the identity of attached bacteria. *Freshw. Biol.* 51, 298–311. <https://doi.org/10.1111/j.1365-2427.2005.01493.x>.
- Emmenegger, L., Schönenberger, R., Sigg, L., Sulzberger, B., 2001. Light-induced redox cycling of iron in circumneutral lakes. *Limnol. Oceanogr.* 46, 49–61. <https://doi.org/10.4319/lno.2001.46.1.0049>.
- Eugster, W., Kling, G., Jonas, T., McFadden, J.P., Wüest, A., MacIntyre, S., Chapin, F.S., 2003. CO₂ exchange between air and water in an Arctic Alaskan and midlatitude Swiss lake: importance of convective mixing. *J. Geophys. Res.: Atmospheres* 108 (D12), 4362. <https://doi.org/10.1029/2002JD002653>.
- Falkner, K.K., Church, M., Measures, C.I., Lebaron, G., Thouron, D., Jeandel, C., Stordal, M.C., Gill, G.A., Mortlock, R., Froelich, P., Chan, L.-H., 1997. Minor and trace element chemistry of Lake Baikal, its tributaries, and surrounding hot springs. *Limnol. Oceanogr.* 42, 329–345. <https://doi.org/10.4319/lno.1997.42.2.0329>.
- Forget, M.-H., Carignan, R., Hudon, C., 2009. Influence of diel cycles of respiration, chlorophyll, and photosynthetic parameters on the summer metabolic balance of temperate lakes and rivers. *Can. J. Fish. Aquat. Sci.* 66, 1048–1058. <https://doi.org/10.1139/F09-058>.
- Gammons, C.H., Nimick, D.A., Parker, S.R., Cleasby, T.E., McCleskey, R.B., 2005. Diel behavior of iron and other heavy metals in a mountain stream with acidic to neutral pH: Fisher Creek, Montana, USA. *Geochim. Cosmochim. Acta* 69, 2505–2516. <https://doi.org/10.1016/j.gca.2004.11.020>.
- Gorham, E., 1991. Northern Peatlands: role in the carbon cycle and probable responses to climatic warming. *Ecol. Appl.* 1, 182–195. <https://doi.org/10.2307/1941811>.
- Guasch, H., Armengol, J., Martí, E., Sabater, S., 1998. Diurnal variation in dissolved oxygen and carbon dioxide in two low-order streams. *Water Res.* 32, 1067–1074. [https://doi.org/10.1016/S0043-1354\(97\)00330-8](https://doi.org/10.1016/S0043-1354(97)00330-8).
- Hamilton-Taylor, J., Willis, M., 1990. A quantitative assessment of the sources and general dynamics of trace metals in a soft-water lake. *Limnol. Oceanogr.* 35, 840–851. <https://doi.org/10.4319/lno.1990.35.4.0840>.
- Hanson, P.C., Carpenter, S.R., Armstrong, D.E., Stanley, E.H., Kratz, T.K., 2006. Lake dissolved inorganic carbon and dissolved oxygen: changing drivers from days to decades. *Ecol. Monogr.* 76, 343–363. [https://doi.org/10.1890/0012-9615\(2006\)076\[0343:LDICAD\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2006)076[0343:LDICAD]2.0.CO;2).
- Johnston, S.E., Bogard, M.J., Rogers, J.A., Butman, D., Striegl, R.G., Dornblaser, M., Spencer, R.G.M., 2019. Constraining dissolved organic matter sources and temporal variability in a model sub-Arctic lake. *Biogeochemistry* <https://doi.org/10.1007/s10533-019-00619-9>.
- Jones, C.A., Nimick, D.A., McCleskey, R.B., 2004. Relative effect of temperature and pH on diel cycling of dissolved trace elements in Prickly Pear Creek, Montana. *Water, Air, Soil Pollution* 153, 95–113. <https://doi.org/10.1023/B:WATE.00000119934.64939.f0>.
- Jonsson, A., Åberg, J., Lindroth, A., Jansson, M., 2008. Gas transfer rate and CO₂ flux between an unproductive lake and the atmosphere in northern Sweden. *J. Geophys. Res. Biogeosci.* 113, G04006. <https://doi.org/10.1029/2008JG000688>.
- Kritzberg, E.S., Cole, J.J., Pace, M.L., Granéli, W., Bade, D.L., 2004. Autochthonous versus allochthonous carbon sources of bacteria: results from whole-lake ¹³C addition experiments. *Limnol. Oceanogr.* 49, 588–596. <https://doi.org/10.4319/lno.2004.49.2.0588>.
- Kritzberg, E.S., Cole, J.J., Pace, M.M., Granéli, W., 2006. Bacterial growth on allochthonous carbon in humic and nutrient-enriched lakes: results from whole-lake ¹³C addition experiments. *Ecosystems* 9, 489–499. <https://doi.org/10.1007/s10021-005-0115-5>.
- Laurion, I., Vincent, W.F., MacIntyre, S., Retamal, L., Dupont, C., Francus, P., Pienitz, R., 2010. Variability in greenhouse gas emissions from permafrost thaw ponds. *Limnol. Oceanogr.* 55, 115–133. <https://doi.org/10.4319/lno.2010.55.1.0115>.
- Liu, H., Zhang, Q., Katul, G.G., Cole, J.J., Chapin, F.S., MacIntyre, S., 2016. Large CO₂ effluxes at night and during synoptic weather events significantly contribute to CO₂ emissions from a reservoir. *Environ. Res. Lett.* 11, 064001. <https://doi.org/10.1088/1748-9326/11/6/064001>.
- Lundin, E.J., Klaminder, J., Giesler, R., Persson, A., Olefeldt, D., Heliasz, M., Christensen, T.R., Karlsson, J., 2016. Is the subarctic landscape still a carbon sink? Evidence from a detailed catchment balance. *Geophys. Res. Lett.* 43, 1988–1995. <https://doi.org/10.1002/2015GL066970>.
- Lynch, J.K., Beatty, C.M., Seidel, M.P., Jungst, L.J., DeGrandpre, M.D., 2010. Controls of riverine CO₂ over an annual cycle determined using direct, high temporal resolution pCO₂ measurements. *J. Geophys. Res. Biogeosci.* 115, G03016. <https://doi.org/10.1029/2009JG001132>.
- Manasyapov, R.M., Vorobyev, S.N., Loiko, S.V., Kritzkov, I.V., Shirokova, L.S., Shevchenko, V.P., Kirpotin, S.N., Kulizhsky, S.P., Kolesnichenko, L.G., Zemtsov, V.A., Sinkinov, V.V., Pokrovsky, O.S., 2015. Seasonal dynamics of organic carbon and metals in thermokarst lakes from the discontinuous permafrost zone of western Siberia. *Biogeosciences* 12, 3009–3028. <https://doi.org/10.5194/bg-12-3009-2015>.
- Marie, D., Partensky, F., Vaulot, D., Brussaard, C., 1999. Enumeration of phytoplankton, bacteria, and viruses in marine samples. *Current Protocols Cytometry* 10 (1), 11111–11115. <https://doi.org/10.1002/0471142956.cy11111s10>.
- Marushchak, M.E., Kiepe, I., Biasi, C., Elsakov, V., Friborg, T., Johansson, T., Soegaard, H., Virtanen, T., Martikainen, P.J., 2013. Carbon dioxide balance of subarctic tundra from plot to regional scales. *Biogeosciences* 10, 437–452. <https://doi.org/10.5194/bg-10-437-2013>.
- McMahon, J.W., 1969. The annual and diurnal variation in the vertical distribution of acid-soluble ferrous and total iron in a small dimictic lake. *Limnol. Oceanogr.* 14, 357–367. <https://doi.org/10.4319/lno.1969.14.3.0357>.
- Mortimer, C.H., 1959. A treatise on limnology. Volume 1. Geography, physics and chemistry. *Limnol. Oceanogr.* 4, 108–114. <https://doi.org/10.4319/lno.1959.4.1.0108>.
- Naiman, R.J., 1983. The annual pattern and spatial distribution of aquatic oxygen metabolism in boreal forest watersheds. *Ecol. Monogr.* 53, 73–94. <https://doi.org/10.2307/1942588>.
- Neal, C., Watts, C., Williams, R.J., Neal, M., Hill, L., Wickham, H., 2002. Diurnal and longer term patterns in carbon dioxide and calcite saturation for the River Kennet, south-eastern England. *Sci. Total Environment* 282–283, 205–231. [https://doi.org/10.1016/S0048-9697\(01\)00952-4](https://doi.org/10.1016/S0048-9697(01)00952-4).
- Nimick, D.A., Gammons, C.H., Parker, S.R., 2011. Diel biogeochemical processes and their effect on the aqueous chemistry of streams: a review. *Chem. Geol.* 283, 3–17. <https://doi.org/10.1016/j.chemgeo.2010.08.017>.
- Nriagu, J.O., Lawson, G., Wong, H.K.T., Cheam, V., 1996. Dissolved trace metals in Lakes Superior, Erie, and Ontario. *Environ. Sci. Technol.* 30, 178–187. <https://doi.org/10.1021/es950221i>.
- Parker, S.R., Gammons, C.H., Poulson, S.R., DeGrandpre, M.D., 2007. Diel variations in stream chemistry and isotopic composition of dissolved inorganic carbon, upper Clark Fork River, Montana, USA. *Appl. Geochem.* 22, 1329–1343. <https://doi.org/10.1016/j.apgeochem.2007.02.007>.
- Parker, S.R., Gammons, C.H., Poulson, S.R., DeGrandpre, M.D., Weyer, C.L., Smith, M.G., Babcock, J.N., Oba, Y., 2010. Diel behavior of stable isotopes of dissolved oxygen and dissolved inorganic carbon in rivers over a range of trophic conditions, and in a mesocosm experiment. *Chem. Geol.* 269, 22–32. <https://doi.org/10.1016/j.chemgeo.2009.06.016>.
- Pavlova, O.A., Pokrovsky, O.S., Manasyapov, R.M., Shirokova, L.S., Vorobyev, S.N., 2016. Seasonal dynamics of phytoplankton in acidic and humic environment in thaw ponds of discontinuous permafrost zone. *Ann. Limnol. - Int. J. Lim.* 52, 47–60. <https://doi.org/10.1051/limn/2016006>.

- Peter, H., Singer, G.A., Preiler, C., Chiffard, P., Stenciczka, G., Battin, T.J., 2014. Scales and drivers of temporal pCO₂ dynamics in an Alpine stream. *J. Geophys. Res. Biogeosci.* 119, 1078–1091. <https://doi.org/10.1002/2013JG002552>.
- Peura, S., Wauthy, M., Domenic Simone, D., Eiler, A., Einarsdóttir, K., Rautio, M., Bertilsson, S., 2019. Ontogenic succession of thermokarst thaw ponds is linked to dissolved organic matter quality and microbial degradation potential. *Limnol. Oceanogr.* 2019, <https://doi.org/10.1002/lno.11349>.
- Podgrajsek, E., Sahlée, E., Rutgersson, A., 2014. Diurnal cycle of lake methane flux. *Journal of Geophysical Research: Biogeosciences* 119, 236–248. <https://doi.org/10.1002/2013JG002327>.
- Pokrovsky, O.S., Shirokova, L.S., 2013. Diurnal variations of dissolved and colloidal organic carbon and trace metals in a boreal lake during summer bloom. *Water Res.* 47, 922–932. <https://doi.org/10.1016/j.watres.2012.11.017>.
- Pokrovsky, O.S., Shirokova, L.S., Kirpotin, S.N., Audry, S., Viers, J., Dupré, B., 2011. Effect of permafrost thawing on organic carbon and trace element colloidal speciation in the thermokarst lakes of western Siberia. *Biogeosciences* 8, 565–583. <https://doi.org/10.5194/bg-8-565-2011>.
- Pokrovsky, O.S., Shirokova, L.S., Kirpotin, S.N., Kulizhsky, S.P., Vorobiev, S.N., 2013. Impact of western Siberia heat wave 2012 on greenhouse gases and trace metal concentration in thaw lakes of discontinuous permafrost zone. *Biogeosciences* 10, 5349–5365. <https://doi.org/10.5194/bg-10-5349-2013>.
- Pokrovsky, O.S., Shirokova, L.S., Kirpotin, S.N., 2014. *Biogeochemistry of Thermokarst Lakes of Western Siberia*. Nova Science Publishers, Inc, New York.
- Pokrovsky, O.S., Manasyrov, R.M., Loiko, S.V., Shirokova, L.S., 2016. Organic and organo-mineral colloids in discontinuous permafrost zone. *Geochim. Cosmochim. Acta* 188, 1–20. <https://doi.org/10.1016/j.gca.2016.05.035>.
- Pokrovsky, O.S., Shirokova, L.S., Payandi-Rolland, D., Lim, A., Manasyrov, R., 2019. Diurnal cycle of C concentration and emission fluxes in small thaw ponds of frozen peatlands. *Mendeley Data V1*. <https://doi.org/10.17632/yvdc63s55.1>.
- Polishchuk, Y.M., Bogdanov, A.N., Muratov, I.N., Polishchuk, V.Y., Lim, A., Manasyrov, R.M., Shirokova, L.S., Pokrovsky, O.S., 2018. Minor contribution of small thaw ponds to the pools of carbon and methane in the inland waters of the permafrost-affected part of the Western Siberian lowland. *Environ. Res. Lett.* 13, 045002. <https://doi.org/10.1088/1748-9326/aab046>.
- Pollard, P.C., 2013. In situ rapid measures of total respiration rate capture the super labile DOC bacterial substrates of freshwater. *Limnol. Oceanography: Methods* 11, 584–593. <https://doi.org/10.4319/lom.2013.11.584>.
- Prokushkin, A.S., Pokrovsky, O.S., Shirokova, L.S., Korets, M.A., Viers, J., Prokushkin, S.G., Amon, R.M.W., Guggenberger, G., McDowell, W.H., 2011. Sources and the flux pattern of dissolved carbon in rivers of the Yenisey basin draining the central Siberian plateau. *Environ. Res. Lett.* 6, 045212. <https://doi.org/10.1088/1748-9326/6/4/045212>.
- Raudina, T.V., Loiko, S.V., Lim, A., Manasyrov, R.M., Shirokova, L.S., Istigechev, G.I., Kuzmina, D.M., Kulizhsky, S.P., Vorobyev, S.N., Pokrovsky, O.S., 2018. Permafrost thaw and climate warming may decrease the CO₂, carbon, and metal concentration in peat soil waters of the Western Siberia Lowland. *Sci. Total Environ.* 634, 1004–1023. <https://doi.org/10.1016/j.scitotenv.2018.04.059>.
- Reiman, J.H., Xu, Y.J., 2019. Diel variability of pCO₂ and CO₂ outgassing from the Lower Mississippi River: implications for riverine CO₂ outgassing estimation. *Water* 11, 43. <https://doi.org/10.3390/w11010043>.
- Rocher-Ros, G., 2019. *Biophysical Controls on CO₂ Evasion from Arctic Inland Waters*. Dissertation for PhD vol. 2019, Department of Ecology and Environmental Science, Umea University.
- Rocher-Ros, G., Sponseller, R.A., Lidberg, W., Mörth, C.-M., Giesler, R., 2019. Landscape process domains drive patterns of CO₂ evasion from river networks. *Limnol. Oceanogr. Lett.* 4, 87–95. <https://doi.org/10.1002/lol2.10108>.
- Roehm, C.L., Giesler, R., Karlsson, J., 2009. Bioavailability of terrestrial organic carbon to lake bacteria: the case of a degrading subarctic permafrost mire complex. *J. Geophys. Res.* 114, G03006. <https://doi.org/10.1029/2008JG000863>.
- Sadro, S., Nelson, C.E., Melack, J.M., 2011. Linking diel patterns in community respiration to bacterioplankton in an oligotrophic high-elevation lake. *Limnol. Oceanogr.* 56, 540–550. <https://doi.org/10.4319/lo.2011.56.2.0540>.
- Sadro, S., Holtgrieve, G.W., Solomon, C.T., Koch, G.R., 2014. Widespread variability in overnight patterns of ecosystem respiration linked to gradients in dissolved organic matter, residence time, and productivity in a global set of lakes. *Limnol. Oceanogr.* 59, 1666–1678. <https://doi.org/10.4319/lo.2014.59.5.1666>.
- Serikova, S., Pokrovsky, O.S., Ala-Aho, P., Kazantsev, V., Kirpotin, S.N., Kopysov, S.G., Krickov, I.V., Laudon, H., Manasyrov, R.M., Shirokova, L.S., Soulsby, C., Tetzlaff, D., Karlsson, J., 2018. High riverine CO₂ emissions at the permafrost boundary of Western Siberia. *Nat. Geosci.* 11, 825–829. <https://doi.org/10.1038/s41561-018-0218-1>.
- Serikova, S., Pokrovsky, O.S., Laudon, H., Krickov, I.V., Lim, A.G., Manasyrov, R.M., Karlsson, J., 2019. High carbon emissions from thermokarst lakes of Western Siberia. *Nat. Commun.* 10, 1–7. <https://doi.org/10.1038/s41467-019-09592-1>.
- Shirokova, L.S., Pokrovsky, O.S., Kirpotin, S.N., Dupré, B., 2009. Heterotrophic bacterioplankton in thawed lakes of the northern part of Western Siberia controls the CO₂ flux to the atmosphere. *Internat. J. Environ. Stud.* 66, 433–445. <https://doi.org/10.1080/00207230902758071>.
- Shirokova, L.S., Pokrovsky, O.S., Viers, J., Klimov, S.I., Moreva, O.Y., Zabelina, S.A., Vorobieva, T.Y., Dupré, B., 2010. Diurnal variations of trace metals and heterotrophic bacterioplankton concentration in a small boreal lake of the White Sea basin. *Ann. Limnol. - Int. J. Lim.* 46, 67–75. <https://doi.org/10.1051/limn/2010011>.
- Shirokova, L.S., Pokrovsky, O.S., Kirpotin, S.N., Desmukh, C., Pokrovsky, B.G., Audry, S., Viers, J., 2013. Biogeochemistry of organic carbon, CO₂, CH₄, and trace elements in thermokarst water bodies in discontinuous permafrost zones of Western Siberia. *Biogeochemistry* 113, 573–593. <https://doi.org/10.1007/s10533-012-9790-4>.
- Shirokova, L.S., Chupakov, A.V., Zabelina, S.A., Neverova, N.V., Payandi-Rolland, D., Causerand, C., Karlsson, J., Pokrovsky, O.S., 2019. Humic surface waters of frozen peat bogs (permafrost zone) are highly resistant to bio- and photodegradation. *Biogeosciences* 16, 2511–2526. <https://doi.org/10.5194/bg-16-2511-2019>.
- Spafford, L., Risk, D., 2018. Spatiotemporal variability in lake-atmosphere net CO₂ exchange in the littoral zone of an oligotrophic lake. *J. Geophys. Res. Biogeosci.* 123, 1260–1276. <https://doi.org/10.1002/2017JG004115>.
- Spencer, R.G.M., Pellerin, B.A., Bergamaschi, B.A., Downing, B.D., Kraus, T.E.C., Smart, D.R., Dahlgren, R.A., Hernes, P.J., 2007. Diurnal variability in riverine dissolved organic matter composition determined by in situ optical measurement in the San Joaquin River (California, USA). *Hydrol. Process.* 21, 3181–3189. <https://doi.org/10.1002/hyp.6887>.
- Squires, M.M., Lesack, L.F.W., Hecky, R.E., Guildford, S.J., Ramlal, P., Higgins, S.N., 2009. Primary production and carbon dioxide metabolic balance of a lake-rich Arctic river floodplain: partitioning of phytoplankton, epipelon, macrophyte, and epiphyton production among lakes on the Mackenzie Delta. *Ecosystems* 12, 853–872. <https://doi.org/10.1007/s10021-009-9263-3>.
- Szyper, J.P., Rosenfeld, J.Z., Piedrahita, R.H., Giovannini, P., 1992. Diel cycles of planktonic respiration rates in briefly incubated water samples from a fertile earthen pond. *Limnol. Oceanogr.* 37, 1193–1201. <https://doi.org/10.4319/lo.1992.37.6.1193>.
- Tank, S.E., Lesack, L.F.W., Gareis, J.A.L., Osburn, C.L., Hesslein, R.H., 2011. Multiple tracers demonstrate distinct sources of dissolved organic matter to lakes of the Mackenzie Delta, western Canadian Arctic. *Limnol. Oceanogr.* 56, 1297–1309. <https://doi.org/10.4319/lo.2011.56.4.1297>.
- Tranvik, L.J., 1988. Availability of dissolved organic carbon for planktonic bacteria in oligotrophic lakes of differing humic content. *Microb. Ecol.* 16, 311–322. <https://doi.org/10.1007/BF02011702>.
- Tranvik, L.J., 1989. Bacterioplankton growth, grazing mortality and quantitative relationship to primary production in a humic and a clearwater lake. *J. Plankton Res.* 11, 985–1000. <https://doi.org/10.1093/plankt/11.5.985>.
- Vachon, D., Prairie, Y.T., Cole, J.J., 2010. The relationship between near-surface turbulence and gas transfer velocity in freshwater systems and its implications for floating chamber measurements of gas exchange. *Limnol. Oceanogr.* 55, 1723–1732. <https://doi.org/10.4319/lo.2010.55.4.1723>.
- Vale, R., Santana, R., Gomes, A.C., Tóta, J., 2018. Increased nocturnal CO₂ concentration during breeze circulation events in a tropical reservoir. *Ambiente & Água Interdisc. J. Appl. Sci.* 13, 1–12. <https://doi.org/10.4136/ambi-agua.2186>.
- Vesala, T., Huotari, J., Rannik, S., Suni, T., Smolander, S., Sogachev, A., Launiainen, S., Ojala, A., 2006. Eddy covariance measurements of carbon exchange and latent and sensible heat fluxes over a boreal lake for a full open-water period. *J. Geophys. Res.: Atmospheres* 111, D11101. <https://doi.org/10.1029/2005JD006365>.
- Viollier, E., Michard, G., Jézéquel, D., Pèpe, M., Sarazin, G., 1997. Geochemical study of a crater lake: Lake Pavin, Puy de Dôme, France. Constraints afforded by the particulate matter distribution in the element cycling within the lake. *Chem. Geol.* 142, 225–241. [https://doi.org/10.1016/S0009-2541\(97\)00093-4](https://doi.org/10.1016/S0009-2541(97)00093-4).
- Vorobyev, S.N., Pokrovsky, O.S., Kolesnichenko, L.G., Manasyrov, R.M., Shirokova, L.S., Karlsson, J., Kirpotin, S.N., 2019. Biogeochemistry of dissolved carbon, major, and trace elements during spring flood periods on the Ob River. *Hydrol. Process.* 33, 1579–1594. <https://doi.org/10.1002/hyp.13424>.
- Wright, J.C., Mills, I.K., 1967. Productivity studies on the Madison River, Yellowstone National Park. *Limnol. Oceanogr.* 12, 568–577. <https://doi.org/10.4319/lo.1967.12.4.0568>.
- Xiao, S., Wang, Y., Liu, D., Yang, Z., Lei, D., Zhang, C., 2013. Diel and seasonal variation of methane and carbon dioxide fluxes at Site Guojaba, the Three Gorges Reservoir. *J. Environ. Sci.* 25 (10), 2065–2071.
- Xiao, S., Wang, C., Wilkinson, R.J., Liu, D., Zhang, C., Xu, W., Yang, Z., Wang, Y., Lei, D., 2016. Theoretical model for diffusive greenhouse gas fluxes estimation across water-air interfaces measured with the static floating chamber method. *Atmos. Environ.* 137, 45–52.
- Yang, R., Chen, B., Liu, H., Liu, Z., Yan, H., 2015. Carbon sequestration and decreased CO₂ emission caused by terrestrial aquatic photosynthesis: insights from diel hydrochemical variations in an epikarst spring and two spring-fed ponds in different seasons. *Appl. Geochem.* 63, 248–260. <https://doi.org/10.1016/j.apgeochem.2015.09.009>.

Yang, R., Xu, Z., Liu, S., Xu, Y.J., 2019. Daily pCO₂ and CO₂ flux variations in a subtropical mesotrophic shallow lake. *Water Res.* 153, 29–38. <https://doi.org/10.1016/j.watres.2019.01.012>.

Yvon-Durocher, G., Allen, A.P., Bastviken, D., Conrad, R., Gudas, C., St-Pierre, A., Thanh-Duc, N., del Giorgio, P.A., 2014. Methane fluxes show consistent temperature dependence across microbial to ecosystem scales. *Nature* 507, 488–491. <https://doi.org/10.1038/nature13164>.

Zirino, A., Fuhrmann, R.A., Oksanen-Gooden, D., Lieberman, S.H., Clavell, C., Seligman, P.F., Mathewson, J.H., Jones, W.D., Kogelschatz, J., Barber, R.T., 1986. pH-temperature-nutrient relationships in the eastern tropical Pacific Ocean. *Sci. Total Environ.* 58, 117–137. [https://doi.org/10.1016/0048-9697\(86\)90082-3](https://doi.org/10.1016/0048-9697(86)90082-3).

UNCORRECTED PROOF