



Long-term exposure to near-future ocean acidification does not affect the expression of neurogenesis- and synaptic transmission-related genes in the olfactory bulb of European sea bass (*Dicentrarchus labrax*)

David Mazurais, Arianna Servili, Nicolas Le Bayon, Sebastien Gislard,
Lauriane Madec, Jose-Luis Zambonino-Infante

► To cite this version:

David Mazurais, Arianna Servili, Nicolas Le Bayon, Sebastien Gislard, Lauriane Madec, et al.. Long-term exposure to near-future ocean acidification does not affect the expression of neurogenesis- and synaptic transmission-related genes in the olfactory bulb of European sea bass (*Dicentrarchus labrax*). *Journal of Comparative Physiology B*, 2020, 190 (2), pp.161-167. 10.1007/s00360-019-01256-2 . hal-03004618

HAL Id: hal-03004618

<https://hal.science/hal-03004618>

Submitted on 21 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Long-term exposure to near-future ocean acidification does not affect the expression of neurogenesis- and synaptic transmission-related genes in the olfactory bulb of European sea bass (*Dicentrarchus labrax*)

Mazurais David ^{1,*}, Servili Arianna ¹, Le Bayon Nicolas ¹, Gislard Sebastien ¹, Madec Lauriane ¹, Zambonino-Infante Jose-Luis ¹

¹ IFREMER, Univ Brest, CNRS, IRD, LEMAR, 29280, Brest, Plouzané, France

* Corresponding author : David Mazurais, email address : dmazurai@ifremer.fr

Abstract :

The decrease in ocean pH that results from the increased concentration of dissolved carbon dioxide (CO₂) is likely to influence many physiological functions in organisms. It has been shown in different fish species that ocean acidification (OA) mainly affects sensory systems, including olfaction. Impairment of olfactory function may be due to a dysfunction of the GABAergic system and to an alteration of neuronal plasticity in the whole brain and particularly in olfactory bulbs. Recent studies revealed that OA-driven effects on sensory systems are partly mediated by the regulation of the expression of genes involved in neurotransmission and neuronal development. However, these studies were performed in fish exposed to acidified waters for short periods, of only a few days. In the present paper, we investigated whether such effects could be observed in adult (4-years old) European sea bass (*Dicentrarchus labrax*) exposed to two hypercapnic and acidified conditions (PCO₂ ≈ 980 μatm; pH total = 7.7 and PCO₂ ≈ 1520 μatm; pH total = 7.5) from the larval stage. In a first approach, we analyzed by qPCR the expression of five genes involved in neurogenesis (DCX) or expressed in GABAergic (Gabra3), glutamatergic (Gria1) or dopaminergic (TH and DDC) neurons in the olfactory bulbs. The tested experimental conditions did not change the expression of any of the five genes. This result would indicate that a potential disruption of the olfactory function of sea bass exposed for a long term to near-future OA, either occurs at a level other than the transcriptional one or involves other actors of the sensory function.

Introduction

The predicted increase in atmospheric PCO₂ level which may reach 420-936 ppm by the end of the century will cause a concurrent ocean acidification (OA: elevated H⁺, reduced pH) up to 0.42 pH units (Field 2014). Several studies during last decade examined the potential consequences of marine organisms' exposure to OA, including fish species (Doney et al. 2009; Stillman and Paganini 2015; Wang et al. 2018). In fish, OA has been shown to disrupt cognitive function and to induce behavioral disorders (Chivers et al. 2014; Domenici et al. 2012; Ferrari et al. 2012; Jutfelt et al. 2013; Nilsson et al. 2012) particularly due to alteration of sensory function such as olfaction (Dixon et al. 2015; Dixon et al. 2010; Munday et al. 2009; Rong et al. 2018; Rossi et al. 2016), vision (Chung et al. 2014; Forsgren et al. 2013) and hearing (Rossi et al. 2016). Changes in brain neurotransmitter function and especially in the functioning of the gamma-Aminobutyric acid (GABA) receptor have been argued to explain these effects (Chivers et al. 2014; Nilsson et al. 2012; Regan et al. 2016). In addition to modifying the neuronal Cl⁻ and HCO₃⁻ gradients that are important for the GABAA receptor function, OA was shown to up-regulate the mRNA expression of α family subunits ($\alpha 1$, $\alpha 3$, $\alpha 4$ and $\alpha 6b$) of the GABAA receptor in brains of three-spined stickleback (*Gasterosteus aculeatus*) (Lai et al. 2017b). Interestingly, recent data obtained in two teleost species [European sea bass (*Dicentrarchus labrax*) and coho salmon (*Oncorhynchus kisutch*)] indicated that OA associated to near future elevated CO₂ induces significant changes in gene expression involved in neurotransmission signal. Among these genes, it can be found glutamate (glutamate receptor and transporters) and dopamine/serotonin [aromatic-L-amino-acid decarboxylase (DDC)] signaling related factors in the olfactory bulbs (Porteus et al. 2018; Williams et al. 2019).

In fish, as in other vertebrates, the olfactory bulbs are brain structures that first receive information by olfactory sensory neuron inputs from the olfactory epithelium (Kermen et al.

2013). The olfactory bulbs are composed of different layers of morphologically and functionally distinct neurons that communicate with each other via synapses, which contribute to the processing of olfactory information. Glutamate, dopamine and GABA are the main neurotransmitters expressed in cells localized in the olfactory bulbs (Edwards and Michel 2002; Kermen et al. 2013). Glutamate is the primary excitatory neurotransmitter for all olfactory sensory neurons in the zebrafish (*Danio rerio*) olfactory bulb (Edwards and Michel 2002). Dopamine, that has been shown to suppress the synaptic transmissions from mitral to granule cells in the goldfish (*Carassius auratus*) olfactory bulb, is considered to play an important role in the negative regulation of olfactory responsiveness (Kawai et al. 2012). Regarding GABAergic neurons, that are located in the internal cellular layer of the olfactory bulb in zebrafish, they have been shown to play a critical role in the synchronization of odor-dependent neuronal signals and the regulation of olfactory bulb output activity (Horzmann and Freeman 2016; Kermen et al. 2013; Tabor et al. 2008).

Additionally to molecular actors involved in neurotransmission, other genes related to brain plasticity and neurogenesis [e.g. Doublecortin (*DCX*)] have also been shown to be regulated by OA (Lai et al. 2017a). Such a neurogenetic effect is characteristic to fish species which, unlike mammals, have a high potential to produce new neurons in the adult central nervous system (Zupanc 2001). Interestingly, in the European sea bass juveniles, genes involved in synaptic plasticity and neuronal excitability were shown to be regulated in the olfactory bulbs after exposure to near future pH (Porteus et al. 2018).

Information gathered on the regulatory mechanisms involved in fish OA response was obtained after relatively short exposure times (from 2 days to 40 days) (Bignami et al. 2014; Dixon et al. 2010; Nilsson et al. 2012; Porteus et al. 2018; Williams et al. 2019). To our knowledge, very few studies have looked at the regulations involved in fish exposed to OA for longer periods (Sundin et al. 2019). Yet, this type of response is ecologically relevant in

order to evaluate the resilience capacity of fish when facing global change related effects such as OA. Recently, studies performed in juvenile spiny chromis damselfish (*Acanthochromis polyacanthus*) showed that long-term acclimation to near-future ocean acidification did not induce any acid-base disruption of energetics (Sundin et al. 2019). This result contrasted with several short-term studies considering that alteration of GABAA receptor function may be a mechanistic explanation for the reported hyperactivity and other behavioural impairments of fishes under elevated CO₂.

In this study, we aimed to examine whether molecular actors differentially expressed by a short exposure to acidification, could also be found regulated in the olfactory bulb of sea bass exposed for 4 years to a near future OA. For this purpose, we quantified mRNA expression levels of genes involved in neurogenesis (DCX) and in glutamatergic (glutamate ionotropic receptor), GABAergic (gamma-aminobutyric acid type A receptor alpha3 subunit) and dopaminergic (tyrosine hydroxylase and DDC) neurons of the olfactory bulbs.

Materials and methods

Animal and experimental conditions

Adult European sea bass used in this study were from the same experimental population used in a previous study that examined the impact of OA during larval development (Crespel et al. 2017). Sea bass were maintained from larval to adult stage (four years post-hatching) in three pH/PCO₂ conditions [i.e. PCO₂ ≈ 590 µatm for Control group at pH total = 7.9; PCO₂ ≈ 980 µatm for low acidity group (LA) at pH total = 7.7; PCO₂ ≈ 1520 µatm for high acidity (HA) group at pH total = 7.5]. The choice of the experimental conditions were based to the IPCC Representative Concentration Pathway (RCP) 6.0 and IPCC RCP8.5 scenario predicting a Δ PCO₂ of ≈ 500 µatm and ≈ 1000 µatm by the end of the present century, respectively (Collins 2014). The rearing conditions during larval stage are detailed in the previous paper (Crespel et al. 2017). The sea water was pumped 500 m from the coastline in the Bay of Brest, from a depth of 20 m approximately, passed through a ~500µm sand filter, heated (tungsten, Plate Heat Exchanger, Vicarb, Sweden), degassed using a column, filtered using a 2µm membrane and finally UV sterilized (PZ50, 75W, Ocene, France). Larvae were maintained at 19 °C, with oxygen concentration above 90%, salinity at 34‰ and the controlled photoperiod was set at 24D during the first week of development and then at 16L:8D until 45dph. Larvae were fed ad libitum with *Artemia* until 28 dph and then with commercial dry pellets (Néo-start, Le Guessant Aquaculture, France) until 45 dph. At juvenile stage, fish were maintained at 18°C, 16L:8D and fed a commercial feed (Neo Grower Extra Marin, Le Guessant Aquaculture, France). From two years post-hatching, adult fish were reared under natural photoperiod and ambient temperature (between 10 to 19°C: natural seasonal differences in the Bay of Brest) and fed a diet that encompasses the specific nutritional requirements of Broodstock (Vitalis Cal, Skretting, Stavanger, Norway). Apart from PCO₂ and associated pH conditions, fish from

the three groups experienced identical experimental conditions (water temperature, salinity, oxygenation, food, etc.) throughout their whole lives.

All fish experiments were conducted under approved protocols in strict compliance with the EU Directive 2010/63/EU for animal experiments. The present work conforms to the French legal requirements concerning welfare of experimental animals (APAFIS permit no. 17132-2018101614401562).

mRNA expression analysis

Total RNA extraction was performed from olfactory bulbs sampled on 40 fish (4 years old; 1240 ± 252 g): 14 individuals from the control group (10 females, 4 males), 13 individuals from the LA group (12 females, 1 male), 13 individuals from the HA group (6 females, 7 males) at post-spawning period. Before sampling, fish were fasted for 24 h. Fish were first lightly anaesthetized (20 mg L^{-1}), and then euthanized with a lethal dose (200 mg. L^{-1}) of Tricaine methane-sulfonate 222 (Pharmaq, Fordingbridge, Hampshire, UK). The fish brain was quickly removed, and then the olfactory bulbs were dissected and stored in RNA Stabilization Reagent (RNAlater, Qiagen, Hilden, Germany) following recommendations from the supplier. Total RNA was extracted using Extract-all reagent (Eurobio, Courtaboeuf, Essonne, France) combined with Nucleospin RNA column that includes one step of DNase treatment (Macherey-Nagel, Düren, Germany) according to the manufacturer's instructions. The concentration, purity and integrity were determined using a ND-1000 NanoDrop® spectrophotometer (Thermo Scientific Inc., Waltham, MA, USA) and by electrophoresis using an Agilent Bioanalyser 2100 (Agilent Technologies Inc., Santa Clara, CA, USA). All samples showed an RNA integrity score (RIN) higher than 8 and thus could be used for

reverse transcription-quantitative PCR (RT-qPCR) analysis. RNA samples were stored at -80°C until use.

Retrotranscription (RT) of cDNA for all samples was carried out using 500 ng of RNA with an iScript™ cDNA Synthesis kit (Bio-Rad Laboratories Inc., Hercules, CA, USA) following strictly recommendations from the supplier. cDNA was stored at -20°C until use. Negative RT controls were performed for all samples.

Quantitative real time PCR was performed for the five genes of interest (*DCX*, *Gria1*, *Gabra3*, *TH*, *DDC*) and two housekeeping genes [elongation factor 1-alpha (*ef1α*) and 28S rRNA] using the primers listed in Table 1.

Table 1: Specific primers used for real time amplification.

Genes	Accession number	Forward primer(5'-3')	Reverse primer(5'-3')
<i>DCX</i>	MN183104	ACACACTTGATGGCAAGCAG	TCATCACCTGCATTCGTTA
<i>Gria1</i>	MN183103	GATGCTCTGGTGGGAGTGAT	TTCAAGAAGGACGCCTCTGT
<i>Gabra3</i>	MN183105	CCGAGAATCAGTTCCTGCTC	TGCGCTTGGTGAAGTAGTTG
<i>TH</i>	MN183107	ATGTCGGAAGCTCAAGGAGA	CTTTTGGGAACCAGTGGAA
<i>DDC</i>	MN183106	CTGAGAGCAGGGAGATCCAC	CCTGCAAAAGCTCAAAGGTC
<i>ef1α</i>	AJ866727.1	CTGGAGGGCAGTGAAAAGAT	CATCAAGAGCCTCCAGCAGT
28S rRNA	AH011863.2	GTCCTTCTGATCGAGGCTCA	CGTGCCGGTATTTAGCCTTA

Gene expression was quantified using the CFX96 Touch Real-Time PCR Detection system (Bio-Rad Laboratories Inc.). Standard curves were estimated for each primer pair using serial dilutions (from 1/10 to 1/270) of a pool of cDNA. Efficiencies of qPCR for each pair of

primers ranged from 95% to 100% with $R^2 > 0.99$. Each sample was analysed in triplicate in a final well volume of 15 μ l containing 5 μ l cDNA (1/30 dilution) and 10 μ l of reaction mix, composed of 0.5 μ l of each primer (10 mmol l⁻¹), 1.5 μ l RNase/DNase-free water and 7.5 μ l iQTM SYBR[®] Green Supermix (Bio-Rad Laboratories Inc.). Negative RT controls were also analysed to check for potential contamination. The qPCR profiles contained an initial activation step at 95°C for 2 min followed by 40 cycles of 5 s at 95°C and 20 s at 60°C. After the amplification phase, a high resolution melting curve was performed to confirm the amplification of a single product in each reaction. For all samples, the corresponding C_q (quantification cycle) value was determined automatically using the Gene Expression Module of the CFX Manager software (Bio-Rad Laboratories Inc.). The relative quantity of messenger was normalized with the $\Delta\Delta C_t$ method using efl α and 28S rRNA as reference genes. These reference genes were used as they did not even show low variation of expression between samples (relative standard deviation <5% among samples).

A two-way ANOVA including sex and acidification status as independent factors was used to test significant differences between treatments. For all analyses, variables were checked for normality (Shapiro test) and equality of variances (Levene test). The level of significance was taken at 0.05.

Results

Long term exposure of European sea bass to the two tested conditions of pH/PCO₂ did not induce significant differences ($p>0.1$) in *DCX*, *Gria1*, *Gabra3*, *TH* and *DDC* mRNA expression levels (table 2, Fig 1). No significant differences in mRNA expression levels were observed between males and females for the examined genes (table 2). There was also no interaction between the acidification and sex factors (table 2).




Figure 1: mRNA expression levels of *DCX*, *Gria1*, *Gabra3*, *TH* and *DDC* genes in European sea bass long term exposed to control, low acidification (LA) and high acidification (HA) conditions. Values are expressed as means (males and females together) $\pm$ s.e.m.

Table 2: Statistical analysis of expression data^a

<i>Gene</i>	<i>Conditions</i>						<i>p value</i>		
	<i>Control</i>		<i>LA</i>		<i>HA</i>		Sex	Acid.	Sex * Acid.
	<i>F(10)</i>	<i>M(4)</i>	<i>F(12)</i>	<i>M(1)</i>	<i>F(6)</i>	<i>M(7)</i>			
<i>DCX</i>	0.62±0.18	0.52±0.08	0.66±0.22	0.66	0.60±0.06	0.67±0.35	<i>p</i> =0.930	<i>p</i> =0.749	<i>p</i> =0.822
<i>Gria1</i>	0.75±0.41	0.52±0.07	0.70±0.29	0.76	0.62±0.08	0.68±0.31	<i>p</i> =0.950	<i>p</i> =0.881	<i>p</i> =0.596
<i>Gabra3</i>	0.66±0.21	0.55±0.08	0.94±0.31	0.61	0.62±0.07	0.69±0.26	<i>p</i> =0.244	<i>p</i> =0.439	<i>p</i> =0.402
<i>TH</i>	0.47±0.50	0.19±0.28	0.33±0.49	0.79	0.20±0.34	0.06±0.03	<i>p</i> =0.757	<i>p</i> =0.192	<i>p</i> =0.223
<i>DDC</i>	0.63±0.27	0.51±0.16	0.77±0.35	0.82	0.66±0.07	0.78±0.42	<i>p</i> =0.825	<i>p</i> =0.510	<i>p</i> =0.788

a: Expression levels for each gene were presented as means ± standard deviation and analyzed using a two-way ANOVA and included sex and acidification status as the independent factors. Numbers in brackets indicate sample sizes. The table shows the p-values for each factor. LA: Low acidification condition; HA: High acidification condition.

Discussion

In an attempt to get information about the long term acclimation of fish to OA, the objective of the present study was to determine whether prolonged experimental exposure of European sea bass (i.e. from larval to adult stage) to near future ocean pH influenced expression of genes that play role in neurogenesis and neurotransmission in the olfactory bulbs. Our results indicate that *DCX* gene involved in neurogenesis and *Gria1*, *Gabra3* and *TH* and *DDC* neurotransmitter-related genes involved in glutamatergic, GABAergic and dopaminergic neurons respectively, were not differentially expressed among all tested pH conditions.

The PCO₂ levels associated to OA that we tested (i.e. ≈ 590 μatm for Control condition at PH total= 7.9; ≈ 980 μatm at pH7.7; ≈ 1520 μatm at pH7.5) were higher than those used in previous studies (Lai et al. 2017a; Porteus et al. 2018). However, the Δ PCO₂ of ≈ 500 μatm and ≈ 1000 μatm that we have chosen to simulate OA were similar and based on IPCC RCP6.0 and RCP8.5 scenario (Collins 2014). Therefore we assume that the absence of effect found in the present work was not related to experimental conditions.

The lack of regulation of *DCX* gene expression contrasts with previous data obtained in stickleback indicating up-regulation of this gene by OA, but is in line with responses observed in cinnamon anemonefish and spiny damselfish (Lai et al. 2017a). Lai et al suggested that interspecies variability in the sensitivity of *DCX* gene expression to OA may reflect differential physiological plasticity between fish species (Lai et al. 2017a). The absence of *DCX* gene regulation observed in the present study may not be related to a low molecular plasticity of European sea bass brain structure with respect to acidified condition. Indeed, expression of a hundred genes including actors involved in neurogenesis (i.e. *NAV3*) were shown to be regulated in the olfactory bulb of European sea bass exposed for 2-7 days to near future pH (pH, 7.82 ± 0.01 ; $1122 \pm 19 \mu\text{atm}$) (Porteus et al. 2018). Moreover, in European sea

bass as in teleost fish, brain is characterized by extensive adult neurogenesis providing an excellent model for adult brain plasticity. We assume that the absence of regulation of *DCX* gene expression in the present study may be related to the long time exposure of fish to acidification conditions and we can only speculate that if *DCX* gene expression is sensitive to OA in European sea bass, this is a transient regulation occurring during the first days/weeks of exposure to low pH. A time series analysis of *DCX* gene expression levels throughout the exposure to acidified environment would constitute a proof of this theory. Moreover, we also can not exclude the possibility that *DCX* gene expression may be regulated in other parts of the central nervous system since other authors found this gene regulated from whole brain samples (Lai et al. 2017a).

We can also speculate a similar regulation control for neurotransmission related genes *Gria1a*, *gabra3* and *DDC* which did not exhibit differential expression in the present study after long term exposure to similar acidification conditions, when they were shown to be regulated during short term exposure to OA in European sea bass or stickleback (Lai et al. 2017b; Porteus et al. 2018). Interestingly, the regulation of these genes was associated with impaired olfaction in fish exposed to OA. The absence of *Gria1a*, *gabra3* and *DDC* gene regulation does not necessarily mean that the olfactory system is not impacted by the long-term exposure to OA but if it is, other molecular actors and mechanisms are likely involved.

The alteration of GABA neurotransmitter function has been proposed as the main mechanism explaining the OA induced behavioral impairment in fish (Nilsson et al. 2012). The absence of *gabra3* gene regulation suggested that, as proposed in juvenile spiny chromis damselfish, the GABA function was not modified in fish long term exposed to OA (Sundin et al. 2019). However, as suggested by Williams et al (2019), we can not exclude in the present study that other molecular actors (such as gamma-aminobutyric acid type B receptor subunit 2 gene) involved in GABA function may be regulated. Finally, the fact that the expression levels of

key genes (*DDC* and *TH*) playing an important role in the physiology of dopaminergic neurons (DA) were not found regulated in the present study, suggested that the known DA neurons-induced negative regulation of olfactory responsiveness is not stimulated in European sea bass exposed for a long time to OA (Kawai et al. 2012).

In conclusion, to our knowledge, the present study is the first that addressed the influence of long term exposure of OA on the expression of genes involved in the processing of olfactory information in a teleost species. In contrast to what has been shown during short exposures to acidified water, long-term exposures do not appear to influence genes involved in the processing of olfactory information. However, this does not necessary suggest a total physiological resilience of European sea bass and particularly that olfaction is not altered by long term exposure to OA. Behavioral experiments and electrophysiological analysis of olfactory sensitivity in the adult European sea bass used in the present study would be necessary to evaluate potential impairment of olfactory function when fish are long term exposed to OA. The present study mainly indicates that mechanisms involved in OA response in the olfactory bulb of European sea bass likely depends on the exposure duration, which could be of great importance from an ecological point of view.

Acknowledgments

This work was supported by LabexMer (ANR-10-LABX-0019,OASYS project), the Ministry of Ecological and Solidarity Transition and the Foundation for Biodiversity Research (Ocean Acidification Program, PACIO project) and the Deutsche Forschungsgemeinschaft, PE 1157/8-1, MA4271/3-1 (the FITNESS project).

References

- Bignami S, Sponaugle S, Cowen RK (2014) Effects of ocean acidification on the larvae of a high-value pelagic fisheries species, mahi-mahi *Coryphaena hippurus* *Aquat Biol* 21:249-260 doi:10.3354/ab00598
- Chivers DP et al. (2014) Impaired learning of predators and lower prey survival under elevated CO₂: a consequence of neurotransmitter interference *Global Change Biol* 20:515-522 doi:10.1111/gcb.12291
- Chung WS, Marshall NJ, Watson SA, Munday PL, Nilsson GE (2014) Ocean acidification slows retinal function in a damselfish through interference with GABAA receptors *J Exp Biol* 217:323-326 doi:10.1242/jeb.092478
- Collins M, R. Knutti, J. Arblaster, J.-L. Dufresne, T. Fichet, P. Friedlingstein, X. Gao, W.J. Gutowski, T. Johns, G. Krinner, M. Shongwe, C. Tebaldi, A.J. Weaver and M. Wehner (2014) Long-term Climate Change: Projections, Commitments and Irreversibility Pages 1029 to 1076. In: Intergovernmental Panel on Climate C (ed) *Climate Change 2013 – The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, pp 1029-1136. doi:DOI: 10.1017/CBO9781107415324.024
- Crespel A et al. (2017) The development of contemporary European sea bass larvae (*Dicentrarchus labrax*) is not affected by projected ocean acidification scenarios *Mar Biol* 164 doi:ARTN 15510.1007/s00227-017-3178-x
- Dixon DL, Jennings AR, Atema J, Munday PL (2015) Odor tracking in sharks is reduced under future ocean acidification conditions *Global Change Biol* 21:1454-1462 doi:10.1111/gcb.12678
- Dixon DL, Munday PL, Jones GP (2010) Ocean acidification disrupts the innate ability of fish to detect predator olfactory cues *Ecol Lett* 13:68-75 doi:10.1111/j.1461-0248.2009.01400.x
- Domenici P, Allan B, McCormick MI, Munday PL (2012) Elevated carbon dioxide affects behavioural lateralization in a coral reef fish *Biol Lett* 8:78-81 doi:10.1098/rsbl.2011.0591
- Doney SC, Fabry VJ, Feely RA, Kleypas JA (2009) Ocean acidification: the other CO₂ problem *Ann Rev Mar Sci* 1:169-192 doi:10.1146/annurev.marine.010908.163834
- Edwards JG, Michel WC (2002) Odor-stimulated glutamatergic neurotransmission in the zebrafish olfactory bulb *J Comp Neurol* 454:294-309 doi:10.1002/cne.10445
- Ferrari MCO et al. (2012) Effects of Ocean Acidification on Learning in Coral Reef Fishes *Plos One* 7 doi:ARTN e3147810.1371/journal.pone.0031478
- Field CB, V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White (2014) Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. In: Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA,, p 1132
- Forsgren E, Dupont S, Jutfelt F, Amundsen T (2013) Elevated CO₂ affects embryonic development and larval phototaxis in a temperate marine fish *Ecol Evol* 3:3637-3646 doi:10.1002/ece3.709
- Horzmann KA, Freeman JL (2016) Zebrafish Get Connected: Investigating Neurotransmission Targets and Alterations in Chemical Toxicity *Toxics* 4:19 doi:10.3390/toxics4030019
- Jutfelt F, Bresolin de Souza K, Vuylsteke A, Sturve J (2013) Behavioural disturbances in a temperate fish exposed to sustained high-CO₂ levels *Plos One* 8:e65825 doi:10.1371/journal.pone.0065825
- Kawai T, Abe H, Oka Y (2012) Dopaminergic neuromodulation of synaptic transmission between mitral and granule cells in the teleost olfactory bulb *J Neurophysiol* 107:1313-1324
- Kermen F, Franco L, Wyatt C, Yaksi E (2013) Neural circuits mediating olfactory-driven behavior in fish vol 7. doi:10.3389/fncir.2013.00062

- Lai F, Fagernes CE, Bernier NJ, Miller GM, Munday PL, Jutfelt F, Nilsson GE (2017a) Responses of neurogenesis and neuroplasticity related genes to elevated CO₂ levels in the brain of three teleost species *Biol Letters* 13 doi:ARTN 2017024010.1098/rsbl.2017.0240
- Lai F, Fagernes CE, Jutfelt F, Nilsson GE (2017b) Expression of genes involved in brain GABAergic neurotransmission in three-spined stickleback exposed to near-future CO₂ (vol 4, pg 1, 2016) *Conserv Physiol* 5 doi:ARTN cox00410.1093/conphys/cox004
- Munday PL, Dixon DL, Donelson JM, Jones GP, Pratchett MS, Devitsina GV, Doving KB (2009) Ocean acidification impairs olfactory discrimination and homing ability of a marine fish *Proc Natl Acad Sci U S A* 106:1848-1852 doi:10.1073/pnas.0809996106
- Nilsson GE, Dixon DL, Domenici P, McCormick MI, Sorensen C, Watson SA, Munday PL (2012) Near-future carbon dioxide levels alter fish behaviour by interfering with neurotransmitter function *Nat Clim Change* 2:201-204 doi:10.1038/Nclimate1352
- Porteus CS, Hubbard PC, Webster TMU, van Aerie R, Canario AVM, Santos EM, Wilson RW (2018) Near-future CO₂ levels impair the olfactory system of a marine fish *Nat Clim Change* 8:737-+ doi:10.1038/s41558-018-0224-8
- Regan MD et al. (2016) Ambient CO₂, fish behaviour and altered GABAergic neurotransmission: exploring the mechanism of CO₂-altered behaviour by taking a hypercapnia dweller down to low CO₂ levels *Journal of Experimental Biology* 219:109-118 doi:10.1242/jeb.131375
- Rong JH et al. (2018) Ocean Acidification Impairs Foraging Behavior by Interfering With Olfactory Neural Signal Transduction in Black Sea Bream, *Acanthopagrus schlegelii* *Front Physiol* 9 doi:ARTN 159210.3389/fphys.2018.01592
- Rossi T, Nagelkerken I, Pistevos JCA, Connell SD (2016) Lost at sea: ocean acidification undermines larval fish orientation via altered hearing and marine soundscape modification *Biol Letters* 12 doi:ARTN 2015093710.1098/rsbl.2015.0937
- Stillman JH, Paganini AW (2015) Biochemical adaptation to ocean acidification *J Exp Biol* 218:1946-1955 doi:10.1242/jeb.115584
- Sundin J, Amcoff M, Mateos-Gonzalez F, Raby GD, Clark TD (2019) Long-term acclimation to near-future ocean acidification has negligible effects on energetic attributes in a juvenile coral reef fish *Oecologia* 190:689-702 doi:10.1007/s00442-019-04430-z
- Tabor R, Yaksi E, Friedrich RW (2008) Multiple functions of GABA A and GABA B receptors during pattern processing in the zebrafish olfactory bulb *Eur J Neurosci* 28:117-127 doi:10.1111/j.1460-9568.2008.06316.x
- Wang M, Jeong CB, Lee YH, Lee JS (2018) Effects of ocean acidification on copepods *Aquatic toxicology* (Amsterdam, Netherlands) 196:17-24 doi:10.1016/j.aquatox.2018.01.004
- Williams CR et al. (2019) Elevated CO₂ impairs olfactory-mediated neural and behavioral responses and gene expression in ocean-phase coho salmon (*Oncorhynchus kisutch*) *Global Change Biol* 25:963-977 doi:10.1111/gcb.14532
- Zupanc GK (2001) Adult neurogenesis and neuronal regeneration in the central nervous system of teleost fish *Brain Behav Evol* 58:250-275 doi:10.1159/000057569