



HAL
open science

Small-scale biophysical couplings in Lake Baikal: new insights from the fractal properties of temperature and phytoplankton distributions

L. Seuront, Y. Tanaka, T. Katano, M. Sugiyama, Y. Watanabe, M. Shimaraev, V. Drucker

► To cite this version:

L. Seuront, Y. Tanaka, T. Katano, M. Sugiyama, Y. Watanabe, et al.. Small-scale biophysical couplings in Lake Baikal: new insights from the fractal properties of temperature and phytoplankton distributions. *Freshwater Biology*, 2020, 4, pp.691-692. 10.31951/2658-3518-2020-A-4-691 . hal-03004804

HAL Id: hal-03004804

<https://hal.science/hal-03004804>

Submitted on 5 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Short communication

ISSN 2658-3518

LIMNOLOGY
FRESHWATER
BIOLOGY

www.limnolwbiol.com

Small-scale biophysical couplings in Lake Baikal: new insights from the fractal properties of temperature and phytoplankton distributions

Seuront L.^{1,2,3*}, Tanaka Y.⁴, Katano T.^{4,6}, Sugiyama M.⁵, Watanabe Y.⁶, Shimaraev M.N.⁷, Drucker V.V.⁷

¹ The University of Lille, CNRS, Univ. Littoral Côte d'Opale, UMR 8187, LOG, Laboratoire d'Océanologie et de Géosciences, F-59000, Lille, France

² Department of Marine Resources and Energy, TUMSAT, Tokyo, Japan

³ Department of Zoology and Entomology, Rhodes University, Grahamstown, South Africa

⁴ Department of Ocean Sciences, TUMSAT, Tokyo, Japan

⁵ Department of Natural Resources, Kyoto University, Kyoto, Japan

⁶ Faculty of Science, Tokyo Metropolitan University, Tokyo, Japan

⁷ Limnological Institute, SB RAS, Irkutsk, Russia

ABSTRACT. Using vertical profiles of temperature and *in vivo* fluorescence sampled between Tanhoy and Listvyanka, we show that temperature and fluorescence fluctuations exhibit specific levels of internal structuration that are independent of the overall vertical structure of the water column, which suggest a previously untapped small-scale biophysical coupling.

Keywords: Couplings, plankton, passive and biologically-active scalars, turbulence, fractals

1. Introduction

Since the pioneering work of the late Trevor Platt (1942-2020) in the seventies, the study of biophysical couplings in lakes and oceans has been a very prolific research topic. Physical processes were often implicitly assumed to be the main factors structuring biological communities, especially in surface waters that are typically under the influence of turbulence. This essentially led to assess the nature of biological couplings through a comparison of the fluctuations of passive scalars (e.g. temperature, salinity, and *a priori* phytoplankton cells), which have widely been described in Fourier space using power spectral analysis as $E(f) \propto f^\beta$ where f is the frequency (s^{-1}) or the wavelength (m^{-1}) and $\beta = 5/3$ for passive scalars, and $\beta \neq 5/3$ for biologically active scalars. From the mid-nineties, fractals and multifractals have been proposed as a new tool to assess the complexity of passive and biologically active scalars; see Seuront (2010) for a review. These methods, together with the increase in both spatial and temporal resolutions of physical and biological sensors, increasingly led to identify distinct structures in the fluctuations of physical and biological scalars. In this context, to illustrate the applicability of the fractal approach in lakes, we used a set of temperature and *in vivo* fluorescence vertical profiles taken across Lake Baikal in August 2003 between Tanhoy and

Listvyanka to compare the nature of temperature and phytoplankton fluctuations under different scenario of vertical distributions (Fig.).

2. Materials and Methods

We investigated the properties of both temperature and fluorescence fluctuations using a method derived from geostatistics and regionalized variables based on the behavior of the semivariance $\gamma(h)$ defined as:

$$\gamma(h) = \frac{1}{2N(h)} \sum_{i=1}^{N(h)} [Z(i) - Z(i+h)]^2$$

where $Z(i+h)$ is the value of the dependent variable $z(i)$ at a depth separated from depth i by the distance, or lag, h , and $N(h)$ is the number of pairs of data points separated by the lag h . Because semivariance analysis is based on the assumption of at least reduced stationarity, our analyses were based on the first difference of both temperature and fluorescence, i.e. $Z(i) = [dx/dz]$, where x is either temperature or fluorescence and z the depth. If the semivariance is a power-law of the scale h , i.e. $\gamma(h) \propto h^m$, where m is the slope of the log-log plot of $\gamma(h)$ versus h , the fractal dimension D ($D = (4 - m)/2$, $D \in [1,2]$) can be used as an index of the degree of spatial structure of $z(i)$. Specifically, the limits

*Corresponding author.

E-mail address: laurent.seuront@cnrs.fr (L. Seuront)






Fig. Study area on the Lake Baikal (A), location (B) and vertical structure of temperature and *in vivo* fluorescence (C) profiles analyzed in the present work, and (D) the fractal dimension D_F of fluorescence fluctuations shown as a function of the fractal dimension D_T of temperature fluctuations. Each of the 9 color codes used in (B), (C) and (D) consistently refer to a specific vertical profile.

$D = 1$ and $D = 2$ characterized vertical distributions where the differences between nearby measurements are respectively higher than and no different from the differences between distant samples.

3. Results and Discussion

All the vertical profiles investigated showed fractal properties, irrespective of the actual vertical structure of the water column (Fig.). The fractal dimensions of temperature fluctuations (D_T) ranged between 1.36 and 1.95 and were significantly ($p < 0.01$) smaller than the fractal dimensions of fluorescence (D_F) distributions, which ranged from 1.72 and 1.93. Noticeably, D_T and D_F were consistently significantly larger than the theoretical value $D = 1.67$ expected for a passive scalar under the influence of three-dimensional fully developed turbulence. Finally,

D_F were significantly positively correlated with D_T , i.e. $D_F = 0.19D_T + 1.48$ ($r^2 = 0.29$, $p < 0.05$).

These preliminary results show that vertical profiles of both physical and biological parameters present specific levels of internal structure that are independent of the overall vertical structure of the water column. More specifically, they indicate (i) a stronger spatial structure in phytoplankton than in temperature, (ii) a lack of control of both temperature and phytoplankton by turbulent velocity fluctuations, and (iii) the existence of a potential causal link between temperature and phytoplankton fluctuations, hence a form of small-scale biophysical coupling.

References

Seuront L. 2010. Fractals and multifractals in ecology and aquatic science. Boca Raton: CRC Press.