



HAL
open science

Comparison of five methodologies to apportion organic aerosol sources during a PM pollution event

D. Srivastava, K.R. Daellenbach, Y. Zhang, N. Bonnaire, B. Chazeau, E. Perraudin, V. Gros, F. Lucarelli, E. Villenave, A.S.H. Prévôt, et al.

► To cite this version:

D. Srivastava, K.R. Daellenbach, Y. Zhang, N. Bonnaire, B. Chazeau, et al.. Comparison of five methodologies to apportion organic aerosol sources during a PM pollution event. *Science of the Total Environment*, 2021, 757, pp.143168. 10.1016/j.scitotenv.2020.143168 . hal-03006211

HAL Id: hal-03006211

<https://hal.science/hal-03006211>

Submitted on 30 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comparison of five methodologies to apportion organic aerosol sources during a PM pollution event

D. Srivastava^{1,2,3,*}, K. R. Daellenbach⁴, Y. Zhang¹, N. Bonnaire⁵, B. Chazeau⁴, E. Perraudin^{2,3}, V. Gros⁵, F. Lucarelli⁶, E. Villenave^{2,3}, A. S. H. Prévôt⁴, I. El Haddad⁴, O. Favez¹, A. Albinet^{1,*}

¹INERIS, Parc Technologique Alata, BP 2, 60550 Verneuil-en-Halatte, France

²CNRS, EPOC, UMR 5805 CNRS, 33405 Talence, France

³Université de Bordeaux, EPOC, UMR 5805 CNRS, 33405 Talence, France

⁴Paul Scherrer Institute (PSI), 5232 Villigen, Switzerland

⁵LSCE - UMR8212, CNRS-CEA-UVSQ, 91191 Gif-sur-Yvette, France

⁶University of Florence, Dipartimento di Fisica Astronomia, 50019, Sesto Fiorentino, Italy

* *Correspondence to:* Alexandre Albinet (alexandre.albinet@gmail.com;

alexandre.albinet@ineris.fr) and Deepchandra Srivastava (deepchandra.srivastava@gmail.com)

Abstract

This study presents a comparison of five methodologies to apportion primary (POA) and secondary organic aerosol (SOA) sources from measurements performed in the Paris region (France) during a highly processed PM pollution event. POA fractions, estimated from EC-tracer method and positive matrix factorization (PMF) analyses, conducted on measurements from PM₁₀ filters, aerosol chemical speciation monitor (ACSM) and offline aerosol mass spectrometry (AMS), were all comparable (2.2 - 3.7 $\mu\text{g m}^{-3}$ as primary organic carbon (POC)). Associated relative uncertainties (measurement + model) on POC estimations ranged from 8 to 50%. The best apportionment of primary traffic OA was achieved using key markers (EC and 1-nitropyrene) in the chemical speciation-based PMF showing more pronounced rush-hour peaks and greater correlation with NO_x than other traffic related POC factors. All biomass burning-related factors were in good agreement, with a typical diel profile and a night-time increase linked to residential heating. If PMF applied to ACSM data showed good agreement with other PMF outputs corrected from dust-related factors (coarse PM), discrepancies were observed between individual POA factors (traffic, biomass burning) and directly comparable SOA factors and highly oxidized OA. Similar secondary organic carbon (SOC) concentrations ($3.3 \pm 0.1 \mu\text{g m}^{-3}$) were obtained from all approaches, except the SOA-tracer method ($1.8 \mu\text{g m}^{-3}$). Associated uncertainties ranged from 14 to 52% with larger uncertainties obtained for PMF-chemical data, EC- and SOA-tracer methods. This latter significantly underestimated total SOA loadings, even including biomass burning SOA, due to missing SOA classes and precursors. None of the approaches was able to identify the formation mechanisms and/or precursors responsible for the highly oxidized SOA fraction associated with nitrate- and/or sulfate-rich aerosols (35% of OA). We recommend the use of a

combination of different methodologies to apportion the POC/SOC concentrations/contributions to get the highest level of confidence in the estimates obtained.

Keywords: Aerosols; EC-tracer; SOA-tracer; PMF; Offline AMS; ACSM

1. Introduction

Airborne organic aerosols (OA) are commonly classified as primary (POA), i.e., directly emitted, or secondary (SOA), i.e., resulting from chemical (trans-) formation processes occurring in the atmosphere (Hallquist et al., 2009). POA and SOA have major impacts on human health, biogeochemical cycles and the Earth's climate (Heal et al., 2012; Boucher et al., 2013). However, determining their origin remains challenging due to the complexity and variability of the processes involved (Donahue et al., 2009). Most common methodologies applied to apportion POA and/or SOA include: elemental carbon-tracer (EC) approach (Grosjean, 1984; Turpin and Huntzicker, 1995; Gray et al., 1986), various statistical receptor models (e.g. chemical mass balance (CMB) and positive matrix factorization (PMF)) (Watson et al., 1990; Paatero, 1997; Paatero and Tapper, 1994), SOA-tracer method (Kleindienst et al., 2007) and radiocarbon (^{14}C) measurements (Szidat et al., 2009; Gelencsér et al., 2007). These methodologies have been successfully applied worldwide and, in general, good agreements have been reported when compared by twos directly (Song et al., 2006; Kim et al., 2004; Kleindienst et al., 2010; Pachon et al., 2010; Heo et al., 2013; Al-Naiema et al., 2018; Srivastava et al., 2018b; Bove et al., 2018; Bae et al., 2019; Antony Chen and Cao, 2018; Hettiyadura et al., 2018; Jiang et al., 2018; Keerthi et al., 2018; Matawle et al., 2018; Shi et al., 2018; Shirmohammadi et al., 2016; Lanzafame et al., 2020). However, large

discrepancies have been also observed when apportioning the SOA fraction, depending on the methods compared, the period of the year considered or atmospheric conditions observed (Srivastava et al., 2018b). As an example, in a recent study conducted in the Paris region (France) by Lanzafame et al. (2020), a good agreement between total secondary organic carbon (SOC) estimations, evaluated by the SOA-tracer method and the filter-based PMF analysis, was observed over the year investigated, but substantial differences were noticed during a secondary PM (particulate matter) pollution event (where ammonium nitrate dominated). Similar observations have been reported by other authors during high pollution episodes or when high secondary contributions were observed (Srivastava et al., 2018b; Zhang et al., 2009; Hu et al., 2010; El Haddad et al., 2011; Feng et al., 2013).

Such high secondary pollution episodes are often observed in Western Europe during late winter–early spring (Petit et al., 2015; Dupont et al., 2016; Petit et al., 2017; Tarrasón, 2016; Hamer, 2017). PM pollution episodes are defined when the daily PM₁₀ concentrations exceeded the European regulatory threshold of 50 µg m⁻³ for at least 3 consecutive days. Emissions from domestic heating, road transport and manure spreading along with anticyclonic atmospheric conditions result in the transport and/or accumulation of pollutants, as well as photochemical processes, within the boundary layer during such events (Waked et al., 2014; Petit et al., 2017; Dupont et al., 2016). Thus, such episodes are often dominated by a high proportion of secondary pollutants (especially ammonium nitrate and SOA) (Petit et al., 2017; Beekmann et al., 2015; Srivastava et al., 2018a; Srivastava et al., 2018c; Waked et al., 2014; Weber et al., 2019; Putaud et al., 2004; Putaud et al., 2010). Similar events have been observed in China in recent years; reduction in SO₂ emissions results in the shift of sulfate to nitrate dominated haze pollution events (Xie et al., 2020; Xu et al., 2019; Tian et al., 2019; Wang et al., 2020). During such events, organic matter (OM) is the second

contributor to the PM mass concentrations observed (Putaud et al., 2004) and, as meteorological and photooxidant conditions promote chemical processes, a significant fraction of OA is probably of secondary origin (Zhang et al., 2019; Huang et al., 2014; Petit et al., 2017; Srivastava et al., 2018a; Srivastava et al., 2019; Srivastava et al., 2018c; Tomaz et al., 2017). However, OA sources during these highly processed PM pollution episodes are still poorly addressed while it is critical in terms of air quality policy management. Subjectivity in the choice of the source apportionment method, final solutions and the assumptions applied, might have strong impacts on the results and conclusions obtained. Comparison of results obtained from different approaches to apportion POA/SOA fractions during such events would highlight discrepancies and uncertainties between each approach and further understanding of OA sources.

In this work, POA and SOA fractions have been resolved using various source apportionment methods, applied to different offline/online datasets collected from a short-term intensive campaign in the Paris area (France) during a springtime PM pollution event. PMF analyses have been conducted using extensive filter-based chemical speciation data, as well as OA mass spectra obtained from online aerosol chemical speciation monitor (ACSM) and offline filter-based aerosol mass spectrometry (AMS) measurements. EC-tracer and SOA-tracer methods have also been applied to the filter-based dataset. Overall, this study evaluates the consistency of outputs retrieved from these methodologies and explores their potential limitations when applied independently to each other in a highly processed environment. To the best of our knowledge, this work is one of the few source apportionment analyses applied to PM pollution episodes to characterize and/or investigate the existing difference between the approaches for OA sources.

2. Experimental

Details about the monitoring site, online measurements, sample collection, chemical speciation analytical procedures as well as backward trajectory analyses have already been reported in two previous articles (Srivastava et al., 2019; Srivastava et al., 2018a) and in the supporting material of the present one (SM). Only the essential information is presented in this section.

2.1. Monitoring site

Measurements were conducted at the ACTRIS-SIRTA observatory (Site Instrumental de Recherche par Télédétection Atmosphérique, 2.15° E; 48.71° N; 150 m; <http://sirta.ipsl.fr>), a well-established facility for the long-term monitoring of physical and chemical aerosol properties in the Paris area (France) (Zhang et al., 2019; Haeffelin et al., 2005). The site is located approximately 25 km southwest of Paris city center and is considered as representative of the background air quality of the Ile-de-France region with influence of the Paris plume under anticyclonic conditions. An intensive campaign was conducted on purpose during a PM pollution event (daily PM₁₀ concentrations >50 µg m⁻³ for several consecutive days) from 6 to 21 March 2015.

2.2. Online instrumentation

PM₁₀ (TEOM 1405F, Thermo), NO_x (T200UP, Teledyne API), O₃ (T400, Teledyne API) and black carbon (BC) (aethalometer AE33, Magee Scientific) concentrations were measured at 15, 1, 1 and 1-min time resolutions, respectively. In addition, meteorological parameters such as temperature, relative humidity (RH), wind direction, and wind speed were also measured at 1-min time resolution. The ACSM (Aerodyne Research) allowed for the measurement of major submicron

(PM₁) non-refractory (NR) chemical species at about 30-min time resolution. Details on these measurements can be found elsewhere (Petit et al., 2017; Zhang et al., 2019). The ACSM dataset used in this work has already been reported previously (Srivastava et al., 2019).

2.3. Filter sample collection and analysis

A high-volume sampler (DA-80, Digital; flow rate: 30 m³ h⁻¹) was used to collect PM₁₀ samples (Tissu-quartz fibre filter, Pallflex, Ø = 150 mm) every 4 hours. A total number of 92 filter samples were collected and analyzed for an extended chemical speciation. Major ions (Cl⁻, NO₃⁻, SO₄²⁻, NH₄⁺, Ca²⁺, Na⁺, Mg²⁺, K⁺) (Guinot et al., 2007; CEN, 2017b), EC and organic carbon (OC) (CEN, 2017a; Cavalli et al., 2010), 7 elemental species (Ca, Ti, Mn, Fe, Ni, Cu, Pb), methanesulfonic acid (MSA), oxalate (C₂O₄²⁻), 3 anhydrosugars (levoglucosan, mannosan and galactosan), 3 polyols/sugar alcohols (arabitol, sorbitol and mannitol) (Verlhac et al., 2013; Yttri et al., 2015), 9 polycyclic aromatic hydrocarbons (PAHs), 14 oxy-PAHs, 8 nitro-PAHs (Albinet et al., 2014; Albinet et al., 2013; Albinet et al., 2006; Tomaz et al., 2016) and 13 SOA markers (e.g., α -methylglyceric acid, pinic acid, and methyl-nitrocatechols) (Albinet et al., 2019; Srivastava et al., 2018c) were analyzed following the protocols already detailed previously. Offline AMS analysis focusing on the bulk composition of the organic aerosol was also performed on PM₁₀ filter samples following the procedure developed previously (Daellenbach et al., 2016).

2.4. Source apportionment approaches

Each source apportionment methodology (EC-tracer, SOA-tracer and PMF analyses performed on the different datasets) is briefly discussed below. Details on the calculation of the overall

uncertainties (measurement + model) linked to POC and SOC estimates using these approaches are presented in the SM. Limitations linked to these approaches are also briefly discussed in the SM and also explained elsewhere (Srivastava et al., 2018b). Note that for measurements, the uncertainties linked to the sampling were not considered. For filter-based methodologies, the uncertainty on the sampling volume collected would have been the same for all the methods and usually accounts for a minor part of the measurement uncertainty (Ringuet et al., 2012; Albinet et al., 2013; Albinet et al., 2014). In addition, as no denuder has been used for the PM₁₀ filter samplings, we are aware that some sampling artifacts (positive, overestimation of the concentrations, or negative, underestimation, due to the sorption or desorption on/from the filter of semi-volatile species and/or due to the degradation/formation of chemical species by heterogeneous processes involving atmospheric oxidants) could induce additional measurement uncertainties, notably for the different organic species quantified here, but that are really difficult to evaluate (Albinet et al., 2010; Goriaux et al., 2006; Mader et al., 2001; Turpin et al., 2000 and references therein). Similarly, uncertainties linked to the measurement artifacts with ACSM or off-line AMS due to the CO₂⁺/NO₃ effect (Pieber et al., 2016; Freney et al., 2019), were not considered in the overall uncertainty estimations.

2.4.1. EC-tracer method

The EC-tracer method has been widely used to estimate the partitioning of measured particulate OC into primary and secondary fractions (Grosjean, 1984; Turpin and Huntzicker, 1995; Castro et al., 1999; Chu, 2005; Saylor et al., 2006; Gray et al., 1986). Briefly, it takes advantage of primary OC and EC co-emissions to estimate SOC from the magnitude of OC-to-EC ratios measured in ambient air (Srivastava et al., 2018b). Here, a primary OC-to-EC ratio ([OC/EC]_p) of 2.9 was

estimated from the measurements during the periods of low photochemical activity. This $[\text{OC}/\text{EC}]_p$ value was used to calculate SOC concentrations (Fig. S1).

2.4.2. PMF-based approaches

Detailed information on PMF principle can be found elsewhere (Paatero and Tapper, 1994; Paatero, 1997). Briefly, this receptor model resolves factor profiles and contributions from a time series of observations using weighted least-squares fitting approach, where the weights are adjusted according to measurement uncertainties. The choice of the optimal solution is notably based on the minimization of the residuals obtained between modeled and observed input species concentrations.

The U.S. Environmental Protection Agency (US-EPA) PMF 5.0 toolkit has been used to perform the source apportionment on the PM_{10} filter chemical dataset (including inorganic species and metals, along with EC, OC and organic markers). As detailed in Srivastava et al. (2018a), OC concentrations obtained for each relevant factor were then used in this study (Figs. S2 and S3).

The information linked to OA source apportionment using the OA mass spectra from ACSM has been given by Srivastava et al. (2019) (Fig. S4). The offline-AMS PMF analyses details are provided in the SM (Figs. S5 to S11). For these analyses, OA mass spectra were treated using the Source Finder toolkit (SoFi) (Canonaco et al., 2013). OC concentrations related to the ACSM factors were further calculated applying OC-to-OA conversion factors specific to each source, i.e., 1.7 for biomass burning (Puxbaum et al., 2007), 1.2 for vehicular emissions (van Drooge and Grimalt, 2015) and 2.0 for secondary organics (Mohr et al., 2009). In the case of offline-AMS, OC-to-OA ratio was determined as explained by Canagaratna et al. (2015) and used to evaluate the OC concentrations of relevant OA factors.

Both US-EPA and SoFi toolkits use the multilinear engine (ME-2) algorithm, allowing the implementation of constraints on the factor chemical profiles and/or time series.

2.4.3. SOA-tracer method

The SOA-tracer method was developed to estimate the contribution of SOA fractions associated with individual gaseous precursors. SOC mass fractions are estimated using conversion factors to calculate SOC loadings from molecular marker concentrations (Kleindienst et al., 2007). A clear limitation of this methodology is related to the fact that only a limited number of organic markers can be accounted (Srivastava et al., 2018b).

SOC mass fractions (anthropogenic and biogenic) were modified according to a subset of markers analyzed following the procedure discussed previously (Rutter et al., 2014) (Tables S1 and S2). In addition, biomass burning SOA fraction was also estimated using the SOA-tracer method. Neglecting this SOA source might lead to significant underestimation of the total wintertime SOC concentrations in Europe due to relatively high contributions of residential wood burning during the cold season (Srivastava et al., 2018b; Petit et al., 2014; Ciarelli et al., 2017; Puxbaum et al., 2007; Daellenbach et al., 2017; Srivastava et al., 2019; Languille et al., 2020). Details on the estimation of biomass burning SOA fraction can be found elsewhere (Lanzafame et al., 2020).

3. Results and discussion

3.1. Overview of the PM pollution event and chemical composition

An overview of the PM chemical composition during the studied period is given in Fig. 1. The daily PM₁₀ concentrations were in the range of 12-130 $\mu\text{g m}^{-3}$, with an average of 49 $\mu\text{g m}^{-3}$. A large predominance of secondary inorganic species, especially ammonium nitrate, was observed during the pollution episode as expected highlighting the significance of secondary processes (Petit et al., 2017). OM concentrations ranged from 2-25 $\mu\text{g m}^{-3}$, with an average value of about 12 $\mu\text{g m}^{-3}$. Slight differences can be observed between the measured and the reconstructed PM₁₀ mass concentrations (slope of 0.83) due to the PM water content and/or some sampling artefacts together with the measurement uncertainties.

3.2. Number of distinguished POA and SOA classes

The identification of each of these factors is summarized in Table 1 and their contributions to total OC in PM₁₀ is shown in Fig. 2. As defined, the EC-tracer method resulted in the estimation of 1 primary and 1 secondary factor. The SOA-tracer method allowed for the quantification of 5 different SOA fractions corresponding to the 5 categories of SOA markers used. PMF analysis conducted on the chemical dataset (PMF-chemical data) led to the discrimination of 10 different OA factors, while analyses on the offline AMS and ACSM datasets (PMF-offline AMS and PMF-ACSM) allowed the identification of 6 and 4 OA factors, respectively. Note that, as ACSM measurements consider only the PM₁, the difference between OC in PM₁₀ (from filter measurements) and in PM₁ was affected by coarse OC (Fig. 2).

Combustion-related factors highly influenced by primary organic molecular markers (for PMF-chemical data) or by hydrocarbon mass fragments (for PMF-offline AMS and PMF-ACSM) were attributed to the POA fraction. The factors mainly influenced by secondary markers or oxygenated mass fragments were ascribed to the SOA fraction. Dust-related OA factors (retrieved from PMF-

chemical data as well as PMF-offline AMS) could be linked to both primary and secondary aerosols due to soil abrasion, resuspension and/or coagulation processes and to condensation of semi-volatile organic species onto mineral dust particles. The dust factor identified by PMF-chemical data contained a significant amount of EC (Fig. S2), along with metals (mainly crustal) and cations (Ti, Ca²⁺, Mg²⁺, Cu and Mn), with negligible amount of secondary species present in the factor profile. This supports the hypothesis of considering this factor to be primary. The SCOA (sulfur-containing OA) factor identified by PMF-offline AMS characterized based on the presence of sulfur containing fragments (CH₃SO₂⁺) in the chemical profile, has been found to be mainly coarse and originate from primary emissions based on previous studies (Vlachou et al., 2018; Daellenbach et al., 2017; Bozzetti et al., 2016). This was also supported as it showed good correlations with crustal metals and therefore this “dust” factor has been attributed to the POA fraction.

Such a disparity in the number of identified factors impeded their direct comparison. However, the comparison of POA and SOA fractions apportioned can be discussed as follows.

3.3. Comparison of the POA factors

Fig. 3 shows the primary OC (POC) concentrations obtained from the different methodologies used for POA apportionment. On average, a good agreement between the outputs from EC-tracer method, PMF-chemical data, and PMF-offline AMS was observed (Fig. 3b). Averaged PM₁₀ POC concentrations obtained from these three methodologies ranged between 3.2 and 3.7 μg m⁻³ representing ~ 50% of the total PM₁₀ OC. Time series were rather similar (datasets normally distributed, paired t-test; p-values > 0.05) except between PMF-chemical data and PMF-offline AMS, as shown in Table S6a and Fig. 3a. However, significant disagreements can be observed during certain days (e.g. 10th, 16th, 19th March) when the PMF-chemical data and EC-tracer showed

high concentrations while both other approaches showed minimum levels. As discussed below, this is mainly due to the retrieval of the traffic-related POC source by the AMS/ACSM based approaches especially during morning rush hours (Figs. 4 and 5). The highest correlation ($r^2=0.64$, $n=92$, $p<0.05$) was observed between PMF-chemical data and EC-tracer method (Table S7a). The overall absolute uncertainties in the POC estimates were in the same range for all methods (about $0.5 - 1.8 \mu\text{g m}^{-3}$) strengthening the good agreement between the different methodologies used. EC-tracer and PMF off-line AMS showed low relative overall uncertainties (32% vs. 15%, respectively). However, larger uncertainties (50% vs 15 % as relative uncertainties) have been obtained for the PMF-chemical data approach (Fig. 3b). It has already been shown that measurement uncertainties, linked to the chemical analysis of the PMF input species (Tables S4 and S5), played a major contribution to the overall uncertainty (Pachon et al., 2010). Here, we tried to consider all the possible sources of uncertainties from the chemical analysis procedures, notably applying a GUM (Guide to the Expression of Uncertainty in Measurement) approach for most of the organic markers (González et al., 2018; White, 2008). In the end, about 90 % of the total estimated uncertainty was associated to the measurement uncertainty and the PMF model uncertainty, estimated following the recommendations of Brown et al. (2015), accounted only for 10%. This highlights that the evaluation of analytical uncertainties must be considered very cautiously given that the input species in the PMF model are weighted by uncertainties.

POC concentrations estimated from the PMF-ACSM analysis were significantly lower (datasets normally distributed, paired t-test; $p\text{-value} < 0.05$) than the three other ones (about $2.2 \mu\text{g m}^{-3}$) (Fig. 3a and Table S6a). This result is mainly related to the difference in the sampling size cut-off between ACSM and filter samplings (PM_1 for ACSM $<$ PM_{10} for filter samples). The time series of POC fraction from the EC-tracer method correlated well with that from PMF-chemical data,

while the POC variations of PMF-offline AMS were much similar with those of PMF-ACSM. These results are likely due to the similarity in chemical composition of the different datasets as the EC-tracer method and PMF-chemical data are based on filter measurements, and on the other side, offline AMS and ACSM both measured the NR-PM species including organics, sulfate, nitrate, ammonium and chloride. As presented in Figs. 2 and S12, the average contribution of the submicron fraction to total OA concentrations was about 77%. The remaining 23% were associated to the coarse OA fraction and possibly attributed to the dust/SCOA factors (20% and 11%, respectively) identified using PMF-chemical data and PMF-offline AMS methodologies.

Considering only the fine OA fraction, total POC concentrations from PMF-ACSM showed a good agreement with both other PMF approaches namely, PMF-offline AMS and PMF-chemical data (Table S7b). However, a significant difference (datasets normally distributed, paired t-test; p-values < 0.05) was found between PMF-ACSM and PMF-offline AMS (so without SCOA) (Table S6a). On average for the whole period of the study, total fine POC concentrations ranged from 2.2 $\mu\text{g m}^{-3}$ (PMF-ACSM) to 2.5 $\mu\text{g m}^{-3}$ (PMF-offline AMS). As observed for the PM_{10} POC, overall uncertainties estimated for the PMF-chemical data method (78%) were much higher than those obtained for both other PMF approaches based on aerosol mass spectrometry measurements (8-19%). Again, the consideration of all the possible sources of uncertainties of measurement by the GUM approach, increased the final uncertainty budget of the PMF-chemical data method. PMF-ACSM and PMF-offline AMS uncertainties were similar though the variability was higher for the PMF-offline AMS analysis. Recovery of OA factors in the offline-AMS analysis which is based on water soluble extraction can be considered responsible for this discrepancy as the hydrophobic nature of the components (e.g., HOA and COA) may lead to such kind of variation. The choice of the chemical profiles used could also have an impact on the overall uncertainty even if it was

supposed to be low, as attempt has been made to choose very specific source specific chemical tracers, based on the literature.

Such a consistency in the fine OA fraction allowed to further compare individual primary PMF factors related to the main combustion sources: biomass burning- and traffic-related OA (Fig. 4). A good agreement was observed between all the identified biomass burning-related factors, with r^2 ranging from 0.48 to 0.61 (Table S8). Similar diel profiles were also obtained from the three PMF approaches, with a significant nighttime increase linked to residential heating activities (Fig. 5). As shown in Figs. 4 and S13, biomass burning-related factor concentration levels ($0.67\text{-}1.19\ \mu\text{g m}^{-3}$) were roughly equivalent throughout the campaign except for a few data points with slightly lower BBOA concentration levels for PMF-ACSM, notably at the beginning of the campaign, inducing a slight underestimation of BBOC by this method. Thus, a possible influence of the super-micrometer ($>1\ \mu\text{m}$) biomass burning-related aerosols cannot be totally ruled-out. It may also be hypothesized that biomass burning OA (ACSM) may be partly included in other factors than BBOA, such as HOA as already shown by some previous ACSM-based studies (Petit et al., 2014; Srivastava et al., 2019; Zhang et al., 2019). This was also supported by significant correlation ($r^2=0.45$, $n=92$, $p<0.05$) observed between BBOA and HOA factors from the PMF-ACSM analysis (Fig. S14).

Consequently, PMF-ACSM traffic related factor frequently showed slightly higher concentrations during biomass burning predominant periods (Figs. 4 and S13). It should also be noted that, on the diel cycle, the morning and evening rush-hours are more pronounced for the PMF-chemical data traffic factor than for the PMF-ACSM and PMF-offline AMS HOA factors (Fig. 5). In addition, traffic factor from PMF-chemical data is clearly better correlated with NO_x and BC_{ff} than the other traffic related POA factors (Fig. S15). These results suggested that the PMF-chemical data analysis

could more accurately describe primary traffic-related OA, which might be related due to the use of key species, such as EC and 1-nitropyrene, in the input chemical data matrix (Srivastava et al., 2019; Srivastava et al., 2018c; Srivastava et al., 2018a; Lanzafame et al., 2020). As mentioned before, the disagreements observed at certain days (e.g. 10th, 16th March) are probably related to the better resolution of the traffic-related POC factor using PMF-chemical data approach. Nevertheless, the relatively good agreement between averaged traffic-related POC concentrations (i.e., about 0.9 $\mu\text{g m}^{-3}$ for both PMF-chemical data and PMF-offline AMS, 1.1 $\mu\text{g m}^{-3}$ for PMF-ACSM) validates the assumption of the predominant vehicular exhaust origin for HOC factors (Lanz et al., 2007; Xu et al., 2014; Ulbrich et al., 2009; Zhang et al., 2007). This suggests that HOA could be a proxy for traffic emissions but less well resolved than using proper markers. For both POA factors, biomass burning and traffic-related, estimated overall uncertainties were in a similar range (0.1 – 2.5 $\mu\text{g m}^{-3}$) strengthening the good agreement between the different methodologies for the estimation of these individual POC factors. Significantly higher uncertainty values were still observed for the PMF-chemical data (228% and 297% for traffic-related POA and for BBOA, respectively) (Fig. S13) suggesting again the significant influence of the measurement uncertainties especially considering all the potential sources of analytical uncertainties in that case. Large differences were also noticed in the relative overall uncertainties for both POA factors estimated using PMF-offline AMS and PMF-ACSM approaches (55% vs. 9% and 52% vs. 22% for HOA and BBOA, respectively) and probably linked to the measurement approaches (off-line vs. on-line measurements).

Finally, for the primary fraction, time series of the PMF-chemical data dust factor and the PMF-offline AMS SCOA factor showed a good agreement for few peaks (Fig. S16). This result suggested that SCOA is probably related in part to primary dust aerosols, in the coarse mode, as

suggested previously (Vlachou et al., 2018; Daellenbach et al., 2017) but additional investigations are required to clearly identify the exact origin of that factor.

3.4. Comparison of the SOA factors

Total SOC concentrations showed a good agreement between all the source apportionment methodologies, except for the SOA-tracer method (Fig. 6 and Table S9). This observation was also confirmed as no statistical difference (datasets normally distributed, paired t-test; p-values > 0.05) was noticed between the different approaches (Table S6b).

Average SOC concentration values ($3.3 \pm 0.1 \mu\text{g m}^{-3}$) were similar for the EC-tracer method, PMF-chemical data, PMF-offline AMS and PMF-ACSM (Fig. 6b). Lower concentrations were derived from the SOA-tracer method (average value of about $1.8 \mu\text{g m}^{-3}$), especially for the second half of the campaign. $\text{SOC}_{\text{SOA-tracer}}$ concentrations obtained here only accounted for SOA formed from the oxidation of isoprene, α -pinene, naphthalene, toluene and phenolic compounds (see section S2 in the SM). Due to the lack of information on other possible SOA hydrocarbon precursors, such as mono- and poly-aromatic compounds, long-chain alkanes and/or alkenes (Srivastava et al., 2018b; Zhao et al., 2014; Tkacik et al., 2012; Lim and Ziemann, 2005), the present SOA-tracer method analysis missed significant additional SOA classes. In addition, relatively high organonitrate and/or organosulfate loadings (Riva et al., 2015; Surratt et al., 2006; Tomaz et al., 2017) could be expected between the 15th and the 21st of March, a period when PM_{10} concentrations were highly dominated by water-soluble inorganic species (Fig. 1) (Srivastava et al., 2018a). However, it should be noted that total SOC concentrations estimated by the SOA-tracer method matched rather well with the outputs from other approaches during the first half of the campaign (before 14th March, Fig. 6a). This suggests relatively fair estimates of SOA-tracer individual factors considered in the present

study and especially for biomass burning linked to the oxidation of phenolic compounds. This is further supported by the comparison between SOA-tracer method and PMF-chemical data outputs calculated over the yearlong (2015) measurement for the same site and showing a good agreement, especially for anthropogenic SOA (Lanzafame et al., 2020). Here, during this PM pollution event, the 3 PMF approaches especially PMF-ACSM and PMF-off-line AMS, were not able to capture the large biomass burning SOC peak observed with the EC- and SOA tracer methods at the beginning of the campaign (08th March) and was considered as POC (or at least in part as for PMF-chemical data). Note this SOC peak showed the same temporal variation as methyl-nitrocatechols, secondary photo-oxidation products of phenolic compounds (i.e., cresols), known to account for major SOA precursors emitted by biomass burning (Inuma et al., 2010), confirming the actual secondary origin of OC at the beginning of the sampling campaign. (Fig. S17).

The absolute uncertainties evaluated for SOC estimates seemed comparable with slightly higher values observed for the EC-tracer and PMF-chemical data methodologies (Fig. 6b) ($0.5\text{-}1.7 \mu\text{g m}^{-3}$). Considering the overall uncertainties obtained, average SOC estimations obtained with all the apportionment methodologies (except SOA-tracer method) used were in very good agreement. The higher uncertainties for the PMF-chemical data (52%) might be linked to the analytical uncertainties of the input species as discussed above (GUM approach for instance). However, the contribution of model uncertainty has increased here by 10% to the overall uncertainty. EC tracer method uncertainty was also high (48%) and mostly related to the primary [OC/EC] ratio evaluation during the cold period, as already reported as a major source of error in the estimation of secondary fraction using this SOC apportionment method (Srivastava et al., 2018b). These observations are consistent with previous results (Srivastava et al., 2018b; Pachon et al., 2010). PMF offline AMS and PMF-ACSM showed lower relative overall uncertainties of about 14%.

Finally, the lower SOC estimations from the SOA-tracer method were also accompanied by comparable relative uncertainties to the other methods (about 27%).

Regarding the individual factors from all approaches, secondary biomass burning contribution resolved using the SOA tracer method and filter-based PMF analysis showed a very good agreement (Fig. S18). Both also showed a reasonable association with primary biomass burning factors resolved based on the mass spectrometry measurements (PMF-offline AMS and PMF-ACSM), indicating that these primary factors may contain some aged aerosols ($r^2=0.44-0.55$, $n=92$, $p<0.05$). SOC from the naphthalene oxidation obtained using the SOA tracer method agreed reasonably well with mixed secondary aerosols obtained by PMF-chemical data analysis ($r^2=0.55$, $n=92$, $p<0.05$).

In addition, highly oxidized OA factors (namely oxidized oxygenated OA (OOA1) and more oxidized oxygenated OA (MO-OOA)), from PMF-offline AMS and PMF-ACSM analyses, respectively) and, marine biogenic SOA and mixed secondary aerosols obtained by PMF-chemical data analysis showed satisfactory correlations (Fig. 7) highlighting the similar origin of the given SOA factors. SOA derived from isoprene, α -pinene and toluene oxidation based on PMF-chemical data and SOA tracer method did not show any direct association with any of the secondary factors, probably due to their low contribution. Nevertheless, LO-OOA (from PMF-ACSM analysis) showed a substantial correlation with primary factors i.e., biomass burning and traffic emissions ($r^2=0.3-0.5$, $n=92$, $p<0.05$), suggesting the influence of anthropogenic emissions.

High concentrations of both highly oxidized OOA factors (OOA1, MO-OOA), marine biogenic SOA + mixed secondary aerosols and SOA from naphthalene oxidation were notably observed during the second half of the campaign (Fig. 8). Along with the significant inorganic loadings (Fig. 1), a probable predominant long-range transport influence might be expected during this period as

reported before (Petit et al., 2017; Srivastava et al., 2018a). A comparable OOA factor has also been reported in winter in other European locations with similar characteristics namely, significant correlation with long range transported secondary inorganic species (Daellenbach et al., 2017; Lanz et al., 2007).

Both highly oxidized OOA factors may include a part of marine aerosols/traces of combustion aerosols (Srivastava et al., 2018a). This was supported as highly oxidized factors showed a good correlation with anthropogenic markers such as phthalic acid and succinic acid ($r^2=0.40-0.50$, $n=92$, $p<0.05$). However, no correlation was observed with sodium/chloride as expected for being considered as an influence of marine origin. Both OOA factors also captured a significant m/z 45 signal, a fragment observed from dicarboxylic acid, supporting the role of highly processed aerosols in ambient air, and has already observed in several studies (Zhang et al., 2005a; Zhang et al., 2005b; Jimenez et al., 2003; Bahreini et al., 2003). Mixed secondary aerosol factor, in the PMF-chemical data analysis, contained also ultimate oxidation by-products like oxalate which is not specific of any precursors or sources, confirming (Srivastava et al., 2018a) the aforementioned hypothesis. This highlighted that a significant amount of SOA could be linked to the aging of the air masses during long range transport, making it challenging to investigate its origins. Finally, none of the used approaches was able to fully identify the specific formation processes and/or the semi-volatile organic precursors responsible for this highly oxidized SOA fraction (accounting for about 35% of total OA in PM_{10} , ($r^2>0.80$, $n=92$, $p<0.05$)). Even combining off-line and on-line data set as reported before, a clear identification of the SOA sources/processes occurring during the last period of the campaign was not possible to achieve (Srivastava et al., 2019). Further studies are needed to investigate the secondary processes associated with nitrate- and/or sulfate-rich aerosols during such highly processed PM pollution events.

4. Conclusions

A comparison of five methodologies to apportion POA and SOA sources, during a PM pollution event in Northern Europe, has been performed.

POA fractions (average POC = 2.2-3.7 $\mu\text{g m}^{-3}$) were well resolved using all approaches. Only lower POC concentrations were estimated by the PMF-ACSM analysis due to the difference in the sampling size cut-off (PM_1 vs PM_{10} for filter-based methods). PMF-chemical data POC estimates were associated with higher relative overall uncertainties (50% vs. 8%) due to the consideration of all the potential sources of errors in the sample chemical analyses.

Approximately half of the OA fraction was secondary (average concentrations = $3.3 \pm 0.1 \mu\text{g m}^{-3}$). However, the estimation of this fraction remains questionable during highly PM processed pollution events. The SOA-tracer method was not able to capture signals from other precursors dominant during the studied period and significantly underestimated SOC ($1.8 \mu\text{g m}^{-3}$). PMF-chemical data presented a good comparison with all other considered approaches and provided valuable information on the SOA contributions from different precursors. Similarly, SOC estimates from PMF-offline AMS and ACSM analysis were comparable, although they did not provide further insight on precursors and/or sources of highly oxidized fractions. The uncertainties of SOC estimates were broadly comparable in absolute value for all methods, but they were higher as relative values (14% vs. 52%) for the EC tracer, PMF-chemical data and SOA-tracer methods.

These results showed that more comprehensive studies focusing on the identification of new SOA tracers from known/unknown class of precursors are needed to fully understand the origins of SOA fractions. Such studies would aid in bridging the gap between SOA contributions from models and measurements and explicitly improve our knowledge on SOA contribution in a highly oxidative environment. In addition, as recommended previously (Srivastava et al., 2018b), this study

demonstrated that to get the highest level of confidence, it is recommended to use a combination of different methodologies, at least more than one, to apportion the POC/SOC concentrations/contributions. Such combination, especially if low and high time resolution measurements are used, could also enhance the understanding of the subtle differences in OA content.

Finally, the use of advanced instruments in near future such as TAG-AMS (thermal desorption aerosol gas chromatograph-AMS) (Williams et al., 2014), soft ionization methods like extractive electrospray (EESI) (Qi et al., 2020; Lopez-Hilfiker et al., 2019; Qi et al., 2019; Stefenelli et al., 2019), or aerosol inlets associated to PTR-MS (proton-transfer reaction-MS) or CIMS (chemical ionization MS) such as FIGAREO (filter inlet for gases and aerosols) (Lopez-Hilfiker et al., 2014), CHARON (chemical analysis of aerosol online) or thermo-desorption (TD) systems (Holzinger et al., 2010; Gkatzelis et al., 2017; Eichler et al., 2015), may provide more in-depth insight into the precursors and SOA formation processes involved during such highly processed PM pollution events.

Acknowledgements

This work has notably been supported by the French Ministry of Environment as part of the activities of the National reference laboratory for air quality monitoring (LCSQA). Aerosol in situ measurements have been performed as part of the H2020 ACTRIS-2 project and partly funded by the ACTRIS-Fr/CLAP initiative at the national level. This work is also the part of the COLOSSAL European COST Action CA16109. The authors gratefully acknowledge François Truong (LSCE) and Robin Aujay-Plouzeau (INERIS) for taking care of samples and instrumentation, as well as

François Kany and Serguei Stavrovski (INERIS) for sample extraction and PAH analyses. They also thank Patrick Bodu for the graphical abstract design.

Author contributions

D.S, O.F., and A.A. proposed the initial idea. O.F. and A.A., designed and led the study. D.S; K.R.D., N.B., B.C and F.L. performed the chemical analyses. D.S., K.R.D and Y.Z conducted the data analyses. I.E.H and A.S.H.P. supported the off-line AMS OA source apportionment analyses. O.F., E.P., E.V. and A.A. supervised D.S.'s PhD work. O.F., V.G. and A.A. supervised Y.Z.'s PhD work. D.S., K.R.D., O.F and A.A. interpreted the data. D.S and A.A. wrote the manuscript with inputs from all co-authors.

Appendix A.

Supplementary material (SM): Supplementary data to this article can be found online.

References

- Al-Naiema, I. M., Hettiyadura, A. P. S., Wallace, H. W., Sanchez, N. P., Madler, C. J., Cevik, B. K., Bui, A. A. T., Kettler, J., Griffin, R. J., and Stone, E. A.: Source apportionment of fine particulate matter in Houston, Texas: insights to secondary organic aerosols, 2018. *Atmos. Chem. Phys.*, 18, 15601-15622, 10.5194/acp-18-15601-2018.
- Albinet, A., Leoz-Garziandia, E., Budzinski, H., and Villenave, E.: Simultaneous analysis of oxygenated and nitrated polycyclic aromatic hydrocarbons on standard reference material 1649a

(urban dust) and on natural ambient air samples by gas chromatography–mass spectrometry with negative ion chemical ionisation, 2006. *J. Chromatogr. A*, 1121, 106-113, <https://doi.org/10.1016/j.chroma.2006.04.043>.

Albinet, A., Papaiconomou, N., Estager, J., Suptil, J., and Besombes, J.-L.: A new ozone denuder for aerosol sampling based on an ionic liquid coating, 2010. *Anal. Bioanal. Chem.*, 396, 857-864.

Albinet, A., Tomaz, S., and Lestremau, F.: A really quick easy cheap effective rugged and safe (QuEChERS) extraction procedure for the analysis of particle-bound PAHs in ambient air and emission samples, 2013. *Sci. Total Environ.*, 450-451, 31-38, 10.1016/j.scitotenv.2013.01.068.

Albinet, A., Nalin, F., Tomaz, S., Beaumont, J., and Lestremau, F.: A simple QuEChERS-like extraction approach for molecular chemical characterization of organic aerosols: application to nitrated and oxygenated PAH derivatives (NPAH and OPAH) quantified by GC–NICIMS, 2014. *Anal. Bioanal. Chem.*, 406, 3131-3148, 10.1007/s00216-014-7760-5.

Albinet, A., Lanzafame, G. M., Srivastava, D., Bonnaire, N., Nalin, F., and Wise, S. A.: Analysis and determination of secondary organic aerosol (SOA) tracers (markers) in particulate matter standard reference material (SRM 1649b, urban dust), 2019. *Anal. Bioanal. Chem.*, 411, 5975-5983, 10.1007/s00216-019-02015-6.

Antony Chen, L. W., and Cao, J.: PM_{2.5} Source Apportionment Using a Hybrid Environmental Receptor Model, 2018. *Environ. Sci. Technol.*, 52, 6357-6369, 10.1021/acs.est.8b00131.

Bae, M.-S., Skiles, M. J., Lai, A. M., Olson, M. R., de Foy, B., and Schauer, J. J.: Assessment of forest fire impacts on carbonaceous aerosols using complementary molecular marker receptor models at two urban locations in California's San Joaquin Valley, 2019. *Environ. Pollut.*, 246, 274-283, <https://doi.org/10.1016/j.envpol.2018.12.013>.

Bahreini, R., Jimenez, J. L., Wang, J., Flagan, R. C., Seinfeld, J. H., Jayne, J. T., and Worsnop, D. R.: Aircraft-based aerosol size and composition measurements during ACE-Asia using an Aerodyne aerosol mass spectrometer, 2003. *J. Geophys. Res.-Atmos.*, 108, 10.1029/2002jd003226.

Beekmann, M., Prévôt, A. S. H., Drewnick, F., Sciare, J., Pandis, S. N., Denier van der Gon, H. A. C., Crippa, M., Freutel, F., Poulain, L., Gherzi, V., Rodriguez, E., Beirle, S., Zotter, P., von der Weiden-Reinmüller, S. L., Bressi, M., Fountoukis, C., Petetin, H., Szidat, S., Schneider, J., Rosso, A., El Haddad, I., Megaritis, A., Zhang, Q. J., Michoud, V., Slowik, J. G., Moukhtar, S., Kolmonen, P., Stohl, A., Eckhardt, S., Borbon, A., Gros, V., Marchand, N., Jaffrezo, J. L., Schwarzenboeck, A., Colomb, A., Wiedensohler, A., Borrmann, S., Lawrence, M., Baklanov, A., and Baltensperger, U.: In situ, satellite measurement and model evidence on the dominant regional contribution to fine particulate matter levels in the Paris megacity, 2015. *Atmos. Chem. Phys.*, 15, 9577-9591, 10.5194/acp-15-9577-2015.

Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., and Lohmann, U.: Clouds and aerosols, in: *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 571-657, 2013.

Bove, M. C., Massabò, D., and Prati, P.: PMF5.0 vs. CMB8.2: An inter-comparison study based on the new European SPECIEUROPE database, 2018. *Atmos. Res.*, 201, 181-188, <https://doi.org/10.1016/j.atmosres.2017.10.021>.

Bozzetti, C., Daellenbach, K. R., Hueglin, C., Fermo, P., Sciare, J., Kasper-Giebl, A., Mazar, Y., Abbaszade, G., El Kazzi, M., Gonzalez, R., Shuster-Meiseles, T., Flasch, M., Wolf, R., Křepelová, A., Canonaco, F., Schnelle-Kreis, J., Slowik, J. G., Zimmermann, R., Rudich, Y., Baltensperger, U., El Haddad, I., and Prévôt, A. S. H.: Size-Resolved Identification, Characterization, and

Quantification of Primary Biological Organic Aerosol at a European Rural Site, 2016. *Environ. Sci. Technol.*, 50, 3425-3434, [10.1021/acs.est.5b05960](https://doi.org/10.1021/acs.est.5b05960).

Bressi, M., Sciare, J., Ghersi, V., Mihalopoulos, N., Petit, J. E., Nicolas, J. B., Moukhtar, S., Rosso, A., Féron, A., Bonnaire, N., Poulakis, E., and Theodosi, C.: Sources and geographical origins of fine aerosols in Paris (France), 2014. *Atmos. Chem. Phys.*, 14, 8813-8839, [10.5194/acp-14-8813-2014](https://doi.org/10.5194/acp-14-8813-2014).

Brown, S. G., Eberly, S., Paatero, P., and Norris, G. A.: Methods for estimating uncertainty in PMF solutions: Examples with ambient air and water quality data and guidance on reporting PMF results, 2015. *Sci. Total Environ.*, 518-519, 626-635, <https://doi.org/10.1016/j.scitotenv.2015.01.022>.

Canagaratna, M. R., Jimenez, J. L., Kroll, J. H., Chen, Q., Kessler, S. H., Massoli, P., Hildebrandt Ruiz, L., Fortner, E., Williams, L. R., Wilson, K. R., Surratt, J. D., Donahue, N. M., Jayne, J. T., and Worsnop, D. R.: Elemental ratio measurements of organic compounds using aerosol mass spectrometry: characterization, improved calibration, and implications, 2015. *Atmos. Chem. Phys.*, 15, 253-272, [10.5194/acp-15-253-2015](https://doi.org/10.5194/acp-15-253-2015).

Canonaco, F., Crippa, M., Slowik, J., Baltensperger, U., and Prévôt, A.: SoFi, an IGOR-based interface for the efficient use of the generalized multilinear engine (ME-2) for the source apportionment: ME-2 application to aerosol mass spectrometer data, 2013. *Atmos. Meas. Tech.*, 6, 3649-3661.

Castro, L. M., Pio, C. A., Harrison, R. M., and Smith, D. J. T.: Carbonaceous aerosol in urban and rural European atmospheres: estimation of secondary organic carbon concentrations, 1999. *Atmos. Environ.*, 33, 2771-2781, [http://dx.doi.org/10.1016/S1352-2310\(98\)00331-8](http://dx.doi.org/10.1016/S1352-2310(98)00331-8).

Cavalli, F., Viana, M., Yttri, K., Genberg, J., and Putaud, J.-P.: Toward a standardised thermal-optical protocol for measuring atmospheric organic and elemental carbon: the EUSAAR protocol, 2010. *Atmos. Meas. Tech.*, 3, 79-89.

CEN: European Committee for Standardization, EN 16909:2017 - Ambient air - Measurement of elemental carbon (EC) and organic carbon (OC) collected on filters. CEN, Brussels (Belgium). 2017a.

CEN: CEN (European Committee for Standardization): EN 16913:2017 - Ambient air - Standard method for measurement of NO_3^- , SO_4^{2-} , Cl^- , NH_4^+ , Na^+ , K^+ , Mg^{2+} , Ca^{2+} in $\text{PM}_{2.5}$ as deposited on filters. CEN, Brussels (Belgium). 2017b.

Chu, S.-H.: Stable estimate of primary OC/EC ratios in the EC tracer method, 2005. *Atmos. Environ.*, 39, 1383-1392, <http://dx.doi.org/10.1016/j.atmosenv.2004.11.038>.

Ciarelli, G., Aksoyoglu, S., El Haddad, I., Bruns, E. A., Crippa, M., Poulain, L., Äijälä, M., Carbone, S., Freney, E., O'Dowd, C., Baltensperger, U., and Prévôt, A. S. H.: Modelling winter organic aerosol at the European scale with CAMx: evaluation and source apportionment with a VBS parameterization based on novel wood burning smog chamber experiments, 2017. *Atmos. Chem. Phys.*, 17, 7653-7669, 10.5194/acp-17-7653-2017.

Daellenbach, K. R., Bozzetti, C., Křepelová, A., Canonaco, F., Wolf, R., Zotter, P., Fermo, P., Crippa, M., Slowik, J. G., Sosedova, Y., Zhang, Y., Huang, R. J., Poulain, L., Szidat, S., Baltensperger, U., El Haddad, I., and Prévôt, A. S. H.: Characterization and source apportionment of organic aerosol using offline aerosol mass spectrometry, 2016. *Atmos. Meas. Tech.*, 9, 23-39, 10.5194/amt-9-23-2016.

Daellenbach, K. R., Stefenelli, G., Bozzetti, C., Vlachou, A., Fermo, P., Gonzalez, R., Piazzalunga, A., Colombi, C., Canonaco, F., Hueglin, C., Kasper-Giebl, A., Jaffrezo, J. L., Bianchi, F., Slowik,

J. G., Baltensperger, U., El-Haddad, I., and Prévôt, A. S. H.: Long-term chemical analysis and organic aerosol source apportionment at nine sites in central Europe: source identification and uncertainty assessment, 2017. *Atmos. Chem. Phys.*, 17, 13265-13282, 10.5194/acp-17-13265-2017.

Donahue, N. M., Robinson, A. L., and Pandis, S. N.: Atmospheric organic particulate matter: From smoke to secondary organic aerosol, 2009. *Atmos. Environ.*, 43, 94-106, 10.1016/j.atmosenv.2008.09.055.

Dupont, J., Haefelin, M., Badosa, J., Elias, T., Favez, O., Petit, J., Meleux, F., Sciare, J., Crenn, V., and Bonne, J. L.: Role of the boundary layer dynamics effects on an extreme air pollution event in Paris, 2016. *Atmos. Environ.*, 141, 571 - 579, 10.1016/j.atmosenv.2016.06.061.

Eichler, P., Müller, M., D'Anna, B., and Wisthaler, A.: A novel inlet system for online chemical analysis of semi-volatile submicron particulate matter, 2015. *Atmos. Meas. Tech.*, 8, 1353-1360, 10.5194/amt-8-1353-2015.

El Haddad, I., Marchand, N., Temime-Roussel, B., Wortham, H., Piot, C., Besombes, J. L., Baduel, C., Voisin, D., Armengaud, A., and Jaffrezo, J. L.: Insights into the secondary fraction of the organic aerosol in a Mediterranean urban area: Marseille, 2011. *Atmos. Chem. Phys.*, 11, 2059-2079, 10.5194/acp-11-2059-2011.

Feng, J., Li, M., Zhang, P., Gong, S., Zhong, M., Wu, M., Zheng, M., Chen, C., Wang, H., and Lou, S.: Investigation of the sources and seasonal variations of secondary organic aerosols in PM_{2.5} in Shanghai with organic tracers, 2013. *Atmos. Environ.*, 79, 614-622, <http://dx.doi.org/10.1016/j.atmosenv.2013.07.022>.

Freney, E., Zhang, Y., Croteau, P., Amodeo, T., Williams, L., Truong, F., Petit, J.-E., Sciare, J., Sarda-Esteve, R., Bonnaire, N., Arumae, T., Aurela, M., Bougiatioti, A., Mihalopoulos, N., Coz,

E., Artinano, B., Crenn, V., Elste, T., Heikkinen, L., Poulain, L., Wiedensohler, A., Herrmann, H., Priestman, M., Alastuey, A., Stavroulas, I., Tobler, A., Vasilescu, J., Zanca, N., Canagaratna, M., Carbone, C., Flentje, H., Green, D., Maasikmets, M., Marmureanu, L., Minguillon, M. C., Prevot, A. S. H., Gros, V., Jayne, J., and Favez, O.: The second ACTRIS inter-comparison (2016) for Aerosol Chemical Speciation Monitors (ACSM): Calibration protocols and instrument performance evaluations, 2019. *Aerosol Sci. Technol.*, 53, 830-842, 10.1080/02786826.2019.1608901.

Gelencsér, A., May, B., Simpson, D., Sánchez-Ochoa, A., Kasper-Giebl, A., Puxbaum, H., Caseiro, A., Pio, C., and Legrand, M.: Source apportionment of PM_{2.5} organic aerosol over Europe: Primary/secondary, natural/anthropogenic, and fossil/biogenic origin, 2007. *Journal of Geophysical Research*, 112, 10.1029/2006jd008094.

Gkatzelis, G. I., Tillmann, R., Hohaus, T., Müller, M., Eichler, P., Xu, K. M., Schlag, P., Schmitt, S. H., Wegener, R., Kaminski, M., Holzinger, R., Wisthaler, A., and Kiendler-Scharr, A.: Comparison of three aerosol chemical characterization techniques utilizing PTR-ToF-MS: A study on freshly formed and aged biogenic SOA, 2017. *Atmos. Meas. Tech. Discuss.*, 2017, 1-31, 10.5194/amt-2017-288.

González, A., Herrador, M., Garcia Asuero, A., and Martin, J.: A Practical Way to ISO/GUM Measurement Uncertainty for Analytical Assays Including In-House Validation Data, in, 2018.

Goriaux, M., Jourdain, B., Temime, B., Besombes, J. L., Marchand, N., Albinet, A., Leoz-Garziandia, E., and Wortham, H.: Field comparison of particulate PAH measurements using a low-flow denuder device and conventional sampling systems, 2006. *Environ. Sci. Technol.*, 40, 6398-6404.

Gray, H. A., Cass, G. R., Huntzicker, J. J., Heyerdahl, E. K., and Rau, J. A.: Characteristics of atmospheric organic and elemental carbon particle concentrations in Los Angeles, 1986. *Environ. Sci. Technol.*, 20, 580-589, 10.1021/es00148a006.

Grosjean, D.: Particulate carbon in Los Angeles air, 1984. *Sci. Total Environ.*, 32, 133-145.

Guinot, B., Cachier, H., Sciare, J., Tong, Y., Xin, W., and Jianhua, Y.: Beijing aerosol: Atmospheric interactions and new trends, 2007. *J. Geophys. Res.-Atmos.*, 112, n/a-n/a, 10.1029/2006JD008195.

Haefelin, M., Barthès, L., Bock, O., Boitel, C., Bony, S., Bouniol, D., Chepfer, H., Chiriaco, M., Cuesta, J., Delanoë, J., Drobinski, P., Dufresne, J. L., Flamant, C., Grall, M., Hodzic, A., Hourdin, F., Lapouge, F., Lemaître, Y., Mathieu, A., Morille, Y., Naud, C., Noël, V., O'Hirok, W., Pelon, J., Pietras, C., Protat, A., Romand, B., Scialom, G., and Vautard, R.: SIRTA, a ground-based atmospheric observatory for cloud and aerosol research, 2005. *Ann. Geophys.*, 23, 253-275, 10.5194/angeo-23-253-2005.

Hallquist, M., Wenger, J., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N., George, C., and Goldstein, A.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, 2009. *Atmos. Chem. Phys.*, 9, 5155-5236.

Hamer, P. D., Tarrasón, L. C., Guerreiro, F., Meleux, L. Rouil Interim Annual Assessment Report for 2016, 2017. http://policy.atmosphere.copernicus.eu/reports/CAMS-71_2016SC2012_D2071.2011.2011.2016_201707_202016IAR_V201704.pdf.

Heal, M. R., Kumar, P., and Harrison, R. M.: Particles, air quality, policy and health, 2012. *Chem. Soc. Rev.*, 41, 6606-6630.

Heo, J., Dulger, M., Olson, M. R., McGinnis, J. E., Shelton, B. R., Matsunaga, A., Sioutas, C., and Schauer, J. J.: Source apportionments of PM_{2.5} organic carbon using molecular marker Positive

Matrix Factorization and comparison of results from different receptor models, 2013. *Atmos. Environ.*, 73, 51-61, <http://dx.doi.org/10.1016/j.atmosenv.2013.03.004>.

Hettiyadura, A. P. S., Xu, L., Jayarathne, T., Skog, K., Guo, H., Weber, R. J., Nenes, A., Keutsch, F. N., Ng, N. L., and Stone, E. A.: Source apportionment of organic carbon in Centreville, AL using organosulfates in organic tracer-based positive matrix factorization, 2018. *Atmos. Environ.*, 186, 74-88, <https://doi.org/10.1016/j.atmosenv.2018.05.007>.

Holzinger, R., Williams, J., Herrmann, F., Lelieveld, J., Donahue, N. M., and Röckmann, T.: Aerosol analysis using a Thermal-Desorption Proton-Transfer-Reaction Mass Spectrometer (TD-PTR-MS): a new approach to study processing of organic aerosols, 2010. *Atmos. Chem. Phys.*, 10, 2257-2267, 10.5194/acp-10-2257-2010.

Hu, D., Bian, Q., Lau, A. K. H., and Yu, J. Z.: Source apportioning of primary and secondary organic carbon in summer PM_{2.5} in Hong Kong using positive matrix factorization of secondary and primary organic tracer data, 2010. *J. Geophys. Res.-Atmos.*, 115, 10.1029/2009JD012498.

Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad, I. E., and Prevot, A. S. H.: High secondary aerosol contribution to particulate pollution during haze events in China, 2014. *Nature*, 514, 218-222, 10.1038/nature13774

<http://www.nature.com/nature/journal/v514/n7521/abs/nature13774.html#supplementary-information>.

Iinuma, Y., Boge, O., Grafe, R., and Herrmann, H.: Methyl-nitrocatechols: atmospheric tracer compounds for biomass burning secondary organic aerosols, 2010. *Environ. Sci. Technol.*, 44, 8453-8459, 10.1021/es102938a.

Jiang, N., Wang, K., Yu, X., Su, F., Yin, S., Li, Q., and Zhang, R.: Chemical Characteristics and Source Apportionment by Two Receptor Models of Size-segregated Aerosols in an Emerging Megacity in China, 2018. *Aerosol. Air Qual. Res.*, 18, 1375-1390, 10.4209/aaqr.2017.10.0413.

Jimenez, J. L., Jayne, J. T., Shi, Q., Kolb, C. E., Worsnop, D. R., Yourshaw, I., Seinfeld, J. H., Flagan, R. C., Zhang, X., and Smith, K. A.: Ambient aerosol sampling using the aerodyne aerosol mass spectrometer, 2003. *J. Geophys. Res.-Atmos.*, 108.

Keerthi, R., Selvaraju, N., Alen Varghese, L., and Anu, N.: Source apportionment studies for particulates (PM10) in Kozhikode, South Western India using a combined receptor model, 2018. *Chemistry and Ecology*, 34, 797-817, 10.1080/02757540.2018.1508460.

Kim, E., Hopke, P. K., and Edgerton, E. S.: Improving source identification of Atlanta aerosol using temperature resolved carbon fractions in positive matrix factorization, 2004. *Atmos. Environ.*, 38, 3349-3362, <https://doi.org/10.1016/j.atmosenv.2004.03.012>.

Kleindienst, T. E., Jaoui, M., Lewandowski, M., Offenberg, J. H., Lewis, C. W., Bhave, P. V., and Edney, E. O.: Estimates of the contributions of biogenic and anthropogenic hydrocarbons to secondary organic aerosol at a southeastern US location, 2007. *Atmos. Environ.*, 41, 8288-8300, <http://dx.doi.org/10.1016/j.atmosenv.2007.06.045>.

Kleindienst, T. E., Lewandowski, M., Offenberg, J. H., Edney, E. O., Jaoui, M., Zheng, M., Ding, X., and Edgerton, E. S.: Contribution of Primary and Secondary Sources to Organic Aerosol and PM2.5 at SEARCH Network Sites, 2010. *J. Air Waste Manage. Assoc.*, 60, 1388-1399, 10.3155/1047-3289.60.11.1388.

Languille, B., Gros, V., Petit, J.-E., Honoré, C., Baudic, A., Perrussel, O., Foret, G., Michoud, V., Truong, F., Bonnaire, N., Sarda-Estève, R., Delmotte, M., Feron, A., Maisonneuve, F., Gaimoz, C., Formenti, P., Kotthaus, S., Haeffelin, M., and Favez, O.: Wood burning: A major source of Volatile Organic Compounds during wintertime in the Paris region, 2020. *Sci. Total Environ.*, 711, 135055, <https://doi.org/10.1016/j.scitotenv.2019.135055>.

Lanz, V. A., Alfarrá, M. R., Baltensperger, U., Buchmann, B., Hueglin, C., and Prévôt, A. S. H.: Source apportionment of submicron organic aerosols at an urban site by factor analytical modelling of aerosol mass spectra, 2007. *Atmos. Chem. Phys.*, 7, 1503-1522, 10.5194/acp-7-1503-2007.

Lanzafame, G. M., Srivastava, D., Favez, O., Bonnaire, V., Alleman, L. Y., Couvidat, F., Bessagnet, B., and Albinet, A.: One-year measurements of secondary organic aerosol (SOA) markers in the Paris region (France): concentrations, gas/particle partitioning and SOA source apportionment, 2020. *Sci. Total Environ.*, Accepted for publication.

Lim, Y. B., and Ziemann, P. J.: Products and mechanism of secondary organic aerosol formation from reactions of n-alkanes with OH radicals in the presence of NO_x, 2005. *Environ. Sci. Technol.*, 39, 9229-9236.

Lopez-Hilfiker, F. D., Mohr, C., Ehn, M., Rubach, F., Kleist, E., Wildt, J., Mentel, T. F., Lutz, A., Hallquist, M., Worsnop, D., and Thornton, J. A.: A novel method for online analysis of gas and particle composition: description and evaluation of a Filter Inlet for Gases and AEROSols (FIGAERO), 2014. *Atmos. Meas. Tech.*, 7, 983-1001, 10.5194/amt-7-983-2014.

Lopez-Hilfiker, F. D., Pospisilova, V., Huang, W., Kalberer, M., Mohr, C., Stefenelli, G., Thornton, J. A., Baltensperger, U., Prevot, A. S. H., and Slowik, J. G.: An extractive electrospray ionization time-of-flight mass spectrometer (EESI-TOF) for online measurement of atmospheric aerosol particles, 2019. *Atmos. Meas. Tech.*, 12, 4867-4886, 10.5194/amt-12-4867-2019.

Mader, B. T., and Pankow, J. F.: Gas/solid partitioning of semivolatile organic compounds (SOCs) to air filters. 3. An analysis of gas adsorption artifacts in measurements of atmospheric SOC and organic carbon (OC) when using Teflon membrane filters and quartz fiber filters, 2001. *Environ. Sci. Technol.*, 35, 3422-3432.

Matawle, J. L., Pervez, S., Deb, M. K., Shrivastava, A., and Tiwari, S.: PM_{2.5} pollution from household solid fuel burning practices in Central India: 2. Application of receptor models for source apportionment, 2018. *Environ. Geochem. Health*, 40, 145-161, [10.1007/s10653-016-9889-y](https://doi.org/10.1007/s10653-016-9889-y).

Mohr, C., Huffman, J. A., Cubison, M. J., Aiken, A. C., Docherty, K. S., Kimmel, J. R., Ulbrich, I. M., Hannigan, M., and Jimenez, J. L.: Characterization of Primary Organic Aerosol Emissions from Meat Cooking, Trash Burning, and Motor Vehicles with High-Resolution Aerosol Mass Spectrometry and Comparison with Ambient and Chamber Observations, 2009. *Environ. Sci. Technol.*, 43, 2443-2449, [10.1021/es8011518](https://doi.org/10.1021/es8011518).

Paatero, P., and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error estimates of data values, 1994. *Environmetrics*, 5, 111-126.

Paatero, P.: Least squares formulation of robust non-negative factor analysis, 1997. *Chemom. Intell. Lab. Syst.*, 37, 23-35, [http://doi.org/10.1016/S0169-7439\(96\)00044-5](http://doi.org/10.1016/S0169-7439(96)00044-5).

Pachon, J. E., Balachandran, S., Hu, Y., Weber, R. J., Mulholland, J. A., and Russell, A. G.: Comparison of SOC estimates and uncertainties from aerosol chemical composition and gas phase data in Atlanta, 2010. *Atmos. Environ.*, 44, 3907-3914, <http://dx.doi.org/10.1016/j.atmosenv.2010.07.017>.

Petit, J.-E., Amodeo, T., Meleux, F., Bessagnet, B., Menut, L., Grenier, D., Pellan, Y., Ockler, A., Rocq, B., and Gros, V.: Characterising an intense PM pollution episode in March 2015 in France

from multi-site approach and near real time data: Climatology, variabilities, geographical origins and model evaluation, 2017. *Atmos. Environ.*, 155, 68-84.

Petit, J. E., Favez, O., Sciare, J., Canonaco, F., Croteau, P., Močnik, G., Jayne, J., Worsnop, D., and Leoz-Garziandia, E.: Submicron aerosol source apportionment of wintertime pollution in Paris, France by double positive matrix factorization (PMF2) using an aerosol chemical speciation monitor (ACSM) and a multi-wavelength Aethalometer, 2014. *Atmos. Chem. Phys.*, 14, 13773-13787, 10.5194/acp-14-13773-2014.

Petit, J. E., Favez, O., Sciare, J., Crenn, V., Sarda-Estève, R., Bonnaire, N., Močnik, G., Dupont, J. C., Haeffelin, M., and Leoz-Garziandia, E.: Two years of near real-time chemical composition of submicron aerosols in the region of Paris using an Aerosol Chemical Speciation Monitor (ACSM) and a multi-wavelength Aethalometer, 2015. *Atmos. Chem. Phys.*, 15, 2985-3005, 10.5194/acp-15-2985-2015.

Pieber, S. M., El Haddad, I., Slowik, J. G., Canagaratna, M. R., Jayne, J. T., Platt, S. M., Bozzetti, C., Daellenbach, K. R., Fröhlich, R., Vlachou, A., Klein, F., Dommen, J., Miljevic, B., Jiménez, J. L., Worsnop, D. R., Baltensperger, U., and Prévôt, A. S. H.: Inorganic Salt Interference on CO₂+ in Aerodyne AMS and ACSM Organic Aerosol Composition Studies, 2016. *Environ. Sci. Technol.*, 50, 10494-10503, 10.1021/acs.est.6b01035.

Putaud, J.-P., Raes, F., Van Dingenen, R., Brüggemann, E., Facchini, M. C., Decesari, S., Fuzzi, S., Gehrig, R., Hüglin, C., Laj, P., Lorbeer, G., Maenhaut, W., Mihalopoulos, N., Müller, K., Querol, X., Rodriguez, S., Schneider, J., Spindler, G., Brink, H. t., Tørseth, K., and Wiedensohler, A.: A European aerosol phenomenology—2: chemical characteristics of particulate matter at kerbside, urban, rural and background sites in Europe, 2004. *Atmos. Environ.*, 38, 2579-2595, <https://doi.org/10.1016/j.atmosenv.2004.01.041>.

Putaud, J. P., Van Dingenen, R., Alastuey, A., Bauer, H., Birmili, W., Cyrys, J., Flentje, H., Fuzzi, S., Gehrig, R., Hansson, H. C., Harrison, R. M., Herrmann, H., Hitzenberger, R., Hüglin, C., Jones, A. M., Kasper-Giebl, A., Kiss, G., Koussa, A., Kuhlbusch, T. A. J., Löschau, G., Maenhaut, W., Molnar, A., Moreno, T., Pekkanen, J., Perrino, C., Pitz, M., Puxbaum, H., Querol, X., Rodriguez, S., Salma, I., Schwarz, J., Smolik, J., Schneider, J., Spindler, G., ten Brink, H., Tursic, J., Viana, M., Wiedensohler, A., and Raes, F.: A European aerosol phenomenology – 3: Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe, 2010. *Atmos. Environ.*, 44, 1308-1320, <https://doi.org/10.1016/j.atmosenv.2009.12.011>.

Puxbaum, H., Caseiro, A., Sánchez-Ochoa, A., Kasper-Giebl, A., Claeys, M., Gelencsér, A., Legrand, M., Preunkert, S., and Pio, C.: Levoglucosan levels at background sites in Europe for assessing the impact of biomass combustion on the European aerosol background, 2007. *J. Geophys. Res.-Atmos.*, 112, n/a-n/a, 10.1029/2006JD008114.

Qi, L., Chen, M., Stefenelli, G., Pospisilova, V., Tong, Y., Bertrand, A., Hueglin, C., Ge, X., Baltensperger, U., Prévôt, A. S. H., and Slowik, J. G.: Organic aerosol source apportionment in Zurich using an extractive electrospray ionization time-of-flight mass spectrometer (EESI-TOF-MS) – Part 2: Biomass burning influences in winter, 2019. *Atmos. Chem. Phys.*, 19, 8037-8062, 10.5194/acp-19-8037-2019.

Qi, L., Vogel, A. L., Esmaeilrad, S., Cao, L., Zheng, J., Jaffrezo, J. L., Fermo, P., Kasper-Giebl, A., Daellenbach, K. R., Chen, M., Ge, X., Baltensperger, U., Prévôt, A. S. H., and Slowik, J. G.: One-year characterization of organic aerosol composition and sources using an extractive electrospray ionization time-of-flight mass spectrometer (EESI-TOF), 2020. *Atmos. Chem. Phys. Discuss.*, 2020, 1-34, 10.5194/acp-2019-1165.

Ringuet, J., Albinet, A., Leoz-Garziandia, E., Budzinski, H., and Villenave, E.: Diurnal/nocturnal concentrations and sources of particulate-bound PAHs, OPAHs and NPAHs at traffic and suburban sites in the region of Paris (France), 2012. *Sci. Total Environ.*, 437, 297-305, <http://dx.doi.org/10.1016/j.scitotenv.2012.07.072>.

Riva, M., Tomaz, S., Cui, T., Lin, Y. H., Perraudin, E., Gold, A., Stone, E. A., Villenave, E., and Surratt, J. D.: Evidence for an unrecognized secondary anthropogenic source of organosulfates and sulfonates: gas-phase oxidation of polycyclic aromatic hydrocarbons in the presence of sulfate aerosol, 2015. *Environ. Sci. Technol.*, 49, 6654-6664, [10.1021/acs.est.5b00836](https://doi.org/10.1021/acs.est.5b00836).

Rutter, A. P., Snyder, D. C., Stone, E. A., Shelton, B., DeMinter, J., and Schauer, J. J.: Preliminary assessment of the anthropogenic and biogenic contributions to secondary organic aerosols at two industrial cities in the upper Midwest, 2014. *Atmos. Environ.*, 84, 307-313, [10.1016/j.atmosenv.2013.11.014](https://doi.org/10.1016/j.atmosenv.2013.11.014).

Saylor, R. D., Edgerton, E. S., and Hartsell, B. E.: Linear regression techniques for use in the EC tracer method of secondary organic aerosol estimation, 2006. *Atmos. Environ.*, 40, 7546-7556, <http://dx.doi.org/10.1016/j.atmosenv.2006.07.018>.

Shi, G., Liu, J., Wang, H., Tian, Y., Wen, J., Shi, X., Feng, Y., Ivey, C. E., and Russell, A. G.: Source apportionment for fine particulate matter in a Chinese city using an improved gas-constrained method and comparison with multiple receptor models, 2018. *Environ. Pollut.*, 233, 1058-1067, <https://doi.org/10.1016/j.envpol.2017.10.007>.

Shirmohammadi, F., Hasheminassab, S., Saffari, A., Schauer, J. J., Delfino, R. J., and Sioutas, C.: Fine and ultrafine particulate organic carbon in the Los Angeles basin: Trends in sources and composition, 2016. *Sci. Total Environ.*, 541, 1083-1096, <http://dx.doi.org/10.1016/j.scitotenv.2015.09.133>.

Song, Y., Zhang, Y., Xie, S., Zeng, L., Zheng, M., Salmon, L. G., Shao, M., and Slanina, S.: Source apportionment of PM_{2.5} in Beijing by positive matrix factorization, 2006. *Atmos. Environ.*, 40, 1526-1537, 10.1016/j.atmosenv.2005.10.039.

Srivastava, D., Favez, O., Bonnaire, N., Lucarelli, F., Perraudin, E., Gros, V., Villenave, E., and Albinet, A.: Speciation of organic fractions does matter for aerosol source apportionment. Part 2: intensive campaign in the Paris area (France), 2018a. *Sci. Total Environ.*

Srivastava, D., Favez, O., Perraudin, E., Villenave, E., and Albinet, A.: Comparison of Measurement-Based Methodologies to Apportion Secondary Organic Carbon (SOC) in PM_{2.5}: A Review of Recent Studies, 2018b. *Atmosphere*, 9, 452.

Srivastava, D., Tomaz, S., Favez, O., Lanzafame, G. M., Golly, B., Besombes, J. L., Alleman, L. Y., Jaffrezo, J. L., Jacob, V., Perraudin, E., Villenave, E., and Albinet, A.: Speciation of organic fraction does matter for source apportionment. Part 1: A one-year campaign in Grenoble (France), 2018c. *Sci. Total Environ.*, 624, 1598-1611, 10.1016/j.scitotenv.2017.12.135.

Srivastava, D., Favez, O., Petit, J. E., Zhang, Y., Sofowote, U. M., Hopke, P. K., Bonnaire, N., Perraudin, E., Gros, V., Villenave, E., and Albinet, A.: Speciation of organic fractions does matter for aerosol source apportionment. Part 3: Combining off-line and on-line measurements, 2019. *Sci. Total Environ.*, 690, 944-955, <https://doi.org/10.1016/j.scitotenv.2019.06.378>.

Stefenelli, G., Pospisilova, V., Lopez-Hilfiker, F. D., Daellenbach, K. R., Hüglin, C., Tong, Y., Baltensperger, U., Prévôt, A. S. H., and Slowik, J. G.: Organic aerosol source apportionment in Zurich using an extractive electrospray ionization time-of-flight mass spectrometer (EESI-TOF-MS) – Part 1: Biogenic influences and day–night chemistry in summer, 2019. *Atmos. Chem. Phys.*, 19, 14825-14848, 10.5194/acp-19-14825-2019.

Surratt, J. D., Kroll, J. H., Kleindienst, T. E., Edney, E. O., Claeys, M., Sorooshian, A., Ng, N. L., Offenberg, J. H., Lewandowski, M., Jaoui, M., Flagan, R. C., and Seinfeld, J. H.: Evidence for Organosulfates in Secondary Organic Aerosol, 2006. *Environ. Sci. Technol.*, 41, 517-527, 10.1021/es062081q.

Szidat, S., Ruff, M., Perron, N., Wacker, L., Synal, H.-A., Hallquist, M., Shannigrahi, A. S., Yttri, K. E., Dye, C., and Simpson, D.: Fossil and non-fossil sources of organic carbon (OC) and elemental carbon (EC) in Göteborg, Sweden, 2009. *Atmos. Chem. Phys.*, 9, 1521-1535.

Tarrasón, L., P.D. Hamer, C. Guerreiro, F. Meleux, L. Rouil Interim Annual Assessment Report for 2015, 2016. CAMS report CAMS71:2016SC2011_D2071.2011.2011.2012_201609_final http://policy.atmosphere.copernicus.eu/reports/CAMS201671_202016SC201601_D201671.201601.201601.201602_201609_final.pdf.

Tian, M., Liu, Y., Yang, F., Zhang, L., Peng, C., Chen, Y., Shi, G., Wang, H., Luo, B., Jiang, C., Li, B., Takeda, N., and Koizumi, K.: Increasing importance of nitrate formation for heavy aerosol pollution in two megacities in Sichuan Basin, southwest China, 2019. *Environ. Pollut.*, 250, 898-905, <https://doi.org/10.1016/j.envpol.2019.04.098>.

Tkacik, D. S., Presto, A. A., Donahue, N. M., and Robinson, A. L.: Secondary Organic Aerosol Formation from Intermediate-Volatility Organic Compounds: Cyclic, Linear, and Branched Alkanes, 2012. *Environ. Sci. Technol.*, 46, 8773-8781, 10.1021/es301112c.

Tomaz, S., Shahpoury, P., Jaffrezo, J. L., Lammel, G., Perraudin, E., Villenave, E., and Albinet, A.: One-year study of polycyclic aromatic compounds at an urban site in Grenoble (France): Seasonal variations, gas/particle partitioning and cancer risk estimation, 2016. *Sci. Total Environ.*, 565, 1071-1083, 10.1016/j.scitotenv.2016.05.137.

Tomaz, S., Jaffrezo, J.-L., Favez, O., Perraudin, E., Villenave, E., and Albinet, A.: Sources and atmospheric chemistry of oxy- and nitro-PAHs in the ambient air of Grenoble (France), 2017. *Atmos. Environ.*, 161, 144-154, <https://doi.org/10.1016/j.atmosenv.2017.04.042>.

Turpin, B. J., and Huntzicker, J. J.: Identification of secondary organic aerosol episodes and quantitation of primary and secondary organic aerosol concentrations during SCAQS, 1995. *Atmos. Environ.*, 29, 3527-3544, [http://dx.doi.org/10.1016/1352-2310\(94\)00276-Q](http://dx.doi.org/10.1016/1352-2310(94)00276-Q).

Turpin, B. J., Saxena, P., and Andrews, E.: Measuring and simulating particulate organics in the atmosphere: problems and prospects, 2000. *Atmos. Environ.*, 34, 2983-3013, [http://dx.doi.org/10.1016/S1352-2310\(99\)00501-4](http://dx.doi.org/10.1016/S1352-2310(99)00501-4).

Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from Positive Matrix Factorization of aerosol mass spectrometric data, 2009. *Atmos. Chem. Phys.*, 9, 2891-2918, 10.5194/acp-9-2891-2009.

van Drooge, B. L., and Grimalt, J. O.: Particle sized-resolved source apportionment of primary and secondary organic tracer compounds at urban and rural locations in Spain, 2015. *Atmospheric Chemistry and Physics Discussions*, 15, 9897-9939, 10.5194/acpd-15-9897-2015.

Verlhac, S., Favez, O., and Albinet, A.: Comparaison inter laboratoires organisée pour les laboratoires européens impliqués dans l'analyse du lévoglucosan et de ses isomères 2013. LCSQA / INERIS <http://www.lcsqa.org/rapport/2013/ineris/comparaison-inter-laboratoires-organisee-laboratoires-europeens-impliques-analys>.

Vlachou, A., Daellenbach, K. R., Bozzetti, C., Chazeau, B., Salazar, G. A., Szidat, S., Jaffrezo, J. L., Hueglin, C., Baltensperger, U., Haddad, I. E., and Prévôt, A. S. H.: Advanced source apportionment of carbonaceous aerosols by coupling offline AMS and radiocarbon size-segregated

measurements over a nearly 2-year period, 2018. *Atmos. Chem. Phys.*, 18, 6187-6206, <https://doi.org/10.5194/acp-18-6187-2018>.

Waked, A., Favez, O., Alleman, L. Y., Piot, C., Petit, J. E., Delaunay, T., Verlinden, E., Golly, B., Besombes, J. L., Jaffrezo, J. L., and Leoz-Garziandia, E.: Source apportionment of PM₁₀ in a north-western Europe regional urban background site (Lens, France) using positive matrix factorization and including primary biogenic emissions, 2014. *Atmos. Chem. Phys.*, 14, 3325-3346, 10.5194/acp-14-3325-2014.

Wang, Y., Chen, Y., Wu, Z., Shang, D., Bian, Y., Du, Z., Schmitt, S. H., Su, R., Gkatzelis, G. I., Schlag, P., Hohaus, T., Voliotis, A., Lu, K., Zeng, L., Zhao, C., Alfarra, M. R., McFiggans, G., Wiedensohler, A., Kiendler-Scharr, A., Zhang, Y., and Hu, M.: Mutual promotion between aerosol particle liquid water and particulate nitrate enhancement leads to severe nitrate-dominated particulate matter pollution and low visibility, 2020. *Atmos. Chem. Phys.*, 20, 2161-2175, 10.5194/acp-20-2161-2020.

Watson, J. G., Robinson, N. F., Chow, J. C., Henry, R. C., Kim, B., Pace, T., Meyer, E. L., and Nguyen, Q.: The USEPA/DRI chemical mass balance receptor model, CMB 7.0, 1990. *Environmental Software*, 5, 38-49.

Weber, S., Salameh, D., Albinet, A., Alleman, L. Y., Waked, A., Besombes, J.-L., Jacob, V., Guillaud, G., Meshbah, B., Rocq, B., Hulin, A., Dominik-Sègue, M., Chrétien, E., Jaffrezo, J.-L., and Favez, O.: Comparison of PM₁₀ Sources Profiles at 15 French Sites Using a Harmonized Constrained Positive Matrix Factorization Approach, 2019. 10, 310.

White, G. H.: Basics of estimating measurement uncertainty, 2008. *The Clinical biochemist. Reviews*, 29 Suppl 1, S53-S60.

Williams, B. J., Jayne, J. T., Lambe, A. T., Hohaus, T., Kimmel, J. R., Sueper, D., Brooks, W., Williams, L. R., Trimborn, A. M., Martinez, R. E., Hayes, P. L., Jimenez, J. L., Kreisberg, N. M., Hering, S. V., Worton, D. R., Goldstein, A. H., and Worsnop, D. R.: The First Combined Thermal Desorption Aerosol Gas Chromatograph—Aerosol Mass Spectrometer (TAG-AMS), 2014. *Aerosol Sci. Technol.*, 48, 358-370, 10.1080/02786826.2013.875114.

Xie, Y., Wang, G., Wang, X., Chen, J., Chen, Y., Tang, G., Wang, L., Ge, S., Xue, G., Wang, Y., and Gao, J.: Nitrate-dominated PM_{2.5} and elevation of particle pH observed in urban Beijing during the winter of 2017, 2020. *Atmos. Chem. Phys.*, 20, 5019-5033, 10.5194/acp-20-5019-2020.

Xu, J., Zhang, Q., Chen, M., Ge, X., Ren, J., and Qin, D.: Chemical composition, sources, and processes of urban aerosols during summertime in northwest China: insights from high-resolution aerosol mass spectrometry, 2014. *Atmos. Chem. Phys.*, 14, 12593-12611, 10.5194/acp-14-12593-2014.

Xu, Q., Wang, S., Jiang, J., Bhattarai, N., Li, X., Chang, X., Qiu, X., Zheng, M., Hua, Y., and Hao, J.: Nitrate dominates the chemical composition of PM_{2.5} during haze event in Beijing, China, 2019. *Sci. Total Environ.*, 689, 1293-1303, <https://doi.org/10.1016/j.scitotenv.2019.06.294>.

Yttri, K. E., Schnelle-Kreis, J., Maenhaut, W., Abbaszade, G., Alves, C., Bjerke, A., Bonnier, N., Bossi, R., Claeys, M., Dye, C., Evtuyugina, M., García-Gacio, D., Hillamo, R., Hoffer, A., Hyder, M., Iinuma, Y., Jaffrezo, J. L., Kasper-Giebl, A., Kiss, G., López-Mahia, P. L., Pio, C., Piot, C., Ramirez-Santa-Cruz, C., Sciare, J., Teinilä, K., Vermeulen, R., Vicente, A., and Zimmermann, R.: An intercomparison study of analytical methods used for quantification of levoglucosan in ambient aerosol filter samples, 2015. *Atmos. Meas. Tech.*, 8, 125-147, 10.5194/amt-8-125-2015.

Zhang, Q., Alfarra, M. R., Worsnop, D. R., Allan, J. D., Coe, H., Canagaratna, M. R., and Jimenez, J. L.: Deconvolution and quantification of hydrocarbon-like and oxygenated organic aerosols based on aerosol mass spectrometry, 2005a. *Environ. Sci. Technol.*, 39, 4938-4952.

Zhang, Q., Canagaratna, M. R., Jayne, J. T., Worsnop, D. R., and Jimenez, J.-L.: Time- and size-resolved chemical composition of submicron particles in Pittsburgh: Implications for aerosol sources and processes, 2005b. *J. Geophys. Res.-Atmos.*, 110, 10.1029/2004jd004649.

Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, 2007. *Geophys. Res. Lett.*, 34, n/a-n/a, 10.1029/2007gl029979.

Zhang, Y., Sheesley, R. J., Schauer, J. J., Lewandowski, M., Jaoui, M., Offenberg, J. H., Kleindienst, T. E., and Edney, E. O.: Source apportionment of primary and secondary organic aerosols using positive matrix factorization (PMF) of molecular markers, 2009. *Atmos. Environ.*, 43, 5567-5574.

Zhang, Y., Favez, O., Petit, J. E., Canonaco, F., Truong, F., Bonnaire, N., Crenn, V., Amodeo, T., Prévôt, A. S. H., Sciare, J., Gros, V., and Albinet, A.: Six-year source apportionment of submicron organic aerosols from near-continuous highly time-resolved measurements at SIRTa (Paris area, France), 2019. *Atmos. Chem. Phys.*, 19, 14755-14776, 10.5194/acp-19-14755-2019.

Zhao, Y., Hennigan, C. J., May, A. A., Tkacik, D. S., de Gouw, J. A., Gilman, J. B., Kuster, W. C., Borbon, A., and Robinson, A. L.: Intermediate-Volatility Organic Compounds: A Large Source of Secondary Organic Aerosol, 2014. *Environ. Sci. Technol.*, 48, 13743-13750, 10.1021/es5035188.

Table 1. List of POA and SOA factors retrieved from each source apportionment methodology.

Method	Size fraction	Temporal resolution	Primary factors		Secondary factors	
			Number	Labelling	Number	Labelling
EC-tracer method	PM ₁₀	4 h	1	POA	1	SOA
SOA-tracer method	PM ₁₀	4 h		Not applicable	5	Isoprene SOA, α -Pinene SOA, Toluene SOA, Naphthalene SOA, Biomass burning (phenolic compounds) SOA
PMF-chemical data	PM ₁₀	4 h	3	Primary traffic emissions; Biomass burning; Dust	7	Biogenic SOA-1; Biogenic SOA-2 Anthropogenic SOA-1; Anthropogenic SOA-2; Anthropogenic SOA-3; Mixed secondary; Nitrate-rich
PMF-offline AMS	PM ₁₀	4 h	4	Traffic POA (hydrocarbon-like OA (HOA), tailpipe/engine); Cooking POA (COA); Biomass burning POA (BBOA); Sulfur-containing OA (SCOA, coarse POA)	2	Oxidized oxygenated OA 1 and 2 (OOA1; OOA2)
PMF-ACSM	PM ₁	0.5 h	2	Traffic POA (HOA, tailpipe/engine); Biomass burning POA (BBOA)	2	More oxidized oxygenated OA (MO-OOA); Low oxidized oxygenated OA (LO-OOA)

Fig. 1. Left: Comparison of the reconstructed PM₁₀ mass from chemical characterization and the PM₁₀ measured using TEOM-FDMS. Right: Relative contribution of the chemical species to PM₁₀ mass and temporal variations of the measured PM₁₀, PM₁ (NR-PM₁ + BC) at Paris-SIRTA, France (March 2015). Sea salt and dust were calculated applying the procedure explained by Bressi et al. (2014).





Fig. 2. Contribution of primary and secondary factors to the total PM₁₀ OC mass (in the case of PMF-ACSM PM₁), retrieved from the different source apportionment approaches. Red border represents the total secondary fractions. Note, colours used for the PMF-chemical and PMF-ACSM pie-charts are the same as already published (Srivastava et al., 2018a; Srivastava et al., 2019).




Fig. 3. Comparison of total POC concentrations obtained from the different source apportionment methodologies. 3a: Time series. 3b: Box-plots show the minimum, first quartile, median, third quartile and maximum values for PM₁₀ OA. 3c: Box-plots for the fine OA fraction with POC*_{PMF-chemical data}: PMF-chemical data without dust (primary traffic emissions + biomass burning); POC*_{PMF-offline AMS}: PMF-offline AMS without SCOC (= hydrocarbon-like OC (HOC) + biomass burning OC (BBOC) + cooking OC (COC)); POC_{PMF-ACSM} = HOC+BBOC. Grey square symbols: POC median overall uncertainties $\pm 2\sigma$.






Fig. 4. Time series of biomass burning- (top panel) and traffic-related (bottom panel) POC factors obtained from the three PMF approaches.




Fig. 5. Diel profiles of median values obtained for the traffic- and biomass burning-related OC factors using the different source apportionment methodologies (local time).




Fig. 6. Comparison of total SOC concentrations obtained from the different methodologies. 4a: Time series. 4b: Box-plots showing the minimum, first quartile, median, third quartile and maximum values. Grey square: error median $\pm 2\sigma$.





Fig. 7. Correlation matrix within individual OA factors resolved using all approaches. 1. EC-tracer method: 1. Total SOC; 2. PMF-offline AMS: 2. OOA1, 2. OOA2 and 2. Total SOC; 3. SOA-tracer: 3. Biomass burning (SOC biomass burning), 3. Isoprene (SOC isoprene), 3. Naphthalene (SOC naphthalene), 3. Toluene (SOC toluene) and 3. Total SOC; 4. PMF-chemical data: 4. BSOA-1 (Biogenic SOA (marine)), 4. BSOA-2 (Biogenic SOA (isoprene)), 4. ASOA-1 (Anthropogenic SOA (oxy-PAHs)), 4. ASOA-2 (Anthropogenic SOA (nitro-PAHs)), 4. ASOA-3 (Anthropogenic SOA (phenolic oxidation)), 4. Mixed secondary, 4Nitrate-rich and 4. Total SOC; 5. PMF-ACSM: 5. MO-OOA, 5. LO-OOA and 5. Total SOC.

- 1. EC-tracer
- 2. PMF-offline AMS
- 3. SOA-tracer
- 4. PMF-chemical data
- 5. PMF-ACSM




Fig. 8. Comparison of temporal variation of highly oxidized OA factors obtained from PMF-offline AMS (OOC1) and PMF-ACSM (MO-OOC), the sum of the mixed secondary aerosol and marine biogenic SOA factors obtained from PMF-chemical data and SOCnaphthalene from SOA-tracer method.

