



HAL
open science

New insights on activity-related bone functional adaptations and alterations in Neolithic Liguria (northwestern Italy)

A. Varalli, Sébastien Villotte, I. Dori, V.S. Sparacello

► **To cite this version:**

A. Varalli, Sébastien Villotte, I. Dori, V.S. Sparacello. New insights on activity-related bone functional adaptations and alterations in Neolithic Liguria (northwestern Italy). *Bulletins et Mémoires de la Société d'anthropologie de Paris*, 2020, <10.3166/bmsap-2020-0072>. <hal-03007251>

HAL Id: hal-03007251

<https://hal.science/hal-03007251v1>

Submitted on 16 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 **Article, reçu le 14/11/2019, accepté le 18/02/2020. 10 figures (Fig. 1, 6, 7, 8, 9**
2 **and 10 une colonne et demi, Fig. 2, 3, 4 and 5 deux colonnes), 7 tableaux, 2**
3 **annexes**

4
5 **Title: New insights into activity-related functional bone adaptations and alterations in**
6 **Neolithic Liguria (northwestern Italy)**

7 **Titre : Nouvelles connaissances sur les adaptations et altérations fonctionnelles osseuses**
8 **liées à l'activité en Ligurie néolithique (nord-ouest de l'Italie)**

9 Authors: Varalli A^{1,2}, Villotte S¹, Dori I^{1,3,4}, Sparacello VS^{1§}.

10 ¹ Univ. Bordeaux, CNRS, MC, PACEA, UMR 5199, 33615 Pessac, France.

11 ² CaSEs Research Group, Department of Humanities, Universitat Pompeu Fabra, c/Triass Fargas
12 25–27, Barcelona 08005, Spain.

13 ³ SABAP Soprintendenza Archeologia, Belle Arti e Paesaggio per le province di Verona, Rovigo
14 e Vicenza, Piazza S. Fermo 3, 37121 Verona, Italy.

15 ⁴ Department of Biology, Laboratory of Anthropology, University of Florence, via del
16 Proconsolo 12, 50122 Florence, Italy.

17 § Corresponding Author: Vitale Stefano Sparacello PACEA - UMR 5199 Université de
18 Bordeaux Bâtiment B8 Allée Geoffroy Saint Hilaire CS 50023 33615 PESSAC
19 CEDEX. vitale.sparacello@u-bordeaux.fr; ritosparacello@gmail.com

20 **Abstract**

21 This study offers a combined analysis of long bone mechanical properties (cross-sectional
22 geometry, CSG), upper limb enthesopathies (ECs) and external auditory exostoses (EAE) among
23 Neolithic people from Liguria (Italy). Previous CSG studies have suggested a high degree of
24 mobility in mountainous terrain and sexual dimorphism in the upper limbs, with males more
25 oriented towards uni-manual activities and females performing strenuous bimanual tasks. The
26 aims of the study were to: 1) increase the sample size of the CSG study via the acquisition of

27 surface 3D models; 2) provide a solid chronological framework through direct dating, to allow
28 for subsampling of individuals dated to the Impresso-Cardial Complex (ICC, c. 5800-5000 BCE)
29 and the Square Mouthed Pottery Culture (SMP, c. 5000-4300 BCE); 3) integrate the results of
30 CSG with information on ECs of the humeral epicondyles; 4) assess possible marine activities
31 through analysis of EAEs. Results from the CSG analysis confirm previous studies, with no
32 significant diachronic change. ECs in the humeral medial epicondyle parallel CSG adaptations:
33 males tend to display more changes, especially unilaterally. Only one individual from the ICC
34 period shows bilateral EAE, suggesting that marine activities were not prevalent. This study adds
35 to our knowledge on activity patterns in the Neolithic in Liguria, and shows that integrating
36 structural adaptations with information from specific enthesal alterations and exostoses can
37 improve reconstructions of past habitual activities.

38 **Abstract French**

39 Nous avons analysé conjointement les propriétés mécaniques des os longs (géométrie des
40 sections transverses, CSG), les enthésopathies des membres supérieurs (ECs) et les exostoses
41 auditives externes (EAE) chez les sujets néolithiques de Ligurie. Des études antérieures des CSG
42 ont suggéré une grande mobilité en terrain montagneux et un dimorphisme sexuel important pour
43 les membres supérieurs, les hommes étant davantage orientés vers des activités uni-manuelles et
44 les femmes effectuant préférentiellement des tâches bimanuelles pénibles. Nous avons 1)
45 augmenté la taille de l'échantillon de l'étude des CSG par l'acquisition de modèles 3D de surface
46 ; 2) fourni un cadre chronologique solide grâce à de nouvelles datations radiocarbones AMS
47 directes, permettant un sous-échantillonnage d'individus datés du complexe Impresso-Cardial
48 (ICC, c. 5800-5000 BC) et de la culture des Vases à Bouche carrée céramique à bec carré (SMP,
49 c. 5000-4300 BC) ; 3) intégré les résultats sur les CSG avec des informations sur les ECs des
50 épicondyles de l'humérus; 4) évalué les activités marines possibles par l'analyse des EAE. Les
51 résultats de l'analyse des CSG confirment les études antérieures, sans changement diachronique
52 significatif. Les résultats pour les ECs dans l'épicondyle médial sont similaires à ceux pour les
53 CSG du membre supérieur : les hommes ont tendance à avoir plus de changements, surtout
54 unilatéralement. Un seul individu (daté de l'ICC) présente une EAE bilatérale, ce qui suggère des
55 activités marines peu répandues. En plus d'enrichir nos connaissances sur les schémas d'activité
56 dans le Néolithique ligurien, cette étude montre que l'intégration des adaptations structurelles

57 avec les informations sur les ECs et les EAE peut améliorer les reconstructions des activités
58 habituelles dans le passé.

59 **Keywords:** habitual activities, subsistence patterns, cross-sectional geometry, enthesal changes,
60 enthesopathies, external auditory meatus exostosis

61 **Mots-clés :** Activités quotidiennes ; modes de subsistance ; géométrie des sections transverses
62 des diaphyses ; changements enthésiques ; enthésopathies ; exostoses du conduit auditif externe.

63

64 **1. Introduction**

65 The evaluation of plastic, epigenetic activity-related morphological adaptations (structural bone
66 properties) and alterations (enthesal changes) in skeletal remains from archaeological contexts
67 provides insights into sociocultural phenomena such as subsistence changes, craft specialization,
68 the origins and development of sexual division of labour and social inequalities [1]. Among these
69 sociocultural phenomena, the Neolithic Transition, i.e. the adoption of a production economy
70 based on the domestication of plants and animals, is “one of the fundamental structural processes
71 of human history” [2,3], and dramatically changed several aspects of the human experience,
72 particularly habitual subsistence-related physical activity. This subsistence shift has been the
73 focus of several studies aiming to reconstruct activity patterns, which generally show a decrease
74 in mobility among farmers when compared to earlier hunter-gatherers (reviews in [4-7]).

75 However, the Neolithisation of Europe was probably a mosaic of different processes of
76 replacement and integration, frontiers and “leap-frog” movements, growth and collapse,
77 coexistence and warfare (e.g. [8-13]). Therefore, the skeletal consequences of the adoption of the
78 Neolithic lifestyle are best explored at a micro-regional level [14]. Few regions in Europe show
79 such a concentration of prehistoric sites from the Upper Palaeolithic to the Metal Ages as
80 western Liguria (north-western Italy): Neolithic funerary sites are especially numerous and
81 concentrated mostly around the municipality of Finale Ligure, where hundreds of karstic caves
82 can be found within a radius of a few kilometres. After the early arrival of the Neolithic
83 Impresso-Cardial Complex (ICC; c. 5800-5000 cal BC; [15-17]), Liguria saw the spread of the
84 Square Mouth Pottery (SMP; 5000-4300 cal BC; [16,18-20]), and later was the access road for
85 the diffusion of the Chassean in northern Italy from France (4300-3700 cal BC; [18,21]). This

86 small strip of land between the mountains and the sea is therefore a pivotal region for our
87 understanding of the cultural and biological dynamics that came into play during the spread and
88 establishment of the Neolithic way of life in the western Mediterranean (e.g. [15,16, 22-25]).

89 Consequently, the functional adaptations of Ligurian Neolithic people have been the subject of
90 several studies, especially via biomechanical analysis of long bones using the cross-sectional
91 geometry method (CSG; review in [26,27]). For the lower limb, one of the main results of
92 previous studies has been the discovery that the “sedentism” of Neolithic communities did not
93 actually result in a low level of mobility, at least as inferred from lower limb biomechanics:
94 Ligurian Neolithic individuals retained CSG properties that were more similar to those shown by
95 Late Upper Palaeolithic hunters than by later agriculturalists, suggesting a high degree of
96 mobility in a mountainous landscape [28,29,31,32,33]. These studies also showed evidence of
97 significant sexual dimorphism in the degree of asymmetry of the upper limb, with males more
98 asymmetrical, and females more symmetrical, than in modern industrialized groups and other
99 agriculturalists [28,29,32,33]. This diverging pattern has been interpreted as indicative of a
100 division of subsistence tasks between the sexes, in which the most strenuous and repetitive
101 activities performed were mostly bi-manual for females, and mostly uni-manual for males [28].
102 This has been tentatively associated with subsistence activities that are apparent in the
103 archaeological record, the most prominent of which may have been cereal processing using large
104 bi-manual querns ([33,34,35]) and woodworking and fodder harvesting with polished stone
105 hatchets [28,33].

106 The Ligurian skeletal series was ideal to study human micro-regional adaptations, thanks to the
107 small area from which it was unearthed. However, its chronological framework is more
108 problematic. Most Neolithic layers had been excavated from the mid-19th century [36], in caves
109 that, in some cases, showed evidence of human occupation from the Late Pleistocene to Roman
110 times (e.g. [37,38]). Following the archaeological methods of the time [39], burials, and
111 especially scattered human remains, were often excavated without keeping an accurate record of
112 their spatial and stratigraphic positions. Probably due to the poor quality of the historic
113 documentation, few burials were considered worthy of direct dating, and the individuals were
114 labelled as likely/probably/possibly “Neolithic” or “Middle Neolithic” (which corresponds to the
115 SMP culture in Liguria) depending on the material available about the associated culture. This

116 limited the number of individuals that could be included in previous bioarchaeological studies,
117 and the lack of a more precise chrono-cultural attribution has constantly prevented
118 anthropologists from providing more than general insights into Ligurian “Neolithic” biocultural
119 adaptations.

120 To begin to overcome the limitations of the skeletal series and its documentation, two research
121 projects (see Acknowledgements section) performed a comprehensive re-assessment of the
122 Ligurian material, including individual burials, partially disturbed burials and assemblages of
123 scattered human remains. Depending on the completeness of the material, the study collected the
124 information available on the biological profile (sex, age, pathology, osteometric and non-metric
125 traits, dental anthropology), as well as funerary information (e.g. [39]). In addition, the study
126 undertook complete direct dating of the Ligurian skeletal collection attributed to the Neolithic (c.
127 180 AMS dates), adding to the results from other recent studies [23,24,39,40]. The results of the
128 dating campaign are reported in detail in another paper [41], but they indicate that the skeletal
129 remains generally attributed to the Neolithic in Liguria span the period from the earliest
130 Neolithic human occupation in the area through to the sixth and fifth millennium BCE, and in
131 some cases belong to the Metal Ages or to historic times. For example, the double burial from
132 Boragni, previously attributed to the Neolithic [42] and used in previous CSG research
133 [30,33,43], actually dates to the Byzantine Period (1460±30 BP; 553-648 CE; Lyon-14599). The
134 vast majority of the burials and skeletal remains fall chronologically within the period in which
135 the Square Mouthed Pottery culture (SMP) has been attested in Liguria (c. 5000-4300 cal BCE;
136 Table 1), with a small but significant presence of individuals belonging to an earlier phase (c.
137 5800-5000 BC, [39]). Thanks to direct dating, this new study was able to include individuals
138 (complete and partial skeletons) that are certainly Neolithic, and to create subsamples based on
139 the various chrono-cultural phases of the Neolithic in Liguria.

140 Furthermore, the new chronological framework makes it possible to assess whether the
141 functional adaptations and behavioural correlates inferred by previous studies still hold when
142 using a better defined and homogenous Neolithic skeletal collection from the SMP, and whether
143 any diachronic change can be detected during the Neolithic. For the lower limb, the hypothesis
144 of high degrees of mobility has important implications in terms of Neolithic adaptive strategies,
145 logistic mobility and energy expenditure [7]; for the upper limb, sexual dimorphism and the

146 hypothesized division of labour tap into issues of societal organization and identity, in addition
147 to land use and energy expenditure to process food [33,34,35]. Confirmation is therefore needed
148 that these results were not influenced by the inclusion in the sample of individuals from earlier
149 pioneering Neolithic communities, or from the Metal Ages, for which technological/societal
150 changes and a greater reliance on pastoralism have been proposed [43].

151 In addition, the project collected data on certain enthesal changes (ECs) that appear to correlate
152 with habitual physical activity and are therefore ideal for integration with previous studies.
153 Enthesal changes (ECs) are visible alterations on the skeleton at sites of tendon or ligament
154 attachment, both fibrous (tendon or ligament attaching directly to the bone) and
155 fibrocartilaginous (tendon or ligament attaching through transitional zones of a different
156 composition) [44,45]. ECs have been associated with activity patterns in bioarchaeological
157 studies, but their reliability as indicators of past behaviour is debated. It has been shown that a
158 significant part of this issue is related to the methods used and the nature of the entheses under
159 study [44,46-48]. Several factors have been mentioned by biological anthropologists to explain
160 the occurrence of ECs at fibrocartilaginous sites, but three main causes can be identified from the
161 medical literature (see [44] for a discussion): age (enthesopathies are more frequent in older
162 individuals, especially after the age of 50), micro and macrotraumas, and systemic diseases
163 (mainly DISH and spondyloarthropathies). Among the ECs that appear to correlate best with
164 habitual activities, the one involving the medial epicondyle of the humerus could be related to
165 strenuous uni-manual activities, especially those associated with the throwing motion [49-52]:
166 ECs indeed appear to be frequent among prehistoric groups for which hunting involving
167 throwing techniques has been hypothesized [51,52]. Given the divergent pattern of asymmetry
168 by sex shown in the Neolithic Ligurian people described above, the study of ECs of the humeral
169 epicondyle could provide further insights into the behavioural correlates of these upper limb
170 adaptations and alterations.

171 Another bone alteration that may be relevant to draw inferences on behavioural correlates are the
172 external auditory exostoses (EAEs), which are osseous exostoses that form in the external
173 auditory canal as a result of an irritation of the periosteum. Several conditions can be responsible
174 for this trait, but contact with cold water appears to be the main cause, and their prevalence and
175 degree of expression has therefore been linked to the use of aquatic resources (for a review of the

176 medical and anthropological literature, see [53,54]). EAE can be considered as one of the most
177 informative of activity-related skeletal morphologies: it has a very well-known etiology, the
178 amount of available clinical data is substantial, and it is possible to compare frequencies between
179 current and past populations, with only minor methodological problems [54]. Although
180 archaeological and faunal evidence suggest continuous use of marine resources throughout the
181 Neolithic, a particular emphasis on terrestrial resources and on the use of the mountainous
182 territory has been proposed for Liguria, based on dietary [55,56], zooarchaeological [57] and
183 biomechanical evidence (see above). A low prevalence of EAEs would therefore be expected
184 among Ligurian Neolithic people.

185 The overall purpose of the study was therefore to 1) perform a new CSG analysis with the
186 benefit of the new chronological framework, and test whether past behavioural inferences still
187 hold; 2) integrate the results from ECs, verifying whether these alterations support or contrast
188 with structural bone adaptations; 3) propose new considerations on land *vs* marine oriented
189 subsistence activities through the analysis of EAEs.

190 **2. Materials and Methods**

191 A total of 57 skeletons from eight nearby Ligurian sites (Finalese and Val Pennavaire; Figure 1)
192 was included in this study, although the sample size is smaller depending on the specific analysis
193 and on the completeness of the remains (Table 1, see also the Excel file named Supplementary
194 Information Tables A). Because EAEs can develop during adolescence [54], individuals from c.
195 12 years old upwards were included in the study of this trait, while only individuals with fused
196 long bone epiphyses were included in the CSG and ECs study. The osteometric measurements of
197 two additional Ligurian Neolithic individuals were taken from the literature [42] to determine
198 their sex via Discriminant Function Analysis (see below). Most individuals are directly dated to
199 either the sixth or fifth millennium BCE; for the few individuals where dating failed or could not
200 be done, the new chronological framework allows for a more robust chrono-cultural attribution
201 based on contextual archaeological evidence, such as the presence of the stone cist for the Square
202 Mouthed Pottery burials (Table 1; [41]).

203 **[Figure 1 (1.5 column) and Table 1 about here]**

204 The attribution of an individual to the “adult” age class was based on the completeness of
205 epiphyseal fusion and dental maturation, integrated with dental wear [58]. The appearance of the
206 pubic symphysis and auricular surface of the ilium was also considered [59-61]. Age at death of
207 immature individuals was based, when possible, on dental maturation and eruption [62-64], and
208 skeletal fusion [65-67], integrated when necessary with bone measurements [65,68].

209 The biological sex of the adult individuals in this study was re-assessed for all individuals via
210 cranial and pelvic morphology, using the standards collected in Buikstra et al. [69], mandibular
211 ramus flexure [70], and Bruzek [71] for the pelvic traits. Compared to previous studies, sex
212 determination was changed for only one individual (Pollera 6246) , from male to female (cf.
213 Table 1 with [32]). For 13 individuals, the fragmentary nature of the remains did not allow
214 attribution based on pelvic and/or cranial features, and sex was assessed via Discriminant
215 Function Analysis (DFA) using the postcranial osteometric measurements of the individuals
216 whose sex was determined from pelvic traits (e.g. [72-75]). Forward and backward stepwise
217 selection of variables was undertaken for each individual where the sex was uncertain, and the
218 preferred model was chosen based on discriminant power, significance, number of individuals
219 used for the equation and number of variables included in the model. Due to the remarkable level
220 of sexual dimorphism displayed by the Ligurian sample, the DFA produced models with a
221 classification rate ranging from 90-100%, and the posterior probabilities for attribution to a given
222 sex was >87% for the 13 uncertain individuals. All the osteometric measurements and equations
223 are given in Supplementary Information Tables A.

224 Biomechanical analysis of functional postcranial adaptations via the cross-sectional geometry
225 method (CSG) is based on the notion that bone tissue optimizes to its mechanical environment so
226 as to maintain physiological strains within normal limits [26,27]. Bone tissue is deposited in the
227 shaft’s cross-section where mechanical loads require it to prevent strain in excess of the elastic
228 limit, whereas below a certain strain threshold, the bone tissue is reabsorbed. By analyzing the
229 cross-sections of the diaphysis, it is therefore possible to obtain variables that correlate with
230 torsional bone rigidity (polar moment of area: J). Although the complexity of the factors
231 influencing mechanical bone competence should always be taken into account when interpreting
232 CSG results [26], it is generally assumed that variations in CSG properties correlate with activity
233 levels and types, once the effect of body size is factored out to obtain a measure of “robusticity”

234 [27]. Recent experimental evidence on modern athletes has substantially confirmed this rationale
235 [76,77,78]. Integration of quantitative data derived from CSG with ethnographic and
236 archaeological information has therefore been widely used to draw inferences about past
237 subsistence strategies, degrees of mobility and other habitual activities [6,7,79].

238 For this comprehensive re-assessment of the skeletal series, all of the preserved long bones were
239 scanned in 3D, using the DAVID SLS-3 structured light scanner, to make a new assessment of
240 structural diaphyseal properties. CSG properties for 20 individuals were added in this study (cf.
241 Table 1 with [28,29,30,32,33,43]), and cross-sections were reconstructed (35% and 50% from the
242 distal end for the humeri, and 50% for the femur and tibia) from the surface scans, which were
243 positioned virtually according to the reference planes following Ruff [80]. The cross-sections
244 were obtained using the “slice” function in Netfabb Standard 2018 for PC (copyright Autodesk
245 2017), and the CSG properties calculated using a version of the SLICE program [81] adapted as
246 a macro routine inserted in Scion Image release Beta 4.03. The “Solid CSG” method was used to
247 estimate actual CSG properties from the periosteal contour via regression equations (provided in
248 [43,82]), as justified in previous research [82-84]. All raw CSG data are given in Supplementary
249 Information Tables A.

250 The variable used to quantify overall mechanical bone strength at a given diaphyseal level is the
251 modulus Z_p section (torsional and (twice) average bending strength [85,86]). True section moduli
252 are calculated by dividing the polar second moment of area J (torsional and (twice) average
253 bending rigidity of the beam) by the distance from the centroid section for torsion to the
254 outermost fibre of the section [86]. Until recently, software packages did not provide true section
255 moduli, which were therefore approximated by dividing J by the average radius of the section or
256 by raising J to the power of 0.73 [81,85,86]. In this study, we use the latter method, which makes
257 the results comparable to a larger body of literature, including previous studies on the same
258 skeletal series. Although $J^{0.73}$ is proportional to rather than strictly equivalent to Z_p , given the
259 substantial equivalency of the variables, this study will refer to $J^{0.73}$ as Z_p , as done in previous
260 research (e.g., [33]).

261 The mechanical loading on long bones is a function of physical activity, bone length and body
262 mass [86]. To obtain a measure of “robusticity”, which is assumed to correlate with the effects of
263 activity, the estimate of overall bone strength Z_p was scaled for size by dividing by mechanical

264 bone length (as defined in Ruff [80]) and body mass [86]. Body mass was estimated from the
265 supero-inferior diameter of the femoral head following the guidelines in Trinkaus et al. [87].
266 Certain osteometric measurements for the determination of the appropriate level of the cross-
267 section and for standardization had to be estimated via regression equations based on the rest of
268 the sample, as commonly done in studies of this kind (e.g. [87,88]). The regression equations for
269 each estimated measurement are provided in Supplementary Information Tables A.

270 In order to characterize the prevalent use of one arm in activities causing strain, the degree of
271 humeral bilateral asymmetry in J was calculated using the formula $[(J_H - J_L)/J_L] \times 100$ (where J_H
272 and J_L are the higher and lower values of J between the two humeri, respectively) and expressed
273 as a percentage, following previous studies [32,33,89,90]. The resulting value represents an
274 absolute (non-directional) asymmetry. Asymmetry was calculated from the absolute values of J
275 (i.e. not standardized by body size), because any prior size standardization would be elided.

276 Given their correspondence with mobility levels [76,78,91], lower limb CSG shape indices were
277 also analyzed. For the femur, the ratio between I_x (second moment of area in the anteroposterior
278 plane) and I_y (second moment of area in the mediolateral plane) was considered. For the tibia, the
279 ratio of I_{max} (maximum second moment of area) to I_{min} , (minimum second moment of area) was
280 used.

281 Enthesal changes (EC) in the humeral medial and lateral epicondyles were recorded. Due to the
282 very small sample size, age-at-death is not controlled for in this study. Therefore, ECs at the
283 lateral epicondyle were scored in order to provide a basis for comparison for the EC frequencies
284 and distribution at the medial epicondyle. Recently, it has been found increasingly relevant to
285 distinguish two areas for the medial epicondyle [92,93]. Thus, unlike in previous studies (i.e.
286 [48,51,52]), the area of the medial epicondyle was divided into two zones: the area of attachment
287 of the common flexor tendon (CFT) and the area of attachment of the anterior band of the ulnar
288 collateral ligament (UCL). ECs at the lateral epicondyle and the area of attachment of the CFT in
289 the medial epicondyle were scored on a 3-stage scale (A, B, C) following Villotte [46]. Several
290 changes at the attachment of the UCL were recorded (as absent or present): erosions and
291 cavitations of the surface, and geodes, were scored as ECG; longitudinal fissures and well-
292 defined pits with no clear interruption of the bone surface were scored as FP (see [92]:24; Figure
293 2). One of these features was enough to consider a change as present at the UCL attachment site.

294 A composite variable for the whole medial epicondyle (ME) was then created, merging the data
295 for the UCL and CFT attachment sites. EC at the ME was considered as present if any change at
296 the attachment sites of the CFT (stage B or C) and/or at the UCL was recorded. All raw data for
297 EC are given in Supplementary Information Tables A. Asymmetry in ECs was assessed as the
298 difference in scores between left and right sites for a given variable in a given individual.
299 Differences between the sexes in terms of the number of asymmetrical individuals were assessed
300 using Fisher's exact test. Differences between left and right sides within each sex in the
301 presence/absence or aggregated score of a certain trait were also assessed using Fisher's exact
302 test, although the independence assumption does not hold for paired humeri.

303 **[Figure 2 (2 columns) about here]**

304 EAEs were investigated through visual inspection of the auditory meatus under appropriate
305 lighting, and recorded using a scoring system of the extent of occlusion previously applied by
306 biological anthropologists (e.g. [94-97] and by clinicians (e.g. [98,99]). All raw data for EAEs
307 are given in Supplementary Information Tables A.

308 The comparative samples for CSG properties consist of European Late Upper Palaeolithic (c.
309 20,000-10,000 BCE) and Italian Iron Age (Orientalizing-Archaic Period, c. 800-500 BCE,
310 Abruzzo region) individuals collected during previous research [31,33,79,88,100,101].

311 **3. Results**

312 *3.1. Cross-sectional geometry*

313 Figures 3 and 4 show the main upper and lower limb CSG properties of the Ligurian sample,
314 divided by sex and by chronological phases (c. 5800-5000 BCE: Impresso-Cardial Complex,
315 ICC; c. 5000-4300 BCE: Square Mouthed Pottery, SMP). Note that the only individual dated to
316 the Chassean (c. 4300-3700 BCE), for which only the left humerus could be analyzed, was not
317 included in the figures. Raw data by individual are given in Supplementary Information.

318 Although the sample size of the ICC individuals does not allow for a clear recognition of
319 diachronic trends, their values are within the range of variability of SMP individuals for all the
320 variables, in both the upper (Figure 3) and lower limbs (Figure 4).

321 **[Figures 3 and 4 (both 2 columns) about here]**

322 Figure 5 and Table 2 show the comparison between upper limb CSG properties of SMP males
323 and females when compared to Late Upper Palaeolithic hunter-gatherers and Iron Age agro-
324 pastoralists. A diachronic trend of increasing robusticity in both humeri can be observed in
325 males, although the SMP sample is not significantly different to LUP and IRON (Table 2).
326 Humeral mid-distal bilateral asymmetry in SMP males is significantly lower than in LUP, but
327 not significantly different to that in IRON males. Conversely, SMP females have the lowest
328 asymmetry, although the comparison with IRON females is not significant at the $\alpha = 0.05$ level
329 after correcting for multiple comparisons (Table 2). Sexual dimorphism in the SMP sample is
330 significant for right humeral robusticity and humeral bilateral asymmetry, with males showing
331 higher values than females.

332 **[Figure 5 (2 columns) and Table 2 about here]**

333 Figures 6 and 7 and Table 2 show the comparison between lower limb CSG properties of SMP
334 males and females when compared to Late Upper Palaeolithic hunter-gatherers and Iron Age
335 agro-pastoralists (for this study the SMP individual Bergeggi 2 was not included, due to lower
336 limb bending deformities probably due to rickets). No significant difference in femoral
337 robusticity is present among the diachronic samples, while SMP females have higher tibial
338 robusticity than IRON females (Figure 6, 7 and Table 2). A trend of temporally declining shape
339 indices is observable in both the femur (Figure 6) and tibia (Figure 7), the decrease being
340 particularly marked between SMP and IRON individuals especially. Both sexes in the SMP
341 sample have significantly higher femoral shape indices than their IRON counterparts after
342 correcting for multiple comparisons (Table 2). Sexual dimorphism in the SMP sample is
343 significant for the shape indices of both the femur and tibia, the males showing higher values
344 than females (Table 2).

345 **[Figures 6 and 7 (both 1.5 column) about here]**

346 *3.2 Humeral enthesal changes*

347 Table 3 shows the results of the evaluation of enthesal changes in the medial and lateral
348 epicondyle of the humerus. For the lateral epicondyle, in both sides and in both sexes, ECs are
349 frequent (ca. 50% of the lateral epicondyles present stage B or C). No significant difference in
350 the pattern of stage frequency could be detected, whether comparing sides within or between

351 sexes. When considering individuals that could be examined bilaterally, the stage of ECs in the
352 lateral epicondyle generally shows symmetry between sides (Table 3).

353 [Table 3 about here]

354 Results for the medial epicondyle differ from the lateral epicondyle in many ways. When
355 considering the area of attachment of the CFT, no individual in the SMP or larger Ligurian
356 Neolithic sample shows stage C. Stage B appears to be more represented in the right CFT area in
357 males, but no variation by side or by sex is statistically significant. However, when considering
358 individuals that could be examined bilaterally, males have consistently higher scores in the right
359 CFT area when compared to females, which are more symmetrical. This result is significant at
360 the $\alpha = 0.05$ level after pooling the SMP and ICC individuals (Table 4).

361 [Table 4 about here]

362 Changes at the UCL attachment site are present in both sexes, but especially on the right side in
363 males. Males have more lesions than females on the right side, but not on the left. Asymmetrical
364 individuals are significantly more frequent in males than in females (Table 5)

365 [Table 5 about here]

366 When the CFT and UCL scores are aggregated, obtaining the “any change” variable for the
367 medial epicondyle (ME any change), the pattern is similar to that described above: males show
368 more changes especially on the right side, and asymmetrical individuals are more frequent in
369 males than in females (Tables 6 and 7). However, the results are no longer statistically
370 significant, probably due to the smaller sample size, when considering only individuals for which
371 both the CFT attachment site and the anterior band of the UCL could be scored (Table 6). When
372 aggregating the scores for individuals to which at least one score could be attributed, the results
373 are statistically significant in the SMP sample (Table 7).

374 [Tables 6 and 7 about here]

375 No significant difference in humeral robusticity (both mid-distal and midshaft Z_p) or humeral
376 bilateral asymmetry (mid-distal and midshaft HUMBA), as described by CSG data, could be
377 detected between the subsamples based on sex and the medial/lateral epicondyle stage of
378 enthesal change (Supplementary Information Tables B). However, SMP individuals (pooled

379 sexes) with asymmetrical CFT stages of enthesal change tend to have more robust right and left
380 humeri as described by mid-distal Z_p (T-test $p < 0.05$; Supplementary Information Table B-2).
381 Furthermore, in both in the pooled Ligurian Neolithic sample and the SMP sample, individuals
382 with changes at the UCL (pooled sexes) show more robust right and left humeri as described by
383 mid-distal Z_p (T-test $p < 0.05$), and more asymmetrical humeri as described by the mid-distal
384 bilateral asymmetry in Z_p (T-test $p < 0.05$; Supplementary Information Tables B-3 and B-4).
385 Although the significance of these results is mostly due to the observed sexual dimorphism in
386 robusticity and asymmetry (Table 2) and in the higher prevalence of changes in the UCL in
387 males (Table 5), it may be noticed that the same pattern is present by sex, although the results are
388 not statistically significant (Figure 8).

389 Furthermore, a correspondence between changes in the UCL (ECG or ECG+FP) and humeral
390 robusticity, asymmetry and laterality can be inferred from the categorized scatterplots (Figure 9).
391 More robust and asymmetrical individuals will tend to have changes in the UCL, and the only
392 individual with ECG on the left humerus is also the most asymmetrical left-dominant individual
393 (Figure 9a). When considering the aggregate score for the medial epicondyle (ME any change)
394 the correspondence between enthesal changes and CSG robusticity and laterality is less
395 apparent, when considering only individuals for which both CFT and UCL could be scored
396 (Figure 9c, and see above), and when including individuals to which only one score could be
397 attributed (Figure 9d).

398 **[Figures 8 and 9 (both 1.5 column) about here]**

399 *3.3 external auditory canal exostosis (EAE)*

400 Among the 30 individuals that could be observed for EAE (Supplementary Information Table
401 A), only one male individual dating to the sixth millennium BCE (Arma dell'Aquila 1 Richard)
402 displays slight bilateral EAEs (Figure 10).

403 **[Figure 10 (1.5 column) about here]**

404 **4. Discussion**

405 This study aimed to reassess functional adaptations in the Ligurian Neolithic skeletal series,
406 based on the new chronological framework recently provided by direct dating [39,41]. In

407 addition to verifying whether the inferences on subsistence-related activity patterns made in the
408 past using CSG (e.g. [28,30]) still hold, this study added new observations on certain enthesal
409 changes that appear to correlate with habitual physical activity, located in the distal humerus
410 [46,92] and in the external auditory meatus [54,97].

411 The direct dating results confirm that most of the skeletal series from Liguria overlap
412 chronologically with the Square Mouthed Pottery culture (SMP), c. 5000-4300 BCE, i.e. the
413 “Middle Neolithic” of Liguria [41]. With the new chronological framework, it became possible
414 to attribute several individuals to this period, for which direct dating had failed or which could
415 not be sampled, based on funerary treatment (e.g. Arene Candide 6PE, Arene Candide II BB,
416 Pollera 32PE; [41]). On the other hand, certain individuals included in previous studies could no
417 longer be assigned to the Neolithic (e.g. Boragni), or were assigned to a cultural tradition earlier
418 than the SMP within the Impresso-Cardial Complex (ICC; e.g. most of the individuals from
419 Arma dell’ Aquila; [39]). Despite the exclusion of these individuals from the SMP sample, the
420 Ligurian “Middle Neolithic” sample size for CSG analysis is still slightly larger than in previous
421 studies, thanks to the inclusion of CSG data from 20 additional individuals that were excluded
422 from previous analyses due to the lack of chrono-cultural attribution.

423 The characterization of functional adaptations of Ligurian SMP people does not change
424 significantly based on this revised sample, which confirms the validity of previous observations
425 and inferences on subsistence patterns [28-30,32,33]. The small size of the ICC sample does not
426 allow for inferences on diachronic trends; however, most ICC individuals do not seem to deviate
427 from the variability of the SMP sample. Lower limb CSG properties are still compatible with
428 high levels of mobility in rugged terrain [30,31], showing values that are closer to Late
429 Palaeolithic hunter-gatherers than later protohistorical agro-pastoralists. This further highlights
430 how subsistence activities, particularly logistic mobility, among early agro-pastoralists in this
431 region may have differed from later, more specialized forms of pastoralism [33]. However, more
432 studies on controlled athlete *vs* sedentary samples are necessary to fully understand how different
433 types of mobility influence femoral and tibial robusticity and lead to specific cross-sectional
434 shapes (cf. our results with [76,78]).

435 The degree of asymmetry in humeral mid-distal mechanical rigidity of SMP males, around 20%,
436 is lower than that shown by LUP hunters, whose subsistence activities probably included a

437 substantial amount of throwing [88], and later Iron Age agro-pastoralists, who performed uni-
438 manual sword training [33,79,100]. However, SMP male asymmetry is still relevant when
439 compared to “physiological” asymmetry (8-12%; [77,89]), and especially when compared to
440 SMP females, which show the lowest asymmetry among the comparative groups. The three
441 females in the ICC sample show very low levels of mid-distal humeral asymmetry (between 2-
442 4%), suggesting that this trait was shared among Neolithic females in Liguria (less relevant for
443 behavioural inferences is their relatively higher asymmetry at midshaft of the humerus, which is
444 often influenced by the shape of the deltoid tuberosity [79]). Previous studies have suggested a
445 possible causal relationship between the diverging pattern of asymmetry in males and females
446 and the division of labour (see Introduction). However, caution is needed when inferring division
447 of labour based on sexual dimorphism in structural properties, given the possible influence of
448 hormonal factors and body size (34,88). In this research, the focus was on determining whether
449 CSG and ECs in the humerus produced compatible results.

450 Both CSG and ECs have been widely used in the past to draw inferences about activity patterns,
451 although many studies have warned against simplistic interpretations of results given the number
452 of concomitant factors that influence mechanical bone competence, enthesal appearance and
453 other “markers of activities” (e.g. [26,102,103]). In particular, age and body size appear to be
454 confounding factors for both methods. For CSG, mechanical properties seem to mostly reflect
455 levels of activity during the pre- and peri-pubertal periods, while the extent of remodelling due to
456 activity in later life is debated [26,27,104]. In addition, senescence does influence CSG
457 properties, due to continued periosteal apposition to mechanically compensate for medullary
458 expansion (especially in males; [105,106]). These limitations are presumed to have a relatively
459 minor impact on the inferences about habitual activity and subsistence drawn from
460 bioarchaeological samples, since it is assumed that individuals in prehistoric societies
461 participated in subsistence activities as from late childhood, and died before reaching senescence.
462 Hormonal and genetic factors may nevertheless confound behavioural interpretations [34,107].
463 In contrast, for ECs, the whole correlation with activity patterns is brought into question once
464 age and body size are taken into account (e.g. [108-110]). Studies integrating CSG and ECs
465 results for bioarchaeological reconstructions of activity patterns have at best found a general
466 correspondence between the two methods (e.g. [111]), and more detailed tests between ECs
467 scores and CSG values have failed to provide a consistent association [110,112-114]. This study

468 has found similar results: individuals with higher scores in medial and lateral humeral epicondyle
469 ECs are not consistently more robust. However, rather than expecting a general correspondence
470 between EC scores and CSG robusticity, it is possible that specific enthesal changes, and
471 especially their asymmetry, may be indicative of certain habitual activities.

472 The shape of the humeral medial epicondyle appears to have an influence on the pronation-
473 supination range during elbow flexion and extension [115], and ECs in this area have been
474 associated with trauma resulting from powerful extensions of the forearm, possibly related to
475 throwing [49,50-52]. Prehistoric groups, especially Upper Palaeolithic hunters, often appear to
476 show lesions in this area [51,52]. Among Upper Palaeolithic hunters, throwing has also been
477 considered to be one of the main factors determining the remarkable degree of asymmetry in
478 humeral mechanical rigidity, which is comparable or greater than that shown in professional
479 tennis players or, indeed, throwers (e.g. [88]). A relationship between ECs in the medial
480 epicondyle and CSG properties, particularly asymmetry, was therefore expected. Accordingly, in
481 this study there is agreement between the inferences that could be drawn based on CSG and
482 medial epicondyle ECs: males tend to be more asymmetrical than females, and tend to have
483 lesions on the right side more frequently. In addition, although the sample size is small, there
484 appears to be, within sexes, a correspondence between the presence of lesions in the dominant
485 arm, in the anterior band of the UCL, and the level of humeral asymmetry. This relationship
486 seems to be consistent with the laterality of CSG asymmetry: the only individual with somewhat
487 relevant left-sided asymmetry is also the only individual with an erosion in the left anterior band
488 of the UCL. However, this possible correspondence between biomechanical properties and ECs
489 becomes less discernible when aggregating other enthesal changes in the medial epicondyle,
490 further suggesting that, among ECs, erosions in the UCL may be particularly relevant for
491 reconstructing activity. The very small sample size, however, makes these considerations
492 speculative. Further research using a large sample with known activity is necessary to confirm
493 the results found here.

494 In the Neolithic Ligurian sample, only one case of bilateral EAE (grade 1) has been recorded, in
495 an individual from the ICC, Arma dell'Aquila 1. It is difficult to discuss frequencies of EAEs in
496 a small sample [54]: although the total sample of Neolithic crania is 30, only three other
497 individuals from the ICC period could be examined for EAE, and only unilaterally

498 (Supplementary Information Table A). More archaeological and biochemical data on Neolithic
499 Ligurian people are needed to understand whether this result could be compatible with a greater
500 focus on marine resource use in the ICC Neolithic. No significant consumption of fish protein
501 during the ICC and SMP periods has been proposed in light of the collagen isotopic composition
502 of human bone [40,55,56]. However, archaeological and faunal evidence (e.g. shells, fish bones,
503 fishing hooks, shell blow horns) are evidence of the continuous use of marine resources
504 throughout the Neolithic [116]. In addition, EAE may be due to causes other than frequent
505 contact with cold water, including infection, eczema, trauma and other pathological conditions
506 affecting the normal homeostasis of the external ear canal [117,118]. Finally, the absence of
507 EAE does not exclude aquatic activities. However, it should be noted that in populations for
508 which isotopic data has indicated significant reliance on aquatic resources, EAEs tend to be very
509 frequent and the occlusion of the ear canal is sometimes severe (i.e.; grades 2 or 3)
510 [54,94,97,119], which is not the case here. The results, although caution is necessary given the
511 small sample size, seem compatible with a subsistence scenario where marine resource use was
512 of relatively minor importance.

513 **5. Conclusions**

514 This study performed a re-assessment of long bone structural properties in the Neolithic Ligurian
515 skeletal series, in the light of the new chronological framework produced by direct dating and
516 new data derived by 3D scanning of the entire assemblage of long bones. As well as considering
517 the plastic adaptation of the diaphyses, this study included an interpretative scenario with new
518 data from skeletal alterations that are considered to be correlated with habitual activities. The
519 analysis included enthesal changes in the humeral epicondyles, which have been associated
520 with vigorous exertion of the upper limb, including the throwing motion, and the presence of
521 exostoses in the external auditory meatus, which are considered to be an indicator of aquatic
522 activities.

523 The biomechanical analysis results confirm previous studies: the data for lower limbs suggest
524 high mobility; the upper limb is robust in both sexes, but males show significantly higher
525 humeral asymmetry than females. No diachronic change was apparent when considering the
526 subsampling based on chrono-cultural complexes within the Neolithic. Enthesal alterations in
527 the humeral medial epicondyle are consistent with the results obtained from structural

528 adaptations: males tend to show more changes than females, especially unilaterally on the more
529 mechanically robust side. In addition, individuals with lesions in the medial epicondyle tend to
530 have higher asymmetry. This result is potentially important because studies on adaptations and
531 alterations have often failed to produce consistent results. However, the sample size is very
532 small, and more research is necessary to confirm this correspondence between results derived
533 from structural adaptations and bone alterations.

534 Only one individual from the sixth millennium BCE shows bilateral external auditory meatus
535 exostosis, supporting the archaeological evidence suggesting that marine activities were not
536 prevalent during the Neolithic in Liguria. However, again, the sample size is too small, and
537 further evidence is necessary to investigate this issue.

538 This study has added to our knowledge on activity patterns in the Neolithic in Liguria, and
539 supports the careful integration of data on structural adaptations with information from specific
540 enthesal alterations and exostoses to draw inferences about past habitual activities.

541

542 **Acknowledgements**

543 The authors would like to thank the Soprintendenza Archeologia, Belle Arti e Paesaggio per la
544 città metropolitana di Genova e le provincie di Imperia, La Spezia e Savona, for granting access
545 to skeletal collections, Superintendent Vincenzo Tiné and Officers Elisabetta Starnini, Marta
546 Conventi, Nico Radi, and Stefano Costa. We are grateful to the directors, curators, and staff of
547 the museums where the skeletal collections are preserved for their continuous assistance during
548 data collection: Daniele Arobba and Andrea De Pascale (Museo Archeologico del Finale, Finale
549 Ligure), Patrizia Garibaldi, Guido Rossi, Irene Molinari (Museo di Archeologia Ligure, Genova),
550 and Monica Zavattaro (Museo di Storia Naturale – Sezione di Antropologia e Etnologia,
551 Università degli Studi di Firenze). Thanks to Camillo Costa and the staff of the Museo Navale
552 Romano di Albenga. Thanks to Chiara Panelli, Stefano Rossi, Roberto Maggi, Vincenzo
553 Formicola, Paolo Biagi, Giovanni Murialdo, Gwenaëlle Goude, Sara Bernardini, Claudia Ferro,
554 Giuseppe “Cisque” Vicino, Maria Tagliafico, Elisa Bianchi, Simona Mordeglia, Walter
555 Siciliano, Jacopo Moggi-Cecchi, Giovanna Stefania, Luca Bachechi, Chiara Bullo and Brunetto
556 Chiarelli for their assistance during data collection and their scientific input.

557 For their continuous assistance during the analysis, we are grateful to all the staff of the
558 University of Groningen Centre for Isotope Research (CIO), to Christine Oberlin, Centre de
559 Datation par le RadioCarbone, Université Claude Bernard Lyon1, and to Lucile Beck,
560 Responsable de la Plateforme Nationale LMC14 Laboratoire de Mesure du Carbone 14
561 (CEA/CNRS/IRD/IRSN/MCC)-LSCE CEA Saclay. The AMS dating performed at the Centre de
562 Datation par le RadioCarbone was supported by the “ARTEMIS” program.

563 The BUR.P.P.H project: Burial practices at the Pleistocene - Holocene transition: the changing
564 role of pathology, violence, and “exceptional events” (PI: VSS) received financial support from
565 the French State under the “Investing for the Future” Program, IdEx Bordeaux, reference ANR-
566 10-IDEX-03-02. The DEN.P.H. project: Dental anthropology at the Pleistocene-Holocene
567 transition – insights on lifestyle and funerary behaviour from Neolithic Liguria (Italy) (PI: ID) is
568 funded by the European Union’s Horizon 2020 research and innovation program under the Marie
569 Skłodowska-Curie grant agreement No 752626.

570

571 **Literature Cited**

- 572 1. Knüsel CJ, Sparacello V (2018) Functional morphology, postcranial, human. In:Trevathan
573 W, Cartmill M, Dufour D, Larsen C, O’Rourke D, Rosenberg K, Strier K (eds) The
574 International Encyclopedia of Biological
575 Anthropologydoi:10.1002/9781118584538.ieba0187
- 576 2. Bocquet-Appel J-P (2011a) When the world’s population took off: the springboard of the
577 Neolithic Demographic Transition. *Science* 333:560–561
- 578 3. Bocquet-Appel J-P (2011b) The agricultural demographic transition during and after the
579 agriculture inventions. *Curr Anthropol* 52:S497–S510
- 580 4. Larsen CS (1995) Biological changes in human populations with agriculture. *Ann Rev*
581 *Anthropol* 24:185–213
- 582 5. Larsen CS (1997) *Bioarchaeology*, Cambridge University Press, Cambridge, 461 p
- 583 6. Pinhasi R, Stock J (eds) (2011) *Human bioarchaeology of the Transition to*
584 *Agriculture*, Wiley-Liss, New York, 488 p
- 585 7. Carlson K, Marchi D (eds) (2014) *Reconstructing mobility: environmental, behavioral, and*
586 *morphological determinants*, Springer, New York, 295 p

- 587 8. Lahr M, Foley R, Pinhasi R (2000) Expected regional patterns of Mesolithic-Neolithic
588 human population admixture in Europe based on archaeological evidence. In: Renfrew C,
589 Boyle K (eds) *Archaeogenetics: DNA and the Population Prehistory of Europe*. McDonald
590 Institute for Archaeological Research, Monographs, Cambridge, pp 81–88
- 591 9. Clare K, Rohling EJ, Weninger B, et al (2008) Warfare in Late Neolithic/Early Chalcolithic
592 Pisidia, southwestern Turkey. Climate induced social unrest in the late 7th millennium cal
593 BC. *Doc Praehist* 25:65–92
- 594 10. Zeder MA (2008) Domestication and early agriculture in the Mediterranean basin: origins,
595 diffusion, and impact. *PNAS* 105(33):11597–11604
- 596 11. Rowley-Conwy P (2011) Westward Ho! The spread of agriculture from Central Europe to
597 the Atlantic. *Curr Anthropol* S52:431–451
- 598 12. Shennan S, Downey SS, Timpson A, et al (2013) Regional population collapse followed
599 initial agriculture booms in mid-Holocene Europe. *Nat Comm* 4:2486
- 600 13. Meyer C, Lohr C, Gronenborn D, et al (2015) The massacre mass grave of Schöneck-
601 Kikianstädten reveals new insights into collective violence in Early Neolithic Central Europe.
602 *PNAS* 112(36):11217–11222
- 603 14. Stock JT, Pinhasi R (2011) Introduction. Changing paradigms in our understanding of the
604 transition to agriculture: human bioarchaeology, behaviour and adaptation. In: Pinhasi R,
605 Stock J (eds) *Human bioarchaeology of the Transition to Agriculture*. Wiley-Liss, New York,
606 pp 1–17
- 607 15. Binder D (2013) *Mésolithique et Néolithique ancien en Méditerranée nord-occidentale entre*
608 *7000 et 5500 cal. BCE: questions ouvertes sur les dynamiques culturelles et les procès*
609 *d’interaction. Actes du XXVIIe Congrès préhistorique de France (Bordeaux–Les Eyzies,*
610 *2010). Société préhistorique française, Paris, pp 341–355*
- 611 16. Binder D, Sénépart I (2010) La séquence de l’Impresso-Cardial de l’abri Pendimoun et
612 l’évolution des assemblages céramiques en Provence. *Mémoire LI de la Société Préhistorique*
613 *française, pp 149–167*
- 614 17. Binder D, Lanos P, Angeli L, et al (2017) Modelling the earliest north-western dispersal of
615 Mediterranean Impressed Wares: new dates and Bayesian chronological model. *Doc Praehist*
616 44:54–77

- 617 18. Maggi R (1997) The Radiocarbon Chronology. In: Maggi R (ed) *Arene Candide: a functional*
618 *and environmental assessment of the Holocene sequence (excavations Bernabò Brea-Cardini*
619 *1940–1950)*. Istituto Italiano di paleontologia umana, Il calamo, Roma, ns 5, pp 31–52
- 620 19. Pearce M (2013) Radiocarbon chronology for the spread of the early Neolithic north through
621 the Tyrrhenian and Ligurian Seas area. In: Pearce M (ed) *Rethinking the North Italian Early*
622 *Neolithic*. Accordia Research Institute, University of London, London, pp 21–84
- 623 20. Del Lucchese A, Starnini E (2015) Aggiornamenti sulla fase antica della Cultura dei Vasi a
624 Bocca Quadrata in Liguria da una revisione dei materiali ceramici in corso. In: Conventi M,
625 Del Lucchese A, Gardini A (eds) *Archeologia in Liguria*, ns5 (2012-2013). Sagep Editrice
626 Genova, pp 27–37
- 627 21. Crepaldi F (2001) Le Chasséen en Ligurie. *Bull Soc Prehistor Fr* 98(3):485–494
- 628 22. Branch NP, Black S, Maggi R, et al (2014) The neolithisation of Liguria (NW Italy): an
629 environmental archaeological and palaeoenvironmental perspective. *Environ Archaeol*
630 19:196–213
- 631 23. Biagi P, Starnini E (2016) La cultura della Ceramica Impressa nella Liguria di Ponente (Italia
632 Settentrionale): distribuzione, cronologia e aspetti culturali. In: Bonet Rosado H (ed) *Del*
633 *neolític a l’edat del bronze en el Mediterrani occidental. Estudis en homenatge a Bernat*
634 *Martí Oliver*. TV SIP 119, València, pp 35-49
- 635 24. Biagi P, Starnini E (2018) Gli scavi all’Arma dell’Aquila (Finale Ligure, Savona): le ricerche
636 e i materiali degli scavi del Novecento. *Quaderno 15*. Società per la Preistoria e Protostoria
637 della Regione Friuli-Venezia Giulia, Trieste, 260 p
- 638 25. Arobba D, Panelli C, Caramiello R, Gabriele M, Maggi R (2017) Cereal remains, plant
639 impressions and 14C direct dating from the Neolithic pottery of Arene Candide Cave (Finale
640 Ligure, NW Italy). *J Archeol Sci: Rep* 12:395–404
- 641 26. Pearson OM, Lieberman DE (2004) The aging of Wolff’s ‘Law’: ontogeny and response to
642 mechanical loading in cortical bone. *Am J Phys Anthropol* 47:63–99
- 643 27. Ruff CB, Holt B, Trinkaus E (2006) Who’s afraid of the big bad Wolff? ‘Wolff’s law’ and
644 bone functional adaptation. *Am J Phys Anthropol* 129:484–498
- 645 28. Marchi D, Sparacello VS, Holt BM, et al (2006) Biomechanical approach to the
646 reconstruction of activity patterns in Neolithic Western Liguria, Italy. *Am J Phys Anthropol*
647 131:447–455

- 648 29. Marchi D (2008) Relationships between lower limb cross-sectional geometry and mobility:
649 the case of a Neolithic sample from Italy. *Am J Phys Anthropol* 137:188–200
- 650 30. Sparacello VS, Marchi D, Shaw CN (2014) The importance of considering fibular robusticity
651 when inferring the mobility patterns of past populations. In: Carlson K, Marchi D (eds)
652 *Reconstructing mobility: environmental, behavioral, and morphological determinants*.
653 Springer, New York, pp 91–111
- 654 31. Sparacello VS, Villotte S, Shaw CN, et al (2018) Changing mobility patterns at the
655 Pleistocene-Holocene transition: the biomechanics of the lower limb of Italian Gravettian and
656 Mesolithic individuals. In: Cristiani E, Borgia V (eds) *Palaeolithic Italy: Advanced studies on*
657 *early human adaptations in the Apennine Peninsula*. Sidestone Press, Leiden, pp 357–396
- 658 32. Sparacello VS, Roberts CA, Canci A, et al (2016) Insights on the paleoepidemiology of
659 ancient tuberculosis from the structural analysis of postcranial remains from the Ligurian
660 Neolithic (northwestern Italy). *Int J Paleopath* 15:50-64.
- 661 33. Sparacello VS, Pearson OM, Coppa A, et al (2011) Changes in robusticity in an Iron Age
662 agropastoral group: the Samnites from the Alfedena necropolis (Abruzzo, Central Italy). *Am*
663 *J Phys Anthropol* 144:119–130
- 664 34. Macintosh AA, Pinhasi R, Stock JT (2014) Divergence in male and female manipulative
665 behaviors with the intensification of metallurgy in Central Europe. *PLoS ONE*
666 9(11):e112116
- 667 35. Sládek V, Ruff CB, Berner M, et al (2016) The impact of subsistence changes on humeral
668 bilateral asymmetry in Terminal Pleistocene and Holocene Europe. *J Hum Evol* 92:37–49
- 669 36. De Pascale A (2008) Le prime esplorazioni nelle caverne ossifere del Finalese: tracce, ipotesi
670 e scoperte ad opera di Issel, Perrando, Morelli, Rovereto, Rossi, Amerano. In: De Pascale A,
671 Del Lucchese A, Raggio O (eds) *La nascita della Paleontologia in Liguria: personaggi,*
672 *scoperte e collezioni tra XIX e XX secolo*. Atti del Convegno (Finale Ligure Borgo, 22-23
673 settembre 2006). Istituto Internazionale di Studi Liguri, Bordighera, pp 223–248
- 674 37. Bernabò Brea L (1946) *Gli Scavi nella Caverna delle Arene Candide. Parte I Gli Strati con*
675 *Ceramiche*. Collezione di Monografie Preistoriche ed Archeologiche, I. Istituto di Studi
676 Liguri, Bordighera

- 677 38. Bernabò Brea L (1956) *Gli Scavi nella Caverna delle Arene Candide (Finale Ligure). Parte*
678 *Prima: Gli Strati con Ceramiche: Campagne di Scavo 1948-50. Collezione di Monografie*
679 *Preistoriche ed Archeologiche, I. Istituto Internazionale di Studi Liguri, Bordighera*
- 680 39. Sparacello VS, Panelli C, Rossi S, et al (2019) *The re-discovery of Arma dell’Aquila (Finale*
681 *Ligure, Italy): New insights on Neolithic funerary behavior from the sixth millennium BCE*
682 *in the north-western Mediterranean. Quat Int 512:67–*
683 *81doi.org/10.1016/j.quaint.2019.02.003.*
- 684 40. Mannino MA, Talamo S, Goude G, et al (2018) *Analisi isotopiche e datazioni sul collagene*
685 *osseo degli inumati dell’Arma dell’Aquila. In: Biagi P, Starnini E, (eds) Gli Scavi nell’Arma*
686 *dell’Aquila (Finale Ligure, Savona): Le Ricerche e i Materiali degli Scavi del Novecento.*
687 *Quaderno 15. Società per la Preistoria e Protostoria della Regione Friuli-Venezia Giulia,*
688 *Trieste, pp 183–188*
- 689 41. Sparacello VS, Varalli A, Rossi S, et al (2019) *Large-scale AMS dating on human skeletal*
690 *series from Ligurian caves (northwestern Italy) and the Neolithic peopling of the*
691 *northwestern Mediterranean. Quat Intdoi.org/10.1016/j.quaint.2019.11.034.*
- 692 42. Parenti R, Messeri P (1962) *I resti scheletrici umani del Neolitico Ligure. Palaeontographia*
693 *Italica 50:5–165*
- 694 43. Marchi D, Sparacello VS, Shaw CN (2011) *Mobility and lower limb robusticity of a*
695 *pastoralist Neolithic population from North-Western Italy. In: Pinhasi R, Stock JY (eds)*
696 *Human bioarchaeology of the Transition to Agriculture. John Wiley & Sons, New York, pp*
697 *317–346*
- 698 44. Villotte S, Knüsel CJ (2013) *Understanding Entheseal Changes: Definition and Life Course*
699 *Changes. Int J Osteoarchaeol 23:135-146*
- 700 45. Villotte S, Assis S, Alves Cardoso F, et al (2016) *In search of consensus: terminology for*
701 *enthesal changes (EC). Int J Paleopath 13:49–55*
- 702 46. Villotte S (2006) *Connaissances médicales actuelles, cotation des enthésopathies: nouvelle*
703 *méthode. Bull Mem Soc Anthropol Paris 18:65–85*
- 704 47. Villotte S (2009) *Enthésopathies et activités des hommes préhistoriques. Recherche*
705 *méthodologique et application aux fossiles européens du Paléolithique Supérieur et du*
706 *Mésolithique. Archaeopress, Oxford, 236 p*

- 707 48. Villotte S, Castex D, Couallier V, et al (2010a) Enthesopathies as occupational stress
708 markers: evidence from the upper limb. *Am J Phys Anthropol*, 142 (2):224–234
- 709 49. Dutour O (1986) Enthesopathies (lesions of muscular insertions) as indicators of the
710 activities of Neolithic Saharan populations. *Am J Phys Anthropol* 71:221–224
- 711 50. Dutour O (2000) Chasse et activités physiques dans la Préhistoire: les marqueurs osseux
712 d’activités chez l’homme fossile. *Anthropol Prehist* 111:156–165
- 713 51. Villotte S, Churchill SE, Dutour OJ, et al (2010b) Subsistence activities and the sexual
714 division of labor in the European Upper Paleolithic and Mesolithic: evidence from upper
715 limb enthesopathies. *J Hum Evol* 59:35–43
- 716 52. Villotte S, Knüsel CJ (2014) “I sing of arms and of a man. . .”: medial epicondylitis and the
717 sexual division of labour in prehistoric Europe. *J Archaeol Sci* 43:168–174
- 718 53. Kennedy GE (1986) The relationship between auditory exostosis and cold water: a latitudinal
719 analysis. *Am J Phys Anthropol* 71:401–415
- 720 54. Villotte S, Knüsel CJ (2016) External auditory exostoses and prehistoric aquatic resource
721 procurement. *J Archaeol Sci: Rep* 6(4):633–6
- 722 55. Le Bras-Goude G, Binder D, Formicola V, et al (2006) Stratégies de subsistance et analyse
723 culturelle de populations néolithiques de Ligurie: approche par l’étude isotopique ($\delta^{13}\text{C}$ et
724 $\delta^{15}\text{N}$) des restes osseux. *Bull Mem Soc Anthropol Paris* 18:45–55
- 725 56. Goude G, Binder D, Del Lucchese A (2014) Alimentation et modes de vie néolithiques en
726 Ligurie. In: Bernabo Brea M, Maggi R, Manfredini A (eds) *Il Pieno Neolitico in Italia* (8-10
727 juin Finale Ligure 2009). *Riv Studi Liguri* 77:371–387
- 728 57. Rowley-Conwy P (1997) The animal bones from Arene Candide (Holocene sequence): final
729 report. In: Maggi R (ed) *Arene Candide: a functional and environmental assessment of the*
730 *Holocene sequence (excavations Bernabò Brea-Cardini 1940–1950)*. *Istituto Italiano di*
731 *paleontologia umana, Il calamo, Roma, ns 5, pp 153–277*
- 732 58. Molnar S (1972) Tooth wear and culture: a survey of tooth functions among some prehistoric
733 population. *Curr Anthropol* 13:511–526
- 734 59. Brooks S, Suchey JM (1990) Skeletal age determination based on the os pubis: a comparison
735 of the Acsádi-Nemeskéri and Suchey-Brooks methods. *Hum Evol* 5:227–238
- 736 60. Buckberry JL, Chamberlain AT (2002) Age estimation from the auricular surface of the
737 ilium: a revised method. *Am J Phys Anthropol* 119:231–239

- 738 61. Schmitt A (2005) Une nouvelle méthode pour estimer l'âge au décès des adultes à partir de la
739 surface sacro-pelvienne iliaque. *Bull Mem Soc Anthropol Paris* 17(1-2):1-15
- 740 62. Ubelaker DH (1989) *Human skeletal remains: excavation, analysis, interpretation,*
741 *Taraxacum, Washington, 172 p*
- 742 63. Smith BH (1991) Standards of human tooth formation and dental age assessment. In:
743 KelleyMA, Larsen CS (eds) *Advances in dental anthropology*. Wiley-Liss, New York, pp
744 143-168
- 745 64. AlQahtani SJ, Hector MP, Liversidge HM (2010) The London atlas of human tooth and
746 eruption. *Am J Phys Anthropol* 142:481-490
- 747 65. Schaefer M, Black S, Scheuer L (2009) *Juvenile osteology – a laboratory and field manual,*
748 *Academic Press, New York, 384 p*
- 749 66. Ríos L, Cardoso HF (2009) Age estimation from stages of union of the vertebral epiphyses of
750 the ribs. *Am J Phys Anthropol* 140:265-274
- 751 67. Cardoso HF, Ríos L (2011) Age estimation from stages of epiphyseal union in the presacral
752 vertebrae. *Am J Phys Anthropol* 144:238-247
- 753 68. Boccone S, Micheletti Cremasco M, Bortoluzzi S, et al (2010) Age estimation in subadults
754 Egyptian remains. *J Comp Hum Biol* 61:337-358
- 755 69. Buikstra JE, Ubelaker DH (1994) *Standards for Data Collection from Human Skeletal*
756 *Remains, Arkansas Archaeological Survey Research Series No 44, Fayetteville, 218 p*
- 757 70. Loth SR, Henneberg M (1996) Mandibular ramus flexure: a new morphologic indicator of
758 sexual dimorphism in the human skeleton. *Am J Phys Anthropol* 99:473-485
- 759 71. Bruzek J (2002) A method for visual determination of sex using the human hip bone. *Am J*
760 *Phys Anthropol* 117:157-168
- 761 72. Thieme FP, Schull WJ (1957) Sex determination of the skeleton. *Hum Biol* 29:242-273.
- 762 73. Richman EA, Michel ME, Schuller-Ellis FP, et al (1979) Determination of sex by
763 discriminant function analysis of postcranial skeletal measurements. *J Forensic Sci* 24:159-
764 167.
- 765 74. Steyn M, İşcan MY (1997) Sex Determination from the Femur and Tibia in South African
766 Whites. *Forensic Sci Int* 90(1-2):111-119
- 767 75. Murail P, Bruzek J, Braga J (1999) A new approach to sexual diagnosis in past populations,
768 Practical adjustments from Van Vark's procedure. *Int J Osteoarchaeol* 9:39-53

- 769 76. Shaw C, Stock J (2009a) Intensity, repetitiveness, and directionality of habitual adolescent
770 mobility patterns influence the tibial diaphysis morphology of athletes. *Am J Phys Anthropol*
771 140:149–159
- 772 77. Shaw C, Stock J (2009b) Habitual throwing and swimming correspond with upper limb
773 diaphyseal strength and shape in modern human athletes. *Am J Phys Anthropol* 140:160–172
- 774 78. Macintosh AA, Stock JT (2019) Intensive terrestrial or marine locomotor strategies are
775 associated with inter- and intra-limb bone functional adaptation in living female athletes. *Am*
776 *J Phys Anthropol* DOI: 10.1002/ajpa.23773
- 777 79. Sparacello VS, d’Ercole V, Coppa A (2015) A bioarchaeological approach to the
778 reconstruction of changes in military organization among Iron Age Samnites (Vestini) from
779 Abruzzo, central Italy. *Am J Phys Anthropol* 156:305–316
- 780 80. Ruff CB (2002) Long bone articular and diaphyseal structure in Old World monkeys and
781 apes. I: locomotor effects. *Am J Phys Anthropol* 119:305–342
- 782 81. Nagurka ML, Hayes WC (1980) An interactive graphics package for calculating cross-
783 sectional properties of complex shapes. *J Biomech* 13:59–64
- 784 82. Sparacello VS, Pearson OM (2010) The importance of accounting for the area of the
785 medullary cavity in cross-sectional geometry: a test based on the femoral midshaft. *Am J*
786 *Phys Anthropol* 143:612–624
- 787 83. Stock JT, Shaw CN (2007) Which measures of skeletal robusticity are robust? A comparison
788 of external methods of quantifying diaphyseal strength to cross-sectional geometric
789 properties. *Am J Phys Anthropol* 134:412–423
- 790 84. Macintosh AA, Davies TG, Ryan TM, et al (2013) Periosteal versus true cross-sectional
791 geometry: a comparison along humeral, femoral, and tibial diaphysis. *Am J Phys*
792 *Anthropol* 150:442–452
- 793 85. Ruff CB (2018) Quantifying skeletal robusticity. In: Ruff CB (ed) *Skeletal variation and*
794 *adaptation in Europeans: Upper Paleolithic to the Twentieth Century*. John Wiley and Sons,
795 Inc., New York, pp 39–47.
- 796 86. Ruff CB (2000) Body size, body shape, and long bone strength in modern humans. *J Hum*
797 *Evol* 38:269–290
- 798 87. Trinkaus E, Ruff CB (2012) Femoral and tibial diaphyseal cross-sectional geometry in
799 Pleistocene Homo. *PaleoAnthropology* 2012:13–62

- 800 88. Sparacello VS, Villotte S, Shackelford LL, et al (2017) Patterns of Humeral Asymmetry
801 among Late Pleistocene Humans. *CR Palevol* 16(5–6):680–689
- 802 89. Trinkaus E, Churchill SE, Ruff CB (1994) Postcranial robusticity in Homo. II. humeral
803 bilateral asymmetry and bone plasticity. *Am J Phys Anthropol* 93:1–34
- 804 90. Rhodes JA, Knüsel CJ (2005) Activity-related skeletal change in medieval humeri: cross-
805 sectional and architectural alterations. *Am J Phys Anthropol* 128:536–546
- 806 91. Holt BM (2003) Mobility in Upper Paleolithic and Mesolithic Europe: evidence from the
807 lower limb. *Am J Phys Anthropol* 122:200–215
- 808 92. Martiarena ML (2016) Analyse d'un marquer d'activité dans une population humaine
809 préhistorique. M2 Thesis, Université libre de Bruxelles, Bruxelles
- 810 93. Polet C, Martiarena ML, Villotte S, et al (2019) Throwing activities among Neolithic
811 populations from the Meuse River Basin (Belgium, 4500–2500 BC) with a focus on
812 adolescents. *Child Past* 12(2):81–95
- 813 94. Crowe F, Sperduti A, O'Connell TC, et al (2010) Water-related occupations and diet in two
814 Roman coastal communities (Italy, first to third century AD): correlation between stable
815 carbon and nitrogen isotope values and auricular exostosis prevalence. *Am J Phys Anthropol*
816 142:355–366
- 817 95. Standen VG, Arriaza BT, Santoro CM (1997) External auditory exostosis in prehistoric
818 Chilean populations: A test of the cold water hypothesis. *Am J Phys Anthropol* 103:119–129
- 819 96. Velasco-Vazquez J, Betancor-Rodriguez A, Arnay-De-La Rosa M, et al (2000) Auricular
820 exostoses in the prehistoric population of Gran Canaria, *Am J Phys Anthropol* 112:49–55
- 821 97. Villotte S, Stefanović S, Knüsel CJ (2014) External auditory exostoses and aquatic activities
822 during the Mesolithic and the Neolithic in Europe: Results from a large prehistoric sample.
823 *Anthropologie LII/1(1):73–89*
- 824 98. Cooper A, Tong R, Neil R, et al (2010) External auditory canal exostoses in white water
825 kayakers. *Br J Sports Medicine* 44:144–147
- 826 99. Hurst W, Bailey M, Hurst B (2004) Prevalence of external auditory canal exostoses in
827 Australian surfboard riders. *J Laryngol Otol* 118:348–351
- 828 100. Sparacello VS (2013) The Bioarchaeology of Changes in Social Stratification, Warfare,
829 and Habitual Activities among Iron Age Samnites of Central Italy. PhD Thesis, University of
830 New Mexico, Albuquerque

- 831 101. Villotte S, Samsel M, Sparacello VS (2017) The paleobiology of the two adult skeletons
832 from Baouso da Torre (Bausu da Ture) (Liguria, Italy): implications for our understanding
833 of Gravettian lifestyle. *Comptes Rendus Palevol* 16:462–473
- 834 102. Meyer C, Nicklisch N, Held P, et al (2011) Tracing patterns of activity in the human
835 skeleton: An overview of methods, problems, and limits of interpretation. *J Comp Hum Biol*
836 62:202–217
- 837 103. Jurmain R, Alves Cardoso F, Henderson C, et al (2012) Bioarchaeology's Holy Grail:
838 The reconstruction of activity. In: Grauer AL (ed) *A Companion to Paleopathology*. Wiley-
839 Blackell, New York, pp 531–552
- 840 104. Lazenby RA (1990) Continuing periosteal apposition II: the significance of peak bone
841 mass, strain equilibrium, and age-related activity differentials for mechanical compensation
842 in human tubular bones. *Am J Phys Anthropol* 82:473–484
- 843 105. Martin RB, Atkinson PJ (1977) Age and sex-related changes in the structure and strength
844 of the human femoral shaft. *J Biomech* 10:223–231
- 845 106. Ruff C, Hayes W (1988) Sex differences in age-related remodeling of the femur and tibia.
846 *J Orthop Res* 6:886–896
- 847 107. Agostini G, Holt BM, Relethford JH (2018) Bone functional adaptation does not erase
848 neutral evolutionary information. *Am J Phys Anthropol* 166.10.1002/ajpa.23460.
- 849 108. Alves Cardoso FA, Henderson CY (2010) Enthesopathy formation in the humerus: Data
850 from known age-at-death and known occupation skeletal collections. *Am J Phys Anthropol*
851 141:550–560
- 852 109. Weiss E (2007) Muscle markers revisited: Activity pattern reconstruction with controls in
853 a Central California Amerind population. *Am J Phys Anthropol* 133:931–940
- 854 110. Nikita E, Xanthopoulou P, Bertsatos A, et al (2019) A three-dimensional digital
855 microscopic investigation of enthesal changes as skeletal activity markers. *Am J Phys*
856 *Anthropol* DOI: 10.1002/ajpa.23850
- 857 111. Lieverse AR, Stock JT, Katzemberg MA, et al (2011) The bioarcheology of habitual
858 activity and dietary change in the Siberian Middle Holocene. In: Pinhasi R, Stock J (eds)
859 *Human bioarchaeology of the Transition to Agriculture*. Wiley-Liss, New York, pp 265–291.
- 860 112. Niinimäki S (2012) The relationship between musculoskeletal stress markers and
861 biomechanical properties of the humeral diaphysis. *Am J Phys Anthropol* 147:618–628

- 862 113. Michopoulou E, Nikita E, Henderson CY (2017) A test of the effectiveness of the
863 Coimbra method in capturing activity-induced enthesal changes. *Int J Osteoarchaeol*
864 27:409–417
- 865 114. Michopoulou E, Nikita E, Valakos ED (2015) Evaluating the efficiency of different
866 recording protocols for enthesal changes in regards to expressing activity patterns using
867 archival data and cross-sectional geometric properties. *Am J Phys Anthropol* 158:557–568
- 868 115. Ibáñez-Gimeno P, Galtés I, Jordana X, et al (2013) Enthesal changes and functional
869 implications of the humeral medial epicondyle. *IntJ Osteoarch* 23:211–220
- 870 116. Desse-Berset N, Desse J (1999) Les poissons. In: Tinè S (ed) *Il Neolitico della caverna*
871 *delle Arene Candide (scavi 1972-1977)*. Istituto Internazionale di Studi Liguri, Bordighera, pp
872 36–50
- 873 117. Di Bartolomeo J, Paparella M, Meyerhoff W (1991) Cysts and tumors of the external ear.
874 In: Shumrick D, Gluckman J, Meyerhoff W (eds) *Otolaryngology*. 3rd edition, 2:1243–1258.
- 875 118. Fowler EP, Osmun PM (1942) New bone growth due to cold water in the ears, *Arch*
876 *Otolaryngol Head Neck Surg* 36:455–466
- 877 119. Kusaka S, Hyodo F, Yumoto T, et al (2010) Carbon and nitrogen stable isotope analysis
878 on the diet of Jomon populations from two coastal regions of Japan. *J Archaeol Sci* 37:1968–
879 1977
- 880
- 881
- 882

Individual ID	Analysis	Age Class	Age Dental	Age Poster.	Sex	AMS date cal. 95.4%	Chrono-cultural attribution
Acqua o Morto 251+254	CSG*, EAE	Adult	30-50	U	M	4797-4695	SMP
Acqua o Morto 252+253	CSG*, ECs	Adult	30-50	U	F	5301-5073	ICC
Arene Candide 6PE 3 Perrando	CSG, ECs	Adult	30-50	30-50	M	failed (prob. V mill. BCE) (5657-4620) ¹	SMP
Arene Candide 1 Tinè	CSG*, EAE	Late adol.	15-18	15-17	M	4704-4374	SMP
Arene Candide 2 Tinè	CSG, EAE, ECs	Adult	30-50	30-50	M	5209-5011	ICC
Arene Candide 6622.1FI + 6730.2FI	CSG*, EAE, ECs	Adult	U	U	F	5208-5000	ICC
Arene Candide 6730.1FI	CSG*, ECs	Young Adult	U	20-30	F	4688-4540	SMP
Arene Candide 7PE	CSG, EAE, ECs	Adult	30-50	30-50	M	4767-4586	SMP
Arene Candide 8PE	CSG, EAE, ECs	Adult	30-50	30-50	M	4708-4555	SMP
Arene Candide II BB	CSG*, ECs	Adult	U	U	F	failed (prob. V mill. BCE)	SMP
Arene Candide III BB	CSG, EAE, ECs	Adult	30-50	30-50	M	4800-4619	SMP
Arene Candide IV BB (+ 6726FI + 6730.7FI)	CSG, EAE, ECs	Young Adult	20-30	20-30	F	4766-4558	SMP
Arene Candide IX BB	CSG, EAE, ECs	Adult	30-50	30-50	F	4779-4611	SMP
Arene Candide VI BB	CSG, EAE, ECs	Adult	30-50	U	M	4581-4457	SMP
Arene Candide VII BB	CSG, EAE, ECs	Adult	30-50	30-50	F	4778-4603	SMP
Arma dell'Aquila 1 Richard	CSG, EAE, ECs	Adult	30-50	30-50	M	5361-5221	ICC
Arma dell'Aquila 1 Zambelli	CSG, EAE, ECs	Adult	30-50	U	F	4723-4551	SMP
Arma dell'Aquila 2 Richard	CSG, EAE, ECs	Adult	>50	U	M	5213-5010	ICC
Arma dell'Aquila 3 Richard	CSG*, ECs	Adult	30-50	U	M	5202-4962	ICC
Arma dell'Aquila 5 Richard	CSG, EAE	Adult	30-50	U	F	5208-4956	ICC
Arma dell'Aquila RS5	CSG*	Adult	U	U	U	5750-5645	ICC
Arma dell'Aquila RS9	CSG*	Adult	U	U	F	5206-4911	ICC
Bergeggi (IV) 3573 + 3573bis	EAE	Adult	U	30-50	F	failed (prob. V mill. BCE)	SMP
Bergeggi 1 Modigliani	EAE	Adolescent	13-19 (12-15?)	12-13	U	4527-4370	SMP
Bergeggi 2 Modigliani	CSG*, EAE, ECs	Young Adult	20-30	30-50	M	4680-4494	SMP
Bergeggi 3 Modigliani	CSG, ECs	Adult	U	U	F	4488-4364	SMP
Bergeggi 4 Modigliani + PE01177, 6893FI	CSG, ECs	Young Adult	U	17-19	M	4680-4494	SMP
Bergeggi 5 Modigliani	CSG, EAE	Adult	U	U	F	4539-4374	SMP
La Matta 01085	EAE	Young Adult	20-30	U	M?	n/a (prob. V mill. BCE)	SMP

La Matta 01086	EAE	Adolescent	11.5-12.5	U	U	n/a (prob. V mill. BCE)	SMP
Nasino 1	CSG*, EAE, ECs	Late adol.	13-19	16-20	M	4232-4000	Chassean
Pian del Ciliegio Adult	CSG, ECs	Young Adult	20-30	20-30	M	4690-4544	SMP
Pipistrelli 3_El Muerto N3	CSG*, ECs	Adult	>20	U	M	4723-4614	SMP
Pipistrelli 5 dep23.I.28 Cirillo	CSG*, ECs	Adult	U	U	M	5016-4844	SMP
Pipistrelli 6 dep 23.II.41 Angelina	CSG*, ECs	Adult	U	30-50	M	4995-4810	SMP
Pollera 1 Tiné	CSG, EAE, ECs	Young Adult	20-30	20-30	F	4712-4552	SMP
Pollera 10PE	CSG, ECs	Young Adult	U	20-30	M	4701-4550	SMP
Pollera 110a	CSG*, ECs	Adult	U	U	F	4690-4544 ²	SMP
Pollera 110b	CSG*	Adult	U	U	F	4690-4544 ²	SMP
Pollera 12PE	CSG, EAE, ECs	Adult	30-50	30-50	F	4794-4687	SMP
Pollera 13PE	CSG, EAE, ECs	Adult	30-50	30-50	M	4686-4527	SMP
Pollera 14PE	CSG, EAE, ECs	Adult	30-50	30-50	F	4786-4616	SMP
Pollera 1PE Issel-Morelli	EAE	Adolescent	10-12	<11	U	4786-4616	SMP
Pollera 22PE	CSG*, ECs	Adult	U	30-50	M	4786-4616	SMP
Pollera 30PE	CSG, EAE, ECs	Adult	30-50	30-50	M	4689-4543	SMP
Pollera 31PE	CSG*	Adult	U	30-50	M	4701-4548	SMP
Pollera 32PE	CSG, ECs	Adult	U	30-50	M	failed (prob. V mill. BCE)	SMP
Pollera 33PE	CSG, EAE, ECs	Adult	30-50	30-50	F	4711-4555	SMP
Pollera 34PE	EAE	Adolescent	13-19	11-16	U	4723-4558	SMP
Pollera 6246PE	CSG, EAE, ECs	Adult	30-50	30-50	F	4650-4462	SMP
Pollera 6673.6FI	CSG*	Adult	U	U	U	4879-4724	SMP
Pollera 6690bis.3FI	CSG*, ECs	Young Adult	U	20-30	F	4794-4687	SMP
Arene Candide 6621.1	Osteometrics only	Adolescent	13-19	12-16	F	4726-4557	SMP
Arene Candide 6634.1+6626.1+6730.2tris	Osteometrics only	Adult	U	U	F	4792-4688	SMP
Arene Candide I BB 6731.1FI + 6627.2FI	Osteometrics only	Adolescent	U	13-17	U	4690-4544	SMP
Arene Candide V BB	Osteometrics only	Adolescent	c. 15	14-16	U	4720-4557	SMP
Pipistrelli 4_dep23.III.73_76	Osteometrics only	Young Adult	20-30	20-30	F	5207-4909	ICC

885 Table 1 – Individuals from the Neolithic in Liguria included in this study, indicating the analyses
886 performed for each. Asterisks indicate the individuals for which CSG data was collected for this study (cf.
887 the last CSG sample [32]). A more detailed list of individuals examined for this study, including their
888 uncalibrated date and the museum where they are preserved, is available as Supplementary Information
889 (excel file “Supplementary Information Tables A”). See also [41].

890 CSG: cross-sectional geometry; EAE: external auditory exostoses; ECs: enthesal changes; U: sex
891 undetermined; M: male; F: female; SMP: Square Mouthed Pottery; ICC; Impresso-Cardial Complex.

892 ¹ date from previous studies with large error (see excel file “Supplementary Information_raw data by
893 analysis”). ² Date belongs to PO 110c, individual found in close association with PO 110a-b.

894 Table 1 – Individus néolithiques de Ligurie inclus dans l’étude, avec pour chacun d’eux une indication
895 des analyses effectuées. Une liste plus détaillée, incluant les données radiocarbone non-calibrées et les
896 lieux de conservation, est disponible dans l’annexe Table A. Voir également [41].

897 U: sexe indéterminé; M: homme; F: femme; ¹ date provenant d’études antérieures avec une large erreur
898 (voir l’annexe “Supplementary Information_raw data by analysis”). ² Date associée à PO 110c, un
899 individu trouvé en étroite relation avec PO 110a-b.

900

	LUP			SMP			IRON			SMP-LUP	SMP-IRON
Males	n	mean	SD	n	mean	SD	n	mean	SD	HSD¹	HSD
Z _p humerus R	13	52.03	7.13	15	54.41	9.71	124	58.82	10.29	NS	NS
Z _p humerus L	13	42.28	10.53	16	48.96	7.65	122	50.80	8.72	NS	NS
HUMBA	14	58.61	28.03	15	18.95	9.11	200	24.67	14.97	p < 0.0001	NS
Z _p femur	18	106.24	10.53	13	102.90	14.47	144	101.62	15.05	NS	NS
I _x /I _y femur	18	1.40	0.25	13	1.30	0.17	156	1.07	0.18	NS	p < 0.0001
Z _p tibia	16	109.41	15.78	13	105.82	19.29	94	102.11	16.67	NS	NS
I _{max} /I _{min} tibia	16	2.85	0.64	13	2.63	0.30	97	2.36	0.47	NS	NS
Females	n	mean	SD	n	mean	SD	n	mean	SD	HSD	HSD
Z _p humerus R	4	49.24	8.06	11	45.20	4.30	53	48.69	6.92	NS	NS
Z _p humerus L	5	48.15	7.51	13	46.28	5.00	52	46.71	6.11	NS	NS
HUMBA	4	12.53	2.81	12	6.4	5.66	97	13.81	10.64	NS	p < 0.05
Z _p femur	7	96.75	13.39	12	94.81	11.40	68	93.75	13.71	NS	NS
I _x /I _y femur	11	1.26	0.27	12	1.15	0.17	71	0.99	0.18	NS	p < 0.05
Z _p tibia	5	101.17	8.45	13	97.31	15.75	39	86.04	13.73	NS	p < 0.05
I _{max} /I _{min} tibia	6	2.14	0.27	13	2.34	0.32	41	2.11	0.40	NS	NS
Sexual Dimorphism		LUP			SMP			IRON			
Z _p humerus R		NS ²			p < 0.01			p < 0.0001			
Z _p humerus L		NS			NS			p < 0.01			
HUMBA		p < 0.01			p < 0.001			p < 0.0001			
Z _p femur		p < 0.1			NS			p < 0.001			
I _x /I _y femur		NS			p < 0.05			p < 0.01			
Z _p tibia		NS			NS			p < 0.0001			
I _{max} /I _{min} tibia		p < 0.05			p < 0.05			p < 0.01			

Table 2 – Main CSG properties of the Ligurian Neolithic sample (SMP: Square Mouthed Pottery), and of the comparative samples (LUP: European Late Upper Palaeolithic; IRON: Orientalizing-Archaic Samnites from Abruzzo, Italy), by sex. The raw CSG data per individual are given in Supplementary Information Tables A. R: right; L: left. ¹ Tukey’s Honest Significant Difference, post-hoc test of ANOVA among groups (LUP-SMP-IRON) by sex. ² T-test between sexes, by period. NS: non-significant.

Table 2 – Principales propriétés géométriques les collections squelettiques néolithiques de Ligurie (SMP: culture des Vases à Bouche carrée) et des échantillons de comparaisons (LUP : fin du Paléolithique supérieur européen ; IRON : Age du Fer des Abruzzes, Italie centrale) suivant le sexe. Les données brutes par individu sont disponibles dans l’annexe Table A. R : droite ; L : gauche. ¹ test des étendues de Tukey, Analyse de variance post-hoc (LUP-SMP-IRON) par sexe. ² T-test suivant le sexe, par période. NS: non significatif.

	LE Left			Fisher Exact test M vs F (A vs B+C)
	A	B	C	
Males	6 (8)	3 (4)	2 (3)	NS
Females	5	2	1	
	LE Right			Fisher Exact test M vs F (A vs B+C)
	A	B	C	
Males	4	4 (5)	1	NS
Females	4 (5)	3	2	
	Fisher Exact test R vs L (A vs B+C)			
Males	NS			
Females	NS			
	LE Asymmetry			Fisher Exact test (R+L vs NO)
	L	R	NO	
Males	0	0	6	NS
Females	0	1	6	

Table 3 – Number of individuals showing entheseal changes (ECs) in the lateral epicondyle (common extensor origin) of the humerus in the Ligurian Neolithic sample, by sex. Numbers and p-values outside of parentheses refer to the SMP sample, numbers and p-values inside the parentheses refer to the pooled Neolithic sample. LE: lateral epicondyle. Scores: A, no change; B, minor changes; C, major changes. L: left; R: right; NO: no asymmetry; M: males; F: females.

Table 3 – Nombre d'individus présentant un changement enthésique (ECs) au niveau de l'épicondyle latéral (origine commune des extenseurs) de l'humérus dans l'échantillon néolithique de Ligurie, suivant le sexe. Les nombres et valeurs p. à l'extérieur des parenthèses font référence à l'échantillon SMP, celles entre parenthèses font référence à l'échantillon total. LE: épicondyle latéral. Scores: A, pas de changement; B, changements mineurs; C, changements majeurs. L: gauche; R: droit; NO: pas d'asymétrie; M: hommes; F: femmes.

	CFT Left			Fisher Exact test M vs F (A vs B+C)
	A	B	C	
Males	7 (9)	3	0	NS
Females	6 (7)	2	0	
	CFT Right			Fisher Exact test M vs F (A vs B+C)
	A	B	C	
Males	4	6	0	NS
Females	5 (6)	4	0	
	Fisher Exact test R vs L (A vs B+C)			
Males	NS			
Females	NS			
	CFT ASYMM			Fisher Exact test M vs F (R+L vs NO)
	L	R	NO	
Males	0	4	2	p < 0.1 (p < 0.05)
Females	0	1	6 (7)	

Table 4 – Number of individuals showing entheseal changes (ECs) in the area of attachment of the common flexor tendon (CFT; medial epicondyle of the humerus) in the Ligurian Neolithic sample, by sex. Numbers and p-values not in parentheses refer to the SMP sample, numbers and p-values in parentheses refer to the pooled Neolithic sample. Scores: A, no change; B, minor changes; C, major changes. L: left; R: right; NO: no asymmetry; M: males; F: females.

Table 4 – Nombre d'individus présentant un changement enthésique (ECs) au niveau de l'insertion commune des fléchisseurs de l'humérus (CFT) dans l'échantillon néolithique de Ligurie, suivant le sexe. Les nombres et valeurs p. à l'extérieur des parenthèses font référence à l'échantillon SMP, celles entre parenthèses font référence à l'échantillon total. Scores: A, pas de changement; B, changements mineurs ; C, changements majeurs. L: gauche; R: droit; NO: pas d'asymétrie; M: hommes; F: femmes.

	UCL_Left		Fisher Exact test M vs F	
	A	P		
Males	11 (16)	1	p = NS	
Females	11 (12)	1		
	UCL_Right		Fisher Exact test M vs F	
	A	P		
Males	3 (6)	7	p < 0.05 (p < 0.1)	
Females	10 (11)	2		
	Fisher Exact test R vs L			
Males	p < 0.01 (p < 0.01)			
Females	NS			
	UCL ASYMM			
	L	R	NO	Fisher Exact test M vs F (R+L vs NO)
Males	0	6	2 (5)	p < 0.01 (p < 0.05)
Females	0	1	10 (11)	

Table 5 – Number of individuals showing erosions/cavitations/geodes and/or fissures/well-defined pits (ECG+FP) in the anterior band of the ulnar collateral ligament in the Ligurian Neolithic sample, by sex. Numbers and p-values not in parentheses refer to the SMP sample, numbers and p-values in parentheses refer to the pooled Neolithic sample. A: no change; P: trait is present; L: left; R: right; M: males; F: females.

Table 5 – Nombre d'individus présentant une érosion/cavité/ géodes et/ou une fissure/dépression(ECG+FP) au niveau de l'insertion du fascia antérieur du ligament collatéral médial dans l'échantillon néolithique de Ligurie, suivant le sexe. Les nombres et valeurs p. à l'extérieur des parenthèses font référence à l'échantillon SMP, celles entre parenthèses font référence à l'échantillon total. Scores: A, pas de changement; P, changements present. L: gauche; R: droit; NO: pas d'asymétrie; M: hommes; F: femmes.

	ME any change Left		Fisher Exact test M vs F	
	A	P		
Males	7 (9)	3	NS	
Females	4 (5)	2		
	ME any change Right		Fisher Exact test M vs F	
	A	P		
Males	1 (2)	8	NS	
Females	3 (4)	6		
	Fisher Exact test R vs L			
Males	p < 0.05 (p < 0.05)			
Females	NS			
	ME any change ASYMM			
	L	R	NO	Fisher Exact test M vs F (R+L vs NO)
Males	0	5	1	NS
Females	0	2	4 (5)	

Table 6 – Number of individuals showing any entheseal change in the common flexor attachment site (scored B or C in Table 4) or in the anterior band of the ulnar collateral ligament (ECG or FP) in the Ligurian Neolithic sample, by sex. The table includes data exclusively from individuals for which both areas could be examined. Numbers and p-values not in parentheses refer to the SMP sample, numbers and p-values in parentheses refer to the pooled Neolithic sample. L: left; R: right; M: males; F: females.

Table 6 – Nombre d'individus présentant un changement enthésique au niveau de l'origine commune des flechisseurs (score B ou C dans le tableau 4) ou au niveau de l'insertion du fascia antérieur du ligament collatéral médial (ECG ou FP) dans l'échantillon néolithique de Ligurie, suivant le sexe. Le tableau inclus uniquement les individus pour lesquels les deux zones ont pu être examinées. Les nombres et valeurs p. à l'extérieur des parenthèses font référence à l'échantillon SMP, celles entre parenthèses font référence à l'échantillon total. L: gauche; R: droit; NO: pas d'asymétrie; M: hommes; F: femmes.

	ME any change Left		Fisher Exact test M vs F	
	A	P		
Males	9 (14)	3	NS	
Females	11 (12)	3		
	ME any change Right		Fisher Exact test M vs F	
	A	P		
Males	2 (5)	10	NS	
Females	6 (7)	6		
	Fisher Exact test R vs L			
Males	p < 0.05 (p < 0.05)			
Females	NS			
	ME any change ASYMM			
	L	R	NO	Fisher Exact test M vs F (R+L vs NO)
Males	0	7	1 (4)	p < 0.05 (p < 0.1)
Females	0	3	9 (10)	

Table 7 – Number of individuals showing any entheseal change in the common flexor attachment site (scored B or C in Table 4) or in the anterior band of the ulnar collateral ligament (ECG or FP) in the Ligurian Neolithic sample, by sex. The table include data from all individuals for which at least one area could be examined. Numbers and p-values not in parentheses refer to the SMP sample, numbers and p-values in parentheses refer to the pooled Neolithic sample. L: left; R: right; M: males; F: females.

Table 7 – Nombre d'individus présentant un changement enthésique au niveau de l'origine commune des flechisseurs (score B ou C dans le tableau 4) ou au niveau de l'insertion du fascia antérieur du ligament collatéral médial (ECG ou FP) dans l'échantillon néolithique de Ligurie, suivant le sexe. Le tableau inclus tous les individus pour lesquels au moins une zone a pu être examinée. Les nombres et valeurs p. à l'extérieur des parenthèses font référence à l'échantillon SMP, celles entre parenthèses font référence à l'échantillon total. L: gauche; R: droit; NO: pas d'asymétrie; M: hommes; F: femmes.

Figures



Figure 1 – Geographical location of the Finalese area in Liguria (northwestern Italy), and location of the cave sites included in this study: 1) Arene Candide; 2) La Matta; 3) Pollera; 4) Arma dell’Aquila; 5) Pipistrelli; 6) Pian del Ciliegio; 7) Bergeggi; 8) Nasino.

Figure 1 – Localisations de la zone de Finale Ligure en Ligurie (Nord-Ouest de l’Italie) et des sites inclus dans cette étude : 1) Arene Candide; 2) La Matta; 3) Pollera; 4) Arma dell’Aquila; 5) Pipistrelli; 6) Pian del Ciliegio; 7) Bergeggi; 8) Nasino.




Figure 2 – Bony defects at the medial epicondyle, in the site of attachment of the anterior fascia of the ulnar collateral ligament. A) Individual Arene Candide 7 PE, despite the concretion obscuring part of the site, clearly shows a cavitation of the cortical surface and the formation of a geode, and was scored 1 for ECG. B) Individual Pollera 30, showing a fissure and well-defined pits in the area, but no interruption of the cortical surface. Scored 0 for ECG, 1 for ECG+FP.

Figure 2 – Défauts osseux au niveau de l'épicondyle médial, dans la zone d'insertion de la fascia antérieure du ligament collatéral médial. A) Individu Arene Candide 7 PE, qui en dépit de la concrétion qui recouvre la zone montre clairement un relief négatif au niveau de la surface et la formation d'une géode, avec un score 1 pour ECG. B) Individu Pollera 30, qui présente une fissure et des dépressions clairement visibles, mais sans rupture de la surface. Scoré 0 pour ECG, 1 pour ECG+FP.




Figure 3 – Boxplots showing the main cross-sectional geometric (CSG) properties of the upper limb in the Ligurian Neolithic skeletal series, subdivided by chronological phase. The boxes indicate the standard error, the whiskers indicate two standard deviations from the mean. ICC: Impresso-Cardial Complex (c. 5800-5000 BCE); SMP: Square Mouthed Pottery (c. 5000-4300 BCE). F: females (circles); M: males (triangles); U: undetermined sex (squares). A) Mid-distal size-standardized right humeral robusticity; B) mid-distal size-standardized left humeral robusticity; C) midshaft size-standardized right humeral robusticity; D) midshaft size-

standardized left humeral robusticity; E) mid-distal humeral asymmetry in torsional rigidity (J) expressed as percentage; F) midshaft humeral asymmetry in torsional rigidity (J) expressed as percentage. All raw data are given in Supplementary Information Tables A.

Figure 3 – Boîtes à moustaches montrant les principales propriétés géométriques des sections transverses (CSG) du membre supérieur pour les collections squelettiques néolithiques de Ligurie, par phase chronologique. La boîte indique l'erreur standard et les moustaches montrent deux écart-types. ICC: complexe impresso-cardial (c. 5800-5000 BCE); SMP: culture des Vases à Bouche carrée (c. 5000-4300 BCE). F: femmes (cercles); M: hommes (triangles); U: sexes indéterminés (carrés). A) Robustesse humérale droite à mi distale de diaphyse, standardisée sur la taille; B) Robustesse humérale gauche à mi distale de diaphyse, standardisée sur la taille ;C) Robustesse humérale droite à mi diaphyse, standardisée sur la taille ; D) Robustesse humérale gauche à mi diaphyse, standardisée sur la taille ; E) Asymétrie humérale du moment polaire de l'aire (J) à mi distal de diaphyse exprimée en pourcentage ; F) Asymétrie humérale du moment polaire de l'aire (J) à mi diaphyse exprimée en pourcentage. Toutes les données brutes sont disponibles dans l'annexe Table A.




Figure 4 – Boxplots showing the main cross-sectional geometric (CSG) properties of the lower limb in the Ligurian Neolithic skeletal series, subdivided by chronological phase. The boxes indicate the standard error, the whiskers indicate two standard deviations from the mean. ICC: Impresso-Cardial Complex (c. 5800-5000 BCE); SMP: Square Mouthed Pottery (c. 5000-4300 BCE). F: females (circles); M: males (triangles); U: undetermined sex (squares). A) Midshaft size-standardized femoral robusticity; B) mid-distal femoral shape index I_x/I_y ; C) midshaft size-

standardized tibial robusticity; D) midshaft tibial shape index I_{\max}/I_{\min} ; E) midshaft size-standardized fibular robusticity; F) midshaft fibular shape index I_{\max}/I_{\min} . All raw data are given in Supplementary Information Tables A.

Figure 4 – Boîtes à moustaches montrant les principales propriétés géométriques des sections transverses (CSG) du membre inférieur pour les collections squelettiques néolithiques de Ligurie, par phase chronologique. La boîte indique l'erreur standard et les moustaches montrent deux écart-types. ICC: complexe impresso-cardial (c. 5800-5000 BCE); SMP: culture des Vases à Bouche carrée (c. 5000-4300 BCE). F: femmes (cercles); M: hommes (triangles); U: sexes indéterminés (carrés). A) Robustesse fémorale à mi diaphyse, standardisée sur la taille ; B) Indice de forme (I_x/I_y) du fémur à mi distal de diaphyse ; C) Robustesse tibiale à mi diaphyse, standardisée sur la taille ; D) Indice de forme (I_{\max}/I_{\min}) du tibia à mi diaphyse ; E) Robustesse fibulaire à mi diaphyse, standardisée sur la taille ; F) Indice de forme (I_{\max}/I_{\min}) de la fibula à mi diaphyse. Toutes les données brutes sont disponibles dans l'annexe Table A.




Figure 5 – Boxplots showing diachronic changes in humeral cross-sectional geometric (CSG) properties from the European Late Upper Palaeolithic (LUP), through the Square Mouthed Pottery Neolithic of Liguria (SMP), to the Orientalizing-Archaic Iron Age of Abruzzo, central Italy (IRON), by sex. The boxes indicate the standard error, the whiskers indicate $1.96 \times SE$. A) Mid-distal size-standardized right humeral robusticity; B) mid-distal size-standardized left humeral robusticity; C) mid-distal humeral asymmetry in torsional rigidity.

Figure 5 – Boîtes à moustaches montrant les variations diachroniques des propriétés géométriques des sections transverses (CSG) de l'humérus suivant le sexe. LUP : fin Paléolithique supérieur européen ; SMP : culture des Vases à Bouche carrée ; IRON : Age du Fer des Abruzzes, Italie centrale. La boîte indique l'erreur standard et les moustaches montrent $1,96 \times$ l'erreur standard. A) Robustesse humérale droite à mi distale de diaphyse, standardisée sur la taille; B) Robustesse humérale gauche à mi distale de diaphyse, standardisée sur la taille ; C) Asymétrie humérale du moment polaire de l'aire (J) à mi distal de diaphyse.




Figure 6 – Boxplots showing diachronic changes in femoral cross-sectional geometric (CSG) properties from the European Late Upper Palaeolithic (LUP), through the Square Mouthed Pottery Neolithic of Liguria (SMP), to the Orientalizing-Archaic Iron Age of Abruzzo, central Italy (IRON), by sex. The boxes indicate the standard error, the whiskers indicate $1.96 \times SE$. A) Midshaft size-standardized femoral robusticity; B) mid-distal femoral shape index I_x/I_y .

Figure 6 – Boîtes à moustaches montrant les variations diachroniques des propriétés géométriques des sections transverses (CSG) du fémur suivant le sexe. LUP : fin du Paléolithique supérieur européen ; SMP : culture des Vases à Bouche carrée ; IRON : Age du Fer des Abruzzes, Italie centrale. La boîte indique l'erreur standard et les moustaches montrent $1,96 \times$ l'erreur standard. A) Robustesse fémorale à mi diaphyse, standardisée sur la taille ;B) Indice de forme (I_x/I_y) du fémur à mi distal de diaphyse.




Figure 7 – Boxplots showing diachronic changes in tibial cross-sectional geometric (CSG) properties from the European Late Upper Palaeolithic (LUP), through the Square Mouthed Pottery Neolithic of Liguria (SMP), to the Orientalizing-Archaic Iron Age of Abruzzo, central Italy (IRON), by sex. The boxes indicate the standard error, the whiskers indicate $1.96 \times SE$.

Italy (IRON), by sex. The boxes indicate the standard error, the whiskers indicate $1.96 \times SE$. A) Midshaft size-standardized tibial robusticity; B) midshaft tibial shape index I_{max}/I_{min} .

Figure 7 – Boîtes à moustaches montrant les variations diachroniques des propriétés géométriques des sections transverses(CSG) du tibia suivant le sexe. LUP : fin du Paléolithique supérieur européen ; SMP : culture des Vases à Bouche carrée ; IRON : Age du Fer des Abruzzes, Italie centrale. La boîte indique l'erreur standard et les moustaches montrent $1,96 * l'$ erreur standard. A) Robustesse tibiale à mi diaphyse, standardisée sur la taille ;B) Indice de forme (I_{max}/I_{min}) du tibia à mi de diaphyse.




Figure 8 – Boxplots showing differences in mid-distal humeral bilateral asymmetry in torsional rigidity in subsamples based on the presence/absence of erosions/cavitations/geodes (ECG) or ECG plus fissures/well-defined pits at the medial epicondyle, in the site of attachment of the anterior fascia of the ulnar collateral ligament. The boxes indicate the standard error, the whiskers indicate $1.96 \times SE$. A) ECG presence/absence, pooled Ligurian Neolithic sample, pooled sexes; B) ECG presence/absence, SMP sample, pooled sexes; C) ECG presence/absence, SMP sample, males; D) ECG presence/absence, SMP sample, females; E) ECG+FP presence/absence, pooled Ligurian Neolithic sample, pooled sexes; F) ECG+FP

presence/absence, SMP sample, pooled sexes; G) ECG+FP presence/absence, SMP sample, males; H) ECG+FP presence/absence, SMP sample, females.

Figure 8 – Boîtes à moustaches montrant les différences d’asymétrie humérale du moment polaire de l’aire (J) à mi distal de diaphyse suivant la présence ou l’absence d’érosions/cavité/géodes (ECG) ou d’ECG associées à des fissures ou des dépressions au niveau de l’épicondyle médial, dans la zone d’insertion du fascia antérieur du ligament collatéral médial. La boîte indique l’erreur standard et les moustaches montrent $1,96 * l’$ erreur standard. A) présence ou absence d’ECG, échantillons néolithiques et sexes confondus ; B) présence ou absence d’ECG pour l’échantillon SMP, sexes confondus ; C) présence ou absence d’ECG pour l’échantillon SMP, hommes ; D) présence ou absence d’ECG pour l’échantillon SMP, femmes ; E) présence ou absence d’ECG+FP, échantillons néolithiques et sexes confondus ; F) présence ou absence d’ECG+FP pour l’échantillon SMP, sexes confondus ; G) présence ou absence d’ECG+FP pour l’échantillon SMP, hommes ; H) présence ou absence d’ECG+FP pour l’échantillon SMP, femmes.




Figure 9 – Scatterplot of right humeral torsional rigidity (J) on left humeral torsional rigidity of the Ligurian Neolithic skeletal series, categorized on the basis of presence/absence and side of occurrence of A) ECG; B) ECG+FP; C) any change in the ME (UCL or CFT), considering only individuals for which both regions could be scored; D) any change in the ME (UCL or CFT),

considering also individuals for which only one region could be scored (see methods). The solid line indicates isometry. AA (black solid circles): left: trait absent, right: trait absent; OA (black circles) left: site not observable, right: trait absent; A0 (black squares) left: trait absent, right: site not observable; OP (red squares) left: not observable, right: trait present; PP (solid red triangles): left: trait present, right: trait present; AP (red triangles): left: trait absent, right: trait present; PA (red diamonds) left: trait present, right: trait absent; P0 (solid red diamonds) left: trait present, right: not observable.

Figure 9 – Diagramme bivarié du moment polaire de l'aire (J) à mi distal de diaphyse du côté gauche par rapport au côté droit pour les collections néolithiques, catégorisées sur la base de la présence ou de l'absence et de la latéralité de A) ECG ; B) ECG+FP ; C) tous les changements au niveau du ME (UCL ou CFT) en ne considérant seulement que les individus pour lesquels les deux zones peuvent être enregistrées ; D) tous les changements au niveau du ME (UCL ou CFT) en considérant tous les individus (voir la section méthode). La ligne indique l'isométrie. AA (cercles noirs pleins) : trait absent à gauche, trait absent à droite ; OA (cercles noirs) : trait non observable à gauche, absent à droite ; A0 (carrés noirs) : trait absent à gauche, non observable à droite ; OP (carrés rouges) : : trait non observable à gauche, présent à droite ; PP (triangles rouges pleins) : trait présent à gauche ; trait présent à droite ; AP (triangles rouges) : trait absent à gauche ; présent à droite ; PA (losanges rouges) : trait présent à gauche ; absent à droite ; P0 (losanges rouges pleins) : trait présent à gauche et non observable à droite.




Figure 10 – Slight (score 1) bilateral external auditory canal exostosis (EAE) in the individual n°1 from Arma dell'Aquila, excavations Richard. A: left side; B: right side.

Figure 10. Exostose du canal auditif externe bilaterale légère (score 1) chez l'individu n°1 d'Arma dell' Aquila, fouilles Richard. A : côté gauche ; B : côté droit.