



**HAL**  
open science

## **The raised ramparts around Titan's northern lakes: Spectral and emissivity analysis**

Anezina Solomonidou, Alice Le Gall, Michael Malaska, Samuel Birch, Rosaly M. C. Lopes, Athena, Coustenis, Sebastien Rodriguez, Stephen D. Wall, Roger J. Michaelides, Maya R. Nasr, et al.

### ► **To cite this version:**

Anezina Solomonidou, Alice Le Gall, Michael Malaska, Samuel Birch, Rosaly M. C. Lopes, et al.. The raised ramparts around Titan's northern lakes: Spectral and emissivity analysis. American Geophysical Union, Fall Meeting 2019, Dec 2019, San Francisco, United States. pp.abstract #P23D-3524. hal-03007610

**HAL Id: hal-03007610**

**<https://hal.science/hal-03007610v1>**

Submitted on 16 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## The raised ramparts around Titan's northern lakes: Spectral and emissivity analysis

A. Solomonidou<sup>1,2,3\*</sup>, A. Le Gall<sup>4</sup>, M.J. Malaska<sup>5</sup>, S.P.D. Birch<sup>6</sup>, R.M.C. Lopes<sup>5</sup>, A. Coustenis<sup>3</sup>, S. Rodriguez<sup>7</sup>, S.D. Wall<sup>5</sup>, R.J. Michaelides<sup>8</sup>, M.R. Nasr<sup>9</sup>, C. Elachi<sup>2</sup>, A.G. Hayes<sup>6</sup>, J.M. Soderblom<sup>9</sup>, A.M. Schoenfeld<sup>10</sup>, C. Matsoukas<sup>11</sup>, P. Drossart<sup>3</sup>, M.A. Janssen<sup>5</sup>, K.J. Lawrence<sup>5</sup>, O. Witasse<sup>12</sup>, J. Radebaugh<sup>13</sup>

<sup>1</sup>European Space Agency (ESA), European Space Astronomy Centre (ESAC), Villanueva de la Canada, Madrid, Spain;

<sup>2</sup>California Institute of Technology (Caltech), Pasadena, California, USA;

<sup>3</sup>LESIA - Observatoire de Paris, CNRS, UPMC Univ. Paris 06, Univ. Paris-Diderot, Meudon, France;

<sup>4</sup>LATMOS/IPSL, UVSQ Université Paris-Saclay, Sorbonne Université, CNRS, Guyancourt, France;

<sup>5</sup>Jet Propulsion Laboratory, California Institute of Technology, California, USA;

<sup>6</sup>Cornell University, Ithaca NY, USA;

<sup>7</sup>Institut de Physique du Globe de Paris (IPGP), CNRS-UMR 7154, Université Paris-Diderot, Paris, France;

<sup>8</sup>Department of Geophysics, Stanford University, Stanford, California, USA;

<sup>9</sup>Department of Earth, Atmospheric and Planetary Sciences, MIT, Cambridge, MA 02139-4307, USA;

<sup>10</sup>Department of Earth, Planetary, and Space Sciences, University of California, Los Angeles, California, USA;

<sup>11</sup>KTH-Royal Institute of Technology, Stockholm, Sweden;

<sup>12</sup>European Space Agency (ESA), European Space Research and Technology Centre (ESTEC), Noordwijk, Netherlands;

<sup>13</sup>Department of Geological Sciences, Brigham Young University, Utah, USA.

We investigate the spectral and microwave properties of the raised ramparts of five liquid-filled lakes [e.g.1;2;3] in Titan's north pole using both Cassini VIMS and RADAR data. Ramparts are radar-bright mounds that extend from the shores of some lakes out for up to tens of kilometers and are distinct features from the raised rims. Dedicated methods have been developed to extract their infrared properties from a high-resolution VIMS cube [4;5] and their 2.2-cm emissivity from radiometry observations [6;7]. Analysis of these combined and co-registered data provides new constraints for the formation mechanisms of raised ramparts that are seen around a subset of Titan's northern lakes. VIMS and emissivity results show that the raised ramparts are made of materials very similar to that of the empty lake floors, which are also part of our investigation. In addition the emissivity of the raised ramparts is close to that of Titan's labyrinths. This suggests that the ramparts, the floors of the lakes, and the labyrinth terrains are made from or are covered by a similar material. In addition, two out of the eight lakes with raised ramparts show spectral differences at three specific wavelengths, between the ramparts and the surrounding terrain. We hypothesize that this could be due to some component, or mixture of components in the ramparts that is less absorbent at these specific wavelengths, or an effect of different grain sizes. A number of theories for the formation of the raised ramparts are discussed [8].

[1] Hayes et al. (2017). GRL 44, 11.745-11.753 [2]Michaelides et al. (2016). Icarus 270, 57-66 [3] Birch et al. (2017). Icarus 282, 214-36 [4] Solomonidou et al. (2016). Icarus 270, 85-99 [5] Solomonidou et al. (2018). JGR 123, 489-507 [6] Janssen et al. (2016) Icarus, 270, 443-459 [7] Le Gall et al. (2013). AGU, P53D-1891. [8] Solomonidou et al. Icarus, in press (2019).