

The causes of replication stress and their consequences on genome stability and cell fate

Indiana Magdalou, Bernard S Lopez, Philippe Pasero, Sarah Lambert

► To cite this version:

Indiana Magdalou, Bernard S Lopez, Philippe Pasero, Sarah Lambert. The causes of replication stress and their consequences on genome stability and cell fate. Seminars in Cell & Developmental Biology, 2014, 30, pp.154-164. 10.1016/j.semcdb.2014.04.035. hal-03008982

HAL Id: hal-03008982 https://hal.science/hal-03008982

Submitted on 29 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	The causes of replication stress and their consequences
2	on genome stability and cell fate
3	Indiana Magdalou ¹ , Bernard S. Lopez ¹ , Philippe Pasero ² and Sarah AE Lambert ³ .
4	
5 6	1: Université Paris Sud, CNRS, UMR 8200 and Institut de Cancérologie Gustave Roussy, équipe labélisée « LIGUE 2014 », Villejuif, France
7 8	2: Institute of Human Genetics, CNRS UPR 1142, équipe labélisée Ligue contre le Cancer, 141 rue de la Cardonille, 34396 Montpellier, France.
9 10	3: Institut Curie, centre de recherche, CNRS UMR338, Bat 110, centre universitaire, 91405, Orsay, France.
11	
12	Correspondence to Sarah AE Lambert, <u>sarah.lambert@curie.fr</u>
13	
14	Number of words: 6832 (without references)
15	Number of figures: 2
16	
17	Key words: replication stress, cancer, senescence, genome instability, catastrophic mitosis.
18	
19	
20	
21	
22	
23	
24	

1 Abstract

Alterations of the dynamics of DNA replication cause genome instability. These alterations known as "replication stress" have emerged as a major source of genomic instability in pre-neoplasic lesions, contributing to cancer development. The concept of replication stress covers a wide variety of events that distort the temporal and spatial DNA replication program. These events have endogenous or exogenous origins and impact globally or locally on the dynamics of DNA replication. They may arise within a short window of time (acute stress) or during each S phase (chronic stress). Here, we review the known situations in which the dynamics of DNA replication is distorted. We have united them in four main categories: i) inadequate firing of replication origins (deficiency or excess), ii) obstacles to fork progression, iii) conflicts between replication and transcription and iv) DNA replication under inappropriate metabolic conditions (unbalanced DNA replication). Because the DNA replication program is a process tightly regulated by many factors, replication stress often appears as a cascade of events. A local stress may prevent the completion of DNA replication at a single locus and subsequently compromise chromosome segregation in mitosis and therefore have a global effect on genome integrity. Finally, we discuss how replication stress drives genome instability and to what extent it is relevant to cancer biology.

1 1 Introduction

2 The maintenance of genome stability is essential to the accurate transmission of genetic information. 3 It relies on the successful duplication of chromosomes, followed by their even segregation during 4 mitosis. Genome instability is not only a hallmark of cancer cells but, maybe more importantly, is also 5 a cause of genomic disorder diseases and cancer predisposition [1]. Errors occurring during DNA 6 replication can impact both on the accurate duplication of chromosomes and on their equal 7 segregation during mitosis (Figure 1) [2]. Thus, replication stress has emerged as a primary source of 8 genome instability [3]. This is clearly the case in cells overexpressing oncogenes such as Cyclin E, 9 Cdc25A, CDC6 or CDT1, which interfere with the DNA replication program and lead to expression of 10 genome instability markers [3-7]. The causes of replication stress are many and it remains 11 challenging to unify them according to their origins or their molecular bases [8]. Pioneering studies 12 on unicellular eukaryotes such as the yeasts S. cerevisiae and S. pombe have unveiled the molecular 13 events linking alterations of the replication dynamics to the expression of genome instability markers 14 [9]. Replication stress arises from endogenous (e.g. inappropriate cellular metabolic conditions) or 15 exogenous (e.g. exposure to DNA-damaging or replication-blocking) sources and acts locally (discrete 16 pause sites) and globally on replication dynamics. Here, we review the known sources of replication 17 stress and make an attempt to classify them into four categories: (i) alterations of origins firing, (ii) 18 impediments to forks progression, (iii) conflicts between the transcription and the DNA replication 19 program and (iv) DNA replication in inappropriate metabolic conditions. We also discuss to what 20 extent these causes of replication stress can jeopardize genome integrity and contribute to human 21 diseases.

22

23 2 The Eukaryotic DNA replication program

24 The replication machinery of eukaryotic cells accurately duplicates the genome and is robust enough to carry out this task in face of numerous obstacles. The DNA replication program is temporally and 25 26 spatially controlled within each individual cell. The initiation of DNA synthesis takes place at 27 thousands of replication origins from which two replication forks progress bi-directionally [10, 11]. 28 The merge of two converging forks occurs randomly within the genome [12]. DNA replication occurs 29 concurrently with many other metabolic processes, such as transcription, mRNA processing and 30 chromatin remodeling. These processes must be tightly coordinated with DNA replication to avoid 31 potential conflicts and ensure the proper transmission of genetic and epigenetic information. The 32 accuracy and robustness of DNA replication relies on key points:

DNA synthesis is highly accurate owing to the fidelity of replicative polymerases and the association
of mismatch repair (MMR) factors to replication factories [13].

Replication origins fire once and only once per cell cycle [14]. They are activated sequentially
throughout the length of the S phase. This sequential activation depends on their accessibility to
limiting initiation factors, for which origins compete [15-18]. Moreover, origins are clustered into
replication timing domains, in which origins fire stochastically [19].

7 - The genome contains a large excess of replication origins. All origins are licensed before S phase,

8 but many of them are not activated in unchallenged growth conditions. These origins are referred to

9 as cryptic or dormant origins. They are used to rescue stalled forks under replication stress

10 conditions [19-22].

The progression of replication forks is escorted by multiples factors that ensure their stability in face
to obstacles and promote their restart when needed [8].

13 Origin-poor regions, such as common fragile sites (CFSs) are particularly at risk and forks travelling

14 through these loci rely on fork-restart mechanism to complete DNA synthesis [23-25]. The same

15 applies to regions that are replicated unidirectionally (such as the ribosomal DNA locus for example),

16 or when two converging forks are simultaneously impeded [26]. Factors promoting the stability and

17 the restart of replication forks are part of the DNA Damage response (DDR) [27]. This cellular

18 response includes the DNA replication checkpoint and the DNA repair and tolerance pathways.

19 Among the DNA repair mechanisms, homologous recombination is particularly important for the

20 robustness of DNA replication [28].

21 Here, we propose that any event that alters the fulfillment of the DNA replication program can be

22 considered as a source of replication stress. These events include alteration of the initiation and

elongation of DNA replication, conflicts between DNA replication and metabolic pathways such as

24 transcription and mRNA processing, as well as inappropriate cellular metabolic conditions, which are

25 not in line with the needs for DNA replication.

26

27 **3** The cellular response to replication stress

28 A hallmark of replication stress is the accumulation of single stranded DNA (ssDNA) at replication

29 forks, even though compromised DNA replication without excess of ssDNA has also been reported

30 [29-31]. Stretches of ssDNA are exposed at replication forks as a consequence of replicative helicases

31 continuing to unwind the parental DNA while the replicative DNA polymerases are stalled [32]. This

1 uncoupling between helicase and polymerase activities is probably not the sole cause of 2 accumulation of ssDNA at stalled forks. It can be also generated by the degradation of newly-3 synthesized DNA through the combined action of nucleases and DNA helicases, such as MRN 4 (MRE11, RAD50, NBS1), S. cerevisiae Sae2 and its human homologue CtIP, EXO1 and the BLM 5 (BLOOM Syndrome Mutated) [33-36]. These enzymes are best known for their role in the resection 6 of DNA ends at DSBs (Double Strand Break), but could also act at stalled forks [37]. Primed ssDNA, 7 together with ssDNA coated by the protein RPA (Replication Protein A), act as a signaling platform to 8 recruit numerous factors involved in the cell response to replication stress and in fine activate the 9 central ATR kinase (Ataxia-Telangectasia mutated and Rad3 related) [38]. ATR orchestrates the 10 cellular response to replication stress by phosphorylating several targets that control the stability of 11 replication forks, the firing of replication origins, mRNA metabolism and cell cycle progression. 12 Altogether, the ATR pathway limits genomic instability during S phase and contributes to cell survival 13 in response to replication stress [39]. In neoplasic lesions, this pathway acts as a barrier to both

14 genomic instability and uncontrolled proliferation [3].

15 Two main functions of ATR can be distinguished: i) stabilization of stalled replication forks (local 16 effect) and *ii*) cell-cycle arrest and repression of late origins (global effects) (Figure 2). At the global 17 level, ATR ensures an optimal management of the resources essential to fork progression, such as 18 dNTP pool and histone supply [39, 40]. When dNTP levels run down, the ATR pathway prevents the 19 activation of late-firing origins but allows the activation of dormant origins to complete DNA 20 synthesis at early domains [19, 41-43]. The local activities of ATR at replication forks remain elusive 21 at the molecular level. Studies in yeast have established that ATR maintains the replisome in a 22 replication-competent state by phosphorylating some of its components [44-50]. Whether similar 23 mechanisms operate in mammalian cells is currently unclear [34, 51]. An emerging target of ATR at 24 mammalian replication forks is the SMARCAL1 helicase involved in the remodeling of stalled forks 25 [52-54]. Inhibition of ATR during an acute replication stress results in fork collapse, associated with 26 the accumulation of ssDNA and DSBs at replication forks (Figure 2). ATR phosphorylates SMARCAL1 27 to prevent fork remodeling and thus fork collapse [34]. Defects in SMARCAL1 cause the human 28 diseases Schimke immunoosseous dysplasia, characterized by growth retardation, immunodeficiency 29 and cancer predisposition [55, 56].

30 An early function of the DDR at replication stress sites is the phosphorylation of the histone H2A in

31 yeasts and H2AX in mammals (named γ -H2A and γ -H2AX, respectively) [57]. In mammalian cells, γ -

- 32 H2AX results from the activities of three kinases: ATM (<u>Ataxia-Telangectasia</u> <u>M</u>utated), ATR and
- 33 DNA-PK (DNA-dependent Protein Kinase) and thus reflects a local activation of the DDR within the
- 34 nucleus. γ-H2AX is frequently used as a replication stress marker, together with the formation of

53BP1 foci, another component of the DDR [58-60]. However, both γ-H2AX and 53BP1 foci also occur
 at DNA damage, in particular DSBs, in a replication-independent manner [57].

3 Other ATR-dependent phosphorylation events, such as the phosphorylation of RPA on Serine 33 and 4 the phosphorylation of the checkpoint kinase Chk1 on serine 345, are also used as markers of 5 replication stress [31]. However, it should be noted that certain types of replication stress remain 6 undetected by the ATR pathway because they do not generate enough ssDNA at forks, either 7 because they affect a small number of forks or because uncoupling between helicases and 8 polymerases does not occur [18, 38, 61, 62]. Replication stress activates the ATR pathway in a 9 stepwise manner. Sensors and mediators are first recruited to forks to control their stability. Then, 10 this local checkpoint signal is amplified through the activation of the Chk1 kinase to delay mitosis [5, 11 61]. Thus, localized alterations of fork progression may escape cell cycle surveillance and jeopardize 12 the stability of the genome.

13 Impediment to fork progression leads to ssDNA gaps and to replication-associated DSBs [63-65]. The 14 origin of these DSBs remains elusive. They might form as a consequence of fork passage through a 15 nick in the parental DNA or as a consequence of enzymatic processing of gapped replication forks 16 [66-69]. In some cases, genome-wide alteration of replication dynamic leads to fork reversion, a 4-17 branched DNA structure in which parental DNA strands are annealed. Genetic data in bacteria are 18 indicative of the existence of such fork structures that have been visualized in yeast and mammalian 19 cells by electronic microscopy (EM) [50, 65, 70]. Fork reversion has also been identified as a major 20 consequence of the topoisomerase 1 inhibitor campthotecin (CPT) on replication dynamics [71]. Fork 21 reversion reflects the activities of replication-restart pathways rather than the formation of 22 pathological intermediates from which DNA synthesis cannot resume [72]. Overexpression of the 23 Cyclin E oncogene leads to the expression of replication stress markers, including slow-down of fork 24 progression, ATR activation, fork reversion and massive chromosome breakages [73]. Remarkably, 25 fork reversion precedes DSB formation, which results from the Mus81-dependent cleavage of 26 reversed forks. Mus81 activity is normally restricted to late G2 or mitosis and conversion of reversed 27 forks into DSBs is a consequence of premature entry of into mitosis [73, 74]. Thus, the concomitant 28 inactivation of the DDR accelerates chromosomes breakages induced by replication stress. 29 Alternatively, the accumulation of ssDNA at replication forks might sensitize them to endonucleases, 30 thus resulting in the formation of replication-associated DSBs.

31

4 The causes of replication stress

Each time a cell enter S phase, it must replicate a large number of loci that represent a challenge to 2 3 fork passage and thus a potential source of local replication stress. Some parts of the genome are 4 also more sensitive to global replication stress of endogenous or exogenous origin. These loci are collectively referred to as "intrinsically-difficult to replicate" sites. In mammals, paradigms for 5 6 difficult to replicate sites are found at CFSs, which are defined as loci that accumulate beaks or gaps, 7 when condensed in response to mild replication stress [75, 76]. CFSs are usually replicated late and 8 thought to remain unreplicated when cells enter mitosis [2]. The molecular bases of CFSs fragility 9 include a defect in replication origins and the presence of non-canonical DNA structures or very large 10 genes [23-25, 77, 78]. The past decade has seen the term "fragile site" used as a generic term to refer to "difficult-to-replicate" loci in many organisms. However, distinct mechanisms are likely underlying 11 12 the genetic instability of these fragile sites, even within a given organism.

13

14 4-1 inappropriate firing of replication origins: excess and deficiency

15 The DNA replication program ensures that origins fire sequentially throughout the length of the S 16 phase. To maintain genome stability during DNA replication, it is critical that: i) a sufficient amount of origins fires to complete DNA replication, *ii*) origins fire in a sequential order and not all at the same 17 time and *iii*) origins fires once and only once per S phase. The sequential firing of replication origins is 18 19 orchestrated by a limited amount of initiation factors and by the availability of dNTP [8, 16, 79]. Both 20 in mammals and yeast, the replication fork speed in a key determinant of the temporal DNA 21 replication program [79-81]. The rates of initiation and elongation are modulated by the level of 22 dNTPs. When dNTP levels are low, replication forks are slower and more origins fire to ensure the 23 completion of DNA replication. If dNTP pools drop below a critical level, ssDNA accumulates at 24 stressed forks, the DNA replication checkpoint is activated and interrupts the replication program 25 [79]. In all eukaryotes, the genome contains an excess of origins to buffer the consequences of fork 26 stalling [10]. As forks slow down, dormant origins fire to offset the completion of DNA replication. In 27 mammals, this compensatory mechanism has been reported in many situations in which DNA 28 replication is challenged and contributes to cell resistance to replication stress [21, 22, 82, 83]. 29 Alterations of the dNTP pools are associated with increased level of genome instability and cancer 30 development during DNA replication [84, 85].

Replication origins are not equally distributed along the genome and their efficiency varies widely.
 Some CFSs are origin-poor over megabases of DNA and are thus unable to compensate for slower

1 forks by activating dormant origins [23-25]. In response to a mild replication stress, duplication of 2 CFSs relies on the ability of slow-progressing forks to synthetize DNA over a long distance. Cells are 3 thus prone to enter mitosis with unreplicated chromosomal segments [2]. As the DNA replication 4 program is cell-type specific, the fragility of CFSs is also cell-type specific [23]. Other CFSs contains 5 elements that can impinge on fork progression [78]. Most CFSs overlap with very long gene (>300 kb) 6 whose transcription might favor collisions with the replication machinery [77]. While the initial cause 7 of replication stress may differ from one CFS to another, the lack of efficient origins could be a 8 common reason of fragility. Importantly, 50 % of recurrent deletions observed in cancer cells 9 originate from CFSs associated with large genes [86].

10 Importantly, an excessive firing of origins is also deleterious to cell viability. In budding yeast,

11 overexpression of limiting initiation factors allows the simultaneously firing of most origins in early S-

12 phase [16, 18]. As a consequence, the pool of dNTP becomes rapidly limiting, with deleterious

13 consequences for cell viability. An excess of origins firing could also consume other limiting

14 replication factors and ultimately destabilize replication forks (Figure 2). Upon replication stress and

15 in the absence of the main checkpoint kinase ATR, a large fraction of replicons are activated and

16 exhibit an increased level of ssDNA covered by RPA at replication forks [34]. As the number of active

17 origins increases within a cell, the amount of RPA becomes limiting, ssDNA remains unprotected and

forks are more susceptible to breakage [87]. The overexpression of RPA appears sufficient to prevent
such fork breakage. Similarly, re-replication of chromosomal segments due to deregulated origins

20 firing also results in fork reversal, massive recruitment of RPA at ssDNA gaps, which are then

21 converted into DSBs during the next round of replication [88]. Many factors involved in origin

22 licensing are overexpressed in cancer cells and re-replication could contribute to cancer progression

23 [89].

24 While the execution of the temporal DNA replication program relies on a limited amount of

25 replication resources, genomes have also evolved an excess of origins and replication factors to

26 buffer replication stress. An excess of origins firing is a factor of acute replication stress by exhausting

27 critical factors that contribute to fork integrity.

28

29 4-2 Obstacles to fork progression

30 A wide variety of obstacles can impede on the progression of replication forks, either by altering the

31 activity of the replicative helicase or the ability of replicative polymerase to incorporate nucleotides.

32 These obstacles include DNA lesions (abasic sites, damaged bases, inter or intra-crosslinks...), DNA-

1 protein complexes, DNA sequences prone to form secondary DNA structures and highly-transcribed 2 genes [8]. These impediments to fork progression are often referred to as RFBs (Replication Fork 3 Barriers). Some of them result from exposure to exogenous genotoxic agents, including anti-cancer 4 chemotherapy. Other RFBs known as natural pause sites or intrinsic RFBs are caused by normal 5 cellular events and occur during each S phase. For example, discrete fork arrests occur at 6 centromeres, telomeres and tRNA genes in yeast. Natural RFBs can also be programmed to avoid 7 collisions between the replication and transcription machineries or to facilitate sexual differentiation 8 in yeast [8]. The biology of intrinsic RFBs has been mainly explored in yeast models but their 9 frequency and importance in mammals remain poorly evaluated. However, a recent study has 10 revealed the existence of specific loci suffering from recurrent replication stress. These loci are mainly replicated early, are enriched for highly-expressed genes, repetitive elements and CpG 11 12 dinucleotide. Importantly, these loci are also hot spots for breaks and genome instability in an ATR-13 dependent manner [90]. The chromatin status has been also proposed to be a source of impediment 14 to fork progression. The histone mark γ -H2AX is enriched at intrinsic RFBs, but also at repressed 15 genes associated with hypo-acetylated histones [58]. Also, the fission yeast heterochromatin at 16 centromeres and telomeres is enriched for γ -H2AX, supporting a link between hypo-acetylated 17 chromatin and replication stress [59].

18 The activity of RFBs and their impact on genome stability are strongly dependent on replication fork 19 direction. In mammals, the DNA replication program is cell-type specific [10]. A given locus may thus 20 be replicated in one direction in one cell type and in the opposite direction in another cell type. 21 Therefore, the activity of some RFBs and their impact on genome stability may be cell-type specific. 22 For example, the expansion of CGG repeats in the FRM1 causes the Fragile X syndrome. In embryonic 23 cells, fork stalling occurs at CGG repeats, and cells that are unable to fire origins upstream CGG 24 repeats show repeat instability. It appears that CGG expansion occurs preferentially when the FRM1 25 gene is replicated such that the CGG strand serves as the lagging strand template [91]. Thus, a 26 developmental switch in the direction of replication fork occurring during cell differentiation may 27 account for the instability of repeated sequences.

The bypass of RFBs requires additional factors than canonical replication components, such as specialized helicases in charge to remove proteins tightly bound to DNA or secondary DNA structures [92]. Also, the bypass of DNA lesions involves additional DNA polymerases activities such as TLS (<u>Trans-Lesions Synthesis</u>) polymerases and PRR (<u>Post-Replication Repair</u>) pathways [93]. An emerging concept in the field of DNA replication is that all genome segments may not be equal in term of factors required for their duplication. The successful duplication of "difficult-to-replicate" loci could be more dependent on accessory factors than other loci and thus components of the replication
 machinery might differ according to the replicated locus.

The implication of natural RFBs in diseases is well established for at-risk motifs that can form secondary DNA structures such as tri-nucleotide repeats whose expansion underline many human diseases [94]. While many of the natural RFBs described so far are associated with hot spots of genome rearrangements and recombination, their potential contributions to cancer progression is not yet clearly established [9]. However, cancer cells may be particularly sensitive to natural RFBs because they try to duplicate their genome in inappropriate metabolic conditions (see below).

9

10 **4-3** Interference between the transcription and the replication programs

11 Transcription represents a major source of replication stress, both in eukaryotes and prokaryotes 12 [95]. This is due to the fact that DNA replication and transcription machineries share the same DNA 13 template and are therefore prone to collisions. Bacterial genomes are generally replicated from a 14 single origin and genes are organized collinearly with replication to prevent frontal collisions [96]. 15 The situation is more complex in eukaryotes due to the presence of multiple replication origins. To 16 limit interference between these processes, DNA replication and transcription are physically and 17 temporally separated within the nucleus [97]. Yet, a large body of evidence indicates that 18 transcription interferes with DNA replication and induces genomic instability in eukaryotes [31, 95]. 19 The mechanism by which transcription impedes fork progression is currently unclear. One likely 20 possibility is that replication stress is caused by RNA/DNA hybrids that form during transcription 21 when nascent RNA anneals to the template DNA strand, leaving the non-template strand unpaired 22 [98]. Formation of these so-called R-loops is favored by GC-rich DNA sequences and increases when 23 assembly of mRNA-particle complexes (mRNPs) is impaired [99, 100]. R-loops are removed by 24 RNaseH and specialized helicases such as Senataxin [101-103]. Depletion of factors that displace R-25 loops or prevent their formation induces genomic instability in yeast and in higher eukaryotes [83, 26 98, 104, 105]. Another source of replication stress is associated to the accumulation of positive 27 supercoiling when transcription and replication converge [83, 106]. In yeast, this is particularly the 28 case of genes physically tethered to the nuclear envelope during mRNA export [107]. In mammalian 29 cells, very large genes are also particularly at risk as they are transcribed throughout the cell cycle 30 [77]. Many of these very large genes are replicated late in S phase and overlap with CFSs. However, 31 their fragility depends on their replication profile and varies widely between cell types [24]. 32 Remarkably, transcription also interferes with DNA replication at early-replicating sequences and a 33 recent study indicates that chromosome breaks occur preferentially at cancer genes under 34 replication stress, for a reason that is still unclear [90, 108]. Altogether, these data indicate that

1 replication and transcription must be tightly coordinated during S phase to prevent genomic

2 instability. Since both processes are perturbed during oncogene-induced tumorigenesis, it is

3 tempting to speculate that replication/transcription interference represents a major source of

4 genomic instability in precancerous lesions. This view is supported by a recent study indicating that

- 5 conflicts between replication and transcription increase in cells overexpressing Cyclin E [109].
- 6

7 4-4 Replication in inappropriate metabolic conditions (unbalanced replication)

8 Accurate chromosome duplication requires adequate resources for DNA replication. As mentioned 9 above, the depletion of dNTP pools results in slow fork progression. While this situation can be 10 artificially induced using replication inhibitors such as hydroxyurea, unbalanced dNTP pools are also associated with genome instability and cancer predisposition in human cells. The expression of HPV-11 12 16 E6/E7 (Human Papillomavirus) proteins forces cells to initiate DNA replication by the aberrant 13 activation of the Rb-E2F pathway. This is accompanied with the expression of replication stress 14 markers (slower forks, firing of dormant origins, chromosome instability and breaks) whose 15 expression is largely due to an insufficient dNTP pools [84]. Such type of replication stress refers to 16 unbalanced DNA replication, a situation in which cells initiate replication in inappropriate metabolic 17 conditions. Similarly, overexpression of the Cyclin E oncogene leads to fork slow down due to an 18 inappropriate dNTP levels. To escape unbalanced replication, cells forced to proliferate upregulate 19 genes involved in nucleotide biosynthesis, including c-myc.

20 The Bloom syndrome, which exhibits a strong association between genome instability and 21 predisposition to all type of cancers, is caused by a mutation in the BLM, a member of the RecQ 22 family of DNA helicases. Patient cells exhibit replication stress markers, including slowed forks and 23 higher number of activated origins [110]. Although these data have been interpreted as a direct 24 involvement of BLM in replication dynamics, patient cells suffer from replication stress due to 25 pyrimidine pool disequilibrium, a consequence of the down-regulation of the CDA gene (Cytidine 26 Deaminase) in BLM-deficient cells [85]. While unbalanced dNTP pools cause replication stress and 27 genome instability, to what extent they are also a general cause of tumor development remains to be 28 determined.

DNA replication also requires a large supply in histones. Parental nucleosomes are disrupted ahead
 of replication forks and restored, together with the deposition of newly-synthetized histones, onto
 daughter strands [111]. Defects in chromatin assembly during DNA replication compromise the
 transmission of epigenetic marks. More surprisingly, it also affects replication dynamics. In the yeast

1 S. cerevisiae, DDR is required for degradation of excess histones, which would otherwise compromise 2 genome stability [40]. Defects in replication-coupled chromatin assembly destabilize moving 3 replication forks that ultimately collapse [112, 113]. In mammals, the supply of newly-synthetized 4 histones regulates the rate of fork elongation, presumably by interfering with the recycling of the 5 replication factor PCNA (Proliferating Cell Nuclear Antigen) [30]. Surprisingly, transient shortage in 6 newly-synthesized histones slows down fork progression without activating of the DDR or firing 7 additional origins, probably because the stretches of ssDNA formed at the forks are too short to 8 activate the ATR pathway. However, prolonged defect in histone deposition at the fork leads to DNA 9 damage. Two developmental human diseases (Wolf-Hirschhorn syndrome and congenital 10 dyserythropoietic anemia type I) are associated with a defective histone supply, but the contribution 11 of such type of replication stress to cancer biology remains unexplored [114, 115].

12

13 5 Consequences of replication stress on genome stability and cell fate

14 Replication stress compromises the completion of chromosomes duplication and impacts on 15 chromosomes segregation at mitosis (figure 1). Incomplete DNA replication, unresolved DNA repair 16 intermediates and intertwined sister chromatid can therefore lead to chromosome breakage due to 17 chromosome non-disjunction. As mentioned above, one of the earliest consequences of replication 18 stress is the formation of RPA-coated ssDNA, which in turn activates the ATR pathway. This pathway 19 ensures that replication forks are able to restart once the source of replication stress is removed. 20 Alternative pathways are required to promote fork recovery at sites of replication stress when the 21 ATR pathway fails to maintain forks in a replication-competent state. The activation of dormant 22 origins is a particularly effective mechanism to complete chromosome duplication both in response 23 to local and global replication stress [19]. However, in regions of the genome that are origin-poor (i.e 24 CFSs) or are replicated unidirectionally (i.e. telomeres), additional fork restart mechanisms are 25 needed to complete DNA replication. Some of these mechanisms are responsible for inheritable 26 genome modifications, from mutations to chromosomal rearrangements.

27

28 5-1 Uneven mitotic chromosome segregation: from a local initiator event to genome-wide effects

29 Cells reaching mitosis with damaged and/or incompletely-replicated chromosomes

As mentioned above, fork slowdown is generally compensated by the firing of dormant replication origins. However, this mechanism does not operate at origin-poor regions of the genome such as at CFSs, leading to incomplete replication [20, 23, 25]. CFSs exhibit unresolved DNA

1 repair/recombination intermediates or intertwined sister chromatid at the end of the S phase. This 2 leads to the formation of atypical chromosome segregation events, such as anaphase bridges (UFB) 3 [116, 117]. The fact that CFSs form anaphase bridges in mitosis indicates that partially-replicated 4 chromosomes remain undetected by checkpoint pathways before cells undergo anaphase [60, 118, 5 119]. This phenomenon could also arise in response to endogenous or low replication stress that 6 escapes cell survey. For example, the phosphorylation of CHK1 is dose dependent and very low doses 7 of HU (such as 10 μ M) decelerate replication forks progression without inducing a significant cell 8 cycle arrest or detectable CHK1 phosphorylation [61, 62].

9 In anaphase, unresolved or unreplicated regions restrain chromosome segregation by creating a 10 physical link between the two sister chromatids, known as anaphase bridges. These structures can be 11 stained with DNA intercalating agents such as DAPI when they result from aberrant chromosomal 12 morphology [2]. In contrast, UFBs do not stain with DAPI but can be detected using immunostaining 13 against UFB-binding proteins such as the PICH or BLM helicases, RPA and some components of the FANC (Fanconi Anemia) complex [116, 117, 120] [121]. A subset of UFBs colocalizes with CFSs and is 14 15 induced by replication stress [116, 117]. Of note, whereas low levels of UFBs are detected under 16 normal growth conditions, they are very abundant in cells defective for BLM, FANC or homologous 17 recombination [117, 121-124].

18 Anaphase separation creates an increasing mechanical tension on partially-replicated chromatids, 19 which are then prone to break. Remarkably, CFSs are prone to chromosome breakage under 20 replicative stress [75, 76]. The breakage of anaphase bridges lead to the uneven segregation of 21 broken chromosome arms and thus to potential translocations. While the nature of unresolved 22 replication intermediates or intertwined sister chromatid junctions formed at unreplicated DNA 23 remains unknown, it is generally believed that these structures are cleaved by nucleases such as 24 MUS81-EME1, GEN1 or dissolved by BLM and topoisomerase III alpha [125-128]. It has been 25 proposed that a failure to cleave unresolved sister chromatids leads to chromosome missegregation 26 [2]. Thus, nuclease attacks of unusual replication intermediates persisting after S phase may not be 27 systematically detrimental to genome stability, but may rather protect the genome from the 28 deleterious effects of replication stress.

29

30 Genome-wide impact of chromosome segregation defects

Replication stress impedes chromosome segregation locally through the formation of anaphase bridges. Besides this local effect, low levels of replication stress can also lead to the formation of extra centrosomes at mitosis [62]. The mechanisms linking S-phase progression and centrosomes duplication have only started to emerge and it remains unclear how chronic replication stress
promotes centrosomes amplification [129]. Since most of mitotic extra centrosomes are functional,
their presence results in multipolar mitosis and unbalanced chromosome segregation [62].
Importantly, extra centrosomes are systematically associated with chromosomes bridges, delayed
chromosome condensation and with prolonged metaphase arrest [130].

Homologous recombination (HR) plays a pivotal role in DNA repair and in replication restart,
contributing therefore to the accuracy of DNA replication [28]. Defect in HR is also a factor of cancer
predisposition. Remarkably, HR-deficient cells display both a genome-wide decrease in replication
fork speed, an increased frequency of extra centrosomes and UFBs, suggesting that HR defect result
in an endogenous replication stress [62, 82, 122, 123, 131]. In support to this, mitotic defects
observed in HR-deficient cells result from slowed fork progression, suggesting that the links between
replication stress and mitotic defect potential underline cancer development [62].

13 Mitotic defects induced by replication stress: an "adaptation-like" process in mammalian cells?

14 While incomplete DNA replication at specific loci is a checkpoint blind event, mitotic defects due to 15 anaphase bridges activate checkpoints and result in prolonged mitotic arrest [62, 132]. The capacity 16 to bypass the mitotic arrest without resolving its cause has been described in yeast as a process 17 called adaptation [133, 134]. We propose that anaphase bridges are revealed in cells bypassing the 18 mitotic arrest, in a process that is reminiscent to the adaptation process described in yeast. It is 19 tempting to speculate that during the mitotic arrest bypass, opposite forces could be transmitted to 20 the centrosomes in an attempt to force segregation, leading to the splitting of centrioles and 21 multipolar segregations. This model is consistent with the facts that (i) extra centrosomes are 22 detected in mitosis but not in interphase, (ii) extra centrosomes are functional, causing multipolar 23 segregations and (ii) almost all mutipolar segregations are associated with chromatin bridges and 24 with prolonged metaphase arrest [62]. Together, these data indicate that incomplete replication 25 results in local chromosome abnormalities at mitosis such as UFBs fragile sites and that the 26 formation of extra centrosomes amplifies the defects to all the genome by generating multipolar 27 attachments and aberrant segregation of fully-replicated chromosomes.

28

29 **5-2** Inheritable genome modifications

30 Replication stress results in inheritable genome modifications, including point mutations and larger

31 chromosomal rearrangements. Some of these rearrangements are particularly complex, including

32 triplication, deletion, inversion and translocation. In cancer cells, the precise mechanisms responsible

for replication stress-induced genome instability remain unclear but they can be determined from
 the sequence of breakpoint junctions [9].

3 Both stretches of ssDNA at stalled forks and replication-associated DSBs are likely initiator events of 4 replication-induced genome instability. When forks collapse and eventually break, replication-restart 5 mechanisms occur and faulty replication restart events are causal mechanisms of genome 6 modifications and chromosome rearrangements. Among the pathways ensuring the continuity of 7 replication, HR is an evolutionary conserved mechanisms involved in DSB repair and replication-8 restart [28]. The purpose of HR is to recombine homologous sequences, usually at sister chromatids, 9 to repair DNA lesions. Occasionally, HR recombines repeated sequences dispersed through the 10 genome and such events refer to NAHR (Non-Allelic Homologous Recombination), a cause of genome 11 rearrangements exhibiting homology at breakpoint junctions [1]. In fission yeast, a single fork arrest 12 is sufficient to induce translocation and genomic deletion in a recombination-dependent manner 13 [135]. Also, slowdown in replication fork progression stimulates fusion between inverted repeats by 14 a recombination pathway in mammals [136]. Replication-borne DSBs can be repaired by BIR (Break-Induced Replication), a type of HR event that can drive translocation in yeast models [137]. 15

16 Single mutations are by-products of TLS-DNA polymerases that bypass DNA lesions in an error-prone 17 manner. The bypass of DNA lesions can occur either at the fork or behind the fork. Indeed, it has be 18 proposed that DNA synthesis can be re-primed downstream the lesion, leaving a gap in the rear of 19 the moving replisome that is then fill-in by either TLS or HR [138-140]. Clustered mutations 20 surrounding the break-points of somatic rearrangements have been also recently reported in cancer 21 cells; such clustered base substitutions refer to as Kataegis [141]. Long stretches of ssDNA generated 22 during the DSB or fork repair are particularly sensitive to DNA damage, in particular to the activities 23 of cytidine deaminase family [142]. In yeast, chronic replication stress induced by exposure to 24 replication-specific damaging agents leads to damaged ssDNA and clustered mutation over hundreds 25 of kilobases [143].

26 Error-prone DNA synthesis associated with repair events likely underlies genome modifications. HR 27 requires a DNA synthesis step that is error-prone, mainly due to a lack of processivity: the elongated 28 strand dissociates frequently from the initial template and anneals to another discontinuous 29 template [144-146]. This phenomenon is called template switch and is driven either by significant 30 length of homology or micro-homology (few bases) [147]. The last refers to as MM-BIR (microhomology mediated break-induced replication) [146, 148]. In fission yeast, replication-restart by HR 31 32 is associated with an error prone DNA synthesis, liable to replication slippage at microhomologies, 33 thus leading to micro-deletions and insertions [135]. When progressing across repeated sequences,

1 forks restarted by HR are prone to fold-back inversion, resulting in large chromosome duplication, a 2 possible mechanism to explain fold back inversions observed in pancreatic cancer cells [149, 150]. 3 DNA synthesis associated with BIR events is particularly error prone, resulting in a 2,800 fold increase 4 in frame-shift, base substitution and template switch when progressing over hundreds of kilobases 5 [144, 151]. Remarkably, overexpression of Cyclin E results in genomic duplication with complex 6 breakpoint junctions [152]. Fork progression relies on two key subunits of the DNA polymerase delta 7 that appear dispensable to replication efficiency in cells not overexpressing Cyclin E. The use of 8 reporter assays demonstrated that BIR events are likely to underlie genomic duplication due to serial 9 template switches of the elongated strand [152]. Thus, under replication stress conditions, the 10 continuity of DNA replication relies on accessory DNA polymerase factors that drive genome instability. 11

CNVs (<u>Copy Number Variation</u>) are the most frequent genome modifications observed from an
 individual to another. CNVs are also common to cancer cells and underlie genomic disorders [1,
 153]. CNVs are induced by replication stress and independently of the DSB repair pathway NHEJ (<u>Non</u>
 <u>Homologous End Joing</u>) [153, 154]. The analysis of breakpoint junctions of CNVs has revealed an
 unexpected complexity, including the presence of microhomologies, small insertions/deletions,
 frame-shifts and base substitutions [155]. Such a complexity has been interpreted as a consequence
 of multiple attempts to restart stalled replication forks.

19

20 **5-3 Cell death, senescence and cancer.**

21 One century ago, Theodor Boveri hypothesized that tumors originate from improper chromosome 22 segregation, thus creating an uploid cells that undergo clonal expansion. He proposed that in some 23 cases, abnormal chromosome segregation and aneuploidy are caused by extra-centrosome 24 generation, leading to multipolar cells [156]. More recently, the formation of active mitotic extra 25 centrosomes has been correlated to replication stress [62]. Consistently, both replication stress and centrosome abnormalities have also been reported at early stages of malignancy [4, 157-159]. 26 27 Replication stress provokes an arrest at mitosis through the activation of the mitotic spindle 28 checkpoint. Strikingly, even low or unprocessed endogenous replication stress results in prolonged 29 metaphase arrest. Prolonged mitotic arrest should then lead to mitotic catastrophe and cell death. 30 These processes could therefore select cells able to escape mitotic arrest and cell death, thus with 31 increased genetic instability (see above) and possibly with de-repressed proliferation program.

1 Endogenous replication stress has been proposed to be involved at the very early steps of 2 senescence and/or malignancy initiation [3, 4, 7, 159, 160]. Beside the induction of tumors, 3 oncogenes can induce senescence in a process called oncogene-induced senescence (OIS). It is 4 proposed that the oncogene activation leads to an hyper-replication state through the stimulation of the cell proliferation program. The hyper-replication increases the endogenous replication stress, 5 6 which activates the DDR. The persistence of the DDR arrests cells in cell cycle, causing senescence [3, 7 161]. Note that senescence is generally considered as a tumor suppressor mechanism through its 8 anti-proliferation consequences. However, senescence is a double-edge sword since it can also 9 induce tumorigenesis through the production of inflammatory cytokines [162, 163]. In addition, the 10 induction of senescence allows to select cells able to bypass the proliferation restriction, thus with 11 stimulated proliferation programs, thus constituting a first step toward neoplasia.

12 The link between replication stress and senescence is supported by several studies. Patients suffering 13 from the Werner syndrome (in which a RecQ-helicase is mutated) or from Fanconi anemia, both 14 syndrome that exhibit markers of endogenous replication stress, also suffer from premature aging 15 and patient cells are prone to senescence [164, 165]. Caloric restriction, which prolongs life span in 16 all eukaryotes, also inhibits DNA replication [166, 167]. Exposure of cells to low doses of HU for two 17 weeks, induce senescence monitored by the lysosomial β -galactosidase activity and the poor ability 18 to incorporate nucleotide analogues [160]. However, the senescence-associated heterochromatin 19 foci (SAHF) were not efficiently induced compared with OIS. Note that OIS-induced SAHFs require 20 DNA replication and ATR [160].

21 Noteworthy, HR protects against spontaneous endogenous replication stress and is affected in many 22 familial breast cancers. More specifically, most of genes identified to be inactivated in familial breast 23 cancer control the replication/recombination interface [168, 169]. In addition, the oncogene AKT1 is 24 activated in 40 to 60% of sporadic breast and ovary cancer [170, 171]. Interestingly, AKT1 stimulates 25 proliferation, represses cell death and HR and thus should generate high endogenous replication 26 stress [170]. Consistently AKT1 over-activation leads to the formation of mitotic extra centrosomes 27 [172]. Such genetic instability should then increase genetic instability favoring thus the tumor 28 progression, in addition to tumor initiation. This shed light on the importance to secure replication 29 completion and to manage replication stress (even endogenous stresses) prior to mitosis.

- 31
- 32

1 6- Concluding remarks

2 The sources of replication stress are many and varied, but different causes results in similar 3 consequences. It is likely that novel mechanisms of genome instability caused by replication stress 4 will emerge, concerning for instance oncogene-induced alterations of the DNA replication program. 5 Our knowledge of the bases of eukaryotic DNA replication program is a pillar to understand the 6 contribution of replication stress to cancer development. The cellular responses to replication stress 7 have been mainly characterized in response to acute and exogenous stress. When the DDR is 8 defective, acute replication stress leads to a massive deregulation of the DNA replication program, 9 catastrophic genome instability and senescence or cell death. Cells escaping apoptosis are prone to 10 accumulate additional mutations and to initiate clonal proliferation. In another scenario, neoplasic 11 lesions suffer from low levels of chronic and endogenous replication stress. Such type of alteration 12 might be in part checkpoint-blind and thus facilitates the transmission of damaged chromosomal 13 segments to the next cell cycle [60]. Also, adaptation processes might take place to face endogenous 14 and chronic replication stress. Tumor cells could "learn" how to cope with for arrest and duplicate 15 their genome in stress conditions. Indeed, replication stress arises in neoplasic lesions but lingers as 16 well in solid tumors in which it contributes to chromosomal instability such as an euploidy [5]. The 17 cascade of DDR activation and its components have been largely characterized in response to acute 18 replication stress, but the cell response to chronic and endogenous replication stress remains poorly 19 described, despite the fact that it likely represents the main source of replication-induced genome 20 instability during cancer development. Understanding to what extent and how cells adapt to subtle 21 replication alterations should help identify new targets and open new avenues for anticancer 22 therapeutic strategies.

- 23
- 24 25 26 27 28 29
- 30

1 Acknowledgements

2 We apologize to our colleagues whose contributions are not cited due to space constraints. Work in

3 the SAEL laboratory is supported by Agence Nationale Recherche grants ANRJCJC10-1203 01 and la

4 Ligue contre le cancer (comité Essonne). IM and BSL were supported by INCa (Institut national du

5 cancer) and by "la Ligue Nationale contre le cancer". PP thanks ANR, INCa and the Ligue contre le

- 6 Cancer for support.

1 References

- 2 [1] Liu P, Carvalho CM, Hastings PJ, Lupski JR. Mechanisms for recurrent and complex human
- 3 genomic rearrangements. Current opinion in genetics & development 2012;22:211-20.
- 4 [2] Mankouri HW, Huttner D, Hickson ID. How unfinished business from S-phase affects mitosis and
- 5 beyond. The EMBO journal 2013;32:2661-71.
- 6 [3] Halazonetis TD, Gorgoulis VG, Bartek J. An oncogene-induced DNA damage model for cancer
- 7 development. Science 2008;319:1352-5.
- 8 [4] Gorgoulis VG, Vassiliou LV, Karakaidos P, Zacharatos P, Kotsinas A, Liloglou T, et al. Activation of
- 9 the DNA damage checkpoint and genomic instability in human precancerous lesions. Nature2005;434:907-13.
- 11 [5] Burrell RA, McClelland SE, Endesfelder D, Groth P, Weller MC, Shaikh N, et al. Replication stress
- 12 links structural and numerical cancer chromosomal instability. Nature 2013;494:492-6.
- 13 [6] Liontos M, Koutsami M, Sideridou M, Evangelou K, Kletsas D, Levy B, et al. Deregulated
- overexpression of hCdt1 and hCdc6 promotes malignant behavior. Cancer research 2007;67:10899-909.
- 16 [7] Bartkova J, Rezaei N, Liontos M, Karakaidos P, Kletsas D, Issaeva N, et al. Oncogene-induced
- 17 senescence is part of the tumorigenesis barrier imposed by DNA damage checkpoints. Nature2006;444:633-7.
- [8] Lambert S, Carr AM. Impediments to replication fork movement: stabilisation, reactivation and
 genome instability. Chromosoma 2013;122:33-45.
- 21 [9] Lambert S, Carr AM. Replication stress and genome rearrangements: lessons from yeast models.
- 22 Current opinion in genetics & development 2013;23:132-9.
- [10] Mechali M. Eukaryotic DNA replication origins: many choices for appropriate answers. Nature
 reviews Molecular cell biology 2010;11:728-38.
- 25 [11] Cayrou C, Coulombe P, Mechali M. Programming DNA replication origins and chromosome
- organization. Chromosome research : an international journal on the molecular, supramolecular and
- evolutionary aspects of chromosome biology 2010;18:137-45.
- 28 [12] McGuffee SR, Smith DJ, Whitehouse I. Quantitative, genome-wide analysis of eukaryotic
- replication initiation and termination. Molecular cell 2013;50:123-35.
- 30 [13] Hombauer H, Srivatsan A, Putnam CD, Kolodner RD. Mismatch repair, but not heteroduplex
- rejection, is temporally coupled to DNA replication. Science 2011;334:1713-6.
- 32 [14] Yoshida K, Poveda A, Pasero P. Time to be versatile: regulation of the replication timing program
- in budding yeast. Journal of molecular biology 2013;425:4696-705.
- 34 [15] Yoshida;, al. e. Mol Cell 2014;under press.
- 35 [16] Mantiero D, Mackenzie A, Donaldson A, Zegerman P. Limiting replication initiation factors
- execute the temporal programme of origin firing in budding yeast. The EMBO journal 2011;30:4805-14.
- 38 [17] Collart C, Allen GE, Bradshaw CR, Smith JC, Zegerman P. Titration of four replication factors is
- 39 essential for the Xenopus laevis midblastula transition. Science 2013;341:893-6.
- 40 [18] Tanaka S, Nakato R, Katou Y, Shirahige K, Araki H. Origin association of Sld3, Sld7, and Cdc45
- 41 proteins is a key step for determination of origin-firing timing. Current biology : CB 2011;21:2055-63.
- [19] Blow JJ, Ge XQ, Jackson DA. How dormant origins promote complete genome replication. Trends
 in biochemical sciences 2011;36:405-14.
- 44 [20] Kawabata T, Luebben SW, Yamaguchi S, Ilves I, Matise I, Buske T, et al. Stalled fork rescue via
- 45 dormant replication origins in unchallenged S phase promotes proper chromosome segregation and
- tumor suppression. Molecular cell 2011;41:543-53.
- 47 [21] Ge XQ, Jackson DA, Blow JJ. Dormant origins licensed by excess Mcm2-7 are required for human
- 48 cells to survive replicative stress. Genes & development 2007;21:3331-41.

- 1 [22] Woodward AM, Gohler T, Luciani MG, Oehlmann M, Ge X, Gartner A, et al. Excess Mcm2-7
- license dormant origins of replication that can be used under conditions of replicative stress. The
 Journal of cell biology 2006;173:673-83.
- 4 [23] Letessier A, Millot GA, Koundrioukoff S, Lachages AM, Vogt N, Hansen RS, et al. Cell-type-specific
- 5 replication initiation programs set fragility of the FRA3B fragile site. Nature 2011;470:120-3.
- 6 [24] Le Tallec B, Dutrillaux B, Lachages AM, Millot GA, Brison O, Debatisse M. Molecular profiling of
- 7 common fragile sites in human fibroblasts. Nature structural & molecular biology 2011;18:1421-3.
- 8 [25] Ozeri-Galai E, Lebofsky R, Rahat A, Bester AC, Bensimon A, Kerem B. Failure of origin activation
- 9 in response to fork stalling leads to chromosomal instability at fragile sites. Molecular cell2011;43:122-31.
- [26] Murray JM, Carr AM. Smc5/6: a link between DNA repair and unidirectional replication? Nature
 reviews Molecular cell biology 2008;9:177-82.
- [27] Ciccia A, Elledge SJ. The DNA damage response: making it safe to play with knives. Molecular cell
 2010;40:179-204.
- [28] Costes A, Lambert SAE. Homologous Recombination as a Replication Fork Escort: Fork-Protectionand Recovery. Biomolecules 2013;3:39-71.
- 17 [29] Groth A, Corpet A, Cook AJ, Roche D, Bartek J, Lukas J, et al. Regulation of replication fork
- 18 progression through histone supply and demand. Science 2007;318:1928-31.
- 19 [30] Mejlvang J, Feng Y, Alabert C, Neelsen KJ, Jasencakova Z, Zhao X, et al. New histone supply
- 20 regulates replication fork speed and PCNA unloading. The Journal of cell biology 2014;204:29-43.
- 21 [31] Zeman MK, Cimprich KA. Causes and consequences of replication stress. Nature cell biology
- 22 2014;16:2-9.
- 23 [32] Byun TS, Pacek M, Yee MC, Walter JC, Cimprich KA. Functional uncoupling of MCM helicase and
- DNA polymerase activities activates the ATR-dependent checkpoint. Genes & development
 2005;19:1040-52.
- 26 [33] Grabarz A, Guirouilh-Barbat J, Barascu A, Pennarun G, Genet D, Rass E, et al. A role for BLM in
- 27 double-strand break repair pathway choice: prevention of CtIP/Mre11-mediated alternative
- 28 nonhomologous end-joining. Cell reports 2013;5:21-8.
- 29 [34] Couch FB, Bansbach CE, Driscoll R, Luzwick JW, Glick GG, Betous R, et al. ATR phosphorylates
- 30 SMARCAL1 to prevent replication fork collapse. Genes & development 2013;27:1610-23.
- 31 [35] Hashimoto Y, Ray Chaudhuri A, Lopes M, Costanzo V. Rad51 protects nascent DNA from Mre11-
- 32 dependent degradation and promotes continuous DNA synthesis. Nature structural & molecular
- 33 biology 2010;17:1305-11.
- 34 [36] Schlacher K, Christ N, Siaud N, Egashira A, Wu H, Jasin M. Double-strand break repair-
- 35 independent role for BRCA2 in blocking stalled replication fork degradation by MRE11. Cell
- 36 2011;145:529-42.
- 37 [37] Symington LS, Gautier J. Double-strand break end resection and repair pathway choice. Annual
- 38 review of genetics 2011;45:247-71.
- 39 [38] MacDougall CA, Byun TS, Van C, Yee MC, Cimprich KA. The structural determinants of checkpoint
- 40 activation. Genes & development 2007;21:898-903.
- 41 [39] Cimprich KA, Cortez D. ATR: an essential regulator of genome integrity. Nature reviews
- 42 Molecular cell biology 2008;9:616-27.
- [40] Gunjan A, Verreault A. A Rad53 kinase-dependent surveillance mechanism that regulates histone
 protein levels in S. cerevisiae. Cell 2003;115:537-49.
- 44 protein levels in S. cerevisiae. Cell 2003;115:537-49.
 45 [41] Zegerman P, Diffley JF. Checkpoint-dependent inhibition of DNA replication initiation by Sld3 and
- 46 Dbf4 phosphorylation. Nature 2010;467:474-8.
- 47 [42] Santocanale C, Sharma K, Diffley JF. Activation of dormant origins of DNA replication in budding
- 48 yeast. Genes & development 1999;13:2360-4.
- 49 [43] Crabbe L, Thomas A, Pantesco V, De Vos J, Pasero P, Lengronne A. Analysis of replication profiles
- 50 reveals key role of RFC-Ctf18 in yeast replication stress response. Nature structural & molecular
- 51 biology 2010;17:1391-7.

- 1 [44] Tercero JA, Diffley JF. Regulation of DNA replication fork progression through damaged DNA by
- 2 the Mec1/Rad53 checkpoint. Nature 2001;412:553-7.
- 3 [45] Lopes M, Cotta-Ramusino C, Pellicioli A, Liberi G, Plevani P, Muzi-Falconi M, et al. The DNA
- 4 replication checkpoint response stabilizes stalled replication forks. Nature 2001;412:557-61.
- 5 [46] De Piccoli G, Katou Y, Itoh T, Nakato R, Shirahige K, Labib K. Replisome stability at defective DNA
- 6 replication forks is independent of S phase checkpoint kinases. Molecular cell 2012;45:696-704.
- 7 [47] Katou Y, Kanoh Y, Bando M, Noguchi H, Tanaka H, Ashikari T, et al. S-phase checkpoint proteins
- 8 Tof1 and Mrc1 form a stable replication-pausing complex. Nature 2003;424:1078-83.
- 9 [48] Sabatinos SA, Green MD, Forsburg SL. Continued DNA synthesis in replication checkpoint
- 10 mutants leads to fork collapse. Molecular and cellular biology 2012;32:4986-97.
- 11 [49] Bailis JM, Luche DD, Hunter T, Forsburg SL. Minichromosome maintenance proteins interact with
- checkpoint and recombination proteins to promote s-phase genome stability. Molecular and cellular
 biology 2008;28:1724-38.
- [50] Sogo JM, Lopes M, Foiani M. Fork reversal and ssDNA accumulation at stalled replication forks
 owing to checkpoint defects. Science 2002;297:599-602.
- 16 [51] Ragland RL, Patel S, Rivard RS, Smith K, Peters AA, Bielinsky AK, et al. RNF4 and PLK1 are
- 17 required for replication fork collapse in ATR-deficient cells. Genes & development 2013;27:2259-73.
- 18 [52] Bansbach CE, Betous R, Lovejoy CA, Glick GG, Cortez D. The annealing helicase SMARCAL1
- 19 maintains genome integrity at stalled replication forks. Genes & development 2009;23:2405-14.
- 20 [53] Betous R, Mason AC, Rambo RP, Bansbach CE, Badu-Nkansah A, Sirbu BM, et al. SMARCAL1
- 21 catalyzes fork regression and Holliday junction migration to maintain genome stability during DNA
- replication. Genes & development 2012;26:151-62.
- 23 [54] Ciccia A, Bredemeyer AL, Sowa ME, Terret ME, Jallepalli PV, Harper JW, et al. The SIOD disorder
- protein SMARCAL1 is an RPA-interacting protein involved in replication fork restart. Genes &
 development 2009;23:2415-25.
- 26 [55] Baradaran-Heravi A, Raams A, Lubieniecka J, Cho KS, DeHaai KA, Basiratnia M, et al. SMARCAL1
- 27 deficiency predisposes to non-Hodgkin lymphoma and hypersensitivity to genotoxic agents in vivo.
- 28 American journal of medical genetics Part A 2012;158A:2204-13.
- 29 [56] Boerkoel CF, Takashima H, John J, Yan J, Stankiewicz P, Rosenbarker L, et al. Mutant chromatin
- 30 remodeling protein SMARCAL1 causes Schimke immuno-osseous dysplasia. Nature genetics 2002;20:215-20
- 31 2002;30:215-20.
- [57] Kinner A, Wu W, Staudt C, Iliakis G. Gamma-H2AX in recognition and signaling of DNA double strand breaks in the context of chromatin. Nucleic acids research 2008;36:5678-94.
- 34 [58] Szilard R, Jacques P, Laramée L, Cheng B, Galicia S, Bataille A, et al. Systematic identification of
- fragile sites via genome-wide location analysis of gamma-H2AX. Nat Struct Mol Biol 2010;17:299-305.
- 36 [59] Rozenzhak S, Mejia-Ramirez E, Williams JS, Schaffer L, Hammond JA, Head SR, et al. Rad3
- 37 decorates critical chromosomal domains with gammaH2A to protect genome integrity during S-Phase
- in fission yeast. PLoS genetics 2010;6:e1001032.
- 39 [60] Lukas C, Savic V, Bekker-Jensen S, Doil C, Neumann B, Pedersen RS, et al. 53BP1 nuclear bodies
- 40 form around DNA lesions generated by mitotic transmission of chromosomes under replication
- 41 stress. Nature cell biology 2011;13:243-53.
- 42 [61] Koundrioukoff S, Carignon S, Techer H, Letessier A, Brison O, Debatisse M. Stepwise activation of
- 43 the ATR signaling pathway upon increasing replication stress impacts fragile site integrity. PLoS
- 44 genetics 2013;9:e1003643.
- 45 [62] Wilhelm T, Magdalou I, Barascu A, Techer H, Debatisse M, Lopez BS. Spontaneous slow
- 46 replication fork progression elicits mitosis alterations in homologous recombination-deficient
- 47 mammalian cells. Proceedings of the National Academy of Sciences of the United States of America
- 48 2014;111:763-8.
- 49 [63] Saintigny Y, Delacote F, Vares G, Petitot F, Lambert S, Averbeck D, et al. Characterization of
- 50 homologous recombination induced by replication inhibition in mammalian cells. The EMBO journal
- 51 2001;20:3861-70.

- 1 [64] Petermann E, Orta ML, Issaeva N, Schultz N, Helleday T. Hydroxyurea-stalled replication forks
- 2 become progressively inactivated and require two different RAD51-mediated pathways for restart
- and repair. Molecular cell 2010;37:492-502.
- [65] Seigneur M, Bidnenko V, Ehrlich SD, Michel B. RuvAB acts at arrested replication forks. Cell
 1998;95:419-30.
- 6 [66] Roseaulin L, Yamada Y, Tsutsui Y, Russell P, Iwasaki H, Arcangioli B. Mus81 is essential for sister
- 7 chromatid recombination at broken replication forks. The EMBO journal 2008;27:1378-87.
- 8 [67] Moriel-Carretero M, Aguilera A. A postincision-deficient TFIIH causes replication fork breakage
- 9 and uncovers alternative Rad51- or Pol32-mediated restart mechanisms. Molecular cell 2010;37:690 701.
- 11 [68] Froget B, Blaisonneau J, Lambert S, Baldacci G. Cleavage of stalled forks by fission yeast
- 12 Mus81/Eme1 in absence of DNA replication checkpoint. Mol Biol Cell 2008;19:445-56.
- 13 [69] Fugger K, Chu WK, Haahr P, Kousholt AN, Beck H, Payne MJ, et al. FBH1 co-operates with MUS81
- in inducing DNA double-strand breaks and cell death following replication stress. Nature
 communications 2013;4:1423.
- 16 [70] Higgins NP, Kato K, Strauss B. A model for replication repair in mammalian cells. Journal of 17 molecular biology 1976;101:417-25.
- 18 [71] Ray Chaudhuri A, Hashimoto Y, Herrador R, Neelsen KJ, Fachinetti D, Bermejo R, et al.
- 19 Topoisomerase I poisoning results in PARP-mediated replication fork reversal. Nature structural &
- 20 molecular biology 2012;19:417-23.
- 21 [72] Berti M, Ray Chaudhuri A, Thangavel S, Gomathinayagam S, Kenig S, Vujanovic M, et al. Human
- RECQ1 promotes restart of replication forks reversed by DNA topoisomerase I inhibition. Nature
 structural & molecular biology 2013;20:347-54.
- 24 [73] Neelsen KJ, Zanini IM, Herrador R, Lopes M. Oncogenes induce genotoxic stress by mitotic
- 25 processing of unusual replication intermediates. The Journal of cell biology 2013;200:699-708.
- 26 [74] Matos J, Blanco MG, Maslen S, Skehel JM, West SC. Regulatory control of the resolution of DNA
- 27 recombination intermediates during meiosis and mitosis. Cell 2011;147:158-72.
- 28 [75] Durkin SG, Glover TW. Chromosome fragile sites. Annual review of genetics 2007;41:169-92.
- 29 [76] Debatisse M, Le Tallec B, Letessier A, Dutrillaux B, Brison O. Common fragile sites: mechanisms
- 30 of instability revisited. Trends in genetics : TIG 2012;28:22-32.
- 31 [77] Helmrich A, Ballarino M, Tora L. Collisions between replication and transcription complexes
- 32 cause common fragile site instability at the longest human genes. Molecular cell 2011;44:966-77.
- 33 [78] Zhang H, Freudenreich CH. An AT-rich sequence in human common fragile site FRA16D causes
- fork stalling and chromosome breakage in S. cerevisiae. Molecular cell 2007;27:367-79.
- 35 [79] Poli J, Tsaponina O, Crabbe L, Keszthelyi A, Pantesco V, Chabes A, et al. dNTP pools determine
- 36 fork progression and origin usage under replication stress. The EMBO journal 2012;31:883-94.
- 37 [80] Courbet S, Gay S, Arnoult N, Wronka G, Anglana M, Brison O, et al. Replication fork movement
- 38 sets chromatin loop size and origin choice in mammalian cells. Nature 2008;455:557-60.
- 39 [81] Anglana M, Apiou F, Bensimon A, Debatisse M. Dynamics of DNA replication in mammalian
- 40 somatic cells: nucleotide pool modulates origin choice and interorigin spacing. Cell 2003;114:385-94.
- 41 [82] Daboussi F, Courbet S, Benhamou S, Kannouche P, Zdzienicka MZ, Debatisse M, et al. A
- 42 homologous recombination defect affects replication-fork progression in mammalian cells. Journal of
- 43 cell science 2008;121:162-6.
- 44 [83] Tuduri S, Crabbe L, Conti C, Tourriere H, Holtgreve-Grez H, Jauch A, et al. Topoisomerase I
- 45 suppresses genomic instability by preventing interference between replication and transcription.
- 46 Nature cell biology 2009;11:1315-24.
- 47 [84] Bester AC, Roniger M, Oren YS, Im MM, Sarni D, Chaoat M, et al. Nucleotide deficiency promotes
- 48 genomic instability in early stages of cancer development. Cell 2011;145:435-46.
- 49 [85] Chabosseau P, Buhagiar-Labarchede G, Onclercq-Delic R, Lambert S, Debatisse M, Brison O, et al.
- 50 Pyrimidine pool imbalance induced by BLM helicase deficiency contributes to genetic instability in
- 51 Bloom syndrome. Nature communications 2011;2:368.

- 1 [86] Le Tallec B, Millot GA, Blin ME, Brison O, Dutrillaux B, Debatisse M. Common fragile site profiling
- 2 in epithelial and erythroid cells reveals that most recurrent cancer deletions lie in fragile sites hosting 3 large genes. Cell reports 2013;4:420-8.
- 4 [87] Toledo LI, Altmeyer M, Rask MB, Lukas C, Larsen DH, Povlsen LK, et al. ATR prohibits replication
- 5 catastrophe by preventing global exhaustion of RPA. Cell 2013;155:1088-103.
- 6 [88] Neelsen KJ, Zanini IM, Mijic S, Herrador R, Zellweger R, Ray Chaudhuri A, et al. Deregulated
- 7 origin licensing leads to chromosomal breaks by rereplication of a gapped DNA template. Genes & 8 development 2013;27:2537-42.
- 9 [89] Blow JJ, Gillespie PJ. Replication licensing and cancer--a fatal entanglement? Nature reviews 10 Cancer 2008;8:799-806.
- 11 [90] Barlow JH, Faryabi RB, Callen E, Wong N, Malhowski A, Chen HT, et al. Identification of early
- 12 replicating fragile sites that contribute to genome instability. Cell 2013;152:620-32.
- 13 [91] Gerhardt J, Tomishima MJ, Zaninovic N, Colak D, Yan Z, Zhan Q, et al. The DNA replication
- 14 program is altered at the FMR1 locus in fragile X embryonic stem cells. Molecular cell 2014;53:19-31.
- 15 [92] Bochman ML, Sabouri N, Zakian VA. Unwinding the functions of the Pif1 family helicases. DNA 16 repair 2010;9:237-49.
- 17 [93] Sale JE, Lehmann AR, Woodgate R. Y-family DNA polymerases and their role in tolerance of
- 18 cellular DNA damage. Nature reviews Molecular cell biology 2012;13:141-52.
- 19 [94] Kim JC, Mirkin SM. The balancing act of DNA repeat expansions. Current opinion in genetics &
- 20 development 2013;23:280-8.
- 21 [95] Aguilera A, Garcia-Muse T. Causes of genome instability. Annual review of genetics 2013;47:1-22 32.
- 23 [96] Lin YL, Pasero P. Interference between DNA replication and transcription as a cause of genomic 24 instability. Current Genomics 2012;13:65-73.
- 25 [97] Wei X, Samarabandu J, Devdhar RS, Siegel AJ, Acharya R, Berezney R. Segregation of
- 26 Transcription and Replication Sites Into Higher Order Domains. Science 1998;281:1502-5.
- 27 [98] Aguilera A, García-Muse T. R Loops: From Transcription Byproducts to Threats to Genome
- 28 Stability. Molecular cell 2012;46:115-24.
- 29 [99] Ginno PA, Lim YW, Lott PL, Korf IF, Chedin F. GC skew at the 5' and 3' ends of human genes links
- 30 R-loop formation to epigenetic regulation and transcription termination. Genome research 2013.
- 31 [100] Huertas P, Aguilera A. Cotranscriptionally Formed DNA:RNA Hybrids Mediate Transcription
- 32 Elongation Impairment and Transcription-Associated Recombination. Molecular cell 2003;12:711-21.
- 33 [101] Alzu A, Bermejo R, Begnis M, Lucca C, Piccini D, Carotenuto W, et al. Senataxin Associates with
- 34 Replication Forks to Protect Fork Integrity across RNA-Polymerase-II-Transcribed Genes. Cell
- 35 2012;151:835-46.
- [102] Mischo HE, Gomez-Gonzalez B, Grzechnik P, Rondon AG, Wei W, Steinmetz L, et al. Yeast Sen1 36
- 37 helicase protects the genome from transcription-associated instability. Molecular cell 2011;41:21-32.
- 38 [103] Stirling PC, Chan YA, Minaker SW, Aristizabal MJ, Barrett I, Sipahimalani P, et al. R-loop-
- 39 mediated genome instability in mRNA cleavage and polyadenylation mutants. Genes & development
- 40 2012;26:163-75.
- 41 [104] Paulsen RD, Soni DV, Wollman R, Hahn AT, Yee MC, Guan A, et al. A genome-wide siRNA screen
- 42 reveals diverse cellular processes and pathways that mediate genome stability. Molecular cell
- 43 2009;35:228-39.
- [105] Wahba L, Amon Jeremy D, Koshland D, Vuica-Ross M. RNase H and Multiple RNA Biogenesis 44
- 45 Factors Cooperate to Prevent RNA:DNA Hybrids from Generating Genome Instability. Molecular cell 46 2011;44:978-88.
- 47 [106] Bermejo R, Capra T, Gonzalez-Huici V, Fachinetti D, Cocito A, Natoli G, et al. Genome-
- 48 Organizing Factors Top2 and Hmo1 Prevent Chromosome Fragility at Sites of S phase Transcription.
- 49 Cell 2009;138:870-84.
- 50 [107] Bermejo R, Capra T, Jossen R, Colosio A, Frattini C, Carotenuto W, et al. The replication
- 51 checkpoint protects fork stability by releasing transcribed genes from nuclear pores. Cell
- 52 2011;146:233-46.

- 1 [108] Crosetto N, Mitra A, Silva MJ, Bienko M, Dojer N, Wang Q, et al. Nucleotide-resolution DNA
- 2 double-strand break mapping by next-generation sequencing. Nat Meth 2013;10:361-5.
- 3 [109] Jones RM, Mortusewicz O, Afzal I, Lorvellec M, Garcia P, Helleday T, et al. Increased replication
- 4 initiation and conflicts with transcription underlie Cyclin E-induced replication stress. Oncogene
- 5 2013;32:3744-53.
- 6 [110] Davies SL, North PS, Hickson ID. Role for BLM in replication-fork restart and suppression of
- 7 origin firing after replicative stress. Nature structural & molecular biology 2007;14:677-9.
- 8 [111] Groth A, Rocha W, Verreault A, Almouzni G. Chromatin challenges during DNA replication and
- 9 repair. Cell 2007;128:721-33.
- 10 [112] Clemente-Ruiz M, Gonzalez-Prieto R, Prado F. Histone H3K56 acetylation, CAF1, and Rtt106
- 11 coordinate nucleosome assembly and stability of advancing replication forks. PLoS genetics
- 12 2011;7:e1002376.
- [113] Clemente-Ruiz M, Prado F. Chromatin assembly controls replication fork stability. EMBO
 reports 2009;10:790-6.
- 15 [114] Ask K, Jasencakova Z, Menard P, Feng Y, Almouzni G, Groth A. Codanin-1, mutated in the
- 16 anaemic disease CDAI, regulates Asf1 function in S-phase histone supply. The EMBO journal
- 17 2012;31:2013-23.
- 18 [115] Kerzendorfer C, Hannes F, Colnaghi R, Abramowicz I, Carpenter G, Vermeesch JR, et al.
- 19 Characterizing the functional consequences of haploinsufficiency of NELF-A (WHSC2) and SLBP
- 20 identifies novel cellular phenotypes in Wolf-Hirschhorn syndrome. Human molecular genetics
- 21 2012;21:2181-93.
- 22 [116] Chan KL, Palmai-Pallag T, Ying S, Hickson ID. Replication stress induces sister-chromatid
- bridging at fragile site loci in mitosis. Nature cell biology 2009;11:753-60.
- 24 [117] Naim V, Rosselli F. The FANC pathway and BLM collaborate during mitosis to prevent micro-
- 25 nucleation and chromosome abnormalities. Nature cell biology 2009;11:761-8.
- 26 [118] Magiera MM, Gueydon E, Schwob E. DNA replication and spindle checkpoints cooperate during
- 27 S phase to delay mitosis and preserve genome integrity. The Journal of cell biology 2014;204:165-75.
- 28 [119] Torres-Rosell J, De Piccoli G, Cordon-Preciado V, Farmer S, Jarmuz A, Machin F, et al. Anaphase
- 29 onset before complete DNA replication with intact checkpoint responses. Science 2007;315:1411-5.
- 30 [120] Baumann C, Korner R, Hofmann K, Nigg EA. PICH, a centromere-associated SNF2 family ATPase,
- 31 is regulated by Plk1 and required for the spindle checkpoint. Cell 2007;128:101-14.
- 32 [121] Vinciguerra P, Godinho SA, Parmar K, Pellman D, D'Andrea AD. Cytokinesis failure occurs in
- Fanconi anemia pathway-deficient murine and human bone marrow hematopoietic cells. The Journal
 of clinical investigation 2010;120:3834-42.
- 35 [122] Laulier C, Cheng A, Stark JM. The relative efficiency of homology-directed repair has distinct
- effects on proper anaphase chromosome separation. Nucleic acids research 2011;39:5935-44.
- 37 [123] Lahkim Bennani-Belhaj K, Rouzeau S, Buhagiar-Labarchede G, Chabosseau P, Onclercq-Delic R,
- Bayart E, et al. The Bloom syndrome protein limits the lethality associated with RAD51 deficiency.
- 39 Molecular cancer research : MCR 2010;8:385-94.
- 40 [124] Rodrigue A, Coulombe Y, Jacquet K, Gagne JP, Roques C, Gobeil S, et al. The RAD51 paralogs
- 41 ensure cellular protection against mitotic defects and aneuploidy. Journal of cell science
- 42 2013;126:348-59.
- 43 [125] Naim V, Wilhelm T, Debatisse M, Rosselli F. ERCC1 and MUS81-EME1 promote sister chromatid
- 44 separation by processing late replication intermediates at common fragile sites during mitosis.
- 45 Nature cell biology 2013;15:1008-15.
- 46 [126] Ying S, Minocherhomji S, Chan KL, Palmai-Pallag T, Chu WK, Wass T, et al. MUS81 promotes
- 47 common fragile site expression. Nature cell biology 2013;15:1001-7.
- 48 [127] Sofueva S, Osman F, Lorenz A, Steinacher R, Castagnetti S, Ledesma J, et al. Ultrafine anaphase
- 49 bridges, broken DNA and illegitimate recombination induced by a replication fork barrier. Nucleic
- 50 acids research 2011;39:6568-84.

- 1 [128] Rouzeau S, Cordelieres FP, Buhagiar-Labarchede G, Hurbain I, Onclercq-Delic R, Gemble S, et al.
- 2 Bloom's syndrome and PICH helicases cooperate with topoisomerase IIalpha in centromere
- disjunction before anaphase. PloS one 2012;7:e33905.
- 4 [129] Hossain M, Stillman B. Meier-Gorlin syndrome mutations disrupt an Orc1 CDK inhibitory
- 5 domain and cause centrosome reduplication. Genes & development 2012;26:1797-810.
- 6 [130] Smith L, Plug A, Thayer M. Delayed replication timing leads to delayed mitotic chromosome
- 7 condensation and chromosomal instability of chromosome translocations. Proceedings of the
- 8 National Academy of Sciences of the United States of America 2001;98:13300-5.
- 9 [131] Bertrand P, Lambert S, Joubert C, Lopez BS. Overexpression of mammalian Rad51 does not
- 10 stimulate tumorigenesis while a dominant-negative Rad51 affects centrosome fragmentation, ploidy
- and stimulates tumorigenesis, in p53-defective CHO cells. Oncogene 2003;22:7587-92.
- 12 [132] Win TZ, Mankouri HW, Hickson ID, Wang SW. A role for the fission yeast Rqh1 helicase in
- 13 chromosome segregation. Journal of cell science 2005;118:5777-84.
- 14 [133] Lee SE, Moore JK, Holmes A, Umezu K, Kolodner RD, Haber JE. Saccharomyces Ku70,
- mre11/rad50 and RPA proteins regulate adaptation to G2/M arrest after DNA damage. Cell
 1998;94:399-409.
- 17 [134] Toczyski DP, Galgoczy DJ, Hartwell LH. CDC5 and CKII control adaptation to the yeast DNA
- 18 damage checkpoint. Cell 1997;90:1097-106.
- 19 [135] Iraqui I, Chekkal Y, Jmari N, Pietrobon V, Freon K, Costes A, et al. Recovery of arrested
- 20 replication forks by homologous recombination is error-prone. PLoS genetics 2012;8:e1002976.
- 21 [136] Hu L, Kim TM, Son MY, Kim SA, Holland CL, Tateishi S, et al. Two replication fork maintenance
- 22 pathways fuse inverted repeats to rearrange chromosomes. Nature 2013;501:569-72.
- 23 [137] Kraus E, Leung WY, Haber JE. Break-induced replication: a review and an example in budding
- 24 yeast. Proceedings of the National Academy of Sciences of the United States of America
- 25 2001;98:8255-62.
- 26 [138] Lopes M, Foiani M, Sogo JM. Multiple mechanisms control chromosome integrity after
- 27 replication fork uncoupling and restart at irreparable UV lesions. Molecular cell 2006;21:15-27.
- 28 [139] Vanoli F, Fumasoni M, Szakal B, Maloisel L, Branzei D. Replication and recombination factors
- contributing to recombination-dependent bypass of DNA lesions by template switch. PLoS genetics2010;6:e1001205.
- 31 [140] Despras E, Daboussi F, Hyrien O, Marheineke K, Kannouche PL. ATR/Chk1 pathway is essential
- 32 for resumption of DNA synthesis and cell survival in UV-irradiated XP variant cells. Human molecular
- 33 genetics 2010;19:1690-701.
- 34 [141] Nik-Zainal S, Alexandrov LB, Wedge DC, Van Loo P, Greenman CD, Raine K, et al. Mutational
- processes molding the genomes of 21 breast cancers. Cell 2012;149:979-93.
- 36 [142] Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SA, Behjati S, Biankin AV, et al. Signatures of
- 37 mutational processes in human cancer. Nature 2013;500:415-21.
- 38 [143] Roberts SA, Sterling J, Thompson C, Harris S, Mav D, Shah R, et al. Clustered mutations in yeast
- and in human cancers can arise from damaged long single-strand DNA regions. Molecular cell2012;46:424-35.
- 41 [144] Smith CE, Llorente B, Symington LS. Template switching during break-induced replication.
- 42 Nature 2007;447:102-5.
- 43 [145] Hicks WM, Kim M, Haber JE. Increased mutagenesis and unique mutation signature associated 44 with mitotic gene conversion. Science 2010;329:82-5.
- 45 [146] Hastings PJ, Ira G, Lupski JR. A microhomology-mediated break-induced replication model for
- the origin of human copy number variation. PLoS genetics 2009;5:e1000327.
- 47 [147] Carvalho CM, Ramocki MB, Pehlivan D, Franco LM, Gonzaga-Jauregui C, Fang P, et al. Inverted
- 48 genomic segments and complex triplication rearrangements are mediated by inverted repeats in the
- 49 human genome. Nature genetics 2011;43:1074-81.
- 50 [148] Payen C, Koszul R, Dujon B, Fischer G. Segmental duplications arise from Pol32-dependent
- 51 repair of broken forks through two alternative replication-based mechanisms. PLoS genetics
- 52 2008;4:e1000175.

- [149] Mizuno K, Miyabe I, Schalbetter SA, Carr AM, Murray JM. Recombination-restarted replication
 makes inverted chromosome fusions at inverted repeats. Nature 2013;493:246-9.
- 3 [150] Campbell PJ, Yachida S, Mudie LJ, Stephens PJ, Pleasance ED, Stebbings LA, et al. The patterns
- 4 and dynamics of genomic instability in metastatic pancreatic cancer. Nature 2010;467:1109-13.
- 5 [151] Deem A, Keszthelyi A, Blackgrove T, Vayl A, Coffey B, Mathur R, et al. Break-induced replication
- 6 is highly inaccurate. PLoS Biol 2011;9:e1000594.
- 7 [152] Costantino L, Sotiriou SK, Rantala JK, Magin S, Mladenov E, Helleday T, et al. Break-induced
- 8 replication repair of damaged forks induces genomic duplications in human cells. Science
- 9 2014;343:88-91.
- [153] Arlt MF, Wilson TE, Glover TW. Replication stress and mechanisms of CNV formation. Current
 opinion in genetics & development 2012.
- 12 [154] Arlt MF, Rajendran S, Birkeland SR, Wilson TE, Glover TW. De novo CNV formation in mouse
- embryonic stem cells occurs in the absence of Xrcc4-dependent nonhomologous end joining. PLoS
 genetics 2012;8:e1002981.
- 15 [155] Carvalho CM, Pehlivan D, Ramocki MB, Fang P, Alleva B, Franco LM, et al. Replicative
- 16 mechanisms for CNV formation are error prone. Nature genetics 2013;45:1319-26.
- 17 [156] Boveri T. Concerning the origin of malignant tumours by Theodor Boveri. Translated and
- 18 annotated by Henry Harris. Journal of cell science 2008;121 Suppl 1:1-84.
- 19 [157] Nigg EA. Centrosome aberrations: cause or consequence of cancer progression? Nature reviews20 Cancer 2002;2:815-25.
- 21 [158] Sluder G, Nordberg JJ. The good, the bad and the ugly: the practical consequences of
- centrosome amplification. Current opinion in cell biology 2004;16:49-54.
- 23 [159] Bartkova J, Horejsi Z, Koed K, Kramer A, Tort F, Zieger K, et al. DNA damage response as a
- 24 candidate anti-cancer barrier in early human tumorigenesis. Nature 2005;434:864-70.
- 25 [160] Di Micco R, Sulli G, Dobreva M, Liontos M, Botrugno OA, Gargiulo G, et al. Interplay between
- 26 oncogene-induced DNA damage response and heterochromatin in senescence and cancer. Nature
- cell biology 2011;13:292-302.
- 28 [161] Darzynkiewicz Z. When senescence masquerades as DNA damage: is DNA replication stress the
- 29 culprit? Cell cycle 2009;8:3810-1.
- [162] Rodier F, Campisi J. Four faces of cellular senescence. The Journal of cell biology 2011;192:54756.
- 32 [163] Campisi J, d'Adda di Fagagna F. Cellular senescence: when bad things happen to good cells.
- 33 Nature reviews Molecular cell biology 2007;8:729-40.
- 34 [164] Bachrati CZ, Hickson ID. RecQ helicases: suppressors of tumorigenesis and premature aging.
- The Biochemical journal 2003;374:577-606.
- 36 [165] Lans H, Hoeijmakers JH. Cell biology: ageing nucleus gets out of shape. Nature 2006;440:32-4.
- 37 [166] Ogura M, Ogura H, Ikehara S, Dao ML, Good RA. Decrease by chronic energy intake restriction
- 38 of cellular proliferation in the intestinal epithelium and lymphoid organs in autoimmunity-prone
- 39 mice. Proceedings of the National Academy of Sciences of the United States of America
- 40 1989;86:5918-22.
- 41 [167] Hursting SD, Lavigne JA, Berrigan D, Perkins SN, Barrett JC. Calorie restriction, aging, and cancer
- 42 prevention: mechanisms of action and applicability to humans. Annual review of medicine
- 43 2003;54:131-52.
- 44 [168] Walsh T, King MC. Ten genes for inherited breast cancer. Cancer cell 2007;11:103-5.
- 45 [169] Walsh T, Casadei S, Lee MK, Pennil CC, Nord AS, Thornton AM, et al. Mutations in 12 genes for
- 46 inherited ovarian, fallopian tube, and peritoneal carcinoma identified by massively parallel
- 47 sequencing. Proceedings of the National Academy of Sciences of the United States of America
- 48 2011;108:18032-7.
- 49 [170] Plo I, Laulier C, Gauthier L, Lebrun F, Calvo F, Lopez BS. AKT1 inhibits homologous
- 50 recombination by inducing cytoplasmic retention of BRCA1 and RAD51. Cancer research
- 51 2008;68:9404-12.

- 1 [171] Guirouilh-Barbat JK, Wilhelm T, Lopez BS. AKT1/BRCA1 in the control of homologous
- recombination and genetic stability: the missing link between hereditary and sporadic breast
 cancers. Oncotarget 2010;1:691-9.
- 4 [172] Plo I, Lopez B. AKT1 represses gene conversion induced by different genotoxic stresses and
- 5 induces supernumerary centrosomes and aneuploidy in hamster ovary cells. Oncogene
- 6 2009;28:2231-7.
- _

1 Figure legend

- 2 Figure 1: Replication stress jeopardizes the completion of chromosome duplication and successful
- 3 mitotic commitment. Left panel: the DNA replication program ensures that origins fire in a sequential
- 4 order, and not all at the same time, thus allowing an appropriate management of the essential
- 5 resources for accurate and complete DNA synthesis. Right panel: replication stress from either
- 6 endogenous or exogenous causes (see text for details) compromises the achievement of
- 7 chromosome duplication, leaving chromosomal segment unreplicated when cells enter mitosis.
- 8 Unreplicated loci form anaphase bridges and challenge the even chromosome segregation.
- 9 Replication stress also favors mitotic extra centrosomes and multipolar mitosis, thus amplifying
- 10 mitotic catastrophes and genome instability to the whole genome.

11 Figure 2: Consequences of replication stress. Top panel: Alteration of the replication fork speed 12 (whatever the cause, see text for detail) generates stretches of ssDNA either because of uncoupling 13 between the replicative helicase and the DNA polymerase or nuclease attacks. Stretches of ssDNA 14 contribute to the activation of the ATR pathway which in turn acts locally to maintain the replisome 15 in a replication-competent state, and acts globally to delay cell progression, inhibits late-replicated 16 domains and probably slows down the speed of activate forks. Of note, ATR, and more generally the 17 DDR, also impact on many other cellular metabolism processes. Bottom panel: acute replication in 18 ATR deficient cells results in an excess of origin firing, a massive amount of chromatin-bound RPA 19 and an excessive consumption of dNTP. Stretches of ssDNA are thus unprotected and sensitive to 20 breakages. Inefficient replication complexes correspond to replisomes that cannot be maintained in 21 a replication-competent state due to ATR deficiency.



Replication stress in ATR proficient cells

