



**HAL**  
open science

## The fate of páramo plant assemblages in the sky islands of the northern Andes

Gwendolyn Peyre, Jonathan Roger Michel Henri Lenoir, Dirk N. Karger,  
Monica Gomez, Alexander Gonzalez, Olivier Broennimann, Antoine Guisan

► **To cite this version:**

Gwendolyn Peyre, Jonathan Roger Michel Henri Lenoir, Dirk N. Karger, Monica Gomez, Alexander Gonzalez, et al.. The fate of páramo plant assemblages in the sky islands of the northern Andes. Journal of Vegetation Science, 2020, 31 (6), pp.967-980. 10.1111/jvs.12898 . hal-03010954

**HAL Id: hal-03010954**

**<https://hal.science/hal-03010954>**

Submitted on 17 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Title page**

2 **Working title:** páramo assemblages Andes

3 **Full title:**

4 **The fate of páramo plant assemblages in the sky islands of the northern Andes**

5 **Authors:**

6 Gwendolyn Peyre<sup>1</sup>, Jonathan Lenoir<sup>2</sup>, Dirk N. Karger<sup>3</sup>, Monica Gomez<sup>1</sup>, Alexander  
7 Gonzalez<sup>1</sup>, Antoine Guisan<sup>4</sup>

8 <sup>1</sup> Department of Civil and Environmental Engineering, University of the Andes, Bogotá,  
9 Colombia

10 <sup>2</sup> Université de Picardie Jules Verne - CNRS, Amiens, France

11 <sup>3</sup> Swiss Federal Research Institute WSL, Birmensdorf, Switzerland

12 <sup>4</sup> Université de Lausanne, Lausanne, Switzerland

13 **Correspondence:**

14 Gwendolyn Peyre, Department of Civil and Environmental Engineering, University of the  
15 Andes, Bogotá, Colombia

16 Email address: [gf.peyre@uniandes.edu.co](mailto:gf.peyre@uniandes.edu.co)

17 **Funding information**

18 GP was financed through the Fondo de Apoyo para Profesores Asistentes – FAPA of the  
19 University of the Andes [PR.3.2016.3708].

20

21 **Abstract**

22 *Aims:* Assessing climate change impacts on biodiversity is a main scientific challenge,  
23 especially in the tropics, therefore, we predicted the future of plant species and communities  
24 on the unique páramo sky islands. We implemented the *Spatially Explicit Species*  
25 *Assemblage Modelling* framework, by i) calculating species' maximum dispersal distance,  
26 ii) modelling species distributions at present up to 2100, iii) assembling models into  
27 communities. Finally, we assessed the vulnerability of sky islands based on richness and  
28 composition changes.

29 *Location:* Ecuadorian super-páramo (>4200 m)

30 *Methods:* Using species trait data, the maximum dispersal distance of 435 species was  
31 calculated. Species distribution models (SDM) were fitted to obtain current and future  
32 distribution predictions based on dispersal and bioclimatic factors. The final assemblages  
33 for present and 2100 were achieved by stacking all probabilistic SDMs and applying the  
34 probability ranking rule. The vulnerability of each sky island was evaluated by quantifying  
35 richness and composition changes.

36 *Results:* Maximum dispersal distances ranged between 0.008-6027 m/year, and across all  
37 scenarios, 70% of models showed a net loss in species distribution while 9% of all species  
38 were predicted to undergo extinction by 2100. Local richness was estimated to decrease by  
39 56.63% on average, and composition changes in each sky island suggested a mean loss of  
40 64.74% of their original species pool against a 12.97% gain. Finally, 5% of the sky island  
41 floras reconverted from high-elevation to low-elevation species. These numbers were  
42 usually more important for high-elevation species and the mountains Pichincha, Ilinizas and  
43 Antisana.

44 *Conclusions:* Our study is methodologically pioneer and provides novel insight on the future  
45 of páramo biodiversity. Significant losses in species distribution and changes in community  
46 richness and composition suggest drastic impacts and call for further study considering  
47 additional factors, such as land-use. Finally, we recommend focusing monitoring and  
48 conservation strategies on the northern sky islands in priority.

49 **Key words:** Andes, community assemblages, climate change, maximum dispersal  
50 distance, páramo, plant species, Species Distribution Models, Spatially Explicit Species  
51 Assemblage Modelling, sky islands.

52

## 53 **Introduction**

54 The climatically-driven redistribution of life on Earth is one of the most challenging  
55 environmental threat humanity faces today (Pecl et al. 2017; Diaz et al. 2019). It is widely  
56 accepted that the current climate change is human-induced (Team et al. 2014) and that its  
57 main impacts on biodiversity include: shifts in phenology (Visser, & Both, 2005); changes  
58 in population densities (Graae et al. 2018), local extinctions (Panetta, Stanton, & Harte,  
59 2018), species range shifts (Lenoir, & Svenning 2015), and accelerated changes in species  
60 richness and assemblages (Bertrand et al. 2011; Steinbauer et al. 2018). Because biodiversity  
61 redistribution implies important changes in ecosystem functioning, human well-being and  
62 climate change itself (Pecl et al. 2017), there is an urgent need to increase our knowledge on  
63 the complex interplay between climate change and biodiversity and implement integrative  
64 scientifically-supported strategies to adapt and mitigate climate change (Diaz et al. 2019).

65 The high-elevation ecosystems in the northern Andes belong to the páramo  
66 biogeographical province, a tropical biodiversity hotspot critically threatened by climate  
67 change (Tovar, Arnillas, Cuesta, & Buytaert, 2013; Cuesta et al. 2019). The páramo is the  
68 youngest biodiversity hotspot and the richest tropical alpine province on Earth, totaling 15  
69 different phytogeographical units and around 5000 plant species, of which 60-80% are  
70 endemic (Sklenář, Hedberg, & Cleef, 2014; Peyre, Balslev, & Font, 2018). It can be  
71 altitudinally divided at present into three main elevation belts: i) the sub-páramo (~ 3000-  
72 3500 m) or lower tree or shrub-dominated ecotone with Andean forests; ii) the mid-páramo  
73 (~ 3500-4200 m), or páramo proper, including vast grasslands and giant rosette  
74 communities; and iii) the super-páramo (> 4200 m), with gradually scattering vegetation  
75 (Cuatrecasas, 1958). Contrarily to the sub-páramo and mid-páramo, which show certain  
76 spatial connectivity and are strongly shaped by human activities, the super-páramo is,  
77 hitherto, geographically isolated and above the current agriculture frontier (~ 4200m). The  
78 super-páramo hence represents the iconic sky islands of the northern Andes and hosts a very  
79 rare and endemic biodiversity with little species migration capacity on mountain tops.

80 Recent estimations predicted a temperature rise of 3°C ( $\pm$  1.5) across the páramo by  
81 the end of the century, accompanied by a general increase up to + 300 mm in vertical  
82 precipitation (rain), however highly variable depending on geography and topography  
83 (Urrutia, & Vuille, 2009; Anderson et al. 2011; Buytaert, Cuesta-Camacho, & Tobón, 2011).  
84 Although there is a general consensus on the need to increase our knowledge on the climate  
85 change impacts on the páramo biota and ecosystems, studies remain very scarce to date  
86 (Tovar et al. 2013; Feeley, Stroud, & Perez, 2017; Anthelme, & Lavergne, 2018; Anthelme,

87 & Peyre, 2019). For instance, Anderson et al. (2011) predicted a general upslope migration  
88 of 600 m for species to track the isotherms throughout the Tropical Andes by 2100. Other  
89 studies have also predicted up to a 50-60% increase in threats and local extinction risks by  
90 the second half of the century in the region (Ramírez-Villegas et al. 2014). At the ecosystem  
91 level, changes in structure and functions are suggested for the Tropical Andes (e.g. Tovar et  
92 al. 2013; Cuesta et al. 2019) but lack proper quantification for the páramo alone, and the  
93 consequent impacts on species assemblages and ecosystem services remain unclear  
94 (Buytaert et al. 2011). In order to fill the important knowledge gap on species redistribution  
95 under anthropogenic climate change and the subsequent structural re-organisation of plant  
96 assemblages in the páramo, new endeavors such as the Institute von Humboldt's Biomodels  
97 (<http://biomodelos.humboldt.org.co>) or the Pontificia Católica Universidad del Ecuador's  
98 Bioweb (<https://bioweb.bio>), as well as individual research initiatives (e.g. Ramírez-Villegas  
99 et al. 2014), looking to model the potential distribution of páramo species under climate  
100 change using species distribution models (SDMs) fitted on bioclimatic data are increasing.

101           Dispersal is key in enabling species to track climate change, essentially through the  
102 mechanisms of seed production, migration over certain dispersal distance and along suitable  
103 habitat routes, germination and establishment (Kammer, Schöb, & Choler, 2007; Vittoz, &  
104 Engler, 2007; Lenoir, & Svenning, 2015). This factor prevails in alpine areas where plants  
105 face many challenges to disperse upslope or across mountains, this due to i) often reduced-  
106 dispersal capacity (Vittoz, Dussex, Wassef, & Guisan, 2009; Morgan, & Venn, 2017), ii) the  
107 presence of biogeographic barriers such as forests and valleys, and iii) steep gradients  
108 favouring certain dispersal paths (Dirnböck & Dullinger 2004; Engler, Hordijk, & Guisan,  
109 2012). Accounting for dispersal when studying changes in species distribution in tropical  
110 mountains over time is crucial, because in comparison to higher latitudes, plants' dispersal  
111 capacities tend to be greater and more diverse (Chen, Tamme, Thomson, & Moles, 2019)  
112 while spatial climatic variations are less pronounced. At the community level, the resulting  
113 persistence vs. migration mismatch between local species, competitive regional pool and  
114 newcomers, such as invasive species, translates in important restructuring and composition  
115 changes (Singer et al. 2016; Alexander et al. 2018). Despite the general agreement on the  
116 need to include dispersal capacity in SDMs to improve current and future species distribution  
117 predictions (Araújo, & Guisan, 2006; Thuiller et al. 2008; Dullinger et al. 2012; Hattab et  
118 al. 2017), this factor remains overlooked today. Moreover, when accounted for, it usually  
119 takes the form of an “all or nothing” parameter, i.e. no dispersal or full dispersal (Guisan, &

120 Thuiller, 2005; Araújo et al. 2006; Holloway et al. 2016), albeit partial dispersal scenarios  
121 based on mean and/or maximum dispersal distance often are more reliable (Dullinger et al.,  
122 2012; Engler et al. 2012; Bateman, Murphy, Reside, Mokany, & VanDerWal, 2013).  
123 Nevertheless, partial dispersal approaches are increasing thanks to new techniques allowing  
124 to calculate significant approximations of dispersal distances based on a suite of plant trait  
125 data (Vittoz & Engler, 2007; Thomson, Moles, Auld, & Kingsford, 2011; Tamme et al. 2014;  
126 Bullock et al. 2017). Among the most dependable traits used to date is plant height, assuming  
127 that a tall plant disperses further than a short one (Muller-Landau, Wright, Calderón, Condit,  
128 & Hubbell, 2008; Thomson et al. 2011). Seed mass has also been evaluated as a significant  
129 trait, generally suggesting that light seeds disperse further than heavy ones (Parolo, & Rossi,  
130 2008), even though this pattern can be significantly influenced by the correlation between  
131 plant height and seed mass (Moles, Falster, Leishman, & Westoby, 2004; Thomson et al.  
132 2011). Finally, dispersal syndrome is usually considered important as, for instance flight-  
133 enhancing structures may increase anemochorous dispersal, and fruit characteristics can  
134 influence zoochorous dispersal (Vittoz, & Engler, 2007; Tamme et al. 2014; Thomson,  
135 Letten, Tamme, Edwards, & Moles, 2017).

136 Yet, to go beyond individual species predictions from SDMs, recent scientific  
137 advances in ecology propose to model entire biotic communities, relying on species  
138 distributions and adding a frame of abiotic and biotic factors such as historical-dispersal  
139 constraints, ecosystem richness capacity and biotic interactions among species (Guisan, &  
140 Rahbek, 2011; Wisz et al. 2013; Mod, le Roux, Guisan, & Luoto, 2015). In this context, the  
141 *Spatially Explicit Species Assemblage Modelling* framework (SESAM; Guisan, & Rahbek,  
142 2011) has been extensively used to predict species assemblages on a wide range of  
143 taxonomic groups, from insects to plants, and for several habitats, from forests to grasslands  
144 (e.g. D'Amen, Pradervand, & Guisan, 2015a; D'Amen et al. 2015b; Mod et al. 2015;  
145 D'Amen, Rahbek, Zimmermann, & Guisan, 2017; Mateo, Mokany, & Guisan, 2017). A  
146 wealth of research efforts has focused on improving the different aspects of SESAM, either  
147 regarding the macroecological constraints that represent the carrying capacity of  
148 communities (D'Amen et al. 2015a; Mateo et al. 2017), or the assembly rules that prioritize  
149 certain species co-occurrences (Wisz et al. 2013; D'Amen et al. 2015b). However, the  
150 historical-dispersal constraints remain overlooked to date and the need persists to improve  
151 techniques and launch more integrative models to refine community modelling at present  
152 and under climate change scenarios (D'Amen et al. 2017; Guisan et al. 2019).

153 The present study aims at implementing the SESAM framework, accounting for  
154 dispersal constraints for the first time, to predict changes in plant distributions and  
155 assemblages of the páramo sky islands during the 21<sup>st</sup> century. First, maximum dispersal  
156 distances were calculated and compared for mid-páramo and super-páramo species. It was  
157 expected that tall species such as shrubs that present either light seeds and dispersal-  
158 enhancing structures, e.g. wings, or heavy seeds with fruits dispersed by efficient zoochory,  
159 e.g. by birds, would be the best dispersing species. Second, the current and future  
160 distributions of each species up to the year 2100 were modelled based on bioclimatic  
161 variables while accounting for dispersal constraints to explore migration and extinction rates  
162 in the páramo. It was assumed that mid-páramo species would migrate further and faster  
163 than super-páramo species, which would in turn present greater extinction risks. Third,  
164 current and future plant communities were assembled on the sky islands by stacking all  
165 previous probabilistic models and applying richness constraints and assembly rules. The  
166 vulnerability of the network of sky islands was assessed based on the magnitude of changes  
167 in species richness and species composition in terms of species numbers and ratio of mid-  
168 páramo and super-páramo species, expecting that locally diverse areas today (Peyre, Balslev,  
169 Font, & Tello, 2019) would suffer faster and more drastic changes.

## 170 **Methods**

### 171 Study area

172 Ecuador was set as study case as a model representant of páramo countries due to its  
173 abundance of páramos and equatorial geolocation. The most representative mountains  
174 carrying the super-páramo belt, or sky islands, in Ecuador are from North to South:  
175 Cayambe, Antisana, Pichincha, Ilinizas, Cotopaxi, Chimborazo, Tungurahua, Sangay and  
176 Cajas (Fig. 1A). The super-páramo is usually found between 4200-4800 m a.s.l, although its  
177 lower and upper limits as well as main vegetation strongly depend on local topography and  
178 topoclimate. For example, low shrublands and cushion communities can be found at lower  
179 elevations under a dense upper cloud elevation zone, while meadows, deserts with shrubs  
180 and periglacial deserts are most likely to occur in the higher areas shaped by volcanic activity  
181 and/or glacier melting (Fig. 1B; Sklenář, & Ramsay, 2001). To conduct the following  
182 modelling analyses, the 4200 m isohypse and mountain tops were assumed as the lower and  
183 upper limits of the actual super-páramo belt respectively. Similarly, the 3000 m isohypse  
184 was set to delimit the broader páramo province (mid-páramo plus super-páramo), in order to



185 consider upward migration of plant species from the mid-páramo to the super-páramo as  
186 climate will warm under future scenarios.

### 187 Vegetation data

188 All vegetation plots corresponding to the Ecuadorian páramo were downloaded from  
189 VegParamo, an open access database for floristic and vegetation páramo data, compiling  
190 information from 40 data sources ([www.vegparamo.com](http://www.vegparamo.com); Peyre et al. 2015). Because  
191 VegParamo comprises the original authors' classification of the plots into sub-páramo, mid-  
192 páramo, super-páramo and azonal vegetation, i.e. *Polylepis forests*, bogs and marshes and  
193 rupicolous vegetation, only the plots belonging to the zonal mid-páramo and super-páramo  
194 elevational belts were retained. Plots located outside the study area, e.g. on Amazonian  
195 mountains, were also removed, as well as plots with a coarse georeferencing precision  
196 superior to 1km in the UTM system. The final plots, all sampled according to the  
197 phytosociological method (Braun-Blanquet, 1964), were rescaled into presence/absence  
198 data.

199 The floristic contents of the plots were checked, and the following groups were  
200 eliminated from the dataset: pteridophytes, bryophytes, lichens and vascular plants  
201 determined at the genus level or higher. When a taxon was determined at the infra-specific  
202 level, such as varieties and subspecies, it was aggregated at the species level. Taxonomic  
203 synonymy was checked using the Plant List ([www.plantlist.com](http://www.plantlist.com)) and Tropicos  
204 ([www.tropicos.org](http://www.tropicos.org)). Because the consequent vegetation dataset showed an important  
205 imbalance of species presences versus absences, additional occurrence data was  
206 complemented for these species from biological and herbarium databases consulted online,  
207 including Tropicos, Aarhus University Herbarium ([www.aubot.dk](http://www.aubot.dk)) and GBIF  
208 ([www.gbif.org](http://www.gbif.org)). The additional occurrence data was revised to remove outlier points and  
209 check duplicates between occurrence data and vegetation plots (based on UTM coordinates).  
210 Decimal spatial coordinates were then obtained for all plots, using the centroid of the  
211 corresponding UTM cell, as well as for the new added occurrence data, relying on the  
212 original georeferencing. Species that still presented less than 6 occurrences were removed  
213 and the complete dataset includes 642 vegetation plots from VegParamo (157 UTM) and  
214 2095 additional occurrences for 435 vascular plant species (Fig. S1; Table S1, S2; Peyre,  
215 2020). Finally, species were classified into super-páramo and mid-páramo species, when at  
216 least 50% of their occurrence data occurred above and below the 4200 m isohypse,  
217 respectively, resulting in 84 super-páramo species and 351 mid-páramo species.

218 Bioclimatic data

219 19 bioclimatic variables were downloaded from the CHELSA V1.2 database ([www.chelsa-](http://www.chelsa-climate.org)  
220 [climate-org](http://www.chelsa-climate.org); Karger et al. 2017) and adjusted to the study area. All variables were obtained  
221 for the period 1979-2013 and averaged across this time interval, considered as the current  
222 conditions (year 2000), which fitted the occurrence and vegetation data (1976-2014).  
223 Multicollinearity between variables was evaluated using a variance inflation factor  
224 correlation analysis (*vif* function, *usdm* R package; Naimi et al. 2014), and all variables  
225 below a threshold value of 0.7 for the Pearson correlation coefficient were retained  
226 (Dormann et al. 2012), resulting in the following selection: mean diurnal temperature range  
227 (bio2); temperature seasonality (bio4); mean temperature of the wettest quarter (bio8);  
228 precipitation seasonality (bio15); and precipitation of the coldest quarter (bio19). This set of  
229 predictor was not fitted per species (D'Amen et al. 2015a; Araújo et al. 2019) but was  
230 considered appropriate for the study area as a whole because of its certain focus on  
231 precipitation-related factors and seasonality, which usually prevail as drivers of plant  
232 diversity variation in tropical mountain areas (Peyre et al. 2019). To represent future  
233 scenarios and in order to encompass sufficient variance while reducing uncertainty in  
234 predictions (Thuiller, Guéguen, Renaud, Karger, & Zimmermann, 2019), a total of 8 climate  
235 change scenarios were selected, based on two representative concentration pathways  
236 (RCPs): CIMC5-RCP45 and CIMC5-RCP85, as well as four global circulation models  
237 (GCMs): bbc-csm1-1, CESM1-BGC, HadGEM2-AO and MRI-CGCM3. These GCMs were  
238 privileged based on their dissimilarity, according to the distance matrix calculated for all  
239 available GCMs presented in Sanderson, Knutti, & Caldwell (2015). Future bioclimatic  
240 predictions for each variable of interest were then downloaded from CHELSA V1.2 for  
241 every scenario at two dates, 2050 and 2070, and later cropped to fit the study area. Because  
242 dispersal is considered here as a key factor to constrain future species distributions, the  
243 bioclimatic data was interpolated by decade to provide frequent steps to include dispersal  
244 limitations (Engler et al. 2012). To do so, simple linear regressions were used to obtain for  
245 each bioclimatic variables the 2010, 2020, 2030 and 2040 values using a coefficient  
246 calculated for the 2000-2050 period following a similar procedure to the one employed by  
247 Adhikari et al. (2018). The same procedure was applied for 2060 based on the 2050-2070  
248 period, and for decades 2080, 2090 and 2100, values by extrapolating the 2050-2070  
249 regression coefficient into the future. All statistical analyses were conducted in R 4.3.1 (R  
250 Core Team, 2019).

## 251 Dispersal capacity

252 The widely-used *dispeRsal* function developed by Tamme et al. (2014) was chosen to  
253 estimate dispersal capacity in form of maximum dispersal distance, which is usually more  
254 suitable to include in SDMs than mean distance (Vittoz, & Engler, 2007; Thomson et al.  
255 2011). Beforehand, a basic trait database including plant height, vegetation stratum, main  
256 dispersion mode and seed mass was compiled for the 435 species. For plant height and  
257 vegetation stratum, information at the species level was retrieved from online herbarium  
258 material, such as the Field Museum (<https://plantidtools.fieldmuseum.org>), COL herbarium  
259 ([www.biovirtual.unal.edu.co](http://www.biovirtual.unal.edu.co)) and Plant JSTOR (<https://plants.jstor.org>), directly measured  
260 and averaged using at least four specimens per species when available. Dispersion mode and  
261 seed mass data at the species level were usually unavailable and therefore, information  
262 provided at the genus level from the Kew Botanical Garden seed collection initiative  
263 (<http://data.kew.org/sid>), and specific páramo literature were used (Frantzen, & Bouman,  
264 1989; Melcher, Bouman, & Cleef, 2000, 2004). Assuming for simplicity reasons that species  
265 would disperse only once yearly, the *dispeRsal* function was run as a kernel-shape  
266 probability density function of dispersal distances to calculate maximum dispersal distances  
267 (in m/yr) (Tamme et al. 2014; Bullock et al. 2017). The raw results were then upscaled to  
268 km per decade to fit the future bioclimatic predictions. Finally, dispersal capacity of mid-  
269 páramo and super-páramo species were sommelly compared by means of a t-Student test.

## 270 Species distribution models

271 The current distribution of each species was modelled based on actual environmental  
272 conditions, represented by the 5 bioclimatic variables selected, and dispersal capacity. The  
273 latter factor was introduced as a new variable, set by log-transforming the species maximum  
274 dispersal distance previously computed from plant traits and scaling the obtained values  
275 between 0 and 1 to rank species from bad to good dispersers (function *iForce*, *iSDM*  
276 package; Hattab et al. 2017). SDMs were then run using two different families of algorithms  
277 fitted for presence-absence data: i) generalized linear models (GLMs), which explore the  
278 linear relationship between explanatory and explained variables through a flexible Gaussian-  
279 identity distribution-link approach and were fitted using second degree polynomial curves  
280 and ii) random forests (RFs) that integrate the data through an ensemble learning, coupling  
281 decision trees with classification and regression settings/approaches (*biomod2* package;  
282 Thuiller et al. 2016). For each species, the data was divided into a 75% training dataset and  
283 a 25% testing dataset, and run with 50 models per algorithm. The true skill statistic (TSS)

284 value was computed for each model to evaluate SDMs' performances and all models with a  
285 TSS value greater than 0.6, either obtained with the GLM or RF algorithms, were retained  
286 (Allouche, Tsoar, & Kadmon, 2006). The final models were then ensembled to obtain the  
287 final probabilistic predictions (Araújo, & New, 2007; Marmion, Parviainen, Luoto,  
288 Heikkinen, & Thuiller, 2009), which were finally transformed into binary values using a  
289 threshold approach that equalizes the sensitivity and specificity metrics (*optimal.threshold*  
290 *function*; *ecospat* package; Liu, Berry, Dawson, & Pearson, 2005; Di Cola et al. 2017).

291 The calibrated current models were then used to project species' distributions into  
292 the future per decade and under each of the 8 different climate-change scenarios. To do so,  
293 the previously-used dispersal covariate was set to 0 and future distributions were projected  
294 against bioclimatic predictors only, in order to alleviate the static species-specific dispersal  
295 constraint. As formerly done, the probabilistic projections for each decade were then  
296 transformed into binary values and constrained in temporal order by the species maximum  
297 dispersal distance (Engler et al. 2012), hence allowing to track progressive changes in  
298 species distributions depending on available bioclimatic conditions and capacity to disperse  
299 from the previous distribution. Additional biogeographical barriers represented by the  
300 Andean forest and anthropogenic activities from the lowlands were also considered by using  
301 a raster mask at each time step that assumed that all pixels below the 3000 m isohypse were  
302 unsuitable. Thus, a species could only cross the matrix of unsuitable conditions in a time-  
303 step if its maximum dispersal distance was greater than the distance separating two suitable  
304 pixels. The set of decadal binary maps obtained for each species separately offered time  
305 series of species redistribution that can be used to assess the species-specific proportion of  
306 areas that were lost, gained or that remained occupied by the focal species between present  
307 and 2100 (functions *migclim.distance* and *migclim.plot*, *migclim* package; Engler et al.  
308 2012). Finally, the 2100 binary prediction was used to fit the 2100 probabilistic prediction  
309 for richness capacity (see below).

310 Changes in species distribution for the 21<sup>st</sup> century were evaluated as a binary result,  
311 net loss or net gain, by comparing and quantifying the number of pixels belonging to the  
312 2000 and 2100 distributions. Finally, extinction risk was considered when a species had at  
313 least one scenario predicting its complete extinction (i.e. the species completely disappeared  
314 from the studied area) by 2100. The severity of the risk was evaluated based on the number  
315 of scenarios predicting extinction by 2100, i) low risk (1-2 scenarios), ii) intermediate risk  
316 (3-4 scenarios), high risk (5-6 scenarios) and extinct (7-8 scenarios).

317

318 Model richness and assemblages

319 The SESAM framework relies on the *predict first assemble later* principle (Guisan, &  
320 Rahbek, 2011), meaning that final community predictions are achieved after applying an  
321 assembly procedure that consists in selecting species from the predicted pool until reaching  
322 the carrying capacity of a community defined as potential richness (Guisan, & Rahbek, 2011;  
323 Mateo et al. 2017). This local carrying capacity (at pixel level) is often estimated by either  
324 stacking the probabilities predicted by the SDMs or by using macroecological models  
325 (D'Amen et al. 2015a; Mateo et al. 2017) but the former has the advantage of being  
326 embedded within a same S-SDM framework and allowing for a varying timeline, so that we  
327 used it here. All SDMs in their probabilistic form were stacked in 2000 and 2100 for each  
328 scenario to evaluate richness. Net richness change at the pixel level was assessed as a  
329 percentage of the current (2000) richness, with positive and negative values meaning a net  
330 gain and loss, respectively, in potential species richness.

331 Finally, species composition at the pixel level was predicted through the probability  
332 ranking rule (PRR; D'Amen et al. 2015b), which consists in selecting species from the  
333 predicted pool in decreasing order of their SDM probability of occurrence until reaching the  
334 previously predicted potential richness. The PRR was applied to the stacks of SDM  
335 predictions for 2000 and 2100 using the *prc.ecospat* function of the *ecospat* package (Di  
336 Cola et al. 2017). Changes in composition, expressed as numbers of species gained and lost,  
337 as well as ratio of mid-páramo and super-páramo species to emulate upward migration and  
338 high-elevation competition, were evaluated for the 10 main páramo sky islands. Lastly,  
339 based on the estimated changes in richness and composition, local vulnerability to climate  
340 change was assessed.

341 **Results**

342 Species varied greatly in terms of their maximum dispersal distance, ranging from 0.008 to  
343 6027 m/yr in a negative exponential manner with a mean of 193 m/yr ( $\pm$  647) (Fig. S2; Table  
344 S2). Mid-páramo and super-páramo species showed significantly different maximum  
345 dispersal distances (t-test: 4.2972, df: 423.26, p-value: 2.148e-05), with mid-páramo species  
346 dispersing relatively well in average at 227.14 m/yr ( $\pm$  714), while super-páramo species  
347 showed shorter dispersal distances at 51.39 m/yr ( $\pm$  136). Long dispersal distances were  
348 usually associated with shrub genera with relatively heavy seeds (10-100 mg) and using  
349 zoochory or assisted anemochory, such as *Monnina* (Polygalaceae) and *Gaiadendron*

350 (Loranthaceae), whereas short dispersal distances was common in small herbaceous plants  
351 with lightweight seeds (0.0001-0.001 mg) and using unassisted anemochory or autochory as  
352 principal dispersal mode, as seen in *Aa* (Orchidaceae) and *Ourisia* (Plantaginaceae).

353 The performance of SDMs varied between species, and the average values were  
354 considered acceptable overall (across models: mean TSS = 0.614, Sensitivity = 79.835,  
355 Specificity = 78.952; Table 1). In general, the RF algorithm performed better than the GLM  
356 algorithm, and more of its models were retained, approximately 29% across all species.  
357 About 70% of the SDMs across all scenarios predicted a net loss in species' spatial  
358 distribution and only 30% predicted a net gain (Fig. 2). A total of 65 species (15% of all  
359 species) presented a degree of extinction risk under at least one scenario by the end of the  
360 century, 18 of which as a low risk, eight as an intermediate risk, 15 as a high risk and finally  
361 39 were predicted to become totally extinct from the studied area by 2100. The most  
362 endangered genera encountered in numbers were *Cerastium* (3 species), *Draba* (3 species),  
363 *Gentianella* (4 species) and *Viola* (3 species). In addition, 22 of the 65 species were super-  
364 páramo species, 5 of which are known endemics at the national scale: *Aetheolaena*  
365 *involutrata*, *Bartsia pumila*, *Festuca chimborazensis*, *Loricaria artisanensis* and *Viola*  
366 *polycephala*. Finally, of the 39 species with the highest extinction risk, 10 were super-  
367 páramo species known as páramo endemics, for example *Cerastium candicans*, *Draba*  
368 *depressa*, *Lachemilla tanacetifolia*, *Lupinus alopecuroides* and *Xenophyllum crassum*.

369 Regarding net richness changes in the páramo sky islands, the general trend at pixel  
370 level (alpha diversity) was a loss of 56.63 % ( $\pm 20.83$ ) of the original species richness  
371 between the present and 2100 (Fig. 3). The most affected areas were the northern mountains  
372 of Pichincha and Ilinizas with an average loss of 82.64 ( $\pm 2.15$ ) and 77.24% ( $\pm 3.95$ )  
373 respectively. Contrarily, the least affected areas were the central mountains of Tungurahua  
374 and Altar with an average loss of 13.52 % ( $\pm 11.12$ ) and 31.90 % ( $\pm 6.52$ ) respectively, and  
375 punctual pixels with positive values hence gaining species, up to 18.73 % on the  
376 Chimborazo.

377 At the sky island level, important significant composition changes in the final plant  
378 communities were observed (Fig. 3; Fig S3; Table S3). Sky islands lost an average 64.74%  
379 ( $\pm 14.12$ ) of their original species pool (gamma diversity), Pichincha and Ilinizas being the  
380 most significantly affected with losses of 85.68% ( $\pm 5.70$ ) and 74.73% ( $\pm 5.12$ ) respectively.  
381 By contrast, Altar and Tungurahua were the least affected, losing 46.49 ( $\pm 10.76$ ) and

382 50.00% ( $\pm 9.74$ ) respectively. Gains at this scale did not compensate losses, with the average  
383 gain reaching 12.97 % ( $\pm 11.41$ ), some páramos gaining more species such as Tungurahua  
384 and Altar with 39.21% ( $\pm 5.03$ ) and 20.86% ( $\pm 6.31$ ), while others gained less, such as  
385 Cayambe and Antisana with 7.17 % ( $\pm 3.02$ ) and 7.47 % ( $\pm 3.05$ ). The most drastic changes  
386 were therefore observed for the northern mountains of Cayambe and Pichincha.

387 On average in 2000, the ratio between mid-páramo and super-páramo species in the  
388 páramo sky islands reached 68.79 / 31.21 ( $\pm 6.09$ ), and changed across all scenarios in 2100  
389 to 73.61 / 26.39 ( $\pm 4.46$ ), meaning that approximately 4.82% of super-páramo species were  
390 replaced with mid-páramo ones (Table 2). The most drastic tendencies were observed for  
391 Ilinizas and Antisana, losing 9.76 and 8.01% of their super-páramo species respectively,  
392 while the least affected sky islands were Cayambe, which gained 0.78% of super-paramo  
393 species, and Sangay that lost 0.48% of super-paramo species. Based on richness at pixel and  
394 sky island level and composition changes, Ilinizas, Pichincha and in lesser means Antisana  
395 were therefore considered as most threatened by climate change.

## 396 **Discussion**

397 Mountains are particularly sensitive to climate change because they are often prone to  
398 accelerated changes and more rapid responses of biota in comparison with lowlands  
399 (Bertrand et al. 2011; Smith, Edmonds, Hartin, Mundra, & Calvin, 2015; Steinbauer et al.  
400 2018). Neotropical mountains in particular are classified as *highly vulnerable* to climate  
401 change (Team et al. 2014) and are expected to suffer drastic changes in biodiversity and its  
402 associated ecosystem services that will affect millions of people on the short to mid-term  
403 (Buytaert et al. 2011; Tovar et al. 2013; Anthelme, & Lavergne, 2018). In addition, the fast  
404 and important changes in rural occupation and land uses that are occurring due to improving  
405 socio-economic conditions and the subsequent population growth, contribute greatly to the  
406 extent and intensity of environmental degradation (Hofstede, Segarra, & Vásconez, 2003).  
407 It is also in this region that predicting the impacts of future global changes becomes  
408 particularly challenging, due to data availability and the lack of knowledge on biodiversity,  
409 climatic variations, anthropogenic dynamics and the responses of biodiversity to these  
410 drivers (Anderson et al. 2011; Ramírez-Villegas et al. 2014; Lenoir, & Svenning 2015;  
411 Feeley et al. 2017).

## 412 Dispersal capacity

413 To date, trait data and functional analyses for the páramo remain scarce, especially regarding  
414 non-leaf characteristics (e.g. Sánchez et al. 2014), and so far few works have focused on  
415 seed-related traits and characterized the dispersal modes of páramo species (Frantzen, &  
416 Bouman, 1989; Melcher et al. 2000; Melcher et al. 2004). Our estimates of species maximum  
417 dispersal distances based on plant dispersal traits highlighted that páramo species disperse  
418 on the lower end of the terrestrial angiosperm dispersal spectrum (Kinlan, & Gaines, 2003)  
419 and in general similarly to other alpine floras (Vittoz, & Engler, 2007; Morgan, & Venn,  
420 2017). Our result showed some support for the hypothesis that low-elevation shrubs,  
421 characterized by heavy seeds and usually using either zoochory (e.g. Ericaceae) or assisted  
422 anemochory (e.g. Asteraceae), disperse best. Indeed, the correlation favouring plant height  
423 over seed mass seemed particularly advantageous for these shrub species, at least at the  
424 landscape level (Parolo, & Rossi, 2008; Thomson et al. 2011, 17), whereas within a same  
425 dispersal syndrome or vegetation type, light-seed species that disperse anemochorously are  
426 usually considered as more efficient (Muller-Landau et al. 2008). Our best dispersers were  
427 often zoochorous and presenting relatively large fruits visible to vertebrate animals, such as  
428 birds, which undoubtedly enhanced local dispersal (Thomson et al. 2011; Tamme et al. 2014)  
429 and also increased frequency of long-distance dispersal events, for example between  
430 mountain tops (Vittoz et al. 2009). Consequently, an upcoming challenge for these plant  
431 species in the future is facing their own responses to climate change but also those of their  
432 animal disperser (Travis et al. 2013). With increasing elevation and distance from the  
433 treeline, Poaceae become progressively dominant in the páramo ecosystems, ensuing a larger  
434 proportion of anemochorous species that often present dispersal-enhancing structures, i.e.  
435 wings or pappi. Such structures are also crucial in helping plants overcome the long-distance  
436 dispersal limitations they face in alpine areas (Vittoz et al. 2009; Morgan, & Venn, 2017;  
437 Thomson et al. 2017). In the super-páramo belt, small characteristic plants such as  
438 *Nototriche*, *Draba* and *Ourisia* (Peyre et al. 2018) are often featuring perennality, small  
439 light-seeds, high autogamy rate and little seed productivity, which might explain the short-  
440 distance dispersal tendency observed. In addition, the yet inhospitable environment and  
441 unavailability of efficient animal dispersers on these sky-islands often forces plant species  
442 to recur to vegetative (clonal) reproduction in priority and limit their sexual reproduction  
443 (Vittoz et al. 2009). Accounting for establishment capacity, the availability of microsites,  
444 good microclimatic and edaphic conditions, and strong species interactions, such as  
445 facilitation of assisted dispersal, is therefore key to finally assess the migration capacity of  
446 these species (Scherrer, & Körner, 2011; Anthelme, Cavieres, & Dangles, 2014; Hupp,



447 Llambí, Ramírez, & Callaway, 2017; Alexander et al. 2018). Finally, considering the  
448 competitiveness of super-páramo species against advantaged shrub and herbaceous species  
449 from lower elevations, including opportunistic and exotic species, is necessary to evaluate  
450 the intensity of the threat (Pauchard et al. 2009; Llambí, Hupp, Saez, & Callaway, 2018).

451 The importance of dispersal capacity in shaping a species distribution has long been  
452 acknowledged and increasingly included to improve SDM performances in the last decade  
453 (Thuiller et al. 2008; Dullinger et al. 2012; Bateman et al. 2013; Hattab et al. 2017).  
454 Community modelling has been considerably improving in parallel, with considerable  
455 efforts being put on better accounting for community assembly processes, including the  
456 carrying capacity of communities and assembly rules (D'Amen et al. 2017; Mateo et al.  
457 2017), nonetheless, dispersal constraints had remained mostly overlooked in community  
458 modelling (D'Amen et al. 2017; Guisan et al. 2019). This study supports that dispersal is a  
459 key factor in defining the future distribution of species and their assemblages, as illustrated  
460 in Fig. 2 by the overall overprediction of potential versus realized distributions quantified as  
461 the *suitable but unoccupied area*. Our finding is therefore in agreement with previous works  
462 maintaining that dispersal capacity is particularly important in tropical mountains, where  
463 climate has had an important but not exclusively prevailing influence on biota distribution  
464 (e.g. Flantua, O'dea, Onstein, Giraldo, & Hooghiemstra, 2019). Our results also suggest that  
465 super-páramo plants might not be able to migrate altitudinally along the same mountain but  
466 also be prevented in their attempts to colonize new mountain ranges, sustaining the key role  
467 of topographic distance and isolation (Sklenář, & Jørgensen, 1999; Flantua et al. 2019).

#### 468 Distribution responses of plant species

469 Of all species, 70% were predicted to show a significant distribution loss by 2100, which  
470 complements previous findings in which 50-60% of tropical Andean species reduced their  
471 distribution in the second half of the 21<sup>st</sup> century (Ramírez-Villegas et al. 2014). Even more  
472 concerning is that 15% of all páramo species showed a significant extinction risk by 2100,  
473 and 25% of all super-páramo species, supporting our hypothesis that super-páramo species  
474 are generally more at risk than mid-páramo ones. Among the vulnerable super-páramo  
475 species are several diagnostic species of super-páramo phytogeographical units, for example  
476 *Ourisia muscosa* of the *Upper humid super-páramo* and *Draba depressa* of the *Upper dry*  
477 *super-páramo* (Peyre et al. 2018). Furthermore, our results posed as threatened many super-  
478 páramo species that are páramo endemics such as *Cerastium candicans*, *Draba depressa* and  
479 *Xenophyllum crassum* or even national páramo endemics, for example *Bartsia pumila*,

480 *Loricaria antisanensis* and *Viola polycephala*. Our study alerts of maximum extinction risk  
481 for 9% of all studied species, some of which were already classified following the IUCN  
482 system as threatened, for example *Bartsia pumila* (VU) and *Gentianella hirculus* (EN), but  
483 also some considered as not threatened, such as *Gentianella limoselloides* (LC) and *Halenia*  
484 *taruga-gasso* (NT) (León-Yáñez, 2012). Because SDMs can be a useful complement to  
485 further conduct UICN evaluations (Breiner, Guisan, Nobis, & Bergamini, 2017; but see  
486 Akcakaya, Butchart, Mace, Stuart, & Hilton-Taylor, 2006), we therefore urge, based on the  
487 SDM results to focus further study on these predicted-extinct species, especially regarding  
488 population health and individual fitness, and encourage ex-situ conservation efforts such as  
489 seed banks to preserve this unique biodiversity.

#### 490 Changes in richness and plant assemblages

491 Local richness at the pixel level has been found to significantly decrease in most sky islands,  
492 even accounting for potential colonization from lower elevations. The northern sky islands  
493 of Ilinizas and Pichincha were the most affected, which contradicts previous findings stating  
494 richness stability in this region (Ramírez-Villegas et al. 2014), and exceptions to the rule  
495 included few central mountains such as Tungurahua and Altar where richness would remain  
496 stable or even increase slightly. As a result, these last sky islands might be experiencing the  
497 worldwide-observed *accelerated colonization process* on mountains (Steinbauer et al.  
498 2018), but buffered by specific environmental conditions and a particularly vulnerable  
499 species pool. Overall, we found no support for the negative relationship hypothesis between  
500 actual richness and future richness changes, as the more diverse (at the local scale, or plot  
501 scale) eastern mountains suffered less drastic changes than the less diverse northern  
502 mountains (Peyre et al. 2019). Because this finding plausibly supposes that the communities'  
503 richness capacity is not reached on these sky islands, we recommend further scientific focus  
504 on that particular macroecological hypothesis.

505         Composition changes were also predicted as very abrupt in the region, estimating  
506 important species losses and little gains at the sky island level (gamma diversity). Northern  
507 mountains were predicted to lose substantial amounts of species as much as 85% in  
508 Pichincha and gain very few species overall. By contrast, central mountains saw moderate  
509 losses and in the case of Tungurahua, a potential even loss and gain, depending on the  
510 climate change scenario considered. Several complementary hypotheses to climate change  
511 might be advanced to explain such a pattern, for example the geographic isolation and  
512 elevation of the massif, which could either facilitate or constrain dispersal and migration

513 (Sklenář, & Jørgensen, 1999). Finally, the ratio of super-páramo vs. mid-páramo species  
514 decreased by 2100, especially in northern mountains, sky islands generally losing 5% of  
515 their super-páramo species to mid-páramo ones. Our results therefore support previous  
516 findings that mid-páramo species are moving upslope and competing with super-páramo  
517 species, although the role of adaptation, microenvironmental refugia and biotic interactions  
518 remain to be considered for more refined plant responses.

519 Our results evidenced that sky islands from northern Ecuador, including Pichincha  
520 and Ilinizas, were evaluated as the most vulnerable to future climate change. These páramos  
521 are located along the inter-Andean valley where an important population resides, i.e. Quito  
522 and Latacunga, making the anthropogenic change expected for the end of the century in the  
523 form of fast and intense land-use changes an even bigger threat than climate change  
524 (Hofstede et al. 2003). Because the agricultural frontier could move upslope too when soil  
525 develops sufficiently at high altitude, entire super-páramo ecosystems could suffer drastic  
526 changes in their structure and functions. Therefore, the predictions provided here, based on  
527 climate-change solely, should be considered as conservative scenarios and even more  
528 forceful impacts should be expected under a global change scenario. In addition, exotic and  
529 efficiently dispersed opportunistic species could take advantage of the anthropogenic change  
530 to migrate and invade the sky islands, competing directly with local species and forming no-  
531 analog communities with little ecological value (Le Roux, & McGeoch, 2008; Anthelme, &  
532 Peyre, 2019). Precise monitoring should therefore be tracking the anthropogenic changes  
533 associated to climate change on these sky islands in priority so to prevent a critical threat to  
534 the sky islands of the northern Andes currently holding the reservoir or museum of super-  
535 páramo species.

#### 536 Study limitations and future recommendations

537 Regarding dispersal estimates, we consider that even though trait-based dispersal distance  
538 methods such as Tamme et al. (2014) are very useful and give reasonable predictions, the  
539 fact that they have been shown to underestimate distances for wind-dispersed species, an  
540 important proportion of our species pool (Herrmann et al. 2016, Morgan, & Venn, 2017),  
541 would support the use of more detailed techniques in the future, for example mechanistic  
542 models that acknowledge wind velocities and dispersal pathways (Skarpaas et al. 2004;  
543 Holloway et al. 2016). Moreover, we did not account for certain crucial variation such as i)  
544 potential changes in biogeographical barriers with climate change such as glacier retreat or

545 forest expansion, which might affect the dispersal reach of species (Caplat et al. 2016), nor  
546 did we consider ii) that specific dispersal distances might change over time (Travis et al.  
547 2013) or iii) the importance of long dispersal events promoted by climate change  
548 consequences such as the frequency and intensity of storms (Hellmann, Byers, Bierwagen,  
549 & Dukes, 2008). Another limitation regards data availability and coverage, a commonly  
550 encountered issue in tropical research. In fact, the species list was recovered from  
551 VegParamo, which is a pioneer database with important representativity of the páramo taxa,  
552 although it still accounts for certain geographic and taxonomic knowledge gaps (Peyre et al.  
553 2015; Bottin et al. 2019). Finally, climatic interpolations in tropical mountain areas that are  
554 topographically and environmentally complex are particularly challenging to obtain,  
555 especially for a broad study area and future climatic scenarios, and more fitted climatic data  
556 with a finer resolution could have improved the models performance and predictions.

557 Future recommendations include complementing the trait data with observed data and  
558 calculate distance on site to provide better estimates of dispersal capacity of plants in the  
559 region. To improve the models' accuracy and ecological meaning, it would be useful to  
560 account for different dynamic factors unavailable at the time (Anthelme et al. 2014; Graae  
561 et al. 2018), for example: i) additional abiotic variables at the macroscale, such as horizontal  
562 precipitation that represents an important water intake for plants, but also meso and  
563 microscale, for example local night freezing, that could balance macroeffects (Scherrer, &  
564 Körner, 2011; Mod et al. 2015), ii) fine biotic interactions to incorporate into the assembly  
565 rules, competition but also facilitation to dispersal or establishment (Llambí et al. 2018), and  
566 iii) land-use changes, whose intensity and expansion might increase under future climate  
567 change (Hofstede et al. 2003; Anderson et al. 2011).

568 **Acknowledgements**

569 We thank Dr. Fabien Anthelme for revising the final form of the paper.

570 **Author contributions**

571 G.P. and A.G. conceived of the research idea; G.P., M.G. and A.G. processed the data;  
572 G.P. performed the statistical analyses and wrote the paper; all authors discussed the  
573 methods, results and commented on the manuscript.

574 **Data accessibility**

575 The vegetation data is freely available from VegParamo ([www.vegparamo.com](http://www.vegparamo.com)), and on  
576 Dryad (<https://datadryad.org/>; Peyre, 2020).

577 **References**

- 578 Adhikari, P., Shin, M. S., Jeon, J. Y., Kim, H. W., Hong, S., & Seo, C. (2018). Potential  
579 impact of climate change on the species richness of subalpine plant species in the mountain  
580 national parks of South Korea. *Journal of Ecology and Environment*, 42(1), 36.  
581 <https://doi.org/10.1186/s41610-018-0095-y>
- 582 Akcakaya, H. R., Butchart, S. H. M., Mace, G. M., Stuart, S. N., & Hilton-Taylor, C. (2006).  
583 Use and misuse of the IUCN Red List Criteria in projecting climate change impacts on  
584 biodiversity. *Global Change Biology*, 12, 2037–2043. <https://doi.org/10.1111/j.1365-2486.2006.01253.x>
- 586 Alexander, J. M., Chalmandrier, L., Lenoir, J., Burgess, T. I., Essl, F., Haider, S., ... &  
587 Pauchard, A. (2018). Lags in the response of mountain plant communities to climate change.  
588 *Global change biology*, 24(2), 563–579. <https://doi.org/10.1111/gcb.13976>
- 589 Allouche, O., Tsoar, A. & Kadmon, R. (2006). Assessing the accuracy of species distribution  
590 models: prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology*, 43,  
591 1223–1232. <https://doi.org/10.1111/j.1365-2664.2006.01214.x>
- 592 Anderson, E. P., Marengo, J., Villalba, R., Halloy, S., Young, B., Cordero, D., ... & Martinez,  
593 R. (2011). *Consequences of climate change for ecosystems and ecosystem services in the*  
594 *tropical Andes*. In S. K. Herzog, R. Martínez, P. M. Jørgensen & H. Tiessen (Eds.) *Climate*  
595 *Change and Biodiversity in the Tropical Andes* (pp. 1–18). São José dos Campos, Brazil:  
596 Inter-American Institute for Global Change Research (IAI) and Scientific Committee on  
597 Problems of the Environment (SCOPE)
- 598 Anthelme, F., Cavieres, L. A., & Dangles, O. (2014). Facilitation among plants in alpine  
599 environments in the face of climate change. *Frontiers in plant science*, 5, 387.  
600 <https://doi.org/10.3389/fpls.2014.00387>
- 601 Anthelme, F., & Lavergne, S. (2018). Alpine and arctic plant communities: a worldwide  
602 perspective. *Perspectives in Plant Ecology Evolution and Systematics*, 30, 1–5. <https://doi.org/10.1016/j.ppees.2017.12.002>
- 604 Anthelme, F., & Peyre, G. (2019). *Biogeography of South American Highlands*. Reference  
605 Module in Earth Systems and Environmental Sciences, Encyclopedia of the World's biomes,  
606 Elsevier. <https://doi.org/10.1016/B978-0-12-409548-9.11811-1>.
- 607 Araújo, M. B., & Guisan, A. (2006). Five (or so) challenges for species distribution  
608 modelling. *Journal of biogeography*, 33(10), 1677–1688. <https://doi.org/10.1111/j.1365-2699.2006.01584.x>
- 610 Araújo, M.B., Thuiller, W. & Pearson, R.G. (2006) Climate warming and the decline of  
611 amphibians and reptiles in Europe. *Journal of Biogeography*, 33, 1712–1728.  
612 <https://doi.org/10.1111/j.1365-2699.2006.01482.x>
- 613 Araújo, M. B., & New, M. (2007). Ensemble forecasting of species distributions. *Trends in*  
614 *Ecology & Evolution*, 22, 42–47. <https://doi.org/10.1016/j.tree.2006.09.010>
- 615 Araújo, M. B., Anderson, R. P., Barbosa, A. M., Beale, C. M., Dormann, C. F., Early, R., ...  
616 & Rahbek, C. (2019). Standards for distribution models in biodiversity assessments. *Science*  
617 *Advances*, 5, eaat4858. <https://doi.org/10.1126/sciadv.aat4858>

- 618 Bateman, B. L., Murphy, H. T., Reside, A. E., Mokany, K., & VanDerWal, J. (2013).  
619 Appropriateness of full-, partial-and no-dispersal scenarios in climate change impact  
620 modelling. *Diversity and Distributions*, 19(10), 1224–1234.  
621 <https://doi.org/10.1111/ddi.12107>
- 622 Bertrand, R., Lenoir, J., Piedallu, C., Riofrío-Dillon, G., de Ruffray, P., Vidal, C., ... &  
623 Gégout, J. C. (2011). Changes in plant community composition lag behind climate warming  
624 in lowland forests. *Nature*, 479, 517. <https://doi.org/10.1038/nature10548>
- 625 Bottin, M., Peyre, G., Vargas, C., Raz, L., Richardson, J. E., & Sanchez, A. (2019).  
626 Phytosociological data and herbarium collections show congruent large-scale patterns but  
627 differ in their local descriptions of community composition. *Journal of Vegetation Science*,  
628 31(1), 208–219. <https://doi.org/10.1111/jvs.12825>
- 629 Braun-Blanquet, J. (1964). *Pflanzensoziologie: Grundzüge der Vegetationskunde*. New  
630 York, NY, US: Springer.
- 631 Breiner, F. T., Guisan, A., Nobis, M. P., & Bergamini, A. (2017). Including environmental  
632 niche information to improve IUCN Red List assessments. *Diversity and Distributions*, 23,  
633 484–495. <https://doi.org/10.1111/ddi.12545>
- 634 Bullock, J. M., Mallada González, L., Tamme, R., Götzenberger, L., White, S. M., Pärtel,  
635 M., & Hooftman, D. A. (2017). A synthesis of empirical plant dispersal kernels. *Journal of*  
636 *Ecology*, 105, 6–19. <https://doi.org/10.1111/1365-2745.12666>
- 637 Buytaert, W., Cuesta-Camacho, F., & Tobón, C. (2011). Potential impacts of climate change  
638 on the environmental services of humid tropical alpine regions. *Global Ecology and*  
639 *Biogeography*, 20, 19–33. <https://doi.org/10.1111/j.1466-8238.2010.00585.x>
- 640 Caplat, P., Edelaar, P., Dudaniec, R. Y., Green, A. J., Okamura, B., Cote, J., ... & Petit, E. J.  
641 (2016). Looking beyond the mountain: dispersal barriers in a changing world. *Frontiers in*  
642 *Ecology and the Environment*, 14(5), 261–268. <https://doi.org/10.1002/fee.1280>
- 643 Chen, S. C., Tamme, R., Thomson, F. J., & Moles, A. T. (2019). Seeds tend to disperse  
644 further in the tropics. *Ecology letters*, 22(6), 954–961. <https://doi.org/10.1111/ele.13255>
- 645 Cuatrecasas, J. (1958). Aspectos de la vegetación natural de Colombia. *Revista de la*  
646 *Academia Colombiana de Ciencias Exactas, Físicas y Naturales*, 10, 221–268.
- 647 Cuesta, F., Llambí, L. D., Huggel, C., Drenkhan, F., Gosling, W. D., Muriel, P., ... & Tovar,  
648 C. (2019). New land in the Neotropics: a review of biotic community, ecosystem, and  
649 landscape transformations in the face of climate and glacier change. *Regional environmental*  
650 *change*, 1–20. <https://doi.org/10.1007/s10113-019-01499-3>
- 651 D'Amen, M., Dubuis, A., Fernandes, R. F., Pottier, J., Pellissier, L., & Guisan, A. (2015a).  
652 Using species richness and functional traits predictions to constrain assemblage predictions  
653 from stacked species distribution models. *Journal of Biogeography*, 42, 1255–1266.  
654 <https://doi.org/10.1111/jbi.12485>
- 655 D'Amen, M., Pradervand, J. N., & Guisan, A. (2015b). Predicting richness and composition  
656 in mountain insect communities at high resolution: a new test of the SESAM framework.  
657 *Global Ecology and Biogeography*, 24, 1443–1453. <https://doi.org/10.1111/geb.12357>

- 658 D'Amen, M., Rahbek, C., Zimmermann, N. E., & Guisan, A. (2017). Spatial predictions at  
659 the community level: from current approaches to future frameworks. *Biological Reviews*,  
660 92, 169–187. <https://doi.org/10.1111/brv.12222>
- 661 Di Cola, V., Broennimann, O., Petitpierre, B., Breiner, F. T., D'amen, M., Randin, C., ... &  
662 Guisan, A. (2017). ecospat: a R package to support spatial analyses and modeling of species  
663 niches and distributions. *Ecography*, 40, 774–787. <https://doi.org/10.1111/ecog.02671>
- 664 Diaz, S., Settele, J., Brondízio, E., Ngo, H., Guèze, M., Agard, J., ... & Chan, K. (2019).  
665 *Summary for policymakers of the global assessment report on biodiversity and ecosystem*  
666 *services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem*  
667 *Services*.
- 668 Dirnböck, T., & Dullinger, S. (2004). Habitat distribution models, spatial autocorrelation,  
669 functional traits and dispersal capacity of alpine plant species. *Journal of Vegetation Science*,  
670 15(1), 77–84. <https://doi.org/10.1111/j.1654-1103.2004.tb02239.x>
- 671 Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., ... & Lautenbach,  
672 S. (2013). Collinearity: a review of methods to deal with it and a simulation study evaluating  
673 their performance. *Ecography*, 36, 27–46. [https://doi.org/10.1111/j.1600-](https://doi.org/10.1111/j.1600-0587.2012.07348.x)  
674 [0587.2012.07348.x](https://doi.org/10.1111/j.1600-0587.2012.07348.x)
- 675 Dullinger, S., Gattlinger, A., Thuiller, W., Moser, D., Zimmermann, N. E., Guisan, A., . . .  
676 Hülber, K. (2012). Extinction debt of high-mountain plants under twenty-first-century  
677 climate change. *Nature Climate Change*, 2, 619–622. <https://doi.org/10.1038/nclimate1514>
- 678 Engler, R., Hordijk, W., & Guisan, A. (2012). The MIGCLIM R package—seamless  
679 integration of dispersal constraints into projections of species distribution models.  
680 *Ecography*, 35, 872–878. <https://doi.org/10.1111/j.1600-0587.2012.07608.x>
- 681 Feeley, K. J., Stroud, J. T., & Perez, T. M. (2017). Most ‘global’ reviews of species’  
682 responses to climate change are not truly global. *Diversity and Distributions*, 23, 231–234.  
683 <https://doi.org/10.1111/ddi.12517>
- 684 Flantua, S. G., O’dea, A., Onstein, R. E., Giraldo, C., & Hooghiemstra, H. (2019). The  
685 flickering connectivity system of the north Andean páramos. *Journal of Biogeography*.  
686 <https://doi.org/10.1111/jbi.13607>
- 687 Frantzen, N. M. L. H. F., & Bouman, F. (1989). Dispersal and growth form patterns of some  
688 zonal páramo vegetation types. *Acta botanica neerlandica*, 38, 449–465.  
689 <https://doi.org/10.1111/j.1438-8677.1989.tb01376.x>
- 690 Graae, B. J., Vandvik, V., Armbruster, W. S., Eiserhardt, W. L., Svenning, J. C., Hylander,  
691 K., ... & Lenoir, J. (2018). Stay or go—how topographic complexity influences alpine plant  
692 population and community responses to climate change. *Perspectives in Plant Ecology,*  
693 *Evolution and Systematics*, 30, 41–50. <https://doi.org/10.1016/j.ppees.2017.09.008>
- 694 Guisan, A., & Thuiller, W. (2005). Predicting species distribution: offering more than simple  
695 habitat models. *Ecology letters*, 8(9), 993–1009. [https://doi.org/10.1111/j.1461-](https://doi.org/10.1111/j.1461-0248.2005.00792.x)  
696 [0248.2005.00792.x](https://doi.org/10.1111/j.1461-0248.2005.00792.x)
- 697 Guisan, A., & Rahbek, C. (2011). SESAM—a new framework integrating macroecological  
698 and species distribution models for predicting spatio-temporal patterns of species



- 699 assemblages. *Journal of Biogeography*, 38(8), 1433–1444. <https://doi.org/10.1111/j.1365->  
700 2699.2011.02550.x
- 701 Guisan, A., Mod, H. K., Scherrer, D., Münkemüller, T., Pottier, J., Alexander, J. M., &  
702 D'Amen, M. (2019). Scaling the linkage between environmental niches and functional traits  
703 for improved spatial predictions of biological communities. *Global Ecology and*  
704 *Biogeography*. <https://doi.org/10.1111/geb.12967>
- 705 Hattab, T., Garzón-López, C. X., Ewald, M., Skowronek, S., Aerts, R., Horen, H., ... &  
706 Lenoir, J. (2017). A unified framework to model the potential and realized distributions of  
707 invasive species within the invaded range. *Diversity and Distributions*, 23, 806–819.  
708 <https://doi.org/10.1111/ddi.12566>
- 709 Hellmann, J. J., Byers, J. E., Bierwagen, B. G., & Dukes, J. S. (2008). Five potential  
710 consequences of climate change for invasive species. *Conservation biology*, 22(3), 534–543.  
711 <https://doi.org/10.1111/j.1523-1739.2008.00951.x>
- 712 Herrmann, J. D., Carlo, T. A., Brudvig, L. A., Damschen, E. I., Haddad, N. M., Levey, D.  
713 J., ... & Tewksbury, J. J. (2016). Connectivity from a different perspective: comparing seed  
714 dispersal kernels in connected vs. unfragmented landscapes. *Ecology*, 97(5), 1274–1282.  
715 <https://doi.org/10.1890/15-0734.1>
- 716 Hofstede, R., Segarra, P., & Vásconez, P. M. (2003). *Los páramos del mundo: Proyecto*  
717 *Atlas Mundial de los Páramos*. Quito, Ecuador: UICN Global Peatland Initiative;  
718 Ecociencia.
- 719 Holloway, P., Miller, J. A., & Gillings, S. (2016). Incorporating movement in species  
720 distribution models: how do simulations of dispersal affect the accuracy and uncertainty of  
721 projections?. *International Journal of Geographical Information Science*, 30(10), 2050–  
722 2074. <https://doi.org/10.1080/13658816.2016.1158823>
- 723 Hupp, N., Llambí, L. D., Ramírez, L., & Callaway, R. M. (2017). Alpine cushion plants have  
724 species-specific effects on microhabitat and community structure in the tropical Andes.  
725 *Journal of vegetation science*, 28, 928–938. <https://doi.org/10.1111/jvs.12553>
- 726 Kammer, P. M., Schöb, C., & Choler, P. (2007). Increasing species richness on mountain  
727 summits: Upward migration due to anthropogenic climate change or re-colonisation? *Journal*  
728 *of Vegetation Science*, 18, 301–306. <https://doi.org/10.1111/j.1654-1103.2007.tb02541.x>
- 729 Karger, D. N., Conrad, O., Böhrer, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., ... &  
730 Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas.  
731 *Scientific data*, 4, 170122. <https://doi.org/10.1038/sdata.2017.122>
- 732 Kinlan, B. P., & Gaines, S. D. (2003). Propagule dispersal in marine and terrestrial  
733 environments: a community perspective. *Ecology*, 84(8), 2007–2020.  
734 <https://doi.org/10.1890/01-0622>
- 735 Le Roux, P. C., & McGeoch, M. A. (2008). Rapid range expansion and community  
736 reorganization in response to warming. *Global Change Biology*, 14, 2950–2962.  
737 <https://doi.org/10.1111/j.1365-2486.2008.01687.x>
- 738 Lenoir, J., & Svenning, J. C. (2015). Climate-related range shifts—a global multidimensional  
739 synthesis and new research directions. *Ecography*, 38, 15–28.  
740 <https://doi.org/10.1111/ecog.00967>

- 741 León-Yáñez, S. (Ed.). (2012). *Libro rojo de las plantas endémicas del Ecuador*. Quito,  
742 Ecuador: Herbario QCA, Pontificia Universidad Católica del Ecuador.
- 743 Liu, C.R., Berry, P.M., Dawson, T.P. & Pearson, R.G. (2005) Selecting thresholds of  
744 occurrence in the 551 prediction of species distributions. *Ecography*, 28, 385–393.  
745 <https://doi.org/10.1111/j.0906-7590.2005.03957.x>
- 746 Llambí, L. D., Hupp, N., Saez, A., & Callaway, R. (2018). Reciprocal interactions between  
747 a facilitator, natives, and exotics in tropical alpine plant communities. *Perspectives in Plant*  
748 *Ecology, Evolution and Systematics*, 30, 82–88. <https://doi.org/10.1016/j.ppees.2017.05.002>
- 749 Marmion, M., Parviainen, M., Luoto, M., Heikkinen, R. K., & Thuiller, W. (2009).  
750 Evaluation of consensus methods in predictive species distribution modelling. *Diversity and*  
751 *Distributions*, 15, 59–69. <https://doi.org/10.1111/j.1472-4642.2008.00491.x>
- 752 Mateo, R. G., Mokany, K., & Guisan, A. (2017). Biodiversity models: what if unsaturation  
753 is the rule?. *Trends in ecology & evolution*, 32, 556–566.  
754 <https://doi.org/10.1016/j.tree.2017.05.003>
- 755 Melcher, I. M., Bouman, F., & Cleef, A. M. (2000). Seed dispersal in páramo plants:  
756 epizoochorous and hydrochorous taxa. *Plant Biology*, 2, 40–52. [https://doi.org/10.1055/s-](https://doi.org/10.1055/s-2000-9146)  
757 [2000-9146](https://doi.org/10.1055/s-2000-9146)
- 758 Melcher, I. M., Bouman, F., & Cleef, A. M. (2004). Seed atlas of the monocotyledonous  
759 genera of the páramo. *Flora-Morphology, Distribution, Functional Ecology of Plants*, 199,  
760 286–308. <https://doi.org/10.1078/0367-2530-00157>
- 761 Mod, H. K., le Roux, P. C., Guisan, A., & Luoto, M. (2015). Biotic interactions boost spatial  
762 models of species richness. *Ecography*, 38, 913–921. <https://doi.org/10.1111/ecog.01129>
- 763 Moles, A.T., Falster, D.S., Leishman, M.R. & Westoby, M. (2004) Small-seeded species  
764 produce more seeds per square metre of canopy per year, but not per individual per lifetime.  
765 *Journal of Ecology*, 92, 384–396. <https://doi.org/10.1111/j.0022-0477.2004.00880.x>
- 766 Morgan, J. W., & Venn, S. E. (2017). Alpine plant species have limited capacity for long-  
767 distance seed dispersal. *Plant Ecology*, 218(7), 813–819. [https://doi.org/10.1007/s11258-](https://doi.org/10.1007/s11258-017-0731-0)  
768 [017-0731-0](https://doi.org/10.1007/s11258-017-0731-0)
- 769 Muller-Landau, H. C., Wright, S. J., Calderón, O., Condit, R., & Hubbell, S. P. (2008).  
770 Interspecific variation in primary seed dispersal in a tropical forest. *Journal of Ecology*,  
771 96(4), 653–667. <https://doi.org/10.1111/j.1365-2745.2008.01399.x>
- 772 Naimi, B., Hamm, N. A., Groen, T. A., Skidmore, A. K., & Toxopeus, A. G. (2014). Where  
773 is positional uncertainty a problem for species distribution modelling?. *Ecography*, 37(2),  
774 191–203. <https://doi.org/10.1111/j.1600-0587.2013.00205.x>
- 775 Panetta, A. M., Stanton, M. L., & Harte, J. (2018). Climate warming drives local extinction:  
776 Evidence from observation and experimentation. *Science advances*, 4, eaaq1819.  
777 <https://doi.org/10.1126/sciadv.aaq1819>
- 778 Parolo, G., & Rossi, G. (2008). Upward migration of vascular plants following a climate  
779 warming trend in the Alps. *Basic and Applied Ecology*, 9(2), 100–107.  
780 <https://doi.org/10.1016/j.baae.2007.01.005>

- 781 Pauchard, A., Kueffer, C., Dietz, H., Daehler, C. C., Alexander, J., Edwards, P. J., ... &  
782 Seipel, T. (2009). Ain't no mountain high enough: plant invasions reaching new elevations.  
783 *Frontiers in Ecology and the Environment*, 7, 479–486. <https://doi.org/10.1890/080072>
- 784 Pecl, G. T., Araújo, M. B., Bell, J. D., Blanchard, J., Bonebrake, T. C., Chen, I. C., ... &  
785 Williams, S. E. (2017). Biodiversity redistribution under climate change: Impacts on  
786 ecosystems and human well-being. *Science*, 355, eaai9214.  
787 <https://doi.org/10.1126/science.aai9214>
- 788 Peyre, G. (2020). The fate of páramo plant assemblages in the sky islands of the northern  
789 Andes - Table S1: Species presence-absence data. Dryad dataset.  
790 <https://doi.org/10.5061/dryad.44j0zpc9z>
- 791 Peyre, G., Balslev, H., Martí, D., Sklenář, P., Ramsay, P., Lozano, P., ... & Font, X. (2015).  
792 VegPáramo, a flora and vegetation database for the Andean páramo. *Phytocoenologia*, 45,  
793 195–201. <https://doi.org/10.1127/phyto/2015/0045>
- 794 Peyre, G., Balslev, H., & Font, X. (2018). Phytoregionalisation of the Andean páramo.  
795 *PeerJ*, 6, e4786. <https://doi.org/10.7717/peerj.4786>
- 796 Peyre, G., Balslev, H., Font, X., & Tello, J. S. (2019). Fine-scale plant richness mapping of  
797 the Andean páramo according to macroclimate. *Frontiers in Ecology and Evolution*, 7, 377.  
798 <https://doi.org/10.3389/fevo.2019.00377>
- 799 R Core Team (2019). R: A language and environment for statistical computing. R  
800 Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- 801 Ramírez-Villegas, J., Cuesta, F., Devenish, C., Peralvo, M., Jarvis, A., & Arnillas, C. A.  
802 (2014). Using species distributions models for designing conservation strategies of Tropical  
803 Andean biodiversity under climate change. *Journal for Nature Conservation*, 22, 391–404.  
804 <https://doi.org/10.1016/j.jnc.2014.03.007>
- 805 Sánchez, A., Posada, J. M., & Smith, W. K. (2014). Dynamic cloud regimes, incident  
806 sunlight, and leaf temperatures in *Espeletia grandiflora* and *Chusquea tessellata*, two  
807 representative species of the Andean Páramo, Colombia. *Arctic, Antarctic, and Alpine*  
808 *Research*, 46, 371–378. <https://doi.org/10.1657/1938-4246-46.2.371>
- 809 Sanderson, B.M., Knutti, R. & Caldwell, P. (2015) A Representative Democracy to Reduce  
810 Interdependency in a Multimodel Ensemble. *Journal of Climate*, 28, 5171–5194.  
811 <https://doi.org/10.1175/jcli-d-14-00362.1>
- 812 Scherrer, D., & Körner, C. (2011). Topographically controlled thermal-habitat  
813 differentiation buffers alpine plant diversity against climate warming. *Journal of*  
814 *biogeography*, 38, 406–416. <https://doi.org/10.1111/j.1365-2699.2010.02407.x>
- 815 Skarpaas, O., Stabbe, O. E., Rønning, I., & Sverdrup, T. O. (2004). How far can a  
816 hawk's beard fly? Measuring and modelling the dispersal of *Crepis praemorsa*. *Journal of*  
817 *Ecology*, 92(5), 747–757. <https://doi.org/10.1111/j.0022-0477.2004.00915.x>
- 818 Sklenář, P., & Jørgensen, P. M. (1999). Distribution patterns of páramo plants in Ecuador.  
819 *Journal of Biogeography*, 26, 681–691. <https://doi.org/10.1046/j.1365-2699.1999.00324.x>

- 820 Sklenář, P., & Ramsay, P. M. (2001). Diversity of zonal páramo plant communities in  
821 Ecuador. *Diversity and Distributions*, 7, 113–124. <https://doi.org/10.1046/j.1472-4642.2001.00101.x>
- 823 Sklenář, P., Hedberg, I., & Cleef, A. M. (2014). Island biogeography of tropical alpine floras.  
824 *Journal of Biogeography*, 41, 287–297. <https://doi.org/10.1111/jbi.12212>
- 825 Singer, A., Johst, K., Banitz, T., Fowler, M. S., Groeneveld, J., Gutiérrez, A. G., ... & Meyer,  
826 K. M. (2016). Community dynamics under environmental change: How can next generation  
827 mechanistic models improve projections of species distributions?. *Ecological Modelling*,  
828 326, 63–74. <https://doi.org/10.1016/j.ecolmodel.2015.11.007>
- 829 Smith, S. J., Edmonds, J., Hartin, C. A., Mundra, A., & Calvin, K. (2015). Near-term  
830 acceleration in the rate of temperature change. *Nature Climate Change*, 5, 333.  
831 <https://doi.org/10.1038/nclimate2552>
- 832 Steinbauer, M. J., Grytnes, J. A., Jurasinski, G., Kulonen, A., Lenoir, J., Pauli, H., ... & Wipf,  
833 S. (2018). Accelerated increase in plant species richness on mountain summits is linked to  
834 warming. *Nature*, 556, 231. <https://doi.org/10.1038/s41586-018-0005-6>
- 835 Tamme, R., Götzenberger, L., Zobel, M., Bullock, J. M., Hooftman, D. A., Kaasik, A., &  
836 Pärtel, M. (2014). Predicting species' maximum dispersal distances from simple plant traits.  
837 *Ecology*, 95, 505–513. <https://doi.org/10.1890/13-1000.1>
- 838 Team, C. W., Pachauri, R. K., & Meyer, L. A. (2014). IPCC, 2014: climate change 2014:  
839 synthesis report. Contribution of Working Groups I, II and III to the Fifth Assessment Report  
840 of the intergovernmental panel on Climate Change. IPCC, Geneva, Switzerland, 151.
- 841 Thomson, F. J., Moles, A. T., Auld, T. D., & Kingsford, R. T. (2011). Seed dispersal distance  
842 is more strongly correlated with plant height than with seed mass. *Journal of Ecology*, 99(6),  
843 1299–1307. <https://doi.org/10.1111/j.1365-2745.2011.01867.x>
- 844 Thomson, F. J., Letten, A. D., Tamme, R., Edwards, W., & Moles, A. T. (2017). Can  
845 dispersal investment explain why tall plant species achieve longer dispersal distances than  
846 short plant species?. *New Phytologist*, 217(1), 407–415. <https://doi.org/10.1111/nph.14735>
- 847 Thuiller, W., Albert, C., Araújo, M. B., Berry, P. M., Cabeza, M., Guisan, A., ... & Sykes,  
848 M. T. (2008). Predicting global change impacts on plant species' distributions: future  
849 challenges. *Perspectives in plant ecology, evolution and systematics*, 9(3-4), 137–152.  
850 <https://doi.org/10.1016/j.ppees.2007.09.004>
- 851 Thuiller, W., Georges, D., Engler, R., Breiner, F., Georges, M. D., & Thuiller, C. W. (2016).  
852 Package 'biomod2'. Species distribution modeling within an ensemble forecasting  
853 framework <https://CRAN.R-project.org/package=biomod2>.
- 854 Thuiller, W., Guéguen, M., Renaud, J., Karger, D. N., & Zimmermann, N. E. (2019).  
855 Uncertainty in ensembles of global biodiversity scenarios. *Nature communications*, 10,  
856 1446. <https://doi.org/10.1038/s41467-019-09519-w>
- 857 Tovar, C., Arnillas, C. A., Cuesta, F., & Buytaert, W. (2013). Diverging responses of tropical  
858 Andean biomes under future climate conditions. *PloS one*, 8, e63634.  
859 <https://doi.org/10.1371/journal.pone.0063634>

- 860 Travis, J. M., Delgado, M., Bocedi, G., Baguette, M., Bartoń, K., Bonte, D., ... &  
861 Saastamoinen, M. (2013). Dispersal and species' responses to climate change. *Oikos*,  
862 122(11), 1532–1540. <https://doi.org/10.1111/j.1600-0706.2013.00399.x>
- 863 Urrutia, R., & Vuille, M. (2009). Climate change projections for the tropical Andes using a  
864 regional climate model: Temperature and precipitation simulations for the end of the 21st  
865 century. *Journal of Geophysical Research: Atmospheres*, 114(D2).  
866 <https://doi.org/10.1029/2008JD011021>
- 867 Visser, M. E., & Both, C. (2005). Shifts in phenology due to global climate change: the need  
868 for a yardstick. *Proceedings of the Royal Society B: Biological Sciences*, 272, 2561–2569.  
869 <https://doi.org/10.1098/rspb.2005.3356>
- 870 Vittoz, P., & Engler, R. (2007). Seed dispersal distances: A typology based on dispersal  
871 modes and plant traits. *Botanica Helvetica*, 117, 109–124. [https://doi.org/10.1007/s00035-](https://doi.org/10.1007/s00035-007-0797-8)  
872 [007-0797-8](https://doi.org/10.1007/s00035-007-0797-8)
- 873 Vittoz, P., Dussex, N., Wassef, J., & Guisan, A. (2009) Diaspore traits discriminate good  
874 from weak colonisers on high-elevation summits. *Basic and Applied Ecology*, 10, 508–515.  
875 <https://doi.org/10.1016/j.baae.2009.02.001>
- 876 Wisz, M. S., Pottier, J., Kissling, W. D., Pellissier, L., Lenoir, J., Damgaard, C. F., ... &  
877 Svenning, J. C. (2013). The role of biotic interactions in shaping distributions and realised  
878 assemblages of species: implications for species distribution modelling. *Biological reviews*,  
879 88, 15–30. <https://doi.org/10.1111/j.1469-185X.2012.00235.x>

880 **Tables**

881 **Table 1:** Evaluation metrics and their standard deviation (in parenthesis) showing the overall  
 882 performance of the GLM and RF models for 435 plant species as well as their comparison  
 883 between algorithm according to a paired t-student test.

	<b>TSS</b>	<b>Sensitivity</b>	<b>Specificity</b>	<b>Percentage of models kept (over 50 runs)</b>
<b>GLM</b>	0.574 (0.17)	79.483 (11.85)	81.520 (13.03)	23.171 (15.79)
<b>RF</b>	0.652 (0.16)	82.171 (10.40)	80.477 (10.40)	28.891 (15.73)
<b>Significant differences between algorithms</b>	SI (t = -14.487, df = 432, p- value < 2.2e-16)	SI (t = -4.1044, df = 432, p-value = 4.846e-05)	NO (t = 1.6303, df = 432, p-value = 0.1038)	SI (t = -11.812, df = 432, p-value < 2.2e-16)

884

885 **Table 2:** Composition changes (in %) in terms of proportions of mid-páramo and super-  
 886 páramo species at the sky island level between 2000 and 2100. The final super-páramo  
 887 balance shows net gain (in green) or loss (in red) of super-páramo species, expressed as a  
 888 percentage.

	2000		2100			Final super-páramo balance
	Mid-páramo species	Super-paramo species	Mid-páramo species	Super-páramo species	Standard deviation	
<b>Altar</b>	66.91	33.09	74.54	25.46	4.15	-7.63
<b>Antisana</b>	67.30	32.70	75.30	27.70	6.67	-8.01
<b>Cajas</b>	81.82	18.18	84.21	15.79	6.10	-2.39
<b>Cayambe</b>	69.33	30.67	68.56	31.44	4.16	+0.78
<b>Chimborazo</b>	64.47	35.53	71.15	28.85	2.40	-6.68
<b>Cotopaxi</b>	67.11	32.89	71.16	28.84	1.72	-4.04
<b>Ilinizas</b>	59.78	40.22	69.55	30.45	7.35	-9.76
<b>Pichincha</b>	69.90	30.10	73.77	26.23	13.44	-3.87
<b>Sangay</b>	75.51	24.49	75.99	24.01	5.33	-0.48
<b>Tungurahua</b>	65.75	34.25	71.92	28.08	6.99	-6.17

889

890

891 **Figure captions**

892 Figure 1: Current potential distribution of the páramo in Ecuador (A): in dark gray, mid-  
893 páramo (3000-4200 m) and in red, super-páramo (> 4200 m); and characteristic super-  
894 páramo vegetation (B), from bottom to top, increasing with elevation.


895 Figure 2: Examples of distribution responses of páramo species to climate change by 2100,  
896 according to the CIMC5-RCP45-CESM1-BGC scenario: A) a mid-páramo species  
897 colonizing higher elevations (*Vicia andicola*) and B) a super-páramo species reducing its  
898 distribution (*Astragalus geminiflorus*).

899 Figure 3: Net richness changes at the pixel level (1 km<sup>2</sup>) between 2000 and 2100 in the 10  
900 páramo sky islands above 4200 m (named in bold). Between parenthesis, balance of gained  
901 and lost species between the present and 2100 (in percentage of the original richness at the  
902 sky island level).




903 **Figures**

904 Figure 1




905

906 Figure 2



907

908 Figure 3



909


910 **Supplementary materials**

911 Table S1: vegetation and occurrence data used in this study (Peyre, 2020)

912 Table S2: List of species included in the analysis, including their representation in the final  
913 dataset, mid-páramo versus super-páramo status and calculated maximum dispersal distance  
914 (in m/yr).

915 Tabla S3: Species composition of the Ecuadorian sky islands for 2000 and 2100 according  
916 to the modelling analyses and under the different climate change scenarios.


917 Figure S1: Distribution of the VegParamo plots (yellow) and additional data points (in blue)  
918 across the study area.



919


920

921 Figure S2: Density plot of the dispersal capacity, as maximum dispersal distance, of the  
922 páramo species (m/yr), differentiating the mid-páramo and super-páramo species.  
923



924

925 Figure S3: Balance of gained and lost species on the Ecuadorian sky islands between the  
926 present and 2100 (in percentage of the original richness at the sky island level). Dark colours  
927 represent lost species and light colours gained species.



928