

Submarine landslides around volcanic islands: A review of what can be learned from the Lesser Antilles Arc

Anne Le Friant, Elodie Lebas, Morgane Brunet, Sara Lafuerza, Matthew Hornbach, Maya Coussens, Sebastien Watt, Michael Cassidy, Peter J. Talling

▶ To cite this version:

Anne Le Friant, Elodie Lebas, Morgane Brunet, Sara Lafuerza, Matthew Hornbach, et al.. Submarine landslides around volcanic islands: A review of what can be learned from the Lesser Antilles Arc. Kei Ogata, Andrea Festa, Gian Andrea Pini. Submarine Landslides: Subaqueous Mass Transport Deposits from Outcrops to Seismic Profiles, American Geophysical Union, 2020, 10.1002/9781119500513.ch17. hal-03014699

HAL Id: hal-03014699 https://hal.science/hal-03014699v1

Submitted on 19 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	
2	Submarine landslides around volcanic islands:
3	A review of what can be learnt from the Lesser Antilles Arc
4	
5	Le Friant, A., Lebas E., Brunet, M., Lafuerza, S., Hornbach, M., Coussens, M, Watt, S.F.L.
6	Cassidy, M.J, Talling, P.J. and IODP 340 Expedition scientists.
7	
8	
9	
LO L1	Corresponding author:
12	A. Le Friant
L3	Institut de Physique du Globe de Paris, Sorbonne Paris Cite, CNRS UMR 7154, 1, rue
L4	Jussieu, FR-75238 Paris, France (lefriant@ipgp.fr)
L5	
L6	
L7 L8	
LO	

Abstract

IODP Expedition 340 successfully drilled, for the first time, large, and likely tsunamigenic, volcanic island-arc landslide deposits. These cores provide evidence and tests of previous hypotheses for the composition, origin and mode of transport of those deposits. Sites in the medial to distal parts of the landslide deposits offshore Montserrat and Martinique recovered seafloor sediment, comprising turbidites and hemipelagic deposits, and lacked the coarse and chaotic subaerial volcanic debris avalanche material. This supports the concepts that i/ the volcanic debris avalanche component of these landslides is restricted to proximal areas, and tend to stop at the slope break, and ii/ emplacement of volcanic debris avalanches in marine settings can trigger widespread and voluminous failures of pre-existing, low-gradient seafloor sediment. The most likely mechanism for generating these large-scale seafloor sediment failures appears to be the propagation of a décollement, from proximal areas that are loaded and incised by a volcanic debris avalanche. These results have implications for the magnitude of tsunami generation by volcanic island landslides. Volcanic island landslides comprised of mainly seafloor sediment may form smaller magnitude tsunamis than equivalent volumes of subaerial block-rich mass flows rapidly entering water.

Key words: Landslide, volcanic island, debris avalanche, seafloor sediment failure, tsunami, IODP

39 1. Introduction

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

Prodigious submarine landslides offshore volcanic islands have been widely recognized worldwide, such as offshore the Hawaiian Archipelago (Lipman et al., 1988; Moore et al., 1989), La Réunion (Labazuy, 1996; Le Friant et al., 2011; Oehler et al., 2008), and in the Canary Islands (Holcomb & Searle, 1991; Krastel et al., 2001; Masson et al., 2002, 2006; Urgeles et al., 1997; Watts & Masson, 1995). Individual landslide deposits can contain hundreds to thousands of cubic kilometers of material and can extend over a distance of hundreds of kilometers (Moore et al., 1989). Over the last decades, increasing numbers of marine geophysical surveys, along with modern marine geophysical techniques, have revealed our ability to identify such deposits around volcanic islands, recognizing flank collapses as a common and important process during the evolution of volcanic edifices (Glicken, 1986; Ida & Voight, 1995; McGuire, 1996; Voight, 2000). The landslide deposits, recognized both in oceanic intraplate setting or subduction zone context, are related to landslides that affected the subaerial and/or submarine parts of the volcanoes, and have similar features suggesting that common processes control volcano instability and landslide behavior across all settings. These catastrophic events represent a range of significant volcanic hazards, which also includes the related explosive eruptions (Siebert, 1984; Glicken, 1986) and tsunamis (Løvholt et al., 2008), which can be triggered by relatively small events. Understanding where submarine landslide deposits are coming from and how they were emplaced offshore is crucial, since emplacement dynamics determine the magnitude of the associated tsunamis. The potential magnitude of volcanic landslide-generated tsunamis is the subject of an intense debate, mainly due to uncertainties associated with slide physics and dynamics to describe how collapsed material enters the ocean (Cassidy et al., 2014; Hunt et al., 2011; Løvholt et al., 2008; Masson et al., 2006; Mattioli et al., 2007; Slingerland & Voight, 1979; Ward & Day, 2001; Watt et al., 2012a,b, 2014). While some studies indicate that volcanic landslides often generate large (10-100 m high) tsunamis (e.g., Frohlich et al., 2009; Nomanbhoy & Satake, 1995; Ward & Day, 2003), other studies (Gardner, 2010; Hornbach et al., 2015; Moore et al., 1994) suggest that many of these volcanic landslides may deform incrementally, via creep, and therefore represent only a limited tsunami hazard. These mechanisms – creep versus catastrophic failure – are not mutually exclusive, as both might occur during volcanic flank collapse (e.g., McGuire, 1996). Unfortunately, the very few direct observations of volcanic island landslide processes are limited to small historical events (e.g., Stromboli, Tinti et al., 2006) and we are not able yet to directly monitor submarine landslide processes. Studies on landslide deposits geometry allow

some authors to propose conceptual models on the emplacement dynamics of submarine landslides. Bull et al. (2009) presented a review of kinematic indicators from mass transport complexes, while Frey-Martinez et al. (2006) defined two categories of submarine landslides depending on the existence of an emergent frontal ramp. Watt et al. (2014) focused on volcanicisland landslides and provided new constraints and models on the emplacement dynamics for down-slope propagation of submarine deformation, for different scenarii: (i) erosion and loading of seafloor sediment by volcanic debris avalanche (e.g., Deplus et al., 2001), (ii) loading of seafloor sediment by an over-running volcanic debris flow (e.g., Watt et al., 2012a,b), (iii) self-loading by down-slope propagating seafloor sediment failure (e.g., Watt et al., 2012a,b, 2014). However, the impacts of volcanic island landslides are not readily assessed from the final landslide deposits without a good knowledge and understanding of the components that make up the deposits.

Previous ODP Legs in Hawaii (ODP Legs 126 and 200: Garcia, 1993; Garcia et al., 1994, 2006) and the Canary Islands (ODP Leg 157: Goldstrand, 1998; Schmincke & Sumita, 1998; Schneider et al., 1997) drilled offshore oceanic intraplate volcanic islands, but the drill sites mainly targeted distal turbidite deposits located away from the seismically chaotic landslide deposits recognized in more proximal settings. The first scientific drilling of volcanic-arc island landslide deposits was done by IODP Expedition 340 offshore the Lesser Antilles Arc (Figure 1). The newly-acquired drilled cores and downhole logging data were combined with seismic reflection data acquired during four previous cruises (AGUADOMAR 1999 – PI: C. Deplus; CARAVAL 2002 – PI: C. Deplus, GWADASEIS 2009 – PI: N. Feuillet, and JC45/46 - PI: P. Talling; e.g., Deplus et al., 2001, Lebas et al. 2011; Le Friant et al., 2003, Watt et al., 2012 a,b) to reinterpret the composition and emplacement dynamics of the large mass-wasting deposits mapped offshore the volcanic islands of Montserrat and Martinique (Figure 1). In this contribution, we discuss the most important outputs of the IODP Expedition 340 concerning (i) the composition of the gigantic chaotic units interpreted as submarine landslide deposits, (ii) the type of deformation and the mechanisms of the landslide generation and propagation onto the seafloor, (iii) the geohazard potential of such systems, and finally (iv) the implications of our results for other volcanic islands. This contribution is mainly based on a review of earlier published material deriving both from the previous seismic and coring campaigns, and the IODP Expedition 340 carried out in 2012 (Brunet, 2015; Brunet et al., 2016, 2017; Cassidy et al., 2013, 2014, 2015; Coussens et al., 2016; Fraas et al., 2016; Hornbach et al., 2015; Jutzeler et al., 2016, 2014; Lafuerza et al., 2014; Le Friant et al., 2013, 2015; Palmer et al., 2016; Wall-Palmer et al., 2014; Watt et al., 2012a,b, 2014).

The following terminology is used in this paper: *Landslide* is used as a general term referring to any type of slope failure and resulting mass-movement, whether initiated onshore or offshore. The term *debris avalanche* is used for a failure comprising material in a mobile flowage-type landslide from a volcanic edifice (Glicken, 1986; Masson et al., 2002, 2006; Siebert, 2002; Voight et al., 1981, 2002, 2012), and the resulting deposits are referred to as *debris avalanches deposits* (DAD). Debris avalanches are distinguished from *slumps*, which are coherent masses of poorly consolidated materials or rocks that move over a relatively short distance downslope, on a concave-up or planar slip surface (e.g., Delcamp et al., 2008). *Debris flows* are water-saturated flows of disaggregated and/or reworked sediment with high sediment concentrations and strong interactions between solid and fluid components (Iverson, 1997). Finally, failures of seafloor sediment located beyond the slope break (i.e., transition between the base of the volcanic edifice and surrounding basin) are referred to as *seafloor sediment failures* (Watt et al., 2012a,b, 2014), and the resulting deposits are referred to as *submarine landslide deposits* (SLD).

2. Previous Work on the Lesser Antilles Arc

The Lesser Antilles Arc results from the subduction of the Atlantic oceanic lithosphere beneath the Caribbean plate (Figure 1), at a recent convergent rate of ~ 2 cm yr⁻¹ (Feuillet et al., 2002). Compared to other subduction zone volcanic chains, magma productivity in the Lesser Antilles Arc is relatively low, and estimated at 3 to 5 km³ Ma⁻¹ km⁻¹ (MacDonald et al., 2000; Sigurdsson et al., 1980; Wadge, 1984). South of Dominica, the volcanic islands are bordered to the west by the 2900 m-deep, back-arc Grenada Basin (Figure 1), which has been a major depocentre for large mass-wasting deposits, volcanogenic turbiditic deposits, pyroclastic flow deposits, and hemipelagic sediment (Boudon et al., 2007; Deplus et al., 2001; Picard et al., 2006; Sigurdsson et al., 1980). Fifty-two individual landslide deposits or flank collapse structures have been identified onshore and offshore on the flanks of the Lesser Antilles Arc volcanoes (Figures 1 and 2) (e.g., Boudon et al., 1984, 1987, 2007; Cassidy et al., 2014; Crutchley et al., 2013; Deplus et al., 2001; Karstens et al., 2013; Komorowski et al., 2005; Le Friant et al., 2002, 2003a,b, 2004, 2009, 2010; Lebas et al., 2011; Trofimovs et al., 2006, 2010, 2013; Vincent et al., 1989; Watt et al., 2012a,b). Combined analysis of marine geophysical datasets collected along the arc has resulted in a series of hypotheses regarding the composition

of the submarine landslide deposits, their origins and emplacement mechanisms. Deplus et al. (2001) and Le Friant et al. (2002, 2003a,b, and 2004) mapped, for the first time, numerous submarine landslide deposits along the arc and proposed that the landslide deposits correspond to DAD originating from flank collapses affecting the volcanoes onshore. The authors suggested that during the flow, the debris avalanche erodes and mobilizes pre-existing seafloor sediment, and thus disturbs underlying pre-existing sedimentary layers. They proposed that the upper part of the seismically chaotic deposits contains both volcanic debris from the original volcano flank collapse and remobilized pre-existing seafloor sediment. In contrast, the lower part would mainly be composed of deformed pre-existing seafloor sediment. Lateral transitions from well-bedded seismic reflections to chaotic facies within the landslide units in the seismic data, indicate that some landslides internally contain substantial components of seafloor-derived sediment (Figure 2). Deplus et al. (2001) also emphasized that a substantial amount of material deriving from the submarine part of the volcanic edifice is commonly involved in the formation of the deposits, as shown by the chutes cutting through the submarine flanks, caused by removal of volcanic debris during avalanches incising the chutes. Le Friant et al. (2004) integrated this information to outline how terrestrial flank-collapse scars, submarine chutes and offshore debris avalanche deposits were linked to the geomorphological evolution of Montserrat. Moreover, Le Friant et al. (2004) and Lebas et al. (2011) also used seismic data to define the extent and volume of the landslide deposits offshore Montserrat. They demonstrated that the volume of seismic chaotic units associated with landslide deposits may be much greater than the volume of the associated subaerial flank-collapse scars, thus requiring incorporation of a significant proportion of the volcano's submerged flank, or of the adjacent seafloor sediment into the deposits. Truncation of in-situ bedded sediment packages was attributed to erosional processes during debris avalanche emplacement. Lebas et al. (2011) also proposed that strong seismic reflectors seen at the base of some of the landslide deposits may result from slide-derived shear zones. Watt et al. (2012a,b, 2014) proposed that the emplacement of volcanic debris avalanches caused erosion and triggered failure of seafloor sediment. Such seafloor sediment failure could be very widespread on low (< 1°) gradients and may dominate the total landslide volume. Their working hypothesis is consistent with the concept that rapid emplacement of volcanic edifice material onto the seafloor led to undrained shear loading and failure of underlying seafloor sedimentary layers (Voight & Elsworth, 1997; Voight et al., 2012). Watt et al. (2012a,b) further remarked on the lack of frontal compression and shear break-out (e.g., Frey Martinez et al., 2006) offshore Montserrat, which suggested that much of the deformed seafloor material they

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

observed did not move over extensive distances. The authors also proposed that some of the landslide deposits may have incorporated large blocks of relatively intact seafloor sediment.

3. IODP Expedition 340

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

The IODP Expedition 340 conducted aboard the R/V JOIDES Resolution (March-April 2012) aimed at documenting the eruptive history and edifice collapses in the Lesser Antilles Arc over the last million years (Le Friant et al., 2013). The main objective of Expedition 340 was to understand the timing, emplacement processes and dynamics of submarine landslide deposits offshore the Lesser Antilles Arc, with implications for tsunami hazards. Nine sites were drilled offshore Montserrat and Martinique during the IODP Expedition 340 (Figure 1), recovering 434 core sections containing 2,384 m of seafloor sediment samples (Table 1 in Le Friant et al. 2015). Drilling directly offshore Dominica through the most spectacular debris avalanche deposit was not done, because of the size of the blocks, and lessons from previous drilling within blocky deposits offshore Montserrat and Martinique (Le Friant et al., 2013). A wide range of physical properties (thermal conductivity, shear strength, natural gamma ray emissions, P-wave velocities, magnetic susceptibility, and density) have been directly acquired on the cores. Additional information was obtained in-situ by downhole logging operations (gamma rays, P-wave velocities, S-wave velocities, magnetic susceptibility, and resistivity; Lafuerza et al., 2014; Le Friant et al., 2013; Manga et al., 2012; Wall-Palmer et al., 2014). The cores contain some of the most sand-rich sequences yet recovered by an IODP Expedition, where piston coring disturbances such as basal flow-in and fall-in are the most likely to occur, thus affecting the original texture of the sediment (Jutzeler et al., 2014). Particular attention has been paid to distinguishing disturbed sediment facies from in-situ facies. To minimize these coring disturbances and provide better stratigraphic correlations between the sections, a minimum of two holes were drilled at each site (cf. Table 1 in Le Friant et al., 2015).

Sites U1393, U1394, U1399, U1400 and U1401, which will be the focus of the paper, were selected to investigate the landslide deposits, their origins and associated mechanisms of emplacement; sites U1396 and U1397 to study the eruptive history and magmatic evolution of the volcanic centres; and sites U1395 and U1398 to study the differences existing in the sedimentation processes in distal environments between the northern and southern part of the arc (Figures 1, 2, and 3; Le Friant et al., 2015; Wall-Palmer et al., 2014). As mentioned above, attempts to drill through the chaotic proximal landslide Deposit 1 offshore Montserrat (U1393) and Deposit 3 offshore Martinique (U1401) were unsuccessful due to the heterogeneous and cohesionless character of the deposits. The only material recovered at these sites consisted of a

combination of hemipelagic mud with interbedded tephra layers and/or volcaniclastic turbiditic deposits, overlying (and therefore post-dating) the chaotic deposits (Le Friant et al., 2013, 2015; Brunet et al., 2016). A more distal site offshore Montserrat (U1394), intersected a widely dispersed landslide. Deposit 2, identified as a chaotic facies in the seismic reflection data with discontinuous local bedded reflections (Figure 2b). The core showed this apparently chaotic deposit to be dominated by a stacked sequence of predominantly thick, massive, and coarsegrained turbiditic deposits, ranging from bioclastic to volcanoclastic in composition. However, it should be noted that core recovery was restricted to the lower half of Deposit 2. No signs of deformation were recognized (Coussens et al., 2016; Le Friant et al., 2015). Sediment retrieved from Sites U1399 and U1400 (both offshore Martinique), in the medial and distal parts of widely dispersed landslide deposits (D1 and D2), consisted of deformed seafloor sediment. This sediment was mainly composed of hemipelagic mud, with interbedded tephra layers and volcaniclastic turbiditic deposits (Figures 2 and 3). Deformed sediment at Site U1400 is characterized by many planar beds inclined in average by 40°, with some dips being as steep as 70°. Interestingly and surprisingly, these widely dispersed landslide deposits D1 and D2 were not unambiguously identified in the cores retrieved in Site 1400, since no difference in composition was observed between the two landslides. Consequently, Brunet et al. (2016) suggested that they belong to the same deposit, and referred to it as SLD. Of particular significance is the absence of debris avalanche deposits comprising blocky material from the subaerial volcanic edifice in the cores.

More details about the lithologies encountered in the deposits both offshore Montserrat and Martinique can be found in Brunet et al. (2016), Coussens et al. (2016), Fraas et al. (2016), Jutzeler et al. (2014, 2016), Lafuerza et al. (2014), Le Friant et al. (2015), Palmer et al. (2016), and Wall-Palmer et al. (2014).

4. New Model of Landslides Generation Offshore the Lesser Antilles Arc

4.1. Two Different Instability Processes

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

Two different types of landslide processes were recognized along the Lesser Antilles Arc: (i) the subaerial instabilities, which correspond to volcanic flank collapses that affected the subaerial part of edifices, generating DAD accumulating offshore, and (ii) the submarine instabilities, which correspond to seafloor sediment failures at the origin of the large SLD identified in the seismic data.

The Deposit 1 offshore Montserrat (Site U1393) and the Deposit 3 offshore Martinique (Site U1401) (Figure 1) display very blocky surface morphologies, which were difficult to drill. They are located on the submarine flanks of the volcanic edifices or near the slope break, and typically characterized by hummocky topography in the bathymetry (e.g., Deplus et al., 2001; Le Friant et al., 2003, 2004, Moore et al., 1989, Urgeles et al., 1999). These landslide deposits correspond to debris avalanches associated to volcanic flank collapses as recognized on land (e.g., Mount St Helens, Glicken, 1986) or offshore (Canary Islands, Urgeles et al., 1999).

The seismically chaotic deposits drilled at Sites U1394 (Deposit 2, Montserrat), U1399 and U1400 (SLD, Martinique) reveal, unexpectedly, the absence of volcanic debris avalanche deposits, though the deposits are located in the prolongation of the flank-collapse structures (horseshoe shapes) observed on the volcanoes. The chaotic deposits correspond to the second type of instabilities recognized along the arc, i.e. submarine landslides. Characteristics of these deposits (i.e., composition, deformation, vertical truncation with adjacent sediments) are consistent with seafloor sediment failure deposits.

A model is proposed to explain how the subaerial volcanic flank collapses are genetically linked to the widespread sediment failures, with respect to the evolution of the volcanoes (Watt et al., 2012b, Brunet et al., 2016; Le Friant et al., 2015, Watt et al., 2012b).

4.2. Décollement Propagation from Debris Avalanche Deposits Loading

Given the spatial relationship between the SLDs, DADs and collapse scars (subaerial and/or submarine), the new information provided by the IODP Expedition 340 allows us to test the previous hypotheses of emplacement dynamics that have been proposed for these submarine landslides (cf. Figure 9 in Le Friant et al., 2015 and Deplus et al., 2001, Watt et al., 2012 a,b,2014). Results of the combined analysis of these datasets (i.e., spatial relationship between the deposits and the collapse structures, composition, type and degree of deformation of the material within the slide units, vertical truncation of the sediments on the seismic reflection profiles, physical properties providing the first insight into the in-situ stress state at the décollement and data on pore pressure) suggest that the mechanisms producing the large seafloor sediment failures observed offshore the Lesser Antilles Arc may have been triggered by subaerial and/or submarine flank-collapses. We proposed a model of décollement propagation from DADs loading to explain the existence of the large-scale seismically chaotic submarine landslides (Figure 4). Previous modelling from Viesca & Rice (2012) suggested that local failure can generate cracks which propagate rapidly over much wider areas than the initial site of failure, similar to thrust-fault and décollement mechanics, and snow slab avalanches.

Lateral propagation can even occur where the load, in our case a debris avalanche, cannot be dissipated due to the low permeability of the seafloor sediment. This undrained load may generate excess pore fluid pressure and trigger sediment failure in slope and basin areas, which were not originally prone to failure. Our IODP studies provide the first insight into the in situ stress state at the décollement surface (Lafuerza et al., 2014; Hornbach et al., 2015) and indicate both undercompacted and elevated pore fluid pressures in clay-rich layers (e.g., Lafuerza et al. 2014), as well as near-lithostatic pressures in sand-rich turbidites at the décollement (Hornbach et al., 2015). From this, we suggest the sediment at the décollement may be near-critically stressed. As a result, one hypothesis is that any small change in pore pressure generated by compaction or minor grain reorganization (due, for example, to shaking via small earthquakes) might trigger additional failure along the décollement (Hornbach et al., 2015). Current underconsolidated sediment at sites U1400 and U1399 would suggest that these excess pore pressures have not been fully dissipated (Figure 6). Lateral loading induced by the debris avalanche, which may have eroded the seafloor, can also promote overpressure. Basal décollement could thus propagate laterally from the emplaced volcanic debris avalanche (Figure 4) to produce widespread deformation of the seafloor sediment above the décollement surface.

This model in which the décollement propagates from proximal regions loaded and incised by the emplacement of a volcanic debris avalanche may also explain the large-scale landslide deposits mapped in the southern part of the Lesser Antilles Arc, from Dominica to St Vincent (Figure 1, Deplus et al., 2001). If so, this process may operate at the scale of the entire arc. Whether these failures occur rapidly and catastrophically over a large area, or, instead, incrementally over a smaller area is unclear, and remains a topic of ongoing lab and numerical research. Ongoing mechanical modelling with BASIN software, and consequent sediment deformation and failure with PLAXIS software (e.g., Llopart et al., 2018), have demonstrated overpressure formation by undrained loading. In addition with analogue modelling (Brunet, 2015), these studies represent an important topic of on-going and future research which will improve the understanding of the décollement propagation in the near future.

4.3. Evolution of the Montagne Pelée Volcano Through the Landslides Processes (Martinique)

The volcanic history of Martinique Island began at 23 Ma (Germa et al., 2010; Westercamp & Traineau, 1983) and has proceeded through different phases that formed several volcanic centres, including the well-known Montagne Pelée, where three major flank collapses removed

much of the western flank (Boudon et al., 2005, 2007; Brunet et al., 2016; Le Friant et al., 2003 a,b). The first flank collapse occurred at approximately 127 ka (Germa et al., 2010, from onland investigations) leaving behind the first horseshoe-shaped structure on the volcanic flank, with a collapsed volume estimated at 25 km³ (Le Friant et al., 2003a) (Figure 5). The associated debris avalanche entered the sea (DAD 1; Figure 5), flowed down the submarine flank and was deposited near the slope break. This first and largest DAD loaded the pre-existing seafloor sediment and initiated failure. The resulting submarine landslide propagated along a décollement surface, deforming an in situ succession of alternating hemipelagic sediment and turbiditic deposits. Distributed shearing at multiple levels within the sequence may have acted as potential sliding surfaces within the SLD (1 and 5; Brunet et al., 2016). The SLD translated across the seafloor and stopped in a confined way.

The second flank collapse that affected the Montagne Pelée followed the construction of a new, voluminous volcanic edifice inside the scar of the first flank collapse, and probably occurred between 45 and 127 ka (Brunet et al., 2016). The collapsed volume has been estimated at about 13 km³ (Le Friant et al., 2003a) and also generated a debris avalanche (DAD2; Figure 5). DAD2 emplaced over DAD1 and may have remobilized unconsolidated surficial sediment within the central part of the SDL deposit, up to the morphological front (Figure 5), and be responsible for the southward drifting of the SLD in its southern part.

During a third event, the newly-formed cone was destroyed again by a flank collapse forming a smaller horseshoe-shaped structure, whose volume of collapsed material was estimated at 2 km³ (Brunet et al., 2016; Le Friant et al., 200 a). The age assigned to the related DAD3 (Figure 5, which corresponds to D3 offshore Martinique in Figure 1), both identified on land and offshore, is still a matter of ongoing discussions and may be dated between 30 and 45 ka based on preliminary measurements of δ^{18} O samples from the IODP Expedition 340 and previously published ages from terrestrial deposits (Table 2 in Le Friant et al., 2015). As for DAD2, DAD3 (clearly identified on the submarine flank by its hummocky morphology) may have locally remobilized sediments within SLD, by exerting an undrained load that supplied the deformation process.

Particularly noteworthy is that the older DAD1 and DAD2 from the conceptual model of Figure 5 were not previously identified offshore on Figure 1. The seismic data in steeper proximal areas, which represent the submarine flank of the island, are of poor quality and equivocal, although hyperbolic reflections could indicate the presence of some blocks (Figure 2a). However, the bathymetric data show a bulge at the base of the volcanic submarine flank,

near the slope break, at the same location as DAD3. Thus, Brunet et al. (2016) proposed that this bulge may result from the accumulation of the older DAD1 and DAD2 related to the first and second flank collapses of the Montagne Pelée (Figure 5).

5. Implications on the Emplacement Mechanisms of the Landslides Deposits

The new model suggests that the large SLDs recognized offshore the Lesser Antilles Arc result from a décollement propagation of the marine sediment due to the emplacement of DADs on top (Le Friant et al., 2015; Watt et al., 2012b, 2014). In the following section, we investigate the emplacement mechanisms of the two types of landslides deposits encountered in the arc (i.e., flank collapses and seafloor sediment failures), by first using a numerical approach and then by integrating geophysical properties and geotechnical constraints resulting from the IODP Expedition 340. We will use the example and the characteristics of the Martinique deposits to present our results.

5.1. Debris avalanches

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

Brunet et al. (2016) and Le Friant et al. (2015) recently proposed that flank collapses having affected the Montagne Pelée volcano produced debris avalanches that flowed down the volcanic edifice and the submarine flank, to stop around the slope break. This new scenario implies a much shorter runout distance of the debris avalanches than previously suggested (Boudon et al., 2007; Deplus et al., 2001; Le Friant et al. 2003a) and it has consequences for understanding debris avalanche dynamics. To confirm this scenario, numerical simulations have been performed by Brunet et al. (2017) using two complementary, depth-averaged thin-layer continuum models, since no complete models were available in the literature at the time of the simulations. The first model used, SHALTOP (Mangeney-Castelnau et al., 2003), describes precisely the topography. It accurately describes dry granular flows over a 3D topography and may be easily extended to describe the submarine avalanche. The second model, HYSEA (Fernandez-Nieto et al., 2008), has the advantage to take into account the interaction with the water column. It describes the subaerial and the submarine parts of the avalanche. These numerical simulations aimed to determine (i) a critical runout distance for the emplacement of the debris avalanches and (ii) the single or multiphase behavior of the associated flank collapse. Numerical simulations have shown that the first and largest debris avalanche from the

Montagne Pelée (DAD 1: 25 km³) may extend up to 50 km from the coastline, which is not an

observation in the IODP data. The simulations have also demonstrated that the runout distance

of the debris avalanche does not increase with successive flows of smaller volume (about 5 km³

each) and stays around 35 km from the coastline. Brunet et al. (2017) noted that the debris avalanche slowed down after losing inertia to accumulate on the slope transition or relatively nearby, at about 35 km from the coastline. These simulations, along with IODP observations showing that the offshore DAD 1 could only be found up to 35 km from the coastline, led Brunet et al. (2017) to suggest that the first and largest flank collapse that affected the Montagne Pelée may have occurred in several phases. Therefore, debris avalanche volumes for each phase probably did not exceed 5 km³, with smaller runout distances than previously thought (Boudon et al., 2007; Deplus et al., 2001; Le Friant et al. 2003a). Complex turbidite stratigraphies correlated with landslide deposits offshore Montserrat suggest that flank collapses may have occurred in multiple stages (e.g., Cassidy et al., 2014). Indeed, analysis of tephra fall and volcaniclastic turbiditic deposits in drill cores at Sites U1394 and U1395, revealed three heightened periods of volcanic activity on the island (930-900 ka, 810-760 ka, and 190-120 ka), which coincides with periods of increased volcano instability and mass-wasting events (Coussens et al., 2016). The youngest period 190–120 ka marks a peak in the volcanic activity of Soufrière Hills. The largest flank collapse having affected the volcano occurred toward the end of this period (~130 ka) and was followed by two younger events occurring during a period of time where the volcanic activity was relatively elevated (Coussens et al., 2016). Multiphase flank collapses have also been described in the Canarian and Hawaiian Islands, where they generated distinctive turbidites that comprise multiple fining-upward subunits (Garcia & Meyerhoff Hull, 1994; Hunt et al., 2011, 2013; Wynn & Masson, 2003). These observations suggest that flank collapses may occur in multiple stages. Recently, Hunt et al. (2018) related volcano flank failures to volcanic activity (i.e. explosive eruptions) in the Canary islands, and suggested that these multiple stages events might have been separated by days to weeks.

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

5.2. Seafloor Sediment Failures Triggered by Debris Avalanches Loading

Using core logging, sedimentological and geotechnical data from the IODP Expedition 340, Lafuerza et al. (2014) analyzed the uppermost 200 m of cores at Sites U1397, U1398, U1399, and U1400 offshore Martinique. The authors found that shear strength-overburden stress ratios (Su/σ'_v) on most of the hemipelagic sediment tested, are lower than 0.17-0.25 (values on normal consolidation conditions, gray shading in Figure 6). Contrasting with consolidation tests, the authors demonstrated that low hydraulic conductivity of hemipelagic sediment prevents their consolidation and causes low rates of dewatering in turbidity current deposits and tephra layers, allowing excess pore fluid pressures to persist at depth (Figure 6). In all cores, the uppermost

200 m showed clear evidence for excess pore fluid pressures, except at Site U1400 where overpressure only occurred after 180 m downwards. Lafuerza et al. (2014) explained this overpressure by a slow escape of pore fluid during the burial of more permeable turbidity current deposits and tephra layers, sandwiched between minimally permeable hemipelagic sediment. Higher overpressures were recognized at Sites U1397, U1398, and U1399, where higher abundance of thicker turbiditic deposits and coarser tephra layers were described. Pore pressures at Site U1399 were also investigated by Hornbach et al. (2015) using integration of physical properties, downhole logging measurements from the IODP Expedition 340, and rock physics models (Hornbach & Manga, 2014). The authors indicated that high pore pressures existed throughout much of the entire cores at the same locations and concluded that unusual, elevated pore fluid pressures occurred at present in multiple sand-rich zones within the SLD deposit, specifying that only small changes in the current stress regime may trigger motion. In their work, Hornbach et al. (2015) also discussed the most likely plausible mechanisms responsible for creating excess pore pressures at Site U1399, and concluded that shearing processes resulting from regional strain and grain reorganization during sediments contraction, may best explain increasing in pore fluid pressures. As also pointed out by Lafuerza et al. (2014), Hornbach et al. (2015) showed that excess pore pressures reside in thicker sand-rich intervals, while intervening mud-rich zones – within and outside the SLD deposit – show very low permeabilities, thus acting as seals. All these observations suggest that highly elevated pore pressures can persist over long time spans within the SLD deposit, long after the initial failure. This implies that deformation within SLD could continue at present or in the near future (Hornbach et al., 2015). The work presented by Hornbach et al. (2015) is coherent with the observations of multiple, post-deposition deformation zones within cores U1399-A and U1400-C, suggesting the existence of a series of failure prone horizons within SLD and not solely a single layer at the base of the deposit (Brunet et al., 2016). These failure prone horizons may coincide with sand layers, or boundaries between sand and mud layers, which exist in many places inside these drill cores. It is also possible that homogenized muddy sand layers result from previous distributed shearing at multiple levels, and that these muddy sand layers act as pressure seals. Ogata et al. (2014) highlighted the fundamental role of shearing-related liquefaction as one of the main factors controlling slide mobility though the "lubrication" of internal and basal friction forces. Thus, these layers may be regarded as multiple décollement surfaces that, when solicited, may act together to promote and enhance deposit mobility, and longer runout distances.

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

6. Implications for Tsunami Hazards

Assessing landslide-related tsunami hazards ultimately requires detailed knowledge of slide physical properties, volumes, water depths, speeds, and evolution with time (e.g., Masson et al., 2006, Ward & Day, 2001, Watt et al., 2012a). Unfortunately, in many instances, these basic physical properties are very poorly constrained. By providing direct observations and measurements of slide physical properties (such as stress state, thickness, and sediment type), IODP drilling data ground-truths seismic interpretations, improves our understanding of slide physical properties, and from this, enhances our understanding of landslide-derived tsunami hazards.

The results derived from the IODP Expedition 340 confirmed that the landslides identified around the Lesser Antilles volcanoes result from two fundamentally different processes: (i) volcanic flank collapses, generating blocky DADs and (ii) seafloor sediment failures, generating large and voluminous SLDs. Brunet et al. (2017) suggested through numerical simulations that the largest flank collapse that affected the Montagne Pelée volcano may have occurred in several phases, with individual debris avalanches volumes not exceeding 5 km³. The multistage character of the volcanic flank collapses is crucial for evaluating the resulting tsunamigenic hazard not only in Martinique, but also throughout the Lesser Antilles Arc in general. Both types of landslide may generate a tsunami, but the magnitude of the landslide-derived tsunamis will be different.

Using Deposits 1 and 2 offshore Montserrat as examples, Watt et al. (2012a) approximated the tsunami magnitudes that might be expected for the deposits, taking into account different possible triggering, i.e., from a failure of: (i) the subaerial flank of the volcanic edifice, (ii) the submerged slopes of the volcano, or (iii) deep-water seafloor sediment neighboring the volcano. Watt et al. (2012a) used a series of relationships based on theory or extrapolation of small-scale laboratory experiments previously proposed in the literature and whose assumptions are discussed in detail by the authors. A key point is that tsunami magnitude strongly depends on the relative landslide and tsunami wave speeds, with tsunami magnitude increasing if the tsunami wave and the submerged landslide travel at comparable speeds. Such a case will typically only occur in shallow water depths (<200 m), implying that seafloor sediment failures occurring in deep water depths tend to produce much smaller tsunami magnitudes (Ward, 2001). Watt et al. (2012a) suggested that a seafloor sediment failure of ~2 km³ would produce a maximum tsunami amplitude approximately 100 times smaller than one involving a subaerial or submerged failure of a volcanic flank of comparable volume (see Watt et al., 2012a more

particularly Figure 7). Even a seafloor sediment failure with a much larger volume, such as the SLD deposit offshore Martinique (25 km³), would likely produce a much smaller tsunami magnitude than a failure of the subaerial or submerged parts of the volcanic edifice, for a similar volume.

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489 490

491

492

493

494

495

496

Moreover, the analysis in Watt et al. (2012a) of a seafloor sediment failure-related tsunami is based on wave-tank experiments with a sliding block (see Watts et al., 2005). Such a model may not capture the way in which the deformation propagates within the sediment offshore Montserrat and Martinique. It may be that deformation in seafloor sediment failures is slow moving compared to a sliding block, and that individual parcels of seafloor sediment move laterally over short distances. If so, such slow velocities and short particle-transport distances could further decrease the tsunami magnitude induced by seafloor sediment failures. We have calculated the tsunami wave height versus water depth for the different submarine slide volumes (5, 13, 25 km³) and velocities offshore Martinique. We estimate wave heights using the nonlinear shallow-water wave approach (e.g., Jiang & LeBlond, 1992). Although we apply a maximum slide velocity of 50 m s⁻¹, turbidity current velocity measurements derived from cable breaks suggest lower values, typically ranging between 5 and 30 m s⁻¹ (Heezen & Ewing, 1952; Hsu et al., 2008; Krause et al., 1970). Both water depth and slide velocity play an important role in tsunami wave height. The largest wave heights, for a given water depth, typically form when slide velocity approaches the natural shallow water wave velocity defined by \sqrt{gd} , where g is the gravitational acceleration and d, the water depth. Figure 7 shows that for an order-ofmagnitude change either in the slide velocity or in the water depth, an order-of-magnitude change also occurs in the tsunami wave height; with deeper and slower slides generating smaller tsunami wave heights. This implies that slower moving slides in deep water offshore the Lesser Antilles may represent a lower tsunamigenic risk. The block of seafloor sediment recovered completely intact at Site U1394 is consistent with relatively slow and limited lateral movement of the landslide. Offshore Martinique, deformation of seafloor sediment appears to occur at different levels within the deposits, which may indicate somewhat more vigorous motion.

A tsunami wave produced by a widespread seafloor sediment failure may thus produce a longer tsunami wavelength than that associated with localized failure of the volcanic edifice. These longer tsunami wavelengths from seafloor sediment failures could increase the runout distance of the tsunami and the regional risk, since longer-wavelength tsunamis can propagate further with less energy loss (Watt et al., 2012a). However, even if the volume of a flank collapse is smaller than the volume of seafloor sediment failures, it may be the most significant in terms of causing tsunami damage on nearby coastlines due to the velocity of the slide

associated with the flank collapse, the shallow water depth along the coastline, and the location of the failure near the coast, leading to less warning time.

7. Comparison with Submarine Landslides around Other Volcanic Islands

Flank collapses on oceanic-island volcanoes have produced some of the largest landslide deposits recognized on Earth. These giant landslides can contain several hundred to several thousands of cubic kilometers of collapsed material (Masson et al., 2002; Moore et al., 1989), exceeding by far the largest known subaerial DAD of Mount Shasta (45 km³, Crandell, 1989). Criteria for identification of landslide deposits in the submarine area have been developed by several authors in different settings over the last decades. Submarine landslide deposits associated with debris avalanches typically display hummocky terrains on the swath bathymetry, speckled pattern on backscatter images, and a hyperbolic facies on echosounder data (e.g., Deplus et al., 2001; Masson et al., 2002; Moore et al., 1989). On seismic data, these landslide deposits show an acoustically chaotic character with strong energy diffraction, which strongly contrasts with the continuous, subhorizontal and subparallel reflectors that define underformed, well-stratified sedimentary successions.

The most block-rich volcanic-island landslides, such as Nuuanu in Hawaii, are dominated by multiple kilometer-scale blocks corresponding essentially to intact fragments of volcanic edifices that have slid to their site of deposition (Moore & Clague, 2002). The prevalence of angular, intact blocks in some DADs suggests a relative damping of fragmentation in the submarine environment. In the Lesser Antilles Arc, such angular blocky deposits have been observed offshore Dominica (Deplus et al., 2001; Le Friant et al., 2002). Watt et al. (2012a, 2014) suggested that in several instance seafloor sediment substrate probably fails during the emplacement of volcanic landslides (e.g., Antilles: Deplus et al., 2001; Papua New Guinea: Silver et al., 2009; Aleutians: Montanaro & Beget, 2011), and would remain almost in situ as a deformed package. Furthermore, limited lateral transport distances are implied by the geometry of these seafloor sediment failures, in which the toe of landslide transitions does not emerge above the seafloor (Frey-Martinez et al., 2006; Watt et al., 2014). When the seafloor sediment becomes more extensively disaggregated and no longer confined within the region of initial failure, a debris flow can propagate for hundreds of kilometers from the initial point of failure, as observed for the Canary Debris Flow induced by the El Golfo debris avalanche (Hunt et al., 2011; Masson et al., 2002).

Drilling through the seismically chaotic deposits offshore the Lesser Antilles volcanoes confirm that chaotic units observed in seismic data do not necessarily represent extensively disaggregated or deformed material that has been transported over long distances from the volcanic edifice (cf. Gafeira et al., 2010) and supports the view that extensive seafloor sediment failures may be relatively commonplace around volcanic islands (Watt et al., 2012a, 2014). The IODP Expedition 340 also provided a valuable opportunity to compare seismic facies with core observations. Such comparison allowed us to understand the amount, style of deformation and sediment lithologies, that produce the different seismic facies. A key issue for such comparison is the substantial differences in resolution of the two types of data, which ranges from millimeters for the sediment cores to tens to hundreds of meters for seismic data (Figure 3). For instance, we observed a good correlation between the upper and lower limits of the SLD deposit offshore Martinique identified using seismic profiles, and the extent of the deformation within drilled holes (Figures 2 and 3). The well-bedded seismic facies corresponds to turbidity current deposits and/or debris flow deposits, or tephra layers interbedded with hemipelagic sediments, which are flat lying and lack deformation in the cores.

New findings from the IODP Expedition 340 thus shed light on the necessity to revisit previous interpretations of large-scale submarine landslide deposits (SLDs) previously interpreted as resulting from catastrophic, large flank collapses of neighboring volcanic islands as these systems are significantly more complex both in space and time than previously believed. Such an approach has recently been taken by Lebas et al. (2018) offshore the volcanic island of La Réunion (Indian Ocean). The presence of alternating undisturbed to slightly deformed seismic reflectors and chaotic reflections (Figure 8) in the seismically chaotic deposit identified offshore the Piton de la Fournaise volcano, led the authors to reinterpret this "chaotic" deposit as widespread, deformed, pre-existing seafloor sediment. This deposit recorded the emplacement of the Piton de la Fournaise DADs on the submarine flank, where hummocky terrains were previously observed on the bathymetry (Le Friant et al., 2011; Lénat et al., 1989; Oehler et al., 2004, 2008). Lebas et al. (2018) concluded that such large seismically "chaotic" deposits may be regarded as evidence of former flank collapse events offshore other volcanic islands that might not be identifiable on the volcanic submarine flanks (due to the age or the resolution of the data), but that clearly deformed marine sediment offshore.

Acknowledgments

We thank the drillers for their relentless efforts to drill through such difficult volcaniclastic material and all R/V *JOIDES Resolution* staff for their collaboration and efficient work on board. We are very grateful to C. Mével who encouraged and helped us to build our drilling proposal. We thank INSU, NERC, ANR-13-BS06-0009, Labex UnivEarth, Interreg Caraibes PREST, JSPS, NSF for funding analyses and IPGP for funding post-docs and PhD. We thank the staff of the Montserrat Volcano Observatory, the Martinique and Guadeloupe Volcanological and Seismological Observatory, for assistance during the cruise and outreach education activities. Data from the IODP Expedition 340 can be obtained through the expedition's implementing organization USIO (http://iodp.tamu.edu/database). IPGP contribution 3960.

570571

560

561

562

563

564

565

566

567

568

569

References

- Boudon, G., Le Friant, A., Komorowski, J.-C., Deplus, C., Semet, M.P (2007). Volcano flank instability in the Lesser Antilles Arc: diversity of scale, processes, and temporal recurrence. *Journal of*
- 575 Geophysical Research, 112, B08205, doi:10.1029/2006JB004674.
- Boudon, G., Le Friant, A., Villemant, B. & Viodé, J.-P. (2005). Martinique. *In*: J.M. Lindsay, R.E.A.
- Robertson, J.B. Shepherd & S. Ali (Eds), Volcanic Atlas of the Lesser Antilles, Seismic Research
- Unit, The University of the West Indies, Trinidad and Tobago, WI, 65-102.
- Boudon, G., Semet, M.P., Vincent, P.M. (1984). Flank failure-directed blast eruption at Soufrière, Guadeloupe, French West Indies: a 3,000-yr-old Mt. St. Helens? *Geology*, 12, 350-353.
- Boudon, G., Semet, M.P., Vincent, P.M. (1987). Magma and hydrothermally driven sector collapses:
- the 3100 and 11500 Y. B.P. eruptions of La Grande Découverte volcano, Guadeloupe, French West
- 583 indies. *J. Volcanol. Geotherm. Res.*, 33, 317-323.
- Brunet, M., 2015. Caractérisation des processus d'instabilité affectant les îles volcaniques : Application
- au volcan de la Montagne Pelée à partir de l'exploitation des forages de l'Expédition IODP 340.
- 586 PhD thesis, Université Paris 7, France.
- Brunet, M., Le Friant, A., Boudon, G., Lafuerza, S., Talling, P., Hornbach, M., Ishizuka, O., Lebas, E.,
- Guyard, H. and IODP Expedition 340 Science Party (2016). Composition, geometry, and
- emplacement dynamics of a large volcanic island landslide offshore Martinique: From volcano
- flank- collapse to seafloor sediment failure?. Geochem. Geophys. Geosyst., 17-3, 699-724,
- 591 doi:10.1002/2015GC006034.

- Brunet, M., Moretti, L., Le Friant, A., Mangeney, A., Fernandez Nieto, E. (2017). Numerical simulations
- 593 of 30-45 ka debris avalanche flow of Montagne Pelee volcano, Martinique: from volcano flank-
- collapse to submarine emplacement, *Natural Hazards*, vol. 87, I2, p1189-1222.
- Cassidy, M., Trofimovs, J., Palmer, M.R., Talling, P.J., Watt, S.F.L., Moreton, S.G., and Taylor, R.N.
- 596 (2013). Timing and emplacement dynamics of newly recognised mass flow deposits at ~8-12 ka
- offshore Soufrière Hills volcano, Montserrat: how submarine stratigraphy can complement subaerial
- eruption histories. *Journal of Volcanology and Geothermal Research*. v. 253, p. 1–14
- Cassidy, M., Trofimovs, J., Watt, S.F.L., Palmer, M.R., Taylor, R.N., Gernon, T.M., Talling, P.J. Le
- Friant, A. (2014). Multi-stage collapse events in the South Soufrière Hills, Montserrat as recorded
- in marine sediment cores. *Geological Society of London*, v. 39, 383-397.
- Cassidy, M., Watt, S.F.L., Talling, P.J. Palmer, M.R., Edmonds M., Jutzeler, M., Wall-Palmer, D.,
- Coussens, M., Gernon, T., Taylor, R.N., Michalik, A., Inglis, E., McCanta, M.C., Manga, M.,
- Boudon, G., Adachi, T., Breitkreuz, C., Hornbach, M.J., Colas, S.L., Endo, D., Fujinawa, A.,
- Kataoka, K.S., Maeno, F., Tamura, Y., Wang, F., Ishizuka, O., Le Friant, A. and Shipboard Science
- Party, (2015). Rapid onset of mafic magmatism facilitated by volcanic edifice collapse. *Geophys*.
- 607 Res. Lett., vol 42, Issue 12, 4778-4785.
- Coussens, M., Wall-Palmer, D., Talling, P., Watt, S., Cassidy, M., Jutzeler, M., Clare, M., Hunt, J.,
- Manga, M., Gernon, T., Palmer, M., Hatter, S., Boudon, G., Endo, D., Fujinawa, A., Hatfield, R.,
- Hornbach, M., Ishizuka, O., Kataoka, K., Le Friant, A., Maeno, F., McCanta, M., Stinton, A. (2016).
- The relationship between eruptive activity, flank collapse and sea-level at volcanic islands: a long-
- term (>1 Ma) record offshore Montserrat, Lesser Antilles. Geochem. Geophys. Geosyst., vol 17,
- 613 Issue 7, 2591-2611.
- 614 Crandell, D.R., 1989. Gigantic debris avalanche of Pleistocene age from ancestral Mount Shasta
- volcano, California, and debris-avalanche hazard zonation. *United State Geological Survey Bulletin*.
- 616 Crutchley, G.J., Karstens, J., Berndt, C., Talling, P.J., Watt, S.F.L., Vardy, M.E., Hühnerbach, V.,
- 617 Urlaub, M., Sarkar, S., Klaeschen, D., Paulatto, M., Le Friant, A., Lebas, E., Maeno, F. (2013).
- Insights into the emplacement dynamics of volcanic landslides from highresolution 3D seismic data
- 619 acquired offshore Montserrat, Lesser Antilles. *Marine Geology*, 335, 1-15.
- doi:10.1016/j.margeo.2012.10.004.
- Delcamp, A., van Wyk de Vries, B., James, M. (2008). The influence of edifice slope and substrata on
- volcano spreading, J. Volcanol. Geotherm. Res. 177, 4, 925-943.
- Deplus, C., Le Friant, A., Boudon, G., Komorowski J-C., Villemant, B., Harford, C., Ségoufin, J.,
- 624 Cheminée, J-L. (2001). Submarine evidence for large-scale debris avalanches in the Lesser Antilles
- 625 arc. Earth & Planetary Science Letters, 192 (2), 145-157.

- 626 Fraass, A.J., Wall-Palmer, D., Leckie, R.M., Hatfield, R.G., Burns, S.J., Le Friant, A., Ishizuka, O.,
- Aljahdali, M., Jutzeler, M., Martinez-Colon, M., Palmer, M., & Talling, P.J., (2016). A revised Plio-
- Pleistocene age model and paleoceanography of the northeastern Caribbean Sea: IODP Site U1396
- off Montserrat, Lesser Antilles. Stratigraphy, vol13, I3, p183-203.
- 630 Fernandez-Nieto E, Bouchut F, Bresch D, Castro-Diaz MJ, Mangeney A (2008). A new Savage-Hutter
- type model for submarine avalanches and generated tsunami. *J Comp Phys* 227(16):7720–7754
- Feuillet, N., Manighetti, I., Tapponnier, P., Jacques, E. (2002). Arc parallel extension and localization
- of volcanic complexes in Guadeloupe, Lesser Antilles. J. Geophys. Res., 107(B12), 2331,
- doi:1029/2001JB000308.
- Frey-Martinez, J., Cartwright, J., James, D. (2006). Frontally confined versus frontally emergent
- submarine landslides: a 3D seismic characterisation. *Marine and Petroleum Geology*, 23, 585-604.
- Frohlich, C., Hornbach, M.J., Taylor, F.W., Shen, C.C., Moala, A., Morton, A.E. and Kruger, J., 2009.
- Huge erratic boulders in Tonga deposited by a prehistoric tsunami. *Geology*, 37(2), pp.131-134.
- 639 Gafeira, J., Long, D., Scrutton, R., Evans, D. (2010). 3D seismic evidence of internal structure within
- Tampen Slide deposits on the North Sea Fan: are chaotic deposits that chaotic? *Journal of the*
- 641 *Geological Society,* 167, 605-616.
- 642 Garcia, M.O., (1993). Pliocene-Pleistocene volcanic sands from site 842: products of giants landslides.
- 643 Proceedings of the Ocean Drilling Program, Scientific Results, 136: 53-63.
- 644 Garcia, M.O. & Meyerhoff Hull, D. (1994), Turbidites from giant Hawaiian landslides: Results from
- Ocean Drilling Program Site 842. *Geology*, 22, 159-162.
- 646 Garcia, M.O., Sherman, S.B., Moore, G.F., Groll, R., Popova-Goll, I., Natland, J.H., Acton, G. (2006).
- Frequent landslides from Koolau Volcano: Results from ODP Hole 1223A. *Journal of Volcanology*
- and Geothermal Research, 151: 251-268.
- 649 Gardner, J.V., 2010. The West Mariana Ridge, western Pacific Ocean: Geomorphology and processes
- from new multibeam data. *Bulletin*, 122(9-10), pp.1378-1388.
- 651 Germa, A., Quidelleur, X., Labanieh, S., Lahitte, P., Chauvel, C. (2010). The eruptive history of Morne
- Jacob volcano (Martinique Island, French West Indies): Geochronology, geomorphology and
- geochemistry of the earliest volcanism in the recent Lesser Antilles arc. *J. Volcanol. Geotherm. Res.*
- 654 198, 297-310.
- Glicken, H. (1986). Study of the rockslide avalanche of May 18 1980, Mount St Helens Volcano, *PhD*
- 656 Thesis, University of California, Santa Barbara, 303 p.
- 657 Goldstrand, P.M. (1998). Provenance and sedimentologic variations of turbidite and slump deposits at
- sites 955 and 956. Proceedings of the Ocean Drilling Program, Scientific Results, 157: 343-360.

- Heezen, B.C. and Ewing, M., 1952. Turbidity currents and submarine slumps, and the 1929 Grand Banks
- earthquake. *American journal of Science*, 250(12), pp.849-873.
- Holcomb, R., and Searle R., (1991). Large landslides from oceanic volcanoes, Mar. Geotechnol., 10,
- 662 19-32.
- Hornbach, MJ; Manga, M; Genecov, M; Valdez, R; Miller, P; Saffer, D; Adelstein, E; Lafuerza, S;
- Adachi, T; Breitkreuz, C; Jutzeler, M; Le Friant, A; Ishizuka, O; Morgan, S; Slagle, A; Tailing,
- PJ; Fraass, A; Fujinawa, A; Hatfield, R; Kataoka, K; Maeno, F; Martinez-Colon, M; McCanta,
- M; Palmer, M; Stinton, A; Subramanyam, KSV; Tamura, Y; Villemant, B; Wall-Palmer, D; Wang,
- F (2015). Permeability and pressure measurements in Lesser Antilles submarine slides: Evidence
- for pressure-driven slow-slip failure, J. Geophys. Res. Solid Earth, 120-12, 7986–8011,
- doi:10.1002/2015JB012061.
- Hornbach, M. J., and M. Manga (2014). The ability of rock physics models to infer marine in situ pore
- pressure, Geochem. Geophys. Geosyst., 15, 4769–4780, doi:10.1002/2014GC005442.
- Hunt, J.E., Cassidy, M., Talling, P.J., (2018). Multi-stage volcanic island flank collapses with coeval
- explosive caldera-forming eruptions. Scientific reports 8 (1), 1146 doi:10.1038/s41598-018-19285-
- 674 2.
- Hunt, J., E., Wynn, R., B., Masson, D., G., Talling, P.J., teagle, D., A., H. (2011). Sedimentological and
- geochemical evidence for multistage failure of volcanic landslides: A case study from Icod landslide
- on north Tenerife, Canary Islands. Geochemistry Geophysics Geosystems, 12. Q12007. 10.,
- 678 1029/2011GC003740.
- Hunt JE, Wynn RB, Talling PJ, Masson DG (2013). Multistage collapse of eight western Canary
- Island landslides in the last 1.5 Ma: sedimentological and geochemical evidence from subunits in
- submarine flow deposits. *Geochem Geophys Geosyst* 14(7):2159–2181.
- 682 Hsu, S.K., Kuo, J., Lo, C.L., Tsai, C.H., Doo, W.B., Ku, C.Y. and Sibuet, J.C. (2008). Turbidity
- 683 Currents, Submarine Landslides and the 2006 Pingtung Earthquake off SW Taiwan. Terrestrial,
- 684 Atmospheric & Oceanic Sciences, 19(6).
- Ida, Y. and Voight, B. (eds.) (1995). Models of magmatic processes and volcanic eruptions. (Special
- Issue) *Journal of Volcanology and Geothermal Research*, 66, 426pp.
- 687 Iverson, R.M. (1997). The physics of debris flows, *Rev. Geophys.*, v. 35, p. 245-296.
- Jiang, L. and LeBlond, P.H., (1992). The coupling of a submarine slide and the surface waves which it
- generates. *Journal of Geophysical Research: Oceans*, 97(C8), pp.12731-12744.
- Jutzeler, M., J. D. L. White, P. J. Talling, M. McCanta, S. Morgan, A. Le Friant, and O. Ishizuka (2014).
- Coring disturbances in IODP piston cores with implications for offshore record of volcanic events

- and the Missoula megafloods, Geochem. Geochemistry Geophysics Geosystems, 15, doi:10.1002/
- 693 2014GC005447.
- Jutzeler, M., Manga, M., White, J.D.L., Talling, P.J., Proussevitch, A.A., Watt, S.F.L., Cassidy, M.,
- Taylor, R.N., Le Friant, A., Ishizuka, O. (2016). Submarine deposits from pumiceous pyroclastic
- density currents traveling over water: An outstanding example from offshore Montserrat (IODP
- 697 340). GSA Bulletin. Vol 129, Issue 3 4, 392-414, Doi:10.1130/B31448.1
- Karstens, J., Crutchley, G.J., Berndt, C., Talling, P.J., Watt, S.F.L., Trofimovs, J., Hühnerbach, V., Le
- Friant, A., and Lebas, E. (2013). Insights into pyroclastic flow emplacement from high-resolution
- 3D seismic data offshore Montserrat, Lesser Antilles. Journal of Volcanological and Geothermal
- 701 *Research*, 257. 1-11.
- 702 Komorowski JC, Boudon G., Semet M., Beauducel F., Anténor-Habazac C., Bazin S., Hammouya G.
- 703 (2005). Guadeloupe. In: J. Lindsay, R. Robertson, J. Shepherd, S. Ali, (Eds.), Volcanic Hazard
- Atlas of the Lesser Antilles, published by University of the West Indies, Seismic Research Unit,
- 705 Trinidad and IAVCEI, pp 65-102.
- 706 Krastel, S., H.U. Schmincke, C.L. Jacobs, R. Rihm, T.M. Le Bas, and B. Alibés (2001). Submarine
- landslides around the Canary Islands, *J. Geophys. Res.*, 106 (B3), 3977-3997.
- Krause, D.C., White, W.C., PIPER, D.J.W. and Heezen, B.C., 1970. Turbidity currents and cable breaks
- in the western New Britain Trench. *Geological Society of America Bulletin*, 81(7), pp.2153-2160.
- Labazuy, P. (1996). Recurrent landslides events on the submarine flank of Piton de la Fournaise volcano
- 711 (Reunion Island), in Volcano Instability on the Earth and Other Planets, Geol. Soc. Spec. Publ., vol.
- 712 110, edited by W.J. McGuire et al., pp. 295-306.
- 713 Lafuerza S., Le Friant, A., Manga, M., Boudon, G., Villemant, B., Stroncik, N., Voight, B., Hornbach,
- M., Ishizuka O. and the Expedition 340 Scientific Party (2014). Geomechanical characterizations of
- submarine volcano flank sediments, Martinique, Lesse Antilles Arc. S. Krastel et al., (Eds.),
- Sumarine mass movements and consequences, Advances in Natural and Technological Hazards
- 717 Research, Springer Inten. Publishing, Switzerland, 2014, 37, doi 10.1007/978-3-319-00972-8 7.
- Lebas, E., Le Friant, A., Boudon, G., Watt, S., Talling, P., Feuillet, N., Deplus, C., Berndt, C., Vardy,
- M. (2011). Multiple widespread landslides during the long-term evolution of a volcanic island:
- 720 insights from high-resolution seismic data, Montserrat Lesser Antilles. *Geochemistry, Geophysics*,
- 721 *Geossystems*, 12, 1, doi: 10:1029/2010GC003451.

- Lebas, E., Le Friant, A., Deplus, C., de Voogd, B., (2018). Understanding the evolution of an oceanic
- 723 intraplate volcano from seismic reflection data: A new model for La Réunion, Indian Ocean. Journal
- of Geophysical Research: Solid Earth, 123, 1035-1059, doi.1002/2017JB014959.
- Le Friant, A., Boudon, G., Arnulf, A., Robertson, R. (2009). Debris avalanche deposits offshore St
- Vincent (West Indies): impact of flank-collapse events on the morphological evolution of the island,
- 727 *J. Volcanol. Geotherm. Res, 179, 1-10.*
- Le Friant, A., Boudon G., Deplus C., Villemant B. (2003a). Large scale flank collapse events during the
- activity of Montagne Pelée, Martinique, Lesser Antilles. Journal of Geophysical Research, 108,
- 730 B1, 2055.
- Le Friant, A., Boudon, G., Komorowski, J-C., Deplus, C. (2002). L'île de la Dominique, à l'origine des
- avalanches de débris les plus volumineuses de l'arc des Petites Antilles. Comptes Rendus,
- 733 *Géosciences*, 334 (4), 235-243.
- Le Friant, A., Deplus, C., Boudon, G., Feuillet, N., Trofimovs, J., Komorowski, J-C., Sparks, R. S. J.,
- Talling, P., Loughlin, S., Palmer, M., Ryan., G. (2010). Eruption of Soufrière Hills (1995-2009)
- from an offshore perspective: insights from repeated swath bathymetry surveys. *Geophys. Res. Lett.*,
- 737 *37, L11307, doi:10.1029/2010GL043580.*
- Le Friant, A., Harford, C., Deplus, C., Boudon, G., Sparks, S., Herd, R., Komorowski, J.-C. (2004).
- Geomorphological evolution of Montserrat (West Indies): importance of flank collapse and
- erosional processes. *Geol. Soc. London*, 161, 147-160.
- Le Friant, A., Heinrich, P., Deplus, C., Boudon, G. (2003b). Numerical simulation of the last flank
- collapse event of Montagne Pelée, Martinique, Lesser Antilles. *Geophysical Research Letters*, 30,
- 743 2, 1034.
- Le Friant, A., Ishizuka, O., Boudon, G., Palmer, M.R., Talling, P., Villemant, B., Adachi, T., Aljahdali,
- M., Breitkreuz, C., Brunet, M., Caron, B., Coussens, M., Deplus, C., Endo, D., Feuillet, N., Fraas,
- A.J., Fujinawa, A., Hart, M. B., Hatfield, R.G., Hornbach, M., Jutzeler, M., Kataoka, K. S.,
- Komorowski, J-C., Lebas, E., Lafuerza, S., Maeno, F., Manga, M., Martínez-Colón, M., McCanta,
- M., Morgan, S., Saito, T., Slagle, A., Sparks, S., Stinton, A., Stroncik, N., Subramanyam, K. S.V.,
- 749 Tamura, Y., Trofimovs, J., Voight, B., Wall-Palmer, D., Wang, F., Watt, S.F.L. (2015). Submarine
- record of volcanic island construction and collapse in the Lesser Antilles arc: First scientific drilling
- of submarine volcanic island landslides by IODP Expedition 340. *Geochem.*, *Geophys.*, *Geosyst*, 16,
- 752 2, 420-442.
- 753 Le Friant, A., Ishizuka, O., Stroncik, N.A., and the Expedition 340 Scientists, Proc. IODP, 340, (2013).
- Tokyo (Integrated Ocean Drilling Program Management International, Inc.). doi:10.2204/
- 755 iodp.proc.340.2013.

- Le Friant, A., Lebas, E., Clément, V., Boudon, G., Deplus, C., de Voogd, B., Bachèlery, P. (2011). A
- 757 new model for the evolution of la Réunion volcanic complex from complete geophysical surveys,
- 758 Geophys. Res. Lett. 38, doi: 10.1029/2011GL047489
- 759 Lénat, J.-F., P. Vincent and P. Bachèlery (1989). The offshore continuation of an active basaltic volcano:
- 760 Piton de la Fournaise (Réunion Island, Indian Ocean); structural and geomorphological
- interpretation from sea beam mapping, J. Volcanol. Geotherm. Res., 36, 1-36.
- Lipman, P.W., W.R. Normark, J.G. Moore, J.B. Wilson, and E. Gutmacher (1988). The giant submarine
- Alika debris slide, Mauna Loa, Hawaii, J. Geophys. Res., 93, 4279-4299.
- Lopart, J., Lafuerza, S. Le Friant, A., Urgeles, R. (2018). Hydrogeological evolution as a pre-
- conditioning factor for slope failure in a back-arc volcanic setting, W off Martinique Island. 8th
- ISSMMTC (International Symposium on Submarine Mass Movements and their consequences), 7-
- 9 May 2018, Victoria, British Columbia, Canada.
- Lovholt, R., Pedersen, G., Gisler, G. (2008). Oceanic propagation of a potential tsunami from the La
- Palma island island. *Journal of Geophysical Research* 113, C09026.
- 770 MacDonald, R., Hawkesworth, C.J. & Heath, E. (2000). The Lesser Antilles volcano chain; a study in
- arc magmatism. Earth Science Reviews, 49, 1-76.
- 772 Manga M., M. Hornbach, A. Le Friant, O. Ishizuka, N. Stroncik, T. Adachi, M. Aljahdali, G. Boudon,
- 773 C. Breithreuz, A. Fraass, A. Fujinawa, R. Hatfield, M. Jutzeler, K. Kataoka, S. Lafuerza, F. Maeno,
- M. Martinez-Colon, M. McCanta, S. Morgan, MR. Palmer, T. Saito, A. Slagle, A. Stinton, K.S.V.
- Subramanyam, Y. Tamura, P. Talling, B. Villemant, D. Wall-Palmer, F. Wang, (2012). Heat Flow
- in the Lesser Antilles island arc and adjacent back arc Grenada basin. *Geochemistry*, *Geophysics*
- 777 and Geosystems (G3), doi: 10.1029/2012GC004260
- 778 Mangeney-Castelnau A, Vilotte JP, Bristeau MO, Perthame B, Bouchut F, Simeoni C, Yernini S (2003).
- Numerical modeling of avalanches based on Saint-Venant equations using a kinetic scheme. J
- 780 *Geophys Res* 108(B11):2527
- Masson, D.G., Harbitz, C.B., Wynn, R.B., Pedersen, G. and Løvholt, F. (2006), Submarine landslides:
- 782 processes, triggers and hazard prediction. *Philosophical Transactions of the Royal Society A*
- 783 364:2009-2039.
- Masson, D.G., Watts, A.B., Gee, M.J.R., Urgeles, R., Mitchell, N.C., Le Bas, T.P. and M. Canals
- 785 (2002). Slope failures on the flanks of the western Canary Islands. Earth-Science Reviews 57:1-
- 786 35.
- 787 Mattioli, G.S., Voight, B., Linde, A T; Sacks, I S; Watts, P; Widiwijayanti, C; Young, S R; Hidayat, D;
- Elsworth, D; Malin, P E; Shalev, E; Van Boskirk, E; Johnston, W; Sparks, R S J; Neuberg, J; Bass,
- V; Dunkley, P; Herd, R; Syers, T; Williams, P; Williams, D. (2007). Unique and remarkable
- dilatometer measurements of pyroclastic flow-generated tsunamis. *Geology* 35, 25-28.

- 791 McGuire, W.J. (1996). Volcano instability: a review of contemporary themes, in Volcano Instability on
- the Earth and Other Planets, Geol. Soc. Spec. Publ., vol. 110, edited by W.J. McGuire et al., pp. 1-
- 793 23.
- Montanaro, C., and Beget (2011). Volcano collapse along the Aleutian Ridge (western Aleutian Arc).
- Natural Hazards and Earth System Sciences, 11:715-730, http://dx.doi.org/10.5194/nhess-11-715-
- 796 2011.
- Moore, J.G., Clague, D.A., Holcomb, R.T., Lipman, P.W., Normark, W.R., and Torresan, M.E. (1989).
- Prodigious submarine landslides on the Hawaiian ridge, *J. Geophys. Res.*, 94(B12), 17465-17484.
- Moore, J.G., and Clague, D.A (2002). Mapping the Nuuanu and Wailau landslides in Hawaii. Pp. 223-
- 244 in Hawaiian Volcanoes: Deep underwater Perspectives. E. Takahashi, P.W. Lipman, M.O.
- Garcia, J. Naka, and S. Aramaki, eds, American Geophysical Union, Washington, DC,
- http://dx.doi.org/10.1029/GM128p0223.
- 803 Moore, J.G., Normark, W.R. and Holcomb, R.T., 1994. Giant hawaiian landslides. Annual Review of
- Earth and Planetary Sciences, 22(1), pp.119-144.
- Nomanbhoy, N. and Satake, K., 1995. Generation mechanism of tsunamis from the 1883 Krakatau
- eruption. *Geophysical Research Letters*, 22(4), pp.509-512.
- 807 Oehler, J.-F., Labazuy, P.and Lénat, J.-F. (2004). Recurrence of major flank landslides during the last
- 2-Ma-history of Reunion Island, Bull. Volcanol., 66, 585-598.
- Oehler, J.F., Lenat, J.F., Labazuy P. (2008). Growth and collapse of the Reunion Island volcanoes.
- 810 *Bulletin of Volcanology*, DOI 10.1007/s00445-007-0163-0.
- Ogata, K., Mountjoy, J.J., Pini, G.A., Festa, A., Tinterri, R. (2014). Shear-zone liquefaction in mass
- 812 transport deposit emplacement: A multi-scale integration of seismic reflection and outcrop data.
- 813 *Marine Geology*, 356, 50-64.
- Palmer, M., Stuart J. Hatter, Thomas M. Gernon, Rex N. Taylor, Michael Cassidy, Peter Johnson, Anne
- Le Friant, and Osamu Ishizuka, (2016). Discovery of a large 2.4 Ma Plinian eruption of Basse-Terre,
- Guadeloupe, from the marine sediment record. *Geology*, 44-2, 123-126, doi:10.1130/G37193.1.
- Picard, M. Schneider, J.-L., & Boudon, G. (2006). Contrasting sedimentary processes along a convergent
- margin: the Lesser Antilles arc system. *Geo-Marine Letters*, 26(6), 397-410.
- 819 Schmincke, H-U, Sumita, M. (1998). Volcanic evolution of Gran Canaria reconstructed from apron
- 820 sediments: synthesis of vicap project drilling. Proceedings of the Ocean Drilling Program, Scientific
- 821 *Results*, 157: 443- 469.

- 822 Schneider, J-L., Gérard, M., Schmincke, H-U., Weaver, P.P.E., Firth, J., Baraza, J., Bristow, J.F.,
- Brunner, C., Carey, S.N., Coakley, B., Fuller, M., Funk, T., Goldstrand, P., Herr, B., Hood, J., Howe,
- R., Jarvis, I., Lebreiro, S., Lindblom, S., Lykke-Andersen, H., Maniscalco, R., Rothwell, G.,
- Sblendorio-Levy, J., Sumita, M., Taniguchi, H., TU, P., Wallace, P. (1997). Du volcan au sediment:
- la dynamique du talus volcanoclastique sous-marin de Gran Canaria, Canaries (Atlantique oriental,
- Leg ODP 157). Comptes rendus de l'Académie des Sciences, Paris, 324, IIa, 891-898.
- 828 Siebert, L. (2002). Landslides resulting from structural failures in volcanoes. In: Evans, S.G., and
- DeGraff, J.V., Catastrophic landslides: Effects, occurrence, and mechanisms. Boulder, Colorado,
- Geol. Soc. America, *Reviews in Engineering Geology*, vol XV, p 209-235,
- 831 Sigurdsson, H., Sparks, R.S.J., Carey, S. & Huang, T.C. (1980). Volcanogenic sedimentation in
- theLesser Antilles Arc. *Journal of Geology*, 88, 523-540.
- 833 Silver, E., Day, S., Ward, S., Hoffmann, G., Llanes, P., Driscoll, N., Appelgate, B., Saunders, S. (2009).
- Volcano collapse and tsunami generation in the Bismark Volcanic Arc, Papua New Guinea. *Journal*
- 835 of Volcanology and Geothermal Research, 186:210-222,
- http://dx.doi.org/10.1016/j.jvolgeores.2009.06.013.
- 837 Slingerland, R., and Voight, B. (1979). Occurrence and predictive models for slide-generated water
- waves. In: *Rockslides and Avalanches*, Elsevier, Amsterdam, v. 2, pp. 317-400.
- 839 Trofimovs, J., Amy, L., G. Boudon, C. Deplus, E. Doyle, N. Fournier, M.M.B. Hart, J-C Komorowski,
- A. Le Friant, E. Lock, C. Pudsey, G. Ryan, R.S.J Sparks, P.J. Talling (2006). What happens when
- pyroclastic flows enter the ocean? *Geology*, 34, 549-552.
- Trofimovs, J., J.K. Fisher, H.A. Macdonald, P.J Talling, R.S.J. Sparks, M.B Hart, C. Smart, G. Boudon,
- 843 C. Deplus, J-C. Komorowski, A. Le Friant, S. Moreton, M. Leng, (2010), Evidence for carbonate
- platform failure during rapid sea-level rise; ca 14 000 year old bioclastic flow deposits in the Lesser
- 845 Antilles. *Sedimentology*, 57 (3), 735-759.
- Trofimovs, J., Talling, P.J., Fisher, J.K., Sparks, R.S.J., Watt, S.F.L., Hart, M.B., Smart, C., Le Friant,
- A., Cassidy, M., Moreton, S.G., Leng, M.J. (2013). Timing, origin and emplacement dynamics of
- mass flows offshore of SE Montserrat in the last 110 ka: implications for landslide and tsunami
- hazards, eruption history, and volcanic island evolution. *Geochemistry, Geophysics, Geosystems*, V.
- 850 14, Issue 2, 385-406, doi:10.1002/ggge.20052.
- 851 Urgeles, R., M. Canals, J. Baraza, B. Alonso, and D. Masson (1997). The most recent megalandslides
- of the Canary Islands: El Golfo debris avalanche and Canary debris flow, west El Hierro Island,
- 853 *Journal of Geophysical Research*, 102 (B9), 20305-20323.

- Urgeles, R., Masson, D.G., Canals, M., Watts, A.B., Le Bas, T. (1999). Recurrent large-scale landsliding
- on the west flank of La Palma, Canary Islands. *Journal of Geophysical Research*, 104 (B11), 25331-
- 856 25348.
- Viesca, R.C. and J.R. Rice (2012). Nucleation of slip-weakening rupture instability in landslides by
- localized increase of pore pressure. *Journal of Geophysical Research* 117, B03104.
- Vincent, P.M., Bourdier, J.-L., Boudon, G. (1989). The primitive volcano of Mount Pelée: its
- construction and partial destruction by flank collapse. J. Volcanol. Geotherm. Res., 38, 1-15.
- Voight, B. (2000). Structural stability of andesite volcanoes and lava domes. *Phil. Trans. Roy. Soc. Lond.*
- 862 A 358 : 1663-1703.
- Voight, B., Janda, R.J., Glicken, H. & Douglass, P.M. (1981). Catastrophic rockslide avalanche of May
- 864 18. *Prof. Paper US Geol. Survey* 1250 : 347-378.
- Voight, B., Komorowski J-C., Norton, G.E., Belousov, A.B., Belousova, M., Boudon, G., Francis, P.W.,
- Franz, W., Heinrich, P., Sparks, R.S.J. & Young, S.R. (2002). The 26 December (Boxing Day)
- 1997 sector collapse and debris avalanche at Soufrière Hills Volcano, Montserrat. *In Druitt*, T.H.,
- and Kokelaar, B.P.(eds) The eruption of Soufriere Hills Volcano, Montserrat, from 1995 to 1999,
- 869 Geol. Soc., London, *Memoirs*, 21, 363-407.
- Voight and Elsworth (1997). Failure of volcano slopes. *Geotechnique* 47, 1-31.
- Voight B., Le Friant, A., Boudon, G., Deplus, C., Komorowski, J.C., Lebas, E., Sparks, R.S.J., Talling,
- P., Trofimovs, J. (2012). Undrained Sediment Loading Key to Long-Runout Submarine Mass
- Movements: Evidence from the Caribbean Volcanic Arc. In: Y. Yamada, K. Kawamura, K. Ikehara,
- Y. Ogawa, R. Urgeles, D. Mosher, J. Chaytor, M. Strasser (Eds), Submarine Mass Movements and
- Their Consequences. Advances in Natural and Technological Hazards Research, 31, 417-428, doi:
- 876 10.1007/978-94-007-2162-3-37.
- Wadge, G. (1984). Comparison of volcanic production rates and subduction rates in the Lesser Antilles
- and Central America. *Geology*, 12, 555-558.
- Wall-Palmer, D., Coussens, M., Talling, P.J., Jutzeler, M., Cassidy, M., Marchant, I., Palmer, M., Watt,
- S., F.L., Smart, C., Fisher, J.K., Hart, M.B., Fraass, A., Trofimovs, J., Le Friant, A., Ishizuka, O.,
- Adachi, T., Aljahdali, M., Boudon, G., Breitkreuz, C., Endo, D., Fujinawa, A., Hatfield, R.,
- Hornbach, M., Kataoka, Kyoko, Lafuerza, S., Maeno, F., Manga, M., Martinez-Colon, M.,
- McCanta, M., Morgan, S., Saito, T., Slage, A., Stinton, A., Subramanyam, K.S.V., Tamura,
- Yoshihiko, Villemant, B., Wang, Fei. (2014). Late Pleistocene stratigraphy of IODP U1396 and
- compiled chronology offshore of south and south west Montserrat, Lesser Antilles. *Geochemistry*,
- 886 *Geophysics, Geosystems*, 15, doi:10.1002/2014GC005402.
- 887 Ward, S. N. (2001). Landslide tsunami. *J. Geophys. Res*, vol 106, n°6 p11, 201-11,215, june 10.

- 888 Ward, S.N. & Day S.J. (2001). Cumbre Vieja volcano potential collapse and tsunami at La Palma,
- 889 Canary Islands. *Geophysical Research Letters*, 28, 3397-3400.
- 890 Ward, S.N. and Day, S., 2003. Ritter Island volcano—lateral collapse and the tsunami of 1888.
- *Geophysical Journal International*, 154(3), pp.891-902.
- Watt, S.F.L., Talling, P.J., and Hunt, J.E. (2014). New insights into the emplacement dynamics of
- volcanic-island landslides. *Oceanography*, v. 27, p. 46-57.
- Watt, S.F.L., Talling, P.J., Vardy, M.E., Heller, V., Hühnerbach, V., Urlaub, M., Sarkar, S., Masson,
- D.G., Henstock, T.J., Minshull, T.A., Paulatto, M., Le Friant, A., Lebas, E., Berndt, C., Crutchley,
- 896 G.J., Karstens, J., Stinton, A.J., Maeno, F. (2012a). Combination of volcanic-flank and seafloor
- sediment failure offshore Montserrat, and their implications for tsunami generation, *EPSL* v. 319,
- 898 2, p. 228-240, doi:116/j.epsl.2011.11.03.
- Watt, S.F.L., Talling, P.J., Vardy, M.E., Masson, D.G., Henstock, T.J., Huehnerbach, T.A., Minshull,
- 900 M., Urlaub, M., Lebas, E., Le Friant, A., Berndt, C., Crutchley, G.J., Karstens, J. (2012b).
- 901 Widespread and progressive seafloor-sediment failure following volcanic debris avalanche
- 902 emplacement: Landslide dynamics and timing offshore Montserrat, Lesser Antilles. Marine
- 903 *Geology*, 323, 69-94. doi: 10.1016/j.margeo.2012.08.002.
- Watts, P., Grilli,, S.T., Tappin, D.R., Fryer, G.J. (2005). Tsunami generation by submarine mass failure.
- 905 II: Predictive equations and case studies. J. Waterw. Port Coastal Ocean Eng. 131, 298-310.
- Watts, A.B., and Masson, D.G. (1995). A giant landslide on the north flank of Tenerife, Canary Islands,
- 907 J. Geophys. Res., 100 (B12), 24487-24498.

- 908 Westercamp, D. and Traineau, H. (1983). Carte géologique au 1/20 000 de la Montagne Pelée, avec
- notice explicative. B.R.G.M. eds, Orléans. Earth Planet. Sci. Lett.
- 910 Wynn R.B., and Masson D.G. (2003). Canary Islands landslides and tsunami generation: can we use
- 911 turbidite deposits to interpret landslide processes? In: Locat J, Mienert J (eds), Submarine mass
- 912 movements and their consequences, Kluwer Academic Publ., Boston, London, pp 325–332


914 **Figures** 915 916 Fig 1: The Lesser Antilles Arc from Montserrat to St Vincent. Extent of the main landslide 917 deposits (chaotic units) on swath bathymetry illuminated from North (AGUADOMAR cruise, 918 1999 and CARAVAL cruise, 2002). Active volcanoes and drilling sites are annotated. Solid 919 lines represent the seismic profiles shown in Figure 2. Modified from Le Friant et al. (2015). 920 921 922 Fig 2: Seismic reflection profiles revealing the presence of chaotic deposits offshore (a) Martinique and (b) Montserrat. Location of the profiles is shown on Figure 1. Deposits are 923 characterized by chaotic reflectors, which contrast with the well-bedded, subhorizontal 924 sedimentary units. Locations of drilling sites are annotated. 925 926 Fig 3: Correlation between seismic and drilling data for (a) Hole U1399-A and (b) Hole U1400-927 928 C. Deformation on both data sets are well correlated despite their large-resolution differences. Adapted from Brunet et al. (2016). 929 930 Fig 4: Schematic figure of processes which can explain the presence of huge landslide deposits 931 offshore volcanic islands (topographic/bathymetric profile from Martinique). Volcanic flank 932 collapses generate debris avalanches that enter the sea, incise the sediments and generate failure, 933 propagation, and deformation of marine sediments. From Le Friant et al. (2015). 934 935 Fig 5: Conceptual model of the relation between aerial instability processes (i.e., volcanic flank 936 collapses) and submarine instability processes (i.e., seafloor sediment failures) offshore 937 Martinique. The first debris avalanche deposit (DAD1) weakened and initiated seafloor 938 sediment failure, that resulted in a submarine landslide propagation. Successive DADs (DAD2 939 and DAD3) remobilized unconsolidated surficial sediments within the proximal part of the 940 941 submarine landslide deposit (SLD). From Brunet et al. (2016). 942 Fig 6: Depth distribution of shear strength-overburden stress ratio S_u/σ'_v where S_u corresponds 943 to undrained shear strength and σ'_{v} to effective vertical stress. Gray shading corresponds to 944 $S_u/\sigma'_v < 0.17$ -0.25 associated with normally consolidated materials. Black and green colors are 945


used to differentiate S_u measurements at holes B (black) and C (green) for each drilling site,


except for site U1397, where hole A is in black and hole C in green. Circles: S_u -handheld penetrometer; crosses: S_u -automated shear vane. From Lafuerza et al. (2014).


Fig. 7: Tsunami wave height versus slide velocity and water depth for three different slide volumes discussed in the text (solid lines: 25 km³, dashed lines: 13 km³, dotted lines: 5 km³). We extend the model to a maximum water depth of 3000 m below sea level (mbsl), as this corresponds to the maximum water depth where we observe slide deposits in the Lesser Antilles Arc. Model results show a peak wave height two minutes after slide initiation, for three different average slide velocities (red: 50 m s⁻¹, black: 10 m s⁻¹, blue: 1 m s⁻¹). All the calculations assume a constant parabolic slide shape (e.g., Jiang & LeBlond, 1992).

Fig. 8: (a) Combined bathymetry (FOREVER and ERODER1 cruises) and topography data of La Réunion volcanic complex illuminated from NW. Dashed thick circle indicates the base of the volcanic complex, with interpretations of mass-wasting processes. The extent of rough and chaotic terrains is mapped in green (Piton des Neiges) and in red (Piton de la Fournaise). Arrows indicates the direction of the instabilities. Secondary front instabilities are mapped in light colors. (b) Seismic profile FOR 63 illustrating the seismic signature of the material composing the large submarine landslide deposit (SLD) identified offshore Piton de la Fournaise volcano. Note the long runout of this large SLD deposit, which extends beyond the base of the volcanic complex. Modified from Le Friant et al. (2011) and Lebas et al. (2018).










2: THE 1st DEBRIS AVALANCHE DEPOSIT (DAD1) ASSOCIATED TO THE 1st FLANK-COLLAPSE EVENT, INITIATED SEAFLOOR SEDIMENT FAILURE AND PROPAGATION OF A SUBMARINE LANDSLIDE



3: THE 2nd AND 3RD DEBRIS AVALANCHE DEPOSITS (DAD2 AND DAD3) ASSOCIATED TO THE 2nd and 3RD FLANK-COLLAPSE EVENTS, REMOBILIZED PROXIMAL PART OF THE SUBMARINE LANDSLIDE DEPOSIT (SLD), ENHANCING DEFORMATION UNTIL THE MORPHOLOGICAL FRONT - CURRENT DEPOSIT OUTLINE

