

Altered fungal communities in contaminated soils from French industrial brownfields

Florian Lemmel, Florence Maunoury-Danger, Corinne Leyval, Aurélie Cebron

▶ To cite this version:

Florian Lemmel, Florence Maunoury-Danger, Corinne Leyval, Aurélie Cebron. Altered fungal communities in contaminated soils from French industrial brownfields. Journal of Hazardous Materials, 2020, pp.124296. 10.1016/j.jhazmat.2020.124296 . hal-03016406

HAL Id: hal-03016406 https://hal.science/hal-03016406

Submitted on 20 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Highlights:

- Fungal diversity of polycyclic aromatic hydrocarbon (PAH) and metal contaminated soils
- Differences at phylum and OTU level among soils of slag heap, settling pond and control groups
- Saccharomycetes, Leotiomycetes and Agaricomycetes favoured in uncontaminated soils
- Higher relative abundance of Cryptomycota in metal-contaminated soils
- Higher relative abundance of Dothideomycetes in PAH-contaminated soils

| | | Controls | Slag heaps | Settling ponds | |
|-----------------------|-------------------------|------------|---|---|--|
| ination els | РАН | Ø | 0 | | |
| Contam lev | Metal (Zn, Pb) | Ø | | | |
| iversity or levels | Abundance & Richness | | \bigcirc | | |
| Alpha-d Indicato | Diversity & Evenness | \bigcirc | © | \bigcirc | |
| ommunity | Repressed fungal ta | аха | Agarico Saccharo Leotion Chytridi | mycetes omycetes nycetes omycota | |
| Fungal co | Favored fungal taxa | | Cryptomycota Pucciniomycetes Dothideomyce Eurotiomycete | | |

Abstract

Polycyclic aromatic hydrocarbons (PAHs) and metals are contaminants of industrial brownfield soils. Pollutants can have harmful effects on fungi, which are major actors of soil functioning. Our objective was to highlight fungal selection following long-term contamination of soils. Fungal diversity was assessed on 30 top-soil samples from ten sites gathered in three groups with different contamination levels and physico-chemical characteristics: 1) uncontaminated controls, 2) slag heaps displaying high PAH and moderate metal contaminations, and 3) settling ponds displaying high metal and intermediate PAH contaminations.

Although fungal abundance and richness were similar among the soil groups, the diversity and evenness indices were lower for the slag heap group. Fungal diversity differed among soil groups at the phylum and OTU levels, and indicator species were identified. The relative abundance of Agaricomycetes, Saccharomycetes, Leotiomycetes and Chytridiomycota was higher in the control soils than in the two groups of contaminated soils. Cryptomycota LKM11 representatives were favoured in the slag heap and settling pond groups, and their relative abundance was correlated to the zinc and lead contamination levels. Dothideomycetes – positively linked to PAH contamination – and Eurotiomycetes were specific to the slag heap group. Pucciniomycetes and especially *Gymnosporangium* members were favoured in the settling pond soils.

1 Altered fungal communities in contaminated soils from French industrial

2 brownfields

3

4 Florian Lemmel¹, Florence Maunoury-Danger², Corinne Leyval¹, Aurélie Cébron^{1,*}

- 5 ¹ Université de Lorraine, CNRS, LIEC, F-54000, Nancy, FRANCE
- 6 ² Université de Lorraine, CNRS, LIEC, F-57000 Metz, France
- 8 * corresponding author: LIEC UMR7360, Faculté des Sciences et Technologies, Bd des Aiguillettes, BP70239, 54506
- 9 Vandoeuvre-les-Nancy; Mail: <u>aurelie.cebron@univ-lorrane.fr</u>; Phone:+33 3 72 74 52 15.
- 10

7

11 Abstract

Polycyclic aromatic hydrocarbons (PAHs) and metals are contaminants of industrial brownfield soils. Pollutants can have harmful effects on fungi, which are major actors of soil functioning. Our objective was to highlight fungal selection following long-term contamination of soils. Fungal diversity was assessed on 30 top-soil samples from ten sites gathered in three groups with different contamination levels and physico-chemical characteristics: 1) uncontaminated controls, 2) slag heaps displaying high PAH and moderate metal contaminations, and 3) settling ponds displaying high metal and intermediate PAH contaminations.

- 18 Although fungal abundance and richness were similar among the soil groups, the diversity and evenness indices
- 19 were lower for the slag heap group. Fungal diversity differed among soil groups at the phylum and OTU levels,
- 20 and indicator species were identified. The relative abundance of Agaricomycetes, Saccharomycetes,
- 21 Leotiomycetes and Chytridiomycota was higher in the control soils than in the two groups of contaminated soils.
- 22 Cryptomycota LKM11 representatives were favoured in the slag heap and settling pond groups, and their relative
- 23 abundance was correlated to the zinc and lead contamination levels. Dothideomycetes positively linked to
- 24 PAH contamination and Eurotiomycetes were specific to the slag heap group. Pucciniomycetes and especially

25 *Gymnosporangium* members were favoured in the settling pond soils.

- 26
- 27
- 28 Keywords: Fungal communities; 18S rDNA sequencing; Soil pollution; Polycyclic aromatic hydrocarbon;
- 29 Metals.
- 30

31 **1. Introduction**

- Fungi are known to play a fundamental role in terrestrial ecosystems: they contribute to carbon cycling through organic matter mineralisation, or mediate plant mineral nutrition(Tedersoo *et al.*, 2014). Soil fungal communities
- 34 and the relative abundance of the different fungal taxa are mostly shaped by climatic and edaphic factors
- 35 (Beauregard *et al.*, 2010; Hannula *et al.*, 2012; Tedersoo *et al.*, 2014). In addition to these natural environmental
- 36 and biogeographic filters, anthropogenic constraints linked to human activities also contribute to shape fungal 37 communities. Due to their toxicity, soil contaminants act as selection factors that shape microbial communities.
- 38 An increase in soil metal (Pb and Zn) concentrations has been shown to decrease arbuscular mycorrhizal fungal
- 39 richness of plant species in pit mine, whereas one taxon was positively selected and developed in the highly
- 40 contaminated area (Zarei et al., 2010). As a consequence, the presence of contaminants and their toxicity over
- 41 the long term could: 1) decrease diversity owing to the extinction of sensitive taxa, 2) disrupt the recycling of
- 42 organic matter, biogeochemical cycles and soil functioning, and 3) select suitable taxa tolerant to pollutants,
- 43 possibly involved in contaminant biodegradation or bioaccumulation (Harms *et al.*, 2011). There is still a need
- 44 nowadays to increase our knowledge on the ecology and adaptation capacities of fungal communities and taxa to
- soil contaminants, and to evidence the selection of certain taxa displaying potentially interesting properties in
- 46 bioremediation.
- 47 Various anthropogenic activities have been negatively affecting the air, water and soil matrices by releasing 48 solid, liquid and gaseous wastes containing diverse pollutants. To date, large brownfield lands, where coking and 49 steel industry took place during the 20th century, are still highly contaminated with metals and organic compounds such as polycyclic aromatic hydrocarbons (PAHs) (Rachwał et al., 2015) that pose serious threats to 50 51 public and environmental health. Only few studies have assessed fungal taxonomic diversity in contaminated 52 soils (Bourceret et al., 2016), especially in soils contaminated for several decades and with various 53 contamination levels and contamination types. PAHs and metals are known for their toxicity, carcinogenicity, or 54 mutagenicity (US EPA, 2014) that can also target soil fungal communities. For example, PAH contamination 55 can reduce fungal diversity (Bell et al., 2014), or species richness (Zhou et al., 2017). Some fungal genera such 56 as Fusarium, Cladosporium, Mucor (Salvo et al., 2005), and the Sordariales and Pleosporales orders (Zhou et 57 al., 2017) have been found more represented in PAH-contaminated soil or sediment than in uncontaminated 58 environments. Ascomycota appear to be more prevalent in metal-contaminated soil, while Basidiomycota 59 dominate in uncontaminated areas (Narendrula and Nkongolo, 2017). More specifically, zinc and cadmium 60 contamination in a forest soil modified fungal community composition, with a dominance of Sistotrema sp., 61 Suillus luteus and Wilcoxina mikolae, but did not affect fungal diversity indices (Op De Beeck et al., 2015). This 62 dominance of various species or genera in contaminated soils suggests fungal selection and adaptation to 63 contamination. For bacterial taxonomic diversity in soils with various PAH and metal contamination levels, the 64 occurrence of certain OTUs was directly correlated to the presence of contaminants (Lemmel et al., 2019). We 65 therefore wondered whether the fungal community response was similar or not in these soils.
- 66 In this context of soils contaminated by PAHs and heavy metals (Zn, Cd, Pb...) for decades following industrial
- activities, we hypothesized that the type and level of contamination led to the selection of microorganisms, and
- 68 to similar fungal community compositions of soils sharing similar contamination types (i.e. with a similar
- 69 industrial history). Fungal species specific to these extremely contaminated soils could be identified as indicators

- and considered for bioremediation. Additionally, a better understanding of fungal community assemblages living
- 71 in historically multi-contaminated soils could improve our knowledge of fungal ecology and soil functioning.
- 72 The aim of this study was to determine the taxonomic diversity and composition of fungal communities from
- 73 long-term contaminated soils presenting both PAH and metal contaminations. Slag heap and settling pond soils
- that evolved from former French steel industries were compared with control uncontaminated soils from the
- same region. The slag heap sites displayed high PAH and medium metal contamination, while the settling pond
- rRNA sites displayed high metal and low PAH contamination. Taxonomic diversity was assessed through 18S rRNA
- 77 gene tag-amplicon sequencing (Illumina MiSeq). The links between soil physico-chemical parameters such as
- 78 contaminants and fungal alpha diversity indices or relative abundance of fungal phyla or OTUs were
- 79 investigated, and indicator species were identified.

81 2. Materials and methods

82 The ten sites and their physico-chemical soil characteristics were previously described in Lemmel et al. (2019), 83 but are briefly summarized in Table 1. These ten sites are located in the same geographical area of northeastern 84 part of France (Grand-Est region; Figure S1) and are located less than 100 km from one another. Briefly, three 85 were considered as the control soils (ctrl), i.e. three forest soils collected at Hémilly (He), Montiers-sur-Saulx 86 (Mo), and Dieulouard (Di) with low anthropogenic influence and without PAH or metal contamination from 87 steel activity. Seven brownfield soils from former steel industries known to be contaminated by metals and/or 88 PAHs were collected from: i) former slag heaps (sh) at Homécourt (Ho), Terville (Te), Uckange (Uc), and 89 Neuves-Maisons (NM), and ii) former settling ponds (sp), at Pompey (Po), Mont-St-Martin (MsM), and 90 Russange-Micheville (RM). All sites presented a similar dense tree cover (dominated by birch and/or beech), 91 except for Uc (sparse birch trees) and NM (colonized neither by trees nor by other plant species, i.e. bare soil). 92 Three independent bulk soil samples (triplicates collected 1 m apart) were collected in November 2015 (early 93 autumn season with a daily mean temperature of 11°C) on each site, as follows: after removing the litter layer, if 94 present, three blocks of 20 cm side and 30 cm depth were sampled, brought back to the laboratory, air-dried, 95 sieved at 2 mm and homogenised before analysis. Aliquots of soil for DNA extraction were immediately frozen 96 and stored at -80°C. The 30 top-soil samples from the ten sites were gathered in three groups (control, slag heap, 97 settling pond). 98 The methods used for soil characterisation are described in Lemmel et al. (2019). The soil collection presented a 99 gradient of PAH contamination ranging from 0.03 to 1,095.90 mg kg⁻¹ dw soil (based on the 16 priority PAHs of 100 the US Environmental Protection Agency, US-EPA) and a gradient of metal contamination mainly represented 101 by Zn (60 to 119,000 mg kg⁻¹), Pb (23 to 39,500 mg kg⁻¹) and Cd (0.09 to 152.00 mg kg⁻¹) (**Table 1**). The slag 102 heap and settling pond soils displayed significantly higher PAH concentrations than the control soils. The 103 settling pond soils displayed significantly higher concentrations of metals (Zn, Cd, and Pb; fluorhydric acid 104 extraction) than the control and slag heap soils. The pH (measured from a soil suspension prepared in distilled 105 water, 1:5 w/v) was rather similar among the soils from all 10 sites (7.2 to 8.0) except for soil He (5.4). Other 106 soil characteristics, namely cation exchange capacity (CEC), carbonates, total organic carbon (24 to 159 g kg⁻¹), 107 total nitrogen (0.56 to 7.19 g kg⁻¹; Dumas method, ISO 10694), and texture (from silty to sandy; ISO 11277 108 method) varied among soils, but no significant statistical difference (Kruskal Wallis test, p>0.05) was found 109 among the three groups of sites (Table 1). To sum up, the slag heap soils were mainly characterised by high 110 PAH contamination and an intermediate level of metal contamination, while the settling pond soils were mainly 111 characterised by high metal contamination and intermediate PAH contamination. The control soils showed the

- lowest PAH and metal contamination levels, but significantly higher concentrations of major elements (Fe, K,
- 113 Al) than all contaminated soils.
- 114

115 2.2. DNA extraction, real-time quantitative PCR, sequencing

Genomic DNA was extracted from *ca*. 0.5 g of the soil samples (30 top-soil samples from the ten sites) using a Fast DNA spin kit for Soil (MP Biomedicals), then diluted to 5 ng μ l⁻¹ to be used as templates for further PCR

- 118 analysis.
- 119 Fungal abundance was estimated by real-time PCR quantification (Thion et al., 2012), using the primer sets
- 120 Fung5F/FF390R (Lueders et al., 2004), as explained in Lemmel et al. (2019).

The V7-V8 region of fungal 18S rRNA genes was amplified (ca. 390 bp) using primers FR1 and FF390 (Vainio 121 122 and Hantula, 2000) and following a previously described dual-index strategy (Kozich et al., 2013) using PCR 123 primers with Illumina adaptor, pad and index sequences (Thomas and Cébron, 2016). PCR reactions were 124 performed on 2 µl of diluted gDNA using Phusion high-fidelity polymerase (Thermo Scientific). PCR reactions 125 consisted of 34 cycles with a touchdown annealing temperature for 20 cycles (63 °C to 53 °C with a decrease of 0.5 °C/cycle) and 14 cycles at 53 °C. Amplification products were checked by 1 % agarose gel electrophoresis, 126 127 and purified using the UltraClean-htp 96 Well PCR Clean-Up kit (Qiagen) following the manufacturer's 128 instructions. After Quant-iT Picogreen ds-DNA assay Kit (Invitrogen) quantification, an amplicon library was 129 prepared (10 nM equimolar pool), purified on a QIAquick PCR purification kit column (Qiagen), and sent for 130 sequencing to Genewiz platform (South Plainfield, NJ, USA) using an Illumina MiSeq V2 Kit for 2 x 250 bp 131 paired-end sequencing. Illumina MiSeq paired-end reads were deposited in the SRA database under BioProject 132 accession number PRJNA601915. Sequence data were analysed following the MiSeq SOP procedure available 133 in March 2017 and described in Kozich et al. (2013), using Mothur v.1.36.0 (Schloss et al., 2009). Paired-end 134 reads were trimmed using the following criteria: QS > 20, 312 bp < length < 320 bp, and no more than 3 ambiguous bases and 8 homopolymers. Chimeras detected using Uchime (Edgar et al., 2011) and singletons 135 136 (sequences occurring only once among all samples) were removed. Unique sequences were aligned and taxa were assigned using the Silva V132 database (cutoff = 80) and following Wang's method (Wang et al. 2007). 137 138 Singletons and sequences not affiliated to fungi were removed. Sequences were clustered in Operational 139 Taxonomic Units (OTUs) at 97 % similarity. Finally, datasets were rarefied to the lowest number of sequences 140 per sample (6,181 reads/sample) to compare the soil diversity levels. The number of reads for each OTU was 141 expressed as a percentage, to express the relative proportion or relative abundance of each OTU within the total 142 community. Alpha diversity was expressed by calculating Chao1 richness, Pielou's evenness J', and Shannon 143 diversity H' indices (Hill et al., 2003) using Mothur.

144

145 2.3. Statistical analyses

- Statistical analyses were performed using RStudio v1.1.463. Significant differences (for fungal abundance, alpha diversity indices, and relative abundance of the different phyla and classes) among the three soil groups (ctrl, sh and sp) were assessed using multiple comparison Kruskal-Wallis test using '*kruskal*' function of the agricolae R package (Mendiburu, 2013), with p < 0.05 and Fisher's least significant difference *post hoc* test.
- 150 Variation in fungal community composition among sites was visualized by nonmetric multidimensional scaling 151 (NMDS) using the function 'metaMDS' on a Bray-Curtis dissimilarity matrix ('vegdist' function), and square-152 root transformed ('sqrt' function) to limit the influence of abundant OTUs. Differences in fungal community 153 OTU composition among *a priori* defined groups of sites were evaluated using permutation-based multivariate analysis of variance (PERMANOVA; Anderson 2001) in the 'adonis2' function of the vegan R package 154 155 (Oksanen et al. 2017), yielding tests of significance using a 999-permutation-based F-test. To validate 156 PERMANOVA, we examined homogeneity of variance using a permutational test of multivariate group 157 dispersions ('betadisper' function).
- 158 Relationships between soil characteristics and soil fungal community diversity (alpha diversity indices, relative
- abundance of the different phyla and of OTUs) were further analysed through three redundancy analyses (RDA;
- 160 'rda' function of the vegan R package; Oksanen et al. 2017) that classify the effects of soil characteristics on the

- 161 mean community composition per site. The carbon to nitrogen concentration ratio (C/N), organic matter (OM),
- 162 dissolved organic carbon (DOC), phosphorus (P₂O₅), carbonates, PAH concentrations (PAH), total metal
- 163 concentrations (Cu, Zn, Pb, Cd), and major element concentrations (Al, Ca, Fe, Na, K), pH, CEC and texture
- 164 (silt and sand percentages) were included to calculate the explained variance in RDA. The relative abundances of
- 165 fungal phyla and fungal OTUs were normalized (log(x+1)), and the most appropriate statistical model for RDA
- 166 was determined using forward selection as implemented in vegan's 'ordistep' and 'anova' functions. Only
- 167 significantly explanatory variables (1,000 Monte Carlo permutations, $\alpha < 0.05$) were included in the final
- 168 selected RDA models.
- 169 To investigate the linear relation between soil parameters and fungal diversity (alpha diversity indices and 170 relative abundance of phyla or classes), we calculated Pearson's r between all pairs of measures using the
- 171 function '*rcorr*' in the Hmisc R package (Harrell, 2012).
- 172 We employed indicator species analysis using 'multipatt' function and 'IndVal.g' attribute of the indicspecies R
- 173 package (Dufrene and Legendre, 1997; De Cáceres *et al.*, 2010) to identify significant (p < 0.05) indicator fungal
- 174 OTUs of the control, slag heap and settling pond sites by combining information of species abundances and
- occurrences in each group. Indicator values were tested using Monte Carlo permutation tests with 9,999
- 176 permutations.
- 177

178 **3. Results**

179 3.1. Fungal community diversity descriptors

180 Fungal abundance (estimated using 18S rRNA gene quantification), the number of OTUs determined at 97% 181 similarity and the taxonomic alpha diversity estimators based on OTU data are shown in Table 2. Fungal 182 abundance, the number of observed OTUs, and Chao1 richness indices were not significantly different among 183 the three groups of soils (Table 2). Slag heap sites harboured less diversified fungal communities, with lower 184 Shannon's diversity and Pielou's evenness indices than the control and settling pond sites. Redundancy analysis 185 was performed to identify the soil parameters that most influenced the alpha diversity indices (Chao1 richness, 186 Shannon diversity H', and Pielou's evenness J'). However, no significant soil characteristic among those 187 measured was identified as contributing significantly to the variation of alpha diversity descriptors.

188

189 3.2. Patterns of fungal phyla and classes

190 We investigated fungal diversity at the phylum level and at the class level for the two major phyla, *i.e.* 191 Basidiomycota and Ascomycota, and explored their relative proportions in each site and in the three groups of 192 soils (Figure 1). The relative proportions of each phylum varied among soils, with a majority of OTUs 193 belonging to Ascomycota, Basidiomycota and Cryptomycota (Figure 1A). Fungal communities from soils 194 classified in the same group had different compositions. For example, the relative abundance of Eurotiomycetes 195 was at least fifteen times higher in sh-Te and sh-Uc than in the other two slag heap soils. However, 196 PERMANOVA analysis indicated that the fungal assemblage at the phylum level was significantly different 197 among the three groups of soils (sum of squares = 0.47, $R^2 = 0.41$, F.model = 9.27, p < 0.001). In general, all 198 soils were dominated by Ascomycota (40.5 % to 90.5 % on average), except those from sites ctrl-He and sp-RM 199 that were dominated by Basidiomycota (61.8 %) and Cryptomycota (73.2 %), respectively (Figure 1A). The

200 relative abundance of Ascomycota was significantly higher in the slag heap sites than in the other 2 sites (Figure

201 1B). The relative abundance of Cryptomycota (98-100 % of the OTUs belonging to this phylum were affiliated 202 to the environmental clade LKM11) was significantly higher in the contaminated sites than in the control sites 203 (Figure 1B). The relative abundance of Basidiomycota representatives was significantly lower in the slag heap sites than in the settling pond sites (Figure 1B). Within the phylum Ascomycota, the relative abundance of fungi 204 205 affiliated to the Leotiomycetes, Pezizomycetes and Saccharomycetes classes was greater in the control sites than in the contaminated sites (Figure 1C). Although Dothideomycetes were not dominant, OTUs affiliated to this 206 207 class represented 21.8 % and 23.3 % on average of fungal community in sites sh-Ho and sh-NM, respectively, and were significantly enriched in the slag heap sites compared with the control sites, where this class 208 209 represented only 2.0 to 4.2 % on average (Figure 1C). Within the Basidiomycota phylum, the relative 210 abundance of the Agaricomycetes, Geminibasidiomycetes and Tremellomycetes classes was significantly higher 211 in the control sites than in the contaminated sites (Figure 1D).

- 212 Redundancy analysis (Figure 2A) was performed to identify the soil parameters that influenced the fungal 213 diversity structure at the phylum level. The first two axes of the RDA accounted for 89.4% of variability in the 214 relative abundance of fungal phyla. This variability was best explained by four of the measured soil parameters (silt, p=0.001; Zn, p=0.001; K, p=0.003; PAH, p=0.035; ANOVA). The three main phyla – Ascomycota, 215 Basidiomycota and Cryptomycota – were largely discriminated in the RDA plot. Ascomycota and Cryptomycota 216 217 were separated on the first axis, mostly explained by soil texture (silt percentage) and metal contamination (total 218 zinc concentration). The plot also showed that the relative abundance of Cryptomycota was positively influenced 219 by zinc concentration, and this observation was confirmed by a significant Pearson correlation between the 220 relative abundance of Cryptomycota and the total zinc and lead concentrations in the soils (p < 0.001, r = 0.93) 221 and r = 0.70, respectively; **Table S1**). The relative abundance of Ascomycota was influenced by soil texture 222 (negatively by the presence of silt) and by the PAH concentration (Figure 2A). We indeed confirmed a 223 significant Pearson correlation between the relative abundance of Ascomycota and the silt percentage (p < 0.001, r = -0.77, Table S1). In addition, the relative proportion of fungi affiliated to Dothideomycetes – a class of the 224 225 Ascomycota phylum – was positively correlated to the PAH concentration (Pearson, p < 0.001, r = 0.96, **Table** 226 S1). The relative abundance of Basidiomycota appeared positively influenced by the potassium (K) 227 concentration (Figure 2A). The Pearson correlation between the relative abundance of Basidiomycota and the soil characteristics evidenced a negative correlation with the soil pH (p < 0.001, r = -84, **Table S1**), and a 228 positive correlation with the dissolved organic carbon content (p < 0.001, r = 0.84, Table S1), two variables not 229 230 considered significant in the RDA model.
- 231

232 3.3. Patterns of fungal OTUs and indicator species

233 The analysis at the OTU level generated 1,614 OTUs, which ranged from 101 (sp-RM) to 191 (sp-Po) mean 234 OTUs per site, without any significant difference among the three groups of soils (Table 2). PERMANOVA analysis indicated that the fungal OTU assemblage structure was significantly different among the three groups 235 of soils (sum of squares = 1.5224, $R^2 = 0.24$, F.model = 4.273, p < 0.001). Redundancy analysis (Figure 2B) was 236 performed to identify the soil parameters that mostly influenced fungal diversity at the OTU level. The first two 237 238 axes of the RDA accounted only for 37.0% of variability in the relative abundance of fungal OTUs, which was 239 significantly explained by two of the measured soil parameters (Al, p=0.002; PAH, p=0.031). This low 240 percentage of variability explained means that there were other non-measured parameters influencing most the fungal diversity at OTU level. However, the control soils were separated from the 2 groups of contaminated soils on the first axis by aluminium concentrations. The two groups of slag heap and settling pond soils were not separated, but two soils of the slag heap group were separated from all the other contaminated soils on the second axis due to their higher PAH concentrations. The RDA plot highlighted a few OTUs explaining most of the variation among soil samples, and some were also identified as indicator species.

246 We searched for indicator species for the three groups of soils. Figure 3 shows the indicator species with the 247 lowest p-values (< 0.01; see **Table S2** for the complete list of indicator species). In the control group, 86 indicator species were identified, with more than half (45 OTUs) belonging to Basidiomycota and one third (27 248 249 OTUs) belonging to Ascomycota. Among the 29 highly significant indicator species of the control soils (Figure 250 3), 11 were abundant OTUs representing 1 to 12% relative abundance of the whole fungal community in some of 251 the control soils. Seventeen were affiliated to Basidiomycota, five to Ascomycota, four to Chytridiomycota, two 252 to Cryptomycota, and one to Mucoromycota. The Basidiomycota indicator species belonged to nine different 253 orders (Agaricales, Pucciniales, Cantharellales, Russulales, Tremellales, Geminibasidiales, Trichosporonales, 254 Kriegeriales, and Cortidiales). In the slag heap group, 22 indicator species were identified, 15 of which belonged 255 to Ascomycota. Among the 7 highly significant ones, four belonged to Ascomycota, one to Basidiomycota, one 256 to Cryptomycota LKM11 division and one to Mucoromycota (Figure 3). The Ascomycota indicator species 257 belonged to two main orders, namely Pleosporales and Chaetothyriales. In the settling pond group, 36 indicator 258 species were identified, 10 of which belonged to Cryptomycota LKM11 division. Among the four highly 259 significant indicator species, two belonged to Ascomycota, one to Basidiomycota and one to Mucoromycota 260 affiliated to the Glomeromycetes class.

261 262

263 **4. Discussion**

We hypothesized that in a context of long-term soil contamination, the type and level of contamination selected 264 265 adapted microorganisms and conducted to similar fungal community composition among soils sharing similar industrial history and contamination type and level. To test this hypothesis, we analysed fungal communities in a 266 267 collection of 30 soils from 10 sampling sites classified in three groups according to their physico-chemical and 268 contamination characteristics. As previously shown in Lemmel et al. (2019), slag heap soils presented a high 269 PAH concentration and an intermediate level of metal contamination, and settling pond soils presented a high 270 level of metal contamination (especially high Zn, Pb and total Cd concentrations) and intermediate PAH 271 concentrations. The control soils were characterised by the lowest level of both PAH and metal contaminations. 272 Although the soils affiliated to one group showed variable fungal community structures, our data confirmed our 273 hypothesis: significant differences were highlighted among the control, slag heap and settling pond soil groups, 274 at various taxonomic levels (from the phylum to the OTU), and indicator species specific to the three soil groups 275 were identified. Fungal abundance and specific richness did not differ among soil groups, indicating that 276 differences were mostly explained by changes in fungal taxonomic composition and/or relative taxon 277 abundances.

280 Our study showed a negative impact of contamination on many fungal taxa. In Basidiomycota phylum, the 281 Agaricomycetes, Geminibasidiomycetes and Tremellomycetes classes were represented in higher relative 282 proportions in the uncontaminated soils of the control group. Among the indicator species identified in the group of control soils, more than half (45 out of 86) belonged to Basidiomycota. Members of the Agaricomycetes class, 283 284 which dominate numerous terrestrial ecosystems (Lauber et al., 2008; Tedersoo et al., 2014), represented the majority (31 out of 45) of the Basidiomycota indicator species, and belonged to the Agaricales, Boletales, 285 286 Cantharellales, Cortitiales, Trechisporales, and Russulales orders. Agaricomycetes fill a wide range of ecological niches, but a very large number of species are wood-decay fungi that play a vital role in carbon cycling (de 287 288 Mattos-Shipley et al., 2016). Furthermore, the relative abundance of Basidiomycota, and among them 289 Agaricomycetes, was positively correlated to different soil fertility indicators, i.e. increased Na, K and DOC 290 concentrations. Canini et al. (2020) also found a positive correlation between the potassium soil concentration 291 and the relative abundance of Basidiomycota and Agaricomycetes, as highlighted by our RDA analysis. A higher 292 DOC concentration could enhance the development of Agaricomycetes or be the result of a higher organic 293 matter degradation rate in these soils. For example, lignin degradation activity can lead to the release of 294 dissolved organic carbon (Kalbitz et al., 2006). Industrial PAH- and metal-contaminated soils are disturbed 295 ecosystems that usually have lower fertility than unanthropised soils (Vetterlein & Hüttle 1999). A decrease of the relative proportion of Agaricomycetes in our contaminated soils could indicate a disruption of the carbon 296 297 cycle and slower organic matter mineralisation due to metal toxicity (Clemente et al., 2006). More globally, 298 Basidiomycota have been shown to be affected by metal contamination, with higher proportions in reference 299 soils than in metal-contaminated ones (Narendrula and Nkongolo, 2017).

300 The relative abundance of representatives of the Saccharomycetes and Leotiomycetes (both belonging to 301 Ascomycota) was higher in the control soils than in the two groups of contaminated soils. At a global scale, the 302 Saccharomycetes and Leotiomycetes classes represent 0.4 % and 7.1 % of fungal communities on average (Tedersoo et al., 2014), respectively, and were present at 1.4 to 5.5 % and 7.7 to 22.0 % in our control soils, 303 304 respectively. OTUs belonging to Saccharomycetes and Leotiomycetes were identified as indicator species of the 305 group of control soils. Among Saccharomycetes, Blastobotrys adenivorans, Debaryomyces hansenii, 306 Sugivamaella lignhabitans, and Ogataea methanolica, identified as control group indicator species, have been 307 described as versatile yeasts able to use a large variety of lignocellulosic and carbohydrate carbon sources mostly 308 on rotting wood or decaying organic matter and could have xerotolerance properties such as halotolerance or 309 thermotolerance (Kunze et al. 2014; Michán et al. 2013). Despite these adaptive properties, we could 310 hypothesize that Saccharomycetes- and Leotiomycetes-affiliated taxa are negatively impacted by soil 311 contamination. Ferrari et al. (2011) indeed found that Saccharomycetes and Leotiomycetes were dominant 312 among pristine-soil isolates but were replaced by other taxa after diesel addition.

Members of the Sordariomycetes class (Hypocreales, Sordariales, Ophiostomatales and Atractospora) were identified among indicator species of the control group, even if no difference appeared in their relative abundance between the groups of soils. Sordariomycetes are the second largest class of Ascomycota that represent 8.0 % of the fungal community in soils at a global scale (Tedersoo *et al.*, 2014) and are known as common soil fungal taxa belonging to many different functional guilds (Maharachchikumbura *et al.*, 2016). In the same line as in the present study, the relative abundance of Sordariomycetes was affected by high soil oil contamination (Bell *et al.*, 2014).

The relative proportion of fungi affiliated to the Chytridiomycota phylum was lower in the settling pond soils 320 321 than in the control soils, where Chytridiomycota OTUs were identified as indicator species. A similar 322 observation was made by Ventorino et al. (2018) who found that even if this phyla did not dominate the fungal 323 community, its proportion decreased in heavily hydrocarbon- and copper-contaminated soils compared with 324 uncontaminated ones. Chytridiomycota can also be inhibited by long-term heavy metal pollution (Chen et al., 325 2014). In the present work, the relative abundance of Chytridiomycota was negatively correlated to the zinc, lead 326 and iron concentrations, but positively correlated to soil nutrient availability (K and Na). The combination of 327 specific physico-chemical characteristics encountered in settling pond soils could explain why members of this 328 ancestral fungal phylum were found less adapted and less competitive than other fungi. For example, 329 Chytridiomycota are the only fungi producing unwalled, asexual spores that swim by means of a single flagellum (Longcore and Simmons, 2020) giving them a better ability to colonize humid or aquatic 330 331 environments, in quite different conditions from those found in metal-contaminated soils.

332

333 4.2. Fungal taxa favoured in slag heap soils

We assessed the impact of high PAH and medium metal contamination on the soil fungal community diversity of the slag heap group. Fungal Shannon diversity and Pielou's evenness indices were lower in the slag heap soils, indicating a disturbance of the relationships among fungal species, while no difference in the bacterial alpha diversity indices was previously found for the same soils (Lemmel *et al.*, 2019). Such reduced Shannon diversity was previously observed in oil-contaminated soil (Bell et al. 2014). Lower diversity and evenness but similar richness indicate that a few species dominate the fungal community, increasing the discrepancy between the abundances of the different taxa.

341 Members of the Dothideomycetes class (Ascomycota) were in higher relative proportions in the slag heap soils 342 than in the control soils. The relative abundance of Dothideomycetes was positively correlated to soil PAH contamination. At the world scale, this class represents 4.5 % of the fungal soil communities on average 343 344 (Tedersoo et al., 2014), but represented 21.8 and 23.3 % of the fungal communities of soils Ho and NM, 345 respectively - the two soils displaying the highest PAH contamination. An increase in culturable Dothideomycetes strains at high hydrocarbon concentrations was found by Ferrari et al. (2011). This class could 346 have been selected in PAH-contaminated soils due to its PAH tolerance or its PAH degradation capacity. In our 347 soils, the Dothideomycetes class was mostly represented by OTUs affiliated to Pleosporales, which was also one 348 349 dominant order found in hydrocarbon-contaminated soils (Bell et al., 2014). Moreover, a quarter of the indicator 350 species identified in the slag heap group belonged to the Pleosporales order, with representatives of the *Preussia*, 351 Corynespora, and Alternaria genera. Preussia is a soft rot fungi with important role in lignin degradation and 352 having ability to cleave aromatic rings compounds (Horwath 2007). Interestingly, several Dothideomycetes 353 genera, including *Preussia*, have been shown to harbour endohyphal bacteria able to biodegrade hydrocarbons 354 (Hoffman and Arnold, 2010). Moreover, members of the Corvnespora genus have been described as steroid and limonene-hydrocarbon degraders (Demyttenaere et al., 2001; Nassiri and Faramarzi, 2015), and found to 355 preferentially develop in a soil spiked with high amounts of phenanthrene (Li et al., 2019). Concerning members 356 357 of the Alternaria genus, some have been isolated for their PAH degradation capacities (Chávez-Gómez et al., 2003). 358

359 Although representatives of the Eurotiomycetes class were detected in variable proportions in our soil collection, 360 they largely dominated fungal communities in two slag heap soils (77.7 % in sh-Uc and 62.7 % sh-Te). This class of Ascomycota represented a high proportion of fungal strains isolated from heavy-metal-contaminated 361 362 sediments (Abdel-Azeem et al., 2015). Among indicator species of the slag heap group, we found 10 OTUs (half 363 of the indicator species) belonging to the Eurotiomycetes class, among which members of the Chaetothyriales 364 Sclerococcales and Eurotiales orders. Chaetothyriales have been detected in a wide range of habitats (Gueidan et 365 al., 2014), so that one cannot be conclusive about their ecological niche. However, six OTUs belonging to Herpotrichiellaceae family (Chaetothyriales order), with three belonging to Exophiala calicioides and one to 366 367 Cladophialophora chaetospira species, were among the 22 indicator species of slag heap soils. Members of 368 Cladophialophora and Exophiliala genera, described as black yeast-like fungi, were largely studied for their 369 bioremediation potential (Prenafeta-Boldú et al. 2004). The genus Exophiliala, is comprised of opportunistic 370 pathogens as well as species recovered from hydrocarbon contaminated environments, such as E. xenobiotica 371 (De Hoog et al. 2006) and E. sideris (also having the ability to tolerate arsenic; Seyedmousavi et al. 2011). 372 Strains with the ability to degrade hydrocarbon (benzene, toluene, ethylbenzene or xylene) were often isolated, 373 such as E. maquariensis (Zhang et al. 2019), E. xenobiotica and E. bergeri (Zhao et al 2010).

374

375 4.3. Fungal taxa favoured in settling pond soils

376 The alpha diversity (richness, evenness and diversity) of the settling pond soils was not different from that of the 377 control soils. However, fungal community structure differences were observed between the settling pond and 378 control soils. Members of the Pucciniomycetes class (Basidiomycota) were in a higher relative proportion in the 379 settling pond soils. The relative abundance of Pucciniomycetes was positively correlated to the Ni, Co and Mo 380 metal concentrations. A quarter (8 out of 36) of the settling ponds indicator species belonged to Pucciniomycetes 381 and were affiliated to the Gymnosporangium genus, whose members are well known phytopathogens causing 382 rust (Yun et al., 2009). Similarly, one of the slag heap indicator species belonged to Rhizoctonia genus known as 383 plant pathogen with a wide host range and worldwide distribution and causing collar, crown or root rots (Lenart-384 Boroń & Boroń, 2014). We can therefore hypothesize that chronic soil pollution induced the selection of these 385 opportunistic plant pathogens, and maybe others, which is worrying for the ecosystem balance. Besides, the settling pond group was the only one that harboured one OTU belonging to the Glomeromycetes class 386 387 (Mucoromycota phylum) and affiliated to Entrophospora infrequens as an indicator species. This fungus is 388 known as an arbuscular mycorrhizal fungus (AMF) beneficial for plants (Koziol and Bever, 2016). Seven OTUs 389 identified as settling pond soil indicator species belonged to Pezizomycotina sub-division of the Ascomycota, 390 but they belong to various classes and genera having various ecological preference, being geophilic fungi such as Dothidea, Parascedosporium, and Helvella genera, being lichenicolous fungi such as Lichina and 391 392 Rhymbocarpus, or entomopathogen such as Metarhizium.

393

394 *4.4. Fungal taxa favoured in the two contaminated soils*

The relative abundance of fungi belonging to the Cryptomycota phylum was higher in the slag heap and settling pond soils than in the control soils. Nearly half of the indicator species of the settling pond group and two indicator species of slag heap group were affiliated to Cryptomycota. This newly proposed phylum is composed of atypical fungi that have been mostly detected in freshwater ecosystems to date. They are single-cell fungi able

399 to form epibiont associations (Jones et al., 2011), they lack a rigid chitin-/cellulose-rich wall, and they are 400 possibly capable of phagotrophy, as described for the Rozella genus (Pöggeler and Wöstemeyer, 2011). Although, Cryptomycota represented 0.3 to 72.7% of the fungal communities in our contaminated soils, they 401 represent only a slight part (0.9 % on average) of fungal soil communities at the world scale (Tedersoo et al., 402 403 2014), but Lazarus and James (2015) found that they were ubiquitous in terrestrial and aquatic ecosystems. 404 Cryptomycota members could be tolerant to metal contamination because their relative abundance was positively 405 correlated to the total zinc and lead concentrations. They were recently detected as dominant fungi colonizing microplastics in aquatic ecosystems (Kettner et al., 2017), which are well-known metal adsorption matrices 406 407 (Cole et al., 2011; Brennecke et al., 2016). In our study, Cryptomycota were mostly represented by sequences 408 belonging to OTUs related to the LKM11 environmental clade. This clade forms a deep branch of true fungi, but 409 its ecology and physiology is still unclear because no culturable representative currently exists (Lara et al., 410 2010). However, based on our data, we can suppose that this LKM11 clade is well adapted to metal 411 contamination in soil ecosystems, even if it was recently detected as dominant in pristine limnetic ecosystems 412 (Rojas-Jimenez et al., 2017). Therefore, its presence in contaminated soils could depend more on other soil 413 parameters such as carbon resources because positive correlation between relative Cryptomycota abundance and 414 organic matter has also been detected.

415

416 5. Conclusions

417 Our study brings the first report about the diversity and structure of fungal communities in aged multi-418 contaminated soils, and identifies fungal taxa affected or favoured in PAH- and/or metal-contaminated soils vs. 419 control soils. Fungal abundance and richness were not significantly different among soil groups, while Shannon 420 diversity and evenness indices were lower in the slag heap soils than in the control soils. The differences among 421 the three groups of soils were confirmed based on fungal diversity at the phylum, class and OTU levels. 422 concerning the occurrence of certain fungal phyla, classes and indicator species depending on the soil physico-423 chemical and contaminant characteristics were highlighted but need to be confirmed on a wider range of sites. 424 Basidiomycota, among which Agaricomycetes, as well as Saccharomycetes and Leotiomycetes (Ascomycota), 425 well known as wood-decaying fungi, were repressed in the two contaminated soil groups compared with the 426 control soils, potentially having an impact on carbon cycle. Cryptomycota representatives, such as the LKM11 427 clade, were favoured in the slag heap and settling pond soils, probably due to the presence of metals such as zinc and lead. Members of the Dothideomycetes (Preussia, Corvnespora, and Alternaria) and Eurotiomycetes 428 429 (Exophiala calicioides and Cladophialophora chaetospira) classes were identified as indicator species of the 430 slag heap soils and could be interesting PAH-degraders as their relative abundance was positively linked to the 431 soil PAH contamination level. Indicator species of the settling pond group mainly belonged to the 432 Gymnosporangium genus (Pucciniomycetes) known as phytopathogen. The contaminated soils could be 433 reservoirs of plant or animal pathogens thus having concern for environmental health but the selection of protective species such as mycorrhizal fungi (Glomeromycetes class) could indicate resistance of the ecosystem. 434 435 Briefly, the fungal communities of contaminated soils could be altered for organic matter recycling functions, 436 and harbour pathogens, but it also potentially contain useful fungi for improving bioremediation strategies.

- 437
- 438

439 *Acknowledgements*

- 440 This work was supported by the French national program EC2CO (Ecobios project), and the OSU-OteLo 441 (TraitMic project). We would like to thank Arcelor Mittal, EPFL, GISFI, ONF (E. Magnier), and LTO of
- 442 Montiers (ANDRA/INRA, M.P. Turpault) for giving us access to the different sampling sites.
- 443 Authors have no conflicts of interest to declare.
- 444
- 445

446 Table 1. Physico-chemical characteristics of the soils from the 10 studied sites and mean values for the

447 three groups: controls ctrl (n = 3), slag heaps sh (n = 4), and settling ponds sp (n = 3). *: soil classification is

448 based on WRB 2016. Significant results of Kruskal-Wallis tests (p < 0.05), testing differences among the 3 soil

449 groups, are indicated by different letters and no letter appears when non-significant differences were found.

- 450 CEC: Cationic Exchange Capacity.
- 451

_

_

| | Soil site group | Soil classification* | рН | CEC | carbonates | organic C | total N | Texture (%) | Fe | Al | К | Zn | Pb | Cd | Cu | РАН |
|--|--------------------------------------|--|---|---|---|--|--|--|---|---|---|---|---|---|---|--|
| | 0.00 | | | (cmol kg ⁻¹) | | (g kg ⁻¹) | | clay:silt:sand | | (g kg ⁻¹) | | | (n | ng kg ⁻¹) | | |
| Mo He Di Uc Te NM Ho Po | ctrl ctrl sh sh sh sp | Cambisol Luvisol Fluvisol Technosol Technosol Technosol Technosol Technosol | 7.0 5.4 7.3 8.0 7.7 7.4 7.5 7.8 7.2 | 35 15 10 4 20 17 10 51 | 5 1 14 8 181 14 114 89 | 81 72 24 30 88 75 159 109 | 4.6 2.8 1.7 0.6 4.3 2.3 2.9 7.2 | 49:46:5 26:69:5 12:14:74 5:18:77 10:21:69 13:28:59 7:12:81 18:53:29 | 44 27 18 452 108 193 66 93 | 67 52 45 20 25 40 40 40 21 | 14.6 18.6 28.6 0.6 5.4 9.2 5.8 5.4 15 | 144 248 60 74 1650 2540 314 29600 55400 | 44 93 31 23 475 653 303 34700 | 0.7 0.3 0.2 0.1 4.5 3.3 1.1 152.0 | 16 13 10 9 146 144 94 148 186 | 3.5 0.1 0.9 0.9 114.7 1095.9 937.7 21.2 49.6 |
| RM | sp | Technosol | 7.6 | 10 | 219 | 149 | 3.8 | 11 : 75 : 14 | 54 | 22 | 3.4 | 119000 | 39500 | 22.0 | 42 | 6.6 |
| Mean per group | ctrl sh sp | | 6.6 7.6 7.6 | 20 13 26 | 7 80 154 | 59 88 126 | 3.1 2.5 5.1 | 29 : 43 : 28 9 : 20 : 71 13 : 67 : 20 | 30 ^b 205 ^a 150 ^a | 55 ^a 31 ^b 27 ^b | 20.6 ^a 5.2 ^b 3.4 ^b | 151 ^b 1144 ^b 68000 ^a | 56 ^b 363 ^b 29533 ^a | 0.4 ^b 2.2 ^b 63.9 ^a | 13 98 125 | 1.5 ^b 537.3 ^a 25.8 ^a |

452 453

Table 2. Fungal abundance and taxonomic alpha diversity indices of the soils from the 10 studied sites. 454

Values are means $(n = 3) \pm$ standard error of the mean. For soil groups, values are means (n=3 for ctrl and sp)455 456 soils, and n = 4 for sh soils) \pm standard error of the mean. Significant results of Kruskal-Wallis tests (p < 0.05),

testing differences among the 3 soil groups, are indicated by different letters, and no letter appears when non-457

458 significant differences were found.

| | Soil site group | 18S rRNA gene copy number (x 10 ⁹ g ⁻¹) | Observed number of OTUs | Chao1 index | Shannon's diversity index H' | Pielou's evenness index J' |
|-------|-----------------|---|----------------------------|-------------|---------------------------------|-------------------------------|
| Mo | ctrl | 48+06 | 162 + 15 | 215 + 31 | 3 40 + 0 02 | 0.669 ± 0.009 |
| He | ctrl | 45.0 + 1.8 | 176 + 10 | 281 + 48 | 3.08 + 0.04 | 0.596 + 0.018 |
| Di | ctrl | 21.8 ± 1.2 | 155 ± 7 | 203 ± 15 | 2.77 ± 0.09 | 0.549 ± 0.014 |
| Uc | sh | 5.1 ± 0.3 | 103 ± 4 | 175 ± 7 | 1.25 ± 0.05 | 0.271 ± 0.011 |
| Te | sh | 34.2 ± 3.0 | 113 ± 2 | 147 ± 15 | 2.08 ± 0.06 | 0.440 ± 0.011 |
| NM | sh | 3.1 ± 0.3 | 150 ± 7 | 179 ± 20 | 2.92 ± 0.03 | 0.581 ± 0.002 |
| Но | sh | 11.8 ± 0.2 | 190 ± 9 | 257 ± 6 | 2.75 ± 0.01 | 0.529 ± 0.005 |
| Po | sp | 18.1 ± 1.6 | 196 ± 8 | 317 ± 22 | 3.14 ± 0.02 | 0.593 ± 0.006 |
| MsM | sp | 10.3 ± 0.9 | 148 ± 7 | 186 ± 9 | 3.11 ± 0.02 | 0.623 ± 0.002 |
| RM | sp | 8.5 ± 0.7 | 101 ± 5 | 158 ± 15 | 1.46 ± 0.05 | 0.313 ± 0.007 |
| Mean | ctrl | 238+59 | 164 + 6 | 233 + 21 | 3 08 + 0 10 ^a | 0 605 + 0 019 ª |
| ner | sh | 135+38 | 139 + 11 | 190 + 14 | 2 25 ± 0 20 b | 0.455 ± 0.015 |
| group | sp | 12.3 ± 1.6 | 148 ± 14 | 220 ± 26 | 2.57 ± 0.28 ° | 0.510 ± 0.049 ° |

460 Figure captions

461

Figure 1 | Taxonomic composition of the fungal communities of soil samples from the 10 studied sites and 462 differences among groups of sites. A) Relative abundance of major fungal phyla and major classes for 463 464 Ascomycota and Basidiomycota in the 10 studied sites. Values are means (n = 3), and error-bars represent standard errors of the mean for each site. The group named 'Others' corresponded to Blastocladiomycota, 465 466 LKM15, Neocallimastigomycota, Zoopagomycota, Entorrhizomycota, unclassified fungi, and taxa affiliated to Basidiomycota and Ascomycota but not shown on the graph due to their minor relative abundances. B) Relative 467 468 abundance of the five major phyla in the three groups of sites, C) Relative abundance of Ascomycota major 469 classes in the three groups of sites, and D) Relative abundance of Basidiomycota major classes in the three groups of sites. Values are means (n=9 for *ctrl* and *sp* group and n=12 for *sh* group). Statistical differences 470 among phylum and class proportions in the soil groups are indicated by different letters (Kruskal-Wallis test, * p 471 < 0.05, ** p < 0.001, ***p < 0.0001, n.s: not significant). 472

- 473
- 474

Figure 2 | Redundancy analysis (RDA) plots showing influence of significant soil characteristics (blue 475 vectors) on the fungal communities of the ten studied sites. A) Fungal phyla relative abundance (in red) and 476 477 B) Fungal OTUs relative abundance (in red; only the names of some OTUs highly discriminating samples on 478 first plan are shown, those circled in blue and green are those from indicator species, cf. Figure 3). Soils (Di: 479 Dieulouard, Ho: Homécourt, He: Hémilly, MsM: Mont St Martin, Mo: Montiers, RM: Russange-Micheville, NM: Neuves-Maisons, Po: Pompey, Te: Terville, Uc: Uckange) were coloured according to the group they 480 481 belong to, ie. settling ponds in red, slag heaps in green and controls in blue. Soil characteristics: PAH, 482 polycyclic aromatic hydrocarbons; K, Zn and Al: total concentration of potassium, zinc and aluminium, 483 respectively; silt: the percentage of silt of the soil texture.

484 485

Figure 3 | Indicator species identified for control, slag heap and settling pond groups of soil (in blue, green and red, respectively), their mean relative abundance in the group and indicator value (Indval). Only indicator species with the lower p-values (p<0.01) are presented (see Table S2 for complete list of indicator species). OTU lowest possible taxonomic classifications are given next to the OTU number with the prefix corresponding to taxonomic rank (s-d = sub-division, c = class, o = order, f = family, g = genus, no prefix = species). Indicator species also appearing as discriminant OTUs on RDA plot (Figure 2B) are indicated with *.

- 492
- 493

494 References

- Abdel-Azeem, A., El-Morsy, E., Nour El-Dein, M., and Rashad, M. (2015) Occurrence and diversity of mycobiota in heavy metal contaminated sediments of Mediterranean coastal lagoon El-Manzala, Egypt.
 Mycosphere 6:228-240. https://doi.org/10.5943/mycosphere/6/2/12
- Beauregard, M.S., Hamel, C., Atul-Nayyar, A., and St-Arnaud, M. (2010) Long-term phosphorus fertilization impacts soil fungal and bacterial diversity but not AM fungal community in alfalfa. *Microb Ecol* 59: 379–389. https://doi.org/10.1007/s00248-009-9583-z
- Bell, T.H., El-Din Hassan, S., Lauron-Moreau, A., Al-Otaibi, F., Hijri, M., Yergeau, E., and St-Arnaud, M.
 (2014) Linkage between bacterial and fungal rhizosphere communities in hydrocarbon-contaminated soils is related to plant phylogeny. *ISME J* 8: 331–343. https://doi.org/10.1038/ismej.2013.149
- Bourceret, A., Cébron, A., Tisserant, E., Poupin, P., Bauda, P., Beguiristain, T., and Leyval, C. (2016) The
 bacterial and fungal diversity of an aged PAH- and heavy metal-contaminated soil is affected by plant
 cover and edaphic parameters. *Microb Ecol* **71**: 711–724. https://doi.org/10.1007/s00248-015-0682-8
- Bourceret, A., Leyval, C., Fouquet, C. de, and Cébron, A. (2015) Mapping the centimeter-scale spatial
 variability of PAHs and microbial populations in the rhizosphere of two plants. *PLOS ONE* 10:
 e0142851. https://doi.org/10.7910/DVN/EFXVQH
- Brennecke, D., Duarte, B., Paiva, F., Caçador, I., and Canning-Clode, J. (2016) Microplastics as vector for heavy
 metal contamination from the marine environment. *Estuar Coast Shelf Sci* 178: 189–195.
 https://doi.org/10.1016/j.ecss.2015.12.003
- Canini, F., Geml, J., D'Acqui, L.P., Selbmann, L., Onofri, S., Ventura, S., and Zucconi, L. (2020) Exchangeable 514 515 cations and pH drive diversity and functionality of fungal communities in biological soil crusts from 516 coastal sites of Victoria Land. Antarctica. Fungal Ecol **45**: 100923. 517 https://doi.org/10.1016/j.funeco.2020.100923
- Chávez-Gómez, B., Quintero, R., Esparza-García, F., Mesta-Howard, A.M., Zavala Díaz de la Serna, F.J.,
 Hernández-Rodríguez, C.H., et al. (2003) Removal of phenanthrene from soil by co-cultures of bacteria
 and fungi pregrown on sugarcane bagasse pith. *Bioresour Technol* 89: 177–183.
 https://doi.org/10.1016/S0960-8524(03)00037-3
- Chen, J., He, F., Zhang, X., Sun, X., Zheng, J., and Zheng, J. (2014) Heavy metal pollution decreases microbial
 abundance, diversity and activity within particle-size fractions of a paddy soil. *FEMS Microbiol Ecol* 87: 164–181. https://doi.org/10.1111/1574-6941.12212
- Clemente, R., Escolar, Á., and Bernal, M.P. (2006) Heavy metals fractionation and organic matter mineralisation
 in contaminated calcareous soil amended with organic materials. *Bioresour Technol* 97: 1894–1901.
 https://doi.org/10.1016/j.biortech.2005.08.018
- Cole, M., Lindeque, P., Halsband, C., and Galloway, T.S. (2011) Microplastics as contaminants in the marine
 environment: A review. Mar Pollut Bull 62: 2588–2597.
 https://doi.org/10.1016/j.marpolbul.2011.09.025
- De Cáceres, M., Legendre, P., and Moretti, M. (2010) De Cáceres M, Legendre P, Moretti M.. Improving
 indicator species analysis by combining groups of sites. Oikos 119: 1674-1684. *Oikos* 119: 1674–1684.
 https://doi.org/10.1111/j.1600-0706.2010.18334.x
- De Hoog, G. S., Zeng, J. S., Harrak, M. J., & Sutton, D. A. (2006). Exophiala xenobiotica sp. nov., an
 opportunistic black yeast inhabiting environments rich in hydrocarbons. *Antonie van Leeuwenhoek*,
 90(3), 257-268. doi: 10.1007/s10482-006-9080-z
- Demyttenaere, J., Belleghem, K., and Kimpe, N. (2001) Biotransformation of (R)-(+)- and (S)-(-)-limonene by
 fungi and the use of solid phase microextraction for screening. *Phytochemistry* 57: 199–208.
 https://doi.org/10.1016/S0031-9422(00)00499-4
- Dufrene, M. and Legendre, P. (1997) Species assemblages and indicator species: the need for a flexible
 asymmetrical approach. *Ecol Monogr* 67: 345–366. https://doi.org/10.1890/0012 9615(1997)067[0345:SAAIST]2.0.CO;2
- Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., and Knight, R. (2011) UCHIME improves sensitivity and
 speed of chimera detection. *Bioinforma Oxf Engl* 27: 2194–2200.
 https://doi.org/10.1093/bioinformatics/btr381
- Ferrari, B.C., Zhang, C., and van Dorst, J. (2011) Recovering greater fungal diversity from pristine and diesel
 fuel contaminated sub-Antarctic soil through cultivation using both a high and a low nutrient media
 approach. *Front Microbiol* 2:217. https://doi.org/10.3389/fmicb.2011.00217
- Gueidan, C., Aptroot, A., Cáceres, M., Badali, H., and Stenroos, S. (2014) A reappraisal of orders and families
 within the subclass Chaetothyriomycetidae (Eurotiomycetes, Ascomycota). *Mycol Prog* 13:990.
 https://doi.org/10.1007/s11557-014-0990-2

- Hannula, S.E., de Boer, W., and van Veen, J. (2012) A 3-year study reveals that plant growth stage, season and
 field site affect soil fungal communities while cultivar and GM-trait have minor effects. *PloS One* 7:
 e33819. https://doi.org/10.1371/journal.pone.0033819
- Harms, H., Schlosser, D., and Wick, L.Y. (2011) Untapped potential: exploiting fungi in bioremediation of
 hazardous chemicals. *Nat Rev Microbiol* 9: 177–192. https://doi.org/10.1038/nrmicro2519
- Hill, T.C.J., Walsh, K.A., Harris, J.A., and Moffett, B.F. (2003) Using ecological diversity measures with
 bacterial communities. *FEMS Microbiol Ecol* 43: 1–11. https://doi.org/10.1111/j.1574 6941.2003.tb01040.x
- Hoffman, M.T. and Arnold, A.E. (2010) Diverse bacteria inhabit living hyphae of phylogenetically diverse
 fungal endophytes. *Appl Environ Microbiol* 76: 4063–4075. https://doi.org/10.1128/AEM.02928-09
- Horwath, W. (2007). Carbon cycling and formation of soil organic matter. In *Soil microbiology, ecology and biochemistry* (pp. 303-339). Academic Press.
- Jones, M.D.M., Forn, I., Gadelha, C., Egan, M.J., Bass, D., Massana, R., and Richards, T.A. (2011) Discovery of
 novel intermediate forms redefines the fungal tree of life. *Nature* 474: 200–203.
 https://doi.org/10.1038/nature09984
- Kalbitz, K., Kaiser, K., Bargholz, J., and Dardenne, P. (2006) Lignin degradation controls the production of
 dissolved organic matter in decomposing foliar litter. *Eur J Soil Sci* 57: 504–516.
 https://doi.org/10.1111/j.1365-2389.2006.00797.x
- Kettner, M.T., Rojas-Jimenez, K., Oberbeckmann, S., Labrenz, M., and Grossart, H.-P. (2017) Microplastics
 alter composition of fungal communities in aquatic ecosystems. *Environ Microbiol* 19: 4447–4459. https://doi.org/10.1111/1462-2920.13891
- Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., and Schloss, P.D. (2013) Development of a dualindex sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq
 Illumina sequencing platform. *Appl Environ Microbiol* **79**: 5112–5120.
 https://doi.org/10.1128/AEM.01043-13
- Koziol, L. and Bever, J. (2016) AMF, phylogeny, and succession: Specificity of response to mycorrhizal fungi
 increases for late-successional plants. *Ecosphere* 7:e01555. https://doi.org/10.1002/ecs2.1555
- Kunze, G., Gaillardin, C., Czernicka, M., Durrens, P., Martin, T., Böer, E., ... & Goffeau, A. (2014). The complete genome of Blastobotrys (Arxula) adeninivorans LS3-a yeast of biotechnological interest. *Biotechnology for biofuels*, 7(1), 66. https://doi.org/10.1186/1754-6834-7-66
- Lara, E., Moreira, D., and López-García, P. (2010) The environmental clade LKM11 and Rozella form the deepest branching clade of fungi. *Protist* 161: 116–121. https://doi.org/10.1016/j.protis.2009.06.005
- Lauber, C.L., Strickland, M.S., Bradford, M.A., and Fierer, N. (2008) The influence of soil properties on the
 structure of bacterial and fungal communities across land-use types. *Soil Biol Biochem* 40: 2407–2415.
 https://doi.org/10.1016/j.soilbio.2008.05.021
- Lazarus, K. L., & James, T. Y. (2015). Surveying the biodiversity of the Cryptomycota using a targeted PCR
 approach. *Fungal Ecology*, 14, 62-70. http://dx.doi.org/10.1016/j.funeco.2014.11.004
- Lemmel, F., Maunoury-Danger, F., Fanesi, A., Leyval, C., and Cébron, A. (2019) Soil properties and multi pollution affect taxonomic and functional bacterial diversity in a range of French soils displaying an
 anthropisation gradient. *Microb Ecol* 77: 993–1013. https://doi.org/10.1007/s00248-018-1297-7
- Lenart-Boroń, A., & Boroń, P. (2014). The effect of industrial heavy metal pollution on microbial abundance
 and diversity in soils—a review. In *Environmental risk assessment of soil contamination*. IntechOpen.
 DOI: 10.5772/57406
- Li, X., Song, Y., Yao, S., Bian, Y., Gu, C., Yang, X., et al. (2019) Can biochar and oxalic acid alleviate the toxicity stress caused by polycyclic aromatic hydrocarbons in soil microbial communities? *Sci Total Environ* 695: 133879. https://doi.org/10.1016/j.scitotenv.2019.133879
- Longcore, J.E. and Simmons, D.R. (2020) Chytridiomycota. In *eLS*. American Cancer Society. https://doi.org/10.1002/9780470015902.a0000349.pub4
- Lueders, T., Wagner, B., Claus, P., and Friedrich, M.W. (2004) Stable isotope probing of rRNA and DNA
 reveals a dynamic methylotroph community and trophic interactions with fungi and protozoa in oxic
 rice field soil. *Environ Microbiol* 6: 60–72. https://doi.org/10.1046/j.1462-2920.2003.00535.x
- Maharachchikumbura, S.S.N., Hyde, K.D., Jones, E.B.G., McKenzie, E.H.C., Bhat, J.D., Dayarathne, M.C., et
 al. (2016) Families of Sordariomycetes. *Fungal Divers* **79**:1-317. https://doi.org/10.1007/s13225-016 0369-6
- de Mattos-Shipley, K.M.J., Ford, K.L., Alberti, F., Banks, A.M., Bailey, A.M., and Foster, G.D. (2016) The
 good, the bad and the tasty: The many roles of mushrooms. *Stud Mycol* 85: 125–157.
 https://doi.org/10.1016/j.simyco.2016.11.002
- Mendiburu, F. de (2013) agricolae: Statistical Procedures for Agricultural Research. R Package Version 1.3-2.
 http://CRAN.R-project.org/package=agricolae

- Michán, C., Martínez, J. L., Alvarez, M. C., Turk, M., Sychrova, H., & Ramos, J. (2013). Salt and oxidative
 stress tolerance in Debaryomyces hansenii and Debaryomyces fabryi. *FEMS yeast research*, *13*(2), 180 188. https://doi.org/10.1111/1567-1364.12020
- Narendrula, R. and Nkongolo, K. (2017) Bacterial and fungal community structure and diversity in a mining
 region under long-term metal exposure revealed by metagenomics sequencing. *Ecol Genet Genomics* 2:
 13–24. https://doi.org/10.1016/j.egg.2016.11.001
- Nassiri, N. and Faramarzi, M. (2015) Recent developments in the fungal transformation of steroids. *Biocatal Biotransformation* 33:1-28. https://doi.org/10.3109/10242422.2015.1022533
- 619 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., et al. (2017) Vegan:
- 620 Community Ecology Package. https://CRAN.R-project.org/package=vegan
- Op De Beeck, M., Lievens, B., Busschaert, P., Rineau, F., Smits, M., Vangronsveld, J., and Colpaert, J.V. (2015)
 Impact of metal pollution on fungal diversity and community structures. *Environ Microbiol* 17: 2035–2047. https://doi.org/10.1111/1462-2920.12547
- Pöggeler, S. and Wöstemeyer, J. eds. (2011) Evolution of Fungi and Fungal-Like Organisms, Berlin Heidelberg:
 Springer-Verlag.
- Prenafeta-Boldú, F. X., Ballerstedt, H., Gerritse, J., & Grotenhuis, J. T. C. (2004). Bioremediation of BTEX
 hydrocarbons: effect of soil inoculation with the toluene-growing fungus Cladophialophora sp. strain
 T1. *Biodegradation*, 15(1), 59-65. https://doi.org/10.1023/B:BIOD.0000009973.53531.96
- Rachwał, M., Magiera, T., and Wawer, M. (2015) Coke industry and steel metallurgy as the source of soil contamination by technogenic magnetic particles, heavy metals and polycyclic aromatic hydrocarbons.
 Chemosphere 138: 863–873. https://doi.org/10.1016/j.chemosphere.2014.11.077
- 632 Rojas-Jimenez, K., Wurzbacher, C., Bourne, E.C., Chiuchiolo, A., Priscu, J.C., and Grossart, H.-P. (2017) Early 633 diverging lineages within Cryptomycota and Chytridiomycota dominate the fungal communities in ice-634 covered McMurdo Dry Valleys, Antarctica. Sci Rep 7: 15348. lakes of the https://doi.org/10.1038/s41598-017-15598-w 635
- Salvo, V.S., Gallizia, I., Moreno, M., and Fabiano, M. (2005) Fungal communities in PAH-impacted sediments
 of Genoa-Voltri Harbour (NW Mediterranean, Italy). *Mar Pollut Bull* 50: 553–559. https://doi.org/10.1016/j.marpolbul.2005.01.001
- Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., et al. (2009) Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* **75**: 7537–7541.
 https://doi.org/10.1128/AEM.01541-09
- Seyedmousavi, S., Badali, H., Chlebicki, A., Zhao, J., Prenafeta-Boldu, F. X., & De Hoog, G. S. (2011).
 Exophiala sideris, a novel black yeast isolated from environments polluted with toxic alkyl benzenes and arsenic. *Fungal biology*, *115*(10), 1030-1037. doi: 10.1016/j.funbio.2011.06.004.
- Tedersoo, L., Bahram, M., Põlme, S., Kõljalg, U., Yorou, N.S., Wijesundera, R., et al. (2014) Global diversity
 and geography of soil fungi. *Science* 346: 1256688. https://doi.org/10.1126/science.1256688
- Thion, C., Cébron, A., Beguiristain, T., and Leyval, C. (2012) Long-term in situ dynamics of the fungal communities in a multi-contaminated soil are mainly driven by plants. *Fems Microbiol Ecol* 82: 169–181. https://doi.org/10.1111/j.1574-6941.2012.01414.x
- Thomas, F. and Cébron, A. (2016) Short-term rhizosphere effect on available carbon sources, phenanthrene
 degradation, and active microbiome in an aged-contaminated industrial soil. *Front Microbiol* 92.
 https://doi.org/10.3389/fmicb.2016.00092
- US EPA, R. (2014) Polycyclic Aromatic Hydrocarbons (PAHs) Fact Sheet. US EPA.
- Vainio, E.J. and Hantula, J. (2000) Direct analysis of wood-inhabiting fungi using denaturing gradient gel
 electrophoresis of amplified ribosomal DNA. *Mycol Res* 104: 927–936.
 https://doi.org/10.1017/S0953756200002471
- Ventorino, V., Pascale, A., Adamo, P., Rocco, C., Fiorentino, N., Mori, M., et al. (2018) Comparative
 assessment of autochthonous bacterial and fungal communities and microbial biomarkers of polluted
 agricultural soils of the Terra dei Fuochi. *Sci Rep* 8: 14281. https://doi.org/10.1038/s41598-018-326885
- Vetterlein D, Hüttl RF (1999) Can applied organic matter fulfilsimilar functions as soil organic matter? Risk benefit analysisfor organic matter application as a potential strategy for reha-bilitation of disturbed
 ecosystems. Plant Soil 213, 1–10. https://doi.org/10.1023/A:1004681506901
- Wang, S., Gutell, R. R., & Miranker, D. P. (2007). Biclustering as a method for RNA local multiple sequence
 alignment. *Bioinformatics*, 23(24), 3289-3296. https://doi.org/10.1093/bioinformatics/btm485
- Yun, H.Y., Hong, S.G., Rossman, A., Lee, S., Lee, K., and Bae, K.S. (2009) The rust fungus Gymnosporangium
 in Korea including two new species, G. monticola and G. unicorne. *Mycologia* 101: 790–809.
 https://doi.org/10.3852/08-221

- Zarei, M., Hempel, S., Wubet, T., Schäfer, T., Savaghebi, G., Jouzani, G.S., et al. (2010) Molecular diversity of
 arbuscular mycorrhizal fungi in relation to soil chemical properties and heavy metal contamination.
 Environ Pollut 158: 2757–2765. https://doi.org/10.1016/j.envpol.2010.04.017
- Zhang, C., Sirijovski, N., Adler, L., & Ferrari, B. C. (2019). Exophiala macquariensis sp. nov., a cold adapted
 black yeast species recovered from a hydrocarbon contaminated sub-Antarctic soil. *Fungal biology*,
 123(2), 151-158. DOI: 10.1016/j.funbio.2018.11.011
- Zhao, J., Zeng, J., De Hoog, G. S., Attili-Angelis, D., & Prenafeta-Boldú, F. X. (2010). Isolation and identification of black yeasts by enrichment on atmospheres of monoaromatic hydrocarbons. *Microbial Ecology*, 60(1), 149-156. DOI: 10.1007/s00248-010-9651-4
- Zhou, Z.-F., Wang, M.-X., Zuo, X.-H., and Yao, Y.-H. (2017) Comparative investigation of Bacterial, Fungal,
 and Archaeal community structures in soils in a typical oilfield in Jianghan, China. Arch Environ
 Contam Toxicol 72: 65–77. https://doi.org/10.1007/s00244-016-0333-1
- 682

9

sp















| | Soil site | Soil | л Ц | CEC | carbonate s | organic C | total N | | |
|------------|-----------|-----------|------------|-----|----------------|-------------|---------|-----------------------|--|
| | group | n | рн | рп | μц | (cmol kg⁻¹) | | (g kg ⁻¹) | |
| Мо | ctrl | Cambisol | 7.0 | 35 | 5 | 81 | 4.6 | | |
| He | ctrl | Luvisol | 5.4 | 15 | 1 | 72 | 2.8 | | |
| Di | ctrl | Fluvisol | 7.3 | 10 | 14 | 24 | 1.7 | | |
| Uc | sh | Technosol | 8.0 | 4 | 8 | 30 | 0.6 | | |
| Те | sh | Technosol | 7.7 | 20 | 181 | 88 | 4.3 | | |
| NM | sh | Technosol | 7.4 | 17 | 14 | 75 | 2.3 | | |
| Но | sh | Technosol | 7.5 | 10 | 114 | 159 | 2.9 | | |
| Ро | sp | Technosol | 7.8 | 51 | 89 | 109 | 7.2 | | |
| MsM | sp | Technosol | 7.2 | 16 | 152 | 119 | 4.2 | | |
| RM | sp | Technosol | 7.6 | 11 | 219 | 149 | 3.8 | | |
| Maan nor | ctrl | | 6.6 | 20 | 7 | 59 | 3.1 | | |
| iviean per | sh | | 7.6 | 13 | 80 | 88 | 2.5 | | |
| group | sp | | 7.6 | 26 | 154 | 126 | 5.1 | | |

| Texture (%) | Fe | AI | К | Zn | Pb | Cd | Cu | РАН |
|--|---|---|--|---|--|--|---|--|
| clay:silt:sand | | (g kg ⁻¹) | | | | (mg kg⁻¹) | | |
| 49:46:5 26:69:5 12:14:74 5:18:77 10:21:69 13:28:59 7:12:81 18:53:29 11:73:16 | 44 27 18 452 108 193 66 93 304 | 67 52 45 20 25 40 40 40 21 | 14.6 18.6 28.6 0.6 5.4 9.2 5.8 5.4 1.5 | 144 248 60 74 1650 2540 314 29600 55400 | 44 93 31 23 475 653 303 34700 14400 | 0.7 0.3 0.2 0.1 4.5 3.3 1.1 152.0 17.6 | 16 13 10 9 146 144 94 148 186 | 3.5 0.1 0.9 114.7 1095.9 937.7 21.2 49.6 |
| 11:75:14 29:43:28 9:20:71 13:67:20 | 30 ^b 205 ^a 150 ^a | 55 ^a 31 ^b 27 ^b | 3.4 20.6 ^a 5.2 ^b 3.4 ^b | 119000 151 ^b 1144 ^b 68000 ^a | 39500 56 ^b 363 ^b 29533 ^a | 0.4 ^b 2.2 ^b 63.9 ^a | 42 13 98 125 | 6.6 1.5 ^b 537.3 ^a 25.8 ^a |

| | Soil site groups | 18S rRNA gene copy number (x 10 ⁹ g ⁻¹) | Observed number of OTUs | Chao1 index |
|----------|---------------------|--|----------------------------|--------------|
| Мо | ctrl | 4.8 ± 0.6 | 162 ± 15 | 215 ± 31 |
| He | ctrl | 45.0 ± 1.8 | 176 ± 10 | 281 ± 48 |
| Di | ctrl | 21.8 ± 1.2 | 155 ± 7 | 203 ± 15 |
| Uc | sh | 5.1 ± 0.3 | 103 ± 4 | 175 ± 7 |
| Те | sh | 34.2 ± 3.0 | 113 ± 2 | 147 ± 15 |
| NM | sh | 3.1 ± 0.3 | 150 ± 7 | 179 ± 20 |
| Но | sh | 11.8 ± 0.2 | 190 ± 9 | 257 ± 6 |
| Ро | sp | 18.1 ± 1.6 | 196 ± 8 | 317 ± 22 |
| MsM | sp | 10.3 ± 0.9 | 148 ± 7 | 186 ± 9 |
| RM | sp | 8.5 ± 0.7 | 101 ± 5 | 158 ± 15 |
| | ctrl | 23.8 + 5.9 | 164 + 6 | 233 + 21 |
| Mean per | sh | 13.5 + 3.8 | 139 + 11 | 190 + 14 |
| group | sn | 12.3 ± 1.6 | 133 ± 11 148 ± 14 | 220 ± 26 |
| | - - | | | |

| Shannon's diversity index <i>H'</i> | Pielou's evenness index J' |
|---|---|
| 3.40 ± 0.02 | 0.669 ± 0.009 |
| 3.08 ± 0.04 | 0.596 ± 0.018 |
| 2.77 ± 0.09 | 0.549 ± 0.014 |
| 1.25 ± 0.05 | 0.271 ± 0.011 |
| 2.08 ± 0.06 | 0.440 ± 0.011 |
| 2.92 ± 0.03 | 0.581 ± 0.002 |
| 2.75 ± 0.01 | 0.529 ± 0.005 |
| 3.14 ± 0.02 | 0.593 ± 0.006 |
| 3.11 ± 0.02 | 0.623 ± 0.002 |
| 1.46 ± 0.05 | 0.313 ± 0.007 |
| 3.08 ± 0.10^{a} 2.25 ± 0.20^{b} 2.57 ± 0.28^{a} | 0.605 ± 0.019^{a} 0.455 ± 0.036^{b} 0.510 ± 0.049^{a} |