



D'hier à aujourd'hui, faire face aux pandémies

Florence Hammer, Patricia Licznar-Fajardo

► To cite this version:

Florence Hammer, Patricia Licznar-Fajardo. D'hier à aujourd'hui, faire face aux pandémies. *Actualités Pharmaceutiques*, 2020, 59 (599), pp.14-17. 10.1016/j.actpha.2020.08.004 . hal-03018311

HAL Id: hal-03018311

<https://hal.science/hal-03018311>

Submitted on 24 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Integrative role of plant mitochondria facing oxidative stress: the case of ozone

This paper is published in memory of Dr Claude Lance, former professor at the University Pierre et Marie Curie Paris 6

Anthony Gandin^a, Pierre Dizengremel^{a,*} and Yves Jolivet^a

^a Université de Lorraine, AgroParisTech, INRAE, Silva, F-54000 Nancy, France

*Corresponding author. E-mail address: pierre.dizengremel@univ-lorraine.fr

Abstract

Ozone is a secondary air pollutant, which causes oxidative stress in plants by producing reactive oxygen species (ROS) starting by an external attack of leaf apoplast. ROS have a dual role, acting as signaling molecules, regulating different physiological processes and response to stress, but also inducing oxidative damage. The production of ROS in plant cells is compartmented and regulated by scavengers and specific enzyme pathways. Chronic doses of ozone are known to trigger an important increase of the respiratory process while decreasing photosynthesis. Mitochondria, which normally operate with usual levels of intracellular ROS, would have to play a prominent role to cope with an enhanced ozone-derived ROS production. It is thus needed to compile the available literature on the effects of ozone on mitochondria to precise their strategy facing oxidative stress. An overview of the mitochondrial fate in three steps is proposed, i) starting with the initial responses of the mitochondria for alleviating the overproduction of ROS by the enhancement of existing antioxidant metabolism and adjustments of the electron transport chain, ii) followed by the setting up of detoxifying processes through exchanges between mitochondria and the cell, and iii) ending by an accelerated senescence initiated by mitochondrial membrane permeability and leading to programmed cell death.

Keywords: Ozone, respiration, mitochondria, oxidative stress, ROS, time-course response steps

Abbreviations – AOX: alternative oxidase, APX: ascorbate peroxidase, ASA: ascorbate, DHA: dehydroascorbate, DHAR: dehydroascorbate reductase, GABA: γ -Aminobutyric acid, GR: glutathione reductase, GSH: glutathione, HAF: Halliwell-Asada-Foyer cycle, MDHAR: monodehydroascorbate reductase, mETC: mitochondrial electron transport chain, O₃: ozone, OAA: oxaloacetate, OG: 2-oxoglutarate, OPPP: oxidative pentose phosphate pathway, PCD: programmed cell death, PEP: phosphoenolpyruvate, PEPcase: phosphoenolpyruvate carboxylase, TCA: tricarboxylic acid, ROS: reactive oxygen species, SA: salicylic acid, SDH: succinate dehydrogenase, UCP: uncoupling protein.

1. Introduction

In photosynthetically active cells, both photosynthesis and respiration produce energy in the form of ATP via photophosphorylation and oxidative phosphorylation, respectively in chloroplasts and mitochondria. Both organelles share the provision of energy whose the balance is tightly regulated under stress conditions especially (Gardestrom and Igamberdiev, 2016). Tropospheric ozone (O_3) is a secondary air pollutant, mainly produced by the photochemical dissociation of nitrogen oxides in the presence of organic compounds issued from car exhaust and industrial fuel combustion. O_3 concentrations increased fourfold during the last century reaching yearly average values of around 50 to 60 ppb but with peak values commonly observed between 100 and 250 ppb (Renaut et al., 2009). These concentrations have harmful impact on crops and forests (Wittig et al., 2009; Ainsworth et al., 2012; Matyssek et al., 2012; Jolivet et al., 2016; Cailleret et al., 2018; Emberson et al., 2018; Franz et al., 2018; Grulke and Heath, 2019). Plants exposed to moderate O_3 concentrations during prolonged time (chronic O_3 exposure) commonly exhibit a reduction in photosynthesis and photorespiration and an increase in respiration (Reich, 1983; Dizengremel, 2001; Bohler et al., 2007; Jolivet et al., 2016). The cellular changes induced by chronic O_3 injury were associated to an accelerated senescence in cell leaves (Pell et al., 1997; Pleijel et al., 1997; Miller et al., 1999; Gielen et al., 2007; Heath, 2008; Yendrek et al., 2017; Emberson et al., 2018; Vollenweider et al., 2019), the chloroplasts being early degraded while mitochondria remain intact until rather late (Sutinen et al., 1990; Holopainen et al., 1996). This delayed response of mitochondria thus confers a prominent role to this organelle in the mechanism of cell response to O_3 injury.

Although occurring in normal conditions, the generation of reactive oxygen species (ROS) at the level of the mitochondrial electron transport chain (mETC) is strongly increased under chronic O_3 which induces oxidative stress (Baier et al., 2005). However, the increase in the respiratory process raises many questions even though it is usually associated to repair, detoxification and synthesis of defense compounds (Dizengremel, 2001; Ainsworth et al. 2012). Respiration not only provides energy to plant cells in the form of ATP and NAD(P)H but also contributes to the production of carbon skeletons and the maintenance of redox balancing. These three functions co-exist, could overlap and thus need to be coordinated to regulate plant respiration (O'Leary et al., 2019). A range of papers describe different aspects of the implication of the mitochondria in the O_3 -induced increase of respiration, but a coordinated synthesis of the results is lacking. This review deepens the role of mitochondria in the response to O_3 , highlighting its ability to catabolize carbon intermediates, balance reducing power and induce antioxidant systems. The time course of mitochondrial events under O_3 exposure, from a successful scavenging of ROS to the initiation of senescence, will be proposed. The integration of the mitochondrial behavior in a coordinated cellular response to O_3 will be discussed.

2. Ozone-increased carbohydrate breakdown in the cytosol to feed mitochondria

The increase in respiration observed in plants submitted to chronic O₃ exposure is associated to an increased activity of a set of enzymes belonging to the glycolytic pathway and the oxidative pentose phosphate pathway (OPPP)(Dizengremel et al., 1994; Sehmer et al. 1998; Dizengremel et al., 2009; Dghim et al., 2012). This increase provides additional reducing power in the form of NADH and/or NADPH, which are useful for detoxification mechanisms (Fig 1.) (Dizengremel, 2001; Dizengremel et al., 2008). It must also be emphasized that several metabolites branch off from glycolysis or/and OPPP (Plaxton and Podesta, 2006) to initiate biosynthetic pathways (Fig 1.). Thus, phosphoenolpyruvate (PEP), end-product of the glycolysis, has a central role under O₃ stress (Dizengremel et al., 2012), and can bind to erythrose-4-phosphate from OPPP, allowing an increased functioning of the phenylpropanoid pathway essential to the production of defense compounds. Furthermore, the carboxylation of PEP into oxaloacetate (OAA) via phosphoenolpyruvate carboxylase (PEPcase) could in part mitigate the loss of CO₂ fixation by Rubisco under O₃ (Fontaine et al., 1999; Renaut et al., 2009). Furthermore, an increase of pyruvate kinase activity, providing more pyruvate to the tricarboxylic acid (TCA) cycle, was also observed under O₃ exposure (Dizengremel et al., 1994, 2009). Another role of PEP will be however emphasized here in the context of an increased mitochondrial respiration under O₃ stress. OAA produced from PEPcase can be transported into the mitochondrion or produce malate in the cytosol (Fig 1.). Malate can thus either produce pyruvate through the O₃-increased activity of the cytosolic NADP-malic enzyme (Dghim et al., 2012; Chen et al., 2019), providing additional NADPH for detoxification purposes, or be transported into the mitochondrion. It cannot be excluded that malate could thus be decarboxylated to pyruvate within the mitochondrion via NAD(P)-malic enzyme (Møller and Rasmusson, 1998; Dizengremel et al., 2012). An increasing offer in OAA, malate and/or pyruvate, would then replenish the TCA cycle (Dizengremel et al., 2012).

3. Deciphering the role of mitochondria under ozone stress

During the first thirty years following the observation of damaging effects of O₃ on plants (Haagen-Smit et al., 1952), few studies were devoted to the effect of O₃ specifically on plant mitochondria (Dizengremel and Citerne, 1988). The concentration of O₃ used was often acute (1 ppm delivered for several hours) and the resulting effect was mainly an inhibition of the oxidative phosphorylation (Lee, 1967). More recent works using chronic O₃ exposure explored the responses of the different mitochondrial pathways contributing to the oxidative degradation of the organic acids.

3.1. A noncyclic anabolic function of TCA to enhance antioxidant defense

The TCA cycle operates in the matrix of mitochondria by fully decarboxylating the organic acids; however, a noncyclic functioning of TCA can also occur (Tcherkez et al., 2009; Sweetlove et al., 2010) allowing the leak of some organic acids (citrate, isocitrate, oxoglutarate) towards amino acid

synthesis (Fig. 1A, O’Leary et al., 2011). This anaplerotic pathway would be solicited under chronic O₃ exposure (Dizengremel et al., 2012). Citrate is exported from the mitochondrion to the cytosol to produce 2-oxoglutarate (OG) via cytosolic isocitrate dehydrogenase (IDH), a reaction that provides NADPH and is increased under O₃ stress (Dghim et al., 2012). Mitochondrial OG can also be directly exported after being produced from isocitrate by the mitochondrial NAD(P)-IDH. OG may thus be integrated into the glutamine synthetase 2 / glutamate oxoglutarate aminotransferase or into the glutamine synthetase 1 (GS1) / aspartate aminotransferase cycles to give glutamate in the chloroplast or cytosol, respectively (Fig 1A). The activity of GS1 was shown to increase in plants fumigated with O₃ (Galant et al., 2012). This could allow to cope with a diminished NH₄⁺ remobilisation due to O₃-decreased photorespiratory activity (Bagard et al., 2008). Interestingly, a higher supply of cytosolic glutamate could also support *de novo* glutathione (GSH) synthesis by γ -glutamylcysteine synthetase and glutathione synthetase and therefore contributes to cell ROS detoxification. In animal cells, a specific OG transporter has been described in the mitochondrial inner membrane (Lash, 2006). This transporter carries out OG to the cytosol in exchange for cytosolic GSH (Fig. 1A). The presence of such a transporter still needs to be investigated in plants in which GSH biosynthesis only occurs in cytosol and chloroplast. Such transporters would therefore insure mitochondrial import of cytosolic GSH during stress. The enhancement of noncyclic TCA and anaplerotic pathway under O₃ would thus feed cytosolic antioxidant synthesis in carbon skeletons and simultaneously import antioxidants in mitochondria.

O₃ enters the leaf through stomata and rapidly generate various ROS (Kangasjarvi et al., 1995) that cause the peroxidation of membrane lipids (Ranieri et al., 1996; Loreto and Velikova, 2001). In this respect, the leak of compounds out of TCA cycle may result from inactivation of some downstream enzymes of the cycle by the lipid peroxidation products, as 4-hydroxy-2-nonenal, which has been shown to specifically inhibit oxoglutarate dehydrogenase and pyruvate dehydrogenase complexes by about 80% (Millar and Leaver, 2000). To maintain the rate of respiration, part of the TCA cycle can be by-passed, as do the γ -aminobutyrate shunt (GABA shunt) for succinate synthesis in bypassing oxoglutarate dehydrogenase and succinyl-coA-synthetase (Fig. 1B, Rhoads et al., 2006; Dizengremel et al., 2012). GABA synthesis is catalyzed by glutamate dehydrogenase and glutamic acid decarboxylase, which both increased in O₃-treated rice leaves (Cho et al., 2008). Increased concentrations of GABA were generally observed in response to stress, conferring on the GABA shunt an adaptive mechanism (Michaeli and Fromm, 2015). Recently, Che-Othman et al. (2019) concluded that increase in GABA shunt activity in wheat leaves overcome the inhibition of the TCA cycle key enzymes, providing an alternative carbon source to support the stress-dependent increase in mitochondrial respiration. In response to O₃, GABA could also act as a transmembrane signal indicating changes in TCA cycle activity from cell to cell (Ramesh et al., 2015; Gilliham and Tyerman, 2016).

The exchange of metabolites between chloroplast and mitochondria allows balancing the cellular redox within plant cell (Hoefnagel et al., 1998; Gardestrom and Igamberdiev, 2016; O’Leary et al., 2019). This is mediated via a system, often called malate valve, which couples a malate/OAA redox shuttle based on mitochondrial NAD-malate dehydrogenase to a chloroplastic NADP-malate dehydrogenase (Hoefnagel et al., 1998; Scheibe, 2004). When the NADPH/NADP ratio is high, malate is transported from the chloroplast into the mitochondria where it is transformed in OAA, resulting in the transfer of chloroplast NADPH in excess to mitochondrial NADH (Fig 1.)(Scheibe, 2004, Yoshida et al., 2007). This system allows to minimize chloroplast endogenous ROS production due to an overly reduced photosynthetic reaction system and a lack of NADP (Finnegan et al., 2004). This could happen under O₃ stress when a transient stromal over-reduction due to damaged Calvin cycle occurs before any impairment of the light reactions (Dizengremel, 2001; Dizengremel et al., 2009). In addition, O₃ is well known to induce stomatal closure (Wittig et al., 2007; Hoshika et al., 2015) that would also drive a NADPH accumulation in chloroplast by reducing CO₂ levels. Malate valve contributes to import reducing power in mitochondria and therefore to detoxification capacity in supporting mitochondrial antioxidant regeneration and to minimize chloroplastic ETC imbalance resulting in ROS generation.

3.2. Role of the mitochondrial electron transport chain

The mETC is a major site of ATP synthesis, but is also a source of ROS (Fig. 1C) through complex I (NADH: ubiquinone oxidoreductase) and complex III (UQH₂: cytochrome c oxidoreductase; cytochrome bc1 complex) (Murphy, 2009). More recently, it was shown that complex II (SDH) can also produce O₂⁻ at the FAD-binding site through the monovalent electron reduction of O₂ (Jardim-Messeder et al., 2015; Huang et al, 2019). Endogenous ROS production by mETC can however be mitigated by the use of alternative pathways. The plant mETC possesses an alternative non-proton pumping pathway, by-passing complexes III and IV, and mediated by alternative oxidase (AOX) (Fig 1C)(Lance et al., 1985; Vanlerberghe and McIntosh, 1997; Millar et al., 2011). AOX might act to maintain a basal ubiquinone pool reduction state, preventing over-reduction of upstream electron-transport components and lowering ROS production (Purvis and Shewfelt, 1993; Rhoads et al., 2006 ; Mittler, 2002; Umbach et al., 2005; Saha et al., 2016). It has been stated that the participation of the alternative oxidase is maintained very low under standard conditions by repressing its gene expression, potentially because of its wasteful nature (Selinski et al. 2018). However, the activity of AOX can be enhanced by several parameters acting at different levels (transcriptional, translational, post-translational) in response to a large range of stressors. The AOX gene expression was shown to be increased by H₂O₂ itself (Vanlerberghe & McIntosh, 1997) or through specific transcription factors (e.g. NAC17; Ng et al., 2013). Furthermore, differential expression of AOX isogenes can lead to the production of heterodimers having varying catalytic properties which potentially offer a fine-tune

mechanism of AOX activity in response to stress. Under O₃ exposure, an activation of AOX at both transcriptional and translational levels was observed in poplar, tobacco and *Arabidopsis* (Jolivet et al., 1997; Dizengremel, 2001, Ederli et al., 2006; Tosti et al., 2006). In tobacco, the O₃ induced activation of gene expression of AOX1a was suggested to be NO-dependent, in cooperation with ethylene signaling, and correlated to an inhibition of cytochrome oxidase by NO (Ederli et al., 2006, Kumari et al., 2019). This is a good example of a disfunctioning of mETC triggering the expression of a nuclear gene encoding a mitochondrial protein (retrograde signaling system, Ng et al., 2014).

AOX activity can also be regulated by redox state. The reduced form of the AOX dimer is more active than the oxidized form, suggesting that AOX activity can be modulated depending on reducing power availability, but may be affected by excessive accumulation of ROS. Thioredoxins belonging to Trxo and Trxh families are able to render AOX operational by reducing disulfide bonds (Trxo family: Marti et al., 2009; Umekawa and Ito, 2019; Trxh family: Gelhaye et al., 2004). The redox state activation of AOX protein is also coupled to ketocarboxylic acids binding that enhance its activity: pyruvate and OG are very effective activators in lowering Km for UQ substrate (Umbach et al., 2002; Selinski et al., 2018). These two systems of AOX activation, redox state and metabolite binding, interact with each other to complete AOX activity regulation. The increased availability of pyruvate under O₃ (Dizengremel et al., 1994, 2009) could explain the enhancing participation of AOX to the mitochondrial electron flow; however this activation would require sufficient reducing power availability to maintain AOX in its reduced form and minimize endogenous ROS production by mETC. In addition, AOX, that has a lower O₂ affinity than cytochrome oxidase (Km of 1-2.5μm vs. 0.1-0.2μm), allows to maintain a low level of O₂ in mitochondria that therefore contributes to minimize ROS production (Gupta et al. 2015).

Alternative pathways of mETC also include the shunt of complexes I and II, implying one of the four alternative non-proton pumping NAD(P)H dehydrogenases located either on the outer (NDex) or inner (NDin) surface of the inner mitochondrial membrane (Fig. 1C). The NDin might largely participate for the reoxidation of matrix NADH under conditions where its concentration is increased (Møller, 2001), which could be the case under O₃ stress as mentioned above. Matrix NAD(P)H can be produced via the NAD(P)-IDH and NAD-malic enzyme or imported through the malate/OAA shuttle. The activity of the NAD(P)-malic enzyme is increased under O₃ (Dizengremel et al., 2009; Dghim et al., 2012. NDin is thus probably competing with NAD(P)H-dependent ROS-detoxifying enzymes for matrix NAD(P)H (Møller, 2001). Although rather inactive in normal conditions, the Ca²⁺-dependent NDex linked to the outer face of the inner mitochondrial membrane could be activated under stress by an increase in Ca²⁺ cytosolic concentration, as it has been previously shown under O₃ (Heath and Castillo, 1988; Clayton et al., 1999). In addition, an increased expression of the NDex was demonstrated to accompany increased AOX expression under several stress treatments (Rhoads et al., 2006). Under

oxidative stress, alternative dehydrogenases could help decrease ROS production linked to a high reduction level of mETC, by keeping the cytosolic NAD(P) pool relatively oxidized (Rhoads et al., 2006). In addition, uncoupling proteins (UCP) cross the inner mitochondrial membrane (Vercesi et al., 2006) and mildly participate to the modulation of the transmembrane potential (Fig 1C)(Rhoads et al., 2006). UCP is also able to decrease ROS formation in reintegrating proton and thus bypassing respiratory control by ATP synthase (Rhoads et al., 2006). While AOX can be inactivated by a strong oxidative stress (Winger et al., 2005), UCP is able to keep operating when ROS level is much higher (Considine et al., 2003; Rhoads et al., 2006). The role of the NADH dehydrogenases and UCP under O₃-driven oxidative stress must however be confirmed, especially in regard to the role of AOX.

Beside electron transport, mETC also hosts the final step of ascorbate (ASA) biosynthesis that converts L-galactono-1,4-lactone to ASA and is catalysed by L-galactono-1,4-lactone dehydrogenase (Fig 1C)(Ôba et al., 1995; Bartoli et al., 2000). This enzyme is located on the inner mitochondrial membrane where it is physically associated to complex I and regulated by electron transport. Furthermore, oxidized cytochrome c is essential to the functioning of L-GLDH in acting as a specific electron acceptor (Fig 1C)(Bartoli et al., 2000). As an example, L-GLDH activity was decreased by 40% in Arabidopsis knockdown mutants under-expressing cytochrome c (Welchen et al. 2016). The question remains as the relative contribution of ASA synthesis to respiratory electron flux compared to that driven by TCA cycle activity and whether this contribution is modulated in O₃ stress conditions. In any case, produced ASA can thereafter be used by cytosolic and mitochondrial antioxidant systems to minimize ROS accumulation. Therefore, the coupling of L-GLDH and mETC activities via cytochrome c could contribute to couple respiration and antioxidant defense to cope with oxidative stress.

4. Antioxidant systems present in mitochondria and known effects of ozone

Several antioxidant mechanisms, mainly present in the soluble parts of the mitochondria (i.e. intermembrane space and matrix), can contribute to mitigate the increased ROS production during O₃ exposure. A matrix-localized Mn-superoxide dismutase was reported in literature, coping with endogenous O₂⁻ production by mETC, and involved in the protection against ROS produced by O₃ attack (Fig 1C)(Alscher et al., 2002). To manage superoxide dismutase products, the occurrence of a Halliwell-Asada-Foyer cycle (HAF, Fig. 1D) has been reported in cytosol, chloroplasts, peroxisomes as well as in mitochondria (Jiménez et al., 1997; Noctor and Foyer, 1998; Foyer and Noctor, 2011) even though the presence of dehydroascorbate reductase (DHAR) in mitochondria remains questionable (Møller, 2001; Noctor et al., 2018). Several ascorbate peroxidases (APX) have been identified in plant mitochondria, one intermembrane APX bound to the inner mitochondrial membrane (De Leonardis et al., 2000) and two others located in the matrix, either soluble or bound to the inner membrane (Mittova et al., 2004). As mentioned above, ASA is delivered to the mitochondrial

intermembrane space where it can be oxidized by the intermembrane APX. Dehydroascorbate (DHA) can then be transported into the matrix via a GLUT-like transporter of the glucose transporter family (Chew et al., 2003; Szarka et al., 2013). In addition, if the lack of DHAR in the matrix were confirmed, DHA will have to be reduced in ASA by the glutaredoxin or directly by GSH (Fig 1D)(Ehrhart and Zeevalk, 2003). Recent work also demonstrated the presence of mitochondrial ascorbate transporter-MAT, a member of the SLC25 family, to directly transport ASA through the mitochondrial inner membrane (Scalera et al., 2018).

The synthesis of GSH only occurs in cytosol and chloroplast (Rausch et al., 2007); however GSH presence has been reported in mitochondria suggesting a crossing of the inner mitochondrial membrane via a still unknown transporter in plants (Fig. 1A; Chen and Lash, 1998; Maughan and Foyer, 2006). The presence of glutathione reductase (GR) was confirmed in mitochondria to allow GSH regeneration where its activity is NADPH dependent (Fig. 1D, Edwards et al., 1990; Jimenez et al., 1997). Under O₃, the synthesis of GSH is increased likely to maintain glutathione redox state (Dumont et al., 2014) and compensates ROS scavenging by glutathione. Similarly, the activities of APX, monodehydroascorbate reductase (MDHAR) and GR have been shown to increase under O₃ (Dusart et al., 2019; Gandin et al., 2019).

As previously mentioned, thioredoxins have been reported in plant mitochondria (Sweetlove and Foyer, 2004; Navrot et al., 2007; Vieira Dos Santos and Rey, 2008). Also, a type II peroxiredoxin (PrxII F), has been identified in plant mitochondria (Rhoads et al., 2006; Finkemeier et., 2005), where it could reduce mitochondrial H₂O₂ by using thioredoxin or GSH (Fig 1D). The latter are then regenerated by thioredoxin reductase or GR (Sweetlove et al., 2004; Rhoads et al., 2006). The type II peroxiredoxin/thioredoxin system could also reduce lipid peroxides (Rouhier et al., 2004) and could therefore contribute to maintain mitochondria membrane integrity under stress. A mitochondrial protein disulfide isomerase functioning together with thioredoxin and NADPH could also reduce disulfides resulting from protein oxidation (Sweetlove et al., 2002; Rhoads et al., 2006). In addition to the above-mentioned effect of thioredoxin on AOX, the association of all these processes would lower the content of ROS in the matrix in O₃ exposed plants.

5. Suggested steps of the response of mitochondria to ozone: from resistance to cell death

O₃ being a powerful oxidant, a plausible scenario of the successive mechanisms coping with the progressively rising intensity of the O₃-related stress is proposed below (Table 1). The sequential presentation is probably an idealized view of more intricate relationships between the responses to O₃. In addition, ROS, responsible for oxidative damages caused by O₃, are also known to play an important role in triggering signaling pathways. Thus, there is a permanent fine-tuning of ROS accumulation to balance beneficial signaling and phytotoxic effect. The steps described below lie on

results obtained from a series of O₃ experiments and, in a minor way, are complement in integrating probable events derived from general oxidative stress.

5.1. Mobilisation of pre-existing mechanisms to cope with ROS

The first impact of O₃ on photosynthesis, and especially on the biochemical components, leads to excessive NADPH accumulation within the chloroplast. A series of metabolic pathways, generally associated to carbohydrate breakdown as glycolysis and OPP, are also enhanced and result in even more NAD(P)H produced (Fig1, Table 1). This additional reducing power allows the existing detoxification processes to function at optimal level. An increase in APX activity has been described under O₃ and shown to be supported by enhanced ASA regeneration by NADH-dependent MDHAR (Gandin et al., 2019). In addition, the cytosolic carboxylating enzyme, PEPcase, is requested, either to cope with the decreased CO₂ fixation by Rubisco and/or to contribute to the anaplerotic pathway (Dizengremel et al., 2012). Several intermediates of catabolism are therefore favoured, either by pyruvate kinase (e.g. pyruvate) or/and by PEPcase (e.g. OAA and malate), and have impact on mitochondria functioning (Dghim et al., 2012). These organic acids will be transported into the mitochondria to feed TCA cycle and finally provide NAD(P)H to the mETC. Additional reducing power is also provided by the transfer of redox equivalents (e.g. through malate/OAA shuttle) either from chloroplasts, in which the redox pressure is transiently high, or from cytosol. However, increasing feeding of mitochondria with NAD(P)H threaten to over-reduce mETC and induce endogenous ROS production.

To maintain the mETC in a sufficiently oxidized state, ATP synthesis and respiratory control could be balanced by the UCP while the electron flow may use the activated AOX and some of the four NAD(P)H dehydrogenases located on both sides of the inner mitochondrial membrane (Table 1). Three NAD(P)H-DH are activated by Ca²⁺ which concentration is known to be increased by O₃ and is involved in phytohormone cascade signals (Short et al., 2012). An increase of the amount of salicylic acid (SA) was shown under O₃ fumigation (Koch et al., 2000). At low concentration, SA was shown to uncouple the mETC (Gleason et al., 2011) and to increase AOX expression (Rhoads and McIntosh, 1992), suggesting a potential role for SA (and may be ethylene) to the signaling process regulating mitochondrial electron flux (Rao et al., 2000; Dizengremel, 2001; Heath, 2008). NO production due to O₃ stress could also be an important regulator of electron partitioning between cytochrome and alternative pathways, in reducing and enhancing cytochrome oxidase and alternative oxidase activities, respectively (Ederli et al. 2006). Therefore, NO impact on mETC could initiate a bottom-up regulation starting with the decrease in mETC ATP yield that would require an increase in glycolysis and TCA cycle activities to fulfill mitochondria ATP demands. Evidence may be seen in the stimulation by O₃

of a mitochondrial phosphate transporter located in the inner membrane, which catalyzes the influx of Pi, essential for the synthesis of ATP in the matrix (Kiiskinen et al., 1997).

5.2. Setting up of detoxifying processes

The enhanced activities of PEPcase as well as of glycolysis and OPP tend to show a plateau for higher doses of O₃, announcing a limit for the capacity of the metabolism to counteract the oxidative stress (Dizengremel et al., 1994; Fontaine et al., 1999; Renaut et al., 2009). With an extended O₃ exposure, the ROS production keeps increasing and more likely starts to accumulate in plant cells. ROS shortly oxidized cell components (e.g. protein, DNA...) among which membrane lipids that are a major target (Rebouças et al., 2017). Lipid peroxidation products are able to inhibit specific TCA cycle enzymes (Millar and Leaver, 2000) and disrupt its functioning (Table 1). As a consequence, TCA cycle would leak intermediates and export isocitrate or OG to the cytosol for amino acid biosynthesis (Dizengremel et al., 2012). This leak in carbon skeletons would also support cytosolic GSH *de novo* biosynthesis, as the enzymes for it have never been reported in mitochondria. GSH can then return to mitochondria through OG-GSH transporters as shown in animal cells (Lash 2006), yet to be described in plants. GSH can therefore support glutathione peroxidase and glutaredoxin activities as well as ASA regeneration through non-catalytic reduction or mediated by DHAR, therefore significantly contributing to antioxidant defense (Fig 1D, Table 1). With ROS accumulation, other antioxidant systems can intervene, as the peroxiredoxin/thioredoxin systems present in mitochondria that could reduce mitochondrial H₂O₂. In addition, it is likely that ASA biosynthesis is also enhanced under O₃ to support HAF functioning. The *de novo* synthesis of ASA however involves additional feeding of the mETC with electrons, increasing even more the reductant pressure on the chain and the risk for endogenous ROS production. In addition, ROS accumulation likely oxidized mETC components as AOX, which is inactive in oxidized form. This runaway effect may lead to increasing endogenous O₂⁻ production by complexes I, II and III and ROS over-accumulation in mitochondria.

Glutamate issued from OG export from mitochondria can also enter the GABA shunt that leads to the production of succinate in mitochondria, bypassing TCA cycle enzymes inactivated by oxidative stress (Fig 1B)(Table 1)(i.e. OG dehydrogenase and succinyl-CoA synthetase). GABA-shunt issued succinate can then be reoxidized by Complex II in mETC that still drives a strong electron flow from the complex I and NAD(P)H-DH. However, complex II can be inhibited at high concentrations of oxidative stress-induced phytohormones as SA (Gleason et al., 2011; Belt et al., 2017). This would lead to the accumulation of GABA shunt intermediates, as succinic semialdehyde that is toxic compound for plants.

5.3. Towards programmed cell death

When the O₃ uptake lasts and oxidative pressure keeps rising, the antioxidant defense is overwhelmed and ROS accumulates in excess. It is admitted that plant mitochondria could then play a role similar to that of animal mitochondria in the PCD process (Table 1)(Xie and Chen, 2000; Tiwari et al., 2002; Vianello et al., 2007). The decrease in ATP production and release of cytochrome c are the first signals of PCD. The depletion of ATP synthesis, above described as accompanying both UCP and AOX implication, can be accelerated by mitochondrial membrane disturbances associated to ROS. The peroxiredoxin/thioredoxin system can help by supplying reducing power to reductases acting on lipid hydroperoxides and disulfides resulting from protein oxidation, but it become insufficient. A collapse of mitochondrial transmembrane potential $\Delta\Psi_m$ would ensue, triggering the opening of the mitochondrial permeability transition pore (Table 1)(Tiwari et al., 2002; Zandalinas and Mittler, 2018).

Cytochrome *c*, a mobile protein attached to the cytosolic face of the inner mitochondrial membrane, acts as an electron carrier between complexes III and IV and is implied in the last step of ASA synthesis. Cytochrome *c* is labile and can be easily detached from the inner mitochondrial membrane followed by its release into the cytosol through a permeabilization of the outer membrane due to oxidative stress (Table 1)(Dizengremel, 1983; Sun et al., 1999). A release of cytochrome *c* was observed upon O₃ exposure of tobacco plants (Pasqualini et al., 2003). The release of cytochrome *c* activates caspases-like proteases that are involved in the execution of the cell death program (Tiwari et al., 2002; Vacca et al., 2006; Welchen and Gonzalez, 2016). A Ca²⁺ overload, admitted to occur upon O₃ exposure, would trigger the entire PCD process (Xiong et al., 2006; Welchen and Gonzalez, 2016). The increased permeability of mitochondria to solutes leads to mitochondrial swelling (Vianello et al., 2007; Zancani et al., 2015) which was early observed in response to O₃ (Lee, 1967, 1968). This ends by overall mitochondria rupture and subsequent cell death.

6. Conclusions and perspectives

In the literature, the role of mitochondria facing O₃ stress is often mentioned as being important but the available information is disseminated. A gathering of these information and a clearer picture of the sequential events following an increasing oxidative pressure is proposed in this review article, based on results mostly obtained with O₃ itself and some others deduced from general oxidative stress (Table 1). As soon as O₃ enters the leaf, delivering ROS, the entire catabolic system is mobilized, partly to compensate the damages and also for providing redox compounds, organic acids and secondary metabolites. These changes allow to optimize detoxification activity and minimize ROS accumulation, in modulating TCA cycle activity and NAD(P)H production, balancing electron transport within mETC, and increasing antioxidant pool in the matrix. In the most severe cases, this wide

reorganization of mitochondria metabolism can be insufficient and mitochondria will be overwhelmed by oxidative pressure, leading its disruption through a PCD-like process.

Even though this scenario is plausible, the action of O₃ on some of the steps needs to be definitively confirmed. The important role of AOX (and potentially other alternative pathways) must be emphasized. AOX has been recently proposed as a functional marker for breeding stress tolerant plant varieties (Polidoros et al., 2009). On the other hand, an elaborate network of anterograde and retrograde signalling between the mitochondria and nucleus is at work in plant cells with redox status and ROS as important actors (Leister, 2005; Rhoads and Subbaiah, 2007; Ng et al, 2014; Farooq et al., 2019). Moreover, it was recently demonstrated that the Ca²⁺-dependent changes in gene expression observed in response to O₃ differ markedly from Ca²⁺ signatures of other oxidative stress, such as H₂O₂ and cold (Short et al., 2012). A more deeply insight concerning the metabolic and signalling networks under O₃ stress thus needs to be studied. In this context, the ambiguous role of hormones raises questions, particularly for ethylene, which is largely implicated in leaf senescence as well as O₃ damage (Moeder et al., 2002; Nunn et al., 2005; Heath, 2008). Finally, the role of antioxidants to cope with O₃ stress is undisputable but very technically complex to assess relative to each intracellular pool. It is therefore essential to develop new methods to estimate the relative contribution of each pool and their potential exchanges within the cell.

Conflicts of interest

None declared

Funding source

This research was supported by the French National Research Agency through the Laboratory of Excellence ARBRE (ANR-12-LABXARBRE-01).

Author Contribution

All authors contributed equally to the writing and editing of the manuscript.

References

- Ainsworth EA, Yendrek CR, Sitch S, Collins WJ, Emberson LD (2012). The effects of tropospheric ozone on net primary productivity and implications for climate change. *Annu Rev Plant Biol* 63: 637–661
- Alscher RG, Erturk N, Heath LS (2002) Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *J Exp Bot* 53: 1331-1341
- Bagard M, Delacote E, Le Thiec D, Hasenfratz-Sauder MP, Dizengremel P, Jolivet Y. (2008) Ozone-induced changes in photosynthesis and photorespiration of hybrid poplar in relation to the developmental stage of the leaves. *Physiol Plant* 134: 559-574
- Baier L, Kandlbinder A, Golldack D, Dietz KJ (2005) Oxidative stress and ozone: perception, signalling and response. *Plant Cell Environ* 28: 1012-1020
- Bartoli CG, Pastori GM, Foyer CH (2000) Ascorbate biosynthesis in mitochondria is linked to the electron transport chain between complexes III and IV. *Plant Physiol* 123: 335-343
- Belt K, Huang S, Thatcher LF, Casarotto H, Singh KB, van Aken O, Millar AH (2017) Salicylic acid-dependent plant stress signalling via mitochondrial succinate dehydrogenase. *Plant Physiol* 173: 2029-2040
- Bohler S, Bagard M, Oufir M, Planchon S, Hoffmann L, Jolivet Y, Hausman JF, Dizengremel P, Renaut J (2007) A DIGE analysis of developing poplar leaves subjected to ozone reveals major changes in carbon metabolism. *Proteomics* 7:1584–1599
- Cailleret M, Ferretti M, Gessler A, Rigling A, Schaub M (2018) Ozone effects on European forest growth – towards an integrative approach. *J Ecology* 106: 1377-1389
- Casolo V, Braidot E, Chiandussi E, Macri F, Vianello A (2000) The role of mild uncoupling and non-coupled respiration in the regulation of hydrogen peroxide generation by plant mitochondria. *FEBS Lett* 474:53–57
- Che-Othman MH, Jacoby RP, Millar AH, Taylor NL (2019) Wheat mitochondrial respiration shifts from the tricarboxylic acid cycle to the GABA shunt under salt stress. *New Phytol* doi.org/10.1111/nph.15713
- Chen Z, Lash LH (1998) Evidence for mitochondrial uptake of glutathione by dicarboxylate and 2-oxoglutarate carriers. *J Pharmacol Exp Ther* 285: 608-618
- Chen Q, Wang B, Ding H, Zhang J, Li S (2019) The role of NADP-malic enzyme in plants under stress. *Plant Sci* 281: 206-212
- Chew O, Whelan J, Millar AH (2003) Molecular definition of the ascorbate-glutathione cycle in *Arabidopsis* mitochondria reveals dual targeting of antioxidant defenses in plants. *J Biol Chem* 278: 46869-46877
- Cho K, Shibato J, Agrawal GK, Jung YH, Kubo A, Jwa NS, Tamogami S, Satoh K, Kikuchi S, Higashi T et al (2008) Integrated transcriptomics, proteomics, and metabolomics analyses to survey ozone responses in the leaves of rice seedlings. *J Proteome Res* 7: 2980-2998

466 Chrobok D, Law SR, Brouwer B, Linden P, Ziolkowska A, Liebsch D, Narsai R, Szal B, Moritz T,
467 Rouhier N, Whelan J, Gardeström P, Keech O (2016) Dissecting the metabolic role of
468 mitochondria during developmental leaf senescence. *Plant Physiol* 172: 2132-2153

469 Clayton H, Knight MR, Knight H, McAinsh MR, Hetherington AM (1999) Dissection of the ozone-
470 induced calcium signature. *Plant J* 17: 575-579

471 Considine MJ, Goodman M, Echtay KS, Laloi M, Whelan J, Brand MD, Sweetlove LJ (2003)
472 Superoxide stimulates a proton leak in potato mitochondria that is related to the activity of
473 uncoupling protein. *J Biol Chem* 278: 22298-22302

474 De Leonardis S, Dipierro N, Dipierro S (2000) Purification and characterization of an ascorbate
475 peroxidase from potato tuber mitochondria. *Plant Physiol Biochem* 38: 773-779

476 Dghim AA, Dumont J, Hasenfratz-Sauder MP, Dizengremel P, Le Thiec D, Jolivet Y (2012) Capacity
477 for NADPH regeneration in the leaves of two poplar genotypes differing in ozone sensitivity.
478 *Physiol Plant* 148:36–50

479 Dizengremel P (1983) Effect of Triton X-100 on electron transport in plant mitochondria. *Physiol Vég*
480 21:731–741

481 Dizengremel P (2001) Effects of ozone on the carbon metabolism of forest trees. *Plant Physiol*
482 *Biochem* 39: 729–742

483 Dizengremel P, Citerne A (1988) Air pollutants effects on mitochondria and respiration. In Schulte-
484 Hostede S, Darrall NM, Blank LW and Wellburn AR (eds) *Air Pollution and Plant Metabolism*,
485 Elsevier, pp 169-188

486 Dizengremel P, Sasek TW, Brown KJ, Richardson CJ (1994) Ozone-induced changes in primary
487 carbon metabolism enzymes of loblolly pine needles. *J Plant Physiol* 144: 300-306

488 Dizengremel P, Le Thiec D, Bagard M, Jolivet Y (2008) Ozone risk assessment for plants: central role
489 of metabolism-dependent changes in reducing power. *Env Poll* 156: 11-15

490 Dizengremel P, Le Thiec D, Hasenfratz-Sauder MP, Vaultier MN, Bagard M, Jolivet Y (2009)
491 Metabolic-dependent changes in plant cell redox power after ozone exposure. *Plant Biol* 11: 35-42

492 Dizengremel P, Vaultier MN, Le Thiec D, Cabané M, Bagard M, Gérant D, Gérard J, Dghim AA,
493 Richet N, Afif D, Pireaux JC, Hasenfratz-Sauder MP, Jolivet Y (2012) Phosphoenolpyruvate is at
494 the crossroads of leaf metabolic responses to ozone stress. *New Phytol* 195: 512-517

495 Dumont J, Keski-Saari S, Keinanen M, Cohen D, Ningre N, Kontunen-Soppela S, Baldet P, Gibon Y,
496 Dizengremel P, Vaultier M-N, Jolivet Y, Oksanen E, Le Thiec D (2014) Ozone affects ascorbate
497 and glutathione biosynthesis as well as amino acid contents in three Euramerican poplar genotypes.
498 *Tree Physiol* 34: 253–266

499 Dusart N, Gérard J, Le Thiec D, Collignon C, Jolivet Y, Vaultier MN (2019) Integrated analysis of the
500 detoxification responses of two Euramerican poplar genotypes exposed to ozone and water deficit:
501 Focus on the ascorbate-glutathione cycle. *Sci Total Environ* 651: 2365–2379

502 Ederli L, Morettini R, Borgogni A, Wasternack C, Miersch O, Reale L, Ferranti F, Tosti N., Pasqualini
503 S. (2006) Interaction between nitric oxide and ethylene in the induction of alternative oxidase in
504 ozone-treated tobacco plants. *Plant Physiol* 142: 595–608

505 Edwards EA, Rawsthorne S, Mullineaux PM (1990) Subcellular distribution of multiple forms of
506 glutathione reductase in leaves of pea (*Pisum sativum* L.). *Planta* 180: 278– 284

507 Ehrhart J, Zeevalk GD (2003) Cooperative interaction between ascorbate and glutathione during
508 mitochondrial impairment in mesencephalic cultures. *J Neurochem* 86: 1487-1497

509 Emberson LD, Pleijel H, Ainsworth E, van den Berg M, Ren W, Osborne S, Mills G, Pandey D,
510 Dentener F, Büker P, Ewert F, Koeble R, van Dingenen (2018) Ozone effects on crops and
511 consideration in crop models. *Europ J Agr* 100: 19-34
512

513 Farooq MA, Niazi AK, Akhtar J , Farooq M, Souri Z, Karimi N, Rengel Z (2019) Acquiring control:
514 The evolution of ROS-Induced oxidative stress and redox signaling pathways in plant stress
515 responses. *Plant Physiol. and Biochem* 141: 353-369

516 Finnegan PM, Soole K, Umbach AL (2004) Alternative mitochondrial electron transport proteins in
517 higher plants. In Day D, Millar AH and Whelan J (eds) *Plant Mitochondria: From Genome to*
518 *Function, Advances in Photosynthesis and Respiration*, Kluwer Academic Press, Dordrecht, vol 17,
519 pp 163-230

520 Finkemeier I, Goodman M, Lamkemeyer P, Kandlbinder A, Sweetlove LJ, Dietz KJ, (2005) The
521 mitochondrial type II peroxiredoxin F is essential for redox homeostasis and root growth of
522 *Arabidopsis thaliana* under stress *J Biol Chem* 280: 12168-12180

523 Fontaine V, Pelloux J, Podor M, Afif D, Gérant D, Grieu P, Dizengremel P (1999) Carbon fixation in
524 *Pinus halepensis* submitted to ozone. Opposite response of ribulose-1,5-bisphosphate
525 carboxylase/oxygenase and phosphoenolpyruvate carboxylase. *Physiol Plant* 105 :187–192

526 Foyer CH, Noctor G (2011) Ascorbate and glutathione: the heart of the redox hub. *Plant Physiol.* 155:
527 2–18

528 Franz M, Alonso R, Arneth A, Büker P, Elvira S, Gerosa G, Emberson L, Feng Z, Le Thiec D,
529 Marzuoli R, Oksanen E, Uddling J, Wilkinson M, Zaehle S (2018) Evaluation of simulated ozone
530 effects in forest ecosystems against biomass damage estimates from fumigation experiments
531 *Biogeosciences* 15: 6941–6957

532 Galant A, Koester RP, Ainsworth EA, Hicks LM, Jez JM (2012) From climate change to molecular
533 response: redox proteomics of ozone-induced responses in soybean. *New Phytologist* 194: 220-
534 229.

535 Gandin A, Davrinche A, Jolivet Y (2019) Deciphering the main determinants of O₃ tolerance in
536 Euramerican poplar genotypes. *Sci Total Environ* 656: 681-690.

537 Gardestrom P, Igamberdiev AU (2016) The origin of cytosolic ATP in photosynthetic cells. *Physiol*
538 *Plant* 157: 367-379

539 Gelhaye E, Rouhier N, Gérard J, Jolivet Y, Gualberto J, Navrot N, Ohlsson PI, Wingsle G, Hirasawa
540 M , Knaff_DB, Wang H, Dizengremel P, Meyer Y, Jacquot JP (2004) A specific form of

541 thioredoxin h occurs in plant mitochondria and regulates the alternative oxidase. PNAS 101:
542 14545-14550

543 Gielen B, Low M, Deckmyn G, Metzger U, Franck F, Heerdts C, Matyssek R, Valcke R, Ceulemans R
544 (2007) Chronic ozone exposure affects leaf senescence of adult beech trees: a chlorophyll
545 fluorescence approach. J Exp Bot 58: 785-795

546 Gilliam M, Tyerman SD (2016) Linking metabolism to membrane signaling: the GABA-malate
547 connection. Trends Plant Sci 21: 295-301

548 Gleason C, Huang S, Thatcher LF, Foley RC, Anderson CR, Carroll AJ, Millar H, Singh K (2011)
549 Mitochondrial complex II has a key role in mitochondrial-derived reactive oxygen species
550 influence on plant stress gene regulation and defense. PNAS 108:10768-10773

551 Grulke NE, Heath RL (2019) Ozone effects on plants in natural ecosystems. Plant Biol
552 doi.org/10.1111/plb.12971

553 Gupta DK, Palma JM Corpas FJ (2015). *Reactive oxygen species and oxidative damage in plants*
554 *under stress* (pp. 1-22). Heidelberg, Germany: Springer

555 Haagen-Smit AJ, Darley EF, Zaitlin M, Hull H, Noble W (1952) Investigation on injury to plants from
556 air pollution in the Los Angeles area. Plant Physiol 27:18–34

557 Heath RL (2008) Modification of the biochemical pathways of plants induced by ozone: What are the
558 varied routes to change? Env Pol 155: 453-463

559 Heath RL, Castillo FJ (1988) Membrane disturbances in response to air pollutants. In Schulte-Hostede
560 S, Darrall NM, Blank LW, Wellburn AR (eds) Air pollution and Plant Metabolism, Elsevier
561 Applied Science, London and New York, pp 55-75

562 Hoefnagel MHN, Atkin OK, Wiskich JT (1998) Interdependence between chloroplasts and
563 mitochondria in the light and the dark. Biochim Biophys Acta 1366: 235-255

564 Holopainen T, Anttonen S, Palomäki V, Kainulainen P, Holopainen JK (1996) Needle ultrastructure
565 and starch content in Scots pine and Norway spruce after ozone fumigation. Can J Bot 74: 67-76

566 Hoshika Y, Watanabe M, Inada N, Koike T (2015) Effects of ozone-induced stomatal closure on
567 ozone uptake and its changes due to leaf age in sun and shade leaves of Siebold's beech. J Agric
568 Meteo 71: 218-226

569 Huang S, Braun HP, Gavryluk RMR, Millar AH (2019) Mitochondrial complex II of plants: subunit
570 composition, assembly, and function in respiration and signaling. The Plant J doi:
571 10.1111/tpj.14227

572 Jardim-Messeder D, Caverzan A, Rauber R, de Souza Ferreira E, Margis-Pinheiro M, Galina A (2015)
573 Succinate dehydrogenase (mitochondrial complexII) is a source of reactive oxygen species in
574 plants and regulates development and stress responses. New Phytol 208: 776–789

575 Jimenez A, Hernandez JA, del Rio LA, Sevilla F (1997) Evidence for the presence of the ascorbate-
576 glutathione cycle in mitochondria and peroxisomes of pea leaves. Plant Physiol 114: 275-284

577 Jolivet Y, Gérant D, Silva Lima M, Gérardin C, Loirat X, Fernandes de Melo D, Dizengremel P
578 (1997) Effect of environmental stresses on plant mitochondrial respiration. International
579 Symposium “Stress of Life, Stress and Adaptation from Molecules to Man” Budapest, Hungary.
580 Abstract

581 Jolivet Y, Bagard M, Cabané M, Vaultier MN, Gandin A, Afif D, Dizengremel P, Le Thiec D (2016)
582 Deciphering the ozone-induced changes in cellular processes: a prerequisite for ozone risk
583 assessment at the tree and forest levels. *Ann For Sci* 73: 923-943

584 Kangasjärvi J, Talvinen J, Utriainen M, Karjalainen R (1994) Plant defense systems induced by ozone.
585 *Plant Cell Environ* 17: 783–794
586

587 Kiiskinen M, Korhonen M, Kangasjärvi J (1997) Isolation and characterization of cDNA for a plant
588 mitochondrial phosphate translocator (*Mpt1*): ozone stress induces *Mpt1* mRNA accumulation in
589 birch (*Betula pendula* Roth) *Plant Mol Biol* 35: 271-279

590 Koch JR, Creelman RA, Eshita SM, Seskar M, Mullet JE, Davis KR (2000) Ozone sensitivity in
591 hybrid poplar correlates with insensitivity to both salicylic acid and jasmonic acid. The role of
592 programmed cell death in lesion formation, *Plant Physiol* 123: 487–496

593 Kumari A, Pathak PK, Bulle M, Igamberdiev AU Gupta KJ (2019) Alternative oxidase is an important
594 player in the regulation of nitric oxide levels under normoxic and hypoxic conditions in plants. *J*
595 *Exp Bot* 70: 4345-4354

596 Lance C, Chauveau M, Dizengremel P (1985) The cyanide-resistant pathway of plant mitochondria.
597 In: Douce R and Day DA (eds) *Encyclopedia of Plant Physiology*, Springer-Verlag, Berlin, vol. 18,
598 pp 202-247

599 Lash LH (2006) Mitochondrial glutathione transport: physiological, pathological and toxicological
600 implications. *Chem Biol Interact* 163: 54-67

601 Laties GG (1982) The cyanide-resistant, alternative path in higher plant respiration. *Annu Rev Plant*
602 *Physiol* 33: 519-555

603 Lee TT (1967) Inhibition of oxidative phosphorylation and respiration by ozone in tobacco
604 mitochondria. *Plant Physiol* 42: 691-696

605 Lee TT (1968) Effect of ozone on swelling of tobacco mitochondria. *Plant Physiol* 43: 133-139

606 Leister D (2005) Genomics-based dissection of the cross-talk of chloroplasts with the nucleus and
607 mitochondria in *Arabidopsis*. *Gene* 354: 110-116

608 Marti MC, Olmos E, Calvete JJ, Diaz I, Barranco-Medina S, Whelan J, Lazaro JJ, Sevilla F, Jimenez
609 A (2009) Mitochondrial and nuclear localization of a novel pea thioredoxin: identification of its
610 mitochondrial target proteins. *Plant Phys* 150: 646-657

611 Matyssek R, Wieser G, Calfapietra C, de Vries W, Dizengremel P, Ernst D, Jolivet Y, Mikkelsen TT,
612 Mohren GMJ, Le Thiec D, Tuovinen JP, Weatherall A, Paoletti E (2012) Forests under climate
613 change and air pollution: gaps in understanding and future directions for research. *Environ Pollut*
614 160: 57–65

615 Maughan S, Foyer CH (2006) Engineering and genetic approaches to modulating the glutathione
616 network in plants. *Physiol Plant* 126:382–397

617 Maxwell DP, Wang Y, McIntosh L (1999) The alternative oxidase lowers mitochondrial reactive
618 oxygen production in plant cultures. *Proc Nat Acad Sci USA* 96: 8271-8276

619 Mehlhorn H, Tabner BJ, Wellburn A (1990) Electron spin resonance evidence for the formation of
620 free radicals in plants exposed to ozone. *Physiol Plant* 79: 377-383.

621 Michaeli S, Fromm H (2015) Closing the loop on the GABA shunt in plants: are GABA metabolism
622 and signaling entwined? *Front Plant Sci* 6: 419. doi: 10.3389/fpls.2015.00419

623 Millar AH, Leaver CJ (2000) The cytotoxic lipid peroxidation product, 4-hydroxy-2-nonenal,
624 specifically inhibits decarboxylating dehydrogenases in the matrix of plant mitochondria. *FEBS*
625 *Letters* 481: 117-121

626 Millar AH, Whelan J, Soole KL, Day DA 2011 Organization and regulation of mitochondrial
627 respiration in plants. *Annu Rev Plant Biol.* 62:79-104.

628 Miller JD, Arteca RN, Pell EJ (1999) Senescence-associated gene expression during ozone-induced
629 leaf senescence in *Arabidopsis*. *Plant Physiol* 120: 1015-1023

630 Mittler R (2002) Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* 7: 405–410

631 Mittova V, Theodoulou FL, Kiddle G, Volokita M, Tal M, Foyer C H Guy M (2004) Comparison of
632 mitochondrial ascorbate peroxidase in the cultivated tomato, *Lycopersicon esculentum*, and its
633 wild, salt-tolerant relative, *L. pennellii*—a role for matrix isoforms in protection against oxidative
634 damage. *Plant Cell Environ* 27: 237-250.

635 Moeder W, Barry CS, Tauriainen AA, Betz C, Tuomainen J, Utriainen M, Grierson D, Sandermann H.
636 Langebartels C, Kangasjarvi J (2002) Ethylene synthesis regulated by biphasic induction of 1-
637 aminocyclopropane-1-carboxylic acid synthase and 1-aminocyclopropane-1-carboxylic acid
638 oxidase genes is required for hydrogen peroxide accumulation and cell death in ozone-exposed
639 tomato. *Plant Physiol* 130: 1918-1926.

640 Møller IM (2001) Plant Mitochondria and oxidative stress: electron transport, NADPH turnover, and
641 metabolism of reactive oxygen species *Annu Rev Plant Physiol Plant Mol Biol* 52: 561-591

642 Møller IM, Rasmusson AG (1998) The role of NADP in the mitochondrial matrix. *Trends Plant Sci* 3:
643 21–27

644 Moller IM, Sweetlove LJ (2010) ROS-signalling specificity is required. *Trends Plant Sci* 15: 370-374

645 Murphy MP (2009) How mitochondria produce reactive oxygen species. *Biochem J* 417: 1-13

646 Navrot N, Rouhier N, Gelhaye E, Jacquot JP (2007) Reactive oxygen species generation and
647 antioxidant systems in plant mitochondria. *Physiol Plant* 129: 185-195

648 Ng S, de Clercq I, van Aken O, Law SR, Ivanova A, Willems P, Giraud E, van Breusegem F, Whelan
649 J (2014) Anterograde and retrograde regulation of nuclear genes encoding mitochondrial proteins
650 during growth, development, and stress. *Mol. Plant* 7: 1075-1093

651 Noctor G, Foyer CH (1998) Ascorbate and glutathione: keeping active oxygen under control. *Ann*
652 *Rev Plant Physiol Plant Mol Biol* 49: 249-279

653 Noctor G, Reichheld JP, Foyer CH (2018) ROS-related redox regulation and signaling in plants.
654 *Semin Cell Dev Biol* 80: 3–12

655 Nunn AJ, Anegg S, Betz G, Simons S, Kalisch G, Seidlitz HK, Grams TEE, Häberle KH, Matyssek R,
656 Bahnweg G, Sandermann H, Langebartels C (2005) Role of ethylene in the regulation of cell death
657 and leaf loss in ozone-exposed European beech. *Plant Cell Environ* 28: 886-897

658 Ôba K, Ishikawa S, Nishikawa M, Mizuno H, Yamamoto T (1995) Purification and properties of l-
659 galactono- γ -lactone dehydrogenase, a key enzyme for ascorbic acid biosynthesis, from sweet
660 potato roots. *J Biochem* 117:120–124

661 O’Leary B, Park J and Plaxton WC (2011) The remarkable diversity of plant PEPC
662 (phosphoenolpyruvate carboxylase): recent insights into the physiological functions and post-
663 translational controls of non-photosynthetic PEPCs. *Biochem J* 436: 15–34

664 O’Leary BM, Asao S, Millar AH, Atkin OK (2019) Core principles which explain variation in
665 respiration across biological scales. *New Phytol* 222: 670–686

666 Pasqualini S, Piccioni C, Reale L, Ederli L, Della Torre G, Ferrant F (2003) Ozone-induced cell death
667 in tobacco cultivar Bel W3 plants. The role of programmed cell death in lesion formation. *Plant*
668 *Physiol* 133: 1122-1134

669 Pell EJ, Schlagnhauser CD, Arteca RN (1997) Ozone-induced oxidative stress: Mechanisms of action
670 and reaction. *Physiol Plant* 100: 264-273

671 Plaxton WC and Podestá FE (2006) The functional organization and control of plant respiration.
672 *Critical Rev Plant Sci* 25: 159–198

673 Pleijel H, Danielsson H (1997) Growth of 27 herbs and grasses in relation to ozone exposure and plant
674 strategy. *New Phytol* 135: 361-367

675 Polidoros AN, Mylona PV, Arnholdt-Schmitt B (2009) Aox gene structure, transcript variation and
676 expression in plants. *Physiol Plant* 137: 342-353

677 Purvis AC, Shewfelt RL (1993) Does the alternative pathway ameliorate chilling injury in sensitive
678 plant tissues? *Physiol Plant* 88: 712–718

679 Ranieri A, D’Urso G, Nali C, Lorenzini G, Soldatini GF (1996) Ozone stimulates apoplastic
680 antioxidant systems in pumpkin leaves. *Physiol Plant* 97: 381-387

681 Rao, M. V., Lee, H. I., Creelman, R. A., Mullet, J. E., & Davis, K. R. (2000) Jasmonic acid signaling
682 modulates ozone-induced hypersensitive cell death. *Plant Cell* 12: 1633-1646

683 Rausch T, Gromes R, Liedschulte V, Müller I, Bogs J, Galovic V, et al. (2007) Novel insight into the
684 regulation of GSH biosynthesis in higher plants. *Plant Biology* 9: 565–572

685 Rebouças DM, De Sousa YM, Bagard M, Costa JH, Jolivet Y, De Melo DF, Repellin A (2017).
686 Combined effects of ozone and drought on the physiology and membrane lipids of two cowpea
687 (*Vigna unguiculata* (L.) Walp) cultivars. *Plants* 6: 14.

688 Reich PB (1983) Effects of low concentrations of O₃ on net photosynthesis, dark respiration, and
689 chlorophyll contents in aging hybrid poplar leaves. *Plant Physiol* 73:291–296

690 Renaut J, Bohler S, Hausman JF, Hoffmann L, Sergeant K, Ashan N, Jolivet Y, Dizengremel P (2009)
691 The impact of atmospheric composition on plants: a case study of ozone and poplar. *Mass*
692 *Spectrometry Rev* 28: 495–516

693 Rhoads DM, Subbaiah CC (2007) Mitochondrial retrograde regulation in plants. *Mitochondrion*
694 7:177–94

695 Rhoads DM, Umbach AL, Subbaiah CC, Siedow JN (2006) Mitochondrial reactive oxygen species.
696 Contribution to oxidative stress and interorganellar signaling. *Plant Physiol* 141: 357– 366

697 Rouhier N, Gelhaye E, Gualberto JM, Jordy MN, de Fay E, Hirasawa M, Duplessis S, Lemaire SD,
698 Frey P, Martin F, Manieri W, Knaff DB, Jacquot JP (2004) Poplar peroxiredoxin Q. A thioredoxin-
699 linked chloroplast antioxidant functional in pathogen defense. *Plant Physiol* 134 : 1027-1038

700 Saha B, Borovskii G, Panda SK (2016) Alternative oxidase and plant stress tolerance. *Plant Signaling*
701 *Behavior* 11:4

702 Scalera V, Giangregorio N, De Leonardis S, Console L, Carulli ES, Tonazzi A (2018)
703 Characterization of a novel mitochondrial ascorbate transporter from rat liver and potato
704 mitochondria. *Frontiers in Mol Biosci.* 5 : 58-65.

705 Scheibe R (2004) Malate valve to balance cellular energy supply. *Physiol Plant* 120: 21-26

706 Sehmer L, Fontaine V, Antoni F, Dizengremel P (1998) Effects of ozone and elevated atmospheric
707 carbon dioxide on carbohydrate metabolism of spruce needles. Catabolic and detoxification
708 pathways. *Physiol Plant* 102:605–611

709 Selinski J, Hartmann A, Deckers-Hebestreit G, Day DA, Whelan J, Scheibe R (2018) Alternative
710 oxidase isoforms are differentially activated by tricarboxylic acid cycle intermediates. *Plant Physiol*
711 176 : 1423-1432

712 Short EF, North KA, Roberts MR, Hetherington AM, Shirras AD, McAinsh MR (2012) A stress-
713 specific calcium signature regulating an ozone -responsive gene expression network in
714 *Arabidopsis*. *Plant J* 71: 948-961

715 Sun YL, Zhao Y, Hong X, Zhai ZH (1999) Cytochrome c release and caspase activation during
716 menadione-induced apoptosis in plants. *FEBS Lett* 462: 317–321

717 Sutinen S, Skärby L, Wallin G, Selldén G (1990) Long-term exposure of Norway spruce, *Picea abies*
718 (L.) Karst., to ozone in open top chambers II. Effects on the ultrastructure of needles. *New Phytol*
719 115: 345-355

720 Sweetlove LJ, Foyer (2004) Roles for reactive oxygen species and antioxidants in plant mitochondria.
721 In Day DA, Millar AH, Whelan J (eds) *Plant Mitochondria: From Genome to Function*, Advances
722 in Photosynthesis and Respiration, Kluwer Academic Press, Dordrecht, vol 17, pp 307–320

723 Sweetlove LJ, Heazlewood JL Herald V, Holtzapffel R, Day DA, Leaver CJ, Millar AH (2002) The
724 impact of oxidative stress on *Arabidopsis* mitochondria. *Plant J* 32: 891-904.

725 Sweetlove LJ, Beard KF, Nunes-Nesi A, Fernie AR, Ratcliffe RG (2010) Not just a circle: flux modes
726 in the plant TCA cycle. *Trends Plant Sci* 15: 462–470

727 Szarka A, Bánhegyi G, Asard H (2013) The inter-relationship of ascorbate transport, metabolism and
728 mitochondrial, plastidic respiration *Antioxid Redox Signal* 19: 1036-1044

729 Tcherkez E, Mahé A, Gauthier P, Mauve C, Gout E, Bligny R, Cornic G, Hodges M (2009) In folio
730 respiratory fluxomics revealed by ¹³C isotopic labeling and H/D isotope effects highlight the
731 noncyclic nature of the tricarboxylic acid “cycle” in illuminated leaves. *Plant Physiol* 151: 620-630

732 Tiwari BS, Belenghi B, A (2002) Oxidative stress increased respiration and generation of reactive
733 oxygen species, resulting in ATP depletion, opening of mitochondrial permeability transition, and
734 programmed cell death. *Plant Physiol* 128: 1271–1281

735 Tosti N, Pasqualini S, Borgogni A, Ederli E, Falistocco E, Crispi S, Paolocci F (2006) Gene
736 expression profiles of ozone treated *Arabidopsis* plants. *Plant Cell Environ* 29: 1686-1702

737 Umbach AL, Gonzalez-Meier MA, Sweet CR, Siedow JN (2002) Activation of the plant
738 mitochondrial alternative oxidase: insights from site-directed mutagenesis. *Biochim Biophys Acta*
739 1554: 118-128

740 Umbach AL, Fiorani F, Siedow JN (2005) Characterization of transformed *Arabidopsis* with altered
741 alternative oxidase levels and analysis of effects on reactive oxygen species in tissue. *Plant Phys*
742 139: 1806-1820

743 Umekawa Y, Ito K (2019) Thioredoxin o-mediated reduction of mitochondrial alternative oxidase in
744 the thermogenic skunk cabbage *Symplocarpus renifolius*. *J Biol Chem* 165: 57-65

745 Vacca RA, Valenti D, Bobba A, Merafina RS, Passarella S, Marra E (2006) Cytochrome c is released
746 in a reactive oxygen species-dependent manner and is degraded via caspase-like proteases in
747 tobacco BY-2 cells en route to heat shock-induced cell death. *Plant Physiol* 141:208-219

748 Vanlerberghe GC, McIntosh L (1997) Alternative oxidase: from gene to function. *Annu. Rev. Plant*
749 *Physiol Plant Mol Biol* 48:703–34

750 Vercesi AE, Borecki J, De Godoy Maia I, Arruda P, Cuccovia IM, Chaimovich H (2006) Plant
751 uncoupling mitochondrial proteins. *Ann Rev Plant Biol* 57: 383-404

752 Vianello A, Zancani M, Peresson C, Petrussa E, Casolo V, Krajnakova J, Patui S, Braidot E, Macrı F
753 (2007) Plant mitochondrial pathway leading to programmed cell death. *Physiol Plant* 129: 242–252

754 Vieira Dos Santos C, Rey P (2006). Plant thioredoxins are key actors in oxidative stress response.
755 *Trends Plant Sci* 11: 329–334

756 Vollenweider P, Günthardt-Goerg S, Menard T, Baumgarten M, Matyssek R, Schaub M (2019)
757 Macro- and microscopic leaf injury triggered by ozone stress in beech foliage (*Fagus sylvatica* L.).
758 *Ann Forest Sci* doi.org/10.1007/s13595-019-0856-5

759 Welchen E, Gonzalez DH (2016) Cytochrome c, a hub linking energy, redox, stress and signaling
760 pathways in mitochondria and other cell compartments. *Physiol Plant* 157:310-321

761 Winger AM, Millar AH, Day DA (2005) Sensitivity of plant mitochondrial terminal oxidases to the
762 lipid peroxidation product 4-hydroxy-2-nonenal (HNE). *Biochem J* 387: 865-870

763 Wittig VE, Ainsworth EA, Long SP (2007) To what extent do current and projected increases in
764 surface ozone affect photosynthesis and stomatal conductance of trees? A meta-analytic review of
765 the last 3 decades of experiments. *Plant Cell Environ* 30:1150–1162

766 Wittig V, Ainsworth E, Naidu S, Karnosky D, Long S (2009) Quantifying the impact of current and
767 future tropospheric ozone on tree biomass, growth, physiology and biochemistry: a quantitative
768 meta-analysis. *Glob Change Biol* 15: 396–424

769 Xie Z, Chen Z (2000) Harpin-induced hypersensitive cell death is associated with altered
770 mitochondrial functions in tobacco cells. *Mol Plant-Microbe Interact* 13: 183-190

771 Xiong TC, Bourque S, Lecourieux D, Amelot N, Grat S, Briere C, Mazars C, Pugin A, Ranjeva R
772 (2006) Calcium signaling in plant cell organelles delimited by a double membrane. *Biochim*
773 *Biophys Acta* 1763 : 1209-1215

774 Yendrek CR, Erice G, Montes CM, Tomaz T, Brown PJ, McIntyre LM, Leakey ADB, Ainsworth
775 EA (2017) Elevated ozone reduces photosynthetic carbon gain by accelerating leaf senescence of
776 inbred and hybrid maize in a genotype-specific manner. *Plant Cell Env* 40: 3088-3100

777 Yoshida K, Terashima I, Noguchi K (2007) Up-regulation of mitochondrial alternative oxidase
778 concomitant with chloroplast over-reduction by excess light. *Plant Cell Physiol* 48: 606-614

779 Zancani M, Casolo V, Petrusa E, Peresson C, Patui S, Bertolini A, De Col V, Braidot E, Boscutti F,
780 Vianello A, Zancani M, Peresson C, Petrusa E, Casolo V, Krajnakova J, Patui S, Braidot E, Macri
781 F (2007) Plant mitochondrial pathway leading to programmed cell death. *Physiol Plant* 129: 242-
782 252

783 Zandalinas SI, Mittler R (2018) ROS-induced ROS release in plant and animal cells. *Free Radical Biol*
784 *Med* 122: 21-27

785

Table

Table 1.

Proposed sequential response of plant mitochondria to O₃ stress (see the text and Fig 1. for a complete explanation); *in italics: events inferred from general oxidative stress.*

Successive steps	Increasing Phytotoxic Ozone Dose
Mobilisation of pre-existing mechanisms to cope with ROS	
<u>Cytosol</u>	
- Enhancement of the catabolic pathways and PEPcase	
- Supply of organic acids and redox power (NADPH)	
- Maintenance of a high detoxification	
- Increase in the phenylpropanoid metabolism	
<u>Mitochondria</u>	
- Increased supply of NAD(P)H to ETC	
- Adjustment of electron flow path in ETC by balancing between high substrate supply (organic acid and NAD(P)H) and ATP requirement	
• Increased activity of UCP and AOX to lower reduction pressure	
• Ca ²⁺ -dependent activation of the NAD(P)H-ND _{ext} and ND _{int}	
• SA-mediated uncoupling of respiratory chain	
• Increased gene expression of AOX	
Setting up of detoxifying processes	
- TCA cycle inhibition by peroxidation products	
• Enhancement of anaplerotic pathway	
• Leak of organic acids to amino acids synthesis	
• Increase synthesis of glutathione and <i>import in the matrix</i>	
• Induction of GABA shunt	
- Conversion of superoxide by Mn-SOD	
- Induction of antioxidant defense	
• Increased <i>de novo</i> synthesis of ascorbate	
• Increased activity of HAF cycle and APX/GR gene expression	
• <i>Enhancement of peroxyredoxin/thioredoxins systems</i>	
- Inactivation of AOX	
- Complex II inhibition by high level of SA	
• <i>Accumulation of toxic intermediates of GABA shunt</i>	
Towards programmed cell death (PCD)	
- Increasing damage to fatty acids and proteins of mitochondrial membranes	
- ATP-synthesis decrease due to uncoupling of electron flow	
- Decrease in ascorbate synthesis	
- <i>Possible occurrence of Mitochondrial Permeability Transition</i>	
- Release of cytochrome c in the intermembrane space	
• <i>Activation of caspases-like proteases</i>	
- Disruption of the outer mitochondrial membrane	
- Swelling of mitochondria	
- Cell death	

838 *Figure legend*

839 *Figure 1. Representative schema of metabolic network involved in plant mitochondria response to O₃-*
840 *induced oxidative stress. Four main focuses of mitochondria response are presented to highlight (A)*
841 *the oxoglutarate-glutathione co-transport between cytoplasm and matrix that allow to replenish*
842 *matrix GSH pool jointly to the leak of OG out of mitochondria; (B) the GABA-shunt bypassing TCA*
843 *cycle enzymes once they are inhibited by ROS and its potential role in cell signalling and/or death;*
844 *(C) the multifaced electron pathway which is modulated to minimize endogenous ROS production and*
845 *its close relationship to ascorbate synthesis and import; (D) the antioxidant system reported in*
846 *mitochondria matrix to scavenge ROS. Dark lines and arrows represent mitochondria-related*
847 *metabolic links. Red lines represent the connectivity between mitochondrial metabolism and ROS*
848 *production or scavenging. ASA: ascorbate, APX: ascorbate peroxidase, AOX: alternative oxidase,*
849 *Cit: citrate, Cyt C: cytochrome C, Er-4-P: erythrose-4-phosphate, GABA: gamma-aminobutyric acid,*
850 *γ-Glu-cys: L-gamma-glutamylcysteine, GLDH: L-galactono-1,4-lactone dehydrogenase, Glu:*
851 *glutamate, GLUT-like Tr: glucose transporter, Glu-6-P: glucose-6-phosphate, GS/GOGAT: glutamine*
852 *synthetase/glutamate synthase, GPX: glutathione peroxidase, GR: glutathione reductase, GRX:*
853 *glutathione reductase, GSH: reduced glutathione, GSSG: oxidized glutathione, iCit: isocitrate, Mal:*
854 *malate, MAT Tr: mitochondrial ascorbic acid transporter, (M)DHA: (mono)dehydroascorbate,*
855 *(M)DHAR: (mono)dehydroascorbate reductase, mETC: mitochondrial electron transport chain,*
856 *MnSOD: Mn-dependent superoxide dismutase, NDin/ex: internal/external NAD(P)H dehydrogenases,*
857 *OAA: oxaloacetate, OG: oxoglutarate, OPPP: oxidative pentose phosphate pathway, PEPc:*
858 *phosphoenolpyruvate carboxylase, PRX/TRX: peroxiredoxin/thioredoxin, Pyr: pyruvate, ROS:*
859 *reactive oxygen species, SSA: succinic semialdehyde, Succ: succinate, UCP: uncoupling protein*

860

Der Doctor Schnabel von Rom



Vos Creditis, als eine fabel,
quod scribitur vom Doctor Schnabel,
der fugit die Contagion
et autert seinen Lohn darvon.
Cadavera sucht er zu fristen,
gleich wie der Corvus auf der Misten.
Ah Credite, zihet nicht dort hin,
dann Romæ regnat die Pestin.

Quis non deberet sehr erschree,
für seiner Virgul oder stecken,
qua loquitur, als wär er stumm,
und deutet sein consilium.
Wie mancher Credit ohne zweiffel
das ihm tentir ein schwarzen Teuffel.
Marsupium heist seine Höll,
und aurum die geholte seel.

I. Columbina, ad vitzum delineavit.

Paulus Fürst. Excudit.

Kleidung wider den Tod zu Rom. Anno 1656.
Also gehen die Doctores Medici dahin zu Rom, wann sie die ander Pest erkrankte Per-
sonen besuchen, sie zu curiren und fragen, sich wider den Ditt zu sichern, ein langes Kleid von ge-
wärtem Tuch ihr Angesicht ist verlarvt, für den Augen haben sie grosse Crystalline Brillen, wider
Nasen einen langen Schnabel voll wolriechender Specereij, in der Hände, welche mit Hand schühern
wol versehen ist, eine lange Kütze und darmit deuten sie, was man thun, und gebrauchte soll.