



HAL
open science

FIRST BAND OF RUELLE RESONANCES FOR CONTACT ANOSOV FLOWS IN DIMENSION 3

Mihajlo Cekić, Colin Guillarmou

► **To cite this version:**

Mihajlo Cekić, Colin Guillarmou. FIRST BAND OF RUELLE RESONANCES FOR CONTACT ANOSOV FLOWS IN DIMENSION 3. *Communications in Mathematical Physics*, 2021, 386, pp.1289–1318. 10.1007/s00220-021-04090-2 . hal-03020967v2

HAL Id: hal-03020967

<https://hal.science/hal-03020967v2>

Submitted on 24 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FIRST BAND OF RUELLE RESONANCES FOR CONTACT ANOSOV FLOWS IN DIMENSION 3

MIHAJLO CEKIĆ AND COLIN GUILLARMOU

ABSTRACT. We show, using semiclassical measures and unstable derivatives, that a smooth vector field X generating a contact Anosov flow on a 3-dimensional manifold \mathcal{M} has only finitely many Ruelle resonances in the vertical strips $\{s \in \mathbb{C} \mid \operatorname{Re}(s) \in [-\nu_{\min} + \varepsilon, -\frac{1}{2}\nu_{\max} - \varepsilon] \cup [-\frac{1}{2}\nu_{\min} + \varepsilon, 0]\}$ for all $\varepsilon > 0$, where $0 < \nu_{\min} \leq \nu_{\max}$ are the minimal and maximal expansion rates of the flow (the first strip only makes sense if $\nu_{\min} > \nu_{\max}/2$). We also show polynomial bounds in s for the resolvent $(-X - s)^{-1}$ as $|\operatorname{Im}(s)| \rightarrow \infty$ in Sobolev spaces, and obtain similar results for cases with a potential. This is a short proof of a particular case of the results by Faure-Tsujii in [FaTs0, FaTs2, FaTs3], using that $\dim E_u = \dim E_s = 1$.

1. INTRODUCTION

In this note, we study the localization of Ruelle resonances for contact Anosov flows in dimension 3 using semiclassical measures, and we show the existence of a first band of resonances under a pinching condition on the maximal and minimal expansion rates of the flow. The fact that a vector field X generating a contact Anosov flow has a band structure (see Figure 1) for its Ruelle resonance spectrum is proved in a series of seminal papers by Faure-Tsujii [FaTs0, FaTs2, FaTs3] using FBI transform techniques and normal forms. In the 3-dimensional case, we give a rather short proof of the existence of the first band using unstable derivatives and semiclassical measures in the spirit of Dyatlov's proof [Dy2] for operators with r -normally hyperbolic trapped set. Thus the aim of this work is to use the 3-dimensional particularity and semiclassical measures to present the mechanism behind this structure.

Let \mathcal{M} be a 3-dimensional closed manifold, equipped with a Riemannian metric g and let X be a smooth vector field on \mathcal{M} such that its flow φ_t is Anosov (see Section 2.1 for a definition). Denote by $E_u \subset T\mathcal{M}$ and $E_s \subset T\mathcal{M}$ the unstable and stable bundles. We shall assume that φ_t is a contact Anosov flow, which means that there is a smooth 1-form α on \mathcal{M} such that $\ker \alpha = E_u \oplus E_s$, $\alpha(X) = 1$ and $\alpha \wedge d\alpha$ is a volume form, and that E_u, E_s are orientable¹ (and hence trivializable) bundles. Define the minimal and maximal expansion rate of φ_t to be²

$$\nu_{\min} := \liminf_{t \rightarrow \infty} \inf_{x \in \mathcal{M}} \frac{1}{t} \log \|d\varphi_t(x)|_{E_u}\|_g, \quad \nu_{\max} := \limsup_{t \rightarrow \infty} \sup_{x \in \mathcal{M}} \frac{1}{t} \log \|d\varphi_t(x)|_{E_u}\|_g.$$

Now, we come to the notion of Ruelle resonances for X . It is proved by Butterley-Liverani [BuLi] and Faure-Sjöstrand [FaSj] that for each $N > 0$, there is a Hilbert space \mathcal{H}^N , called anisotropic Sobolev space of order N , satisfying³ $H^N(\mathcal{M}) \subset \mathcal{H}^N \subset H^{-N}(\mathcal{M})$ so that for $\operatorname{Re}(s) > -\nu_{\min}N$

$$-X - s : \{u \in \mathcal{H}^N \mid -Xu \in \mathcal{H}^N\} \rightarrow \mathcal{H}^N$$

is a Fredholm operator of index 0, implying that $-X$ has discrete spectrum in the half-plane $\{s \in \mathbb{C} \mid \operatorname{Re}(s) > -\nu_{\min}N\}$. Moreover, there is no spectrum in $\operatorname{Re}(s) > 0$, the spectrum in $\operatorname{Re}(s) > -\nu_{\min}N_0$ on \mathcal{H}^N does not depend on $N \geq N_0$ or the choice of the space \mathcal{H}^N and the resolvent

$$R(s) := (-X - s)^{-1} : C^\infty(\mathcal{M}) \rightarrow C^{-\infty}(\mathcal{M})$$

¹For instance, this is satisfied for Anosov geodesic flows on orientable Riemannian surfaces.

²The limit exists by Fekete's lemma, and the same holds as well for V_{\max}, V_{\min} defined later.

³ $H^N(\mathcal{M}) = (1 + \Delta_g)^{-N/2} L^2(\mathcal{M})$ denotes the usual Sobolev space of order N on \mathcal{M} .




FIGURE 1. Left: resonance spectrum of a closed hyperbolic surface Σ [DFG]; colouring describes the nature of the resonance. Right: band structure of a contact Anosov flow on a 3-manifold given by Theorem 1. In dashed blue regions we have only finitely many resonances, with semiclassical resolvent bounds written using \mathcal{O} and o notation; gray regions are not covered by the theorem.

is meromorphic in \mathbb{C} . An $s \in \mathbb{C}$ is a pole of the resolvent if and only if it is an eigenvalue of $-X$ in \mathcal{H}^N for some $N > -\text{Re}(s)/\nu_{\min}$, and such s are called *Ruelle resonances*, and $u \in \mathcal{H}^N$ with $(X + s)u = 0$ are called *resonant states*.

Since we want to prove absence of Ruelle resonances and bounds on the resolvent $R(s)$ for large $\text{Im}(s)$, it is convenient to use a semiclassical rescaling, that is to take a small parameter $h > 0$ so that $h\text{Im}(s)$ is uniformly bounded. For such small parameter $h > 0$, we denote by H_h^N the $H^N(\mathcal{M})$ Sobolev space equipped with the norm $\|u\|_{H_h^N} := \|(1 + h^2\Delta_g)^{N/2}u\|_{L^2}$.

We prove the following result:

Theorem 1. *Assume that X is a smooth contact Anosov flow on a closed 3-manifold \mathcal{M} with E_u, E_s orientable, and let ν_{\min}, ν_{\max} be the minimal and maximal expansion rates of the flow. For any $\varepsilon > 0$, there are only finitely many Ruelle resonances in the regions (see Figure 1)*

$$\mathcal{S}_0(\varepsilon) := \left\{ s \in \mathbb{C} \mid \text{Re } s > -\frac{\nu_{\min}}{2} + \varepsilon \right\} \quad \text{and} \quad \mathcal{S}_1(\varepsilon) := \left\{ s \in \mathbb{C} \mid \text{Re } s \in (-\nu_{\min} + \varepsilon, -\frac{\nu_{\max}}{2} - \varepsilon) \right\},$$

and the following bounds hold, as $|s| \rightarrow \infty$

$$\|(-X - s)^{-1}\|_{H^{\frac{1}{2}}(\mathcal{M}) \rightarrow H^{-\frac{1}{2}}(\mathcal{M})} = o(|s|), \quad s \in \mathcal{S}_0(\varepsilon),$$

$$\|(-X - s)^{-1}\|_{H^1(\mathcal{M}) \rightarrow H^{-1}(\mathcal{M})} = \mathcal{O}(|s|), \quad s \in \mathcal{S}_1(\varepsilon).$$

More precisely, if $h \in (0, h_0)$ is a small parameter, for any $N > 0$ there exist Hilbert spaces \mathcal{H}_h^N satisfying uniform bounds

$$\exists C > 0, \forall u \in C^\infty(\mathcal{M}), \forall h \in (0, h_0), \quad C^{-1}\|u\|_{H_h^{-N}} \leq \|u\|_{\mathcal{H}_h^N} \leq C\|u\|_{H_h^N},$$

such that the following bounds hold: for all $\varepsilon > 0$ and $A > 1$, and $\lambda \in \mathbb{C}$ satisfying $|\text{Im}(\lambda)| \in [A^{-1}, A]$, as $h \rightarrow 0$

$$\|(-hX - \lambda)^{-1}\|_{\mathcal{H}_h^N \rightarrow \mathcal{H}_h^N} = o(h^{-2}), \quad h^{-1}\lambda \in \mathcal{S}_0(\varepsilon), \quad N \geq \frac{1}{2},$$

$$\|(-hX - \lambda)^{-1}\|_{\mathcal{H}_h^N \rightarrow \mathcal{H}_h^N} = \mathcal{O}(h^{-2}), \quad h^{-1}\lambda \in \mathcal{S}_1(\varepsilon), \quad N \geq 1.$$

As mentioned above, this result is contained in the set of results of Faure-Tsujii [FaTs0, FaTs2, FaTs3]. In fact, the second strip we obtain with finitely many resonances is not optimal and should be $\text{Re}(s) \in [-\frac{3\nu_{\min}}{2} + \varepsilon, -\frac{\nu_{\max}}{2} - \varepsilon]$ for all $\varepsilon > 0$ small. A band structure for Ruelle

resonances was first obtained by Faure-Tsujii in [FaTs1, FaTs2, FaTs3] for prequantum maps and Anosov flows using FBI transform methods and normal forms, including the case of a geometric potential (multiples of the unstable Jacobian). In the case of constant curvature or locally symmetric spaces, the bands are actually lines [FaTs0, DFG, GHW, KuWe] (see Figure 1). In a related context, for operators with r -normally hyperbolic trapped set, Dyatlov [Dy1, Dy2] also proved some band structure for resonances.

Our proof is adapted to the case of a 1-dimensional bundles E_u, E_s , and should be quite accessible for a reader with a semiclassical background (pseudo-differential calculus). The aim of this note is to exhibit the mechanism behind the band structure in this simpler 3-dimensional case. The existence of the first region $\{s \in \mathbb{C} \mid \operatorname{Re}(s) > -\nu_{\min}/2 + \varepsilon\}$ with only finitely many Ruelle resonances for contact Anosov flows was first shown by Tsujii [Ts1, Ts2] using FBI transform, and then by Nonnenmacher-Zworski [NoZw] using normal hyperbolicity of the trapped set; both results also hold in higher dimension where ν_{\min} is replaced by $\lim_{t \rightarrow \infty} \inf_{x \in \mathcal{M}} \frac{1}{t} \log \det(d\varphi_t(x)|_{E_u})$. Here, the method of proof is more similar to Dyatlov's approach [Dy1, Dy2] for r -normally hyperbolic trapped sets, and uses semiclassical measures and propagation estimates. It was also used in the setting of frame flows in constant curvature [GuKu]. We also mention that the existence of a (non-explicit) small strip without Ruelle resonances in the contact Anosov setting was first proved by Dolgopyat [Do] and Liverani [Li], and that there are two recent breakthroughs for 3-dimensional Anosov flows: first for volume preserving Anosov flows by Tsujii [Ts3], and finally Tsujii-Zhang [TsZh] recently proved the existence of a strip with no Ruelle resonances for all topologically mixing Anosov flows in dimension 3.

The main idea of our proof is based on an observation made in a paper of the second author and Faure [FaGu], namely that the Ruelle resonant states $u \in \mathcal{H}^1$ with resonance in $\operatorname{Re}(s) > -\nu_{\min}$ satisfy $U_- u = 0$ where U_- is a vector field tangent to the unstable foliation (with regularity $C^{2-\varepsilon}(\mathcal{M}; T\mathcal{M})$). The outline of our proof is as follows. First, the existence of a sequence of Ruelle resonances s_n (or of quasimodes) satisfying $\operatorname{Re}(s_n) \rightarrow -\gamma$ for some $-\gamma > -\nu_{\min} + \varepsilon$ and $\operatorname{Im}(s_n) \rightarrow \infty$ implies the existence of a non-zero semiclassical measure μ on $T^*\mathcal{M}$, which, in microlocal terms, is the weak limit of a sequence of Ruelle resonant states (or quasimodes) $u_n \in \mathcal{H}^1$. By microlocal ellipticity it is supported in $\{(x, \xi) \in T^*\mathcal{M} \mid \xi(X) = -1\}$, and using the propagation of singularities estimate, one can see that for all $t \in \mathbb{R}$

$$(\Phi_t)_* \mu = e^{-2\gamma t} \mu, \quad (1.1)$$

where $\Phi_t(x, \xi) = (\varphi_t(x), (d\varphi_t(x))^{-1}{}^T \xi)$ is the symplectic lift of the flow φ_t on $T^*\mathcal{M}$. Next, using radial point propagation estimates related to the hyperbolicity of Φ_t , we can see that

$$\operatorname{supp}(\mu) \subset \Gamma_+ := E_u^* \oplus E_0^*,$$

where $E_0^* := \mathbb{R}\alpha$ and E_u^* is the annihilator of $\mathbb{R}X \oplus E_u \subset T\mathcal{M}$. The last step is to use a version of the horocyclic invariance of the resonant states u_n proved in [FaGu], that is $U_- u_n = 0$ (or small in h). This invariance of u_n by U_- can be understood as an extra invariance of the measure μ , which implies (by propagation estimates again) some regularity of μ at the trapped set E_0^* of Φ_t . This is where the contact assumption is important, it appears as a transversality of the symplectic lift of the horocyclic flow (i.e. the flow of U_-) with the trapped set $E_0^* = \mathbb{R}\alpha$. Namely one shows that for all $\delta > 0$ small and \mathcal{U}_δ a δ -neighborhood of $E_0^* \cap \{\xi(X) = -1\}$ in $T^*\mathcal{M}$

$$\mu(\mathcal{U}_\delta) \in (\delta/C, C\delta)$$

for some uniform $C > 0$. Combining with (1.1) and using the hyperbolicity of Φ_t :

$$\forall t \geq 0, \quad (\mathcal{U}_{\delta e^{-(\nu_{\max} + \varepsilon)t}} \cap \Gamma_+) \subset \Phi_{-t}(\mathcal{U}_\delta \cap \Gamma_+) \subset (\mathcal{U}_{\delta e^{-(\nu_{\min} - \varepsilon)t}} \cap \Gamma_+),$$

which implies by an elementary argument that $-\gamma$ must be in $[-(\nu_{\max} + \varepsilon)/2, -(\nu_{\min} - \varepsilon)/2]$. One of the main issues in this argument is that U_- is only a $C^{2-\varepsilon}(\mathcal{M}; T\mathcal{M})$ vector field for all $\varepsilon > 0$, which complicates the use of microlocal methods. To circumvent the problem, we regularize U_- at scale $h^\rho = |\operatorname{Im}(s_n)|^{-\rho}$ for some $0 < \rho < 1$, to make it a smooth, h -dependent, vector field U_-^h in some exotic class of differential operators. The resonant states (or quasimodes) are

not killed anymore by U_-^h but they are good enough quasimodes to apply the reasoning above. We have to show, in particular, that the propagation estimates still hold in this exotic class (see Appendix A).

In fact, since it does not involve more difficulties, we prove the more general result involving a smooth real potential V . We introduce the quantities

$$V_{\max} := \lim_{t \rightarrow \infty} \sup_{x \in \mathcal{M}} \frac{1}{t} \int_0^t V(\varphi_s(x)) ds, \quad V_{\min} := \lim_{t \rightarrow \infty} \inf_{x \in \mathcal{M}} \frac{1}{t} \int_0^t V(\varphi_s(x)) ds. \quad (1.2)$$

Theorem 2. *Let \mathcal{M} and X satisfy the same assumptions as in Theorem 1, and let $V \in C^\infty(\mathcal{M})$ be a real valued potential. For any $\varepsilon > 0$, there are only finitely many Ruelle resonances of $-X + V$ in the regions*

$$\begin{aligned} \mathcal{S}_0(\varepsilon) &:= \left\{ s \in \mathbb{C} \mid \operatorname{Re} s > -\frac{\nu_{\min}}{2} + V_{\max} + \varepsilon \right\} \quad \text{and} \\ \mathcal{S}_1(\varepsilon) &:= \left\{ s \in \mathbb{C} \mid \operatorname{Re} s \in \left(-\nu_{\min} + V_{\max} + \varepsilon, -\frac{\nu_{\max}}{2} + V_{\min} - \varepsilon\right) \right\}, \end{aligned}$$

and the following bounds hold, as $|s| \rightarrow \infty$

$$\begin{aligned} \|(-X + V - s)^{-1}\|_{H^{\frac{1}{2}}(\mathcal{M}) \rightarrow H^{-\frac{1}{2}}(\mathcal{M})} &= o(|s|), \quad s \in \mathcal{S}_0(\varepsilon), \\ \|(-X + V - s)^{-1}\|_{H^1(\mathcal{M}) \rightarrow H^{-1}(\mathcal{M})} &= \mathcal{O}(|s|), \quad s \in \mathcal{S}_1(\varepsilon). \end{aligned} \quad (1.3)$$

More precisely, with the notation of Theorem 1, then for all $\varepsilon > 0$ and $A > 1$, and for all $\lambda \in \mathbb{C}$ satisfying $|\operatorname{Im}(\lambda)| \in [A^{-1}, A]$, we have as $h \rightarrow 0$

$$\begin{aligned} \|(-hX + hV - \lambda)^{-1}\|_{\mathcal{H}_h^N \rightarrow \mathcal{H}_h^N} &= o(h^{-2}), \quad h^{-1}\lambda \in \mathcal{S}_0(\varepsilon), \quad N \geq \frac{1}{2}, \\ \|(-hX + hV - \lambda)^{-1}\|_{\mathcal{H}_h^N \rightarrow \mathcal{H}_h^N} &= \mathcal{O}(h^{-2}), \quad h^{-1}\lambda \in \mathcal{S}_1(\varepsilon), \quad N \geq 1. \end{aligned} \quad (1.4)$$

Here again the existence of a band structure for Ruelle resonances in the case of a potential is proved in [FaTs0, FaTs2, FaTs3].

Acknowledgements. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 725967). We thank Semyon Dyatlov for several useful discussions and for suggesting to apply the method of [Dy2] in this setting.

2. ANOSOV FLOWS, ANISOTROPIC SPACES AND RUELLE RESONANCES

2.1. Dynamical background. Let \mathcal{M} be a 3-dimensional closed manifold equipped with a Riemannian metric g , which also induces a metric on $T^*\mathcal{M}$, the Sasaki metric. We will consider a smooth vector field X on \mathcal{M} and we assume that the flow φ_t of X is Anosov: there is a $d\varphi_t$ invariant splitting

$$T\mathcal{M} = E_0 \oplus E_u \oplus E_s$$

where $E_0 = \mathbb{R}X$ is the flow direction, E_s is the stable bundle defined by

$$v \in E_s \iff \exists C > 0, \nu > 0, \forall t \geq 0, \quad \|d\varphi_t(v)\| \leq Ce^{-\nu t} \|v\|$$

and E_u is the unstable bundle defined by

$$v \in E_u \iff \exists C > 0, \nu > 0, \forall t \leq 0, \quad \|d\varphi_t(v)\| \leq Ce^{-\nu|t|} \|v\|.$$

The bundles E_u and E_s are 1-dimensional vector bundles, which have Hölder regularity. We say that the flow is a *contact* Anosov flow if there is a smooth 1-form α such that $\alpha(X) = 1$, $\ker d\alpha = E_u \oplus E_s$ and $d\alpha$ is non-degenerate on $\ker \alpha$, or equivalently $\alpha \wedge d\alpha$ is a smooth volume form on \mathcal{M} . If the flow of X is a contact Anosov flow, then the stable and unstable bundles E_s, E_u have regularity $C^{2-\varepsilon}(\mathcal{M})$ for all $\varepsilon > 0$ by the result of Hurder-Katok [HuKa]. For

notational convenience we will denote $C^{k-}(\mathcal{M}) := \cap_{\varepsilon>0} C^{k-\varepsilon}(\mathcal{M})$ for $k \in \mathbb{N}$. We shall assume that E_s and E_u are orientable, which means that there are two non-vanishing vector fields $U_{\pm} \in C^{2-}(\mathcal{M}; T\mathcal{M})$ such that

$$E_s = \mathbb{R}U_+, \quad E_u = \mathbb{R}U_-.$$

Note that the orientability condition for E_u, E_s is satisfied for Anosov geodesic flows on orientable surfaces. We can define a dual Anosov decomposition of $T^*\mathcal{M}$

$$T^*\mathcal{M} = E_0^* \oplus E_u^* \oplus E_s^*, \quad \text{with } E_u^*(E_u \oplus \mathbb{R}X) = 0, \quad E_s^*(E_s \oplus \mathbb{R}X) = 0$$

and $E_0^* = \mathbb{R}\alpha$ is the annihilator of $E_u \oplus E_s$. We will also define

$$\Gamma_+ := E_u^* \oplus E_0^*, \quad \Gamma_- = E_s^* \oplus E_0^*, \quad K = \Gamma_+ \cap \Gamma_- = E_0^* \quad (2.1)$$

that we call the outgoing tail, the incoming tail and the trapped set for X , respectively.

Let us now recall a couple of properties about U_{\pm} from [FaGu, Lemma 2.2]: we have the following commutation formula

$$[X, U_{\pm}] = \pm r_{\pm} U_{\pm}, \quad (2.2)$$

and $r_{\pm}, U_{\pm} \in C^{2-}$. From (2.2), we conclude for all $t \in \mathbb{R}$

$$\begin{aligned} d\varphi_{-t}(x)U_-(x) &= e^{-\int_{-t}^0 \varphi_s^* r_-(x) ds} U_-(\varphi_{-t}(x)), \\ d\varphi_t(x)U_+(x) &= e^{-\int_0^t \varphi_s^* r_+(x) ds} U_+(\varphi_t(x)). \end{aligned} \quad (2.3)$$

We note that both U_- and r_- are not uniquely defined but the large time average of r_- along orbits are intrinsic to X . By Fekete's lemma, we may define the following finite, positive quantities

$$\nu_{\min} := \liminf_{t \rightarrow \infty} \inf_{x \in \mathcal{M}} \frac{1}{t} \int_0^t r_-(\varphi_{-s}(x)) ds, \quad \nu_{\max} := \limsup_{t \rightarrow \infty} \sup_{x \in \mathcal{M}} \frac{1}{t} \int_0^t r_-(\varphi_{-s}(x)) ds. \quad (2.4)$$

Note that in the contact case, one also has

$$\nu_{\min} = \liminf_{t \rightarrow \infty} \inf_{x \in \mathcal{M}} \frac{1}{t} \int_0^t r_+(\varphi_s(x)) ds, \quad \nu_{\max} = \limsup_{t \rightarrow \infty} \sup_{x \in \mathcal{M}} \frac{1}{t} \int_0^t r_+(\varphi_s(x)) ds. \quad (2.5)$$

Then for each $\varepsilon > 0$, there exists $C_{\varepsilon} > 0$ such that for all $t \geq 0$ and $x \in \mathcal{M}$

$$C_{\varepsilon}^{-1} e^{-(\nu_{\max} + \varepsilon)t} \leq |d\varphi_{\pm t} U_{\pm}(x)| \leq C_{\varepsilon} e^{-(\nu_{\min} - \varepsilon)t}.$$

2.2. Anisotropic space and extension of the resolvent of X . Let $V \in C^{\infty}(\mathcal{M})$ be a potential. We want to prove a spectral gap with resolvent estimates at high frequency for $-X + V$. It is convenient to make the semiclassical rescaling

$$P_h := -ihX + ihV, \quad P_h(\lambda) = P_h - i\lambda \quad (2.6)$$

where $h > 0$ is a small parameter and $\lambda \in \mathbb{C}$. The semiclassical principal symbol of P_h is given by

$$p(x, \xi) = \xi(X). \quad (2.7)$$

We will denote by Φ_t the Hamiltonian flow at time t of p : notice that

$$\Phi_t(x, \xi) = (\varphi_t(x), (d\varphi_t(x))^{-1})^T \xi).$$

The Hamiltonian vector field of p will be denoted by H_p so that $\Phi_t = e^{tH_p}$. In [FaSj], Faure and Sjöstrand construct a family of Hilbert spaces, called anisotropic Sobolev spaces, using variable order pseudo-differential operators. Another presentation is given by Dyatlov-Zworski [DyZw], where a semiclassical parameter is included. We recall a few results about these spaces and the spectral properties of P_h acting on them, we refer to [FRS, FaSj, DyZw] for more details.

If $m \in S^0(T^*\mathcal{M})$ is a symbol of degree 0 on $T^*\mathcal{M}$, i.e. satisfying the bounds in local coordinates (for some constants $C_{\alpha, \beta} > 0$)

$$|\partial_x^{\alpha} \partial_{\xi}^{\beta} m(x, \xi)| \leq C_{\alpha, \beta} \langle \xi \rangle^{-|\beta|},$$

we denote by $\Psi_h^m(\mathcal{M})$ the space of semiclassical pseudo-differential operators of order m as defined in [FRS, Appendix] (see also [DyZw] for the h -dependent version): these are operators which have the form in local coordinates

$$Au(x) = (2\pi h)^{-3} \int e^{\frac{i}{h}(x-y)\xi} a(x, \xi) u(y) dy d\xi$$

where $a \in S^m(T^*\mathcal{M})$ is a symbol of order m , ie. it satisfies local bounds for each $\varepsilon > 0$ small

$$|\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \leq C_{\alpha, \beta, \varepsilon} \langle \xi \rangle^{m(x, \xi) - (1-\varepsilon)|\beta|}.$$

We also fix a semiclassical quantization Op_h on \mathcal{M} mapping symbols to operators acting on L^2 (see [Zw] and [DyZw, Appendix E]). The space $\Psi_h^{\text{comp}}(\mathcal{M})$ denotes the space of compactly microsupported operators, defined as the space of operators of the form $\text{Op}_h(a) + h^\infty \Psi_h^{-\infty}(\mathcal{M})$ with $a \in C_c^\infty(T^*\mathcal{M})$. We shall denote by $\sigma : \Psi_h^m(\mathcal{M}) \rightarrow S^m(T^*\mathcal{M})/hS^{m-1+\varepsilon}(T^*\mathcal{M})$ the semiclassical principal symbol map, satisfying $\sigma(\text{Op}_h(a)) - a \in hS^{m-1+\varepsilon}(T^*\mathcal{M})$ for all $\varepsilon > 0$. Finally, let $H_h^s := (1 + h^2 \Delta_g)^{-s/2} L^2(\mathcal{M})$ be the semiclassical Sobolev space of order s (here Δ_g is the Riemannian Laplacian of g on \mathcal{M}).

First, by [FaSj] there exists an *escape function* $G \in C^\infty(T^*\mathcal{M})$ of the form

$$G(x, \xi) = m(x, \xi) \log f(x, \xi)$$

where $m \in C^\infty(T^*\mathcal{M}; [-1, 1])$ is homogeneous of degree 0 in the fiber variable ξ for $|\xi| > 1$, $f > 0$ is homogeneous of degree 1 in the fiber variable for $|\xi| > 1$ and satisfies the following properties: there is a $C_m > 0$ such that

$$\begin{cases} m(x, \xi) = -1 \text{ in a conical neighborhood } V_u \text{ of } E_u^*, \\ m(x, \xi) = 1 \text{ in a conical neighborhood } V_s \text{ of } E_s^*, \\ H_p m \leq 0 \text{ on } T^*\mathcal{M}, \\ H_p G < -C_m \text{ outside a conical neighborhood } V_0 \text{ of } E_0^*. \end{cases}$$

We can thus define the anisotropic Sobolev space of order $N \in \mathbb{R}$ to be, for h small enough

$$\mathcal{H}_h^{NG} := \text{Op}_h(e^{NG})^{-1} L^2(\mathcal{M}).$$

Here $\text{Op}_h(e^{NG})$ belongs to the class of semiclassical pseudodifferential operators $\Psi_h^{Nm}(\mathcal{M})$ of variable order m , and it is invertible in the class of semiclassical pseudodifferential operators with inverse $\text{Op}_h(e^{NG})^{-1} \in \Psi_h^{-Nm}(\mathcal{M})$ if $h > 0$ is small enough. We notice that, since $\text{Op}_h(\langle \xi \rangle^{-N}) \text{Op}_h(e^{NG}) \in \Psi_h^0(\mathcal{M})$, there is $C > 0$ such that for all $u \in C^\infty(\mathcal{M})$ and $h > 0$

$$\frac{1}{C} \|u\|_{H_h^{-N}} \leq \|u\|_{\mathcal{H}_h^{NG}} \leq C \|u\|_{H_h^N}. \quad (2.8)$$

The following result was proved in [FaSj] (see also [DyZw])

Proposition 2.1. *There is⁴ $c_X > 0$, $c_V \in \mathbb{R}$ such that the operator $P_h(h\lambda) : \mathcal{D}(P_h) \subset \mathcal{H}_h^{NG} \rightarrow \mathcal{H}_h^{NG}$ is Fredholm in the region $\text{Re}(\lambda) > c_V - c_X N$, where $\mathcal{D}(P_h) = \{u \in \mathcal{H}_h^{NG} \mid P_h u \in \mathcal{H}_h^{NG}\}$. Its inverse $(P_h - i\lambda h)^{-1}$, called the resolvent of X , is a meromorphic family of bounded operators on \mathcal{H}_h^{NG} and the poles are called Ruelle resonances.*

3. SEMICLASSICAL MEASURES ASSOCIATED TO SEQUENCES OF RESONANCES AND QUASIMODES

In this Section, we shall see that the presence of an infinite number of Ruelle resonances λ_n with $\text{Re}(\lambda_n) \rightarrow -\gamma$ and $|\lambda_n| \rightarrow \infty$ implies the existence of a non-trivial measure $T^*\mathcal{M}$, with certain invariance properties with respect to H_p , and more generally the same holds for good quasimodes. This will allow us to prove gaps of resonances and bounds on the resolvent. The idea is to rescale the equation $(-X + V - \lambda_n)u_n = r_n$ with r_n small in some appropriate norm by setting $h_n = 1/|\lambda_n|$ so that, according to (2.6), $(P_{h_n} - ih_n \lambda_n)u_n = ih_n r_n$ and to view this

⁴It can be checked, by inspecting the proof, that one can choose $c_X = \nu_{\min}$ and $c_V = V_{\max}$.

last equation as a semiclassical problem. This method follows some ideas developed by Dyatlov in [Dy2] for operators with r -normally hyperbolic trapped set.

Assumption 1. For some $\gamma > 0$ and $N > 0$, consider a sequence

$$\operatorname{Re} \lambda_h = -h\gamma + o(h), \quad \operatorname{Im} \lambda_h = 1 + o(1), \quad -\gamma > V_{\max} - N\nu_{\min},$$

where $h > 0$ is a parameter going to 0, which can be discrete or continuous. Let $u_h \in \mathcal{D}(P_h) \subset \mathcal{H}_h^{NG}$ be a sequence (as $h \rightarrow 0$) of quasimodes for $P_h(\lambda_h)$: (A1)

$$\|u_h\|_{\mathcal{H}_h^{NG}} = 1, \quad \|P_h(\lambda_h)u_h\|_{\mathcal{H}_h^{NG}} = o(h^\beta),$$

for some $\beta > 0$.

First, by applying [DyZw, Theorem E.42], we directly have the

Lemma 3.1. *Under the assumptions of (A1), there is a subsequence u_{h_j} and a Radon measure μ on $T^*\mathcal{M}$ such that for each $A_h \in \Psi_h^{\text{comp}}(\mathcal{M})$*

$$\langle A_{h_j}u_{h_j}, u_{h_j} \rangle_{L^2} \rightarrow \int_{T^*\mathcal{M}} \sigma(A) d\mu \text{ as } j \rightarrow \infty.$$

We can also consider the sequence $\tilde{u}_{h_j} = \operatorname{Op}_{h_j}(e^{NG})u_{h_j}$ which satisfies $\|\tilde{u}_{h_j}\|_{L^2} = 1$ and thus one can consider its semiclassical measure $\tilde{\mu}$ satisfying, as $j \rightarrow \infty$,

$$\langle A_{h_j}\tilde{u}_{h_j}, \tilde{u}_{h_j} \rangle_{L^2} \rightarrow \int_{T^*\mathcal{M}} \sigma(A) d\tilde{\mu}.$$

3.1. Support and first invariance properties of the semiclassical measure. We first remark that the measure μ must satisfy some support properties and some invariance under the Hamiltonian flow of the $p(x, \xi)$ (defined in (2.7)). These are consequences of elliptic regularity and propagation of singularities.

Lemma 3.2. *The measures $\tilde{\mu}$ and μ satisfy $e^{2NG}d\mu = d\tilde{\mu}$, and*

$$\operatorname{supp}(\mu) \subset \{(x, \xi) \in T^*\mathcal{M} \mid \xi(X) = -1\}.$$

Assume that $\beta \geq 1$ in (A1). If $p(x, \xi) = \xi(X)$ and H_p is the Hamiltonian vector field of p , then for each $a \in C_c^\infty(T^\mathcal{M})$*

$$\int_{T^*\mathcal{M}} (H_p + 2(\gamma + V))a d\mu = 0. \quad (3.1)$$

Proof. The first identity follows from the definition of $\mu, \tilde{\mu}$ for $A \in \Psi_h^{\text{comp}}(\mathcal{M})$

$$\begin{aligned} \langle A_{h_j}\tilde{u}_{h_j}, \tilde{u}_{h_j} \rangle_{L^2} &= \langle \operatorname{Op}_{h_j}(e^{NG})^* A \operatorname{Op}_{h_j}(e^{NG})u_j, u_j \rangle_{L^2} \\ &\rightarrow_{j \rightarrow \infty} \int_{T^*\mathcal{M}} \sigma(\operatorname{Op}_{h_j}(e^{NG})^* A \operatorname{Op}_{h_j}(e^{NG})) d\mu = \int_{T^*\mathcal{M}} \sigma(A) e^{2NG} d\mu. \end{aligned}$$

The semiclassical principal symbol of $P_h(\lambda_h)$ is $\xi(X) + 1 + o(1)$ thus

$$\operatorname{supp}(\mu) \subset \{(x, \xi) \in T^*\mathcal{M} \mid \xi(X) = -1\}$$

using microlocal ellipticity [DyZw, Theorem E.43]. We write $\operatorname{Im}(P_h(\lambda_h)) = (P_h(\lambda_h) - P_h(\lambda_h)^*)/2i \in h\Psi_h^0(\mathcal{M})$, with

$$h^{-1}\operatorname{Im}(P_h(\lambda_h)) = \gamma + V + o(1).$$

By [DyZw, Theorem E.44] (or Proposition A.4), we have that for each $a \in C_c^\infty(T^*\mathcal{M})$, (3.1) holds. \square

Next, we shall use propagation estimates to obtain information on the support of μ and establish that $\mu \neq 0$.

Lemma 3.3. *Assume $\beta \geq 1$ in (A1), then the semiclassical measure μ satisfies*

$$\text{supp}(\mu) \subset \Gamma_+ = E_0^* \oplus E_u^*. \quad (3.2)$$

Moreover, if $K_R := \{(x, \xi) \in T^*\mathcal{M} \mid d_g((x, \xi), E_0^*) \leq R\}$, we have $\mu(K_R) > 0$ for any $R > 0$.

Proof. We fix $N_1 > 0$ an arbitrarily large constant. Let us consider the fiber compactification $\overline{T^*}\mathcal{M}$ of $T^*\mathcal{M}$ (see [DyZw, Appendix E] for the definition). The bundles E_u^* , E_s^* and E_0^* extend to $\overline{T^*}\mathcal{M}$ naturally (by taking their closures) and we let $L := E_s^* \cap \partial\overline{T^*}\mathcal{M}$ and $L' := E_u^* \cap \partial\overline{T^*}\mathcal{M}$. The set L is a radial source and L' a radial sink in the terminology of [DyZw, Definition E.50]. First, we will show that μ has no mass for ξ large enough in a conic neighborhood V_s of E_s^* . This can be proved using the radial point estimates of [DyZw, Theorem E.52]. There is $s_0 \in \mathbb{R}$ such that for all $s > s_0$ and $N_1 > 0$, and $B_1 \in \Psi_h^0(\mathcal{M})$ with $L \subset \text{ell}_h(B_1)$, there is $A_0 \in \Psi_h^0(\mathcal{M})$ elliptic in a neighborhood U of L in $\overline{T^*}\mathcal{M}$, and $C > 0$ so that for all $u \in H_h^s(\mathcal{M})$ such that $B_1 P_h(\lambda_h)u \in H_h^s$

$$\|A_0 u\|_{H_h^s} \leq Ch^{-1} \|B_1 P_h(\lambda_h)u\|_{H_h^s} + Ch^{N_1} \|u\|_{H_h^{-N_1}}. \quad (3.3)$$

Note that by (A1) we may choose $s_0 = N - \delta_0$ for some $\delta_0 > 0$. Now, since $u_h \in \mathcal{H}_h^{NG}$ and $P_h(\lambda_h)u_h \in \mathcal{H}_h^{NG}$, we can choose B_1 so that $\text{WF}_h(B_1) \subset V_s$ and, since $B_1 \text{Op}_h(e^{NG})^{-1} \in \Psi_h^{-N}(\mathcal{M})$ by the fact that $m = 1$ in V_s , we see that $B_1 P_h(\lambda_h)u_h \in H_h^N$ and $\|B_1 P_h(\lambda_h)u_h\|_{H_h^N} = o(h^\beta)$. We thus choose $s = N$ with $N > s_0$ and get that there is $A_0 \in \Psi_h^0(\mathcal{M})$ elliptic in a neighborhood U of L such that

$$\|A_0 u_h\|_{H_h^N} = o(h^{\beta-1}). \quad (3.4)$$

This shows that $\mu(\chi|\sigma(A_0)|^2) = 0$ for all $\chi \in C_c^\infty(T^*\mathcal{M})$ and thus μ vanishes in U . By the Anosov property of X (L is an attractor for the backward Hamiltonian flow $\Phi_{-t} = e^{-tH_p}$) for all $(x, \xi) \in T^*\mathcal{M} \setminus E_u^* \oplus E_0^*$, there is $T > 0$ such that $\Phi_{-T}(x, \xi) \in U$. We can then use the invariance (3.1) to deduce that $\mu = 0$ outside $E_u^* \oplus E_0^*$ so that (3.2) holds.

To prove that $\mu \neq 0$, we also need to have H_h^{-N} estimates on $Q u_h$ for some appropriate $Q \in \Psi_h^0(\mathcal{M})$ microsupported outside E_u^* .

For an operator $Y \in \Psi_h^m(\mathcal{M})$, denote by $Y^{(N)} := \text{Op}_h(e^{NG})Y \text{Op}_h(e^{NG})^{-1}$ the conjugated operator. By [FRS, Appendix] we then have $Y^{(N)} \in \Psi_h^m(\mathcal{M})$ and $\sigma_h(Y^{(N)}) = \sigma_h(Y)$. First, the elliptic estimate of [DyZw, Theorem E.33] gives that for each $A_1 \in \Psi_h^0(\mathcal{M})$ with $\text{WF}_h(A_1) \cap \{p = -1\} = \emptyset$ and each $N_1 > 0$, there is $C > 0$ so that for all $h > 0$

$$\|A_1^{(N)} \tilde{u}_h\|_{L^2} \leq C \|(P_h(\lambda_h))^{(N)} \tilde{u}_h\|_{L^2} + Ch^{N_1} \|u_h\|_{H_h^{-N}}. \quad (3.5)$$

Here we used the above facts for $Y = A_1, P_h(\lambda_h)$. Therefore, for each cone $\mathcal{C}_0 \subset \overline{T^*}\mathcal{M}$ satisfying $\mathcal{C}_0 \cap (E_u^* \oplus E_s^*) = \emptyset$, there is $R > 0$ large and $A_1 \in \Psi_h^0(\mathcal{M})$ satisfying $\sigma(A_1) = 1$ in $\mathcal{C}_0 \cap \{|\xi| > R\}$ such that, for some $N_1 > \beta$

$$\|A_1 u_h\|_{\mathcal{H}_h^{NG}} \leq C \|P_h(\lambda_h)u_h\|_{\mathcal{H}_h^{NG}} + Ch^{N_1} \|u_h\|_{H_h^{-N}} = o(h^\beta). \quad (3.6)$$

Next, we remark from the Anosov property of X that for each cone $\mathcal{C}_s \subset \overline{T^*}\mathcal{M} \setminus (E_u^* \oplus E_0^*)$ there is $T > 0$ so that $\Phi_{-T}(\mathcal{C}_s) \subset U$ where U is the neighborhood of L used before, we can then use the propagation of singularity estimate of [DyZw, Theorem E.47] to deduce the following. For each cone \mathcal{C}_s as above, there is $A_2 \in \Psi_h^0(\mathcal{M})$ with $\sigma(A_2) = 1$ on \mathcal{C}_s , $B \in \Psi_h^0(\mathcal{M})$ with $\text{WF}_h(B) \subset U$, so that for all $N_1 > 0$, there is a constant $C > 0$ such that

$$\|A_2 u_h\|_{\mathcal{H}_h^{NG}} \leq Ch^{-1} \|P_h(\lambda_h)u_h\|_{\mathcal{H}_h^{NG}} + C \|B u_h\|_{\mathcal{H}_h^{NG}} + Ch^{N_1} \|u_h\|_{H_h^{-N}}.$$

Note that, strictly speaking, we applied the propagation of singularities estimate to $(A_2)^{(N)}$, $(P_h(\lambda_h))^{(N)}$, $B^{(N)}$ and \tilde{u}_h , similarly to (3.5). Using (3.4) and elliptic estimates, $\|B u_h\|_{\mathcal{H}_h^{NG}} \leq$

$C\|A_0u_h\|_{H_h^N} + \mathcal{O}(h^\infty) = o(h^{\beta-1})$, and combining with (A1), we deduce that

$$\|A_2u_h\|_{\mathcal{H}_h^{NG}} = o(h^{\beta-1}). \quad (3.7)$$

Next, by [DyZw, Theorem E.54], there is s_1 such that for all $s < s_1$, $N_1 > 0$, there is $A_3 \in \Psi_h^0(\mathcal{M})$ elliptic near $L' := E_u^* \cap \partial \bar{T}^* \mathcal{M}$, $B' \in \Psi_h^0(\mathcal{M})$ with $\text{WF}_h(B') \cap L' = \emptyset$, $B_1 \in \Psi_h^0(\mathcal{M})$ elliptic on $\text{WF}_h(A_3)$ and $C > 0$ such that for all $v \in H_h^s(\mathcal{M})$ such that $B_1 P_h(\lambda_h)v \in H_h^s$, all $h > 0$

$$\|A_3v\|_{H_h^s} \leq C\|B'v\|_{H_h^s} + Ch^{-1}\|B_1P_h(\lambda_h)v\|_{H_h^s} + Ch^{N_1}\|v\|_{H_h^{-N}}. \quad (3.8)$$

Note that, by possibly applying another time the propagation of singularities estimate, one can choose B' to be microsupported also close to L' ; we can assume that A_3, B' are microsupported where $C_1\langle \xi \rangle^{-N} \leq e^{NG} \leq C_2\langle \xi \rangle^{-N}$. Note that by (A1), we may choose $s_1 = -N + \delta_1$ for some $\delta_1 > 0$. Let $\tilde{A}_3 \in \Psi_h^0(\mathcal{M})$ elliptic so that $\text{WF}_h(\tilde{A}_3 - A_3) \cap L' = \emptyset$, i.e. \tilde{A}_3 is microlocally equal to A_3 near L' . We can now write for some $N_1 > \beta$ and some $C, C' > 0$, using (3.8) with $v = u_h$ and $s = -N$ in the third line

$$\begin{aligned} 1 &= \|u_h\|_{\mathcal{H}_h^{NG}} \leq C\|\tilde{A}_3u_h\|_{\mathcal{H}_h^{NG}} + \mathcal{O}(h^{N_1}) \\ &\leq C\|A_3u_h\|_{H_h^{-N}} + C\|(\tilde{A}_3 - A_3)u_h\|_{\mathcal{H}_h^{NG}} + \mathcal{O}(h^{N_1}) \\ &\leq C'\|B'u_h\|_{H_h^{-N}} + C\|(\tilde{A}_3 - A_3)u_h\|_{\mathcal{H}_h^{NG}} + o(h^{\beta-1}). \end{aligned} \quad (3.9)$$

Note that B' and $(\tilde{A}_3 - A_3)$ satisfy the same property that their wavefront set does not intersect L' in $\bar{T}^* \mathcal{M}$. But if $B'' \in \Psi_h^0(\mathcal{M})$ is such that $\text{WF}_h(B'') \cap L' = \emptyset$, then one can choose \mathcal{C}_s and \mathcal{C}_0 above so that $\text{WF}_h(B'') \subset \mathcal{C}_s \cup \mathcal{C}_0 \cup \{|\xi| \leq 2R\}$ for some $R > 1$ large enough, and thus there exists $Q \in \Psi_h^{\text{comp}}(\mathcal{M})$ such that, for any $N_1 > 0$

$$\|B''u_h\|_{\mathcal{H}_h^{NG}} \leq C(\|B''A_1u_h\|_{\mathcal{H}_h^{NG}} + \|B''A_2u_h\|_{\mathcal{H}_h^{NG}} + \|B''Qu_h\|_{H_h^{NG}}) + Ch^{N_1}\|u_h\|_{H_h^{-N}}.$$

Applying this to $B'' = B'$ and $B'' = \tilde{A}_3 - A_3$ and using (3.7), (3.6) and (3.9), we deduce that there is $Q \in \Psi_h^{\text{comp}}(\mathcal{M})$ elliptic on $\{|\xi| \leq R\}$ and $C > 0$ such that

$$1 \leq C\|Qu_h\|_{\mathcal{H}_h^{NG}} + \mathcal{O}(h^{N_1}) + o(h^{\beta-1}).$$

Taking some $N_1 > \beta$ and since $\beta \geq 1$, the right hand side converges to 0 as $h \rightarrow 0$ and we deduce that $\mu(|\sigma(Q)|^2) > 0$, showing that $\mu(\{|\xi| \leq R\}) > 0$ for $R > 0$ large enough. Using the flow invariance (3.1) and the fact that e^{tH_p} is expanding linearly in ξ in the E_u^* direction where μ is supported, we deduce that $\mu(K_R) > 0$ for all $R > 0$. \square

We remark that the proof of this Lemma shows that $u = o(h^{\beta-1})$ microlocally on compact sets outside $\Gamma_+ \cap \{p = -1\}$ (use (3.5) and (3.7)).

4. UNSTABLE DERIVATIVES AND EXOTIC CALCULUS

If $V \in C^\infty(\mathcal{M})$ is a potential, the proof of [FaGu, Theorem 2] gives that there is $\alpha_V \in C_{\nu_{\max}^-}^{\nu_{\min}^-}(\mathcal{M})$ such that $(-X - r_-)\alpha_V = U_-(V)$ and

$$[-X + V, U_- + \alpha_V] = r_-(U_- + \alpha_V), \quad r_- \in C^{2^-}(\mathcal{M}), \quad X\alpha_V \in C_{\nu_{\max}^-}^{\nu_{\min}^-}(\mathcal{M}). \quad (4.1)$$

Recall that u is called a *generalized resonant state* at s if $(X + s)^k u = 0$ for some $k \geq 0$ and $u \in \mathcal{H}^N$ for some N large enough. We recall the result of [FaGu]:

Proposition 4.1 (Faure-Guillarmou [FaGu]). *If X is a contact 3-dimensional Anosov flow, the Ruelle generalized resonant states u with resonance in $\text{Re}(s) > -\nu_{\min}$ satisfy $U_-u = 0$.*

This extra property of resonant states will give us a new identity on the semiclassical measures associated to sequences of resonances: roughly speaking μ will be invariant by the Hamiltonian flow of $\sigma(U_-)$. More generally we can use this unstable derivative even for quasimodes. However, there is a technical drawback, which is that U_- is not smooth. For microlocal methods, this complicates the argument, and we will have to regularize U_- with an h -dependent scale. This leads us to use a slightly exotic class of pseudo-differential operators and symbols.

4.1. Regularization and exotic pseudo-differential calculus. If $m \in \mathbb{R}$, $\rho \in (0, 1)$ and $k \in \mathbb{R}^+$, we define the exotic pseudo-differential calculus $\Psi_{h,\rho,k}^m(\mathcal{M})$ to be the set of operators of the form $\text{Op}_h(a)$ where the symbol a in the class $S_{h,\rho,k}^m(T^*\mathcal{M})$, which is the space of smooth functions on $T^*\mathcal{M}$ such that in local coordinates (here $x_+ := \max(x, 0)$ for $x \in \mathbb{R}$)

$$\forall \alpha, \beta, \quad |\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \leq C_{\alpha,\beta,\varepsilon} \langle \xi \rangle^{m-|\beta|} h^{-\rho(|\alpha|-k)_+}. \quad (4.2)$$

We will also write $\Psi_{h,\rho}^m(\mathcal{M})$ when $k = 0$. The calculus defined by this property has all the good properties of a semiclassical pseudo-differential calculus, and we refer to Appendix A for a summary of the needed properties that we shall use freely in this section.

We want to work with smooth coefficients and will thus regularize Hölder functions at scale h^ρ as follows. We fix a partition of unity $(\psi_j)_j$ associated to a family of local charts $(\mathcal{O}_j)_j$ (i.e. $\text{supp}(\psi_j) \subset \mathcal{O}_j$), and denote by $(x_1, x_2, x_3) : \mathcal{O}_j \rightarrow \mathbb{R}^3$ the local coordinates in each chart. Without loss of generality, we shall choose the coordinate systems in the charts so that $X = \partial_{x_1}$. Then if $u \in C^k(\mathcal{M})$ for $k \geq 0$, define $u^h := \sum_j \psi_j u_j^h$ where $u_j = u|_{\mathcal{O}_j}$ and the h^ρ regularization in the chart \mathcal{O}_j , identified to an open set of \mathbb{R}^3 , is given by

$$\forall a \in L_{\text{comp}}^2(\mathbb{R}^3), \quad a^h(x) := h^{-3\rho} \int_{\mathbb{R}^3} \chi((x-x')/h^\rho) a(x') dx' \quad (4.3)$$

where $\chi \in C_c^\infty(\mathbb{R}^3; \mathbb{R}^+)$ has integral 1 and is chosen to be a radial function, i.e. a function of $|x|$. Note that $\psi_j u_j^h$ depends only on the value of u on an h^ρ -neighborhood of $\text{supp}(\psi_j)$. Notice also in that $1^h = 1$. Similarly for a semiclassical differential operator P with $C^k(\mathcal{M})$ coefficients, that is an operator which in each chart \mathcal{O}_j has the form

$$\sum_{\alpha} a_{j,\alpha}(x) h^{|\alpha|} \partial_x^\alpha$$

with $a_{j,\alpha} \in C^k(\mathcal{O}_j)$, we define $P^h = \sum_j \psi_j P_j^h$ where $P_j = P|_{\mathcal{O}_j}$, the regularization in each chart \mathcal{O}_j , is given by

$$P_j^h = \sum_{\alpha} a_{j,\alpha}^h(x) h^{|\alpha|} \partial_x^\alpha.$$

Moreover, if P is a differential operator (i.e. non-semiclassical), we define P^h analogously as an h -dependent differential operator. For instance, for a vector field Y we then have $(hY)^h = hY^h$. We notice that this process depends a priori on the charts and a partition of unity, but this will not cause us any problems. The following holds true:

Lemma 4.2. *Let $k \in (0, 2)$ and let $a \in C^k(\mathcal{M})$. For $m \in [0, k)$ integer and $\ell \in \mathbb{N}$, one has*

$$\|a^h - a\|_{C^m} = \mathcal{O}(h^{\rho(k-m)}) \|a\|_{C^k}, \quad \|a^h\|_{C^\ell} = \mathcal{O}(h^{-\rho(\ell-k)_+}),$$

$$\forall j, \quad \|\psi_j a_j^h - \psi_j a^h\|_{C^m} = \mathcal{O}(h^{(k-m)\rho}).$$

In particular one has $a^h \in S_{h,\rho,k}^0(\mathcal{M})$ and $\psi_j a_j^h = \psi_j a^h + \mathcal{O}_{S_{h,\rho,0}^0}(h^{\rho k})$. If $k < 1$ but $Xa \in C^k(\mathcal{M})$, one also has

$$Xa^h = Xa + \mathcal{O}_{C^0}(h^{k\rho}).$$

If P is a semiclassical differential operator of order ℓ on \mathcal{M} with $C^k(\mathcal{M})$ coefficients, then $P^h \in \Psi_{h,\rho,k}^\ell(\mathcal{M})$ and

$$P^h = P + \mathcal{O}_{C^0}(h^{\rho k}). \quad (4.4)$$

where the norm is the C^0 norm of the coefficients in the local bases ∂_{x_i} .

Proof. We work out the case $k \geq 1$, the case $k < 1$ is similar and indeed simpler. We write in the chart \mathcal{O}_j (on $\text{supp}(\psi_j)$)

$$a_j^h(x) = \int_{\mathbb{R}^3} \chi(y) a_j(x + h^\rho y) dy$$

and use that $a_j(x + h^\rho y) = a_j(x) + h^\rho da_j(x) \cdot y + \mathcal{O}_{C^0}(h^{k\rho} \|a\|_{C^k})$. But since χ is radial, one has $\int \chi(y) da_j(x) \cdot y dy = 0$, and this shows that

$$a_j^h(x) = a_j(x) + \mathcal{O}(h^{k\rho} \|a\|_{C^k}).$$

One has a similar estimate for the derivative

$$\partial_{x_i} a_j^h = \partial_{x_i} a_j + \mathcal{O}_{C^0}(h^{(k-1)\rho} \|a\|_{C^k}).$$

In general, for all $|\alpha| \geq 2$ we get $\partial_x^\alpha a_j^h = \mathcal{O}_{C^0}(h^{-\rho(|\alpha|-k)_+})$ since

$$\begin{aligned} \partial_x^\alpha a_j^h(x) &= (-1)^{|\alpha|} h^{-|\alpha|\rho} \int_{\mathbb{R}^3} \partial_y^\alpha \chi(y) a_j(x + h^\rho y) dy \\ &= (-1)^{|\alpha|-1} h^{(1-|\alpha|)\rho} \int_{\mathbb{R}^3} \partial_y^{\alpha'} \chi(y) \partial_{x_i} a_j(x + h^\rho y) dy \end{aligned}$$

for some α' satisfying $|\alpha'| = |\alpha| - 1$, with i so that $\alpha_i > 0$, and using that $\partial_{x_i} a_j(x + h^\rho y) = \partial_{x_i} a_j(x) + \mathcal{O}((h^\rho |y|)^{k-1})$ and $\int \partial_y^{\alpha'} \chi(y) dy = 0$. Thus, if $V \in C^\infty(\mathcal{M}; T\mathcal{M})$, one has

$$\begin{aligned} Va^h &= \sum_j (V(\psi_j) a_j^h + \psi_j V(a_j^h)) = \sum_j (V(\psi_j)(a_j^h - a_j) + \psi_j V(a_j^h - a_j) + V(\psi_j) a_j + \psi_j V(a_j)) \\ &= \sum_j (V(\psi_j) a_j + \psi_j V(a_j)) + \mathcal{O}_{C^0}(h^{(k-1)\rho}) = Va + \mathcal{O}_{C^0}(h^{(k-1)\rho}). \end{aligned}$$

Next, we also get from this analysis that for $m \leq k$, in \mathcal{O}_j

$$\partial_{x_i}^m (\psi_j (a_j^h - a_j)) = \psi_j \partial_{x_i}^m (a_j - a_j) + \partial_{x_i}^m (\psi_j) (a_j - a_j) + \mathcal{O}_{C^0}(h^{(k-m)\rho}) = \mathcal{O}_{C^0}(h^{(k-m)\rho}).$$

Next, if $a \in C^k$ with $Xa \in C^k$, by using that $X = \partial_{x_1}$ in each chart \mathcal{O}_j so that $X(a_j^h) = (Xa_j)^h$, we can write as above

$$Xa^h = \sum_j (X(\psi_j) a_j^h + \psi_j X(a_j^h)) = \sum_j (X(\psi_j)(a_j^h - a_j) + \psi_j (Xa_j)^h) = (Xa)^h + \mathcal{O}_{C^0}(h^{k\rho}),$$

where we used that $\sum_j \psi_j = 1$. The analysis above also implies that for a semi-classical differential operator P the full local symbol of P^h in charts satisfies

$$|\partial_x^\alpha \partial_\xi^\beta \sigma_{\text{full}}(P^h)(x, \xi)| \leq C_{\alpha, \beta} \langle \xi \rangle^{\ell - |\beta|} h^{-\rho(|\alpha|-k)_+}$$

and that (4.4) holds. \square

Applying the preceding lemma to hU_- , we get for any $\varepsilon > 0$

$$hU_-^h \in \Psi_{h, \rho, 2-\varepsilon}^1(\mathcal{M}), \quad hU_-^h = hU_- + \mathcal{O}_{C^0}(h^{1+\rho(2-\varepsilon)} \|U_-\|_{C^{2-\varepsilon}}). \quad (4.5)$$

For technical purposes, we need the following claim on the size of $(ab)^h - a^h b^h$:

Lemma 4.3. *Let $a \in C^k(\mathcal{M})$ and $b \in C^\ell(\mathcal{M})$ with $k, \ell \in (0, 2)$. Then, if $a_j = a|_{\mathcal{O}_j}$ and $b_j = b|_{\mathcal{O}_j}$, one has*

$$a^h b^h = (ab)^h + \mathcal{O}_{S_{h, \rho, 0}^0}(h^{\rho \cdot \min(k, \ell)}), \quad \psi_j a_j^h b_j^h = \psi_j (a_j b_j)^h + \mathcal{O}_{S_{h, \rho, 0}^0}(h^{\rho \cdot \min(k, \ell)}).$$

Proof. First we work in the chart \mathcal{O}_j . Using $b_j(x) = b_j^h(x) + \mathcal{O}(h^{\ell\rho})$ and $a_j(x) = a_j^h(x) + \mathcal{O}(h^{k\rho})$, the bound $|\partial_x^\alpha b_j^h| = \mathcal{O}(h^{-\rho(|\alpha|-\ell)_+})$ in $\text{supp}(\psi_j) \subset \mathcal{O}_j$, and the Taylor expansion

$$(a_j^h b_j^h)(x + h^\rho y) = (a_j^h b_j^h)(x) + d(a_j^h b_j^h)(x) \cdot h^\rho y + R(x, y),$$

where $R(\cdot, y) = \mathcal{O}_{S_{h,\rho,0}^0}(h^{\rho \cdot \min(k,\ell)})$ uniformly in y , we obtain

$$\begin{aligned} (a_j b_j)^h(x) &= \int_{\mathbb{R}^3} \chi(y) a_j(x + h^\rho y) b_j(x + h^\rho y) dy \\ &= \int_{\mathbb{R}^3} \chi(y) a_j^h(x + h^\rho y) b_j^h(x + h^\rho y) dy + h^{-3\rho} \int_{\mathbb{R}^3} \chi\left(\frac{x-x'}{h^\rho}\right) a_j(x') (b_j - b_j^h)(x') dx' \\ &\quad + h^{-3\rho} \int_{\mathbb{R}^3} \chi\left(\frac{x-x'}{h^\rho}\right) (a_j - a_j^h)(x') b_j^h(x') dx' \\ &= a_j^h(x) b_j^h(x) + h^\rho d(a_j^h b_j^h)(x) \cdot \underbrace{\int_{\mathbb{R}^3} y \chi(y) dy}_{=0} + \mathcal{O}_{S_{h,\rho,0}^0}(h^{\rho \cdot \min(k,\ell)}). \end{aligned}$$

This gives the local result by using that χ is radial (so the h^ρ factor vanishes). Now for the global result, by the first part of the lemma

$$(ab)^h = \sum_j \psi_j a_j^h b_j^h + \mathcal{O}_{S_{h,\rho,0}^0}(h^{\rho \cdot \min(k,\ell)}) = a^h b^h + \sum_j \psi_j a_j^h (b_j^h - b^h) + \mathcal{O}_{S_{h,\rho,0}^0}(h^{\rho \cdot \min(k,\ell)})$$

which yields the result since $\psi_j (b_j^h - b^h) = \mathcal{O}_{S_{h,\rho,0}^0}(h^{\rho\ell})$ by Lemma 4.2. \square

We next compute the regularized commutator of $h(X - V)$ and $h(U_- + \alpha_V)^h$ where α_V is the function appearing in (4.1):

Lemma 4.4. *Assume $\beta_0 \in (0, \frac{\nu_{\min}}{\nu_{\max}})$ and $\rho > \frac{1}{2-\beta_0}$. If $V \in C^\infty(\mathcal{M})$ is h -independent, the following commutation relation holds*

$$[h(X - V), h(U_- + \alpha_V)^h] = -r_-^h h^2 (U_- + \alpha_V)^h + h^{2+\beta_0\rho} R_h,$$

where the upper index h denotes the regularization at scale h^ρ and $R_h \in \Psi_{h,\rho,0}^1(\mathcal{M})$.

Proof. For $V \in C^\infty(\mathcal{M})$, let $U_-^V := U_- + \alpha_V$, then

$$[h(X - V), h(U_-^V)^h] = \sum_j \psi_j [h(X - V), h(U_{-,j}^V)^h] + h^2 X(\psi_j) (U_{-,j}^V)^h$$

where $U_{-,j}^V = U_-^V|_{\mathcal{O}_j}$. We start by computing $[h(X - V), h(U_{-,j}^V)^h]$ in \mathcal{O}_j . Writing $X = \partial_{x_1}$ as before and $U_{-,j} = \sum_i a_i(x) \partial_{x_i}$ in \mathcal{O}_j we have, using $\partial_{x_i} a^h = (\partial_{x_i} a)^h$,

$$[h(X - V), h(U_{-,j}^h + \alpha_{V,j}^h)] = h^2 \left(X(\alpha_{V,j}^h) + \sum_{i=1}^3 \left((X a_i)^h \partial_{x_i} + a_i^h \partial_{x_i}(V) \right) \right).$$

Recall by (4.1) that $X\alpha_V + U_-(V) = -r_- \alpha_V$ and that $\alpha_V, X\alpha_V \in C^{\beta_0}(\mathcal{M})$ for all $\beta_0 \in (0, \nu_{\min}/\nu_{\max})$; we can then use Lemma 4.3 and Lemma 4.2 to deduce that on $\text{supp}(\psi_j)$, for any $\varepsilon > 0$

$$\begin{aligned} [h(X - V), h(U_{-,j}^h + \alpha_{V,j}^h)] &= h^2 [X, U_-]_j^h + h^2 (X\alpha_V)_j^h + h^2 (U_-(V))_j^h + \mathcal{O}_{\Psi_{h,\rho,0}^0}(h^{2+\rho(2-\varepsilon)}) \\ &= -h^2 r_{-,j}^h (U_{-,j}^h + \alpha_{V,j}^h) + \mathcal{O}_{\Psi_{h,\rho,0}^0}(h^{2+\rho\beta_0}) + \mathcal{O}_{\Psi_{h,\rho,0}^1}(h^{1+\rho(2-\varepsilon)}). \end{aligned}$$

Now $(U_{-,j}^V)^h = (U_{-,j})^h + \alpha_{V,j}^h$ with $h^2\psi_j(U_{-,j})^h \in h\Psi_{h,\rho,2-\varepsilon}^1(\mathcal{M})$ and $h^2\psi_j\alpha_{V,j}^h \in h^2\Psi_{h,\rho,\beta_0}^0(\mathcal{M})$ satisfy, using again that $\sum_j \psi_j = 1$ and Lemma 4.2, for any $\varepsilon > 0$

$$\begin{aligned} \sum_j hX(\psi_j)\alpha_{V,j}^h &= \sum_j hX(\psi_j)[\alpha_{V,j}^h - \alpha_V] = \mathcal{O}_{C^0}(h^{1+\beta_0\rho}), \\ \sum_j hX(\psi_j)(U_{-,j})^h &= \sum_j hX(\psi_j)[(U_{-,j})^h - U_-] = \mathcal{O}_{C^\ell}(h^{1+\rho(2-\varepsilon-\ell)}), \end{aligned}$$

for $\ell = 0, 1$, where C^ℓ denotes the norm on vector fields. In particular, $\sum_j hX(\psi_j)(U_{-,j})^h \in h^{\rho(2-\varepsilon)}\Psi_{h,\rho,0}^1(\mathcal{M})$ and $\sum_j hX(\psi_j)\alpha_{V,j}^h \in h^{1+\beta_0\rho}\Psi_{h,\rho,0}^0(\mathcal{M})$. Thus the error term belongs to $h^{\min(2+\rho\beta_0, 1+\rho(2-\varepsilon))}\Psi_{h,\rho,0}^1(\mathcal{M})$ and using the assumption on ρ completes the proof. \square

4.2. A propagation estimate and regularity of the semiclassical measure in the unstable direction. In this section, we are going to use the regularized version of the commutation formula $[X, U_-] = -r_-U_-$ proved in Lemma 4.4 to deduce that the semiclassical measure μ enjoys some extra regularity in the (lifted) unstable direction.

Before proceeding, we define the semiclassical principal symbol of $-ihU_\pm$

$$p_\pm(x, \xi) := \sigma(-ihU_\pm)(x, \xi) = \xi(U_\pm(x)).$$

Recall here U_- spans E_u and U_+ spans E_s , so we have

$$p_-^{-1}(0) = E_0^* \oplus E_u^* = \Gamma_+, \quad p_+^{-1}(0) = E_0^* \oplus E_s^* = \Gamma_-.$$

Since $p_- \in C^{2-}$, the Hamiltonian vector field H_{p_-} has C^{1-} coefficients, which would a priori not be sufficient to define its flow as it is not Lipschitz. However, since this is a Hamiltonian vector field, the flow equation in local coordinates reads

$$\dot{x}(t) = U_-(x(t)), \quad \dot{\xi}(t) = -\partial_x U_-(x(t)) \cdot \xi(t). \quad (4.6)$$

Since $U_- \in C^{2-}$, we see that the first equation with initial condition $x(0) = x_0$ has a unique solution given by the (horocycle) flow $\phi_t(x_0) = e^{tU_-}(x_0)$ of U_- , and $(t, x) \mapsto \phi_t(x)$ is C^{2-} but it is C^2 in the t -variable. Since $x(t)$ is well-defined (in a unique way), the second equation for $\xi(t)$ with $\xi(0) = \xi_0$ can now obviously be solved for each x since it is linear, the solution is unique, and it is given by $\xi(t) = (d\phi_t(x)^{-1})^T \xi$. We will thus define $e^{tH_{p_-}}$ to be the symplectic lift of ϕ_t

$$e^{tH_{p_-}}(x, \xi) := (\phi_t(x), (d\phi_t(x)^{-1})^T \xi)$$

which is the unique solution of (4.6) with initial condition $(x, \xi) \in T^*\mathcal{M}$.

Let $\delta > 0$ be small and define

$$\mathcal{U}_\delta := \{(x, \xi) \in T^*\mathcal{M} : |p_-(x, \xi)| < \delta, |p_+(x, \xi)| < \delta, |p(x, \xi) + 1| < \delta\}. \quad (4.7)$$

We now state a technical propagation Lemma, which is comparable to [Dy2, Lemma 2.7].

Lemma 4.5. *Let $k \geq 1$. Fix $\delta > 0$ small and assume $A, B, B_1 \in \Psi_h^{\text{comp}}(\mathcal{M})$ satisfy (see Figure 2)*

1. $\text{WF}_h(A) \subset \mathcal{U}_{3\delta/2}$ and $A = 1 + \mathcal{O}(h^\infty)$ on $\overline{\mathcal{U}_\delta}$.
2. $\text{WF}_h(B) \subset \mathcal{U}_{3\delta} \cap \{|p_-| > \delta/2\}$ and $B = 1 + \mathcal{O}(h^\infty)$ on $\overline{\mathcal{U}_{2\delta}} \cap \{|p_-| \geq \delta\}$.
3. $\text{WF}_h(B_1) \subset \mathcal{U}_{3\delta}$ and $B_1 = 1 + \mathcal{O}(h^\infty)$ on $\overline{\mathcal{U}_{2\delta}}$.

Let $W \in \Psi_{h,\rho,k}^0(\mathcal{M})$ with real principal symbol $\sigma(W)$ so that $\sigma(W) \rightarrow \sigma(W)_0 \in C^0(T^*\mathcal{M})$ uniformly as $h \rightarrow 0$, and assume that for $T \gg 1$ and some $c > 0$

$$\frac{1}{T} \int_0^T (\sigma(W)_0 - V - \gamma) \circ e^{-tH_p} dt \geq c \quad \text{on } \mathcal{U}_\delta \cap \Gamma_+. \quad (4.8)$$

Then for all $u_h \in \mathcal{D}'(\mathcal{M})$ and all $N_1 \gg 1$, there is a $C > 0$ such that

$$\|Au_h\|_{L^2} \leq Ch^{-1} \|B_1(P_h(\lambda_h) - ihW)u_h\|_{L^2} + C \|Bu_h\|_{L^2} + Ch^{N_1} \|u_h\|_{H_h^{-N_1}}. \quad (4.9)$$




FIGURE 2. The wavefront set setup in Lemma 4.5. In shaded blue we have $A = 1$, in red is $B = 1$, in yellow is $B_2 = 1$, in gray we have $B_1 = 1$, while the dark green rectangle bounds the set where $B'_2 = 1$, all taken modulo $\mathcal{O}(h^\infty)$.

Proof. We split the proof in two steps.

Step 1. Take $B_2 \in \Psi_h^{\text{comp}}(\mathcal{M})$ such that $\text{WF}_h(B_2) \subset \mathcal{U}_{2\delta}$ and $B_2 = 1 + \mathcal{O}(h^\infty)$ on $\mathcal{U}_{3\delta/2}$. We claim that, for every fixed $\varepsilon_0 > 0$, there is $C > 0$ such that for all $u \in L^2(\mathcal{M})$

$$\|Au\|_{L^2} \leq Ch^{-1}\|B_2(P_h(\lambda_h) - ihW)u\|_{L^2} + C\|Bu\|_{L^2} + \varepsilon_0\|u\|_{L^2}. \quad (4.10)$$

We prove this by contradiction: assume there is a sequence $u_h \in L^2(\mathcal{M})$ with $h \rightarrow 0$ and $\|u_h\|_{L^2} = 1$, such that

$$\|B_2(P_h(\lambda_h) - ihW)u_h\|_{L^2} = o(h), \quad \|Bu_h\|_{L^2} = o(1), \quad \|Au_h\|_{L^2} \geq \varepsilon_0. \quad (4.11)$$

By [DyZw, Theorem E. 42] there is a semiclassical (Radon) measure ν associated to u_h . Now use propagation of singularities estimate Proposition A.2 (item 8): for every $(x, \xi) \in \mathcal{U}_{3\delta/2} \setminus \Gamma_+$, there is a $t_0 \geq 0$ with $e^{-t_0 H_p}(x, \xi) \in \{\sigma(B) = 1\}$ and $e^{-t H_p}(x, \xi) \in \mathcal{U}_{3\delta/2}$ for $t \in [0, t_0]$. Thus for all $Q \in \Psi_h^{\text{comp}}(\mathcal{M})$ with $\text{WF}_h(Q) \subset \mathcal{U}_{3\delta/2} \setminus \Gamma_+$,

$$\|Qu\|_{L^2} \leq C\|Bu\|_{L^2} + Ch^{-1}\|B_2(P_h(\lambda_h) - ihW)u\|_{L^2} + \mathcal{O}(h^\infty).$$

By (4.11) we obtain $\|Qu\|_{L^2} = o(1)$ and so $\nu = 0$ on $\mathcal{U}_{3\delta/2} \setminus \Gamma_+$.

Next, (4.11) implies $(P_h(\lambda_h) - ihW)u = o(h)$ microlocally in $\mathcal{U}_{3\delta/2}$, so by Proposition A.4 we have for all $a \in C_0^\infty(\mathcal{U}_{3\delta/2})$

$$\int_{T^*\mathcal{M}} H_p a \, d\nu = 2 \int_{T^*\mathcal{M}} (\sigma(W)_0 - V - \gamma) a \, d\nu.$$

Set $f := 2(\sigma(W)_0 - V - \gamma)$. Equivalently, since for $t \geq 0$ we have $e^{-t H_p} : \mathcal{U}_{3\delta/2} \cap \Gamma_+ \rightarrow \mathcal{U}_{3\delta/2} \cap \Gamma_+$ and since ν is zero on $\mathcal{U}_{3\delta/2} \setminus \Gamma_+$, we may write an evolution equation for $\nu' = \nu|_{\mathcal{U}_{3\delta/2} \cap \Gamma_+}$,

$$(e^{-t H_p})^* d\nu' = e^{\int_0^t f \circ e^{-r H_p} dr} d\nu', \quad t \geq 0. \quad (4.12)$$

Applying this relation to $0 \leq a \in C_0^\infty(\mathcal{U}_{3\delta/2})$ and as ν is a Radon measure, we get for $t \gg 1$

$$\int_{\mathcal{U}_{3\delta/2} \cap \Gamma_+} a \circ e^{t H_p} d\nu = \int_{\mathcal{U}_{3\delta/2} \cap \Gamma_+} e^{\int_0^t f \circ e^{-r H_p} dr} a \, d\nu \geq e^{2ct} \int_{\mathcal{U}_{3\delta/2} \cap \Gamma_+} a \, d\nu,$$

by the condition (4.8). The left hand side of this equation is bounded from above by $\|a\|_{L^\infty} \nu(\mathcal{U}_{3\delta/2})$, while the right hand side is growing exponentially fast. Thus, $\nu \equiv 0$ on $\mathcal{U}_{3\delta/2}$, contradicting the last point of (4.11) and proving the claim.

Step 2. For every $B'_2 \in \Psi_h^{\text{comp}}(\mathcal{M})$ with $B'_2 = 1 + \mathcal{O}(h^\infty)$ microlocally on $\text{WF}_h(B_2)$, we may apply inequality (4.10) to $B'_2 u$ to get

$$\|Au\|_{L^2} \leq Ch^{-1} \|B_2(P_h(\lambda_h) - ihW)u\|_{L^2} + C\|Bu\|_{L^2} + \varepsilon_0 \|B'_2 u\|_{L^2} + Ch^{N_1} \|u\|_{H_h^{-N_1}}. \quad (4.13)$$

Choose B'_2 such that for every $(x, \xi) \in \text{WF}_h(B'_2)$, there is a $t_0 \geq 0$ with $e^{-t_0 H_p}(x, \xi) \in \{\sigma(A) \neq 0\} \cup \{\sigma(B) \neq 0\}$, and for all $t \in [0, t_0]$ we have $e^{-t H_p}(x, \xi) \in \{\sigma(B_1) \neq 0\}$. Thus by the propagation of singularities estimate, Proposition A.2 (item 8), we obtain

$$\|B'_2 u\|_{L^2} \leq Ch^{-1} \|B_1(P_h(\lambda_h) - ihW)u\|_{L^2} + C\|Bu\|_{L^2} + C\|Au\|_{L^2} + Ch^{N_1} \|u\|_{H_h^{-N_1}}. \quad (4.14)$$

By using (4.14) to estimate the $\|B'_2 u\|_{L^2}$ term in (4.13), the elliptic estimate [DyZw, Theorem E.33] and the fact that $\text{WF}_h(B_2) \subset \text{WF}_h(B_1)$

$$\|Au\|_{L^2} \leq Ch^{-1} \|B_1(P_h(\lambda_h) - ihW)u\|_{L^2} + C\|Bu\|_{L^2} + \varepsilon_0 C \|Au\|_{L^2} + Ch^{N_1} \|u\|_{H_h^{-N_1}}.$$

Taking ε_0 small enough, we absorb the $\varepsilon_0 C \|Au\|_{L^2}$ term to the left hand side, thus completing the proof. \square

We will now use this propagation estimate to deduce some regularity of the semiclassical measure μ in the direction of H_{p_-} . Recall the definition of V_{\min}, V_{\max} in (1.2).

Proposition 4.6. *Assume $\gamma < \nu_{\min} - V_{\max}$ in (A1). If additionally $\beta \geq 2$ in (A1), for $\delta > 0$ small enough and any $a \in C_c^\infty(\mathcal{U}_\delta)$, we have*

$$\int_{\mathcal{U}_\delta} (H_{p_-} a - (-\text{div } U_- + 2\alpha_V) a) d\mu = 0, \quad (4.15)$$

where div denotes the divergence with respect to the contact measure $\alpha \wedge d\alpha$. Moreover, if (A1) holds with $\|P_h(\lambda_h)u_h\|_{\mathcal{H}_h^{NG}} = \mathcal{O}(h^2)$ there is a $C > 0$ such that

$$\left| \int_{\mathcal{U}_\delta} H_{p_-}(a) d\mu \right| \leq C \|a\|_{L^\infty}. \quad (4.16)$$

Proof. Denoting $U_-^V := U_- + \alpha_V$, the commutation relation in Lemma 4.4 reads:

$$[h(X - V), h(U_-^V)^h] = -r_-^h h^2 (U_-^V)^h + h^{2+\rho\beta_0} R_h, \quad R_h \in \Psi_{h,\rho}^1(\mathcal{M}),$$

for some $\beta_0 > 0$. Therefore, we have the relation

$$(P_h(\lambda_h) - i h r_-^h) h(U_-^V)^h = h(U_-^V)^h P_h(\lambda_h) - i h^{2+\beta_0\rho} R_h. \quad (4.17)$$

Next, for each $\varepsilon > 0$ so that $\nu_{\min} - V_{\max} - \gamma > 3\varepsilon$, there is $T_0 > 0$ such that for all $T > T_0$

$$\frac{1}{T} \int_0^T (r_- - V) \circ \varphi_{-t} dt > \nu_{\min} - V_{\max} - 2\varepsilon > \gamma + \varepsilon.$$

Thus, we can apply Lemma 4.5 with the functions $h(U_-^V)^h u_h \in H_h^{-N-1}(\mathcal{M})$ and with $W := r_-^h \in \Psi_{h,\rho,2-\varepsilon}^0(\mathcal{M})$, which gives

$$\|Ah(U_-^V)^h u_h\|_{L^2} \leq Ch^{-1} \|B_1(P_h(\lambda_h) - i h r_-^h) h(U_-^V)^h u_h\|_{L^2} + C \|Bh(U_-^V)^h u_h\|_{L^2} + \mathcal{O}(h^\infty) \|h(U_-^V)^h u_h\|_{H_h^{-N-1}}, \quad (4.18)$$

where $A, B, B_1 \in \Psi_h^{\text{comp}}(\mathcal{M})$ satisfy the conditions of Lemma 4.5.

We analyse (4.18) by studying each term on the right hand side separately. Firstly, since $\text{WF}_h(B)$ does not intersect $\Gamma_+ \cap \{p = -1\}$, using the remark after Lemma 3.3, we have

$$\|Bh(U_-^V)^h u_h\|_{L^2} = o(h^{\beta-1}), \quad h \rightarrow 0.$$

Here we also used that $h(U_-^V)^h \in \Psi_{h,\rho,0}^1(\mathcal{M})$ is suitably bounded by Proposition A.2 (items 1 and 6). By the same boundedness properties

$$\|h(U_-^V)^h u_h\|_{H_h^{-N-1}} \leq C \|u_h\|_{H_h^{-N}} = \mathcal{O}(1),$$

which shows that the last term of (4.18) equals $\mathcal{O}(h^\infty)$. Finally, for the first term it suffices to estimate, by (4.17)

$$\begin{aligned} h^{-1} \|B_1(h(U_-^V)^h P_h(\lambda_h) - ih^{2+\rho\beta_0} R_h)u_h\|_{L^2} &\leq h^{-1} \|B_1 h(U_-^V)^h P_h(\lambda_h)u_h\|_{L^2} + h^{1+\rho\beta_0} \|B_1 R_h u_h\|_{L^2} \\ &\leq h^{-1} \|P_h(\lambda_h)u_h\|_{H_h^{-N}} + h^{1+\rho\beta_0} \|u_h\|_{H_h^{-N}} + \mathcal{O}(h^\infty) \\ &= o(h^{\beta-1}) + \mathcal{O}(h^{1+\rho\beta_0}) + \mathcal{O}(h^\infty). \end{aligned}$$

We used that $\|R_h\|_{H_h^1 \rightarrow L^2} = \mathcal{O}(1)$ by $\sigma(R_h) = \mathcal{O}(1)$ and Proposition A.2 (item 6). Therefore, by (4.18)

$$\|Ah(U_-^V)^h u_h\|_{L^2} = o(h^{\beta-1}) + \mathcal{O}(h^{1+\rho\beta_0}),$$

and so $h(U_-^V)^h u_h = o(h)$ microlocally in \mathcal{U}_δ if $\beta = 2$. Now note that

$$h^{-1} \operatorname{Im}(-ih(U_-^V)^h) = \frac{-ih(U_-^V)^h - (-ih(U_-^V)^h)^*}{2ih} = \frac{1}{2} \operatorname{div} U_-^h - \alpha_V^h = \mathcal{O}_{\Psi_{h,\rho,0}^0(\mathcal{M})}(1),$$

by Lemma 4.2. The main result (4.15) then follows by Proposition A.4, since $\operatorname{div} U_-^h \rightarrow \operatorname{div} U_-$, $\alpha_V^h \rightarrow \alpha_V$ and $H_{\sigma(-ih(U_-^V)^h)} a \rightarrow H_{p_-} a$ uniformly.

Finally, if $\|P_h(\lambda_h)u_h\|_{\mathcal{H}_h^{NG}} = \mathcal{O}(h^2)$, a similar argument gives $h(U_-^V)^h u_h = \mathcal{O}(h)$ microlocally in \mathcal{U}_δ . The final conclusion follows again by applying Proposition A.4. \square

We now show that Proposition 4.6 implies some Lipschitz regularity of μ inside Γ_+ .

Lemma 4.7. *Assume that $\gamma < \nu_{\min} - V_{\max}$ in (A1). If additionally $\beta \geq 2$ in (A1), then for $\delta > 0$ small enough, there is a $C > 0$ such that for every $\delta_0 > 0$ small enough*

$$\frac{\delta_0}{C} \leq \mu(\mathcal{U}_\delta \cap \{|p_+| < \delta_0\}) \leq C\delta_0. \quad (4.19)$$

Moreover, if (A1) is valid with $\|P_h(\lambda_h)u_h\|_{\mathcal{H}_h^{NG}} = \mathcal{O}(h^2)$, then the upper bound in (4.19) holds.

Proof. First, we make the following observation using the contact structure α : we have

$$H_{p_-}(p_+)(x, \xi) = \xi([U_-, U_+](x)), \quad (4.20)$$

and $[U_-, U_+]$ is a C^{1-} vector field that does not vanish such that $0 \neq \alpha([U_-, U_+]) = -d\alpha(U_-, U_+)$ (as $U_\pm \in \ker \alpha$), by the relation

$$0 \neq \alpha \wedge d\alpha(X, U_-, U_+) = 2d\alpha(U_-, U_+). \quad (4.21)$$

This means that either $H_{p_-}(p_+)(x, \xi) > c_1$ or $-H_{p_-}(p_+)(x, \xi) > c_1$ for some $c_1 > 0$ on \mathcal{U}_δ if $\delta > 0$ is small enough. Let $\tilde{a} \in C_c^\infty(\mathbb{R})$ with $\operatorname{supp}(\tilde{a}) \subset (-2\delta_0, \delta_1)$ satisfying

$$\|\tilde{a}\|_\infty \leq 1, \quad \partial_s \tilde{a} \geq -\frac{2}{\delta_1}, \quad \partial_s \tilde{a} \geq \frac{1}{3\delta_0} \quad \text{for } |s| \leq \delta_0, \quad (4.22)$$

for some small fixed $\delta_1 \in (0, \delta)$, independent of δ_0 (here we take $\delta_0 < \delta_1/2$). Let $\chi \in C_c^\infty(-\delta, \delta)$ be equal to 1 in $(-\delta/2, \delta/2)$. Consider $a(x, \xi) := \tilde{a}(p_+(x, \xi))\chi(p_-(x, \xi))\chi(p(x, \xi) + 1)$ which is a $C_c^{2-}(\mathcal{U}_\delta)$ function. If $H_{p_-}(p_+) > c_1$ on \mathcal{U}_δ , using that $\operatorname{supp}(\mu) \subset (\Gamma_+ \cap \{p = -1\}) \subset \{|p_-| \leq \delta/2, |p_+ + 1| \leq \delta/2\}$, we get

$$\int_{\mathcal{U}_\delta} H_{p_-}(a) d\mu = \int_{\mathcal{U}_\delta} \partial_s \tilde{a}(p_+) H_{p_-}(p_+) d\mu \geq \frac{c_1}{3\delta_0} \mu(\{|p_+| \leq \delta_0\}) - \frac{2c_1}{\delta_1} \mu(\{|p_+| \in (\delta_0, \delta_1)\}).$$

The result of Proposition 4.6 (in the case $\|P_h(\lambda_h)u_h\|_{\mathcal{H}_h^{NG}} = \mathcal{O}(h^2)$ as well) also holds for $a \in C_c^1(\mathcal{U}_\delta)$ by using a density argument, so we can apply it to our function a and, since $\mu(\{|p_+| \in (\delta_0, \delta_1)\}) \leq \mu(\mathcal{U}_\delta)$, we get that there is a $C > 0$ such that for all $\delta_0 > 0$ small

$$\mu(\{|p_+| \leq \delta_0\}) \leq C\delta_0.$$

In the case where $H_{p_-}(p_+)(x, \xi) < 0$ we can do the same reasoning by bounding below the integral $-\int_{\mathcal{U}_\delta} H_{p_-}(a) d\mu$.

To obtain a lower bound for the measure of $\mathcal{U}_\delta \cap \{|p_+| \leq \delta_0\}$, we can proceed as follows. Without loss of generality, we can assume that $c_2 > H_{p_-}(p_+) > c_1 > 0$ as above in \mathcal{U}_δ – this precisely means that p_+ is increasing along $e^{tH_{p_-}}$. Moreover, we compute $H_{p_-}p(x, \xi) = \xi([U_-, X](x)) = r_-(x)p_-(x, \xi)$, and thus both p and p_- are constant along $e^{tH_{p_-}}$ on Γ_+ .

We integrate (4.15) to get, for each $a \in C_c^\infty(\mathcal{U}_{\delta/2})$ and $|t| \leq \frac{\delta}{2c_2}$ (so $\text{supp}(a \circ e^{tH_{p_-}}) \subset \mathcal{U}_\delta$)

$$\int_{\mathcal{U}_\delta} a \circ e^{tH_{p_-}} d\mu = \int_{\mathcal{U}_\delta} e^{\int_0^t (-\text{div}(U_-) + 2\alpha_V) \circ e^{-sH_{p_-}} ds} a d\mu. \quad (4.23)$$

We thus get, using that $|-\text{div}(U_-) + 2\alpha_V|$ is uniformly bounded, that there is $C > 0$ independent of δ_0 such that

$$\int_0^{\frac{\delta}{2c_2}} \int_{\mathcal{U}_\delta} a \circ e^{tH_{p_-}} d\mu dt \leq C \int_{\mathcal{U}_\delta} a d\mu. \quad (4.24)$$

We choose $a = \tilde{a}(p_+)\chi(p+1)\chi(p_-)$ with χ as above, $\tilde{a} \in C_c^\infty((-\delta_0, \delta_0); \mathbb{R}^+)$ satisfying $\tilde{a} = 1$ on $\{|p_+| \leq \delta_0/2\}$. For each $z \in \mathcal{U}_{\delta/2}$, the map $t \mapsto p_+(e^{tH_{p_-}}(z))$ is a C^1 -diffeomorphism for $|t| \leq \frac{\delta}{2c_2}$ since $H_{p_-}(p_+) \in [c_1, c_2]$. Thus we can perform the change of variable $q = p_+(e^{tH_{p_-}}(z))$

$$\begin{aligned} \int_0^{\frac{\delta}{2c_2}} \int_{\mathcal{U}_{\delta/2}} a(e^{tH_{p_-}}(z)) d\mu(z) dt &= \int_{\mathcal{U}_{\delta/2}} \int_{p_+(z)}^{p_+(e^{\frac{\delta}{2c_2}H_{p_-}}(z))} \frac{\tilde{a}(q)}{H_{p_-}(p_+)(e^{t(q)H_{p_-}}(z))} dq d\mu(z) \\ &\geq \frac{1}{c_2} \int_{\mathcal{U}_{\delta/2}} \int_{p_+(z)}^{p_+(e^{\frac{\delta}{2c_2}H_{p_-}}(z))} \tilde{a}(q) dq d\mu(z) \\ &\geq \frac{\delta_0}{c_2} \underbrace{\mu(\{z \in \mathcal{U}_{\delta/2} \mid (-\delta_0/2, \delta_0/2) \subset [p_+(z), p_+(e^{\frac{\delta}{2c_2}H_{p_-}}(z))]\})}_{A:=}. \end{aligned}$$

Since $H_{p_-}p_+ \in [c_1, c_2]$ on \mathcal{U}_δ , the set A contains $p_+^{-1}([\frac{\delta_0}{2} - \frac{\delta c_1}{2c_2}, \frac{\delta_0}{2}])$. Note first that by the Lipschitz bound and Lemma 3.3, for any $\delta > 0$ we have $\mu(\mathcal{U}_\delta \setminus E_0^*) > 0$. Then the horocyclic invariance (4.23) implies that for all $R_1, R_2 \in \mathbb{R}$ small enough, the strip $S_{R_1, R_2} = \{z \in T^*\mathcal{M} \mid R_1 \leq p_- \leq R_2\}$ satisfies $\mu(S_{R_1, R_2}) > 0$. Thus for δ_0 small enough, $\mu(A) \geq C'$ for some constant $C' > 0$ depending only on c_1, c_2 and δ (and not on δ_0). We therefore obtain, by combining with (4.24), that there is $C'' > 0$ such that for all $\delta_0 > 0$ small

$$\mu(\mathcal{U}_\delta \cap \{|p_+| \leq \delta_0\}) \geq \int_{\mathcal{U}_\delta} a d\mu \geq C''\delta_0,$$

concluding the proof. \square

5. PROOF OF THE MAIN THEOREM

We proceed to the proof of the main result. Note that Theorem 1 is an immediate corollary of Theorem 2 for the case $V = 0$.

Proof of Theorem 2. We proceed by contradiction. Assume that the estimates in (1.4) do not hold, i.e. there is a sequence $h_n \rightarrow 0$, $\lambda_n \in \mathbb{C}$ and $u'_n, f_n \in \mathcal{H}_h^{NG}$ with norm $\|f_n\|_{\mathcal{H}_h^{NG}} = 1$, such that $(-ih_n X + ih_n V - i\lambda_n)u'_n = f_n$ satisfies $\|u'_n\|_{\mathcal{H}_h^{NG}} > Ch_n^{-2}$ for some $C > 0$ (resp. $\|u'_n\|_{\mathcal{H}_h^{NG}} >$

nh_n^{-2}) if $h_n^{-1} \operatorname{Re} \lambda_n \in \mathcal{S}_0(\varepsilon)$ (resp. $h_n^{-1} \operatorname{Re} \lambda_n \in \mathcal{S}_1(\varepsilon)$). Up to extracting a subsequence and rescaling h_n , we can assume that $\operatorname{Im}(\lambda_n) \rightarrow 1$ and $h_n^{-1} \operatorname{Re}(\lambda_n) \rightarrow -\gamma$ as $n \rightarrow \infty$ with $-\gamma \in \mathcal{S}_0(\varepsilon)$ (resp. $-\gamma \in \mathcal{S}_1(\varepsilon)$). Define $u_n := u'_n / \|u'_n\|_{\mathcal{H}_h^{NG}} \in \mathcal{H}_h^{NG}$ which satisfies

$$\|P_{h_n}(\lambda_n)u_n\|_{\mathcal{H}_h^{NG}} = \frac{\|f_n\|_{\mathcal{H}_h^{NG}}}{\|u'_n\|_{\mathcal{H}_h^{NG}}} = \begin{cases} \mathcal{O}(h_n^2), & -\gamma \in \mathcal{S}_0(\varepsilon), \\ o(h_n^2), & -\gamma \in \mathcal{S}_1(\varepsilon). \end{cases}$$

This means that we are in the case of (A1) with $\|P_h(\lambda_h)u_h\|_{\mathcal{H}_h^{NG}} = \mathcal{O}(h^2)$ and $N \geq \frac{1}{2}$, if $-\gamma \in \mathcal{S}_0(\varepsilon)$, or with $\beta = 2$ and $N \geq 1$, if $-\gamma \in \mathcal{S}_1(\varepsilon)$.

By Lemma 3.1 there is a semiclassical measure μ associated to u_h , which by Lemmas 3.2 and 3.3 has support in $\Gamma_+ \cap \{p = -1\}$. By Lemma 3.2, we have for every $a \in C_c^\infty(\mathcal{U}_\delta)$ and $t \geq 0$ (see (4.7) for the definition of \mathcal{U}_δ)

$$\int_{\mathcal{U}_\delta} a \circ e^{tH_p} d\mu = \int_{\mathcal{U}_\delta} e^{-2\gamma t - 2 \int_0^t V \circ e^{-rH_p} dr} a d\mu.$$

Therefore for all $\varepsilon' > 0$ and $t \geq T_{\varepsilon'}$ large enough, we have $e^{-tH_p}(\Gamma_+ \cap \mathcal{U}_\delta) \subset \Gamma_+ \cap \mathcal{U}_\delta$ and

$$e^{-2t(\gamma + V_{\max} + \varepsilon')} \mu(\Gamma_+ \cap \mathcal{U}_\delta) \leq \mu(e^{-tH_p}(\Gamma_+ \cap \mathcal{U}_\delta)) \leq e^{-2t(\gamma + V_{\min} - \varepsilon')} \mu(\Gamma_+ \cap \mathcal{U}_\delta). \quad (5.1)$$

Moreover, we claim that for each $\varepsilon' > 0$ and $t \geq T_{\varepsilon'}$ large enough

$$\{|p_+| \leq e^{-(\nu_{\max} + \varepsilon')t} \delta\} \cap \Gamma_+ \cap \mathcal{U}_\delta \subset e^{-tH_p}(\Gamma_+ \cap \mathcal{U}_\delta) \subset \{|p_+| \leq e^{-(\nu_{\min} - \varepsilon')t} \delta\} \cap \Gamma_+ \cap \mathcal{U}_\delta. \quad (5.2)$$

To see this, first note that $H_p(p_+)(x, \xi) = \xi(|X, U_+|(x)) = r_+(x)p_+(x, \xi)$ by (4.1) and thus

$$p_+(e^{-tH_p}(x, \xi)) = e^{-\int_0^t r_+(\varphi_{-r}(x)) dr} p_+(x, \xi), \quad t \in \mathbb{R},$$

and we obtain, by (2.5), that for $t \geq T_{\varepsilon'}$ large enough and $(x, \xi) \in \mathcal{U}_\delta$

$$e^{-(\nu_{\max} + \varepsilon')t} |p_+|(x, \xi) \leq |p_+|(e^{-tH_p}(x, \xi)) \leq e^{-(\nu_{\min} - \varepsilon')t} \delta,$$

thus giving (5.2). If $-\gamma \in \mathcal{S}_1(\varepsilon)$, by (5.2) and Lemma 4.7 (note it is here that we use $-\gamma > -\nu_{\min} + V_{\max}$), we obtain that there is $C = C(\varepsilon') > 0$ such that for $\varepsilon' > 0$ and $t \geq T_{\varepsilon'}$

$$C^{-1} e^{-(\nu_{\max} + \varepsilon')t} \leq \mu(e^{-tH_p}(\Gamma_+ \cap \mathcal{U}_\delta)) \leq C e^{-(\nu_{\min} - \varepsilon')t}. \quad (5.3)$$

Combining these inequalities with (5.1), we obtain for all $t \geq T_{\varepsilon'}$

$$C^{-1} e^{(2\gamma + 2V_{\min} - \nu_{\max} - 3\varepsilon')t} \leq \mu(\Gamma_+ \cap \mathcal{U}_\delta) \leq C e^{(2\gamma + 2V_{\max} - \nu_{\min} + 3\varepsilon')t}. \quad (5.4)$$

Since $-2\gamma < -\nu_{\max} + 2V_{\min} - 2\varepsilon$ we get a contradiction by choosing ε' small and letting $t \rightarrow \infty$.

Next, if $-\gamma \in \mathcal{S}_0(\varepsilon)$, by (5.2) and Lemma 4.7 we similarly get the upper bound of (5.3), which combined with (5.1) yields the upper bound in (5.4). Since $-2\gamma > -\nu_{\min} + 2V_{\max} + 2\varepsilon$, we can choose ε' small enough and by letting $t \rightarrow \infty$ this would force to have $\mu(\mathcal{U}_\delta) = 0$, contradicting Lemma 3.3.

Finally, the classical estimates (1.3) follow by introducing a semiclassical parameter $h := |\operatorname{Im}(s)|^{-1}$ and applying the semiclassical estimates (1.4), as well as (2.8). \square

APPENDIX A. AN EXOTIC SYMBOL CLASS

Let \mathcal{M} be a closed n -manifold equipped with a Riemannian metric g , let $k \geq 0$ and $0 < \rho < 1$. We use the usual notation $\langle \xi \rangle = (1 + |\xi|^2)^{1/2}$, $h > 0$ will be a small semiclassical parameter, we let $\bar{T}^* \mathcal{M}$ be the fiber radial compactification of $T^* \mathcal{M}$ as defined in [DyZw, Section E.1.3] and H_p will denote the Hamiltonian vector field of $p \in C^\infty(T^* \mathcal{M})$. Given an operator P , we write $\operatorname{Im} P = \frac{P - P^*}{2i}$ for the imaginary part of P and $\operatorname{Re} P = \frac{P + P^*}{2}$ for the real part; then $P = \operatorname{Re} P + i \operatorname{Im} P$. Recall that for $x \in \mathbb{R}$ we write $x_+ = \max(x, 0)$.

Most of the results we gather in this appendix are simple extensions of classical results in semiclassical analysis that can be found in the books [Zw] and [DyZw, Appendix E]. We shall only point out the main differences with our setting.

For each $m \in \mathbb{R}$, we define the exotic pseudo-differential calculus $\Psi_{h,\rho,k}^m(\mathcal{M})$ by saying that $A \in \Psi_{h,\rho,k}^m(\mathcal{M})$ if its Schwartz kernel K_A is in $\mathcal{O}_{C^N}(\mathcal{M} \times \mathcal{M})(h^N)$ for all $N > 0$ outside a neighborhood of the diagonal, and near the diagonal can be written in local coordinates as

$$K_A(x, y) = (2\pi h)^{-n} \int_{\mathbb{R}^n} e^{\frac{i}{h}(x-y)\xi} a(x, \xi) d\xi,$$

where the local symbols are in the class $S_{h,\rho,k}^m(\mathbb{R}^{2n})$ defined by the property: $a \in S_{h,\rho,k}^m(\mathbb{R}^{2n})$ if $a \in C^\infty(\mathbb{R}^{2n})$ is an h -dependent function and satisfies in local coordinates (for some $C_{\alpha,\beta}$ uniform in h)

$$|\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \leq C_{\alpha,\beta} \langle \xi \rangle^{m-|\beta|} h^{-\rho(|\alpha|-k)_+}. \quad (\text{A.1})$$

We notice that it is important here, for the calculus, that the loss of $h^{-\rho}$ happens only in the x -derivatives and not in the ξ derivatives. First, observe the basic properties for $0 \leq k' \leq k$

$$\begin{aligned} a \in S_{h,\rho,k}^m(\mathbb{R}^{2n}), b \in S_{h,\rho,k'}^{m'}(\mathbb{R}^{2n}) &\implies ab \in S_{h,\rho,k'}^{m+m'}(\mathbb{R}^{2n}), \\ a \in S_{h,\rho,k}^m(\mathbb{R}^{2n}) &\implies \partial_x^\alpha a \in h^{-\rho(|\alpha|-k)_+} S_{h,\rho,(k-|\alpha|)_+}^m(\mathbb{R}^{2n}), \partial_\xi^\alpha a \in S_{h,\rho,k}^{m-|\alpha|}(\mathbb{R}^{2n}), \\ \forall j \in [0, k], \quad h^{j\rho} S_{h,\rho,k-j}^m(\mathbb{R}^{2n}) &\subset S_{h,\rho,k}^m(\mathbb{R}^{2n}). \end{aligned} \quad (\text{A.2})$$

We define $S_{h,\rho,k}^m(T^*\mathcal{M})$ to be $C^\infty(T^*\mathcal{M})$ functions that, using local coordinates on \mathcal{M} , are in $S_{h,\rho,k}^m(\mathbb{R}^{2n})$. First, one directly sees from the formula of symbols under a change of coordinates [Zw, Theorem 9.9] that the symbol in local coordinates being in $S_{h,\rho,k}^m(\mathbb{R}^{2n})$ is invariant by change of coordinates, and moreover there is a principal symbol map

$$\sigma : \Psi_{h,\rho,k}^m(\mathcal{M}) \rightarrow S_{h,\rho,k}^m(T^*\mathcal{M})/hS_{h,\rho,k}^{m-1}(T^*\mathcal{M}).$$

Using local charts and a partition of unity, we fix a semi-classical quantization $\text{Op}_h : S_{h,\rho,k}^m(T^*\mathcal{M}) \rightarrow \Psi_{h,\rho,k}^m(\mathcal{M})$, which satisfies

$$\sigma(\text{Op}_h(a)) = a \bmod hS_{h,\rho,k}^{m-1}(T^*\mathcal{M}).$$

We first check that symbols in this class are closed under composition. Recall from [Zw, Theorem 4.14] that if $A \in \Psi_{h,\rho,k}^m(\mathbb{R}^n)$ and $B \in \Psi_{h,\rho,k}^{m'}(\mathbb{R}^n)$ have full symbol a, b then AB has full symbol (as an oscillatory integral)

$$a\#b(x, \xi) = (2\pi h)^{-n} \int_{\mathbb{R}^{2n}} e^{-\frac{i}{h}(x'\cdot\xi')} a(x, \xi + \xi') b(x + x', \xi) dx' d\xi' \quad (\text{A.3})$$

which has expansion

$$a\#b = \sum_{|\alpha| \leq N} \frac{(-ih)^{|\alpha|}}{\alpha!} \partial_\xi^\alpha a \partial_x^\alpha b + \mathcal{O}_{S_{h,\rho,0}^{m+m'-N-1}(\mathbb{R}^{2n})}(h^{(N+1)(1-\rho)}). \quad (\text{A.4})$$

Here $|\partial_\xi^\alpha a \partial_x^\alpha b| \leq C_\alpha h^{-|\alpha|\rho} \langle \xi \rangle^{m+m'-|\alpha|}$ so that higher order terms in the expansion are higher powers of h and of $\langle \xi \rangle^{-1}$.

Lemma A.1. *Let $a \in S_{h,\rho,k}^{m_1}(\mathbb{R}^{2n})$ and $b \in S_{h,\rho,k}^{m_2}(\mathbb{R}^{2n})$. Then $a\#b \in S_{h,\rho,k}^{m_1+m_2}(\mathbb{R}^{2n})$.*

Proof. This follows from (A.4) and (A.2). \square

For $A \in \Psi_{h,\rho,k}^m(\mathcal{M})$, we say that $(x_0, \xi_0) \in \overline{T^*\mathcal{M}}$ is not in $\text{WF}_h(A)$ if there is a small neighborhood U of (x_0, ξ_0) in $\overline{T^*\mathcal{M}}$ so that the full local symbol of A restricted to U is in $h^N S_{h,\rho,0}^{-N}(U)$ for all $N > 0$. We also define the elliptic set $\text{ell}_h(A)$ of $A \in \Psi_{h,\rho,k}^m(\mathcal{M})$ to be the set of points $(x_0, \xi_0) \in \overline{T^*\mathcal{M}}$ so that for a neighborhood U of (x_0, ξ_0) there is $c_0 > 0$ so that $\langle \xi \rangle^{-m} |\sigma(A)(x, \xi)| \geq c_0$, for $(x, \xi) \in U$. We finally list some properties of the $\Psi_{h,\rho,k}^m(\mathcal{M})$ calculus.

Proposition A.2. *The following properties hold:*

1. Let $A \in \Psi_{h,\rho,k}^m(\mathcal{M})$. If $B, B' \in \Psi_h^{\text{comp}}(\mathcal{M})$ with $\text{WF}_h(B) \cap \text{WF}_h(B') = \emptyset$, then $BAB' \in h^\infty \Psi_h^{\text{comp}}(\mathcal{M})$.

2. The principal symbol is well-defined as a map

$$\sigma : \Psi_{h,\rho,k}^m(T^*\mathcal{M}) \rightarrow S_{h,\rho,k}^m(T^*\mathcal{M})/hS_{h,\rho,k}^{m-1}(T^*\mathcal{M})$$

with kernel $h\Psi_{h,\rho,k}^{m-1}(\mathcal{M})$, and it satisfies for any $A \in \Psi_{h,\rho,k}^m(\mathcal{M})$ and $B \in \Psi_{h,\rho,k}^{m'}(\mathcal{M})$

$$\sigma(AB) = \sigma(A)\sigma(B) \text{ mod } h^{1-\rho}S_{h,\rho,k}^{m+m'-1}(T^*\mathcal{M}),$$

$$\sigma(AB) = \sigma(A)\sigma(B) \text{ mod } hS_{h,\rho,k-1}^{m+m'-1}(T^*\mathcal{M}) \text{ if } k \geq 1.$$

3. If $A \in \Psi_{h,\rho,k}^{m_1}(\mathcal{M})$ and $B \in \Psi_{h,\rho,k}^{m_2}(\mathcal{M})$, then

$$[A, B] \in h^{1-\rho}\Psi_{h,\rho,k}^{m_1+m_2-1}(\mathcal{M}),$$

$$h^{-1}[A, B] \in \Psi_{h,\rho,k-1}^{m_1+m_2-1}(\mathcal{M}) \text{ if } k \geq 1,$$

and $\sigma(h^{-1}[A, B]) = -i\{\sigma(A), \sigma(B)\}$.

4. If $P = -ihX$ for X a vector field on \mathcal{M} , $\Theta \in \Psi_{h,\rho,k}^m(\mathcal{M})$ and $H_p\sigma(\Theta) \in S_{h,\rho,k}^m(\mathcal{M})$, where $p = \sigma(P) = \xi(X)$, then $[P, \Theta] \in h\Psi_{h,\rho,k}^m(\mathcal{M})$.

5. If $A \in \Psi_{h,\rho,k}^m(\mathcal{M})$, then $A^* \in \Psi_{h,\rho,k}^m(\mathcal{M})$ and

$$\sigma(A^*) = \overline{\sigma(A)} \text{ mod } h^{1-\rho}S_{h,\rho,k}^{m+m'-1}(T^*\mathcal{M}),$$

$$\sigma(A^*) = \overline{\sigma(A)} \text{ mod } hS_{h,\rho,k-1}^{m-1}(T^*\mathcal{M}) \text{ if } k \geq 1.$$

6. Each $A \in \Psi_{h,\rho,0}^0(\mathcal{M})$ is bounded $L^2 \rightarrow L^2$ and for each $\varepsilon > 0$

$$\|A\|_{L^2 \rightarrow L^2} \leq (1 + \varepsilon) \sup_{h,x,\xi} |\sigma_h(A)(x, \xi)| + \mathcal{O}_\varepsilon(h^\infty).$$

Moreover, for any $A \in \Psi_{h,\rho,k}^m(\mathcal{M})$ and any $s \in \mathbb{R}$ we have

$$\|A\|_{H_h^s \rightarrow H_h^{s-m}} \leq (1 + \varepsilon) \sup_{h,x,\xi} |\langle \xi \rangle^{-m} \sigma(A)| + \mathcal{O}_\varepsilon(h^\infty).$$

7. Let $P \in \Psi_{h,\rho,k}^p(\mathcal{M})$, $A \in \Psi_{h,\rho,k}^m(\mathcal{M})$ and $B_1 \in \Psi_{h,\rho,k}^l(\mathcal{M})$. Assume $\text{WF}_h(A) \subset \text{ell}_h(P) \cap \text{ell}_h(B_1)$. Then for all $s \in \mathbb{R}$, $N > 0$, and u with $B_1Pu \in H_h^{s-p-l}(\mathcal{M})$

$$\|Au\|_{H_h^{s-m}} \leq C\|B_1Pu\|_{H_h^{s-p-l}} + \mathcal{O}(h^\infty)\|u\|_{H_h^{-N}}.$$

8. Assume $k \geq 1$ and let $P \in \Psi_{h,\rho,k}^1(\mathcal{M})$ with $\text{Re } P \in \Psi_h^1(\mathcal{M})$ and $\text{Im } P \in h\Psi_{h,\rho,k}^0(\mathcal{M})$.

Denote $p := \sigma(P)$ and assume that for each $(x, \xi) \in \text{WF}_h(A) \subset \overline{T^*\mathcal{M}}$, there is $T > 0$ such that $e^{-TH_p}(x, \xi) \in \text{ell}_h(B)$ and $e^{-tH_p}(x, \xi) \in \text{ell}_h(B_1)$ for $t \in [0, T]$. Then for each $u \in L^2$ with $Pu \in L^2(\mathcal{M})$, and every $N > 0$ there is a $C > 0$ such that

$$\|Au\|_{L^2} \leq C\|Bu\|_{L^2} + Ch^{-1}\|B_1Pu\|_{L^2} + Ch^N\|u\|_{H_h^{-N}}.$$

Proof. 1. This follows from the composition formula (A.4).

2. This was discussed above.

3. From the composition formula (A.4), locally we have

$$a\#b - b\#a \sim h(D_\xi a \partial_x b - D_\xi b \partial_x a) + \frac{1}{2}h^2(D_\xi^2 a \partial_x^2 b - D_\xi^2 b \partial_x^2 a) + \frac{1}{6}h^2(D_\xi^3 a \partial_x^3 b - D_\xi^3 b \partial_x^3 a) + \dots$$

where, after taking the expansion to a high enough order, the remainder is in $h^N S_{h,\rho,0}^{m-N}(\mathbb{R}^{2n})$ for some large $N > 0$. By (A.2), all these terms are in $h^{1-\rho} S_{h,\rho,k}^{m_1+m_2-1}(\mathbb{R}^{2n}) \cap h S_{h,\rho,k-1}^{m_1+m_2-1}(\mathbb{R}^{2n})$ if $k \geq 1$, and the principal symbol of $h^{-1}[A, B]$ is $-i(\partial_\xi a \partial_x b - \partial_x a \partial_\xi b)$.

4. This follows from the composition formula (A.4), the fact that $\partial_\xi^\alpha \sigma(P) = 0$ for $|\alpha| \geq 2$ and item 3 above.

5. This follows from the fact that, if A has full local symbol a in local coordinates, the full symbol a^* of A^* is

$$a^*(x, \xi) \sim \sum_{\alpha} \frac{h^{|\alpha|}}{\alpha!} \partial_{\xi}^{\alpha} \overline{D_x^{\alpha} a(x, \xi)}.$$

6. The argument is standard (see [GrSj, Theorem 4.5]) and we just make a brief summary. Let $a := \sigma(A) \in S_{h,\rho,k}^0(\mathcal{M})$. For $M = \|a\|_{\infty}$, we can construct $B_0 \in \Psi_{h,\rho,0}^0(\mathcal{M})$ with principal symbol $b_0 := \sqrt{(1+\varepsilon)^2 M^2 - |a|^2} \geq \varepsilon M > 0$, so that

$$C := (1+\varepsilon)^2 M^2 - A^* A = B_0^* B_0 + h^{1-\rho} R_0,$$

for some $R_0 = R_0^* \in \Psi_{h,\rho,0}^{-1}(\mathcal{M})$ by item 2 (and the fact that $a \in S_{h,\rho,0}^m$ and $a > 0$ implies $\sqrt{a} \in S_{h,\rho,0}^{m/2}$). Next we can choose $B_1 = B_0 + \frac{h^{1-\rho}}{2} \text{Op}_h(b_0^{-1})^* R_0$ so that $B_1^* B_1 = C - h^{2(1-\rho)} R_1$ with $R_1 \in \Psi_{h,\rho,0}^{-2}(\mathcal{M})$. We iterate this procedure to find $B_N \in \Psi_{h,\rho,0}^0(\mathcal{M})$ such that $(1+\varepsilon)^2 M^2 - A^* A = B_N^* B_N + h^{(N+1)(1-\rho)} R_N$ with $R_N \in \Psi_{h,\rho,0}^{-N-1}(\mathcal{M})$. By Schur's lemma we have $\|R_N\|_{L^2 \rightarrow L^2} = \mathcal{O}(h^{(N+1)(1-\rho)-n})$ and thus for any $u \in L^2$

$$\|Au\|_{L^2}^2 = (1+\varepsilon)^2 M^2 \|u\|_{L^2}^2 - \|B_N u\|_{L^2}^2 + \mathcal{O}(h^{2(N+1)(1-\rho)-n}) \|u\|_{L^2}^2$$

which shows the desired estimate by choosing N large. Similarly the case of arbitrary k follows and the Sobolev bound is an easy consequence of this.

7. This follows from the parametrix construction in the elliptic set, the main thing to notice is that if $B \in \Psi_{h,\rho,k}^m(\mathcal{M})$, $A \in \Psi_{h,\rho,k}^l(\mathcal{M})$ and $\text{WF}_h(A) \subset \text{ell}_h(B)$, then

$$\frac{\sigma(A)}{\sigma(B)} \in S_{h,\rho,k}^{l-m}(\mathcal{M}).$$

8. We follow the proof of [DyZw, Theorem E.47] and divide the proof into steps.

Step 0: an escape function. Fix $\beta \geq 0$. There is a $g \in C^{\infty}(\overline{T^* \mathcal{M}})$ with $\text{supp } g \subset \text{ell}_h(B_1)$, such that

$$g \geq 0, \quad g > 0 \text{ on } \text{WF}_h(A), \quad H_p g \leq -\beta g,$$

where the last condition holds outside $\text{ell}_h(B)$.

Step 1. Note that $g \in S^0(T^* \mathcal{M})$ and define

$$G := \text{Op}_h(\langle \xi \rangle^s g) \in \Psi_h^s(\mathcal{M}), \quad \text{WF}_h(G) \subset \text{ell}_h(B_1).$$

We can that $u \in C^{\infty}(\mathcal{M})$. If we write $f = Pu$,

$$\text{Im} \langle f, G^* Gu \rangle = \underbrace{\text{Im} \langle (\text{Re } P)u, G^* Gu \rangle}_{\text{term T}_1} + \underbrace{\text{Re} \langle (\text{Im } P)u, G^* Gu \rangle}_{\text{term T}_2}.$$

We will bound the two terms on the right separately.

Step 2: term T₁. Since $\text{Re}(P) \in \Psi_h^1(\mathcal{M})$ is non exotic, this step is exactly the same as in the proof of [DyZw, Theorem E.47] and we get

$$T_1 \leq (C_1 - \beta)h \|Gu\|_{L^2}^2 + Ch \|Bu\|_{H_h^s}^2 + Ch^2 \|B_1 u\|_{H_h^{s-\frac{1}{2}}}^2 + \mathcal{O}(h^{\infty}) \|u\|_{H_h^{-N}}^2.$$

Step 3: term T₂. Write

$$T_2 = \text{Re} \langle (\text{Im } P)u, G^* Gu \rangle = \langle (\text{Im } P)Gu, Gu \rangle + \text{Re} \langle [G, \text{Im } P]u, Gu \rangle.$$

We estimate the two terms on the right hand side separately. Firstly

$$|\langle (\text{Im } P)Gu, Gu \rangle| = h |\langle (h^{-1} \text{Im } P)Gu, Gu \rangle| \leq C_2 h \|Gu\|_{L^2}^2 + \mathcal{O}(h^{\infty}) \|u\|_{H_h^{-N}}^2,$$

where we used the boundedness property (item 6) for the exotic operator $h^{-1} \text{Im } P$. For the second term, we need to deal with a commutator. Note first that

$$\text{Re } G^* [G, \text{Im } P] = h \text{Re } G^* [G, h^{-1} \text{Im } P] \in h^2 \Psi_{h,\rho,k-1}^{2s-1}(\mathcal{M}),$$

by item 3 above. As $\text{WF}_h(G^*[G, \text{Im } P]) \subset \text{ell}_h(B_1)$, by the elliptic estimate

$$\begin{aligned} |\langle \text{Re}(G^*[G, \text{Im } P])u, u \rangle| &= h^2 |\langle h^{-2} \text{Re}(G^*[G, \text{Im } P])u, Yu \rangle| + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}}^2 \\ &\leq Ch^2 \|B_1 u\|_{H_h^{s-\frac{1}{2}}}^2 + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}}^2, \end{aligned}$$

where $Y \in \Psi_h^0(\mathcal{M})$ is such that $Y = 1 + \mathcal{O}(h^\infty)$ microlocally on $\text{WF}_h(\text{Re}(G^*[G, \text{Im } P]))$ and $\text{WF}_h(Y) \subset \text{ell}_h(B_1)$. Here we used the boundedness in the exotic class item 6, item 1, the elliptic estimate in the exotic class (item 7). Adding the two estimates we finally obtain

$$|\text{Re}\langle (\text{Im } P)u, G^*Gu \rangle| \leq C_2 h \|Gu\|_{L^2}^2 + Ch^2 \|B_1 u\|_{H_h^{s-\frac{1}{2}}}^2 + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}}^2.$$

Step 4. Adding the estimates in Steps 2 and 3, we obtain

$$\text{Im}\langle f, G^*Gu \rangle \leq (C_1 + C_2 - \beta)h \|Gu\|_{L^2}^2 + Ch \|Bu\|_{H_h^s}^2 + Ch^2 \|B_1 u\|_{H_h^{s-\frac{1}{2}}}^2 + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}}^2.$$

By ellipticity of B on $\text{WF}_h(G)$, there is $Q \in \Psi_h^s(\mathcal{M})$ such that $G = QB_1 + R$, where $R \in h^\infty \Psi_h^{-\infty}(\mathcal{M})$, thus

$$|\langle f, G^*Gu \rangle| \leq |\langle QB_1 f, Gu \rangle| + |\langle Rf, Gu \rangle| \leq C \|B_1 f\|_{H_h^s} \|Gu\|_{L^2} + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}}^2.$$

Now choose $\beta = C_1 + C_2 + 1$ to get

$$\|Gu\|_{L^2}^2 \leq C \|Bu\|_{H_h^s}^2 + Ch^{-1} \|B_1 f\|_{H_h^s} \|Gu\|_{L^2} + Ch \|B_1 u\|_{H_h^{s-\frac{1}{2}}}^2 + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}}^2.$$

We can absorb the $\|Gu\|_{L^2}$ term to the left hand side, at the cost of the additional term $Ch^{-2} \|B_1 f\|_{H_h^s}^2$ on the right hand side. Next, use the condition $\text{WF}_h(A) \subset \text{ell}_h(G)$ and elliptic estimates to derive

$$\|Au\|_{H_h^s} \leq C \|Bu\|_{H_h^s} + Ch^{-1} \|B_1 f\|_{H_h^s} + Ch^{\frac{1}{2}} \|B_1 u\|_{H_h^{s-\frac{1}{2}}} + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}}. \quad (\text{A.5})$$

Step 5. Here, one can use the same induction procedure as in [DyZw, Proof of Th. E.47] to show that for each $\ell \in \mathbb{N}$

$$\|Au\|_{H_h^s} \leq C \|Bu\|_{H_h^s} + Ch^{-1} \|B_1 f\|_{H_h^s} + Ch^{\frac{\ell}{2}} \|B_1 u\|_{H_h^{s-\frac{\ell}{2}}} + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}},$$

where the first step of the induction is exactly where we arrived in (A.5). \square

Next we discuss semiclassical defect measures in the setting of the exotic calculus.

Proposition A.3. *Assume that $u_h \in L^2(\mathcal{M})$ is a family satisfying $\|u_h\|_{L^2} = \mathcal{O}(1)$. Then there exists a Radon measure μ , called semiclassical measure, and a sequence $h_j \rightarrow 0$, such that for any $A \in \Psi_{h,\rho,0}^{\text{comp}}(\mathcal{M})$ with $\lim_{h \rightarrow 0} \sigma(A)(h; x, \xi) = a_0(x, \xi)$ in $C_c^0(T^*\mathcal{M})$, it holds that*

$$\lim_{j \rightarrow \infty} \langle Au_{h_j}, u_{h_j} \rangle_{L^2} = \int_{T^*\mathcal{M}} a_0 d\mu.$$

Proof. We follow the proof of [DyZw, Theorem E.42]. By Proposition A.2, we have that $A - \text{Op}_h(a_h) \in h\Psi_{h,\rho,0}^{\text{comp}}(\mathcal{M})$ for some symbol $a_h \in S_{h,\rho,0}^{\text{comp}}(T^*\mathcal{M})$ so that $a_h \rightarrow a_0$ in $C_c^0(T^*\mathcal{M})$; it suffices to prove the claim for $A = \text{Op}_{h_j}(a_{h_j})$. We write $I_h(a_h) := \langle \text{Op}(a_h)u_h, u_h \rangle_{L^2}$ and claim

$$\limsup_{h \rightarrow 0} |I_h(a_h)| \leq C \limsup_{h \rightarrow 0} \|a_h\|_\infty \leq C \sup_h \|a_h\|_\infty. \quad (\text{A.6})$$

Indeed, by Cauchy-Schwarz and Proposition A.2 (item 6), we have

$$|I_h(a_h)| \leq C \|a_h\|_\infty + \mathcal{O}_A(h^\infty), \quad (\text{A.7})$$

where $C = C(\sup_h \|u_h\|_{L^2}) > 0$. Take a countable, dense subset $(a_h^\ell)_{\ell \in \mathbb{N}} \subset S_{h,\rho,0}^{\text{comp}}(T^*\mathcal{M})$, where $S_{h,\rho,0}^{\text{comp}}(T^*\mathcal{M})$ is equipped with the inductive limit topology from the seminorms in (A.1). By a

diagonal argument and since by (A.7) $I_h(a_h^\ell)$ is bounded for all ℓ , we may extract a sequence $h_j \rightarrow 0$ such that $I_{h_j}(a_{h_j}^\ell)$ converges for all ℓ . For each $a_h \in S_{h,\rho,0}^{\text{comp}}(T^*\mathcal{M})$ and ℓ , we get by (A.7)

$$\limsup_{j,j' \rightarrow \infty} |I_{h_j}(a_{h_j}) - I_{h_{j'}}(a_{h_{j'}})| \leq \limsup_{j,j' \rightarrow \infty} |I_{h_j}(a_{h_j}^\ell) - I_{h_{j'}}(a_{h_{j'}}^\ell)| + C \sup_h \|a_h - a_h^\ell\|_\infty.$$

Using the density of a_h^ℓ , we obtain that $I_{h_j}(a_{h_j})$ is a Cauchy sequence and we may define for $a_h \in S_{h,\rho,0}^{\text{comp}}(T^*\mathcal{M})$

$$I(a_h) := \lim_{j \rightarrow \infty} I_{h_j}(a_{h_j}).$$

By (A.7), the map I satisfies for each $a_h \in S_{h,\rho,0}^{\text{comp}}(T^*\mathcal{M})$

$$|I(a_h)| \leq C \limsup_{j \rightarrow \infty} \|a_{h_j}\|_\infty \leq C \sup_{h \in (0, h'_0)} \|a_h\|, \quad (\text{A.8})$$

for any $h'_0 > 0$. In particular, I extends to a continuous linear functional on $C_c^0(T^*\mathcal{M})$ (h -independent functions). Given $a_h \in S_{h,\rho,0}^{\text{comp}}(T^*\mathcal{M})$ with $\lim_{h \rightarrow 0} a_h = a_0 \in C_c^\infty(T^*\mathcal{M})$ in the $C_c^0(T^*\mathcal{M})$ topology, we get by (A.8)

$$|I(a_0 - a_h)| \leq C \sup_{h \in (0, h'_0)} \|a_0 - a_h\|_\infty \rightarrow 0 \text{ as } h'_0 \rightarrow 0,$$

and thus $I(a_h) = I(a_0)$. By Gårding's inequality $I(a_0) \geq 0$ when $a_0 \geq 0$ and so by the Riesz-Markov representation theorem there is a Radon measure μ such that for each $a_0 \in C_c^\infty(T^*\mathcal{M})$

$$I(a_0) = \lim_{j \rightarrow \infty} \langle \text{Op}_{h_j}(a_0)u_{h_j}, u_{h_j} \rangle_{L^2} = \int_{T^*\mathcal{M}} a_0 d\mu.$$

The main claim follows from this by using $I(a_h) = I(a_0)$ under the given assumptions. \square

Now we prove a version of a propagation estimate for the semiclassical measure in the exotic calculus. This is a slight extension of [DyZw, Theorem E.44]. Note that if $\sigma(P)$ is real valued and $k \geq 1$, then $\text{Im } P \in h\Psi_{h,\rho,k-1}^{m-1}(\mathcal{M})$ if $P \in \Psi_{h,\rho,k}^m(\mathcal{M})$ by Proposition A.2 (item 5).

Proposition A.4. *Assume $\|u_h\|_{L^2} = \mathcal{O}(1)$ and u_h converges to a semiclassical measure μ . Let $P \in \Psi_{h,\rho,k}^m(\mathcal{M})$ with $k \geq 1$, denote $p := \sigma(P)$ and assume that p is real-valued for all h , and define $b := \sigma(h^{-1} \text{Im } P)$. Assume that for each $a \in C_c^\infty(T^*\mathcal{M})$*

$$(H_p a)_0 = \lim_{h \rightarrow 0} H_p a \quad \text{and} \quad b_0 = \lim_{h \rightarrow 0} b \quad \text{exist in } C_c^0(T^*\mathcal{M}).$$

Then there is $C > 0$ such that for all $a \in C_c^\infty(T^\mathcal{M})$ and $Y \in \Psi_h^{\text{comp}}(\mathcal{M})$ with $Y = 1 + \mathcal{O}(h^\infty)$ microlocally on $\text{supp}(a)$*

$$\left| \int_{T^*\mathcal{M}} ((H_p a)_0 + 2b_0 a) d\mu \right| \leq C \|a\|_\infty \limsup_{h \rightarrow 0} (h^{-1} \|Y P u_h\|_{L^2} \|Y u_h\|_{L^2}).$$

Proof. Assume without loss of generality that a is real valued. Let $A \in \Psi_h^{\text{comp}}(\mathcal{M})$ be such that $\sigma(A) = a$ and $A^* = A$. We compute

$$\begin{aligned} h^{-1} \text{Im} \langle P u_h, A u_h \rangle &= (2i)^{-1} h^{-1} \langle A P u_h, u_h \rangle - (2i)^{-1} h^{-1} \langle P^* A u_h, u_h \rangle \\ &= (2i)^{-1} \langle h^{-1} [A, P] u_h, u_h \rangle + \langle (h^{-1} \text{Im } P) A u_h, u_h \rangle. \end{aligned}$$

Now by Proposition A.2 (item 3), we have $h^{-1} [A, P] \in \Psi_{h,\rho,k-1}^{\text{comp}}(\mathcal{M})$ with $-i\sigma(h^{-1} [A, P]) = H_p a$ and by assumptions $\sigma((h^{-1} \text{Im } P) A) = ba$. Thus by Proposition A.3 there is a semiclassical measure μ such that the right hand side converges to, after extracting a subsequence $h_j \rightarrow 0$

$$\int_{T^*\mathcal{M}} \left(\frac{1}{2} (H_p a)_0 + b_0 a \right) d\mu.$$

Moreover, the left hand side equals (using Proposition A.2, item 1)

$$(2ih)^{-1} (\langle P Y u_h, A Y u_h \rangle - \langle A Y u_h, P Y u_h \rangle) + \mathcal{O}(h^\infty),$$

from which we easily deduce the main estimate. □

REFERENCES

- [BuLi] O. Butterley, C. Liverani, *Smooth Anosov flows: correlation spectra and stability*. J. Mod. Dyn. **1** (2) (2007), 301–322.
- [DiSj] M. Dimassi, J. Sjöstrand, *Spectral asymptotics in the semi-classical limit*. London Mathematical Society Lecture Note Series, 268. Cambridge University Press, Cambridge, 1999. xii+227 pp.
- [Do] D. Dolgopyat, *On decay of correlations in Anosov flows*, Annals of Mathematics **147** (1998) 357–390.
- [Dy1] S. Dyatlov, *Resonance projectors and asymptotics for r -normally hyperbolic trapped sets*. J. Amer. Math. Soc. **28** (2015), no. 2, 311–381.
- [Dy2] S. Dyatlov, *Spectral gaps for normally hyperbolic trapping*. Ann. Inst. Fourier (Grenoble) **66** (2016), no. 1, 55–82.
- [DFG] S. Dyatlov, F. Faure, C. Guillarmou, *Power spectrum of the geodesic flow on hyperbolic manifolds*. Anal. PDE **8** (2015), no. 4, 923–1000.
- [DyZw] S. Dyatlov, M. Zworski, *Mathematical theory of scattering resonances*, AMS Graduate Studies in Mathematics 200.
- [FaGu] F. Faure, C. Guillarmou, *Horocyclic invariance of Ruelle resonant states for contact Anosov flows in dimension 3*. Math Research Letters **25** (2018), no 5, 1405–1427.
- [FRS] F. Faure, N. Roy, J. Sjöstrand, *Semi-classical approach for Anosov diffeomorphisms and Ruelle resonances*, Open Math. Journal, vol. **1** (2008), 35–81.
- [FaSj] F. Faure, J. Sjöstrand, *Upper bound on the density of Ruelle resonances for Anosov flows*, Comm. Math. Phys. **308** (2011), no. 2, 325–364.
- [FaTs0] F. Faure, M. Tsujii, *Band structure of the Ruelle spectrum of contact Anosov flows*, Comptes rendus Mathématique **351** (2013), 385–391.
- [FaTs1] F. Faure, M. Tsujii, *Prequantum transfer operator for Anosov diffeomorphism*, Astérisque **375**, (2015), SMF.
- [FaTs2] F. Faure, M. Tsujii, *The semiclassical zeta function for geodesic flows on negatively curved manifolds* Inventiones mathematicae **208** (2017), 851–998.
- [FaTs3] F. Faure, M. Tsujii, *Micro-local analysis of contact Anosov flows and band structure of the Ruelle spectrum*, preprint arXiv:2102.11196.
- [GrSj] A. Grigis, J. Sjöstrand, *Microlocal analysis for differential operators. An introduction*. London Mathematical Society Lecture Note Series, 196. Cambridge University Press, Cambridge, 1994. iv+151 pp.
- [GHW] C. Guillarmou, J. Hilgert, T. Weich, *Classical and quantum resonances for hyperbolic surfaces*. Math. Annalen **370** (2018), Issue 3-4, pp 1231–1275.
- [GuKu] C. Guillarmou, B. Küster, *Spectral theory of the frame flow on hyperbolic 3-manifolds* (with an Appendix by Charles Hadfield), Preprint arXiv:2005.08387.
- [HuKa] S. Hurder, A. Katok, *Differentiability, rigidity and Godbillon-Vey classes for Anosov flows*, Publications IHES, **72** (1990), 5–61.
- [KuWe] B. Küster, T. Weich, *Quantum-classical correspondence on associated vector bundles over locally symmetric spaces*, IMRN April 2019 rnz068.
- [Li] C. Liverani, *On contact Anosov flows*, Ann. of Math. **159** (2004), no 2, 1275–1312.
- [NoZw] S. Nonnenmacher, M. Zworski, *Decay of correlations for normally hyperbolic trapping*, Inventiones mathematicae **200** (2015), no 2, 345–438.
- [Ts1] M. Tsujii, *Quasi-compactness of transfer operators for contact anosov flows*, Nonlinearity **23** (2010), 1495–1545.
- [Ts2] M. Tsujii, *Contact Anosov flows and the FBI transform*, Erg. Th. Dyn. Syst., **32** (2012), 2083–2118.
- [Ts3] M. Tsujii, *Exponential mixing for generic volume-preserving Anosov flows in dimension three*, Journal of the Mathematical Society of Japan. **70** (2018), no2, 757–821.
- [TsZh] M. Tsujii, Z. Zhang, *Smooth mixing Anosov flows in dimension three are exponential mixing*, arXiv 2006.04293.
- [Zw] M. Zworski, *Semiclassical analysis*. Graduate Studies in Mathematics, 138. American Mathematical Society, Providence, RI, 2012. xii+431 pp.

Email address: mihajlo.cekic@universite-paris-saclay.fr

LABORATOIRE DE MATHÉMATIQUES D’ORSAY, UNIVERSITÉ PARIS-SACLAY, CNRS, 91405 ORSAY, FRANCE

Email address: colin.guillarmou@math.u-psud.fr

LABORATOIRE DE MATHÉMATIQUES D’ORSAY, UNIV. PARIS-SUD, CNRS, UNIVERSITÉ PARIS-SACLAY, 91405 ORSAY, FRANCE