



Surface temperatures and their influence on the permafrost thermal regime in high Arctic rock walls on Svalbard

Juditha U Schmidt, Bernd Etzelmüller, Thomas V Schuler, Florence Magnin, Julia Boike, Moritz Langer, Sebastian Westermann

► To cite this version:

Juditha U Schmidt, Bernd Etzelmüller, Thomas V Schuler, Florence Magnin, Julia Boike, et al.. Surface temperatures and their influence on the permafrost thermal regime in high Arctic rock walls on Svalbard. 2020. <hal-03024157>

HAL Id: hal-03024157

<https://hal.science/hal-03024157v1>

Preprint submitted on 25 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Surface temperatures and their influence on the permafrost thermal regime in high Arctic rock walls on Svalbard

5 Juditha U. Schmidt¹, Bernd Etzelmüller¹, Thomas V. Schuler¹, Florence Magnin², Julia Boike^{3,4}, Moritz Langer^{3,4}, Sebastian Westermann¹

¹Department of Geosciences, University of Oslo, Oslo, 0316, Norway

²EDYTEM Lab, Université Savoie Mont Blanc, CNRS, Le Bourget-du-Lac, France

10 ³Alfred Wegener Institute (AWI), Helmholtz Centre for Polar and Marine Research, Potsdam, 14473, Germany

⁴Department of Geography, Humboldt-Universität zu Berlin, Berlin, 12489, Germany

Correspondence to: Juditha U. Schmidt (juditha.schmidt@geo.uio.no)

Abstract.

Permafrost degradation in steep rock walls and associated slope destabilization have been studied
15 increasingly in recent years. While most studies focus on mountainous and sub-Arctic regions, the occurring thermo-mechanical processes play an important role also in the high Arctic. A more precise understanding is required to assess the risk of natural hazards enhanced by permafrost warming in high Arctic rock walls.

This study presents rock surface temperature measurements of coastal and non-coastal rock walls in a
20 high Arctic setting on Svalbard. We applied the surface energy balance model CryoGrid 3 for evaluation, including adjusted radiative forcing to account for vertical rock walls.

Our measurements and model results show that rock surface temperatures at coastal cliffs are up to 1.5 °C higher than non-coastal rock walls when the fjord is ice-free in the winter season, resulting from additional energy input due to higher air temperatures at the coast and radiative warming by relatively warm
25 seawater. An ice layer on the fjord counteracts this effect, leading to similar rock surface temperatures as in non-coastal settings. Our results include a simulated surface energy balance with short-wave radiation as the dominant energy source during spring and summer, and long-wave radiation being the main energy

loss. While sensible heat fluxes can both warm and cool the surface, latent heat fluxes are mostly insignificant. Simulations for future climate conditions result in a warming of rock surface temperatures and a deepening of active layer thickness for both coastal and non-coastal rock walls. Our field data presents a unique data set of rock surface temperatures in steep high Arctic rock walls, while our model can contribute towards the understanding of factors influencing coastal and non-coastal settings and the associated surface energy balance.

1 Introduction

As a response to a warming climate, degradation of permafrost in steep rock walls can contribute to slope destabilization (Gruber and Haeberli, 2007; Krautblatter et al., 2013). Increased frequencies of slope failures have been observed in recent years (Fischer et al., 2012; Gruber et al., 2004; Raveland et al., 2010, 2017). These natural hazards can damage infrastructure and cause casualties in downslope regions (Harris et al., 2001, 2009). Permafrost in rock walls has been studied in mountainous regions (Allen et al., 2009; Krautblatter et al., 2010; Magnin et al., 2015; Noetzli and Gruber, 2009) as well as in sub-arctic areas (Blikra and Christiansen, 2014; Lewkowicz et al., 2012; Magnin et al., 2019). However, permafrost dynamics in steep rock walls in the high Arctic are poorly understood. In this study, we will focus on rock surface temperatures in steep coastal and fluvial cliffs at a high Arctic site near Ny-Ålesund, Svalbard (Figure 1).

Svalbard is located in the northern part of the warm North-Atlantic current and therefore, it is very sensitive to atmospheric and oceanic changes (Walczowski and Piechura, 2011). Increasing air temperatures are observed for more than a century (Nordli et al., 2020). Climate models predict an increase in precipitation and a warming of air temperature with the most pronounced air temperature change in winter season (Hanssen-Bauer et al., 2019; Isaksen et al., 2016). The climatic changes are also apparent in permafrost temperatures on Svalbard as observed in boreholes over the last decades (Boike et al., 2018; Christiansen et al., 2010; Isaksen et al., 2007). Simulated thermal conditions in Svalbard show an increase of ground temperatures and indicate a significant warming and an increase in active layer thickness over the 21st century (Eitzelmüller et al., 2011). Besides large-scale climate changes, local

55 conditions can play an important role in the surface energy budget, resulting in an amplification or dampening of the large-scale signal (Westermann et al., 2009). Besides sensible and latent heat fluxes, short-wave and long-wave radiation are crucial factors as they have a strong impact on the energy transfer processes from the atmosphere to the ground, effectively modulated in presence of insulating snow cover. The terrain exposure induces significant spatial variability of short-wave radiation that should be
60 considered when modelling thermal conditions in steep rock walls (Fiddes and Gruber, 2014; Magnin et al., 2015).

Besides down-welling radiation, long-wave radiation emitted by water bodies as well as reflected short-wave radiation on snow and ice can influence the rock surface temperature. Therefore, sea ice coverage
65 plays an important role for the surface energy balance of coastal cliffs. According to observations since 1997, Kongsfjorden was characterized by sea ice cover during winter season (Gerland and Hall, 2006). Since 2006, the sea ice extent has been reduced significantly and the ice thickness and snow cover on ice have become thinner (Johansson et al., 2020). This could also affect coastal erosion as sea ice and development of an ice foot protect the cliffs by absorbing ocean wave energy and control the removal of
70 weathered material from the base of the cliff (Ødegård and Sollid, 1993). With shorter or absent fast ice periods, coastal cliffs are exposed to waves and tides for longer durations. Climate models predict a further reduction of sea ice cover in the western fjords of Svalbard (Hanssen-Bauer et al., 2019). Thus, thermal models for steep rock walls have to consider the influence of aspect and slope angle on radiative forcing as well as additional heat sources like open seawater and reflection of short-wave radiation on sea
75 ice.

In this study, we applied a full energy balance model to evaluate the role of the different radiative forcing elements on the thermal regime in steep slopes at a high Arctic site. In doing so, we extended the parametrization of radiative forcing in the thermal model CryoGrid 3 to account for effects governing
80 steep rock walls, and validated the model with measured rock wall temperatures in the study area. Our objectives were to analyze the effect of coastal and non-coastal settings (i) on rock surface temperatures

of vertical rock walls, (ii) the surface energy balance throughout the seasons and (iii) to estimate future developments of the thermal regime until 2100 for these settings.

2 Study site

85 The observation site is situated near the village of Ny-Ålesund, Kongsfjorden, located at the west coast of Spitsbergen. We measured rock surface temperatures in steep coastal and non-coastal rock walls (Figure 1). Carbonate rocks of Permian to Carboniferous age with an apparent joint system are the dominant bedrocks (Figure 2). The surrounding of the study area is a strandflat and characterized by tundra vegetation, while the surface sediments are dominated by fine to medium-grained glacial and
90 marine deposits (Hop and Wiencke, 2019; Westermann et al., 2009).

Long-term records of climatological parameters are evidence of ongoing changes in the Arctic climate system with an increase of mean annual temperature by $+1.3 \pm 0.7^{\circ}\text{C}$ per decade and a rise during winter months by $+3.1 \pm 2.6^{\circ}\text{C}$ per decade. The winter warming is linked to a change in downward long-wave
95 radiation of $+15.6 \pm 11.6 \text{ Wm}^{-2}$ per decade (Maturilli et al., 2015), while the net short-wave radiation is altered by a decrease in reflection caused by reduced snow cover duration (Hop and Wiencke, 2019).

The main surface wind direction is along the axis of Kongsfjorden from the inland to the coast throughout all seasons. The mountains cause complex wind fields (Maturilli and Kayser, 2017) and a south-easterly
100 wind flow occurs as a result of channeled winds from the Kongsvegen glacier (Beine et al., 2001). Measured mean annual precipitation in Ny-Ålesund in the period 1971-2000 was 409 mm (Hanssen-Bauer et al., 2019). It can occur as both rain and snow throughout the year, but the snow-free season is typically from June to October (Hop and Wiencke, 2019).

3 Methods

3.1 Surface rock temperature monitoring

In this study, we used eight iButton (© Maxim) temperature loggers with an accuracy of ± 0.5 °C for this study (Table 1). They were installed during summers and springs of 2016 and 2017 in different locations, labelled with RW01 to RW08 (Figure 1) near Ny-Ålesund in Svalbard and were placed on north- to east-facing rock walls. They represented three different settings: (i) non-coastal rock walls in the canyon of Bayelva (three locations), (ii) coastal cliffs at the open fjord next to Ny-Ålesund airport (four locations) and (iii) a coastal cliff in the bay of Thiisbukta (one location). The settings allowed the analysis of permafrost temperatures in non-coastal rock walls and coastal cliffs affected by seawater (Figure 2).

3.2 Model description

We adapted the CryoGrid 3 ground thermal model (Westermann et al., 2016), originally designed for horizontal surfaces, to account for conditions in steep rock walls (Magnin et al., 2017). CryoGrid 3 calculates rock temperatures by solving the heat equation, uses the surface energy balance as an upper boundary condition, and considers latent heat effects depending on water content of the substrate. The heat transfer to the ground is calculated by heat conduction. The surface energy balance is derived from time series of air temperature, specific humidity, wind speed at a known height above the ground, incoming short-wave and long-wave radiation, air pressure and rates of snowfall and rainfall (Westermann et al., 2016). In the standard version designed for horizontal surfaces, turbulent fluxes between the surface and the atmosphere are controlled by vertically moving air parcel as defined in the Monin-Obukhov similarity theory (Monin and Obukhov, 1954). As a consequence, movement of air parcels at a vertical wall would be parallel to the surface rather than perpendicular. Therefore, we assumed in all model calculations, that the near-surface wind profile follow a neutral atmospheric stratification (Magnin et al., 2017).

Besides the analysis of rock surface temperatures (RST), we used CryoGrid 3 to determine the active layer thickness (ALT). We applied a small grid spacing in the upper layers (0.1 m between 0 m and 1 m

130 depth) and gradually increased the grid spacing to the lower layers of the model (10 m between 50 m and 100 m depth) to account for detailed ground temperature calculations in the active layer near the surface.

3.3 Preprocessing

As the energy input of short-wave and long-wave radiation is depending on varying aspects and slope angles of the rock walls, we modified the model to account for the different physical settings of the logger
 135 locations. We calculate incoming short-wave radiation as the sum of direct, diffuse and reflected short-wave radiation, while incoming long-wave radiation includes atmospheric long-wave radiation as well as heat emission of the close environment.

We divided short-wave radiation into direct and diffuse components (Fiddes and Gruber, 2014). It required the determination of the atmospheric clearness index k_t , the ratio between solar radiation arriving
 140 at the surface S_{in} and the radiation at the top of the atmosphere S_{TOA} :

$$k_t = \frac{S_{in}}{S_{TOA}}. \quad \text{Eq. (1)}$$

The fraction of diffuse short-wave radiation k_d was computed based on the clearness index k_t

$$k_d = 0.952 - 1.041e^{-\exp(2.300 - 4.702 * k_t)} \quad \text{Eq. (2)}$$

145

and taking into account the sky view factor SVF. As we applied the model to vertical rock walls, we assumed a SVF of 0.5 for all locations (Kastendeuch, 2013). Therefore, the amount of diffuse short-wave radiation S_{diff} can be expressed as

$$S_{diff} = SVF * k_d * S_{in}. \quad \text{Eq. (3)}$$

150 Consequently, the amount of direct short-wave radiation S_{dir} is the remaining fraction $(1 - k_d) * S_{in}$. After we determined the azimuth α_{sun} and elevation β_{sun} of the sun for every time step depending on latitude, longitude and altitude of each location, we projected direct short-wave radiation S_{dir} on inclined slopes (Appendix A).

155 Besides direct and diffuse short-wave radiation, we implemented reflected short-wave radiation in the model to account for diffuse reflection on ice and snow surfaces as well as on snow-free terrain. Assuming Lambertian reflectance, we derived reflected short-wave radiation S_{ref} , taking into account the albedo of the surface α and the sky view factor SVF:

$$S_{ref} = S_{in} * \alpha * (1 - SVF). \quad \text{Eq. (4)}$$

160

We used the sum of diffuse, direct and reflected short-wave radiation as a driving variable for the model CryoGrid 3 on vertical rock walls.

Moreover, we modified long-wave radiation by using the implemented sky view factor SVF. For
 165 simplicity, we assumed an SVF of 0.5 for all locations, so 50 % of the long-wave radiation is given by the forcing data L_{in_forc} , representing the atmospheric long-wave radiation, while the rest is derived from the ambient temperature T_{amb} applying the Stefan-Boltzman law with Stefan-Boltzman constant σ (Fiddes and Gruber, 2014):

$$L_{in} = SVF * L_{in_forc} + (1 - SVF) * \sigma * (T_{amb} + 273.15)^4 \quad \left[\frac{W}{m^2}\right]. \quad \text{Eq. (5)}$$

170

With this approach, we assumed that incoming long-wave radiation is isotropic. The ambient temperature T_{amb} was given by either the air temperature, or the sea surface temperature in case the logger is located directly above the sea. If the seawater was covered by ice and could not emit any heat, we used air temperature for deriving the long-wave radiation.

175

Apart from the modification of incoming short-wave and long-wave radiation, we included the water balance in the model by implementing a water bucket approach. Due to the vertical alignment of the rock walls and its consistency of hard bedrock, precipitation does not infiltrate into the material and evaporation of moisture at the rock surface dominated the latent heat flux. Therefore, the latent heat flux

180 had only minor influence on the total surface energy balance and a simplistic water bucket approach was sufficient for the required model setup (Appendix B).

We did not consider snow cover in the model, which, according to field observations in April and May, was adequate for the majority of measurement sites. An exception is displayed in Figure 3, showing the
185 damped signal of RW01 due to snow cover. The data of these time periods were analyzed but excluded from the comparison with the model.

We define the surface as the interface between the atmosphere and the rock wall. Fluxes, which transport energy away from the surface have a negative sign, while fluxes, which transport energy towards the
190 surface are denoted positive.

3.4 Model parameters and forcing data

The study focused on rock surface temperatures, thus simplified subsurface properties were implemented (Table 2). We considered the bedrock to have a volumetric mineral content of 97 % and a volumetric water content of 3 %, which implied saturated conditions during the entire simulation. Albedo for
195 limestones was found between approximately 0.22 and 0.32 (Blumthaler and Ambach, 1988) and as light-colored carbonates build up the cliffs, we assumed an albedo of 0.3 in the model setup. An important fitting parameter was the roughness length z_0 as performed in Magnin et al., (2017). We set it to a value of 0.018 m, which represents roughly 1/10 of the height of the surface roughness elements. This fitted well to the rugged rock surface characterized by joint systems (Figure 2). We set the albedo for the
200 horizontal ground surface to 0.15, for water surfaces to 0.1 and the albedo for ice and snow to a relatively low value of 0.55, as the highest influence of reflected short-wave radiation was expected for spring, when snowmelt decreases the albedo (Table 2).

Atmospheric forcing was provided by the AROME-Arctic weather model, which is a regional high-
205 resolution, non-hydrostatic numerical weather prediction system for the European Arctic (Müller et al., 2017). It is based on HARMONIE-AROME as part of the ALADIN-HIRLAM system, which provides

short-range weather forecasts for Northern and Southern European countries (Bengtsson et al., 2017; Seity et al., 2011). Archive files of atmospheric data are available since October 2015. In 2017, updates were implemented to improve high-resolution weather forecasts over the Nordic regions (Müller et al., 210 2017). AROME-Arctic operates on a resolution of ~2.5 km grid spacing at 65 vertical levels. We used time series ranging from October 2015 to August 2020 from AROME-Arctic as forcing data for the model. The nearest 2.5 km grid cell of the AROME-Arctic to the required locations was located northeast of Ny-Ålesund in Kongsfjorden (78.9N, 11.98E, 20 m a.s.l.). The selected grid cell represents the transition between the fjord and the land and therefore, it provides suitable forcing data for the loggers 215 located directly or within a short distance to the shoreline. The driving variables absolute humidity, wind speed, down-welling short-wave and long-wave radiation, air pressure and rates of snowfall and rainfall for this grid cell have been extracted from the archive. Incident solar radiation at the top of the atmosphere was provided by ERA5 (Hersbach, 2016).

220 The spatial resolution of air temperature given by AROME-Arctic was not sufficient to capture small-scale variabilities. Therefore, we used records from two climate stations to force the model (Boike et al., 2018, 2019; Maturilli, 2020a, 2020b, 2020c, 2020d). The Baseline surface radiation network (BSRN) station in Ny-Ålesund is located in the village center (78.9250N, 11.9300E) with a distance of about 300 m to the coast (Maturilli et al., 2013). The second station at the Bayelva site is located on top of the 225 Leirhaugen hill, which is in 1.3 km distance to the coast (Boike et al., 2018). Using records of two different stations allowed us to estimate gradients in air temperatures from the coast to environments further inland. For simplicity, we interpolated linearly between the two stations and estimated the air temperature at the rock walls subject to their distance to the open water body of the fjord. As sea ice coverage enlarges the distance to the open water body, we added an additional mean distance (Table 3), 230 estimated by analysis of the webcam time series from the mountain Zeppelinfjellet (Pedersen, 2013).

We used water temperature of Kongsfjorden recorded by the AWIPEV underwater observatory in 12 m depth to estimate the long-wave heat emission of the water body. The data provides a time series of water

temperatures for the entire period from October 2015 to August 2020 with a resolution of one hour
235 (Fischer et al., 2018a, 2018b, 2018c, 2019).

We simulated long-term climate impacts of three different representative concentration pathways (RCP) RCP2.6, RCP4.5 and RCP8.5 (van Vuuren et al., 2011) for coastal and non-coastal settings. For the period 1980 to 2019, we used forcing data of the ERA Interim Reanalysis, while the years 2020 to 2100 were
240 created using an anomaly approach based on CMIP5 projections of Community Climate System Model (CCSM4) (NCAR, 2016). Therefore, decadal monthly anomalies were derived from the CCSM4 projections using a reference period from 2009 to 2019 which were then applied to the Reanalysis data of the same period (see Koven et al., 2015).

To account for small-scale variabilities between coastal and non-coastal rock walls, we calculated a linear
245 regression using the AROME forcing data. Others parameters were simplified due to a lack of information: sea temperature was set to a constant value of 2.53 °C, which is the mean sea temperatures of the analyzed period 2016 to 2020. Sea ice was assumed in the months February to May until the year 2005 (Gerland and Hall, 2006). Despite these uncertainties, these steps allowed us to analyze possible trajectories for the future developments of the rock wall thermal regime.

250 **3.5 Model scenarios**

In this study, we considered several scenarios to represent the thermal conditions at the selected rock walls (Table 4). We varied the source of forcing air temperature, the source for heat emission and the albedo of the foot of the slope to account for the different locations. The *non-coastal scenario* was controlled by conditions either with or without snow at the foot of the slope resulting in temporarily
255 changing albedos for snow and terrain. The *frozen bay scenario* showed temporarily frozen seawater leading to changes in the source of heat emission and albedo of the foot of the slope. The *open fjord scenario* was characterized by a predominantly unfrozen fjord during the entire year and had consequently no varying parameters in most of the simulation time. Short periods of a frozen layer occur temporarily so that the model parameters were modified the same way as in the *frozen bay scenario*. Daily frozen

260 conditions were estimated by analyzing webcam time series from the mountain Zeppelinfjellet, which provides photos of Ny-Ålesund and the adjacent coastline every ten minutes (Pedersen, 2013).

4 Results

4.1 Measurements of rock surface temperature

265 Mean annual temperatures (Sep – Aug) as well as mean temperatures in the winter season (Dec – Feb) are given in Table 6. For the measurement period 2016 to 2020, all logger record below-freezing mean annual rock surface temperatures (MARST) with values between -0.6 °C (RW06 in 2017/18) and -4.3 °C (RW02 in 2019/20). The MARST typically vary up to several degrees between the recorded years, with 2017/18 being the warmest year. Measurements in 2019/20 show the lowest MARST, which is related to a comparatively cold winter (Dec – Feb) and spring (Mar – May) season (Table 6).

270 Minimum and maximum daily RST are found between -24.2 °C (RW03) and 18.9 °C (RW06). The variability of daily RST show a higher frequency in summer and higher amplitudes in winter. Fluctuations of RST are especially pronounced for non-coastal rock walls during the cold periods of the year, while the signal at coastal cliffs at the open fjord is dampened in the same time.

275 We emphasize that loggers at the coastal cliffs record higher MARST than loggers in non-coastal rock walls with a mean difference of MARST of 1.0 °C (Table 6), although the loggers are located in just about 1.5 km distance (Figure 1) at similar elevations. The setting is especially important in winter and spring season and RST differences account for 1.5 °C to 2.2 °C in these time periods. The lower the temperatures, the larger is the temperature difference between these two settings, which is apparent in Figure 4a.

280 Besides these observation in RST, time series of station data show that higher air temperatures are recorded for the BSRN station in Ny-Ålesund compared to the Bayelva site further inland. During the year 2017/18, the air temperature difference was 0.9 °C with the highest differences of 1.6 °C during winter season and 1.5 °C during spring season.

285 We highlight that RST values in the Thiisbukta bay are significantly lower in winter than RST at the coastal cliffs even though they are all located at the shoreline. This is especially true for periods where the bay is characterized by an ice layer on the water (Figure 4a), but can also be observed for unfrozen conditions in the bay. If not only the bay is frozen, but widespread sea ice occurs in the fjord, RST values in all three settings show about the same temperatures (Figure 4b).

290 **4.2 Model validation**

We compared monthly average values of measured rock surface temperature RST to the model results of the *non-coastal scenario* (RW01, RW02, RW03), the *open fjord scenario* (RW04, RW05, RW06, RW07) and the *frozen bay scenario* (RW08). The measured RST was reproduced closely with the applied model setup, especially for temperatures near freezing point (Figure 5). Besides the visually good agreement of
295 Figure 5, a root-mean-square error (RMSE) below 1.2 °C, the bias (-0.5 °C to 0.4 °C), the coefficient of determination R^2 (above 0.97) and the Nash-Sutcliffe efficiency NSE (above 0.96) for all locations confirmed a good reproduction of the measured data (Table 5).

4.3 The influence of open water and sea ice on RST

The model results are in good agreement with the in situ measurements and corroborate the pattern of
300 open water and sea ice influence on RST.

Below-freezing MARST are modelled for all locations, ranging from -0.3 °C (RW06 in 2017/18) to -3.9 °C (RW02 in 2019/20) with the warmest year being 2017/18 and the coldest year being 2019/20. The lowest MARST are modelled at rock walls in the *non-coastal scenario*, while the *open fjord scenario* produces the highest MARST.

305 The model results show differences in MARST according to the exposition of the rock wall: In the *open fjord scenario*, the lowest MARST in 2017/18 is found on the north-facing rock wall RW05 (-0.9 °C), while the highest MARST is calculated for RW06 facing east-north-east (-0.3 °C). Accordingly, the model results with a similar model setup are dependent on the orientation of the rock wall.

310 In the model results, we find that temporarily occurring ice cover on the fjord results in lower RST at the nearby rock walls. For time periods with a frozen bay, but no sea ice in the open fjord, only RW08 is affected. Model results show ca. 1 – 1.5 °C colder RST compared to the other rock walls at the shoreline but they are still warmer than the modelled RST in non-coastal settings. However, the results indicate that days with a widespread sea ice extent in the fjord lead to similar RST in all locations (Table 7).

315 Lower RST values under frozen conditions can be traced back to three different factors: While (1) lower air temperature and (2) the lack of heat emission from the ocean lead to a cooling of RST, (3) the reflection of short-wave radiation on the ice layer increases RST as an additional energy source. The amount to which these factors influence the decrease in RST between the *open fjord scenario* and the *frozen bay scenario* is given in Figure 6. Between December and February, air temperature and the lack of heat emission are the dominant factors, while reflected short-wave radiation plays no role. In March, the influence of reflected short-wave radiation increases as polar night conditions end.

4.4 The surface energy balance

Individual fluxes of the surface energy budget in the different seasons are given in Figure 7 (Winter = 325 Dec – Feb; Spring = Mar – May; Summer = Jun – Aug; Fall = Sep – Nov) with positive fluxes directed towards the surface. For comparison of the different scenarios, the fluxes are calculated for vertical rock walls with an aspect of 40° (~NE), which comprises model runs of RW02, RW04 and RW08.

During winter which mostly coincides with polar night conditions (25 Oct - 14 Feb), short-wave radiation 330 is zero or only reaches very small values. During this period, the system loses energy mainly due to negative net long-wave radiation, which is especially pronounced for the *non-coastal scenario*, followed by the *frozen bay scenario*. The loss of energy is opposed by positive sensible heat fluxes, representing a warming of the surface and a cooling of the atmosphere. Strong sensible heat fluxes are associated with high wind speeds and high temperature differences of air and rock wall. Compared to the other terms of the SEB, the negative ground heat flux leading to ground cooling is only small.

In spring, net short-wave radiation increases significantly and becomes the dominant energy source with highest energy input for the *non-coastal scenario* and the lowest for the *open fjord scenario*. Long-wave radiation counteracts this process and cool the surface with highest fluxes in the *non-coastal scenario*.
340 Besides, sensible heat fluxes contribute to the energy loss with slight differences in the scenarios. However, sensible heat fluxes and emitted long-wave radiation cannot compensate the incoming energy by short-wave radiation and RST as well as ground heat fluxes start to increase, especially as no energy is used for melting due to snow-free conditions.

345 The summer period is characterized by similar fluxes of the surface energy balance in all scenarios. The warming of RST continues due to strong short-wave radiation as the main energy source. Energy is lost by long-wave radiation and sensible heat fluxes but also latent heat fluxes cool the surface. However, these fluxes cannot compensate the energy input. Consequently, the ground heat flux increases even more, leading to a seasonal thawing of the active layer.

350 During fall, net short-wave radiation decreases rapidly due to shorter days, but sensible heat fluxes turn positive, acting as an energy source again. The loss of energy by long-wave radiation is slightly higher for *non-coastal scenarios* and ground heat fluxes are close to zero, indicating the turn to refreezing of the active layer.

355 In the course of a year, short-wave radiation is naturally the main source of energy to the system, while most energy is lost by long-wave radiation. Sensible heat fluxes warm the surface in fall and winter, while they cool in spring and summer. Latent heat fluxes are of minor importance, reflecting the small water holding capacity of the rock surface assumed in the model. Net ground heat fluxes are close to zero.

360 **4.5 Simulations of future climate change scenarios**

The past and future simulations of different RCPs show an increase in MARST for both the *non-coastal* and the *open fjord scenario* (Figure 8). Between 1980 and 2020, MARST increases by several degrees and the MARST difference of the *non-coastal scenario* and the *open fjord scenario* is significant. We

emphasize that in this period winter sea ice loss has been drastic in Kongsfjorden, going from a normally
365 frozen fjord to a normally open fjord. Thus, in reality, the actual warming may have even been higher
than either of the scenarios suggests. Figure 8 clarifies, that the current measurement period represents
relatively warm years in the occurring fluctuations of MARST and a drop to colder MARST in 2020.
During the years 2020 to 2080, all three pathways show slightly increasing MARST. The influence of the
different rock wall locations decreases with time and consequently, MARST values of the *non-coastal*
370 and *open fjord scenario* become more aligned. After 2080, RCP2.6 and RCP4.5 are stabilized with
MARST between -4 °C and 0 °C, while RCP8.5 shows further increasing MARST reaching mean annual
values up to 2 °C.

The model results suggest that a large part of the warming in RST has already happened until the year
2020, while the prospective increase of RST in the 21st century will get the permafrost close to thawing.

375

Furthermore, simulations of the RCPs suggest an effect on the active layer thickness ALT (Figure 9).
Between 1980 and 2010, ALT is about 2 m, with a slightly deeper active layer for the *open fjord scenario*.
After 2010, the ground begins to thaw deeper during summer season, but to various extents in the three
simulated RCPs: RCP2.6 is characterized by a slight increase in ALT and a stabilization between 2.5 m
380 and 3.5 m after 2080. RCP4.5 shows a similar trend with ALTs between 3.0 m and 4.5 m at the end of
the century. However, the simulation of RCP8.5 results in a significant increase of ALT below 8 m and
with no apparent stabilization effect. Moreover, a talik is developed after 2095, implying that the cold
winter seasons do not lead to a freezing of the entire ground column anymore.

5 Discussion

385 5.1 Model uncertainties

Our model setup contains uncertainties regarding unknown model parameters, which were estimated
using literature and calibrating the model. Especially rock (surface) parameters could be improved by a
more precise analysis of the lithological characteristics.

390 A critical point is the assumption of a neutral atmospheric stratification perpendicular to the vertical rock wall which must be regarded as a first order approximation. This leads to uncertainties in near surface turbulent exchange of the vertical wall as micro-topography and changing weather conditions can influence the movement of the air parcels. Wind profiles perpendicular to the wall as well as measurements of air temperature and humidity could help to estimate the importance of this error source.
395 However, we scaled the roughness length z_0 to compensate these assumption and to fit the modelled surface temperatures to the observed values.

Another error source is the estimation of air temperature at the logger location. The linear interpolation between the two stations and the dependency on the distance to the open water body of the fjord is a
400 coarse approximation at best which does not take the wind direction, the atmospheric boundary layer structure and local micro-climate into account. Air temperature measurements at the shoreline could help to improve the quality of the forcing data.

5.2 The influence of open water and sea ice on RST

In the time period 2016 to 2020, all records display negative MARST which indicates permafrost
405 conditions at all locations (Table 6). Svalbard lies in the continuous permafrost zone (Brown et al., 1997; Obu et al., 2019) and deep permafrost is observed in the area, e.g. within the abandoned mine shafts (Liestøl, 1977) and in boreholes (Christiansen et al., 2010).

Two main factors control the RST difference during winter season between the relatively warm *open fjord*
410 *scenario* and the relatively cold *non-coastal scenario*: (i) Air temperature gradients from the coast to the inland play an important role. They have a pronounced effect on the surface energy balance, especially on turbulent fluxes, and result in higher RST at the coastal settings. Furthermore, (ii) incoming long-wave radiation leads to higher RST values at the coastal cliffs. The rock walls receive energy by long-wave radiation emitted by the surfaces in the field of view, which is controlled by seawater temperature for
415 open fjord settings and by air temperature for non-coastal rock walls. As the seawater has significantly higher temperatures than the air during winter, the energy input is larger at the coastal cliffs.

Thick snow deposits (> 1 m depth) can effectively insulate the ground resulting in higher RST during winter (Haberkorn et al., 2015). This is also true for Ny-Ålesund where ground surface measurements and model results indicate higher mean annual ground surface temperatures (MAGST) for planes with thick snow covers as documented by Gislås et al., (2014). However, thin snow cover (< 0.5 m depth) can lead to a lowering of RST as low air temperatures can still affect the rock while the high albedo of the snow reflects large parts of solar radiation (Haberkorn et al., 2015; Magnin et al., 2017). Both the warming and the cooling effects are apparent in RST measurements at those logger positions, which are temporarily covered by snow (Figure 3). Furthermore, a direct comparison of MARST in the largely snow-free cliffs with MAGST measured in near-horizontal tundra settings close to the Bayelva station (using the measurement setup of Gislås et al., (2014)) suggest that MARST at coastal cliffs can be even warmer than MAGST under a thick snow cover.

While an ice cover in the bay leads to locally decreased RST in the bay, a widespread sea ice extent results in lower RST for all settings. Maximum sea ice coverage is significantly reduced since 2006 and a shorter sea ice season is observed since 2002 (Johansson et al., 2020). Hence, it can be assumed that RST at the coastal cliff has increased since then in winter and spring season. Three main factors could be identified which influence the model results: During polar night conditions, (i) air temperature gradients between the open water body and inland and (ii) radiative heating by comparatively warm seawater strongly affect RST, leading to lower RST for frozen conditions. When incoming solar radiation increases again in March, (iii) the increased reflection of short-wave radiation on the ice cover can counteract these processes to a certain extent.

5.3 The surface energy balance

The components of the surface energy balance are estimated for different seasons of the year. In summer and fall, fluxes are largely similar for the different scenarios, while significant differences can be noticed in winter and spring (Figure 7).

Net short-wave radiation is the dominant source of energy for all scenarios in spring and summer due to
445 midnight sun conditions. The flux is especially strong, when solar radiation is strongly reflected by
surrounding sea ice or snow cover, in addition to the direct and diffuse short-wave radiation. This can be
seen in spring in the *non-coastal scenario* (reflection on snow) and the *frozen bay scenario* (reflection on
ice). As the bay is not continuously frozen, the effect is less pronounced in the *frozen bay scenario*. In the
open fjord scenario, reflection of short-wave radiation plays a minor role as a result of the low albedo of
450 seawater.

The system loses energy by net long-wave radiation during the entire year and the differences between
the scenarios are most pronounced in winter and spring. The small net long-wave radiation in the *open
fjord scenario* can be explained by higher incoming long-wave radiation through emission of the
455 relatively warm seawater. During summer, the temperature difference between seawater and air is smaller
which limits the influence compared to the cold seasons. However, sensible heat fluxes are a major
component and play an important role during the entire year. In winter and fall, they warm the surface
and cool the atmosphere, while this process is reversed in spring and summer. In winter, sensible heat
fluxes are larger in the *non-coastal* and *frozen bay scenario* compared to the *open fjord scenario*. As
460 roughness lengths and wind speed are assumed to be the same, this effect can be traced back to larger
temperature differences of air and rock wall. Therefore, the air temperature gradients in the surroundings
of Ny-Ålesund intensify the sensible heat transfer to the surface and influence the surface energy budget
of the rock walls significantly. Latent heat fluxes play a minor role in the energy budget as just a small
amount of water can be stored in the vertical bedrock that is available for evaporation.

465 **5.4 Future climate change scenarios**

At the costal cliff sites, our simulations suggest that MARST have significantly increased in the last 40
years (Figure 8). The increase of MARST is especially pronounced since the year 2000. This is in line
with increasing ground temperatures, observed from 1998 to 2017 at the Bayelva station close to the
setting described in this study (Boike et al., 2018). A further increase of MARST is predicted for all

470 pathways with positive values for RCP8.5 at the end of the century. The main reasons are increasing air temperatures and long-wave radiation in the course of the 21st century.

Another important effect is the reduced difference of the *non-coastal scenario* and the *open fjord scenario* with ongoing permafrost warming. This can be explained by a convergence of seawater temperature and air temperature during the winter season in the assumed model forcing, which likely reflects true
475 conditions in a future ice-free Arctic. As a consequence, the influence of the relatively warm seawater on the radiation budget and coastal air temperatures becomes less important under a warmer climate and RST of non-coastal and open fjord settings become more similar.

Moreover, the impact of a changing climate becomes visible in deeper layers of the ground, e.g. through
480 a deepening of the active layer. However, these model results must be interpreted carefully as additional warming from the top of the cliff must be considered, depending on the geometry of the cliffs. Therefore, the presented model results for deeper layers are likely biased for the investigated small rock walls, while they might be applicable to higher cliffs in the close surroundings of Ny-Ålesund.

5.5 Coastal cliffs in the high Arctic – a future geohazard?

485 Our model results indicate a significant warming of permafrost temperatures and a deepening of ALT in the 21st century, a trend that can affect the mechanical erosion of the rock walls. The time and depth spend in the optimal thermal range for frost cracking between -3 °C and -8 °C may change (Anderson, 1998; Hales and Roering, 2007) and longer durations of open-water season and a correlated interaction of water with the shoreline can enhance coastal erosion (Barnhart et al., 2014). Besides, warming of permafrost
490 temperatures can lead to further destabilization as rock- and ice-mechanical properties are modified (Krautblatter et al., 2013). At the seashore, loss of sea ice may intensify the expected degradation of permafrost and result in enhanced slope instabilities.

In this study, the thermal regime of relatively low coastal cliffs is investigated. Indeed, similar processes
495 can also affect much higher coastal cliffs on Svalbard, for example in the Mesozoic rocks in the Isfjorden area (Etzel Müller et al., 2003). As they are mostly not entirely vertical, corresponding sky view factors

might be larger, resulting in a slightly smaller influence of long-wave radiation emitted by seawater and reflected short-wave radiation from sea ice. However, warmer coastal air temperatures will have an impact independent on inclination of the slope. As a consequence, shorter durations of sea ice cover will
500 likely lead to an enhanced warming of permafrost in high rock cliffs in Svalbard.

Failures of coastal rock slopes can impact the water body and trigger displacement waves along shorelines (Hermanns et al., 2006). Prominent examples are the landslide in Paatuut / West Greenland in 2000, creating a tsunami with a run-up height of up to 50 m (Dahl-Jensen et al., 2004) and the rock avalanche
505 in Tåfjord / Norway in 1934, producing an up to 62 m high displacement wave (Hermanns et al., 2006). Despite different geological and climatological settings, an occurrence of rock slope failures causing tsunamis cannot be excluded for Svalbard. Large active rock slope instabilities over water bodies have been detected, like the Forkastningsfjellet rockslide in Isfjorden of circa 300.000 m³, which is located in just 8 km distance of Longyearbyen with 2200 inhabitants (Kuhn et al., 2019). Due to permafrost
510 degradation, rock slope failures in the high Arctic might become more likely in future.

6 Conclusions

In this study, we present measurements of rock surface temperatures (RST) from steep coastal and non-coastal cliffs in the high Arctic setting of Ny-Ålesund, Svalbard, comprising data from 2016 to 2020. The permafrost model CryoGrid 3 is applied for this thermal regime with an adapted parametrization of
515 radiative forcing: The slope angles and aspects of the rock walls have been taken into account as well as additional heat sources like long-wave emission from seawater and reflecting short-wave radiation on snow and ice covers. With our measurements and model results, we can draw the following conclusions:

- Measured RSTs in coastal cliffs are up to 2 °C higher during the winter season than the non-
520 coastal rock walls. Model results suggest that this results from slightly higher air temperatures at the coast compared to inland locations as well as from the continuous energy input by long-wave radiation from the relatively warm seawater.

- When the sea adjacent to the coastal cliff is covered by sea ice, coastal RSTs are decreased and closely match RST at inland locations. This can be explained by lower air temperatures and disabled long-wave emission of seawater. Reflection of short-wave radiation on the ice cover counteracts this process, but is only effective when polar night conditions end. As a consequence, sea ice loss in Kongsfjorden is expected to increase RST on coastal cliffs during winter.
- Simulations for future climate conditions show an increase of mean annual rock surface temperatures MARST with a stabilization between -4 °C and 0 °C for RCP2.6 and RCP4.5 and a further warming to above-freezing MARST for RCP8.5 at the end of the century. Furthermore, the model predicts a deepening of the active layer for all RCPs.

Appendix

Appendix A: Projection of direct short-wave radiation on inclined planes

We determined the direct short-wave radiation perpendicular to an inclined slope S_{dir_slope} by projecting the direct short-wave radiation on the horizontal S_{dir} as following:

$$S_{dir_slope} = \frac{S_{dir}}{\cos(\delta_{sun_hor} * \frac{\pi}{180})} * \cos(\delta_{sun_slope} * \frac{\pi}{180}) \quad [\frac{W}{m^2}] \quad \text{Eq. (6)}$$

with δ_{sun_hor} being the angle between the solar rays and the normal on the horizontal plane and δ_{sun_slope} being the angle between the solar rays and the normal on the inclined slope. The valid solar zenith angle was set to 85° excluding the very early sunrise and sunset, as this would lead to highly overestimated energy input on a vertical wall due to nearly vertical solar rays.

Appendix B: The water bucket approach

We based the infiltration of water on a water bucket approach. As long as the uppermost grid cell is unfrozen, water could infiltrate and the water content of the grid cell increased. After reaching saturation, excess water was removed by surface runoff. We calculated the potential evaporation with the turbulent latent heat flux based on the Monin-Obukhov similarity theory (Monin and Obukhov, 1954). We weighted

the amount of evaporated water with the water content of the grid cell, taking into account that water can be evaporated more easily under saturated conditions.

References

- 550 Allen, S. K., Gruber, S. and Owens, I. F.: Exploring steep bedrock permafrost and its relationship with recent slope failures in the Southern Alps of New Zealand, *Permafrost and Periglacial Processes*, 20(4), 345–356, doi:10.1002/ppp.658, 2009.
- Anderson, R. S.: Near-surface Thermal Profiles in Alpine Bedrock: Implications for the Frost Weathering of Rock, *Arctic and Alpine Research*, 30(4), 362–372, doi:10.1080/00040851.1998.12002911, 1998.
- Barnhart, K. R., Anderson, R. S., Overeem, I., Wobus, C., Clow, G. D. and Urban, F. E.: Modeling erosion of ice-rich
555 permafrost bluffs along the Alaskan Beaufort Sea coast, *Journal of Geophysical Research: Earth Surface*, 119, 1155–1179, doi:10.1002/2013JF002845, 2014.
- Beine, H. J., Argentini, S., Maurizi, A., Mastrantonio, G. and Viola, A.: The local wind field at Ny-Ålesund and the Zeppelin mountain at Svalbard, *Meteorology and Atmospheric Physics*, 78(1), 107–113, doi:10.1007/s007030170009, 2001.
- Bengtsson, L., Andrae, U., Aspelien, T., Batrak, Y., Calvo, J., de Rooy, W., Gleeson, E., Hansen-Sass, B., Homleid, M., Hortal,
560 M., Ivarsson, K.-I., Lenderink, G., Niemelä, S., Nielsen, K. P., Onvlee, J., Rontu, L., Samuelsson, P., Muñoz, D. S., Subias, A., Tijm, S., Toll, V., Yang, X. and Køltzow, M. Ø.: The HARMONIE–AROME Model Configuration in the ALADIN–HIRLAM NWP System, *Monthly Weather Review*, 145(5), 1919–1935, doi:10.1175/MWR-D-16-0417.1, 2017.
- Blikra, L. H. and Christiansen, H. H.: A field-based model of permafrost-controlled rockslide deformation in northern Norway, *Geomorphology*, 208, 34–49, doi:10.1016/j.geomorph.2013.11.014, 2014.
- 565 Blumthaler, M. and Ambach, W.: Solar Uvb-Albedo of Various Surfaces, *Photochemistry and Photobiology*, 48(1), 85–88, doi:10.1111/j.1751-1097.1988.tb02790.x, 1988.
- Boike, J., Juszak, I., Lange, S., Chadburn, S., Burke, E., Overduin, P. P., Roth, K., Ippisch, O., Bornemann, N., Stern, L., Gouttevin, I., Hauber, E. and Westermann, S.: A 20-year record (1998–2017) of permafrost, active layer and meteorological conditions at a high Arctic permafrost research site (Bayelva, Spitsbergen), *Earth System Science Data*, 10(1),
570 355–390, doi:https://doi.org/10.5194/essd-10-355-2018, 2018.
- Boike, J., Grünberg, I., Bornemann, N. and Lange, S.: Meteorological observations at Bayelva Station in 2018 (level 1), *PANGAEA*, doi:10.1594/PANGAEA.898683, 2019.
- Brown, J., Ferrians Jr., O. J., Heginbottom, J. A. and Melnikov, E. S.: Circum-Arctic map of permafrost and ground-ice conditions, Report., 1997.
- 575 Christiansen, H. H., Etzelmüller, B., Isaksen, K., Juliusen, H., Farbrøt, H., Humlum, O., Johansson, M., Ingeman-Nielsen, T., Kristensen, L., Hjort, J., Holmlund, P., Sannel, A. B. K., Sigsgaard, C., Åkerman, H. J., Foged, N., Blikra, L. H., Pernosky, M. A. and Ødegård, R. S.: The thermal state of permafrost in the nordic area during the international polar year 2007–2009, *Permafrost and Periglacial Processes*, 21(2), 156–181, doi:10.1002/ppp.687, 2010.

- 580 Dahl-Jensen, T., Larsen, L. M., Pedersen, S. A. S., Pedersen, J., Jepsen, H. F., Pedersen, G., Nielsen, T., Pedersen, A. K., Von Platen-Hallermund, F. and Weng, W.: Landslide and Tsunami 21 November 2000 in Paatuut, West Greenland, *Natural Hazards*, 31(1), 277–287, doi:10.1023/B:NHAZ.0000020264.70048.95, 2004.
- Etzelmlüller, B., Ødegård, R. S. and Sollid, J. L.: The spatial distribution of coast types on Svalbard, *Arctic Coastal Dynamics – Report of the 3rd International Workshop*, 33–40, 2003.
- 585 Etzelmlüller, B., Schuler, T. V., Isaksen, K., Christiansen, H. H., Farbrot, H. and Benestad, R.: Modeling the temperature evolution of Svalbard permafrost during the 20th and 21st century, *The Cryosphere*, 5(1), 67–79, doi:https://doi.org/10.5194/tc-5-67-2011, 2011.
- Fiddes, J. and Gruber, S.: TopoSCALE v.1.0: downscaling gridded climate data in complex terrain, *Geoscientific Model Development*, 7(1), 387–405, doi:https://doi.org/10.5194/gmd-7-387-2014, 2014.
- 590 Fischer, L., Purves, R. S., Huggel, C., Noetzli, J. and Haeberli, W.: On the influence of topographic, geological and cryospheric factors on rock avalanches and rockfalls in high-mountain areas, *Natural Hazards and Earth System Sciences*, 12(1), 241–254, doi:https://doi.org/10.5194/nhess-12-241-2012, 2012.
- Fischer, P., Schwanitz, M., Brand, M., Posner, U., Brix, H. and Baschek, B.: Hydrographical time series data of the littoral zone of Kongsfjorden, Svalbard 2015, Alfred Wegener Institute - Biological Institute Helgoland, PANGAEA, doi:10.1594/PANGAEA.896771, 2018a.
- 595 Fischer, P., Schwanitz, M., Brand, M., Posner, U., Brix, H. and Baschek, B.: Hydrographical time series data of the littoral zone of Kongsfjorden, Svalbard 2016, Alfred Wegener Institute - Biological Institute Helgoland, PANGAEA, doi:10.1594/PANGAEA.896770, 2018b.
- Fischer, P., Schwanitz, M., Brand, M., Posner, U., Brix, H. and Baschek, B.: Hydrographical time series data of the littoral zone of Kongsfjorden, Svalbard 2017, Alfred Wegener Institute - Biological Institute Helgoland, PANGAEA, doi:10.1594/PANGAEA.896170, 2018c.
- 600 Fischer, P., Schwanitz, M., Brand, M., Posner, U., Gattuso, J.-P., Alliouane, S., Brix, H. and Baschek, B.: Hydrographical time series data of the littoral zone of Kongsfjorden, Svalbard 2018, Alfred Wegener Institute - Biological Institute Helgoland, doi:10.1594/PANGAEA.897349, 2019.
- Gerland, S. and Hall, R.: Variability of fast-ice thickness in Spitsbergen fjords, *Annals of Glaciology*, 44, 231–239, doi:10.3189/172756406781811367, 2006.
- 605 Gisnås, K., Westermann, S., Schuler, T. V., Litherland, T., Isaksen, K., Boike, J. and Etzelmlüller, B.: A statistical approach to represent small-scale variability of permafrost temperatures due to snow cover, *The Cryosphere*, 8, 2063–2074, doi:10.5194/tc-8-2063-2014, 2014.
- Gruber, S. and Haeberli, W.: Permafrost in steep bedrock slopes and its temperature-related destabilization following climate change, *Journal of Geophysical Research: Earth Surface*, 112(F2), doi:10.1029/2006JF000547, 2007.
- 610 Gruber, S., Hoelzle, M. and Haeberli, W.: Permafrost thaw and destabilization of Alpine rock walls in the hot summer of 2003, *Geophysical Research Letters*, 31(13), doi:10.1029/2004GL020051, 2004.
- Haberkorn, A., Phillips, M., Kenner, R., Rhyner, H., Bavay, M., Galos, S. P. and Hoelzle, M.: Thermal regime of rock and its relation to snow cover in steep alpine rock walls: gemsstock, central swiss alps, *Geografiska Annaler: Series A, Physical Geography*, 97(3), 579–597, doi:10.1111/geoa.12101, 2015.
- 615

- Hales, T. C. and Roering, J. J.: Climatic controls on frost cracking and implications for the evolution of bedrock landscapes, *Journal of Geophysical Research: Earth Surface*, 112(F2), doi:10.1029/2006JF000616, 2007.
- Hanssen-Bauer, I., Førland, E., Hisdal, H., Mayer, S., Sandø, A., Sorteberg, A., Adakudlu, M., Andresen, J., Bakke, J., Beldring, S., Benestad, R., van der Bilt, W., Bogen, J., Borstad, C., Breili, K., Breivik, O., Børsheim, K., Christiansen, H., Dobler, A. and Wong, W.: Climate in Svalbard 2100 - a knowledge base for climate adaptation, NCCS report., 2019.
- Harris, C., Davies, M. C. R. and Etzelmüller, B.: The assessment of potential geotechnical hazards associated with mountain permafrost in a warming global climate, *Permafrost and Periglacial Processes*, 12(1), 145–156, doi:10.1002/ppp.376, 2001.
- Harris, C., Arenson, L. U., Christiansen, H. H., Etzelmüller, B., Frauenfelder, R., Gruber, S., Haeberli, W., Hauck, C., Hölzle, M., Humlum, O., Isaksen, K., Kääb, A., Kern-Lütschg, M. A., Lehning, M., Matsuoka, N., Murton, J. B., Nötzli, J., Phillips, M., Ross, N., Seppälä, M., Springman, S. M. and Vonder Mühll, D.: Permafrost and climate in Europe: Monitoring and modelling thermal, geomorphological and geotechnical responses, *Earth-Science Reviews*, 92(3), 117–171, doi:10.1016/j.earscirev.2008.12.002, 2009.
- Hermanns, R. L., Blikra, L. H., Naumann, M., Nilsen, B., Panthi, K. K., Stromeyer, D. and Longva, O.: Examples of multiple rock-slope collapses from Köfels (Ötztal valley, Austria) and western Norway, *Engineering Geology*, 83(1), 94–108, doi:10.1016/j.enggeo.2005.06.026, 2006.
- Hersbach, H.: The ERA5 Atmospheric Reanalysis., in AGU Fall Meeting Abstracts, vol. 33. [online] Available from: <http://adsabs.harvard.edu/abs/2016AGUFMNG33D.01H> (Accessed 2 November 2020), 2016.
- Hop, H. and Wiencke, D.: The Ecosystem of Kongsfjorden, Svalbard, Springer International Publishing, Cham., 2019.
- Isaksen, K., Sollid, J. L., Holmlund, P. and Harris, C.: Recent warming of mountain permafrost in Svalbard and Scandinavia, *Journal of Geophysical Research: Earth Surface*, 112(F2), doi:10.1029/2006JF000522, 2007.
- Isaksen, K., Nordli, Ø., Førland, E. J., Łupikasza, E., Eastwood, S. and Niedźwiedź, T.: Recent warming on Spitsbergen—Influence of atmospheric circulation and sea ice cover, *Journal of Geophysical Research: Atmospheres*, 121(20), 11,913–11,931, doi:10.1002/2016JD025606, 2016.
- Johansson, A. M., Malnes, E., Gerland, S., Cristea, A., Doulgeris, A. P., Divine, D. V., Pavlova, O. and Lauknes, T. R.: Consistent ice and open water classification combining historical synthetic aperture radar satellite images from ERS-1/2, Envisat ASAR, RADARSAT-2 and Sentinel-1A/B, *Annals of Glaciology*, 1–11, doi:10.1017/aog.2019.52, 2020.
- Kastendeuch, P. P.: A method to estimate sky view factors from digital elevation models, *International Journal of Climatology*, 33(6), 1574–1578, doi:10.1002/joc.3523, 2013.
- Koven, C. D., Schuur, E. a. G., Schädel, C., Bohn, T. J., Burke, E. J., Chen, G., Chen, X., Ciais, P., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Jafarov, E. E., Krinner, G., Kuhry, P., Lawrence, D. M., MacDougall, A. H., Marchenko, S. S., McGuire, A. D., Natali, S. M., Nicolsky, D. J., Olefeldt, D., Peng, S., Romanovsky, V. E., Schaefer, K. M., Strauss, J., Treat, C. C. and Turetsky, M.: A simplified, data-constrained approach to estimate the permafrost carbon–climate feedback, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 373(2054), 20140423, doi:10.1098/rsta.2014.0423, 2015.
- Krautblatter, M., Verleysdonk, S., Flores-Orozco, A. and Kemna, A.: Temperature-calibrated imaging of seasonal changes in permafrost rock walls by quantitative electrical resistivity tomography (Zugspitze, German/Austrian Alps), *Journal of Geophysical Research: Earth Surface*, 115(F2), doi:10.1029/2008JF001209, 2010.

- Krautblatter, M., Funk, D. and Günzel, F. K.: Why permafrost rocks become unstable: a rock–ice-mechanical model in time and space, *Earth Surface Processes and Landforms*, 38(8), 876–887, doi:10.1002/esp.3374, 2013.
- 655 Kuhn, D., Redfield, T. F., Hermanns, R. L., Fuchs, M., Torizin, J. and Balzer, D.: Anatomy of a mega-rock slide at Forkastningsfjellet, Spitsbergen and its implications for landslide hazard and risk considerations, *Norwegian Journal of Geology*, 99(1), 41–61, doi:10.17850/njg99-1-03, 2019.
- Lewkowicz, A. G., Bonnaventure, P. P., Smith, S. L. and Kuntz, Z.: Spatial and thermal characteristics of mountain permafrost, northwest Canada, *Geografiska Annaler: Series A, Physical Geography*, 94(2), 195–213, doi:10.1111/j.1468-0459.2012.00462.x, 2012.
- 660 Liestøl, O.: Pingos, springs and permafrost in Spitsbergen, *Norsk Polarinstitutt Årbok* 1975, 7–29, 1977.
- Magnin, F., Deline, P., Ravanel, L., Noetzli, J. and Pogliotti, P.: Thermal characteristics of permafrost in the steep alpine rock walls of the Aiguille du Midi (Mont Blanc Massif, 3842 m a.s.l.), *The Cryosphere*, 9(1), 109–121, doi:https://doi.org/10.5194/tc-9-109-2015, 2015.
- 665 Magnin, F., Westermann, S., Pogliotti, P., Ravanel, L., Deline, P. and Malet, E.: Snow control on active layer thickness in steep alpine rock walls (Aiguille du Midi, 3842ma.s.l., Mont Blanc massif), *CATENA*, 149, 648–662, doi:10.1016/j.catena.2016.06.006, 2017.
- Magnin, F., Etzelmüller, B., Westermann, S., Isaksen, K., Hilger, P. and Hermanns, R. L.: Permafrost distribution in steep rock slopes in Norway: measurements, statistical modelling and implications for geomorphological processes, *Earth Surface Dynamics*, 7(4), 1019–1040, doi:https://doi.org/10.5194/esurf-7-1019-2019, 2019.
- 670 Maturilli, M.: Continuous meteorological observations at station Ny-Ålesund (2015-10 to 2019-12), reference list of 51 datasets, Alfred Wegener Institute - Research Unit Potsdam, PANGAEA, doi:10.1594/PANGAEA.914207, 2020a.
- Maturilli, M.: Continuous meteorological observations at station Ny-Ålesund (2020-01), Alfred Wegener Institute - Research Unit Potsdam, PANGAEA, doi:10.1594/PANGAEA.914805, 2020b.
- 675 Maturilli, M.: Continuous meteorological observations at station Ny-Ålesund (2020-02), Alfred Wegener Institute - Research Unit Potsdam, PANGAEA, doi:10.1594/PANGAEA.914807, 2020c.
- Maturilli, M.: Continuous meteorological observations at station Ny-Ålesund (2020-03), Alfred Wegener Institute - Research Unit Potsdam, PANGAEA, doi:10.1594/PANGAEA.914808, 2020d.
- 680 Maturilli, M. and Kayser, M.: Arctic warming, moisture increase and circulation changes observed in the Ny-Ålesund homogenized radiosonde record, *Theor Appl Climatol*, 130(1), 1–17, doi:10.1007/s00704-016-1864-0, 2017.
- Maturilli, M., Herber, A. and König-Langlo, G.: Climatology and time series of surface meteorology in Ny-Ålesund, Svalbard, *Earth System Science Data*, 5(1), 155–163, doi:https://doi.org/10.5194/essd-5-155-2013, 2013.
- Maturilli, M., Herber, A. and König-Langlo, G.: Surface radiation climatology for Ny-Ålesund, Svalbard (78.9° N), basic observations for trend detection, *Theoretical and Applied Climatology*, 120(1), 331–339, doi:10.1007/s00704-014-1173-4, 2015.
- 685 Monin, A. S. and Obukhov, A. M.: Basic laws of turbulent mixing in the surface layer of the atmosphere, *Tr. Akad. Nauk SSSR Geophiz. Inst.*, 24(151), 163–187, 1954.

- Müller, M., Homleid, M., Ivarsson, K.-I., Køltzow, M. A. Ø., Lindskog, M., Midtbø, K. H., Andrae, U., Aspelien, T., Berggren, L., Bjørge, D., Dahlgren, P., Kristiansen, J., Randriamampianina, R., Ridal, M. and Vignes, O.: AROME-MetCoOp: A Nordic Convective-Scale Operational Weather Prediction Model, *Weather and Forecasting*, 32(2), 609–627, doi:10.1175/WAF-D-16-0099.1, 2017.
- NCAR: CLM4.0 Offline Model Forcing Data Archived from CCSM4 historical and RCP simulations, [online] Available from: https://www.cesm.ucar.edu/models/cesm1.0/clm/clm_ccsm4forcingdata_esg.html (Accessed 7 July 2020), 2016.
- Noetzli, J. and Gruber, S.: Transient thermal effects in Alpine permafrost, *The Cryosphere*, 3(1), 85–99, doi:<https://doi.org/10.5194/tc-3-85-2009>, 2009.
- Nordli, Ø., Wyszyński, P., Gjeltén, H. M., Isaksen, K., Łupikasza, E., Niedźwiedź, T. and Przybylak, R.: Revisiting the extended Svalbard Airport monthly temperature series, and the compiled corresponding daily series 1898–2018, *Polar Research*, 39, doi:10.33265/polar.v39.3614, 2020.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H. H., Dashtseren, A., Delaloye, R., Elberling, B., Etzelmüller, B., Kholodov, A., Khomutov, A., Kääb, A., Leibman, M. O., Lewkowicz, A. G., Panda, S. K., Romanovsky, V., Way, R. G., Westergaard-Nielsen, A., Wu, T., Yamkhin, J. and Zou, D.: Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale, *Earth-Science Reviews*, 193, 299–316, doi:10.1016/j.earscirev.2019.04.023, 2019.
- Ødegård, R. S. and Sollid, J. L.: Coastal cliff temperatures related to the potential for cryogenic weathering processes, western Spitsbergen, Svalbard, *Polar Research*, 12(1), 95–106, doi:10.3402/polar.v12i1.6705, 1993.
- Pedersen, C.: Zeppelin Webcam Time Series [Data set], Norwegian Polar Institute, doi:10.21334/npolar.2013.9fd6dae0, 2013.
- Ravanel, L., Allignol, F., Deline, P., Gruber, S. and Ravello, M.: Rock falls in the Mont Blanc Massif in 2007 and 2008, *Landslides*, 7(4), 493–501, doi:10.1007/s10346-010-0206-z, 2010.
- Ravanel, L., Magnin, F. and Deline, P.: Impacts of the 2003 and 2015 summer heatwaves on permafrost-affected rock-walls in the Mont Blanc massif, *Science of The Total Environment*, 609, 132–143, doi:10.1016/j.scitotenv.2017.07.055, 2017.
- Seity, Y., Brousseau, P., Malardel, S., Hello, G., Bénard, P., Bouttier, F., Lac, C. and Masson, V.: The AROME-France Convective-Scale Operational Model, *Monthly Weather Review*, 139(3), 976–991, doi:10.1175/2010MWR3425.1, 2011.
- van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J. and Rose, S. K.: The representative concentration pathways: an overview, *Climatic Change*, 109(1), 5, doi:10.1007/s10584-011-0148-z, 2011.
- Walczowski, W. and Piechura, J.: Influence of the West Spitsbergen Current on the local climate, *International Journal of Climatology*, 31(7), 1088–1093, doi:10.1002/joc.2338, 2011.
- Westermann, S., Lüers, J., Langer, M., Piel, K. and Boike, J.: The annual surface energy budget of a high-arctic permafrost site on Svalbard, Norway, *The Cryosphere*, 3(2), 245–263, doi:<https://doi.org/10.5194/tc-3-245-2009>, 2009.
- Westermann, S., Langer, M., Boike, J., Heikenfeld, M., Peter, M., Etzelmüller, B. and Krinner, G.: Simulating the thermal regime and thaw processes of ice-rich permafrost ground with the land-surface model CryoGrid 3, *Geoscientific Model Development*, 9(2), 523–546, doi:<https://doi.org/10.5194/gmd-9-523-2016>, 2016.

725

Table 1: Settings of surface temperature loggers used in this study at eight different locations RW01 – RW08 in the surroundings of Ny-Ålesund / Svalbard.

Location	Site	Time period	Aspect
RW01	Non-coastal	27.08.2017 – 27.08.2020	NE
RW02	Non-coastal	27.08.2016 – 27.08.2020	NE
RW03	Non-coastal	27.08.2016 – 27.08.2020	NE
RW04	Open fjord	27.08.2017 – 27.08.2020	NE
RW05	Open fjord	31.08.2016 – 27.08.2020	N
RW06	Open fjord	12.05.2017 – 27.08.2020	ENE
RW07	Open fjord	12.05.2017 – 27.08.2020	NE
RW08	Bay	31.08.2016 – 27.08.2020	NE

730 **Table 2: Model parameters assumed in the simulations.**

Parameter		Value	Unit
Albedo rock wall	α	0.30	[-]
Albedo ground	α_g	0.15	[-]
Albedo open water	α_w	0.1	[-]
Albedo melting snow / ice	α_s	0.55	[-]
Emissivity	ε	0.97	[-]
Roughness length	z_0	0.018	[m]
Mineral fraction	$mineral$	0.97	[-]
Water and ice fraction	$waterIce$	0.03	[-]
Water bucket depth	d	0.001	[m]

Table 3: Distances to the open water body used for the linear interpolation of air temperature and logger locations where the distances are applied to.

Site	Logger	Distance [m]
Station Bayelva		1300 m
Station Ny-Ålesund		300 m

735

Non-coastal loggers	RW01-RW03	600 m
Coastal loggers	RW04-RW08	0 m
Ice cover in the bay	RW08	+ 300 m
Ice cover in the fjord	RW01-RW08	+ 600 m

Table 4: Model scenarios and corresponding settings with the three varying parameters source for air temperature, source for long-wave heat emission and albedo of the foot of the slope.

Scenario name	Representation	State	Albedo	Long-wave radiation source
Non-coastal scenario	RW01, RW02, RW03	No snow	Terrain = 0.15	T_{air}
		Snow	Ice / snow = 0.55	
Open fjord scenario	RW04, RW05, RW06, RW07	Unfrozen	Water = 0.1	T_{sea}
		Frozen	Ice / snow = 0.55	T_{air}
Frozen bay scenario	RW08	Unfrozen	Water = 0.1	T_{sea}
		Frozen	Ice / snow = 0.55	T_{air}

740

Table 5: Summary statistics of the model validation including RMSE, bias, R^2 and NSE. The statistics were calculated for all locations RW01 to RW08, comprising each entirely recorded month for the measurement periods stated in Table 1.

Location	Site	RMSE	bias	R^2	NSE
RW01	Non-coastal	0.8	0.0	0.989	0.986
RW02	Non-coastal	1.0	0.3	0.984	0.981
RW03	Non-coastal	0.9	0.4	0.989	0.986
RW04	Open fjord	0.7	0.3	0.991	0.988
RW05	Open fjord	1.1	-0.5	0.978	0.968
RW06	Open fjord	1.0	-0.1	0.980	0.980
RW07	Open fjord	1.1	0.2	0.980	0.969
RW08	Bay	1.2	0.3	0.992	0.977

745

Table 6: Measured MARST and mean RST in the winter season (Dec – Feb) for all locations RW01 to RW08. Lack of data results from either snow cover on the logger or missing records. MARST are coldest in non-coastal settings (RW01 – RW03). Mean RST in winter season are found to be coldest in non-coastal settings, closely followed by settings in the bay (RW08), while settings at the open fjord show highest RST (RW04 – RW07). For colour code see Figure 1.


Location	Entire year				Winter: Dec - Feb			
	2016/17	2017/18	2018/19	2019/20	2016/17	2017/18	2018/19	2019/20
RW01	-	-1.5	-	-	-8.1	-6.7	-9.0	-13.1
RW02	-2.2	-1.8	-2.4	-4.3	-8.5	-6.6	-9.6	-13.5
RW03	-1.8	-2.0	-2.4	-4.1	-9.2	-7.2	-10.0	-13.8
RW04	-	-1.0	-2.1	-3.6	-	-5.1	-8.7	-12.0
RW05	-0.9	-0.8	-	-	-6.6	-5.2	-7.2	-10.4
RW06	-	-0.6	-	-	-	-5.0	-8.5	-11.5
RW07	-	-0.8	-	-3.6	-	-4.6	-7.6	-11.1
RW08	-1.4	-0.9	-2.1	-	-8.6	-5.8	-9.7	-13.1

750

Table 7: Modelled mean RST with frozen conditions in the bay or sea ice in the fjord. Frozen conditions in the bay lead to a local cooling of RW08, while sea ice result in similar RST for all settings. RW02, RW04 and RW08 were selected as they have the same aspect but different settings. For colour code see Figure 1.

Location	Site	Dec	Jan	Feb	Mar	Apr
Mean RST of the days with a frozen bay						
RW02	Non-coastal	-11.8	-13.5	-13.9	-15.0	-7.6
RW04	Open fjord	-10.2	-10.6	-11.7	-13.4	-6.8
RW08	Bay	-11.2	-12.3	-13.0	-14.4	-7.2
Mean RST of the days with widespread sea ice in the fjord						
RW02	Non-coastal	-	-	-17.3	-15.0	-10.9
RW04	Open fjord	-	-	-17.2	-15.0	-10.9
RW08	Bay	-	-	-17.2	-15.0	-10.9

Figures



760 **Figure 1: locations of installed logger in the canyon (blue labels: RW01 to RW03), at the coastal cliffs (red labels: RW04 to RW07) and in the bay Thiisbukta (green label: RW08).**
Source: NP_Basiskart_Svalbard_WMTS_25833 / FKB_Svalbard_WMTS_25833, ETRS 89 UTM 33 © Norsk Polarinstitutt (npolar.no).

765




Figure 2: Locations of rock wall loggers used in this study: (a) coastal cliffs at the open fjord next to Ny-Ålesund airport (tidal zone visible in bottom); (b) non-coastal rock walls in the canyon of Bayelva; (c) close-up of a rock wall logger.

770






Figure 3: Mean measured daily RST in winter 2017. RW01 shows a temporarily damped signal due to snow cover.



26. January 2020



03. March 2020





Figure 4: Mean measured daily RST for two different conditions in Kongsfjorden: (a) The time period 01.01.-31.01.2020 was mainly characterized by a frozen bay and an open fjord. RST at coastal cliffs show higher values than RST at the non-coastal rock walls and at rock walls in the bay. The most pronounced differences are found with RST below -10° C. (b) At the time period 15.02.-15.03.2020, the bay and the fjord were predominantly frozen. In this case, no clear differences could be observed between the three settings. RW02, RW04 and RW08 were selected as they have the same aspect but different settings.



785 **Figure 5: Mean modelled vs. measured monthly RST for all locations RW01 to RW08, comprising each entirely recorded month for the measurement periods stated in Table 1.**




Figure 6: Temperature difference between the *open fjord scenario* (reference) and different model scenarios. “Colder air temperature”: as reference, but using colder air temperature as distance to the open water body is enlarged; “No seawater long-wave emission”: as reference but using air temperature instead of seawater temperature; “Reflected short-wave radiation”: as reference but assuming ice albedo for the surrounding terrain; “Total difference”: *frozen bay scenario* combining all three effects.




Figure 7: Surface energy balance for the seasons of the year for a vertical rock wall with an aspect of 40° (RW02, RW04 and RW08). Most pronounced differences in the scenarios are found in winter and spring, while summer and fall show similar fluxes. SW = net short-wave radiation; LW = net long-wave radiation; Qe = latent heat flux; Qh = sensible heat flux; G = ground heat flux.




Figure 8: MARST of past and future simulations with the settings of the *non-coastal scenario* and the *open fjord scenario* with RCP2.6, RCP4.5 and RCP8.5. The grey box shows the measurement period.




Figure 9: Active layer thickness of past and future simulations with the settings of the *non-coastal scenario* and the *open fjord scenario* with RCP2.6, RCP4.5 and RCP8.5.