

Divergence of riparian forest composition and functional traits from natural succession along a degraded river with multiple stressor legacies

Philippe Janssen, John C. Stella, Hervé Piégay, Bianca Räpple, Bernard Pont, Jean-Michel Faton, Johannes Hans C. Cornelissen, André Evette

► To cite this version:

Philippe Janssen, John C. Stella, Hervé Piégay, Bianca Räpple, Bernard Pont, et al.. Divergence of riparian forest composition and functional traits from natural succession along a degraded river with multiple stressor legacies. Science of the Total Environment, 2020, 721, pp.15/137730. 10.1016/j.scitotenv.2020.137730. hal-03026265

HAL Id: hal-03026265 https://hal.science/hal-03026265

Submitted on 26 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

| 1 | Divergence of riparian forest composition and functional traits from natural succession along a degraded |
|----|--|
| 2 | river with multiple stressor legacies |
| 3 | |
| 4 | |
| 5 | |
| 6 | Janssen, Philippe ^{1,2} *, Stella, John C. ³ , Piégay, Hervé ¹ , Räpple, Bianca ¹ , Pont, Bernard ⁴ , Faton, Jean-Michel ⁴ , |
| 7 | Cornelissen, Johannes Hans C. ⁵ , Evette, André ² |
| 8 | |
| 9 | ¹ Univ. Lyon, UMR 5600 Environnement Ville société, CNRS, Site of ENS Lyon, Lyon, France |
| 10 | ² Univ. Grenoble Alpes, INRAE, LESSEM, St-Martin-d'Hères, France |
| 11 | ³ Department of Forest and Natural Resources Management, State University of New York College of |
| 12 | Environmental Science and Forestry, Syracuse, NY, 13210 USA |
| 13 | ⁴ Réserves Naturelles de France, Quétigny, France |
| 14 | ⁵ Department of Ecological Science, Faculty of Science, Vrije Universiteit Amsterdam, Amsterdam, The |
| 15 | Netherlands |
| 16 | |
| 17 | * Corresponding author, e-mail philippe.janssen@inrae.fr, phone +33 476762879 |
| 18 | |
| 19 | Email addresses of other authors: |
| 20 | J.C. Stella (<u>stella@esf.edu</u>), H. Piégay (<u>herve.piegay@ens-lyon.fr</u>), B. Rapple (<u>b.raepple@yahoo.de</u>), B. Pont |
| 21 | (bernard.pont@cen-isere.org), J-M. Faton (jmfaton@lagaredesramieres.com), J.H.C. Cornelissen |
| 22 | (j.h.c.cornelissen@vu.nl), A. Evette (andre.evette@inrae.fr) |
| 23 | |

24 Running head: Accelerated succession on degraded river

25 Abstract

26 Prolonged exposure to human induced-stressors can profoundly modify the natural trajectory of ecosystems. 27 Predicting how ecosystems respond under stress requires understanding how physical and biological properties 28 of degraded systems parallel or deviate over time from those of near-natural systems. Utilizing comprehensive 29 forest inventory datasets, we used a paired chronosequence modelling approach to test the effects of long-30 term channelization and flow regulation of a large river on changes in abiotic conditions and related riparian 31 forest attributes across a range of successional phases. By comparing ecological trajectories between the highly 32 degraded Rhône and the relatively unmodified Drôme rivers, we demonstrated a rapid, strong and likely 33 irreversible divergence in forest succession between the two rivers. The vast majority of metrics measuring life 34 history traits, stand structure, and community composition varied with stand age but diverged significantly 35 between rivers, concurrent with large differences in hydrologic and geomorphic trajectories. Channelization 36 and flow regulation induced a more rapid terrestrialization of the river channel margins along the Rhône River 37 and accelerated change in stand attributes, from pioneer-dominated stands to a mature successional phase 38 dominated by non-native species. Relative to the Drôme, dispersion of trait values was higher in young forest 39 stands along the Rhône, indicating a rapid assembly of functionally different species and an accelerated 40 transition to post-pioneer communities. This study demonstrated that human modifications to the hydro-41 geomorphic regime have induced acute and sustained changes in environmental conditions, therefore altering 42 the structure and composition of riparian forests. The speed, strength and persistence of the changes suggest 43 that the Rhône River floodplain forests have strongly diverged from natural systems under persistent multiple 44 stressors during the past two centuries. These results reinforce the importance of considering historical 45 changes in environmental conditions to determine ecological trajectories in riparian ecosystems, as has been 46 shown for old fields and other successional contexts.

47

Key-words: riparian forests, ecological succession, community-weighted traits, human induced-stressors,
 ecological trajectories

50 Introduction

51 The concept of ecological succession dates almost to the origins of the field of ecology itself. Over the 52 last century, conceptual frameworks of how plant communities assemble (or re-assemble) following large 53 perturbations have evolved from a focus on pattern to the underlying mechanisms, notably the feedbacks that occur between abiotic and biotic ecosystem components as they develop over time (Connell and Slatyer 1977, 54 55 Pickett et al. 1987, Meiners et al. 2015). This shift in focus has come with an increasing appreciation of 56 ecological complexity, particularly the effects of multiple interacting drivers and stressors, thresholds and other 57 non-linear responses, and the importance of contingent factors (e.g., variation in historical management 58 strategies or in the intensity and duration of management) in driving community trajectories (Clark et al. 2019, 59 Chang and Turner 2019). Nevertheless, synthesizing general principles and testing predictions of ecological 60 succession across diverse ecosystems remain a challenge. The recent focus on monitoring change in ecological 61 traits, rather than species composition, in plant assemblages over time shows promise in distilling patterns and 62 drivers of community dynamics in response to shifts in both extrinsic factors (e.g., disturbance, climate) and 63 intrinsic ones (e.g., soil properties, competition) (Meiners et al. 2015, Chang and Turner 2019).

64 The trait-based approach is particularly well-suited for studying natural communities that are severely 65 degraded by multiple, interacting human stressors; these include resource exploitation (e.g., overfishing), 66 exotic species invasions, and changes in land use, climate and disturbance regimes due to human activities 67 (Mouillot et al. 2013). As with natural disturbances, human-induced stressors can greatly alter ecological 68 trajectories of both terrestrial and aquatic ecosystems, and lead to rapid and potentially irreversible shifts in 69 ecosystems properties (Folke et al. 2004). Numerous studies document divergences in successional trajectories 70 of species assemblages and functional traits between altered and reference ecosystems (e.g., Odion et al. 2010, 71 Sfair et al. 2016, Clark et al. 2019), in many cases leading to alternative ecosystem states (Beisner et al. 2003). 72 Strong shifts in species dominance during succession due to native biodiversity loss and/or invasion by exotic 73 species may impede the ecological resilience of ecosystems, and compromise efforts to restore them or 74 preserve their functions and services (Suding 2011).

Riparian ecosystems are among the most modified by human activities (Tonkin et al. 2018) and
comprise some of the most vulnerable biomes to ongoing global change (Perry et al. 2012, Stella et al. 2013b).
More than most forest ecosystems, riparian woodlands comprise a fragmented patch mosaic of diverse
composition and age, which are related to strong physical gradients and flood history (Naiman and Decamps

79 1997, Scott et al. 1997, Bendix and Stella 2013). Abiotic conditions and plant species composition can change 80 markedly within only a few decades during riparian succession (Johnson et al. 1976, Osterkamp and Hupp 81 2010), and community trajectories can be interrupted and superimposed by new disturbances. Nevertheless, 82 riparian disturbance regimes present an excellent opportunity to study succession at local scale. Frequent flood disturbance and the close proximity of forest stands that have colonized floodplain surfaces of different ages 83 84 allow the use of chronosequences, or space-for-time proxies, to study successional processes in riparian 85 communities (Stella et al. in review, Schnitzler 1995, Scott et al. 1997, Fierke and Kauffman 2005). In the 86 absence of long-term longitudinal datasets, which with few exceptions are virtually absent for riparian areas 87 (but see the case of the Missouri River, USA in Johnson et al. 2012), this approach is particularly suited to 88 understand how community dynamics along highly modified rivers may diverge from historical or reference 89 conditions (Prach and Walker 2011).

90 To date, the effect of human induced-modifications on riparian communities have mostly been studied 91 by adopting a comparative approach. By comparing channelized versus unchannelized rivers (Dufour, Barsoum, 92 Muller, & Piégay, 2007; Nakamura, Jitsu, Kameyama, & Mizugaki, 2002; Oswalt & King, 2005) or regulated 93 versus unregulated rivers (Merritt and Cooper 2000, Kui et al. 2017, Bejarano et al. 2018), negative impacts to 94 riparian communities have been documented, including reduced native species and functional trait diversity, 95 increased invasion by exotic species, and homogenised forest stand structure. Despite widespread degradation 96 to riparian systems and the substantial investigations into the ecological mechanisms that drive succession 97 (Chang and Turner 2019), few studies have examined how modifications to river flow and sediment regimes 98 impact the successional trajectory of riparian plant communities, and none to our knowledge have analysed 99 these patterns in terms of functional trait dynamics. Yet, understanding both the current status and future 100 trajectory of degraded riparian forest ecosystems is critical for defining and prioritizing effective strategies to 101 conserve and restore them.

In this context, we aimed to understand how the current ecological trajectory of riparian forests along
 a large, channelized and regulated river in eastern France either paralleled or deviated from a relatively
 unmodified local reference system, across a range of successional phases. Using datasets of abiotic drivers –
 flow and sediment regimes – and biotic responses that included functional, structural and compositional
 attributes, we modeled chronosequences on both rivers to test for deviation in their ecological trajectories.
 The chronosequence approach is well-suited for assessing community shifts during plant succession, especially

if these changes are predictable over time (Walker et al. 2010). Moreover, comparing trends in species and
trait composition over time can help to identify the mechanisms causing divergence in ecological trajectories,
and thus potential barriers to restoration (Suding 2011, Chang and Turner 2019). Based on this scheme we
addressed the following two questions: (i) How do abiotic and biotic conditions vary with forest stand age along
a heavily impacted river compared to a more natural reference ecosystem? (ii) Are successional trajectories of
riparian forests different between both river systems, in terms of species composition and functional trait
trends over a long chronosequence?

115

116 Materials and methods

117 Study area

118 The study was carried out along the mainstem Rhône River (total length = 810 km, catchment area = 119 96,500 km², mean annual discharge = 1,700 m³ s⁻¹) and the Drôme River (total length = 110 km, catchment area = 1,663 km², mean annual discharge = 20 m³ s⁻¹), a tributary of the Rhône (Figure 1). Both rivers are located in 120 121 SE France and experience a temperate climate with mean annual temperature and precipitation of 13.6°C and 122 755 mm in the southern part and of 11.9°C and 782 mm in the northern part. Despite differences in catchment 123 size and discharge magnitude, the rivers prior to human modification shared similar snowmelt flow regimes 124 and sediment dynamics. The riparian forests of both rivers contain a mix of early successional trees including 125 Populus nigra and Salix alba, post-pioneers such as Fraxinus excelsior and Acer platanoides, and understorey species including Crataegus monogyna and Sambucus nigra. The most abundant newcomers to the species 126 127 pool include the non-native trees Acer negundo and Robinia pseudoacacia (See Appendix S1 for further details 128 on tree species' characteristics).

The Drôme River is a free-flowing river with a channel mostly unconstrained by human infrastructure and a shifting braided pattern throughout its downstream portion, which constitutes the study reach. Bedload is still actively transported in the lower section despite channel incision due to sediment mining that occurred during the 1970s. Since the second half of the 20th century, the Drôme River has also been subject to land-use changes in the floodplain (Liébault and Piégay 2002). Both mining and land-use changes have favored establishment of native riparian woodlands within the braided section, which is subject to periodic floods that scour older forest patches and induce sediment deposition and subsequent initiation of new stands. As a

consequence, the riparian forest is characterized by a complex mosaic of patches of different ages (Räpple etal. 2017).

138 The lower Rhône River, downstream of the city of Lyon, is a highly modified river, which has shifted 139 from a braided pattern to a series of impounded reaches within the span of approximatively one century 140 (Olivier et al. 2009, Bravard 2010). Two main historical management phases have greatly modified the river during this period. During the 19th Century the river was engineered and channelized of longitudinal 141 142 submersible dikes and lateral dikes, forming more or less rectangular compartments, called "casiers Girardon" 143 (see 1954 photo in Figure 1). These structures were constructed within the river's main channel and arranged 144 sequentially along extensive reaches of the river to concentrate flow into a single narrow channel, thus 145 facilitating navigation (see Thorel et al. 2018 for further details). In the second management phase, nineteen 146 hydropower plants were built along the French part of the Rhône River during the second half of the 20th 147 Century, among which sixteen plants were built on artificial channels that bypass 162 km out of 522 km of the 148 river (Lamouroux et al. 2015). Each of these hydropower works contains a diversion dam that impounds a 149 reservoir upstream and conveys a large part of the river discharge into the diversion canal on which the power 150 plant is located for energy production (for further details see, Appendix S2 and Vázquez-Tarrío et al. 2019). As a 151 consequence, the historical bypassed channels, also called the old Rhône, receives a minimum discharge most 152 of the year, on average 5% of the natural discharge (Bravard and Gaydou 2015). Thus, these bypassed channels 153 which had been previously channelized are also significantly impacted by river regulation, which have induce 154 channel incision, lateral stabilization and armouring of the channel bed. Along the highly artificial, stable river 155 margins of the bypassed channels, many of the Girardon structures filled with overbank fine sediment and 156 were subsequently colonized by woody pioneer species and succeeded to mature riparian forests (Räpple

157 2018).

158 Riparian forest inventory sampling design

Forest inventories were conducted along both the Drôme and Rhône rivers and confined to established stands in the active and abandoned floodplain of each river. To ensure independence and avoid edge influences, all inventory plots were established > 60 m away from any other plot, were located in forested patches > 0.5 ha and were > 20 m from the nearest stand edge. Along the Drôme River, 69 plots were sampled in the summer of 1994 within the Natural Reserve of the Ramières, in which large patches of riparian forest remain. The plots were distributed between two braided reaches, west (n = 19 plots) and east (n = 50)

(Appendix S3). Along the Rhône River, 65 plots were surveyed along four bypassed channels (*i.e.*, the former
active channel) in summer 2015. From upstream to downstream, these reaches were: Pierre-Bénite (PBN, n =
16 plots), Péage de Roussillon (PDR, n = 14), Montélimar (MON, n = 18) and Donzère-Mondragon (DZM, n = 17)
(Appendix S3). Currently, the Girardon structures in these four bypassed channels support the largest extent of
riparian forests remaining along the lower Rhône River downstream of Lyon.

170 On the Drôme River, stands were characterized using a 13.8-m-radius circular plot (area: 597 m²), 171 centered on a systematic 130 x 130 m grid covering the whole forest area of the Natural Reserve. Within each 172 plot, all standing trees with a diameter at breast height (DBH) ≥ 7.5 cm were recorded. On the Rhône River, a 173 stratified sampling scheme was implemented, with plots randomly located within floodplain areas of known 174 age. Rhône inventories used two concentric plot sizes, with a 10-m radius plot (area: 314 m²) for recording 175 standing trees with a DBH \ge 7.5 cm, and a larger plot of 20-m radius (area: 1,256 m²) used to record standing 176 trees with DBH \ge 30 cm. For each tree, species, DBH and health class were recorded, comprising live trees (< 177 50% dead crown and branches), dying trees (>50% dead crown and branches), or dead trees.

178 To accurately compare stand attributes between the two rivers, which were subject to different forest 179 inventory campaigns, data were standardized to a per hectare basis and common metrics in both systems 180 describing riparian forest composition and structure were used (Table 1). Basal area (m² ha⁻¹), mean diameter (cm) and stem density (n ha^{-1}) were calculated for total live and total dead standing trees, as well as subsets for 181 182 total exotic trees, and standing live trees of several genera: Populus spp. (i.e., P. alba, P. x canescens, P. nigra, 183 P. tremula and P. x canadensis), Fraxinus spp. (i.e., F. angustifolia and F. excelsior) and Acer spp. (i.e., A. 184 campestre, A. monspessulanum, A. negundo, A. opalus, A. platanoides and A. pseudoplatanus). Diversity 185 metrics of tree diameter classes (5-cm ranges across n = 14 classes) and tree species (n = 31) were computed 186 using the Shannon diversity index with integer values of basal area as the abundance weighting measure. 187 Overbank fine sediment depth was measured at the center of each plot using a soil corer that was 188 inserted into the fine sediment until the gravel interface was reached (Table 1). Mean and standard-deviation 189 of annual flow were calculated using daily data recorded between 1966 and 2017 for the Drôme River 190 (http://hydro.eaufrance.fr) and between 1920 and 2010 for the Rhône River (provided by the Compagnie 191 Nationale du Rhône). Because forest inventories on the two rivers were conducted in different periods (i.e., 192 Drôme = 1994, Rhône = 2015), we used a common time frame, fixed between 1940 and 2000, to compare 193 changes in hydrology but also to account for different switch-on dates of the diversion canals along the Rhône

River (i.e., DZM = 1952, MON = 1957, PBN = 1966, PDR = 1977). We used data from the Saillans measurement station on the Drôme River, 15 km upstream of the east sampling area, and from two stations on Rhône River, at Ternay within the PBN bypassed channel, and at Viviers within the MON bypassed channel. These two stations' data were then averaged to estimate unregulated flow prior to the diversions in each reach, whereas regulated flow levels were estimated using dam operation rules imposed following the diversions.

199 Trait data

200 Exotic tree species were distinguished from native tree species, using data available from the Baseflor 201 database (Julve 1998). For functional traits, we focused on three traits that typically vary across different life 202 history strategies as plants response to shifts in limiting resources and competitors during ecological 203 succession: specific leaf area (SLA; leaf area per dry mass), wood density, and dry seed mass. SLA is related to 204 resource acquisition and conservation strategies, contrasting "conservative" species with long-lived leaves to 205 "acquisitive" species with rapid turnover of plant leaves (Wright et al. 2004). Wood density is related to growth 206 strategy, contrasting fast-growing species with low wood density against slow-growing but stress-tolerant 207 species with high wood density (Chave et al. 2009). Seed mass is related to dispersal ability, contrasting species 208 with light seeds but high seedling mortality from species with large seeds but high long-term seedling survival 209 (Westoby 1998). In riparian environments, where large and rapid changes in abiotic stressors and available 210 resources drive community dynamics, this set of traits is expected to vary greatly among tree species and with 211 forest stand age. Due to contrasting life-history strategies between pioneer and post-pioneer riparian species, 212 we expected to see increasing trends across young to old forest stands in community weighted averages of SLA 213 (from rapid to longer leaf turnover), wood density (from faster to slower growth) and seed mass (from long-214 distance to more local dispersal).

215 Trait data were extracted from the TRY database, from which we calculated a mean trait value per 216 species, after removing all values with an error risk > 4 (Kattge et al. 2011, see Appendix S4). Missing values in 217 the wood density data were completed using the global wood density database (Zanne et al. 2009, Chave et al. 218 2009). For each trait value, we computed community-weighted means (CWM) and functional dispersion (FDis) 219 using basal area as a measure of relative abundance. CWM is defined as the mean of trait values weighted by 220 the relative abundance of each species bearing each value (Lavorel et al. 2008). FDis is defined as the mean 221 distance of individual species to the weighted centroid of all species in the assemblage and is unaffected by 222 difference in species richness among plots (Laliberté and Legendre, 2010).

223 Forest stand age reconstruction

224 Forest stand age was characterized using geo-referenced aerial photograph series in a Geographic 225 Information System (QGIS Development Team 2015). For the Drôme River, seven series of aerial photographs 226 were available, taken from 1932 to 1991. For the Rhône River, available information varied among reaches and 227 among plot positions within reach, with four to six series of aerial photographs available per reach, taken from 228 1938 to 2002 (see Appendix S3 for details). For each plot, the physiognomic unit was characterized visually 229 from aerial photographs into four categories: water, bare ground, shrub (i.e., sparse patches of low vegetation) 230 and forest units (i.e., dense patches of tall vegetation). The effective timing of forest establishment for each 231 plot was determined by averaging the last unforested series date (t = no forest cover) and the next one where 232 forest was evident (t+x = forest cover). When a time-series was too fragmented, i.e., > 20 years between two 233 superimposed aerial photographs, and when physiognomic units were "shrub" on the old aerial photograph 234 series and "forest" on the next series, the date of forest cover was approximated in between the two dates. To 235 confirm the continuity of the forest cover since the first date of appearance, posterior aerial photograph series 236 were also inspected (Appendix S3).

237 Statistical analysis

238 We used linear models to test how abiotic drivers and biotic responses varied across the forest stand 239 age (or years), and whether these trends differed significantly between rivers. For all models, we considered 240 forest stand age ("Age" in tables and figures) as a continuous predictor, as well as sediment depth 241 ("Sed depth"). Both of these variables were standardized prior to analysis. River (noted "River") was 242 considered as an independent, categorical factor. Abiotic drivers included the fine sediment depth as well as 243 the mean and standard deviation (SD) of annual flow. Because the discharge was largely different between the 244 two rivers, mean and SD of annual flow were standardized according to the range of their own values prior to 245 analysis. Biotic response included the community-weighted mean (CWM) and the dispersion (FDis) of 246 functional trait values, the basal area, the mean diameter and the stem density of stand attributes and of tree 247 genera as well as the diversity indices (Table 1). Because the seed mass of two tree species – Juglans regia and 248 Quercus pubescens – was very large, seed mass values were log-transformed before CWM and FDis calculation. 249 To investigate how abiotic drivers and biotic responses varied over time between the two rivers, we 250 fitted normal linear models (LMs) or log-normal LMs for skewed response variables (i.e., sediment depth, SLA, wood density, basal area, diameter and stem density). We developed a set of a priori models testing the 251

252 individual, additive and interactive effects of predictor variables, plus a null (i.e., intercept-only) model (see 253 Appendices S5 and S6). In all candidate models the variance inflation factor was < 3. The most parsimonious 254 model was identified using Akaike's information criterion corrected for small sample sizes (Burnham and 255 Anderson 2002) and goodness of fit was estimated using the adjusted coefficient of determination. For all of 256 the response variables, Moran's I values in the top-ranked model residuals was non-significant, indicating that 257 spatial patterns were accounted for by independent variables (Appendix S7). For each response variable, 258 parameter estimates and associated unconditional standard errors were extracted from the top-ranked model 259 and we checked for consistency among parameter estimates and confidence intervals on the subset of top 260 ranking models for which the delta AICc was < 7 (Appendix S5 and S6).

261 To investigate whether the forest structure and composition varied with forest stand age between the 262 two rivers we used multivariate generalized linear models (GLMs). We fitted the full model (Age x River + 263 Sed depth) to each diameter class/tree species, using basal area as the abundance measure, and summed 264 across the univariate responses to estimate their multivariate response with a negative binomial distribution. 265 The significance of independent variables in the multivariate GLM was assessed using an analysis of variance 266 with the PIT-trap method and 999 bootstrap resamples (Warton et al. 2017). To determine which diameter 267 classes/tree species best contribute to the overall model deviation, we calculated univariate test statistics and 268 p-values, and adjusted to correct for multiple testing for each diameter classes and tree species (Wang et al. 269 2012). Finally, to provide a graphical representation of the interaction between forest stand age and river, we 270 used a canonical analysis of principal coordinates (Anderson and Willis 2003), using a Bray-Curtis distance. All 271 analyses were performed with R version 3.5.3. (R Core Team 2019).

272

273 Results

274 Contrasting trends in abiotic drivers among rivers

Of the abiotic drivers analysed, sediment depth and mean annual flow had pronounced trends over time that differed significantly between the Rhône and Drôme rivers (Table 2; Figure 2). Thus, models that included an interaction term between river and forest stand age (for sediment depth) or river and year (for mean discharge) best predicted the respective response. Sediment depth increased more rapidly with forest stand age along the Rhône compared to the Drôme (Figure 2), whereas the mean annual flow strongly decreased over time along the Rhône, with no clear trend on the Drôme (Appendix S8). The standard-deviation

of annual flow decreased significantly over time (Figure 2) but did not differ significantly between the two

rivers (Appendix S8).

283 Trends in forest functional, structural and compositional attributes

Most of the whole-stand and genera-specific forest attributes, community-weighted functional traits, and diversity indices varied strongly with riparian forest stand age. Similar to the abiotic drivers, many of the trends in forest development also differed significantly between rivers. For a large fraction of the response variables, the top-ranked model included the interaction term between stand age and river, and some of these included sediment depth as a covariate (Table 2). The goodness of fit of these models ranged greatly, from 0.08 for the basal area of *Populus spp.*, to 0.71 for the mean diameter of *Populus* spp. (Table 2).

Parameters retained in the top ranked models indicate that only a few trends in forest characteristics were equivalent between rivers; these were overall mean seed mass, and for *Fraxinus* spp. only, their basal area and stem density (Table 3). Some characteristics varied at the same rate (i.e., parallel slopes for both rivers), but differed significantly in magnitude between the Drôme and Rhône. These included stem density for live and dead trees, which were consistently higher along the Drôme River. In contrast, stem density and mean tree diameter for exotic trees, mean tree diameter for *Fraxinus* spp. and *Acer* spp., as well as functional dispersion of SLA were significantly higher along the Rhône River (Table 1; Figure 3 and 4).

297 All others trait and stand-level responses were mediated by the interaction terms between river and 298 forest stand age (Tables 3). For example, the mean value for SLA and wood density (Figure 3), and the basal 299 area of exotic trees (Figure 4) increased more steeply with forest stand age along the Rhône River. Conversely, 300 the Drôme River riparian forest showed steeper trends over time for the trait dispersion of wood density and 301 seed mass (Figure 3), Acer spp. basal area and stem density (Figure 5), as well as the diversity of tree species 302 and of stem diameter classes (Figure 6). For some responses, the two rivers showed opposite trends. In 303 particular, the basal area and mean diameter of live and dead trees (Figure 4), as well as the basal area of 304 native Populus spp. (Figure 5), increased with forest stand age along the Drôme River but, counter to most 305 patterns of stand development, decreased along the Rhône River. The mean diameter of Populus trees was 306 much larger along the Rhône, and they achieved their maximum size much faster, within two decades (Figure 307 5). This, combined with the decreasing density of these large, dominant trees later in the successional 308 sequence had the effect of lowering Populus and overall stand basal area with time, in contrast to the Drôme, 309 where both metrics increased.

Finally, sediment depth was a significant predictor for all stand basal area fractions (live, dead, and exotic trees), for *Fraxinus* density, and for *Acer spp*. basal area, mean diameter and density. However, the effect of sediment was positive for *Acer* spp. basal area and density but negative for *Fraxinus* spp. density and *Acer* spp. mean diameter (Table 3).

314 Differences in forest composition and structure between rivers

315 The multivariate GLMs showed that forest structure was significantly influenced by forest stand age 316 (Dev = 14.5, P = 0.031), river (Dev = 73.8, P = 0.001), sediment depth (Dev = 22.3, P = 0.034) and the interaction 317 term between age and river (Dev = 20.7, P = 0.043). Based on the deviation explained by parameters, most of 318 the variation in forest structure was related to differences between rivers; this dichotomy is well-represented 319 by the separation of Rhône and Drôme plots along the first axis of the canonical analysis of principal 320 coordinates (CAP 1, Figure 7A). The second axis was more evidently related to forest age, though some sites 321 appeared to be well outside the predominant distribution. Univariate tests for each diameter class showed that 322 the largest trees (DBH > 70 cm) had the highest test statistic values for the effect size between rivers (Dev = 323 25.15, P_{adi} < 0.001) (Appendix S9). Consequently, divergence in forest structure between the two rivers was 324 mostly due to the greatest abundance of very large diameter trees within the Rhône River forests (mean basal 325 area/plot = 8.17 m² ha⁻¹) as compared to the Drôme River (mean basal area/plot = $0.11 \text{ m}^2 \text{ ha}^{-1}$). 326 As with forest structure, community composition was significantly influenced by river (Dev = 340.7, P = 327 0.001) and the interaction term between age and river (Dev = 49.6, P = 0.040); this is similarly evident in the 328 plot clustering along the first CAP axis (Figure 7B). However, community composition was much less influenced 329 than forest structure by stand age (Dev = 40.2, P = 0.063) and sediment depth (Dev = 33.2, P = 0.437). The CAP 330 plot shows that age classes were quite dispersed and randomly distributed along the second axis. Furthermore, 331 the Drôme River plots were more clustered than the Rhône River plots, indicating greater similarity in tree 332 species composition within the Drôme River corridor compared to the Rhône. Univariate tests for each tree 333 species showed that Acer negundo has the largest effect size between the rivers (Dev = 130.30, P_{adi} < 0.001) 334 (Appendix S10). Consequently, divergence in forest composition between the two rivers was in large part due 335 to the absence of this exotic species within the Drôme River forests (mean plot basal area = $0.00 \text{ m}^2 \text{ ha}^{-1}$) as 336 compared to the Rhône River (mean plot basal area = $5.37 \text{ m}^2 \text{ ha}^{-1}$).

337

338 Discussion

339 Overall, the successional trajectories of riparian forests along the channelized and regulated bypassed 340 channels of the Rhône River deviate from the expected historical trajectories, as represented by the 341 unchannelized and unregulated reaches of the Drôme River . Though the riparian forests of the Drôme River 342 are also under the influence of multiple human-induced stressors (Liébault and Piégay 2002), and though many 343 trends in species composition and community functional traits followed expected forest stand trajectories, 344 differences between the two rivers were stronger than the age gradient, which spanned 70 years in this study. 345 Because the biogeographic setting and potential species pools are similar, these results strongly suggest that 346 large differences in local environmental conditions are driving riparian forest development patterns (Clark et al. 347 2019). Many of the shifts along the Rhône River chronosequence were more pronounced and rapid compared 348 to the Drôme, in particular for foundation species (e.g., Populus spp.) and invasive exotic species (e.g., Acer 349 negundo) (Figure 8). In light of these rapid and profound divergences in community composition and structure, 350 mostly related to cumulative effect of human-induced stressors, it may be argued that the resulting riparian 351 forests correspond to the definition of a "novel ecosystem" (Hobbs et al. 2009). Indeed, the riparian forests of 352 Rhône River are rapidly dominated by non-native species, and the greatly supressed disturbance regime 353 (Vázquez-Tarrío et al. 2019) and lack of pioneer forest regeneration along bypassed channels, impedes 354 community dynamics processes evident in more natural reference systems (e.g., Stella et al. in review, Fierke 355 and Kauffman 2005, Balian and Naiman 2005). In this severely degraded system, it therefore seems more 356 pragmatic to prioritize management and restoration measures toward desired human benefits rather than 357 target a return to a potential reference state (Dufour and Piégay 2009).

358 Flow regulation and channelization induce rapid changes in abiotic conditions

359 Strongly divergent trends in abiotic conditions were evident between the two river systems. While it 360 cannot be ruled out that natural differences in physical and ecological conditions between the Drôme and the 361 Rhône watersheds could be a driver behind this observed effect, we inferred that the different management 362 histories of the Rhône and Drôme is one of the most important ecological drivers. Along the highly artificial, 363 stable river margins of the Rhône River, the Girardon dike structures were very effective in trapping suspended 364 sediment, with very high accumulation rates of overbank fine sediment, especially immediately after the initial 365 channelization and prior to the dam construction period (for further details see, Tena et al. 2020). Large 366 quantities of fine sediment trapped along channelized river margins have been shown to increase ecosystem 367 productivity by maintaining higher nutrient pools (Franklin et al. 2009). In addition, high overbank

368 sedimentation rates induce strong differences in soil properties (e.g., texture and moisture retention), which 369 can also promote a resource-rich environment. These factors may explain the rapid and vigorous growth of 370 pioneer trees, particularly Populus species that reached maximum tree sizes very quickly, i.e., at a very early 371 stage in the succession sequence. Overall, the increase in bank accretion, the decrease in overbank flows 372 (Figure 2) as well as the lowering of the water table due to river entrenchment, has induced gradual 373 disconnection between the channelized margins and the main channel. This has ultimately led to changes in 374 channel morphology and physical conditions on the Rhône, i.e., a shift from anabranching reaches to narrow, 375 very stable, single thread channels (Tena et al. 2020), as has been shown on other rivers (Wilcock and Essery 376 1991, Marston et al. 1995).

377 Disconnection of the riparian zone from the river channel is even more exacerbated by changes in the 378 hydrologic regime, as evidenced by the decrease in the mean annual flow along the Rhône River since the 379 1940s. This decrease is mainly due to water diversion into lateral canals following the construction of multiple 380 hydropower dams in the post-war period, which were commissioned at different dates along the Rhône 381 corridor (i.e., DZM = 1952, MON = 1957, PBN = 1966, PDR = 1977, see also Figure 2). The combined effects of 382 high overbank sedimentation rates and pronounced decreases in the mean annual discharge induced a loss of 383 up to 60% of the active channel surface of the Rhône River (Tena et al. 2020). Tena et al. (2020) found that 384 along the bypassed channels the terrestrialization process and associated vegetation encroachment were 385 primarily driven by the Girardon dike structures (i.e., channelization) rather than by flow regulation. The 386 cumulative effects of channel correction and regulation can also induce a rapid decline in the groundwater 387 table and thus a more limited access to soil water for existing riparian trees (Franklin et al. 2009), which on the 388 Rhône River may have been amplified locally by gravel extraction and pumping for irrigation (Thorel et al. 389 2018). Finally, channelization and flow regulation may reduce the variability of hydrologic conditions, especially 390 lower intensity and frequency of flooding on higher riparian floodplains, as well as bedload transport capacities 391 (Vázquez-Tarrío et al. 2019). Conversely, along the Drôme River, low rate of fine sediment accumulation and 392 maintenance of a natural flow regime likely promote a dynamic equilibrium across successional phases without 393 the rapid change in floodplain conditions that both correction and regulation induced on the Rhône (Figure 8). 394 Overall, our results pointed out profound changes in hydro-geomorphic processes between the two river 395 systems. In this view, we suggest that the artificially stabilizing effect of the combined Girardon structures and

flow regulation on the Rhône River enabled the development of a distinct and potentially unique riparian forestcommunity (Thorel et al. 2018).

Changes in abiotic conditions strongly alter the functional, structural and compositional attributes of riparian
 forests

400 Consistent with changes in abiotic conditions over time, our results demonstrate rapid divergence in 401 riparian forest dynamics between the two river systems, reflecting environmental selection toward a pool of 402 traits and species best adapted to local conditions (Figure 8). Although successional shifts in plant functional 403 traits followed predictable successional trajectories along both rivers, (e.g., Fierke and Kauffman 2005, Van Pelt 404 et al. 2006, Muñoz-Mas et al. 2017), increases in wood density and SLA values within the riparian forest were 405 faster along the Rhône River (Figure 3). Specifically, our results show that channelization and flow regulation 406 accelerate the transition from a disturbance-dependent pioneer community, i.e., fast-growing tree species with 407 high resource acquisition strategies, to a post-pioneer community dominated by slow-growing, competitive 408 species with high resource conservation and retention strategies (Marston et al. 1995, Oswalt and King 2005). 409 This accelerated shift in plant ecological strategy is likely related to both a more limited access to groundwater 410 and high overbank sedimentation rates promoting resource-rich environments along the Rhône, both of which 411 were mediated by changes to the hydrologic and disturbance regimes (Figure 2) (see also, Vázquez-Tarrío et al. 412 2019). Consequently, there has been a major divergence in plant habitat niches between the two river systems, 413 and large corresponding differences in stand attributes (Figure 4), plant assemblages (Figure 7) and forest 414 successional pathways. These results reinforce the importance of contingent factors in determining ecological 415 trajectories, as has been shown for old fields and other successional contexts (e.g., Clark et al. 2019) 416 Prolonged and rapid decline in the groundwater table has been shown to greatly alter growth and 417 survival of pioneer species requiring permanent access to soil water (Reily and Johnson 1982, Scott et al. 1999, 418 Lite and Stromberg 2005, Stella et al. 2013a), and favor more competitive species such as post-pioneer and 419 exotic species (Merritt and Poff 2010, González et al. 2010, Nadal-Sala et al. 2017). Within the region 420 encompassed by our study, radial growth of Populus nigra, an obligate phreatophyte that dominates early 421 successional forest stands, is more sensitive to groundwater manipulation compared to the mesophytic species 422 Fraxinus excelsior, which can use both soil moisture and deeper groundwater opportunistically to maintain

423 more constant inter-annual growth (Singer et al. 2013). Relative basal area and mean diameter of post-pioneer

424 species, including *Fraxinus* spp. and *Acer* spp., increased over time in both rivers. For *Populus* species, however,

425 while tree diameter and stem density varied consistently between the two rivers, the basal area increased 426 along the Drôme but decreased along the Rhône. In addition, most of the divergence in species composition 427 between the two river systems was due to the great abundance of the exotic invasive species Acer negundo 428 along the Rhône River. Compared to native pioneer and post-pioneer tree species, Acer negundo shows higher level of plasticity in foliage allocation (i.e., larger specific leaf area and total leaf area) when resources are not 429 430 limiting, thus allowing for better growth and survival (Porté et al. 2011). Thus, the temporal trends in whole-431 stand (Figure 4) and genus-specific (Figure 5) tree distributions suggest that channelization and flow regulation 432 have promoted a resource-rich environment within the Girardon structures (Tena et al. 2020), leading to an 433 acceleration of ecological succession due to faster growth and mortality of native pioneer species and to better 434 survival and establishment of stress tolerant species, including exotic species (Catford et al. 2014). 435 Resource-rich environments also benefit native species, in particular high growth rates of Salicaceae 436 species (Karrenberg et al. 2002) and other, more competitive species (e.g., Porté et al. 2011, Nadal-Sala et al. 437 2017). The Rhône River experienced much faster biomass accumulation, mostly due to the greater abundance 438 of very large trees that achieved their maximum size within 20 years. Contrary to expectations, however, basal 439 area and mean tree diameter of both live and dead fractions decreased with stand age along the Rhône River. 440 The anomalous pattern on the Rhône was mostly due to the scarcity of very large pioneer trees (primarily Salix 441 spp. and Populus spp.) in mature forest stands. Though they are generally short-lived trees, their rapid 442 reduction in density along the Rhône meant that overall basal area decreased on that river with forest age, 443 whereas it continued to increase on the Drôme, as expected. Since biomass usually increases along the initial

successional phases in riparian forests (e.g., Fierke and Kauffman 2005, Balian and Naiman 2005, Van Pelt et al.

445 2006), even along regulated rivers (Stella et al. in review), these results suggest that several interacting

446 stressors may be responsible for the unusual patterns. One explanation may be the rapid dewatering and

447 hydric stress that followed flow reductions in the bypassed channels with the operation of the post-war

449

448 hydropower canals. This would have preferentially affected older, established trees that could not adjust to the

new groundwater regime, increasing mortality of pioneer trees and reducing the growth rate of survivors in the

450 oldest stands. Another potential factor is selective cutting of larger trees along levees and dikes, which was

451 sometimes conducted by farmers and/or river managers of the "Compagnie Nationale du Rhône" in order to

452 protect the structural integrity of dikes and limit large wood inputs to the main channel. As such, patterns

453 observed along the Rhône River most likely result from cumulative multiple stressors related to human activity,

454 mainly channelization and flow regulation but also occasionally selective cutting, which are difficult to
455 disentangle from other drivers (Stella and Bendix 2018).

456 The increase in trait dispersion over time (Figure 3) indicates that older floodplain forests increasingly 457 support co-occurring species with contrasting ecological strategies (Figure 3). Hydro-geomorphic changes over 458 time not only drive changes in abiotic conditions at the floodplain scale but may also induce greater habitat 459 heterogeneity at finer scale, potentially due to greater variability in topographic and hydrologic conditions. 460 Accordingly, several authors have reported an increase in tree species richness with later successional stages in 461 riparian forests (Schnitzler 1995, Nakamura et al. 1997, Van Pelt et al. 2006). Here, we showed that ecological 462 succession in riparian forests not only promotes taxonomic diversity (Figure 6) but also functional diversity 463 (Figure 3). The contrasts in functional traits between rivers suggest that the riparian forests of the Rhône River 464 underwent a more rapid change to functionally different species assemblages compared to the Drôme. Indeed, 465 our results show that the shift in community-weighted SLA and wood density was accelerated along the Rhône 466 River (Figure 3). In contrast, the trait dispersion for seed mass was more pronounced along the Drôme River, 467 indicating a higher occurrence of tree species with small seed mass (i.e., mostly Salicaceae) in older stands. 468 Given the importance of floods and hydrochory in maintaining early successional forest stands (Scott et al. 469 1997, Fraaije et al. 2017), the greater representation of pioneer species within more successional stages on the 470 Drôme River further confirms the influence of dynamic fluvial processes (e.g., flooding, channel migration) in 471 riparian successional processes on intact rivers, compared to a stabilized and controlled environment as on the 472 Rhône .

473

474 Conclusion

475 In comparing successional trajectories between the heavily modified Rhône and more natural Drôme 476 rivers, we found that the legacy of cumulative impacts of river channelization and flow regulation strongly 477 altered the structure, composition and ecological trajectory of riparian forests. These findings accord with prior 478 studies along heavily modified rivers (Merritt and Cooper 2000, Nakamura et al. 2002, Oswalt and King 2005, 479 Dufour et al. 2007, González et al. 2010, Johnson et al. 2012). By using a chronosequence modelling approach, 480 we illustrated the parallel changes in river conditions, riparian forest structure and composition, and functional 481 trait expression within the plant community along a successional gradient. Thus, although the earliest phase of 482 stand development (< 20 years) on both rivers was characterized by the dominance of pioneer and

483 disturbance-dependent species, the shift to a mature forest stage dominated by post-pioneer and exotic 484 species was greatly accelerated (within 20-50 years) along the Rhône River (Figure 8). Beyond differences not 485 related to human-induced stressors, this accelerated shift was likely linked to the fact that the Rhône River 486 margins have experienced a more rapid terrestrialization than natural floodplains, due to the combined effect 487 of less frequent and lower magnitude flooding events, greater overbank fine sediment deposition, and a 488 decline in groundwater tables due to flow diversion. Thus, by disrupting natural hydro-geomorphological 489 processes at different spatial and temporal scales, channelization and flow regulation have induced severe and 490 long-term changes in environmental conditions, in turn driving riparian plant communities far from their 491 historical reference trajectories (Figure 7). Due to the strong differences in the biotic community, as well as 492 their abiotic drivers, we suggest that restoration of the Rhône River riparian forests to a more natural (i.e., pre-493 development) state is highly improbable, even if physical conditions such as flooding and bank revetment are 494 mitigated by restoration actions. The novel structural, compositional and functional attributes of the Rhône 495 riparian forest, including the dominance of non-native species in the regeneration fraction and understory, 496 suggest that the forest will be highly resistant to change and that process-based restoration actions would 497 likely have a long and persistent hysteresis (Suding et al. 2004). In this system as in other severely degraded 498 ones, riparian management may benefit from a more pragmatic approach focusing on desired human benefits 499 (Dufour and Piégay 2009) and specific ecological targets (Palmer et al. 2005). For the Rhône River's riparian 500 forest, which is increasingly valued for its ecological and cultural legacy, these approaches may include active 501 stand management by selective cutting to remove exotic species and enhance native tree diversity, promoting 502 through active and passive means and pioneer stages which are no longer rejuvenated, as well as enhancing 503 other ecosystem services such as flood and pollutant mitigation, carbon storage, and habitat quality for wildlife 504 and fish.

505

506 Authors' contributions

P.J., J.C.S., H.P. and A.E. conceived and designed the experiments. J.C.S., B.R., P.B., J-M.F. and C.J. collected the
data. P.J. analysed the data and wrote the first draft of the manuscript. All authors contributed critically to the
drafts and gave final approval for publication.

510

511 Acknowledgements

- The study has been supported by the TRY initiative on plant traits (http://www.try-db.org). The TRY initiative
 and database is hosted, developed and maintained by J. Kattge and G. Boenisch (Max Planck Institute for
 Biogeochemistry, Jena, Germany). TRY is currently supported by Future Earth/bioDISCOVERY and the German
 Center for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig. This work was cofunded by the Labex
 DRIIHM, which is managed by the ANR, French program "Investissements d'Avenir" (ANR-11-LABX-0010), the
 OHM Vallée du Rhône and IRSTEA. It was also performed within the framework of the EUR H2O'Lyon (ANR-17-
- 519

518

520 References

- Anderson, M. J., and T. J. Willis. 2003. Canonical analysis of principal coordinates: a useful method of
 constrained ordination for ecology. Ecology 84:511–525.
- 523 Balian, E. V., and R. J. Naiman. 2005. Abundance and production of riparian trees in the lowland floodplain of

524 the Queets River, Washington. Ecosystems 8:841–861.

EURE-0018) of Université de Lyon (UdL).

- Beisner, B., D. Haydon, and K. Cuddington. 2003. Alternative stable states in ecology. Frontiers in Ecology and
 the Environment 1:376–382.
- Bejarano, M. D., C. Nilsson, and F. C. Aguiar. 2018. Riparian plant guilds become simpler and most likely fewer
 following flow regulation. Journal of Applied Ecology 55:365–376.
- 529 Bendix, J., and J. C. Stella. 2013. Riparian vegetation and the fluvial environment: a biogeographic perspective.
- 530 Pages 53–74 in D. Butler and C. Hupp, editors. Treatise on Geomorphology, Vol 12,
- 531 Ecogeomorphology. Elsevier.
- Bravard, J.-P. 2010. Discontinuities in braided patterns: the River Rhône from Geneva to the Camargue delta
 before river training. Geomorphology 117:219–233.
- 534 Bravard, J.-P., and P. Gaydou. 2015. Historical development and integrated management of the Rhône River
- floodplain, from the Alps to the Camargue delta, France. Pages 289–320 in P. F. Hudson and H.
- 536 Middelkoop, editors. Geomorphic approaches to integrated floodplain management of lowland fluvial
- 537 systems in North America and Europe. Springer New York, New York, NY.
- 538 Burnham, K. P., and D. R. Anderson. 2002. Model selection and multi-model inference: a practical information-
- 539 theoretic approach, 2nd ed. Springer-Verlag, New York.

- 540 Catford, J. A., W. K. Morris, P. A. Vesk, C. J. Gippel, and B. J. Downes. 2014. Species and environmental
- 541 characteristics point to flow regulation and drought as drivers of riparian plant invasion. Diversity and
 542 Distributions 20:1084–1096.
- 543 Chang, C. C., and B. L. Turner. 2019. Ecological succession in a changing world. Journal of Ecology 107:503–509.
- Chave, J., D. Coomes, S. Jansen, S. L. Lewis, N. G. Swenson, and A. E. Zanne. 2009. Towards a worldwide wood
 economics spectrum. Ecology Letters 12:351–366.
- 546 Clark, A. T., J. M. H. Knops, and D. Tilman. 2019. Contingent factors explain average divergence in functional
 547 composition over 88 years of old field succession. Journal of Ecology 107:545–558.
- 548 Connell, J. H., and R. O. Slatyer. 1977. Mechanisms of succession in natural communities and their role in

549 community stability and organization. The American Naturalist 111:1119–1144.

- 550 Dufour, S., N. Barsoum, E. Muller, and H. Piégay. 2007. Effects of channel confinement on pioneer woody
- 551 vegetation structure, composition and diversity along the River Drôme (SE France). Earth Surface
- 552Processes and Landforms 32:1244–1256.
- 553 Dufour, S., and H. Piégay. 2009. From the myth of a lost paradise to targeted river restoration: forget natural
 554 references and focus on human benefits. River Research and Applications 25:568–581.
- Fierke, M. K., and J. B. Kauffman. 2005. Structural dynamics of riparian forests along a black cottonwood
 successional gradient. Forest Ecology and Management 215:149–162.
- Folke, C., S. Carpenter, B. Walker, M. Scheffer, T. Elmqvist, L. Gunderson, and C. S. Holling. 2004. Regime shifts,
 resilience, and biodiversity in ecosystem management. Annual Review of Ecology, Evolution, and
 Systematics 35:557–581.
- Fraaije, R. G. A., S. Moinier, I. van Gogh, R. Timmers, J. J. van Deelen, J. T. A. Verhoeven, and M. B. Soons. 2017.
 Spatial patterns of water-dispersed seed deposition along stream riparian gradients. PLOS ONE
 12:e0185247.
- Franklin, S. B., J. A. Kupfer, S. R. Pezeshki, R. Gentry, and R. D. Smith. 2009. Complex effects of channelization
 and levee construction on western Tennessee floodplain forest function. Wetlands 29:451–464.
- 565 González, E., M. González-Sanchis, Á. Cabezas, F. A. Comín, and E. Muller. 2010. Recent changes in the riparian
- 566 forest of a large regulated mediterranean River: implications for management. Environmental
 567 Management 45:669–681.

Hobbs, R. J., E. Higgs, and J. A. Harris. 2009. Novel ecosystems: implications for conservation and restoration.

569 Trends in Ecology & Evolution 24:599–605.

- Johnson, W. C., R. L. Burgess, and W. R. Keammerer. 1976. Forest overstory vegetation and environment on the
 Missouri River floodplain in North Dakota. Ecological Monographs 46:59–84.
- Johnson, W. C., M. D. Dixon, M. L. Scott, L. Rabbe, G. Larson, M. Volke, and B. Werner. 2012. Forty years of

573 vegetation change on the Missouri River floodplain. BioScience 62:123–135.

- Julve, P. 1998. Baseflor. Index botanique, écologique et chorologique de la flore de France. Version : 2017.
 http://perso.wanadoo.fr/philippe.julve/catminat.htm.
- Karrenberg, S., P. J. Edwards, and J. Kollmann. 2002. The life history of Salicaceae living in the active zone of
 floodplains. Freshwater Biology 47:733–748.
- 578 Kattge, J., S. Díaz, S. Lavorel, I. C. Prentice, P. Leadley, G. Bönisch, E. Garnier, M. Westoby, P. B. Reich, I. J.

579 Wright, J. H. C. Cornelissen, C. Violle, S. P. Harrison, P. M. Van Bodegon, M. Reichstein, B. J. Enquist, N.

580 A. Soudzilovskaia, D. D. Ackerly, M. Anand, O. Atkin, M. Bahn, T. R. Baker, D. Baldocchi, R. Bekker, C. C.

- 581 Blanco, B. Blonder, W. J. Bond, R. Bradstock, D. E. Bunker, F. Casanoves, J. Cavender-Bares, J. Q.
- 582 Chambers, F. S. Chapin III, J. Chave, D. Coomes, W. K. Cornwell, J. M. Craine, B. H. Dobrin, L. Duarte, W.
- 583 Durka, J. Elser, G. Esser, M. Estiarte, W. F. Fagan, J. Fang, F. Fernández-Méndez, A. Fidelis, B. Finegan,
- 584 O. Flores, H. Ford, D. Frank, G. T. Freschet, N. M. Fyllas, R. V. Gallagher, W. A. Green, A. G. Gutierrez, T.
- 585 Hickler, S. I. Higgins, J. G. Hodgson, A. Jalili, S. Jansen, C. A. Joly, A. J. Kerkhoff, D. Kirkup, K. Kitajima, M.
- 586 Kleyer, S. Klotz, J. M. H. Knops, K. Kramer, I. Kühn, H. Kurokawa, D. Laughlin, T. D. Lee, M. Leishman, F.
- 587 Lens, T. Lenz, S. L. Lewis, J. Lloyd, J. Llusià, F. Louault, S. Ma, M. D. Mahecha, P. Manning, T. Massad, B.
- 588 E. Medlyn, J. Messier, A. T. Moles, S. C. Müller, K. Nadrowski, S. Naeem, U. Niinemets, S. Nöllert, A.
- 589 Nüske, R. Ogaya, J. Oleksyn, V. G. Onipchenko, Y. Onoda, J. Ordoñez, G. Overbeck, W. A. Ozinga, S.
- 590 Patiño, S. Paula, J. G. Pausas, J. Peñuelas, O. L. Phillips, V. Pillar, H. Poorter, L. Poorter, P. Poschlod, A.
- 591 Prinzing, R. Proulx, A. Rammig, S. Reinsch, B. Reu, L. Sack, B. Salgado-Negret, J. Sardans, S. Shiodera, B.
- 592 Shipley, A. Siefert, E. Sosinski, J.-F. Soussana, E. Swaine, N. Swenson, K. Thompson, P. Thornton, M.
- 593 Waldram, E. Weiher, M. White, S. White, S. J. Wright, B. Yguel, S. Zaehle, A. E. Zanne, and C. Wirth.
- 594 2011. TRY a global database of plant traits. Global Change Biology 17:2905–2935.

- Kui, L., J. C. Stella, P. B. Shafroth, P. K. House, and A. C. Wilcox. 2017. The long-term legacy of geomorphic and
 riparian vegetation feedbacks on the dammed Bill Williams River, Arizona, USA. Ecohydrology
 10:e1839.
- Laliberté, E., and P. Legendre. 2010. A distance-based framework for measuring functional diversity from
 multiple traits. Ecology 91:299–305.
- Lamouroux, N., J. A. Gore, F. Lepori, and B. Statzner. 2015. The ecological restoration of large rivers needs
 science-based, predictive tools meeting public expectations: an overview of the Rhône project.
 Freshwater Biology 60:1069–1084.
- 603 Lavorel, S., K. Grigulis, S. McIntyre, N. S. G. Williams, D. Garden, J. Dorrough, S. Berman, F. Quétier, A. Thébault,
- 604and A. Bonis. 2008. Assessing functional diversity in the field methodology matters! Functional605Ecology 22:134–147.
- Liébault, F., and H. Piégay. 2002. Causes of 20th century channel narrowing in mountain and piedmont rivers of
 southeastern France. Earth Surface Processes and Landforms 27:425–444.
- Lite, S. J., and J. C. Stromberg. 2005. Surface water and ground-water thresholds for maintaining Populus–Salix
 forests, San Pedro River, Arizona. Biological Conservation 125:153–167.
- 610 Marston, R. A., J. Girel, G. Pautou, H. Piégay, J.-P. Bravard, and C. Arneson. 1995. Channel metamorphosis,
- 611 floodplain disturbance, and vegetation development: Ain River, France. Geomorphology 13:121–131.
- 612 Meiners, S. J., M. W. Cadotte, J. D. Fridley, S. T. A. Pickett, and L. R. Walker. 2015. Is successional research
- 613 nearing its climax? New approaches for understanding dynamic communities. Functional Ecology
 614 29:154–164.
- Merritt, D. M., and D. J. Cooper. 2000. Riparian vegetation and channel change in response to river regulation:
 a comparative study of regulated and unregulated streams in the Green River basin, USA. Regulated
 Rivers: Research and management 16:543–564.
- 618 Merritt, D. M., and N. L. R. Poff. 2010. Shifting dominance of riparian *Populus* and *Tamarix* along gradients of 619 flow alteration in western North American rivers. Ecological Applications 20:135–152.
- 620 Mouillot, D., N. A. J. Graham, S. Villéger, N. W. H. Mason, and D. R. Bellwood. 2013. A functional approach
- 621 reveals community responses to disturbances. Trends in Ecology & Evolution 28:167–177.
- 622 Muñoz-Mas, R., V. Garófano-Gómez, I. Andrés-Doménech, D. Corenblit, G. Egger, F. Francés, M. T. Ferreira, A.
- 623 García-Arias, E. Politti, R. Rivaes, P. M. Rodríguez-González, J. Steiger, F. J. Vallés-Morán, and F.

- 624 Martínez-Capel. 2017. Exploring the key drivers of riparian woodland successional pathways across
 625 three European river reaches. Ecohydrology 10:e1888.
- Nadal-Sala, D., S. Sabaté, E. Sánchez-Costa, S. Poblador, F. Sabater, and C. Gracia. 2017. Growth and water use
 performance of four co-occurring riparian tree species in a Mediterranean riparian forest. Forest
 Ecology and Management 396:132–142.
- Naiman, R. J., and H. Decamps. 1997. The ecology of interfaces: riparian zones. Annual review of Ecology and
 Systematics 28:621–658.
- Nakamura, F., M. Jitsu, S. Kameyama, and S. Mizugaki. 2002. Changes in riparian forests in the Kushiro Mire,
 Japan, associated with stream channelization. River Research and Applications 18:65–79.
- 633 Nakamura, F., T. Yajima, and S. Kikuchi. 1997. Structure and composition of riparian forests with special
- reference to geomorphic site conditions along the Tokachi River, northern Japan. Plant Ecology
 133:209–219.
- Odion, D. C., M. A. Moritz, and D. A. DellaSala. 2010. Alternative community states maintained by fire in the
 Klamath Mountains, USA. Journal of Ecology 98:96–105.
- Olivier, J.-M., M.-J. Dole-Olivier, C. Amoros, G. Carrel, F. Malard, N. Lamouroux, and J.-P. Bravard. 2009. The
 Rhône River basin. Pages 247–295 Rivers of Europe (Tockner K., Robinson C.T. & Uehlinger U. eds).
 Elsevier, Amsterdam.
- Osterkamp, W. R., and C. R. Hupp. 2010. Fluvial processes and vegetation Glimpses of the past, the present,
 and perhaps the future. Geomorphology 116:274–285.
- Oswalt, S. N., and S. L. King. 2005. Channelization and floodplain forests: Impacts of accelerated sedimentation
 and valley plug formation on floodplain forests of the Middle Fork Forked Deer River, Tennessee, USA.
 Forest Ecology and Management 215:69–83.
- 646 Palmer, M. A., E. S. Bernhardt, J. D. Allan, P. S. Lake, G. Alexander, S. Brooks, J. Carr, S. Clayton, C. N. Dahm, J.
- 647 Follstad Shah, D. L. Galat, S. G. Loss, P. Goodwin, D. D. Hart, B. Hassett, R. Jenkinson, G. M. Kondolf, R.
- 648 Lave, J. L. Meyer, T. K. O'Donnell, L. Pagano, and E. Sudduth. 2005. Standards for ecologically
- 649 successful river restoration: Ecological success in river restoration. Journal of Applied Ecology 42:208–
- 650

217.

- 651 Perry, L. G., D. C. Andersen, L. V. Reynolds, S. M. Nelson, and P. B. Shafroth. 2012. Vulnerability of riparian
- ecosystems to elevated CO2 and climate change in arid and semiarid western North America. Global
 Change Biology 18:821–842.
- Pickett, S. T. A., S. L. Collins, and J. J. Armesto. 1987. Models, mechanisms and pathways of succession.
 Botanical Review 53:335–371.
- Porté, A. J., L. J. Lamarque, C. J. Lortie, R. Michalet, and S. Delzon. 2011. Invasive Acer negundo outperforms
 native species in non-limiting resource environments due to its higher phenotypic plasticity. BMC
 Ecology 11:28.
- Prach, K., and L. R. Walker. 2011. Four opportunities for studies of ecological succession. Trends in Ecology &
 Evolution 26:119–123.
- QGIS Development Team. 2015. QGIS Geographic Information System. Open Source Geospatial Foundation
 Project.
- R Core Team. 2019. R: A language and environment for statistical computing. R Foundation for Statistical
 Computing, Vienna, Austria.
- Räpple, B. 2018. Sedimentation patterns and riparian vegetation characteristics in novel ecosystems on the
 Rhône River, France: a comparative approach to identify drivers and evaluate ecological potentials.

667 Thèse de Doctorat, Université de Lyon, Lyon.

- 668 Räpple, B., H. Piégay, J. C. Stella, and D. Mercier. 2017. What drives riparian vegetation encroachment in
- braided river channels at patch to reach scales? Insights from annual airborne surveys (Drôme River,
- 670 SE France, 2005-2011). Ecohydrology 10:e1886.
- Reily, P. W., and W. C. Johnson. 1982. The effects of altered hydrologic regime on tree growth along the
 Missouri River in North Dakota. Canadian Journal of Botany 60:2410–2423.
- 673 Schnitzler, A. 1995. Successional status of trees in gallery forest along the river Rhine. Journal of Vegetation
 674 Science 6:479–486.
- Scott, M. L., G. T. Auble, and J. M. Friedman. 1997. Flood dependency of Cottonwood establishment along the
 Missouri River, Montana, USA. Ecological Applications 7:14.
- Scott, M. L., P. B. Shafroth, and G. T. Auble. 1999. Responses of riparian Cottonwoods to alluvial water table
 declines. Environmental Management 23:347–358.

- 679 Sfair, J. C., V. Arroyo-Rodríguez, B. A. Santos, and M. Tabarelli. 2016. Taxonomic and functional divergence of 680 tree assemblages in a fragmented tropical forest. Ecological Applications 26:1816–1826.
- 681 Singer, M. B., J. C. Stella, S. Dufour, H. Piégay, R. J. S. Wilson, and L. Johnstone. 2013. Contrasting water-uptake 682 and growth responses to drought in co-occurring riparian tree species. Ecohydrology 6:402–412.

683

Stella, J. C., and J. Bendix. 2018. Multiple stressors in riparian ecosystems. Pages 81–110 Multiple stressors in

- 684 river ecosystems: status, impacts and prospects for the future (Sabater, S., Elosegi, A., Ludwig, R.,
- 685 eds.). Elsevier, San Diego.
- 686 Stella, J. C., L. Kui, G. H. Golet, and F. Poulsen. in review. Modeling riparian forest structure to estimate large 687 wood inputs and other ecosystem services on geomorphically active floodplains. Earth Surface Processes and Landforms. 688

689 Stella, J. C., J. Riddle, H. Piégay, M. Gagnage, and M.-L. Trémélo. 2013a. Climate and local geomorphic

- 690 interactions drive patterns of riparian forest decline along a Mediterranean Basin river.
- 691 Geomorphology 202:101–114.
- 692 Stella, J. C., P. M. Rodríguez-González, S. Dufour, and J. Bendix. 2013b. Riparian vegetation research in 693 Mediterranean-climate regions: common patterns, ecological processes, and considerations for

694 management. Hydrobiologia 719:291–315.

695 Suding, K. N. 2011. Toward an era of restoration in ecology: successes, failures, and opportunities ahead.

696 Annual Review of Ecology, Evolution, and Systematics 42:465–487.

697 Suding, K. N., K. L. Gross, and G. R. Houseman. 2004. Alternative states and positive feedbacks in restoration 698 ecology. Trends in Ecology & Evolution 19:46–53.

699 Tena, A., H. Piégay, G. Seignemartin, A. Barra, J. F. Berger, B. Mourier, and T. Winiarski. 2020. Cumulative

700 effects of channel correction and regulation on floodplain terrestrialisation patterns and connectivity. 701 Geomorphology 354:107034.

- 702 Thorel, M., H. Piégay, C. Barthélémy, B. Räpple, C. R. Gruel, P. Marmonier, T. Winiarski, J. P. Bedell, F. Arnaud,
- 703 G. Roux, J. Stella, G. Seignemartin, A. Tena-Pagan, V. Wawrzyniak, D. Roux-Michollet, B. Oursel, S.
- 704 Fayolle, C. Bertrand, and E. Franquet. 2018. Socio-environmental implications of process-based
- 705 restoration strategies in large rivers: should we remove novel ecosystems along the Rhône (France)?
- 706 Regional Environmental Change 18:2019–2031.

- Tonkin, J. D., David. M. Merritt, J. D. Olden, L. V. Reynolds, and D. A. Lytle. 2018. Flow regime alteration
 degrades ecological networks in riparian ecosystems. Nature Ecology & Evolution 2:86–93.
- Van Pelt, R., T. C. O'Keefe, J. J. Latterell, and R. J. Naiman. 2006. Riparian forest stand development along the
 Queets River in Olympic National Park, Washington. Ecological Monographs 76:277–298.

711 Vázquez-Tarrío, D., M. Tal, B. Camenen, and H. Piégay. 2019. Effects of continuous embankments and

- 512 successive run-of-the-river dams on bedload transport capacities along the Rhône River, France.
- 713 Science of The Total Environment 658:1375–1389.
- Walker, L. R., D. A. Wardle, R. D. Bardgett, and B. D. Clarkson. 2010. The use of chronosequences in studies of
 ecological succession and soil development: Chronosequences, succession and soil development.

716Journal of Ecology 98:725–736.

- Wang, Y., U. Naumann, S. T. Wright, and D. I. Warton. 2012. mvabund an R package for model-based analysis
 of multivariate abundance data. Methods in Ecology and Evolution 3:471–474.
- Warton, D. I., L. Thibaut, and Y. A. Wang. 2017. The PIT-trap—A "model-free" bootstrap procedure for
 inference about regression models with discrete, multivariate responses. PloS one 12:e0181790.

721 Westoby, M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. Plant and Soil 199:213–227.

722 Wilcock, D. N., and C. I. Essery. 1991. Environmental impacts of channelization on the river Main, County

723 Antrim, Northern Ireland. Journal of Environmental Management 32:127–143.

- 724 Wright, I. J., P. B. Reich, M. Westoby, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-Bares, T. Chapin, J. H.
- 725 Cornelissen, M. Diemer, J. Flexas, E. Garnier, P. K. Groom, J. Gulias, K. Hikosaka, B. B. Lamont, T. Lee,
- 726 W. Lee, C. Lusk, J. J. Midgley, M.-L. Navas, ü. Niinemets, J. Oleskyn, N. Osada, H. Poorter, P. Poot, L.
- 727 Prior, V. Pyankov, C. Roumet, S. C. Thomas, M. G. Tjoekler, E. J. Veneklaas, and R. Villar. 2004. The
- 728 worldwide leaf economics spectrum. Nature 428:821–827.
- Zanne, A. E., G. Lopez-Gonzales, D. A. Coomes, J. Ilic, S. Jansen, S. L. Lewis, R. B. Miller, N. G. Swenson, M. C.
 Wiemann, and J. Chave. 2009. Data from: Towards a worldwide wood economics spectrum. Dryad
 Digital Repository https://doi.org/10.5061/dryad.234.

| 733 | Table 1. Variation in inde | pendent and depender | nt variables between | the rivers Drôme | (n = 69 p | lots) and Rhône |
|-----|----------------------------|----------------------|----------------------|------------------|-----------|-----------------|

734 (n = 65 plots), France (CWM = community-weighted mean, FDis = functional dispersion, SLA = specific leaf

735 area).

| Maagura | Variable | Drôme | River | Rhône River | | |
|------------------|------------------------------------|---------------|--------------|----------------|----------------|--|
| Measure | variable | mean (±SD) | range | mean (±SD) | range | |
| Plot descriptio | n | | | | | |
| | Forest stand age (years) | 31.3 (±20.3) | 5.0 - 62.0 | 44.6 (±18.6) | 16.0 - 76.0 | |
| | Sediment depth (cm) | 42 (±36) | 0 - 140 | 249 (±121) | 24 - 490 | |
| | Mean annual flow (m ³) | 18.4 (±4.6) | 7.3 - 31.3 | 527.2 (±402.4) | 77.7 - 1550.4 | |
| Functional trai | ts | | | | | |
| CWM | SLA | 12.63 (±1.66) | 10.0 - 17.0 | 14.87 (±3.00) | 11.1 - 21.8 | |
| | Wood density | 0.41 (±0.05) | 0.3 - 0.6 | 0.47 (±0.10) | 0.3 - 0.7 | |
| | Seed mass | 1.40 (±0.90) | 0.4 - 4.2 | 1.63 (±1.04) | 0.2 - 3.8 | |
| FDis | SLA | 0.24 (±0.20) | 0.0 - 0.7 | 0.40 (±0.26) | 0.0 - 1.0 | |
| | Wood density | 0.51 (±0.39) | 0.0 - 1.2 | 0.69 (±0.44) | 0.0 - 1.4 | |
| | Seed mass | 0.41 (±0.36) | 0.0 - 1.4 | 0.40 (±0.25) | 0.0 - 1.1 | |
| Stand attribute | es | | | | | |
| Basal area | Live trees | 15.47 (±9.76) | 0.1 - 38.7 | 27.65 (±10.85) | 6.3 - 57.8 | |
| (m² ha) | Dead trees | 3.72 (±3.69) | 0.0 - 18.0 | 5.13 (±4.43) | 0.0 - 22.3 | |
| | Exotic trees | 0.58 (±1.10) | 0.0 - 5.2 | 8.22 (±9.92) | 0.0 - 42.8 | |
| Mean | Live trees | 17.49 (±4.91) | 8.7 - 30.7 | 24.91 (±8.02) | 10.7 - 47.7 | |
| diameter | Dead trees | 15.60 (±5.63) | 9.0 - 44.2 | 28.96 (±16.97) | 9.9 - 76.6 | |
| (cm) | Exotic trees | 13.7 (±4.23) | 8.0 - 24.4 | 17.49 (±6.11) | 7.5 - 35.4 | |
| Stem density | Live trees | 504 (±262.65) | 16 - 1066 | 463 (±241.09) | 71 - 1169 | |
| (n na) | Dead trees | 157 (±107.85) | 0 - 500 | 98 (±99.12) | 0 - 453 | |
| | Exotic trees | 38 (±80.39) | 0 - 466 | 276 (±225.97) | 0 - 1122 | |
| Tree genera | | | | | | |
| Basal area | Populus spp. | 11.44 (±8.87) | 0.00 - 38.59 | 14.83 (±11.90) | 0.00 - 43.97 | |
| (m² ha) | Fraxinus spp. | 1.03 (±1.76) | 0.00 - 7.28 | 1.59 (±2.71) | 0.00 - 14.08 | |
| | Acer spp. | 1.43 (±2.54) | 0.00 - 11.74 | 4.04 (±4.52) | 0.00 - 19.10 | |
| Mean | Populus spp. | 21.22 (±7.27) | 8.01 - 43.60 | 54.15 (±16.20) | 26.78 - 113.00 | |
| diameter | Fraxinus spp. | 13.64 (±3.99) | 8.00 - 23.33 | 23.54 (±12.82) | 8.50 - 68.50 | |
| (cm) | Acer spp. | 14.58 (±3.65) | 8.00 - 22.00 | 20.62 (±10.34) | 7.50 - 60.50 | |
| Stem density | Populus spp. | 268 (±180.38) | 0 - 833 | 97 (±129.87) | 0 - 604 | |
| (n ha) | Fraxinus spp. | 53 (±93.35) | 0 - 600 | 58 (±113.75) | 0 - 652 | |
| | Acer spp. | 72 (±128.49) | 0 - 633 | 157 (±162.13) | 0 - 636 | |
| Diversity indice | es | | | | | |
| | Diameter classes | 1.59 (±0.43) | 0.0 - 2.1 | 1.94 (±0.29) | 0.8 - 2.4 | |
| | Tree species | 0.79 (±0.56) | 0.0 - 1.7 | 0.96 (±0.37) | 0.0 - 1.7 | |

Table 2. Top-ranking models predicting variations in abiotic drivers and biotic responses over time along the
rivers Drôme and Rhône, France, as assessed with Akaike's information criterion corrected for small sample
size (AIC_c). Other model information provided include the number of estimated parameters including the
intercept (k), AIC_c weight (*W*) and adjusted coefficient of determination (R²). Asterisks in the model formulas
indicate an interaction term (CWM = community-weighted mean, FDis = functional dispersion, SLA = specific
leaf area).

| Measure | Dependent variable | Top-ranked model | k | W | R ² |
|-------------------|--------------------|-------------------------|---|-------|----------------|
| Abiotic drivers | | | | | |
| | Sediment depth | Age * River | 5 | 0.823 | 0.564 |
| | Mean annual flow | Year * River | 5 | 0.941 | 0.476 |
| | SD annual flow | Year | 5 | 0.425 | 0.108 |
| Functional traits | | | | | |
| CWM | SLA | Age * River | 5 | 0.423 | 0.387 |
| | Wood density | Age * River | 5 | 0.399 | 0.430 |
| | Seed mass | Age | 3 | 0.282 | 0.376 |
| FDis | SLA | Age + River | 4 | 0.373 | 0.232 |
| | Wood density | Age * River | 5 | 0.305 | 0.228 |
| | Seed mass | Age * River | 5 | 0.500 | 0.300 |
| Stand attributes | | | | | |
| Basal area | Live trees | Age * River + Sed_depth | 6 | 0.775 | 0.290 |
| | Dead trees | Age * River + Sed_depth | 6 | 0.978 | 0.174 |
| | Exotic trees | Age * River + Sed_depth | 6 | 0.962 | 0.578 |
| Mean diameter | Live trees | Age * River | 5 | 0.632 | 0.332 |
| | Dead trees | Age*River + Sed_depth | 6 | 0.857 | 0.155 |
| | Exotic trees | Age*River | 5 | 0.378 | 0.183 |
| Stem density | Live trees | Age + River + Sed_depth | 5 | 0.612 | 0.139 |
| | Dead trees | Age * River + Sed_depth | 6 | 0.297 | 0.117 |
| | Exotic trees | Age + River | 4 | 0.417 | 0.490 |
| Tree genera | | | | | |
| Basal area | Populus spp. | Age * River | 5 | 0.720 | 0.079 |
| | Fraxinus spp. | Age + Sed_depth | 4 | 0.351 | 0.228 |
| | Acer spp. | Age * River + Sed_depth | 6 | 0.584 | 0.344 |
| Mean diameter | Populus spp. | Age*River | 5 | 0.482 | 0.706 |
| | Fraxinus spp. | Age + River | 4 | 0.407 | 0.295 |
| | Acer spp. | Age + River + Sed_depth | 5 | 0.574 | 0.182 |
| Stem density | Populus spp. | Age * River | 5 | 0.617 | 0.328 |
| | Fraxinus spp. | Age + Sed_depth | 4 | 0.457 | 0.180 |
| | Acer spp. | Age * River + Sed_depth | 6 | 0.944 | 0.397 |
| Diversity indices | | | | | _ |
| | Diameter classes | Age * River | 5 | 0.435 | 0.241 |
| | Tree species | Age * River | 5 | 0.731 | 0.247 |

744 Table 3. Standardized coefficient estimates (±SE) and confidence intervals (95% CI) for each variable used to predict mean and dispersion of functional trait values, basal

745 area, mean diameter and stem density of stand attributes and of tree genera as well as diversity indices to forest stand age along the rivers Drôme and Rhône, France. Bold

font indicates coefficient estimates for which the 95% confidence interval excludes zero (CWM = community-weighted mean, FDis = functional dispersion, SLA = specific leaf

747 area).

| | Variable | Age | | River | | Age*River | | Sed_depth | |
|---------------|---------------|-----------------|-----------------|-----------------|------------------|-----------------|------------------|-----------------|-----------------|
| weasure | | Estimate (±SE) | (95% CI) | Estimate (±SE) | (95% CI) | Estimate (±SE) | (95% CI) | Estimate (±SE) | (95% CI) |
| Functional tr | raits | | | | | | | | |
| CWM: | SLA | 0.052 (±0.017) | (0.019; 0.085) | 0.095 (±0.026) | (0.044; 0.146) | 0.069 (±0.026) | (0.018; 0.120) | NA | NA |
| | Wood density | 0.075 (±0.017) | (0.042; 0.108) | 0.065 (±0.025) | (0.016; 0.114) | 0.060 (±0.025) | (0.011; 0.109) | NA | NA |
| | Seed mass | 0.603 (±0.067) | (0.472; 0.734) | NA | NA | NA | NA | NA | NA |
| FDis: | SLA | 0.073 (±0.014) | (0.046; 0.100) | 0.062 (±0.029) | (0.005; 0.119) | NA | NA | NA | NA |
| | Wood density | 0.162 (±0.029) | (0.105; 0.219) | 0.039 (±0.044) | (-0.047; 0.125) | -0.092 (±0.044) | (-0.178; -0.006) | NA | NA |
| | Seed mass | 0.167 (±0.022) | (0.124; 0.210) | -0.065 (±0.034) | (-0.132; 0.002) | -0.113 (±0.034) | (-0.180; -0.046) | NA | NA |
| Stand attribu | utes | | | | | | | | |
| Basal area: | Live trees | 0.365 (±0.100) | (0.169; 0.561) | 0.413 (±0.218) | (-0.014; 0.840) | -0.567 (±0.153) | (-0.867; -0.267) | 0.256 (±0.119) | (0.023; 0.489) |
| | Dead trees | 0.270 (±0.087) | (0.099; 0.441) | -0.343 (±0.188) | (-0.711; 0.025) | -0.452 (±0.132) | (-0.711; -0.193) | 0.379 (±0.103) | (0.177; 0.581) |
| | Exotic trees | -0.003 (±0.086) | (-0.172; 0.166) | 0.798 (±0.186) | (0.433; 1.163) | 0.461 (±0.130) | (0.206; 0.716) | 0.304 (±0.101) | (0.106; 0.502) |
| Mean tree | Live trees | 0.064 (±0.034) | (-0.003; 0.131) | 0.373 (±0.051) | (0.273; 0.473) | -0.209 (±0.051) | (-0.309; -0.109) | NA | NA |
| diameter: | Dead trees | 0.077 (±0.051) | (-0.023; 0.177) | 0.374 (±0.107) | (0.164; 0.584) | -0.356 (±0.077) | (-0.507; -0.205) | 0.141 (±0.058) | (0.027; 0.255) |
| | Exotic trees | 0.033 (±0.053) | (-0.071; 0.137) | 0.159 (±0.069) | (0.024; 0.294) | 0.123 (±0.070) | (-0.014; 0.260) | NA | NA |
| Stem | Live trees | 0.147 (±0.061) | (0.027; 0.267) | -0.560 (±0.165) | (-0.883; -0.237) | NA | NA | 0.280 (±0.089) | (0.106; 0.454) |
| density: | Dead trees | 0.513 (±0.181) | (0.158; 0.868) | -1.449 (±0.394) | (-2.221; -0.677) | -0.409 (±0.276) | (-0.950; 0.132) | 0.391 (±0.215) | (-0.030; 0.812) |
| | Exotic trees | -0.024 (±0.154) | (-0.326; 0.278) | 3.327 (±0.307) | (2.725; 3.929) | NA | NA | NA | NA |
| Tree genera | | | | | | | | | |
| Basal area: | Populus spp. | 0.143 (±0.115) | (-0.082; 0.368) | 0.248 (±0.172) | (-0.089; 0.585) | -0.594 (±0.173) | (-0.933; -0.255) | NA | NA |
| | Fraxinus spp. | 0.372 (±0.060) | (0.254; 0.490) | NA | NA | NA | NA | -0.095 (±0.060) | (-0.213; 0.023) |
| | Acer spp. | 0.476 (±0.089) | (0.302; 0.650) | 0.181 (±0.192) | (-0.195; 0.557) | -0.347 (±0.135) | (-0.612; -0.082) | 0.212 (±0.105) | (0.006; 0.418) |
| Mean tree | Populus spp. | 0.142 (±0.039) | (0.066; 0.218) | 0.909 (±0.058) | (0.795; 1.023) | -0.124 (±0.059) | (-0.240; -0.008) | NA | NA |

| diameter: | Fraxinus spp. | 0.131 (±0.048) | (0.037; 0.225) | 0.392 (±0.092) | (0.212; 0.572) | NA | NA | NA | NA |
|--------------|------------------|-----------------|-----------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| | Acer spp. | 0.177 (±0.051) | (0.077; 0.277) | 0.448 (±0.127) | (0.199; 0.697) | NA | NA | -0.135 (±0.063) | (-0.258; -0.012) |
| Stem | Populus spp. | -0.004 (±0.151) | (-0.300; 0.292) | -1.380 (±0.226) | (-1.823; -0.937) | -0.606 (±0.228) | (-1.053; -0.159) | NA | NA |
| density: | Fraxinus spp. | 1.062 (±0.191) | (0.688; 1.436) | NA | NA | NA | NA | -0.605 (±0.191) | (-0.979; -0.231) |
| | Acer spp. | 1.506 (±0.228) | (1.059; 1.953) | 0.159 (±0.496) | (-0.813; 1.131) | -1.483 (±0.347) | (-2.163; -0.803) | 0.755 (±0.270) | (0.226; 1.284) |
| Diversity in | dexes | | | | | | | | |
| | Diameter classes | 0.159 (±0.043) | (0.075; 0.243) | 0.284 (±0.065) | (0.157; 0.411) | -0.133 (±0.065) | (-0.260; -0.006) | NA | NA |
| | Tree species | 0.330 (±0.052) | (0.228; 0.432) | 0.048 (±0.077) | (-0.103; 0.199) | -0.277 (±0.078) | (-0.430; -0.124) | NA | NA |

Figure 1. Map of the study area within the Rhône River basin (A); location of the study reaches along the Drôme
and Rhône rivers (B); and examples of aerial photograph series showing the riparian forest development along
the two rivers (C, aerial photographs: IGN).



Figure 2. Variation in fine sediment depth with forest stand age along the rivers Drôme and Rhône and trends in the mean and standard deviation (SD) of annual flow since 1940. Annual flow variables were standardized to account for differences in basin size between rivers (n.s. interaction = non-significant Year x River interaction; see Appendix S8). To facilitate interpretation, the dates of commissioning of the diversion canals are shown with arrows in the figure (i.e., DZM = 1952, MON = 1957, PBN = 1966, PDR = 1977).



Figure 3. Variation in the mean (CWM) and dispersion (FDis) of specific leaf area (SLA, m² kg⁻¹), wood density (g cm³) and seed mass (mg) with forest stand age along the rivers Drôme and Rhône (n.s. interaction = nonsignificant Age x River interaction; see Table 3). Trait values and their variation increased more rapidly along the Rhône River for SLA and wood density, compared to the Drôme.



- 760 Figure 4. Variation in the basal area, mean diameter and stem density of live, dead and exotic tree species with
- 761 forest stand age along the rivers Drôme and Rhône (n.s. interaction = non-significant Age x River interaction;
- see Table 3). Basal area increased much more rapidly within the Rhône riparian forest for live trees and for the



763 exotic species fraction, largely due to more large trees in early to mid-successional phases.

Figure 5. Variation in the basal area, mean diameter and stem density of *Populus* spp., *Fraxinus*, spp. and *Acer*spp. with forest stand age along the rivers Drôme and Rhône (n.s river = non-significant River effect; see Table
3). Whereas development of *Fraxinus* spp., a native post-pioneer taxon, was comparable among rivers, the
overall divergence in successional trajectories (Figure 4) was driven by greater *Populus* spp. tree size and basal
area, as well as the higher density and basal area of non-native *Acer* species, primarily *Acer negundo*, within the



769 Rhône River riparian forest community.

770 Figure 6. Variation in the diversity of tree diameter classes and tree species with forest stand age along the



771 rivers Drôme and Rhône.

Figure 7. Constrained canonical analysis of principal coordinates of A) tree diameter classes and B) tree species
in riparian forests along the rivers Drôme and Rhône. To facilitate graphical interpretation, data are grouped
into three classes of forest stand age (<25, 25-50 and >50 years). In addition, the centroids in principle
coordinate space of the three most influential diameter classes ("70" is DBH > 70 cm; "52.5" is 51 cm < DBH <
56 cm; "12.5" is 11 cm < DBH < 16 cm) and tree species (Acer.neg is *Acer negundo*, Sal.alb is *Salix alba*, Sam.nig
is *Sambucus nigra*) are provided (see Appendix S9 and S10).



- 778 Figure 8. Conceptual model of ecological trajectory of riparian forests between the channelized and regulated
- 779 Rhône River and the relatively unmodified Drôme River, across a range of successional phases in eastern



780 France.