



## The Buchweitz set of a numerical semigroup

Shalom Eliahou, Juan Ignacio García-García, Daniel Marín-Aragón, Alberto Vigneron-Tenorio

### ► To cite this version:

Shalom Eliahou, Juan Ignacio García-García, Daniel Marín-Aragón, Alberto Vigneron-Tenorio. The Buchweitz set of a numerical semigroup. *Boletim da Sociedade Brasileira de Matemática / Bulletin of the Brazilian Mathematical Society*, 2023, 54 (4), 10.1007/s00574-022-00322-8 . hal-03032801

HAL Id: hal-03032801

<https://hal.science/hal-03032801v1>

Submitted on 1 Dec 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The Buchweitz set of a numerical semigroup

S. Eliahou, J.I. García-García,  
D. Marín-Aragón, A. Vigneron-Tenorio

## Abstract

Let  $A \subset \mathbb{Z}$  be a finite subset. We denote by  $\mathcal{B}(A)$  the set of all integers  $n \geq 2$  such that  $|nA| > (2n-1)(|A|-1)$ , where  $nA = A + \cdots + A$  denotes the  $n$ -fold sumset of  $A$ . The motivation to consider  $\mathcal{B}(A)$  stems from Buchweitz's discovery in 1980 that if a numerical semigroup  $S \subseteq \mathbb{N}$  is a Weierstrass semigroup, then  $\mathcal{B}(\mathbb{N} \setminus S) = \emptyset$ . By constructing instances where this condition fails, Buchweitz disproved a longstanding conjecture by Hurwitz (1893). In this paper, we prove that for any numerical semigroup  $S \subset \mathbb{N}$  of genus  $g \geq 2$ , the set  $\mathcal{B}(\mathbb{N} \setminus S)$  is finite, of unbounded cardinality as  $S$  varies.

*Keywords and phrases.* Weierstrass numerical semigroup; gapset; additive combinatorics; sumset growth; Freiman's  $3k-3$  theorem.

*To the memory of our friend and colleague  
Fernando Torres (1961-2020)*

## 1 Introduction

Denote  $\mathbb{N} = \{0, 1, 2, 3, \dots\}$  and  $\mathbb{N}_+ = \mathbb{N} \setminus \{0\} = \{1, 2, 3, \dots\}$ . For  $a, b \in \mathbb{Z}$ , let  $[a, b[ = \{z \in \mathbb{Z} \mid a \leq z < b\}$  and  $[a, \infty[ = \{z \in \mathbb{Z} \mid a \leq z\}$  denote the integer intervals they span. A *numerical semigroup* is a subset  $S \subseteq \mathbb{N}$  containing 0, stable under addition and with finite complement in  $\mathbb{N}$ . Equivalently, it is a subset of  $\mathbb{N}$  of the form  $S = \langle a_1, \dots, a_n \rangle = \mathbb{N}a_1 + \cdots + \mathbb{N}a_n$  where  $\gcd(a_1, \dots, a_n) = 1$ . The set  $\{a_1, \dots, a_n\}$  is then called a *system of generators* of  $S$ , and the smallest such  $n$  is called the *embedding dimension* of  $S$ .

For a numerical semigroup  $S$ , its corresponding *gapset* is the complement  $G = \mathbb{N} \setminus S$ , its *genus* is  $g = |G|$ , its *multiplicity* is  $m = \min S^*$  where  $S^* = S \setminus \{0\}$ , its

*Frobenius number* is  $f = \max(\mathbb{Z} \setminus S)$  and its *conductor* is  $c = f + 1$ . Thus  $[c, \infty[ \subseteq S$  and  $c$  is minimal for this property. Finally, the *depth* of  $S$  is  $q = \lceil c/m \rceil$ .

Given a finite subset  $A \subset \mathbb{N}$ , we denote by  $nA = A + \cdots + A$  the *n-fold sumset* of  $A$ . See Section 2 for more details.

**Definition 1.1.** Let  $A \subset \mathbb{Z}$  be a finite subset. We associate to  $A$  the function  $\beta = \beta_A : \mathbb{N}_+ \rightarrow \mathbb{Z}$  defined for all  $n \geq 1$  by

$$\beta_A(n) = |nA| - (2n - 1)(|A| - 1).$$

**Notation 1.2.** We denote by  $\mathcal{B}(A)$  the positive support in  $2 + \mathbb{N}$  of the function  $\beta_A$ , i.e.

$$\mathcal{B}(A) = \{n \geq 2 \mid \beta_A(n) \geq 1\}.$$

For instance,  $2 \in \mathcal{B}(A)$  if and only if  $|2A| \geq 3|A| - 2$ . Interestingly, the failure of this condition, namely the inequality  $|2A| \leq 3|A| - 3$ , is the key hypothesis of the famous Freiman's  $3k - 3$  Theorem in additive combinatorics [5].

**Example 1.3.** If  $|A| = 0$  or  $1$ , then  $\mathcal{B}(A)$  is infinite. Indeed, if  $A = \emptyset$ , then  $|nA| = 0$  and so  $\beta_\emptyset(n) = 2n - 1$  for all  $n \geq 1$ . Thus  $\mathcal{B}(\emptyset) = 2 + \mathbb{N}$  in that case. Similarly, if  $|A| = 1$ , then  $\beta_A(n) = 1$  for all  $n \geq 1$ . So here again  $\mathcal{B}(A) = 2 + \mathbb{N}$ .

In sharp contrast, Theorem 3.3 below states that if  $S \subset \mathbb{N}$  is a numerical semigroup of genus  $g \geq 2$ , then  $\mathcal{B}(\mathbb{N} \setminus S)$  is finite.

**Example 1.4.** Let  $S = \langle 3, 7 \rangle$ . Then  $\mathbb{N} \setminus S = \{1, 2, 4, 5, 8, 11\}$  and  $\beta_{\mathbb{N} \setminus S}(n) = 0$  for all  $n \geq 2$  as easily seen. In particular,  $\mathcal{B}(\mathbb{N} \setminus S) = \emptyset$ .

More generally, it was shown in [8, 10] that  $\beta_{\mathbb{N} \setminus S}(n) = 0$  for all symmetric numerical semigroups  $S$  of multiplicity  $m \geq 3$  and all  $n \geq 2$ . We shall not use this result below, but instead give a short self-contained proof of an immediate consequence, namely that  $\mathcal{B}(\mathbb{N} \setminus S)$  is empty in that case.

In fact,  $\mathcal{B}(\mathbb{N} \setminus S)$  is empty in most cases. Indeed, Buchweitz discovered in 1980 that the condition  $\mathcal{B}(\mathbb{N} \setminus S) = \emptyset$  is necessary for  $S$  to be a Weierstrass semigroup. By constructing instances where this condition fails, Buchweitz [1] was able to negate the longstanding conjecture by Hurwitz [6] according to which all numerical semigroups of genus  $g \geq 2$  are Weierstrass semigroups. His first counterexample was  $S = \langle 13, 14, 15, 16, 17, 18, 20, 22, 23 \rangle$ , with corresponding gapset

$$G = \mathbb{N} \setminus S = [1, 12] \cup \{19, 21, 24, 25\}$$

of cardinality 16. Then  $2G = [2, 50] \setminus \{39, 41, 47\}$ , so that  $|2G| = 46$  and  $\beta_G(2) = 46 - 3 \cdot 15 = 1$ , implying  $2 \in \mathcal{B}(G)$  and thus impeding  $S$  to be a Weierstrass semi-group. For more information on Buchweitz's condition and Weierstrass semi-groups, see e.g. [3, 7].

Here are the contents of this paper. In Section 2, we recall a result of Nathanson in additive combinatorics and we use it to study the asymptotic behavior of the function  $\beta_A(n)$ . In Section 3, we introduce the Buchweitz set of a numerical semi-group and we prove our main results. Section 4 concludes the paper with open questions on the possible shapes of the sets  $\mathcal{B}(A)$ .

## 2 Sumset growth

Given finite subsets  $A, B$  of a commutative monoid  $(M, +)$ , we denote as usual

$$A + B = \{a + b \mid a \in A, b \in B\},$$

the *sumset* of  $A, B$ , and  $2A = A + A$ . More generally, if  $n \geq 2$ , we denote  $nA = A + (n-1)A$ , where  $1A = A$ . The set  $nA$  is called the  $n$ -fold sumset of  $A$ .

A classical question in additive combinatorics is, how does  $|nA|$  grow with  $n$ ? Here we only consider the case  $M = \mathbb{Z}$ . We shall need the following result of Nathanson [9, Theorem 1.1].

**Theorem 2.1.** *Let  $A_0 \subset \mathbb{N}$  be a finite subset of cardinality  $k \geq 2$ , containing 0 and such that  $\gcd(A_0) = 1$ . Let  $a_0 = \max(A_0)$ . Then there exist integers  $c, d$  and subsets  $C \subseteq [0, c-2], D \subseteq [0, d-2]$  such that*

$$nA_0 = C \sqcup [c, a_0n-d] \sqcup (a_0n-D)$$

for all  $n \geq \max\{(|A_0|-2)(a_0-1)a_0, 1\}$ .

As pointed out in [9], the hypotheses  $0 \in A_0$  and  $\gcd(A_0) = 1$  are not really restrictive. Indeed, for any finite set  $A \subset \mathbb{Z}$  with  $|A| \geq 2$ , the simple transformation  $A \mapsto A_0 = (A - \alpha)/d$ , where  $\alpha = \min(A)$  and  $d = \gcd(A - \alpha)$ , yields a set  $A_0$  satisfying these hypotheses and such that  $|nA_0| = |nA|$  for all  $n$ . In view of our applications to gaps, we shall need the following version.

**Corollary 2.2.** *Let  $A \subset \mathbb{N}_+$  be a finite subset containing  $\{1, 2\}$ . Let  $a = \max(A)$ . Then there is an integer  $b \leq 1$  such that*

$$|nA| = (a-1)n + b$$

for all  $n \geq (|A|-2)(a-2)(a-1)$ .

*Proof.* Set  $A_0 = A - 1$  and  $a_0 = a - 1$ . Then  $A_0$  contains  $\{0, 1\}$ , hence it satisfies the hypotheses of Theorem 2.1. Using the same notation, its conclusion implies

$$(1) \quad |nA_0| = a_0 n + b$$

for all  $n \geq \max\{|A_0| - 2, (a_0 - 1)a_0, 1\}$ , where

$$b = |C| + |D| - c - d + 1.$$

Note that  $b \leq 1$  since  $|C| \leq \max(0, c - 1)$ ,  $|D| \leq \max(0, d - 1)$ . The desired statement follows from (1) since  $|nA| = |nA_0|$  for all  $n \geq 0$ .  $\square$

## 2.1 Asymptotic behavior of $\beta_A(n)$

We now study the evolution of  $\beta_A(n)$  as  $n$  grows.

**Theorem 2.3.** *Let  $A \subset \mathbb{N}_+$  be a finite set containing  $\{1, 2\}$ . Let  $f = \max(A)$  and  $g = |A|$ . Then*

$$\lim_{n \rightarrow \infty} \beta_A(n) = \begin{cases} -\infty & \text{if } f \leq 2g - 2, \\ +\infty & \text{if } f \geq 2g. \end{cases}$$

Finally if  $f = 2g - 1$ , then  $\beta_A(n)$  is constant and nonpositive for  $n$  large enough.

*Proof.* By Corollary 2.2, we have  $|nA| = (f - 1)n + b$  for some integer  $b \leq 1$  and for  $n$  large enough. Hence

$$\begin{aligned} \beta_A(n) &= (f - 1)n + b - (2n + 1)(g - 1) \\ &= (f - 2g + 1)n + b + 1 - g \end{aligned}$$

for  $n$  large enough. The claims for  $f \leq 2g - 2$  and  $f \geq 2g$  follow. If  $f = 2g - 1$ , then  $\beta_A(n) = b + 1 - g \leq 0$  for  $n$  large enough, since  $b \leq 1$  and  $g \geq 2$ .  $\square$

**Corollary 2.4.** *Let  $A \subseteq \mathbb{N}_+$  be a finite set containing  $\{1, 2\}$ . Let  $f = \max(A)$  and  $g = |A|$ . Then  $\mathcal{B}(A)$  is finite if and only if  $f \leq 2g - 1$ .*

*Proof.* If  $f \geq 2g$ , then  $\lim_{n \rightarrow \infty} \beta_A(n) = \infty$  by the theorem, whence  $\beta_A(n) \geq 1$  for all large enough  $n$ . Thus  $\mathcal{B}(A)$  is infinite in this case. If  $f \leq 2g - 1$ , the theorem implies  $\beta_A(n) \leq 0$  for  $n$  large enough, whence  $\mathcal{B}(A)$  is finite in that case.  $\square$

### 3 Application to numerical semigroups

**Definition 3.1.** Let  $S \subseteq \mathbb{N}$  be a numerical semigroup. We define the Buchweitz set of  $S$  as  $\text{Buch}(S) = \mathcal{B}(\mathbb{N} \setminus S)$ . Explicitly, setting  $G = \mathbb{N} \setminus S$ , we have

$$\begin{aligned}\text{Buch}(S) &= \{n \geq 2 \mid |nG| > (2n-1)(|G|-1)\} \\ &= \{n \geq 2 \mid \beta_G(n) \geq 1\}.\end{aligned}$$

In this section, we first prove that  $\text{Buch}(S)$  is *finite* for all numerical semigroups  $S$  of genus  $g \geq 2$ . We then show, by explicit construction, that the cardinality of  $\text{Buch}(S)$  may be arbitrarily large.

#### 3.1 Finiteness of $\text{Buch}(S)$

We start with a well known inequality linking the Frobenius number and the genus of a numerical semigroup.

**Proposition 3.2.** Let  $S \subset \mathbb{N}$  be a numerical semigroup with Frobenius number  $f$  and genus  $g \geq 1$ . Then  $f \leq 2g-1$ .

*Proof.* Let  $x \in S \cap [0, f]$ . Then  $f - x \notin S$  since  $S$  is stable under addition and  $x + (f - x) = f \notin S$ . Hence, the map  $x \mapsto f - x$  induces an injection

$$S \cap [0, f] \hookrightarrow \mathbb{N} \setminus S.$$

Since  $|S \cap [0, f]| = (f+1) - g$ , it follows that  $f \leq 2g-1$ , as claimed.  $\square$

Recall that  $S$  is said to be *symmetric* if  $|S \cap [0, f]| = |\mathbb{N} \setminus S|$ , i.e. if  $f = 2g-1$ . A classical result of Sylvester states that any numerical semigroup of the form  $S = \langle a, b \rangle$  with  $\gcd(a, b) = 1$  is symmetric.

**Theorem 3.3.** Let  $S \subseteq \mathbb{N}$  be a numerical semigroup of genus  $g \geq 2$ . Then  $\text{Buch}(S)$  is finite.

*Proof.* Let  $G = \mathbb{N} \setminus S$ . Then  $\text{Buch}(S) = \mathcal{B}(G)$  by definition. We have  $g = |G| \geq 2$ . Let  $f = \max(G)$  be the Frobenius number of  $S$ . Let  $m = \min(S \setminus \{0\})$  be the multiplicity of  $S$ . Then  $m \geq 2$  since  $g \geq 2$ , and  $[1, m-1] \subseteq G$ .

Assume first  $m \geq 3$ . Then  $\{1, 2\} \subseteq G$ . Hence Corollary 2.4 applies, and since  $f \leq 2g-1$  by Proposition 3.2, it yields that  $\mathcal{B}(G)$  is finite, as desired.

Assume now  $m = 2$ . Then  $S = \langle 2, b \rangle$  with  $b$  odd and  $b \geq 5$  since  $|G| \geq 2$ . At this point, we might conclude the proof right away using what is known in

the symmetric case [8, 10]. However, for the convenience of the reader, let us give a short self-contained argument. We have  $G = \{1, 3, \dots, b-2\}$ , i.e. all odd numbers from 1 to  $b-2$ . Hence  $G-1 = \{0, 2, \dots, b-3\}$  and  $\gcd(G-1) = 2$ . Set  $A = (G-1)/2 = [0, k]$ , where  $k = (b-3)/2$ . For all  $n \geq 1$ , we have

$$|nG| = |nA| = |nk + 1| = n(|G| - 1) + 1.$$

Therefore  $\beta_G(n) = -(n-1)|G| + n$ , whence  $\beta_G(n) \leq 0$  for all  $n \geq 2$ . It follows that  $\mathcal{B}(G) = \emptyset$  and we are done.  $\square$

### 3.2 Unboundedness of $|\text{Buch}(S)|$

We show here, by explicit construction, that  $|\text{Buch}(S)|$  may be arbitrarily large.

**Proposition 3.4.** *For any integer  $b \geq 3$ , there exists a numerical semigroup  $S$  such that  $\text{Buch}(S) = [2, b]$ .*

*Proof.* Let  $k = b-2$ , and let  $S$  be the numerical semigroup of multiplicity  $m = 6k+15$  and depth  $q = 2$  whose corresponding gapset  $G = \mathbb{N} \setminus S$  is given by

$$(2) \quad G = [1, m-1] \sqcup \{2m-7, 2m-5, 2m-2, 2m-1\}.$$

We claim that  $\text{Buch}(S) = [2, k+2]$ . Indeed, we will show a more precise statement, namely

$$\beta_G(n) = \begin{cases} 1 & \text{if } n = 2, \\ 2 & \text{if } 3 \leq n \leq k+2, \\ -6(n-k-3) & \text{if } n \geq k+3. \end{cases}$$

Let  $A = (2m-1) - G$ . Then  $\beta_G(n) = \beta_A(n)$  since  $|nG| = |nA|$  for all  $n \geq 1$ . We have

$$A = [0, 1] \sqcup \{4, 6\} \sqcup [m, 2(m-1)].$$

Let us compute  $2A$  and  $3A$ . We obtain

$$\begin{aligned} 2A &= [0, 2] \sqcup [4, 8] \sqcup \{10, 12\} \sqcup [m, 4(m-1)], \\ 3A &= [0, 14] \cup \{16, 18\} \cup [m, 6(m-1)]. \end{aligned}$$

In general, we have

$$(3) \quad nA = ([0, 6n-4] \sqcup \{6n-2, 6n\}) \cup [m, 2n(m-1)]$$

for all  $n \geq 3$ , as easily verified by induction on  $n$ .

Let us determine  $|nA|$  for all  $n \geq 1$ . Note first that *the union in (3) is disjoint if and only if  $6n+1 \leq m$* . Moreover, as  $m = 6k+15$ , we have

$$6n+1 \leq m \iff n \leq k+2.$$

In contrast, if  $n \geq k+3$ , i.e. if  $6n-3 \geq m$ , then the union in (3) collapses to a single interval and we get

$$nA = [0, 2n(m-1)].$$

Summarizing, we have

$$|nA| = \begin{cases} m+3 & \text{if } n=1, \\ 3m+7 & \text{if } n=2, \\ (2n-1)(m-1) + 6n-1 & \text{if } 3 \leq n \leq k+2, \\ 2n(m-1)+1 & \text{if } n \geq k+4. \end{cases}$$

The stated formula for  $\beta_G(n) = \beta_A(n) = |nA| - (2n-1)(|A|-1)$  follows. Hence  $\text{Buch}(S) = [2, k+2]$ , as claimed.  $\square$

This family of numerical semigroups was inspired by the *PF-semigroups* introduced in [4].

### 3.3 More intervals

What are the possible shapes of  $\text{Buch}(S)$  when  $S$  varies? We do not know in general. By Proposition 3.4, any finite integer interval  $I$  with  $|I| \geq 2$  and  $\min(I) = 2$  may be realized as  $I = \text{Buch}(S)$  for some numerical semigroup  $S$ . Here we present families of numerical semigroups  $S$  realizing as  $\text{Buch}(S)$  all finite integer intervals  $I$  with  $|I| \geq 2$  and  $\min(I) \in \{3, 4, 5, 6\}$ .

**Proposition 3.5.** *Let  $k \geq 1$ . Let  $S$  be the numerical semigroup of multiplicity  $m = 6k+19$  and depth  $q = 2$  whose corresponding gapset  $G = \mathbb{N} \setminus S$  is given by*

$$(4) \quad G = [1, m-1] \sqcup \{2m-7, 2m-6, 2m-2, 2m-1\}.$$

*Then  $\text{Buch}(S) = [3, k+3]$ .*

*Proof.* Let again  $A = (2m-1) - G = [0,1] \sqcup \{5,6\} \sqcup [m,2(m-1)]$ . We then have

$$\begin{aligned} 2A &= [0,2] \sqcup [5,7] \sqcup [10,12] \sqcup [m,4(m-1)], \\ 3A &= [0,3] \sqcup [5,8] \sqcup [10,13] \sqcup [15,18] \sqcup [m,6(m-1)], \\ 4A &= [0,24] \sqcup [m,8(m-1)]. \end{aligned}$$

It follows that  $nA = [0,6n] \cup [m,2n(m-1)]$  for all  $n \geq 4$ . In particular, if  $6n \geq m$  then  $nA = [0,2n(m-1)]$ . Therefore,

$$\beta_G(n) = \beta_A(n) = \begin{cases} 0 & \text{if } n = 2, \\ 1 & \text{if } n = 3, \\ 4 & \text{if } 4 \leq n \leq k+3, \\ 6k-6n+22 & \text{if } n \geq k+4. \end{cases}$$

Hence  $\text{Buch}(S) = [3, k+3]$ , as claimed.  $\square$

**Proposition 3.6.** *For  $k \geq 1$  and  $i \in \{1,2,3\}$ , let  $S_i$  be the numerical semigroup with  $G_i = \mathbb{N} \setminus S_i$  given by*

$$\begin{aligned} G_1 &= [1, m_1-1] \sqcup \{2m_1-6, 2m_1-2, 2m_1-1\}, \\ G_2 &= [1, m_2-1] \sqcup \{2m_2-10, 2m_2-4, 2m_2-3, 2m_2-2\}, \\ G_3 &= [1, m_3-1] \sqcup \{2m_3-10, 2m_3-9, 2m_3-2\}, \end{aligned}$$

where  $m_1 = 4k+22$ ,  $m_2 = 7k+44$  and  $m_3 = 5k+55$ , respectively. Then

$$\text{Buch}(S_1) = [4, k+4], \quad \text{Buch}(S_2) = [5, k+5], \quad \text{Buch}(S_3) = [6, k+6].$$

*Proof.* Similar to the proofs of Propositions 3.4 and 3.6. We omit it here.  $\square$

Having realized all finite integer intervals  $I$  with  $|I| \geq 2$  and  $\min(I) \in [2,6]$  as  $I = \text{Buch}(S)$  for a suitable numerical semigroups  $S$ , is it possible to do the same for all finite integer intervals  $I$  with  $\min(I) \geq 7$ ? We do not know in general. But here is a particular case where  $\min(I)$  can be arbitrarily large. It is based on a family of numerical semigroups found in [8].

**Proposition 3.7.** *For any integer  $k \geq 1$ , there is a numerical semigroup  $S$  such that  $\text{Buch}(S) = [7+2k, 7+4k]$ .*

*Proof.* For  $k \geq 1$ , let  $S$  be the numerical semigroup minimally generated by the set  $T_1 \cup T_2 \cup T_3$ , where

$$\begin{aligned} T_1 &= [44+27k+4k^2, 79+51k+8k^2], \\ T_2 &= [81+51k+8k^2, 84+53k+8k^2], \\ T_3 &= [87+53k+8k^2, 87+54k+8k^2]. \end{aligned}$$

The corresponding gapset  $G = \mathbb{N} \setminus S$  is then given by

$$G = [1, 43 + 27k + 4k^2] \cup \{80 + 51k + 8k^2, 85 + 53k + 8k^2, 86 + 53k + 8k^2\}.$$

Let  $A = (86 + 53k + 8k^2) - G$ . Then

$$A = [0, 1] \sqcup \{6 + 2k\} \sqcup [43 + 26k + 4k^2, 85 + 53k + 8k^2],$$

of cardinality  $|A| = 46 + 27k + 4k^2$ . The  $n$ -fold sumsets of  $A$  are then given by

$$(5) \quad nA = [0, n] \cup \left( \bigcup_{i=1}^n [i(6+2k), i(6+2k) + n - i] \right) \cup [43 + 26k + 4k^2, (85 + 53k + 8k^2)n].$$

• Assume first  $2 \leq n < 6 + 2k$ . In this case, we have

$$\begin{aligned} 0 < n < 6 + 2k < 6 + 2k + n - 1 < \dots < (n-1)(6+2k) \\ &< (n-1)(6+2k) + n - 1 < n(6+2k) < (7+2k)(6+2k) + 1 \\ &= 43 + 26k + 4k^2 < (85 + 53k + 8k^2)n. \end{aligned}$$

Thus, all the sets appearing in (5) are disjoint and the cardinality of  $nA$  is equal to

$$\begin{aligned} (n+1) + \sum_{i=1}^n i + ((8k^2 + 53k + 85)n - (4k^2 + 26k + 43) + 1) = \\ -41 - 26k - 4k^2 + (173n)/2 + 53kn + 8k^2n + n^2/2. \end{aligned}$$

Thus,

$$(6) \quad \begin{aligned} \beta_G(n) &= (-41 - 26k - 4k^2 + (173n)/2 + 53kn + 8k^2n + n^2/2) \\ &\quad - (4k^2 + 27k + 46 - 1)(2n - 1) = \\ &\quad (4 + k) - \left(\frac{7}{2} + k\right)n + \frac{1}{2}n^2 \end{aligned}$$

for every  $n \in [2, 5 + 2k]$ .

The only difference between the case  $n = 6 + 2k$  and the previous one is that the sets  $[0, n]$  and  $[6 + 2k, 6 + 2k + n - 1]$  have a nonempty intersection, equal to  $\{6 + 2k\}$ . Replacing  $n$  by  $6 + 2k$  and subtracting one, we obtain  $\beta_G(6 + 2k) = 0$ .

- Assume now  $6 + 2k < n \leq 11 + 4k$ . The sequence of sets  $[0, n]$  and

$$[6 + 2k, 6 + 2k + n - 1], \dots, [(n - 5 - 2k)(6 + 2k), (n - 5 - 2k)(6 + 2k) + (5 + 2k)]$$

verifies that the intersection of any two consecutive terms is nonempty. Moreover, their union is the interval  $[0, (n - 5 - 2k)(6 + 2k) + (5 + 2k)]$  whose cardinality is equal to  $(n - 5 - 2k)(6 + 2k) + (5 + 2k) + 1$ . For  $i = n - 4 - 2k, \dots, 6 + 2k$  the intervals are disjoint with all the others sets appearing in the expression (5); the cardinality of the union of these sets is equal to  $\sum_{i=n-2k-5}^{2k+5} i$ . For every  $i = 7 + 2k, \dots, n$  the intersection

$$[i(6 + 2k), i(6 + 2k) + n - i] \cap [43 + 26k + 4k^2, (8k^2 + 53k + 85)n]$$

is nonempty, except for  $n = 7 + 2k$ . Since  $(7 + 2k)(6 + 2k) = 42 + 26k + 4k^2$ , the set

$$\left( \bigcup_{i=7+2k}^n [i(6 + 2k), i(6 + 2k) + n - i] \right) \cup [43 + 26k + 4k^2, (85 + 53k + 8k^2)n]$$

is equal to  $[42 + 26k + 4k^2, (85 + 53k + 8k^2)n]$ , and the cardinality of this set is  $((8k^2 + 53k + 85)n - 4k^2 - 26k - 42) + 1$ . Putting all the above together, we have that if  $6 + 2k < n \leq 11 + 4k$ , the set  $nA$  has cardinality equal to

$$\begin{aligned} & ((n - 5 - 2k)(6 + 2k) + (5 + 2k) + 1) + \\ & \sum_{i=n-2k-5}^{2k+5} i + ((8k^2 + 53k + 85)n - 4k^2 - 26k - 42) + 1 = \\ & - 65 - 46k - 8k^2 + (193n)/2 + 57kn + 8k^2n - n^2/2, \end{aligned}$$

and therefore

$$\begin{aligned} (7) \quad \beta_G(n) &= -65 - 46k - 8k^2 + (193n)/2 + 57kn + 8k^2n - n^2/2 \\ &\quad - (2n - 1)(46 + 27k + 4k^2 - 1) = \\ &\quad (-20 - 19k - 4k^2) + (3k + \frac{13}{2})n - \frac{n^2}{2} \end{aligned}$$

for every  $n \in [6 + 2k, 11 + 4k]$ .

- Finally, assume  $11 + 4k < n$ . The set

$$[0, n] \cup \left( \bigcup_{i=1}^{6+2k} [i(6+2k), i(6+2k) + n - i] \right) \cup [43 + 26k + 4k^2, (85 + 53k + 8k^2)n]$$

is equal to  $[0, (85 + 53k + 8k^2)n]$  and the remaining intervals are contained in this union. So we have  $nA = [0, (85 + 53k + 8k^2)n]$  and therefore

$$(8) \quad \beta_G(n) = - (4k^2 + 27k + 46 - 1)(2n - 1) + (8k^2 + 53k + 85)n + 1 = \\ (4k^2 + 27k + 46) - (k + 5)n$$

for every  $n > 11 + 4k$ .

Combining  $\beta_G(6+2k) = 0$  with the formulation of (6), (7) and (8) for  $\beta_G(n)$ , we get the following formulas:

$$\beta_G(n) = \begin{cases} (4+k) - (\frac{7}{2} + k)n + \frac{1}{2}n^2 & \text{if } 2 \leq n < 6+2k, \\ 0 & \text{if } n = 6+2k, \\ (-20 - 19k - 4k^2) + (3k + \frac{13}{2})n - \frac{n^2}{2} & \text{if } 6+2k < n \leq 11+4k, \\ (4k^2 + 27k + 46) - (k+5)n & \text{if } 11+4k < n. \end{cases}$$

Let  $k \geq 1$  be fixed. For  $2 \leq n < 6+2k$ , the formula of  $\beta_G(n)$  is a degree two polynomial in  $n$  with positive leading coefficient such that  $\beta_G(2) = -1 - k < 0$  and  $\beta_G(5+2k) = -1 - k < 0$ . We have therefore  $\beta_G(n) < 0$  for every  $n = 2, \dots, 5+2k$ .

If  $n > 11 + 4k$ , we now have that  $\beta_G(n)$  is a degree one polynomial with negative leading coefficient and such that  $\beta_G(12 + 4k) = -14 - 5k < 0$ . So  $\beta_G(n) < 0$  for every  $n > 11 + 4k$ .

Finally, if  $6+2k < n \leq 11+4k$  the function  $\beta_G(n)$  is a degree two polynomial in  $n$  with negative leading coefficient. As in addition  $\beta_G(7+2k) = 1+k$ ,  $\beta_G(7+4k) = 1$  and  $\beta_G(8+4k) = -k$ , the only positive values that we have in this part are for  $n \in [7+2k, 7+4k]$ .

Since  $\beta_G(n) \leq 0$  except for  $n \in [7+2k, 7+4k]$ , the set  $\text{Buch}(S)$  is equal to  $[7+2k, 7+4k]$ .  $\square$

## 4 Concluding remarks

The current knowledge on the structure of  $\mathcal{B}(A)$  for finite subsets  $A \subset \mathbb{Z}$  is very scarce, even for gapsets. Do they have some special shape or property? We end this paper with three questions based on the few currently available observations.

**Question 4.1.** Let  $A \subset \mathbb{Z}$  be a finite subset, or more specifically a gapset. Is the set  $\mathcal{B}(A)$  always an interval of integers?

**Question 4.2.** Even more so, is the function  $\beta_A(n)$  unimodal?

**Question 4.3.** In sharp contrast with the above questions, let  $T \subset 2 + \mathbb{N}$  be any finite subset. Does there exist a finite subset  $A \subset \mathbb{N}$ , or more specifically a gapset, such that  $\mathcal{B}(A) = T$ ?

**Acknowledgement.** Part of this paper was written during a visit of the first-named author to the Universidad de Cádiz (Spain) which was partially supported by *Ayudas para Estancias Cortas de Investigadores* (EST2019-039, Programa de Fomento e Impulso de la Investigación y la Transferencia en la Universidad de Cádiz). The second, third and fourth-named authors were partially supported by Junta de Andalucía research group FQM-366 and by the project MTM2017-84890-P (MINECO/FEDER, UE).

## References

- [1] R.-O. BUCHWEITZ, On Zariski's criterion for equisingularity and non-smoothable monomial curves. Preprints 115s, Universität Hannover, Institut für Mathematik (1980).
- [2] M. DELGADO, P.A. GARCÍA-SÁNCHEZ AND J. MORAIS, “Numericalsgps”: a GAP package on numerical semigroups. <http://www.gap-system.org/Packages/numericalsgps.html>
- [3] D. EISENBUD AND J. HARRIS, Existence, decomposition, and limits of certain Weierstrass points. *Invent. math.* 87 (1987) 495–515.
- [4] J.I. GARCÍA-GARCÍA, D. MARÍN-ARAGÓN, F. TORRES AND A. VIGNERON-TENORIO, On reducible non-Weierstrass semigroups. [arXiv:2005.12896](https://arxiv.org/abs/2005.12896) [math.AG].
- [5] G.A. FREIMAN, The addition of finite sets. I. (Russian) *Izv. Vysš. Učebn. Zaved. Matematika* no. 6 (13) (1959) 202–213.
- [6] A. HURWITZ, Über algebraischer Gebilde mit eindeutigen Transformationen in sich. *Math. Ann.* 41 (1893) 403–442.
- [7] N. KAPLAN AND L. YE, The proportion of Weierstrass semigroups. *J. Algebra* 373 (2013) 377–391.

- [8] J. KOMEDA, Non-Weierstrass numerical semigroups. *Semigroup Forum* 57 (1998) 157–185.
- [9] M.B. NATHANSON, *Additive Number Theory, Inverse Problems and the Geometry of Sumsets*. Graduate Texts in Mathematics, vol. 165, Springer, New York, 1996.
- [10] G. OLIVEIRA, Weierstrass semigroups and the canonical ideal of non-trigonal curves. *Manuscripta math.* 71 (1991) 431–450.

**Authors' addresses:**

Shalom Eliahou<sup>a,b</sup>

<sup>a</sup>Univ. Littoral Côte d'Opale, UR 2597 - LMPA - Laboratoire de Mathématiques Pures et Appliquées Joseph Liouville, F-62100 Calais, France

<sup>b</sup>CNRS, FR2037, France

**e-mail:** eliahou@univ-littoral.fr

Juan Ignacio García-García<sup>c,d</sup>

**e-mail:** ignacio.garcia@uca.es

Daniel Marín-Aragón<sup>c</sup>

<sup>c</sup>Departamento de Matemáticas, Universidad de Cádiz, E-11510 Puerto Real (Cádiz, Spain).

**e-mail:** daniel.marin@uca.es

Alberto Vigneron-Tenorio<sup>d,e</sup>

<sup>e</sup>Departamento de Matemáticas, Universidad de Cádiz, E-11406 Jerez de la Frontera (Cádiz, Spain).

<sup>d</sup>INDESS (Instituto Universitario para el Desarrollo Social Sostenible), Universidad de Cádiz, E-11406 Jerez de la Frontera (Cádiz, Spain).

**e-mail:** alberto.vigneron@uca.es