



# Disentangling the response of fishes to recreational fishing over 30 years within a fringing coral reef reserve network

A.K. Cresswell, T.J. J Langlois, S.K. Wilson, Joachim Claudet, D.P. Thomson, M. Renton, C.J. Fulton, R. Fisher, M.A. Vanderklift, R.C. Babcock, et al.

## ► To cite this version:

A.K. Cresswell, T.J. J Langlois, S.K. Wilson, Joachim Claudet, D.P. Thomson, et al.. Disentangling the response of fishes to recreational fishing over 30 years within a fringing coral reef reserve network. Biological Conservation, 2019, 237, pp.514-524. 10.1016/j.biocon.2019.06.023 . hal-03033674

**HAL Id: hal-03033674**

**<https://hal.science/hal-03033674v1>**

Submitted on 1 Dec 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Disentangling the response of fishes to recreational fishing over 30 years within a fringing coral reef reserve network

Cresswell, A.K.<sup>1,2,3</sup>, Langlois, T.J.<sup>1,3</sup>, Wilson, S.K.<sup>3,4</sup>, Claudet, J.<sup>5,6</sup>, Thomson, D.P.<sup>2</sup>, Renton, M.<sup>1,7</sup>, Fulton, C.J.<sup>8</sup>, Fisher, R.<sup>3,9</sup>, Vanderklift, M.A.<sup>2</sup>, Babcock, R.C.<sup>3,10</sup>, Stuart-Smith, R. D.<sup>11</sup>, Haywood, M.D.E.<sup>10</sup>, Depczynski, M.<sup>3,9</sup>, Westera, M.<sup>12</sup>, Ayling, A.M.<sup>13</sup>, Fitzpatrick, B.<sup>14</sup>, Halford, A. R.<sup>15</sup>, McLean, D.L.<sup>3,4,9</sup>, Pillans, R. D.<sup>10</sup>, Cheal, A.J.<sup>9</sup>, Tinkler, P.<sup>16</sup>, Edgar, G. J.<sup>11</sup>, Graham, N.A.J.<sup>17</sup>, Holmes, T.H.<sup>4</sup>

<sup>1</sup> School of Biological Sciences, The University of Western Australia, Crawley, WA, Australia

<sup>2</sup> CSIRO Oceans & Atmosphere, Indian Ocean Marine Research Centre, Crawley, WA, 6009, Australia

<sup>3</sup> The UWA Oceans Institute, The University of Western Australia, Crawley, WA, Australia

<sup>4</sup> Marine Science Program, Western Australia Department of Biodiversity, Conservation and Attractions

<sup>5</sup> National Center for Scientific Research, PSL Université Paris, CRILOBE, USR 3278 CNRS-EPHE-UPVD, Maison des Océans, 195 rue Saint-Jacques 75005 Paris, France

<sup>6</sup> Laboratoire d'Excellence CORAIL, Moorea, French Polynesia.

<sup>7</sup> School of Agriculture and Environment, The University of Western Australia, Crawley, WA, Australia

<sup>8</sup> Research School of Biology, The Australian National University, Canberra ACT 2601, Australia

<sup>9</sup> Australian Institute of Marine Science, Indian Ocean Marine Research Centre, Crawley, WA, 6009, Australia

<sup>10</sup> CSIRO Oceans & Atmosphere, Queensland Biosciences Precinct, St Lucia, QLD, 4067, Australia

<sup>11</sup> Institute for Marine and Antarctic Studies, University of Tasmania. Private bag 49, Hobart, Tasmania 7001

<sup>12</sup> BMT, Level 4/20 Parkland Rd, Osborne Park, WA, 6017, Australia

<sup>13</sup> Sea Research, 20 Rattray Ave, c, QLD, 4800, Australia

<sup>14</sup> Oceanwise Australia Pty Ltd, Floreat, Perth, Western Australia 6014, Australia

<sup>15</sup> SPC, B.P. D5 – 98848 Noumea Cedex, New Caledonia

<sup>16</sup> Deakin University, School of Life and Environmental Sciences, Warrnambool, Vic, Australia

<sup>17</sup> Lancaster Environment Centre, Lancaster University, Lancaster, UK

**Corresponding author:** Anna Katrina Cresswell

+614 09 629 885

[anna.cresswell@csiro.au](mailto:anna.cresswell@csiro.au)

Indian Ocean Marine Research Centre

64 Fairway - Level 4, The University of Western Australia

Crawley WA 6009

Australia

## Abstract

Few studies on the effects of recreational fishing in isolation from commercial fishing exist. We used meta-analysis to synthesise 4,444 samples from 30 years (1987-2017) of fish surveys inside and outside a large network of highly protected reserves in the Ningaloo Marine Park, Western Australia, where the major fishing activity is recreational. Data were collected by different agencies, using varied survey designs and sampling methods. We contrasted the relative abundance and biomass of target and non-target fish groups between fished and reserve locations. We considered the influence of, and possible interactions between, seven additional variables: age and size of reserve, one of two reserve network configurations, reef habitat type, recreational fishing activity, shore-based fishing regulations and survey method. Taxa responded differently: the abundance and biomass inside reserves relative to outside was higher for targeted lethrinids, while other targeted (and non-targeted groups) were indistinguishable. Reef habitat was important for

explaining lethrinid response to protection, and this factor interacted with reserve size, such that larger reserves were only demonstrably more effective in the back reef and lagoon habitats. There was little evidence of changes in relative abundance and biomass of fishes with reserve age, or before and after rezoning and expansion of the reserve network. Our study demonstrates the complexities in quantifying fishing effects, highlighting some of the key factors and interactions that likely underlie the varied results to date in reserve assessments that should be considered in future reserve design and assessment.

## Keywords

Marine protected area; MPA; fisheries; coral reef; Ningaloo; adaptive management; recreational fishing; *Lethrinus*

## 1. Introduction

Anthropogenic activities continue to expand worldwide, particularly in the tropics, threatening natural systems and the ecosystem services they provide (Barlow *et al.*, 2018). As a result, ‘protected areas’ that seek to balance extractive activities with other socio-ecological values are increasingly being used to manage terrestrial and marine systems (Jenkins & Joppa, 2009; Sala *et al.*, 2018). Many studies have assessed the conservation effects of marine reserves (reviewed by Mosquera *et al.*, 2000; Russ, 2002), including quantitative syntheses of regional and global studies, with most finding higher abundance and size of targeted species within reserve boundaries in the case of ‘no-take’, or highly protected reserves (Lester *et al.*, 2009). The large majority of these findings are from regions with commercial fisheries, and less is documented about the impacts of recreational fisheries, despite several studies flagging the potentially high impacts of these fisheries (McPhee *et al.*, 2002; Coleman *et al.*, 2004; Cowx & Cooke, 2004; Lewin *et al.*, 2006). No-take reserves are a key tool for assessing the impacts of fishing (Ballantine, 2014) and while there are a handful of empirical studies that have demonstrated the effects of fishing, using inside outside comparisons, on targeted invertebrates (Shears *et al.*, 2006; Babcock *et al.*, 2007) and finfish (Denny *et al.*, 2004) a comprehensive assessment including reserves with different characteristics over long time frames is lacking. The magnitude of differences inside to outside reserves has been correlated with their design, in particular size and age, with larger and older reserves typically resulting in greater abundance and/or size of targeted fishes than reserves that are smaller or newly established (Claudet *et al.*, 2008; Edgar *et al.*, 2014). The effects of reserves vary among biomes, locations and taxa of interest (Côté *et al.*, 2005; Claudet *et al.*, 2010; Mora & Sale, 2011) and there are examples of reserves having negligible effects on targeted fish communities (McLaren *et al.*, 2015). In addition to size and age of reserves, explanations for this variability include high levels of cross-boundary movement by fishes (Pillans *et al.*, 2014) and minimal to no difference in fishing activity across no-take and fished areas due to accessibility and/or non-compliance by fishers (Bergseth *et al.*, 2017), all of which make disentangling the true effects of fishing more complicated.

Ideally assessments of the influence of reserves are based on replicated studies across multiple comparable reserves with long time series of biological data before and after reserve establishment (Underwood, 1993; Russ, 2002; Osenberg *et al.*, 2011). Yet such data are typically beyond the scope of single research programs, necessitating the integration of multiple datasets. ‘Adaptive management,’ involving changes to the number, size or boundaries of reserves in response to new scientific information, changes in fishing pressure or changing social attitudes (McCook *et al.*, 2010) further complicates long-term assessments. Ongoing improvement of ecological sampling methods and technologies has resulted in new survey methods being introduced to monitoring (Goetze *et al.*, 2015): video based methods (baited remote underwater video (BRUV) and diver operated video (DOV)) are now commonly used alongside or in place of the previously more common underwater visual census (UVC) (Mallet & Pelletier, 2014). Therefore, evaluations of reserves that have long-term datasets must have the capacity to incorporate and evolve with changes in reserve design and survey methods (Claudet & Guidetti, 2010). Other factors, including differences in habitat and benthic structure, have been shown to affect outcomes of reserve evaluation (Miller & Russ, 2014; Rees *et al.*, 2018b) and while these factors have been studied independently, few assessments consider multiple factors simultaneously, including possible interactions (Edgar *et al.*, 2014). Differences in fishing pressure outside of reserves will also directly impact inside to outside comparisons, yet data that quantify

103 localised variation in fishing activity at the scale of marine parks and typical reserve networks are rarely  
104 available (Lewin *et al.*, 2006).

105  
106 Here, we synthesise a unique 30 year dataset from within a multiple-use marine park at Ningaloo Reef,  
107 Western Australia. The type of fishing activity at Ningaloo Reef (almost exclusively recreational) in  
108 combination with a highly protected and regulated network of reserves that have undergone significant  
109 expansion during the study period, offers the opportunity to advance on previous studies and inform on the  
110 potential impacts of recreational fisheries. We integrate data from numerous agencies with varied survey  
111 designs and methods, and therefore use a meta-analytical approach to compare the abundance and biomass  
112 of select targeted and non-targeted tropical reef fish inside reserves with adjacent fished areas. We tested  
113 two hypotheses: (1) the relative abundance/biomass of targeted fish taxa will be greater inside reserves than  
114 outside due to recreational fishing activity; and (2) the observed relative abundance and biomass will vary  
115 with survey method, age and size of reserve, spatial variability in fishing activity (including shore-based  
116 fishing) and/or habitat. Our study offers four main novelties. First, the effect of recreational fishing on  
117 targeted species is assessed in isolation from commercial fishing. Second, we explicitly consider potential  
118 interactions between variables. Third, the influence of changes in the reserve network are considered in the  
119 context of the increasingly common adaptive management. Fourth, we consider the influence of shore-based  
120 fishing, which has rarely been investigated. We therefore provide advances on previous work that are of  
121 importance for future planning and assessment of protected areas.

## 122 123 124 **2. Material and methods**

### 125 126 *2.1 Study region*

127 Data for this study are from the Ningaloo Marine Park (NMP) on the western Australian coastline (22°S,  
128 113°E; Fig. 1). The Park covers the majority of Ningaloo Reef (a World Heritage site) which is a fringing  
129 coral reef almost 300 km in length. The reef encompasses a sheltered lagoon that is highly accessible by  
130 shore-based fishers and those operating recreational vessels (Smallwood & Beckley, 2012). Despite a  
131 relatively small permanent human population, this area is a popular tourism destination for recreational  
132 fishers (Sumner *et al.*, 2002; Smallwood & Beckley, 2012; Mitchell *et al.*, 2018). There have not been any  
133 major commercial fishing activities within the marine park since the 1970s, (for summary see pg. 78, CALM  
134 (2005) and pg. 70, DPIRD (2017)). Recreational spearfishing has additional restrictions of varying degrees  
135 outside of the reserves, with spearfishing prohibited along a 70km stretch of coast between Tantabiddi Well  
136 and Winderabandi Point, and spearfishing for Labridae and Serranidae prohibited throughout the Park  
137 (DPIRD, 2018) (Fig. 1).

138  
139 A network of eight no-take marine reserves was established as part of the Park in April 1987 to cover 10%  
140 of the total marine park area ~22,400 ha, (Fig. 1a) (CALM, 1989). In 2005, the majority of the existing eight  
141 reserves were expanded in size and 10 new reserves were added (Fig. 1b), increasing the reserve coverage to  
142 88,365 ha (34% of the NMP). At the same time, three reserves, covering 1,929 ha, were established as a part  
143 of the 28,616 ha Muiron Islands Marine Management Area (MIMMA), immediately adjacent to the northern  
144 boundary of the NMP (CALM, 2005). Together, the NMP and MIMMA form a continuous network  
145 (CALM, 2005). There is some variation in the regulations along the boundaries of the 21 current reserves,  
146 complicating terminology and analysis, with eight reserves allowing shore-based fishing from their coastal  
147 boundaries (Appendix A, CALM, 2005). According to recent classifications of marine reserves, the two  
148 forms of reserves in the present study, those with shore-based fishing prohibited and those where it is  
149 allowed, would classify as Fully Protected Areas and Highly Protected Areas, respectively (Horta e Costa *et al.*  
150 *et al.*, 2016), both of which would be expected to provide protection for fished species (Zupan *et al.*, 2018).  
151 We explicitly include consideration of the effect of shore-based fishing in our analyses.

### 152 153 *2.2 Survey data*

154 Data from all major research and monitoring programs surveying fish in the NMP over the last 30 years  
155 (1987 – 2017) were collated (Appendix B) to create a very large synthesis of information. Locations of

individual samples are given as Appendix C. Three different survey methods were used to census fish: Baited Remote Underwater stereo-Video (BRUV), Diver Operated stereo-Video (DOV) and Underwater Visual Census (UVC) (Langlois *et al.*, 2010; Murphy & Jenkins, 2010). The majority (90%) of surveys also estimated the length of fish (an *in situ* estimate of total length for UVC, and fork length measured from stereo-video for DOV and BRUV), which allowed estimates of biomass using formulae from FishBase (Froese, 2018) (Appendix B).

Data were organised hierarchically with a sample (individual UVC or DOV transect or a BRUV deployment) being the lowest level of replication. Samples were classified to the next hierarchical level and termed a 'comparison pair', based on the criteria: (i) that there were at least two samples inside and two samples outside a given reserve, (ii) these samples were collected within 2 weeks of each other, (iii) samples were collected more than 200 m from within or outside of the reserve boundaries (excluding one reserve, the small size of which meant this was not a logical rule), (iv) samples were collected using the same survey method within one of four *habitat* categories (see Table 1). Data satisfying these conditions consisted of 4,444 samples classified into 305 relative abundance comparison pairs and 3,892 samples classified into 268 relative biomass comparison pairs. These data covered seven of the initial eight reserves and 16 of the 21 current reserves (Appendix B).

### 2.3 Fish groups

We consider three main fish groups common at Ningaloo Reef, at family or subfamily and species level (Appendix D) which differ in terms of their behaviour and representation in fisheries catch reports. This included: parrotfishes (Scarinae), which are not typically targeted by fishers in Australia, and two groups which are highly targeted by recreational fishers in the region (Ryan *et al.*, 2013; Ryan *et al.*, 2015) that have different behaviours; emperors (Lethrinidae; mobile roving predators) and groupers (Epinephelinae; mostly site-attached ambush predators). Previous work has indicated both Epinephelinae and Lethrinidae are vulnerable to fishing and many species in both subfamilies are targeted across the Indo-Pacific (Abesamis *et al.*, 2014). Species level analyses included two species from Lethrinidae: the spangled emperor, *Lethrinus nebulosus*, which is recognised as the most highly targeted species in the region, consistently featuring at the top of the estimated catch for the bioregion over the 30-year study period, and the yellow-tailed emperor, *L. atkinsoni*, a species that is anecdotally retained by fishers and featured as the 6<sup>th</sup> most common species recorded in the 1998/9 catch survey, but was a minor component in subsequent surveys (Sumner *et al.*, 2002; Ryan *et al.*, 2013; Ryan *et al.*, 2015). The Chinaman Rockcod, *Epinephelus rivulatus* (Epinephelinae) was also considered, with catches comparable to those of *L. nebulosus* across the catch reports (Ryan *et al.*, 2013; Ryan *et al.*, 2015). Individual species were not considered from the Scarinae subfamily due to inconsistencies in the accuracy of identification of species from this family.

### 2.4 Meta-analysis

We used a mixed-effects meta-analytical approach to assess the effect of the reserves on fish abundance and biomass. We calculated effect sizes as log-ratios for each of the comparison pairs inside to outside the reserves (Claudet *et al.*, 2008) (see Appendix E for formulas). A constant was added to the mean abundance ( $c = 0.5$ ) and mean biomass ( $c = 100$  g) to allow calculation of the log ratio in cases where fish were absent either inside or outside (i.e. zero values). We ran a sensitivity analysis on the value of the constant (Appendix F) to determine an appropriate value. The size of the constant impacted the magnitude of the effect size, but in general did not influence the significance. Nonetheless, the exact magnitude of the overall effect size should be interpreted with caution. In cases where both the inside and outside mean count were zero, the samples were excluded from the analysis. Effect sizes were weighted by the inverse of the sum of the within- and among-study variances (Appendix E). Weighted effect sizes and variances were calculated using the *metafor* package (Viechtbauer, 2010) in the statistical program R (R Core Team, 2017) with the variance estimator set to "REML," restricted maximum likelihood estimator. Overall effect sizes were comparable for both abundance and biomass and for simplicity we presented the abundance results as these were available for a larger dataset, providing biomass results in Appendix H.

## 2.5 Sources of variability

We considered seven variables that might mediate the response of fish abundance and biomass to the presence of the reserves (Table 1): (i) the number of years between when a sample was collected and when the zoning went into place; (ii) initial or current zoning scheme (see Figure A1); (iii) survey method; (iv) four coarse *habitats* with distinct coral and algae assemblages: ‘exposed reef slope’, ‘reef flat’, ‘back reef & lagoon coral’, and ‘lagoon algae’; (v) spatial area of a reserve; (vi) an estimate of fishing pressure outside of individual reserves; (vii) on the presence of shore-based fishing zones adjacent to some reserves. Data were explored following the protocol of Zuur *et al.* (2010) and transformed to normalise their distribution where appropriate (see Table 1).

As all effect sizes were heterogeneous (Appendix G), we first explored the influence of the seven variables using weighted mixed-effects categorical meta-analyses and meta-regression, considering each variable as a moderator in isolation to determine which variables explained significant heterogeneity in the overall effect size (see Appendix E for formulas). We also investigated *reserve identity* to allow comparison between individual reserves. Given there were correlations among the variables and potential interactions and non-linear effects, we used weighted full-subsets generalised additive mixed modelling (*FSSgam*) (Fisher *et al.* 2018 A simple function for full-subsets multiple regression in ecology with R. *Ecology and Evolution*) to investigate the relative importance of each variable in explaining variability in the overall effect size for each fish group. The response variable, effect size  $e$ , was modelled with a Gaussian distribution using *gam()* in the *mgcv* package in R (Wood, 2011). *Years protection* and *boat fishing* were included as continuous smoothers in the *FSSgam* to allow for non-linear relationships. The distribution of *reserve size* was not much improved by transformation and  $\sqrt{\text{reserve size}}$  was therefore included in the model set as a linear predictor. *Reserve identity* was highly collinear with other variables (in particular *reserve size*), and therefore, rather than including this as a random effect, a smoother of the mean *latitude* of comparison pairs was included in all models (and as part of the null model). This yielded comparable results to including *reserve identity* as a random effect. Interactions between factor variables *habitat* and *shore fishing* and the continuous variables *reserve size* and *years protection* were tested. In all models the smoothing parameter was limited to a simple spline, allowing only monotonic relationships ( $k=3$ ) for all continuous variables except for *latitude*, which was unlimited. Summed AICc weights were used as a metric of variable importance to investigate the relative importance of each predictor variable across the full set of models (Anderson & Burnham, 2002). Variables included in the most parsimonious model (fewest variables and lowest estimated degrees of freedom within two units of the AICc) were plotted to visualise the shape and direction of relationships between the variables and the effect size. We interpret results of variable importance and the top models with caution and consider the results of the mixed-effects meta-analyses and meta-regression alongside the results of the *FSSgam*.

Lastly, given the importance of temporal patterns in investigations of protected areas, we explicitly investigate data from the Osprey reserve (see Fig. 1), the best temporally replicated reserve in the dataset. Using available and relatively consistently collected UVC and DOV data we estimated mean fish density as count per transect area. We tested for significant linear and quadratic relationships and fitted generalised additive mixed models between the density of *L. nebulosus* and survey year.

## 3. Results

When compared to areas open to fishing, Lethrinidae were on average 57% more abundant (78% more biomass) inside the reserves ( $e = 0.45 \pm 0.12$ , 95% CI, Fig. 2a), however the effect was heterogeneous ( $Q_T = 2002.6$ ,  $df = 301$ ,  $p < 0.001$ , Table G1). The most parsimonious model for Lethrinidae abundance consisted of an interaction between *habitat* and *reserve size* (Table 2), with the same true for biomass (Appendix H). The categorical meta-analysis supported the importance of *habitat* for relative abundance; showing it explained significant heterogeneity among effect sizes ( $Q_M = 39.5$ ,  $df = 3$ ,  $p < 0.001$ , Table G2) with the most positive effect identified in *back reef & lagoon coral* sites with an average of 93% more Lethrinidae inside the reserves ( $e = 0.66 \pm 0.14$ , 95% CI) (Fig. 2a, Fig. G1) in this habitat. On the *reef flat* Lethrinidae were 53%

more abundant inside the reserves ( $e = 0.42 \pm 0.32$ , 95%CI) while there was no significant effect on the *exposed reef slope* and a negative effect in the *lagoon algae* habitat (Fig. G1). The interaction of *reserve size* and *habitat* was evident as an increase in effect size with increasing *reserve size* in the *back reef & lagoon coral* habitat versus no clear trends in the other habitats.

*Lethrinus nebulosus* were on average 42% more abundant (86% more biomass) inside reserves than outside ( $e = 0.35 \pm 0.15$ , 95%CI, Fig 2a). The effect was heterogeneous ( $Q_T = 1971.1$ ,  $df = 256$ ,  $p < 0.001$ , Table G1). The most parsimonious model included the interaction between *habitat* and *reserve size* with these two variables also having the highest variable importance across the full-subsets model set (Table G3, Fig. 2b). The same was true in the biomass analysis (Appendix H). *Habitat* explained significant heterogeneity for relative fish abundance ( $Q_M = 32.5$ ,  $df = 3$ ,  $p < 0.001$ , Table G2) and *L. nebulosus* were on average 84% more abundant within *back reef & lagoon coral* sites inside the reserves ( $e = 0.61 \pm 0.17$ , 95%CI), whereas no differences were observed for the *reef flat* or *exposed reef slope* sites and a negative effect was observed for *lagoon algae* sites (Fig. G1). As for Lethrinidae, the interaction of *reserve size* and *habitat* was evident by an increase in the effect size with increasing *reserve size* in the *back reef & lagoon coral* habitat and no clear effects in the other habitats.

On average, the abundance of *L. atkinsoni* was 40% more abundant (60% more biomass) inside reserves than outside ( $e = 0.34 \pm 0.09$ , 95%CI). The effect was heterogeneous ( $Q_T = 1739.7$ ,  $df = 279$ ,  $p < 0.001$ , Table G1). The most parsimonious model included *zoning scheme* and *method*, which also had the highest importance according to weighted AICc. These two variables explained significant heterogeneity according to the categorical mixed-effects meta-analyses. Predictions indicated that the BRUV *method* contributed the most to the positive effect size of *L. atkinsoni* (Fig. 3c), though this was not significant, nor were the differences between initial and current zoning, showing a slightly higher effect size from the older zoning scheme. Multiple variables explained significant heterogeneity for *L. atkinsoni* according to the categorical meta-analysis and the meta-regression (Table G2), including *habitat* ( $Q_M = 14.6$ ,  $df = 3$ ,  $p < 0.001$ , Table G2). *Reef flat* sites had 94% higher abundance, ( $e = 0.66 \pm 0.26$ , 95%CI) and *back reef & lagoon coral* sites 43% higher abundance ( $e = 0.36 \pm 0.12$ , 95%CI) inside the reserves. There were no significant effects for the other habitats (Fig. G1). The biomass analysis for *L. atkinsoni* indicated that *years protection* may interact with *habitat*, and that on the reef flat the effect size was higher and showed a parabolic pattern with *years protection* (Fig. H2).

The effect size for Epinephelinae abundance was significantly negative with 9% fewer fishes inside than outside the reserves ( $e = -0.09 \pm 0.08$ , 95%CI), although this result was heterogeneous ( $Q_T = 1125.7$ ,  $df = 276$ ,  $p < 0.001$ , Table G1). Variable importance scores showed no variables with high importance relative to the Lethrinidae and *L. nebulosus* model sets. *Reserve size* and *years protection* were present in the most parsimonious model, while for the biomass it was *method* and *boat fishing* (Appendix H). There were weak increasing trends for both *reserve size* and *years protection*, however the lack of strongly important or consistent variables in these model sets means the results should be interpreted cautiously.

On average there was no significant difference inside to outside the reserves for *E. rivulatus* abundance ( $e = -0.06 \pm 0.09$ , 95%CI), though the effect was heterogeneous ( $Q_T = 477.3$ ,  $df = 166$ ,  $p < 0.001$ , Table G1). *Zoning scheme* and *boat fishing* had the highest variable importance across the model set and featured in the most parsimonious model. The effect size transitioned from no effect for low boat fishing activity, to a positive effect when there was high boat fishing activity, but the confidence intervals did not show this trend to be significant. The initial reserve network (in place longer) had a more positive effect than the newer reserves, but again this was not significant (Fig. 3e).

The control fish group, Scarinae, showed no significant difference inside to outside the reserves ( $e = -0.01 \pm 0.11$ , 95%CI) and this effect was heterogeneous ( $Q_T = 1701.1$ ,  $df = 260$ ,  $p < 0.001$ , Table G1). All variables had low importance according to AICc (Fig 2b, Table 2) and while *boat fishing* and *shore fishing* appear in the most parsimonious model we interpret this with caution. In the biomass analysis *habitat* made up the most parsimonious model (Appendix H).

In the full-subsets analysis *reserve size* and *habitat* appeared with the highest variable importance (for Lethrinidae and *L. nebulosus*) while other variables - survey method, years protection, zoning scheme and shore fishing - had low importance across all six fish groups. In many cases the heterogeneity statistics from the mixed-effect meta-analysis models supported the findings of the full-subsets analysis, but for some variables such as shore fishing, the meta-analysis indicated this variable explained significant heterogeneity for all fish groups, except for *L. nebulosus*, while the full-subsets gave this variable low relative importance.

The temporal investigation of effect sizes for the most highly targeted fish, *L. nebulosus*, at Osprey reserve gave results that generally confirmed what was found in the full meta-analysis for *L. nebulosus*, showing effect sizes that are mostly positive though time, with higher abundance and biomass inside than outside (Fig 4). There were not show strong or significant patterns with time, except for the abundance density outside of the reserve, which had a significantly negative linear trend ( $P=0.032$ ). Generalised additive model fits indicate that, particularly in the latter half of the study period, both abundance and biomass may have declined both inside and outside the reserve, while there is some indication that abundance initially increased inside of the reserve following establishment. However, confidence in these trends is low and the gam fits were not statistically different from null models (except for abundance density outside the reserve,  $P = 0.048$ ).

#### 4. Discussion

Across the 30 year synthesis higher abundance and biomass of certain targeted fish taxa inside the reserves suggests that recreational fishing can have significant effects in isolation from commercial harvest, as also shown in some previous studies (Denny *et al.*, 2004; Shears *et al.*, 2006; Babcock *et al.*, 2007). We found the extent of this effect was variable among targeted taxa and influenced by a range of other factors. While our analyses revealed higher relative abundance and biomass of lethrinids (*Lethrinus nebulosus* and *L. atkinsoni*) inside reserves, no significant effect was found for the abundance of *Epinephelus rivulatus*, and a small negative effect was detected for the epinephelids as a group. All effects were heterogeneous, which was not surprising given the size and complexity of the synthesised dataset (including differences in size and age of reserves) and given that fish responses to reserves are known to vary with taxon-specific, ecological and zoning factors (Barrett *et al.*, 2007; Claudet *et al.*, 2010; Edgar *et al.*, 2014). Here we advance previous findings with the largest meta-analysis on recreational fishing in isolation from commercial fishing, illustrating the new information that can be gained from synthesising existing data, though we do not discount the advantages of strategic and consistent monitoring data. We show that it is important for assessments of reserves to take into account habitat effects, and potential interactions with factors such as reserve size or age, as well as variability in fishing activity, or differences in survey method in order to avoid oversimplified conclusions on how fish abundance and biomass respond to management.

Recreational fishing, specifically angling, is the only major fishing activity within the marine park. Some previous studies in the Park have linked higher abundance and biomass of targeted species inside reserves to protection from fishing (Westera, 2003; Babcock *et al.*, 2008; Fitzpatrick *et al.*, 2015); though results of other studies are more equivocal (Wilson *et al.*, 2012; Wilson *et al.*, 2018a). The reasons behind the disparate conclusions are unclear, but may be due to limited and/or varied spatial and temporal scales of the individual studies, different survey methodologies, the confounding influence of habitat, or high variability in target species abundance distributions. We also investigated the regulations on shore-based fishing on the coastal boundaries of reserves, with the hypothesis that this may influence the ability of the reserves to maintain higher abundance and biomass of fishes. There were mixed results with the full-subsets analysis indicating this variable had low importance, while the meta-analysis showed it did explain significant heterogeneity, and indicated effect sizes were larger (though not significantly) when shore fishing was prohibited. However, this factor was likely correlated with other variables not available in the present study, such as accessibility to reserves, which would influence our ability to test this hypothesis. High correlation between fish recruitment and larger natural cycles (El Niño Southern Oscillation) has also been suggested as a reason for inconsistencies in fishes response to reserves (Wilson *et al.*, 2018b). In the present study we

found high variability in the relative fish abundances of lethrinids among the different reserves, which can at least partly account for the varied conclusions of previous studies at smaller spatial scales (Fig. G2). Nonetheless, when all data were pooled the average effect was clearly positive for abundance and biomass of the three lethrinid groups. The magnitudes of the positive effects were small (max 57% higher inside) relative to studies in other parts of the world (Watson & Ormond, 1994; Russ *et al.*, 2015). A significant positive response for *L. atkinsoni* (40% higher), similar to that of *L. nebulosus* (42%) was not expected, given *L. atkinsoni* does not feature highly in catch reports (Ryan *et al.*, 2017), suggesting it may be more susceptible to recreational angling than previously recognised.

Known differences in behaviour between lethrinid and serranid taxa did not correlate with their response to reserves as expected. Lethrinids are known to have large home ranges relative to many epinephelids, including *E. rivulatus*, and are therefore more likely to move across reserve boundaries (Mackie & Black, 1999; Pillans *et al.*, 2014; Babcock *et al.*, 2017), with the expectation that they may experience lower levels of protection than epinephelids. However, we only observed positive responses for the lethrinids. It is possible that higher counts of lethrinids than epinephelids in the dataset may have reduced the power to detect an effect in the latter group, or there are other factors that have not been captured in our analyses.

The age of no-take reserves has been shown to be a significant positive correlate of relative fish abundance for targeted species (Claudet *et al.*, 2008; Edgar *et al.*, 2014; Zupan *et al.*, 2018) and demonstrated increases in effect size with time help attribute positive effect sizes to the presence of a protected area, rather than other factors (Russ *et al.*, 2015). In the present study there was negligible evidence of changes in effect sizes with age of reserve. Where relationships were present, the shape of the trend was generally parabolic, showing an increase initially, before subsequent decrease around 2005, though no relationships were significant. This was supported by examining data for *L. nebulosus*, from the best temporally replicated reserve, Osprey, where again no clear temporal patterns were found. Potentially of concern for managers was the significantly negative decline in *L. nebulosus* density outside of the Osprey reserve, and a slight increase followed by a decrease inside this reserve. However the confidence intervals on all temporal patterns were large. These findings are in contrast with previous studies, for example Russ *et al.* (2015) showed lethrinids continued to increase in density inside reserves in the Philippines on time scales of 8-30 years. In the present study rezoning in 2005 made temporal analyses more complex, though by including zoning scheme as a variable we partly addressed this. Effect sizes were not strongly influenced by this variable, implying that the effect sizes were broadly consistent across the initial and current reserve networks. Where zoning scheme did feature for *L. atkinsoni*, the older reserves had a more positive effect, as expected.

The absence of a strong temporal link with effect size must be considered when interpreting the positive effect sizes, however there are various factors which may have contributed to the absence of a strong relationship. First, while there is limited evidence of a reduction in fishing activity within the Park (Ryan *et al.*, 2015, 2017) a shift in fishing activity to areas offshore (>100m depth) (West *et al.*, 2015; Mitchell *et al.*, 2018), which are not part of the current survey data, is likely. Second, the mobile behaviour of lethrinid fishes may be capping the levels of the observed effect size, if a proportion of their population is travelling further than the reserve boundaries. Pillans *et al.* (2014) found that approximately 60% of lethrinid individuals move at scales greater than the average reserve size over a year period. Third, illegal fishing within the reserves may also limit a temporal increase in effect size, as Smallwood and Beckley (2012) found 8-12% of observed vessels were fishing inside reserves in the Park in 2007. Fourth, we do not discount that the unevenness of sampling through time, with some years being more highly sampled than others (Fig. B2) potentially influenced our capacity to detect a trend if it were present. The analysis of *L. nebulosus* density at Osprey showed that the temporal patterns inside and outside reserves can be complex and not always captured by the overall effect size. Parallel declines or increases in density occurring both inside and outside are masked from the effect size, and such declines have been observed in other fisheries closures on the western Australian coast (Bornt *et al.*, 2015).

Though our study only had a very coarse level of habitat classification available, our results support previous studies (Miller & Russ, 2014; Rees *et al.*, 2018a; Rees *et al.*, 2018b), showing the importance of habitat when assessing the ability of reserves to support target species abundance. We further demonstrate interactions between habitat and reserve size, showing that conclusions on both the magnitude and direction (positive or negative) of observed effects for the relative abundance of Lethrinidae and *L. nebulosus* are influenced by this interaction. In the case of *L. atkinsoni* biomass we also found an interaction between habitat and reserve age, though the models were not as strong. Previous studies have demonstrated the positive influence of larger and older reserves (Halpern & Warner, 2002; Claudet *et al.*, 2008; Edgar *et al.*, 2014; Zupan *et al.*, 2018), however the interaction with habitat has not previously been explored. Furthermore, it is noteworthy that effect sizes were greatest in the *back reef & lagoon coral* habitat for *L. nebulosus*, while for *L. atkinsoni*, the effect was greatest on the *reef flat*, a result that may be attributed to these habitats being preferred by the adults of each species respectively (Babcock *et al.*, 2008; Wilson *et al.*, 2017). This is important when considering potential changes to habitat inside or outside of reserves, as Russ *et al.* (2015), showed that changes in benthic habitat due to disturbance could markedly influence the effect of reserves for lethrinids. We advise that reserves must incorporate adequate amounts of the essential habitats of the species or communities they are designed to protect, and assessment of reserve effectiveness must account for possible interactions between habitat and reserve size and age.

While habitat was particularly important for the lethrinid groups, it was not found to be an important predictor for Epinephelinae or *E. rivulatus*. Again, this was contrary to expectations given the often high site fidelity of Epinephelinae (Mackie & Black, 1999). However, the relatively coarse habitat classification available for our analyses likely did not adequately capture the habitat requirements for this group. Previous work has shown *E. rivulatus* is strongly associated with macroalgal habitats at Ningaloo Reef (Wilson *et al.*, 2012) but that variability in the quality of macroalgal habitats can be substantial and have major implications for fish abundance (Fulton *et al.*, 2014; Wilson *et al.*, 2014; Lim *et al.*, 2016). Furthermore, Beckley and Lombard (2012) found that deeper habitats seaward of the reef have relatively lower spatial protection from recreational fishing, despite these habitats potentially supporting a high biomass of epinephelids (Babcock *et al.*, 2008). It is thus plausible that habitats outside of the reserves were more appropriate for Epinephelinae, particularly prior to re-zoning in 2005, which could explain the overall negative and null effects for these groups. A much better understanding of the habitat requirements, electivity and movement across seascapes by targeted taxa and appropriate ‘micro-habitat’ classifications are needed to more fully understand these results.

Where the boat fishing variable appeared in models for *E. rivulatus*, there were subtle positive trends in effect size as fishing activity increased, i.e. where boat fishing was most prevalent the effect size was greater. Our metric for fishing activity is unlikely to be representative across the 30 years of data, as it was an estimate from 2007 (Smallwood & Beckley, 2012), yet still showed some importance. We think this is a particularly important factor when assessing reserves, as variability in fishing activity (spatially and temporally) makes it very difficult to disentangle the true effect of the reserves if this variability is not quantified. We suggest that finer-scale spatiotemporal data on the pressures outside, and indeed inside, of reserves would clarify reserve assessments, both in the case of the present study but also more generally in any assessment of spatial protection. In the case of marine reserves, quantitative standardised data on fishing activity at the scale of individual reserves should be prioritised alongside the collection of ecological data.

Synthesizing data from multiple survey methods leads to larger datasets, and the advent of video-based methods in the last decades (e.g. BRUV and DOV) has increased the diversity of methods used to monitor fish. Contrary to expectations, in general, survey method did not strongly influence the effect size. The strongest effect sizes (Lethrinidae and *L. nebulosus*) were consistently detected regardless of the survey method. *L. atkinsoni* exhibited a more positive effect when surveyed by remote video as compared to diver-based methods, which may be partly explained by fish behaviours associated with both the attraction to bait and avoidance of divers (Watson *et al.*, 2005; Goetze *et al.*, 2015), particularly on SCUBA (Radford *et al.*, 2005). On balance, we did not distinguish a single survey method as optimal, and in most cases it was appropriate to compare data from the three methods for the effect size calculation. This is likely possible

because of the nature of our effect size, which, as a ratio, is more robust to different units of measurement. However, this cannot provide the same level of information as standardised temporal data on fish density inside to outside, as shown by density patterns inside and outside at Osprey, underlying the overall effect size for this reserve. We therefore suggest that monitoring programs should prioritise resurveying existing monitoring sites with comparable methodology to build more robust time-series data, else adopt the method(s) that are best suited to surveying the taxa of interest.

## 5. Conclusions

There were two major challenges in addressing the aims of this study. The first stemmed from the nature of the available data, as while we showed that new information can be gained from collaboration and the synthesis of disparate data, a lack of consistent temporal data meant it was not possible to understand the temporal changes to the fish populations. This was demonstrated by some complex trends in the estimated fish density inside and outside the Osprey reserve that underlay the overall effect size. Therefore, the value of consistent monitoring across time and space is unequivocal, particularly given a likely increase in adaptive management complicating temporal assessments. Indeed at Ningaloo, a new Australian Marine Park, in commonwealth waters directly seaward of the Ningaloo Marine Park of the present study has recently been implemented (1/1/18). Our findings suggest that consistent monitoring, producing data that can be compared to that of the present study should be implemented for this new Park. The second challenge was explored by Underwood (1995), who argued that ecological research can better aid management if management interventions are treated as testable hypotheses. No-take marine reserves can provide experiments with which to test hypotheses regarding the effects of fishing (Ballantine & Langlois, 2008). However, our study has highlighted that variability in ‘experimental design,’ resulting from a range of complexities including spatial and temporal variability in fishing activity, shore fishing zones adjacent to no-takes areas and modifications to reserve design over time, make determining the long-term outcomes of these experiments. We suggest that in order to best analyse across such complicated experimental designs it is necessary to account for (i) habitat; (ii) potential interactions between habitat and reserve size and age; and (iv) variability in fishing activity outside of reserves and compliance inside reserves. Regarding the last point, integration of the collection of fishing activity data with the collection of ecological data is likely to help interpret the true effects of reserves. The two are clearly intertwined and having data on both the pressure and the response is essential for holistic assessments of the efficacy of spatial management interventions.

## Acknowledgements

We thank the Department of Biodiversity, Conservation and Attractions and the Department of Primary Industries and Regional Development staff in Exmouth, in particular we are grateful for comments from Claire Smallwood with expertise on recreational fishing in the study region and Peter Barnes for expertise on the management regimes of the Ningaloo Marine Park. We would like to thank: Claire Butler for assistance and advice handling spatial data in QGIS; Emma Lawrence for discussion on the sensitivity analysis; Glenn Moore (WA Museum) for advising on the current taxonomy of Serranidae and Epinephelinae; George Cresswell, Susan Blackburn and Brett Molony for comments and editing. Field data are extremely time intensive to collect and we appreciatively acknowledge the many people who collected the data and provided logistical support for the numerous studies included in this data synthesis: **UWA**: Todd Bodd, Matt Birt, Brigit Vaughan, Isabella Lindgren; **CSIRO**: Geordie Clapin, Nicole Murphy, David Kozak, Julia Phillips, Ryan Downie, Fiona Graham, Kylie Cook, Catherine Seytre, Auriane Jones, Monique Grol, Andrea Zabala Peres, Helene Boulloche-Sabine, Lydiane Mattio, Cindy Bessey, Melanie Trapon, Doug Bearham, James McLaughlin, Ryan Crossing, Mark Wilson, Margaret Miller, Darren Dennis, David Milton, Rodrigo Bustamante, Tim Skewes; **RLS**; all volunteer citizen scientists involved in data collection, especially Paul Day, Kevin Smith and Ben Jones, as well as Antonia Cooper and Just Berkhout for database support; **DBCA**: Peter Barnes, Huw Dilley, David Lierich, Matt Smith, Teresa Edgecombe, Dani Robb, Jutta Wildforster, Joe Morgan, Shannon Armstrong, George Shedrawi; **ANU**: Mae Noble; **AIMS**: Conrad Speed, Ben Radford; **Sea Research**: Ian Parker, Gerry Allen, Avril Ayling

## Funding

The study was possible through funding provided by numerous organisations and programs: CSIRO, the Western Australian Marine Science Institution, WA Department of Biodiversity, Conservation and Attractions, The Australian National University, the Australian Institute of Marine Science, Caring for our Country, and the Gorgon Barrow Island Net Conservation Benefits Fund, BHP-CSIRO Industry-Science Ningaloo Outlook Marine Research Partnership, AIMS-Woodside Energy, WA State NRM program and Royalties for Regions program, Coastwest, the University of Western Australia, Edith Cowan University School of Natural Sciences, the Jean Rogerson Postgraduate Scholarship

## Appendices

**Appendix A** – Additional information on individual reserves and rezoning

**Appendix B** – Data summary

**Appendix C** – Geographic locations of individual surveys

**Appendix D** – Information on fish groups

**Appendix E** – Formulas used for calculating effect sizes

**Appendix F** – Sensitivity analysis for cases of one-armed zero events

**Appendix G** – Meta-analysis statistics

**Appendix H** – Biomass results

## References

- Abesamis, R.A., Green, A.L., Russ, G.R. & Jadloc, C.R.L. (2014) The intrinsic vulnerability to fishing of coral reef fishes and their differential recovery in fishery closures. *Reviews in Fish Biology and Fisheries*, **24**, 1033-1063.
- Anderson, D.R. & Burnham, K.P. (2002) Avoiding pitfalls when using information-theoretic methods. *The Journal of Wildlife Management*, 912-918.
- Babcock, R., Pillans, R. & Rochester, W. (2017) Environmental and individual effects on the behaviour and spawning movements of *Lethrinus nebulosus* on a coral reef. *Marine and Freshwater Research*, **68**, 1422-1437.
- Babcock, R., Phillips, J., Lourey, M. & Clapin, G. (2007) Increased density, biomass and egg production in an unfished population of Western Rock Lobster (*Panulirus cygnus*) at Rottnest Island, Western Australia. *Marine and Freshwater Research*, **58**, 286-292.
- Babcock, R., Haywood, M., Vanderklift, M., Clapin, G., Kleczkowski, M., Dennis, D., Skewes, T., Milton, D., Murphy, N. & Pillans, R. (2008) Ecosystem Impacts of Human Usage and the Effectiveness of Zoning for Biodiversity conservation: Broad-scale Fish Census. Final Analysis and Recommendations 2007. In: *Final Analysis and Recommendations*, p. 99. CSIRO Marine and Atmospheric Research, Hobart.
- Ballantine, B. (2014) Fifty years on: lessons from marine reserves in New Zealand and principles for a worldwide network. *Biological Conservation*, **176**, 297-307.
- Barlow, J., França, F., Gardner, T.A., Hicks, C.C., Lennox, G.D., Berenguer, E., Castello, L., Economo, E.P., Ferreira, J. & Guenard, B. (2018) The future of hyperdiverse tropical ecosystems. *Nature*, **559**, 517.
- Barrett, N.S., Edgar, G.J., Buxton, C.D. & Haddon, M. (2007) Changes in fish assemblages following 10 years of protection in Tasmanian marine protected areas. *Journal of Experimental Marine Biology and Ecology*, **345**, 141-157.
- Beckley, L.E. & Lombard, A.T. (2012) A systematic evaluation of the incremental protection of broad-scale habitats at Ningaloo Reef, Western Australia. *Marine and Freshwater Research*, **63**, 17-22.
- Bergseth, B.J., Williamson, D.H., Russ, G.R., Sutton, S.G. & Cinner, J.E. (2017) A social-ecological approach to assessing and managing poaching by recreational fishers. *Frontiers in Ecology and the Environment*, **15**, 67-73.
- Bornt, K.R., McLean, D.L., Langlois, T.J., Harvey, E.S., Bellchambers, L.M., Evans, S.N. & Newman, S.J. (2015) Targeted demersal fish species exhibit variable responses to long-term protection from fishing at the Houtman Abrolhos Islands. *Coral Reefs*, **34**, 1297-1312.

- CALM (1989) Ningaloo Marine Park Management Plan 1989 - 1999. In: *Department of Conservation and Land Management*, p. 108. Department of Conservation and Land Management, Perth, WA, Australia.
- CALM (2005) Management Plan for the Ningaloo Marine Park and Muiron Islands Marine Management Area 2005-2015: Management Plan No 52. In: *Department of Conservation and Land Management, Fremantle*, p. 115. Western Australia Department of Conservation and Land Management, Perth, WA, Australia.
- Claudet, J. & Guidetti, P. (2010) Improving assessments of marine protected areas. *Aquatic conservation: marine and freshwater ecosystems*, **20**, 239-242.
- Claudet, J., Osenberg, C.W., Benedetti- Cecchi, L., Domenici, P., García- Charton, J.A., Pérez- Ruzafa, Á., Badalamenti, F., Bayle- Sempere, J., Brito, A. & Bulleri, F. (2008) Marine reserves: size and age do matter. *Ecology letters*, **11**, 481-489.
- Claudet, J., Osenberg, C., Domenici, P., Badalamenti, F., Milazzo, M., Falcón, J.M., Bertocci, I., Benedetti- Cecchi, L., García-Charton, J.A. & Goñi, R. (2010) Marine reserves: fish life history and ecological traits matter. *Ecological applications*, **20**, 830-839.
- Coleman, F.C., Figueira, W.F., Ueland, J.S. & Crowder, L.B. (2004) The Impact of United States Recreational Fisheries on Marine Fish Populations. *Science*, **305**, 1958.
- Collins, L.B., Zhu, Z.R., Wyrwoll, K.-H. & Eisenhauer, A. (2003) Late Quaternary structure and development of the northern Ningaloo Reef, Australia. *Sedimentary Geology*, **159**, 81-94.
- Côté, I.M., Mosqueira, I. & Reynolds, J.D. (2005) Effects of marine reserve characteristics on the protection of fish populations: a meta- analysis. *Journal of Fish Biology*, **59**, 178-189.
- Cowx, I.G. & Cooke, S.J. (2004) The Role of Recreational Fishing in Global Fish Crises. *BioScience*, **54**, 857-859.
- Denny, C.M., Willis, T.J. & Babcock, R.C. (2004) Rapid recolonisation of snapper *Pagrus auratus*: Sparidae within an offshore island marine reserve after implementation of no-take status. *Marine Ecology Progress Series*, **272**, 183-190.
- DPIRD, D.o.P.I.a.R.D. (2018) *Ningaloo Marine Park*. Available at: <https://parks.dpaw.wa.gov.au/sites/default/files/downloads/parks/2462-13%20Spearfishing%20in%20Ningaloo%20WEB.pdf> (accessed 4/11/2018 2018).
- DPIRD, D.o.P.I.a.R.D.W.A. (2017) *Gascoyne Coast Bioregion, Status reports of the fisheries and aquatic resources of Western Australia 2016/17*. Available at: (accessed
- Edgar, G.J., Stuart-Smith, R.D., Willis, T.J., Kininmonth, S., Baker, S.C., Banks, S., Barrett, N.S., Becerro, M.A., Bernard, A.T. & Berkhout, J. (2014) Global conservation outcomes depend on marine protected areas with five key features. *Nature*, **506**, 216-220.
- Fisher, R., Wilson Shaun, K., Sin Tsai, M., Lee Ai, C. & Langlois Tim, J. (2018) A simple function for full- subsets multiple regression in ecology with R. *Ecology and Evolution*, **0**
- Fitzpatrick, B., Harvey, E., Langlois, T., Babcock, R. & Twiggs, E. (2015) Effects of fishing on fish assemblages at the reefscape scale. *Marine Ecology Progress Series*, **524**, 241-253.
- Froese, R.a.D.P. (2018) *FishBase*. Available at: [www.fishbase.org](http://www.fishbase.org) (accessed 15/1/2018 2018).
- Fulton, C.J., Depczynski, M., Holmes, T.H., Noble, M.M., Radford, B., Wernberg, T. & Wilson, S.K. (2014) Sea temperature shapes seasonal fluctuations in seaweed biomass within the Ningaloo coral reef ecosystem. *Limnology and Oceanography*, **59**, 156-166.
- Goetze, J., Jupiter, S., Langlois, T., Wilson, S., Harvey, E., Bond, T. & Naisilisili, W. (2015) Diver operated video most accurately detects the impacts of fishing within periodically harvested closures. *Journal of Experimental Marine Biology and Ecology*, **462**, 74-82.
- Halpern, B.S. & Warner, R.R. (2002) Marine reserves have rapid and lasting effects. *Ecology Letters*, **5**, 361-366.
- Horta e Costa, B., Claudet, J., Franco, G., Erzini, K., Caro, A. & Gonçalves, E.J. (2016) A regulation-based classification system for Marine Protected Areas (MPAs). *Marine Policy*, **72**, 192-198.
- Jenkins, C.N. & Joppa, L. (2009) Expansion of the global terrestrial protected area system. *Biological conservation*, **142**, 2166-2174.


- Langlois, T.J., Harvey, E.S., Fitzpatrick, B., Meeuwig, J.J., Shedrawi, G. & Watson, D.L. (2010) Cost-efficient sampling of fish assemblages: comparison of baited video stations and diver video transects. *Aquatic biology*, **9**, 155-168.
- Lester, S.E., Halpern, B.S., Grorud-Colvert, K., Lubchenco, J., Ruttenberg, B.I., Gaines, S.D., Aïramé, S. & Warner, R.R. (2009) Biological effects within no-take marine reserves: a global synthesis. *Marine Ecology Progress Series*, **384**, 33-46.
- Lewin, W.-C., Arlinghaus, R. & Mehner, T. (2006) Documented and Potential Biological Impacts of Recreational Fishing: Insights for Management and Conservation. *Reviews in Fisheries Science*, **14**, 305-367.
- Lim, I.E., Wilson, S.K., Holmes, T.H., Noble, M.M. & Fulton, C.J. (2016) Specialization within a shifting habitat mosaic underpins the seasonal abundance of a tropical fish. *Ecosphere*, **7**, e01212.
- Mackie, M. & Black, R. (1999) *Research on Two Serranid Species (Serranidae: Epinephelinae) in Western Australian Waters*. University of Western Australia/Fisheries Research & Development Corporation.
- Mallet, D. & Pelletier, D. (2014) Underwater video techniques for observing coastal marine biodiversity: a review of sixty years of publications (1952–2012). *Fisheries Research*, **154**, 44-62.
- McCook, L.J., Ayling, T., Cappo, M., Choat, J.H., Evans, R.D., De Freitas, D.M., Heupel, M., Hughes, T.P., Jones, G.P., Mapstone, B., Marsh, H., Mills, M., Molloy, F.J., Pitcher, C.R., Pressey, R.L., Russ, G.R., Sutton, S., Sweatman, H., Tobin, R., Wachenfeld, D.R. & Williamson, D.H. (2010) Adaptive management of the Great Barrier Reef: A globally significant demonstration of the benefits of networks of marine reserves. *Proceedings of the National Academy of Sciences*, **107**, 18278.
- McLaren, B.W., Langlois, T.J., Harvey, E.S., Shortland-Jones, H. & Stevens, R. (2015) A small no-take marine sanctuary provides consistent protection for small-bodied by-catch species, but not for large-bodied, high-risk species. *Journal of Experimental Marine Biology and Ecology*, **471**, 153-163.
- McPhee, D.P., Leadbitter, D. & Skilleter, G.A. (2002) Swallowing the bait: is recreational fishing in Australia ecologically sustainable? *Pacific Conservation Biology*, **8**, 40-51.
- Miller, K.I. & Russ, G.R. (2014) Studies of no-take marine reserves: Methods for differentiating reserve and habitat effects. *Ocean & Coastal Management*, **96**, 51-60.
- Mitchell, J., McLean, D., Collin, S., Taylor, S., Jackson, G., Fisher, R. & Langlois, T. (2018) Quantifying shark depredation in a recreational fishery in the Ningaloo Marine Park and Exmouth Gulf, Western Australia. *Marine Ecology Progress Series*, **587**, 141-157.
- Mora, C. & Sale, P.F. (2011) Ongoing global biodiversity loss and the need to move beyond protected areas: a review of the technical and practical shortcomings of protected areas on land and sea. *Marine ecology progress series*, **434**, 251-266.
- Mosquera, I., Côté, I.M., Jennings, S. & Reynolds, J.D. (2000) Conservation benefits of marine reserves for fish populations. *Animal Conservation forum* (ed by), pp. 321-332.
- Murphy, H.M. & Jenkins, G.P. (2010) Observational methods used in marine spatial monitoring of fishes and associated habitats: a review. *Marine and Freshwater Research*, **61**, 236-252.
- Osenberg, C.W., Shima, J.S., Miller, S.L. & Stier, A.C. (2011) Ecology: assessing effects of marine protected areas: confounding in space and possible solutions. *Marine protected areas: a multidisciplinary approach*, 143-167.
- Pillans, R.D., Bearham, D., Boomer, A., Downie, R., Patterson, T.A., Thomson, D.P. & Babcock, R.C. (2014) Multi year observations reveal variability in residence of a tropical demersal fish, *Lethrinus nebulosus*: implications for spatial management. *PLoS One*, **9**, e105507.
- R Core Team (2017) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Radford, C.A., Jeffs, A.G., Tindle, C.T., Cole, R.G. & Montgomery, J.C. (2005) Bubbled waters: The noise generated by underwater breathing apparatus. *Marine and freshwater behaviour and physiology*, **38**, 259-267.
- Rees, M.J., Knott, N.A. & Davis, A.R. (2018a) Habitat and seascape patterns drive spatial variability in temperate fish assemblages: implications for marine protected areas. *Marine Ecology Progress Series*, **607**, 171-186.

- Rees, M.J., Knott, N.A., Neilson, J., Linklater, M., Osterloh, I., Jordan, A. & Davis, A.R. (2018b) Accounting for habitat structural complexity improves the assessment of performance in no-take marine reserves. *Biological Conservation*, **224**, 100-110.
- Russ, G.R. (2002) Yet another review of marine reserves as reef fishery management tools. *Coral reef fishes: dynamics and diversity in a complex ecosystem*, **24**, 421.
- Russ, G.R., Miller, K.I., Rizzari, J.R. & Alcala, A.C. (2015) Long-term no-take marine reserve and benthic habitat effects on coral reef fishes. *Marine Ecology Progress Series*, **529**, 233-248.
- Ryan, K., Wise, B., Hall, N., Pollock, K., Sulin, E. & Gaughan, D.J. (2013) *An integrated system to survey boat-based recreational fishing in Western Australia 2011/12*. Fisheries Research Division, Western Australian Fisheries and Marine Research Laboratories.
- Ryan, K., Hall, N., Lai, E., Smallwood, C., Taylor, S. & Wise, B. (2015) *State-wide survey of boat-based recreational fishing in Western Australia 2013/14*. Fisheries Research Division.
- Ryan, K., Hall, N., Lai, E., Smallwood, C., Taylor, S. & Wise, B. (2017) State-wide survey of boat-based recreational fishing in Western Australia 2015/16. In. Department of Primary Industries and Regional Development
- Sala, E., Lubchenco, J., Grorud-Colvert, K., Novelli, C., Roberts, C. & Sumaila, U.R. (2018) Assessing real progress towards effective ocean protection. *Marine Policy*, **91**, 11-13.
- Shears, N.T., Grace, R.V., Usmar, N.R., Kerr, V. & Babcock, R.C. (2006) Long-term trends in lobster populations in a partially protected vs. no-take Marine Park. *Biological conservation*, **132**, 222-231.
- Smallwood, C.B. & Beckley, L.E. (2012) Spatial distribution and zoning compliance of recreational fishing in Ningaloo Marine Park, north-western Australia. *Fisheries Research*, **125**, 40-50.
- Sumner, N.R., Williamson, P.C. & Malseed, B.E. (2002) *A 12-month survey of recreational fishing in the Gascoyne bioregion of Western Australia during 1998-99*. Department of Fisheries, Western Australia.
- Underwood, A. (1993) The mechanics of spatially replicated sampling programmes to detect environmental impacts in a variable world. *Australian Journal of ecology*, **18**, 99-116.
- Viechtbauer, W. (2010) Conducting meta-analyses in R with the metafor package. *J Stat Softw*, **36**, 1-48.
- Watson, D.L., Harvey, E.S., Anderson, M.J. & Kendrick, G.A. (2005) A comparison of temperate reef fish assemblages recorded by three underwater stereo-video techniques. *Marine Biology*, **148**, 415-425.
- Watson, M. & Ormond, R. (1994) Effect of an artisanal fishery on the fish and urchin populations of a Kenyan coral reef. *Marine Ecology Progress Series*, **115**, 115-129.
- West, L., Stark, K., Murphy, J., Lyle, J. & Ochwada-Doyle, F. (2015) Survey of recreational fishing in New South Wales and the ACT, 2013/14.
- Westera, M.B. (2003) The effect of recreational fishing on targeted fishes and trophic structure, in a coral reef marine park.
- Wilson, S., Fulton, C., Depczynski, M., Holmes, T., Noble, M., Radford, B. & Tinkler, P. (2014) Seasonal changes in habitat structure underpin shifts in macroalgae-associated tropical fish communities. *Marine biology*, **161**, 2597-2607.
- Wilson, S.K., Graham, N.A.J., Holmes, T., MacNeil, M.A. & Ryan, N. (2018a) Visual versus video methods for estimating reef fish biomass. *Ecological Indicators*, **85**, 146-152.
- Wilson, S.K., Babcock, R.C., Fisher, R., Holmes, T.H., Moore, J.A.Y. & Thomson, D.P. (2012) Relative and combined effects of habitat and fishing on reef fish communities across a limited fishing gradient at Ningaloo. *Marine Environmental Research*, **81**, 1-11.
- Wilson, S.K., Depczynski, M., Holmes, T.H., Noble, M.M., Radford, B.T., Tinkler, P. & Fulton, C.J. (2017) Climatic conditions and nursery habitat quality provide indicators of reef fish recruitment strength. *Limnology and Oceanography*, **62**, 1868-1880.
- Wilson, S.K., Depczynski, M., Fisher, R., Holmes, T.H., Noble, M.M., Radford, B.T., Rule, M., Shedrawi, G., Tinkler, P. & Fulton, C.J. (2018b) Climatic forcing and larval dispersal capabilities shape the replenishment of fishes and their habitat-forming biota on a tropical coral reef. *Ecology and Evolution*, **8**, 1918-1928.
- Wood, S.N. (2011) Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **73**, 3-36.

- 731 Zupan, M., Fragkopoulou, E., Claudet, J., Erzini, K., Horta e Costa, B. & Gonçalves, E.J. (2018) Marine  
732 partially protected areas: drivers of ecological effectiveness. *Frontiers in Ecology and the*  
733 *Environment*, **16**, 381-387.
- 734 Zuur, A.F., Ieno, E.N. & Elphick, C.S. (2010) A protocol for data exploration to avoid common statistical  
735 problems. *Methods in ecology and evolution*, **1**, 3-14.
- 736
- 737


738  
739

Figure and Tables



740  
741  
742  
743  
744  
745  
746  
747  
748

**Figure 1** The Ningaloo Marine Park (NMP) and Muiron Islands Marine Management Area boundaries (dotted lines) with the location of sanctuary zones (referred to as reserves in the present study) shown in green along the Ningaloo coast of Western Australia under the a) initial (1987 – 2005) and b) current (2005 - 2017) zoning schemes. Tantabiddi Well and Winderabandi Point are indicated with red markers spearfishing is prohibited between these locations. The Osprey reserve is also indicated. In b) blue regions indicate zones on the coastal boundaries of the reserves where shore-based fishing is allowed.



**Figure 2.** a) Relative fish abundance inside to outside the reserves (back-transformed weighted mean effect sizes) with 95% confidence intervals), for the six fish groups: Lethrinidae, *Lethrinus nebulosus*, *L. atkinsoni*, Epinephelinae, *Epinephelus rivulatus* and Scarinae. Effect sizes are significant when the confidence intervals do not overlap 1.0. Open dots correspond to non-significant effects (i.e. no effect). Sample sizes are given in Table F1. Triangular points show the predicted effect size when *habitat* was included as a moderator variable in the meta-analysis, for the habitat with the largest mean effect (orange represents the back reef & lagoon coral, and blue represents the reef flat). b) Importance scores (based on summed Akaike weights corrected for finite samples (AICc)) from full-subsets analyses exploring the influence of seven variables on the overall effect size for each fish taxa: 1 is highly important while 0 is not important. Red X symbols mark the variables that were included in the most parsimonious models for each fish taxa (also see Table 2 and Fig. 3).














Figure 3. Predicted relative fish abundance inside to outside reserves (back-transformed predicted weighted effect sizes) with 95% confidence intervals) for the six fish groups – a) Lethrinidae; b) *Lethrinus nebulosus* c) *Lethrinus atkinsoni*; d) Epinephelinae; e) *Epinephelus rivulatus*; f) Scarinae for abundance– as a function of variables present in the most parsimonious models (Table 2) from full-subsets GAMM analysis. Ribbons represent 95% confidence intervals




Figure 4. Effect sizes for a) abundance and b) biomass from comparison pairs for the Osprey reserve through time and estimated density of c) abundance and d) biomass inside and outside the reserve through time. Ribbons indicate 95% confidence intervals on generalised additive models.

**Table 1.** Description and summary of the seven variables used in analysis

Variable (transformation used in analyses)	Description	Description of variable levels		Source
<b>Years protection</b>	Years between zoning and survey data collection	A survey is classified to a single reserve based on its location. In cases where rezoning means that a reserves size was increased, a survey falling inside the old area is classified as the pre-zoning reserve, while if the survey falls in the extended area it is classified as the post-zoning reserve. Years since protection is calculated on the same principal relating to the initial zoning or the rezoning dependent on survey location. 0 - 30 years from time of survey to reserve implementation.		(CALM 2005) (CALM 1989)
<b>Zoning scheme</b>	Factor describing the two major spatial zonings implemented in the Ningaloo Marine Park	<i>Initial</i>	1987-2005, 8 no-take zones, see Fig. 1a	(CALM 2005) (CALM 1989) (Appendix A)
		<i>Current</i>	2005-present, 18 no-take (excepting shore fishing) and 3 no-take zones in the Muiron Islands Marine Management Area, see Fig. 1b	
<b>Survey method</b>	Factor describing major survey methods used to collect the fish count and size data	<i>UVC</i>	Underwater visual census, collected along transect lines of set length and width (25 x 5 m, 50 x 5 m or 100 x 10 m, 250 x 10 m). Most on SCUBA, some via snorkel. Fish counted and length estimated <i>in situ</i> .	For more information on methods see Appendix B
		<i>BRUV</i>	Baited remote underwater stereo-video deployments, (30-60 minutes) point location, fish counted and length estimated <i>post hoc</i> from video	
		<i>DOV</i>	Diver operated stereo-video, collected along a transect line of set length and width (5 m in width and varying between 25 and 50 m length), fish counted and length estimated <i>post hoc</i> from video	
<b>Habitat</b>	Factor describing four major habitat types which have differences both in the dominant benthic community and wave exposure	<i>Exposed reef slope</i>	The ocean side of the fringing reef, where the reef slopes to deeper water and the majority of wave energy is received	Classified by authors, (see Collins et al. 2003)
		<i>Reef flat</i>	Shallow (~2-3m deep), shoreward from the reef crest for tens to hundreds of meters, typically dominated by the plate coral <i>Acropora spicifera</i> on limestone bedrock	
		<i>Back reef &amp; lagoon coral</i>	From where the reef flat breaks into more patchy reef and sand environments, sheltered from wave energy and including some large coral bommies	
		<i>Lagoon algae</i>	Sheltered shallow water lagoon, dominated by fleshy canopy forming seaweed of the genera <i>Sargassum</i> and <i>Sargassopsis</i> .	
<b>Reserve size (square –root)</b>	Area (ha) of each no-take reserve at time of survey	50 – 44752 hectares Mean: 6031 ha; Median: 1756 ha		(CALM 2005) (CALM 1989)
<b>Boat fishing *</b> (log-transformation)	A mean estimate of the number of vessels recreationally fishing at the outside reserve survey sites	Mean density of vessels observed fishing during aerial surveys in peak season in 2007. Each survey sample latitude and longitude was assigned the value of the underlying spatial data in Fig. 4 of Smallwood and Beckley (2012). For surveys inside the reserves it was assumed that fishing activity was 0. 0 - 0.625 vessels fishing per 9 km <sup>2</sup> Mean: 0.12; Median: 0.11 * Not available for the Muiron Islands Other estimates of fishing activity exist (Sumner et al. 2002) but this metric was deemed the most detailed		Smallwood and Beckley (2012)
<b>Shore fishing</b>	Factor describing whether or not a reserve has shore fishing zones on its coastal boundary	<i>Allowed</i>	Shore fishing is allowed along the entire, or part of the coastal boundary of the reserve (26% of data)	(CALM 2005)
		<i>Prohibited</i>	No shore fishing is permitted anywhere in the reserve (74% of data)	

**Table 2.** Top Generalised Additive Mixed Models (GAMMs) for predicting the response ratio inside to outside reserves,  $\bar{E}$ , for abundance from full subset analyses for the abundance of the six fish groups. Difference between the lowest reported corrected Akaike Information Criterion ( $\Delta\text{AICc}$ ), AICc weights ( $\omega\text{AICc}$ ), variance explained ( $R^2$ ) and estimated degrees of freedom (EDF) are reported for model comparison. Model selection was based on the most parsimonious model (fewest variables and lowest EDF) within two units of the lowest AICc. This model is shown in bold text.

Fish group	Model	$\Delta\text{AICc}$	$\omega\text{AICc}$	$R^2$	EDF
LETHRINIDAE	Habitat + Years protection by Habitat + Size by Habitat	0.00	0.31	0.14	14.6
	Years protection + Habitat + Size by Habitat	0.17	0.28	0.13	10.7
	<b>Habitat + Size by Habitat</b>	0.19	0.28	0.12	9.0
<i>L. nebulosus</i>	<b>Habitat + Size by Habitat</b>	0.00	0.57	0.17	9.0
<i>L. atkinsoni</i>	<b>Method + Zoning scheme</b>	0.00	0.19	0.08	6.3
	Habitat + Method + Zoning scheme	0.09	0.18	0.09	9.0
	boat.log + Method + Zoning scheme	0.71	0.14	0.08	7.4
	Habitat + Method + Size	1.41	0.10	0.08	8.0
	Habitat + Size + Years protection by Habitat	0.00	0.18	0.11	14.4
	Years protection + Boat fishing + Size	0.92	0.11	0.08	8.3
EPINEPHELINAE	<b>Years protection + Size</b>	1.51	0.08	0.08	7.3
<i>E. rivulatus</i>	<b>Boat fishing + Zoning scheme</b>	0.00	0.60	0.17	8.9
SCARINAE	<b>Boat fishing + Shore fishing</b>	0.00	0.16	0.03	4.0
	Boat fishing + Zoning scheme	0.80	0.11	0.03	4.2
	Years protection + Habitat + Size by Habitat	1.25	0.08	0.05	10.5
	Habitat + Size by Habitat	1.84	0.06	0.04	9.0

827 **Supplementary Information**


828 **Appendix A – Additional information on individual reserves and rezoning**

829 Maps of individual reserves can be found in CALM (1989), pgs. 55, for the 1987-2005 zoning and (CALM  
830 2005), pgs. 89-96, for the 2005-current zoning. Table A1 details individual reserves, their year of  
831 establishment, size, and the regulations on shore-based fishing on the coastal edges of the reserves. As well,  
832 the number of inside outside comparisons and the total number of surveys are given for each reserve.

833  
834 **Table A1.** Features of historical and current reserves in the Ningaloo Marine Park

	Reserve	Management area	Year of Establishment	Size (ha)	Shore fishing	Total number comparison pairs	Total number of samples
2005 – PRESENT MANAGEMENT (CURRENT)	3 Mile	Ningaloo Marine Park	2005	395	N	0	0
	Bateman	Ningaloo Marine Park	2005	1111	N	1	12
	Bundegi rezoned	Ningaloo Marine Park	1987/2005	696	N	3	24
	Cape Farquhar	Ningaloo Marine Park	2005	5326	N	3	31
	Cloates/ Dugong rezoned	Ningaloo Marine Park	1987/2005	44752	P	17	268
	Gnaraloo Bay	Ningaloo Marine Park	2005	1021	N	2	39
	Jurabi	Ningaloo Marine Park	2005	754	Y	9	78
	Lakeside	Ningaloo Marine Park	2005	8	N	0	0
	Lighthouse Bay	Ningaloo Marine Park	2005	763	P	9	136
	Mandu rezoned	Ningaloo Marine Park	1987/2004	1349	N	12	110
	Mangrove rezoned	Ningaloo Marine Park	1987/2005	1135	N	27	429
	Maud rezoned	Ningaloo Marine Park	1987/2005	2151	P	10	133
	Murat	Ningaloo Marine Park	2005	490	Y	0	0
	North Muiron	Muiron Islands Marine Management Size	2005	828	N	4	66
	Osprey rezoned	Ningaloo Marine Park	1987/2005	9513	P	7	54
	Pelican rezoned	Ningaloo Marine Park	1987/2006	10864	P	28	454
	South Muiron	Muiron Islands Marine Management Size	2005	784	N	4	54
	Sunday Island	Muiron Islands Marine Management Size	2005	317	N	0	0
	Tantabiddi	Ningaloo Marine Park	2005	50	N	8	98
	Turtles	Ningaloo Marine Park	2005	2461	N	0	0
	Winderabandi	Ningaloo Marine Park	2005	5526	Y	7	52
1987 – 2005 MANAGEMENT (INITIAL)	Bundegi	Ningaloo Marine Park	1987	297	N	20	333
	Cloates	Ningaloo Marine Park	1987	6257	N	16	186
	Dugong	Ningaloo Marine Park	1987	8852	N	0	0
	Mandu	Ningaloo Marine Park	1987	1163	N	50	1001
	Mangrove	Ningaloo Marine Park	1987	403	N	2	14
	Maud	Ningaloo Marine Park	1987	1806	N	36	652
	Osprey	Ningaloo Marine Park	1987	1756	N	44	701
	Pelican	Ningaloo Marine Park	1987	908	N	7	66
	3 Mile	Ningaloo Marine Park	2005	395	N	0	0

835  
836 The rezoning of the eight reserves in 2005, with the addition of new reserves and the expansion of the  
837 existing reserves required that data be clearly classified to account for this. Figure A1 shows an example of  
838 how samples were referenced based on their spatial location to the initial or current reserves and the time  
839 period of sampling.



**Figure A1.** Example classification of ‘initial’ and ‘current’ zoning. The boundaries of the initial 1987 – 2005 reserves, Cloates and Dugong, are indicated by a green outline, while the current reserve (2005 – 2017) is shown in black which combined the two initial reserves into one larger reserve, ‘Cloates/Dugong rezoned’ (also see Table 1). Pre 2005, surveys were classified as inside/outside based on their spatial relation to initial reserves (green outline). Post 2005, surveys were classified as inside/outside based on their spatial relation to the current reserves (black). If an inside reserve is within the initial boundaries, but surveyed post 2005, it was referenced as initial as shown in c). If a sample was collected at a site within the new area of the rezoned reserve post the rezoning, it was referenced as current as seen in d).

## Appendix B– Data summary

Suitable data were identified through searches on Google Scholar and research databases compiled by the Department of Biodiversity, Conservation and Attractions. In addition, relevant researchers from universities, research institutions, industry and citizen science programs operating in Western Australia were contacted to source unpublished data. Nine major custodians contributed data to this study. Table B1 summarises the survey methods used by each custodian, the number of samples collected, the temporal span of data, the reserves surveyed and directs to further reading for more information. Figure B2 graphically illustrates the important data constraints.

**Table B1.** Summary of data contributions

Data custodian	Total number of surveys	Reserves surveyed	Maximum temporal span of surveys	Main survey methods	More information available:
Australian Institute of Marine Science – Woodside Energy (AIMS-Woodside)	183	Bundegi, Bundegi rezoned, Cloates, Cloates/Dugong rezoned, Mandu, Mangrove rezoned, Maud, North Muiron, Osprey, Pelican rezoned, Tantabiddi	1993 -2014	Underwater Visual Census : 50 x 5 m	(Depczynski et al. 2015)
Ben Fitzpatrick	345	Mandu, Osprey	2006 - 2007	Baited Remote Underwater stereo-Video	(Fitzpatrick et al. 2015)
Commonwealth Industrial and Scientific Research Organisation (CSIRO)	18	Bundegi, Bundegi rezoned, Cape Farquhar, Cloates, Cloates/Dugong rezoned, Gnarlaloo Bay, Jurabi, Lighthouse Bay, Mandu, Mandu rezoned, Mangrove, Mangrove rezoned, Maud, North Muiron, Osprey, Osprey rezoned, Pelican, Pelican rezoned, South Muiron	2006 - 2017	Underwater Visual Census, 4 main sizes: 1) Either a singular, or three, 25 × 5 m transects per site. 2) 30 × 5 m transects 3) 50 × 5 m transects 4) 100 × 10 m transects at a site	(Babcock et al. 2008)
Western Australian Department of Biodiversity, Conservation and Attractions (DBCA)	1237	Bundegi , Cape Farquhar, Cloates, Cloates/Dugong rezoned, Jurabi, Lighthouse Bay, Mandu, Mangrove rezoned, Maud, Maud rezoned, North Muiron, Osprey, Pelican, Pelican rezoned, Tantabiddi, Winderabandi	2010 - 2016	Diver Operated stereo-Video: six replicate 50 x 5 m belt transects per site; or nine replicate 30 x 5 m belt transects per site	(Wilson et al. 2012; Holmes et al. 2013; Wilson et al. 2018b)
Mark Westera	257	Mandu, Maud, Osprey	1999 - 2000	Baited Remote Underwater stereo-Video, 30 minute deployments at 12 replicate locations in each zone, Underwater visual census using snorkel, 250 x 10 m transects	(Westera 2003a, b)
Reef Life Survey (RLS)	291	Bateman, Bundegi, Cloates, Maud, Maud rezoned, Pelican rezoned	2010 - 2017	Underwater Visual Census, 50 x 5 m transects	(see <a href="http://reeflifesurvey.com/files/2008/09/rils-reef-monitoring-procedures.pdf">http://reeflifesurvey.com/files/2008/09/rils-reef-monitoring-procedures.pdf</a> ).
Tony Ayling	60	Osprey	1987		(Ayling & Ayling 1987)
The University of Western Australia	325	Cloates, Cloates/Dugong rezoned, Mandu rezoned, Mangrove rezoned, Osprey, Osprey rezoned, Pelican rezoned, Winderabandi	2014 - 2015	Baited Remote Underwater stereo-Video, generally 60 minute deployments	(McLean et al. 2016)
Joint data custodians WA Department of Biodiversity, Conservation and Attractions, Australian Institute of Marine Science, Australian National University	587	Cloates/Dugong rezoned, Mandu, Mangrove rezoned, Maud, Maud rezoned, Pelican, Pelican rezoned,	2013 - 2015	Underwater Visual Census, 4 main sizes: 1) Either a singular, or three, 25 × 5 m transects per site. 2) 30 × 5 m transects 3) 50 × 5 m transects 4) 100 × 10 m transects at a site	(Fulton et al. 2014)

In most surveys biomass had been calculated by the respective data custodians from the estimated or measured fish length. In these cases the provided values were used (with the reasoning that different survey methods may warrant slightly different biomass calculations). In cases where length data was available but biomass had not been calculated the fish counts and length estimates were converted to biomass (kg) using constants and formulas from Fishbase ([www.fishbase.org](http://www.fishbase.org)).


$$Biomass = e^{\ln(a)+b \times \ln(L)}$$

where  $L$  is the estimated total length of the fish and  $a$  and  $b$  are constants for the species or family in question. In cases where data provided fork length measurements, these were converted to total length using the formula from FishBase:

$$TL = c + d \times FL$$

where  $TL$  is estimated total fish length,  $FL$  is the measured fork length from BRUV or DOV video and  $c$  and  $d$  are parameters specific to the fish species in question.

More data was available for some NTMRs and years, with increased sampling after 2005 (Appendix B).



**Figure B2.** Temporal and spatial distribution of samples: a) number of samples by year, with shade indicating the type of survey method (BRUV, Baited Remote Underwater stereo-Video, DOV, Diver Operated stereo-Video and UVC, Underwater Visual Census); b) distribution of samples inside and outside reserves for each of the four major habitats; c) distribution of samples inside and outside for each reserve under both the initial and current zoning.

## References

- Ayling T, Ayling AL 1987. Ningaloo Marine Park: preliminary fish density assessment and habitat survey: with information on coral damage due to *Drupella cornus* grazing: a report prepared for the Department of Conservation and Land Management, Western Australia. Department of Conservation and Land Management.
- Babcock R, Haywood M, Vanderklift M, Clapin G, Kleczkowski M, Dennis D, Skewes T, Milton D, Murphy N, Pillans R. 2008. Ecosystem Impacts of Human Usage and the Effectiveness of Zoning for Biodiversity conservation: Broad-scale Fish Census. Final Analysis and Recommendations 2007., Hobart.
- Babcock RC, Shears NT, Alcala AC, Barrett NS, Edgar GJ, Lafferty KD, McClanahan TR, Russ GR. 2010. Decadal trends in marine reserves reveal differential rates of change in direct and indirect effects. *Proceedings of the National Academy of Sciences* **107**:18256.
- Depczynski MM, Tinkler PP, Cheal AA, Speed CC. 2015. Ningaloo fish communities. Page 121 in Miller KK, Depczynski MM, Cappo MM, Wakeford MM, Speed CC, Stowar MM, Colquhoun JJ, Tinkler PP, Cheal AA, and Fisher RR, editors. Ningaloo and Outer Shark Bay Environmental Baseline Survey 2014. Report prepared by the Australian Institute of Marine Science for Woodside Energy Ltd. Australian Institute of Marine Science, Townsville, Australia.
- Fitzpatrick B, Harvey E, Langlois T, Babcock R, Twiggs E. 2015. Effects of fishing on fish assemblages at the reefscape scale. *Marine Ecology Progress Series* **524**:241-253.
- Holmes TH, Wilson SK, Travers MJ, Langlois TJ, Evans RD, Moore GI, Douglas RA, Shedrawi G, Harvey ES, Hickey K. 2013. A comparison of visual-and stereo-video based fish community assessment methods in tropical and temperate marine waters of Western Australia. *Limnology and Oceanography, Methods* **11**:337-350.
- Westera MB. 2003a. The effect of recreational fishing on targeted fishes and trophic structure, in a coral reef marine park.
- Westera MB. 2003b. The effect of recreational fishing on targeted fishes and trophic structure, in a coral reef marine park. Edith Cowan University, Perth, Western Australia.

918    ***Appendix C – Geographic locations of individual samples***  
919    Kml file for viewing samples in GoogleEarth  
920  
921

## Appendix D: Fish groups

923

924 Analyses were conducted at the family/subfamily level and species level. Investigation of patterns at the  
925 family/subfamily level allowed the inclusion of data from targeted but rare species.

926

927 Table D1. Total count of fish groups across all data

Taxa	Total count across synthesised data
LETHRINIDAE	10307
<i>L. nebulosus</i>	4183
<i>L. atkinsoni</i>	4765
EPINEPHELINAE	4012
<i>E. rivulatus</i>	1119
SCARINAE	44931

928

929

930

931 Genera included in the family/ sub-family analysis that were sampled in the synthesised data are  
932 summarised below.

933

### 934 Lethrinidae

935 • <i>Gnathodentex aureolineatus</i>	946 • <i>Lethrinus microdon</i>
936 • <i>Gymnocranius euanus</i>	947 • <i>Lethrinus miniatus</i>
937 • <i>Gymnocranius grandoculis</i>	948 • <i>Lethrinus nebulosus</i>
938 • <i>Gymnocranius griseus</i>	949 • <i>Lethrinus obsoletus</i>
939 • <i>Gymnocranius spp.</i>	950 • <i>Lethrinus olivaceus</i>
940 • <i>Lethrinus amboinensis</i>	951 • <i>Lethrinus ravus</i>
941 • <i>Lethrinus atkinsoni</i>	952 • <i>Lethrinus rubrioperculatus</i>
942 • <i>Lethrinus genivittatus</i>	953 • <i>Lethrinus semicinctus</i>
943 • <i>Lethrinus harak</i>	954 • <i>Lethrinus variegatus</i>
944 • <i>Lethrinus laticaudis</i>	955 • <i>Lethrinus spp.</i>
945 • <i>Lethrinus lentjan</i>	956 • <i>Monotaxis grandoculis</i>

957

### 958 Epinephelinae \*

959 • <i>Aethaloperca rogae</i>	968 • <i>Cephalopholis sonnerati</i>
960 • <i>Anyperodon leucogrammicus</i>	969 • <i>Cromileptes altivelis</i>
961 • <i>Cephalopholis spp.</i>	970 • <i>Epinephelus spp.</i>
962 • <i>Cephalopholis argus</i>	971 • <i>Epinephelus areolatus</i>
963 • <i>Cephalopholis boenak</i>	972 • <i>Epinephelus bilobatus</i>
964 • <i>Cephalopholis cyanostigma</i>	973 • <i>Epinephelus coeruleopunctatus</i>
965 • <i>Cephalopholis formosa</i>	974 • <i>Epinephelus coioides</i>
966 • <i>Cephalopholis miniata</i>	975 • <i>Epinephelus corallicola</i>
967 • <i>Cephalopholis sexmaculata</i>	976 • <i>Epinephelus fasciatus</i>

977	•	<i>Epinephelus fuscoguttatus</i>	986	•	<i>Epinephelus quoyanus</i>
978	•	<i>Epinephelus hexagonatus</i>	987	•	<i>Epinephelus rivulatus</i>
979	•	<i>Epinephelus lanceolatus</i>	988	•	<i>Epinephelus tauvina</i>
980	•	<i>Epinephelus macrospilos</i>	989	•	<i>Epinephelus tukula</i>
981	•	<i>Epinephelus maculatus</i>	990	•	<i>Plectropomus leopardus</i>
982	•	<i>Epinephelus malabaricus</i>	991	•	<i>Plectropomus maculatus</i>
983	•	<i>Epinephelus melanostigma</i>	992	•	<i>Plectropomus</i> spp.
984	•	<i>Epinephelus merra</i>	993	•	<i>Variola albimarginata</i>
985	•	<i>Epinephelus polyphekadion</i>	994	•	<i>Variola louti</i>
995	•	996			
997					

998 \* Note that there is current discussion of whether Epinephelinae may be better classified as its own family,  
999 Epinephelidae (Ma & Craig 2018), however we have chosen to name it as its subfamily here.

000

001

## 002 **Scarinae**

1003	•	<i>Calotomus carolinus</i>
1004	•	<i>Calotomus spinidens</i>
1005	•	<i>Cetoscarus ocellatus</i>
1006	•	<i>Chlorurus bleekeri</i>
1007	•	<i>Chlorurus microrhinos</i>
1008	•	<i>Chlorurus sordidus</i>
1009	•	<i>Hipposcarus longiceps</i>
1010	•	<i>Leptoscarus vaigiensis</i>
1011	•	<i>Scarus chameleon</i>
1012	•	<i>Scarus dimidiatus</i>
1013	•	<i>Scarus flavipectoralis</i>
1014	•	<i>Scarus frenatus</i>
1015	•	<i>Scarus ghobban</i>
1016	•	<i>Scarus globiceps</i>
1017	•	<i>Scarus niger</i>
1018	•	<i>Scarus oviceps</i>
1019	•	<i>Scarus prasiognathos</i>
1020	•	<i>Scarus psittacus</i>
1021	•	<i>Scarus rivulatus</i>
1022	•	<i>Scarus rubroviolaceus</i>
1023	•	<i>Scarus schlegeli</i>
1024	•	<i>Scarus</i> spp.

1025

1026   **References**

1027   Ma KY, Craig MT. 2018. An inconvenient monophyly: an update on the taxonomy of the  
1028                   groupers (Epinephelidae). *Copeia* **106**:443-456.

1029

1030

## **Appendix E: Formulas used for calculating effect sizes**

For each inside/ outside comparison pair a log response ratio ( $e$ ) (Hedges et al. 1999) was calculated as the ratio of the mean abundance/ biomass inside to outside a reserve for comparison pair  $i$  and fish group  $j$  as

$$e_{ij} = \ln \left( \frac{\bar{X}_{ij,I}}{\bar{X}_{ij,O}} \right)$$

where  $\bar{X}_{ij,I}$  and  $\bar{X}_{ij,O}$  are the mean abundance or biomass inside (I) and outside (O) a reserve. Therefore, a positive  $e_{ij}$  implies a greater fish abundance/ biomass inside the reserves than outside. A log response ratio was appropriate because it is independent of the actual unit of measurement across the different survey methods.

The variance of  $e_{ij}$  was also quantified (i.e. the within-study variance, where a study is a comparison pair), given that sampling error plays an important role in introducing variability in the overall outcome of a meta-analysis:

$$v_{ij} = \frac{\sigma_{ij,I}^2}{(n_{ij,I} \bar{X}_{ij,I}^2)} + \frac{\sigma_{ij,O}^2}{(n_{ij,O} \bar{X}_{ij,O}^2)}$$

Here,  $v_{ij}$  is calculated from the standard deviation,  $\sigma_{ij}$ , sample size,  $n_{ij}$ , and mean  $\bar{X}_{ij}$  for inside (I) and outside (O) for comparison pair  $i$  and fish group  $j$ .

Effect sizes were weighted by the inverse of the sum of the within- and among-study variances,  $w_{ij}$ . The within-study variance (where a study is a comparison pair) was the sum of the variances associated with each mean in the log-ratio. The among-study variance was calculated using the *metafor* package (Viechtbauer 2010) in the statistical program R (R Core Team 2017).

$$w_{ij} = \frac{1}{v_{ij}}$$

This weighting minimized the influence of studies with low statistical power, and increased the influence of studies with high statistical power, meaning that each inside/ outside replicate did not contribute equally to the final pooled outcome. Weighted meta-analyses of this sort are considered to increase the precision and power of meta-analyses (Osenberg et al.

1999) and was an appropriate approach in the present study as there was a large distribution of sample sizes and variance associated with the inside/ outside comparison pairs.

The weighted cumulative effect size for fish group  $j$ ,  $\bar{E}_j$ , was obtained as

$$\bar{E}_j = \frac{\sum_{i=1}^{n_{ij}} w_{ij} e_{ij}}{\sum_{i=1}^{n_{ij}} w_{ij}} \text{ with associated variance, } \bar{v}_j = \frac{1}{\sum_{i=1}^{n_{ij}} w_{ij}}.$$

where  $w_{ij}$  and  $e_{ij}$  are defined above. The overall heterogeneity ( $Q_t$ ) for fish group  $j$  was calculated as

$$Q_{t,j} = \sum_{i=1}^{n_{ij}} w_{ij} (e_{ij} - \bar{E}_j)^2$$

and its significance was tested against the  $\chi^2$  distribution with  $n_{ij} - 1$  degrees of freedom.

We used a random-effects model to calculate  $\bar{E}_j$  and  $Q_{t,j}$  using the *metafor* package in R through the call

`rma(eij, vij, method = "REML", data = )` following the suggestions in Viechtbauer (2005)

that the REML, restricted maximum likelihood estimator for variance, strikes a good balance between unbiasedness and efficiency.

We ran mixed-effects categorical analyses and meta-regression to examine how the seven additional variables (Table 1, main text) mediated the overall effect size  $\bar{E}_j$ . For a given level ( $L$ ) of the factor, a weighted cumulative effect size was calculated as

$$\bar{E}_{L,j} = \frac{\sum_{i=1}^{n_{Lj}} w_{ij} e_{ij}}{\sum_{i=1}^{n_{Lj}} w_{ij}}$$

Where  $n_{Lj}$  is the number of comparison pairs in level,  $L$ , of the factor, and  $e_{ij}$  and  $w_{ij}$  are defined above. The heterogeneity of the model explained by the factor ( $Q_m$ ) was calculated as

$$Q_{m,j} = \sum_{i=1}^{n_{Lj}} w_{ij} (\bar{E}_{L,j} - \bar{E}_j)^2$$

The significance of  $Q_m$  was tested against the  $\chi^2$  distribution with  $n_{Lj} - 1$  degrees of freedom

## References

1082 Hedges LV, Gurevitch J, Curtis PS. 1999. The meta- analysis of response ratios in  
1083 experimental ecology. Ecology **80**:1150-1156.  
1084 Osenberg CW, Sarnelle O, Cooper SD, Holt RD. 1999. Resolving ecological questions  
1085 through meta- analysis: goals, metrics, and models. Ecology **80**:1105-1117.  
1086  
1087  
1088

## Appendix F - Sensitivity analysis for case of one-armed zero events

Cheng et al. (2016) define two scenarios relating to zero events: zero-event and double-zero-event. In the present study there were both double-zero-events (a zero mean abundance/biomass for both inside and outside) and zero-events (zero mean abundance/biomass for either the inside or the outside), summarised in Table 1. In both cases,  $\ln R$  cannot be computed due to division by zero and/or logarithm of zero. The former case are often removed from meta-analyses, while the latter is often handled via the addition of a small constant (Bradburn et al. 2007; Spittal et al. 2015). The total number of comparisons ( $n$ ) used in the final calculated of the weighted average effect size is given.


In the present study the species examined can be encountered infrequently on a survey. As well, one of the key species, *Lethrinus nebulosus*, is a schooling species which, when it occurs, can be present in larger numbers. Consequently there are many zero counts in the data which needed to be explicitly considered.

**Table F1. Summary of total number of comparisons and zero events**

	Fish Group	Total number of possible comparisons	Double-zero events	Zero-events	Final sample size (n) in calculation of mean weighted effect size
ABUNDANCE	LETHRINIDAE	305	4	151	301
	<i>L. nebulosus</i>	305	48	162	257
	<i>L. atkinsoni</i>	305	25	101	280
	EPINEPHELINAE	305	28	110	277
	<i>E. rivulatus</i>	305	138	210	167
	SCARINAE	305	44	46	261
BIOMASS	LETHRINIDAE	268	4	49	264
	<i>L. nebulosus</i>	268	45	143	223
	<i>L. atkinsoni</i>	268	21	86	247
	EPINEPHELINAE	268	15	87	253
	<i>E. rivulatus</i>	268	112	180	156
	SCARINAE	268	45	48	223

Double-zero-events were removed from analysis. For the remaining data a constant of 0.5 fish (half the smallest unit in the abundance analysis) was added to both the inside and outside mean counts, while for biomass a constant of 100 grams was added to both the inside and outside mean biomass.

Given there is no clear protocol for the size of the constant used to deal with zeros in a  $\ln R$  meta-analysis, we conducted a sensitivity analysis on the value of the constant. We ran the sensitivity analysis for the abundance data for the six taxa examined. We tested constant values,  $c$ , ranging from  $c = 0.01$  to  $c = 1$  in 0.01 increments.



**Figure F1.** Sensitivity analysis for the size of the constant added to inside/outside mean abundance showing the resulting transformed mean weighted effect size, exponential of  $\bar{E}$ , and 95% confidence interval (CI) for values of constant  $c$  ranging from 0.01 to 1 in 0.01 increments. a) Lethrinidae, b) *L. nebulosus*, c) *L. atkinsoni*, d) Epinephelinae, e) *E. rivulatus*, f) Scarinae. The mean effect sizes are considered significant when the confidence intervals do not include one.

It is clear that the size of the constant used influenced the mean overall effect size. For example, the magnitude of  $\bar{E}$  for *L. nebulosus* total abundance varied from  $\bar{E} = 0.73$  (0.54) when  $c = 0.01$  to  $\bar{E} = 0.28$  (0.08) when  $c = 1$ . However, while the constant size impacted the magnitude of the effect size, it did not influence the significance, except for *E. rivulatus*, which had the highest count of zero-events, and transitioned from marginally negative to not significantly different from one. From this analysis we decided a constant of 0.5 would be an adequate, and conservative addition for the calculation of  $\ln R$  in this analysis. Given the high levels of inherent variability expected in fish count data (Samoilys et al. 1995; Cappo & Brown 1996) and additional variation from uncontrolled variables, even with addition of a constant, overall differences in abundance would have to be consistent in order to observe statistical significance. Nonetheless, we urge caution in the interpretation of the magnitude of the overall effect.

## References

- Abesamis RA, Green AL, Russ GR, Jadloc CRL. 2014. The intrinsic vulnerability to fishing of coral reef fishes and their differential recovery in fishery closures. *Reviews in Fish Biology and Fisheries* **24**:1033-1063.
- Anderson DR, Burnham KP. 2002. Avoiding pitfalls when using information-theoretic methods. *The Journal of Wildlife Management*:912-918.
- Ayling T, Ayling AL 1987. Ningaloo Marine Park: preliminary fish density assessment and habitat survey: with information on coral damage due to *Drupella cornus* grazing: a report prepared for the Department of Conservation and Land Management, Western Australia. Department of Conservation and Land Management.
- Babcock R, Haywood M, Vanderklift M, Clapin G, Kleczkowski M, Dennis D, Skewes T, Milton D, Murphy N, Pillans R. 2008. Ecosystem Impacts of Human Usage and the Effectiveness of Zoning for Biodiversity conservation: Broad-scale Fish Census. Final Analysis and Recommendations 2007., Hobart.
- Babcock R, Phillips J, Lourey M, Clapin G. 2007. Increased density, biomass and egg production in an unfished population of Western Rock Lobster (*Panulirus cygnus*) at Rottnest Island, Western Australia. *Marine and Freshwater Research* **58**:286-292.
- Babcock R, Pillans R, Rochester W. 2017. Environmental and individual effects on the behaviour and spawning movements of *Lethrinus nebulosus* on a coral reef. *Marine and Freshwater Research* **68**:1422-1437.
- Ballantine B. 2014. Fifty years on: lessons from marine reserves in New Zealand and principles for a worldwide network. *Biological Conservation* **176**:297-307.
- Barlow J, França F, Gardner TA, Hicks CC, Lennox GD, Berenguer E, Castello L, Economo EP, Ferreira J, Guenard B. 2018. The future of hyperdiverse tropical ecosystems. *Nature* **559**:517.
- Barrett NS, Edgar GJ, Buxton CD, Haddon M. 2007. Changes in fish assemblages following 10 years of protection in Tasmanian marine protected areas. *Journal of Experimental Marine Biology and Ecology* **345**:141-157.
- Beckley LE, Lombard AT. 2012. A systematic evaluation of the incremental protection of broad-scale habitats at Ningaloo Reef, Western Australia. *Marine and Freshwater Research* **63**:17-22.
- Bergseth BJ, Williamson DH, Russ GR, Sutton SG, Cinner JE. 2017. A social–ecological approach to assessing and managing poaching by recreational fishers. *Frontiers in Ecology and the Environment* **15**:67-73.
- Bornt KR, McLean DL, Langlois TJ, Harvey ES, Bellchambers LM, Evans SN, Newman SJ. 2015. Targeted demersal fish species exhibit variable responses to long-term protection from fishing at the Houtman Abrolhos Islands. *Coral Reefs* **34**:1297-1312.
- Bradburn MJ, Deeks JJ, Berlin JA, Russell Localio A. 2007. Much ado about nothing: a comparison of the performance of meta- analytical methods with rare events. *Statistics in medicine* **26**:53-77.
- CALM. 1989. Ningaloo Marine Park Management Plan 1989 - 1999. Perth, WA, Australia.
- CALM. 2005. Management Plan for the Ningaloo Marine Park and Muiron Islands Marine Management Area 2005-2015: Management Plan No 52. Perth, WA, Australia.
- Cappo MM, Brown II. 1996. Evaluation of sampling methods for reef fish populations of commercial, recreational interest. CRC Reef Research Technical report No. 6.
- Cheng J, Pullenayegum E, Marshall JK, Iorio A, Thabane L. 2016. Impact of including or excluding both-armed zero-event studies on using standard meta-analysis methods for rare event outcome: a simulation study. *BMJ Open* **6**.

1188 Claudet J, Guidetti P. 2010. Improving assessments of marine protected areas. *Aquatic*  
1189 *conservation: marine and freshwater ecosystems* **20**:239-242.

1190 Claudet J, Osenberg C, Domenici P, Badalamenti F, Milazzo M, Falcón JM, Bertocci I,  
1191 Benedetti-Cecchi L, García-Charton JA, Goñi R. 2010. Marine reserves: fish life  
1192 history and ecological traits matter. *Ecological applications* **20**:830-839.

1193 Claudet J, Osenberg CW, Benedetti- Cecchi L, Domenici P, García- Charton JA,  
1194 Pérez- Ruzafa Á, Badalamenti F, Bayle- Sempere J, Brito A, Bulleri F. 2008. Marine  
1195 reserves: size and age do matter. *Ecology letters* **11**:481-489.

1196 Coleman FC, Figueira WF, Ueland JS, Crowder LB. 2004. The Impact of United States  
1197 Recreational Fisheries on Marine Fish Populations. *Science* **305**:1958.

1198 Collins LB, Zhu ZR, Wyrwoll K-H, Eisenhauer A. 2003. Late Quaternary structure and  
1199 development of the northern Ningaloo Reef, Australia. *Sedimentary Geology* **159**:81-  
1200 94.

1201 Côté IM, Mosqueira I, Reynolds JD. 2005. Effects of marine reserve characteristics on the  
1202 protection of fish populations: a meta- analysis. *Journal of Fish Biology* **59**:178-189.

1203 Cowx IG, Cooke SJ. 2004. The Role of Recreational Fishing in Global Fish Crises.  
1204 *BioScience* **54**:857-859.

1205 Denny CM, Willis TJ, Babcock RC. 2004. Rapid recolonisation of snapper *Pagrus auratus*:  
1206 Sparidae within an offshore island marine reserve after implementation of no-take  
1207 status. *Marine Ecology Progress Series* **272**:183-190.

1208 Depczynski MM, Tinkler PP, Cheal AA, Speed CC. 2015. Ningaloo fish communities. Page  
1209 121 in Miller KK, Depczynski MM, Cappo MM, Wakeford MM, Speed CC, Stowar  
1210 MM, Colquhoun JJ, Tinkler PP, Cheal AA, and Fisher RR, editors. Ningaloo and  
1211 Outer Shark Bay Environmental Baseline Survey 2014. Report prepared by the  
1212 Australian Institute of Marine Science for Woodside Energy Ltd. Australian Institute  
1213 of Marine Science, Townsville, Australia.

1214 DPIRD DoPIaRD. 2018. Ningaloo Marine Park, Available from  
1215 [https://parks.dpaw.wa.gov.au/sites/default/files/downloads/parks/2462-](https://parks.dpaw.wa.gov.au/sites/default/files/downloads/parks/2462-13%20Spearfishing%20in%20Ningaloo%20WEB.pdf)  
1216 [13%20Spearfishing%20in%20Ningaloo%20WEB.pdf](https://parks.dpaw.wa.gov.au/sites/default/files/downloads/parks/2462-13%20Spearfishing%20in%20Ningaloo%20WEB.pdf) (accessed 4/11/2018 2018).

1217 DPIRD DoPIaRDWA. 2017. Gascoyne Coast Bioregion, Status reports of the fisheries and  
1218 aquatic resources of Western Australia 2016/17.

1219 Edgar GJ, Stuart-Smith RD, Willis TJ, Kininmonth S, Baker SC, Banks S, Barrett NS,  
1220 Becerro MA, Bernard AT, Berkhout J. 2014. Global conservation outcomes depend  
1221 on marine protected areas with five key features. *Nature* **506**:216-220.

1222 Fisher R, Wilson Shaun K, Sin Tsai M, Lee Ai C, Langlois Tim J. 2018. A simple function  
1223 for full- subsets multiple regression in ecology with R. *Ecology and Evolution* **0**.

1224 Fitzpatrick B, Harvey E, Langlois T, Babcock R, Twiggs E. 2015. Effects of fishing on fish  
1225 assemblages at the reefscape scale. *Marine Ecology Progress Series* **524**:241-253.

1226 Froese RaDP. 2018. FishBase, Available from [www.fishbase.org](http://www.fishbase.org) (accessed 15/1/2018 2018).

1227 Fulton CJ, Depczynski M, Holmes TH, Noble MM, Radford B, Wernberg T, Wilson SK.  
1228 2014. Sea temperature shapes seasonal fluctuations in seaweed biomass within the  
1229 Ningaloo coral reef ecosystem. *Limnology and Oceanography* **59**:156-166.

1230 Goetze J, Jupiter S, Langlois T, Wilson S, Harvey E, Bond T, Naisilisili W. 2015. Diver  
1231 operated video most accurately detects the impacts of fishing within periodically  
1232 harvested closures. *Journal of Experimental Marine Biology and Ecology* **462**:74-82.

1233 Halpern BS, Warner RR. 2002. Marine reserves have rapid and lasting effects. *Ecology*  
1234 *Letters* **5**:361-366.

1235 Hedges LV, Gurevitch J, Curtis PS. 1999. The meta- analysis of response ratios in  
1236 experimental ecology. *Ecology* **80**:1150-1156.

1237 Holmes TH, Wilson SK, Travers MJ, Langlois TJ, Evans RD, Moore GI, Douglas RA,  
 1238 Shedrawi G, Harvey ES, Hickey K. 2013. A comparison of visual-and stereo-video  
 1239 based fish community assessment methods in tropical and temperate marine waters of  
 1240 Western Australia. *Limnology and Oceanography, Methods* **11**:337-350.  
 1241 Horta e Costa B, Claudet J, Franco G, Erzini K, Caro A, Gonçalves EJ. 2016. A regulation-  
 1242 based classification system for Marine Protected Areas (MPAs). *Marine Policy*  
 1243 **72**:192-198.  
 1244 Jenkins CN, Joppa L. 2009. Expansion of the global terrestrial protected area system.  
 1245 *Biological conservation* **142**:2166-2174.  
 1246 Langlois TJ, Harvey ES, Fitzpatrick B, Meeuwig JJ, Shedrawi G, Watson DL. 2010. Cost-  
 1247 efficient sampling of fish assemblages: comparison of baited video stations and diver  
 1248 video transects. *Aquatic biology* **9**:155-168.  
 1249 Lester SE, Halpern BS, Grorud-Colvert K, Lubchenco J, Ruttenberg BI, Gaines SD, Aïramé  
 1250 S, Warner RR. 2009. Biological effects within no-take marine reserves: a global  
 1251 synthesis. *Marine Ecology Progress Series* **384**:33-46.  
 1252 Lewin W-C, Arlinghaus R, Mehner T. 2006. Documented and Potential Biological Impacts of  
 1253 Recreational Fishing: Insights for Management and Conservation. *Reviews in*  
 1254 *Fisheries Science* **14**:305-367.  
 1255 Lim IE, Wilson SK, Holmes TH, Noble MM, Fulton CJ. 2016. Specialization within a  
 1256 shifting habitat mosaic underpins the seasonal abundance of a tropical fish. *Ecosphere*  
 1257 **7**:e01212.  
 1258 Ma KY, Craig MT. 2018. An inconvenient monophyly: an update on the taxonomy of the  
 1259 groupers (Epinephelidae). *Copeia* **106**:443-456.  
 1260 Mackie M, Black R 1999. Research on Two Serranid Species (Serranidae: Epinephelinae) in  
 1261 Western Australian Waters. University of Western Australia/Fisheries Research &  
 1262 Development Corporation.  
 1263 Mallet D, Pelletier D. 2014. Underwater video techniques for observing coastal marine  
 1264 biodiversity: a review of sixty years of publications (1952–2012). *Fisheries Research*  
 1265 **154**:44-62.  
 1266 McCook LJ, et al. 2010. Adaptive management of the Great Barrier Reef: A globally  
 1267 significant demonstration of the benefits of networks of marine reserves. *Proceedings*  
 1268 *of the National Academy of Sciences* **107**:18278.  
 1269 McLaren BW, Langlois TJ, Harvey ES, Shortland-Jones H, Stevens R. 2015. A small no-take  
 1270 marine sanctuary provides consistent protection for small-bodied by-catch species, but  
 1271 not for large-bodied, high-risk species. *Journal of Experimental Marine Biology and*  
 1272 *Ecology* **471**:153-163.  
 1273 McLean DL, et al. 2016. Distribution, abundance, diversity and habitat associations of fishes  
 1274 across a bioregion experiencing rapid coastal development. *Estuarine, Coastal and*  
 1275 *Shelf Science* **178**:36-47.  
 1276 McPhee DP, Leadbitter D, Skilleter GA. 2002. Swallowing the bait: is recreational fishing in  
 1277 Australia ecologically sustainable? *Pacific Conservation Biology* **8**:40-51.  
 1278 Miller KI, Russ GR. 2014. Studies of no-take marine reserves: Methods for differentiating  
 1279 reserve and habitat effects. *Ocean & Coastal Management* **96**:51-60.  
 1280 Mitchell J, McLean D, Collin S, Taylor S, Jackson G, Fisher R, Langlois T. 2018.  
 1281 Quantifying shark depredation in a recreational fishery in the Ningaloo Marine Park  
 1282 and Exmouth Gulf, Western Australia. *Marine Ecology Progress Series* **587**:141-157.  
 1283 Mora C, Sale PF. 2011. Ongoing global biodiversity loss and the need to move beyond  
 1284 protected areas: a review of the technical and practical shortcomings of protected  
 1285 areas on land and sea. *Marine ecology progress series* **434**:251-266.

- Mosquera I, Côté IM, Jennings S, Reynolds JD. 2000. Conservation benefits of marine reserves for fish populations. Pages 321-332. Animal Conservation forum. Cambridge University Press.
- Murphy HM, Jenkins GP. 2010. Observational methods used in marine spatial monitoring of fishes and associated habitats: a review. *Marine and Freshwater Research* **61**:236-252.
- Osenberg CW, Sarnelle O, Cooper SD, Holt RD. 1999. Resolving ecological questions through meta- analysis: goals, metrics, and models. *Ecology* **80**:1105-1117.
- Osenberg CW, Shima JS, Miller SL, Stier AC. 2011. Ecology: assessing effects of marine protected areas: confounding in space and possible solutions. *Marine protected areas: a multidisciplinary approach*:143-167.
- Pillans RD, Bearham D, Boomer A, Downie R, Patterson TA, Thomson DP, Babcock RC. 2014. Multi year observations reveal variability in residence of a tropical demersal fish, *Lethrinus nebulosus*: implications for spatial management. *PLoS One* **9**:e105507.
- R Core Team. 2017. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Radford CA, Jeffs AG, Tindle CT, Cole RG, Montgomery JC. 2005. Bubbled waters: The noise generated by underwater breathing apparatus. *Marine and freshwater behaviour and physiology* **38**:259-267.
- Rees MJ, Knott NA, Davis AR. 2018a. Habitat and seascape patterns drive spatial variability in temperate fish assemblages: implications for marine protected areas. *Marine Ecology Progress Series* **607**:171-186.
- Rees MJ, Knott NA, Neilson J, Linklater M, Osterloh I, Jordan A, Davis AR. 2018b. Accounting for habitat structural complexity improves the assessment of performance in no-take marine reserves. *Biological Conservation* **224**:100-110.
- Russ GR. 2002. Yet another review of marine reserves as reef fishery management tools. *Coral reef fishes: dynamics and diversity in a complex ecosystem* **24**:421.
- Russ GR, Miller KI, Rizzari JR, Alcalá AC. 2015. Long-term no-take marine reserve and benthic habitat effects on coral reef fishes. *Marine Ecology Progress Series* **529**:233-248.
- Ryan K, Hall N, Lai E, Smallwood C, Taylor S, Wise B 2015. State-wide survey of boat-based recreational fishing in Western Australia 2013/14. Fisheries Research Division.
- Ryan K, Hall N, Lai E, Smallwood C, Taylor S, Wise B. 2017. State-wide survey of boat-based recreational fishing in Western Australia 2015/16.
- Ryan K, Wise B, Hall N, Pollock K, Sulin E, Gaughan DJ 2013. An integrated system to survey boat-based recreational fishing in Western Australia 2011/12. Fisheries Research Division, Western Australian Fisheries and Marine Research Laboratories.
- Sala E, Lubchenco J, Grorud-Colvert K, Novelli C, Roberts C, Sumaila UR. 2018. Assessing real progress towards effective ocean protection. *Marine Policy* **91**:11-13.
- Samoilys M, Fuentes H, Tuwai I, Tikomainiusiladi B, Leqata J, Oreihaka E, Mobiha A, Potuku T, Die D, Connell S. 1995. Application of underwater visual census to assessing coral reef fish stocks in the tropical Pacific. Report prepared for the Australian Centre for International Agricultural Research (ACIAR). ACIAR Project.
- Shears NT, Grace RV, Usmar NR, Kerr V, Babcock RC. 2006. Long-term trends in lobster populations in a partially protected vs. no-take Marine Park. *Biological conservation* **132**:222-231.
- Smallwood CB, Beckley LE. 2012. Spatial distribution and zoning compliance of recreational fishing in Ningaloo Marine Park, north-western Australia. *Fisheries Research* **125**:40-50.
- Spittal MJ, Pirkis J, Gurrin LC. 2015. Meta-analysis of incidence rate data in the presence of zero events. *BMC medical research methodology* **15**:42.

- Sumner NR, Williamson PC, Malseed BE 2002. A 12-month survey of recreational fishing in the Gascoyne bioregion of Western Australia during 1998-99. Department of Fisheries, Western Australia.
- Underwood A. 1993. The mechanics of spatially replicated sampling programmes to detect environmental impacts in a variable world. *Australian Journal of ecology* **18**:99-116.
- Viechtbauer W. 2005. Bias and efficiency of meta-analytic variance estimators in the random-effects model. *Journal of Educational and Behavioral Statistics* **30**:261-293.
- Viechtbauer W. 2010. Conducting meta-analyses in R with the metafor package. *J Stat Softw* **36**:1-48.
- Watson DL, Harvey ES, Anderson MJ, Kendrick GA. 2005. A comparison of temperate reef fish assemblages recorded by three underwater stereo-video techniques. *Marine Biology* **148**:415-425.
- Watson M, Ormond R. 1994. Effect of an artisanal fishery on the fish and urchin populations of a Kenyan coral reef. *Marine Ecology Progress Series*:115-129.
- West L, Stark K, Murphy J, Lyle J, Ochwada-Doyle F. 2015. Survey of recreational fishing in New South Wales and the ACT, 2013/14.
- Westera MB. 2003a. The effect of recreational fishing on targeted fishes and trophic structure, in a coral reef marine park.
- Westera MB. 2003b. The effect of recreational fishing on targeted fishes and trophic structure, in a coral reef marine park. Edith Cowan University, Perth, Western Australia.
- Wilson S, Fulton C, Depczynski M, Holmes T, Noble M, Radford B, Tinkler P. 2014. Seasonal changes in habitat structure underpin shifts in macroalgae-associated tropical fish communities. *Marine biology* **161**:2597-2607.
- Wilson SK, Babcock RC, Fisher R, Holmes TH, Moore JAY, Thomson DP. 2012. Relative and combined effects of habitat and fishing on reef fish communities across a limited fishing gradient at Ningaloo. *Marine Environmental Research* **81**:1-11.
- Wilson SK, Depczynski M, Fisher R, Holmes TH, Noble MM, Radford BT, Rule M, Shedrawi G, Tinkler P, Fulton CJ. 2018a. Climatic forcing and larval dispersal capabilities shape the replenishment of fishes and their habitat-forming biota on a tropical coral reef. *Ecology and Evolution* **8**:1918-1928.
- Wilson SK, Depczynski M, Holmes TH, Noble MM, Radford BT, Tinkler P, Fulton CJ. 2017. Climatic conditions and nursery habitat quality provide indicators of reef fish recruitment strength. *Limnology and Oceanography* **62**:1868-1880.
- Wilson SK, Graham NAJ, Holmes T, MacNeil MA, Ryan N. 2018b. Visual versus video methods for estimating reef fish biomass. *Ecological Indicators* **85**:146-152.
- Wood SN. 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **73**:3-36.
- Zupan M, Fragkopoulou E, Claudet J, Erzini K, Horta e Costa B, Gonçalves EJ. 2018. Marine partially protected areas: drivers of ecological effectiveness. *Frontiers in Ecology and the Environment* **16**:381-387.
- Zuur AF, Ieno EN, Elphick CS. 2010. A protocol for data exploration to avoid common statistical problems. *Methods in ecology and evolution* **1**:3-14.

1384

1385

1386

1387

1388

## Appendix G - Meta-analysis statistics

**Table G1. Total heterogeneity statistics**

Fish group	ABUNDANCE			BIOMASS		
	Q <sub>T</sub>	df	P	Q <sub>T</sub>	df	P
LETHRINIDAE	2002.57	300	<0.001	2318.26	263	<0.001
<i>L. nebulosus</i>	1971.07	256	<0.001	2886.99	222	<0.001
<i>L. atkinsoni</i>	1739.71	279	<0.001	1928.65	246	<0.001
EPINEPHELINAE	1125.65	276	<0.001	1307.49	252	<0.001
<i>E. rivulatus</i>	477.33	166	<0.001	590.76	155	<0.001
SCARINAE	1701.09	260	<0.001	1224.57	222	<0.001

Table G2 and G3 summarise the results of weighted mixed-effects meta-analyses for all seven variables and for *reserve identity* modelled individually. Figure 1 shows the predicted effect size for the cases where the moderator reserve identity explained a significant amount of heterogeneity for *habitat* so that it can be directly compared to the overall effect sizes in Figure 2 (main text).

**Table G2. Mixed-effects models heterogeneity statistics for abundance data**

	Fish group	ABUNDANCE					
		Model heterogeneity*			Residual heterogeneity		
		Q <sub>m</sub>	df	P	Q <sub>e</sub>	df	P
HABITAT	LETHRINIDAE	39.46	3	<0.001	1622.38	297	<0.001
	<i>L. nebulosus</i>	32.51	3	<0.001	1574.07	253	<0.001
	<i>L. atkinsoni</i>	14.55	3	<0.001	1614.93	276	<0.001
	EPINEPHELINAE	0.31	3	0.96	1117.33	273	<0.001
	<i>E. rivulatus</i>	5.39	3	0.15	467.33	163	<0.001
	SCARINAE	6	3	0.11	1589.43	257	<0.001
METHOD	LETHRINIDAE	4.09	2	0.13	1975.79	298	<0.001
	<i>L. nebulosus</i>	0.71	2	0.7	1946.56	254	<0.001
	<i>L. atkinsoni</i>	7.85	2	0.02	1676.16	277	<0.001
	EPINEPHELINAE	4.35	2	0.11	1094.35	274	<0.001
	<i>E. rivulatus</i>	0.98	2	0.61	474.68	164	<0.001
	SCARINAE	0.01	2	0.99	1633.03	258	<0.001
RESERVE SIZE	LETHRINIDAE	3.8	1	0.05	1959.06	299	<0.001
	<i>L. nebulosus</i>	0.01	1	0.94	1970.56	255	<0.001
	<i>L. atkinsoni</i>	8.11	1	<0.001	1688.35	278	<0.001
	EPINEPHELINAE	1.58	1	0.21	1112.23	275	<0.001
	<i>E. rivulatus</i>	0.47	1	0.49	477.08	165	<0.001
	SCARINAE	0.02	1	0.88	1701.09	259	<0.001
YEARS PROTECTION	LETHRINIDAE	0	1	0.95	2001.07	299	<0.001
	<i>L. nebulosus</i>	0.92	1	0.34	1949.89	255	<0.001
	<i>L. atkinsoni</i>	1.86	1	0.17	1736.82	278	<0.001
	EPINEPHELINAE	0.42	1	0.52	1115.08	275	<0.001
	<i>E. rivulatus</i>	5.43	1	0.02	462.46	165	<0.001
	SCARINAE	0.83	1	0.36	1689.03	259	<0.001
ZONING SCHEME	LETHRINIDAE	0.83	1	0.36	2002.31	299	<0.001
	<i>L. nebulosus</i>	0.87	1	0.35	1961.63	255	<0.001
	<i>L. atkinsoni</i>	8.43	1	<0.001	1696.16	278	<0.001
	EPINEPHELINAE	1.1	1	0.29	1100.82	275	<0.001
	<i>E. rivulatus</i>	13.43	1	<0.001	440.94	165	<0.001
	SCARINAE	3.82	1	0.05	1680.66	259	<0.001
ENTIRE DATA SET	LETHRINIDAE	1.15	1	0.28	1987.99	295	<0.001

	<i>L. nebulosus</i>	0.7	1	0.4	1960.92	251	<0.001
	<i>L. atkinsoni</i>	0.55	1	0.46	1698.56	275	<0.001
	EPINEPHELINAE	0.01	1	0.92	1099.86	271	<0.001
	<i>E. rivulatus</i>	4.79	1	<b>0.03</b>	457.62	162	<0.001
	SCARINAE	5.9	1	<b>0.02</b>	1670.41	255	<0.001
SHORE FISHING	LETHRINIDAE	7.09	1	<b>0.01</b>	1953.98	299	<0.001
	<i>L. nebulosus</i>	0.16	1	0.69	1966.75	255	<0.001
	<i>L. atkinsoni</i>	9.03	1	<b>&lt;0.001</b>	1689.1	278	<0.001
	EPINEPHELINAE	3.9	1	<b>0.05</b>	1060.95	275	<0.001
	<i>E. rivulatus</i>	6.16	1	<b>0.01</b>	461.9	165	<0.001
RESERVE IDENTITY	SCARINAE	5.59	1	<b>0.02</b>	1674.58	259	<0.001
	LETHRINIDAE	43.22	21	<b>&lt;0.001</b>	1746.07	279	<0.001
	<i>L. nebulosus</i>	48.82	21	<b>&lt;0.001</b>	1665.43	235	<0.001
	<i>L. atkinsoni</i>	35.29	20	<b>0.02</b>	1494.98	259	<0.001
	EPINEPHELINAE	54.33	21	<b>&lt;0.001</b>	880.04	255	<0.001
SURVEY YEAR	<i>E. rivulatus</i>	68.95	20	<b>&lt;0.001</b>	344.61	146	<0.001
	SCARINAE	27.81	21	0.15	1484.39	239	<0.001
	LETHRINIDAE	2.39	1	0.12	1994.3	299	<0.001
	<i>L. nebulosus</i>	0	1	0.99	1962.36	255	<0.001
	<i>L. atkinsoni</i>	5.75	1	<b>0.02</b>	1674.53	278	<0.001
	EPINEPHELINAE	0.5	1	0.48	1115.22	275	<0.001
	<i>E. rivulatus</i>	3.59	1	0.06	468.95	165	<0.001
	SCARINAE	21.29	1	<b>&lt;0.001</b>	1579.86	259	<0.001

\* Total heterogeneity is provided in Table G1

**Table G3. Mixed-effects models heterogeneity statistics for abundance data**

		BIOMASS					
		Model heterogeneity*			Residual heterogeneity		
		<b>Q<sub>m</sub></b>	<b>df</b>	<b>P</b>	<b>Q<sub>e</sub></b>	<b>df</b>	<b>P</b>
HABITAT	LETHRINIDAE	41.75	3	<b>&lt;0.001</b>	2082.5	260	<0.001
	<i>L. nebulosus</i>	29.38	3	<b>&lt;0.001</b>	2743.83	219	<0.001
	<i>L. atkinsoni</i>	17.13	3	<b>&lt;0.001</b>	1714.19	243	<0.001
	EPINEPHELINAE	1.21	3	0.75	1296.97	249	<0.001
	<i>E. rivulatus</i>	2.53	3	0.47	517.38	152	<0.001
	SCARINAE	22.86	3	<b>&lt;0.001</b>	1038.84	219	<0.001
METHOD	LETHRINIDAE	2.46	2	0.29	2140.53	261	<0.001
	<i>L. nebulosus</i>	0.63	2	0.73	2829.86	220	<0.001
	<i>L. atkinsoni</i>	7.79	2	<b>0.02</b>	1720.61	244	<0.001
	EPINEPHELINAE	12.37	2	<b>&lt;0.001</b>	1258.74	250	<0.001
	<i>E. rivulatus</i>	5.9	2	0.05	586.26	153	<0.001
	SCARINAE	1.16	2	0.56	1193.63	220	<0.001
RESERVE SIZE	LETHRINIDAE	1.3	1	0.25	2318.18	262	<0.001
	<i>L. nebulosus</i>	0.1	1	0.75	2886.72	221	<0.001
	<i>L. atkinsoni</i>	1.97	1	0.16	1925.76	245	<0.001
	EPINEPHELINAE	3.78	1	0.05	1307.41	251	<0.001
	<i>E. rivulatus</i>	1.32	1	0.25	571.52	154	<0.001
	SCARINAE	0.59	1	0.44	1220.28	221	<0.001
YEARS PROTECTION	LETHRINIDAE	0.01	1	0.94	2312.18	262	<0.001
	<i>L. nebulosus</i>	0.71	1	0.4	2882.59	221	<0.001
	<i>L. atkinsoni</i>	3.13	1	0.08	1924.77	245	<0.001
	EPINEPHELINAE	0.12	1	0.73	1307.3	251	<0.001
	<i>E. rivulatus</i>	5.01	1	<b>0.03</b>	587.67	154	<0.001
	SCARINAE	0.02	1	0.89	1218.81	221	<0.001
ZONING SCHEME	LETHRINIDAE	0.44	1	0.51	2314.92	262	<0.001
	<i>L. nebulosus</i>	0.54	1	0.46	2883.99	221	<0.001
	<i>L. atkinsoni</i>	9.59	1	<b>&lt;0.001</b>	1925.41	245	<0.001
	EPINEPHELINAE	0.04	1	0.83	1307.27	251	<0.001
	<i>E. rivulatus</i>	9.96	1	<b>&lt;0.001</b>	587.78	154	<0.001
	SCARINAE	3.09	1	0.08	1222.86	221	<0.001
BOAT FISHING	LETHRINIDAE	0.18	1	0.67	2313.23	258	<0.001
	<i>L. nebulosus</i>	0.04	1	0.84	2879.67	217	<0.001
	<i>L. atkinsoni</i>	0.14	1	0.7	1923.96	243	<0.001
	EPINEPHELINAE	0.82	1	0.37	1287.94	247	<0.001
	<i>E. rivulatus</i>	3.75	1	0.05	578.7	151	<0.001

SHORE FISHING	SCARINAE	0.57	1	0.45	1193.33	217	<0.001
	LETHRINIDAE	2.19	1	0.14	2295.06	262	<0.001
	<i>L. nebulosus</i>	0	1	0.96	2606.71	221	<0.001
	<i>L. atkinsoni</i>	5.25	1	<b>0.02</b>	1927.59	245	<0.001
	EPINEPHELINAE	0.38	1	0.54	1268.07	251	<0.001
	<i>E. rivulatus</i>	3.05	1	0.08	580.64	154	<0.001
RESERVE IDENTITY	SCARINAE	5.78	1	<b>0.02</b>	1168.18	221	<0.001
	LETHRINIDAE	29.09	21	0.11	2019.62	242	<0.001
	<i>L. nebulosus</i>	27.02	21	0.17	1941.89	201	<0.001
	<i>L. atkinsoni</i>	33.93	20	<b>0.03</b>	1846.92	226	<0.001
	EPINEPHELINAE	58.1	21	<b>&lt;0.001</b>	987.83	231	<0.001
	<i>E. rivulatus</i>	49.47	20	<b>&lt;0.001</b>	503.04	135	<0.001
SURVEY YEAR	SCARINAE	33.11	21	<b>0.04</b>	1046.15	201	<0.001
	LETHRINIDAE	2.39	1	0.12	1994.3	299	<0.001
	<i>L. nebulosus</i>	0	1	0.99	1962.36	255	<0.001
	<i>L. atkinsoni</i>	5.75	1	<b>0.02</b>	1674.53	278	<0.001
	EPINEPHELINAE	0.5	1	0.48	1115.22	275	<0.001
	<i>E. rivulatus</i>	3.59	1	0.06	468.95	165	<0.001
	SCARINAE	21.29	1	<b>&lt;0.001</b>	1579.86	259	<0.001

\* Total heterogeneity is provided in Table G1

Figure G1 shows the transformed predicted effect sizes as relative fish abundance by *habitat* while Figure G2 shows this for *reserve identity*.






Figure G1. Relative fish abundance inside to outside reserves (back-transformed weighted mean effect sizes) with 95% confidence intervals) for the Lethrinidae, *Lethrinus nebulosus*, and *L. atkinsoni*, the fish groups for which *habitat* explained a significant amount of variation. Effect sizes are considered significant when the confidence intervals do not include one. Black dots correspond to significant effects while grey dots correspond to non-significant effects.

Figure G2 is included to show that there is heterogeneity between individual reserves and because it may be useful to managers of the Ningaloo Marine Park.



1425



1426


1427

Figure G2. Relative fish abundance (Transformed weighted mean effect sizes ( $\exp(e)$ ) with 95% confidence intervals ( $CI$ )), for the Lethrinidae, *Lethrinus nebulosus*, *Lethrinus atkinsoni*, Epinephelinae and *Epinephelus rivulatus*, the fish groups for which *reserve identity* explained a significant amount of variation. Results are categorised into panels based on the initial or current zoning. Point shape indicates whether shore fishing is allowed or prohibited along the coastal edge of the reserve (circle = prohibited; triangle = allowed). Effect sizes are considered significant when the confidence intervals do not include one. Black dots correspond to significant effects while grey dots correspond to non-significant effects.

## Appendix H – Biomass results


**Table H1.** Top Generalised Additive Mixed Models (GAMMs) for predicting the response ratio inside to outside reserves,  $\bar{E}$ , for biomass from full subset analyses for the abundance of the six fish groups. Difference between the lowest reported corrected Akaike Information Criterion ( $\Delta AICc$ ),  $AICc$  weights ( $\omega AICc$ ), variance explained ( $R^2$ ) and estimated degrees of freedom (EDF) are reported for model comparison. Model selection was based on the most parsimonious model (fewest variables and lowest EDF) within two units of the lowest  $AICc$ . This model is shown in bold text.

Fish group	Model	$\Delta AICc$	$\omega AICc$	$R^2$	EDF
LETHRINIDAE	<b>Habitat + Size by Habitat</b>	0	0.855	0.17399	9
	<i>L. nebulosus</i> Habitat + Years protection by Habitat + Size by Habitat	0	0.443	0.19863	14.51
<i>L. atkinsoni</i>	<b>Habitat + Size by Habitat</b>	0.857	0.289	0.17016	9
	Years protection + Habitat + Size by Habitat	1.301	0.231	0.17594	10.66
	Habitat + Method + Years protection by Habitat	0	0.29	0.10828	12.98
	<b>Habitat + Years protection by Habitat</b>	0.39	0.238	0.09662	10.67
	Habitat + Method + Zoning scheme	1.273	0.153	0.08162	8
	Years protection + Boat fishing + Method	0	0.196	0.09835	7.81
EPINEPHELINAE	Boat fishing + Method	0.516	0.152	0.09171	6.68
	Years protection + Method	1.092	0.114	0.09192	7.24
	<b>Method</b>	1.198	0.108	0.08585	5.88
	Boat fishing + Method + Shore fishing	1.751	0.082	0.09098	7.61
	Boat fishing + Method + Zoning scheme	1.955	0.074	0.09064	7.71
	<b>Boat fishing + Method + Zoning scheme</b>	0	0.546	0.17358	12.31
<i>E. rivulatus</i>	Years protection + Boat fishing + Method	1.457	0.264	0.16292	11.96
	Years protection + Habitat + Size by Habitat	0	0.271	0.12181	16.44
SCARIDAE	<b>Habitat</b>	1.354	0.138	0.05738	5
	Years protection + Habitat + Size	1.406	0.134	0.09791	13.04
	Habitat + Zoning scheme	1.698	0.116	0.06077	6
	Habitat + Size by Habitat	1.889	0.106	0.07496	9
	Habitat + Size	1.952	0.102	0.06103	6.28



**Figure H1.** a) Relative fish biomass (Transformed weighted mean effect sizes ( $\exp(\bar{E})$ ) with 95% confidence intervals (CI)), for the relative abundance of the six fish taxa: Lethrinidae, *Lethrinus nebulosus*, *L. atkinsoni*, Epinephelinae, *Epinephelus rivulatus* and Scarinae. Effect sizes are considered significant when the confidence intervals do not include one. Open dots correspond to non-significant effects. Sample sizes are given in Appendix G. b) Importance scores (based on summed Akaike weights corrected for finite samples (AICc)) from full-subsets analyses exploring the influence of seven variables on the overall effect size for each fish taxa. *Reserve size* was square-root transformed and *boat fishing* was log-transformed in all models. Red X symbols mark the variables that were included in the most parsimonious models for each fish taxa (also see Table H1 and Fig. H2).

1473  
1474



1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483

Figure H2. Predicted relative fish biomass inside to outside reserves (back-transformed predicted weighted effect sizes) with 95% confidence intervals) for the six fish groups – a) Lethrinidae; b) *Lethrinus nebulosus* c) *Lethrinus atkinsoni*; d) Epinephelinae; e) *Epinephelus rivulatus*; f) Scarinae for abundance– as a function of variables present in the most parsimonious models (Table G4) from full-subsets GAMM analysis. Ribbons represent 95% confidence intervals

1484  
1485  
1486  
1487  
1488