



Connecting the data landscape of long-term ecological studies: the SPI-Birds data hub

Antica Culina, Frank Adriaensen, Liam Bailey, Malcolm Burgess, Anne Charmantier, Ella Cole, Tapio Eeva, Erik Matthysen, Chloé Nater, Ben Sheldon, et al.

► To cite this version:

Antica Culina, Frank Adriaensen, Liam Bailey, Malcolm Burgess, Anne Charmantier, et al.. Connecting the data landscape of long-term ecological studies: the SPI-Birds data hub. *Journal of Animal Ecology*, 2021, 90 (9), pp.2147-2160. 10.1111/1365-2656.13388 . hal-03035065

HAL Id: hal-03035065

<https://hal.science/hal-03035065>

Submitted on 2 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DR ANTICA CULINA (Orcid ID : 0000-0003-2910-8085)
 MR LIAM D. BAILEY (Orcid ID : 0000-0002-8226-9454)
 MS CHLOÉ REBECCA NATER (Orcid ID : 0000-0002-7975-0108)
 DR BEN C. SHELDON (Orcid ID : 0000-0002-5240-7828)
 PROFESSOR BERNT-ERIK SÆTHER (Orcid ID : 0000-0002-0049-9767)
 DR LUCY APLIN (Orcid ID : 0000-0001-5367-826X)
 ELENA ANGULO (Orcid ID : 0000-0001-5545-4032)
 MS JOSEFA BLEU (Orcid ID : 0000-0002-3403-8272)
 SANDRA BOUWHUIS (Orcid ID : 0000-0003-4023-1578)
 DR DAVID CANAL (Orcid ID : 0000-0003-2875-2987)
 DR ALEJANDRO CANTARERO (Orcid ID : 0000-0002-5816-701X)
 DR NIELS JEROEN DINGEMANSE (Orcid ID : 0000-0003-3320-0861)
 DAVIDE MICHELANGELO DOMINONI (Orcid ID : 0000-0003-2063-9955)
 DR DAMIEN ROGER FARINE (Orcid ID : 0000-0003-2208-7613)
 DR GERGELY HEGYI (Orcid ID : 0000-0002-4906-3550)
 MS CAROLINE ISAKSSON (Orcid ID : 0000-0002-6889-1386)
 DR BART KEMPENAERS (Orcid ID : 0000-0002-7505-5458)
 DR MILOŠ KRIST (Orcid ID : 0000-0002-6183-686X)
 MR JAANIS LODJAK (Orcid ID : 0000-0001-8089-948X)
 DR MARKO MÄGI (Orcid ID : 0000-0003-0736-5638)
 DR MARK C MAINWARING (Orcid ID : 0000-0002-0427-9673)
 DR SHINICHI NAKAGAWA (Orcid ID : 0000-0002-7765-5182)
 DR JOHAN NILSSON (Orcid ID : 0000-0001-6744-6486)
 DR JAIME POTTI (Orcid ID : 0000-0002-2284-0022)
 DR JOHN LEO QUINN (Orcid ID : 0000-0001-9363-3146)
 DR JUAN CARLOS SENAR (Orcid ID : 0000-0001-9955-3892)
 DR GÁBOR SERESS (Orcid ID : 0000-0001-9828-4934)
 PROFESSOR MARCEL E. E. VISSER (Orcid ID : 0000-0002-1456-1939)

Article type : Research Article

Handling Editor: Roberto Salguero-Gómez

Connecting the data landscape of long-term ecological studies: the SPI-Birds data hub

Antica Culina^{1#}, Frank Adriaensen², Liam D. Bailey^{1,3}, Malcolm D. Burgess^{4,5}, Anne

Charmantier⁶, Ella F. Cole⁷, Tapio Eeva^{8,9}, Erik Matthysen², Chloé R. Nater¹⁰, Ben C. Sheldon⁷,

© 2020 The Authors. Journal of Animal Ecology published by John Wiley & Sons Ltd on behalf of British Ecological Society

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Bernt-Erik Sæther¹⁰, Stefan J.G. Vriend¹⁰, Zuzana Zajkova¹, Peter Adamík¹¹, Lucy M. Aplin¹², Elena Angulo¹³, Alexandr Artemyev¹⁴, Emilio Barba¹⁵, Sanja Barišić¹⁶, Eduardo Belda¹⁷, C. Can Bilgin¹⁸, Josefa Bleu¹⁹, Christiaan Both²⁰, Sandra Bouwhuis²¹, Claire Branston²², Juli Broggi¹³, Terry Burke²³, Andrey Bushuev²⁴, Carlos Camacho²⁵, Daniela Campobello^{26,27}, David Canal²⁸, Alejandro Cantarero⁸, Samuel P. Caro⁶, Maxime Cauchoix²⁹, Alexis Chaine²⁹, Mariusz Cichoń³⁰, Davor Ćiković¹⁶, Camillo A. Cusimano²⁶, Caroline Deimel³¹, André A. Dhondt^{2,32}, Niels J. Dingemanse³³, Blandine Doligez³⁴, Davide M. Dominoni²², Claire Doutrelant⁶, Szymon M. Drobnik^{35,36}, Anna Dubiec³⁷, Marcel Eens², Kjell Einar Erikstad³⁸, Silvia Espín³⁹, Damien R. Farine^{40,41,42}, Jordi Figuerola⁴³, Pınar Kavak Gülbeyaz^{18,44}, Arnaud Grégoire⁶, Ian R. Hartley⁴⁵, Michaela Hau³¹, Gergely Hegyi⁴⁶, Sabine Hille⁴⁷, Camilla A. Hinde⁴⁸, Benedikt Holtmann³³, Tatyana Ilyina²⁴, Caroline Isaksson²⁵, Arne Iserbyt², Elena Ivankina⁴⁹, Wojciech Kania⁵⁰, Bart Kempenaers⁵¹, Anvar Kerimov²⁴, Jan Komdeur²⁰, Peter Korsten⁵², Miroslav Král¹¹, Miloš Krist¹¹, Marcel Lambrechts⁶, Carlos E. Lara⁵³, Agu Leivits⁵⁴, András Liker⁵⁵, Jaanis Lodjak⁵⁶, Marko Mägi⁵⁶, Mark C. Mainwaring⁵⁷, Raivo Mänd⁵⁶, Bruno Massa²⁶, Sylvie Massemin¹⁹, Jesús Martínez-Padilla⁵⁸, Tomasz D. Mazgajski³⁷, Adele Mennerat⁵⁹, Juan Moreno⁶⁰, Alexia Mouchet³³, Shinichi Nakagawa³⁶, Jan-Åke Nilsson²⁵, Johan Nilsson²⁵, Ana Cláudia Norte⁶¹, Kees van Oers^{1,62}, Markku Orell⁶³, Jaime Potti¹³, John L. Quinn⁶⁴, Denis Réale⁶⁵, Tone Kristin Reiertsen³⁸, Balázs Rosivall⁴⁶, Andrew F Russel⁶⁶, Seppo Rytkönen⁶³, Pablo Sánchez-Virosta³⁹, Eduardo S. A. Santos⁶⁷, Julia Schroeder⁶⁸, Juan Carlos Senar⁶⁹, Gábor Seress⁵⁵, Tore Slagsvold⁷⁰, Marta Szulkin⁷¹, Céline Teplitsky⁶, Vallo Tilgar⁵⁶, Andrey Tolstoguzov¹⁴, János Török⁴⁶, Mihai Valcu⁵¹, Emma Vatka⁷², Simon Verhulst²⁰, Hannah Watson²⁵, Teru Yuta^{73,74}, José M. Zamora-Marín⁷⁵ & Marcel E. Visser¹

¹ Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Wageningen, The Netherlands

² Department of Biology, University of Antwerp, Antwerp, Belgium

³ Department of Evolutionary Genetics, Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany

4 RSPB Centre for Conservation Science, The Lodge, Sandy, UK
 5 Centre for Research in Animal Behaviour, University of Exeter, Exeter, UK
 6 CEFE, Univ Montpellier, CNRS, EPHE, IRD, Univ Paul Valéry Montpellier 3, Montpellier, France
 7 Edward Grey Institute, Department of Zoology, University of Oxford, Oxford, UK
 8 Department of Biology, University of Turku, Turku, Finland
 9 Kevo Subarctic Research Institute, University of Turku, Turku, Finland
 10 Centre for Biodiversity Dynamics, Department of Biology, Norwegian University of Science and
 Technology, Trondheim, Norway
 11 Department of Zoology, Faculty of Science, Palacký University, Olomouc, Czech Republic
 12 Max Planck Institute of Animal Behavior, Radolfzell, Germany
 13 Department of Ethology and Biodiversity Conservation, Estación Biológica de Doñana-CSIC,
 Seville, Spain
 14 Institute of Biology of the Karelian Research Centre of the Russian Academy of Sciences,
 Petrozavodsk, Russia
 15 Cavanilles Institute of Biodiversity and Evolutionary Biology, University of Valencia, Paterna,
 Spain
 16 Institute of Ornithology, Croatian Academy of Sciences and Arts, Zagreb, Croatia
 17 Universitat Politècnica de València, Gandia, Valencia, Spain
 18 Biodiversity and Conservation Lab, Department of Biology, METU, Ankara, Turkey
 19 Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France
 20 Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen
 21 Institute of Avian Research, Wilhelmshaven, Germany
 22 Institute of Biodiversity, Animal Health & Comparative Medicine, Univ. of Glasgow, Glasgow,
 UK
 23 Department of Animal & Plant Sciences, University of Sheffield, Sheffield, UK
 24 Department of Vertebrate Zoology, Faculty of Biology, Lomonosov Moscow State Univ., Moscow,
 Russia
 25 Department of Biology, Lund University, Lund, Sweden
 26 Stazione Ornitologica, Monreale, Italy
 27 Department STEBICEF, Università degli Studi di Palermo, Palermo, Italy
 28 Institute of Ecology and Botany, Centre for Ecological Research, Vácrátót, Hungary
 29 Station d'Ecologie Théorique et Expérimentale du CNRS (UMR5321), Moulis, France
 30 Institute of Environmental Sciences, Jagiellonian University, Kraków
 31 Max Planck Institute for Ornithology, Evolutionary Physiology Group, Seewiesen, Germany
 32 Lab of Ornithology, Cornell University, Ithaca, USA
 33 Behavioural Ecology, Department of Biology, Ludwig-Maximilians University of Munich,
 Planegg-Martinsried, Germany
 34 CNRS, Department of Biometry & Evolutionary Biology, University of Lyon, Villeurbanne, France
 35 Institute of Environmental Sciences, Jagiellonian University, Krakow, Poland
 36 Evolution & Ecology Research Centre and School of Biological, Earth and Environmental
 Sciences, University of New South Wales, Sydney, Australia
 37 Museum and Institute of Zoology, Polish Academy of Sciences, Warsaw, Poland
 38 Norwegian Institute for Nature Research (NINA), Tromsø, Norway
 39 Area of Toxicology, Department of Health Sciences, University of Murcia, Murcia, Spain
 40 Department of Collective Behavior, Max Planck Institute of Animal Behavior, Konstanz, Germany

41 Department of Biology, University of Konstanz, Konstanz, Germany
 42 Centre for the Advanced Study of Collective Behaviour, University of Konstanz, Konstanz,
 Germany
 43 Department of Wetland Ecology. Estación Biológica de Doñana (CSIC), Sevilla, Spain
 44 Environmental Engineering Department, Hacettepe University, Ankara, Turkey
 45 Lancaster Environment Centre, Lancaster University, Lancaster, UK
 46 Behavioural Ecology Group, Department of Systematic Zoology and Ecology, ELTE Eötvös
 Loránd University, Budapest, Hungary
 47 Institute of Wildlife Biology and Game Management, University of Natural Resources and Life
 Sciences Vienna, Vienna, Austria
 48 Behavioural Ecology Group, Biological Sciences, Anglia Ruskin University, Cambridge, UK
 49 Zvenigorod Biological Station, Faculty of Biology, Lomonosov Moscow State University,
 Moscow, Russia
 50 Ornithological Station, Museum & Institute of Zoology, Polish Academy of Sciences, Gdańsk,
 Poland
 51 Department Behavioural Ecology & Evolutionary Genetics, Max Planck Inst. for Ornithology,
 Seewiesen, Germany
 52 Department of Animal Behaviour, Bielefeld University, Bielefeld, Germany
 53 Department of Zoology, University of Otago, Dunedin, New Zealand
 54 Department of Nature Conservation, Environmental Board, Estonia
 55 MTA-PE Evolutionary Ecology Research Group, University of Pannonia, Veszprém, Hungary
 56 Department of Zoology, Inst. of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia
 57 Division of Biological Sciences, University of Montana, Missoula, USA
 58 Department of Biodiversity Conservation and Ecosystem Restoration, Pyrenean Institute of
 Ecology (CSIC), Jaca, Spain
 59 Department of Biological Sciences, University of Bergen, Bergen, Norway
 60 Museo Nacional de Ciencias Naturales (CSIC), Depart. de Ecología Evolutiva, Madrid, Spain
 61 University of Coimbra, MARE - Marine and Environmental Sciences Centre, Department of Life
 Sciences, Coimbra, Portugal
 62 Behavioural Ecology Group, Department of Animal Sciences, Wageningen University & Research,
 Wageningen, The Netherlands
 63 Ecology and Genetics Research Unit, University of Oulu, Finland
 64 School of Biological, Earth and Environmental Sciences, University College Cork, Ireland
 65 Département des Sciences Biologiques, Université du Québec A Montréal, Montréal, Canada
 66 Centre for Ecology and Conservation, Univ. of Exeter, Tremough Campus, Penryn, Cornwall, UK
 67 BECO do Departamento de Zoologia, Universidade de São Paulo, São Paulo, Brazil
 68 Department of Life Sciences, Imperial College London, Silwood Park Campus, Ascot, UK
 69 Evolutionary and Behavioural Ecology Research Unit, Museu de Ciències Naturals de Barcelona,
 Barcelona, Spain
 70 Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biosciences, University
 of Oslo, Oslo, Norway
 71 Centre of New Technologies, University of Warsaw, Warsaw, Poland
 72 Ecological Genetics Research Unit, Organismal and Evolutionary Biology Research Programme,
 Faculty of Biological & Environmental Sciences, University of Helsinki, Finland
 73 Yamashina Institute for Ornithology, Abiko, Japan
 74 Graduate School of Environment Science, Hokkaido University, Sapporo, Japan

Abstract

1. The integration and synthesis of the data in different areas of science is drastically slowed and hindered by a lack of standards and networking programmes. Long-term studies of individually marked animals are not an exception. These studies are especially important as instrumental for understanding evolutionary and ecological processes in the wild. Further, their number and global distribution provides a unique opportunity to assess the generality of patterns and to address broad-scale global issues (e.g. climate change).
2. To solve data integration issues and enable a new scale of ecological and evolutionary research based on long-term studies of birds, we have created the SPI-Birds Network and Database (www.spibirds.org) – a large-scale initiative that connects data from, and researchers working on, studies of wild populations of individually recognizable (usually ringed) birds. Within year and a half since the establishment, SPI-Birds has recruited over 120 members, and currently hosts data on almost 1,5 million individual birds collected in 80 populations over 2000 cumulative years, and counting.
3. SPI-Birds acts as a data hub and a catalogue of studied populations. It prevents data loss, secures easy data finding, use and integration, and thus facilitates collaboration and synthesis. We provide community-derived data and meta-data standards and improve data integrity guided by the principles of Findable, Accessible, Interoperable, and Reusable (FAIR), and aligned with the existing metadata languages (e.g. ecological meta-data language).

4. The encouraging community involvement stems from SPI-Bird's decentralized approach: research groups retain full control over data use and their way of data management, while SPI-Birds creates tailored pipelines to convert each unique data format into a standard format. We outline the lessons learned, so that other communities (e.g. those working on other taxa) can adapt our successful model. Creating community-specific hubs (such as ours, COMADRE for animal demography, etc.) will aid much-needed large-scale ecological data integration.

Keywords: database | birds | research network | data standards | meta-data standards | long-term studies | data hub | data management | FAIR data

Introduction

The importance of long-term individual-based studies

Long-term individual-based studies of animals in their natural environment underpin our understanding of evolutionary and ecological patterns and processes in wild populations (Clutton-Brock & Sheldon 2010). These studies considerably increase our ability to establish the links among genes, individual traits (including physiology and behaviour), fitness, and the environment (Broggi et al. 2005, Schroeder et al. 2015, Johnston et al. 2016, Bonnet et al 2019). They further document the responses of natural populations to changing environments (Grant & Grant 2002, Réale et al. 2003, Espín et al. 2016, Mennerat et al. 2019, Paniw et al. 2019), and facilitate evidence-based conservation (Stokes 2008, Tylianakis et al. 2008, Vátka et al. 2014, Festa-Bianchet et al. 2019).

The first large-scale individual-based field studies of vertebrates were conducted on birds and birds remain the most commonly studied group (Clutton-Brock & Sheldon 2010, Radschuk et al. 2019). Several types of birds (e.g. hole-nesting passerines, colonially breeding seabirds, or fairy-

wrens) have proven to be highly suitable for long-term individual-based monitoring of reproduction and survival. Some of the longest-running field studies with over 65 years of non-interrupted time series focus on hole-nesting birds (e.g. Kluijver 1951, Lack 1954, Lack 1966, Ahola et al. 2007). Hole-nesters are well suited to detailed study as they often breed at high densities in nest-boxes (Dhondt 2007, Lambrechts et al. 2010), which allows for easy monitoring of the breeding performance (e.g. lay date, clutch size, nesting success) and capture of a large number of birds. Up to now studies cover species with different life-histories over a wide latitudinal and longitudinal range, and in a variety of habitat types, including urban habitats (Andersson et al. 2015, Charmantier et al. 2017, Corsini et al. 2017, Senar et al. 2017, Seress et al. 2018). Importantly, these long-term datasets make it possible to answer questions that were not anticipated at the onset of data collection (e.g. influence of global warming on phenology, Visser et al. 1998; effects of habitat fragmentation, Dhondt 2007).

The main asset of individual-based bird studies is not only the long temporal scale, but also the high degree of spatial replication provided by multiple studies conducted simultaneously (Korsten et al. 2010, Dingemanse et al. 2012). The amount of information available when studies are combined has the potential to bring our understanding of ecological and evolutionary processes to entirely new levels, and has, not surprisingly, led to a number of collaborative projects (e.g. Both et al. 2004, Sæther et al. 2007, Eeva et al. 2011, Laine et al. 2016, Vaugoyeau et al. 2016, Wilkins et al. 2016, Keogan et al. 2018, Samplonius et al. 2018, Loukola et al. 2020), and we provide some examples in more detail in Box 1. This large-scale synthesis (including the meta-analysis context, Siepielski et al. 2017, Culina et al. 2018a, Siepielski et al. 2019) is especially important for capturing the diversity of biological systems and the variation in ecological conditions that are experienced by different populations. Which processes may be described as being general? Which processes can be identified as being more specific to certain environmental conditions? Only when we have answers to these questions, we can make

Accepted Article

predictions and tackle global issues, such as habitat degradation, animal welfare, or global warming, and gain insights into reproducibility of findings based on ecological time-series.

Over time, individual-based studies have become more complex and in addition to data on breeding parameters, other types of data have been collected (e.g. morphological, behavioural, physiological, genetic and genomic). Further, the number of potential relational links to other sources, such as biological samples, climatic data and individual movement data has increased. With the increasing extent and complexity of datasets we urgently need to address data archiving, standards and integration, not only for individual based-studies but in all branches where many independent research groups collect similar but differently managed, and consequently under-exploited, data (the long-tail of science, Box 2, Palmer et al. 2007, Wallis et al. 2013). In these branches, transition to Findable, Accessible, Interoperable, and Reusable (FAIR) data (Wilkinson 2016; Box 2) is more urgent, but also more challenging compared to fields where data standards have been set up at the very start (e.g. genomics).

Scientific collaborations that involve large-scale sharing of standardized data, even when access to data is not fully open but restricted, have been shown to generate significant insights, but we can only guarantee this with adequate mechanisms in place to align, store, and advertise the data that are available for such endeavours. Examples of projects that successfully integrate animal data across a large number of studies are EURING Data Bank (<https://euring.org/>, du Feu et al. 2016) that stores encounter records of ringed birds, Movebank database (<https://www.movebank.org/>, Kranstauber et al. 2011) on animal movement data, and COMADRE database on animal demography (Salguero-Gómez et al. 2016).

Barriers to collaboration

Ideally, data should be openly archived in a way that supports FAIR principles (Wilkinson 2016), and as increasingly mandated by funders (Roche et al. 2014, Culina et al. 2018b). All data should be in a single, standard format, and accompanied by rich meta-data that include the description of the data collection protocols, and support data finding and reuse. In practice, this is difficult to achieve.

The core *cultural/sociological reasons* that prevent open data are the lack of incentives, the fear of being scooped, and worries about losing control and overview over the way the data are interpreted and used (Roche et al. 2014, Evans 2016). The latter is not without good reason: it is easy to misinterpret data collected under specific ecological conditions and to misunderstand how variables were derived (Nelson 2009, Mills et al. 2016). Further, even when researchers are willing to adopt common data standards, they might lack the technical knowledge or time. Yet, inspiring examples of overcoming these barriers exist. For example, all national bird ringing schemes originally used their own data storing format, but ultimately agreed on one common output format, creating the European Union for Bird Ringing (EURING, du Feu et al. 2016). Now, all bird ringing data can be brought together at the European level.

The core *practical obstacles* to effective data reuse and collaboration are the lack of: i) meta-data standards to describe populations, ii) data standards, and iii) a central registry of all the populations (Culina et al. 2018b). To find datasets, researchers commonly search the published literature, then contact data owners (who are not always readily reachable, e.g. if they change institution or retire) to determine whether the data are suited for an intended project and whether their owner is willing to share them. This process can take up to a year, and sometimes it fails (personal experience of the authors). If the data are obtained, the user needs to understand the

specific conditions of data collection (e.g. specific field protocols, ecologically relevant conditions), the data structure and vocabularies. Groups/researchers store data in different types of databases and formats, use different vocabularies to name data elements (e.g. different languages) or different coding for the same data element (e.g. some record hatching date as day 1, others as day 0). Thus, data owners usually require much time to extract and compile the data and provide meta-data to meet the user's needs. This process needs to be repeated for each new collaborative project. Reformatting data is not only time consuming, but may also increase the risk of introducing errors.

Cultural and technical barriers must be solved in parallel; and until open data practices become the norm and researchers recognise their benefits, it is crucial to encourage and enable proper data archiving and establish meta-data and data standards. To achieve FAIR data, and to increase and facilitate collaboration and data synthesis, we created Studies of Populations of Individuals – Birds (SPI – Birds) Network and Database. To overcome cultural barriers, we opted for an approach where data owners can decide to keep full control over the use of their data or can make their data open access. This approach where some data are open access and some are not has also been previously successful with the Movebank, (Kranstauber et al. 2011). Data owners also keep their way of data management (i.e. how they organize their data) and SPI-Birds converts these primary data into a standard (FAIR) format. All meta-data stored by SPI Birds are open access, as well as the code to convert primary data into the standard format.

SPI-Birds: Connecting researchers and data

SPI-Birds Network and Database (www.spibirds.org) is a grassroots initiative that connects researchers working on populations of birds in which individuals are uniquely marked, and thus can be recognized (at capture, or by sight). The main goals are to: (1) increase the coordination

Accepted Article

and collaboration between research groups; (2) host the registry of populations and equalise the visibility among research groups; (3) buffer against data loss and provide long-term access to datasets; (4) ensure data quality and integrity; and (5) facilitate data use, and give appropriate credit for data use. To achieve these goals we: (a) derive meta-data attributes that describe populations (Box 2); (b) centrally archive version-controlled primary data from research groups, with attached conditions of data use; (c) derive data standards with controlled vocabularies and convert primary data format into a standard format; (d) conduct data quality checks; (e) run a series of technical reports on the impact that protocols for data collection may have on derived variables; (f) provide expert advice to researchers setting up new populations; and (g) provide an online interface to find and request data, and maintain outreach activities.

To date, we count more than 120 members from 21 countries, monitoring over 80 populations of 19 species (Fig. 1, Fig. 2A). Currently, the majority of the species are hole-nesting passerines (Table 1), but as a part of our long-term goal, we are actively reaching out to researchers that work on other species groups across the world, and so far have incorporated some of these into the database (e.g. owls, seabirds, dunnocks). The only requirement that needs to be met is that most of the birds in a population can be individually recognized (which is commonly achieved by a metal or coloured rings) and that at least one component of the breeding success of these individuals (e.g. laydate, clutch size) has been measured over at least two years.

[Insert Fig1 here]

Figure 1. A map showing the location of the populations with the data hosted in the SPI-Birds database as of August 2020.

Table 1. Number of unique breeders, breeding attempts, and ringed nestlings per species, as hosted at SPI-Birds Database

Species	N° unique breeders	N° breeding attempts	N° ringed nestlings
<i>Parus major</i>	60,501	90,882	590,157
<i>Cyanistes caeruleus</i>	44,878	59,386	438,840
<i>Ficedula albicollis</i>	35,088	45,933	247,141
<i>Ficedula hypoleuca</i>	22,406	26,099	116,662
<i>Sterna hirundo</i>	3,063	10,097	NA
<i>Poecile montanus</i>	3,000	3,673	12,945
<i>Poecile palustris</i>	2,319	1,537	3,495
<i>Periparus ater</i>	1,462	2,160	13,662
<i>Passer domesticus</i>	1021	2,690	2890
<i>Corvus monedula</i>	852	2,120	NA
<i>Prunella modularis</i>	305	466	432
<i>Lophophanes cristatus</i>	195	164	465
<i>Emberiza melanocephala</i>	186	66	221
<i>Poecile cinctus</i>	184	253	NA
<i>Sitta europea</i>	133	743	456
<i>Parus minor</i>	120	232	745
<i>Strix aluco</i>	63	84	168
<i>Phoenicurus phoenicurus</i>	45	487	1586
<i>Parus varius</i>	10	24	76
Overall	175,831	247,096	1,416,293

Community data standards

To facilitate data compatibility and integration, SPI-Birds has already created data standard for storing breeding-season data on individually monitored birds. This standard format is described in detail on the SPI-Birds GitHub repository (Culina et al. 2019). It is designed to cover the data fields that are common across most individual-based bird studies, and is aligned with the standards suggested by the Ecological Meta-Data Language (EML, Jones et al. 2019) and the principles of FAIR data (Wilkinson 2016). The standard format is dynamic and can be further

extended or adjusted to accommodate the breeding biology (e.g. cooperative breeders) of species yet to be included into the database.

SPI-Birds creates tailored pipelines to convert data from each research group/contributor (i.e. primary data format, Box 2) into the standard format. We hope that this standard format will be adopted by both new and existing research groups to archive their data. The existing groups will be more likely to start using the standard format once their old data have been converted into it by SPI-Birds. We further plan to extend this format (and create new standards) to accommodate other information (e.g. genetic, hormonal, colouration, ecotoxicological, behavioural data). Currently, each population's meta-data clearly indicate whether this additional information has been collected, and the corresponding data can be stored at SPI-Birds (although not yet standardized). For example, physiological or personality data has been collected in almost 30 populations (Fig. 2B)

[Insert Fig2 here]

Figure 2. Summary information on the number of populations hosted at SPI-Birds that (A) collect data on a certain species; (B) collect different types of data on individuals (alongside basic breeding parameters); (C) have been studied for a certain period of time.

Data processing: integration, quality checks, yearly updates

Fig. 3A provides an overview of the SPI-Birds data flow (data collection, standardization, request, and provisioning). First, data owners upload their primary data (Box 2) to SPI-Birds. Primary data come in various storing formats of different complexity, from spreadsheet files (e.g. MS Excel) and simple self-contained databases (e.g. MS Access), to dedicated database servers (e.g. MySQL). Tailored pipelines are then constructed (Fig. 3B) for each dataset to convert primary data into the standard format. The pipeline code is discussed with the data owners (e.g.

discussion of how fields in the primary data are coerced into corresponding fields in the standard format) to ensure maximum accuracy. Pipeline construction sometimes requires several iterations before an accurate pipeline is created. At this point, the pipeline can be confidently applied. The primary data and data in the standard format are stored within the secure SPI-Birds data hub (Fig. 3A), at the file server cluster of the Netherlands Institute of Ecology (NIOO-KNAW) and backed up seven times a week. The pipelines are publicly available via GitHub.

[Insert Fig3 here]

Figure 3. Overview of SPI-Birds infrastructure. (A) Main data workflow that consists of provisioning of primary data, data processing (standardization and quality checks), and data request and provisioning. Panel (B) describes data processing, and panel (C) describes data request and provisioning process. The internal part (not accessible for users) of the SPI-Birds data hub stores versioned data in the primary and the standard format, with an accompanying quality report for each dataset. Users can search meta-data and request data (C) via the external part of the SPI-Bird data hub (i.e. website). Data is sent to the user (if approved by the data owner, or if data are fully open access) in the community standard format, together with the data-quality report(s). Prim. data = data in the primary format, as stored by a data owner; Stand. data = data in the standard format; Qual. report = a report produced by the standard quality check.

Standard quality check is applied to the standardized data. It involves automated checks for missing data, formats of variables (e.g. date, integer), inconsistencies between variables (e.g. false brood assignment), and unexpected values within variables. The output of the standard quality check are two types of flags: ‘warnings’ (i.e. values that are uncommon or unusual) and ‘likely errors’ (i.e. values that should be impossible). In discussion with the data owner, warnings and likely errors are resolved, if at all possible, and the quality check is updated. If data owners decide to address the ‘warnings’ and ‘likely errors’ and update their own primary data, these updated primary data will then be stored in the SPI-Birds data hub, under the version control

system. Finally, any remaining unresolved flagged records appear in the quality check report that is sent to the user. The ‘warnings’ and ‘likely errors’ list is part of the meta-data for each version of the dataset.

For all populations with ongoing data collection primary data are updated to a new version every year and may include not only additional data collected over the additional year but also corrections of errors found in earlier data. We store all versions of the primary data following these yearly updates. This way, we aid to the reproducibility of results based on the version of the data used for the analysis.

Data use: discovery, provisioning, terms of use

Populations hosted at SPI-Birds can be searched via SPI-Birds website based on meta-data (e.g. species studied, country, length of data collection, variables measured). Once the relevant populations have been identified, data can be requested using the SPI-Birds request form (Fig 3C). Unless the data owners have made their data fully open access, data requests are sent for approval to the data owner. If approval is given, standardized data from the requested population(s), accompanied by the standard quality check report, are delivered to the user, and the data owner is informed about the data sharing. When the user requests multiple datasets, all datasets (in a standard format) are compiled and sent to the user. Each dataset comes with a specific terms of data use, and these are stated in its meta-data. To give credit to those who have been collecting/managing the raw data, as a minimal requirement for data use (i.e. even when the data owner does not request any other conditions of data use) we ask that the data owner(s) and funding source(s) that they state in the meta-data, are explicitly acknowledged upon data use (e.g. in the acknowledgment section of an article). We also require acknowledgment of the SPI-Birds Network and Database, and citation of this paper. Further, SPI-Birds encourage citation of the

dataset source (or related publication) via DOIs (digital object identifiers). The detailed Data Access Policy can be found on our website.

Understanding data and their limitations

As discussed above, datasets come with errors and limitations. While SPI-birds increases data integrity and quality, the standard data may still contain errors, and data from different populations might still not be entirely comparable. To enable users to understand how the primary data and standardized data were derived and to highlight potential limitations in the dataset, we provide several documents (as a part of the meta-data). These include the description of the study site (e.g. location, size, habitat type), data collection protocols, and the list of any initial quality checks on the primary data conducted by a data owner. This way, users can better understand how the primary data were derived. Next, we provide a detailed description of decisions and assumptions made during the conversion of data from primary to standard format (with all the pipelines openly available via GitHub), details on quality checks conducted by SPI-Birds and the resulting quality report. Finally, we publish a series of ‘technical reports’ on the SPI-Birds website, where we discuss a range of topics related to methodological conventions (e.g. conversion from one type of tarsus measurement method to another type) and potential biases induced by methodological approaches to data collection (e.g. impact of the frequency at which nests are checked on the estimation of laying date, the impact of nest box design on the vital rates, such as survival of young, Lambrechts et al. 2010).

Lessons learned – creating a community data hub

The need to adopt global meta-data and data standards in ecology and evolution is growing (Poisot et al. 2019, Schneider et al. 2019). We strongly believe, and our example supports, that the best way to achieve the adoption of global standards is to first create standards for well-defined communities (Poisot et al. 2019). When research communities that work on a similar type of data have established their own standards, it becomes easier to scale up to even larger, more global standards (e.g. EML, Jones et al. 2019). Lessons learned from the SPI-Birds example can be useful to research communities where many researchers (groups) collect data of a comparable type (or purpose), but where research protocols and data management are not uniform (i.e. the long-tail of science, Box 2, Palmer et al. 2007, Wallis et al. 2013).

We suggest four key points in establishing a common database and community data standards in the long-tail of science: (1) How to start. Aim to identify researchers/groups that belong to your research community. This is largely a snowballing process – once you locate several members, ask them to identify others. Ideally, organize a kick-off meeting to discuss the aims, distribution of tasks, and further steps. From our experience, it is important to have at least several research groups keen on the project at the start. Further, it is important to consider needs and fears of your research community when deciding on the best working model. For example, our success in mobilizing members largely comes from a decentralized approach: data owners keep full control (i.e. ownership) over their data, and over their data management practices. (2) Keep the community engaged and informed. We found it essential to enable all the members to have the opportunity to contribute to the decisions made. For example, all of our members can provide feedback on any component of the project. Second, it is important, especially at the start, to show that the project is active. We suggest publishing a newsletter every month or two, and creating a social media account. We tweet about each data set we receive, keeping the community informed

of our continuous growth. Third, organise workshops/meetings where the community physically (or virtually) comes together. (3) Funding. Plan to allow for different funding scenarios. We find that it is best to plan finances in steps (if no long-term large funding is available at the very start). Make sure that the first step – what you want to achieve at the minimum – is financially covered at the start. This must include securing a permanent, long-term platform to archive the datasets. After that, plan in five-year (or similar) steps. Here, make sure that in the worst-case scenario (no further funding secured) each step is maintainable with a minimal financial and personnel commitment. For example, our first step was to integrate data on hole-nesting passerines in Eurasia, and this period was financially covered by a grants held by participating individuals, and volunteer contributions from several members. After this initial phase the SPI-Birds database can be kept functional with a minimal investment (e.g. storage capacity). In the next step, we plan to increase our scope. At this stage, our project has already proven successful, which makes it more attractive for longer-term support (e.g. European open science funds). Finally, we ask (but do not mandate) that those whose research plans rely on the collective power of datasets hosted at SPI-Birds to allocate some of their resources to the SPI-Birds initiative.

Vision for an integrated future

SPI-Birds is a large-scale initiative that integrates data on individual-based studies of breeding birds and connects researchers that collect data in these populations. With this paper we also call for additional members to join our fast growing community. To join, please use the contact details as given on the SPI-Birds website (www.spibirds.org). We are inviting contributions from anyone who monitors a population of a bird species, where birds are individually recognizable (usually this would be numbered or colour rings), and where breeding success (at least one component of the breeding success, e.g. clutch size) is recorded over years (at least two years). In

Accepted Article

further developments of the database we plan to: (1) cover additional populations, species, and a wider geographical area; (2) integrate and standardize other data types (e.g. hormonal, behavioural); (3) connect with ongoing centralized efforts to map the full spectrum of different types of data on birds that can complement each other. Here, the main collaborators are scientific groups that centralize the collection of complementary types of data (e.g. Movebank, Kranstauber et al. 2011, Fiedler and Davidson 2012; EURING, du Feu et al. 2016; the great tit HapMap project, Spurgin et al. 2019). Within this scope, we can connect individual-level data hosted at SPI-Birds to other types of data on the same individuals based on their unique ID and provide even more comprehensive information on individuals across their full life-cycle. A second target group are citizen science projects such as Nestkast (de Jong et al. 2018), or the Woodland Trust phenology network (<https://naturescalendar.woodlandtrust.org.uk/>).

SPI-Birds can also serve as a platform to enable better resource allocation between research groups. For example, while a data owner might have the data, they might lack funds to analyse them. On the other hand, a data user might have funds or even apply for funds based on these data. SPI-Birds can thus help pull the resources (data and funds) together, thereby enabling scientific projects, and progress, where it may otherwise be unlikely to occur. We also encourage use of SPI-Birds data in student projects. Finally, during the unforeseen international crisis, such as caused by a novel corona virus during writing of this contribution, SPI-Birds provided an excellent platform to update and document field situations and to mitigate the unbalanced effects of the crisis on research groups. We hope that initiatives such as SPI-Birds can truly help a transition to a new level of ecological synthesis.

Box 1 –Examples of using multiple wild populations

(a) Assessing the ability to substitute space-for-time

Within the scope of understanding and predicting ecological and evolutionary responses to climate change, sampling and studying multiple populations of the same species across latitudinal or altitudinal gradient may provide insights into adaptation to climate variation, if we assume that time can be substituted by space in the processes involved (Phillimore et al. 2010, Blois et al. 2013). For example, Bay et al. (2018) sampled yellow warblers (*Setophaga petechia*) across their breeding range to analyse genomic variation across space and environments (climate, vegetation type, and elevation). Assuming that the current spatial variation in traits of this species may provide information on temporal variation in the future, this study suggested that those yellow warbler populations that have already experienced the largest population declines, require the greatest shifts in allele frequencies to keep pace with future climate change (i.e. are most genetically vulnerable). Similarly, urban-driven evolutionary adaptation is a fascinating process that not only can be followed in time, but also across space, and fostering long-term ecological and evolutionary monitoring in urban areas is key (Szulkin et al. 2020a). In urban evolutionary biology, the spatial dimension is particularly valuable from an empirical perspective as it allows researchers to take advantage of replicated urbanisation gradients, where each city or urban area acts as independent urbanisation replicate (Vaugoyeau et al. 2016, Szulkin et al. 2020b, Santangelo et al. 2020).

(b) Using spatial replication to infer causal relationships

Spatial variation in local temperature trends across long-term population studies allows researchers to separate effects of climate change from confounding correlates which may also be changing over time. As we, unfortunately, have no replicate world without climate change, it is often difficult to attribute changes in local phenotypic distributions to temperature change, rather than to the multitude of other environmental changes that may happen simultaneously. For example, based on local trends of spring temperatures and laying dates in 25 long-term populations of Ficedula flycatchers across Europe, Both et al. (2004) showed that many populations did not exhibit a trend towards earlier breeding, but altogether, there was a clear negative population-level correlation between the trend in laying date and the trend in temperature. In a similar analysis on great tits (*Parus major*) and blue tits (*Cyanistes caeruleus*), such an effect of local temperature was not found: rather, populations originally having a low frequency of second broods did advance, whereas populations in which second broods used to be common did not advance their laying dates (Visser et al., 2003). These examples nicely illustrate how both within- and between-species comparisons of long-term studies deepen our understanding of how organisms may adapt to

climate change.

(c) Comparisons of evolutionary potential

Evolutionary potential depends on the genetic architecture of traits. From a quantitative genetics perspective, this architecture is summarized in G , the additive genetic (co)variance matrix. Comparisons of evolutionary potential across populations or species enable us to evaluate the generality of evolutionary constraints (Agrawal & Stinchcombe, 2009) and to gain insight into the evolution of the underlying genetic architecture (Steppan, Phillips, & Houle, 2002, McGlothlin et al., 2018). For example, using long-term datasets with pedigree information, Teplitsky et al. (2014) assessed the expected constraints on evolutionary responses of morphological traits in ten populations of seven wild bird species. Based on estimated G matrices and selection gradients for four morphological traits, their results suggest that genetic correlations may reduce the expected rate of evolution by 28% on average, even for traits such as morphological traits, that are generally thought to have a high evolutionary potential.

In terms of the evolution of genetic architecture, Delahaie et al. (2017) showed that the genetic architecture of life history and morphological traits is relatively conserved across populations of blue tits inhabiting contrasting habitats. Additionally, Martínez-Padilla et al. (2017) compiled all published estimates of additive genetic variation of morphological traits quantified from 20 long-term and individually-monitored populations of 12 wild European bird species. They found that the evolutionary potential of morphological traits decreases as environmental conditions approaches the extremes, either being favourable or unfavourable. Stronger selection pressures that erode additive genetic variation when environmental conditions were unfavourable or high intraspecific competition in favourable environmental conditions may explain the pattern. These examples illustrate the need of larger-scale studies, both in terms of geography and phylogeny, to fully address the question of the evolution of genetic architecture in wild populations.

(d) Resolving methodological issues

Long-term individual level studies often vary in protocols, applied methodologies and approaches to data collection. Using many long-term datasets may help identify such variation, and point towards those variables that can have potentially significant impacts on how results are interpreted, especially at the between-study level. Møller et al. (2014) targeted one important, strongly varying component of long-term hole-nesting bird studies: nest-box design. Their study included reproductive data of four bird species: blue and great tits, and pied and collared flycatchers. They have found a positive relationship between nest-box floor area and clutch size in great tits, and between box material (wood vs. concrete) and clutch size in blue tits. These results indicate that variation in study design at the between-population level should

always be included as it may prove an important predictor of some of the observed inter-population variation.

Box 2 – Glossary

Individual-based studies of birds – Individual birds are marked with rings engraved with a unique identifying number. Birds are captured (or observed), often over subsequent years, and data on individual characteristics and/or breeding parameters (e.g. laying date, clutch size, number of hatchlings and fledglings, partner) are collected. This information directly links to fitness because it provides data on breeding success and on survival of individuals between years, and thus can be used to study different ecological and evolutionary processes, such as selection on individual traits or population-dependent processes (e.g. density-dependent selection). Other types of data are also increasingly collected, e.g. behavioural, hormonal, genetic or genomic, fine-scale environmental data (including e.g. pollutant data).

Long tail of science – Dispersed scientific research that is conducted by many individual researchers/teams. Data produced in the long tail tend to be small in volume, and less standardized within the same field of study. The majority of scientific funding is spent on this type of research.

Open Data – Data that anyone is free to use, reuse and redistribute — subject, at most, to the requirement to attribute or share-alike, <https://creativecommons.org/licenses/by-sa/2.5/>

FAIR data – FAIR data are equivalent to open data. FAIR data are structured and described in a way that supports their Findability, Accessibility, Interoperability, and Reusability, for both machines and humans.

Meta-data – Data that describe datasets. Meta-data comprise information explaining the purpose and origin of data, methods used to acquire them, the structure of the data, time references, geographical location, brief description of the study site(s), creator, access conditions and terms of use.

Data owner – A person or institution that has collected the raw data and/or is hosting the primary data.

Data user – A person interested in using the data owned by the data owner. Data owners can be data users of someone else's data.

Raw data – Data as collected in the field.

Primary data – Data stored locally by each research group. Primary data might differ from raw data because of (1) errors made during transcribing raw data into primary data or (2) correction of obvious errors in raw data during transcribing them into primary data (3) primary data contain some derivative of

primary data (e.g. average value for a repeated measurement of an individual)

Primary data format – A format in which primary data are stored. This includes the way that data are divided among different tables, the variables recorded, names of these variables, and how values of these variables are expressed.

Standard data format – A format agreed upon within the research community to record and archive data. The standard format defines the way data are organized among different tables, the vocabularies used to describe the data elements (names of the variables), and conventions used to express the values of the variables.

Standard quality check – A range of checks to test the quality and integrity of the primary data converted into the standard format. Each check differentiates between two main types of flags: ‘warnings’ (values that are uncommon or unusual) and ‘likely errors’ (values that seem impossible).

Data hub – A central location to physically store (archive) all data for a certain domain.

Pipeline – A set of code functions and commands used to convert data provided in the primary format into the standard format. A pipeline usually has a hierarchical structure (outputs of one component of the processing sequence are fed to the next step) and often is modular (non-necessary components can be removed or changed to modify the final structure of output data).

Funding

The SPI-Birds have been supported by an NWO personal grant (grant number 016.Veni.181.054) to AC, and a Research Council of Norway grant: 223257 (SFF-III) and 267511 (EVOCLIM).

Acknowledgments

We would like to thank Alastair Wilson and Lisette Senerpont Domis for their valuable comments on the manuscript. We would also like to acknowledge the enormous efforts of all current and past fieldworkers that have contributed to data collection in the populations we list.

Data availability

Data and code to recreate Figure 2 are available from Dryad Digital Repository: <https://doi.org/10.5061/dryad.51c59zw6r> [Culina et al., 2020]

Authors contributions

A. Culina, F. Adriaensen, L.D. Bailey, M.D. Burgess, A. Charmantier, E.F. Cole, T. Eeva, E. Matthysen, C. R. Nater, B.C. Sheldon, B-E. Sæther, S.J.G. Vriend, M.E. Visser conceptualized the ideas. A. Culina wrote the manuscript draft and supervised the project. Z.Zajkova, A.Culina and S. Drobniak produced the Figures. A.Culina and M.E. Visser enabled overall data curation. All the co-authors collected datasets hosted at SPI-Birds, critically reviewed and edited the manuscript.

References

- Agrawal, A. F. & Stinchcombe J.R. How much do genetic covariances alter the rate of adaptation? *Proceedings of the Royal Society B-Biological Sciences* **276**(1659), 1183-1191 (2009).
- Ahola, M.P., Laaksonen, T., Eeva, T. & Lehikoinen, E. Climate change can alter competitive relationships between resident and migratory birds. *Journal of Animal Ecology*, **76**, 1045-1052 (2007).
- Andersson, M. N., et al. Composition of physiologically important fatty acids in great tits differs between urban and rural populations on a seasonal basis. *Frontiers in Ecology and Evolution* **3**, 93 (2015).
- Bay R.A., Harrigan, R.J., Underwood, V.L, Gibbs, H.L., Smith, T.B. & Ruegg. K. Genomic signals of selection predict climate-driven population declines in a migratory bird. *Science* **359**, 83-86 (2018).
- Bonnet, T., et al. The role of selection and evolution in changing parturition date in a red deer population. *PLOS Biol.* **17**, e3000493 (2019).
- Both, C., et al. Large-scale geographical variation confirms that climate change causes birds to lay earlier. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **271**(1549), 1657–1662 (2004).
- Broggi, J., Hohtola, E., Orell, M. & Nilsson, J-Å. Local adaptation to winter conditions in a passerine spreading north: a common-garden approach. *Evolution* **59**, 1600-1603 (2005).
- Charmantier, A., Demeyrier, V., Lambrechts, M., Perret, S. & Grégoire, A. Urbanization Is associated with divergence in pace-of-life in great tits. *Front. Ecol. Evol.* **5**, 53 (2017).
- Class, B., Brommer, J. & van Oers, K. Exploratory behavior undergoes genotype–age interactions in a wild bird. *Ecology and Evolution* **9**, 8987–8994 (2019).
- Clutton-Brock, T. & Sheldon, B.C. Individuals and populations: the role of long-term, individual-based studies of animals in ecology and evolutionary biology. *Trends in Ecology & Evolution* **25**(10), 562–73 (2010).
- Corsini, M., Dubiec, A., Marrot, P. & Szulkin, M. Humans and tits in the city: quantifying the effects of human presence on great tit and blue tit reproductive trait variation. *Frontiers in Ecology and Evolution* **5**, 82 (2017).
- Culina, A., Crowther, T. W., Ramakers, J. J. C., Gienapp, P., & Visser, M. E. How to do meta-analysis of open datasets: comment. *Nature Ecology and Evolution* **2**, 1053-1056 (2018a).
- Culina, A., et al. Navigating the unfolding open data landscape in ecology and evolution. *Nature Ecology & Evolution* **2**(3), 420–26 (2018b).

Culina, A., Bailey, L.D, Vriend, S.J.G, & Visser, M.E. *Standard protocol for the collection of individual level data.*
https://github.com/SPI_Birds/documentation/blob/master/standard_protocol/SPI_Birds_ProtoPro_v1.0.0.pdf (2019).

Culina, A., Zajkova, Z., Vriend, S, Nater, C, Bailey, L. (2020). Data from: Connecting the data landscape of long-term ecological studies: the SPI-Birds data hub. Dryad Digital Repository. <https://doi.org/10.5061/dryad.51c59zw6r>

Delahaie, B., et al. "Conserved G-matrices of morphological and life-history traits among continental and island blue tit populations." *Heredity* **119** (2), 76-87 (2017).

Dhondt, A. A. in *Ecology and behavior of chickadees and titmice: an integrated approach* (KA Otter, Editor), What drives differences between North American and Eurasian tit studies (Oxford University Press, 2007).

Dingemanse, N. J., et al. Variation in personality and behavioural plasticity across four populations of the great tit *Parus major*. *Journal of Animal Ecology* **81**, 116– 126 (2012).

du Feu, C. R., Clark, J. A., Schaub, M., Fiedler, W., & Baillie, S. R. The EURING Data Bank – A critical tool for continental-scale studies of marked birds. *Ringling & Migration* **31**, 1– 18 (2016).

Eeva, T., et al. Geographical trends in the yolk carotenoid composition of the pied flycatcher (*Ficedula hypoleuca*). *Oecologia* **165**, 277-287 (2011).

Espín et al. Tracking pan-continental trends in environmental contamination using sentinel raptors-what types of samples should we use? *Ecotoxicology* **25**(4), 777-801 (2016).

Evans, S.R. Gauging the purported costs of public data archiving for long-term population studies. *PLoS Biol* **14**(4), e1002432 (2016).

Festa-Bianchet, M., Cote, S., Hamel, S. & Pelletier, F. Long-term studies of bighorn sheep and mountain goats reveal fitness costs of reproduction. *J Animal Ecology* **88**, 1118-1133 (2019).

Fiedler W. & Davidson, S. Movebank: eine offene Internetplattform für Tierwanderungsdaten [Movebank: an open internet platform for animal movement data]. *Vogelwarte* **50**(1), 15–20 (2012).

Grant, P. R. & Grant, B. R. Unpredictable evolution in a 30-year study of Darwin's finches. *Science* **296**, 707–117 (2002).

de Jong, M., et al. Timing of avian breeding in an urbanized world. *Ardea* **106**, 31 – 38 (2018).

Johnston, S. E., Béréños, C., Slate, J., & Pemberton, J.M. Conserved genetic architecture underlying individual recombination rate variation in a wild population of Soay sheep (*Ovis aries*). *Genetics* **203**, 583-598 (2016).

Kattge J., et al. TRY – a global database of plant traits. *Global Change Biology* **17**, 2905–2935 (2011).

Keogan, K., et al. Global phenological insensitivity to shifting ocean temperatures among seabirds. *Nature Climate Change* **8**, 313–318 (2018).

Kluijver, H.N..The population ecology of the great tit, *Parus m. major* L. *Ardea* **39**(1-3), 1–135 (1951).

Korsten, P., et al. Association between DRD4 gene polymorphism and personality variation in great tits: a test across four wild populations. *Molecular Ecology* **19**, 832–843 (2010).

Kranstauber, B., et al. The Movebank data model for animal tracking. *Environmental Modelling & Software* **26**(6), 834–835 (2011).

Jones, M.B., O'Brien, M., Mecum, B., Boettiger, C. Schildhauer, M. Et al. Ecological Metadata Language version 2.2.0. KNB Data Repository (2019). doi:10.5063/F11834T2

Lack, D. *The Natural Regulation of Animal Numbers* (Clarendon Press, 1954).

Lack, D. *Population Studies of Birds* (Oxford University Press, 1966).

Laine, V., et al. Evolutionary signals of selection on cognition from the great tit genome and methylome. *Nat. Commun.* **7**, 10474 (2016).

Lambrechts, M.M., et al. The design of artificial nestboxes for the study of secondary hole-nesting birds: a review of methodological inconsistencies and potential biases. *Acta Ornithol.* **45**, 1–26 (2010).

Loukola, O. J., et al. The roles of temperature, nest predators and information parasites for geographical variation in egg covering behaviour of tits (*Paridae*). *Journal of Biogeography* (2020).

Martínez-Padilla, J., et al. Evolvability meets biogeography: evolutionary potential decreases at high and low environmental favourability. *Proceedings of the Royal Society B: Biological Sciences* **284**(1856), 20170516 (2017).

McGlothlin, J. W., et al. Adaptive radiation along a deeply conserved genetic line of least resistance in *Anolis* lizards. *Evolution Letters* **2**(4), 310–322 (2018).

Mennerat, A., Charmantier, A., Perret, P., Hurtrez-Boussès, S. & Lambrechts, M.M. Parasite intensity is driven by temperature in a wild bird. Preprint at: <https://www.biorxiv.org/content/10.1101/323311v1> (2019).

Mills, J.A., et al. Archiving primary data: solutions for long-term studies. *Trends in Ecology & Evolution* **30**(10), 581–89 (2015).

Møller, A. P., et al. Clutch-size variation in Western Palaearctic secondary hole-nesting passerine birds in relation to nest box design. *Methods Ecol Evol.* **5**, 353–362 (2014).

Nelson, B. Empty archives. *Nature* **461**, 160-163 (2009).

Paniw, M.,*Maag, N., Cozzi, G., Clutton-Brock, T., & Ozgul, A. Life history responses of meerkats to seasonal changes in extreme environments. *Science* **363**, 631-635 (2019).

Poisot, T., Bruneau, A., Gonzalez, A., Gravel, D., & Peres-Neto, P. Ecological data should not be so hard to find and reuse. *Trends in Ecology & Evolution* **34**(6), 494–496 (2019).

Radchuk, V., et al. Adaptive responses of animals to climate change are most likely insufficient. *Nature Communications* **10**(1), 1–14 (2019).

Réale, D., McAdam, A., Boutin, S., & Berteaux, D. Genetic and plastic responses of a northern mammal to climate change. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **270**(1515), 591-596 (2003).

Roche, D. G., et al. Troubleshooting public data archiving: suggestions to increase participation. *PLoS Biol* **12**(1), e1001779 (2014).

Sæther, B-E., et al. The extended moran effect and large-scale synchronous fluctuations in the size of great tit and blue tit populations. *Journal of Animal Ecology* **76**, 315-325 (2007).

Salguero-Gómez, R., Jones, O.R., Archer, C.R., Bein, C., de Buhr, H et al. COMADRE: a global data base of animal demography. *Journal of Animal Ecology* **85**, 371-384 (2016).

Salmón, P., Nilsson, J. F., Nord, A., Bensch, S., & Isaksson, C. Urban environment shortens telomere length in nestling great tits, *Parus Major*. *Biology Letters* **12**(6), 20160155 (2016).

Samplonius, J. M., et al. Phenological sensitivity to climate change is higher in resident than in migrant bird populations among European cavity breeders. *Glob. Change Biol.* **24**, 3780– 3790 (2018).

Santangelo, J. S., et al. In: *Urban Evolutionary Biology* (ed. Szulkin M., Munshi-South J., & Charmantier, A.) Urban environments as a framework to study parallel evolution. (Oxford University Press, 2020).

Schneider et al. Towards an ecological trait-data standard. *Methods Ecol. Evol.* **10**, 2006–2019 (2019).

Schroeder, J., Rees, M., Nakagawa, S., & Burke, T. Reduced fitness in progeny from old parents in a wild population. *PNAS* **112**, 4021-4025 (2015).

Senar, J.C., et al. Urban great tits (*Parus major*) show higher distress calling and pecking rates than rural birds across Europe. *Front. Ecol. Evol.* **5**, 163 (2017).

Seress, G., et al. Impact of urbanization on abundance and phenology of caterpillars and consequences for breeding in an insectivorous bird. *Ecological Applications* **28**, 1143-1156 (2018).

Siepielski, A. M., et al. Precipitation drives global variation in natural selection. *Science* **355**(6328), 959-962 (2017).

Siepielski, A. M., et al. No evidence that warmer temperatures are associated with selection for smaller body sizes. *Proceedings of the Royal Society B: Biological Sciences* **286**(1907), 20191332 (2019).

Spurgen et al. The great tit HapMap project: a continental-scale analysis of genomic variation in a songbird. Preprint at: <https://www.biorxiv.org/content/10.1101/561399v1> (2019).

Steppan, S. J., Phillips, P. C., & Houle, D. Comparative quantitative genetics: evolution of the G-matrix. *Trends in Ecology & Evolution*, **17**(7), 320-327 (2002).

Szulkin, M., Munshi-South, J., & Charmantier, A. In: *Urban Evolutionary Biology* (ed. Szulkin, M, Munshi-South J, & Charmantier A.) Introduction. (Oxford University Press, 2020a).

Szulkin, M., et al. In: *Urban Evolutionary Biology* (ed. Szulkin, M, Munshi-South J, & Charmantier A.) How to quantify urbanisation when testing for urban evolution? (Oxford University Press, 2020b).

Teplitsky, C., et al. Assessing multivariate constraints to evolution across ten long-term avian studies. *PLoS ONE* **9**(3), e90444 (2014).

Tylianakis, J.M., Raphael K. Didham, Bascompte J., Wardle, D.A. Global change and species interactions in terrestrial ecosystems. *Ecology Letters*, **11**, 1351–1363 (2008).

Vatka, E., et al. Nest site selection of a primary hole-nesting passerine reveals means to developing sustainable forestry. *Journal of Avian Biology* **45**(2), 187-196 (2014).

Vaugoyeau, E., et al. Interspecific variation in the relationship between clutch size, laying date and intensity of urbanisation in four species of hole-nesting birds. *Ecology & Evolution* **6**, 5907–5920 (2016).


Visser M. E., van Noordwijk, A.J., Tinbergen J. M., & Lessells C. M. Warmer springs lead to mis-timed reproduction in great tits (*Parus major*). *Proc. R. Soc. Lond. B.* **265**, 1867-1870 (1998).

Visser, M. E., et al. Variable responses to large-scale climate change in European *Parus* populations. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **270**(1513), 367–72 (2003).

Wallis, J. C., Rolando E., & Borgman C. L. If we share data, will anyone use them? Data sharing and reuse in the long tail of science and technology. *PLoS ONE* **8**(7), e67332 (2013).


Wilkins, M. R., et al. Phenotypic differentiation is associated with divergent sexual selection among closely related barn swallow populations. *J. Evol. Biol.* **29**, 2410-2421 (2016).

Wilkinson, M. D. The FAIR Guiding Principles for scientific data management and stewardship. *Sci. Data* **2**, 1–9 (2016).




A

Species




B

Type of data collected




C


Length of the study (years)



(A) Main workflow



(B) Data processing workflow



(C) Data request/provisioning workflow

