



HAL
open science

Dynamics of a tracer particle interacting with excitations of a Bose-Einstein condensate

Jonas Lampart, Peter Pickl

► **To cite this version:**

Jonas Lampart, Peter Pickl. Dynamics of a tracer particle interacting with excitations of a Bose-Einstein condensate. 2020. hal-03035282v1

HAL Id: hal-03035282

<https://hal.science/hal-03035282v1>

Preprint submitted on 2 Dec 2020 (v1), last revised 9 Jul 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Dynamics of a tracer particle interacting with excitations of a Bose-Einstein condensate

Jonas Lampart ^{*†}, Peter Pickl^{*‡}

December 2, 2020

Abstract

We consider the quantum dynamics of a large number N of interacting bosons coupled a tracer particle, i.e. a particle of another kind, on a torus. We assume that in the initial state the bosons essentially form a homogeneous Bose-Einstein condensate, with some excitations. With an appropriate mean-field scaling of the interactions, we prove that the effective dynamics for $N \rightarrow \infty$ is generated by the Bogoliubov-Fröhlich Hamiltonian, which couples the tracer particle linearly to the excitation field.

1 Introduction

The reaction of a many-particle system to small perturbations – for example the introduction of a different particle – carries detailed information about its global properties. In physics, the system of a Bose-Einstein condensate and an impurity particle is of particular interest. Apart from the obvious motivation of understanding this state of matter, Bose-Einstein condensates in the laboratory have the advantage that many of their parameters are controllable, which allows for comparison of theory and experiment over a large range parameter values. Systems of Bose-Einstein condensates with few impurities can now be realised using cold atoms and promise numerous applications [44]. Theoretical approaches to their description are discussed in [21].

In this manuscript we will perform a mathematical analysis of the dynamics of such an impurity entering a Bose gas at zero temperature. In keeping with the convention of [13, 10], and since one motivation is to use the impurity to probe the condensate, we will call it “tracer” particle in what follows.

We will show that, for a certain choice of parameters and for a Bose-Einstein condensate of constant density, the tracer particle will effectively interact with the excitations in the gas,

^{*}Ludwig-Maximilians-Universität, Mathematisches Institut, Theresienstr. 39, 80333 München, Germany.

[†]CNRS & Laboratoire Interdisciplinaire Carnot de Bourgogne (UMR 6303), 9. Av. A. Savary, 21078 Dijon Cedex. E-mail: jonas.lampart@u-bourgogne.fr

[‡]E-mail: pickl@math.lmu.de

which will be of finite number and are governed by the dynamics proposed by Bogoliubov in his seminal work [4]. Thus the system is – for a large number of bosons – in good approximation described by the Bogoliubov-Fröhlich Hamiltonian, given in (9) below. Our result is related to the findings in a recent paper by Mysliwy and Seiringer, where spectral properties the same system are analyzed [29].

In order to get an interaction which is governed by the excitations in the condensate we have to assume that the condensate state has constant density – at least in the interaction region of the tracer particle. Otherwise, the effect coming from particles in the condensate state – which is at zero temperature the vast majority – would be dominant. Assuming constant density, this leading contribution of the interaction will also be a constant which simply leads to a global change in phase for the dynamics, and thus does not influence the motion of the tracer. To simplify the technicalities we assume that our system lives on a three dimensional torus of fixed length. However, we expect that this can be generalized. This generalization has to ensure, of course, that density stays constant in the range of the tracer particle for the full time scale of our observation, which amounts to considering gases of large volume.

The configuration space of one particle is the d -dimensional torus of side-length one, $T^d = \mathbb{R}^d/\mathbb{Z}^d$. The wave function of a system of N bosons and one tracer particle in T^d is then an element of

$$\mathcal{H}_N = L^2(T^d) \otimes L^2(T^d)^{\otimes_s N}, \quad (1)$$

where \otimes_s denotes the symmetric tensor product.

The dynamics of the system is described by the Schrödinger equation on \mathcal{H}_N

$$\begin{cases} i \frac{d}{dt} \Psi(t) = H_N \Psi(t) \\ \Psi(0) = \Psi_N, \end{cases} \quad (2)$$

with Hamiltonian

$$H_N = -\frac{1}{2m} \Delta_x - \sum_{j=1}^N \Delta_{y_j} + g_B \sum_{j \neq k=1}^N V(y_j - y_k) + g_I \sum_{j=1}^N W(x - y_j), \quad (3)$$

where x denotes the position of the tracer and y_1, \dots, y_n those of the bosons.

Without loss of generality we can assume that $\int V(x) dx = 0 = \int W(x) dx$. In the general setting all our results are up to a time dependent global phase given by the respective energy shift.

As an N -boson initial state Ψ_N one can think of an N -fold product of the normalized constant on the torus times an L^2 -function on the torus describing the initial state of the tracer. More generally, this could be modified by adding some “excitations”, i.e. factors orthogonal to the constant function, as long as their number stays finite as $N \rightarrow \infty$ (see (20) below for the precise condition).

The coupling constants g_B of the boson interaction and g_I of the interaction with the impurity will depend on the number N of bosons. The precise form of the dependence has to be chosen appropriately to get the desired effect, i.e. a system where the interaction with excitations is of order one.

1.1 Mean-field equations and corrections

The effective description of many-body systems is an old and yet vivid research area of mathematical physics. Hepp [22] and later Ginibre and Velo [14, 15] as well as Spohn [43] proved that the dynamics of a many body Bose gas at zero temperature is, in the weak coupling limit, described by the Hartree equation in the sense that the one-particle marginal density of the N -body wave function converges to a pure state given by the solution of the Hartree equation as N tends to infinity. The latter can be derived by replacing the interaction of the many body system by the respective mean-field, effectively turning the N -body description to a one-body dynamics. Subsequently, many improvements have been found, for example generalizations to a larger class of interactions, and different notions of convergence (see [25] for a review). The dilute (Gross-Pitaevskii) limit, where interactions are of order one but rare, is a variation of this theme that has also been studied extensively (see [1]).

More recently, there have also been many works on the fluctuations around the mean-field behavior. Grillakis, Margetis and Machedon proved the validity of Bogoliubov theory in a weak-coupling situation with mathematical rigor [19, 20, 17]. They showed that the number of excitations is finite in this case and that the convergence of the N -body wave function towards the respective solution of the Bogoliubov time evolution holds in L^2 -norm. This implies also that the errors are small compared to the effects of Bogoliubov's theory. Their results have been generalized by many other authors in different situations [8, 18, 23, 26, 28, 30, 31, 32, 33, 36]. Corrections beyond the Bogoliubov dynamics were studied in [35, 5], including explicit formulas for higher contributions [6].

In parallel to these results concerning the dynamics, there are also numerous results related to the static question, which show closeness of the N -body ground state to the minimizer of an effective functional (see [25]). Bogoliubov theory has also been justified in this context [41, 16, 11, 42, 27, 37, 38, 39, 2], and allows for a resolution of the spectrum close to the ground state [11, 16, 41, 27, 3]. Recently, higher order corrections [7] and the interaction with a tracer particle [29] have also been considered in the static context.

1.1.1 Heuristics of the mean-field scaling

In this article we consider the mean-field scaling for the boson interactions, i.e. the situation where g_B is of order N^{-1} . With a weak coupling of this form we can expect the interaction energy of the N bosons, which is roughly of order $N^2 g_B$, to be of the same order as their kinetic energy. In models for dilute gases this is complemented by a scaling of the potential V_N such that V_N tends to δ weakly as $N \rightarrow \infty$. We will not consider this situation here. We only remark that, in addition to the difficulties in the derivation of the Bogoliubov Hamiltonian, it will be necessary to take into account the renormalization of the limiting model, see remark 2.

As explained above, we are interested in a tracer particle entering the Bose gas. A similar system has already been considered in [10, 9], but we will be interested in a different regime. We want to consider a scaling, respectively coupling, which is such that the interaction of the tracer is with the excitations from the ground state. Since the vast majority of the bosons will be part of the condensate, the leading contribution to the energy of the tracer will generally come from the condensate and this should be of order $g_I N_0 \sim g_I N$, where N_0 is the number of

bosons in the condensate. However, since the condensate density is assumed to be constant, this contribution will be trivial, i.e. a constant. The next order would be the back-reaction on the tracer of small changes to the condensate, i.e. excitations. Since this is a second order effect one can expect it to be of order $g_I^2 N$. We thus assume that g_I is of order $N^{-1/2}$, which will also ensure that the number of excitations coming from the interaction with the tracer is of the same order as the number of excitations we have without tracer particle (i.e. of order one). This is exactly the same scaling as in [29].

The articles [10, 9] considered the different situation where the interaction of the tracer with the Bose gas is such that the density of the gas is changed significantly. To achieve this, the coupling constant between tracer and gas is chosen to be one. Thus the back-reaction of the gas particles and the tracer will be of order N (respectively of the order of the density if the gas has large volume). In order to have a nice limiting regime the mass of the tracer-particle in [10, 9] was assumed to be of order N , resulting in an acceleration of order one.

1.1.2 Heuristics of Bogoliubov theory

We now briefly explain the heuristic derivation of the Bogoliubov-Fröhlich Hamiltonian (see also [21, Sect.3.3]), which we will subsequently make rigorous. Let $\varphi_0 \equiv 1$ denote the constant function on the torus, which we think of as the condensate state, and extend this to a complete orthogonal system $(\varphi_j)_{j \in \mathbb{N}_0}$ of the one-particle Hilbert space $L^2(T^d)$ (of course, one can take the φ_j to be plane waves, but for our purposes this is not necessary and we prefer to retain this generality). Let $a_j^* := a^*(\varphi_j)$ and $a_j := a(\varphi_j)$ be the creation, respectively annihilation, operators for the state φ_j (see [40, Sect.X.7] for definitions of these, and related, objects).

The boson-Hamiltonian with coupling $g_B = N^{-1}$ can then be written as

$$H_B = d\Gamma(-\Delta) + N^{-1} \sum_{j,k,\ell,m=0}^{\infty} V_{jklm} a_j^* a_k^* a_\ell a_m \quad (4)$$

with

$$V_{jklm} = \int_{T^d} \int_{T^d} \bar{\varphi}_j(y) \bar{\varphi}_k(y') V(y - y') \varphi_\ell(y') \varphi_m(y) dy' dy . \quad (5)$$

Bogoliubov's idea [4] is that, since the vast majority of particles will be in the condensate state, the operators a_0^* and a_0 will give contributions of order \sqrt{N} , whereas a_j^* and a_j with $j \geq 1$ give contributions of order one. Consequently, all terms with less than two of the indices j, k, ℓ, m equal to zero are small. Replacing then a_0^* and a_0 by \sqrt{N} in the remaining ones and assuming that V is even and $\int_{T^d} V = 0$ (which eliminates all terms where more than two of the indices j, k, ℓ and m are equal to zero – see also Lemma 1.1 below), one arrives at the Bogoliubov Hamiltonian

$$H^{\text{Bog}} = d\Gamma(-\Delta) + 2 \sum_{j,k=1}^{\infty} V_{j0k0} a_j^* a_k + \sum_{j,k=1}^{\infty} V_{jk00} a_j^* a_k^* + \sum_{j,k=1}^{\infty} V_{00jk} a_j a_k . \quad (6)$$

If we apply the same reasoning to the interaction of the bosons with the tracer, we write

$$\sum_{n=1}^N W(x - y_n) = \sum_{j,k=0}^{\infty} a_j^* a_k (W_x)_{j,k}, \quad (7)$$

with $W_x(y) = W(x - y)$, $(W_x)_{j,k} = \int \bar{\varphi}_j(y) \varphi_k(y) W_x(y) dy$. Using that that that $(W_x)_{0,0}(x) = \int W = 0$, and then setting $g_I = N^{-1/2}$, dropping the terms with $j \neq 0 \neq k$ and replacing a_0^* and a_0 by \sqrt{N} as above, we arrive at (see also Lemma 1.2 below)

$$\sum_{j=1}^{\infty} \left(a_j(W_x)_{0,j} + a_j^*(W_x)_{j,0} \right) = a(W_x) + a^*(W_x). \quad (8)$$

Adding this interaction as well as the kinetic energy of the tracer to the Bogoliubov Hamiltonian, we obtain the Bogoliubov-Fröhlich Hamiltonian

$$H^{\text{BF}} = -\frac{1}{2m} \Delta_x + H^{\text{Bog}} + a^*(W_x) + a(W_x). \quad (9)$$

Remark 1. An important feature of the original Bogoliubov Hamiltonian is that it is a quadratic expression in the operators a, a^* . There is a unitary Bogoliubov transformation U_{Bog} (that amounts to changing the creation/annihilation operators) such that

$$U_{\text{Bog}} H^{\text{Bog}} U_{\text{Bog}}^* = d\Gamma(E) + E_0 \quad (10)$$

for a one-particle operator E and a constant E_0 . The excitations can thus be described by a non-interacting theory and, in the case of the torus, the energy levels can be computed explicitly (see e.g. [29]).

We emphasize that the Bogoliubov-Fröhlich Hamiltonian does not share this feature. Even though the expression (9) seems to be quadratic (i.e. polynomial of degree two), the interaction depends on the position x of the tracer particle, which also appears in the Laplacian. Performing an x -dependent Bogoliubov transformation will thus not yield a simple result (alternatively, introducing a field for the impurities and viewing H^{BF} as the restriction to the one-impurity space, one sees that the interaction is in fact cubic in the creation and annihilation operators, cf. [21, Sect.3.2]). One can, of course, transform H^{BF} using U_{Bog} , which yields an operator of the form

$$(1 \otimes U_{\text{Bog}}) H^{\text{BF}} (1 \otimes U_{\text{Bog}}^*) = -\frac{1}{2m} \Delta_x + d\Gamma(E) + a^*(\tilde{W}_x) + a(\tilde{W}_x) + E_0, \quad (11)$$

with a suitably transformed interaction \tilde{W} (see [21, Sect.3.3] and [29, Eq.(1.10)] for explicit expressions). Operators of the form (11), with general E and W , which feature the interaction of a particle (or several) with a quantum field by a linear coupling have been studied in many variants and are sometimes referred to as Fröhlich or polaron Hamiltonians. They include the original Fröhlich model [12] and the Nelson model [34]. One of the main motivations of the present work was to show rigorously that such a model arises as an effective description of a many-particle system.

Remark 2. If, instead of the mean-field regime, we were to consider a dilute N -particle system (as in [2]) then, as an intermediate step, we would find a Hamiltonian of the same form as H^{BF} , but with $W = W_N$ depending on N and converging to δ weakly as $N \rightarrow \infty$. Since $a^*(\delta)$ is not a densely defined operator, it is not immediately clear how to define the limiting

dynamics. It was shown in [24] that a renormalized version of H^{BF} with $W = \delta$ can be constructed in this case and that its unitary group can be approximated using operators with ultraviolet cut-off, up to a divergent phase. In a dilute system the particle number N would function as an effective ultraviolet cutoff, so we expect that in this case the phase would have to be modified accordingly in order to obtain the effective dynamics generated by the renormalized Bogoliubov-Fröhlich Hamiltonian.

1.2 The condensate-excitation representation

In order to make the heuristics of the previous section rigorous, we will represent functions in \mathcal{H}_N by decomposing them into their components along the (constant) condensate wave-function φ_0 and in the orthogonal complement. The rigorous implementation of this idea was pioneered by Lewin, Nam, Serfaty and Solovej [27] and we closely follow their presentation. It naturally gives rise to Fock space and thus the representation of the excitations by a quantum field.

Recall that $\varphi_0 \equiv 1$ denotes the constant function on T^d and $(\varphi_j)_{j \in \mathbb{N}_0}$ a complete orthonormal system in $L^2(T^d)$. Let

$$P\psi = \int_{T^d} \psi(y) dy$$

be the projection to $\text{span}\{\varphi_0\} =: \mathfrak{H}_0$ in $L^2(T^d)$ and $Q = 1 - P$. The function φ_0 plays the role of the condensate wave-function, while $\mathfrak{H}_+ := QL^2(T^d)$ is the Hilbert space of an excitation. The functions $(\varphi_j)_{j \in \mathbb{N}}$, with $j > 0$, thus form a complete orthonormal system in \mathfrak{H}_+ . Let $\Psi \in L^2(T^d)^{\otimes_s N}$, then we can write

$$\Psi = (P + Q)^{\otimes N} \Psi = \sum_{j=0}^N \varphi_0^{\otimes(N-j)} \otimes_s \Phi^{(j)} \quad (12)$$

with $\Phi^{(j)} \in \mathfrak{H}_+^{\otimes_s j}$. We can thus represent $\Psi \in L^2(T^d)^{\otimes_s N}$ uniquely by the sequence

$$(\Phi^{(n)})_{n \leq N} \in \bigoplus_{n=0}^N \mathfrak{H}_+^{\otimes_s n} =: \mathcal{F}_+^{\leq N}. \quad (13)$$

We denote by $\mathcal{F}_+ = \Gamma(\mathfrak{H}_+)$ the Fock space of excitations, and by $\mathcal{F}_+^{\leq N}$ the truncated Fock space, defined as above.

The N -boson space \mathcal{H}_N also contains an additional factor for the tracer particle. We define

$$\mathcal{H}_+ =: L^2(T^d) \otimes \mathcal{F}_+, \quad \mathcal{H}_+^{\leq N} =: L^2(T^d) \otimes \mathcal{F}_+^{\leq N}. \quad (14)$$

Leaving the first tensor factor untouched, we obtain a unitary

$$U_N : \mathcal{H}_N \rightarrow \mathcal{H}_+^{\leq N} \subset \mathcal{H}_+. \quad (15)$$

By viewing $L^2(T^d)^{\otimes_s N}$ as embedded in the Fock space $\mathcal{F} = \Gamma(L^2(T^d))$, the unitary can be expressed by the formula (see [27, Eq.(4.6)])

$$U_N \Psi = \bigoplus_{j=0}^N Q^{\otimes j} \left(\frac{a_0^{N-j}}{\sqrt{(N-j)!}} \Psi \right). \quad (16)$$

Denote by $\mathcal{N}_+ := d\Gamma(Q)$ the number operator in \mathcal{F}_+ . Then from (16) one easily deduces the following identities for $f, g \in \mathfrak{H}_+$ (see [27, Prop.14]):

$$\begin{aligned}
U_N a_0^* a_0 U_N^* &= N - \mathcal{N}_+ \\
U_N a^*(f) a_0 U_N^* &= a^*(f) \sqrt{N - \mathcal{N}_+} \\
U_N a_0^* a(f) U_N^* &= \sqrt{N - \mathcal{N}_+} a(f) \\
U_N a(f)^* a(g) U_N^* &= a(f)^* a(g)
\end{aligned} \tag{17}$$

Note that on the left hand side we only consider number-preserving combinations of creation and annihilation operators, so these expressions can be viewed as operators on \mathcal{H}_N .

We now apply this transformation to the interaction terms in the Hamiltonian.

Lemma 1.1. *Let $(\varphi_n)_{n \in \mathbb{N}_0}$ be as above and assume that $V \in L^\infty(T^d)$ is an even function with $\int_{T^d} V(y) dy = 0$. The following identity holds*

$$\begin{aligned}
U_N \left(\sum_{j,k=1}^N V(y_j - y_k) \right) U_N^* &= 2 \sum_{j,k \in \mathbb{N}} V_{j0k0} a_j^* (N - \mathcal{N}_+) a_k \\
&+ \sum_{j \in \mathbb{N}} V_{jk00} a_j^* \sqrt{N - \mathcal{N}_+} a_k^* \sqrt{N - \mathcal{N}_+} + \sum_{j \in \mathbb{N}} V_{00jk} \sqrt{N - \mathcal{N}_+} a_j \sqrt{N - \mathcal{N}_+} a_k \\
&+ 2 \sum_{j,k,\ell \in \mathbb{N}} \left(V_{jk\ell 0} a_j^* \sqrt{N - \mathcal{N}_+} a_k^* a_\ell + V_{0jk\ell} a_j^* a_k \sqrt{N - \mathcal{N}_+} a_\ell \right) + \sum_{j,k,\ell,m \in \mathbb{N}} V_{jklm} a_j^* a_k^* a_\ell a_m
\end{aligned}$$

with V_{jklm} defined by (5).

Proof. We write the multiplication operator $\sum_{j,k} V(y_j - y_k)$ on $L^2(T^d)$ as in (4). We then apply the transformation U_N using the identities (17). Note that, since V is even, $V_{0j0k} = V_{j0k0}$ are real. In order to transform the quartic expressions in a_j, a_j^* , these must be grouped into number-preserving pairs, e.g. for $j, k, \ell > 0$,

$$U_N a_j^* a_k^* a_\ell a_0 U_N^* = U_N a_j^* a_0 U_N^* U_N a_k^* a_\ell U_N^* = a_j^* \sqrt{N - \mathcal{N}_+} a_k^* a_\ell. \tag{18}$$

The claim then follows by noting that $V_{jk\ell 0} = V_{k j 0 \ell}$ and that $\int V = 0$ implies $V_{j00k} = 0 = V_{0jk0}$ for $j, k \in \mathbb{N}_0$. \square

Lemma 1.2. *Assume that $W \in L^\infty(T^d)$ with $\int_{T^d} W(y) dy = 0$ and set $W_x(y) := W(x - y)$. The following identity holds*

$$U_N \left(\sum_{j=1}^N W(x - y_j) \right) U_N^* = \sqrt{N - \mathcal{N}_+} a(W_x) + a^*(W_x) \sqrt{N - \mathcal{N}_+} + d\Gamma(QW_x Q).$$

Proof. We can rewrite W_x as in (7). The claim then follows from the identities (17) by noting

that

$$\begin{aligned}
(W_x)_{00} &= PW_xP = \int_{T^d} W(x-y)dy = 0, \\
\sum_{k=1}^{\infty} (W_x)_{k,0} a_k^* &= \sum_{k=1}^{\infty} a^*(\varphi_k) \langle \varphi_k, W_x \varphi_0 \rangle = a^*(QW_x \varphi_0) = a^*(W_x), \\
\sum_{k,\ell=1}^{\infty} a_k^* a_\ell (W_x)_{k,\ell} &= d\Gamma(QW_xQ).
\end{aligned} \tag{19}$$

□

2 Main Result

We assume that the N -particle initial state $\Psi_N \in \mathcal{H}_N$ is a member of a sequence such that

$$\lim_{N \rightarrow \infty} U_N \Psi_N =: \Phi \tag{20}$$

exists in \mathcal{H}_+ . This assumption means that the number of excitations out of the condensate in the initial states Ψ_N remains finite *with probability one* as $N \rightarrow \infty$. Note, however, that the *expected number* of excitations $\langle \mathcal{N}_+ U_N \Psi_N, U_N \Psi_N \rangle$ may diverge.

Recall that the expression of H^{BF} in (9) is given by $H_0 := -\frac{1}{2m}\Delta_x + d\Gamma(-\Delta)$ plus several terms that are linear or quadratic in creation and annihilation operators. If $V, W \in L^2(T^d)$, the latter are bounded relative to \mathcal{N}_+ (see Proposition A.3). Consequently, H^{BF} is a well defined operator on $D(H_0)$, since on \mathcal{F}_+

$$d\Gamma(-\Delta) \geq 4\pi^2 \mathcal{N}_+. \tag{21}$$

In fact, H^{BF} is essentially self-adjoint on $D(H_0)$, see Proposition A.3. Thus the time evolution under H^{BF} , $e^{-iH^{\text{BF}}t}\Phi$, is well-defined and we may compare it to the time evolution with H_N using U_N .

If we make the assumption that $\Phi \in D(\mathcal{N}_+)$, then we obtain a quantitative estimate proving the closeness of the two evolutions for large N .

Theorem 2.1. *Assume that $V, W \in L^\infty(T^d)$ with $V(-y) = V(y)$ and $\int_{T^d} V = 0 = \int_{T^d} W$. There exists $v > 0$ such that for all $\Phi \in D(\mathcal{N}_+)$ and every sequence $\Psi_N \in \mathcal{H}_N$, $N \in \mathbb{N}$, with $\lim_{N \rightarrow \infty} U_N \Psi_N = \Phi$ in \mathcal{H}_+ there is $K > 0$ such that for all $N \in \mathbb{N}$ and $t \in \mathbb{R}$*

$$\|U_N e^{-iH_N t} \Psi_N - e^{-iH^{\text{BF}}t} \Phi\|_{\mathcal{H}_+} \leq K e^{v|t|} N^{-1/4}.$$

Note that in this statement only K depends on the initial states. By density of $D(\mathcal{N}_+)$ in \mathcal{H}_+ , the statement can be extended to any $\Phi \in \mathcal{H}_+$, but without an explicit rate of convergence.

Theorem 2.2. *Assume that $V, W \in L^\infty(T^d, \mathbb{R})$ with $V(-y) = V(y)$ and $\int_{T^d} V = 0 = \int_{T^d} W$. Then for every $\Phi \in \mathcal{H}_+$ and every sequence Ψ_N , $N \in \mathbb{N}$, with $\lim_{N \rightarrow \infty} U_N \Psi_N = \Phi$ we have*

$$\lim_{N \rightarrow \infty} \|U_N e^{-iH_N t} \Psi_N - e^{-iH^{\text{BF}}t} \Phi\|_{\mathcal{H}_+} = 0$$

uniformly in t on compact subsets of \mathbb{R} .

Proof. This follows from Theorem 2.1 above by an approximation argument. Let $\Phi \in \mathcal{H}_+$ be given and $T > 0$, $\varepsilon > 0$. Let $\tilde{\Phi} \in D(\mathcal{N}_+)$ with $\|\Phi - \tilde{\Phi}\|_{\mathcal{H}_+} < \varepsilon$ and set $\tilde{\Psi}_N = U_N^* \tilde{\Phi}^{\leq N}$. Since $\lim_{N \rightarrow \infty} U_N \Psi_N = \Phi$, we have for sufficiently large N

$$\|\Psi_N - \tilde{\Psi}_N\|_{\mathcal{H}_N} \leq \|U_N \Psi_N - \Phi\|_{\mathcal{H}_+} + \|\Phi - \tilde{\Phi}\| + \|\tilde{\Phi}^{> N}\| < 3\varepsilon. \quad (22)$$

By unitarity of the time evolutions we thus have

$$\|U_N e^{-iH_N t} \Psi_N - e^{-iH^{\text{BF}} t} \Phi\|_{\mathcal{H}_+} < 4\varepsilon + \|U_N e^{-iH_N t} \tilde{\Psi}_N - e^{-iH^{\text{BF}} t} \tilde{\Phi}\|_{\mathcal{H}_+}. \quad (23)$$

By Theorem 2.1 the last term is bounded for $|t| \leq T$ by

$$\|U_N e^{-iH_N t} \tilde{\Psi}_N - e^{-iH^{\text{BF}} t} \tilde{\Phi}\|_{\mathcal{H}_+} \leq K e^{vT} N^{-1/4}, \quad (24)$$

so it is smaller than ε for sufficiently large N . \square

The proof of Theorem 2.1 will be the content of the next section.

3 Proof of convergence

As a first step we will prove that the time evolution $U_N e^{-iH_N t} U_N^*$ is well approximated by the time evolution generated by a truncation of $U_N H_N U_N^*$, where terms that are more than quadratic in the creation and annihilation operators are neglected. We collect the leading-order terms in an auxiliary operator H^{aux} . This operator acts on $\mathcal{H}_+^{\leq N}$ and is naturally extended to \mathcal{H}_+ by zero. Since $\langle \varphi_k, \Delta \varphi_0 \rangle = 0 = \langle \varphi_0, \Delta \varphi_k \rangle$ for $k \in \mathbb{N}_0$, we have

$$-U_N \sum_{j=1}^N \Delta_{y_j} U_N^* = d\Gamma^{\leq N}(-\Delta),$$

where $d\Gamma^{\leq N}(-\Delta)$ denotes the projection of $d\Gamma(-\Delta)$ to $\mathcal{H}_+^{\leq N}$. In view of Lemmas 1.2, 1.1, we define H_N^{aux} by the expression

$$\begin{aligned} H_N^{\text{aux}} := & -\frac{1}{2m} \Delta_x + d\Gamma^{\leq N}(-\Delta) + \frac{2}{N} \sum_{j,k>1} V_{j0k0} a_j^* (N - \mathcal{N}_+) a_k \\ & + \frac{1}{N} \sum_{j,k>1} V_{jk00} a_j^* \sqrt{N - \mathcal{N}_+} a_k^* \sqrt{N - \mathcal{N}_+} \\ & + \frac{1}{N} \sum_{j,k>1} V_{00jk} \sqrt{N - \mathcal{N}_+} a_k \sqrt{N - \mathcal{N}_+} a_j \\ & + \frac{1}{\sqrt{N}} a^*(W_x) \sqrt{N - \mathcal{N}_+} + \frac{1}{\sqrt{N}} \sqrt{N - \mathcal{N}_+} a(W_x) \end{aligned} \quad (25)$$

on $D(H_N^{\text{aux}}) = U_N D(H_N) \subset \mathcal{H}_+^{\leq N}$. Note that H_N^{aux} is self-adjoint by Lemma A.2 and that H^{BF} is obtained from H_N^{aux} by simply replacing $\sqrt{1 - \mathcal{N}_+/N}$ by one and extending to \mathcal{H}_+ .

To prove closeness of $e^{-iH_N^{\text{aux}} t}$ and $U_N e^{-iH_N t} U_N^*$, and later $e^{-iH^{\text{BF}} t}$ and $e^{-iH_N^{\text{aux}} t}$, the crucial ingredient is control of the number of excitations for all times.

Lemma 3.1. For any $\Phi \in D(\mathcal{N}_+)$ set

$$\begin{aligned}\alpha(t) &:= \|\mathcal{N}_+ U_N e^{-iH_N t} U_N^* \Phi^{\leq N}\|^2 \\ \alpha^{\text{aux}}(t) &:= \|\mathcal{N}_+ e^{-iH_N^{\text{aux}} t} \Phi^{\leq N}\|^2 \\ \alpha^{\text{BF}}(t) &:= \|\mathcal{N}_+ e^{-iH^{\text{BF}} t} \Phi\|^2.\end{aligned}$$

There exists a constant v such that for all $\Phi \in D(\mathcal{N}_+)$, $N \in \mathbb{N}$ and $t \in \mathbb{R}$ we have

$$\alpha^\bullet(t) \leq \alpha^\bullet(0) e^{v|t|}$$

for $\bullet \in \{\emptyset, \text{aux}, \text{BF}\}$.

Proof. Since $\Phi \in D(\mathcal{N}_+)$, the statement clearly holds for $t = 0$. We will use Grönwall's Lemma to obtain a bound for all $t \geq 0$ (the proof for $t \leq 0$ is the same).

We first prove the claim for α and α^{aux} (since H_N^{aux} is obtained from $U_N H_N U_N^*$ by dropping some terms the proof for α^{aux} is contained in the one for α). We have for $\Phi \in U_N D(H_N)$

$$\begin{aligned}\frac{d\alpha}{dt}(t) &= \frac{d}{dt} \left\langle U_N e^{-iH_N t} U_N^* \Phi^{\leq N}, \mathcal{N}_+^2 U_N e^{-iH_N t} U_N^* \Phi^{\leq N} \right\rangle \\ &= \left\langle U_N e^{-iH_N t} U_N^* \Phi^{\leq N}, i[U_N H_N U_N^*, \mathcal{N}_+^2] U_N e^{-iH_N t} U_N^* \Phi^{\leq N} \right\rangle.\end{aligned}\quad (26)$$

With $[U_N H_N U_N^*, \mathcal{N}_+^2] = \mathcal{N}_+[U_N H_N U_N^*, \mathcal{N}_+] + [U_N H_N U_N^*, \mathcal{N}_+] \mathcal{N}_+$ and Cauchy-Schwarz we obtain

$$\left| \frac{d\alpha}{dt}(t) \right| \leq 2\sqrt{\alpha(t)} \|[U_N H_N U_N^*, \mathcal{N}_+] e^{-iH_N t} U_N^* \Phi^{\leq N}\|.\quad (27)$$

We can thus prove the claim by bounding the latter norm in terms of α .

To do this, we start with the terms from H_N^{aux} . Using that

$$[a_j^* a_j, a_k^* a_\ell^*] = (\delta_{jk} + \delta_{j\ell}) a_j^* a_j^* \quad (28)$$

$$[a_j^* a_j, a_k a_\ell] = -(\delta_{jk} + \delta_{j\ell}) a_j a_j \quad (29)$$

and $\mathcal{N}_+ = \sum_{j=1}^{\infty} a_j^* a_j$ we find

$$\left[\mathcal{N}_+, \sum_{j,k=1}^{\infty} V_{jk00} a_j^* a_k^* \right] = 2 \sum_{j,k=1}^{\infty} V_{jk00} a_j^* a_k^* \quad (30)$$

$$\left[\mathcal{N}_+, \sum_{j,k=1}^{\infty} V_{00jk} a_j a_k \right] = -2 \sum_{j,k=1}^{\infty} V_{00jk} a_j a_k. \quad (31)$$

We thus have

$$\begin{aligned}[\mathcal{N}_+, H_N^{\text{aux}}] &= \frac{2}{N} \sum_{j,k=1}^{\infty} V_{jk00} a_j^* \sqrt{N - \mathcal{N}_+} a_k^* \sqrt{N - \mathcal{N}_+} \\ &\quad - \frac{2}{N} \sum_{j,k=1}^{\infty} V_{00jk} \sqrt{N - \mathcal{N}_+} a_k \sqrt{N - \mathcal{N}_+} a_j \\ &\quad + \frac{1}{\sqrt{N}} a^*(W_x) \sqrt{N - \mathcal{N}_+} - \frac{1}{\sqrt{N}} \sqrt{N - \mathcal{N}_+} a(W_x).\end{aligned}\quad (32)$$

This operator is bounded relative to \mathcal{N}_+ by Lemma A.1 and (59), which gives the desired bound. The expression for $U_N H_N U_N^*$ also contains the term $N^{-1/2} d\Gamma(QW_x Q)$ from the interaction (see Lemma 1.2), which commutes with \mathcal{N}_+ . Additionally, there are the terms from the boson interaction (see Lemma 1.1)

$$\frac{2}{N} \sum_{j,k,\ell=1}^{\infty} \left(V_{jk\ell 0} a_j^* \sqrt{N - \mathcal{N}_+} a_k^* a_\ell + V_{0jk\ell} a_j^* a_k \sqrt{N - \mathcal{N}_+} a_\ell \right), \quad (33)$$

$$\frac{1}{N} \sum_{j,k,\ell,m=1}^{\infty} V_{jk\ell m} a_j^* a_k^* a_\ell a_m. \quad (34)$$

The latter also commutes with \mathcal{N}_+ , so it remains to bound the commutator of \mathcal{N}_+ with the first line. We have from the canonical commutation relations (for the first term – the second one yields minus the adjoint)

$$\left[\mathcal{N}_+, \frac{2}{N} V_{jk\ell 0} a_j^* \sqrt{N - \mathcal{N}_+} a_k^* a_\ell \right] = -\frac{2}{N} \sum_{j,k,\ell=1}^{\infty} V_{jk\ell 0} a_j^* \sqrt{N - \mathcal{N}_+} a_k^* a_\ell. \quad (35)$$

In order to bound this operator, let $\Psi, \Xi \in \mathcal{H}_+^{\leq N}$, and rewrite (with $V_y(\cdot) := V(y \cdot)$, cf. (19))

$$\begin{aligned} & \sum_{j,k,\ell=1}^{\infty} \langle \Psi, V_{jk\ell 0} a_j^* \sqrt{N - \mathcal{N}_+} a_k^* a_\ell \Xi \rangle \\ &= \int_{T^d} \sum_{j,k,\ell=1}^{\infty} \bar{\varphi}_j(y) \langle \sqrt{N - \mathcal{N}_+} a_j \Psi, (V_y)_{k\ell} a_k^* a_\ell \Xi \rangle dy \\ &= \int_{T^d} \left\langle \sqrt{N - \mathcal{N}_+} \sum_{j=1}^{\infty} \varphi_j(y) a_j \Psi, d\Gamma(QV_y Q) \Xi \right\rangle dy. \end{aligned} \quad (36)$$

By the Cauchy-Schwarz inequality and the fact that $\|N^{-1/2} \sqrt{N - \mathcal{N}_+}\|_{\mathcal{H}_+^{\leq N} \rightarrow \mathcal{H}_+^{\leq N}} \leq 1$ we thus have

$$\begin{aligned} & \frac{1}{N} \left| \sum_{j,k,\ell=1}^{\infty} \langle \Psi, V_{jk\ell 0} a_j^* \sqrt{N - \mathcal{N}_+} a_k^* a_\ell \Xi \rangle \right| \\ & \leq \left(\int_{T^d} N^{-1} \left\| \sum_{j=1}^{\infty} \varphi_j(y) a_j \Psi \right\|^2 dy \right)^{1/2} \left(\int_{T^d} \left\| d\Gamma(QV_y Q) \Xi \right\|^2 dy \right)^{1/2}. \end{aligned} \quad (37)$$

Now

$$\int_{T^d} \left\| d\Gamma(QV_y Q) \Xi \right\|^2 dy \leq \int_{T^d} \|QV_y Q\|_{\mathfrak{H}_+ \rightarrow \mathfrak{H}_+}^2 \|\mathcal{N}_+ \Xi\|^2 \leq \|V\|_{L^\infty}^2 \|\mathcal{N}_+ \Xi\|^2, \quad (38)$$

and since $\sum_{j=1}^{\infty} \varphi_j(y) \varphi_j(y')$ is the integral kernel of Q , which acts on \mathfrak{H}_+ as the identity, we have for $n < N$

$$\int_{T^d} N^{-1} \left\| \sum_{j=1}^{\infty} \varphi_j(y) a_j \Psi^{(n+1)} \right\|_{\mathfrak{H}_+^{\otimes s_n}}^2 dy = \int_{T^d} N^{-1} \left\| \sqrt{n+1} \Psi^{(n+1)}(y, \cdot) \right\|_{\mathfrak{H}_+^{\otimes s_n}}^2 dy \leq \|\Psi^{(n+1)}\|_{\mathfrak{H}_+^{\otimes s(n+1)}}^2. \quad (39)$$

This proves that the commutator (35) is \mathcal{N}_+ -bounded uniformly in N , so, by our earlier reasoning, there is a constant C such that

$$\frac{d\alpha}{dt}(t) \leq \left| \frac{d\alpha}{dt}(t) \right| \leq C\alpha(t).$$

Thus by Grönwall's Lemma $\alpha(t) \leq \alpha(0)e^{Ct}$, which proves the claim for $\alpha, \alpha^{\text{aux}}$.

The proof for α^{BF} follows from the same argument, since $[\mathcal{N}_+, H^{\text{BF}}]$ is \mathcal{N}_+ -bounded by the identities (30), (31) and Lemma A.1. Taking the growth rate v to be the maximum of the constants for $\alpha, \alpha^{\text{BF}}$ proves the claim. \square

3.1 Proof of Theorem 2.1

Lemma 3.2. *Let \mathcal{H} be a Hilbert space and $D \subset \mathcal{H}$ a dense subspace. If $(H_1, D), (H_2, D)$ are self-adjoint operators, then for $\Psi_j \in D, j = 1, 2$, and $t \geq 0$*

$$\|e^{-iH_1 t}\Psi_1 - e^{-iH_2 t}\Psi_2\|_{\mathcal{H}}^2 \leq \|\Psi_1 - \Psi_2\|_{\mathcal{H}}^2 + 2 \int_0^t \left| \langle e^{-iH_1 s}\Psi_1, (H_1 - H_2)e^{-iH_2 s}\Psi_2 \rangle \right| ds.$$

Proof. We have

$$\frac{d}{dt} \|e^{-iH_1 t}\Psi_1 - e^{-iH_2 t}\Psi_2\|_{\mathcal{H}}^2 = -2\text{Re}\langle e^{-iH_1 t}\Psi_1, i(H_1 - H_2)e^{-iH_2 t}\Psi_2 \rangle, \quad (40)$$

so the claim follows from the fundamental theorem of calculus. \square

Lemma 3.3. *Assume the hypothesis of Theorem 2.2 and let $\Phi \in D(\mathcal{N}_+)$, $\Psi_N = U_N^* \Phi^{\leq N}$. There exists $K > 0$ such that for all $N \in \mathbb{N}$ and $t \in \mathbb{R}$*

$$\|e^{-iH_N t}\Psi_N - U_N^* e^{-iH_N^{\text{aux}} t} U_N \Psi_N\| \leq K N^{-1/4} e^{v|t|},$$

where v is the constant of Lemma 3.1.

Proof. First assume that $\Psi_N \in D(H_N) = U_N^* D(H^{\text{aux}})$ (see Lemma A.2). Using Lemma 3.2 it follows that

$$\begin{aligned} & \|e^{-iH_N t}\Psi_N - U_N^* e^{-iH_N^{\text{aux}} t} U_N \Psi_N\|^2 \\ & \leq 2 \int_0^t \left| \langle e^{-iH_N s}\Psi_N, (H_N - U_N^* H^{\text{aux}} U_N) U_N^* e^{-iH_N^{\text{aux}} s} U_N \Psi_N \rangle \right| ds \\ & = 2 \int_0^t \left| \langle U_N e^{-iH_N s}\Psi_N, (U_N H_N U_N^* - H^{\text{aux}}) e^{-iH_N^{\text{aux}} s} \Phi^{\leq N} \rangle \right| ds. \end{aligned} \quad (41)$$

In view of Lemmas 1.2, 1.1, we have

$$U_N H_N U_N^* - H^{\text{aux}} = \frac{1}{\sqrt{N}} d\Gamma(QW_x Q) \quad (42)$$

$$+ \frac{2}{N} \sum_{j,k,\ell=1}^{\infty} \left(V_{j k \ell 0} a_j^* \sqrt{N - \mathcal{N}_+} a_k^* a_{\ell} + V_{0 j k \ell} a_j^* a_k \sqrt{N - \mathcal{N}_+} a_{\ell} \right) \quad (43)$$

$$+ \frac{1}{N} \sum_{j,k,\ell,m=1}^{\infty} V_{j k \ell m} a_j^* a_k^* a_{\ell} a_m. \quad (44)$$

Since $d\Gamma(QW_xQ)$ is \mathcal{N}_+ -bounded, we have

$$\left| \langle U_N e^{-iH_N t} \Psi_N, d\Gamma(QW_xQ) e^{-iH_N^{\text{aux}} t} \Phi^{\leq N} \rangle \right| \leq \|W\|_{L^\infty} \|\Psi_N\| \alpha^{\text{aux}}(t). \quad (45)$$

The first term in (43) satisfies, by the reasoning of (37),

$$\begin{aligned} & \left| \frac{2}{N} \left| \sum_{j,k,\ell=1}^{\infty} \langle U_N e^{-iH_N t} \Psi_N, V_{j k \ell 0} a_j^* \sqrt{N - \mathcal{N}_+} a_k^* a_\ell e^{-iH_N^{\text{aux}} t} \Phi^{\leq N} \rangle \right| \right| \\ & \leq \frac{2}{\sqrt{N}} \|V\|_{L^\infty} \|\sqrt{\mathcal{N}_+} U_N e^{-iH_N t} \Psi_N\| \|\mathcal{N}_+ e^{-iH_N^{\text{aux}} t} \Phi^{\leq N}\| \\ & \leq \frac{2}{\sqrt{N}} \|V\|_{L^\infty} (\|\Psi_N\| + \alpha(t)) \alpha^{\text{aux}}(t). \end{aligned} \quad (46)$$

The adjoint term from (43) satisfies the same bound.

The quartic term (44) can be bounded using similar arguments. First, we may rewrite

$$\begin{aligned} & \sum_{j,k,\ell,m=1}^{\infty} \langle U_N e^{-iH_N t} \Psi_N, V_{j k \ell m} a_j^* a_k^* a_\ell a_m e^{-iH_N^{\text{aux}} t} \Phi^{\leq N} \rangle \\ & = \sum_{j,m=1}^{\infty} \int_{T^d} \overline{\varphi_j}(y) \varphi_m(y) \langle a_j U_N e^{-iH_N t} \Psi_N, d\Gamma(QV_yQ) a_m e^{-iH_N^{\text{aux}} t} \Phi^{\leq N} \rangle. \end{aligned} \quad (47)$$

Then, using the Cauchy-Schwarz inequality and the same reasoning as in (38), (39),

$$\begin{aligned} & \left| \frac{1}{N} \left| \sum_{j,k,\ell,m=1}^{\infty} \langle U_N e^{-iH_N t} \Psi_N, V_{j k \ell m} a_j^* a_k^* a_\ell a_m e^{-iH_N^{\text{aux}} t} \Phi^{\leq N} \rangle \right| \right| \\ & \leq \frac{1}{N} \left(\int_{T^d} \left\| (\mathcal{N}_+ + 1)^{1/2} \sum_{j=1}^{\infty} \varphi_j(y) a_j U_N e^{-iH_N t} \Psi_N \right\|^2 dy \right)^{1/2} \\ & \quad \times \left(\int_{T^d} \left\| d\Gamma(QV_yQ) (\mathcal{N}_+ + 1)^{-1/2} \sum_{m=1}^{\infty} \varphi_m(y) a_m e^{-iH_N^{\text{aux}} t} \Phi^{\leq N} \right\|^2 dy \right)^{1/2} \\ & \leq \frac{1}{N} \|V\|_{L^\infty} \|\mathcal{N}_+ U_N e^{-iH_N t} \Psi_N\| \|\mathcal{N}_+ e^{-iH_N^{\text{aux}} t} \Phi^{\leq N}\| = \frac{1}{N} \|V\|_{L^\infty} \alpha(t) \alpha^{\text{aux}}(t). \end{aligned} \quad (48)$$

Inserting these estimates into (41) we find by application of Lemma 3.1

$$\begin{aligned} & \left\| e^{-iH_N t} \Psi_N - U_N^* e^{-iH_N^{\text{aux}} t} U_N \Psi_N \right\|^2 \\ & \leq N^{-1/2} \int_0^t 2 \left(\|W\|_{L^\infty} + \|V\|_{L^\infty} (4\|\Psi_N\| + 5\alpha(s)) \right) \alpha^{\text{aux}}(s) ds \\ & \leq N^{-1/2} \int_0^t 2 \left(\|W\|_{L^\infty} + \|V\|_{L^\infty} (4\|\Psi_N\| + 5\|\mathcal{N}_+ U_N \Psi_N\| e^{vs}) \right) \|\mathcal{N}_+ U_N \Psi_N\| e^{vs} ds \\ & \leq N^{-1/2} K^2 e^{2vt} \end{aligned} \quad (49)$$

for an appropriate constant K . This proves the claim for $\Psi_N \in D(H_N)$. Since the right hand side of (49) depends continuously on $U_N \Psi_N \in \mathcal{H}_+^{\leq N}$, the claim for general $\Psi_N = U_N^* \Phi^{\leq N}$ follows by density. \square

To complete the proof of Theorem 2.1, it remains to prove the following Lemma on the approximation of $e^{-iH_N^{\text{aux}}t}$ by $e^{-iH^{\text{BF}}t}$, which essentially amounts to removing the restriction to $\mathcal{H}_+^{\leq N}$.

Lemma 3.4. *Assume the hypothesis of Theorem 2.2 and let $\Phi \in D(\mathcal{N}_+)$. There exists a constant K such that for all $N \in \mathbb{N}$ and $t \in \mathbb{R}$*

$$\|e^{-iH_N^{\text{aux}}t}\Phi^{\leq N} - e^{-iH^{\text{BF}}t}\Phi\| \leq Ke^{v|t|}N^{-1/4},$$

where v is the constant of Lemma 3.1.

Proof. Assume first that $\Phi \in D(H_0) \subset D(H^{\text{BF}})$ (see Proposition A.3), so $\Phi^{\leq N} \in D(H_N^{\text{aux}})$ by Lemma A.2. Note that

$$\|\Phi - \Phi^{\leq N}\|^2 = \sum_{n=N+1}^{\infty} \|\Phi^{(n)}\|^2 \leq (N+1)^{-2} \|\mathcal{N}_+ \Phi\|^2, \quad (50)$$

so, by Lemma 3.2, we have

$$\|e^{-iH_N^{\text{aux}}t}\Phi^{\leq N} - e^{-iH^{\text{BF}}t}\Phi\|^2 \leq (N+1)^{-2} \|\mathcal{N}_+ \Phi\|^2 + \int_0^t \left| \left\langle e^{-iH_N^{\text{aux}}s}\Phi^{\leq N}, (H_N^{\text{aux}} - H^{\text{BF}})e^{-iH^{\text{BF}}s}\Phi \right\rangle \right| ds. \quad (51)$$

The difference of the operators can be written as

$$\begin{aligned} H_N^{\text{aux}} - H^{\text{BF}} &= -\frac{2}{N} \sum_{j,k>1} V_{j0k0} a_j^* \mathcal{N}_+ a_k \\ &+ \frac{1}{N} \sum_{j,k>1} V_{jk00} a_j^* (\sqrt{N - \mathcal{N}_+} - \sqrt{N}) a_k^* (\sqrt{N - \mathcal{N}_+} - \sqrt{N}) \\ &+ \frac{1}{N} \sum_{j,k>1} V_{00jk} (\sqrt{N - \mathcal{N}_+} - \sqrt{N}) a_k (\sqrt{N - \mathcal{N}_+} - \sqrt{N}) a_k \\ &+ \frac{1}{\sqrt{N}} a^*(W_x) (\sqrt{N - \mathcal{N}_+} - \sqrt{N}) + \frac{1}{\sqrt{N}} (\sqrt{N - \mathcal{N}_+} a(W_x) - \sqrt{N}). \end{aligned}$$

Using that

$$a_k^* \mathcal{N}_+ = (\mathcal{N}_+ - 1) a_k^* \quad (52)$$

$$a_k^* \sqrt{N - \mathcal{N}_+} = \sqrt{N + 1 - \mathcal{N}_+} a_k^* \quad (53)$$

$$a_k \sqrt{N - \mathcal{N}_+} = \sqrt{N - \mathcal{N}_+ - 1} a_k \quad (54)$$

we can move all of the \mathcal{N}_+ -dependent factors to the left. The factors on the right are then terms that also appear in H^{BF} and are \mathcal{N}_+ -bounded by Lemma A.1 and (59). With

$$\begin{aligned} & N^{-1} \|(\sqrt{N+1-\mathcal{N}_+}\sqrt{N+2-\mathcal{N}_+}-N)\Psi\|^2 \\ & \leq N^{-1} \langle ((N+2)^2 + N^2)\Psi, \Psi \rangle - 2N^{-1} \text{Re} \langle N(N+1-\mathcal{N}_+)\Psi, \Psi \rangle \\ & \leq 2N^{-1} \langle (2N+16)\Psi, \Psi \rangle + 2 \langle \Psi, \mathcal{N}_+\Psi \rangle \leq C(\|\Psi\| + \|\mathcal{N}_+\Psi\|)^2, \end{aligned} \quad (55)$$

we then obtain

$$\left| \left\langle e^{-iH_N^{\text{aux}}s} \Phi^{\leq N}, (H_N^{\text{aux}} - H^{\text{BF}}) e^{-iH^{\text{BF}}s} \Phi \right\rangle \right| \leq CN^{-1/2} (\|\Phi\| + \alpha^{\text{aux}}(s)) \alpha^{\text{BF}}(s) \quad (56)$$

with some constant $C > 0$. Using Lemma 3.1 completes the proof for $\Phi \in D(H_0)$, and the inequality extends to $\Phi \in D(\mathcal{N}_+)$ by density. \square

A Self-adjointness of the Hamiltonians

Here we prove the relevant self-adjointness and domain properties of H^{aux} and H^{BF} , as well as a useful general Lemma.

Lemma A.1. *Let $(M_{jk})_{j,k \in \mathbb{N}} \in \ell^2(\mathbb{N} \times \mathbb{N})$. Then for any $\Phi \in D(\mathcal{N}_+)$*

$$\begin{aligned} \left\| \sum_{j,k \in \mathbb{N}} M_{j,k} a_j a_k \Phi \right\|_{\mathcal{F}_+} & \leq \|M\|_{\ell^2} \|\mathcal{N}_+ \Phi\|_{\mathcal{F}_+} \\ \left\| \sum_{j,k \in \mathbb{N}} M_{j,k} a_j^* a_k^* \Phi \right\|_{\mathcal{F}_+} & \leq \|M\|_{\ell^2} \|(\mathcal{N}_+ + 2)\Phi\|_{\mathcal{F}_+}. \end{aligned}$$

Proof. We only prove the first inequality, the second can be proved in a similar way. We have for any $\Psi \in \mathcal{F}_+$ by the Cauchy-Schwarz inequality

$$\left| \left\langle \Psi, \sum_{j,k \in \mathbb{N}} M_{j,k} a_j a_k \Phi \right\rangle \right| \leq \|M\|_{\ell^2} \left(\sum_{j,k \in \mathbb{N}} |\langle \Psi, a_j a_k \Phi \rangle|^2 \right)^{1/2}. \quad (57)$$

Using that $\sum_{k \in \mathbb{N}} a_k^* a_k = \mathcal{N}_+$, we have

$$\begin{aligned} \sum_{j,k \in \mathbb{N}} |\langle \Psi, a_j a_k \Phi \rangle|^2 & \leq \|\Psi\|^2 \sum_{j,k \in \mathbb{N}} \langle a_j a_k \Phi, a_j a_k \Phi \rangle \\ & = \|\Psi\|^2 \sum_{j,k \in \mathbb{N}} \langle \Phi, (a_j^* a_j a_k^* a_k + a_j^* [a_k^*, a_j] a_k) \Phi \rangle \\ & = \|\Psi\|^2 \langle \Phi, \mathcal{N}_+(\mathcal{N}_+ - 1)\Phi \rangle. \end{aligned} \quad (58)$$

This yields the claim. \square

Lemma A.2. *The operator H_N^{aux} defined by the expression (25) is self-adjoint on $U_N D(H_N)$ and essentially self-adjoint on $D(H_0)$.*

Proof. Since H_N^{aux} acts non-trivially only on $\mathcal{H}_+^{\leq N}$, the differences with $U_N H_N U_N^*$ and the projection of H_0 to $\mathcal{H}_+^{\leq N}$ are bounded operators. For the quadratic terms this follows from Lemma A.1 and Parseval's identity:

$$\begin{aligned}
\|V_{jk00}\|_{\ell^2(\mathbb{N}\times\mathbb{N})}^2 &= \|V_{00jk}\|_{\ell^2(\mathbb{N}\times\mathbb{N})}^2 \\
&\leq \sum_{j,k\in\mathbb{N}_0} \left| \int_{T^d} \int_{T^d} dy \, dy' \bar{\varphi}_j(y) \bar{\varphi}_k(y') V(y-y') \right|^2 \\
&= \|V(y-y')\|_{L^2(T^d\times T^d)}^2 = \|V\|_{L^2(T^d)}^2, \\
\|V_{j0k0}\|_{\ell^2(\mathbb{N}\times\mathbb{N})}^2 &\leq \sum_{j,k\in\mathbb{N}_0} |\langle \varphi_j \otimes \varphi_0, V(y_1-y_2) \varphi_0 \otimes \varphi_k \rangle|^2 = \|V\|_{L^2(T^d)}^2 \leq \|V\|_{L^\infty}^2. \tag{59}
\end{aligned}$$

For the cubic and quartic terms, it follows in an analogy with the bounds (37), (48). The remaining interaction terms are bounded by $\|a(W_x)\Psi\|_{\mathcal{H}_+} \leq \|W\|_{L^2} \|\sqrt{\mathcal{N}_+}\Psi\|_{\mathcal{H}_+}$. The claim thus follows from the Kato-Rellich theorem. \square

Proposition A.3. *Let $V, W \in L^2(T^d)$, then $H^{\text{BF}} - H_0$ is \mathcal{N}_+ -bounded and H^{BF} is essentially self-adjoint on $D(H_0)$.*

Proof. The first statement follows from Lemma A.1 as above. Consequently, there exists a constant c such that $H^{\text{BF}} + c\mathcal{N}_+$ is self-adjoint on $D(H_0) = D(H_0 + c\mathcal{N}_+)$ by the Kato-Rellich theorem, since $H^{\text{BF}} - H_0$ is $H_0 + c\mathcal{N}_+$ -bounded with bound less than one.

Essential self-adjointness of H^{BF} can now be obtained by applying the commutator theorem [40, Thm.X.36], with $H^{\text{BF}} + c\mathcal{N}_+$ as a comparison operator. For this, it is sufficient to prove that

$$|\langle \Phi, [\mathcal{N}_+, H^{\text{BF}}]\Phi \rangle| = |\langle \Phi, [\mathcal{N}_+, H^{\text{BF}} - H_0]\Phi \rangle| \leq C \langle \Phi, \mathcal{N}_+ \Phi \rangle,$$

for some constant $C > 0$ and all $\Phi \in D(H_0^{1/2})$. Since the commutator of \mathcal{N}_+ with H^{Bog} is again a quadratic operator, composed of the same terms up to signs, this follows from Lemma A.1 as above. \square

References

- [1] N. Benedikter, M. Porta, and B. Schlein. *Effective Evolution Equations from Quantum Dynamics*. Springer Briefs in Mathematical Physics. Springer, 2016.
- [2] C. Boccato, C. Brennecke, S. Cenatiempo, and B. Schlein. Complete Bose–Einstein condensation in the Gross–Pitaevskii regime. *Commun. Math. Phys.*, 359(3):975–1026, 2018.
- [3] C. Boccato, C. Brennecke, S. Cenatiempo, and B. Schlein. Bogoliubov theory in the Gross–Pitaevskii limit. *Acta Math.*, 222(2):219–335, 2019.
- [4] N. Bogoliubov. On the theory of superfluidity. *J. Phys.*, 11(1):23, 1947.

- [5] L. Boßmann, N. Pavlović, P. Pickl, and A. Soffer. Higher order corrections to the mean-field description of the dynamics of interacting bosons. *J. Stat. Phys.*, 178(6):1362–1396, 2020.
- [6] L. Boßmann, P. Petrat, P. Pickl, and A. Soffer. Beyond Bogoliubov dynamics. *arXiv:1912.11004*, 2019.
- [7] L. Boßmann, P. Petrat, and R. Seiringer. Asymptotic expansion of the low-energy excitation spectrum for weakly interacting bosons. *arXiv:2020.09825*, 2020.
- [8] C. Brennecke, P.T. Nam, M. Napiórkowski, and B. Schlein. Fluctuations of N-particle quantum dynamics around the nonlinear Schrödinger equation. *Ann. Inst. H. Poincaré C, Anal. Non Linéaire*, 36(5):1201 – 1235, 2019.
- [9] T. Chen and A. Soffer. Mean field dynamics of a quantum tracer particle interacting with a boson gas. *J. Funct. Anal.*, 276(3):971–1006, 2019.
- [10] D.-A. Deckert, J. Fröhlich, P. Pickl, and A. Pizzo. Dynamics of sound waves in an interacting Bose gas. *Adv. Math.*, 293:275–323, 2016.
- [11] J. Dereziński and M. Napiórkowski. Excitation spectrum of interacting bosons in the mean-field infinite-volume limit. *Ann. Henri Poincaré*, 15(12):2409–2439, 2014.
- [12] H. Fröhlich. Theory of electrical breakdown in ionic crystals. *Proc. R. Soc. Lond. A*, 160(901):230–241, 1937.
- [13] J. Fröhlich, Z. Gang, and A. Soffer. Some Hamiltonian models of friction. *J. Math. Phys.*, 52:083508, 2011.
- [14] J. Ginibre and G. Velo. The classical field limit of scattering theory for non-relativistic many-boson systems. I. *Commun. Math. Phys.*, 66(1):37–76, 1979.
- [15] J. Ginibre and G. Velo. The classical field limit of scattering theory for non-relativistic many-boson systems. II. *Commun. Math. Phys.*, 68(1):45–68, 1979.
- [16] P. Grech and R. Seiringer. The excitation spectrum for weakly interacting bosons in a trap. *Commun. Math. Phys.*, 322(2):559–591, 2013.
- [17] M. Grillakis and M. Machedon. Pair excitations and the mean field approximation of interacting bosons, I. *Commun. Math. Phys.*, 324(2):601–636, 2013.
- [18] M. Grillakis and M. Machedon. Pair excitations and the mean field approximation of interacting bosons, II. *Commun. PDE*, 42(1):24–67, 2017.
- [19] M. Grillakis, M. Machedon, and D. Margetis. Second-order corrections to mean field evolution of weakly interacting bosons, I. *Commun. Math. Phys.*, 294(1):273, 2010.
- [20] M. Grillakis, M. Machedon, and D. Margetis. Second-order corrections to mean field evolution of weakly interacting bosons, II. *Adv. Math.*, 228(3):1788–1815, 2011.

- [21] F. Grusdt and E. Demler. New theoretical approaches to Bose polarons. In M. Inguscio, W. Ketterle, S. Stringari, and G. Roati, editors, *Proceedings of the international school of physics "Enrico Fermi"*, pages 325–411. Società Italiana di Fisica, 2016.
- [22] K. Hepp. The classical limit for quantum mechanical correlation functions. *Commun. Math. Phys.*, 35(4):265–277, 1974.
- [23] E. Kuz. Exact evolution versus mean field with second-order correction for bosons interacting via short-range two-body potential. *Differ. Integral Equ.*, 30(7/8):587–630, 2017.
- [24] J. Lampart. The renormalised Bogoliubov-Fröhlich Hamiltonian. *J. Math. Phys.*, 61(10):101902, 2020.
- [25] M. Lewin. Mean-field limit of Bose systems: rigorous results. *arXiv:1510.04407*, 2015.
- [26] M. Lewin, P. T. Nam, and B. Schlein. Fluctuations around Hartree states in the mean field regime. *Amer. J. Math.*, 137(6):1613–1650, 2015.
- [27] M. Lewin, P. T. Nam, S. Serfaty, and J. P. Solovej. Bogoliubov spectrum of interacting Bose gases. *Commun. Pure Appl. Math.*, 68(3):413–471, 2015.
- [28] D. Mitrouskas, S. Petrat, and P. Pickl. Bogoliubov corrections and trace norm convergence for the Hartree dynamics. *Rev. Math. Phys.*, 31(8), 2019.
- [29] K. Mysliwy and R. Seiringer. Microscopic derivation of the Fröhlich Hamiltonian for the Bose polaron in the mean-field limit. *arXiv preprint arXiv:2003.12371*, 2020.
- [30] P. T. Nam. Bogoliubov theory and bosonic atoms. *arXiv:1109.2875*, 2011.
- [31] P. T. Nam and M. Napiórkowski. Bogoliubov correction to the mean-field dynamics of interacting bosons. *Adv. Theor. Math. Phys.*, 21(3):683–738, 2017.
- [32] P. T. Nam and M. Napiórkowski. A note on the validity of Bogoliubov correction to mean-field dynamics. *J. Math. Pures Appl.*, 108(5):662–688, 2017.
- [33] P. T. Nam and M. Napiórkowski. Norm approximation for many-body quantum dynamics: focusing case in low dimensions. *Adv. Math.*, 350:547–587, 2019.
- [34] E. Nelson. Interaction of nonrelativistic particles with a quantized scalar field. *J. Math. Phys.*, 5(9):1190–1197, 1964.
- [35] T. Paul and M. Pulvirenti. Asymptotic expansion of the mean-field approximation. *Discrete Contin. Dyn. Syst. A*, 39(4):1891–1921, 2019.
- [36] S. Petrat, P. Pickl, and A. Soffer. Derivation of the Bogoliubov time evolution for a large volume mean-field limit. *Ann. Henri Poincaré*, 21(2):461–498, 2020.
- [37] A. Pizzo. Bose particles in a box I. A convergent expansion of the ground state of a three-modes Bogoliubov Hamiltonian. *arXiv:1511.07022*, 2015.

- [38] A. Pizzo. Bose particles in a box II. A convergent expansion of the ground state of the Bogoliubov Hamiltonian in the mean field limiting regime. *arXiv:1511.07025*, 2015.
- [39] A. Pizzo. Bose particles in a box III. A convergent expansion of the ground state of the Hamiltonian in the mean field limiting regime. *arXiv:1511.07026*, 2015.
- [40] M. Reed and B. Simon. *Methods of modern mathematical physics:II Fourier analysis, self-adjointness*. Academic Press, 1975.
- [41] R. Seiringer. The excitation spectrum for weakly interacting bosons. *Commun. Math. Phys.*, 306(2):565–578, 2011.
- [42] R. Seiringer. Bose gases, Bose–Einstein condensation, and the Bogoliubov approximation. *J. Math. Phys.*, 55(7):075209, 2014.
- [43] H. Spohn. Kinetic equations from Hamiltonian dynamics: Markovian limits. *Rev. Modern Phys.*, 52(3):569–615, 1980.
- [44] C. Zipkes, S. Palzer, C. Sias, and M. Köhl. A trapped single ion inside a Bose–Einstein condensate. *Nature*, 464(7287):388–391, 2010.