



# **Does lambing season affect mother-young relationships and lamb vigor in D'man sheep reared in oases?**

Amel Dhaoui, Mohamed Chniter, F Levy, Raymond Nowak, M. Hammadi

## **► To cite this version:**

Amel Dhaoui, Mohamed Chniter, F Levy, Raymond Nowak, M. Hammadi. Does lambing season affect mother-young relationships and lamb vigor in D'man sheep reared in oases?. *Animal*, 2020, 14 (11), pp.2363-2371. <10.1017/S1751731120001342>. <hal-03038594>

**HAL Id: hal-03038594**

**<https://hal.science/hal-03038594v1>**

Submitted on 3 Dec 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC-ND 4.0 - Attribution - Non-commercial use - No Derivative Works - International License

**Does lambing season affect mother-young relationships and lamb vigor in  
D'man sheep reared in oases?**

A. Dhaoui<sup>1, 2</sup>, M Chniter<sup>1, 3</sup>, F Lévy<sup>4</sup>, R Nowak<sup>4</sup>, M Hammadi<sup>1, 5</sup>

<sup>1</sup> LR16IRA04, Institut des Régions Arides, Institution de la Recherche et de  
l'Enseignement Supérieur Agricoles, 4119 Médenine, Tunisie.

<sup>2</sup> Faculté des Sciences de Gabès, Université de Gabès, Cité Erriadh 6072 Gabès,  
Tunisie

<sup>3</sup> Département des sciences animales, Institut National Agronomique de Tunisie, 43  
Av Charles Nicolle 1082 Tunis, Tunisia

<sup>4</sup> PRC, INRAE, CNRS, IFCE, Université de Tours, 37380 Nouzilly, France.

<sup>5</sup> Ecole Doctorale SIS, Université de Gabès Cité Erriadh 6072 Gabès, Tunisie

Corresponding author: Amel Dhaoui. Email: [amel.ghiloufi@yahoo.com](mailto:amel.ghiloufi@yahoo.com)

Short title: **Does season affect ewe and lamb behaviors?**

## Abstract

Although sheep are known to be seasonal breeders and give birth in winter, not all of them follow this trend. A few breeds can be mated and give birth all year round, meaning that mothers and neonates will have to face contrasting climatic conditions. The aim of this study was to test whether lambing season affects maternal and neonatal behaviors in D'man sheep. During four different lambing seasons (winter, autumn, summer, spring), periparturient ewes ( $n = 111$ ) and their lambs ( $n = 213$ ) were kept under 24-h-video surveillance in order to record post-partum behaviors. Mother-young preference was tested around 48 h after parturition. Lamb vigor was studied by the determination of birth weight, early postnatal behavior and rectal temperature at birth and 48 h later. Litter expulsion time was not affected by lambing season but birth weight was biased against summer and winter born lambs. Ewes provided a higher intensity of care to their offspring in winter: latency for grooming was shorter and time spent grooming was longer compared to lambing in spring and summer ( $P = 0.01$  in all cases). On the other hand, lambs were the most active in spring as they were faster to extend their hind legs ( $P = 0.01$ ), stand up ( $P = 0.04$ ) and reach the udder ( $P = 0.04$ ). Rectal temperature at 48 h was affected by season of birth ( $P < 0.001$ ) with higher values observed in summer. Glycaemia variation between birth and 48 h was the lowest in spring born lambs and plasma levels increased less in spring born lambs than in winter ( $P < 0.0001$ ), autumn ( $P < 0.0001$ ) and summer born lambs ( $P < 0.0001$ ). In the choice test, mothers clearly preferred their own young and no season effect was detected except that in the first minute of the test they spent less time near their own young in winter than in the other seasons ( $P = 0.04$ ). Lambs also chose their mother successfully without any major effect of the season however, but winter born lambs were the least vocal ( $P = 0.01$ ). Overall,

this study show that maternal care, lamb behavior and vigor vary lightly according to seasons, albeit not in a consistent manner. In conclusion, a season is no more detrimental than another for the onset of mother-young relationships.

**Keywords:** parturition, maternal behavior, neonate behavior, bonding, glycaemia

## **Implications**

D'man ewes give birth all year round in oases, which may have a major impact on mother-young interactions and lamb survival due to weather fluctuations (heat/cold), especially since mortality is reported to be higher in winter. This study showed that maternal care, lamb behavior and neonatal physiology may vary according to seasons but only mildly and not in a consistent manner. We concluded that winter is no more detrimental than another season for the onset of mother-young relationships in intensive rearing conditions providing shelter.

## Introduction

In mammals giving birth to precocial young, maternal care from birth to weaning leads to a strong bond between the mother and her offspring. This bond is essential not only for the survival of the young but also for the establishment of its emotional behaviors and fitness (Dwyer, 2014). In sheep, behaviors displayed around and after lambing have a major impact on lamb survival but also on its growth rate and weaning weight, and consequently on global productivity (O'Connor et al., 1985). Lamb survival is crucially dependent on the ability to surmount transition difficulties from intra- to extra-uterine existence. This ability depends on the quality of the interactions with the mother especially in large litters. Maternal efforts to maintain proximity and to provide an adequate social environment will also strengthen the mother-young bond by providing juveniles with the abilities to cope with stressful situations (Nowak et al., 2000). Consequently, the expression of maternal care and the establishment of maternal-filial bond throughout lactation are dependent on the initial onset of maternal behavior (Pickup and Dwyer, 2011). Early maternal care, which appears abruptly around parturition, consists of grooming and nursing activities to provide the newborn with colostrum to satisfy its metabolic needs, and protection to face various elements of the environment (Nowak et al., 2000). Grooming the birth fluids limits heat loss by drying the coat, and prevents suffocation by removal of the membranes from the lamb's face. Grooming other parts of the body such as the rump stimulates the newborn to stand up, helps it to focus on the maternal body and guides the young toward her mammary region as the mother adopts a specific nursing posture which makes the teats easily accessible (Nowak, 2006). Moreover, during grooming, the ewe learns the smell of her newborn and an olfactory memory takes place, which is a crucial step for maternal bonding (Lévy et al., 2004). Nursing,

on the other hand, calms and comforts the lamb, and plays a reinforcing role in the development of the mother young bond (Nowak et al., 2000). Infant behavior plays also an important part in the establishment of the mutual mother-young bond and survival (Nowak and Lindsay, 1992). Thermoregulation, which depends on birth weight and milk availability, is the most vital goals to be reached after birth (Dwyer, 2008). In cold and windy weather, neonates lose heat quickly as they have to surmount the abrupt drop in ambient temperature. Maintenance of body temperature relies on a balance between heat loss and production (Dwyer et al., 2015). Therefore, success or failure to surmount cold conditions will depend on the lambs' capacity to suckle and ingest colostrum as soon as possible after birth.

Lamb mortality still remains a major problem both for economic and animal welfare issues as the highest rate occurs within the first three post-natal days (Dwyer, 2008; 2014; Dwyer et al., 2015). It is particularly important amongst lambs born in large litters and with low birth weights (Chniter et al., 2011). Lambs with low birth weight have lower rectal temperature and glucose sera concentration than lambs with heavier birth weight (Chniter et al., 2013). Therefore, they are behaviorally slow and are more inclined to suffer from hypothermia (Dwyer et al., 2005). This difficulty to thermoregulate impairs their ability to suckle rapidly and to bond with their mothers (Chniter et al., 2017). Under intensive lambing conditions, Chniter et al. (2011) observed that multiple born D'man lambs had a higher mortality rate in winter especially within the first 72 h. This cannot be due to low milk production because D'man ewes produce more milk in winter and autumn than in summer (Dhaoui et al., 2018). If we exclude milk as a causal factor, a possible explanation could be found in the early mother-young interactions that may be influenced by season. There is no

evidence in the literature on this issue in sheep but it has been shown in Mountain goats that maternal characteristics affect winter survival of kids (Théoret-Gosselin and Hamel, 2015). In sheep, maternal care, neonatal behavior, and lamb survival are intimately linked (Dwyer et al., 2015). Should lambing take place under harsh weather conditions, neonatal behavior is penalized and lamb losses increase in the following 3 days. Therefore, we hypothesized that season of lambing affects the onset of maternal care and neonatal behavior in D'man sheep which would be weakened by winter conditions. Characterization of mother-young interactions at birth over different seasonal conditions should help us understand the increased winter loss reported by Chniter et al. (2011) and lead to more adequate management strategies during lambing.

## Material and methods

This study was carried out in the Experimental Station of the Institut des Régions Arides located in Chenchou (Gabès), approximately 20 km West of Gabès (N33°53' E9°53', Tunisia). This region is characterized by a low arid Mediterranean climate with an irregular rainfall over the year (Dhaou et al., 2018). An automatic weather station (CR510 Campbell Scientific data logger) provided average ambient temperatures, minima and maxima, and rainfall data throughout the study. Meteorological data are summarized in Table 1.

**Insert table 1 here**

## Animals

A total of 111 D'man ewes (80 multiparous and 31 primiparous) and their 213 lambs were used in this study. Details about maternal experience and litter size are presented in table 2.

**Insert table 2 here**

Animals were reared under intensive production system, kept as a single flock in a communal yard for each season (4.5 m<sup>2</sup>/ewe), and managed as previously indicated in Dhaoui et al. (2018). Briefly, the yard had a North-East to South-West orientation and possessed a roof to protect the animals from adverse weather conditions providing more light in winter and more shade in summer. Feeding regimes and management procedures were adapted to the physiological stage of the ewes. They received a fixed diet based on Lucerne hay (*Medicago sativa*) and commercial concentrate mixture. Animals had access to fresh water and trace-mineral salt blocks *ad-libitum*.

Lambing periods spread from mid-November 2014 to January 2015 for winter (22 ewes weighing 35.6 ± 6.7 kg; 38 lambs weighing 2.9 ± 0.9 kg), September to October 2015 for autumn (25 ewes weighing 40.9 ± 6.8 kg; 51 lambs weighing 2.7 ± 0.8 kg), July to September 2016 for summer (31 ewes weighing 45.8 ± 6.8 kg; 65 lambs weighing 2.6 ± 0.8 kg) and March to May 2017 for spring season (33 ewes weighing 43.6 ± 9.1 kg; 59 lambs weighing 3.1 ± 0.5 kg).

Once born, lambs were identified, ear-tagged and weighted. Newborn lambs were allowed 2 h to stand and suck successfully unaided. At 3 h, the ewe and her litter were transferred from the communal yard into a 1.5 m<sup>2</sup> individual pen and stayed in it



until the behavioral tests were terminated. Any lamb that had not sucked by 2 h after birth was assisted and excluded from the study.

#### *Behaviors on birth site*

Ewes were kept under 24-h-video surveillance (Xenyum Pro Surveillance System), and 2 h video sessions were chosen to assess maternal and neonatal behaviors on the birth site based on Dwyer et al. (2005). An ethogram was defined for coding the duration of, litter expulsion, latency of first grooming of each neonate and time spent grooming during the 30 min following first lamb expulsion (Table 3). Behavioral data were videotaped and analyzed by one expert person using Boris recording software (Friard and Gamba, 2016).

**Insert table 3 here**

#### *Two-choice tests of preference*

Mother-young preference was assessed during a 5-min test approximately 48 h after parturition. The procedures and testing pens were similar as previously described by Chniter et al. (2017). For both lambs and ewes, the location of the testing pen was in an isolated indoor room. The pen consisted of an equilateral triangular arena (6 m × 6 m × 6 m). One corner ended with a starting pen for the subject to be tested and the two others consisted of individual pens for stimulus subjects (2 m × 1 m). These were separated by an empty pen (2 m × 1 m) which prevented subjects from being too close to each other. A 50-cm-wide zone next to the stimulus subjects was considered as a contact zone.

The preference for the lamb by its mother was assessed first by placing the own and alien lambs in individual pens at the opposite stimulus corners and the ewe in the starting pen. In the case of large litters, only one lamb was used as a stimulus to minimize any bias due to differences in litter sizes.

The preference for the mother by the lamb was assessed immediately after. Mothers and alien recent parturient ewes (a maximum delay of 3 days after birth) were separately kept at the opposite stimulus corners of the testing arena. The lamb was carefully placed in the starting pen. Each ewe was used only twice: first with one of her own lamb and then as an alien mother in the following test before being released and returned to the individual pen with her litter. During the test, the siblings were kept away (15 m) from the testing arena to avoid vocal communication.

Both for young or mothers, the location of stimulus subjects was balanced so that own and alien conspecifics were allocated to left and right corners equally. The following data were taken: latency to reach the stimulus corners (both forelegs in the contact zone), time spent in the contact zones near the own or alien partner during the 1<sup>st</sup> min, and the 5 min of the test, and the number of bleats during the 5 min of the test. In addition, ewes and lambs were considered to make a correct choice if they spent more than 60 % of the time near the two stimulus subjects and at least two-thirds of this time near its familiar partner (Chniter et al., 2017).

#### *Rectal temperature and glycaemia*

Rectal temperature (RT, °C) was recorded at 1 h and 48 h of age in an area unexposed to solar radiation, using a digital probe thermometer (range: 32.00 ~ 43.00°C, accuracy: +/- 0.10°C (35.50°C ~ 42.00°C). Blood samples were taken from 190 lambs by jugular venipuncture into hemolysis vacutainer tubes for glucose

measurement. A fully trained person, who intervened calmly, caught the lamb, restrained it on its side, and collected the rectal temperature and the blood sample. Human intervention had no detrimental effect on subsequent ewe-lamb interactions. After the blood samples were centrifuged ( $3000 \times g$ ,  $4^{\circ}\text{C}$ ) for 20 min, the separated sera were stored at  $-20^{\circ}\text{C}$  until analysis of glucose using a colorimetric method (Kit Ref 20121, Biomaghreb, Tunisia).

## Statistical analyses

All the data were statistically analyzed and graphically represented using GraphPad Prism 6.0 software. Tests of normality (Shapiro-Wilk test) were performed and showed non-normal distributions. Consequently, two-tailed nonparametric tests were used to study the season effect on maternal and neonatal behavior, and on lamb vigor. Overall, the effect of the season was performed by Kruskal-Wallis test with Dunn's correction for multiple comparisons. Inter-season comparisons were performed using the Mann-Whitney test. Preference toward a partner (own vs alien) was analyzed by the Wilcoxon signed-rank test. In all cases, P-values less than 0.05 were considered as statistically significant. Results for behavioral traits and vigor are presented in terms of median and lower and upper quartiles [LQ-HQ].

## Results

### *Behaviors on the birth site*

*Lambing process and maternal care.* Season of lambing did not affect (Kruskal-Wallis = 4.94;  $P = 0.18$ ) litter expulsion time. However, the latency required to groom each neonate was affected by season of lambing (Kruskal-Wallis = 8.81;  $P = 0.03$ ; Figure

1a). It was shorter in winter compared to lambing in spring (Mann-Whitney  $U = 243.0$ ;  $P = 0.01$ ) and summer (Mann-Whitney  $U = 309.5$ ;  $P = 0.01$ ). In autumn, ewes had an intermediate value.

**Insert Figure1 here**

During the first 30 min following birth, ewes lambing in winter spent significantly more time grooming their lambs than ewes lambing in spring (Mann-Whitney  $U = 89.0$ ;  $P = 0.01$ ) and in summer (Mann-Whitney  $U = 69.0$ ;  $P = 0.04$ ). Ewes lambing in autumn spent an intermediate value grooming their lambs (Mann-Whitney  $U = 97.0$ ;  $P = 0.16$ ) (Figure 1b).

*Neonatal behavior.* Immediate postnatal behavior was affected significantly by season of lambing. Lambs born in spring pushed upon their knees (Kruskal-Wallis = 10.36;  $P = 0.01$ ; Figure 2a), stood up (Kruskal-Wallis = 8.26;  $P = 0.04$ ; Figure 2b) and reached the udder earlier (Kruskal-Wallis = 8.15;  $P = 0.04$ ; Figure 2c) than those born in the other seasons.

**Insert Figure 2 here**

*Preference for own lamb at 48 h postpartum*

All the ewes reached the contact zone before the end of the test without any effect of the season, and 78/111 made a correct choice. Ewes reached their own lamb's side significantly more rapidly than the side of the alien lamb in all four seasons (Wilcoxon tests = Spring: 5.0 s [4.0-14.0] vs 43.0 s [14.0-300.0],  $Z = 3.86$ ,  $P < 0.001$ ; Summer: 4.0 s [3.0-8.0] vs 49.0 s [18.0-112.0]  $Z = 4.06$ ,  $P < 0.001$ ; Autumn: 7.0 s [3.0-6.0] vs

265 39.0 s [11.0-222.0],  $Z = 3.41$ ,  $P < 0.001$ ; Winter: 9.0 s [4.0-17.0] vs 25.5 s [5.0-74.0],  
266  $Z = 2.07$ ,  $P = 0.04$ ). They also spent significantly more time near their own than near  
267 the alien lamb during the first minute of the test (Wilcoxon tests = Spring: 42.0 s  
268 [17.0-50.0] vs 3.0 s [0.0-8.0],  $Z = 3.85$ ,  $P < 0.001$ ; Summer: 46.0 s [34.0-51.0] vs 2.0  
269 s [0.0-6.0],  $Z = 4.84$ ,  $P < 0.0001$ ; Autumn: 44.0 s [35.0-48.0] vs 1.0 s [0.0-4.0],  $Z =$   
270 4.37,  $P < 0.0001$ ; Winter : 25.5 s [18.0-42.0] vs 2.0 s [0.0-10.0],  $Z = 3.04$ ,  $P = 0.002$ ).  
271 These differences were confirmed for the whole duration of the test (Wilcoxon tests =  
272 Spring: 224.0 s [95.0-278.0] vs 10.0 s [0.0-29.0],  $Z = 4.37$ ,  $P < 0.0001$ ; Summer:  
273 211.0 s [184.0-270.0] vs 11.0 s [2.0-39.0],  $Z = 4.62$ ,  $P < 0.0001$ ; Autumn: 233.0  
274 [148.0-271.0] vs 9.0 s [1.0-16.0],  $Z = 4.34$ ,  $P < 0.0001$ ; Winter: 175.0 s [82.0-228]  
275 [6.0-39.0] vs 16.0 s,  $Z = 3.45$ ,  $P < 0.001$ ).

276 During the first minute, ewes lambing in winter spent significantly less time near their  
277 own lamb than those lambing in autumn (25.5 s [18.0-42.0] vs 44.0 s [35.0-48.0],  
278 Mann-Whitney  $U = 156$ ;  $P = 0.01$ ) or in summer (25.5 s [18.0-42.0] vs 46.0 s [34.0-  
279 51.0], Mann-Whitney  $U = 186.5$ ;  $P = 0.01$ ). This effect was lost over the whole  
280 duration of the test even though winter lambing ewes still had the lowest median  
281 value. There was no difference between seasons on the latency to reach the contact  
282 zone next to a lamb, on the time spent near the alien lamb during the first minute or  
283 during the whole duration of the test, nor on the number of bleats emitted by ewe.

284

#### 285 *Preference for the mother at 48 h of age*

286 Lambs reached more rapidly their own mothers than the alien ewes (Wilcoxon tests =  
287 Spring: 30.0 s [10.0-82.0] vs 78.0 s [33.0-176.0],  $Z = 2.66$ ,  $P < 0.01$ ; Summer: 43.0 s  
288 [19.0-136.0] vs 147.0 s [41.0-300.0],  $Z = 3.30$ ,  $P < 0.001$ ; Autumn: 31.0 s [12.0-  
289 126.0] vs 83.0 s [26.0-238.0],  $Z = 2.08$ ,  $P = 0.03$ ; Winter: 22.0 s [12.0-88.0] vs 90.5 s

[20.0-147.0],  $Z = 1.57$ ,  $P = 0.01$ ). Latencies to reach a ewe were not affected by the season. They spent more time near their own mothers than alien ewes during the first minute (Wilcoxon tests = Spring: 26.0 s [0.0-42.0] vs 0.0 s [0.0-13.0],  $Z = 3.23$ ,  $P < 0.01$ ; Summer: 8.0 s [0.0-39.0] vs 0.0 s [0.0-8.0],  $Z = 3.39$ ,  $P < 0.001$ ; Autumn: 21.0 s [0.0-40.0] vs 0.0 s [0.0-15.0],  $Z = 2.95$ ,  $P = 0.003$ ; Winter: 29.5 s [0.0-42.0] vs 0.0 s [0.0-14.0],  $Z = 2.24$ ,  $P = 0.02$ ) and for the whole duration of the test (Wilcoxon tests = Spring: 175.0 s [95.0-241.0] vs 50.0 s [9.0-113.0],  $Z = 4.25$ ,  $P < 0.0001$ ; Summer: 151.0 s [10.0-238.0] vs 24.0 s [0.0-101.0],  $Z = 3.74$ ,  $P < 0.001$ ; Autumn: 145.0 s [85.0-217.0] vs 52.0 s [14.0-98.0],  $Z = 3.94$ ,  $P < 0.0001$ ; Winter: 159.5 s [116.0-210.0] vs 47.0 s [17.0-118.0],  $Z = 3.16$ ,  $P < 0.01$ ).

Overall, season of birth did not affect the capacity of lambs to discriminate between their own and alien mothers at 48 h of age. However, lambs born in winter emitted significantly less bleats (13.5) than lambs born in autumn (13.5 [9.0-25.0] vs 21.0 [12.0-33.0], Mann-Whitney  $U = 711$ ;  $P = 0.03$ ), in summer (13.5 [9.0-25.0] vs 23.0 [16.0-40.0], Mann-Whitney  $U = 833.5$ ;  $P = 0.01$ ) and in spring (13.5 [9.0-25.0] vs 26.0 [14.0-41.0], Mann-Whitney  $U = 671.5$ ;  $P = 0.001$ ).

#### *Birth weight, rectal temperature and glycaemia*

Lambs born in summer were lighter than those born in autumn (Mann-Whitney  $U = 527.0$ ;  $P = 0.04$ ) and spring (Mann-Whitney  $U = 1210$ ;  $P = 0.001$ ; Figure 3). Winter born lambs had an intermediate birth weight.

**Insert here Figure 3.**

One hour after birth, the median rectal temperature of D'man lamb was 38.7 °C [38.40 – 39.08]. It rose to 39.20 °C [39.00 – 39.60] at 48 h of age with a median

difference of + 0.6 °C. Season of birth did not affect the temperatures recorded around birth but it did 48 h later (Kruskal-Wallis = 19.74;  $P < 0.001$ ; Figure 4a and 4b). It was higher in summer than in winter (Mann-Whitney U = 763.5;  $P = 0.001$ ) and in spring born lambs (Mann-Whitney U = 1064;  $P < 0.0001$ ). Autumn born lambs had an intermediate value. Season of birth had a significant effect on rectal temperature variation from 1 to 48 h of age (Kruskal-Wallis = 12.67;  $P = 0.01$ ; Figure 4c). Spring born lambs had a lower rectal temperature increase than winter (Mann-Whitney U = 735;  $P = 0.01$ ), autumn (Mann-Whitney U = 1041;  $P = 0.01$ ) and summer born lambs (Mann-Whitney U = 1260;  $P = 0.001$ ).

**Insert here Figure 4.**

Median glycaemia was 2.72 mmol / L [2.24 – 3.12] within the hour following birth and reached 4.63 mmol / L [3.92 – 6.34] 48 h later with a rise of 1.67 mmol / L [1.17 – 4.08]. Season of birth had a significant effect both at 1 h (Kruskal-Wallis = 11.09;  $P = 0.01$ ; Figure 4a) and 48 h of age (Kruskal-Wallis = 54.63;  $P < 0.0001$ ; Figure 4b), and also on the increase between 1 h and 48 h of age (Kruskal-Wallis = 48.65;  $P < 0.0001$ ; Figure 4c). At birth, summer born lambs had significantly lower glycaemia compared to spring born lambs (Mann-Whitney U = 293;  $P = 0.0001$ ). Winter and autumn born lambs had an intermediate value. At 48 h age, spring born lambs had the lowest concentrations compared to winter (Mann-Whitney U = 35;  $P < 0.0001$ ), autumn (Mann-Whitney U = 46;  $P < 0.0001$ ) and summer born lambs (Mann-Whitney U = 571;  $P < 0.0001$ ). In fact, glycaemia increased significantly less in spring born lambs than in winter (Mann-Whitney U = 25;  $P < 0.0001$ ), autumn (Mann-Whitney U = 29.5;  $P < 0.0001$ ) and summer born lambs (Mann-Whitney U = 281.5;  $P < 0.0001$ ).

340

## 341 **Discussion**

342 In this study, immediately after parturition ewes provided a higher intensity of care to  
343 their offspring in winter, but lambs were the most active in spring getting up and  
344 reaching the udder more rapidly. Forty-eight hours later, season of lambing did not  
345 affect the partners' ability to express a preference for each other even though in  
346 winter ewes spent less time near their own lambs during the first minute of the test  
347 and their lambs were less vocal. In terms of neonatal physiology, body temperature  
348 and glycaemia usually went up from birth to 48 h in all four seasons but spring born  
349 lambs had the lowest increase in both parameters, and summer born lambs reached  
350 the highest body temperature at 2 days of age. Clearly, while some behavioral and  
351 physiological parameter fluctuate according to seasons, the effects are minor and  
352 inconsistent. We cannot conclude that winter or any specific season (nor a  
353 temperature range) has detrimental consequences on mother-young relationships.  
354 The literature shows that lambs are at risk in winter when environmental conditions  
355 combine wind, rain and cold which prevents them from maintaining homeothermy  
356 and normal behavior (review: Pollard, 2006). This was not the case in our study even  
357 though we must keep in mind the limitations of the recording method concerning  
358 meteorological data. The database was based on average temperatures, minima and  
359 maxima for each lambing season, it did not provide specific local conditions for each  
360 birth. We cannot exclude however that some very cold winter nights (or even very hot  
361 summer days) may have influenced the expression of early behavior patterns; this  
362 should be taken into account in future work. Nonetheless, our average recording  
363 does not reveal hostile winter climatic conditions especially in terms of rainfall and we  
364 cannot conclude that lambs were submitted to cold exposure. In addition, the fact



that the yard was enclosed, possessed a roof, and the animals were put in individual pens could have protected them from adverse conditions.

D'man ewes started to groom their neonates sooner and for longer during the first 30 min postpartum in winter than in spring and summer. Although smaller lambs lead to an easier birth (Grommers et al., 1985) and an easy birth is associated with better maternal care (Poindron et al., 1984), neither lambing process nor litter size and lambs weight are satisfactory explanations for the difference in the expression of grooming behavior. Maternal experience on the other hand, instead of season, is a more likely causal factor. The number of primiparous ewes was lower in winter than in spring and in summer and parity, and not just age, is known to influence the expression of maternal care (Alexander et al 1993). It appears quite logical that the grooming activity reflecting maternal motivation should be more pronounced in winter lambing ewes. Some authors report that maternal attention stimulates neonatal behavior progression (Alexander and William, 1964; McGlone and Stobart, 1986; O'Connor et al., 1992) while others suggest that maternal attention slows down neonatal behavior progression (Vince, 1993; Dwyer and Lawrence, 1999). Our results are more in favor of the second findings since winter born lambs were the least active, while it was in spring and summer that lambs were the quickest to stand up and reach the udder. According to Vince (1993) grooming the head and shoulders, which is where the mother usually starts from, has soothing properties. Some studies also report that the lamb's neonatal activity is impaired in winter (Scotland: Dwyer and Morgan, 2006; Brazil: Fonsêca et al., 2014) but this does not seem an unanimous fact (Wassmuth et al. 2001). Local conditions as well as breed differences may account for the differences reported between authors.

Mother-young discrimination was well established at 48 h postpartum with no main season effect. All D'man ewes and lambs reached the contact zone and in most cases chose their own partner in comparison to an unfamiliar one. Nonetheless, the fact that winter parturient ewes spent less time next to their own young at the beginning of the test might suggest that their lambs were not initially as attractive or the mothers not as motivated to stay nearby. Limitation in the recording equipment did not allow us to analyze the behavioral interactions and find potential differences in animals' reactivity during the test. Acoustic in complement to olfactory cues from the lamb trigger maternal motivation and are important in the recognition of the young (Sèbe et al., 2007, Morton et al., 2018). This could have been valuable information especially because when lambs were submitted to the choice test it was found that, winter born subjects were the least vocal. Whether a lesser vocal activity is a characteristic of lambs born in winter has to be confirmed.

The fact that summer born lambs had a higher body temperature than winter born lambs at 48 h is quite logical as they were born during the warmest season, a difference that had already been found by Chniter et al. (2013). Our results on rectal temperature and glycaemia suggest that overall D'man neonates were not susceptible to oases climatic conditions and could maintain body temperature within normal range through two mechanisms. The first is mobilization of brown adipose tissue, a mechanism that helps newborn lambs adapt to cold immediately after being born (Dwyer and Morgan, 2006; Dwyer et al., 2015). The second is based on glucose plasma levels. Glucose is an important source of energy originating from colostrum and is required to maintain homeothermy beyond the neonatal period (Thompson et al., 2006; Miller et al., 2010). After birth, D'man lambs rapidly found the udder and

sucked successfully making glycaemia increased as a consequence which has also been reported by Chniter et al. (2016).

While in the present study, the overall rise in glycaemia and body temperature is a clear sign of the newborns' ability to adapt to their rearing conditions, it does not imply that D'man lambs are insusceptible to hypothermia. Chniter et al. (2011; 2013) had shown that winter born lambs were more fragile and suffered a higher mortality rate. However, in the present work we only took into account the lambs that survived until the time of the preference tests, excluding the weakest subjects more prone to suffer from hypothermia or postnatal trauma. Under such experimental conditions, confirming differences in neonatal physiology like in the studies by Chniter et al. (2011; 2013) was a more arduous task. Spring born lambs appeared anyhow to be the most vigorous at birth; they were also amongst the heaviest. The most important birth weight and vigor in spring born D'man lambs could reflect optimal climatic conditions during gestation with enhancing fetal growth in late pregnancy as suggested by Chniter et al. (2011). Their lesser glycaemic variability and their lower postnatal rise in body temperature suggest easier adaptation to the moderate spring weather conditions. Spring born lambs did not have to metabolize as much energy as autumn and winter born lambs, and could thermoregulate better than summer born lambs.

While our data indicate that lambing season affects some aspects of maternal and neonatal behavior in D'man sheep as well as neonatal physiology, we cannot conclude that one season has more impact than another does. Spring born lambs appeared the most vigorous at birth but mothers showed more intense maternal care in winter. Unsurprisingly, born lambs had a higher body temperature in summer but it was in spring that physiological factors were the least variable suggesting better

neonatal adaptation to mild climatic conditions. Some behavioral aspects involved in the identification of mother or young were suboptimal in winter but overall the ewe-lamb bond was clearly established in all four seasons by 48 h postpartum. Should lamb survival be at more at risk in winter as reported by Chniter et al. (2011), there is no reason to think that early mother-young interactions and bonding can be hold responsible for it in the conditions of the study.

### **Acknowledgements**

The present work was carried out in the “Laboratoire d’Elevage et Faune Sauvage (LR16IRA04), Institut des Régions Arides de Medenine”. It was funded by the Tunisian Ministry of Higher Education and Scientific Research, Gabès University. The analysis of the data was performed during Amel Dhaoui’s internship at the “Unité Mixte de Recherche de Physiologie de la Reproduction & des Comportements, INRA-PRC Val de Loire-France” and gave birth to the present manuscript. The first author thanks all INRA-PRC team for the warm welcome.

### **Declaration of interest**

The authors declare that they have no conflicts of interest.

### **Ethical statement**

The authors followed institutional and national guidelines for the care and use of laboratory animals. While there is no ethic committee in Tunisia, experimental conditions in the present work complied with the European Directive 2010/63/EU on the protection of animals used for scientific purposes.

**Software and data repository resources**

None of the data were deposited in an official repository.

**References**

- Alexander G, Bradley LR and Stevens D 1993. Effect of age and parity on maternal behaviour in single-bearing Merino ewes. *Australian Journal of Experimental Agriculture* 33, 721–728.
- Alexander G and Williams D 1964. Maternal facilitation of sucking drive in newborn lambs. *Science* 146, 665–666.
- Chniter M, Hammadi M, Khorchani T, Krit R, Lahsoumi B, Ben Sassi M, Nowak R and Ben Hamouda M 2011. Phenotypic and seasonal factors influence birth weight, growth rate and lamb mortality in D'man sheep maintained under intensive management in Tunisian oases. *Small Ruminant Research* 99, 166–170.
- Chniter M, Hammadi M, Khorchani T, Ben Sassi M, Ben Hamouda M and Nowak R 2013. Aspects of neonatal physiology have an influence on lambs' early growth and survival in prolific D'man sheep. *Small Ruminant Research* 111, 162–170.
- Chniter M, Salhi I, Harrabi H, Khorchani T, Lainé AL, Nowak R and Hammadi M 2016. Physiological changes in the peri-partum period and colostral IgG transfer in prolific D'man sheep: effects of parity and litter size. *Tropical Animal Health and Production* 48, 387–394.
- Chniter M, Dhaoui A, Hammadi M, Khorchani T, Ben Hamouda M, Poindron P, Cornilleau F, Lévy F and Nowak R 2017. Mother–young bonding in prolific D'man and Romanov sheep. *Journal of Ethology* 35, 297–305.
- Dhaou H, Jarray L and Ouessar M 2018. Bulletin climatique annuel 2017 Station de Chenchou, Edition Institut des régions arides, 5–16.
- Dhaoui A, Chniter M, Atigui M, Dbara M, Seddik MM and Hammadi M 2018. Factors affecting the milk yield and composition over lactation of prolific D'man ewes in Tunisian oases. *Tropical Animal Health and Production* 51, 507–518.

- 490 Dwyer CM and Lawrence AB 1999. Does the behaviour of the neonate influence the  
491 expression of maternal behaviour in sheep? Behaviour 136, 367–389.
- 492 Dwyer CM, Calvert SK, Farish M, Donbavand J and Pickup HE 2005. Breed, litter and parity  
493 effects on placental weight and placentome number, and consequences for the neonatal  
494 behaviour of the lamb. Theriogenology 63, 1092–1110.
- 495 Dwyer CM and Morgan CA 2006. Maintenance of body temperature in the neonatal lamb:  
496 Effects of breed, birth weight, and litter size. American Society of Animal Science 84,  
497 1093–1101.
- 498 Dwyer CM 2008. The welfare of the neonatal lamb. Small Ruminant Research 76, 31–41.
- 499 Dwyer CM 2014. Maternal behaviour and lamb survival: From neuroendocrinology to  
500 practical application. Animal 8, 102–112.
- 501 Dwyer CM, Conington J, Corbiere F, Holmoy IH, Muri K, Nowak R, Rooke J, Vipond J and  
502 Gautier JM 2015. Invited review: Improving neonatal survival in small ruminants: Science  
503 into practice. Animal 449–459.
- 504 Fonsêca VFC, Saraiva EP, Filho ECP, Furtado DA, Mariz TMA, Silva AL, Almeida MEV and  
505 Pinheiro AC 2014. Influence of the climatic environment and duration of labor on the  
506 mother-offspring interaction in Morada Nova sheep. Journal of animal science 92, 4123–  
507 4129.
- 508 Friard O and Gamba M 2016. BORIS: a free, versatile open-source event-logging software  
509 for video/audio coding and live observations. Methods in Ecology and Evolution 7, 1325–  
510 1330.
- 511 Grommers FJ, Elving L and Van Eldik P 1985. Parturition difficulties in sheep. Animal  
512 Reproduction Science 9, 365–374.
- 513 Lévy F, Keller M and Poindron P 2004. Olfactory regulation of maternal behavior in  
514 mammals. Hormones and Behavior 46, 284–302.
- 515 McGlone JJ and Stobart RH 1986. A quantitative ethogram of behavior of yearling ewes  
516 during two hours post-parturition. Applied Animal Behaviour Science 16, 157–164.

- 517 Miller DR, Blache D, Jackson RB, Downie EF and Roche JR 2010. Metabolic maturity at birth  
518 and neonate lamb survival: Association among maternal factors, litter size, lamb birth  
519 weight, and plasma metabolic and endocrine factors on survival and behavior. *Journal of*  
520 *Animal Science* 88, 581–592.
- 521 Morton CL, Hinch G, Small A and McDonald PG 2018. Flawed mothering or infant signaling?  
522 The effects of deficient acoustic cues on ovine maternal response. *Developmental*  
523 *Psychobiology* 60, 975–988.
- 524 Nowak RF and Lindsay DR 1992. Discrimination of merino ewes by their newborn lambs:  
525 important for survival? *Applied Animal Behaviour Science* 34, 61–74.
- 526 Nowak R, Porter RH, Lévy F, Orgeur P and Schaa B 2000. Role of mother-young  
527 interactions in the survival of offspring in domestic mammals. *Reviews of Reproduction* 5,  
528 153–163.
- 529 Nowak R 2006. Suckling, Milk, and the Development of Preferences toward Maternal Cues  
530 by Neonates: From Early Learning to Filial Attachment? *Advances in the Study of*  
531 *Behavior* 36, 1–58.
- 532 O'Connor, CE, Jay NP, Nicol M and Beatson PR 1985. Ewe maternal behaviour score and  
533 lamb survival. *Proceedings of the New Zealand Society of Animal Production*. 45, 159–  
534 162.
- 535 O'Connor CE and Lawrence AB 1992. Relationship between lamb vigour and ewe behaviour  
536 at parturition. *Animal Production* 54, 361–366.
- 537 Pickup HE and Dwyer CM 2011. Breed differences in the expression of maternal care at  
538 parturition persist throughout the lactation period in sheep. *Applied Animal Behaviour*  
539 *Science* 132, 33–41.
- 540 Poindron P, IldikoRaksanyi, Orgeur P and Le Neindre P 1984. Comparaison du  
541 comportement maternel en bergerie à la parturition chez des brebis primipares ou  
542 multipares de race Romanov, Préalpes du Sud et Ile-de-France. *Genetics Selection*  
543 *Evolution* 16, 503–522.

- 544 Pollard JC. 2006. Shelter for lambing in New Zealand: a review. New Zealand Journal of  
545 Agricultural Research 49, 395-404.
- 546 Sèbe F, Nowak R, Poindron P and Aubin T 2007. Establishment of vocal communication and  
547 discrimination between ewes and their lamb in the first two days after parturition.  
548 Developmental Psychobiology 49, 375–386.
- 549 Théoret-gosselin R, Hamel S and Côté SD 2015. The role of maternal behavior and offspring  
550 development in the survival of mountain goat kids. Oecologia 178, 175–186.
- 551 Thompson MJ, Briegel JR, Thompson AN and Adams NR 2006. Differences in survival and  
552 neonatal metabolism in lambs from flocks selected for or against staple strength.  
553 Australian Journal of Agricultural Research 57, 1221–1228.
- 554 Vince MA 1993. Newborn Lambs and Their Dams: The Interaction That Leads to Sucking.  
555 Advances in the Study of Behavior 22, 239–268.
- 556 Wassmuth R, Löer A and Langholz HJ 2001. Vigour of lambs newly born to outdoor wintering  
557 ewes. Animal Science 72, 169-178.
- 558



**Table 1** Mean monthly temperature values and rainfall during lambing seasons of D'man ewes.

	Autumn	Winter	Spring	Summer
Minimum temperature (°C)	8.5	2	5.9	15.6
Average temperature (°C)	19.5	11.8	18.9	28.4
Maximum temperature (°C)	32.9	23.4	35.4	46.7
Rainfall (mm)	17.3	8.3	1.2	0.1

565

566 **Table 2** Number of ewes giving birth at day or night time, maternal experience, and number  
 567 of lambs involved in the study according to lambing seasons

568

Parturition period			Maternal experience		Litter size		
Season	Day	Night	Multiparous	Primiparous	Single	Twin	Triplet +
Autumn	13	12	21	4	9	15	14
Winter	13	9	20	2	7	25	19
Spring	20	13	20	13	8	29	28
Summer	17	14	19	12	16	20	23

569

570

571

572

573 **Table 3** Behaviors of mother and young recorded during the first 2 h postpartum in D'man

574 sheep

---

**Ewes**

---

Litter expulsion time(s)	Interval between appearance of the first lamb at the vulva and expulsion of the last one.
--------------------------	---

First grooming (s)	Time interval between full expulsion of each lamb and first grooming given to each of them.
--------------------	---

Grooming (s)	Time spent grooming during the first 30 min after giving birth of the first lamb.
--------------	---

---

**Lambs**

---

Gets up on forelegs(s)	Lamb on chest, pushing up on knees, supporting part of body off the ground.
------------------------	---

Stands up(s)	Lamb supporting itself on all 4 feet for at least 5 s.
--------------	--

Attempts to suckle(s)	Lamb with head under the ewe's udder region.
-----------------------	--

---

575

576

577

**Figure 1** Postnatal grooming activity of D'man ewes according to lambing season: (a) time interval between full expulsion of each lamb and first grooming given each of them; (b) and total duration of grooming during the first 30 min after giving birth to the first lamb. Values are medians (bar within the box) and upper and lower quartiles (boarders of box) with 10 and 90 percentile. a,b :  $P < 0.05$

**Figure 2** D'man lambs' neonatal behavior according to season of birth: (a) latency to get up on forelegs, (b) latency to stand up, (c) latency to reach the udder. Values are medians (bar within the box) and upper and lower quartiles (boarders of box) with 10 and 90 percentile. a,b :  $P < 0.05$

**Figure 3** Birth weight of D'man lambs according to season of birth. Values are medians (bar within the box) and upper and lower quartiles (boarders of box) with 10 and 90 percentile. a,b :  $P < 0.05$

**Figure 4** Rectal temperature and glycaemia in D'man lambs according to season of birth: (a) within 1 h and (b) 48 h of birth, and (c) variations between these two times. Values are medians (bar within the box) and upper and lower quartiles (boarders of box) with 10 and 90 percentile. a,b,c :  $P < 0.05$ .




Figure 1




Figure 2




Figure 3




Figure 4