

Regardless of the deposition pathway, aminoacid 31 in histone variant H3 is essential at gastrulation in Xenopus

David Sitbon, Ekaterina Boyarchuk, Geneviève Almouzni

▶ To cite this version:

David Sitbon, Ekaterina Boyarchuk, Geneviève Almouzni. Regardless of the deposition pathway, aminoacid 31 in histone variant H3 is essential at gastrulation in Xenopus. 2020. hal-03041039

HAL Id: hal-03041039 https://hal.science/hal-03041039

Preprint submitted on 4 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

Regardless of the deposition pathway, aminoacid 31 in histone variant H3 is essential at gastrulation in Xenopus

3 4

5 Authors: David Sitbon^{1,2}, Ekaterina Boyarchuk^{1,2} and Geneviève Almouzni^{1,2,*}

- 6 ¹Institut Curie, PSL Research University, CNRS, UMR3664, Equipe Labellisée Ligue contre
- 7 le Cancer, 75005, Paris, France.
- ²Sorbonne Universités, UPMC Univ Paris 06, CNRS, UMR3664, 75005, Paris, France.
- 9 * Email corresponding author: genevieve.almouzni@curie.fr
- 10

11 Corresponding author: Dr. Geneviève Almouzni, 26 rue d'Ulm 75005 Paris,
12 +33156246701, genevieve.almouzni@curie.fr

13

14 Abstract

15 The closely related replicative H3 and non-replicative H3.3 variants show specific 16 requirement during development in vertebrates. Whether it involves distinct mode of 17 deposition or unique roles once incorporated into chromatin remains unclear. To disentangle 18 the two aspects, we took advantage of the Xenopus early development combined with 19 chromatin assays. Our previous work showed that in Xenopus, depletion of the non-20 replicative variant H3.3 impairs development at gastrulation, without compensation through 21 provision of the replicative variant H3.2. We systematically mutated H3.3 at each four 22 residues that differ from H3.2 and tested their ability to rescue developmental defects. 23 Surprisingly, all H3.3 mutated variants functionally complemented endogenous H3.3, 24 regardless of their incorporation pathways, except for one residue. This particular residue, the 25 serine at position 31 in H3.3, gets phosphorylated onto chromatin in a cell cycle dependent 26 manner. While the alanine substitution failed to rescue H3.3 depletion, a phosphomimic 27 residue sufficed. We conclude that the time of gastrulation reveals a critical importance of the 28 H3.3S31 residue independently of the variant incorporation pathway. We discuss how this 29 single evolutionary conserved residue conveys a unique property for this variant in 30 vertebrates during cell cycle and cell fate commitment.

- 31
- 32

33 Introduction

34 The organization of DNA into chromatin provides not only a means for compaction but also a 35 versatile landscape contributing to cell fate and plasticity (Schneider and Grosschedl 2007; 36 Yadav et al. 2018). The basic unit, the nucleosome core particle, is composed of a histone 37 tetramer (H3-H4)₂ flanked by two histone dimers H2A-H2B, around which 147bp of DNA 38 are wrapped as shown by crystal structure (Luger et al. 1997). Importantly, modulation of this 39 unit exploits the choice of histone variants and reversible PTMs (Post-translational 40 modification) to impact cell functions (Kouzarides 2007; Gurard-Levin and Almouzni 2014). 41 Three out of the four-histone families possess histone variants (Franklin and Zweidler 1977; 42 Sarma and Reinberg 2005; Buschbeck and Hake 2017; Sitbon et al. 2017; Talbert and 43 Henikoff 2017). In the H3 family, while the centromere-specific histone CENP-A 44 (Centromere protein A) is very distinct and marks the centromere, the other well-45 characterized H3 histone variants are closely related and thought to ensure similar functions 46 (Maze et al. 2014). Indeed, while they show similar structural features at the core particle 47 level (Tachiwana et al. 2011), an important difference concerned their cell cycle regulation 48 and distinct mode of incorporation (Mendiratta et al. 2018). In human, the two replicative 49 variants H3.1 and H3.2 are incorporated into chromatin in a DSC (DNA-synthesis coupled) 50 manner via the histone chaperone complex CAF-1 (Chromatin assembly factor 1) (Stillman 51 1986; Smith and Stillman 1989; Gaillard et al. 1996; Tagami et al. 2004; Polo et al. 2006; 52 Groth et al. 2007; Latreille et al. 2014). By contrast, the non-replicative form H3.3, which 53 differs from H3.1 and H3.2 by five and four residues respectively, is incorporated in a DSI 54 (DNA-synthesis independent) manner (Ahmad and Henikoff 2002b; Ray-Gallet et al. 2002; 55 Tagami et al. 2004; Ray-Gallet et al. 2011). This incorporation depends on the histone 56 chaperone complex HIRA (Histone regulator A) in euchromatin regions (Lamour et al. 1995; 57 Ray-Gallet et al. 2002; Tagami et al. 2004; Goldberg et al. 2010; Ray-Gallet et al. 2011), 58 while the presence of H3.3 at telomeres and pericentric heterochromatin rather relies on the 59 histone chaperone complex DAXX (Death domain-associated protein)/ATRX (Alpha 60 thalassemia/mental retardation syndrome X-linked) (Gibbons et al. 1995; Yang et al. 1997; 61 Drane et al. 2010; Goldberg et al. 2010; Lewis et al. 2010). Thus, the dynamics of the 62 different histone variants with their deposition is critically linked to dedicated histone 63 chaperones (Gurard-Levin et al. 2014; Hammond et al. 2017). Inspired by the quote "nothing 64 in biology makes sense except in the light of evolution" (Dobzhansky 1973), we considered 65 H3 variants in light of their conservation in different organisms. In Saccharomyces cerevisiae, 66 the only non-centromeric histone H3 is mostly related to H3.3, and achieves both replicative 67 and non-replicative variant functions, illustrating the capacity of survival with a single variant 68 (Dion et al. 2007; Jamai et al. 2007; Rufiange et al. 2007). Intriguingly, however, humanized

69 S. cerevisiae with all histories replaced by human orthologs survived better with hH3.1 than 70 hH3.3 (Truong and Boeke 2017). The better adaptation to H3.1 could argue for non-essential 71 roles of H3.3 in S. cerevisae, a unicellular organism. However, in metazoans like Drosophila 72 melanogaster, the replicative variant can compensate for the loss of H3.3 during development 73 in somatic tissues, although the adults are sterile (Loppin et al. 2005; Bonnefoy et al. 2007; 74 Hodl and Basler 2009; Sakai et al. 2009; Orsi et al. 2013). Since the sterility could simply 75 reflect a shortage of maternal H3.3 to replace protamine from sperm chromatin after 76 fertilization, the most parsimonious hypothesis suggests that the nature of the variant itself 77 might not necessarily be essential. Similarly, H3.3 is dispensable in *Caenorhabditis elegans*, 78 since its removal does not result in lethality but rather to reduced fertility and viability in 79 response to stress (Delaney et al. 2018). In Arabidopsis thaliana, in contrast, replicative and 80 non-replicative H3 variants are clearly essential. The absence of H3.3 leads to embryonic 81 lethality and also partial sterility due to defective male gametogenesis (Wollmann et al. 82 2017). In vertebrate models like *Mus musculus*, the deletion of one of the two copies of the 83 H3.3 gene leads not only to sterility but also to developmental defects at E12.5 (Couldrey et 84 al. 1999; Bush et al. 2013; Tang et al. 2015). Finally in human, while a complete absence is 85 not reported, dominant effects with substitutions in H3.3, like H3.3 K27M and H3.3 G34R/V, 86 along with mutations affecting their chaperones like DAXX/ATRX, have been involved in 87 distinct aggressive cancers (Heaphy et al. 2011; Jiao et al. 2011; Schwartzentruber et al. 2012; 88 Wu et al. 2012; Behjati et al. 2013; Lewis et al. 2013; Lindroth and Plass 2013; Fontebasso et 89 al. 2014; Taylor et al. 2014; Wu et al. 2014). Furthermore, not only mutation but also 90 expression of histone chaperones is often found altered in aggressive cancer (Corpet et al. 91 2011; Montes de Oca et al. 2015; Gurard-Levin et al. 2016). Thus, the developmental defects, 92 along with situation encountered in particular cancers, underline the importance of each H3 93 variant and their chaperones in vertebrates. Given the limited sequence difference between 94 H3 variants, a major puzzle is whether the need for a particular histone variant could reflect 95 either (i) a unique mode of incorporation and provision or (ii) a distinct identity once 96 incorporated into chromatin to drive their functions. While, the first hypothesis had been 97 largely favored, based on previous work including ours, the issue has never been formally 98 addressed. To disentangle these two possibilities, we decided to use *Xenopus laevis* as it is an 99 ideal model system to tackle this issue. Indeed, extensively characterized both in 100 developmental biology and chromatin studies (Laskey et al. 1977; Almouzni and Mechali 101 1988; Almouzni et al. 1990; Almouzni and Wolffe 1993), its external development permits direct access to embryos for observation and manipulation (Nieuwkoop and Faber 1994). 102 103 With retention of H3 variants in sperm (Katagiri and Ohsumi 1994) and only one replicative 104 histone H3.2, it offers a simpler situation while retaining amino acid sequence conservation 105 with human variants for both H3.2 and H3.3. Furthermore, following fertilization,

106 development starts by twelve rapid embryonic cell divisions, which include only S and M 107 phases (Newport and Kirschner 1982; Etkin 1988; Masui and Wang 1998). At the MBT 108 (midblastula transition), zygotic activation occurs concomitantly with a progressive 109 lengthening of the cell cycle, to reach, at gastrulation, a typical cell cycle with two gap 110 phases. In addition, cells begin to engage into defining different types with acquisition of 111 migration properties. Importantly, previous work in our laboratory revealed a specific 112 requirement of H3.3 during Xenopus laevis early development at the time of gastrulation 113 (Szenker et al. 2012). This work showed that depletion of endogenous H3.3 (further referred 114 to as H3.3) leads to severe gastrulation defects that cannot be rescued by providing the 115 replicative counterpart H3.2 but only by H3.3 itself. Interestingly, there are only four residues 116 that diverge between these two forms. A first region of divergence now known as the AIG 117 motif, in the globular domain of H3.3, accounts for histone variant recognition by dedicated 118 histone chaperones (Ahmad and Henikoff 2002a; Elsasser et al. 2012; Liu et al. 2012; 119 Ricketts et al. 2015). This first region is thus key for the choice of deposition pathways. The 120 other distinctive residue, a serine only present in H3.3, is located on the histone N-terminus, 121 at position 31, and can be phosphorylated (Hake et al. 2005; Wong et al. 2009; Hinchcliffe et 122 al. 2016). In our study, we systematically mutated the H3.3 histone variant in its distinct 123 residues to assess their ability to rescue the gastrulation defects and examined their 124 incorporation mode. In our assays, we find that mutations affecting the incorporation pathway 125 proved neutral to set specific H3.3 functions at the time of gastrulation. By contrast, the serine 126 at position 31 is critical in the rescue experiments after H3.3 depletion. In Xenopus, this 127 residue, which is notably conserved in human cells, gets phosphorylated during the cell cycle 128 with a peak in mitosis. Remarkably, in our rescue experiments, the phosphomimic form, 129 which cannot be dynamically modified, fully rescued. We discuss how this residue and the 130 charge at this site can contribute to establish particular chromatin states important in the 131 context of somatic cell cycle and development.

132

133 *Results*

134 Conservation of histone H3 variants and their chaperones to assay H3.3 defects at135 gastrulation in Xenopus.

While *Homo sapiens* presents two replicative H3 variants, H3.1 and H3.2, *Xenopus laevis* only possesses one replicative variant, H3.2. Both H3.2 and H3.3 are conserved with their human orthologs (Figure 1 A). Interestingly, the two H3 variants are almost identical and conserved through evolution (Waterborg 2012) (Figure 1 B). Two regions show differences in H3.2 and H3.3. The first one encompasses positions 87, 89 and 90 with a serine, a valine, and a methionine, known as the SVM motif in H3.2. Instead, these positions correspond to an

142 alanine, an isoleucine and a glycine, known as the AIG motif in H3.3. The second difference 143 lies at position 31 where H3.2 shows an alanine and H3.3 a serine. Considering sequences for 144 H3.2 and H3.3 histone variants from five different model organisms in which the function 145 along with the deposition pathways of H3 variants have been studied, the replicative variant 146 H3.2 shows around 3% (four variable residues out of 136). In the case of the non-replicative 147 variant H3.3, it varies of 4% differences, with six variable residues. Remarkably, the region 148 responsible for histone chaperone recognition shows the highest variations, in line with 149 possible coevolution with their respective histone chaperones that are not closely conserved 150 (Supplemental figure 1 A). To examine how deposition pathways and the role for histone 151 variants can be related, targeting these regions could thus be considered. Using Xenopus 152 *laevis* embryos, we had previously found that a morpholino against H3.3 leads to defects 153 during late gastrulation (Szenker et al. 2012) (Figure 1 C and Supplemental figure 1 B and 154 Film 1). While the blastopore forms and invaginates during gastrulation, depletion of H3.3 155 leads to an arrest of the blastopore closure. Exogenous HA-tagged H3.3, but not HA-tagged 156 H3.2 mRNAs (further referred as eH3s) when co-injected with the morpholino, can rescue the 157 phenotype (Szenker et al. 2012). To optimize the assay conditions, we decided to titrate the 158 concentration of morpholino (Supplemental figure 1 C). The lowest concentration enabled the 159 blastopore to start to invaginate without complete closure and lead to late gastrulation defects. 160 consistent with our previous findings (Szenker et al. 2012). By increasing by two to fourfold 161 morpholino concentrations, gastrulation defects coincide with an even earlier phenotype, 162 where the blastopore closure does not happen at all. These data are in line with a titration 163 effect, whereby gastrulation time allows to readily reveal H3.3 requirements. We could thus 164 use it as a read out to assess the role for a distinct mode of incorporation for H3 variants or a 165 need of the histone variant itself once incorporated.

166

Permuting individually each amino acid of the H3.3 AIG motif towards H3.2 still ensures H3.3 functions during *Xenopus laevis* early development.

169 To investigated the importance of the deposition pathway, we first look at the H3.3 histone 170 chaperone recognition motif (Figure 2 A). Incorporation into chromatin of the non-replicative 171 variant H3.3 occurs at any time of the cell cycle (DSI), and involves the HIRA complex in 172 gene-rich regions (Ahmad and Henikoff 2002b; Ray-Gallet et al. 2002; Tagami et al. 2004; 173 Ray-Gallet et al. 2011) and the DAXX/ATRX complex in heterochromatin regions (Gibbons 174 et al. 1995; Yang et al. 1997; Drane et al. 2010; Goldberg et al. 2010; Lewis et al. 2010). In 175 contrast, incorporation of the replicative variants is coupled to DNA synthesis (DSC) and is 176 mediated by the CAF-1 complex (Stillman 1986; Smith and Stillman 1989; Gaillard et al. 177 1996; Tagami et al. 2004; Polo et al. 2006; Groth et al. 2007; Latreille et al. 2014). 178 Importantly, structural studies enabled the identification of how histone chaperones

179 discriminate the distinct histone variants through a motif located in the globular domain of 180 histones (Elsasser et al. 2012; Liu et al. 2012; Ricketts et al. 2015). Both DAXX and HIRA 181 complexes bind to the AIG motif of H3.3, with a particular affinity for the glycine at position 182 90. Therefore, we mutated H3.3 in the histone chaperone recognition motif to test their ability 183 to rescue the loss of H3.3. We first designed single mutants for each residue of the AIG motif 184 (Figure 2 B). To our surprise, every single mutant for the motif, *i.e.* eH3.3 A87S, eH3.3 I89V 185 and eH3.3 G90M could rescue loss of H3.3, and embryo development occurred with the same 186 efficiency. We verified that all constructs expressed at similar levels in the embryo 187 (Supplemental figure 2 A) and incorporated into chromatin (Supplemental figure 2 B). Since 188 it was possible that early development may allow looser interactions with the dedicated 189 chaperones, we decided to completely change the recognition motif and assess both 190 development and incorporation means.

191

Swapping the H3.3 histone chaperone recognition motif with the one from H3.2 still enables *Xenopus laevis* early development yet switching the mode of incorporation.

194 Since the individual substitution in the H3.3 motif did not affect the developmental rescue in 195 vivo, we assessed whether mutation of all three residues of the motif would then affect H3.3 196 functions (Figure 3). To our surprise, in this context, this eH3.3 SVM hybrid form proved still 197 able to rescue the loss of H3.3 as eH3.3 WT (wild-type), while eH3.2 WT cannot. In addition, 198 such mutant form is expressed as well as incorporated into chromatin at similar levels in the 199 embryo (Supplemental figure 3 A and B). Thus, by substituting the H3.3 recognition motif for 200 its histone chaperone with the one from H3.2, we could ensure the rescue *in vivo*. Thus, the 201 next question was whether in this context, the histone chaperone recognition motif actually 202 determined the respective deposition pathways for each histone variant. The H3.3 dedicated 203 chaperone complex HIRA recognized best H3 carrying the H3.3 AIG motif (Figure 4 A). 204 However, p60, one subunit of the CAF-1 complex dedicated to H3.2, could perfectly 205 recognize eH3.3 SVM, *i.e.* when the whole H3.3 motif is substituted to the one of H3.2. This 206 shows that eH3.3 SVM can be recognized by CAF-1, arguing for a possible swap in the 207 means for incorporation. We confirmed this finding *in vivo* by immunoprecipitation of eH3 208 forms directly from embryos at gastrulation stage (Supplemental figure 4 A). We found that 209 p60 again only interacted with eH3 carrying the H3.2 SVM motif. Therefore, changing the 210 motif alters the chaperone interactions. We thus explored the potential impact on the mode of 211 histone variant incorporation. To test this aim, we performed chromatin assembly assays using extracts derived from Xenopus laevis eggs (Smythe and Newport 1991; Almouzni and 212 213 Wolffe 1993). We supplemented Xenopus egg extracts with eH3.3 WT, eH3.2 WT or eH3.3 214 SVM and monitored their incorporation into chromatin using sperm nuclei under conditions 215 allowing or not DNA synthesis (Figure 4 B). In interphase extracts, sperm DNA forms nuclei,

216 can replicate and reassemble chromatin whereas mitotic extracts lack this DNA replication 217 capacity. We isolated and analyzed sperm chromatin nuclei from interphase extracts, in 218 presence or absence of the DNA synthesis inhibitor aphidicolin. Regardless of DNA 219 synthesis, the incorporation of eH3.3 WT using sperm nuclei occurred with a similar 220 efficiency in the presence or absence of DNA synthesis. By contrast, eH3.2 WT incorporation 221 was severely affected by the presence of aphidicolin. Importantly, eH3.3 SVM incorporation 222 shows the same dependency on DNA replication, arguing that its incorporation mode is 223 switched toward a DNA-synthesis coupled mechanism. Consistently, p60 recruitment to 224 chromatin is, as expected, highly reduced when DNA synthesis is inhibited. We further 225 confirmed in mitotic extracts that only variant with the AIG motif could get incorporated 226 independently of DNA synthesis (Supplemental figure 4 B). Based on these data, we 227 conclude that the H3.3 histone variant could be efficiently provided regardless of the pathway for incorporation. Thus, defects at the time of gastrulation do not arise from a need for a 228 229 distinct incorporation pathway outside S phase independently of DNA synthesis. It rather 230 reflects an inherent feature of the variant when incorporated into chromatin. We thus 231 examined more closely if the only remaining specific residue of H3.3, aminoacid 31, could 232 account for this unique feature.

233

H3.3S31 residue is critical for *Xenopus laevis* early development and is phosphorylated during the cell cycle.

236 To address the role of the H3.3 residue at position 31, we first constructed a new H3.3 237 mutant, H3.3 S31A, containing an alanine instead of a serine at position 31 while keeping its 238 original AIG motif (Figure 5 A and Supplemental figure 5 A). Remarkably, eH3.3 S31A was 239 not able to rescue H3.3 depletion during Xenopus early development, indicating that the 240 specific serine of eH3.3 at position 31 cannot be substituted by an alanine, which is notably 241 the corresponding residue of H3.2, to fulfill H3.3 dedicated functions at this window of time 242 during development. Interestingly, this particular residue has first been found phosphorylated 243 in human cell line during mitosis (Hake et al. 2005). Furthermore, a threonine replaces this 244 serine in Arabidopsis, a residue that can possibly also undergo phosphorylation, albeit this has 245 not yet been documented. A key question was thus whether the actual need for H3.3 was 246 linked to the capacity of H3.3S31 to be phosphorylated. To this end, we examined H3.3S31 247 phosphorylation in the Xenopus system using a Xenopus A6 cell line derived from kidney. 248 By immunofluorescence, we detected a strong enrichment of H3.3S31p during mitosis 249 (Figure 5 B). Interestingly, when compared to a mitotic modification, H3S10p, common to 250 both H3 variant forms, its pattern was different, indicating a possible distinct function. While 251 H3S10p covers the edges of mitotic chromosomes, H3.3S31p is enriched at centric and

252 pericentric heterochromatin. This pattern, similar to previous observations in HeLa B cells 253 (Supplemental figure 5 B) as in (Hake et al. 2005), is consistent with observations in mouse 254 and monkey cell lines (Wong et al. 2009; Hinchcliffe et al. 2016). Kinases responsible for 255 H3.3S31 phosphorylation are still unclear, currently implicating either CHK1 (Chang et al. 256 2015) or Aurora B (Li et al. 2017). Using Xenopus sperm chromatin, we further characterized 257 the acquisition of this mark prior to or after replication in interphase extract or in extracts 258 pushed into mitosis (Figure 5 C). We did not detect any significant signal for H3.3S31p in the 259 soluble pool of H3.3 in either mitotic or interphase extracts, indicating that the modification is 260 likely acquired once the variant is incorporated into chromatin. In addition, we could not 261 detect any signal for this mark on non-remodeled sperm chromatin or on chromatin 262 assembled in interphase extracts. This mark was most enriched in the fraction corresponding 263 to isolated mitotic chromatin, in a profile similar to the H3S10p mark. We therefore conclude 264 that H3.3S31p showed predominantly a mitotic chromatin-dependent mark in both somatic 265 cells and reconstituted chromatin in early embryonic extracts, a mark imposed within 266 chromatin and not prior to deposition. The next question was whether this modification or 267 charge was also critical at the time of gastrulation in our rescue experiments.

268

The negative charge of H3.3S31 is essential during *Xenopus laevis* early development at gastrulation.

271 In order to explore a potential need for the phosphorylation of H3.3S31, we designed another 272 H3.3 mutant for the serine carrying an aspartic acid, eH3.3 S31D, which acts as a 273 phosphomimic version of this residue and carries a constitutive negative charge at position 31 274 that cannot be dynamically regulated by kinases and phosphatases (Figure 6 A). Remarkably, 275 this mutant form could rescue the depletion of H3.3 comparable to the eH3.3 WT. Notably, 276 all mutants were expressed and incorporated into chromatin in the same proportion 277 (Supplemental Figure 6 A and B) and we verified that these mutations on the H3.3 tail did not 278 indirectly alter the ability to interact with specific histone chaperones (Figure 6 B). 279 Especially, neither H3.3 S31A nor S31D mutations affected the mode of incorporation of 280 H3.3 in the Xenopus egg extracts-base chromatin assembly assays (Figure 6 C and 281 Supplemental figure 6 C). This enabled us to discard any defects that could be related to 282 inefficient incorporation. We can thus conclude that the residue at position 31 in H3.3, either 283 as a serine or as a negatively charged residue (phosphomimic), is specifically needed for the 284 function of the H3.3 variant once incorporated into chromatin as revealed at the time of 285 gastrulation during early development in the vertebrate Xenopus.

286

287 Discussion

By exploiting depletion/complementation assays in *Xenopus laevis* and monitoring the capacity to undergo gastrulation, we disentangle the critical role of H3.3 within chromatin through its aminoacid 31 independently of its mode of incorporation.

291

H3.3 incorporation into chromatin is important during Xenopus early development regardless of its mode of incorporation.

294 Replicative and non-replicative H3 variants do show distinct genome wide distributions 295 (Clement et al. 2018), with enrichment of H3.3 at enhancers, or proximal to telomeres and 296 centromeres as observed in several models in somatic cells and embryonic cells (Filipescu et 297 al. 2014). How these distinct patterns arise and evolve dynamically during development and 298 are then maintained in given lineages remains to be established. Moreover, how the unique 299 properties of H3.3 influence cell cycle related functions and/or cell fate programming is still 300 an open question. We first examined whether the mutations of the residues assigned to a key 301 role in the choice of the deposition pathway (Ahmad and Henikoff 2002a; Elsasser et al. 302 2012; Liu et al. 2012; Ricketts et al. 2015) affect the capacity to rescue the depletion of H3.3 303 (Figure 2 and Figure 3). Surprisingly, an eH3.3 form containing H3.2 recognition motif still 304 complements H3.3 functions in our developmental assay (Figure 3) even though this swap 305 between motifs does effectively alter the chaperone interactions and the incorporation 306 pathway (Figure 4). These data underline the fact that neither the chaperone interaction nor 307 the mode of incorporation of the variant is critical to enable H3.3 specific roles at this time of 308 development. Therefore, in the context of Xenopus early development, in sharp contrast with 309 what one would have anticipated, it is the presence *per se* of H3.3 into chromatin that proves 310 most important *in vivo*, regardless of the means to get incorporated, (Figure 7). This discovery 311 shed light on unique roles performed by the single amino acid S31 on H3.3, highlighting the 312 idea that every amino acid matters (Maze et al. 2014)!

313

314 H3.3S31 and a constitutive negative charge are essential for H3.3 functions during 315 Xenopus early development.

We showed by replacing the residue at position 31 in the N-terminal tail of H3.3 with the alanine found in H3.2 that such eH3.3 S31A mutant could not replace H3.3 (Figure 5 A). This proves that H3.3S31 is essential in our complementation assay after H3.3 depletion. Moreover, this serine on H3.3 can undergo phosphorylation while the alanine on H3.2 cannot. This specific phosphorylation of H3.3 regulated during the cell cycle shows enrichment during metaphase (Hake et al. 2005) (Figure 5 B). In our experiments, remarkably a phosphomimic form with a negative charge on the serine at position 31 could readily rescue

323 the depletion of H3.3 (Figure 6 A). However, the exact role of H3.3S31 remains enigmatic. 324 Although, we (Figure 5 and Supplemental figure 5) and others (Hake et al. 2005; Wong et al. 325 2009; Hinchcliffe et al. 2016) detected strong H3.3S31p signals primilary in mitosis, we 326 cannot exclude that discrete sites of transcription could be marked but undetected in our 327 assays. Indeed, this mark has been associated with particular transcribed regions in activated 328 macrophages (Thorne et al. 2012). Furthermore, in mammals, a function for H3.3S31p has 329 been linked to intron retention and pre-mRNA processing by preventing binding of the tumor 330 suppressor BS69, also known as ZMYND11 (Zinc finger MYND domain-containing protein 331 11), an H3.3K36me3-specific reader (Guo et al. 2014; Wen et al. 2014). Our assays 332 demonstrate the importance of H3.3S31 at gastrulation (Figure 5 and Figure 6), a critical time 333 for lineage commitment accompanied by dramatic cell cycle changes and modifications in 334 gene expression patterns. Taking into account the specific importance given to H3.3 that 335 ranges from transcription to reprogramming (Ahmad and Henikoff 2002b; Schwartz and 336 Ahmad 2005; Ng and Gurdon 2008; Jullien et al. 2012), the observed developmental defects 337 could arise from a role for H3.3S31 and its phosphorylation in transcription initiation or 338 maintenance. Importantly, MBT in Xenopus also leads to changes in the cell cycle 339 progression: establishment of the longer cell cycle "somatic-type" with the gap phases and 340 acquisition of checkpoints (Etkin 1988; Masui and Wang 1998). Moreover, histories are key 341 component in regulating the start of the MBT, possibly through titration mechanisms 342 (Almouzni and Wolffe 1995; Amodeo et al. 2015). It should be valuable to explore the role of 343 specific variants into this context. Thus, H3.3S31 function could equally depend on a specific 344 role of H3.3 role during the cell cycle. Indeed, H3.3 has been involved in chromosome 345 segregation as shown in double H3.3 KO embryonic stem cells that leads to an increase in 346 anaphase bridges and lagging chromosomes (Jang et al. 2015). Furthermore, this mark can 347 coat lagging chromosomes in parallel to p53 activation in order to prevent aneuploidy 348 (Hinchcliffe et al. 2016). Considering H3.3S31p accumulation at centromeres in mitosis 349 (Figure 5 B) (Hake et al. 2005), it is thus tempting to speculate that H3.3S31p could crosstalk 350 with CENP-A incorporation at late mitosis and the beginning of G1. Interestingly, CENP-A 351 incorporation at the centromere has been shown to be dependent on both the presence of H3.3 352 as a placeholder (Dunleavy et al. 2011) and on transcription (Ohkuni and Kitagawa 2011; 353 Bobkov et al. 2018). To reconcile both aspects, we could envision that H3.3S31p may act as a 354 phosphoswitch in mitosis to regulate transcription at critical chromosomal landmarks and in 355 interphase to control/maintain a transcription program. Alternatively, one could also consider 356 that the alanine present on H3.2 could prevent particular interactions specific to H3.3S31 and 357 permitted by an aspartic acid. Notably in Arabidopsis, the replicative form cannot be rescued 358 by H3.3 (i.e. A31T mutation) (Jiang and Berger 2017). Indeed, ATXR5 (Arabidopsis 359 trithorax-related protein 5), a plant specific H3K27 methyltransferase, specifically recognizes

360 H3.1 and not H3.3 (Jacob et al. 2014). Therefore, the alanine at position 31 on replicative 361 histone H3.1 is prevents the heterochromatinization of H3.3-rich regions during replication. 362 Thus, in plants, the replicative histone H3 variant avoids the presence of a negative charge at 363 position 31. In our work, we show that there is potentially no need for the dynamics of the 364 mark. Rather, we demonstrate a need to provide a negative charge or to avoid an alanine 365 during early development. Indeed, our data indicate that the dynamics of the H3.3S31p may 366 be dispensable, since the phosphomimic mutant is able to fully rescue H3.3 depletion (Figure 367 6). We can foresee several possible explanations. On the one hand, the negative charge at 368 position 31 could be an absolute requirement for H3.3 functions regardless of any dynamics. 369 On the other hand, the negative charge on this residue could be associated with mechanisms 370 that occur solely in mitosis and the dynamics may simply be ensured by removal of the 371 variant without invoking a particular phosphatase. More work, specifically exploring eviction 372 mechanisms is needed to disentangle these two possibilities. Thus, the residue at position 31 373 of the replicative and non-replicative H3 variant may be key to promote/allow or exclude the 374 presence of a negative charge with different outcomes for critical binding partners. 375 Conversely, other modifications on H3 tail may impact S31 and in this respect the existence 376 of mutations affecting neighboring residues will be interesting to explore. Notably, H3.3S31 377 is close to residues often found substituted in aggressive cancers like pediatric glioblastoma 378 (H3 K27M and G34R) (Heaphy et al. 2011; Jiao et al. 2011; Schwartzentruber et al. 2012; 379 Wu et al. 2014). Thus, it will be very interesting to evaluate the connection between these 380 particular cancers involving oncohistones with this H3.3 specific phosphorylation. Altogether, 381 we show that H3.3S31 is the key residue that confers specific functions to H3.3 within 382 chromatin. It establishes for the first time the importance of a distinct histone variant residue 383 for the proper development of a vertebrate during gastrulation. Future work should explore 384 whether a similar requirement also occurs in mammals to provide a comprehensive view of 385 the importance of the non-replicative variant H3.3 and its role during vertebrate development 386 and in disease states.

387

388 "Qu'importe le flacon, pourvu qu'on ait l'ivresse", Alfred de Musset

390 Material & Methods

391 Xenopus laevis embryo manipulation

392 We used Xenopus laevis adults (2 years old) from the Centre de Ressource Biologie Xenope. 393 We prepared embryos at 18°C as in (Almouzni et al. 1994) and staged them according to 394 (Nieuwkoop and Faber 1994). We acquired embryos during gastrulation with a MZFLIII 395 magnifier (Leica) and the SPOT software. Animal care and use for this study were performed 396 in accordance with the recommendations of the European Community (2010/63/UE) for the 397 care and use of laboratory animals. Experimental procedures were specifically approved by 398 the ethics committee of the Institut Curie CEEA-IC #118 (Authorization APAFIS#11226-399 2017091116031353-v1 given by National Authority) in compliance with the international 400 guidelines. David Sitbon, Ekaterina Boyarchuk and Geneviève Almouzni possess the 401 Authorization for vertebrates' experimental use.

402

403 Plasmid cloning and mRNA transcription

404 We cloned all H3 cDNAs in the p β RN3P vector (Zernicka-Goetz et al. 1996). This vector 405 stabilizes RNA and improves their translation efficiency while injected into Xenopus eggs. In 406 addition, an HA tag is inserted in the C-terminal of H3. We obtained mRNAs by in vitro 407 transcription of PCR-amplified fragments of the different pßRN3P vectors (Forward: 5'-408 gtaaaacgacggccagt-3' and Reverse: 5'-ggaaacagctatgaccatga-3'). We transcribed mRNAs starting with 5ng of PCR-amplified fragment, 10µL Buffer 5X, 5µL of DTT 100mM, 0.25µL 409 410 bovine serum albumin (BSA) 10mg/mL (NEB), 5µL of ATP, CTP, UTP 10mM, 1.65µL GTP 10mM (Sigma-Aldrich), 3.35µL Me7GTP 10mM (NEB), 2µL of RNasin Plus RNase 411 412 Inhibitor (Promega), and 50µL H2O qsp. After 10min on ice, we added 2µL of T3 RNA 413 Polymerase (Promega) to each sample as well as after 30min at 37°C for another 1h30. After 414 DNA digestion, we extracted mRNAs with phenol-chloroform and purified them through 415 Sephadex G-50 Quick Spin Column for radiolabeled RNA purification (Sigma-Aldrich), 416 previously equilibrated 6 times with 1mL of TE 10mM.

417

418 Morpholino and mRNA microinjection into *Xenopus laevis* embryo

We microinjected 2-cell embryos using a Brinkmann micromanipulator and a Drumond microinjector on two-cell stage eggs with an injection volume set to 9.2nL to deliver the appropriate quantity of morpholino and mRNAs (1X=4.6ng). Between 20 and 30 embryos are injected per condition, with at least 3 biological replicates. We knocked down H3.3 with morpholino designed to bind to the initiation region of *Xenopus laevis* H3.3 mRNAs (Szenker et al. 2012).

425

426 Xenopus laevis embryo protein extract preparations and Western blotting

We prepared total protein extracts from *Xenopus laevis* embryos using the CelLytic Express reagent (Sigma-Aldrich) and centrifuging at full speed the extracts after 30min incubation. We analyzed protein samples by electrophoresis on 4%–12% NuPAGE SDS-PAGE gels with MES SDS Running buffer (Life Technologies) and corresponding LDS buffer NuPage (Invitrogen) with DTT. Primary antibodies were detected using horse-radish-peroxidaseconjugated secondary antibodies (Jackson Immunoresearch Laboratories) and ECL2 kit (Pierce).

434

435 Xenopus laevis embryo immunoprecipitation

436 We collected embryos at stage 12 and centrifuged them at 25000g after 3 washes in TE buffer 437 (10mM Hepes, 75mM KCl and 50mM Sucrose). After ultracentrifugation at 150000g of the 438 supernatant, we used 100µg of each condition for every immunoprecipitation overnight at 439 4°C in 400µL of IP buffer (10mM Hepes, 150mM KCl, 0:05% NP-40, 5mM DTT and 440 protease inhibitors). After 3 washes, we eluted proteins and analyzed samples by 441 electrophoresis on 4%-12% NuPAGE SDS-PAGE gels with MOPS SDS Running buffer 442 (Life Technologies) and corresponding LDS buffer NuPage (Invitrogen) with DTT. Primary 443 antibodies were detected using horse-radish-peroxidase-conjugated secondary antibodies 444 (Jackson Immunoresearch Laboratories) and ECL2 kit (Pierce).

445

446 Xenopus laevis embryo fractionation

447 We collected 40 embryos at stage 12 and performed fractionation as in (Szenker et al. 2012). 448 We lysed the embryos in 200µL of Lysis Buffer 1 (10mM Tris-HCl pH 7.5, 200mM NaCl, 449 5mM MgCl2, 0.5% NP40, protease inhibitors) or Lysis Buffer 2 (10mM Tris-HCl pH 7.5, 450 10mM NaCl, 5mM MgCl2, 0;5% NP40, protease inhibitors) and centrifuged them at 1000g 451 for 2min at 4°C. We obtained the clear soluble fraction after ultracentrifugation of the 452 supernatant at 160000g for 1 hour at 4°C from Lysis Buffer 1. We washed the chromatin 453 fraction from Lysis Buffer 2 after centrifugation in Buffer A (10mM Tris-HCl pH 7.5, 15mM 454 NaCl, 60mM KCl, 0.34M sucrose, 1mM DTT, protease inhibitors). We performed MNase 455 (Roche Diagnostics, #1010792100) digestion during 10min with 11.25U/mL final 456 concentration, after addition of CaCl2 (2mM final) and RNase A (Roche Diagnostics, 457 #10109142001) treatment (75µg/ml final, 5min at 37°C). We stopped the reaction by EDTA 458 (50mM final) and recovered solubilized chromatin fraction after a centrifugation at 1000 g for 459 2min.

- 460
- 461
- 462

463 Xenopus laevis egg extract preparation

We prepared *Xenopus laevis* sperm nuclei and low-speed extracts arrested by CSF (cytostatic factor) of *Xenopus laevis* eggs as previously described (Kornbluth et al. 2001). Briefly, we collected eggs freshly and centrifuged them at low speed (16000g) to conserve the mitotic phase, below lipids. We induced interphase by the addition of CaCl₂ at the final concentration 0.06mM to CSF-arrested egg extracts. We added sperm chromatin at a concentration 1000– 4000 nuclei/µl. After DNA replication, a 2/3 volume of the corresponding CSF-arrested extract was added to induce mitosis.

471

472 Xenopus laevis egg extract immunoprecipitation

Briefly, we produced recombinant H3 mutant proteins from mRNAs using rabbit reticulocyte
lysate (Promega L4600). After 3h of incubation at 4°C in interphase extracts followed by
another 3h incubation with anti-HA beads, we pulled-down and washed the proteins in 0.8x
CSF-XB buffer (10mM Hepes-KOH, pH 7.7, 100mM KCl, 2mM MgCl2, 5mM EGTA)
containing 5% glycerol, 0.5% Triton X-100 and protease and phosphatase inhibitors.

478

479 Xenopus laevis sperm chromatin purification

480 We diluted fivefold 100 μ alignots of each reaction with 0.8× CSF-XB buffer containing 481 20mM β -glycerophosphate, 5% glycerol and 0.5% Triton X-100, which we incubated for 1 482 min at RT. We then layered the samples onto a 35% glycerol-containing CSF-XB cushion 483 and centrifuged them at 10000g for 5 min at 4°C. We resuspended the pellets in the same 484 buffer, and repeated the centrifugation step. For purification of interphase chromatin, we 485 diluted 100µl aliquots of extract with $0.8 \times$ CSF-XB buffer containing 20 mM β -486 glycerophosphate and 5% glycerol, which we incubated for 1 min at RT, followed by 487 centrifugation through the cushion at 10000g for 5 min at 4°C. We analyzed protein samples 488 by electrophoresis on 4%-12% NuPAGE SDS-PAGE gels with MES SDS Running buffer 489 (Life Technologies) and corresponding LDS buffer NuPage (Invitrogen) NuPage reducing 490 agent (Invitrogen). Primary antibodies were detected using horse-radish-peroxidase-491 conjugated secondary antibodies (Jackson Immunoresearch Laboratories). We revealed the 492 signal by chemiluminescence with SuperSignal West Dura Extended Duration Substrate 493 (ThermoFisher). The signal was acquired using ChemiDoc Imager (Biorad).

494

495 Chromatin assembly assays

496 We added 100000 sperm nuclei to each 150μ L of corresponding extracts with or without 497 aphidicolin (50μ g/mL). We then supplied 15μ L of the rabbit reticulocyte lysate used to 498 produce recombinant H3 mutant proteins from mRNAs. After 40min of incubation at room

499 temperature, we purified chromatin and we analyzed protein samples by electrophoresis as

500 described above.

501

502 WebLogo sequence alignment

- 503 After performing multiple sequence alignment using MUSCLE (Edgar 2004), we displayed
- the alignments using WebLogo3 (Crooks et al. 2004) with probability units.
- 505

506 Antibody list

Antibody List	WB	IF
HA (3F10, Roche)	1/1000	/
H3.3S10p (Mabi0312, Wako)	1/1000	1/1000
H3.3S31p (ab92628, Abcam)	1/1000	1/200
H3.3 (M01, Abnova)	1/200	/
H4 (05-858, Upstate)	1/1000	/
Tubulin (T9026, Sigma)	1/1000	/
H3 (ab1791, Abcam)	1/5000	/
xEmi2 (From Mary Dasso)	1/1000	/
xHIRA (Agro Bio)	1/1000	/
xDAXX (From Mary Dasso)	1/500	/
xp60 (Agro Bio)	1/1000	/
xCENP-A (From Aaron Straight)	1/1000	/

507

508 Immunofluorescence and epifluorescence microscopy

We first fixed A6 and HeLa B cells on coverslips for 20 min in 4% and 2% paraformaldehyde, respectively, before to permeabilize them with 0.2% Triton X-100. We blocked them for 45min with 5% bovine serum albumin. We then incubated coverslips with primary and secondary antibodies and stained them with DAPI. We mounted the coverslips in Vectashield medium. We used a Confocal Zeiss LSM780, and we acquired images using $63 \times /1.4$ NA under Zen blue software (Zeiss Germany).

515

682 Acknowledgments

We are grateful to Dominique Ray-Gallet, Shauna Katz and Jean-Pierre Quivy for critical 683 684 reading and to members of UMR3664 for helpful discussions. We thank Mary Dasso and 685 Aaron Straight for Xenopus histone chaperone antibodies. This work was supported by la 686 Ligue Nationale contre le Cancer (Equipe labellisée Ligue), ANR-11-LABX-0044 DEEP and ANR-10-IDEX-0001-02 PSL, ANR-12-BSV5-0022-02 "CHAPINHIB", ANR-14-CE16-0009 687 688 "Epicure", ANR-14-CE10-0013 "CELLECTCHIP", EU project 678563 "EPOCH28", ERC-689 2015-ADG- 694694 "ChromADICT", ANR-16-CE15-0018 "CHRODYT", ANR-16-CE12-690 0024 "CHIFT", "Parisian Alliance of Cancer Research Institutes". DS was supported by PSL. 691

091

692 Author contributions

D.S., E.B. and G.A. conceived the overall strategy and wrote the paper. G.A. supervised the
work. D.S. performed *Xenopus laevis* and cellular experiments and analyzed data. E.B.
performed *Xenopus laevis egg* extract experiments and analyzed data. Critical reading and
discussion of data involved all authors.

697

698 Competing interests

- 699 The authors declare no competing interests.
- 700

701 **References**

- Ahmad K, Henikoff S. 2002a. Histone H3 variants specify modes of chromatin assembly. *Proc Natl Acad Sci U S A* 99 Suppl 4: 16477-16484.
- -. 2002b. The Histone Variant H3.3 Marks Active Chromatin by Replication Independent Nucleosome Assembly. *Mol Cell* 9: 1191-1200.
- Almouzni G, Clark DJ, Mechali M, Wolffe AP. 1990. Chromatin assembly on replicating DNA in vitro. *Nucleic Acids Res* 18: 5767-5774.
- Almouzni G, Khochbin S, Dimitrov S, Wolffe AP. 1994. Histone acetylation
 influences both gene expression and development of Xenopus laevis. *Dev Biol*165: 654-669.
- Almouzni G, Mechali M. 1988. Xenopus egg extracts: a model system for chromatin
 replication. *Biochim Biophys Acta* 951: 443-450.
- Almouzni G, Wolffe AP. 1993. Nuclear assembly, structure, and function: the use of
 Xenopus in vitro systems. *Exp Cell Res* 205: 1-15.
- -. 1995. Constraints on transcriptional activator function contribute to transcriptional quiescence during early Xenopus embryogenesis. *Embo J* 14: 1752-1765.
- Amodeo AA, Jukam D, Straight AF, Skotheim JM. 2015. Histone titration against the
 genome sets the DNA-to-cytoplasm threshold for the Xenopus midblastula
 transition. *Proc Natl Acad Sci U S A* 112: E1086-1095.
- Behjati S, Tarpey PS, Presneau N, Scheipl S, Pillay N, Van Loo P, Wedge DC, Cooke
 SL, Gundem G, Davies H et al. 2013. Distinct H3F3A and H3F3B driver
 mutations define chondroblastoma and giant cell tumor of bone. *Nat Genet* 45: 1479-1482.
- Bobkov GOM, Gilbert N, Heun P. 2018. Centromere transcription allows CENP-A to
 transit from chromatin association to stable incorporation. *J Cell Biol* 217: 1957-1972.
- Bonnefoy E, Orsi GA, Couble P, Loppin B. 2007. The essential role of Drosophila
 HIRA for de novo assembly of paternal chromatin at fertilization. *PLoS Genet*3: 1991-2006.
- Buschbeck M, Hake SB. 2017. Variants of core histones and their roles in cell fate
 decisions, development and cancer. *Nat Rev Mol Cell Biol.*
- Bush KM, Yuen BT, Barrilleaux BL, Riggs JW, O'Geen H, Cotterman RF, Knoepfler
 PS. 2013. Endogenous mammalian histone H3.3 exhibits chromatin-related
 functions during development. *Epigenetics Chromatin* 6: 7.
- Chang FT, Chan FL, JD RM, Udugama M, Mayne L, Collas P, Mann JR, Wong LH.
 2015. CHK1-driven histone H3.3 serine 31 phosphorylation is important for
 chromatin maintenance and cell survival in human ALT cancer cells. *Nucleic Acids Res* 43: 2603-2614.
- Clement C, Orsi GA, Gatto A, Boyarchuk E, Forest A, Hajj B, Mine-Hattab J, Garnier
 M, Gurard-Levin ZA, Quivy JP et al. 2018. High-resolution visualization of
 H3 variants during replication reveals their controlled recycling. *Nat Commun*9: 3181.
- Corpet A, De Koning L, Toedling J, Savignoni A, Berger F, Lemaitre C, O'Sullivan
 RJ, Karlseder J, Barillot E, Asselain B et al. 2011. Asf1b, the necessary Asf1
 isoform for proliferation, is predictive of outcome in breast cancer. *Embo J* 30: 480-493.
- Couldrey C, Carlton MBL, Nolan PM, Colledge WH, Evans MJ. 1999. A retroviral
 Gene Trap Insertion into the Histone 3.3A Gene Causes Partial Neonatal

- Lethality, Stunted Growth, Neuromuscular Deficits and Male Sub-fertility in
 Transgenic Mice. *Hum Mol Genet* 8: 2489-2495.
- Crooks GE, Hon G, Chandonia JM, Brenner SE. 2004. WebLogo: A sequence logo generator. *Genome Res* 14: 1188-1190.
- Delaney K, Mailler J, Wenda JM, Gabus C, Steiner FA. 2018. Differential Expression
 of Histone H3.3 Genes and Their Role in Modulating Temperature Stress
 Response in Caenorhabditis elegans. *Genetics* 209: 551-565.
- Dion MF, Kaplan T, Kim M, Buratowski S, Friedman N, Rando OJ. 2007. Dynamics
 of Replication-Independent Histone Turnover in Budding Yeast. *Science* 315: 1405-1408.
- Dobzhansky T. 1973. Nothing in Biology Makes Sense Except in the Light of
 Evolution. American Biology Teacher 35: 125-129.
- Drane P, Ouararhni K, Depaux A, Shuaib M, Hamiche A. 2010. The death-associated
 protein DAXX is a novel histone chaperone involved in the replication independent deposition of H3.3. *Genes Dev* 24: 1253-1265.
- Dunleavy E, Almouzni G, Karpen GH. 2011. H3.3 is deposited at centromeres in S
 phase as a placeholder for newly assembled CENP-A in G1 phase. *Nucleus* 2:
 1-12.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and
 high throughput. *Nucleic Acids Res* 32: 1792-1797.
- 769 Elsasser SJ, Huang H, Lewis PW, Chin JW, Allis CD, Patel DJ. 2012. DAXX
 770 envelops a histone H3.3-H4 dimer for H3.3-specific recognition. *Nature* 491: 560-565.
- Etkin LD. 1988. Regulation of the mid-blastula transition in amphibians. *Dev Biol (N Y 1985)* 5: 209-225.
- Filipescu D, Muller S, Almouzni G. 2014. Histone H3 variants and their chaperones
 during development and disease: contributing to epigenetic control. *Annu Rev Cell Dev Biol* 30: 615-646.
- Fontebasso AM, Papillon-Cavanagh S, Schwartzentruber J, Nikbakht H, Gerges N,
 Fiset PO, Bechet D, Faury D, De Jay N, Ramkissoon LA et al. 2014.
 Recurrent somatic mutations in ACVR1 in pediatric midline high-grade
 astrocytoma. *Nat Genet* 46: 462-466.
- Franklin SG, Zweidler A. 1977. Non-allelic variant of histone 2a, 2b and 3 in
 mammals. *Nature* 266: 273-275.
- Gaillard PH, Martini EM, Kaufman PD, Stillman B, Moustacchi E, Almouzni G.
 1996. Chromatin assembly coupled to DNA repair: a new role for chromatin assembly factor I. *Cell* 86: 887-896.
- Gibbons RJ, Picketts DJ, Villard L, Higgs DR. 1995. Mutations in a putative Global
 Transcriptional Regulator Cause X-Linked Mental Retardation with a Thalassemia (ATR-X Syndrome) *Cell* 80: 837-845.
- Goldberg AD, Banaszynski LA, Noh KM, Lewis PW, Elsaesser SJ, Stadler S, Dewell
 S, Law M, Guo X, Li X et al. 2010. Distinct factors control histone variant
 H3.3 localization at specific genomic regions. *Cell* 140: 678-691.
- Groth A, Corpet A, Cook AJ, Roche D, Bartek J, Lukas J, Almouzni G. 2007.
 Regulation of replication fork progression through histone supply and demand. *Science* 318: 1928-1931.
- Guo R, Zheng L, Park JW, Lv R, Chen H, Jiao F, Xu W, Mu S, Wen H, Qiu J et al.
 2014. BS69/ZMYND11 Reads and Connects Histone H3.3 Lysine 36
 Trimethylation-Decorated Chromatin to Regulated Pre-mRNA Processing.
 Mol Cell 56.

700	
/99	Gurard-Levin ZA, Almouzni G. 2014. Histone modifications and a choice of variant:
800	a language that helps the genome express itself. F1000Prime Rep 6: 76.
801	Gurard-Levin ZA, Quivy JP, Almouzni G. 2014. Histone chaperones: assisting
802	histone traffic and nucleosome dynamics. Annu Rev Biochem 83: 487-517.
803	Gurard-Levin ZA, Wilson LO, Pancaldi V, Postel-Vinay S, Sousa FG, Reyes C,
804	Marangoni E, Gentien D, Valencia A, Pommier Y et al. 2016. Chromatin
805	Regulators as a Guide for Cancer Treatment Choice. Mol Cancer Ther 15:
806	1768-1777.
807	Hake SB, Garcia BA, Kauer M, Baker SP, Shabanowitz J, Hunt DF, Allis CD. 2005.
808	Serine 31 phosphorylation of histone variant H3.3 is specific to regions
809	bordering centromeres in metaphase chromosomes. Proc Natl Acad Sci USA
810	102 : 6344-6349.
811	Hammond CM, Stromme CB, Huang H, Patel DJ, Groth A. 2017. Histone chaperone
812	networks shaping chromatin function. Nat Rev Mol Cell Biol 18: 141-158.
813	Heaphy CM, de Wilde RF, Jiao Y, Klein AP, Edil BH, Shi C, Bettegowda C,
814	Rodriguez FJ. Eberhart CG. Hebbar S et al. 2011. Altered telomeres in tumors
815	with ATRX and DAXX mutations <i>Science</i> 333 . 425
816	Hinchcliffe EH Day CA Karanieet KB Fadness S Langfald A Vaughan KT Dong
817	7. 2016 Chromosome missegregation during anaphase triggers n53 cell cycle
818	arrest through histone H3 3 Ser31 phosphorylation Nat Cell Riol
819	Hodl M Basler K 2009 Transcription in the absence of histore H3 3 <i>Curr Riol</i> 19 :
820	1221_1226
821	Iacob V Bergamin E. Donoghue MT. Mongeon V. LeBlanc C. Voigt P. Underwood
822	CL Brunzelle IS Michaels SD Reinberg D et al 2014 Selective methylation
822	of histone H3 variant H3 1 regulates heterochromatin realization Science 343:
823	$12/0_{-}1253$
825	Jamai A. Imoberdorf RM. Strubin M. 2007. Continuous histore H2B and
825	transcription_dependent histore H3 exchange in yeast cells outside of
820	replication Mol Call 25: 345-355
828	Jang CW Shihata V Starmer I Vee D Magnuson T 2015 Histore H3 3 maintains
820	genome integrity during mammalian development Ganas Day 29 : 1377-1392
820	Jiang D. Berger F. 2017. DNA replication coupled histone modification maintains
830	Delycomb gene silencing in plants. Science
837	Jiao V Shi C Edil BH de Wilde RE Klimstra DS Maitra A Schulick PD Tang I H
832	Walfgang CL Chati MA at al 2011 DAXY/ATRY MENI and mTOP
835	nothway genes are frequently altered in paperentic neuroendocrine tumors
835	Science 331 : 1100 1203
835	Jullion I. Astrond C. Szonkor F. Corrott N. Almouzni G. Gurdon IP. 2012 HIPA
830 827	dependent H2 2 deposition is required for transcriptional representation
02/	following nuclear transfer to Vancous acoutes. Enigenetics Charactin 5:17
020	Keta sini C. Ohanni K. 1004. Dama daling of many shrangetin in decad in and strategy
839	Katagiri C, Onsumi K. 1994. Remodeling of sperm chromatin induced in egg extracts
840	of amphibians. Int J Dev Biol 38 : 209-216.
841	Kornbluth S, Yang J, Powers M. 2001. Analysis of the cell cycle using Xenopus egg
842	extracts. Current Protocols in Cell Biology.
843	Kouzarides 1. 2007. Chromatin modifications and their function. <i>Cell</i> 128 : 693-705.
844	Lamour V, Lecluse Y, Desmaze C, Spector M, Bodescot M, Aurias A, Osley MA,
845	Lipinski M. 1995. A human homolog of the S. cerevisiae HIR1 and HIR2
846	transcriptional repressors cloned from the DiGeorge syndrome critical region.
847	<i>Hum Mol Genet</i> 4 : 791-799.

- Laskey RA, Mills AD, Morris NR. 1977. Assembly of SV40 chromatin in a cell-free
 system from Xenopus eggs. *Cell* 10: 237-243.
- Latreille D, Bluy L, Benkirane M, Kiernan RE. 2014. Identification of histone 3
 variant 2 interacting factors. *Nucleic Acids Res* 42: 3542-3550.
- Lewis PW, Elsaesser SJ, Noh KM, Stadler SC, Allis CD. 2010. Daxx is an H3.3specific histone chaperone and cooperates with ATRX in replicationindependent chromatin assembly at telomeres. *Proc Natl Acad Sci U S A* 107: 14075-14080.
- Lewis PW, Muller MM, Koletsky MS, Cordero F, Lin S, Banaszynski LA, Garcia
 BA, Muir TW, Becher OJ, Allis CD. 2013. Inhibition of PRC2 activity by a
 gain-of-function H3 mutation found in pediatric glioblastoma. *Science* 340:
 859 857-861.
- Li M, Dong Q, Zhu B. 2017. Aurora Kinase B Phosphorylates Histone H3.3 at Serine
 31 during Mitosis in Mammalian Cells. *J Mol Biol* 429: 2042-2045.
- Lindroth AM, Plass C. 2013. Recurrent H3.3 alterations in childhood tumors. *Nat Genet* 45: 1413-1414.
- Liu CP, Xiong C, Wang M, Yu Z, Yang N, Chen P, Zhang Z, Li G, Xu RM. 2012.
 Structure of the variant histone H3.3-H4 heterodimer in complex with its chaperone DAXX. *Nat Struct Mol Biol* 19: 1287-1292.
- Loppin B, Bonnefoy E, Anselme C, Laurencon A, Karr TL, Couble P. 2005. The
 histone H3.3 chaperone HIRA is essential for chromatin assembly in the male
 pronucleus. *Nature* 437: 1386-1390.
- Luger K, Mäder AW, Richmond RK, Sargent DF, Richmond TJ. 1997. Crystal
 structure of the nucleosome core particle at 2.8 Å resolution. *Nature* 389: 251560.
- Masui Y, Wang P. 1998. Cell cycle transition in early embryonic development of
 Xenopus laevis. *Biol Cell* 90: 537-548.
- Maze I, Noh KM, Soshnev AA, Allis CD. 2014. Every amino acid matters: essential
 contributions of histone variants to mammalian development and disease. *Nat Rev Genet* 15: 259-271.
- Mendiratta S, Gatto A, Almouzni G. 2018. Histone supply: Multitiered regulation
 ensures chromatin dynamics throughout the cell cycle. *J Cell Biol*:
 ijcb.201807179.
- Montes de Oca R, Gurard-Levin ZA, Berger F, Rehman H, Martel E, Corpet A, de
 Koning L, Vassias I, Wilson LO, Meseure D et al. 2015. The histone
 chaperone HJURP is a new independent prognostic marker for luminal A
 breast carcinoma. *Mol Oncol* 9: 657-674.
- Newport J, Kirschner M. 1982. A major developmental transition in early Xenopus
 embryos: I. characterization and timing of cellular changes at the midblastula
 stage. *Cell* 30: 675-686.
- Ng RK, Gurdon JB. 2008. Epigenetic memory of an active gene state depends on
 histone H3.3 incorporation into chromatin in the absence of transcription. *Nat Cell Biol* 10: 102-109.
- 891 Nieuwkoop PD, Faber J. 1994. Normal Table of *Xenopus laevis*. *Daudin*.
- Ohkuni K, Kitagawa K. 2011. Endogenous transcription at the centromere facilitates
 centromere activity in budding yeast. *Curr Biol* 21: 1695-1703.
- Orsi GA, Algazeery A, Meyer RE, Capri M, Sapey-Triomphe LM, Horard B, Gruffat
 H, Couble P, Ait-Ahmed O, Loppin B. 2013. Drosophila Yemanuclein and
 HIRA cooperate for de novo assembly of H3.3-containing nucleosomes in the
 male pronucleus. *PLoS Genet* 9: e1003285.

- Polo SE, Roche D, Almouzni G. 2006. New histone incorporation marks sites of UV
 repair in human cells. *Cell* 127: 481-493.
- Ray-Gallet D, Quivy JP, Scamps C, Martini EM, Lipinski M, Almouzni G. 2002.
 HIRA Is Critical for a Nucleosome Assembly Pathway Independent of DNA
 Synthesis. *Mol Cell* 9: 1091-1100.
- Ray-Gallet D, Woolfe A, Vassias I, Pellentz C, Lacoste N, Puri A, Schultz DC,
 Pchelintsev NA, Adams PD, Jansen LE et al. 2011. Dynamics of histone H3
 deposition in vivo reveal a nucleosome gap-filling mechanism for H3.3 to
 maintain chromatin integrity. *Mol Cell* 44: 928-941.
- Ricketts DM, Frederick B, Hoff H, Tang Y, Schultz DC, Singh Rai T, Grazia Vizioli
 M, Adams PD, Marmorstein R. 2015. Ubinuclein-1 confers histone H3.3specific-binding by the HIRA histone chaperone complex. *Nat Commun* 6: 7711.
- Rufiange A, Jacques PE, Bhat W, Robert F, Nourani A. 2007. Genome-wide
 replication-independent histone H3 exchange occurs predominantly at
 promoters and implicates H3 K56 acetylation and Asf1. *Mol Cell* 27: 393-405.
- 914 Sakai A, Schwartz BE, Goldstein S, Ahmad K. 2009. Transcriptional and
 915 developmental functions of the H3.3 histone variant in Drosophila. *Curr Biol*916 19: 1816-1820.
- 917 Sarma K, Reinberg D. 2005. Histone variants meet their match. *Nat Rev Mol Cell Biol*918 6: 139-149.
- Schneider R, Grosschedl R. 2007. Dynamics and interplay of nuclear architecture,
 genome organization, and gene expression. *Genes Dev* 21: 3027-3043.
- Schwartz BE, Ahmad K. 2005. Transcriptional activation triggers deposition and
 removal of the histone variant H3.3. *Genes Dev* 19: 804-814.
- 923 Schwartzentruber J, Korshunov A, Liu XY, Jones DT, Pfaff E, Jacob K, Sturm D,
 924 Fontebasso AM, Quang DA, Tonjes M et al. 2012. Driver mutations in histone
 925 H3.3 and chromatin remodelling genes in paediatric glioblastoma. *Nature* 482:
 926 226-231.
- 927 Sitbon D, Podsypanina K, Yadav T, Almouzni G. 2017. Shaping Chromatin in the
 928 Nucleus: The Bricks and the Architects. *Cold Spring Harb Symp Quant Biol*929 82: 1-14.
- Smith S, Stillman B. 1989. Purification and Characterization of CAF-I, a Human Cell
 Factor Required for Chromatin Assembly during DNA Replication In Vitro.
 Cell 58: 15-25.
- Smythe C, Newport JW. 1991. Systems for the study of nuclear assembly, DNA
 replication, and nuclear breakdown in Xenopus laevis egg extracts. *Methods Cell Biol* 35: 449-468.
- Stillman B. 1986. Chromatin Assembly during S_40 DNA Replication In Vitro. *Cell*45: 555-565.
- Suzuki Y, Horikoshi N, Kato D, Kurumizaka H. 2016. Crystal structure of the
 nucleosome containing histone H3 with crotonylated lysine 122. *Biochem Biophys Res Commun* 469: 483-489.
- Szenker E, Lacoste N, Almouzni G. 2012. A developmental requirement for HIRAdependent H3.3 deposition revealed at gastrulation in Xenopus. *Cell Rep* 1: 730-740.
- Tachiwana H, Osakabe A, Shiga T, Miya Y, Kimura H, Kagawa W, Kurumizaka H.
 2011. Structures of human nucleosomes containing major histone H3 variants. *Acta Crystallogr D Biol Crystallogr* 67: 578-583.

- Tagami H, Ray-Gallet D, Almouzni G, Nakatani Y. 2004. Histone H3.1 and H3.3
 Complexes Mediate Nucleosome Assembly Pathways Dependent or
 Independent of DNA Synthesis. *Cell* 116: 51-61.
- Talbert PB, Henikoff S. 2017. Histone variants on the move: substrates for chromatin
 dynamics. *Nat Rev Mol Cell Biol* 18: 115-126.
- Tang MC, Jacobs SA, Mattiske DM, Soh YM, Graham AN, Tran A, Lim SL, Hudson
 DF, Kalitsis P, O'Bryan MK et al. 2015. Contribution of the Two Genes
 Encoding Histone Variant H3.3 to Viability and Fertility in Mice. *PLoS Genet*11: e1004964.
- Taylor KR, Mackay A, Truffaux N, Butterfield YS, Morozova O, Philippe C, Castel
 D, Grasso CS, Vinci M, Carvalho D et al. 2014. Recurrent activating ACVR1
 mutations in diffuse intrinsic pontine glioma. *Nat Genet* 46: 457-461.
- Thorne JL, Ouboussad L, Lefevre PF. 2012. Heterochromatin protein 1 gamma and
 IkappaB kinase alpha interdependence during tumour necrosis factor gene
 transcription elongation in activated macrophages. *Nucleic Acids Res* 40:
 7676-7689.
- Truong DM, Boeke JD. 2017. Resetting the Yeast Epigenome with Human
 Nucleosomes. *Cell* 171: 1508-1519 e1513.
- Waterborg JH. 2012. Evolution of histone H3: emergence of variants and
 conservation of post-translational modification sites. *Biochem Cell Biol* 90:
 79-95.
- Wen H, Li Y, Xi Y, Jiang S, Stratton S, Peng D, Tanaka K, Ren Y, Xia Z, Wu J et al.
 2014. ZMYND11 links histone H3.3K36me3 to transcription elongation and tumour suppression. *Nature* 508: 263-268.
- Wollmann H, Stroud H, Yelagandula R, Tarutani Y, Jiang D, Jing L, Jamge B,
 Takeuchi H, Holec S, Nie X et al. 2017. The histone H3 variant H3.3 regulates
 gene body DNA methylation in Arabidopsis thaliana. *Genome Biol* 18: 94.
- Wong LH, Ren H, Williams E, McGhie J, Ahn S, Sim M, Tam A, Earle E, Anderson
 MA, Mann J et al. 2009. Histone H3.3 incorporation provides a unique and
 functionally essential telomeric chromatin in embryonic stem cells. *Genome Res* 19: 404-414.
- Wu G, Broniscer A, McEachron TA, Lu C, Paugh BS, Becksfort J, Qu C, Ding L,
 Huether R, Parker M et al. 2012. Somatic histone H3 alterations in pediatric
 diffuse intrinsic pontine gliomas and non-brainstem glioblastomas. *Nat Genet*44: 251-253.
- Wu G, Diaz AK, Paugh BS, Rankin SL, Ju B, Li Y, Zhu X, Qu C, Chen X, Zhang J et
 al. 2014. The genomic landscape of diffuse intrinsic pontine glioma and
 pediatric non-brainstem high-grade glioma. *Nat Genet* 46: 444-450.
- Yadav T, Quivy JP, G. A. 2018. Chromatine plasticity: A versatile landscape that underlies cell fate and identity. *Science* 361: 1332-1336.
- Yang X, Khosravi-Far R, Chang HY, Baltimore D. 1997. Daxx, a novel Fas-binding
 protein that activates JNK and apoptosis. *Cell* 89: 1067-1076.
- Zernicka-Goetz M, Pines J, Siemering KR, Haseloff J, Evans MJ, Gurdon JB. 1996.
 An indelible lineage marker for Xenopus using a mutated green fluorescent protein. *Development* 122: 3719-3724.

992

516 Figure Legends

517 Figure 1: H3.3 is essential for gastrulation of Xenopus laevis. A) Best-studied non-518 centromeric H3 histone variants in Homo sapiens and Xenopus laevis. The two well-519 characterized forms of non-centromeric H3 variants correspond to the replicative histones 520 H3.1 and H3.2 and the non-replicative histone H3.3, depicted in purple and green 521 respectively. In human, the replicative H3 form is comprised of H3.1 and H3.2 that differ by 522 only one residue at position 96, a cysteine and a serine respectively. The non-replicative form 523 H3.3 shares more than 96% of identity with replicative forms, with five and four residue 524 differences with H3.1 and H3.2 respectively. Additionally, Xenopus laevis embryos possess 525 only one replicative histone variant, H3.2. Finally, histone sequences are conserved between 526 Homo sapiens and Xenopus laevis. B) Logo for H3 variant sequence differences of Homo 527 sapiens, Mus musculus, Drosophila melanogaster, Xenopus laevis, and Arabidopsis thaliana 528 after performing multiple sequence alignment using MUSCLE. We display this alignment 529 using WebLogo3. Specific histone variants are highlighted in green and purple. Histone 530 variants show an extreme conservation between species. C) Developmental assay to monitor 531 H3.3 functions. Morpholino and eH3.3 mRNA are injected at the 2-cell stage and associated 532 defects can be observed at the gastrulation stage if there is no rescue. Scale bar corresponds to 533 500µm. See film 1.

534

535 Figure 2: eH3.3 AIG single mutants rescue depletion of H3.3 during Xenopus laevis 536 early development. A) Highlights of the histone chaperone recognition motif residues of H3 537 variants. Dedicated histone chaperones recognize histone variants by the H3.2 SVM and H3.3 538 AIG motifs. Although both motifs are structurally similar, the main difference appears for the 539 residue 90 that is critical for histone chaperone binding. Crystal structure adapted from PDB 540 ID codes: 5B0Z (Suzuki et al. 2016) and 3AV2 (Tachiwana et al. 2011). B) Rescue assays 541 with H3.3 AIG single mutants. Injections are performed at 2-cell stage. The morpholino 542 against H3.3 induces gastrulation defects that are rescued by all eH3.3 with single mutations 543 of the AIG motif. Scale bar corresponds to 500µm. Quantification of properly developed 544 embryos after injections of the different H3.3 AIG mutant forms shows no difference of 545 rescue efficiency after H3.3 depletion. Each experiment has been reproduced at least 3 times 546 with a minimum of 30 embryos.

547

548 Figure 3: eH3.3 AIG triple mutant rescues depletion of H3.3 during *Xenopus laevis* early 549 development. Rescue assays with eH3.3 triple AIG mutant. Injections are performed at 2-cell 550 stage. The morpholino against H3.3 induces gastrulation defects that are not rescued by 551 eH3.2. In contrast, a triple mutant of H3.3 for the AIG motif, eH3.3 SVM, rescues the

phenotype while carrying the same SVM motif than H3.2. Scale bar corresponds to 500µm.
Quantification of properly developed embryos after injection of the eH3.3 triple AIG mutant
form shows similar rescue efficiency than eH3.3 WT after H3.3 depletion. Each experiment
has been reproduced at least 3 times with a minimum of 30 embryos.

556

557 Figure 4: Swapping the AIG motif to the SVM motif leads to changes in chaperone 558 interactions and histone modes of incorporation into chromatin in vivo A) 559 Immunoprecipitation of eH3 variant forms in interphase extract. Recombinant proteins are 560 produced in rabbit reticulocyte lysates and pulled down by their HA-tag after incubation. 561 HIRA only recognizes the AIG motif while p60 recognizes solely the SVM motif. * 562 corresponds to a non-specific band. B) Incorporation of eH3 variant forms into sperm 563 chromatin in interphase extract. Purified nuclei remodeled in the interphase extracts 564 supplemented with indicated eH3.3 in the presence or absence of aphidicolin were analyzed 565 by WB with indicated antibodies. While all eH3 variant forms are incorporated when both 566 DSC and DSI pathways are available, only the eH3 variant forms with the AIG motif can be 567 when only the DSI is available.

568

569 Figure 5: H3.3S31 is critical *Xenopus laevis* early development and is phosphorylated in

570 this model organism. A) Rescue assays with eH3.3 S31A mutant. Injections are performed at 571 2-cell stage. Compared with eH3.3 WT, eH3.3 S31A mutant form cannot rescue the 572 phenotype. Scale bar corresponds to 500µm. Quantification of properly developed embryos 573 after injection of the eH3.3 triple AIG mutant form shows similar rescue efficiency than 574 eH3.3 WT after H3.3 depletion. Each experiment has been reproduced at least 3 times with a 575 minimum of 30 embryos. B) 3D-distribution and timing of H3.3S31 phosphorylation in A6 576 cell line. H3.3S31p follows the same dynamics than H3S10p, but does not localize to the 577 same physical places. Scale bar represents 10 µm. C) Characterization of H3.3S31 578 phosphorylation in Xenopus cell-free extract. Non-remodeled sperm nuclei and nuclei 579 purified after remodeling in interphase or mitotic extracts, H3 PTMs are analyzed by WB. 580 H3.3S31p and H3S10p are enriched exclusively in sperm nuclei purified from mitotic 581 extracts. The marks are prevalently found in the chromatin and not in H3.3 soluble form. 582 Emi2 (Early mitotic inhibitor) serves as a control to confirm that extracts are in mitosis. CSF 583 (Cytostatic factor).

584

585 Figure 6: H3.3S31 negative charge is essential to rescue depletion of H3.3 during 586 Xenopus laevis early development.

587 A) Rescue assays with eH3.3 S31D mutant. Injections are performed at 2-cell stage. In 588 contrast to eH3.3 S31A, the use of an exogenous phosphomimic mutant, eH3.3 S31D,

589 rescues, as the eH3.3 WT, the defects during gastrulation. Scale bar corresponds to 500µm. 590 Ouantification of properly developed embryos after injection of the H3.3 S31D mutant form 591 shows similar rescue efficiency than eH3.3 WT after H3.3 depletion. In contrast, eH3.3 S31A 592 mutant form leads to the same proportion of defects than eH3.2 after H3.3 depletion. Each 593 experiment has been reproduced at least 3 times with a minimum of 30 embryos. B) 594 Immunoprecipitation of eH3 S31 mutant forms in interphase extract. Recombinant proteins 595 are produced in rabbit reticulocyte lysates and pulled down by their HA-tag after incubation. 596 Mutations of the residue 31 do not alter histone chaperone interactions. * corresponds to a 597 non-specific band. C) Incorporation of eH3 S31 mutant forms into sperm chromatin in 598 interphase extract. Purified nuclei remodeled in the interphase extracts supplemented with 599 indicated eH3.3 in the presence or absence of aphidicolin were analyzed by WB with 600 indicated antibodies. Mutations of the residue 31 do not change the mode of incorporation of 601 these forms.

602

Figure 7: Graphical abstract. Defects associated with H3.3 depletion can be rescued with
H3 histone variants carrying a potential negatively charge residue regardless of the mode of
incorporation.

606

Film 1: Endogenous H3.3 depletion leads to gastrulation defects and stops *Xenopus laevis* early development. Left embryo is injected with eH3.3 WT mRNA while the middle embryo is injected with 4.6ng of morpholino against H3.3. Right embryo is injected with both morpholino and eH3.3 WT mRNA. Injection of the morpholino leads to a gastrulation defects followed by developmental arrest that can be rescued by co-injection of eH3.3 WT mRNA.

612

Film 2: Dose-dependent H3.3 depletion defects. From left to right, the three embryos have
been injected with 4.6ng, 9.2ng, 18.4ng of morpholino against H3.3. Right embryo has not
been injected. Two replicates for each condition are showed. Injection of increased
morpholino concentration leads to earlier gastrulation defects than previously observed.

617

618 Supplemental figure 1: Histone chaperone sequence conservation and morpholino 619 strategy and titration. A) Logo for UBN1, DAXX and p60 partial sequences of Homo 620 sapiens, Mus musculus, Drosophila melanogaster, Xenopus laevis after performing multiple 621 sequence alignment with MUSCLE. Complete histone-binding domain HRD (131-171) of 622 UBN1 is presented, while only a part (from 300 to 339) of the DAXX histone-binding domain 623 HBD (178-389) is displayed. No crystallographic data are available regarding the CAF1 624 complex but the overall sequence shows variation as exemplified by the 41 first residues. 625 Histone chaperones show little conservation between species. Final alignment was made

626 using WebLogo3. B) Sequences of the morpholino against endogenous H3.3 and its targeted 627 sequence. The corresponding sequences matching the targeted sequence are highlighted in 628 green for the first 10 codons of Xenopus laevis H3F3B and eH3 sequences. In contrast, 629 differences with the targeted sequence are emphasized in red. C) Titration of the morpholino 630 against H3.3. Increased concentrations of morpholino lead to earlier developmental defects. 631 Morpholino against H3.3 is injected at the 2-cell stage with 3 different doses. While the lower 632 one previously used in (Szenker et al. 2012) induces a defect at late gastrulation, two and four 633 times more morpholino induce a defect appearing during early gastrulation. See film 2.

634

635 Supplemental figure 2: Single mutant forms of eH3.3 AIG motif are all expressed and 636 incorporated in a similar fashion in vivo. A) Western blot to control eH3.3 AIG single 637 mutant expression in embryos. Every eH3.3 AIG single mutants are expressed at similar 638 levels in whole embryo extracts at stage 12. B) Western blots to control eH3.3 AIG single 639 mutant incorporation in embryos. Fractionation of stage 12 embryos shows that all mutants 640 are insoluble at 600nm NaCl, and therefore most likely incorporated into chromatin. Note that 641 both eH3.3 WT and Control conditions are reused in Supplemental Figure 3 B and 6 B since 642 each condition used comes from the same western blot.

643

644 Supplemental figure 3: Triple mutant form of eH3.3 AIG motif is efficiently expressed 645 and incorporated in vivo. A) Western blot to control eH3.3 AIG triple mutant expression in 646 embryos. eH3.3 AIG triple mutant expression compares with the other eH3 conditions in 647 whole embryo extracts at stage 12. B) Western blots to control eH3.3 AIG triple mutant 648 incorporation in embryos. Fractionation of stage 12 embryos shows that this mutant remains insoluble at 600nm NaCl, and therefore most likely incorporated into chromatin. Note that 649 650 both eH3.3 WT and Control conditions are reused in Supplemental Figure 2 B and 6 B since 651 each condition used comes from the same western blot.

652

653 Supplemental figure 4: p60 histone chaperone interaction in vivo and mode of 654 incorporation of eH3.3 AIG triple mutant in mitotic extract in vitro. A) 655 Immunoprecipitation of eH3 variant forms from stage 12 embryos. Only the SVM motif is 656 recognized by p60 in vivo. B) Incorporation of eH3 variant forms into sperm chromatin in 657 mitotic extract. Purified nuclei remodeled in the CSF-arrested extracts supplemented with 658 eH3.3 mutants were analyzed by WB with indicated antibodies. In mitotic extract where the 659 only pathway available is the DSI, only the eH3 variant form with the AIG motif, *i.e.* eH3.3 660 WT, is incorporated.

661

662 Supplemental figure 5: H3.3S31 phosphorylation in human has the same dynamics that

in *Xenopus laevis*. A) Highlights of the histone tail specific residue of H3 variants. H3.3 tail
possesses a serine at position 31 that can be phosphorylated, while the alanine of H3.2 cannot.
Crystal structure adapted from PDB ID codes: 5B0Z (Suzuki et al. 2016) and 3AV2
(Tachiwana et al. 2011). B) 3D-distribution and timing of H3.3S31 phosphorylation in HeLa
B cell line. As in the Xenopus, H3.3S31p follows the same dynamics than H3S10p, but does
not localize to the same physical places. Scale bar represents 10 µm.

669

670 Supplemental figure 6: Mutant forms of H3.3S31 are all expressed and incorporated in

671 **a similar fashion.** A) Western blot to control eH3.3S31 mutant expression in embryos.

672 eH3.3S31 mutant expression compares with the other eH3 conditions in whole embryo 673 extracts at stage 12. B) Western blots to control eH3.3S31 mutant incorporation in embryos. 674 Fractionation of stage 12 embryos shows that these mutant remains insoluble at 600nm NaCl, 675 and therefore most likely incorporated into chromatin. Note that both eH3.3 WT and Control 676 conditions are reused in Supplemental Figure 2 B and 3 B since each condition used comes 677 from the same western blot. C) Incorporation of eH3.3S31 mutant forms into sperm 678 chromatin in mitotic extract. Purified nuclei remodeled in the CSF-arrested extracts 679 supplemented with eH3.3 mutants were analyzed by WB with indicated antibodies. In mitotic 680 extract where the only pathway available is the DSI, mutations of the residue 31 do not

681 change the mode of incorporation of these forms.

bioRxiv preprint first posted online Apr. 17, 2019; doi: http://dx.doi.org/10.1101/612515. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. Figure 1: H3.3 IS essential tord cost to display the preprint in perpetuity.





bioRxiv preprint first posted online Apr. 17, 2019; doi: http://dx.doi.org/10.1101/612515. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. Figure 3: eH3.3 AIC is the period of the second second

MO H3.3 +

100

XX

√

 \checkmark

In vivo morpholino perturbation and rescue with the eH3.3 AIG triple mutant









Western blot

Western blot

bioRxiv preprint first posted online Apr. 17, 2019; doi: http://dx.doi.org/10.1101/612515. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. Figure 6: H3.3S31 negative on arge 18 negative on a second second second and the preprint in perpetuity. H3.3 during Xenopus laevis early development

A In vivo morpholino perturbation and rescue with the eH3.3S31 phosphomimic mutant





B Interaction of histone chaperones with the eH3.3S31 mutants



Mode of eH3.3S31 mutants incorporation into sperm chromatin in interphase extract







bioRxiv preprint first posted online Apr. 17, 2019; doi: http://dx.doi.org/10.1101/612515. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. Supplemental figure is 2 add any for the preprint of the second secon

expressed and incorporated in a similar fashion in vivo





B Incorporation into chromatin of eH3.3 AIG single mutants



bioRxiv preprint first posted online Apr. 17, 2019; doi: http://dx.doi.org/10.1101/612515. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity if is efficiently Supplemental figure is date appendice of the author/funder. The copyright holder for this preprint is perpetuity if is efficiently expressed and incorporated in vivo



В Incorporation into chromatin of eH3.3 AIG triple mutant





Supplement dynamics t	int first posted only peer-reviewed) tal figur hat in Xe	ine Apr. 17, 2019; do is the author/funder Bis Dade al alabi ENOPUS Ia	bi: http://dx.doi.org/1 , who has granted bi DE CONCE NEVIS	0.1101/612515. The ioRxiv a license to di 30100194011	copyright holder for splay the preprint in ON IN NUI	this preprint perpetuity, nan has	the same
A Aminoacid H3.2 H3.3	31 in H3 varia N-ter tail <section-header></section-header>	ants 31 KSAPA KSAPS	TGG —— TGG —— ∞ Site of a vai	riant-specific m	odification		
B Dynamics	of H3.3S31p i	ة ° o n HeLa B cell lir	ne	·			
-	H3S10p	H3.3S31p	DAPI	Merge	1		
Interphase							
Prophase							
Metaphase							
Anaphase			No.				

bioRxiv preprint first posted online Apr. 17, 2019; doi: http://dx.doi.org/10.1101/612515. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. Supplemental figure is ded with one for PNS of and is 31 are all expressed and incorporated in a similar fashion



C Mode of eH3.3S31 mutant incorporation into sperm chromatin in mitotic extract

