



HAL
open science

A Lagrangian Analysis of Water Vapor Sources and Pathways for Precipitation in East China in Different Stages of the East Asian Summer Monsoon

Yi Shi, Zhihong Jiang, Zhengyu Liu, Laurent Li

► **To cite this version:**

Yi Shi, Zhihong Jiang, Zhengyu Liu, Laurent Li. A Lagrangian Analysis of Water Vapor Sources and Pathways for Precipitation in East China in Different Stages of the East Asian Summer Monsoon. *Journal of Climate*, 2020, 33 (3), pp.977-992. 10.1175/JCLI-D-19-0089.1 . hal-03047303

HAL Id: hal-03047303

<https://hal.science/hal-03047303>

Submitted on 8 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 A Lagrangian analysis of water vapor sources and pathways for
2 precipitations in East China in different stages of the East Asian
3 summer monsoon

4
5 YI SHI AND ZHIHONG JIANG*

6 *Key Laboratory of Meteorological Disaster of Ministry of Education, Collaborative*
7 *Innovation Center on Forecast and Evaluation of Meteorological Disasters, Nanjing*
8 *University of Information Science & Technology, Nanjing, 210044, China*

9 ZHENGYU LIU

10 *Atmospheric Science Program, Department of Geography, The Ohio State University,*
11 *Columbus, Ohio, USA*

12 LAURENT LI

13 *Laboratoire de Météorologie Dynamique, CNRS, Sorbonne Université, Ecole Normale*
14 *Supérieure, Ecole Polytechnique, Paris, France*

15
16 *Corresponding author address: Zhi-Hong Jiang, Key Laboratory of Meteorological Disaster
17 of Ministry of Education, Collaborative Innovation Center on Forecast and Evaluation of
18 Meteorological Disasters, Nanjing University of Information Science and Technology, 219
19 Ningliu Rd., Nanjing 210044, China.

20 Email: zhjiang@nuist.edu.cn

21

Abstract

22
23 The Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) platform is used to
24 simulate Lagrangian trajectories of air parcels in East China during the summer monsoon. The
25 investigation includes four distinct stages of the East Asian Summer Monsoon (EASM)
26 during its seasonal migration from south to north. Correspondingly, the main water vapor
27 channel migrates from the West Pacific Ocean (PO) for the pre-monsoon in South China (SC)
28 to the Indian Ocean (IO) for the monsoon in SC and in the Yangtze-Huaihe River Basin, and
29 finally back to PO for the terminal stage of monsoon in North China. Further calculations
30 permit to determine water vapor source regions and water vapor contribution to precipitations
31 in East China. To a large extent, moisture leading to precipitation doesn't come from the
32 strongest water vapor pathways. For example, the proportions of trajectories from IO are
33 larger than 25% all the time, but moisture contributions to actual precipitations are smaller
34 than 10%. This can be explained by the large amount of water vapor lost in the pathways
35 across moisture-losing areas such as the Indian Peninsula and Indochina Peninsula. Local
36 water vapor recycling inside East China (EC) contributes significantly to regional
37 precipitations, with contributions mostly over 30%, although the trajectory proportions from
38 sub-regions in EC are all under 10%. This contribution rate can even exceed 55% for the
39 terminal stage of monsoon in North China. Such a result provides an important guidance to
40 understand the role of land surface conditions in modulating rainfall in North China.

41 **Key words:** Lagrangian trajectory, East Asian Summer Monsoon, Moisture transport pathways,
42 Moisture source

43

44 1. Introduction

45 Unlike tropical summer monsoons, such as the Indian monsoon or the West African
46 monsoon, the East Asian summer monsoon (EASM) is characterized by the presence of a
47 front between warm air masses from the south and cold air masses from the north (Wang,
48 2006; Chang, 2004). A rain belt is formed along the front. During the seasonal course in
49 summer, the rain belt marks a clear migration from south to north, making different rainy
50 periods in different regions in East China. It is well recognized that three major periods of
51 rainfall take place in South China, the Yangtze River basin and North China (Zhou and Yu,
52 2005; Ding et al., 2008; Shi et al., 2009; Sun et al., 2011; Zhu et al., 2011; Sun and Wang,
53 2015). The rain belt, whatever its geographic position during its migration course from south
54 to north, is always strongly related to moisture transport which is a necessary condition
55 leading to precipitation (Huang et al. 1998; Jiang et al., 2006; Sun et al., 2011). It is of great
56 significance to quantitatively determine the moisture source of the monsoon rainfall for its
57 different phases.

58 Early researches mainly used the Eulerian approach to analyze moisture transport
59 pathways. For the rainy period in South China, Lin et al. (2014) showed that southwesterly
60 water vapor transports carrying moisture from the Bay of Bengal, West Pacific and from the
61 southwest side of the Tibetan Plateau are major pathways. For the Meiyu rainy season in the
62 Yangtze River Valley, Jiang et al. (2009) showed that water vapor is mainly from the southern
63 Indian Ocean, the East African coast and the Arabian Sea. For rainfalls in North China, Xie
64 and Ren (2008) indicated that water vapor mainly comes from the Bay of Bengal, South
65 China Sea/west Pacific Ocean and the mid latitude westerlies. These researches mainly focus

66 on the moisture transport channels during EASM, and indicated the significant changes for
67 different stages of the summer monsoon.

68 However, the simple analysis of moisture transport with an Eulerian vision is unable to
69 assess moisture origins from remote regions. It generally ignores any moisture changes during
70 the transport, although some Eulerian variants with an online implementation of moisture
71 tagging technique in global or regional climate models were proved to be very useful to
72 understand water vapor sources (see Insua-Costa et al. 2018 and the relevant references there).
73 When such a water vapor tracer scheme was incorporated into the regional model WRF,
74 Dominguez et al. (2016) and Eiras-Barca et al. (2017) were able to investigate the origin of
75 moisture for the North American monsoon and for two events of atmospheric river landfall
76 causing catastrophic precipitations.

77 The most advanced methods that are widely used in recent years to determine moisture
78 origin are based on Lagrangian models (Stohl and James, 2004, 2005; Dominguez et al. 2006;
79 Dirmeyer et al. 2009), capable of calculating backward or forward trajectories of air parcels
80 residing over the target region. They can trace pathways of the moisture transport and changes
81 of physical quantities along the pathways. The Lagrangian methodology has been highly
82 valued in investigating global moisture transport and regional moisture recycling. We can cite
83 a few works to show the diversity of research topics that used the Lagrangian approach.
84 Firstly, at global scale, Gimeno et al. (2010) presented results on the moisture contribution for
85 each continent from the global ocean in different seasons. At regional scale, Brubaker et al.
86 (2001) and Diem and Brown (2006) calculated the moisture sources for precipitations in the
87 Mississippi River Basin and in South West America. Bertò et al. (2004), Perry et al. (2007),

88 Sodemann and Stohl (2009), Bottyán et al. (2014) demonstrated that water vapor for rainfalls
89 in South Europe come, for a large proportion, from the Mediterranean Sea.

90 In East China, there are also numerous studies using Lagrangian models. Drumond et al
91 (2011a) revealed the main moisture source in different regions in China. Sun and Wang (2014,
92 2015) detected the origins of moisture over East China and semiarid grassland. However,
93 different researches are still quite divergent. Chen et al. (2013) showed that moisture of
94 summer rainfall in the Yangtze-Huaihe River Basin is mainly from the South China Sea and
95 the Bay of Bengal. Drumond et al. (2011a) indicated that the moisture of East China is mainly
96 from the East China Sea, the moisture of South China is mainly from the Bay of Bengal and
97 the Arabian Sea. In recent works, Sun and Wang (2015) affirmed that the water vapor for
98 precipitations in the Yangtze-Huaihe River Basin and in North China is mainly from local
99 evaporation and from the West Pacific Ocean.

100 The divergence of results in the scientific literature has two major sources. First, the
101 target area in East China is different among different publications. There has been a lack of
102 systematic investigation for different regions and for the whole seasonal course when the East
103 Asian summer monsoon migrates from south to north. Second, since the calculations of
104 Lagrangian trajectories require heavy computation, many existing studies have been limited to
105 short periods, which may impact the stability and robustness of their results. It is thus
106 desirable to study different phases during the seasonal migration course of the rainfalls in East
107 China for a long time period.

108 In this article, based on the HYSPLIT platform of Lagrangian trajectories, we
109 systematically analyze the variation of water vapor transport in East China during different

110 phases of EASM. Our analysis covers the period from 1961 to 2010. Advanced diagnostics
111 are performed to assess moisture transport pathways, moisture sources and contributions to
112 precipitation in each phase of EASM.

113 The article is organized as follows: the data and analyzing methods are described in
114 Section 2. Section 3 presents main results on the water vapor sources and contributions from
115 different regions during the northward propagation of EASM. It furthermore quantifies the
116 proportion of trajectories and the contribution of water vapor to precipitations during the
117 seasonal course of EASM. Conclusions are drawn in the last Section.

118 **2. Data and methods**

119 *2.1 Data*

120 We used the reanalysis data taken from NOAA-National Centers for Environmental
121 Prediction (NCEP)/National Center for Atmospheric Research (NCAR). Global pressure-level
122 data were reprocessed into the HYSPLIT compatible format in the Air Resources Laboratory
123 (ARL), NOAA, U.S. They are available since 1948 at 6 hourly temporal resolution and 2.5°
124 (latitude/longitude grid) spatial resolution with 17 levels in the vertical. Data have been
125 downloaded from the ARL's archives (http://ready.arl.noaa.gov/gbl_reanalysis.php).

126 *2.2 Determination of the rainy season course in East China*

127 The rainy season course in East China is directly taken from the *National Climate*
128 *Bulletin* published annually by the National Climate Center and the Central meteorological
129 observatory (<http://cmdp.ncc-cma.net/cn/monitoring.htm#>). As commonly practiced (Jiang et
130 al., 2006, Sun and Wang, 2014, 2015, Zhou and Yu, 2005 and Ding et al., 2008), the rainy
131 season course in East China is divided into three periods, and characterized by abundant

132 rainfalls in three geographical regions which are, from south to north, South China (SC), the
133 Yangtze-Huaihe River (YHR) basin and North China (NC). Data period covers from 1961 to
134 2010. The three areas are shown in Figure 1 with the following coordinates: 20°N–26°N,
135 106°E–120°E; 28°N–34°N, 110°E–123°E; 35°N–43°N, 110°E–120°E.

136 The situation in SC can be divided into two distinct sub-periods separated by the onset of
137 the South China Sea (SCS) summer monsoon which induces different characteristics of
138 atmospheric circulation in South China and associated water vapor transport (Shen et al.,
139 1982; Chen and Zhu et al., 1991; Lin et al., 2014). Before the onset of SCS summer monsoon,
140 the water vapor transport in South China is associated with westerly wind belt and South
141 China Sea sea-level pressure high. With the onset and strengthening of the SCS monsoon, the
142 cross-equatorial flow from the Southern Hemisphere intensifies and connects with the water
143 vapor channel from the Bay of Bengal. Regarding the onset date of SCS monsoon, we use that
144 defined by Wang et al. (2004), that is, the day when the 850 hPa zonal winds averaged over
145 the central SCS (5°–15°N, 110°–120°E) shift from westerlies to easterlies. Figure 2 shows the
146 beginning and ending dates of the four periods characterizing the rainy season course of East
147 China from 1961 to 2010. The average start days of the four periods are Apr 6, May 17, Jun 8
148 and July 16. The average end days are May 16, Jun 1, July 17 and Aug 18. As shown in Fig. 2,
149 there is a large interannual variability for the timing and duration of these four sub-periods,
150 which obligates us to use precise beginning and end dates in our statistics of trajectories.

151 *2.3 Methods*

152 2.3.1 HYSPLIT water vapor tracking simulation

153 The National Oceanic and Atmospheric Administration (NOAA) Hybrid Single-Particle

154 Lagrangian Integrated Trajectory (HYSPLIT) model (Draxler and Hess, 1998; Draxler and
155 Rolph, 2011) is used in this paper for trajectories and tracks of air parcels. The back-trajectory
156 algorithm is applied to all stations inside the three target areas. From their initial position,
157 back-trajectories are calculated to cover all times between the beginning and the end of each
158 of the four sub-periods of the monsoon propagation course. We use six levels at 100, 500,
159 1500, 3000, 5000 and 9000 m as our initial height of particles release. They roughly
160 correspond to the near surface layer, levels at 925, 850, 700, 500 and 300 hPa respectively.
161 All parcels were integrated backward in time, until 10 days. Outputs were recorded every 6
162 hours with variables indicating the position (latitude, longitude, and altitude) and
163 meteorological conditions (temperature, specific humidity, and geopotential height) for
164 further analysis and diagnostics (Jiang et al. 2017). In particular, we averaged all trajectories
165 in each of the five main moisture source areas, which helps us to get actual moisture transport
166 pathways.

167 2.3.2 Evaporation-minus-Precipitation diagnosis method (E-P) to deduce source/sink regions
168 of water vapor

169 Water vapor in an air parcel changes when evaporation or precipitation takes place,
170 increasing with the former and decreasing with the latter. It is thus clear that a correct
171 determination of the source and sink characteristics of the water vapor transport has to take
172 into account processes of evaporation and precipitation following the trajectory. Stohl and
173 James (2004, 2005) proposed such a diagnostic method, commonly called
174 Evaporation-minus-Precipitation method to consider the change of moisture content with time
175 as the air parcel travels. The budget equation can be written as follows:

176
$$e - p = m \frac{dq}{dt} \quad (1)$$

177 where e and p are the rates of evaporation and precipitation, respectively, q is the specific
 178 humidity and m is the mass of the air parcel. For a given air parcel, when q decreases,
 179 precipitation is larger than evaporation, and the parcel loses water vapor. When q increases,
 180 evaporation is larger than precipitation, and the parcel gains water vapor. Summing the $e-p$ of
 181 all air parcels residing in the atmospheric column over an area A , the surface net water flux in
 182 the area can be given by the following equation:

183
$$E - P \approx \frac{\sum_{k=1}^K (e-p)}{A} \quad (2)$$

184 where E and P are the total evaporation and precipitation in the column and K is the number
 185 of particles over the area. Equation (2) can be regarded as a Lagrangian transformation of the
 186 Eulerian atmospheric moisture budget equation:

187
$$E - P = \frac{\partial W}{\partial t} + \nabla \cdot \frac{1}{g} \int_0^{P_s} q \vec{v} dp \quad (3)$$

188 where $W = \frac{1}{g} \int_0^{P_s} q dp$ is the vertically-integrated water vapor or precipitable water in the
 189 column, g is the gravitational acceleration, \vec{v} is wind. p is atmospheric pressure and P_s is
 190 pressure at surface. The last term in Eq. 3 represents the divergence of vertically-integrated
 191 water vapor flux. Stohl and James (2004, 2005) showed that when using a large enough
 192 number of particles, the Lagrangian and Eulerian methods give almost identical results. So,
 193 the E-P results can be obtained, and the overall water vapor budget in a certain region can be
 194 further determined. The case $E-P > 0$ indicates that the area is source of water vapor to the
 195 precipitation in target region, whereas when $E-P < 0$ the area is sink to the precipitation to the
 196 target region. More details of this Evaporation-minus-Precipitation method can be found in
 197 Stohl and James (2004, 2005) and Sodemann and Zubler (2010).

198 2.3.3 The improved areal source–receptor attribution method

199 The Evaporation-minus-Precipitation diagnosis can be used to determine the regional
200 source and sink characteristics of the water vapor, but it cannot quantify the composition of
201 actual contribution of various water vapor sources to the precipitation in the considered area.
202 For example, a region of high positive Evaporation-minus-Precipitation value can almost
203 certainly be considered as a moisture source region. However, the air parcels passing over this
204 region to a target region may not carry a large amount of moisture from this source region into
205 the target region. This issue of water vapor contribution from a source point to a target point
206 can be assessed if one performs a further simple (forward) operation on the identified
207 trajectories with a consideration of water vapor evolution (evaporation and precipitation)
208 along the trajectories. Sodemann et al. (2008) and Martius et al. (2013) reported interesting
209 results about moisture source attribution within this simple framework.

210 However, this attribution calculation is only valid from a source point to a target point. It
211 fails when one wants to evaluate an area for either the source or the target (or both). As a
212 remedy of this drawback, Sun and Wang (2014) introduced a method called the areal
213 source-receptor attribution to evaluate the contribution of a specific water vapor source region
214 to the precipitation in a target region. When the air parcel is between the source and target
215 regions, a normal operation is applied to calculate water vapor changes in function of
216 evaporation (precipitation) that increases (decreases) water vapor. The specificity of the areal
217 source-receptor attribution method is in its special treatments when the air parcel is in the
218 source and target regions. Details can be found in Step 5 of the algorithm which is
219 decomposed into seven steps and presented in Sun and Wang (2014).

220 In our study, we further improve the methodology with two new aspects. First, we take
221 into account the water vapor in the initial air parcels (when the back trajectories reach a level
222 below 50 m and cannot go further), while Sun and Wang (2014) put this initial water vapor
223 content at zero. This enhances the overall contribution rate of water vapor. Second, as also
224 suggested in Huang et al. (2018), we consider a gaining of water vapor when parcels flow
225 below the boundary layer top, where vertical mixing is dominant. Air parcels in this case are
226 considered gaining water vapor as from the ground, while an increase of specific humidity
227 above the boundary layer is mainly affected by a phase change of water vapor. In the latter
228 case, we cannot directly assign the region as a source region.

229 A third improvement that we implemented in the initial algorithm of Sun and Wang
230 (2014) is on the consideration of source region. The initial algorithm took into account only a
231 well-determined source/target duality. In our work here, we extend the source region to an
232 ensemble of unit surface areas of 1 by 1 latitude/longitude. This permits us to introduce the
233 concept of Contribution Density Function (CDF), while Sun and Wang (2014) could calculate
234 only the contribution proportion from source to target.

235 To summarize, we can now calculate, for each trajectory j departing from the grid i , the
236 quantity $R_j(i)$ representing the moisture gained from the unit surface area i and then released
237 in the target region. Let R_{total} be the precipitation falling down in the target. It should be the
238 general sum of $R_j(i)$, i.e. the total moisture release from all trajectories (Sodemann et al., 2008;
239 Sun and Wang, 2014, 2015; Chu et al., 2017). But it is actually a little larger than the general
240 sum, due to the lack of trajectory calculations at very high layers. The contribution of water
241 vapor from each unit area can be assessed with the Contribution Density Function (CDF),

242 defined and calculated as:

$$243 \quad \text{CDF}(i) = \frac{\sum_{j=1}^n R_j(i)}{R_{total}} \quad (4)$$

244 where n is the number of trajectories departing from the area i . CDF behaving as a 2-D
245 probability density function can serve as a measure to evaluate the moisture contribution rate
246 of a unit area i to the precipitations falling into the target area.

247 With the improvement that we implemented, the global integration of CDF is
248 significantly increased, which indicates that we can more accurately identify water vapor
249 sources for the precipitation. For example, the global sum of CDF for North China monsoon
250 rainfall is about 0.8 with the original method (Sun and Wang, 2015), and about 0.9 with our
251 modifications.

252 **3. Results**

253 Based on the methodology described in Section 2.3, we simulated the backward
254 trajectory of parcels for EASM. From their initial positions shown in Figure 1,
255 back-trajectories are calculated to cover all times during the four sub-periods of monsoon
256 propagation in Figure 2 for 10 days.

257 *3.1 Pre-monsoon in South China*

258 For the pre-monsoon, rainfall mainly occurs in South China. Although it is before the
259 onset of the South China Sea (SCS) summer monsoon, rainfall is still considered as part of
260 EASM. We first analyze the moisture transport pathways (or channels) obtained as the mean
261 trajectory which is the average of all back-trajectories departing from our pre-defined
262 geographic sectors. Figure 3a shows the main water vapor transport pathways (or channels)
263 during the pre-monsoon in South China. The trajectories proportions are also shown in

264 relation to the channels. For the pre-monsoon in South China, the most important water vapor
265 transport channel is from the West Pacific Ocean accounting for 33.3% of all trajectories. Air
266 parcels come from the West Pacific Ocean, move northwestward across the Philippines, and
267 finally enter the target area at the south boundaries. The average specific humidity of air
268 parcels increases from 10 to 14 g/kg, which indicates that a significant portion of moisture is
269 absorbed along the trajectories. The secondary moisture transport channel is from the Indian
270 Ocean, which accounts for 24.6% of all trajectories. The average trajectories move across the
271 Bay of Bengal, Indochina Peninsula and then move into South China. Contrary to the West
272 Pacific channel, the average specific humidity of air parcels from the Indian Ocean decreases
273 from 12 to 8 g/kg, which indicates a great deal of water vapor loss. 14.2% of the trajectories
274 are from the Westerlies in the north, and their average specific humidity is small, increasing
275 from 2 to 7 g/kg. Besides, the proportion of local trajectories is 6.5% and the South China Sea
276 channel accounts for about 7.9% and the average specific humidity is 14 g/kg, the largest
277 among all the channels. These results are consistent with our general expectation, and in
278 agreement with Chen and Luo (2018) who also concluded that the main water vapor transport
279 channel during pre-monsoon period in South China is from the Pacific Ocean. Nevertheless,
280 the trajectory proportion obtained from our research is a bit higher for the Pacific Ocean and
281 lower for the South China Sea.

282 The vertically-integrated water vapor transports within the Eulerian framework are
283 shown in Figure 3b. The pre-monsoon in South China is mainly influenced by the West
284 Pacific moisture transport channel. The circulation from southeast over the West Pacific
285 carries abundant water vapor into the Indochina Peninsula and then affects South China.

286 Compared to the water vapor channel from the West Pacific Ocean, the one from the Indian
287 Ocean is relatively weak, but it also plays an important role for the pre-monsoon in South
288 China. These results are consistent with the existing literature for this regard. For example,
289 Chi et al. (2005) and Lin et al. (2014) showed already that before the onset of SCS summer
290 monsoon, the subtropical high dominates the South China Sea and moisture reaching South
291 China mainly comes from west winds over the Arabian Sea, east winds at the southwest flank
292 of the Western Pacific Subtropical High and southwest winds over the Bay of Bengal.

293 However, the transport pathways cannot give precise moisture source/sink characteristics.
294 Based on the evaporation-minus-precipitation diagnosis, Figure 3c shows the spatial
295 distribution of average E-P from 1 to 10 days before air parcels reaching South China. It
296 clearly reveals that the West Pacific is the most important moisture source region for the
297 pre-monsoon in South China, and the maximum value is located in offshore areas of East
298 China. East China and the South China Sea are also important moisture source regions.
299 Moisture sink region for the precipitation in South China is mainly in the Indochina Peninsula,
300 which indicates great moisture loss from the Indian Ocean.

301 The spatial distribution of moisture contribution leading to precipitations of
302 pre-monsoon in South China (water vapor CDF) is shown in Figure 3d, obtained from the
303 improved areal source–receptor attribution calculation. Large contribution density values are
304 mainly located near the coasts of Southeast China, and the adjacent seas, with an average
305 contribution density about 2×10^{-3} . In the West Pacific Ocean, the CDF of 1.0×10^{-5} extends to
306 170°E , and that of 1.0×10^{-4} extends to 150°E . However, CDF in the Indian Ocean monsoon
307 region is relatively small, only the Bay of Bangel, the Indian subcontinent and the Indochina

308 Peninsula show CDF values larger than 1.0×10^{-5} .

309 In summary, for the pre-monsoon in South China, the main moisture transport channel
310 originates from the West Pacific which is consistent with a previous work of Chen and Luo
311 (2018). The most important moisture source is also from the West Pacific providing abundant
312 water vapor that is actually converted to precipitation in South China.

313 *3.2 Monsoon in South China*

314 After the onset of SCS summer monsoon, the entire regional atmospheric circulation
315 jumps to a new state, EASM actually starts and enters into a stronger period. But main
316 monsoonal precipitations remain in South China. Main moisture transport channels are shown
317 in Figure 4. EASM is remarkably different from what it is in the precedent period. We can see,
318 in Figure 4a, that the Indian Ocean moisture channel becomes the strongest one. The
319 trajectories proportion increases from 24.6% (before the onset of SCS summer monsoon) to
320 53.9%. More than half of the trajectories come from the Indian Ocean. The average specific
321 humidity increases from 10 to 14 g/kg, which also indicates that water vapor for this period is
322 mainly from the Indian Ocean. At the same time, the West Pacific Ocean shows an obviously
323 weaker contribution. Only 22.0% of the trajectories come from the West Pacific Ocean and
324 the corresponding pathway is a bit northward shifted, compared to the former period. Besides,
325 trajectories from the westerlies and from local areas are also weakened, their trajectories
326 proportions are only 6.5% and 4.9% respectively. These results are also consistent with
327 previous works of Chen and Luo (2018) and Chu et al (2017). They divided the Indian Ocean
328 channel into two channels and the sum of these two channels makes the total trajectories
329 proportion at a level of about 50% which is very close to our result. The vertically-integrated

330 water vapor transport, calculated within the Eulerian framework, confirms the conclusion
331 (Figure 4b). During this period, the southwest monsoonal flow from the Indian Ocean
332 strengthens rapidly and the whole South Asian monsoon system goes quickly into its mature
333 stage, with a strong Somalian cross-equatorial flow, and monsoonal flows over the Arabian
334 Sea, the Indian Peninsula, the Bay of Bengal, and the Indochina Peninsula, and finally over
335 the South China Sea. This monsoonal flow inducing rainfalls in South China is now the main
336 moisture channel. The West Pacific Ocean channel is rapidly weakened and its influence
337 reduced. Such behaviors are in total agreement with what described in the current literature
338 (e.g. Chi et al. 2005, and Li et al. 2014, among many others) which also show that after the
339 onset of SCS summer monsoon, the subtropical high withdraws out of the South China Sea,
340 and the powerful cross-equatorial flow crosses the Bay of Bengal and transports moisture
341 towards South China. So, the Indian Ocean channel is the most important moisture transport
342 pathway in this stage.

343 The spatial distribution of E-P is also calculated and shown in Figure 4c. We can see that
344 after the onset of SCS summer monsoon, the maximum value of E-P is in north part of the
345 South China Sea, moisture source from the Indian Ocean is stronger and that from the Pacific
346 Ocean is weaker. As for moisture CDF (Figure 4d), the biggest value is in South China and
347 north part of the South China Sea where the CDF value is larger than 2.0×10^{-3} . CDF in the
348 South China Sea is about 5.0×10^{-4} . Compared to the rainy season before the onset of SCS
349 summer monsoon, intensity of CDF is decreased in the West Pacific Ocean. Contours CDF of
350 1.0×10^{-5} and 1.0×10^{-4} extend to 165°E and 135°E , respectively. This shows a clear westward
351 shift. On the contrary, CDF in the Indian Ocean is obviously increased. CDF values for most

352 part of the Indian Ocean are larger than 1.0×10^{-5} . CDF in the Bay of Bangel is larger than
353 1.0×10^{-4} .

354 During the monsoon in South China, the Indian Ocean channel is the most important
355 moisture transport channel, which includes more than half of the trajectories, and the
356 contribution rate also increases a lot. Meanwhile the West Pacific Ocean impact is reduced.
357 Besides, the biggest value of the moisture contribution is in South China and north part of the
358 South China Sea.

359 *3.3 Monsoon (Meiyu) in the Yangtze-Huaihe River basin*

360 For the Meiyu period, the EASM propagates to the Yangtze-Huaihe River Basin, and
361 leads to persistent rainfalls in this region. The moisture transport characters for this period
362 have both similarities and differences with the former monsoon stage. During Meiyu, the IO
363 channel is still the most important one and the portion of trajectories reaches to 42.1%. The
364 PO channel (22.7%) and Westerly channel (13.5%) are a bit strengthening. The spatial
365 distribution of the vertically-integrated water vapor transport (Figure 5b) shows that the IO
366 channel, the SCS channel and the PO channel converge in South China and move northward
367 to converge in the Yangtze-Huaihe River Basin region, resulting in plum rain in this period.
368 Xu et al. (2004, 2008), Zuo et al. (2009), Wei et al. (2012) and Jiang et al. (2013) also indicate
369 that IO channel is the most important water vapor channel in Yangtze-Huaihe River basin.

370 The spatial distribution of E-P (Figure 5c) also shows some similarities with the former
371 monsoon stage, main moisture source is found in the West Pacific Ocean, East China, the
372 South China Sea and the Indian Ocean. It is interesting to note that the Indian Ocean channel
373 is the main moisture pathway, but the Indian Ocean does not constitute the main moisture

374 source. In fact, as shown in Figure 5c, the pathway from the Indian Ocean goes through the
375 India Peninsula and the Indochina Peninsula, a significant part of moisture from the Indian
376 Ocean is lost along the trajectories. By consequence, in the spatial distribution of CDF (Figure
377 5d), the Indian Ocean is not the most important moisture source with a contribution density to
378 precipitation only about 1.0×10^{-5} to 1.0×10^{-4} . The maximum of the contribution density, larger
379 than 1.0×10^{-3} , is in the Yangtze-Huaihe River Basin itself and in South China. Compared to
380 the monsoon in South China, the Meiyu period shows CDF decreasing in south and increasing
381 in north. CDF in the Indian Ocean and South China Sea is in decrease, but the region of CDF
382 exceeding 1.0×10^{-5} extends to 50°N .

383 For the Meiyu stage of EASM in the Yangtze-Huaihe River Basin, the most important
384 moisture channel is still that from the Indian Ocean, which comprises 43.4% of the
385 trajectories. But it has great moisture loss along the trajectories, so the moisture contribution
386 for the precipitation is relatively small. The most important moisture contribution region is the
387 Yangtze-Huaihe River Basin and South China.

388 *3.4 Terminal stage of monsoon in North China*

389 North China rainy season is the terminal stage of EASM. As the monsoon moves to
390 North China, the rainy belt also shifts to North China and can stay for a few weeks. During
391 the monsoon terminal stage in North China, the most important moisture transport paths
392 change from IO to mid-latitude westerlies and to PO (figure 6a), the proportions for these two
393 channels being 30.3% and 26.8% respectively. As shown in Figure 6b, the IO channel, the
394 SCS channel and the PO channel move northward to North China, affecting the local rainfall.
395 Mid-latitude westerlies also influence the rain belt. It should be noticed that, although the

396 number of trajectories in mid-latitude westerlies is the biggest, specific humidity of this
397 channel is small, so the vertically-integrated water vapor transport in this channel is relatively
398 small. Our results confirm Jiang et al. (2017) who showed that Eurasia has the maximum
399 parcels and the Pacific Ocean has the second largest number of parcels during the rainy
400 season in North China.

401 Figure 6c is the spatial distribution of E-P for the terminal stage of EASM in North
402 China. The main moisture source regions for the rainfall in North China are the eastern
403 coastal areas of China, the northern part of the South China Sea and the Yangtze-Huaihe River
404 Basin. Main moisture sink regions are North China, South China and the Indochina Peninsula.
405 The spatial distribution of CDF to precipitation in North China indicates that the maximum
406 value is still near the target area, and the average contribution density exceeds 5.0×10^{-3} . CDF
407 in the south is further reduced, only the South China Sea and small areas east of Taiwan show
408 CDF values higher than 1.0×10^{-4} . Meanwhile, CDF in the north is significantly increased. The
409 region contour 1.0×10^{-5} reaches to Balkhash Lake in the west and 60°N in the north. On the
410 whole, the main moisture transport channels during the terminal stage of monsoon in North
411 China are those from the mid-latitude westerlies and the West Pacific Ocean, with trajectories
412 rates at 30.3% and 26.8% respectively. But there is low specific humidity from the westerlies
413 channel, so the westerlies channel has little influence for the monsoon in North China. The
414 most important moisture sources are from East China.

415 *3.5 Moisture transport evolution during the northward propagation of EASM*

416 From what shown above, it is clear that, accompanying the northward propagation of
417 EASM, the main water vapor channel experiences profound changes. It is sourced in the West

418 Pacific Ocean (PO) for the pre-monsoon in South China (SC). It changes then to the Indian
419 Ocean (IO) when the monsoon sets up in SC and in the Yangtze-Huaihe River Basin. Finally
420 the main water vapor channel is back to PO for the terminal stage of monsoon in North China.
421 As shown in Fig. 7, the proportion of trajectories shows quite distinct behaviors for their
422 seasonal evolution. The two channels from south and southwest (SCS channel and IO channel)
423 increase firstly, and then decrease with northward shift of the rain-belt. In the contrary, the
424 water vapor channels from the mid-latitude westerly, the West Pacific Ocean and the local
425 moisture recycling show opposite characteristics, with an initial decrease followed by an
426 increase.

427 Not only water vapor channels change profoundly following the seasonal course of the
428 summer monsoon, the spatial distribution of water vapor contribution leading to rainfall also
429 shows strong changes from one period to another. Figure 8 shows difference maps of CDF
430 between two sequential stages. From the pre-monsoon to the onset of monsoon in South
431 China, the moisture contribution from the Indian Ocean and the south part of the South China
432 Sea has an obvious increase. Meanwhile, there are decreases of CDF in the north part of the
433 South China Sea and in the West Pacific Ocean. This situation is largely expected since the
434 onset of the monsoon in India and in South China implies much more water vapor coming
435 from southwest. A large decreasing zone covering the Bay of Bengal, South China and the
436 adjacent South China Sea reflects strong monsoon rainfalls in these areas. It is to be noted that
437 there is a zone of increasing CDF in the northwest area of our target region, certainly related
438 to the special topography in the lee side of the Tibetan Plateau.

439 When the monsoon rain belt reaches the Yangtze-Huaihe River Basin, we observe an

440 opposite situation as shown in Figure 8b. There is a clear dipole with decreasing contribution
441 from the south and increasing contribution from the north. The demarcation line is located
442 around 25° - 30° N. As we mentioned earlier, the northward shift of the research region
443 reflects the seasonal migration of the summer monsoon which induces a systematic northward
444 shift of the water vapor contribution. This phenomenon is also visible in Figure 8c displaying
445 the changes of CDF when the monsoon shifts from the Yangtze-Huaihe River Basin to its
446 terminal stage in North China. The water vapor from north is more important and that from
447 south is less important. It should be noticed that the moisture contribution from the Pacific
448 Ocean is larger, showing a more important role played by the West Pacific Ocean.

449 *3.6 Contribution from different source regions*

450 From what described previously, we can see that our Lagrangian trajectories can be
451 quantitatively used to assess the intensity of moisture transport channels and determine the
452 main pathway in each stage of the monsoonal northward propagation. The
453 Evaporation-minus-Precipitation diagnosis considering the absorption and release of moisture
454 along the trajectories can further determine the source characteristics of water vapor. Finally,
455 the improved areal source–receptor attribution method allows us to quantify the moisture
456 contribution of each source for the precipitation in the target region. With the northward
457 propagation of EASM, significant changes take place in water vapor channels, including
458 moisture sources and water vapor contributions to rainfall. In this section, we study
459 furthermore the properties of CDF which is a new measure with powerful diagnostic abilities.
460 We want to focus on its dynamic evolution in function of the monsoon northward
461 propagation.

462 To be consistent with the above-shown moisture transport channel, we use the same five
463 moisture source regions: (1) East China (EC), (2) West Pacific Ocean (PO), (3) South China
464 Sea (SCS), (4) Indian Ocean (IO) and (5) Eurasia (EA) (Figure 9, background map). We can
465 now calculate the moisture contribution to precipitation for each of these source regions by
466 just integrating the Contribution Density Function (CDF) for each region. It is also relevant to
467 compare such regional contributions to the intensity of each moisture transport channel.

468 As shown in Figure 9, the moisture contribution from East China (EC) to regional
469 precipitations is very high for all the four stages of the summer monsoon. This regional
470 moisture contribution is well above the proportion of trajectories. In fact, all proportions of
471 trajectories are under 10%, but the moisture contributions are mostly over 30%. This result
472 clearly indicates that more than 30% of the moisture for precipitations in East China is
473 originated from local evaporations. The local water recycling plays an important role for
474 precipitation in each stage of the rainy season. Obviously, this kind of local evaporation is
475 closely related to the hydrological conditions and vegetation behaviors of the underlying
476 surface. In fact, it is closely related to early precipitations. Previous studies have also shown
477 that soil moisture is an important factor for monthly and seasonal forecasting (Walker and
478 Rowntree, 1977; Tuttle and Salvucci, 2016). Zuo and Zhang (2007) already pointed out that
479 spring precipitation and soil moisture in East China exert an important effect on summer
480 precipitation. It is remarkable that, after the onset of SCS summer monsoon, soil moisture
481 accumulation in East China due to earlier precipitation enhances local evaporation and then
482 precipitation. This is especially true for the terminal stage of monsoon in North China with a
483 regional moisture contribution up to 55%, reflecting that local water vapor recycling and

484 precipitation in early rainy season play a very important role in the regional monsoon strength.

485 In general, evaporation in East China plays an important role for the rainfall in EASM.

486 PO is also an important moisture source region contributing to precipitations in East
487 China. The regional contributions from PO are all larger than 10%. It is the highest (25%) for
488 the pre-monsoon in South China. The second largest contribution ($> 20\%$) is found for the
489 terminal stage of monsoon in North China.

490 The moisture contribution from SCS shows a big difference between the rainfall stage in
491 South China and other two stages in the Yangtze River valley and in North China during the
492 monsoon seasonal course. In South China, no matter before or after the onset of the SCS
493 summer monsoon, the moisture contributions from SCS are both larger than 20%, which
494 constitutes an important moisture source. But when the monsoon goes to the Yangtze River
495 basin and further to North China, this moisture contribution diminishes rapidly, until 5.1% in
496 the terminal stage of monsoon in North China.

497 For the Indian Ocean, if we examine the proportion of trajectories, we can see that the IO
498 water vapor transport channel is an important one for the whole monsoon course. This is in
499 agreement with our general believing. In particular, for rainfall period in South China and the
500 Meiyu in the Yangtze-Huaihe River valley, we can account about half of the trajectories.
501 However, if we examine the moisture contribution to the actual precipitations, a different
502 image appears. The contribution from IO is actually quite small. For the four sub-periods of
503 the monsoon seasonal course, it is only 7.2%, 18.0%, 9.5% and 2.4% respectively. The main
504 reason is that the pathway of the IO channel goes through the Indian Peninsula and the
505 Indochina Peninsula which are important moisture sink regions to the rainfall in target region.

506 Moisture is thus lost along the trajectories, which induces a low contribution from the Indian
507 Ocean for precipitations in East China.

508 Eurasian land is another region for which the “true” moisture contribution rates are all
509 lower than the proportions of accounted trajectories. As is shown in Figure 8, the trajectories
510 rates are all larger than 10% except when the monsoon is in South China, but enhanced when
511 the monsoon reached North China. For the latter case, the proportion of trajectories from
512 mid-latitude westerlies reaches its peak value of 30.2%. In terms of moisture contribution to
513 precipitations, all contributions from EA are smaller than 3.5%. The main reason is that the
514 specific humidity of air parcels from EA is small (Figure 3a). Therefore, the impact on
515 precipitation of EASM is also small.

516 In summary, with the northward migration of EASM during its seasonal course, the
517 moisture transport channels and moisture source regions show continuous and significant
518 changes. The main water vapor channel is firstly from PO for the pre-monsoon in South China.
519 It is then from IO for the starting monsoon in South China and Meiyu in the Yangtze-Huaihe
520 River Basin. Finally, it returns to PO in the terminal stage of monsoon in North China. And
521 based on the evaporation-minus-precipitation diagnosis and the areal source–receptor
522 attribution method, the main moisture source during pre-monsoon in South China is PO and
523 EC and the contribution rates are 34.3% and 27.7%, respectively. After the onset of South
524 China Sea summer monsoon, EC and SCS are the most important moisture sources in this
525 period, their contribution rates being both larger than 23%. During Meiyu in the
526 Yangtze-Huaihe River Basin and terminal stage of monsoon in North China, the moisture of
527 this period mainly comes from EC, the contribution rate reaching to 40.9% and 55.3%. And

528 our results on water vapor channels are consistent with published literature, such as, Lin et al.
529 (2014), Chen and Luo (2018), Zuo et al. (2009) and Jiang et al. (2017). But it is clear that
530 main moisture actually contributing to EASM precipitations doesn't necessarily come from
531 the strongest water vapor pathways. For example, trajectory proportions for IO are larger than
532 25% especially in the stage of monsoon in South China (53%), but the moisture contributions
533 are generally smaller than 10%. This is mainly due to the fact that a large amount of water
534 vapor is actually lost in the transport pathways through important moisture sink areas such as
535 the Indian Peninsula and Indochina Peninsula. On the contrary, local water vapor recycling
536 inside EC plays an important role for the regional precipitations, with contributions mostly
537 over 30%, although the trajectory proportions for EC are all under 10%. The contribution rate
538 can even exceed 55% for the terminal stage of monsoon in North China.

539 Our results seem qualitatively in agreement with Sun and Wang (2015) who emphasized
540 the role of land evaporation in precipitations falling in the Yangtze-Huaihe river basin and in
541 North China. Similar conclusions are also reported in Drumond et al. (2011a) for
542 precipitations in different regions of China. One can also mention the work of Drumond et al.
543 (2011b) investigating precipitations in the Mediterranean area. It was convincingly shown that
544 local sources provide moisture for the Eastern Mediterranean and Western North Africa.
545 However, if we quantitatively compare our results to those reported in Sun and Wang (2015),
546 our results seem to give larger weights on evaporation along the trajectories. This is certainly
547 due to the fact that we incorporated a rule of boundary layer water vapor source in our
548 calculations.

549 **4. Conclusion and Discussion**

550 In this paper, we used the HYSPLIT platform to calculate Lagrangian trajectories of air
551 parcels that reach East China during the northward propagation of the East Asian Summer
552 Monsoon. Our study, with NCEP/NCAR reanalysis data as driving conditions, covers a long
553 period from 1961 to 2010 to ensure the significance of results. In Supplementary materials S1
554 and S2, we explored the sensitivity of our results to two other datasets and to the use of a
555 second Lagrangian trajectory model, FLEXPART. The difference is estimated below 5 %
556 among different datasets and below 10% between the two Lagrangian models. To fully
557 explore the Lagrangian trajectories, we calculate the mean moisture trajectories to get the
558 main moisture channel. Since water vapor is not a conservative tracer following the
559 trajectories, we also elaborated the Evaporation-minus-Precipitation diagnosis and the areal
560 source–receptor attribution of water vapor contribution. We can thus obtain a robust picture
561 for the water vapor transport, including paths, moisture sources and moisture contribution
562 from different regions to precipitations in EASM. Main findings are summarized as follows:

563 Four main water vapor channels and their contribution have significant changes while
564 the rain-belt in East China propagates from south to north. These water vapor channels are
565 from PO (West Pacific Ocean), IO (Indian Ocean), SCS (South China Sea) and the
566 mid-latitude westerlies, respectively. And local transportation also plays an important role in
567 each rainy season. In the pre-monsoon stage with rainfall in South China, the most important
568 moisture transport channels are from PO and IO, the proportion of the trajectories being 33.3%
569 and 24.6% respectively, consistent with our general expectation. But main moisture sources
570 are from East China and PO, with moisture contribution rates at 34.3% and 25.7%
571 respectively, highlighting the role of local evaporation for precipitations. After the onset of the

572 SCS summer monsoon, but when the monsoon rain-belt is still in South China, the IO channel
573 strengthens, with the trajectories proportion increasing to 53.9%. This reflects the onset of the
574 Indian monsoon which enters into the South China Sea. Despite the Indian monsoon
575 strengthening, EC and SCS are, however, the most important moisture sources in this period,
576 their contribution rates to precipitations being both larger than 23%. For Meiyu in the
577 Yangtze-Huaihe River basin, the IO channel is still the most important water vapor channel,
578 which includes 42.1% of all the trajectories. But the contribution rate from East China to
579 precipitations is the highest (40.9%). East China becomes the most important moisture source
580 region. When the monsoon is in its terminal stage in North China, the most important
581 moisture channels are the mid-latitude westerlies channel and PO channel, their trajectories
582 proportions being 30.3% and 26.8%, and the local water vapor recycling in East China has the
583 greatest contribution to rainfall in North China during this stage, its contribution rate
584 exceeding 55%.

585 It is worthy to emphasize that main moisture for the EASM precipitation doesn't
586 necessarily come from the strongest water vapor pathways. For example, trajectory
587 proportions for IO are all larger than 25% (exceeding 53% in the stage of monsoon in South
588 China), but moisture contributions to rainfall are always smaller than 10%. This
589 counter-intuitive result is in fact quite logic, since a large amount of water vapor is lost in the
590 transport pathways crossing important moisture sink areas such as the Indian Peninsula and
591 Indochina Peninsula. On the contrary, local water vapor recycling inside East China plays an
592 important role for the regional precipitations, with contributions mostly over 30%, especially
593 in the terminal stage of monsoon in North China the contribution rate can even exceed 55%,

594 although the trajectory proportions for East China are all under 10%.

595 Finally, we can see that our analysis based on Lagrangian trajectories provides very
596 useful information on water vapor transport, its source-sink regions and its contributions to
597 rainfall. Results are generally in agreement with Eulerian diagnosis of water vapor transport.
598 The focus of this study was put on precipitations in East China during the northward
599 migration course of the regional summer monsoon. We examined only climatological fields. It
600 is necessary to extend the current study to investigating interannual and interdecadal
601 variations of moisture transport.

602 **Acknowledgments.** Constructive comments from three anonymous reviewers were very helpful to improve
603 an earlier version of this paper. We thank Dr. Sun Bo for his great help on technology guidance. We
604 acknowledge the National Climate Center (NCC) of China (<http://ncc.cma.gov.cn>) for the observations and
605 NOAA, the Air Resources Laboratory (ARL), from ARL's archives
606 (http://ready.arl.noaa.gov/gbl_reanalysis.php) for NCEP/NCAR reanalysis data. This work is supported by
607 the National Natural Science Foundation of China (41675081), and the National Key Research and
608 Development Program of China (grant 2016YFA0600402). Zhengyu Liu is partly supported by the U.S.
609 NSF and DOE. Laurent Li is partly supported by the French ANR project China-Trend-Stream.

610 **References**

611 Bertò, A., A. Buzzi, and D. Zardi, 2004: Back-tracking water vapour contributing to a
612 precipitation event over Trentino: a case study. *Meteorologische Zeitschrift*, **13**, 189-200,
613 <https://doi.org/10.1127/0941-2948/2004/0013-0189>.

614 Brubaker, K. L., P. A. Dirmeyer, A. Sudrajat, and B. S. Levy, 2001: A 36-yr climatological
615 description of the evaporative of warm-season precipitation in the Mississippi river basin.

- 616 *J. Hydrometeor.*, **2**, 537-557,
617 [https://doi.org/10.1175/1525-7541\(2001\)002<0537:AYCDOT>2.0.CO;2](https://doi.org/10.1175/1525-7541(2001)002<0537:AYCDOT>2.0.CO;2).
- 618 Bottyán, E., G. Czuppon, K. Kármán, T. Weidinger, and L. Haszpra, 2014: Moisture source
619 diagnostic for Hungary based on trajectory analysis and stable isotopic composition of
620 precipitation. *EGU General Assembly Conference Abstracts*, Vol. **16**.
- 621 Chang, C. P., 2004: *East Asian Monsoon*. World Scientific. 564pp.
- 622 Chen, B., X. Xu, and T. Zhao, 2013: Main moisture sources affecting lower Yangtze River
623 Basin in boreal summers during 2004-2009. *Int. J. Climatol.*, **33**,1035-1046,
624 <https://doi.org/10.1002/joc.3495>.
- 625 Chen, L., and Coauthors, 1991: East Asia Monsoon. *China Meteorological Press*, 362. (in
626 Chinese)
- 627 Chen, Y., and Y. Luo, 2018: Analysis of Paths and Sources of Moisture for the South China
628 Rainfall during the Pre-summer Rainy Season of 1979–2014. *Journal of Meteorological
629 Research*, **32**, 744-757, <https://doi.org/10.1007/s13351-018-8069-7>.
- 630 Chi, Y., J. He, and W. Wu, 2005: Features analysis of the different precipitation periods in the
631 pre-flood season in South China. *Journal of Nanjing Institute of Meteorology*, **28**,
632 163-171, <https://doi.org/10.3969/j.issn.1674-7097.2005.02.003>.
- 633 Chu, Q., Q. Wang, and G. Feng, 2017: Determination of the major moisture sources of
634 cumulative effect of torrential rain events during the preflood season over South China
635 using a Lagrangian particle model, *J. Geophys. Res. Atmos.*, **122**, 8369–8382,
636 <https://doi.org/10.1002/2016JD026426>.
- 637 Connolley, W. M., J. C. King, 1993: Atmospheric water - vapour transport to Antarctica

- 638 inferred from radiosonde data. *Quarterly Journal of the Royal Meteorological Society*,
639 **119**(510): 325-342, <https://doi.org/10.1002/qj.49711951006>.
- 640 Diem, J. E., and D. P. Brown, 2006: Tropospheric moisture and monsoonal rainfall over the
641 southwestern United States. *J. Geophys. Res. Atmos.*, **111**(D16),
642 <https://doi.org/10.1029/2005JD006836>.
- 643 Dirmeyer, P. A., K. L. Brubaker, and T. DelSole, 2009: Import and export of atmospheric
644 water vapor between nations, *J. Hydrol.*, **365**, 11–22,
645 <https://doi.org/10.1016/j.jhydrol.2008.11.016>.
- 646 Ding, Y., Z. Wang, and Y. Sun, 2008: Inter-decadal variation of the summer precipitation in
647 East China and its association with decreasing Asian summer monsoon. Part I: Observed
648 evidences. *Int. J. Climatol.*, **28**: 1139–1161, <https://doi.org/10.1002/joc.1615>.
- 649 Dominguez, F., P. Kumar, X. Liang, and M. Ting, 2006: Impact of atmospheric moisture
650 storage on precipitation recycling. *J. Climate*, **19**, 1513–1530,
651 <https://doi.org/10.1175/JCLI3691.1>.
- 652 Dominguez, F., G. Miguez-Macho, and H. Hu, 2016: WRF with water vapor tracers: A study
653 of moisture sources for the North American monsoon. *Journal of Hydrometeorology*,
654 **17**(7), 1915-1927. <https://doi.org/10.1175/jhm-d-15-0221.1>
- 655 Draxler, R. R., and G. Hess, 1998: An overview of the HYSPLIT_4 modelling system for
656 trajectories, *Aust. Meteorol. Mag.*, **47**, 295–308.
- 657 Draxler, R. R., and G. D. Rolph, 2011: HYSPLIT (HYbrid Single-Particle Lagrangian
658 Integrated Trajectory) Model; National Oceanic and Atmospheric Administration, Air
659 Resources Laboratory READY Web site. <https://ready.arl.noaa.gov/HYSPLIT.php>.

- 660 Drumond, A., R. Nieto, and L. Gimeno, 2011a: Sources of moisture for China and their
661 variations during drier and wetter conditions in 2000–2004: a Lagrangian approach.
662 *Climate Research*, **50**: 215-225. <https://doi.org/10.3354/cr01043>.
- 663 Drumond, A., R. Nieto, E. Hernandez, and L. Gimeno, 2011b: A Lagrangian analysis of the
664 variation in moisture sources related to drier and wetter conditions in regions around the
665 Mediterranean Basin, *Nat. Hazards Earth Syst. Sci.*, **11**: 2307-2320,
666 <https://doi.org/10.5194/nhess-11-2307-2011>,.
- 667 Eiras-Barca, J., F. Dominguez, H. Hu, D. Garaboa-Paz, and G. Miguez-Macho, 2017:
668 Evaluation of the moisture sources in two extreme landfalling atmospheric river events
669 using an Eulerian WRF tracers tool. *Earth System Dynamics*, **8**(4), 1247,
670 <https://doi.org/10.5194/esd-8-1247-2017>.
- 671 Gimeno, L., A. Drumond, R. Nieto, R. M. Trigo, and A. Stohl, 2010: On the origin of
672 continental precipitation. *Geophysical Research Letters*, **37**(13),
673 <https://doi.org/10.1029/2010GL043712>.
- 674 Huang, R., Z. Zhang, G. Huang, and B. Ren, 1998: Characteristics of the water vapor
675 transport in East Asian monsoon region and its difference from that in South Asian
676 monsoon region in summer. *Chinese J. Atmos. Sci.*, **22**, 469-479,
677 <https://doi.org/10.3878/j.issn.1006-9895.1998.04.08>.
- 678 Huang, W., X. He, Z. Yang, T. Qiu, J. S. Wright, B. Wang, and D. Lin, 2018: Moisture sources
679 for wintertime extreme precipitation events over South China during 1979–2013. *J.*
680 *Geophys. Res. Atmos.*, **123**, 6690-6712, <https://doi.org/10.1029/2018JD028485>.
- 681 Insua-Costa, D., and G. Miguez-Macho, 2018: A new moisture tagging capability in the

- 682 Weather Research and Forecasting model: formulation, validation and application to the
683 2014 Great Lake-effect snowstorm. *Earth System Dynamics*, **9**(1), 167-185,
684 <https://doi.org/10.5194/esd-9-167-2018>.
- 685 Jiang, X., Y. Li, and X. Wang, 2009: Water vapor transport over China and its relationship
686 with drought and flood in Yangtze River Basin. *Journal of Geographical Sciences*, **19**,
687 153-163, <https://doi.org/10.1007/s11442-009-0153-6>.
- 688 Jiang, Z., J. He, J. Li, J. Yang, and J. Wang, 2006: Northerly Advancement Characteristics of
689 the East Asian Summer Monsoon with Its Interdecadal Variations. *Acta Geographica
690 Sinica*. **61**, 675-686, <https://doi.org/10.3321/j.issn:0375-5444.2006.07.001>. (in Chinese)
- 691 Jiang, Z., W. Ren, Z. Liu, and H. Yang, 2013: Analysis of water vapor transport characteristics
692 during the Meiyu over the Yangtze-Huaihe River valley using the Lagrangian method.
693 *Acta Meteorologica Sinica*, **271**, 295-304, <https://doi.org/10.11676/qxxb2013.017>. (in
694 Chinese)
- 695 Jiang, Z., S. Jiang, Y. Shi, Z. Liu, W. Li, and L. Li, 2017: Impact of moisture source variation
696 on decadal-scale changes of precipitation in North China from 1951 to 2010, *J. Geophys.
697 Res. Atmos.*, **122**, 600–613, <https://doi.org/10.1002/2016JD025795>.
- 698 Lin, A., D. Gu, B. Zheng, C. Li, and J. Zhang, 2014: Anomalous transport of water vapor for
699 sustained torrential rain and its variation. *J. Trop. Meteor.*, **30**, 1001-1010,
700 <https://doi.org/10.3969/j.issn.1004-4965.2014.06.001>. (in Chinese)
- 701 Martius, O., and Coauthors, 2013: The role of upper-level dynamics and surface processes for
702 Pakistan flood of July 2010. *Quart. J. Roy. Meteor. Soc.*, **139**, 1780–1797,
703 <https://doi.org/10.1002/qj.2082>.

- 704 Perry, L. B., C. E. Konrad, and T. W. Schmidlin, 2007: Antecedent upstream air trajectories
705 associated with northwest flow snowfall in the southern Appalachians. *Wea. Forecasting*,
706 **22**, 334-352, <https://doi.org/10.1175/WAF978.1>.
- 707 Shen, R., and Coauthors, 1982: Low latitude circulation variation in the high and low level of
708 convective layer and precipitation during the pre-rainy season in South China.
709 *Proceedings of the National Conference on tropical summer monsoon*. Yunnan People's
710 Publishing House, 10-21 (in Chinese)
- 711 Shi, Y., X. J. Gao, Y. G. Wang, F. Giorgi, 2009: Simulation and projection of monsoon rainfall
712 and rain patterns over eastern China under global warming by RegCM3. *Atmos. Oceanic*
713 *Sci. Lett.*, **2**, 1–6, <https://doi.org/10.1080/16742834.2009.11446816>.
- 714 Sodemann, H., C. Schwierz, and H. Wernli, 2008: Interannual variability of Greenland winter
715 precipitation sources: Lagrangian moisture diagnostic and North Atlantic Oscillation
716 influence, *J. Geophys. Res.*, **113**, D03107, <https://doi.org/10.1029/2007JD008503>.
- 717 Sodemann, H., and A. Stohl, 2009: Asymmetries in the moisture origin of Antarctic
718 precipitation. *Geophysical research letters*, **36**, 273-289,
719 <https://doi.org/10.1029/2009GL040242>.
- 720 Sodemann, H., and E. Zubler, 2010: Seasonal and inter-annual variability of the moisture
721 sources for Alpine precipitation during 1995–2002. *Int. J. Climatol.*, **30**, 947-961.
722 <https://doi.org/10.1002/joc.1932>.
- 723 Stohl, A., and P. James, 2004: A Lagrangian analysis of the atmospheric branch of the global
724 water cycle. Part I: Method description, validation, and demonstration for the August
725 2002 flooding in central Europe. *J. Hydrometeorol.*, **5**, 656-678,

- 726 [https://doi.org/10.1175/1525-7541\(2004\)005<0656:alaota>2.0.co;2](https://doi.org/10.1175/1525-7541(2004)005<0656:alaota>2.0.co;2).
- 727 Stohl, A., and P. James, 2005: A Lagrangian analysis of the atmospheric branch of the global
728 water cycle. Part II: Moisture transports between earth's ocean basins and river
729 catciunents. *J. Hydrometeorol*, **6**, 961-984, <https://doi.org/10.1175/JHM470.1>.
- 730 Sun, B., and H. Wang, 2014: Moisture sources of semi-arid grassland in China using the
731 Lagrangian Particle Model FLEXPART. *J. Clim.*, **27**, 2457-2474,
732 <https://doi.org/10.1175/JCLI-D-13-00517.1>.
- 733 Sun, B., and H. Wang, 2015: Analysis of the major atmospheric moisture sources affecting
734 three sub-regions of east China. *Int. J. Climatol.* **35**, 2243-2257,
735 <https://doi.org/10.1002/joc.4145>.
- 736 Sun, B., Y. L. Zhu, H. J. Wang, 2011: The recent interdecadal and interannual variation of
737 water vapor transport over eastern China. *Adv. Atmos. Sci.*, **28**, 1039–1048,
738 <https://doi.org/10.1007/s00376-010-0093-1>.
- 739 Tuttle, S., and G. Salvucci, 2016: Empirical evidence of contrasting soil moisture–
740 precipitation feedbacks across the United States. *Science*, **352**, 825-828,
741 <https://doi.org/10.1126/science.aaa7185>.
- 742 Walker, J., and P. R. Rowntree, 1977: The effect of soil moisture on circulation and rainfall in
743 a tropical model. *Quarterly Journal of the Royal Meteorological Society*, **103**, 29-46,
744 <https://doi.org/10.1002/qj.49710343503>.
- 745 Wang, B., H. Lin, Y. Zhang, and M. M. Lu, 2004: Definition of South China Sea monsoon
746 onset and commencement of the East Asia summer monsoon. *J. Clim.*, **17**, 699-710,
747 [https://doi.org/10.1175/1520-0442\(2004\)017<0699%3ADOSCSM>2.0.CO%3B2](https://doi.org/10.1175/1520-0442(2004)017<0699%3ADOSCSM>2.0.CO%3B2)

- 748 Wang B. 2006: *The Asian monsoon*. Springer Science & Business Media. 787pp.
- 749 Wei, J., P. A. Dirmeyer, M. G. Bosilovich, and R. Wu, 2012: Water vapor sources for Yangtze
750 River Valley rainfall: Climatology, variability, and implications for rainfall forecasting, *J.*
751 *Geophys. Res.*, **117**, D05126, <https://doi.org/10.1029/2011JD016902>.
- 752 Xie, K., and X. Ren. 2008: Climatological characteristics of atmospheric water vapor
753 transport and its relation with rainfall over north China in summer. *Scientia*
754 *Meteorologica Sinica*, **28**, 5508-5514, <https://doi.org/10.3724/SP.J.1047.2008.00014>. (in
755 Chinese)
- 756 Xu, X., L. Chen, X. Wang, Q. Miao, and S. Tao, 2004: Moisture transport source/ sink
757 structure of the meiyu rain belt along the yangtze river valley. *Chinese Science Bulletin*,
758 **49**, 181-188, <https://doi.org/10.1360/03wd0047>.
- 759 Xu, X., X. Shi, Y. Wang, S. Peng, and X. Shi, 2008: Data analysis and numerical simulation
760 of moisture source and transport associated with summer precipitation in the yangtze
761 river valley over china. *Meteorology and Atmospheric Physics*, **100**, 217-231,
762 <https://doi.org/10.1007/s00703-008-0305-8>.
- 763 Zhou, T., and R. Yu, 2005: Atmospheric water vapor transport associated with typical
764 anomalous summer rainfall patterns in China. *J. Geophys. Res.* **110**, D08104,
765 <https://doi.org/10.1029/2004JD005413>.
- 766 Zhu, Y., H. J. Wang, W. Zhou, J. G. Ma, 2011: Recent changes in the summer precipitation
767 pattern in East China and the background circulation. *Clim. Dyn.*, **36**, 1463–1473,
768 <https://doi.org/10.1007/s00382-010-0852-9>.
- 769 Zuo, J., H. Ren, W. Li, P. Zhang, and M. Yang, 2009: Intraseasonal Characteristics of Water

770 Vapor Transport Associated with Low-Frequency Rainfall Regimes over Southern China
771 in Summer. *Chinese Journal of Geophysics*, **52**, 922-935,
772 <https://doi.org/10.1002/cjg2.1417>.

773 Zuo, Z., and R. Zhang, 2007: The spring soil moisture and the summer rainfall in eastern
774 China. *Chinese Science Bulletin*, **52**, 3310-3312,
775 <https://doi.org/10.1007/s11434-007-0442-3>.

776

777 **Figure Captions**

778 **Figure 1.** Location of the three rectangular target domains in East China, it is North China
779 region (35° – 43° N, 110° – 120° E), Yangtze-Huaihe River Basin region (28° – 34° N, 110° – 123° E)
780 and South China region (20° – 26° N, 106° – 120° E) from north to south. Dots indicate locations
781 of the observational stations in these three regions. Superimposed are three major rivers in
782 East China: the Pearl river in the south, the Yangtze river in the middle and the Yellow river
783 in the north.

784
785 **Figure 2.** Beginning and end dates of pre-monsoon in South China (dark blue, PSC),
786 monsoon in South China (light blue, MSC), monsoon (or Meiyu) in the Yangtze-Huai River
787 Basin (red, YHR) and monsoon in North China (green, NC) from 1961 to 2010.

788
789 **Figure 3.** Characteristics of moisture transport, moisture source and sink for the pre-monsoon
790 in South China (SC). (a) Moisture transport channels based on average trajectories. Five
791 transport channels are identified, from East China (EC), the West Pacific Ocean (PO), the
792 South China Sea (SCS), the Indian Ocean (IO) and the Eurasian westerly (EA), respectively.
793 Colors on the pathways indicate the average specific humidity of air parcels along the
794 trajectories (units: g/kg). The thickness of the pathways represents the percentage of
795 trajectories, also marked in numbers. Grey shadings are the number of particles weighted by
796 specific humidity (units: g/kg) arriving in the target region (the rectangular zone) for day 10.
797 (b) The climatology of vertically integrated atmospheric water vapor transport (vectors, unit:
798 $\text{kg m}^{-1} \text{s}^{-1}$) under the Eulerian view, only vector more than $50 \text{kg m}^{-1} \text{s}^{-1}$ are shown, and

799 shadings represent amount of the water vapor transport. (c) Mean $E - P$ (unit: mm day^{-1}) of air
800 parcels in 1–10 days before reaching target region (the rectangular zone). (d) Water vapor
801 Contribution Density Function (CDF) showing the contribution of water vapor source regions
802 (unit: 10^{-5} , areas of 1 by 1 in latitude/longitude) to the precipitation in the target region of
803 South China.

804

805 **Figure 4.** Same as in Figure 3 but for the starting phase of the monsoon in South China.

806

807 **Figure 5.** Same as in Figure 3, but for the phase of the monsoon (Meiyu) in the
808 Yangtze-Huaihe River Basin

809

810 **Figure 6.** Same as in Figure 3, but for the terminal stage of the monsoon in North China

811

812 **Figure 7.** (a) Schematic showing the five main moisture channels from East China (EC) itself,
813 the West Pacific Ocean (WPO), the South China Sea (SCS), the Indian Ocean (IO) and the
814 Eurasian westerly (EA) that affect precipitation in East China. (b) Proportions of trajectories
815 from different channels in function of the four major stages of the East Asian summer
816 monsoon: pre-monsoon in South China (PSC), monsoon in South China (MSC) Meiyu in the
817 Yangtze-Huaihe River basin (YHR) and terminal stage of monsoon in North China (NC).

818

819 **Figure 8.** Sequential changes of the water vapor Contribution Density Function (CDF) among
820 the four stages of the summer monsoon course: stage 2 – stage 1 (left), stage 3 – stage 2


821 (middle) and stage 4 – stage 3 (right) (unit: 10^{-5} , areas of 1 by 1 in latitude/longitude).

822


823 **Figure 9.** Bar charts showing the proportion of trajectories (red bars in %) and moisture
824 contributions (blue bars in %) leading to precipitations in different stages of the summer
825 monsoon (PSC: pre-monsoon in South China; MSC: monsoon in South China; YHR:
826 monsoon or Meiyu in the Yangtze-Huaihe River basin; NC: terminal monsoon in North
827 China). Background map shows the division of geographic sectors (black boxes: East China,
828 South China Sea, Indian Ocean, West Pacific Ocean and Eurasia) used to account the
829 trajectories and moisture contributions.

830

831



832
833 Figure 1. Location of the three rectangular target domains in East China, it is North China
834 region (35° – 43° N, 110° – 120° E), Yangtze-Huaihe River Basin region (28° – 34° N, 110° – 123° E)
835 and South China region (20° – 26° N, 106° – 120° E) from north to south. Dots indicate locations
836 of the observational stations in these three regions. Superimposed are three major rivers in
837 East China: the Pearl river in the south, the Yangtze river in the middle and the Yellow river
838 in the north.
839




840

841 Figure 2. Beginning and end dates of pre-monsoon in South China (dark blue, PSC), monsoon

842 in South China (light blue, MSC), monsoon (or Meiyu) in the Yangtze-Huai River Basin (red,

843 YHR) and monsoon in North China (green, NC) from 1961 to 2010.

844



845

846 Figure 3. Characteristics of moisture transport, moisture source and sink for the pre-monsoon

847 in South China (SC). (a) Moisture transport channels based on average trajectories. Five

848 transport channels are identified, from East China (EC), the West Pacific Ocean (PO), the

849 South China Sea (SCS), the Indian Ocean (IO) and the Eurasian westerly (EA), respectively.

850 Colors on the pathways indicate the average specific humidity of air parcels along the

851 trajectories (units: g/kg). The thickness of the pathways represents the percentage of

852 trajectories, also marked in numbers. Grey shadings are the number of particles weighted by


853 specific humidity (units: g/kg) arriving in the target region (the rectangular zone) for day 10.

854 (b) The climatology of vertically integrated atmospheric water vapor transport (vectors, unit:

855 $\text{kg m}^{-1} \text{s}^{-1}$) under the Eulerian view, only vector more than $50 \text{ kg m}^{-1} \text{s}^{-1}$ are shown, and856 shadings represent amount of the water vapor transport. (c) Mean $E - P$ (unit: mm day^{-1}) of air

857 parcels in 1–10 days before reaching target region (the rectangular zone). (d) Water vapor
858 Contribution Density Function (CDF) showing the contribution of water vapor source regions
859 (unit: 10^{-5} , areas of 1 by 1 in latitude/longitude) to the precipitation in the target region of
860 South China.


861



862

863 Figure 4. Same as in Figure 3 but for the starting phase of the monsoon in South China.

864




865

866 Figure 5. Same as in Figure 3, but for the phase of the monsoon (Meiyu) in the

867 Yangtze-Huaihe River Basin


868



869

870 Figure 6. Same as in Figure 3, but for the terminal stage of the monsoon in North China

871



872

873 Figure 7. (a) Schematic showing the five main moisture channels from East China (EC) itself,

874 the West Pacific Ocean (WPO), the South China Sea (SCS), the Indian Ocean (IO) and the

875 Eurasian westerly (EA) that affect precipitation in East China. (b) Proportions of trajectories


876 from different channels in function of the four major stages of the East Asian summer

877 monsoon: pre-monsoon in South China (PSC), monsoon in South China (MSC) Meiuyu in the

878 Yangtze-Huaihe River basin (YHR) and terminal stage of monsoon in North China (NC).

879

880




881

882 Figure 8. Sequential changes of the water vapor Contribution Density Function (CDF) among
883 the four stages of the summer monsoon course: stage 2 – stage 1 (left), stage 3 – stage 2
884 (middle) and stage 4 – stage 3 (right) (unit: 10^{-5} , areas of 1 by 1 in latitude/longitude).

885

886



887

888 Figure 9. Bar charts showing the proportion of trajectories (red bars in %) and moisture
 889 contributions (blue bars in %) leading to precipitations in different stages of the summer
 890 monsoon (PSC: pre-monsoon in South China; MSC: monsoon in South China; YHR:
 891 monsoon or Meiyu in the Yangtze-Huaihe River basin; NC: terminal monsoon in North
 892 China). Background map shows the division of geographic sectors (black boxes: East China,

893 South China Sea, Indian Ocean, West Pacific Ocean and Eurasia) used to account the
894 trajectories and moisture contributions.
895