



**HAL**  
open science

# The Emerging Role of the Prokineticins and Homeobox Genes in the Vascularization of the Placenta: Physiological and Pathological Aspects

Nadia Alfaidy, Sophie Brouillet, Gayathri Rajaraman, Bill Kalionis, Pascale Hoffmann, Tiphaine Barjat, Mohamed Benharouga, Padma Murthi

## ► To cite this version:

Nadia Alfaidy, Sophie Brouillet, Gayathri Rajaraman, Bill Kalionis, Pascale Hoffmann, et al.. The Emerging Role of the Prokineticins and Homeobox Genes in the Vascularization of the Placenta: Physiological and Pathological Aspects. *Frontiers in Physiology*, 2020, 11, pp.591850. 10.3389/fphys.2020.591850 . hal-03058339

**HAL Id: hal-03058339**

**<https://hal.science/hal-03058339>**

Submitted on 18 Jun 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# The Emerging Role of the Prokineticins and Homeobox Genes in the Vascularization of the Placenta: Physiological and Pathological Aspects

Nadia Alfaidy<sup>1,2,3\*</sup>, Sophie Brouillet<sup>4</sup>, Gayathri Rajaraman<sup>5</sup>, Bill Kalionis<sup>6</sup>, Pascale Hoffmann<sup>1,2,3</sup>, Tiphaine Barjat<sup>7</sup>, Mohamed Benharouga<sup>8</sup> and Padma Murthi<sup>6,9</sup>

<sup>1</sup> Unité 1036, Institut National de la Santé et de la Recherche Médicale, Grenoble, France, <sup>2</sup> Department of Biology, University of Grenoble Alpes, Grenoble, France, <sup>3</sup> Commissariat à l'Energie Atomique et aux Energies Alternatives (CEA), Biosciences and Biotechnology Institute of Grenoble, Grenoble, France, <sup>4</sup> INSERM U1203, Department of Reproductive Biology, University of Montpellier, Montpellier, France, <sup>5</sup> Faculty of Health and Biomedicine, First Year College, Victoria University, St. Albans, VIC, Australia, <sup>6</sup> Department of Maternal-Fetal Medicine, Obstetrics and Gynaecology, Pregnancy Research Centre, Royal Women's Hospital, The University of Melbourne, Parkville, VIC, Australia, <sup>7</sup> Unité 1059, Saint-Etienne Hospital, Institut National de la Santé et de la Recherche Médicale, Saint-Étienne, France, <sup>8</sup> Unité Mixte de Recherche 5249, Laboratoire de Chimie et Biologie des Métaux, Centre National de la Recherche Scientifique (CNRS), Grenoble, France, <sup>9</sup> Department of Pharmacology, The Ritchie Centre, Monash Biomedicine Discovery Institute, Hudson Institute of Medical Research, Monash University, Clayton, VIC, Australia

## OPEN ACCESS

### Edited by:

Luca Munaron,  
University of Turin, Italy

### Reviewed by:

Martina Rudnicki,  
York University, Canada  
Martha Sosa-Macías,  
National Polytechnic Institute (IPN),  
Mexico

### \*Correspondence:

Nadia Alfaidy  
nadia.alfaidy-benharouga@cea.fr

### Specialty section:

This article was submitted to  
Vascular Physiology,  
a section of the journal  
Frontiers in Physiology

**Received:** 05 August 2020

**Accepted:** 13 October 2020

**Published:** 12 November 2020

### Citation:

Alfaidy N, Brouillet S, Rajaraman G, Kalionis B, Hoffmann P, Barjat T, Benharouga M and Murthi P (2020) The Emerging Role of the Prokineticins and Homeobox Genes in the Vascularization of the Placenta: Physiological and Pathological Aspects. *Front. Physiol.* 11:591850. doi: 10.3389/fphys.2020.591850

Vasculogenesis and angiogenesis are key processes of placental development, which occur throughout pregnancy. Placental vasculogenesis occurs during the first trimester of pregnancy culminating in the formation of hemangioblasts from intra-villous stem cells. Placental angiogenesis occurs subsequently, forming new blood vessels from existing ones. Angiogenesis also takes place at the fetomaternal interface, allowing essential spiral arteriole remodeling to establish the fetomaternal circulation. Vasculogenesis and angiogenesis in animal models and in humans have been studied in a wide variety of *in vitro*, physiological and pathological conditions, with a focus on the pro- and anti-angiogenic factors that control these processes. Recent studies revealed roles for new families of proteins, including direct participants such as the prokineticin family, and regulators of these processes such as the homeobox genes. This review summarizes recent advances in understanding the molecular mechanisms of actions of these families of proteins. Over the past decade, evidence suggests increased production of placental anti-angiogenic factors, as well as angiogenic factors are associated with fetal growth restriction (FGR) and preeclampsia (PE): the most threatening pathologies of human pregnancy with systemic vascular dysfunction. This review also reports novel clinical strategies targeting members of these family of proteins to treat PE and its consequent effects on the maternal vascular system.

**Keywords:** vessel development, endothelial cells, angiogenesis, prokineticins, homeobox genes, vascularization, pregnancy, EG-VEGF

## INTRODUCTION

The placenta is the key organ for a successful pregnancy. It acts as a semi-permeable barrier to control nutrient and gasses exchanges and regulate waste produced by developing fetus. This hemochorial villous organ develops earlier during pregnancy, following the erosion of nearby maternal capillaries by the lytic activity of the syncytiotrophoblast. Around the 11th-12th day post-conception, the primitive uteroplacental circulation is launched (Gude et al., 2004). The establishment of this circulation is based on two key processes, the first one takes place within the placental villi and is governed by the intra-villi vasculogenesis and angiogenesis; the second one occurs at the fetomaternal interface and is governed by the extravillous trophoblasts (EVT) that remodel maternal spiral arteries to establish the fetomaternal circulation (Velicky et al., 2016).

Villous vascularization is an important process in organogenesis and is essential for the placenta to function efficiently (Zygmunt et al., 2003). At the end of the 3rd week post-conception, intra-villous stem cells differentiate into connective tissue culminating in the formation of hemangioblasts. The differentiation of the hemangioblasts into angioblasts and hematopoietic cells allows the formation of new blood vessels that connect with those of the embryo to form the primitive capillary network (Gude et al., 2004). Placental vasculogenesis is followed by two phases of angiogenesis; branching angiogenesis in immature villi where new vessels form by sprouting, and non-branching angiogenesis in the stem villi, where capillary loops form through elongation. These processes increase the surface area to volume ratio (Charnock-Jones, 2002; Chaddha et al., 2004; Zou et al., 2011) and enhance diffusional exchange between the maternal and fetal circulations (Kingdom et al., 2000; Kaufmann et al., 2004). The second type of angiogenesis, occurring at the fetomaternal interface is mainly ensured by EVTs. By the end of first trimester of pregnancy, the EVTs acquire an invasive phenotype along with markers of endothelial cells. These features allow them to colonize maternal spiral arteries, through the replacement of maternal endothelial cells, culminating in an increase in diameter of these vasculature, which allow more oxygenated maternal blood in the intervillous space (Burton and Jauniaux, 2001; Burton, 2009).

## IMPAIRED PLACENTAL ANGIOGENESIS AND PREGNANCY-ASSOCIATED DISORDERS

The spectrum of vascular defects associated with clinically significant pregnancy disorders including fetal growth restriction (FGR) and preeclampsia (PE), attests to the close relationship between the placental angiogenesis and embryonic development. Villi from placentae where intrauterine embryonic death and blighted ova exhibit aberrant vascular characteristics with significantly decreased vascular density, fibrosis and hydropic degeneration (Meegdes et al., 1988).

Studies using stereological techniques have reported that in placentae from FGR pregnancies, the number, surface area

and volume of terminal villi were significantly reduced in FGR compared with placentae from uncomplicated pregnancies. Villous vessels exhibited fewer branches with slender and uncoiled vessels (Chen et al., 2002; Mayhew, 2003; Mayhew et al., 2003). Reduced capability of branching angiogenesis (non-branching angiogenesis) in FGR was strongly associated with a reduced supply of oxygen and nutrients to the fetus, and a subsequent delay in fetal growth (Kingdom et al., 2000; Salafia et al., 2006). Whether the vascular defects cause human FGR, or whether these changes are a consequence of aberrant biological mechanisms in the placentae from FGR pregnancies (Maulik et al., 2006) is unknown. To address this, it is vital to understand the molecular regulation of angiogenesis in the human placenta.

## MOLECULAR REGULATION OF ANGIOGENESIS

The processes of angiogenesis involve distinct changes in the phenotype of endothelial cells (ECs), which comprise the basic organizational units of vascular structures. The stimuli for these complex processes of placental angiogenesis are temporally coordinated by the microenvironment surrounding the EC surface (Patel et al., 2005). At the molecular level, *in vitro* and *in vivo* studies reported several growth factors and receptors that activate critical signaling pathways (Arderiu et al., 2007). Vascular endothelial growth factor (VEGF), placental growth factor (PlGF), and angiopoietins together with their primary receptors, VEGF receptor-1 (VEGFR-1) and VEGF receptor-2 (VEGFR-2) and PlGF that binds only to VEGFR-1, were identified as key candidates by Patel et al. (2005). VEGF is a potent stimulator of EC proliferation, migration, and production of plasminogen activators that are required for degradation of the basement membrane (Regnault et al., 2002).

In uncomplicated pregnancies, placental expression of key growth factors correlates with their established roles. For example, expression of VEGF and VEGFR-2 is highest during early gestation, which coincides with vasculogenesis and branching angiogenesis, but expression declines with advancing pregnancy (Jackson et al., 1994). Conversely, PlGF and VEGFR-1 expression is highest toward term, coinciding with non-branching angiogenesis (Clark et al., 1996).

Although differential expression of these angiogenic factors have been implicated in the development in PE and FGR (Ahmed and Perkins, 2000; Tsatsaris et al., 2003), new families of proteins have been identified and reported to play key roles in the control of these angiogenic processes. These include two important families of proteins; the prokineticins and the nuclear transcription factors including homeobox genes/homeodomain proteins.

## PROKINETICIN FAMILY IN THE CONTROL OF PLACENTAL ANGIOGENESIS

Recent studies from our group provided evidence for the direct role of a new placental angiogenic called prokineticin 1 (PROK1)

in normal and FGR pregnancies (Hoffmann et al., 2006, 2009; Brouillet et al., 2012b, 2013b; Murthi et al., 2015; Sergent et al., 2016). Due to its similarities of action with VEGF (LeCouter et al., 2001), PROK1 is also called as endocrine gland derived-vascular endothelial growth factor (EG-VEGF). EG-VEGF belongs to the prokineticin family that includes two key members PROK1 and PROK2, also called BV8 ( $\approx 8$  kDa) (LeCouter et al., 2001; Lin et al., 2002). These circulating ligands show differential expression patterns in humans depending on the organ/tissue types. For example, PROK1/EG-VEGF is highly expressed in peripheral tissues (specifically in steroidogenic organs such as human ovary, placenta and adrenal gland), whereas PROK2/Bv8 is widely expressed in the central nervous system and non-steroidogenic cells of the testes (LeCouter et al., 2001; Lin et al., 2002; Traboulsi et al., 2015). EG-VEGF and BV8 activate two G-protein linked receptors namely prokineticin receptor 1 (PROKR1) and -2 (PROKR2). The signaling pathways include, cAMP, Akt, and p42-p44 MAP-kinases phosphorylation and calcium mobilization. PROKs regulate a stunning array of biological functions such as gastrointestinal motility (LeCouter et al., 2001; Lin et al., 2002), circadian rhythm regulation, neurogenesis, angiogenesis, pain perception, mood regulation, and reproduction (Brouillet et al., 2010, 2012b; Alfaidy et al., 2016; Zhao et al., 2019). Dysregulation of PROKs/PROKR signaling pathways have been reported in a variety of diseases, such as cancer, abnormal angiogenesis and pregnancy pathologies (Brouillet et al., 2010, 2012b; Alfaidy et al., 2016; Traboulsi et al., 2017; Zhao et al., 2019).

Throughout normal human pregnancy, a dynamic expression of circulating EG-VEGF is found in the serum of pregnant women, with a five-fold increase during the first trimester ( $\approx 250$  pg/ml) (Hoffmann et al., 2009). The placental expression of EG-VEGF is high during the first trimester of human pregnancy, with a peak at 8–11 weeks of gestation (Hoffmann et al., 2009). PROKR1 and PROKR2 are highly expressed in villous cytotrophoblasts (VCT) as well as micro- and macrovascular placental endothelial cells (Brouillet et al., 2010). At the cellular level, EG-VEGF is expressed in the syncytiotrophoblast (ST), VCT, fetal endothelium, and Hofbauer cells (Ho) (Hoffmann et al., 2006; Holloway et al., 2014; Garnier et al., 2015).

## PROKINETICINS EXPRESSION IN MICRO- AND MACROVASCULAR PLACENTAL ECs

Lang et al. (2003) reported distinct morphogenetic, antigenic and functional differences between the two EC types present in the placenta, with respect to the secretion of vasoactive substances and the proliferative response to cytokines. The dissimilar responses of micro- and macrovascular ECs to various stimuli (Lang et al., 2003; Brouillet et al., 2010) most likely reflect differences in the activation of transcription factors, **Figure 1**. Despite these differences, HUVEC are the predominant cell type used to model placental vasculogenic and angiogenic processes (Demir et al., 1989). Microvascular ECs, despite being


the predominant cell type that vascularize the placental villi are less well understood. Importantly, these cells play critical role in placental disorders such as FGR and PE (Demir et al., 1989; Kingdom et al., 2000; Lang et al., 2003).

Endocrine gland derived-vascular endothelial growth factor enhances angiogenesis within the placental villi during the first trimester of human pregnancy (Brouillet et al., 2010). It controls various angiogenic processes including endothelial cell proliferation, survival, migration, tube organization, sprouting, permeability, and paracellular transport (Brouillet et al., 2010). Interestingly, EG-VEGF also displays stronger effects on the placental microvascular endothelial cells, the PLEC cells compare to its effects on the macrovascular cells, the HUVEC (Human umbilical vessel endothelial cells) (Brouillet et al., 2010). Via its PROKR1 receptor, EG-VEGF increases PLEC proliferation, migration and sprouting and controls their permeability via PROKR2. Importantly, EG-VEGF effects on PLEC cells were stronger compare to the VEGF effects on the same cells (Brouillet et al., 2010). At the fetomaternal interface, we demonstrated that EG-VEGF controls extra-villi angiogenesis through the inhibition of precocious invasion of EVT into the maternal spiral arteries (Hoffmann et al., 2009).

Recent studies from our own group and from others demonstrated that increased expression of EG-VEGF is associated with PE and FGR development (Hoffmann et al., 2006; Alfaidy, 2016; Inan et al., 2018). Our group also demonstrated that the concentration of EG-VEGF is significantly increased in both pathologies (Hoffmann et al., 2006; Murthi et al., 2015; Alfaidy, 2016). More recently, a study confirmed the increase in EG-VEGF levels in PE and proposed this factor as a biomarker for the diagnosis of PE patients (Inan et al., 2018).

## KEY REGULATORS OF PROKINETICINS FOR SUCCESSFUL PLACENTAL ANGIOGENESIS

Since its identification EG-VEGF has been associated with the control of placental angiogenesis during the first trimester of pregnancy and the reactivation of its receptors in the placenta of patients with pregnancy pathologies such as PE and FGR (Brouillet et al., 2010, 2012a, 2013a, 2014a,b; Garnier et al., 2015; Alfaidy, 2016; Sergent et al., 2016). The reactivation of angiogenic processes was meant to compensate for the associated deleterious vascularization. In these pathologies, EG-VEGF and receptors contribute to neoangiogenesis processes that allow pregnancy to progress. To fulfill these functions, EG-VEGF and its receptors have been reported to be regulated by key actors of vasculogenesis and angiogenesis within the placenta. The first supposed regulator of EG-VEGF expression was oxygen. This was reported by Ferrara et al. (LeCouter et al., 2001), as HIF1 $\alpha$  response element was identified in the promoter region of EG-VEGF and BV8. In 2006, we demonstrated that EG-VEGF and its PROKR1 receptor were upregulated by hypoxia in the human placenta (Hoffmann et al., 2006). EG-VEGF upregulation by hypoxia substantiated its role during the first trimester, as placental development and vascularization occur in hypoxic



environment during this period (Burton, 2009). In 2012, we demonstrated that another key actor of placental development, the human chorionic gonadotropin ( $\beta$ hCG) upregulates the expression of EG-VEGF and its receptors (Brouillet et al., 2012b). Importantly, in 2013, glycosylated-hCG has been reported to regulate another actor of angiogenesis, the TGF $\beta$  (Berndt et al., 2013). These findings suggest that EG-VEGF belongs to a complex of placenta proteins that are controlled by the master hormone,  $\beta$ hCG to fulfill a well-orchestrated angiogenesis. In the same line, we have also demonstrated in 2015 that part of the effects of the transcription factor PPAR $\gamma$  on placental angiogenesis are mediated by EG-VEGF, suggesting that the placental defects observed in the PPAR $\gamma$  knockout mice might well be due to deregulations in the EG-VEGF/PROKR functions (Garnier et al., 2015).

## TRANSCRIPTIONAL FACTORS IN THE CONTROL OF PLACENTAL ANGIOGENESIS

Cells respond to cues from growth factors and signaling molecules, allowing them to either maintain or alter their state of differentiation during angiogenesis (Irving and Lala, 1995). However, EC nuclear transcription factors determine how these cues are interpreted and drive the cellular response. Many different types of transcription factors play essential roles in placental cell differentiation, including endothelial and trophoblast cells (Cross et al., 2002). Most transcription factors

have common protein structural motifs allowing them to be placed into a few large families (e.g., zinc finger, leucine zipper, helix-loop-helix and helix-turn-helix) (Johnson and McKnight, 1989; Woodside et al., 2004). This review will mainly focus on members of the “homeobox” gene family of transcription factors.

## HOMEBOX GENES AND HOMEODOMAIN PROTEINS

Homeobox genes are frequently present as clusters of related homeobox genes called “HOX” cluster genes, but there are also individual, divergent HOX-like genes. Homeobox genes contain a highly conserved 180 base pair DNA sequence, which encodes a 60 amino acid “homeodomain” and contains a helix-turn-helix DNA binding motif. Although homeodomain proteins have similar DNA binding specificity, they regulate highly diverse and context-dependent cellular functions (Levine and Hoey, 1988), which includes the processes of vasculogenesis and angiogenesis.

## HOMEBOX GENES IN MURINE AND HUMAN PLACENTAL DEVELOPMENT

Targeted deletion of specific homeobox genes in murine models provides genetic proof of homeobox gene regulation of placental development during pregnancy (Rossant and Cross, 2001; Myers and Capper, 2002; Gorski and Leal, 2003; Gorski and Walsh, 2003). For example, targeted deletion of *Esx1*

(Fohn and Behringer, 2001) and *Dlx3* (Morasso et al., 1999) resulted in disruption of the vascular network in the placental labyrinthine layer (Cross et al., 2003). Mutant embryos in both cases were growth-restricted, and *Dlx3*<sup>-/-</sup> mutants were embryonic lethal due to adequate placental circulation (Morasso et al., 1999). These studies show homeobox genes are specific regulators of placental vascular development.

Recent studies from our laboratory provided comprehensive analyses of homeobox genes in human placental pathologies. We carried out the first screening of a 32-weeks placental cDNA library for homeobox genes, which led to the isolation of *DLX4*, *MSX2*, *GAX* and *HLX* (Quinn et al., 1997). Immunohistochemical analyses localized these homeobox genes/homeodomain proteins to trophoblasts and ECs (Murthi et al., 2006; Rajaraman et al., 2008; Chui et al., 2010). We also reported decreased homeobox gene *HLX* expression in ECs and trophoblast cells in FGR-affected placentae (Murthi et al., 2006). Homeobox genes *DLX4* and *DLX3* showed increased expression in FGR-affected placentae (Murthi et al., 2006), whereas *GAX* and *MSX2* showed no significant difference in expression.

## HOMEBOX GENE EXPRESSION IN MICRO- AND MACROVASCULAR PLACENTAL ECs

In our studies, primary PLEC were used to identify homeobox genes expressed in the placental microvasculature, and to compare that with macrovascular HUVEC. We detected mRNA expression of homeobox genes *HLX*, *MSX2*, *DLX3*, *DLX4*, and *GAX* and in both PLEC and HUVEC. Notably, *HLX* mRNA in HUVEC was significantly lower compared with PLEC (Murthi et al., 2007). These data provided evidence of heterogeneity in homeobox gene expression between microvascular PLEC and macrovascular HUVEC, which most likely reflects significant differences in EC function in the two different cellular environments, **Table 1**.

## FUNCTIONAL SIGNIFICANCE OF HOMEBOX GENES IN MICRO- AND MACROVASCULAR ECs

We detected high *HLX* mRNA expression in PLEC, which are also proliferative cell types compared with their macrovascular counterparts (Murthi et al., 2007). Moreover, in response to placental growth factor (PIGF), PLEC have more proliferative activity compared with HUVEC (Lang et al., 2003; Brouillet et al., 2010). Together, these data suggest a possible role for *HLX* in the proliferative capacity of microvascular ECs. The role(s), if any, of homeobox genes *HLX*, *MSX2*, *DLX3*, *DLX4*, and *GAX* in the transcriptional regulation of other PLEC function such as migration and invasion is yet to be explored. Quinn et al. (2000) proposed that co-expression of a combination of homeobox genes (i.e., *HLX*, *MSX2*, and *GAX*) may play a role in the regulation of epithelial-mesenchymal interactions in the placenta. Thus, co-expression of the homeobox genes in both trophoblast

and endothelium may also be important in coordinating villous outgrowth and angiogenesis within the terminal villi. Other studies showed that Homeobox genes regulate numerous key genes such as, *CDKN1*, *RB*, *GATA2*, *PPARG*, *ITGAV*, *NRP-1*, *ANGPT-1*, and *2* (Rajaraman et al., 2008; Chui et al., 2013; Novakovic et al., 2017; Harris et al., 2019; **Table 1**).

The repertoire of homeobox genes expressed in PLEC, was further investigated by microarray expression profiling of ECs (Murthi et al., 2008). We identified homeobox genes *TLX1*, *TLX2*, *TGIF*, *HEX*, *PHOX1*, *MEIS2*, *HOXB7*, and *LIM6* in PLEC. Importantly, our studies reported that these homeobox genes were differentially expressed in macro- compared with microvascular ECs, **Figure 1**. Functional studies in cultured ECs are underway in our laboratory to determine the role of these novel endothelial homeobox genes.

## GROWTH FACTOR REGULATION OF HOMEBOX GENES IN THE PLACENTA

Many EC growth factors and signal transduction pathways are involved in the maintenance of an efficient uteroplacental vasculature (Thaete et al., 2004). Our previous studies have demonstrated that *HLX* expression in *in vitro* models of human EVT, HTR8-SV neo and SGH-PL4 was significantly upregulated by HGF (Rajaraman et al., 2010) and CSF-1 (Rajaraman et al., 2007, 2008). Recent studies also showed that the homeobox gene *HOXB7* is regulated by *DKK1* and the *Wnt-1/b-catenin* (Huang et al., 2019; **Table 1**).

In 2015, our group demonstrated that the canonical member of the prokineticin family, EG-VEGF, controls homeobox genes expression in normal human placenta and in placenta from FGR pregnancies (Murthi et al., 2015). This regulation was observed in whole placenta explants, including endothelial, stroma and trophoblast cells. In particular, EG-VEGF up-regulated the following homeobox genes, *HOXA9*, *HOXC8*, *HOXC10*, *HOXD1*, *HOXD8*, *HOXD9*, and *HOXD11*, while downregulating the expression of *NKX 3.1*. Further investigations using an *in vitro* model of trophoblast cells, we demonstrated that reduced *NKX3.1* expression significantly enhanced premature differentiation and apoptosis in the syncytiotrophoblast cell line, the BeWo, while significantly reduced migration and invasive potential of the HTR8-SV neo cells (Murthi et al., 2015). This study was the first to demonstrate that the new placental angiogenic factor exhibits part of its effects on trophoblast invasion and differentiation through the *NKX3.1* homeobox gene. The demonstration that growth factors such as EG-VEGF regulate homeobox genes in the trophoblast lineage, especially the EVT, involved in extravillous angiogenesis, opens new perspectives into the potential involvement of homeobox genes in the cross talk between trophoblast and extra-villi and intra-villi endothelial cells to fulfill placental angiogenesis throughout pregnancy. Ongoing validations of the newly discovered EG-VEGF-regulated homeobox genes should bring more insights into the role of these homeobox genes in the control of angiogenic processes at both intra-villi and extra-villi sites, throughout human pregnancy, **Figure 1**.

**TABLE 1** | Summarizes the localizations of the prokineticin and HOX members within the placenta, their respective roles in the control of the angiogenesis in macro and microvascular blood vessels and lists their local regulators.

Placental prokinetins and homeobox genes	Role in the placenta	Type of placenta associated structure	Regulated genes in the placenta		Placental regulators	Associated placental vascular pathologies	References
EG-VEGF, PROKR1 and PROKR2	Angiogenesis, inflammation Placental development	Trophoblast cells, micro and macrovascular systems, Hofbauer cells	HOXD1, 8, 9, and 11 HOX A9 and HOXC8, 10		Nicotine, Hypoxia, $\beta$ hCG, PPARg	FGR, Hydatidiform moles, Choriocarcinoma and PE	Hoffmann et al., 2006, 2009; Brouillet et al., 2012a,b, 2013b; Holloway et al., 2014; Garnier et al., 2015; Murthi et al., 2015; Traboulsi et al., 2015, 2017; Sergent et al., 2016
HLX, DLX3, DLX4, MSX2, GAX, TLX1, TLX2	Cell Invasion, migration and proliferation Stem cell proliferation and differentiation	Microvascular system	CDKN1C (+) RB (+) GATA2, PPAR g ITGAV, NRP-1, ANPGT-1 and 2	VEGF, PLGF, HGF, CSF-1 Angiopoietins, PPARg, IGF-II, Endoglin, TGFb	IGFR2 (-) PLGF (+)	FGR and PE	Murthi et al., 2007, 2008; Rajaraman et al., 2008, 2010; Chui et al., 2013; Liang et al., 2016; Novakovic et al., 2017; Harris et al., 2019
TGIF		Micro and macrovascular systems				FGR (+)	Pathirage et al., 2013; Gunatillak et al., 2016
HEX	Role in hematopoiesis	Vascular system					Unpublished data
PHOX1		Macrovascular system					Unpublished data
MEIS2	Control of mouse placental vascularization	Macrovascular system	Activin and Inhibin				Unpublished data
LIM6		Macrovascular system	VEGF				Unpublished data
HOXB7	Endothelial differentiation	Macrovascular system			DKK1 (-) Wnt1/b catenin (-)	FGR (+)	Huang et al., 2019
NKX3.1	Role in trophoblast differentiation and proliferation	Trophoblast lineage			EG-VEGF (-)		Murthi et al., 2015
HOXD1, 8, 9, and 11 HOX A9 and HOXC8, 10			HOXA9 upregulates MMP14, EphB4, eNOS, VEGFR2		EG-VEGF (+)	FGR (+)	Murthi et al., 2015

*Also reports their associated placental vascular pathologies.*

## PROKINETICINS AND/OR HOMEBOX GENES AS POTENTIAL TARGETS IN FGR AND PE PATHOLOGIES

Numerous studies from our team have clearly demonstrated that EG-VEGF and its receptors are directly involved in normal placental vascularization and angiogenesis during the first trimester of pregnancy and that maintenance of increased circulating levels of placental EG-VEGF over that trimester is associated with the development of PE and FGR (Hoffmann et al., 2006, 2009; Brouillet et al., 2012b; Murthi et al., 2015; Traboulsi et al., 2015). These findings strongly suggest that antagonism of EG-VEGF signaling might contribute to the attenuation of vascular-pregnancy pathologies. Importantly, we have recently demonstrated that treatment of animal model of choriocarcinoma with PROKR2 antagonist significantly reduced tumor growth, vascularization and metastasis (Traboulsi et al., 2017). Hence, one can speculate that the benefit upon the antagonism of the prokineticin signaling might well trigger an upstream regulation of the EG-VEGF-dependent homeobox genes including NKX3.1, HOXA9, HOXC8, HOXC10, HOXD1, HOXD8, HOXD9, and HOXD11 as potential target genes for aberrant angiogenesis associated with the pathogenesis of PE

and FGR. Further studies are needed to better characterize the relationship between placental angiogenic factors and the homeobox genes to fulfill successful pregnancy outcomes.

## AUTHOR CONTRIBUTIONS

NA and PM designed the manuscript, supervised the progress of the review, and verified the English aspects. SB, GR, and BK wrote the different parts of the review. MB, TB, and PH helped with the clinical aspects and performed the figure and table. All authors contributed to the article and approved the submitted version.

## FUNDING

We acknowledge the following sources of funding: Institut National de la Santé et de la Recherche Médicale (U1036), University of Grenoble Alpes, Commissariat à l'Énergie Atomique (DSV/iRTSV/BCI), Région Auvergne-Rhône-Alpes "CLARA, Oncostarter," Ligue Nationale Contre le Cancer and Ligue Départementale (Savoie) Contre le Cancer, Inserm Transfert.

## REFERENCES

- Ahmed, A., and Perkins, J. (2000). Angiogenesis and intrauterine growth restriction. *Baillieres Best Pract. Res. Clin. Obstet. Gynaecol.* 14, 981–998. doi: 10.1053/beog.2000.0139
- Alfaidy, N. (2016). Prokineticin1 and pregnancy. *Ann. Endocrinol.* 77, 101–104. doi: 10.1016/j.ando.2016.04.014
- Alfaidy, N., Hoffmann, P., Gillois, P., Gueniffey, A., Lebayle, C., Garcin, H., et al. (2016). PROK1 level in the follicular microenvironment: a new noninvasive predictive biomarker of embryo implantation. *J. Clin. Endocrinol. Metab.* 101, 435–444. doi: 10.1210/jc.2015-1988
- Arderiu, G., Cuevas, I., Chen, A., Carrio, M., East, L., and Boudreau, N. J. (2007). HoxA5 stabilizes adherens junctions via increased Akt1. *Cell Adh. Migr.* 1, 185–195. doi: 10.4161/cam.1.4.5448
- Berndt, S., Blacher, S., Munaut, C., Detilleux, J., Perrier d'Hauterive, S., Huhtaniemi, I., et al. (2013). Hyperglycosylated human chorionic gonadotropin stimulates angiogenesis through TGF-beta receptor activation. *FASEB J.* 27, 1309–1321. doi: 10.1096/fj.12-213686
- Brouillet, S., Dufour, A., Prot, F., Feige, J. J., Equy, V., Alfaidy, N., et al. (2014a). Influence of the umbilical cord insertion site on the optimal individual birth weight achievement. *Biomed. Res. Int.* 2014:341251. doi: 10.1155/2014/341251
- Brouillet, S., Hoffmann, P., Alfaidy, N., and Feige, J. J. (2014b). [Prokineticins: new regulatory peptides in human reproduction]. *Med. Sci.* 30, 274–279. doi: 10.1051/medsci/20143003015
- Brouillet, S., Hoffmann, P., Benharouga, M., Salomon, A., Schaal, J. P., Feige, J. J., et al. (2010). Molecular characterization of EG-VEGF-mediated angiogenesis: differential effects on microvascular and macrovascular endothelial cells. *Mol. Biol. Cell* 21, 2832–2843. doi: 10.1091/mbc.E10-01-0059
- Brouillet, S., Hoffmann, P., Chauvet, S., Salomon, A., Chamboredon, S., Sergent, F., et al. (2012a). Revisiting the role of hCG: new regulation of the angiogenic factor EG-VEGF and its receptors. *Cell Mol. Life Sci.* 69, 1537–1550. doi: 10.1007/s00018-011-0889-x
- Brouillet, S., Hoffmann, P., Feige, J. J., and Alfaidy, N. (2012b). EG-VEGF: a key endocrine factor in placental development. *Trends Endocrinol. Metab.* 23, 501–508. doi: 10.1016/j.tem.2012.05.006
- Brouillet, S., Hoffmann, P., Thomas-Cadi, C., Bergues, U., Feige, J. J., Alfaidy, N., et al. (2013a). [PROK1, prognostic marker of embryo implantation?]. *Gynecol. Obstet. Fertil.* 41, 562–565. doi: 10.1016/j.gyobfe.2013.07.007
- Brouillet, S., Murthi, P., Hoffmann, P., Salomon, A., Sergent, F., De Mazancourt, P., et al. (2013b). EG-VEGF controls placental growth and survival in normal and pathological pregnancies: case of fetal growth restriction (FGR). *Cell Mol. Life Sci.* 70, 511–525. doi: 10.1007/s00018-012-1141-z
- Burton, G. J. (2009). Oxygen, the Janus gas; its effects on human placental development and function. *J. Anat.* 215, 27–35. doi: 10.1111/j.1469-7580.2008.00978.x
- Burton, G. J., and Jauniaux, E. (2001). Maternal vascularisation of the human placenta: does the embryo develop in a hypoxic environment? *Gynecol. Obstet. Fertil.* 29, 503–508. doi: 10.1016/s1297-9589(01)00179-5
- Chaddha, V., Viero, S., Huppertz, B., and Kingdom, J. (2004). Developmental biology of the placenta and the origins of placental insufficiency. *Semin. Fetal Neonatal Med.* 9, 357–369. doi: 10.1016/j.siny.2004.03.006
- Charnock-Jones, D. S. (2002). Soluble flt-1 and the angiopoietins in the development and regulation of placental vasculature. *J. Anat.* 200, 607–615. doi: 10.1046/j.1469-7580.2002.00063.x
- Chen, C. P., Bajoria, R., and Aplin, J. D. (2002). Decreased vascularization and cell proliferation in placentas of intrauterine growth-restricted fetuses with abnormal umbilical artery flow velocity waveforms. *Am. J. Obstet. Gynecol.* 187, 764–769. doi: 10.1067/mob.2002.125243
- Chui, A., Kalionis, B., Abumaree, M., Cocquebert, M., Fournier, T., Evain-Brion, D., et al. (2013). Downstream targets of the homeobox gene DLX3 are differentially expressed in the placentae of pregnancies affected by human idiopathic fetal growth restriction. *Mol. Cell. Endocrinol.* 377, 75–83. doi: 10.1016/j.mce.2013.06.032
- Chui, A., Pathirage, N. A., Johnson, B., Cocquebert, M., Fournier, T., Evain-Brion, D., et al. (2010). Homeobox gene distal-less 3 is expressed in proliferating and differentiating cells of the human placenta. *Placenta* 31, 691–697. doi: 10.1016/j.placenta.2010.05.003
- Clark, D. E., Smith, S. K., Sharkey, A. M., and Charnock-Jones, D. S. (1996). Localization of VEGF and expression of its receptors flt and KDR in human placenta throughout pregnancy. *Hum. Reprod.* 11, 1090–1098. doi: 10.1093/oxfordjournals.humrep.a019303



- Cross, J. C., Baczyk, D., Dobric, N., Hemberger, M., Hughes, M., Simmons, D. G., et al. (2003). Genes, development and evolution of the placenta. *Placenta* 24, 123–130. doi: 10.1053/plac.2002.0887
- Cross, J. C., Hemberger, M., Lu, Y., Nozaki, T., Whiteley, K., Masutani, M., et al. (2002). Trophoblast functions, angiogenesis and remodeling of the maternal vasculature in the placenta. *Mol. Cell. Endocrinol.* 187, 207–212. doi: 10.1016/s0303-7207(01)00703-1
- Demir, R., Kaufmann, P., Castellucci, M., Erben, T., and Kotowski, A. (1989). Fetal vasculogenesis and angiogenesis in human placental villi. *Acta Anat.* 136, 190–203. doi: 10.1159/000146886
- Fohn, L. E., and Behringer, R. R. (2001). ESX1L, a novel X chromosome-linked human homeobox gene expressed in the placenta and testis. *Genomics* 74, 105–108. doi: 10.1006/geno.2001.6532
- Garnier, V., Traboulsi, W., Salomon, A., Brouillet, S., Fournier, T., Winkler, C., et al. (2015). PPARgamma controls pregnancy outcome through activation of EG-VEGF: new insights into the mechanism of placental development. *Am. J. Physiol. Endocrinol. Metab.* 309, E357–E369. doi: 10.1152/ajpendo.00093.2015
- Gorski, D. H., and Leal, A. J. (2003). Inhibition of endothelial cell activation by the homeobox gene Gax. *J. Surg. Res.* 111, 91–99. doi: 10.1016/s0022-4804(03)00042-8
- Gorski, D. H., and Walsh, K. (2003). Control of vascular cell differentiation by homeobox transcription factors. *Trends Cardiovasc. Med.* 13, 213–220. doi: 10.1016/s1050-1738(03)00081-1
- Gude, N. M., Roberts, C. T., Kalionis, B., and King, R. G. (2004). Growth and function of the normal human placenta. *Thromb. Res.* 114, 397–407. doi: 10.1016/j.thromres.2004.06.038
- Harris, L. K., Pantham, P., Yong, H. E. J., Pratt, A., Borg, A. J., Crocker, I., et al. (2019). The role of insulin-like growth factor 2 receptor-mediated homeobox gene expression in human placental apoptosis, and its implications in idiopathic fetal growth restriction. *Mol. Hum. Reprod.* 25, 572–585. doi: 10.1093/molehr/gaz047
- Hoffmann, P., Feige, J. J., and Alfaidy, N. (2006). Expression and oxygen regulation of endocrine gland-derived vascular endothelial growth factor/prokineticin-1 and its receptors in human placenta during early pregnancy. *Endocrinology* 147, 1675–1684. doi: 10.1210/en.2005-0912
- Hoffmann, P., Saoudi, Y., Benharouga, M., Graham, C. H., Schaal, J. P., Mazouni, C., et al. (2009). Role of EG-VEGF in human placenta: physiological and pathological implications. *J. Cell Mol. Med.* 13, 2224–2235. doi: 10.1111/j.1582-4934.2008.00554.x
- Holloway, A. C., Salomon, A., Soares, M. J., Garnier, V., Raha, S., Sergent, F., et al. (2014). Characterization of the adverse effects of nicotine on placental development: in vivo and in vitro studies. *Am. J. Physiol. Endocrinol. Metab.* 306, E443–E456. doi: 10.1152/ajpendo.00478.2013
- Huang, L., Ying, H., Chen, Z., Zhu, Y. L., Gu, Y., Hu, L., et al. (2019). Down-regulation of DKK1 and Wnt1/beta-catenin pathway by increased homeobox B7 resulted in cell differentiation suppression of intrauterine fetal growth retardation in human placenta. *Placenta* 80, 27–35. doi: 10.1016/j.placenta.2019.03.001
- Inan, C., Varol, F. G., Erzincan, S. G., Uzun, I., Sutcu, H., and Sayin, N. C. (2018). Use of prokineticin-1 (PROK1), pregnancy-associated plasma protein A (PAPP-A) and PROK1/PAPP-A ratio to predict adverse pregnancy outcomes in the first trimester: a prospective study. *J. Matern. Fetal Neonatal Med.* 31, 2685–2692. doi: 10.1080/14767058.2017.1351536
- Irving, J. A., and Lala, P. K. (1995). Functional role of cell surface integrins on human trophoblast cell migration: regulation by TGF-beta, IGF-II, and IGFBP-1. *Exp. Cell Res.* 217, 419–427. doi: 10.1006/excr.1995.1105
- Jackson, M. R., Carney, E. W., Lye, S. J., and Ritchie, J. W. (1994). Localization of two angiogenic growth factors (PDECGF and VEGF) in human placenta throughout gestation. *Placenta* 15, 341–353. doi: 10.1016/0143-4004(94)90002-7
- Johnson, P. F., and McKnight, S. L. (1989). Eukaryotic transcriptional regulatory proteins. *Annu. Rev. Biochem.* 58, 799–839. doi: 10.1146/annurev.bi.58.070189.004055
- Kaufmann, P., Mayhew, T. M., and Charnock-Jones, D. S. (2004). Aspects of human fetoplacental vasculogenesis and angiogenesis. II. Changes during normal pregnancy. *Placenta* 25, 114–126. doi: 10.1016/j.placenta.2003.10.009
- Kingdom, J., Huppertz, B., Seaward, G., and Kaufmann, P. (2000). Development of the placental villous tree and its consequences for fetal growth. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 92, 35–43. doi: 10.1016/s0301-2115(00)00423-1
- Lang, I., Pabst, M. A., Hiden, U., Blaschitz, A., Dohr, G., Hahn, T., et al. (2003). Heterogeneity of microvascular endothelial cells isolated from human term placenta and macrovascular umbilical vein endothelial cells. *Eur. J. Cell Biol.* 82, 163–173. doi: 10.1078/0171-9335-00306
- LeCouter, J., Kowalski, J., Foster, J., Hass, P., Zhang, Z., Dillard-Telm, L., et al. (2001). Identification of an angiogenic mitogen selective for endocrine gland endothelium. *Nature* 412, 877–884. doi: 10.1038/35091000
- Levine, M., and Hoey, T. (1988). Homeobox proteins as sequence-specific transcription factors. *Cell* 55, 537–540. doi: 10.1016/0092-8674(88)90209-7
- Liang, H., Zhang, Q., Lu, J., Yang, G., Tian, N., Wang, X., et al. (2016). MSX2 Induces Trophoblast Invasion in Human Placenta. *PLoS One* 11:e0153656. doi: 10.1371/journal.pone.0153656
- Lin, D. C., Bullock, C. M., Ehlert, F. J., Chen, J. L., Tian, H., and Zhou, Q. Y. (2002). Identification and molecular characterization of two closely related G protein-coupled receptors activated by prokineticins/endocrine gland vascular endothelial growth factor. *J. Biol. Chem.* 277, 19276–19280. doi: 10.1074/jbc.M202139200
- Maulik, D., Frances Evans, J., and Ragolia, L. (2006). Fetal growth restriction: pathogenic mechanisms. *Clin. Obstet. Gynecol.* 49, 219–227. doi: 10.1097/00003081-200606000-00005
- Mayhew, T. M. (2003). Changes in fetal capillaries during preplacental hypoxia: growth, shape remodelling and villous capillarization in placenta from high-altitude pregnancies. *Placenta* 24, 191–198. doi: 10.1053/plac.2002.0895
- Mayhew, T. M., Ohadike, C., Baker, P. N., Crocker, I. P., Mitchell, C., and Ong, S. S. (2003). Stereological investigation of placental morphology in pregnancies complicated by pre-eclampsia with and without intrauterine growth restriction. *Placenta* 24, 219–226. doi: 10.1053/plac.2002.0900
- Meegdes, B. H., Ingenhoves, R., Peeters, L. L., and Exalto, N. (1988). Early pregnancy wastage: relationship between chorionic vascularization and embryonic development. *Fertil. Steril.* 49, 216–220. doi: 10.1016/s0015-0282(16)59704-0
- Morasso, M. I., Grinberg, A., Robinson, G., Sargent, T. D., and Mahon, K. A. (1999). Placental failure in mice lacking the homeobox gene *Dlx3*. *Proc. Natl. Acad. Sci. U.S.A.* 96, 162–167. doi: 10.1073/pnas.96.1.162
- Murthi, P., Brouillet, S., Pratt, A., Borg, A., Kalionis, B., Goffin, F., et al. (2015). An EG-VEGF-dependent decrease in homeobox gene *NKX3.1* contributes to cytotrophoblast dysfunction: a possible mechanism in human fetal growth restriction. *Mol. Med.* 21, 645–656. doi: 10.2119/molmed.2015.00071
- Murthi, P., Hiden, U., Rajaraman, G., Liu, H., Borg, A. J., Coombes, F., et al. (2008). Novel homeobox genes are differentially expressed in placental microvascular endothelial cells compared with macrovascular cells. *Placenta* 29, 624–630. doi: 10.1016/j.placenta.2008.04.006
- Murthi, P., Said, J. M., Doherty, V. L., Donath, S., Nowell, C. J., Brennecke, S. P., et al. (2006). Homeobox gene *DLX4* expression is increased in idiopathic human fetal growth restriction. *Mol. Hum. Reprod.* 12, 763–769. doi: 10.1093/molehr/gal087
- Murthi, P., So, M., Gude, N. M., Doherty, V. L., Brennecke, S. P., and Kalionis, B. (2007). Homeobox genes are differentially expressed in macrovascular human umbilical vein endothelial cells and microvascular placental endothelial cells. *Placenta* 28, 219–223. doi: 10.1016/j.placenta.2006.02.012
- Myers, L. J., and Capper, W. L. (2002). A transmission line model of the human foetal circulatory system. *Med. Eng. Phys.* 24, 285–294. doi: 10.1016/s1350-4533(02)00019-x
- Novakovic, B., Fournier, T., Harris, L. K., James, J., Roberts, C. T., Yong, H. E. J., et al. (2017). Increased methylation and decreased expression of homeobox genes *TLX1*, *HOXA10* and *DLX5* in human placenta are associated with trophoblast differentiation. *Sci. Rep.* 7:4523. doi: 10.1038/s41598-017-04776-5
- Patel, N. S., Li, J. L., Generali, D., Poulosom, R., Cranston, D. W., and Harris, A. L. (2005). Up-regulation of delta-like 4 ligand in human tumor vasculature and the role of basal expression in endothelial cell function. *Cancer Res.* 65, 8690–8697. doi: 10.1158/0008-5472.CAN-05-1208
- Quinn, L. M., Johnson, B. V., Nicholl, J., Sutherland, G. R., and Kalionis, B. (1997). Isolation and identification of homeobox genes from the human placenta including a novel member of the Distal-less family, *DLX4*. *Gene* 187, 55–61. doi: 10.1016/s0378-1119(96)00706-8

- Quinn, L. M., Latham, S. E., and Kalionis, B. (2000). The homeobox genes MSX2 and MOX2 are candidates for regulating epithelial-mesenchymal cell interactions in the human placenta. *Placenta* 21(Suppl. A), S50–S54. doi: 10.1053/plac.1999.0514
- Rajaraman, G., Murthi, P., Brennecke, S. P., and Kalionis, B. (2010). Homeobox gene HLX is a regulator of HGF/c-met-mediated migration of human trophoblast-derived cell lines. *Biol. Reprod.* 83, 676–683. doi: 10.1095/biolreprod.109.078634
- Rajaraman, G., Murthi, P., Leo, B., Brennecke, S. P., and Kalionis, B. (2007). Homeobox gene HLX1 is a regulator of colony stimulating factor-1 dependent trophoblast cell proliferation. *Placenta* 28, 991–998. doi: 10.1016/j.placenta.2007.03.011
- Rajaraman, G., Murthi, P., Quinn, L., Brennecke, S. P., and Kalionis, B. (2008). Homeodomain protein HLX is expressed primarily in cytotrophoblast cell types in the early pregnancy human placenta. *Reprod. Fertil. Dev.* 20, 357–367. doi: 10.1071/rd07159
- Regnault, T. R., Galan, H. L., Parker, T. A., and Anthony, R. V. (2002). Placental development in normal and compromised pregnancies— a review. *Placenta* 23(Suppl. A), S119–S129. doi: 10.1053/plac.2002.0792
- Rossant, J., and Cross, J. C. (2001). Placental development: lessons from mouse mutants. *Nat. Rev. Genet.* 2, 538–548. doi: 10.1038/35080570
- Salafia, C. M., Charles, A. K., and Maas, E. M. (2006). Placenta and fetal growth restriction. *Clin. Obstet. Gynecol.* 49, 236–256. doi: 10.1097/00003081-200606000-00007
- Sergent, F., Hoffmann, P., Brouillet, S., Garnier, V., Salomon, A., Murthi, P., et al. (2016). Sustained endocrine gland-derived vascular endothelial growth factor levels beyond the first trimester of pregnancy display phenotypic and functional changes associated with the pathogenesis of pregnancy-induced hypertension. *Hypertension* 68, 148–156. doi: 10.1161/HYPERTENSIONAHA.116.07442
- Thaete, L. G., Dewey, E. R., and Neerhof, M. G. (2004). Endothelin and the regulation of uterine and placental perfusion in hypoxia-induced fetal growth restriction. *J. Soc. Gynecol. Investig.* 11, 16–21. doi: 10.1016/j.jsigi.2003.07.001
- Traboulsi, W., Brouillet, S., Sergent, F., Boufettal, H., Samouh, N., Aboussaouira, T., et al. (2015). Prokineticins in central and peripheral control of human reproduction. *Horm. Mol. Biol. Clin. Investig.* 24, 73–81. doi: 10.1515/hmbci-2015-0040
- Traboulsi, W., Sergent, F., Boufettal, H., Brouillet, S., Slim, R., Hoffmann, P., et al. (2017). Antagonism of EG-VEGF Receptors as Targeted Therapy for Choriocarcinoma Progression In Vitro and In Vivo. *Clin Cancer Res* 23, 7130–7140. doi: 10.1158/1078-0432.CCR-17-0811
- Tsatsaris, V., Goffin, F., Munaut, C., Brichtant, J. F., Pignon, M. R., Noel, A., et al. (2003). Overexpression of the soluble vascular endothelial growth factor receptor in preeclamptic patients: pathophysiological consequences. *J. Clin. Endocrinol. Metab.* 88, 5555–5563. doi: 10.1210/jc.2003-030528
- Velicky, P., Knofler, M., and Pollheimer, J. (2016). Function and control of human invasive trophoblast subtypes: intrinsic vs. maternal control. *Cell Adh. Migr.* 10, 154–162. doi: 10.1080/19336918.2015.1089376
- Woodside, K. J., Shen, H., Muntzel, C., Daller, J. A., Sommers, C. L., and Love, P. E. (2004). Expression of Dlx and Lhx family homeobox genes in fetal thymus and thymocytes. *Gene Expr. Patterns* 4, 315–320. doi: 10.1016/j.modgep.2003.10.003
- Zhao, Y., Wu, J., Wang, X., Jia, H., Chen, D. N., and Li, J. D. (2019). Prokineticins and their G protein-coupled receptors in health and disease. *Prog. Mol. Biol. Transl. Sci.* 161, 149–179. doi: 10.1016/bs.pmbts.2018.09.006
- Zou, J., Luo, H., Zeng, Q., Dong, Z., Wu, D., and Liu, L. (2011). Protein kinase CK2alpha is overexpressed in colorectal cancer and modulates cell proliferation and invasion via regulating EMT-related genes. *J. Transl. Med.* 9:97. doi: 10.1186/1479-5876-9-97
- Zygmunt, M., Herr, F., Munstedt, K., Lang, U., and Liang, O. D. (2003). Angiogenesis and vasculogenesis in pregnancy. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 110(Suppl. 1), S10–S18. doi: 10.1016/s0301-2115(03)00168-4

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Alfaidy, Brouillet, Rajaraman, Kalionis, Hoffmann, Barjat, Benharouga and Murthi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.