



HAL
open science

Electrochemical Intramolecular Oxytrifluoromethylation of N -Tethered Alkenyl Alcohols: Synthesis of Functionalized Morpholines

Aurélie Claraz, Thibaut Courant, Geraldine Masson

► **To cite this version:**

Aurélie Claraz, Thibaut Courant, Geraldine Masson. Electrochemical Intramolecular Oxytrifluoromethylation of N -Tethered Alkenyl Alcohols: Synthesis of Functionalized Morpholines. *Organic Letters*, 2020, 22 (4), pp.1580-1584. 10.1021/acs.orglett.0c00176 . hal-03058500

HAL Id: hal-03058500

<https://hal.science/hal-03058500v1>

Submitted on 13 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Electrochemical Intramolecular Oxytrifluoromethylation of *N*-tethered Alkenyl Alcohols: Synthesis of Functionalized Morpholines.

Aurélie Claraz,* Thibaut Courant, and Géraldine Masson*

Institut de Chimie des Substances Naturelles, CNRS UPR 2301, Université Paris-Sud, Université Paris-Saclay, 1, av. de la Terrasse, Gif-sur-Yvette 91198 Cedex, France

Supporting Information Placeholder




ABSTRACT: An electrochemical intramolecular oxytrifluoromethylation of *N*-tethered alkenyl alcohols was developed providing straightforward access to CF₃-containing morpholines derivatives. The method features mild reaction conditions with direct anodic oxidation of Langlois reagent as cheap and easy to handle trifluoromethylating reagent. Various substituted 2-(2,2,2-trifluoroethyl)morpholines were obtained in moderate to high yields under constant current electrolysis in an undivided cell.

Tetrahydro-1,4-oxazines, namely morpholine derivatives, are privileged core skeletons that continually appear in numerous pharmaceuticals and agrochemicals.^{1,2} It belongs to the top 25 most frequent *N*-heterocycles in US FDA approved drugs.³ Recently, trifluoromethylated morpholines have received substantial interest because the incorporation of a trifluoromethyl (CF₃) group can significantly improve their physical, biological and chemical properties.⁴ Therefore, several efficient methods have been developed to construct trifluoromethylated morpholines that rely on two main strategies (Scheme 1a). The first one involves the construction of the morpholine core *via* either cyclization of trifluoromethylated linear compounds⁵ or the ring expansion of CF₃-containing heterocycles.⁶ The second one is based on trifluoromethylation of existing morpholines.⁷ Although both approaches have their own particular advantages, exploring more efficient synthetic methodologies for valuable trifluoromethylated morpholines is still highly desired.

Over past decades, the intramolecular difunctionalization of alkenes involving the simultaneous formation of C–CF₃ and C–O bonds has proven to be a powerful tool for synthesis of trifluoromethylated oxacycles.⁸ This approach involves the generation of a CF₃ radical via a single-electron transfer (SET) process which reacts with an alkene to provide a β-CF₃-substituted radical intermediate. The resulting radical then undergoes a stepwise SET oxidation and subsequent intramolecular O-nucleophilic attack to produce oxygen-containing heterocycles (Scheme 1b). Surprisingly, to the best of our knowledge, such an intramolecular oxytrifluoromethylation has not been investigated for the preparation of CF₃-containing morpholines. The development of such a cyclization reaction would provide an efficient synthesis of novel family of trifluoromethylated morpholines (Scheme 1c).

Scheme 1. Synthesis of CF₃-containing Morpholines




In recent years, electrochemistry⁹ has emerged as a powerful tool to introduce a CF₃ group into alkenes¹⁰ through the anodic oxidation of the cheap and easy to handle Langlois reagent (CF₃SO₂Na).^{11,12} It has been applied for the construction of pyrrolidines,¹³ oxindoles,¹⁴ indoline or hydroisoquinoline-1,3-diones,^{14b} quinolinones and lactones⁸ⁿ *via* intramolecular processes. In this context and in line with our long-standing interest in rad-

ical trifluoromethylation of alkenes,^{15,81} we speculated that electrochemical trifluoromethylation/cyclization of *N*-tethered alken-6-ols is a worthwhile approach to accessing trifluoromethylated morpholines. Herein we report our preliminary efforts toward the construction of new type of 2-(2,2,2-trifluoroethyl)morpholine derivatives.

Our investigations began by studying the cyclization of alken-6-ol **1a** with Langlois reagent **2** in an undivided cell using graphite carbon cathode, a platinum cathode under constant current electrolysis and a mixture of CH₃CN and H₂O (v/v 1:1) (Table 1). With LiClO₄ as the electrolyte, the desired morpholine **3a** was isolated in 62% yield (entry 1). After testing other organic solvents such as acetone and THF (entries 2 and 3), CH₃CN was considered to be the best one. Replacing H₂O by methanol had no significant effect. When the electrolysis was conducted in pure CH₃CN to reduce the formation of intermolecular oxytrifluoromethylated product,¹⁶ **3a** was isolated with an improved yield of 79%. Pleasingly, switching platinum plated anode for cheap carbon graphite or nickel electrodes gave similar results (entries 6 and 7). Further experiments have shown that a change of the electrolyte or current intensity (from 15 to 30 mA) leads to a significantly decrease of yield (entries 8 and 9).¹⁷

Table 1. Optimization of Reaction Conditions^a




entry	electrodes	solvent	yield ^b (%)
1	C(+) Pt(-)	CH ₃ CN/H ₂ O (10:1)	62
2	C(+) Pt(-)	acetone/H ₂ O (10:1)	48
3	C(+) Pt(-)	THF/H ₂ O (10:1)	39
4	C(+) Pt(-)	CH ₃ CN/MeOH (10:1)	63
5	C(+) Pt(-)	CH ₃ CN	79
6	C(+) C(-)	CH ₃ CN	78
7	C(+) Ni(-)	CH ₃ CN	81
8 ^c	C(+) Ni(-)	CH ₃ CN	50
9 ^d	C(+) Ni(-)	CH ₃ CN	69
10 ^e	C(+) Ni(-)	CH ₃ CN	0

^aReaction conditions: undivided cell, **1a** (0.25 mmol), **2** (0.5 mmol), solvent (2.5 mL), LiClO₄ (0.2 M), CCE (15 mA), 3.2 F.mol⁻¹, rt. ^bIsolated yield. ^c*n*-Bu₄NBF₄ instead of LiClO₄. ^d30 mA instead of 15 Ma. ^eNo current.

With the best reaction conditions in hand, we next explored the scope of the electrolytic oxytrifluoromethylation (Scheme 2). A set of representative *N*-tethered alken-6-ols with various aryl groups on the alkene proved to be suitable substrates, generating the desired morpholines in good to high yields (**3a-3f**). In general, electron-poor aromatic rings with halogen substituents reacted smoothly whereas slightly lower yield was obtained with a *para*-*tert*-butyl substituent. This is mainly due to a competitive trifluoromethylation of the arene ring. The *N*-2-nitrophenylsulfonyl substrates **1g** and **1h** underwent the oxytrifluo-

romethylation reaction with equal efficiency. Pleasingly secondary alcohol **1k** was well tolerated furnishing the corresponding 2,2,6-trisubstituted morpholine **3k** in good yield as a mixture of diastereoisomers. Moreover, enantioenriched alcohols **1i** and **1j** were successfully engaged affording the corresponding morpholines **3i** and **3j** in 36% and 65% yield, respectively, as a mixture of diastereoisomers. The lower yield with **1i** is presumably due to overoxidation of the benzylic carbon. The polycyclic compounds **3l** and **3m** were produced in descent yields albeit with moderate diastereoselectivity. To our delight, this method can also be applied to the synthesis of a difluoromethylated morpholine **3n** under slightly modified reaction conditions with 3 equivalents of CF₂HOSO₂Na¹⁸ as a source of CF₂H radical¹⁹ in CH₃CN/H₂O (10:1). To evaluate the potential application of this protocol, a 1 mmol scale reaction of **1a** was performed under the standard reaction conditions. Delightfully, the desired morpholine product (**3a**) was obtained in a comparable yield (79%).

Scheme 2. Intramolecular Trifluoromethylation of *N*-tethered Alken-6-ols **1^a**




^aReaction conditions: undivided cell, C(+)|Ni(-), **1a** (0.25 mmol), **2** (0.5 mmol), CH₃CN (2.5 mL), LiClO₄ (0.2 M), CCE (15 mA), 3.2 F.mol⁻¹, rt. ^bdr were determined by ¹H NMR analysis of the reaction mixtures. ^cReaction conducted on 1 mmol scale.

A further investigation revealed that the formation of seven-membered oxacycle **5** by intramolecular oxytrifluoromethylation way of *N*-tethered alken-7-ol **4** was also possible (Scheme

3). This provides a rapid access to 1,4-benzoxazepine, an important class of seven-membered rings heterocycles featuring interesting biological properties and found in numerous drugs and preclinical leads.²⁰


Scheme 3. Intramolecular Trifluoromethylation of *N*-tethered Alken-7-ols 4^a



^aReaction conditions: undivided cell, C(+)|Ni(-), **4** (0.25 mmol), **2** (0.5 mmol), CH₃CN (2.5 mL), LiClO₄ (0.2 M), CCE (15 mA), 3.2 F.mol⁻¹, rt.

To gain more insight into the mechanism of the oxytrifluoromethylation of *N*-tethered alken-6-ols, some preliminary control experiments were performed. It is noteworthy that there is no reaction in the absence of electric current (Table 1, entry 10). Cyclic voltammetry (CV) analysis of Langlois reagent (**2**) in CH₃CN with LiClO₄ as electrolyte showed irreversible anodic oxidation waves (See the Supporting Information, Figure S1) at a potential much lower than that of substrate **1a** (1.48 and 2.07 V vs Ag/AgCl respectively). On the basis of the above results as well as other reports,¹⁰ a plausible reaction mechanism is depicted in scheme 4. The SET anodic oxidation of CF₃SO₂Na would produce the CF₃ radical through the release of SO₂ from CF₃SO₂ radical. Subsequent regioselective addition of electrophilic CF₃ radical to alkene **1** would lead to the radical intermediate **6**, which would be further oxidized into cation **7** by a second SET anodic oxidation. Final intramolecular nucleophilic addition of the alcohol moiety would lead to the corresponding trifluoromethylated morpholine **3** along with the release of H⁺. Reductions of H⁺ and residual trace of water would be the counter-reactions taking place at the cathode.


Scheme 4. Plausible Mechanism



To further demonstrate the usefulness of our method to access morpholine derivatives, the tosyl group of **3a** was easily removed under mild conditions by treatment with magnesium in

a sonicated methanol solution affording unprotected morpholine **8** in high yield (Scheme 5).

Scheme 5. Deprotection of Morpholine **3a**



To conclude, we have developed an intramolecular radical oxytrifluoromethylation of *N*-tethered alken-6-ols by using environmentally friendly electrochemical reaction conditions with Langlois reagent as cheap and easy to handle source of trifluoromethyl radical. The procedure is suitable for the synthesis of a wide variety of functionalized oxytrifluoromethylated morpholine derivatives. This protocol was extended to a *N*-tethered alken-7-ol providing the synthesis of a CF₃-containing benzoxazepine. Extension of this method to the preparation of other trifluoromethylated heterocycles is currently underway in our laboratory and will be reported in due course.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/>.

Experimental procedures and spectroscopic data for all compounds (PDF)

AUTHOR INFORMATION

Corresponding Author

Aurélien Claraz, <https://orcid.org/0000-0002-0692-7830>, E-mail: aurelie.claraz@cnrs.fr

Géraldine Masson, Institut de Chimie des Substances Naturelles CNRS <https://orcid.org/0000-0003-2333-7047>, E-mail: geraldine.masson@cnrs.fr

Other Authors

Thibaut Courant, Institut de Chimie des Substances Naturelles

Note

The authors declare no competing financial interest.

ACKNOWLEDGMENT

We thank ICSN and CNRS for financial support. TC thanks ICSN for a postdoctoral fellowship.

REFERENCES

- (1) (a) Arshad, F.; Khan, M. F.; Akhtar, W.; Alam, M. M.; Nainwal, L. M.; Kaushik, S. K.; Akhter, M.; Parvez, S.; Hasan, S. M.; Shaquiquzaman, M. Revealing quinquennial anticancer journey of morpholine: A SAR based review. *Eur. J. Med. Chem.* **2019**, *167*, 324. (b) Andrs, M.; Korabecny, J.; Jun, D.; Hodny, Z.; Bartek, J.; Kuca, K. Phosphatidylinositol 3-kinase (PI3K) and phosphatidylinositol 3-kinase-related kinase (PIKK) inhibitors: Importance of the morpholine ring. *J. Med. Chem.* **2015**, *58*, 41. (c) Pal'chikov, V. A. Morpholines. Synthesis and biological activity. *Russ. J. Org. Chem.* **2013**, *49*, 787. (d) Lamberth, C. Morpholine Fungicides for the Treatment of Powdery Mildew. In *Bioactive Heterocyclic Compound Classes: Agrochemicals*; Lamberth, C., Dinges, J., Eds.; WILEY-VCH Verlag: Weinheim, 2012, Chapter 10.
- (2) (a) For a recent review on the synthesis of morpholines: Palchikov, V. A.; Chebanov, V. A. Recent progress in the synthesis of morpholines. *Chem. Heterocycl. Compd.* **2019**, *55*, 324. For selected

examples: (b) Bissonnette, N. B.; Ellis, J. M.; Hamann, L. G.; Romanov-Michailidis F. Expedient access to saturated nitrogen heterocycles by photoredox cyclization of imino-tethered dihydropyridines. *Chem. Sci.* **2019**, *10*, 9591. (c) Aubineau, T.; Cossy, J. A One-Pot Reaction toward the Diastereoselective Synthesis of Substituted Morpholines. *Org. Lett.* **2018**, *20*, 7419. (d) Zhai, H.; Borzenko, A.; Lau, Y. Y.; Ahn, S. H.; Schafer, L. L. Catalytic asymmetric synthesis of substituted morpholines and piperazines. *Angew. Chem. Int. Ed.* **2012**, *51*, 12219. (e) Luescher, M. U.; Bode, J. W. Catalytic Synthesis of N-Unprotected Piperazines, Morpholines, and Thiomorpholines from Aldehydes and SnAP Reagents. *Angew. Chem. Int. Ed.* **2015**, *54*, 10884. (f) Deka, M. J.; Indukuri, K.; Sultana, S.; Borah, M.; Saikia, A. K. Synthesis of Five-, Six-, and Seven-Membered 1,3- and 1,4-Heterocyclic Compounds via Intramolecular Hydroalkoxylation/Hydrothioalkoxylation of Alkenols/Thioalkenols. *J. Org. Chem.* **2015**, *80*, 4349. (g) Sequeira, F. C.; Chemler, S. R. Stereoselective synthesis of morpholines via copper-promoted oxyamination of alkenes. *Org. Lett.* **2012**, *14*, 4482. (h) Patil, P.; Madhavachary, R.; Kurpiewska, K.; Kalinowska-Thuscik, J.; Dömling, A. De Novo Assembly of Highly Substituted Morpholines and Piperazines. *Org. Lett.* **2017**, *19*, 642. (i) Kishorevandavasi, J.; Hu, W. P.; Chandrusenadi, G.; Chen, H. T.; Chen, H. Y.; Hsieh, K. C.; Wang, J. J. Aryl λ^3 -Iodane-Mediated 6-exo-trig Cyclization to Synthesize Highly Substituted Chiral Morpholines. *Adv. Synth. Catal.* **2015**, *357*, 2788.

(3) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the structural diversity, substitution patterns, and frequency of nitrogen heterocycles among U.S. FDA approved pharmaceuticals. *J. Med. Chem.* **2014**, *57*, 10257.

(4) (a) Hagemann, W. K. The Many Roles for Fluorine in Medicinal Chemistry. *J. Med. Chem.* **2008**, *51*, 4359. (b) Ilardi, E. A.; Vitaku, E.; Njardarson, J. T. Data-mining for sulfur and fluorine: An evaluation of pharmaceuticals to reveal opportunities for drug design and discovery. *J. Med. Chem.* **2014**, *57*, 2832. (c) Müller, K.; Faeh, C.; Diederich, F. Fluorine in Pharmaceuticals: Looking Beyond Intuition. *Science* **2007**, *317*, 1881.

(5) (a) Rulev, A. Y.; Romanov, A. R.; Kondrashov, E. V.; Ushakov, I. A.; Vashchenko, A. V.; Muzalevskiy, V. M.; Nenajdenko, V. G. Domino Assembly of Trifluoromethylated N,O-Heterocycles by the Reaction of Fluorinated α -Bromoones with Amino Alcohols. *J. Org. Chem.* **2016**, *81*, 10029. (b) Rulev, A. Y.; Romanov, A. R.; Kondrashov, E. V.; Ushakov, I. A.; Muzalevskiy, V. M.; Nenajdenko, V. G. Assembly of Trifluoromethylated Morpholines through Cascade Reactions of Bromoenones with Secondary Amino Alcohols. *Eur. J. Org. Chem.* **2018**, 4202. (c) Chaume, G.; Van Severen, M. C.; Marinkovic, S.; Brigaud, T. Straightforward synthesis of (S)- and (R)- α -trifluoromethyl proline from chiral oxazolindines derived from ethyl trifluoropyruvate. *Org. Lett.* **2006**, *8*, 6123. (d) Chaume, G.; Van Severen, M. C.; Ricard, L.; Brigaud, T. Concise access to enantiopure (S)- and (R)- α -trifluoromethyl pyroglutamic acids from ethyl trifluoropyruvate-based chiral CF₃-oxazolindines (Fox). *J. Fluorine Chem.* **2008**, *129*, 1104. (e) Zaporozhets, O. A.; Zozulya, S.; Shishkin, O. V.; Kravchuk, O.; Shcherbatiuk, A. V.; Yarmoliuk, D. V.; Shishkina, S. V.; Tolmachev, A. A.; Moriev, R.; Manoilenko, O.; et al. Synthesis of 2- and 3-trifluoromethylmorpholines: useful building blocks for drug discovery. *Tetrahedron* **2013**, *69*, 3796.

(6) Kenis, S.; D'Hooghe, M.; Verniest, G.; Reybroeck, M.; Dang Thi, T. A.; Pham The, C.; Thi Pham, T.; Törnroos, K. W.; Van Tuyen, N.; De Kimpe, N. Nucleophile-directed selective transformation of cis-1-tosyl-2-tosyloxymethyl-3-(trifluoromethyl)aziridine into aziridines, azetidines, and benzo-fused dithianes, oxathianes, dioxanes, and (thio)morpholines. *Chem. Eur. J.* **2013**, *19*, 5966.

(7) (a) Fustero, S.; Albert, L.; Aceña, J. L.; Sanz-Cervera, J. F.; Asensio, A. An efficient entry to optically active anti- and syn- β -amino- α -trifluoromethyl alcohols. *Org. Lett.* **2008**, *10*, 605. (b) Takata, T.; Hirano, K.; Miura, M. Synthesis of α -Trifluoromethylamines by Cu-Catalyzed Regio- and Enantioselective Hydroamination of 1-Trifluoromethylalkenes. *Org. Lett.* **2019**, *21*, 4284. (c) Philippe, C.; Kaffy, J.; Milcent, T.; Bonnet-Delpon, D. Synthesis and Evaluation of Fluorinated Analogues of Monoamine Reuptake Inhibitors. *J. Fluor. Chem.* **2012**, *134*, 136.

(8) For selected examples under oxidative conditions: (a) Yu, J.; Yang, J.; Fu, H. Transition Metal-Free Trifluoromethylation of N-Alkylamides with Sodium Trifluoromethanesulfonate: Synthesis of Trifluoromethyl-Containing Oxazolines. *Adv. Synth. Catal.* **2014**, *356*, 3669. Under copper-catalysis: (b) Wang, Y.; Jiang, M.; Liu, J.-T. Copper-Catalyzed Diastereoselective Synthesis of Trifluoromethylated Tetrahydrofurans. *Adv. Synth. Catal.* **2016**, *358*, 1322. (c) He, Y.-T.; Li, L.-H.; Yang, Y.-F.; Wang, Y.-Q.; Luo, J.-Y.; Liu, X.-Y.; Liang, Y.-M. Copper-catalyzed synthesis of trifluoromethyl-substituted isoxazolines. *Chem. Commun.* **2013**, *49*, 5687. (d) Cheng, Y.-F.; Dong, X.-Y.; Gu, Q.-S.; Yu, Z.-L.; Liu, X.-Y. Achiral Pyridine Ligand-Enabled Enantioselective Radical Oxytrifluoromethylation of Alkenes with Alcohols. *Angew. Chem., Int. Ed.* **2017**, *56*, 8883. (e) Zhu, R.; Buchwald, S. L. Enantioselective Functionalization of Radical Intermediates in Redox Catalysis: Copper-Catalyzed Asymmetric Oxytrifluoromethylation of Alkenes. *Angew. Chem., Int. Ed.* **2013**, *52*, 12655. (f) Zhu, R.; Buchwald, S. L. Versatile Enantioselective Synthesis of Functionalized Lactones via Copper-Catalyzed Radical Oxyfunctionalization of Alkenes. *J. Am. Chem. Soc.* **2015**, *137*, 8069. (g) Zhu, R.; Buchwald, S. L. Copper-Catalyzed Oxytrifluoromethylation of Unactivated Alkenes. *J. Am. Chem. Soc.* **2012**, *134*, 30. Under photocatalysis: (h) Kim, E.; Choi, S.; Kim, H.; Cho, E. J. Generation of CF₃-containing epoxides and aziridines by visible-light-driven trifluoromethylation of allylic alcohols and amines. *Chem. Eur. J.* **2013**, *19*, 6209. (i) Yasu, Y.; Arai, Y.; Tomita, R.; Koike, T.; Akita, M. Highly regio- and diastereoselective synthesis of CF₃-substituted lactones via photoredox-catalyzed carbolactonization of alkenoic acids. *Organic Lett.* **2014**, *16*, 780. (j) Deng, Q.-H.; Chen, J.-R.; Wei, Q.; Zhao, Q.-Q.; Lu, L.-Q.; Xiao, W.-J. Visible-light-induced photocatalytic oxytrifluoromethylation of N-allylamides for the synthesis of CF₃-containing oxazolines and benzoxazines. *Chemical Commun.* **2015**, *51*, 3537. (k) Noto, N.; Miyazawa, K.; Koike, T.; Akita, M. Anti-diastereoselective synthesis of CF₃-containing spirooxazolines and spirooxazines via regioselective trifluoromethylative spirocyclization by photoredox catalysis. *Organic Lett.* **2015**, *17*, 3710. (l) Jarrige, L.; Carboni, A.; Dagousset, G.; Levitre, G.; Magnier, E.; Masson, G. Photoredox-Catalyzed Three-Component Tandem Process: An Assembly of Complex Trifluoromethylated Phthalans and Isoindolines. *Org. Lett.* **2016**, *18*, 2906. (m) Noto, N.; Tanaka, Y.; Koike, T.; Akita, M. Strongly Reducing (Diarylamino)anthracene Catalyst for Metal-Free Visible-Light Photocatalytic Fluoroalkylation. *ACS Catal.* **2018**, *8*, 9408. Under electrochemical conditions: (n) Zhang, S.; Li, L.; Zhang, J.; Zhang, J.; Xue, M.; Xu, K. Electrochemical Fluoromethylation Triggered Lactonizations of Alkenes under Semi-Aqueous Conditions. *Chem. Sci.* **2019**, *10*, 3181.

(9) Selected review: (a) Sperry, J. B.; Wright, D. L. The application of cathodic reductions and anodic oxidations in the synthesis of complex molecules. *Chem. Soc. Rev.* **2006**, *35*, 605. (b) Jutand, A. Contribution of Electrochemistry to Organometallic Catalysis. *Chem. Rev.* **2008**, *108*, 2300. (c) Yoshida, J.-i.; Kataoka, K.; Horcajada, R.; Nagaki, A. Modern Strategies in Electroorganic Synthesis. *Chem. Rev.* **2008**, *108*, 2265. (d) Frontana-Urbe, B. A.; Little, R. D.; Ibanez, J. G.; Palma, A.; Vasquez-Medrano, R. Organic Electrosynthesis: A Promising Green Methodology in Organic Chemistry. *Green Chem.* **2010**, *12*, 2099. (e) Horn, E. J.; Rosen, B. R.; Baran, P. S. Synthetic Organic Electrochemistry: An Enabling and Innately Sustainable Method. *ACS Cent. Sci.* **2016**, *2*, 302. (f) Yan, M.; Kawamata, Y.; Baran, P. S. Synthetic Organic Electrochemical Methods since 2000: On the Verge of a Renaissance. *Chem. Rev.* **2017**, *117*, 13230. (g) Feng, R.; Smith, J. A.; Moeller, K. D. Anodic Cyclization Reactions and the Mechanistic Strategies That Enable Optimization. *Acc. Chem. Res.* **2017**, *50*, 2346. (h) Wiebe, A.; Gieshoff, T.; Möhle, S.; Rodrigo, E.; Zirbes, M.; Waldvogel, S. R. Electrifying Organic Synthesis. *Angew. Chem., Int. Ed.* **2018**, *57*, 5594. (e) Waldvogel, S. R. Modern Electrochemical Aspects for the Synthesis of Value-Added Organic Products. *Angew. Chem., Int. Ed.* **2018**, *57*, 6018. (f) Jiang, Y.; Xu, K.; Zeng, C. Use of Electrochemistry in the Synthesis of Heterocyclic Structures. *Chem. Rev.* **2018**, *118*, 4485. (g) Jiang, Y.; Xu, K.; Zeng, C. Use of Electrochemistry in the Synthesis of Heterocyclic Structures. *Chem. Rev.* **2018**, *118*, 4485. (h) Sauer, S. G.; Lin, S. An Electrocatalytic Approach to the Radical Difunctionalization of Alkenes. *ACS Catal.* **2018**, *8*, 5175. (i) Sauermann, N.; Meyer, T. H.; Qiu, Y.; Ackermann, L.

Electrocatalytic C–H Activation. *ACS Catal.* **2018**, *8*, 7086. (j) Kärkäs, M. D. Electrochemical strategies for C–H functionalization and C–N bond formation. *Chem. Soc. Rev.* **2018**, *47*, 5786. (k) Wang, H.; Gao, X.; Lv, Z.; Abdelilah, T.; Lei, A. Recent Advances in Oxidative R1-H/R2-H Cross-Coupling with Hydrogen Evolution via Photo-/Electrochemistry. *Chem. Rev.* **2019**, *119*, 6769. (l) Yuan Y, Lei A. Electrochemical Oxidative Cross-Coupling with Hydrogen Evolution Reactions. *Acc. Chem. Res.* **2019**, *52*, 3309

(10) Selected examples: (a) Jud, W.; Kappe, C. O.; Cantillo, D. Catalyst-Free Oxytrifluoromethylation of Alkenes through Paired Electrolysis in Organic-Aqueous Media. *Chem. Eur. J.* **2018**, *24*, 17234. (b) Zhang, L.; Zhang, G.; Wang, P.; Li, Y.; Lei, A. Electrochemical Oxidation with Lewis-Acid Catalysis Leads to Trifluoromethylative Difunctionalization of Alkenes Using $\text{CF}_3\text{SO}_2\text{Na}$. *Org. Lett.* **2018**, *20*, 7396. (c) Ye, K. Y.; Pombar, G.; Fu, N.; Sauer, G. S.; Keresztes, I.; Lin, S. Anodically Coupled Electrolysis for the Heterodifunctionalization of Alkenes. *J. Am. Chem. Soc.* **2018**, *140*, 2438. (d) Huang, Y.; Hong, H.; Zou, Z.; Liao, C.; Lu, J.; Qin, Y.; Li, Y.; Chen, L. Electrochemical Vicinal Aminotrifluoromethylation of Alkenes: High Regioselective Acquisition of β -Trifluoromethylamines. *Org. Biomol. Chem.* **2019**, *17*, 5014. (e) Sun, X.; Ma, H. X.; Mei, T. S.; Fang, P.; Hu, Y. Electrochemical Radical Formyloxylation-Bromination, -Chlorination, and -Trifluoromethylation of Alkenes. *Org. Lett.* **2019**, *21*, 3167. (f) Kang, J.-C.; Tu, Y.-Q.; Dong, J.-W.; Chen, C.; Zhou, J.; Ding, T.M.; Zai, J.T.; Chen, Z.-M.; Zhang, S.-Y.; Electrochemical Semipinacol Rearrangements of Allylic Alcohols: Construction of All-Carbon Quaternary Stereocenters. *Org. Lett.* **2019**, *21*, 2536. (g) Guan, Z.; Wang, H.; Huang, Y.; Wang, Y.; Wang, S.; Lei, A. Electrochemical Oxidative Aryl(alkyl)trifluoromethylation of Allyl Alcohols via 1,2-Migration. *Org. Lett.* **2019**, *21*, 12, 4619.

(11) (a) Langlois, B.; Laurent, E.; Roidot, N., Trifluoromethylation of aromatic compounds with sodium trifluoromethanesulfinate under oxidative conditions. *Tetrahedron Lett.* **1991**, *32*, 7525. (b) Laurent, E.; Roidot, N. Trifluoromethylation of Organic Disulfides with Sodium Trifluoromethanesulfinate under Oxidative Conditions: Synthesis of Trifluoromethyl Thioethers. *Phosphorus, Sulfur Silicon Relat. Elem.* **1991**, *59*, 169. (c) Wang, D.; Deng, G.; Chen, S.; Gong, H. Catalyst-free direct C–H trifluoromethylation of arenes in water–acetonitrile. *Green Chem.* **2016**, *18*, 5967. (d) Lefebvre, Q. Toward Sustainable Trifluoromethylation Reactions: Sodium Trifluoromethanesulfinate under the Spotlight. *Synlett* **2016**, *28*, 19. (e) Guyon, H.; Chachignon, H.; Cahard, D. $\text{CF}_3\text{SO}_2\text{X}$ (X = Na, Cl) as Reagents for Trifluoromethylation, Trifluoromethylsulfenyl-, -sulfinyl- and -sulfonylation. Part 1: Use of $\text{CF}_3\text{SO}_2\text{Na}$. *Beilstein J. Org. Chem.* **2017**, *13*, 2764.

(12) For pioneering work on the electrochemical oxidation of Langlois reagent: Tommasino, J. B.; Brondex, A.; Médebielle, M.; Thomalla, M.; Langlois, B. R.; Billard, T. Trifluoromethylation Reactions with Potassium Trifluoromethanesulfinate under Electrochemical Oxidation. *Synlett* **2002**, 1697.

(13) Ye, K. Y.; Song, Z.; Sauer, G. S.; Harenberg, J. H.; Fu, N.; Lin, S. Synthesis of Chlorotrifluoromethylated Pyrrolidines by Electrocatalytic Radical Ene-Yne Cyclization. *Chem. Eur. J.* **2018**, *24*, 12274.

(14) (a) Jiang, Y.-Y.; Dou, G.-Y.; Xu, K.; Zeng, C.-C. Bromide-Catalyzed Electrochemical Trifluoromethylation/Cyclization of N-Arylacrylamides with Low Catalyst Loading. *Org. Chem. Front.* **2018**, *5*,

2573. (b) Zhang, Z.; Zhang, L.; Cao, Y.; Li, F.; Bai, G.; Liu, G.; Yang, Y.; Mo, F. Mn-Mediated Electrochemical Trifluoromethylation/ $\text{C}(\text{sp}^2)$ -H Functionalization Cascade for the Synthesis of Azaheterocycles. *Org. Lett.* **2019**, *21*, 762. (c) Ruan, Z.; Huang, Z.; Xu, Z.; Mo, G.; Tian, X.; Yu, X. Y.; Ackermann, L. Catalyst-Free, Direct Electrochemical Tri- and Difluoroalkylation/Cyclization: Access to Functionalized Oxindoles and Quinolinones. *Org. Lett.* **2019**, *21*, 1237.

(15) (a) Courant T.; Masson, G. Recent Progress in Visible-Light Photoredox-Catalyzed Intermolecular 1,2- Difunctionalization of Double Bonds via an ATRA-Type Mechanism. *J. Org. Chem.* **2016**, *81*, 6945. (b) Carboni, A.; Dagousset, G.; Magnier, E.; Masson, G. One pot and selective intermolecular aryl- and heteroaryltrifluoromethylation of alkenes by photoredox catalysis. *Chem. Commun.* **2014**, *50*, 14197. (c) Dagousset, G.; Carboni, A.; Magnier, E.; Masson, G. *Org. Lett.* **2014**, *16*, 4340. (d) Carboni, A.; Dagousset, G.; Magnier, E.; Masson, G. Three-Component Photoredox-Mediated Chloro-, Bromo-, or Iodo-trifluoromethylation of Alkenes. *Synthesis* **2015**, *47*, 2439. (e) Levitre, G.; Dagousset, G.; Anselmi, E.; Tuccio, B.; Magnier, E.; Masson, G. Four-Component Photoredox-Mediated Azidoalkoxy-trifluoromethylation of Alkenes. *Org. Lett.* **2019**, *21*, 6005. (f) Daniel, M.; Dagousset, G.; Diter, P.; Klein, P.-A.; Tuccio, B.; Goncalves, A.-M.; Masson, G.; Magnier, E. Fluorinated Sulfilimino Iminiums: Efficient and Versatile Sources of Perfluoroalkyl Radicals under Photoredox Catalysis. *Angew. Chem. Int. Ed.* **2017**, *56*, 3997.

(16) About 16% of alcohol corresponding to intermolecular oxytrifluoromethylation is detected through analysis of the ^1H NMR spectrum of the crude material in a mixture of $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (10:1).

(17) For more details on the optimization study with other electrolytes, combination of organic solvents and electrodes, see the SI.

(18) He, Z.; Tan, P.; Ni, C.; Hu, J. Fluoroalkylative aryl migration of conjugated N-arylsulfonylated amides using easily accessible sodium di- and monofluoroalkanesulfonates. *Org. Lett.* **2015**, *17*, 1838.

(19) For other examples of electrochemical generation of CF_2H radical from $\text{CF}_2\text{HSO}_2\text{Na}$: see ref 14c; 8n; 10d; from other reagents: (a) Xiong, P.; Xu, H. H.; Song, J.; Xu, H. C. Electrochemical Difluoromethylation of Alkynes. *J. Am. Chem. Soc.* **2018**, *140*, 2460. (b) O'Brien, A. G.; Maruyama, A.; Inokuma, Y.; Fujita, M.; Baran, P. S.; Blackmond, D. G. Radical C–H functionalization of heteroarenes under electrochemical control. *Angew. Chem. Int. Ed.* **2014**, *53*, 11868.

(20) (a) Popp, T. A.; Tallant, C.; Rogers, C.; Fedorov, O.; Brennan, P. E.; Müller, S.; Knapp, S.; Bracher, F. Development of Selective CBP/P300 Benzoxazepine Bromodomain Inhibitors. *J. Med. Chem.* **2016**, *59*, 8889. (b) Takeuchi, C. S.; Kim, B. G.; Blazey, C. M.; Ma, S.; Johnson, H. W. B.; Anand, N. K.; Arcalás, A.; Baik, T. G.; Buhr, C. A.; Cannoy, J.; et al. Discovery of a novel class of highly potent, selective, ATP-competitive, and orally bioavailable inhibitors of the mammalian target of rapamycin (mTOR). *J. Med. Chem.* **2013**, *56*, 2218. (c) Miki, T.; Kori, M.; Mabuchi, H.; Tozawa, R.-I.; Nishimoto, T.; Sugiyama, Y.; Teshima, K.; Yukimasa, H. Synthesis of novel 4,1-benzoxazepine derivatives as squalene synthase inhibitors and their inhibition of cholesterol synthesis. *J. Med. Chem.* **2002**, *45*, 4571.