



**HAL**  
open science

## **Processive Steps in the Reverse Direction Require Uncoupling of the Lead Head Lever Arm of Myosin VI**

Julie Ménétreay, Tatiana Isabet, Virginie Ropars, Monalisa Mukherjea, Olena Pylypenko, Xiaoyan Liu, Javier Perez, Patrice Vachette, H Lee Sweeney, Anne Houdusse

► **To cite this version:**

Julie Ménétreay, Tatiana Isabet, Virginie Ropars, Monalisa Mukherjea, Olena Pylypenko, et al.. Processive Steps in the Reverse Direction Require Uncoupling of the Lead Head Lever Arm of Myosin VI. *Molecular Cell*, 2012, 48 (1), pp.75 - 86. 10.1016/j.molcel.2012.07.034 . hal-03059379

**HAL Id: hal-03059379**

**<https://hal.science/hal-03059379>**

Submitted on 25 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Processive Steps in the Reverse Direction Require Uncoupling of the Lead Head Lever Arm of Myosin VI

Julie Ménétrey,<sup>1,2,6,7</sup> Tatiana Isabet,<sup>1,2,6</sup> Virginie Ropars,<sup>1,2</sup> Monalisa Mukherjea,<sup>3</sup> Olena Pylypenko,<sup>1,2</sup> Xiaoyan Liu,<sup>3</sup> Javier Perez,<sup>4</sup> Patrice Vachette,<sup>5</sup> H. Lee Sweeney,<sup>3,\*</sup> and Anne M. Houdusse<sup>1,2,\*</sup>

<sup>1</sup>Structural Motility, Institut Curie, Centre de Recherche, Paris, F-75248 France

<sup>2</sup>CNRS, UMR144, 26 rue d'Ulm, 75248 Paris Cedex 05, France

<sup>3</sup>Department of Physiology, University of Pennsylvania School of Medicine, 3700 Hamilton Walk, Philadelphia, PA 19104-6085, USA

<sup>4</sup>Synchrotron SOLEIL, BP 48, 91192 Gif-sur-Yvette Cedex, France

<sup>5</sup>IBBMC, Université Paris Sud CNRS UMR8619, Bâtiment 430, F-91405 Orsay Cedex, France

<sup>6</sup>These authors contributed equally to this work

<sup>7</sup>Present address: Laboratoire d'Enzymologie et Biochimie Structurales, Centre de Recherche de Gif, CNRS, 91198 Gif-sur-Yvette, France

\*Correspondence: [anne.houdusse@curie.fr](mailto:anne.houdusse@curie.fr) (A.M.H.), [lsweeney@mail.med.upenn.edu](mailto:lsweeney@mail.med.upenn.edu) (H.L.S.)

<http://dx.doi.org/10.1016/j.molcel.2012.07.034>

## SUMMARY

Myosin VI is the only known reverse-direction myosin motor. It has an unprecedented means of amplifying movements within the motor involving rearrangements of the converter subdomain at the C terminus of the motor and an unusual lever arm projecting from the converter. While the average step size of a myosin VI dimer is 30–36 nm, the step size is highly variable, presenting a challenge to the lever arm mechanism by which all myosins are thought to move. Herein, we present structures of myosin VI that reveal regions of compliance that allow an uncoupling of the lead head when movement is modeled on actin. The location of the compliance restricts the possible actin binding sites and predicts the observed stepping behavior. The model reveals that myosin VI, unlike plus-end directed myosins, does not use a pure lever arm mechanism, but instead steps with a mechanism analogous to the kinesin neck-linker uncoupling model.


## INTRODUCTION

Myosin VI is unique in the superfamily of myosin motors in its ability to perform processive steps that are large on average, but highly variable in size, and in the reverse direction as compared to all other characterized myosin motors on actin (Rock et al., 2001; Nishikawa et al., 2002). Even though its directionality has been reversed, myosin VI undergoes the same kinetic cycle (Figure 1) for chemomechanical transduction as for all other characterized myosins (Sweeney and Houdusse, 2010). The cycle begins when myosin binds to actin in a state known as the pre-powerstroke state that has hydrolyzed ATP, but has trapped the hydrolysis products, MgADP and P<sub>i</sub>. The interaction with actin causes a conformational change in the myosin that releases P<sub>i</sub> and allows a strong binding interface to be created. This induces further conformational changes that increase the binding affinity for actin and trigger the release of

MgADP to form a rigor conformation on actin. Concomitant with the changes in binding affinity and product release, these conformational changes cause a swing of the C-terminal subdomain of the motor, which is known as the converter. This drives the movement of an extended alpha-helix that binds multiple calmodulin (CaM) or calmodulin-like light chains. Both the converter and the CaM-/light chain-binding region form the myosin lever arm. The lever arm amplifies the conformational changes in the motor into a movement on the order of nanometers known as the powerstroke. The powerstroke ends with the formation of the rigor state in which myosin is strongly bound to actin and has released MgADP. ATP binding to the rigor state on actin induces conformational changes in the myosin motor that causes a large cleft at the actin interface to open, destroying high affinity actin binding and creating the post-rigor state, which rapidly dissociates from actin. Once the myosin has dissociated, there is a rapid and reversible isomerization between the post-rigor state, which cannot hydrolyze ATP, and the pre-powerstroke state that can rapidly hydrolyze ATP due to repositioning of a nucleotide-binding element known as switch II (Figures 1 or 2A). This movement of switch II is part of a conformational change in the motor that repositions the myosin lever arm, repriming the lever arm for movement on actin.

Although the motor domain of myosin VI undergoes similar rearrangements in its ATPase cycle as for plus-end motors, the movement of its lever arm is in the opposite direction. This is accomplished by a repositioning of the CaM-binding region on the converter. In order to reposition the myosin VI lever arm, myosin VI contains an insert (insert-2) found only in Class VI and XXX myosins (Odrionitz and Kollmar, 2007) at the junction between the converter and the CaM-binding region. The first part of insert-2 interacts with the converter while the distal part of the insert-2, and the IQ motif that follows, each bind a calmodulin to create an unusual and repositioned CaM-binding region (Ménétrey et al., 2005). The lever arm in turn is followed by a proximal tail region (PTD), which folds as a triple helix bundle in the monomeric head but opens in the dimeric molecule as an elongated lever arm extension (Mukherjea et al., 2009).

A further adaptation of myosin VI to enable a larger powerstroke to occur in the reverse direction is that during the transition from pre-powerstroke the converter subdomain undergoes



**Figure 1. ATPase Cycle of the Myosin VI Motor**

The different structural states that the myosin VI motor adopts along the ATP hydrolysis cycle are indicated with detached states (Post-rigor, 2VAS, and Pre-powerstroke [PPS], 4ANJ) and actin-binding states (PPS-4ANJ and Rigor, 2BKH). The motor is composed of important connectors (SwII, orange – relay, yellow – SH1 helix, red) and four subdomains (Nter, gray – U50, blue – L50, white – converter, green). Insert-2 and its bound Ca<sup>2+</sup>-calmodulin are in purple and pink.

rearrangement creating two distinct conformations of the converter (Ménétreay et al., 2007), which we refer to herein as the R-fold and P-fold. The R-fold is found in the rigor state (R) at the end of the powerstroke, and is similar to the conformation that has been seen in plus-end-directed myosins. The P-fold found in the pre-powerstroke state (PPS) differs significantly. In the rigor state, the P-fold cannot be explored due to steric hindrance, but in the PPS state, modeling of a R-fold of the converter is possible without creating steric collision.

Myosin VI can dimerize upon interaction with its cargo proteins creating a dimer that moves processively along an actin filament with steps that are highly variable in length (Phichith et al., 2009). The lever arm extension appears to be less rigid than that of well characterized myosins such as those of class II and V, which initially was proposed to be the basis for variability of the myosin VI step size (Rock et al., 2005). However later work suggested that the variability of the step sizes appears to be derived from properties of the myosin converter itself, or to the nature of the converter interactions with the motor domain and/or the rest of the lever arm, since chimeric molecules with rigid lever arms after insert-2 also display the same degree of step size variability (Park et al., 2007; Bryant et al., 2007). We report herein structural investigations that provide insights into how myosin VI moves on actin. They demonstrate the location of compliance within the lever arm and show that it differs in different states of the motor. These structures explain the observed processive, but highly variable, movement of myosin VI in the reverse direction on actin.


## RESULTS AND DISCUSSION

To attempt to explain the variability of the myosin VI step size on actin, we examined the possibility that there are multiple possible structures of the pre-powerstroke (PPS) state, the structural state that the lead head of a processive dimer is in when it rebinds to actin (Figure 1). We did this by solving six structures of the pre-powerstroke state of myosin VI in different crystal packing environments. Based on these structures, we modeled how variability in the converter conformation and/or in the lever arm position in the PPS state might influence where a lead head can dock on actin as well as the subsequent power stroke, thus contributing to the variability of the myosin VI dimer's step size.

### The Myosin VI Pre-Powerstroke State—Overall Structure

As discussed above and depicted in Figure 1, the so-called pre-powerstroke (PPS) state of myosin is the state that reattaches to actin with MgADP and P<sub>i</sub> trapped in the active site, initiating the powerstroke and force generation. The first structure of the pre-powerstroke state of myosin VI was derived from a construct truncated prior to the CaM-binding site of insert-2 and thus was a motor domain (MD) construct without the CaM-binding region (Ménétreay et al., 2007). Interestingly the converter, the C-terminal subdomain of the MD that controls the lever arm orientation, had adopted a new conformation (P-fold) compared to that previously described for the nucleotide-free state (R-fold). The R-fold conformation of the myosin VI converter is the same as seen for all other myosin structures to date. As described in the Supplemental Information, the transition between these two configurations of the converter is made possible due to conformational changes of loops, called hereafter hinge regions that connect together the converter secondary elements: the β sheet to the helices and the helices to each other (Figure S1).

The six structures of the PPS state allowed identification of compliant elements in the motor in this state. In particular, the influence of the lever arm on the converter conformation was visualized at 2.6 Å resolution in a PPS crystal structure of



**Figure 2. Variability in the Converter of Myosin VI in Different Pre-Powerstroke State Structures**

(A) The structure of the myosin VI MD<sup>ins2-GFP</sup> prior to force generation (pre-powerstroke state) is shown with the motor subdomains and important structural motifs indicated in colors (Switch II in orange, relay in yellow and SH1 helix in red). Insert-2 and calmodulin are shown in purple and pink, while the GFP fusion is in green.

(B) (Right) The converter and its contacts with the Relay and the N-terminus subdomain are shown with the same color code as in A (except for the last helix shown in orange). (Left) A ribbon structure of the converter is displayed with the

construct encompassing the first part of the lever arm (insert-2 bound to its CaM with four bound Ca<sup>2+</sup>) fused to a Green Fluorescent Protein (GFP) molecule (MD<sup>ins2-GFP</sup> construct; Figure 2A).

In order to generate new crystal packing forms of the myosin VI PPS, we created a new construct (MD<sup>PM</sup>) with modified surface residues (D23R, I24R and R569E), which would be predicted to prevent the packing seen in earlier MD crystals. We solved ~2.3 Å resolution structures of both the MD and MD<sup>PM</sup> constructs in two new crystal forms (see Table 1 and Experimental Procedures) with two independent molecules in each asymmetric unit. In total, this allowed structural comparisons of six structures in the PPS state (PDB code 2V26 (Ménétreay et al., 2007); and five new structures) crystallized in different packing environments. In all these structures, there are no significant differences to report for the regions of the motor domain prior to the converter (residues 5-705) with an average root mean square deviation (rmsd) lower than 0.68 Å on 662 C $\alpha$  atoms. The MD structure is very similar to that found for plus-end motors in this state (Ménétreay et al., 2007), except for the conformation of the converter. The motor is in a conformation that positions the converter appropriately for a “primed” lever arm position, characteristic of the PPS.

### The Converter Always Adopts a P-Fold Conformation in the Pre-Powerstroke State

In all five of the new PPS structures of myosin VI, the converter adopts the P-fold conformation, as determined previously (Ménétreay et al. 2007) (rmsd < 0.85 Å on 68 C $\alpha$  atoms; residues 706-773) (Figure 2D). This conformation is markedly different from the one observed in the rigor state structure, which we refer to as the R-fold (rmsd > 3.5 Å on 68 C $\alpha$  atoms; residues 706-773) (PDB code 2BKI, Ménétreay et al., [2005]) (Figure S1). It is important to note that one of the molecules of the MD<sup>PM</sup> structure adopts a well-defined P-fold even though the R-fold would be compatible with the packing environment of the crystal and there is sufficient space to allow the transition in the crystal (Figure S2A).

To remove any possible crystal packing constraints, we investigated the conformation of the MD<sup>ins2-GFP</sup> construct in solution using Small-angle X-ray Scattering (SAXS). The molecule was trapped in the PPS state using MgADP.VO4. The SAXS studies clearly showed that, in solution, the converter adopts a P-fold in the PPS state and does not populate the R-fold in the PPS state (Figure S2).

While the position of the motor domain structural elements that control the overall position of the converter (known as the

conserved regions indicated in blue and the variable regions in red. The hinge and the insert-2 are indicated in gray and purple, respectively.

(C) Cross-eyed stereo view of the variable interactions between insert-2 and the rest of the converter in different PPS structures superimposed on insert-2 (4ANJ in green (converter) and purple (insert-2), 2V26 in black, 4E7Z\_molB in blue and 4E7Z\_molA in red).

(D) The converter in the pre-powerstroke state of myosin VI. A superposition on the converter of the six different PPS structures is shown with the same color code as in C. Note that overall the variation in the P-fold conformation of the converters result in different orientations of the insert-2 proximal helix. Modeling of the full insert-2 helix is shown to highlight the degree of variability in the insert-2 orientation.

**Table 1. Data Collection and Refinement Statistics**

	MD <sup>ins2-GFP</sup>	MD <sup>PM</sup>	MD <sup>crystal form 2</sup>
Data Collection			
Space group	C2	P2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	P2 <sub>1</sub>
Cell dimensions			
<i>a</i>	193.09	81.95	98.15
<i>b</i>	62.66	135.14	93.32
<i>c</i> (Å)	156.04	196.54	101.87
$\beta$ (°)	117.96	90.00	90.60
Molecules per asym. unit	1	2	2
Resolution (Å)	48.8-2.6	30-2.25	30-2.3
(High. Res. Shell)	(2.74-2.60)	(2.30-2.25)	(2.35-2.3)
R <sub>meas</sub>	11.4 (42.8)	11.1 (46.2)	7.1 (47.9)
<i>I</i> / $\sigma$ <i>I</i>	7.2 (2.0)	12.86 (3.44)	13.61 (3.25)
Completeness (%)	100 (100)	98.8 (98.0)	98.3 (98.3)
Redundancy	4.8 (4.8)	4.8 (3.68)	4.0 (3.81)
Refinement			
Resolution (Å)	2.60	2.25	2.30
No. reflections	51,178	102,983	80,429
R <sub>work</sub> /R <sub>free</sub>	23.8/28.7	17.9/21.6	17.6/23.0
No. atoms			
Protein	9062	11964	11881
Heterogen	36	107	102
Water	91	469	456
B-wilson (Å <sup>2</sup> )	41.8	47.21	40.0
R.m.s. deviations			
Bond lengths (Å)	0.007	0.010	0.009
Bond angles (°)	1.038	1.215	1.191

Data were collected from a single crystal. Values in parentheses are for the highest resolution shell.

relay and the SH1 helix; shown in Figure 2A) and the beta sheet of the converter have interactions and orientations that are conserved in all of the PPS structures, variability in the relative position of the converter helices demonstrates that the P-fold conformation of the converter is not rigid and is in fact a compliant structure (Figure 2C). The second helix which is parallel to the surface of the beta sheet is found translated by more than 3.2 Å in these different PPS structures. Large variation also occurs for the loop that precedes and the proximal part of insert-2 which interacts with it. In contrast, the last helix of the converter adopts a conserved position in these structures allowing conserved interactions with the N-terminal subdomain of the motor to be maintained (Figure 2B and Table S1). The variability in the P-fold of the converter indicates that the insert-2 orientation that defines the orientation of the lever arm of myosin VI differs by at least 18° in these structures (Figure 2D). This is likely an under estimate of the possible variability in lever arm orientation in a functioning myosin VI molecule, since strain would further perturb the lever arm position given the compliance of this region.

In contrast, comparison of the rigor and post-rigor state structures, which both have converters in the R-fold conformation

(Ménétrety et al., 2005, 2008), demonstrates that the proximal part of the insert-2 helix (residues 774-787) interacts with both the converter in this conformation as well as with the insert-2/CaM module. These multiple interactions should make the converter/insert2/CaM module much stiffer when the converter adopts the R-fold compared to when it adopts a P-fold, and thus greatly constrain the lever arm position (Figure S3). These structures further reveal that upon converter rearrangement from P-fold to R-fold, the proximal part of insert-2 undergoes a 90° rotation, but maintains interactions with the converter through hydrophobic interactions (Ménétrety et al., 2007). However, these interactions do not restrict the insert 2 proximal helix to a precise position in the P-fold conformation (Figure 2D).


While steric hindrance with the N-terminal subdomain prevents the converter from adopting the P-fold when the motor domain is found in the rigor or in the post-rigor state (Figure S4D), the two converter conformations are theoretically compatible with the motor domain when it adopts the pre-powerstroke state (Ménétrety et al., 2007). However, only the P-fold is populated in solution for myosin VI in the PPS state and all of the six PPS crystal structures we obtained adopt a P-fold. There are important interactions between the N-terminal subdomain of the motor, the end of the SH1 helix and the last helix of the converter that are conserved in all these PPS structures. These conserved interactions stabilize the P-fold of the converter (Figure S4 and Table S1). A detailed description of these interactions is presented in the Supplemental Information.

Altogether, these observations reveal that the atypical conformation of the converter first described in the MD structure (Ménétrety et al., 2007) is the sole conformation adopted in the PPS state and was not simply trapped in the original structure by the crystal packing or the lack of the myosin VI lever arm. Furthermore, these structures highlight that there is considerable compliance within the PPS conformation of the converter that will allow variability in lever arm positioning. Directed stress that would be generated between the distal part of the lever arm and the motor domain when an unbound lead head of a dimer is docking on actin during processive stepping can thus be partially accommodated by compliance associated with changes in the position of the converter helices.

#### Lever Arm Orientation and Stroke Size

The MD<sup>ins2-GFP</sup> structure reveals that the lever arm orientation in the pre-powerstroke state is directed toward the barbed-end (plus-end) of the actin filament as previously modeled (Ménétrety et al., 2007) (Figure 3A). The axis defined by the two positions observed for the pre- and post-powerstroke states of myosin VI is very similar to that of the actin filament axis. This PPS structure and the nucleotide-free (rigor like) structure of myosin VI (Ménétrety et al., 2005) thus could allow a 135° swing of the lever arm, parallel to the actin filament, during the powerstroke in the absence of strain. As indicated in Figure 3B, this degree of lever arm rotation predicts the measured stroke sizes seen with different myosin VI length lever arms (Bryant et al., 2007; Rock et al., 2005; Sivaramakrishnan et al., 2008).

A model of the pre-powerstroke state in which the converter would adopt a R-fold can easily be constructed by superimposition of the beta sheet of the converter. In this case, the



**Figure 3. The Lever Arm Orientation in the Pre-Powerstroke State**

(A and B) Superposition on the lower 50 kDa subdomain of the MD<sup>ins2-GFP</sup> pre-powerstroke (this study; blue) and the MD<sup>ins2</sup> rigor-like (2BKH, (Ménétreay et al., 2005); red) structures is shown with two different orientations (A) and (B). The calmodulin molecules were omitted for clarity while the IQ and PTD region were modeled based on the structure of the lever arm (3GN4). (A) Relative rotations of the lever arms are indicated. (B) Distances at the extremity of the lever arm prior and after the powerstroke (measured at the end of insert-2 (816), at the end of the IQ motif (835) and at the end of the folded triple helix bundle (912) are indicated. (C and D) In the same view as in (A) and (B), is a model of the pre-powerstroke state if the converter would adopt the rigor fold in its primed position (orange), as compared to the rigor state of myosin VI (2BKH, Ménétreay et al., [2005]; red) to evaluate the stroke size that would be produced if there was no rearrangement in the fold of the converter upon the powerstroke.

lever arm would be directed perpendicularly compared to the actin filament axis (Figures 3C and 3D). The predicted stroke would then be of ~3.1 nm for a lever arm consisting of both the insert-2 CaM and IQ CaM, and ~6.6 nm with the addition of the folded 3-helix bundle. No movement would be predicted for a lever arm of only the insert-2 CaM. The stroke size measured for these various constructs are much larger than these predictions, but similar to the predictions based on a P-fold, consistent with the converter being solely in a P-fold conformation in the PPS at the beginning of the powerstroke. It is unlikely that any isomerization of the converter conformation occurs in the PPS state since this would manifest as a much wider distribution and a smaller average of stroke sizes. This is an important insight, because it implies that variability in the myosin VI step size is not due to an isomerization of the converter conformation occurring in the PPS state prior to the lead head of a dimer binding to the actin filament.


#### A Pliant Region within Insert-2 Further Increases the Compliance of the Myosin VI Lever Arm in the PPS State

The proximal part of the myosin VI lever arm, consisting of the converter and the beginning of the insert-2 helix, is followed by

a part of insert-2 to which a Ca<sup>2+</sup>-bound calmodulin is bound. In the PPS conformation, the MD<sup>ins2-GFP</sup> structure reveals that the converter makes no interactions with the calmodulin bound to the distal part of insert-2 (Figure 4A). This is in contrast to what has been described for the rigor-like and post-rigor states, in which strong interactions with the R-fold converter maintain a specific position for the insert-2/CaM (Ménétreay et al., 2005; Figures 4B and S3). For these interactions to occur, insert-2 must adopt a kink at position Trp787, at the junction between

the proximal helix that interacts with the converter and the distal helix that binds calmodulin. The converter and the insert-2/CaM form a rigid module in the rigor state that would minimize the compliance of this proximal region of the lever arm in the rigor state, which may be critical for stepping while bearing load.

In contrast, the absence of interactions between the converter and the insert-2 bound calmodulin in the pre-powerstroke state (Figure 4A) creates flexibility in the positioning of the lever arm by freeing a pliant region in insert-2. No interactions stabilize the insert-2 helix residues N785 to R792, allowing highly variable bending of the helix. This in turn allows multiple positions of the myosin VI lever arm (CaM-Ins2/CaM + IQ/CaM + 3-helix bundle) for a particular MD/converter pre-powerstroke state (Figure 4C). We also performed experiments to monitor the exchange of the calmodulin bound to insert-2, which would be greatly inhibited if interaction occur between this CaM and the converter. These experiments confirm that almost no exchange occurs in the rigor-like state of the motor while CaM exchange does occur in the pre-powerstroke state (see Table S2). This is consistent with the absence of interactions involving this calmodulin and the P-fold of the converter and thus the presence of a pliant region within insert-2 in the PPS state. Thus the exact



**Figure 4. The Converter-Insert2-CaM Structural Module**

(A–D) Structures of the converter-insert-2-CaM module for the pre-powerstroke (A) and the rigor-like (B) states are indicated with the same color code as in Figure 1. These show the different conformations of the converter (green), the different bends of the insert-2 helix and the different position of the N-lobe of the CaM relative to the converter. Note that the insert-2 helix is straight in the pre-powerstroke state (A) and no interactions between the converter and the N-terminal CaM takes place. Eight residues (N785 to R792, yellow) of the insert-2 helix are not involved in any interactions. (B) In contrast, the helix is kinked at residue Trp787 in the rigor state, allowing interactions between the converter and the N-lobe of CaM to take place. Some of the salt bridges between the converter and the calmodulin are also shown – see also Figure S3 for comparison with the post-rigor state. (C) Incorporating the observed bending in this pliant region allows modeling of lever arm positions (two extremes in light blue) that differ by as much as 60° without creating any steric hindrance. The resulting variability in the stroke size is indicated for a MD-Ins2-IQ-PTD fragment. (D) Using the pre-powerstroke (blue) positions of the lever from (C), the post-stroke state (green) is modeled using a converter that has rotated to adopt a position close to that found in MD<sup>ins2</sup> rigor-like (2BKH, Ménétrey et al., 2005), but in which the PPS fold was maintained. The results of bending of the pliant region are modeled in light green. Note that the converter rotation without a converter rearrangement allows the lever arm to stay close to its position in the pre-powerstroke state (very small stroke size), while a larger azimuthal component would occur for the stroke. However this azimuthal component could be greatly diminished by changes in the pliant region since a ~60° amplitude of movement (light green) is possible. This model shows that the lead head can undergo this converter rotation without much change in the lever arm orientation and position when the rear head imposes strain on the lead head (E).

orientation of the lever arm of the head that rebinds to actin can vary greatly compared to the rest of the motor domain due to large variations in this pliant region. As modeled below, this contributes even more variability to the myosin VI step size than does the variable positioning of the converter helices orientations that gives rise to the compliance within the P-fold of the converter described above.

### Compliance in the Lever Arm Greatly Differs in the Lead Head and the Rear Head

For a monomeric motor, the strokes measured are consistent with a large swing of the lever arm direction derived from a combination of a converter swing coupled to a change in the converter conformation (P-fold to R-fold). The converter swing itself is similar to that of plus-end motors, but insert-2 redirects the lever arm so that the rotation axis of the lever arm differs from that of plus-end motors and results in a large azimuthal component for the lever arm swing. The lever arm would rotate by  $\sim 70^\circ$  due solely to a converter swing in the P-fold conformation, to adopt an orientation that would be perpendicular from the actin axis (Figure 4E). A subsequent isomerization of the converter toward the R-fold would further rotate the lever arm by  $\sim 90^\circ$  in the same plane. Thus a large azimuthal torque develops during the stroke of myosin VI.

For a dimeric molecule, compliance in the myosin VI converter and at the junction with the lever arm could allow the lead head to attach strongly to actin without developing strain on the lever arm. The new structures presented herein highlight that dynamics in the converter as well as at the pliant region within insert-2 controls the relative position of the converter and the rest of the lever arm. Lever arm compliance is much greater in the pre-powerstroke state (P-fold) than in the rigor or post-rigor states, since these adopt a R-fold converter, which has interactions that markedly constrain the lever arm position. The dynamics in the myosin VI lever arm are thus quite different in the lead head of a stepping dimer (that rebinds to actin with a converter in the P-fold), as compared to that of the rear head (that is strongly bound to actin in a rigor state). In the rear head, the CaM/converter interactions limit the compliance in the lever arm and would enable it to bear load as well as to efficiently direct the lever arm (and thus the other head) toward the minus end of the actin filament.

### Modeling Step Sizes and Converter Conformations

To ascertain to what extent the compliance due to the pliant region of insert-2 and the variable positioning of the converter helices can explain the observed stepping behavior of a myosin VI dimer, we first modeled a myosin VI lead head docked on actin with the range of lever arm positions that would be specified by the variable lever arm positions found in our six PPS structures (Figure 5). For modeling of dimers with both heads docked on actin, we used the general model of a myosin VI dimer proposed by Mukherjee et al., (2009), which has a lever arm that is extended by an unfolded three-helix bundle and dimerizes using the sequence immediately following this region. (See Supplemental Information for details).

The compliance we observed in the lead head (PPS) of myosin VI allows converter rotation in the P-fold conformation to occur

with minimal change in the orientation of the distal part of the lever arm (Figure 4D). Further modeling reveals that the isomerization between the P- and R-fold of the lead head converter cannot occur while the rear head is strongly attached, since this would result in considerable torque and strain within the lever arm (Figure 5A). Modeling of the lead head undergoing 19–36 nm steps (Figure 5) reveals that the lever arm compliance we detect is sufficient to allow the converter to swing to the post-stroke position if it remains in the P-fold conformation. This explains why the motor domain of the lead head is not slowed in releasing  $P_i$  and ADP by the attached rear head. Instead, the lead head undergoes the necessary conformational rearrangements promoted by strong binding to actin and release its products as if it is not influenced by the bound rear head (Sweeney et al., 2007). This P-fold converter swing is also consistent with the fact that reverse strain has been shown to facilitate strong binding of a myosin VI head (Iwaki et al., 2009) since the center of mass of the converter is moved toward the plus-end of the filament in this transition (Figure 5A).

Note that the rear head prevents the lead head from undergoing the further  $\sim 90^\circ$  rotation that would occur upon transition to the R-fold of the converter after its swing (Figure 5A). Although the lead head is not prevented from releasing or rebinding MgADP with the rear head attached, it is prevented in binding ATP until the rear head detaches (Sweeney et al., 2007). A possible implication is that rapid binding of ATP can only occur if the converter assumes its R-fold conformation. How this converter isomerization is coupled to elements of the nucleotide-binding site is unclear. However these coordinated changes in the converter and nucleotide binding site must underlie the slow isomerization that was detected and required to allow ATP binding to the lead head following detachment of the rear head (Sweeney et al., 2007).

With modeling, we next investigated what structural states and properties of myosin VI account for the variable step sizes that have been observed for the processive dimer (see Figure 5B). We considered both the possibility that the unfolded 3-helix bundle is flexible and the possibility that it is somewhat rigid, since there is no detailed characterization of this region. For both possibilities we report the possible size steps that can be produced for a myosin VI dimer depending on the converter fold that the lead head adopts. Note that in comparing these values to published step sizes obtained for myosin VI, in many cases experiments have been done by placing a fluorescent probe on only one head of a myosin VI dimer in order to use the FIONA (Yildiz et al., 2004) technique to track its position with ultra-high resolution. This means that what is actually observed is movement achieved by two sequential steps, which is then reported as either the summation of these steps (i.e., two sequential steps of 30 nm would be reported as 60 nm) or as their average. We comment on this point when relevant below.

First we modeled the rear head in the rigor conformation with a  $\sim 18$  nm long lever arm (2.5 nm for the converter/insert-2 module, 3.5 nm for the IQ motif/CaM and  $\sim 12$  nm for the unfolded 3-helix bundle) assuming that they constitute a rather rigid lever arm. We consider this the most likely possibility, since replacement of the IQ-CaM and 3-helix bundle of myosin VI with the relatively stiff lever arm of myosin V does not alter the



observed step size distribution of the dimer (Park et al., 2007). Our modeling revealed that if the converter of the lead head adopts a P-fold conformation, as in all of our PPS structures, then three actin-binding sites can be easily reached. This would allow the swing of the converter to take place without much strain due the pliant region of insert-2. 30 nm (11 actin subunits) away from the rear head seems to be the most easily reached and would position the lever of the lead head parallel to the actin axis, but 36 nm (13 subunits) and 25 nm (9 subunits) would be allowed and have only a small azimuthal component (see Figure 5B). The actin-binding site 19 nm (7 subunits) from the rear head might also be reached if a large bend is applied in the pliant region of insert-2. This would orient the lead head lever arm perpendicularly to the actin filament axis. The actin binding site 13.5 nm (5 subunits) from the rear head cannot be reached since it would be too far from the end of the rigid lever arm of the rear head. Note that the even numbered actin binding sites are not accessible since they are found on the opposite side of the actin filament helix. (In many cases experiments with myosin VI dimers have been performed with a probe on one head only, and thus the observed steps are the summation of two successive stepping events. Since modeling suggests that the favored spacing between heads will be 25, 30 or 36 nm (see Figure 5B), the modeling accounts for the 50–72 nm steps that have been observed experimentally with probes on one of the two heads).

We also modeled the situation in which the distal part of the myosin VI lever arm, created by unfolding of the three-helix bundle, is compliant rather than rigid. The main impact of this would be to decrease the average step size by favoring attachment of the lead head to actin-binding sites 19, 25 or 30 nm from the rear head (see Supplemental Discussion and Figure S5A). As shown in Supplemental Discussion, note also that the formal possibility that the lead head would adopt a R-fold converter upon attachment must be excluded since the head could not release its products without creating a large torque on the lever arm (Figures S5B, S5C, and S5D).

### Generation of Very Small Steps

Very small steps (~5.5 nm) have also been measured for myosin VI and led to the suggestion that myosin VI has an inch worm-like component to its stepping behavior that alternates with more conventional hand-over-hand stepping (Nishikawa et al., 2010). (By inch worm mechanism, the authors suggest that the rear head sometimes moves to an actin binding site adjacent to the former lead head, much as an inch worm pulls the rear half of his body toward the front half.) Modeling indicates a simple explanation for this observed behavior. These small steps can occur if rebinding of the lead head takes place while the rear head converter is still in a P-fold conformation, and thus its lever arm is not directed toward the minus end. While this maintains

compliance and variability in the position of the rear head lever arm, the lever arm is nonetheless mainly oriented perpendicular to the actin filament axis by the overall converter position in the post-stroke position. Modeling shows that in this case, binding at 5.5 nm (a+2) from this attached rear head followed by a converter swing would be possible if the lead head would have its converter in the P-fold thus mainly oriented parallel to the actin filament axis (Figure 5C). Both heads would then likely swing together and end up with both converters in the R-fold conformation. This would explain why the next step following a 5.5 nm step has to be a large step, as has been experimentally observed (Nishikawa et al., 2010).

Note that if such small steps occur and one is tracking the movements of only one head of a dimer experimentally, the step size observed will depend on the distance between the heads prior to detachment to which 5.5 nm would be added. If we assume the prior distance to be 25–36 nm, which is the favored spacing (see Figure 5B), this modeling accounts for the 30–42 nm steps that have been observed. The fact that the frequency of small steps increases when the ADP concentration is raised (Nishikawa et al., 2010) indicates that the isomerization from P-fold to R-fold of the rear head may be slower if ADP is bound to the head, which is consistent with the converter rearrangement toward the R-fold being promoted by the rear head being free of nucleotide. [Note that, based on the kinetic measurements on myosin VI dimers, when the lead head is constrained by an attached rear head, ADP can freely dissociate and rebind to the lead head (Sweeney et al., 2007). Thus as ADP concentration increases, the lead head is increasingly likely to be an ADP-bound state when the rear head detaches. While this lowers the likelihood of a processive run on actin being terminated due to ATP binding causing detachment of the lead head prior to strong reattachment of the rear head (Sweeney et al., 2007), it also will increase the frequency of small steps.

### Predicted Lever Arm Angles during Stepping

From the modeling presented it is clear that compared to myosin V, the lever arms of myosin VI are oriented so that they stay closer to the actin filament, which had been suggested to allow myosin VI to tend to stay on the same actin track (Sweeney and Houdusse, 2010). When the lead head of myosin VI rebinds to actin with a P-fold with its preferred step size of 25–36 nm, the IQ/CaM module of the lever arm would tend to stay mostly parallel to the actin filament axis during the converter swing. This point has been addressed experimentally, with two different groups examining the distribution of the lever arm angles relative to the actin filament axis during processive stepping of myosin VI (Reifenberger et al., 2009, Sun et al., 2007). However, their conclusions differ in terms of whether the lever arm is always parallel to the actin filament axis or not (i.e., whether or not there

head are indicated with a MD<sup>ins2-IQ-PTD</sup> construct (with converter in the P-fold). Only the site at 5.5 nm would allow the converter rotation of the lead and a transition of both heads together toward the rigor state (red). The docking sites of the lead head in black or gray cannot lead to a productive step. (Right) Same model in a different orientation to highlight the azimuthal components of the various dockings.

(D) Different phases for the step of a dimeric myosin VI molecule: (1) Docking of the lead head; (2) Swing with a PPS converter of the lead head maintains the orientation of its lever arm without significant strain, allowing strong binding to actin and phosphate and MgADP release (MgADP can rebind, but MgATP cannot bind); (3) Detachment of the rear head by MgATP binding allows the converter fold transition and recoupling of the lever arm in the lead head to bias rebinding of the former rear head toward the minus end.

is a significant azimuthal component to the lever arm swing). From our analysis, when the most favored steps of 25–36 nm occur, they would not lead to a significant azimuthal component to the lever arm position, as reported by Reifengerger et al., (2009). (Note that in the experiments of Reifengerger et al. [2009] a probe on only one head of a dimer was observed, thus they were measuring the distance of two sequential steps. For two sequential steps of 25–36 nm, they would have observed steps of 50–72 nm, which is what they reported.) However, when the distance between the heads is 19 nm, then significant azimuthal angles would be predicted to be measurable for the lead head lever arm, since these distances correspond to binding to actin sites on the side of the filament. One could thus speculate that the differences in the data reported by Reifengerger et al. (2009) and Sun et al. (2007) might be accounted for by differing in vitro geometries if the assay of Sun et al. (2007) allowed myosin to spiral around the actin filament while the assay system of Reifengerger et al. (2009) did not (Figure 5B). In this case, the “short” steps (30–42 nm) observed by Reifengerger et al. (2009) are likely occurring when the attached rear head converter has not undergone the P-fold to R-fold transition, promoting redocking of the head in an orientation parallel to the actin filament axis at the site 5.5 nm from this uncoupled head (as discussed above). In the case of Sun et al. (2007), the small steps could in part correspond to stepping on the side of the actin filament with 19 nm spacing between heads and this stepping would also explain the frequent observation of azimuthal angles for the lever arm in this assay (Figure 5B).

#### State-Dependent Compliance in the Myosin VI Lever Arm Is Essential for Processive Movement in the Reverse Direction

Based on this modeling, the compliance in the converter and pliant region insert-2 found in our PPS structures can completely account for the variable step sizes of the myosin VI dimer by allowing different actin binding sites to be reached (Figure 5). However, this step size variability may simply be a consequence of the design that is necessary to allow the reattached lead head to bind strongly to actin without developing significant intramolecular strain so that the release of P<sub>i</sub> and ADP can occur even though the lead head is constrained by the rear-head (Sweeney et al., 2007). Progression through the strong-binding states of the motor (accompanying product release) requires MD rearrangements that must involve the rotation of the converter due to movements of the relay. If no compliance were found in the lead head, this would result in an azimuthal torque on the lever arm that would likely either inhibit strong binding of the lead head until the rear head detaches or lead to high degrees of intramolecular strain if both heads were attached strongly. While intramolecular strain can be used to gate plus-end directed myosin motors, such as myosin V (Rosenfeld and Sweeney 2004; Veigel et al. 2005) by greatly slowing ADP release from the lead head. We have argued that due to the repositioning of the lever arm in myosin VI, intramolecular strain would not slow ADP release from the lead head, and could in fact slow ADP release from the rear head. Thus, processive movement would necessitate a new type of gating via a design that allows stepping without the development of intramolecular strain as well

as a new means of preventing release of the lead head prior to release of the rear head (Sweeney et al., 2007; Sweeney and Houdusse, 2010). We have presented evidence that gating is accomplished in myosin VI by preventing ATP binding to the lead head until the rear head has detached (Sweeney et al., 2007).

#### Dimeric Myosin VI Does Not Generate Force Using a Swinging Lever Arm Mechanism

Based on our new structures and the prior structural knowledge of myosin VI, a mechanism emerges that involves the motor domain of myosin VI undergoing similar conformational changes as those proposed for plus-end motors, while the compliant properties of the P-fold converter and insert-2 elements are essential to decouple the rotation of the lever arm from that of the converter at the beginning of the myosin VI powerstroke. This ensures committed attachment of the lead head upon processive movement without the development of intramolecular strain in the lever arm, which allows the ATP hydrolysis products to be released from the lead head at the unstrained rate. Since the lever arm reorientation is not coupled to the swing of the converter, the converter swing does not impose a large azimuthal torque.

Thus these results lead us to propose that dimeric myosin VI does not strictly direct its lever arm upon force generation (as postulated by the swinging lever arm mechanism), but rather has an initial phase of force generation in which its lever arm is essentially uncoupled from motor domain rearrangements due to lever arm compliance, which is then followed by a second phase in which the lever arm is directed by further MD rearrangement that includes a recoupling of the lever arm upon strain release when the rear head detaches (see Figure 5D). Fluctuations in the position of the IQ/CaM module have been reported when myosin VI walks (Yildiz et al., 2004) and are consistent with this model since it would correspond to the variability in lever arm position of the lead head upon the converter swing while in the P-fold compensated by a pliant region. In this regard, the processive movement of myosin VI is similar in mechanism to that proposed for kinesin motors (Rice et al., 1999; Vale and Milligan, 2000). This kinesin mechanism involves uncoupling of the neck linker/lever arm to allow strong attachment of the lead head and recoupling of the linker/lever arm within the lead head upon dissociation of the rear head.

#### Summary

This study highlights the importance of the converter isomerization in myosin VI and the variable, state-dependent compliance within the converter subdomain and within the adjacent pliant region of insert-2. The puzzling stepping behavior of the myosin VI dimer can be easily explained by the variability in actin attachment created by these converter/lever arm properties. Compliance in the converter of a plus end motor such as myosin II is also likely to play a role for force generation of the motor under load, while lack of compliance in the lever arm and lack of global conformational changes in the converter would allow for more precise stepping behavior. Interestingly, mutations in the converter of cardiac myosin II that are linked to hypertrophic cardio-myopathies have been reported to influence force

generation (Seebohm et al., 2009). Sequence variations for the converter in different muscle isoforms of *Drosophila* myosin II are also likely to be a way of tuning the motor properties of these myosins (Miller et al., 2003). For myosin VI, it appears that a high degree of compliance is necessary to prevent the generation of intramolecular strain within the dimer in order to allow processive movement in the opposite direction. Analogous to conventional kinesin motors, the mechanism involves strong attachment of the lead head with a lever arm poorly coupled to its motor, which after rear head detachment is followed by a recoupling of the lever arm to bias the rebinding of the new lead head toward the minus end of the track.

### EXPERIMENTAL PROCEDURES

Details of the experimental procedures are indicated in [Supplemental Information](#). Briefly, A series of truncations of porcine myosin VI cDNA allowed to purify different constructs of myosin VI with C-terminal truncations corresponding to amino acids 816 fused to GFP (MD<sup>ins2-GFP</sup>), 839 (MD<sup>ins2-IQ</sup>) and 789 (MD and the packing mutant MD<sup>PM</sup>). Crystals were obtained using the hanging drop method and were cryo-cooled prior to data collection at either the European Synchrotron Radiation Facility beamlines or the SOLEIL Proxima-1 beamline. The data sets were integrated and scaled with XDS and XSCALE (Kabsch, 2010) or Mosflm and SCALA (Collaborative Computational Project, Number 4, 1994). Statistics on the data collection for these four new pre-powerstroke state myosin VI structures are indicated in [Table 1](#). The structures were solved by molecular replacement with the myosin VI MD pre-powerstroke model (PDB code 2V26, Ménétrey et al. [2007]) by using the program AmoRe (Navaza, 1994) or Phaser (McCoy et al., 2007). Model building was done as mentioned in [Supplemental Information](#). Refinement was performed with Coot (Emsley and Cowtan, 2004) and Refmac5 (for MD<sup>ins2-GFP</sup> and MD<sup>ins2-IQ</sup>) (Collaborative Computational Project, Number 4, 1994) or Phenix (for MD<sup>crystal form 2</sup> and MD<sup>PM</sup>) (Adams et al., 2010). Crystallographic statistics of the final models are summarized in [Table 1](#). All diagrams for the figures were computed using MOLSCRIPT/Raster3D (Kraulis, 1991; Merritt and Bacon, 1997) or pymol (DeLano, 2002).

### ACCESSION NUMBERS

Atomic coordinates and structure factors for the pre-powerstroke structures of the myosin VI constructs have been deposited in the Protein Data Bank under the accession numbers 4ANJ for the MD<sup>ins2-GFP</sup>, 4E7Z for the P2<sub>1</sub> crystal form of the MD, and 4E7S for the MD<sup>PM</sup> (packing mutant) construct, respectively.

### SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Results and Discussion, two tables, five figures, Supplemental Experimental Procedures, and Supplemental References and can be found with this article online at <http://dx.doi.org/10.1016/j.molcel.2012.07.034>.

### ACKNOWLEDGMENTS

We thank Pierre Legrand and Andy Thomson at the PROXIMA 1 beamline of the SOLEIL synchrotron for assistance with data collection. O.P. is the recipient of an ARC postdoctoral fellowship. M.M. is the recipient of an American Heart Association fellowship. Author contributions: A.H. and H.L.S. designed research; J.M., T.I., V.R., M.M., O.P., and X.L. performed research; V.R., J.P., and P.V. analyzed the SAXS data; A.H. performed the modeling; J.M. and A.H. made the figures; and A.H. and H.L.S. wrote the paper. This work was supported by grants from the NIH (DC009100 and GM094229) (H.L.S.), the CNRS (A.H.), the ARC subvention fixe 1067XA0830F, and the ANR BLAN 2010 1504 01 (A.H.).

Received: March 19, 2012

Revised: July 2, 2012

Accepted: July 19, 2012

Published online: August 30, 2012

### REFERENCES

- Adams, P.D., Afonine, P.V., Bunkóczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W., et al. (2010). PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr. D Biol. Crystallogr.* 66, 213–221.
- Bryant, Z., Altman, D., and Spudich, J.A. (2007). The power stroke of myosin VI and the basis of reverse directionality. *Proc. Natl. Acad. Sci. USA* 104, 772–777.
- Collaborative Computational Project, Number 4. (1994). The CCP4 suite: program for protein crystallography. *Acta Crystallographica D Biol. Crystallogr.* 50, 760–763.
- DeLano, W.L. (2002) PyMOL molecular graphics system. <http://www.pymol.org>.
- Emsley, P., and Cowtan, K. (2004). Coot: model-building tools for molecular graphics. *Acta Crystallogr. D Biol. Crystallogr.* 60, 2126–2132.
- Iwaki, M., Iwane, A.H., Shimokawa, T., Cooke, R., and Yanagida, T. (2009). Brownian search-and-catch mechanism for myosin-VI steps. *Nat Chem Biol.* 5, 403–405.
- Kabsch, W. (2010). XDS. *Acta Crystallogr. D Biol. Crystallogr.* 66, 125–132.
- Kraulis, P.J. (1991). MOLSCRIPT: a program to produce both detailed and schematic plots of protein structures. *J. Appl. Cryst.* 24, 946–950.
- McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C., and Read, R.J. (2007). Phaser crystallographic software. *J. Appl. Cryst.* 40, 658–674.
- Ménétrey, J., Bahloul, A., Wells, A.L., Yengo, C.M., Morris, C.A., Sweeney, H.L., and Houdusse, A. (2005). The structure of the myosin VI motor reveals the mechanism of directionality reversal. *Nature* 435, 779–785.
- Ménétrey, J., Llinas, P., Cicolari, J., Squires, G., Liu, X., Li, A., Sweeney, H.L., and Houdusse, A. (2008). The post-rigor structure of myosin VI and implications for the recovery stroke. *EMBO J.* 27, 244–252.
- Ménétrey, J., Llinas, P., Mukherjee, M., Sweeney, H.L., and Houdusse, A. (2007). The structural basis for the large powerstroke of myosin VI. *Cell* 131, 300–308.
- Merritt, E.A., and Bacon, D.J. (1997). Raster3D: photorealistic molecular graphics. *Methods Enzymol.* 277, 505–524.
- Miller, B.M., Nyitrai, M., Bernstein, S.I., and Geeves, M.A. (2003). Kinetic analysis of *Drosophila* muscle myosin isoforms suggests a novel mode of mechanochemical coupling. *J. Biol. Chem.* 278, 50293–50300.
- Mukherjee, M., Llinas, P., Kim, H.-J., Travaglia, M., Safer, D., Ménétrey, J., Franzini-Armstrong, C., Selvin, P.R., Houdusse, A., and Sweeney, H.L. (2009). Myosin VI dimerization triggers an unfolding of a three-helix bundle in order to extend its reach. *Mol. Cell* 35, 305–315.
- Navaza, J. (1994). AMoRe: an automated package for molecular replacement. *Acta Crystallogr. Section A* 50, 157–163.
- Nishikawa, S., Homma, K., Komori, Y., Iwaki, M., Wazawa, T., Hikikoshi Iwane, A., Saito, J., Ikebe, R., Katayama, E., Yanagida, T., and Ikebe, M. (2002). Class VI myosin moves processively along actin filaments backward with large steps. *Biochem. Biophys. Res. Commun.* 290, 311–317.
- Nishikawa, S., Arimoto, I., Ikezaki, K., Sugawa, M., Ueno, H., Komori, T., Iwane, A.H., and Yanagida, T. (2010). Switch between large hand-over-hand and small inchworm-like steps in myosin VI. *Cell* 142, 879–888.
- Odronitz, F., and Kollmar, M. (2007). Drawing the tree of eukaryotic life based on the analysis of 2,269 manually annotated myosins from 328 species. *Genome Biol.* 8, R196.
- Park, H., Li, A., Chen, L.Q., Houdusse, A., Selvin, P.R., and Sweeney, H.L. (2007). The unique insert at the end of the myosin VI motor is the sole determinant of directionality. *Proc. Natl. Acad. Sci. USA* 104, 778–783.

- Phichith, D., Travaglia, M., Yang, Z., Liu, X., Zong, A.B., Safer, D., and Sweeney, H.L. (2009). Cargo binding induces dimerization of myosin VI. *Proc. Natl. Acad. Sci. USA* *106*, 17320–17324.
- Reifenberger, J.G., Toprak, E., Kim, H., Safer, D., Sweeney, H.L., and Selvin, P.R. (2009). Myosin VI undergoes a 180 degrees power stroke implying an uncoupling of the front lever arm. *Proc. Natl. Acad. Sci. USA* *106*, 18255–18260.
- Rice, S., Lin, A.W., Safer, D., Hart, C.L., Naber, N., Carragher, B.O., Cain, S.M., Pechatnikova, E., Wilson-Kubalek, E.M., Whittaker, M., et al. (1999). A structural change in the kinesin motor protein that drives motility. *Nature* *402*, 778–784.
- Rock, R.S., Rice, S.E., Wells, A.L., Purcell, T.J., Spudich, J.A., and Sweeney, H.L. (2001). Myosin VI is a processive motor with a large step size. *Proc. Natl. Acad. Sci. USA* *98*, 13655–13659.
- Rock, R.S., Ramamurthy, B., Dunn, A.R., Beccafico, S., Rami, B.R., Morris, C., Spink, B.J., Franzini-Armstrong, C., Spudich, J.A., and Sweeney, H.L. (2005). A flexible domain is essential for the large step size and processivity of myosin VI. *Mol. Cell* *17*, 603–609.
- Rosenfeld, S.S., and Sweeney, H.L. (2004). A model of myosin V processivity. *J. Biol. Chem.* *279*, 40100–40111.
- Seebohm, B., Matinmehr, F., Köhler, J., Francino, A., Navarro-Lopéz, F., Perrot, A., Ozcelik, C., McKenna, W.J., Brenner, B., and Kraft, T. (2009). Cardiomyopathy mutations reveal variable region of myosin converter as major element of cross-bridge compliance. *Biophys. J.* *97*, 806–824.
- Sivaramakrishnan, S., Spink, B.J., Sim, A.Y., Doniach, S., and Spudich, J.A. (2008). Dynamic charge interactions create surprising rigidity in the ER/K alpha-helical protein motif. *Proc. Natl. Acad. Sci. USA* *105*, 13356–13361.
- Sun, Y., Schroeder, H.W., 3rd, Beausang, J.F., Homma, K., Ikebe, M., and Goldman, Y.E. (2007). Myosin VI walks “wiggly” on actin with large and variable tilting. *Mol. Cell* *28*, 954–964.
- Sweeney, H.L., and Houdusse, A. (2010). Myosin VI rewrites the rules for myosin motors. *Cell* *141*, 573–582.
- Sweeney, H.L., Park, H., Zong, A.B., Yang, Z., Selvin, P.R., and Rosenfeld, S.S. (2007). How myosin VI coordinates its heads during processive movement. *EMBO J.* *26*, 2682–2692.
- Vale, R.D., and Milligan, R.A. (2000). The way things move: looking under the hood of molecular motor proteins. *Science* *288*, 88–95.
- Veigel, C., Schmitz, S., Wang, F., and Sellers, J.R. (2005). Load-dependent kinetics of myosin-V can explain its high processivity. *Nat. Cell Biol.* *7*, 861–869.
- Yildiz, A., Park, H., Safer, D., Yang, Z., Chen, L.Q., Selvin, P.R., and Sweeney, H.L. (2004). Myosin VI steps via a hand-over-hand mechanism with its lever arm undergoing fluctuations when attached to actin. *J. Biol. Chem.* *279*, 37223–37226.