



**HAL**  
open science

## Pathogens Shape Sex Differences in Mammalian Aging

Morgane Tidière, Adèle Badruna, David Fouchet, Jean-Michel Gaillard,  
Jean-François Lemaître Lemaître, Dominique Pontier

► **To cite this version:**

Morgane Tidière, Adèle Badruna, David Fouchet, Jean-Michel Gaillard, Jean-François Lemaître Lemaître, et al.. Pathogens Shape Sex Differences in Mammalian Aging. Trends in Parasitology, 2020, 36 (8), pp.668-676. 10.1016/j.pt.2020.05.004 . hal-03060299

**HAL Id: hal-03060299**

**<https://hal.science/hal-03060299>**

Submitted on 18 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# 1 Pathogens shape sex differences in mammalian aging

2 Morgane Tidière<sup>1,2\*</sup>, Adèle Badruna<sup>1,2</sup>, David Fouchet<sup>1,2</sup>, Jean-Michel Gaillard<sup>1,2</sup>, Jean-  
3 François Lemaître<sup>1,2</sup>, and Dominique Pontier<sup>1,2</sup>

4 <sup>1</sup> Université de Lyon, F-69000, Lyon; Université Lyon 1; CNRS, UMR5558, Laboratoire de  
5 Biométrie et Biologie Evolutive, F-69622, Villeurbanne, France.

6 <sup>2</sup> LabEx ECOFECT, Université de Lyon, F-69000, Lyon, France.

7 \*Correspondence: [mtidiere@gmail.com](mailto:mtidiere@gmail.com) (M. Tidière)

8

9 KEYWORDS: pathogen exposure; immunosenescence; lifespan; life history traits; medicine

10

## 11 ABSTRACT

12 Understanding the origin of sex differences in lifespan and aging patterns remains a salient  
13 challenge in both biogerontology and evolutionary biology. Different factors have been  
14 studied but the potential influence of pathogens has never been investigated. Sex differences  
15 especially in hormones and resource allocation generate a differential response to pathogens  
16 and thereby shape sex differences in lifespan or aging. We provide an integrative framework  
17 linking host pathogenic environment with both sex-specific selections on immune  
18 performance and mortality trajectories. We propose future directions to fill existing  
19 knowledge gaps about mechanisms that link sex differences, not only to exposition and  
20 sensitivity to pathogens, but also to mortality patterns, whilst emphasizing the urgent need to  
21 consider the role of sex in medicine.

## 22 **Sex differences in mammalian lifespan**

23 In almost all human populations, and consistently throughout history, women live longer than  
24 men [1]. This pattern is pervasive in mammals [2], although the difference in **lifespan** (see  
25 Glossary) between sexes varies strikingly across species and populations facing contrasted  
26 environmental conditions [2,3]. Currently, a large body of research is devoted to the  
27 understanding of evolutionary, genetic, and physiological mechanisms that govern these sex  
28 differences in lifespan [4,5]. Differences in life-history strategies between sexes driven by  
29 **sexual selection** are among the most common explanations [6]. However, evidence that  
30 sexual selection mostly shapes sex differences in lifespan remains equivocal [2]. Here, we  
31 highlight that sexual selection, by selecting for traits and behaviors that increase male  
32 reproductive success, also leads males to be more exposed to **pathogens** than females.  
33 Moreover, earlier **aging** of the immune system can occur in males. This can lead to sex  
34 differences in aging, which can be exacerbated in pathogen-rich environments.

35 Up to now, the literature on the deleterious consequences of infectious diseases has focused  
36 on the deterioration of the immune system with increasing age (i.e. the **immunosenescence**  
37 process; Box 1). In humans and wild vertebrate populations, aging of the **adaptive immune**  
38 **system** may occur earlier in males [7,8]. These different patterns of immunosenescence could  
39 occur between sexes due to a variety of interactions between physiological factors and the  
40 local environmental context. For instance, more intense stimulation of immune defenses  
41 through a greater pathogen exposure could contribute to earlier aging of the male immune  
42 system, leading to an earlier onset of their **actuarial senescence**. In support, HIV-1-infected  
43 individuals experience immunologic changes similar to uninfected elderly persons as a result  
44 of the continuous stimulation of their immune system [9]. However, the contribution of  
45 immunosenescence to the observed sex differences in lifespan and aging within and across  
46 mammals is yet to be deciphered. We argue that sex-specific immune responses and sex

47 differences in energy allocation strategies driven by sexual selection, by leading to differences  
48 in both exposure and susceptibility to pathogens, should result in an accelerated  
49 immunosenescence in males through immune exhaustion. If this hypothesis is correct, sex  
50 differences in lifespan, through either direct (host mortality) or indirect (faster  
51 immunosenescence) effects, could be modulated according to the pathogenic environment of  
52 populations [2] (Figure 1) and have direct consequences in human as well as veterinarian  
53 medicine (Box 2). Although the sample size was limited, a preliminary analysis of currently  
54 available data supports this hypothesis with sex differences of lifespan increasing in favor of  
55 females when pathogen richness increases (Box 3).

## 56 **How biological differences between males and females contribute to disease** 57 **dimorphism?**

58 As a general rule, **prevalence**, **intensity**, and **pathogenesis** of infections caused by diverse  
59 pathogens (i.e. viruses, bacteria, parasites, and fungi) are higher in males than in females in  
60 humans, domesticated animals, companion animals, laboratory rodents, and numerous species  
61 in the wild (see review in [10]). For example, in adult mice (*Mus musculus*) males display  
62 more symptoms of sickness than females when they are exposed to bacteria that cause an  
63 illness with symptoms similar to the flu [11]. Women have more than 40% less HIV-1 RNA  
64 loads than men [12], while Italian men infected by the SARS-CoV-2 are approximately twice  
65 more likely to die than infected women [13]. Beyond obvious morphological, anatomical, and  
66 behavioral differences (e.g. aggressiveness) that might affect exposure to pathogens, many  
67 other biological differences may contribute to observed sex differences in the susceptibility or  
68 disease progression [14]. Yet these differences remain little studied and data on the  
69 underlying biological mechanisms are surprisingly scarce, while they lead to different  
70 responsiveness to treatment between sexes [14] (see Box 2).

71 *Sex, hormones and immunity*

72 It has become increasingly clear that males and females differ in immune responses [15]. As a  
73 general rule, females exhibit greater capability of producing antibodies than males [16]. They  
74 are less susceptible to infectious diseases [15] but can develop a stronger predisposition to  
75 autoimmune or inflammatory diseases than males [15]. However, the consequences in terms  
76 of sex differences in lifespan and aging are yet to be accurately quantified [17].

77 Differences in disease prevalence and expression between males and females are mainly  
78 attributed to sex steroids (e.g. estrogens, testosterone, progesterone), which, by binding to  
79 hormone receptors on the surface of immune cells, modulate the phenotype of immunological  
80 cells [7,15]. Most of them express multiple sex hormone receptors that drive sex-specific  
81 immune responses following antigen stimulation [7]. Males and females differ in steroid  
82 concentrations: males have a higher concentration of testosterone and females a greater  
83 concentration of estrogen and progesterone. Estrogen receptors are detected in immune cell  
84 populations, including lymphocytes, monocytes and macrophages [18], and have an immuno-  
85 enhancing effect as well as diverse protective effects [19]. On the contrary, progesterone and  
86 testosterone have mainly immunosuppressive effects [20]. Folstad and Karter [21] proposed  
87 that testosterone has, in fact, a double-sword effect that increases the probability of mating for  
88 a male while decreasing its ability to fight over pathogens. For instance, territorial male  
89 chamois (*Rupicapra rupicapra*) have almost six times more fecal androgen metabolites and  
90 are three times more parasitized during the rut period than non-territorial males [22]. Hence,  
91 we may expect that males will develop stronger disease symptoms resulting in a lower  
92 probability to survive and thereby increased between-sex differences of lifespan in pathogen-  
93 rich environments. However, inconsistencies among studies of vertebrate species (e.g. Cape  
94 ground squirrels, *Xerus inauris* [23]) suggest that testosterone is not all story.

95 *Immune system and reproduction*

96 Major physiological changes occur during reproductive seasons or cycles. Typically,  
97 testosterone production is increased in males when competing for fecund females, while  
98 pregnancy induces a decrease in estrogen and an increase in progesterone concentration to  
99 avoid the immunologic aggression of the fetus [24]. However, infectious diseases that are  
100 acquired during pregnancy or lactation are often associated with lower birth mass and reduced  
101 breast-milk in humans, suggesting that females reactivate the formally quiescent immune  
102 system [25] to combat infection at the expense of reproduction. For example, women who  
103 contract malaria during pregnancy have higher circulating levels of pro-inflammatory  
104 cytokines, which in turn are associated with lower birth mass [26]. This suggests that females  
105 should show a lower **lifetime reproductive success** in an environment with high pathogenetic  
106 load, but their lifespan should be much less affected than that of males.

107 **The evolutionary roots of sex differences in response to pathogens**

108 From an evolutionary viewpoint, the immune response is a key fitness-related trait, but the  
109 energy it requires should inevitably be traded for allocation to life-history traits governing  
110 other biological functions [27]. For instance, the growth and maintenance of secondary sexual  
111 traits impair male immune performance through a resource-based allocation trade-off [28],  
112 making them more sensitive to pathogens [29] (Figure 1). Many observations in rodents,  
113 birds, and insects document substantial energetic, reproductive, and survival costs of immune  
114 activation [30]. Hence, organismal responses to pathogens should involve sex differences  
115 given that the optimal solution to the trade-off between reproduction and survival differs  
116 between sexes, with females allocating more than males to their immune responses to not  
117 jeopardize their survival and increase thereby their lifetime reproductive success [27]. We can  
118 even expect that iteroparous females should rapidly redirect their energy to immune functions

119 at the expense of fetus/offspring survival when exposed to a pathogen during pregnancy. The  
120 optimal strategy should be radically different in semelparous females who only have one  
121 opportunity to reproduce. For example, parasite removal reduces reproducing female survival  
122 in the Taiwan field mouse (*Apodemus semotus*), possibly by allowing breeding females to  
123 increase maternal investment (i.e. allocation at the cost of their future survival [31]). By  
124 contrast, in males, immune performance is impaired by the growth and maintenance of  
125 secondary sexual traits, which makes them more sensitive to pathogens [28]. In mice,  
126 increased aggression is costly and is associated with reduced resistance to disease [32]. As a  
127 consequence, sex differences of lifespan should increase in favor of females when pathogen  
128 load increases (Box 3).

129 Finally, males and females correspond to markedly different environments for pathogens,  
130 which may shape their evolution. Hence, the observed sex-biased disease prevalence and/or  
131 severity might be the result of the parasite having adapted to grow in specific host sex [33].  
132 Úbeda and Jansen [34] formalized this idea and suggested that natural selection can act  
133 differently on pathogens in males and females depending on the transmission route of the  
134 pathogen. In Japan, where the transmission of HTLV-1 (Human T-cell Lymphotropic Virus  
135 Type 1) occurs through breast-feeding rather than through sexual transmission, the  
136 progression to Adult T-cell Leukemia is slower in women than men. Sex-specific adaptation  
137 of HTLV-1 to preserve women as a viral route could be a potential explanation for this  
138 puzzling observation [34].

139 **The need to consider sex-specific immune performance and pathogen exposure in**  
140 **evolutionary biology**

141 *Challenges within species*

142 The immune system includes many different immune cell types, each having its unique  
143 function, and collectively protecting the host against pathogens. The reliable measurement of  
144 multiple markers of both immunity (e.g. cellular components, T cell repertoire) and aging  
145 (e.g. epigenetic markers of biological age [35]) remains challenging. However, such metrics  
146 are required to allow a better understanding of the ecological (pathogen exposure) and  
147 evolutionary forces that shape the sex-specific decline of immune responses and between-sex  
148 differences in lifespan. Ideally, these measures should be made in wild populations, which  
149 display a large variation in the magnitude of sex differences in lifespan (see [2]) because of  
150 differential risk-taking behavior, food requirements, mortality due to direct sexual  
151 competition, or exposition to commensal and pathogenic organisms. Longitudinal studies are  
152 thus required to assess how immunosenescence patterns are shaped according to sex (see for  
153 example [36], and references in [8]) and also to identify mortality causes, which allows  
154 deciphering the different ways through which pathogens reduce individuals' lifespan (through  
155 early deaths from lethal diseases or through advanced immunosenescence due to immune  
156 exhaustion). This has yet rarely been investigated in wild populations (but see [37]).

157 Even though studies of rodents (reviewed in [38]) show that reaching this aim is possible in  
158 wild populations, it remains a complex task. Populations living in protected conditions (e.g.  
159 wild species in zoos, domestic animals and companion animals living with humans,  
160 laboratory animals) would offer excellent simplified systems to explore more deeply the  
161 deterioration of immune functions with increasing age and its role in between-sex mortality  
162 patterns (see [39] for an example in mice). This would allow an investigation into how the  
163 level of pathogen exposure as well as the virulence of pathogens to which hosts are exposed  
164 shape the immunosenescence profile and thus the mortality rate and lifespan of males and  
165 females.



166 Captive mammalian populations could also provide insightful information on the potential  
167 effect of chronic and putatively asymptomatic infections on the differential rates of  
168 immunosenescence between sexes, which is currently totally underestimated. Research on  
169 human health has provided important and somewhat unexpected results in this field. Notably,  
170 the chronic infection by the common CMV (cytomegalovirus) is involved in the remodeling  
171 of the immune system. Hence chronic exposition to any microbial agent (i.e. not only known  
172 pathogens) could be both implied in shaping sex-specific immunosenescence [40]. Only long-  
173 term monitoring of individuals from birth to death, in which molecular and cellular markers of  
174 adaptive and innate functions can be recorded repeatedly throughout their lifetime, and a  
175 reliable estimate of the infection date becomes available at the individual level, will allow a  
176 deeper understanding of immunosenescence according to age and sex concerning microbial  
177 agent exposure.

178 Additionally, the comparative analysis of populations within a single species occupying  
179 multiple habitat types can offer important insights into the intraspecific variation in  
180 immunosenescence and its consequences on life history traits such as lifespan in both sexes.  
181 Selection is expected to produce the immune response that maximizes individual fitness, in  
182 interaction with other selection pressures imposed by the environmental context (i.e. resource  
183 availability, weather conditions), the social and mating systems, and the pathogen exposure.  
184 Thus, the immune system of rodents from different populations and environments in the wild  
185 differs from that of laboratory rodents, the former being continuously exposed to commensal  
186 and pathogenic organisms (reviewed in [38]). It would be thus interesting to evaluate how  
187 such excessive energetic demands to activate the immune system compromise other fitness  
188 components of laboratory rodents according to their pathogenic environment, notably the  
189 differential actuarial senescence rates between males and females.

190 *Inter-specific comparisons*

191 Taking into account the evolutionary history of mammals, their pathogen species richness,  
192 which varies largely across species [41], may help explain between-sex differences in  
193 lifespan. In presence of numerous pathogens, males from species subjected to strong sexual  
194 selection should be more exposed to pathogens compared to females than males from species  
195 with a low sexual selection [42]. This could partly explain sex differences in  
196 immunosenescence (see for example [8]) and highlight how these differences are likely  
197 determined by fine scale interactions between sex-specific physiological pathways and the  
198 local environment in pathogens [2].

199 Variation in pathogen richness, infection rate, and pathogen load correlates with numerous  
200 morphological and ecological traits, mostly driven by sexual selection, which are also related  
201 to the lifespan variation in mammals [43]. Among these traits, variation in host body size  
202 [44], mating system and sexual size dimorphism [42], MHC allelic diversity [45],  
203 geographical range [46], social group size [47], population density [48], and phylogeny [6,41]  
204 have been identified as playing a role. However, most cross-species studies that investigate  
205 the interaction between pathogens and host ecological and life-history traits performed to date  
206 did not include between-sex differences in pathogens nor their consequences in terms of sex-  
207 specific lifespan (e.g. [49]). Although male-biased parasitism is positively correlated with the  
208 level of polygyny in mammals [42], the level of polygyny also correlates with a shorter  
209 lifespan and a more pronounced aging in males compared to females, at least across captive  
210 populations of large herbivores [6].

211 All these challenges will be made possible thanks to databases gathering longitudinal data that  
212 measure accurately sex differences in lifespan, which are becoming increasingly available  
213 (e.g. [2]). Besides, several databases recording the number of pathogen species found in  
214 mammals, such as the World Register of Marine Species [50], Pathogen-Host Interactions  
215 (Phi-Base) [51] or the Database of Bat-associated Viruses (DBatVir) [52] have been

216 developed. These databases provide a large amount of information about the diversity and the  
217 richness of parasites across mammalian species displaying contrasted life styles. Combining  
218 these databases should allow better understanding the role of pathogens in between-sex  
219 differences of lifespan and aging (e.g. in terms of both onset and rate of actuarial and/or  
220 **reproductive senescence** patterns) across mammalian species (see Box 3). In our view, this  
221 information should inform on how pathogens shape sex differences rather than how sex  
222 differences shape responses to pathogens because the magnitude of sex differences seems to  
223 be context-specific (see [2] for a study of mortality patterns). It might be thus difficult to  
224 envision mammals evolving differential responses to parasites mediated by sex differences if  
225 the magnitude of sex differences is so labile.

## 226 **Concluding remark**

227 Understanding the determinants of differences in male and female outcomes is becoming a  
228 crucial recognized challenge in infectious disease research in humans as well as in veterinary  
229 clinical research, such as identifying the true contribution of pathogens to overall mammalian  
230 mortality (see Outstanding Questions). Available information so far calls for an appropriate  
231 inclusion of sex differences over the whole human life span to anchor medicine in the real  
232 world (see Box 2). Longitudinal studies combining detailed immunological and non-  
233 immunological follow-ups for individuals of each sex should help to understand aging of the  
234 immune processes and how it differs between sexes and environments (see Outstanding  
235 Questions). By taking into account the evolutionary history of mammalian species, such a  
236 holistic approach could help explain between-sex differences in lifespan. For instance, studies  
237 should focus on consequences of resistance and tolerance of pathogens on lifespan across  
238 mammals (see Outstanding Questions). We encourage the scientific community to adopt this  
239 holistic approach, which is necessary to embrace all the methodological, theoretical and

240 conceptual challenges raised by pathogens regarding sex differences in lifespan. The  
241 increasing availability of required data offers great opportunities to formulate and test new  
242 hypotheses.

## 243 **Acknowledgments**

244 This work was supported by the LabEx ECOFECT (ANR-11-LABX-0048) of Université de  
245 Lyon. We thank three anonymous referees for their helpful comments.

## 246 **REFERENCES**

- 247 1 Zarulli, V. *et al.* (2018) Women live longer than men even during severe famines and  
248 epidemics. *Proc. Natl. Acad. Sci. U. S. A.* 115, E832–E840
- 249 2 Lemaître, J.-F. *et al.* (2020) Sex differences in longevity and aging rates across wild  
250 mammals. *Proc. Natl. Acad. Sci.* 117, 8546–8553
- 251 3 Tidière, M. *et al.* (2016) Comparative analyses of longevity and senescence reveal  
252 variable survival benefits of living in zoos across mammals. *Sci. Rep.* 6, 36361
- 253 4 Ortona, E. *et al.* (2019) Editorial: sex hormones and gender differences in immune  
254 responses. *Front. Immunol.* 10, 1076
- 255 5 Xirocostas, Z.A. *et al.* (2020) The sex with the reduced sex chromosome dies earlier: a  
256 comparison across the tree of life. *Biol. Lett.* 16, 20190867
- 257 6 Tidière, M. *et al.* (2015) Does sexual selection shape sex differences in longevity and  
258 senescence patterns across vertebrates? A review and new insights from captive ruminants.  
259 *Evolution* 69, 3123–3140
- 260 7 Gubbels Bupp, M.R. *et al.* (2018) The confluence of sex hormones and aging on  
261 immunity. *Front. Immunol.* 9, 1269
- 262 8 Peters, A. *et al.* (2019) Immunosenescence in wild animals: meta-analysis and  
263 outlook. *Ecol. Lett.* 22, 1709–1722
- 264 9 Appay, V. and Sauce, D. (2017) Assessing immune aging in HIV-infected patients.  
265 *Virulence* 8, 529–538
- 266 10 Zuk, M. and McKean, K.A. (1996) Sex differences in parasite infections: patterns and  
267 processes. *Int. J. Parasitol.* 26, 1009–1024
- 268 11 Cai, K.C. *et al.* (2016) Age and sex differences in immune response following LPS  
269 treatment in mice. *Brain. Behav. Immun.* 58, 327–337
- 270 12 Griesbeck, M. and Altfeld, M. (2015) Sex differences in the manifestations of HIV-1  
271 infection. In *Sex and Gender Differences in Infection and Treatments for Infectious Diseases*  
272 (Klein, S. L. and Roberts, C. W., eds), pp. 103–181, Springer International Publishing
- 273 13 Promislow, D.E.L. (2020) A geroscience perspective on COVID-19 mortality. *J.*  
274 *Gerontol. Ser. A* DOI: 10.1093/gerona/glaa094
- 275 14 Bongen, E. *et al.* (2019) Sex differences in the blood transcriptome identify robust  
276 changes in immune cell proportions with aging and influenza infection. *Cell Rep.* 29, 1961-  
277 1973.e4
- 278 15 Klein, S.L. and Flanagan, K.L. (2016) Sex differences in immune responses. *Nat. Rev.*

279 *Immunol.* 16, 626–638

280 16 Markle, J.G. and Fish, E.N. (2014) Sex matters in immunity. *Trends Immunol.* 35,

281 97–104

282 17 Ngo, S.T. *et al.* (2014) Gender differences in autoimmune disease. *Front.*

283 *Neuroendocrinol.* 35, 347–369

284 18 Kovats, S. (2015) Estrogen receptors regulate innate immune cells and signaling

285 pathways. *Cell. Immunol.* 294, 63–69

286 19 Brooks, R.C. and Garratt, M.G. (2017) Life history evolution, reproduction, and the

287 origins of sex-dependent aging and longevity. *Ann. N. Y. Acad. Sci.* 1389, 92–107

288 20 Taneja, V. (2018) Sex hormones determine immune response. *Front. Immunol.* 9,

289 1931

290 21 Folstad, I. and Karter, A.J. (1992) Parasites, bright males, and the immunocompetence

291 handicap. *Am. Nat.* 139, 603–622

292 22 Corlatti, L. *et al.* (2019) Parasitism and alternative reproductive tactics in Northern

293 chamois. *Ecol. Evol.* DOI: 10.1002/ece3.5427

294 23 O’Brien, K.A. *et al.* (2018) Trade-offs between immunity and testosterone in male

295 African ground squirrels. *J. Exp. Biol.* 221, jeb177683

296 24 Klein, S.L. (2004) Hormonal and immunological mechanisms mediating sex

297 differences in parasite infection. *Parasite Immunol.* 26, 247–264

298 25 Abrams, E.T. and Meshnick, S.R. (2009) Malaria during pregnancy in endemic areas:

299 a lens for examining maternal-fetal conflict. *Am. J. Hum. Biol.* 21, 643–650

300 26 Rogerson, S.J. *et al.* (2003) Placental tumor necrosis factor alpha but not gamma

301 interferon is associated with placental malaria and low birth weight in Malawian women.

302 *Infect. Immun.* 71, 267–270

303 27 Zuk, M. and Stoehr, A.M. (2002) Immune defense and host life history. *Am. Nat.* 160,

304 S9–S22

305 28 Sheldon, B.C. and Verhulst, S. (1996) Ecological immunology: costly parasite

306 defences and trade-offs in evolutionary ecology. *Trends Ecol. Evol.* 11, 317–321

307 29 Metcalf, C.J.E. and Graham, A.L. (2018) Schedule and magnitude of reproductive

308 investment under immune trade-offs explains sex differences in immunity. *Nat. Commun.* 9,

309 4391

310 30 Lochmiller, R.L. and Deerenberg, C. (2000) Trade-offs in evolutionary immunology:

311 just what is the cost of immunity? *Oikos* 88, 87–98

312 31 Lo, H.-Y. and Shaner, P.-J.L. (2015) Sex-specific effects of parasitism on survival and

313 reproduction of a rodent host in a subtropical montane region. *Oecologia* 177, 657–667

314 32 Barnard, C.J. *et al.* (1997) Modulation of behaviour and testosterone concentration in

315 immunodepressed male laboratory mice (*Mus musculus*). *Physiol. Behav.* 61, 907–917

316 33 Duneau, D. and Ebert, D. (2012) Host sexual dimorphism and parasite adaptation.

317 *PLOS Biol.* 10, e1001271

318 34 Úbeda, F. and Jansen, V.A.A. (2016) The evolution of sex-specific virulence in

319 infectious diseases. *Nat. Commun.* 7, 13849

320 35 Horvath, S. *et al.* (2016) An epigenetic clock analysis of race/ethnicity, sex, and

321 coronary heart disease. *Genome Biol.* 17, 171

322 36 Cheynel, L. *et al.* (2017) Immunosenescence patterns differ between populations but

323 not between sexes in a long-lived mammal. *Sci. Rep.* 7, 13700

324 37 Froy, H. *et al.* (2019) Senescence in immunity against helminth parasites predicts

325 adult mortality in a wild mammal. *Science* 365, 1296–1298

326 38 Viney, M. and Riley, E.M. (2017) The immunology of wild rodents: current status and

327 future prospects. *Front. Immunol.* 8, 1481

328 39 Lippens, C. *et al.* (2019) Early *Plasmodium* -induced inflammation does not accelerate

329 aging in mice. *Evol. Appl.* 12, 314–323

330 40 Jergović, M. *et al.* (2019) Impact of CMV upon immune aging: facts and fiction. *Med.*  
331 *Microbiol. Immunol. (Berl.)* 208, 263–269

332 41 Guy, C. *et al.* (2019) Phylogeny matters: revisiting ‘a comparison of bats and rodents  
333 as reservoirs of zoonotic viruses.’ *R. Soc. Open Sci.* 6, 181182

334 42 Moore, S.L. and Wilson, K. (2002) Parasites as a viability cost of sexual selection in  
335 natural populations of mammals. *Science* 297, 2015–2018

336 43 Morand, S. and Harvey, P.H. (2000) Mammalian metabolism, longevity and parasite  
337 species richness. *Proc. R. Soc. Lond. B Biol. Sci.* 267, 1999–2003

338 44 Preisser, W. (2019) Latitudinal gradients of parasite richness: a review and new  
339 insights from helminths of cricetid rodents. *Ecography* 42, 1315–1330

340 45 Zhang, L. *et al.* (2015) Major histocompatibility complex alleles associated with  
341 parasite susceptibility in wild giant pandas. *Heredity* 114, 85–93

342 46 Gay, N. *et al.* (2014) Parasite and viral species richness of Southeast Asian bats:  
343 Fragmentation of area distribution matters. *Int. J. Parasitol. Parasites Wildl.* 3, 161–170

344 47 Nunn, C.L. *et al.* (2015) Infectious disease and group size: more than just a numbers  
345 game. *Philos. Trans. R. Soc. B Biol. Sci.* 370, 20140111

346 48 Nunn, C.L. *et al.* (2003) Comparative tests of parasite species richness in primates.  
347 *Am. Nat.* 162, 597–614

348 49 Cooper, N. *et al.* (2012) Host longevity and parasite species richness in mammals.  
349 *PLoS ONE* 7, e42190

350 50 WoRMS World Register of Marine Species. .

351 51 Urban, M. *et al.* (2017) PHI-base: a new interface and further additions for the multi-  
352 species pathogen–host interactions database. *Nucleic Acids Res.* 45, D604–D610

353 52 Chen, L. *et al.* (2014) DBatVir: the database of bat-associated viruses. *Database* 2014,  
354 bau021–bau021

355 53 Pawelec, G. (2018) Age and immunity: What is “immunosenescence”? *Exp. Gerontol.*  
356 105, 4–9

357 54 Zeng, Y. *et al.* (2018) Sex differences in genetic associations with longevity. *JAMA*  
358 *Netw. Open* 1, e181670

359 55 Wald, C. and Wu, C. (2010) Of mice and women: the bias in animal models. *Science*  
360 327, 1571–1572

361 56 Klein, S.L. and Poland, G.A. (2013) Personalized vaccinology: one size and dose  
362 might not fit both sexes. *Vaccine* 31, 2599–2600

363 57 Klein, S.L. *et al.* (2015) Sex-based differences in immune function and responses to  
364 vaccination. *Trans. R. Soc. Trop. Med. Hyg.* 109, 9–15

365 58 Clayton, J.A. and Collins, F.S. (2014) NIH to balance sex in cell and animal studies.  
366 *Nature* 509, 282–283

367 59 Abolins, S. *et al.* (2017) The comparative immunology of wild and laboratory mice,  
368 *Mus musculus domesticus*. *Nat. Commun.* 8, 14811

369 60 Perlman, R.L. (2016) Mouse models of human disease: an evolutionary perspective.  
370 *Evol. Med. Public Health* DOI: 10.1093/emph/eow014

371 61 Shi, J. *et al.* (2020) Susceptibility of ferrets, cats, dogs, and other domesticated  
372 animals to SARS–coronavirus 2. *Science* DOI: 10.1126/science.abb7015

373

374 GLOSSARY

375 **Adaptive immune system:** antigen-specific immune responses characterized by  
376 immunological memory that makes future responses against a specific antigen more efficient.

377 **Aging:** decline in the age-specific contribution to fitness throughout life. As a consequence,  
378 the aging process is mostly studied in the two main fitness components: survival and  
379 reproduction. Survival ageing (also called **actuarial senescence**) corresponds to the decrease  
380 in survival rates with increasing age and reproductive ageing (also called **reproductive**  
381 **senescence**) corresponds to the decrease in reproductive rates with increasing age.

382 **Immunosenescence:** decline of immune parameters with increasing age, associated with  
383 detrimental clinical outcome (e.g. high risk of contracting and dying from infectious diseases).

384 **Intensity of infection:** pathogen load within an individual at a given time.

385 **Lifetime reproductive success:** number of offspring alive produced by an individual at some  
386 life stage (usually birth, weaning, or one year of age) during its entire lifetime.

387 **Lifespan:** time interval between the birth and the death of a given individual.

388 **Pathogens:** organisms that cause disease to the host. They include in particular ectoparasites,  
389 protozoan, helminths, virus, bacteria and fungi.

390 **Pathogenesis:** progression of the disease. It encompasses the complete sequence of events  
391 accompanying acute and persistent infections.

392 **Prevalence:** proportion of infected individuals within a population at a given time, which  
393 encompasses both old and new cases.

394 **Sexual selection:** evolutionary process arising either through mating choice (i.e. preference  
395 by one sex for certain characteristics (e.g. form, color, or behavior) in individuals of the other  
396 sex) or through intra-sex competition (i.e. competition among mature individuals of one sex to  
397 monopolize access to mate with individuals of the other sex). Both ways select for a large  
398 sexually size dimorphism and extravagant sexually secondary characters such as ornaments or  
399 weapons when sexual selection is strong.



400 **Box 1. Sex-specific immunosenescence**

401 A decline in immune defense with increasing age, defining the immunosenescence process,  
402 has been highlighted in numerous species across the tree of life [8]. However, the few results  
403 available so far do not support that a sex difference of immunosenescence pattern would  
404 cause the observed sex differences of adult mortality [8,36].

405 Progressive deterioration of the organism results from concomitant retention (or exacerbation)  
406 of innate immunity coupled with a dysregulation (or dysfunction) of adaptive immunity [53].  
407 For now, in all studies conducted so far in humans, a universal age-associated immune  
408 alteration is consistently observed. The numbers and proportions of naive peripheral blood  
409 CD8+ T cells are reduced as a consequence of the developmentally-programmed thymic  
410 involution [53]. Moreover, as the establishment of the innate immune response causes  
411 inflammation and ROS production that induce collateral tissue damage, we can hypothesize  
412 that the more an organism is subject to a repeated innate immune response (e.g. due to high  
413 exposure to pathogens), the more permanent the damage will be. This potentially accelerates  
414 an organism's deterioration, ultimately leading to its death. Similarly, the adaptive immune  
415 system is particularly affected by the effect of long-term exposure to a variety of antigenic  
416 stimuli. Accordingly, an adaptive immune system highly exposed to pathogens will quickly  
417 lose efficiency, leading it to exhaustion with dramatic consequences such as high mortality.

418 As increased solicitation of the immune system leads to an acceleration of its dysregulation  
419 and efficiency, we suggest that in presence of pathogens, males undergo stronger  
420 immunosenescence than females, because of their higher exposure and tolerance to infectious  
421 agents. This results in the observed higher mortality rate and shorter lifespan of males  
422 compared to females. For example, Zeng et al. [54] highlighted that two immune pathways,  
423 the cytokine IL-6 and TLR3 proinflammatory signaling pathways, are positively associated

424 with the lifespan of centenarian men but not women. This suggests that dysregulation of these  
425 proinflammatory pathways with age makes elderly men more susceptible to infectious  
426 pathogens than elderly women. However, while numerous researchers asked for more  
427 longitudinal studies to define more accurately immunosenescence profiles and identify the  
428 underlining mechanisms [53], sex remains overlooked in biological research despite its  
429 critical consequences in the veterinarian and human medicine.

430 **Box 2. The urgent need to consider sex in medicine**

431 Investigating the role of pathogens on sex differences in aging highlighted how much males  
432 and females differ regarding their immune system and their response to infectious diseases. In  
433 spite of this, physicians still tend to prescribe the same treatment to both male and female  
434 patients for a given diagnostic. One reason for this is that sex differences in immune functions  
435 are not well understood yet.

436 During the last thirty years, most biomedical research has routinely used only males in both  
437 cohort and animal model studies because the cyclic hormonal fluctuations of females  
438 introduce additional experimental variation [55]. This could explain the higher number of  
439 secondary effects observed in women than in men following the commercialization of a given  
440 drug [56]. Upon vaccination, women develop a higher antibody immune response, but also  
441 more frequent and severe adverse side effects than men [57]. The application of sex-specific  
442 medicine is thus urgently required [56]. The American National Institute of Health recently  
443 declared that clinical trials not taking sex-specific responses into account will no longer be  
444 funded [58].

445 A great deal of knowledge about sex differences in immune functions comes from laboratory  
446 animals, notably the mouse model, which have been used extensively to develop research and  
447 test therapies before they are used in humans. However, very little is known about how much  
448 information from inbred and laboratory-adapted mice can be extrapolated to mammalian  
449 immune responses in the wild [59]. First, the selection of laboratory mice has resulted in the  
450 alteration of life-history traits (such as reproduction or lifespan, see [60]) and immunological  
451 traits of individuals. Second, as argued above, animals' immune responses are just one  
452 dimension of a wider life-history strategy to maximize fitness within the constraints of the  
453 environmental context. Laboratory conditions strongly differ from the environment

454 individuals face in the wild, which can have serious consequences on the immune response  
455 they mount. Comparison between male and female immune functions in wild and laboratory  
456 animals is thus crucially needed to reveal both the relevance and limitations of laboratory  
457 animals as immunological models. Linking wild and laboratory animal immunology by using  
458 tools and concepts of immunology, but also of ecology and evolutionary biology, is badly  
459 needed. In that respect, companion animals, which live in the same environment as their  
460 owners and are exposed to similar pathogens [61], may potentially serve as bridges between  
461 laboratory and wild species. The affordability of new 'omic' approaches and the availability  
462 of new trusted biomarkers (e.g. antibodies, cytokines, cellular responses) and immunological  
463 reagents (e.g. monoclonal antibodies) will help to quantify male and female exposition to  
464 microbes and the dysregulation of immune parameters with increasing age in a wider range of  
465 mammalian species, aiding immunologists, ecologists and evolutionary biologists to work  
466 altogether.

467 **Box 3. Relationship between sex differences in mean adult lifespan and increased**  
468 **pathogen richness**

469 Using between-sex differences of mean adult lifespan in 13 mammalian species (8 carnivores  
470 and 5 primates, see more details in supplemental Table S1), a preliminary analysis (Figure I)  
471 reveals that between-sex differences of lifespan increases ( $0.335 \pm 0.139$ ,  $p = 0.03$ ) in favor of  
472 females when pathogen richness increases. The mean adult lifespan for the 13 species was  
473 obtained from [2]. Pathogen richness is defined as the cumulative number of pathogen species  
474 inventoried (arthropods, helminths, protozoans, virus, and bacteria) for each carnivore and  
475 primate species, males and females combined (extracted from literature and standardized for a  
476 given sampling effort). Linear models have been fitted with between-sex differences of mean  
477 adult lifespan as the response variable, and pathogen richness and ‘family’ (Carnivora or  
478 Primates as factors) as explanatory variables. For each species considered, the number of  
479 studies inventorying one or several pathogen species was recorded as an indication of the  
480 sampling effort exerted on the species and included in the analysis: pathogen richness values  
481 correspond to the residuals of the linear model with real pathogen richness in response  
482 variable and the sampling effort as an explanatory variable.

483  
484 **Figure I. Increase of sex differences of mean adult lifespan in favor of females when**  
485 **pathogen richness increases for 8 carnivore (full circle) and 5 primate (open circle)**  
486 **species.**

487

488 **Figure legend**

489 **Figure 1. Different selective strength between sexes driving the evolution of sex**  
490 **differences.** These differences include those in life-history traits (blue boxes), in physiology  
491 and the immune response to pathogens (yellow boxes), and in aging and lifespan (red boxes).  
492

