



HAL
open science

Asynchrony of actuarial and reproductive senescence: a lesson from an indeterminate grower

Hugo Cayuela, Jean-François Lemaître Lemaître, Lorenzo Rugiero, Massimo Capula, Luca Luiselli

► **To cite this version:**

Hugo Cayuela, Jean-François Lemaître Lemaître, Lorenzo Rugiero, Massimo Capula, Luca Luiselli. Asynchrony of actuarial and reproductive senescence: a lesson from an indeterminate grower. *Biological Journal of the Linnean Society*, 2020, 131 (3), pp.667-672. 10.1093/biolinnean/blaa127. hal-03060309

HAL Id: hal-03060309

<https://hal.science/hal-03060309v1>

Submitted on 14 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Asynchrony of actuarial and reproductive senescence: a lesson from an**
2 **indeterminate grower**

3 Hugo Cayuela¹, Jean-François Lemaître², Lorenzo Rugiero³, Massimo Capula⁴, Luca Luiselli^{3,5,6}

4

5 ¹Département de Biologie, Institut de Biologie Intégrative et des Systèmes (IBIS), Université
6 Laval, Pavillon Charles-Eugène-Marchand, Québec, QC G1V 0A6, Canada

7 ²Université Lyon 1, CNRS, UMR 5558, Laboratoire de Biométrie et Biologie Evolutive, F-
8 69622, Villeurbanne, France

9 ³Institute for Development, Ecology, Conservation and Cooperation, via G. Tomasi di
10 Lampedusa 33, I-00144 Rome, Italy

11 ⁴Museo Civico di Zoologia, Rome, Italy

12 ⁵Department of Applied and Environmental Biology, Rivers State University of Science and
13 Technology, P.M.B. 5080 Nkpolu, Port Harcourt, Rivers State, Nigeria

14 ⁶University of Lomé, Faculty of Sciences, Department of Zoology, BP: 6057 Lome Togo

15

16 Running head: **Senescence in an indeterminate grower**

17

18 **Abstract**

19

20 Aging evolutionary theories predict that patterns of actuarial and reproductive senescence should
21 be aligned, with a common onset of senescence set at the age of first reproduction. However, a few
22 empirical studies reported asynchrony between actuarial and reproductive senescence. This
23 asynchrony is expected to be particularly pronounced in organisms with indeterminate growth. Yet,
24 this process is still poorly documented due to the lack of long-term demographic data on known-
25 aged individuals. We investigated the asynchrony of actuarial and reproductive senescence in the

42 **Introduction**

43

44 Although a few studies have challenged the idea of an inevitable senescence in nature (e.g. Martinez
45 1998, Cayuela et al. 2020), it is now broadly recognized that senescence, the decline in the
46 contribution to fitness with increasing age, is a widespread phenomenon across the animal kingdom
47 (Nussey et al. 2013; Gaillard & Lemaître 2020). Senescence can be due to an age-specific decrease
48 in survival probabilities (i.e., actuarial senescence) and/or reproductive performance (i.e.,
49 reproductive senescence), two processes usually anchored in the same theoretical framework
50 (Gaillard & Lemaître 2017). Evolutionary theories predict that patterns of actuarial and
51 reproductive senescence should be aligned (Williams 1957, Maynard-Smith 1962), with a common
52 onset of senescence at the age of first reproduction (Williams 1957). However, few recent studies,
53 performed in both laboratory and free-ranging conditions, reported asynchrony between actuarial
54 and reproductive senescence patterns (reviewed in Gaillard & Lemaître 2017). The factors
55 modulating the degree of synchrony are still poorly understood.

56 The asynchrony between actuarial and reproductive senescence could be particularly
57 pronounced in organisms with indeterminate (i.e., continuous) growth (e.g., fishes, amphibians,
58 and reptiles) where survival and reproductive traits (e.g., fecundity, offspring survival) are often
59 positively correlated with maternal body size, and thus indirectly determined by female age
60 (Hoekstra et al. 2020). This asynchrony could result from non-linear relationships between fitness
61 components and age. The fecundity of females with indeterminate growth recurrently increases
62 with body size and age in a linear way (e.g., in reptiles, Jenkins et al. 2009, Setser et al. 2010, Rose
63 et al. 2018), whereas the strength and the form of the relationship between mortality and age after
64 sexual maturity can be highly variable among species (Colchero et al. 2019, Cayuela et al. 2019a)
65 and populations (Cayuela et al. 2020).

66 To date, the asynchrony of actuarial and reproductive senescence in indeterminate growers
67 is still poorly documented, mainly due to the lack of long-term demographic data on known-aged

94 day lasting from 0800 to 1800), by two or three people who searched intensively for snakes in the
95 dilapidated building and neighbouring area of about 100× 80 m. Snakes were captured by hand,
96 and individually marked by ventral scale clipping for future identification. At their first capture,
97 female snakes were measured for snout-vent-length (SVL). We did not measure them on further
98 captures to minimize disturbance and hence risks of their departure from the nesting area, as
99 permanent emigration would bias capture-recapture inferences. We were able to assess the age of
100 16 females that were captured at juvenile stage over the capture-recapture survey. Detailed
101 information about the number of captures performed each year is given in **Table S1**.

102 Several females (N = 16) on their first capture were palpated in the abdomen to verify their
103 pregnancy status and to count the number of eggs (Filippi et al. 2007). We also captured 44
104 additional females in the surroundings of the study area to increase the number of observations of
105 egg clutch size (N = 60 in total).

106

107 *Reproductive senescence: assessing age-dependent variation in clutch size*

108

109 First, we verified that SVL was a good proxy of female age by quantifying the correlation between
110 age and SVL in a small set of females for which the age was known (N = 16). Then, we examined
111 how annual egg clutch size was associated with SVL in a larger sample of females (N = 60). In
112 both cases, we used regression models where SVL or egg clutch size were included as the response
113 variable and body size or age were included as explanatory variables. We compared null (i.e.
114 without covariate), linear, and polynomial models using their AIC. We graphically verified the fit
115 of the model by examining model residuals. The analyses were conducted in R program (R
116 Development Core Team 2018).

117

118 *Modeling age-dependent mortality using BaSTA*

119

146

147 *Reproductive senescence pattern*

148

149 Our results confirmed that female SVL linearly increased with age (**Fig.1A** and **Table S2** for model
150 selection procedure) and was thus a robust proxy of their age ($R^2 = 0.43$). Next, we showed that
151 annual egg clutch size increased with SVL (**Fig.1B**). Although the polynomial model was close
152 (less than 2 AIC points between the two models), the linear model was selected R^2 (0.74) (**Table**
153 **S3**). Overall, our analyses highlighted that female annual fecundity increased linearly with SVL
154 and age.

155

156 *Actuarial senescence pattern*

157

158 BaSTA models revealed that capture-recapture data were best described by a logistic function
159 without any refinement of the shape (**Table 1**). The cumulative probability of surviving was 0.75
160 until age six, 0.50 until age seven, and 0.25 until age eight (**Fig.1C**). Furthermore, the model
161 showed a strong positive effect of age on hazard rate (**Fig.1D**), which indicates an early and strong
162 actuarial senescence in *H. viridiflavus*.

163

164 **Discussion**

165

166 Using capture-recapture data collected over a 29-year period, we showed that *H. viridiflavus* female
167 did not experience any fecundity loss late in life, suggesting a negligible reproductive senescence.
168 In contrast, they suffered from an early and severe actuarial senescence. Taken together, those
169 results revealed a pronounced asynchrony in the senescence process of two main fitness
170 components.

197 general inferences about the ecological and biological factors modulating the asynchrony between
198 aging components. At the intraspecific level, abiotic factors such as temperature and habitat
199 predictability regulate lifespan and actuarial senescence (Sears 2005, Hjernquist et al. 2012, Zhang
200 et al. 2018, Cayuela et al. 2019b), as well as reproductive investment (Morrison et al. 2003, Cayuela
201 et al. 2016, Hughes et al. 2019), in ectotherm vertebrates with an indeterminate growth, which could
202 ultimately modify the asynchrony of senescence patterns. In our study system, the low abundance
203 of natural predators compared to other habitats occupied by *H. viridiflavus* (Philippi et al. 2007)
204 could also buffer actuarial senescence by reducing predation-induced mortality late in life, which
205 might lead to intraspecific variation in the degree of asynchrony between aging components.
206 Furthermore, covariation between mortality patterns and several phenotypic traits of indeterminate
207 growers could result in interspecific variation in the asynchrony of actuarial and reproductive
208 senescence. In particular, the magnitude of the asynchrony could be diminished in species with
209 size-dependent fecundity and negligible or even “negative” actuarial senescence (*sensu* Vaupel et
210 al. 2004, Jones & Vaupel 2017) allowed by high regeneration capacities (e.g., salamanders, Cayuela
211 et al. 2019a), or chemical (e.g., venomous snakes, Cayuela et al. 2019a, Colchero et al. 2019) and
212 armour protection (e.g., turtles, Warner et al. 2016). Overall, our study raises important questions
213 about the ecological and evolutionary mechanisms allowing the uncoupling of reproductive and
214 actuarial senescence processes.

215

216 **Acknowledgments**

217 We thank Christopher R. Friesen and two other anonymous reviewers for their helpful comments.

218

219 **References**

220

246 Du, W., Ji, X., & Shine, R. (2005). Does body-volume constrain reproductive output in lizards?
247 Biology Letters, 1, 98-100.

248 Filippi, E., Anibaldi, C., Capizzi, D., Ceccarelli, A., Capula, M. & Luiselli, L. (2007). Long-term
249 fidelity to communal oviposition sites in *Hierophis viridiflavus*. Herpetological Journal 17, 7-
250 13.

251 Fornasiero, S., Bonnet, X., Dendi, F., & Zuffi, M. A. (2016). Growth, longevity and age at maturity
252 in the European whip snakes, *Hierophis viridiflavus* and *H. carbonarius*. Acta Herpetologica,
253 11, 135-149.

254 Gaillard, J. M., & Lemaître, J. F. (2017). The Williams' legacy: A critical reappraisal of his nine
255 predictions about the evolution of senescence. Evolution, 71, 2768-2785.

256 Gaillard, J. M., & Lemaître, J. F. (2020). An integrative view of senescence in nature. Functional
257 Ecology, 34, 4-16.

258 Gompertz, B. (1825) On the nature of the function expressive of the law of human mortality, and
259 on a new mode of determining the value of life contingencies. Philosophical Transactions of the
260 Royal Society of London, 115, 513–583.

261 Hjernquist, M. B., Söderman, F., Jönsson, K. I., Herczeg, G., Laurila, A., & Merilä, J. (2012).
262 Seasonality determines patterns of growth and age structure over a geographic gradient in an
263 ectothermic vertebrate. Oecologia, 170, 641-649.

264 Hoekstra, L. A., Schwartz, T. S., Sparkman, A. M., Miller, D. A., & Bronikowski, A. M. (2020).
265 The untapped potential of reptile biodiversity for understanding how and why animals age.
266 Functional Ecology, 34, 38-54.

267 Hughes, D. F., Meshaka Jr, W. E., Lieb, C. S., & Pechmann, J. H. (2019). Latitudinal variation in
268 life history reveals a reproductive advantage in the Texas Horned Lizard (*Phrynosoma*
269 *cornutum*). Copeia, 107, 736-747.

295 Rose, J. P., Ersan, J. S., Wylie, G. D., Casazza, M. L., & Halstead, B. J. (2018). Reproductive
296 frequency and size-dependence of fecundity in the giant gartersnake (*Thamnophis gigas*).
297 Herpetological Conservation and Biology, 13, 80-90.

298 Sears, M. W. (2005). Geographic variation in the life history of the sagebrush lizard: the role of
299 thermal constraints on activity. *Oecologia*, 143, 25-36.

300 Setser, K., Mociño-Deloya, E., Pleguezuelos, J. M., Lazcano, D., & Kardon, A. (2010).
301 Reproductive ecology of female Mexican lance-headed rattlesnakes. *Journal of Zoology*, 281,
302 175-182.

303 Shine, R. (2003). Reproductive strategies in snakes. *Proceedings of the Royal Society of London*.
304 Series B: Biological Sciences, 270, 995-1004.

305 Siler, W., (1979). A competing-risk model for animal mortality. *Ecology*, 60, 750-757.

306 Sparkman, A. M., Arnold, S. J., & Bronikowski, A. M. (2007). An empirical test of evolutionary
307 theories for reproductive senescence and reproductive effort in the garter snake *Thamnophis*
308 *elegans*. *Proceedings of the Royal Society B: Biological Sciences*, 274, 943-950.


309 Spiegelhalter, D. J., Best, N. G., Carlin, B. P., & Van Der Linde, A. (2002). Bayesian measures of
310 model complexity and fit. *Journal of the Royal Statistical Society: Series B*, 64, 583-639.

311 Stanford, K. M., & King, R. B. (2004). Growth, survival, and reproduction in a northern Illinois
312 population of the plains gartersnake, *Thamnophis radix*. *Copeia*, 2004, 465-478.

313 Vaupel, J., Manton, K., & Stallard, E. (1979). The impact of heterogeneity in individual frailty on
314 the dynamics of mortality. *Demography*, 16, 439-454. URL <http://www.jstor.org/stable/206122>

315 Vaupel, J. W., Baudisch, A., Dölling, M., Roach, D. A., & Gampe, J. (2004). The case for negative
316 senescence. *Theoretical population biology*, 65, 339-351.

317 Warner, D. A., Miller, D. A., Bronikowski, A. M., & Janzen, F. J. (2016). Decades of field data
318 reveal that turtles senesce in the wild. *Proceedings of the National Academy of Sciences*, 113,
319 6502-6507.



329

330 **Fig.1.** Reproductive and actuarial senescence in females of *Hierophis viridiflavus*. (A) Linear
 331 relationship between snout-vent-length (SVL) and age. (B) Linear relationship between egg clutch
 332 size and SVL. (C) Relationship between survival and age described by a logistic function. (D)
 333 Relationship between hazard rate and age described by a logistic function.

334

335