

Anatoxin-a: overview on a harmful cyanobacterial neurotoxin from the environmental scale to the molecular target

Simon Colas, Benjamin Marie, Emilie Lance, Catherine Quiblier, Hélène Tricoire-Leignel, César Mattei

▶ To cite this version:

Simon Colas, Benjamin Marie, Emilie Lance, Catherine Quiblier, Hélène Tricoire-Leignel, et al.. Anatoxin-a: overview on a harmful cyanobacterial neurotoxin from the environmental scale to the molecular target. Environmental Research, 2021, 193, pp.110590. 10.1016/j.envres.2020.110590. hal-03065970

HAL Id: hal-03065970 https://hal.science/hal-03065970

Submitted on 15 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Anatoxin-a: overview on a harmful cyanobacterial neurotoxin from the environmental scale to the molecular target

Simon Colas^{1,2}, Benjamin Marie¹, Emilie Lance¹, Catherine Quiblier^{1,3}, Hélène Tricoire-Leignel^{2*}, César Mattei^{2*}

¹ UMR 7245 CNRS/MNHN "Molécules de Communication et Adaptations des Microorganismes", Muséum National d'Histoire Naturelle, Paris, France

² Mitochondrial and Cardiovascular Pathophysiology – MITOVASC, UMR CNRS
6015, INSERM U1083, UBL/Angers University, Angers, France

³ Université de Paris – Paris Diderot, 5 rue Thomas Mann, Paris, France

* Correspondence: <u>cesar.mattei@univ-angers.fr</u>; <u>helene.leignel@univ-angers.fr</u> – Mitochondrial and Cardiovascular Pathophysiology – MITOVASC, UMR CNRS 6015, INSERM U1083, UBL/Angers University, Angers, France tel: +33 244688274

Abstract

Anatoxin-a (ATX-a) is a neurotoxic alkaloid, produced by several freshwater planktonic and benthic cyanobacteria (CB). Such CB have posed human and animal health issues for several years, as this toxin is able to cause neurologic symptoms in humans following food poisoning and death in wild and domestic animals. Different episodes of animal intoxication in the wild have incriminated ATX-a, as confirmed by the presence of ATX-a-producing CB in the consumed water or biofilm and/or the observation of neurotoxic symptoms, which match experimental toxicity *in vivo*. Regarding toxicity parameters, toxicokinetics knowledge is currently incomplete and needs to be improved. The toxin is able to cross passively biological membranes and act rapidly on nicotinic receptors, its main molecular target. *In vivo* and *in vitro* acute effects of ATX-a have been studied and make possible to draw its mode of action, highlighting its deleterious effects on the nervous systems and its effectors, namely muscles, heart and vessels, and the respiratory apparatus. However, very little is known about its putative chronic toxicity. This review updates available data on ATX-a, from the ecodynamic of the toxin to its physiological and molecular targets.

Keywords: cyanobacteria · anatoxin-a · nicotinic acetylcholine receptor · nervous system · algal blooms

This work was supprted by ANSES project "Cyanariv" (2019-CRD-06; SJ N°364/19).

1. Introduction¹

In the context of global changes, with - among other consequences - the increase of N and P inputs due to agricultural activities, and the rise of temperatures and low-water periods, the phytobenthic and phytoplanktonic communities are largely dominated by cyanobacteria (CB) (O'Neil et al., 2012).

One of the deleterious consequences of the rapid proliferation of CB, often due to weak water circulation, is the production and secretion of cyanotoxins which develop a large range of toxicity mechanisms, the main ones being hepatotoxicity, dermatotoxicity and/or neurotoxicity caused by hepatotoxins (*eg* microcystins, nodularins), dermatotoxins (*eg* lyngbiatoxins) and/or neurotoxins (anatoxins, saxitoxins), respectively (Sivonen and Jones, 1999; Osswald et al., 2007; Aráoz et al., 2010; Huang and Zimba, 2019). However, the ecological role of these toxins, in terms of biological benefits for the CB, remains still poorly understood. Some of the possible roles explored in the literature are (i) to limit CB predation by other organisms such as zooplankton and nekton, and/or (ii) to serve as chemical mediators involved in cellular communication – allelopathy and chemotaxis – which can establish ecological relationships with other CB or other micro- or macro-

¹Abbreviations used: ACh Acetylcholine; ATX-a Anatoxin-a; CB Cyanobacteria; CNS Central nervous system; dhATX-a Dihydroanatoxin-a; dhHTX-a Dihydrohomoanatoxin-a; FW Fresh weight; HTX-a Homoanatoxin-a; IP Intraperitoneal; IV Intravenous; LD₅₀ Lethal dose 50%; LD₉₀ Lethal dose 90%; LOAEL Lowest observed adverse effect level; mAChR Muscarinic acetylcholine receptor; MC Microcystin; nAChR Nicotinic acetylcholine receptor; NOAEL No observable adverse effect level; PNS Peripheral nervous system; PKS Polyketide synthase; TE Thioesterase

organisms and reduce the development of competitors (Kaebernick and Neilan, 2001; Holland and Kinnear, 2013).

Figure 1 near here

In freshwater, the significant concentrations of cyanotoxins are consecutive to the proliferation of cyanotoxin-producing CB. Anatoxin-a (ATX-a) and related molecules are mainly secreted by freshwater benthic CB (Figure 1) (*e.g.*, *Phormidium*, *Kamptonema*, *Microcoleus*, *Oscillatoria*) in weak water currents (Edwards et al., 1992; Gugger et al., 2005; Wood et al., 2018), and planktonic CB (*e.g.*, *Anabaena*, *Aphanizomenon*, *Cuspidothrix*, *Cylindrospermopsis*, *Cylindrospermum*, *Dolichospermum*, *Planktothirx*, *Raphidiopsis*, *Tychonema*) in stagnant water (Sivonen et al., 1989; Viaggiu et al., 2004; Vehovszky et al., 2009; Hodoki et al., 2013; Shams et al., 2015; Trainer and Hardy, 2015; Ballot et al., 2018). This harmful cyanotoxin causes neurotoxic symptoms, sometimes lethal, in livestock, fish, pets and humans (Osswald et al., 2007). (Osswald et al., 2007).

2. Chemistry of ATX-a

ATX-a (2-acetyl-9-azabycyclo[4.2.1.]non-2-ene – MW 165 Da, $C_{10}H_{15}NO$) is a watersoluble secondary bicyclic amine, structurally related to alkaloids (Figure 2). It mainly exists in cationic form (pKa 9.6) and proved to be photosensitive at alkaline pH, where it degrades into non-toxic compounds (see section 3.2; Stevens and Krieger, 1991). The natural isomer is (+)-ATX-a, and the production of (-)-ATX-a by cyanobacteria has not been described to date. In this review, we specify - when available - which toxin isomer is used experimentally.

Figure 2 near here

ATX-a should not be confounded with ATX-a(s), which is also a neurotoxic cyanotoxin, but structurally related to organophosphates. ATX-a(s) will not be considered in this review. Several analogues of ATX-a have been detected in different field samples or in CB cultures (Skulberg et al., 1992; James et al., 1998). The most common are homoanatoxin-a (HTX-a), dihydroanatoxin-a (dhATX-a) and dihydrohomoanatoxin-a (dhHTX-a) (Figure 2). HTX-a (2-(propan-1-oxo-1-yl)-9-azabicyclo [4.2.1] non-2-ene – MW 179 Da, C₁₁H₁₈NO) is a methylated homologue of ATX-a with a comparable toxicity (Devlin et al., 1977; Wonnacott et al., 1992; Osswald, 2007; Aráoz et al., 2010), whereas dhATX is ten times less potent than ATX-a (Wonnacott et al., 1991). These four congeners can be found simultaneously in the environment in variable proportions. According to recent studies, dhATX-a seems to be the most common congener in New Zealand rivers followed by dhHTX-a (McAllister et al., 2016; Wood et al., 2018).

3. Biosynthesis and transformation

3.1. Biosynthesis

On the biosynthesis of ATX-a and HTX-a, a partial pathway using glutamate as substrate and involving polyketide synthases (PKS) was firstly described (Hemscheidt et al., 1995). As such, the production of ATX-a results from the condensation to three units of acetate on the (S)-1-pyrroline-5carboxylate, which derived from glutamate, followed by a decarboxylation. HTX-a is thought to result from the consecutive methylation of ATX-a. Thereafter, the genes encoding PKS enzymes involved in ATX-a biosynthesis have been identified (Cadel-Six et al., 2009). A nucleotidic sequence of Oscillatoria sp. strain PCC 6506 encoding a PKS has been found only in the genome of strains producing ATX-a or HTX-a, and surprisingly, within a cluster of genes encoding probably various enzymes involved in the biosynthesis of secondary ATX-a metabolites (Méjean et al., 2010). Bioinformatics analysis of the cluster showed that these genes encoded three proteins - namely AnaB (Prolyl-ACP oxidase), AnaC and AnaD - which are involved in the production of a substrate for three PKS enzymes (AnaE, AnaF, AnaG). Predicting the function of each of these gene products has allowed the description of a full biosynthetic pathway for ATX-a and HTX-a (Méjean et al., 2009, 2014). The cyclic amino acid proline is activated and attached to an acyl-carrying protein (AnaD) by an adenylation protein (AnaC). Then, the oxidation of this prolyl-ACP to 1-pyrroline-5-carboxyl-ACP (P5CACP) is performed by the AnaB oxidase. This newly formed imine is processed by a first PKS (AnaE), which adds a reduced acetate unit. The chain is then modified by the second PKS (AnaF), which probably adds an acetate unit and performs a Michael-type cyclization leading to the anatoxinic thioester linked to the ACP domain. The latest PKS (AnaG) is supposed to add an unreduced acetate unit. Subsequently, methylation is carried out to form the precursor of HTX-a. Without this methylation step, the latter PKS is the precursor of

 ATX-a. In fact, since the latter PKS does not have a thioesterase (TE) domain, TE type II (AnaA) performs hydrolysis to form carboxyATX-a or carboxy-homoATX-a, which lead to the corresponding compounds through spontaneous decarboxylation (Méjean et al., 2014).

3.2. Transformation

ATX-a and HTX-a are unstable in natural condition, where they are partially or totally degraded and converted into low- or non-toxic metabolites dhATX-a and epoxyATX-a (Stevens et Kriger, 1991; James et al., 1998). ATX-a can be degraded through photolysis and non-photochemical reactions, each of them affecting its chemical stability. It is also less bioavailable by dilution and adsorption (Stevens and Krieger, 1991). ATX-a degrades more quickly at higher temperatures (-76% after 1-hour exposure at 100°C, pH 7) and alkaline pH (-48% after 60-days exposure at pH 9.5°C, 20°C) (Stevens and Krieger, 1991; Kaminski et al., 2013). Based on analysis of reaction products by gas chromatography-chemical ionization mass spectrometry (GC/MS) from various experimental photolysis, several products were identified weighing 165 Da, corresponding to the MW of ATX-a. It was concluded that photolysis of ATX-a generated a complex mixture of rearrangement products involving different bridgehead positions (Stevens and Krieger, 1991; Kaminski et al., 2013). The breakdown products resulting from both photolytic and non-photolytic degradation are less toxic than ATX-a, some even being inactive (Stevens and Krieger, 1991). However, UV-B irradiation and high temperatures degraded ATX-a in the same way, leading to similar metabolites (Afzal et al., 2010; Kaminski et al., 2013). Its half-life in freshwater is estimated to be 5 days at neutral pH (Smith and

Sutton, 1993), but under conditions of sunlight intensity and at pH 9, ATX-a has a half-life of 1-2 h (Stevens and Krieger, 1991).

Micro-organisms are also able to degrade ATX-a, in different type of lake sediments which undergo cyanobacterial blooms, but also in lake sediments with no history of proliferation. In the Finnish eutrophic Lake Tuusulanjärvi the toxin concentration in water drops from 2.4 μ g.mL⁻¹ to 0 within 4 days (Rapala et al., 1994). It also has been shown that *Pseudomonas* sp. degrades ATX-a at a rate of 6-30 μ g.mL⁻¹ per 3 days (Kiviranta et al., 1991).

4. Factors influencing ATX-a production

Environmental conditions have a significant impact on cyanotoxin biosynthesis (Neilan et al., 2013; Boopathi and Ki, 2014). The amount of toxin can be up-regulated through an increase of the CB biomass, and/or an increase in toxin biosynthesis. Both mechanisms have been observed simultaneously.

Average concentrations of ATX-a measured range from 0.1 to 0.6 µg/L of water samples (Cerasino and Salmaso, 2012). These concentrations can somehow reach much higher levels such as 172,640 µg.L⁻¹ in 2008 in the Anderson lake (USA) (Hamel, 2009). In CB biofilms, ATX-a concentrations reach 1233 µg.g⁻¹ dw in the Tarn River (France) and 712 µg.g⁻¹ in the Oreti River (New Zealand) (McAllister et al., 2016; Echenique-Subiabre et al., 2018). In natural conditions, the cellular concentration of ATX-a was 2.5 pg.cell⁻¹ (Wood and Puddick, 2017). This cell content of ATX-a can reach maximal values and 1683 µg.g⁻¹ dw, and a production of 146 µg.L⁻¹ in an *Aphanizomenon issatschenkoi* strain under moderate levels of

nitrogen (Gagnon and Pick, 2012), but experimental conditions influence this level. A 23-day culture of the same CB ended up with a maximum of 1105 μ g.L⁻¹ production (Selwood et al., 2007). These values, among others, illustrate the heterogeneous nature of ATX-a production.

Early studies demonstrated that light, temperature and growth phase influenced ATXa production by *Anabaena flos-aquae* NRC-44-1 strains (Peary and Gorham, 1966). The highest toxicity was observed after 14 growing days at 22°C and 2000-7500 lux. Similar results confirmed the relationship between physicochemical parameters, growth phase and ATX-a production by cells (Rapala et al., 1993; Rapala and Sivonen, 1998). The amount of ATX-a produced by cell appears to be correlated with the exponential growth phase (Gagnon and Pick, 2012). In fact, the highest ATX-a intracellular levels are found during the first two weeks of CB growth before reaching a plateau and halving within a week (Rapala et al., 1993; Gallon et al., 1994; Harland et al., 2013).

In addition, the synthesis of ATX-a has been shown to be down-regulated at high temperatures, regardless of cyanobacterial cell growth, in *Anabaena* and *Aphanizomenon* strains (Rapala et al., 1993). Temperature has a significant impact on toxin concentrations. The highest intracellular ATX-a concentrations have been recorded between 19°C and 21°C (Rapala et al., 1993; Rapala and Sivonen, 1998). When temperatures rise too much, ATX-a production decreases to a minimum observed at 30°C (Rapala et al., 1993). N and P concentrations also influence the production of ATX-a. It is suggested that cellular production of ATX-a may be stimulated by moderate N deficiency (Gagnon and Pick, 2012). Besides, the available form of N may also play a role in ATX-a production. The toxin concentration is slightly higher when CB are in a medium with N₂ or NH₄-N compared to a medium

enriched with NO₃ or urea-nitrogen. However, no correlation between levels of toxin produced by CB and P concentrations has been observed. A deficiency in P seems only to limit the cyanobacterial growth (Rapala et al., 1993).

In addition to influencing toxin production, temperature and environmental composition also modulate the preferential biosynthesis of ATX-a or HTX-a (Aráoz et al., 2005). For instance, when grown on BG11 medium at 22°C without CO₂ enrichment, *Oscillatoria* PCC 6506 preferentially produces ATX-a. Conversely, when grown in BG1110 (with 10 nM NaHCO3) at 25°C with CO₂ enrichment, HTX-a is the dominant congener. Different strains of the same species may respond in multiple ways to the same environmental conditions. Two strains of *Oscillatoria* sp. (PCC 6506 and PCC 9029), although deriving from the same original isolate, exhibit different toxin production profiles. Strain PCC 6506, grown under standard conditions in non-gassed flasks, in both BG11 and 2N media, exclusively produced ATX-a. In contrast, PCC 9029 grown under the same conditions, produced mainly HTX-a (Aráoz et al., 2005).

Finally, biotic interactions can also influence the production of ATX-a, or at least the abundance of CB possessing the *ana* gene cluster. Microbial assemblages in mat samples with the ATX-a biosynthetic gene cluster consistently present a lower relative proportion of *Burkholderiales* (Betaproteobacteria) species than did mats without ATX-producing genes, suggesting that certain non-cyanobacterial organisms select CB species with genes for toxin production (Bouma-Gregson et al., 2019). In conclusion, abiotic and biotic parameters seem to play a crucial role in toxigenic CB biomass and toxin production.

5. Bioaccumulation and depuration

There is still very little work on the accumulation and excretion of ATX-a, so it does not provide a clear picture. Although ATX-a is found in the environment – water and biofilms samples –, naturally exposed organisms do not seem so far to prominently bioaccumulate the toxin (Osswald et al., 2007; Ferrão-Filho and Kozlowsky-Suzuki, 2011). However, it is important to point out that the hypothesis of ATX-a bioaccumulation is recent and has been little explored. Toxico-kinetics in fish appears to be fairly rapid. Following oral exposure of adult female medaka fish to a No Observable Adverse Effect Level (NOAEL = $6.67 \mu g.g^{-1}$ of (±)-ATX-a), the toxin could not be detected in the liver after 12h, and in the gut and muscles after 3 days (Colas et al., 2020). Besides, the depuration rate for the 12 first hours is 57, 100 and 90% in guts, livers and muscles, respectively, leading to the rapid excretion of ATX-a. This suggests that fish do not bio-accumulate ATX-a in these tissues.

In 2009 and 2010, environmental and biological samples (water, fish, plants) were investigated for the presence of ATX-a in various watering places in Nebraska and Washington (USA). Extracellular ATX-a was found but no trace detected in all fish samples studied, while other cyanotoxins – namely β -N-methylamino-L-alanine (BMAA) and 2,4-diaminobutyric acid dihydrochloride (DABA) – were present in the same analysed tissues (Al-Sammak et al., 2014; Hardy et al., 2015). When testing the ability of the rainbow trout (*Oncorhynchus mykiss*) to bioconcentrate ATX-a by balneation in water containing dissolved ATX-a up to 5 mg.L⁻¹, no significant difference between bioconcentration factors (BCF) of ATX-a in juvenile subjects exposed to different doses of the toxin for 96 h was observed: in fact, ATX-a accumulates in fish body with BCF ranging from 30 to 47 (Osswald et al., 2011).

The exposure of the mussel *Mytilus galloprovincialis* to cultures of ATX-a-producing CB for 15 days revealed very low assimilation rates, despite the great capacity of this bivalve to bioaccumulate different toxicants, such as heavy metals, drug residues or even microcystins (MCs) (Vasconcelos, 1995; Osswald et al., 2008). Moreover, only one day after being placed in a CB-free environment, mussels presented no detectable ATX-a in their tissues. This toxin, which was mainly found in the digestive tract, seems to be quickly eliminated within 24h.

Under natural conditions, ATX-a was detected for the first time in fish harvested in a Polish lake by Pawlik-Skowrońska et al. (2012). The highest amounts were mainly found in livers compared to muscles, omnivorous fish accumulating roughly the same amounts of ATX-a as carnivorous fish: omnivorous roach (*Rutilus rutilus*) exhibited concentrations of 11 to 18 µg/g FW in liver in autumn 2008. Moreover, the bream *Abramis brama* from a dam reservoir was found to accumulate 6.2 to 18.4 µg/g FW in liver in 2010 (Pawlik-Skowrońska et al., 2013). The contents of ATX-a detected in fish tissues seemed to increase with the occurrence of the toxin in the environment. Both intracellular and extracellular ATX-a were found and the intracellular form was predominant, i.e. 10 times in spring and 3 times in summer (Pawlik-Skowrońska et al., 2013).

ATX-a has also been detected in aquatic plants of the genus *Myriophyllum* (Al-Sammak et al., 2014). This contamination may be due to the endosymbiotic relationships of CB with some other higher plants, but may also occur via the presence of cyanotoxins in the aquatic environment and their absorption by vegetal organisms (Al-Sammak et al., 2014). Other aquatic organisms were challenged for

ATX-a bioaccumulation, such as benthic *Chironomus* which, fed on MC- and/or ATXa- producing CB, accumulated cyanotoxins (Toporowska et al., 2014).

6. Accidental exposure

6.1. Animal intoxications

Aquatic organisms can be naturally exposed to ATX-a by the oral, dermal or respiratory route (Kuiper-Goodman et al., 1999; Carmichael et al., 2001; Draisci et al., 2001). The ingestion of cyanotoxin-contaminated water is the cause of frequent cases of acute animal intoxication causing neurologic symptoms up to death (Quiblier et al. 2013, Wood et al, 2020). Water originating from a source with a proliferation of CB may contain toxins released during cell lysis (e.g. MCs) or excreted by CB (e.g. ATX-a) (Chorus and Bartram, 1999). Acute risks to animal health increase with the direct ingestion of CB biomass or biofilms, or of water containing large quantities of cyanotoxins. Alternatively, chronic intoxication may happen with the frequent ingestion of small doses of cyanotoxins over a long period of time through drinking water or food consumption (Svirčev et al., 2010). The irrefutable imputation of a poisoning to a single toxin often lacks evidence because the same species of CB can produce different kind of toxins. For instance, *Anabaena spiroids* produces ATX-a and also MCs; some *Kamptonema* species can produce both ATX-a and cylindrospermopsin (Méjean et al., 2009; Mazmouz et al., 2010).

Table 1 near here

Nevertheless, ATX-a has been implicated in numerous animal poisonings worldwide (Table 1). Notably, the death of dogs due to ATX-a poisoning has been reported in North America, in New Zealand, but also in France, Kenya and Germany among others (Edwards et al., 1992; James et al., 1997; Hamill et al., 2001; Gugger et al., 2005; Wood et al., 2007; Pushner et al., 2008, 2010; Faassen et al., 2012; Fastner et al., 2018). In addition, ATX-a and ATX-a-secreting CB have been found in tanks used for domestic water supply (Fawell et al, 1999).

6.2. Human intoxication

The presence of cyanotoxins in drinking water is not uncommon and has been mentioned in Asia, Europe and Oceania in the last ten years (Buratti et al., 2017). The consumption of cyanotoxin-containing water poses a health issue as it arises acutely in regions where the supply of drinking or irrigation water is made by storage – notably dams, reservoirs, wells, retention basins – likely to accumulate toxin-producing CB or cyanotoxins, and no or insufficient treatment of drinking water. Human intoxications might happen through the consumption of water, but other routes of exposure could be implicated.

With only one exception (Biré et al., 2020), there is no epidemiological data on human poisoning formally incriminating ATX-a, although in several cases it is suspected to be the cause of symptoms in humans, due to their neurotoxic type (Stewart et al., 2006; Weirich and Miller, 2014). Human accidental ingestion of water during aquatic activities may result in the ingestion of dissolved cyanotoxins or cyanotoxin-producing CB. A variety of symptoms have been described, in relation to cyanotoxin exposure through water drinking.

 The time to onset of symptoms - depending on the toxin involved, its dose and the route of exposure - are highly variable: a few minutes or even a few hours for skin symptoms and neurological disorders, several hours for hepatotoxins. The most commonly reported symptoms are gastrointestinal signs (diarrhoea, vomiting), fever and skin irritations. Headache, sore throat, and dry cough can be observed. Considering only the ingestion or inhalation of CB, hepatic toxicities and neurotoxicities are more frequently described (Wood, 2016; Svirčev et al., 2019).

Despite this risk associated with such intoxication, no recurrent cases of human death associated with swimming have been reported. The only exception occurred in July 2002, following a swim in water with a bloom of ATX-a producing CB, in Wisconsin in the United States: five teens developed gastrointestinal symptoms 48 hours after swimming, and one died from heart failure. Stool and blood tests revealed the presence of *Anabaena flos-aquae* CB. ATX-a has been incriminated, despite the delay between exposure and death. However, this case has not been the subject of a scientific report (Weirich and Miller, 2014), and the identification of ATX-a was subsequently questioned (Carmichael et al., 2004; James et al., 2005). To date, there is only one human poisoning case formally involving ATX-a: the consumption of ascidians of the *Microcosmus* genus in the south of France which intoxicated 26 people. The symptoms observed, mainly neurotoxic (blurred vision, ataxia, dizziness, asthenia, headache, muscle cramps, paresthesia) are entirely compatible with exposure to ATX-a. This work identified ATX-a at concentrations of 193.7 to 1240.2 µg .kg⁻¹ in the seafood consumed (Biré et al., 2020).

7. Toxicity

7.1. Acute toxicity

The symptoms observed experimentally after acute exposure to ATX-a are predominantly neurotoxic (Jakubowska and Szeląg-Wasielewska., 2015). There are no formal kinetic data for ATX-a in mammals. However, toxicity studies indicate that the toxin is rapidly absorbed after ingestion and widely distributed in highly vascularized organs such as liver, gut and muscles. Because it is a secondary amine, it rather crosses the blood-brain barrier in rats, which contributes to its acute toxicity, namely a decrease of locomotor activity (Rowell and Wonnacott, 1990; Stolerman et al., 1992).

Table 2 near here

ATX-a generally causes paralysis of intoxicated organisms. In rodents, ATX-a is lethal for doses of about 0.1 to 15 mg.kg⁻¹, depending on the route of exposure and the enantiomer tested (Table 2). In mice, ATX-a IP exposure causes after few minutes' hyperventilation, tremors, mild convulsions, urination, hypersalivation, hyperactivity, muscle contractions, then a decrease in locomotor activity, muscle paralysis and respiratory arrest preceding death. The acute toxicity shows rapid effects (2 to 6 min after administration) with muscular and respiratory paralysis. At a sub-lethal dose, there is a retrocession of all symptoms observed. However, and although animals can recover, changes in behaviour could have significant effects on the reproduction and interactions of intoxicated animals in their biotope (Osswald et al., 2007). There is a difference in toxicity between the two enantiomers and the route of exposure: (+)-ATX-a being significantly more toxic than (-)-ATX-a (Table 1), while

the LD₅₀ is about 60 times smaller through the IP route than through the oral route (Wonnacott and Gallagher, 2006; Metcalf and Codd, 2014).

Finally, a study carried out by administering CB extracts containing ATX-a shows a great difference in sensitivity of various organisms to the toxin: rodent mammals are much less sensitive to ATX-a than avian species (ratio oral LD₉₀ 2.1 - 5.1), and the goldfish displays a much higher sensitivity than avian and rodent species (ratio oral LD₉₀ 2.9 - 15.0) (Carmichael and Biggs, 1978). This heterogeneity could explain the fact that certain species may naturally accumulate ATX-a, without effect on their behaviour and fitness and constitute subsequent animal reservoirs of ATX-a, capable of transferring higher amount of this toxin through the food web (see Table 2).

7.2. Subchronic and chronic toxicity

Subchronic toxicity by ATX-a force-feeding or added to drinking water has been investigating (Osswald et al., 2007). Due to mortalities observed in groups with higher doses, the oral NOAEL has been estimated at 98 µg.kg⁻¹.d⁻¹ in mice, but the uncertainty on this value remains very high (Fawell et al., 1999). This study provides data on daily chronic oral intoxication with ATX-a in mice (0 to 2.46 mg of ATX-a.kg⁻¹ during 4 weeks), with no particular symptom in chronic intoxication been retrieved (Fawell et al., 1999). Subcutaneous injection of (+)-ATX-a (0.05 to 0.2 mg/kg) to rats produced tolerance with weekly administration in a food-reinforcement paradigm (Jarema et al., 2008).

Hence, no data are available on a putative subsequent genotoxicity or carcinogenesis of ATX-a. Reprotoxic effects in male mice have been observed with the racemic mixture of ATX-a. An intraperitoneal Lowest Observed Adverse Effect

Level (LOAEL) value of 50 µg.kg⁻¹.d⁻¹ was estimated in male mice after 7 days, on sperm count (Yavasoglu et al., 2008). Furthermore, maternal toxicity is reported at 200 µg.kg⁻¹ IP in mice but without generating malformation or mortality in foetuses. Conversely, ATX-a effects on foetuses are observed in the hamster: foetal stunting was observed, without skeletal or soft tissue malformations (Astrachan et al., 1980; Fawell et al., 1999; MacPhail et al., 2005; Rogers et al., 2005). In pregnant female mice, no sign of toxicity was observed in the offspring. The NOAEL (teratogenicity at 28 days) is established at 2.46 mg of ATX-a.kg⁻¹.d⁻¹ (Fawell et al., 1999), as no adverse effects were deduced from exposure of pregnant mices by gavage with this ATX-a dose (2.46 mg.kg⁻¹ bw per day), nor abnormal foetal development.

This toxin seems in fact to exert very little effects on the development of other vertebrate organisms. Embryos of zebrafish (*Danio rerio*) exposed to 400 μ g.L⁻¹ ATX-a exhibit variations in heart rate, but these variations are differently modulated depending on the stage of development, without being significantly different from the control group. At the pec-fin stage (55 h), corresponding to the pectoral fin blade development, heart rate was decreased to a mean of 9 % of control embryos, while at the protruding-mouth stage (80 h), it was increased to a mean of 12 %. All these observations were temporary, and the toxin did not seem to cause chronic effects on the development of embryos (Oberemm et al., 1999).

It can however be noticed that there is a difference between the effects on early developmental stages induced by the exposure to pure ATX-a or to cyanobacterial extracts of an ATX-a producing strain (Osswald et al., 2009). A challenge with pure ATX-a has almost no effect on the development of carp eggs (*Cyprinus carpio*). Only a decrease in larval length was observed at the highest concentration (640 µg.L⁻¹), while all the parameters studied (time to mortality, mortality rate, time to hatching,

hatching rate, skeletal malformations rate and larval length) were not affected by the increasing concentrations of CB extracts (Osswald et al., 2009).

8. Molecular target

8.1. Acetylcholine and cholinergic pathways

ATX-a acts as an agonist of nicotinic acetylcholine receptors expressed in cholinergic synapses, which are widely distributed in the nervous system of vertebrates (Thomas et al., 1993; Swanson et al., 1986; Bertrand and Terry, 2018). Acetylcholine (ACh) is the main neurotransmitter of the peripheral nervous system (PNS) and one of the most important in the central nervous system (CNS). In the CNS, ACh plays a crucial role in a very large number of functions, including cognition or central motor control (Bertrand and Terry 2018). The peripheral cholinergic synapses are found at neuromuscular junctions and in the autonomic nervous system and its effectors (heart, viscera, exocrine and endocrine glands). ACh ensures (i) the transmission of motor pathways which results in the contraction of the skeletal muscles, (ii) the transmission of an excitatory message in ganglia of the parasympathetic and orthosymapthic systems and (iii) the post-ganglionic transmission of the parasympathetic pathway. Two types of acetylcholine receptors mediate its action in the postsynaptic cell: ionotropic receptors (called nicotinic receptors, nAChRs) and metabotropic receptors (called muscarinic receptors, denoted mAChRs). After activating these receptors, ACh is hydrolysed by acetylcholinesterase, an enzyme present in the synaptic cleft. This reaction makes it possible to recycle choline in the presynaptic cell (Amenta and Tayebati, 2008). Activation of nAChRs by ACh opens a central pore permeable to cations (Na⁺, Ca²⁺ and K⁺), which promotes the excitation

of the postsynaptic cell. From a pharmacological point of view, all nAChRs can be exogenously activated by the plant alkaloid nicotine. Throughout the discovery of new ligands, several pharmacological profiles have been characterized so far, which can be linked to their structure. Belonging to the ligand-gated ion channel family, nAChRs are expressed as homo- or heteropentamers. To date, 17 subunits have been identified in vertebrates: α_{1-10} , β_{1-4} , γ , δ , and ϵ (Pedersen, 2019). They can organize in not less than 10 different subunit combinations, always including an α subunit which is crucial for the ligand binding. There are generally neuronal and muscle specific nAChR structures. The muscle nAChR stoechiometry is $\alpha_1(2)\beta\gamma\delta$ (in embryo) or $\alpha_1(2)\beta\epsilon\delta$ (in adult). It is activated by two ACh molecules which bind to two sites located at α_1/ϵ and α_1/δ interfaces. The activation of nAChRs ends up in depolarization and eventually drives the muscle fibre contraction (Martyn et al., 2009).

The neuronal nAChRs are heteropentamers with a stoichiometry $\alpha_x(2)\beta_y(3)$ or homopentamers such as $\alpha_7(5)$. Heteropentamers and homopentamers carry two and five ACh binding sites at the α/β and α/α interface respectively (Papke, 2014). The most widely expressed neuronal nAChRs are $\alpha_4\beta_2$ and α_7 in the brain and $\alpha_3\beta_4$ in the peripheral nervous system.

Neuronal and muscle nAChRs can also be expressed in non-neuronal and nonmuscle tissues, such as vascular endothelium, bones, adrenal medulla, immune cells... (Zoli, 2018). The large distribution of nAChRs in the body potentially explains the variety of symptoms associated with an inhibition or an over-activation of these receptors.

8.2. ATX-a activation of nAChRs

ATX-a is a potent agonist of nAChRs and to a lesser extent mAChRs (Aronstam and Wiktop, 1981). The agonist activity of ATX-a for mAChRs is weak and requires micromolar concentrations of ATX-a to compete with muscarinic agonists in binding assays (Aronstam and Witkop, 1981; Aracava et al., 1987), although, at neuromuscular junctions, (+)-ATX-a is a more powerful agonist than acetylcholine for stimulating muscle contraction, due to a greater affinity for nAChRs and a much a slower desensitization kinetics. On neuronal a7 receptors expressed in Xenopus oocytes, (+)-ATX-a exhibits an EC₅₀ of 0.58 μ M, much lower than acetylcholine EC₅₀ (320 μ M) and (-)-nicotine EC₅₀ (24 μ M), illustrating the greater potency of the toxin to interact with the receptor (Amar et al., 1993; Thomas et al., 1993). Competitive inhibition assays on membranes of Torpedo ocellata electric organs - containing muscle-type nAChRs - evidenced that ATX-a bind at sub-micromolar concentrations to the ACh site in a competitive way towards both ACh and tubocurarine, making the toxin an efficient agonist of the receptor (Aronstam and Witkop, 1981). Despite its capacity to cross the blood-brain barrier, ATX-a lethal effect can be primarily explained by its potent binding to muscle nAChRs, inducing a sustained depolarizing neuromuscular block (Carmichael et al., 1975).

Figure 3 near here

Indeed, in cholinergic synapses, ATX-a is not hydrolyzed by acetylcholinesterase (which naturally degrades ACh) and the nAChRs are therefore overstimulated (Figure 3). This excessive stimulation disrupts the response of the effector organ, which can be a nervous ganglion (autonomous nervous system), a neuron (CNS) or a muscle fibre (sustained contraction) (Spivak et al., 1980; Wonnacott et al., 1992;

muscle nAChR $\alpha_1(2)\beta_{\epsilon\delta}$. Through its binding to these nicotinic receptors, ATX-a causes the opening of their cationic pore, generating an influx of Na⁺/Ca²⁺ ions into the postsynaptic cell and an efflux of K⁺ ions, which leads to membrane depolarization. Prolonged exposure to ATX-a causes desensitization of nAChRs, and blockade of neuromuscular transmission (Spivak et al., 1980). HTX-a acts in a similar way as ATX-a: their affinities for nAChRs are comparable (Wonnacott et al., 1992). In addition to these blocking effects of neuromuscular transmission and taking into account the fact that nAChRs are present in autonomic nervous system, their activation at interneurons of parasympathetic ganglia induces the release of ACh which stimulates post-synaptic muscarinic receptors on organs. Consequently, ATXa affects the cardiovascular system: it causes a decrease in heart rate, an increase in blood pressure, and disruptions of gas exchanges (pO₂ and pCO₂), which leads to hypoxia and respiratory arrest, as well as acidosis observed in animal deaths (Adeymo and Sire'n, 1992). In the adrenergic system, ATX-a acts on post-ganglionic and adrenal nAChRs and stimulates the secretion of catecholamines (dopamine, adrenaline and noradrenaline) which display multiple effects in the CNS and PNS (cardiac function, metabolic functions). ATX-a crosses the blood-brain barrier and exerts its nAChR agonist activity (Figure 3), but less effectively than on neuromuscular junctions (Carmichael et al., 1977; Stolerman et al., 1992). The most deleterious effects of ATX-a are therefore due to excessive stimulation of the skeletal muscles of the respiratory tract, which can lead to respiratory arrest (Spivak et al., 1980; Aronstam and Witkop, 1981; Swanson et al., 1986).

Thomas et al., 1993). At the molecular level, ATX-a very effectively activates the

9. Regulatory thresholds

Numerous animal intoxications involving CB-producing ATX-a have been published and show that it is a harmful toxin well disseminated in the environment. The symptoms observed experimentally or on field emphasize its neurotoxic potency, but also its cardiovascular effects in acute exposure. Despite the risk, ATX-a is scarcely regulated. Guideline values have been proposed locally in some countries: 5 µg.g¹ in fish flesh in California; 40 and 100 μ g L⁻¹ in drinking water for livestock and dogs, respectively, in California; 3.7 µg.L⁻¹ in drinking water in Quebec (used as a temporary acceptable maximum dose); 6 µg.L⁻¹ and 2 µg.L⁻¹ in drinking water for ATX-a and HTX-a respectively, in New Zealand (used as the provisional maximum permissible dose); 20 μ g.L⁻¹ and 3 μ g.L⁻¹ in drinking water in Ohio and Oregon, respectively (Farrer et al., 2015). No threshold currently exists in Europe for ATX-a or HTX-a. Recently, ATX-a producers were also evidenced for the first time in Australia, highlighting the need for more frequent investigation of the distributions of toxinproducing CB in this region (John et al., 2019). Given its harmful character, its growing frequency in CB blooms and its involvement in animal poisoning, regulatory ATX-a thresholds aim at being more carefully investigated.

10.Concluding remarks

Studies on ATX-a and its structural analogues have resulted in a relatively satisfactory characterization of the ecological causes and circumstances under which this toxin is produced by CB, present in the water and the food web. These factors, in particular of anthropogenic origin, which favour CB proliferation, comprise eutrophication and perturbations of river hydric regime. The acute toxicity of ATX-a has been the subject of fairly old work, and its neurotoxic tropism has been evidenced, even if the toxicokinetic data are still quite fragmented. ATX-a, by its

chemical structure and its physicochemical properties, seems to be able to cross most biological membranes - including the gastro-intestinal tract, the blood-brain barrier and the placenta - and quickly reaches its molecular target. After causing deleterious effects in the nervous system, it is rapidly excreted, and no sequels are evidenced at sublethal doses. The toxin binds with high affinity to nAChR, which is considered so far as its sole target. This ionotropic receptor is expressed in the central and peripheral cholinergic pathways. When bound, ATX-a causes nAChR persistent opening, compromising the communication between neuronal and postsynaptic cells, at the level of the skeletal muscle, cardiovascular and CNS. The symptoms observed in animals on field or through experimental exposure can therefore be explained by this ATX-a/nAChR interaction. There is no human epidemiological data sufficiently reliable to link human intoxication to exposure to this toxin, with the exception of one case recently described in the south of France. We can anticipate that a human acute intoxication linked to ATX-a would be managed through symptomatic treatments. The chronic toxicity of ATX-a remains however not characterized, and a long-lasting exposure is still possible, due to the recurrent CB bloom occurrence. Finally, the differential sensitivity to ATX-a of various animal species still specifically deserves to be the subject of future studies. Indeed, some animals seem to resist to this toxin better than others and the toxicokinetic or toxicodynamic determinants of this "resistance" are not yet known. Regulatory thresholds for water contamination should be proposed to consider and prevent possible human intoxication.

Declaration of competing interest

The authors declare no conflicts of interest.

Funding

This work was supported by ANSES project "Cyanariv" (2019-CRD-06; SJ N°364/19).

References

Adeyemo, O.M., Sirén, A.L., 1992. Cardio-respiratory changes and mortality in the conscious rat induced by (+)- and (+/-)-anatoxin-a. Toxicon 30 (8), 899-905. https://doi.org/10.1016/0041-0101(92)90388-I

Afzal, A., Oppenländer, T., Bolton, J.R., and El-Din, M.G., 2010. Anatoxin-a degradation by Advanced Oxidation Processes: Vacuum-UV at 172 nm, photolysis using medium pressure UV and UV/H2O2. Water Research 44 (1), 278-286. https://doi.org/10.1016/j.watres.2009.09.021

Al-Sammak, M.A., Hoagland, K.D., Cassada, D., Snow, D.D., 2014. Co-occurrence of the cyanotoxins BMAA, DABA and anatoxin-a in Nebraska reservoirs, fish, and aquatic plants. Toxins 6 (2), 488-508. <u>https://doi.org/10.3390/toxins6020488</u>

Amar, M., Thomas, P., Johnson, C., Lunt, G.G., Wonnacott, S., 1993. Agonist pharmacology of the neuronal alpha 7 nicotinic receptor expressed in Xenopus oocytes. FEBS Lett. 327 (3), 284-288. <u>https://doi.org/10.1016/0014-5793(93)81005-k</u>

Amenta, F., Tayebati, S.K., 2008. Pathways of acetylcholine synthesis, transport and release as targets for treatment of adult-onset cognitive dysfunction. Curr. Med. Chem. 15(5): 488-498. https://doi.org/10.2174/092986708783503203

Aracava, Y., Deshpande, S.S., Swanson, K.L., Wonnacott, S., Lunt, G., Albuquerque, E.X., 1987. Nicotinic acetylcholine receptors in cultured neurons from the hippocampus and brain stem of the rat characterized by single channel recording. FEBS Lett. 222 (1), 63-70. https://doi.org/10.1016/0014-5793(87)80192-8

Aráoz, R., Nghiêm, H.O., Rippka, R., Palibroda, N., de Marsac, N.T., Herdman, M., 2005. Neurotoxins in axenic oscillatorian cyanobacteria: coexistence of anatoxin-a and homoanatoxin-a determined by ligand-binding assay and GC/MS. Microbiology 151(4), 1263-1273. <u>https://doi.org/10.1099/mic.0.27660-0</u>

Aráoz, R., Molgó, J., Tandeau de Marsac, N., 2010. Neurotoxic cyanobacterial toxins. Toxicon 56 (5), 813-828. <u>https://doi.org/10.1016/j.toxicon.2009.07.036</u>

Aronstam, R.S., Witkop, B., 1981. Anatoxin-a interactions with cholinergic synaptic molecules. Proc. Natl. Acad. Sci. (USA) 78 (7), 4639-4643. https://doi.org/10.1073/pnas.78.7.4639

Astrachan, N.B., Archer, B.G., Hilbelink D.R., 1980. Evaluation of the subacute toxicity and teratogenicity of anatoxin-a. Toxicon 18 (5-6), 684-688. https://doi.org/10.1016/0041-0101(80)90100-2

Ballot, A., Krienitz, L., Kotut, K., Wiegand, C., Metcalf, J.S., Codd, G.A., Pflugmacher, S., 2004. Cyanobacteria and cyanobacterial toxins in three alkaline Rift Valley lakes of Kenya—Lakes Bogoria, Nakuru and Elmenteita. J. Plank. Res. 26 (8), 925-935. https://doi.org/10.1093/plankt/fbh084 Ballot, A., Scherer, P.I., Wood, S.A., 2018. Variability in the anatoxin gene clusters of *Cuspidothrix issatschenkoi* from Germany, New Zealand, China and Japan. PLoS One 13(7), e0200774. https://doi.org/10.1371/journal.pone.0200774

Bertrand, D., Terry, A.V. Jr., 2018. The wonderland of neuronal nicotinic acetylcholine receptors. Biochem. Pharmacol. 151, 214-225. https://doi.org/10.1016/j.bcp.2017.12.008

Biré, R., Bertin, T., Dom, I., Hort, V., Schmitt, C., Diogène J., Lemée R., De Haro L., Nicolas, M., 2020. First evidence of the presence of anatoxin-A in sea figs associated with human food poisonings in France. Mar. Drugs 18 (6), E285. https://doi.org/10.3390/md18060285

Boisseleau, D., Peigner, P., Polato, T., 2018. Cas groupés d'intoxications de chiens par des cyanobactéries dans la Loire. Bulletin épidémiologique, santé animale et alimentation 82 (7). Available at: <u>https://be.anses.fr/sites/default/files/N-046_2018-01-</u> <u>2 Cyanobacteries-chiens_Boisseleau_MaqVF_0.pdf</u>, Accessed date: 28 July 2020.

Boopathi, T., Ki, J.S., 2014. Impact of environmental factors on the regulation of cyanotoxin production. Toxins 6 (7), 1951-1978. https://doi.org/10.3390/toxins6071951

Bouma-Gregson, K., Olm, M.R., Probst, A.J., Anantharaman, K., Power, M.E., Banfield, J.F., 2019. Impacts of microbial assemblage and environmental conditions on the distribution of anatoxin-a producing cyanobacteria within a river network. The ISME journal 13(6), 1618-1634. <u>https://doi.org/10.1038/s41396-019-0374-3</u>

Buratti, F.M., Manganelli, M., Vichi, S., Stefanelli, M., Scardala, S., Testai, E., Funari, E., 2017. Cyanotoxins: producing organisms, occurrence, toxicity, mechanism of

 action and human health toxicological risk evaluation. Arch. Toxicol. 91 (3), 1049-1130. <u>https://doi.org/10.1007/s00204-016-1913-6</u>

Cadel-Six, S., Iteman, I., Peyraud-Thomas, C., Mann, S., Ploux, O., Mejean, A., 2009. Identification of a polyketide synthase coding sequence specific for anatoxin-a-producing *Oscillatoria* cyanobacteria. Appl. Environ. Microbiol. 75(14), 4909-4912. https://doi.org/10.1128/AEM.02478-08

Carmichael, W.W., Biggs, D.F., Gorham, P.R., 1975. Toxicology and pharmacological action of anabaena flos-aquae toxin. Science 187: 542-544. https://doi.org/10.1126/science.803708

Carmichael, W.W., Gorham, P.R., Biggs, D.F., 1977. Two laboratory case studies on the oral toxicity to calves of the freshwater cyanophyte (blue-green alga) *Anabaena* flos-aquae NRC-44-1. Can. Vet. J. 18 (3), 71-75. PMID: 404019

Carmichael, W.W., Biggs, D.F., 1978. Muscle sensitivity differences in two avian species to anatoxin-a produced by the freshwater cyanophyte *Anabaena* flos-aquae NRC-44-1. Can. J. Zool. 56, 510-512. <u>https://doi.org/10.1139/z78-071</u>

Carmichael, W.W., Azevedo, S.M., An, J.S., Molica, R.J., Jochimsen, E.M., Lau, S., Rinehart, K.L., Shaw, G.R., Eaglesham, G.K., 2001. Human fatalities from cyanobacteria: chemical and biological evidence for cyanotoxins. Environ. Health Perspect. 109 (7), 663-668. <u>https://doi.org/10.1289/ehp.01109663</u>

Carmichael, W.W., Yuan, M., Friday, C.F., 2004. Human mortality from accidental ingestion of toxic cyanobacteria – a case re-examined. Sixth International Conference on Toxic Cyanobacteria, Bergen: pp 61-62.

Chorus, I., Bartram, J., 1999. Toxic cyanobacteria in water: a guide to their public health consequences, monitoring, and management. WHO. London: E&FN Spon. Available at <u>https://apps.who.int/iris/handle/10665/42827</u>, Accessed date: 28 July 2020.

Colas, S., Duval, C., Marie, B., 2020. Toxicity, transfer and depuration of anatoxin-a (cyanobacterial neurotoxin) in medaka fish exposed by single-dose gavage. Aquat. Toxicol. *222*, 105422. <u>https://doi.org/10.1016/j.aquatox.2020.105422</u>

Devlin, J.P., Edwards, O.E., Gorham, P.R., Hunter, N.R., Pike, R.K., Stavric, B., 1977. Anatoxin-a, a toxic alkaloid from *Anabaena flos-aquae* NRC-44h. Can. J. Chem. 55, 1367-1371. <u>https://doi.org/10.1139/v77-189</u>

Draisci, R., Ferretti, E., Palleschi, L., Marchiafava, C., 2001. Identification of anatoxins in blue-green algae food supplements using liquid chromatography-tandem mass spectrometry. Food Addit. Contam. 18 (6), 525-531. https://doi.org/10.1080/02652030118558

Echenique-Subiabre, I., Tenon, M., Humbert, J. F., Quiblier, C., (2018). Spatial and temporal variability in the development and potential toxicity of *Phormidium* biofilms in the Tarn River, France. Toxins, 10 (10), 418. https://doi.org/10.3390/toxins10100418

Edwards, C., Beattie, K.A., Scrimgeour, C.M., Codd, G.A., 1992. Identification of anatoxin-a in benthic cyanobacteria (blue-green algae) and in associated dog

Faassen, E.J., Harkema, L., Begeman, L., Lurling, M., 2012. First report of (homo)anatoxin-a and dog neurotoxicosis after ingestion of benthic cyanobacteria in The Netherlands. Toxicon 60 (3), 378-384. https://doi.org/10.1016/j.toxicon.2012.04.335

Farrer, D., Counter, M., Hillwig, R., Cude, C., 2015. Health-based cyanotoxin guideline values allow for cyanotoxin-based monitoring and efficient public health response to cyanobacterial blooms. Toxins 7, 457–77. https://doi.org/10.3390/toxins7020457

Fastner, J., Beulker, C., Geiser, B., Hoffmann, A., Kröger, R., Teske, K., Hoppe, J., Mundhenk, L., Neurath, H., Sagebiel, D., Chorus, I., 2018. Fatal neurotoxicosis in dogs associated with tychoplanktic, anatoxin-a producing *Tychonema* sp. in mesotrophic lake Tegel, Berlin. Toxins 10(2), 60. <u>https://doi.org/10.3390/toxins10020060</u>

Fawell, J.K., Mitchell, R.E., Hill, R.E., Everett, D.J., 1999. The toxicity of cyanobacterial toxins in the mouse: II Anatoxin-a. Hum Exp Toxicol. 18, 168-173. https://doi.org/10.1177/096032719901800306

Ferrão-Filho, A.D.S., Kozlowsky-Suzuki, B. (2011). Cyanotoxins: bioaccumulation and effects on aquatic animals. Mar. Drugs, 9(12), 2729-2772. https://doi.org/10.3390/md9122729 Gagnon, A., Pick, F.R., 2012. Effect of nitrogen on cellular production and release of the neurotoxin anatoxin-a in a nitrogen-fixing cyanobacterium. Front. Microbiol. 3, 211. https://doi.org/10.3389/fmicb.2012.00211

Gallon, J.R., Kittakoop, P., Brown, E.G., 1994. Biosynthesis of anatoxin-a by *Anabaena* flos-aquae: examination of primary enzymic steps. Phytochemistry 35 (5), 1195-1203. <u>https://doi.org/10.1016/S0031-9422(00)94821-0</u>

Gugger, M., Lenoir, S., Berger, C., Ledreux, A, Druart, J.-C., Humbert, J.-F., Guette, C., Bernard, C., 2005. First report in a river in France of the benthic cyanobacterium *Phormidium favosum* producing anatoxin-a associated with dog neurotoxicosis. Toxicon 45 (7), 919-928. <u>https://doi.org/10.1016/j.toxicon.2005.02.031</u>

Hamel, K., 2009. Freshwater Algae Control Program: Report to the Washington StateLake Legislature (2008-2009). Washington State Department of Ecology, Olympia,WA(PublicationNo.09-10-082).Availableat:https://app.leg.wa.gov/ReportsToTheLegislature/Home/GetPDF?fileName=0910082cd324181-a42d-441f-91ac-2ca987fe5b12.pdf, Accessed date: 28 July 2020.

Hamill, K.D., 2001. Toxicity in benthic freshwater cyanobacteria (blue-green algae): First observations in New Zealand. N.Z. J. Mar. Freshwater Res. 35 (5): 1057-1059. https://doi.org/10.3390/toxins5122504

Hardy, F.J., Johnson, A., Hamel, K., Preece, E., 2015. Cyanotoxin bioaccumulation in freshwater fish, Washington State, USA. Environ. Monit. Assess.187 (11), 667. https://doi.org/10.1007/s10661-015-4875-x Harland, F.M., Wood, S.A., Moltchanova, E., Williamson, W.M., Gaw, S., 2013. *Phormidium autumnale* growth and anatoxin-a production under iron and copper stress. Toxins 5(12), 2504-2521. <u>https://doi.org/10.3390/toxins5122504</u>

Hemscheidt, T., Rapala, J., Sivonen, K., Skulberg, O.M., 1995. Biosynthesis of anatoxin-a in *Anabaena* flos-aquae and homoanatoxin-a in *Oscillatoria formosa*. J. Chem. Soc., Chem. Comm. 13, 1361-1362. <u>https://doi.org/10.1039/C39950001361</u>

Hodoki, Y., Ohbayashi, K., Kobayashi, Y., Takasu, H., Okuda, N., Nakano, S. I., 2013. Anatoxin-a-producing *Raphidiopsis mediterranea* Skuja var. *grandis* Hill is one ecotype of non-heterocytous *Cuspidothrix issatschenkoi* (Usačev) Rajaniemi et al. in Japanese lakes. Harmful algae, 21, 44-53. <u>https://doi.org/10.1016/j.hal.2012.11.007</u>

Holland, A., Kinnear, S., 2013. Interpreting the possible ecological role (s) of cyanotoxins: compounds for competitive advantage and/or physiological aide? Mar. Drugs 11 (7), 2239-2258. <u>https://doi.org/10.3390/md11072239</u>

Huang, I.S., Zimba, P.V., 2019. Cyanobacterial bioactive metabolites—A review of their chemistry and biology. Harmful Algae 86, 139-209. https://doi.org/10.1016/j.hal.2019.05.001

Jakubowska, N., Szeląg-Wasielewska, E., 2015. Toxic picoplanktonic cyanobacteria--review. Mar. Drugs 13, 1497-1518. <u>https://doi.org/10.3390/md13031497</u>

James, K.J., Sherlock, I.R., Stack, M.A., 1997. Anatoxin-a in Irish freshwater and cyanobacteria, determined using a new fluorimetric liquid chromatographic method. Toxicon 35 (6), 963-971. <u>https://doi.org/10.1016/s0041-0101(96)00201-2</u>

James, K.J., Furey, A., Sherlock, I.R., Stack, M.A., Twohig, M., Caudwell, F.B., Skulberg, O.M., 1998. Sensitive determination of anatoxin-a, homoanatoxin-a and

their degradation products by liquid chromatography with fluorimetric detection. J. Chromatogr. A. 798 (1-2),147-157. https://doi.org/10.1016/s0021-9673(97)01207-7

James, K.J., Crowley, J., Hamilton, B., Lehane, M., Skulberg, O., Furey, A., 2005. Anatoxins and degradation products, determined using hybrid quadruple time-of-flight and quadruple ion-trap mass spectrometry: forensic investigations of cyanobacterial neurotoxin poisoning. Rapid Commun Mass Spectrom 19: 1167-1175. https://doi.org/10.1002/rcm.1894

Jarema, K.A., Poling, A., MacPhail, R.C., 2008. Effects of weekly exposure to anatoxin-a and nicotine on operant performance of rats. Neurotoxicol. Teratol. 30 (3), 220-227. https://doi.org/10.1016/j.ntt.2008.02.001.

John, N., Baker, L., Ansell, B.R.E., Newham, S., Crosbie, N.D., Jex, A.R., 2019. First report of anatoxin-a producing cyanobacteria in Australia illustrates need to regularly up-date monitoring strategies in a shifting global distribution. Sci. Rep. 9 (1), 10894. https://doi.org/10.1038/s41598-019-46945-8

Kaebernick, M., Neilan, B.A., 2001. Ecological and molecular investigations of cyanotoxin production. FEMS Microbiol. Ecol. 35 (1), 1-9. https://doi.org/10.1111/j.1574-6941.2001.tb00782.x

Kaminski, A., Bober, B., Lechowski, Z., Bialczyk J., 2013. Determination of anatoxina stability under certain abiotic factors. Harmful Algae 28, 83-87. https://doi.org/10.1016/j.hal.2013.05.014

Kiviranta, J., Sivonen, K., Lahti, K., Luukkainen, R., Niemelä, S. I., 1991. Production and biodegradation of cyanobacterial toxins-a laboratory study. Archiv für Hydrobiologie, 121(3), 281-294. Krienitz, L., Ballot, A., Kotut, K., Wiegand, C., Pütz, S., Metcalf, J.S., Codd, G., Pflugmacher, P., 2003. Contribution of hot spring cyanobacteria to the mysterious deaths of Lesser Flamingos at Lake Bogoria, Kenya. FEMS Microbiol. Ecol. 43 (2), 141-148. <u>https://doi.org/10.1111/j.1574-6941.2003.tb01053.x</u>

Kuiper-Goodman, T., Falconer, I., Fitzgerald, J., 1999. Human health aspects. In "Toxic Cyanobacteria in Water: A guide to their public health consequences, monitoring and management". WHO, ed. by Chorus I. and Bartram J., 113-153. Available at

https://www.who.int/water_sanitation_health/resourcesquality/toxcyanchap4.pdf?ua= 1, Accessed date: 28 July 2020.

Macphail, R.C., Farmer, J.D., Jarema, K.A., Chernoff, N., 2005. Nicotine effects on the activity of mice exposed prenatally to the nicotinic agonist anatoxin-a. Neurotoxicol. Teratol. 27, 593-598. <u>https://doi.org/10.1016/j.ntt.2005.05.004</u>

Martyn, J.A., Fagerlund, M.J., Eriksson, L.I., 2009. Basic principles of neuromuscular transmission. Anaesthesia 64 (S1), 1-9. <u>https://doi.org/10.1111/j.1365-</u>2044.2008.05865.x

Mazmouz, R., Chapuis-Hugon, F., Mann S., Pichon, V., Méjean, A., Ploux, O., 2010. Biosynthesis of cylindrospermopsin and 7-epicylindrospermopsin in *Oscillatoria* sp. strain PCC 6506: identification of the cyr gene cluster and toxin analysis. Appl. Environ. Microbiol. 76 (15), 4943-4949. https://doi.org/10.1128/AEM.00717-10

McAllister, T.G., Wood, S.A., Hawes, I., 2016. The rise of toxic benthic *Phormidium* proliferations: a review of their taxonomy, distribution, toxin content and factors regulating prevalence and increased severity. Harmful Algae 55, 282-294. https://doi.org/10.1016/j.hal.2016.04.002

Méjean, A., Mann, S., Maldiney, T., Vassiliadis, G., Lequin, O., Ploux, O., 2009. Evidence that biosynthesis of the neurotoxic alkaloids anatoxin-a and homoanatoxina in the cyanobacterium *Oscillatoria* PCC 6506 occurs on a modular polyketide synthase initiated by L-proline. J. Am. Chem. Soc. 131, 7512-7513. https://doi.org/10.1021/ja9024353

Méjean, A., Mazmouz, R., Mann, S., Calteau, A., Medigue, C., Ploux, O., 2010. The genome sequence of the cyanobacterium *Oscillatoria* sp. PCC 6506 reveals several gene clusters responsible for the biosynthesis of toxins and secondary metabolites. J. Bacteriol. 192 (19), 5264-5265. <u>https://doi.org/10.1128/JB.00704-10</u>

Méjean, A., Paci, G., Gautier, V., Ploux, O., 2014. Biosynthesis of anatoxin-a and analogues (anatoxins) in cyanobacteria. Toxicon 91, 15-22. https://doi.org/10.1016/j.toxicon.2014.07.016

Metcalf, J.S., Morrison, L.F., Krienitz, L., Ballot, A., Krause, E., Kotut, K., Pütz, S., Wiegand, C., Pflugmacher, S., Codd, G.A., 2006. Analysis of the cyanotoxins anatoxin-a and microcystins in Lesser Flamingo feathers. Toxicol. Environ. Chem. 88 (1), 159-167. <u>https://doi.org/10.1016/j.chemosphere.2012.09.094</u>

Metcalf, J.S., Codd, G.A., 2014. Cyanobacterial toxins (Cyanotoxins) in water. Review of current knowledge. Foundation for Water Research. 43 pp. Available at <u>http://www.fwr.org/cyanotox.pdf</u> http://www.fwr.org/cyanotox.pdf, Accessed date: 28 July 2020

Neilan, B.A., Pearson, L.A., Muenchhoff, J., Moffitt, M.C., Dittmann, E., 2013. Environmental conditions that influence toxin biosynthesis in cyanobacteria. Environ. Microbiol. 15 (5), 1239-1253. <u>https://doi.org/10.1111/j.1462-2920.2012.02729.x</u> Nowruzi, B., Blanco, S., Nejadsattari, T., 2018. Chemical and molecular evidences for the poisoning of a duck by anatoxin-a, nodularin and cryptophycin at the coast of lake Shoormast (Mazandaran province, Iran). Algologia, 28 (4), 409-427. https://doi.org/10.15407/alg28.04.409

Oberemm, A., Becker, J., Codd, G.A., Steinberg, C., 1999. Effects of cyanobacterial toxins and aqueous crude extracts of cyanobacteria on the development of fish and amphibians. Environ. Toxicol. 14 (1), 77-88. <u>https://doi.org/10.1002/(sici)1522-7278(199902)14:1<77::aid-tox11>3.0.co;2-f</u>

O'Neil, J.M., Davis, T.W., Burford, M.A., Gobler, C.J., 2012. The rise of harmful cyanobacteria blooms: The potential roles of eutrophication and climate change. Harmful Algae 14, 313–334. <u>https://doi.org/10.1016/j.hal.2011.10.027</u>

Osswald, J., Rellán, S., Gago, A., Vasconcelos, V., 2007. Toxicology and detection methods of the alkaloid neurotoxin produced by cyanobacteria, anatoxin-a. Environ. Int. 33, 1070-1089. <u>https://doi.org/10.1016/j.envint.2007.06.003</u>

Osswald, J., Rellán, S., Gago, A., Vasconcelos, V., 2008. Uptake and depuration of anatoxin-a by the mussel *Mytilus galloprovincialis* (Lamarck, 1819) under laboratory conditions. Chemosphere 72 (9), 1235-1241. https://doi.org/10.1016/j.chemosphere.2008.05.012

Osswald, J., Carvalho, A. P., Claro, J., Vasconcelos, V., 2009. Effects of cyanobacterial extracts containing anatoxin-a and of pure anatoxin-a on early developmental stages of carp. Ecotoxicol. Environ. Saf. 72 (2), 473-478. https://doi.org/10.1016/j.ecoenv.2008.05.011 Osswald, J., Azevedo, J., Vasconcelos, V., Guilhermino, L., 2011. Experimental determination of the bioconcentration factors for anatoxin-a in juvenile rainbow trout (*Oncorhynchus mykiss*). Proc. Int. Acad. Ecol. Env. Sci. 1 (2), 77-86. Available at http://www.iaees.org/publications/journals/piaees/articles/2011-1(2)/Experimental-determination-bioconcentration-factors.pdf, Accessed date: 28 July 2020

Papke, R.L., 2014. Merging old and new perspectives on nicotinic acetylcholine receptors. Biochem. Pharmacol. 89 (1), 1-11. <u>https://10.1016/j.bcp.2014.01.029</u>

Pawlik-Skowrońska, B., Toporowska, M., Rechulicz, J., 2012. Simultaneous accumulation of anatoxin-a and microcystins in three fish species indigenous to lakes affected by cyanobacterial blooms. Ocean. and Hydro. 41 (4), 53-65. https://doi.org/10.2478/s13545-012-0039-6

Pawlik-Skowrońska, B., Kalinowskaa, R., Skowrońskiac, T., 2013. Cyanotoxin diversity and food web bioaccumulation in a reservoir with decreasing phosphorus concentrations and perennial cyanobacterial blooms. Harmful Algae 28, 118-125. https://doi.org/10.1016/j.hal.2013.06.002

Peary, J.A., Gorham, P.R., 1966. Influence of light and temperature on growth and toxin production by *Anabaena flos-aquae*. J Phycol 2 (Suppl.), 3

Pedersen, J.E., Bergqvist, C.A., Larhammar, D., 2019. Evolution of vertebrate nicotinic acetylcholine receptors. BMC Evol. Biol. 19 (1), 38. https://doi.org/10.1186/s12862-018-1341-8

Puschner, B., Hoff, B., Tor, E.R., 2008. Diagnosis of anatoxin-a poisoning in dogs from North America. J. Vet. Diagn. Invest. 20 (1), 89-92. https://doi.org/10.1177/104063870802000119 Puschner, B., Pratt C., Tor E.R., 2010. Treatment and diagnosis of a dog with fulminant neurological deterioration due to anatoxin-a intoxication. J. Vet. Emerg. Crit. Care 20, 518-522. <u>https://doi.org/10.1111/j.1476-4431.2010.00578.x</u>

Pybus, M.J., Hobson, D.P., Onderka, D.K., 1986. Mass mortality of bats due to probable blue-green algal toxicity. J. Wildl. Dis. 22 (3), 449-450. https://doi.org/10.7589/0090-3558-22.3.449

Quiblier, C., Wood, S., Echenique-Subiabre, I., Heath, M., Villeneuve, A., Humbert, J.-F., 2013. A review of current knowledge on toxic benthic freshwater cyanobacteriaecology, toxin production and risk management. Water Res. 47 (15), 5464-5479. <u>https://doi.org/10.1016/j.watres.2013.06.042</u>

Rapala, J., Sivonen, K., Luukkainen, R., Niemelä S., 1993. Anatoxin-a concentration in *Anabaena* and *Aphanizomenon* under different environmental conditions and comparison of growth by toxic and non-toxic *Anabaena*-strains - a laboratory study. J. Appl. Phycol. 5, 581-591. <u>https://doi.org/10.1007/BF02184637</u>

Rapala, J., Lahti, K., Sivonen, K., Niemelä, S.I., 1994. Biodegradability and adsorption on lake sediments of cyanobacterial hepatotoxins and anatoxin-a. Lett. Appl. Microbiol. 19 (6), 423-428. <u>https://doi.org/10.1111/j.1472-765x.1994.tb00972.x</u>

Rapala, J., Sivonen, K., 1998. Assessment of environmental conditions that favor hepatotoxic and neurotoxic *Anabaena* spp. strains cultured under light limitation at different temperatures. Microb. Ecol. 36 (2), 181-192. <u>https://doi.org/10.1007/s002489900105</u>

Rogers, E.H., Hunter, E.S. 3rd, Moser, V.C., Phillips, P.M., Herkovits, J., Muñoz, L., Hall, L.L., Chernoff, N., 2005. Potential developmental toxicity of anatoxin-a, a cyanobacterial toxin. J. Appl. Toxicol. 25 (6), 527-534. https://doi.org/10.1002/jat.1091

Rowell, R.R., Wonnacott, S., 1990. Evidence for functional activity of up-regulated nicotine binding sites in rat striatal synaptosomes. J. Neurochem. 55 (6), 2105-2110. https://doi.org/10.1111/j.1471-4159.1990.tb05802.x

Selwood, A.I., Holland, P.T., Wood, S.A., Smith, K.F., Mcnabb, P.S., 2007. Production of anatoxin-a and a novel biosynthetic precursor by the cyanobacterium *Aphanizomenon issatschenkoi*. Environ. Sci. Technol. 41, 506–510. https://doi.org.10.1021/es0619830

Shams, S., Capelli, C., Cerasino, L., Ballot, A., Dietrich, D. R., Sivonen, K., Salmaso, N., 2015. Anatoxin-a producing *Tychonema* (Cyanobacteria) in European waterbodies. Water Res. 69, 68-79. <u>https://doi.org/10.1016/j.watres.2014.11.006</u>

Sivonen, K., Himberg, K., Luukkainen, R., Niemelä, S. I., Poon, G. K., Codd, G. A., 1989. Preliminary characterization of neurotoxic cyanobacteria blooms and strains from Finland. Environ. Toxicol. Water Qual. 4 (3), 339-352. https://doi.org/10.1002/tox.2540040310

Sivonen, K., Niemelä, S. I., Niemi, R. M., Lepistö, L., Luoma, T. H., Räsänen, L. A., 1990. Toxic cyanobacteria (blue-green algae) in Finnish fresh and coastal waters. Hydrobiologia 190, 267-275. <u>https://doi.org/10.1007/BF00008195</u>

Sivonen, K., Jones, G., 1999. Cyanobacterial toxins. In "Toxic Cyanobacteria in Water: A guide to their public health consequences, monitoring and management". WHO, ed. by Chorus I. and Bartram J., 43-112. Available at https://www.who.int/water_sanitation_health/resourcesquality/toxcyanchap3.pdf?ua= 1, Accessed date: 28 July 2020

 Skulberg, O.M., Skulberg, R., Carmichael, W.W., Andersen, R.A., Matsunaga, S., Moore, R.E., 1992. Investigations of a neurotoxic oscillatorialean strain (Cyanophyceae) and its toxin. Isolation and characterization of homoanatoxin-a. Environ. Toxicol. Chem. 11 (3), 321-329. <u>https://doi.org/10.1002/etc.5620110306</u>

Smith, R.A., Lewis, D., 1987. A rapid analysis of water for anatoxin a, the unstable toxic alkaloid from *Anabaena* flos-aquae, the stable non-toxic alkaloids left after bioreduction and a related amine which may be nature's precursor to anatoxin a. Vet. Hum. Toxicol. 29 (2), 153-154. PMID: 3107204.

Smith, C., Sutton, A., 1993. The persistence of anatoxin-a in reservoir water. Foundation for Water Research report no. FR0427. available at <u>http://www.fwr.org/waterq/fr0427.htm</u>, Accessed date: 28 July 2020

Spivak, C.E., Witkop, B., Albuquerque, E.X., 1980. Anatoxin-a: a novel, potent agonist at the nicotinic receptor. Mol. Pharmacol. 18 (3): 384-394. https://doi.org/10.1016/0922-4106(95)90153-1

Stepanova, N., Nikitin, O., Latypova, V., Kondratyeva, T., 2018. Cyanotoxins as a possible cause of fish and waterfowl death in the Kazanka river (Russia). In Proceedings of the 18th International Multidisciplinary Scientific GeoConference SGEM, Albena, Bulgaria, 2-8 July 2018; Vol. 18: 229-236.

Stevens, D.K., Krieger, R.I., 1988. Analysis of anatoxin-a by GC/ECD. J. Anal. Toxicol. 12 (3), 126-131. <u>https://doi.org/10.1093/jat/12.3.126</u>

Stevens, D.K., Krieger, R.I., 1991. Stability studies on the cyanobacterial nicotinic alkaloid snatoxin-A. Toxicon 29, 167–179. <u>https://doi.org/10.1016/0041-0101(91)90101-v</u>

Stewart, I., Webb, P.M., Schluter, P.J., Shaw, G.R., 2006. Recreational and occupational field exposure to freshwater cyanobacteria--a review of anecdotal and case reports, epidemiological studies and the challenges for epidemiologic assessment. Environ. Health. 5, 6. <u>https://doi.org/10.1186/1476-069x-5-6</u>

Stolerman, I.P., Albuquerque, E.X., Garcha, H.S., 1992. Behavioural effects of anatoxin, a potent nicotinic agonist, in rats. Neuropharmacology 31 (3), 311-314. <u>https://doi.org/10.1016/0028-3908(92)90182-0</u>

Svirčev, Z., Baltić, V., Gantar, M., Juković, M., Stojanović, D., Baltić, M., 2010. Molecular aspects of microcystin-induced hepatotoxicity and hepatocarcinogenesis. J. Environ. Sci. Health C. Environ. Carcinog. Ecotoxicol. Rev. 28 (1), 39-59. https://doi.org/10.1080/10590500903585382

Svirčev, Z., Lalić, D., Bojadžija Savić, G., Tokodi, N., Backović, D.D., Chen, L., Meriluoto, J., Codd, G.A., 2019. Global geographical and historical overview of cyanotoxin distribution and cyanobacterial poisonings. Arch Toxicol. 93 (9), 2429-2481. <u>https://doi-org.10.1007/s00204-019-02524-4</u>

Swanson, K.L., Allen, C.N., Aronstam, R.S., Rapoport, H., Albuquerque, E.X., 1986. Molecular mechanisms of the potent and stereospecific nicotinic receptor agonist (+)anatoxin-a. Mol. Pharmacol. 29 (3), 250-257. PMID: 2419745

Thomas, P., Stephens, M., Wilkie, G., Amar, M., Lunt, G.G., Whiting, P., Gallagher T., Pereira, E., Alkondon, M., Albuquerque, E.X., Wonnacott S., 1993. (+)-Anatoxin-a is a potent agonist at neuronal nicotinic acetylcholine receptors. J. Neurochem. 60 (6), 2308-2311. https://doi.org/10.1111/j.1471-4159.1993.tb03519.x

Toporowska, M, Pawlik-Skowrońska, B, Kalinowska, R., 2014. Accumulation and effects of cyanobacterial microcystins and anatoxin-a on benthic larvae of

Trainer, V.L., Hardy, F.J., 2015. Integrative monitoring of marine and freshwater harmful algae in Washington State for public health protection. Toxins 7(4), 1206-1234. <u>https://doi.org/10.3390/toxins7041206</u>

Valentine, W.M., Schaeffer, D.J., Beasley, V.R., 1991. Electromyographic assessment of the neuromuscular blockade produced in vivo by anatoxin-a in the rat. Toxicon 29 (3), 347-357. <u>https://doi.org/10.1016/0041-0101(91)90288-3</u>

Vasconcelos, V.M., 1995. Uptake and depuration of the heptapeptide toxin microcystin-LR in *Mytilus galloprovincialis*. Aqua. Toxicol. 32 (2-3), 227-237. https://doi.org/10.1016/0166-445X(94)00085-5

Vehovszky, Á., Ács, A., Kovacs, A. W., Szabó, H., Gyori, J., Farkas, A., 2009. Isolated strains of *Cylindrospermopsis raciborskii* from Lake Balaton (Hungary) produce anatoxin-a like neurotoxins. Comp. Biochem. Physiology Part. Mol. Integr. Physiology 153, S88. <u>https://doi.org/10.1016/j.cbpa.2009.04.083</u>

Viaggiu, E., Melchiorre, S., Volpi, F., Di Corcia, A., Mancini, R., Garibaldi, L., Cricchigno, G., Bruno, M., 2004. Anatoxin- a toxin in the cyanobacterium *Planktothrix rubescens* from a fishing pond in northern Italy. Environmental Toxicology: An International Journal, 19 (3), 191-197. <u>https://doi-org.buadistant.univ-angers.fr/10.1002/tox.20011</u>.

Weirich, C.A., Miller, T.R., 2014. Freshwater harmful algal blooms: toxins and children's health. Curr. Probl. Pediatr. Adolesc. Health Care 44 (1), 2-24. https://doi.org/10.1016/j.cppeds.2013.10.007. Wonnacott, S., Jackman, S., Swanson, K.L., Rapoport, H., Albuquerque, E.X., 1991. Nicotinic pharmacology of anatoxin analogs. II. Side chain structure-activity relationships at neuronal nicotinic ligand binding sites. J. Pharmacol. Exp. Ther. 259 (1), 387-391. PMID: 1920126.

Wonnacott, S., Swanson, K.L., Albuquerque, E.X., Huby, N.J.S., Thompson, P., Gallagher, T., 1992. Homoanatoxin: a potent analogue of anatoxin-a. Biochem. Pharmacol. 43 (3), 419-423. <u>https://doi.org/10.1016/0006-2952(92)90558-z</u>

Wonnacott, S., Gallagher, T., 2006. The chemistry and pharmacology of anatoxin-a and related homotropanes with respect to nicotinic acetylcholine receptors. Mar. Drugs 4 (3), 228-254. <u>https://doi.org/10.3390/md403228</u>

Wood, S.A., Selwood, A.I., Rueckert, A., Holland, P.T., Milne, J.R., Smith, K.F., Smits, B., Watts, L.F., Cary, C.S., 2007. First report of homoanatoxin-a and associated dog neurotoxicosis in New Zealand. Toxicon 50 (2), 292-301. https://doi.org/10.1016/j.toxicon.2007.03.025

Wood, R., 2016. Acute animal and human poisonings from cyanotoxin exposure - A review of the literature. Environ. Int., 91, 276-282. https://doi.org/10.1016/j.envint.2016.02.026

Wood, S.A., Puddick, J., Fleming, R., Heussner, A.H., 2016. Detection of anatoxinproducing *Phormidium* in a New Zealand farm pond and an associated dog death. N.-Z. J. Bot. 55, 1-11. <u>https://doi.org/10.1080/0028825X.2016.1231122</u>

Wood, S. A., Puddick, J., 2017. The abundance of toxic genotypes is a key contributor to anatoxin variability in *Phormidium*-dominated benthic mats. Marine Drugs, 15 (10), 307. <u>https://doi.org/10.3390/md15100307</u>.

Wood, S.A., Biessy, L., Puddick, J., 2018. Anatoxins are consistently released into the water of streams with *Microcoleus autumnalis*-dominated (cyanobacteria) proliferations. Harmful algae 80, 88-95. <u>https://doi.org/10.1016/j.hal.2018.10.001</u>

Wood, S.A., Kelly L.T., Bouma-Gregson K., Humbert J.-F., Laughinghouse, IV H.D., Lazorchak, J., McAllister, T.G., McQueen, A., Pokrzywinski, K., Puddick, J., Quiblier, C., Reitz, L.A., Ryan, K.G., Vadeboncoeur, Y., Zastepa, A., Davis, T.W., 2020. Toxic benthic freshwater cyanobacterial proliferations: Challenges and solutions for enhancing knowledge and improving monitoring and mitigation. Freshwater Biol. (under the press). https://doi.org/10.1111/fwb.13532

Yavasoglu, A., Karaaslan, M.A., Uyanikgil, Y., Sayim, F., Ates, U., Yavasoglu, NU., 2008. Toxic effects of anatoxin-a on testes and sperm counts of male mice. Exp. Toxicol. Pathol. 60, 391-396. https://doi.org/10.1016/j.etp.2008.04.001

Zoli, M., Pucci, S., Vilella, A., Gotti, C., 2018. Neuronal and extraneuronal nicotinic acetylcholine receptors. Curr. Neuropharmacol. 16 (4), 338-349. https://doi.org/10.2174/1570159X15666170912110450

Figure captions

Figure 1 : ATX-a producing cyanobacteria. (A) *Phormidium* sp bloom in the Cher river, France (© Jean-François Humbert). (B) *Phormidium* sp biofilm (© Jean-François Humbert). (C) *Phormidium favosum* from the Tarn river, France, observed with scanning electron microscopy. Note the diatoms at the centre (© Isidora Echenique-Subiabre)

Figure 2: Structures of ATX-a and related molecules.

Figure 3: Cellular and molecular mode of action of ATX-a. (A) ATX-a binds and opens nAChRs, but unlike their natural ligands – ACh – the toxin does not get hydrolyzed by resident acetylcholinesterases. (B) ATX-a targets cholinergic synapses where nAChRs are expressed, in both CNS and PNS. (C) Through its rapid absorption, ATX-a exerts deleterious effects on the brain, the cardiovascular system, the respiratory tract and the skeletal muscles.

Table 1: Main episodes of ATX-a intoxication in the wild. DAD, Diode Array Detector, DW Dry Weight; ELISA, Enzyme Linked ImmunoSorbent Assay; ESI, Electrospray Ionisation; FW, Fresh Weight; GC/ECD, GC Gas Chromatography with Electron Capture Detector; MS, Mass Spectrometry; HPLC, High-Performance Liquid Chromatography; MALDI-TOF, Matrix Assisted Laser Desorption Ionization - Time Of Flight; NI, not indicated; PDA, Photometric Diode Array.

Year, Country and Location	Intoxication and mortalities	Method of detection	Matrix	Congener and maximum concentration	Cyanobacteria species	Reference
	episode				(benthic/planktonic)	
1981, USA, Black	11 cows died	GC/ECD	Water sample	2.5 g.kg ⁻¹ (DW) of ATX-a	Anabaena flos-aquae	Stevens and Krieger,
Lake	2 dogs died				(planktonic)	1988
	1000 bats died, 24	GC-MS	Green slime from	ATX-a	Anabaena flos-aquae	Pybus et al., 1986
1985, Canada,	mallards and		the carcasses		(planktonic)	
Steele Lake	American wigeons					
	died					
1985-1986,	5 cows died	HPLC, GC-	Water sample	3.7 g.kg ⁻¹ (DW) of ATX-a	Anabaena spp.	Sivonen et al., 1990
Finland, Lakes		MS			(Planktonic)	
Sääskjärvi and						
Säyhteenjärvi						
1987, Canada,	16 cows died	GC-MS	NI	ATX-a	Anabaerna flos-aquae	Smith and Lewis, 1987
Alberta					(Planktonic)	
1990-1991,	4 dogs died and	HPLC-DAD,	Water sample	ATX-a	Oscillatoria spp.	Edwards et al., 1992
Scotland, Loch	one intoxicated	GC-MS	Dog stomach	ATX-a	(benthic)	
Insh			content			
1992-1993-1994,	6 dogs died	HPLC-FL,	Water sample	444 µg.L⁻¹ (FW) of ATX-a	Oscillatoria spp.	James et al., 1997
Ireland, Caragh		GC-MS			(benthic)	
Lake						
1998-1999-2000,	12 dogs died	HPLC	Cyanobacterial	ATX-a	Oscillatoria spp.	Hamill, 2001
New Zealand,			mat sample		(benthic)	
Waikanae and						
Mataura Rivee						
2001, Kenya, Lake	Mortalities of	HPLC-PDA,	Cyanobacterial	18 mg.kg ⁻¹ (DW) of ATX-a	Phormidium	Krienitz et al., 2003
Bogoria	Lesser Flamingos	MALDI-TOF	mat sample		terebriformis (benthic),	

			Flamingo	4.349 mg.kg ⁻¹ (FW) of ATX-a	Oscillatoria willei	
			stomach content		(benthic), <i>Spirulina</i>	
			Flamingo intestine	0.762 mg.kg ⁻¹ (FW) of ATX-a	subsalsa (benthic) and	
			Flamingo pellet	0.245 mg.kg ⁻¹ (FW) of ATX-a	Synechococcus	
			0 1		<i>bigranulatus</i> (benthic)	
2001-2002 Kenya,	Mortalities of	HPLC-PDA,	Water sample	223 mg.kg ⁻¹ (DW) of ATX-a	Arthrospira fusiformis	
Lakes Bogoria and	Lesser Flamingos	MALDI-TOF			(planktonic)	Ballot et al., 2004
Nakaru						
			Cyanobacterial	8 g.kg ⁻¹ (DW) of ATX-a		
2003, France, La	2 dogs died	ESI-MS/MS	mat sample		Phormidium favosum	Gugger et al., 2005
Loue River			Dog liver extracts	0.6 g.kg ⁻¹ (DW) of ATX-a	(benthic)	
2005, Kenya,	Mortalities of	HPLC-PDA,	Feathers	800 µg.kg⁻¹ (DW) of ATX-a	n.r.	
Lakes Bogoria and	Lesser Flamingos	MALDI-TOF				Metcalf et al., 2006
Nakuru						
2005, New	5 dogs died	LC-MS		27 µg.kg⁻¹ (FW) of ATX-a	Phormidium	
Zealand, Hutt			Cyanobacterial	4400 µg.kg ⁻¹ (FW) of HTX-a	autumnale	Wood et al., 2007
River			mat sample		(benthic)	
2006, Canada,	3 dogs died	LC-	Dog stomach	10 µg.kg-1 (?) of ATX-a	Planktothrix spp.	Puschner et al., 2008
pond in Ontario		MS/MS/MS	content		(benthic)	
2011,			Cyanobacterial	272 µg.g⁻¹ (DW) of ATX-a		
Netherlands, Lake	3 dogs died and 1	LC-MS/MS	mat sample		Phormidium sp.	Faassen et al., 2012
IJmeer	dog intoxicated		Dog stomach	9.5 µg.g⁻¹ (DW) of ATX-a	(benthic)	
			content			
0044 N				25 µg.L ⁻⁺ (FW) of ATX-a		
2014, New			Environmental	2118 µg.L ⁻¹ (FW) of dhATX-a	Dh a ma i dia ma	
Zeleand, pond	1 dog diod		sample		Phormialum	Weed at al. 2010
near Kaikoura	i dog died	LC-1015/1015	Dog stomacn	36 µg.kg ⁻ (FVV) of ATX-a	autumnale (honthio)	wood et al., 2016
			content		(Dentinc)	
					Nastas an (hanthis)	
2015 Iron	Ducks diad		Duck stomach	ATA-a	Anabaona sp	Nowruzi et al. 2018
2015, Itali, Shoormost Lako	Ducks uleu	LC-IVIS	Duck Stornach		(planktonic) and	Nowiuzi et al., 2018
Shoomast Lake			Content		Nodularia sp	
					(planktonic)	
2016 Russia	1000 gibel carps	FLISA kit	Water sample	0.294 ug L ⁻¹ (FW) of ATX-a	Planktolvnobva	Stepanova et al 2018
Kazanka River	died		Fish mussles	$E_{211} = \frac{1}{100} E_{10} = \frac$	<i>limnetica</i> (planktonic)	
				5.211 µg.kg (FVV) 01 ATX-a	and	
			Duck muscles	1.559 µg.kg⁻¹ (FW) of ATX-a		

	10 ducks died		Duck Liver	5.193 µg.kg⁻¹ (FW) of ATX-a	Aphanizomenon flos- aquae (planktonic)	
2017, Germany, Lake Tegel	3 dogs dieds	LC-MS/MS	Water sample	1 μg.L ⁻¹ (FW) of ATX-a	<i>Tychonema</i> sp. (planktonic)	Fastner et al., 2018
			Sand sample	1 μg.L ⁻¹ (FW) of ATX-a		
			Cyanobacterial mat sample	1870 μg.L ⁻¹ (FW) of ATX-a		
			Dog stomach content	8870 µg.L ⁻¹ (FW) of ATX-a		
2017, France, La Loire and Le Cher rivers	12 dogs died			ATX-a	Phormidium sp. (benthic)	Boisseleau et al., 2018

Table 2: Acute in vivo toxicity of ATX-a in vertebrates

Toxin used	Route of	Species	LD ₅₀ (mg.kg ⁻¹)	LD ₉₀ (mg.kg ⁻¹)	References
or CB extract	administration				
(+)-ATX-a		mouse	16.2		Stevens and Krieger, 1991
(+)-ATX-a	Oral	medaka fish	5.75		Colas et al., 2020
(±)-ATX-a		medaka fish	11.5		Colas et al., 2020
(+)-ATX-a	IP	mouse	0.26		Stevens and Krieger, 1991
(+)-ATX-a		mouse	0.39		Valentine et al., 1991
(±)-ATX-a		mouse	0.91		Valentine et al., 1991
(+)-ATX-a	IV	rat	0.08		Adeyemo and Sirén, 1992
(±)-ATX-a		rat	0.4		Adeyemo and Sirén, 1992
		mouse		1800	
Anabena floS-		rat		1500	
aquae NRC-44-1	Oral	pheasant		850	Carmichael and Biggs, 1978

duck	350	
gold fish	120	







С

Declaration of interests

 \boxtimes The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

□The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: